

On the number of A -transversals in hypergraphs

János Barat*

Alfréd Rényi Institute of Mathematics, and
University of Pannonia, Department of Mathematics
barat@renyi.hu

Dániel Gerbner†

Alfréd Rényi Institute of Mathematics
gerbner@renyi.hu

Anastasia Halfpap

University of Montana
anastasia.halfpap@umontana.edu

Abstract

A set S of vertices in a hypergraph is *strongly independent* if every hyperedge shares at most one vertex with S . We prove a sharp result for the number of maximal strongly independent sets in a 3-uniform hypergraph analogous to the Moon-Moser theorem.

Given an r -uniform hypergraph \mathcal{H} and a non-empty set A of non-negative integers, we say that a set S is an A -*transversal* of \mathcal{H} if for any hyperedge H of \mathcal{H} , we have $|H \cap S| \in A$. Independent sets are $\{0, 1, \dots, r-1\}$ -transversals, while strongly independent sets are $\{0, 1\}$ -transversals. Note that for some sets A , there may exist hypergraphs without any A -transversals. We study the maximum number of A -transversals for every A , but we focus on the more natural sets, e.g., $A = \{a\}$, $A = \{0, 1, \dots, a\}$ or A being the set of odd or the set of even numbers.

*Research supported by ERC Advanced Grant "GeoScape" and the National Research, Development and Innovation Office, grant K-131529.

†Research supported by the National Research, Development and Innovation Office - NKFIH under the grants SNN 129364, FK 132060, and KKP-133819.

1 Introduction

We use standard notations of (hyper)graph theory. In particular, the degree of a vertex v is the number of (hyper)edges containing v . Recall that a vertex set in a (hyper)graph is *independent* if it contains no (hyper)edge. An independent set is *maximal* if it is not a proper subset of a larger independent set. Let $MIS(G)$ denote the number of maximal independent sets in a graph G . Miller and Muller [7] and independently Moon and Moser [8] showed that for all n -vertex graphs G , $MIS(G) \leq 3^{n/3}$, which is sharp as given by the vertex-disjoint union of triangles. More precisely, let $MIS(n)$ denote the largest $MIS(G)$ where G has n vertices. Then the following holds.

Theorem 1.1.

$$MIS(n) = \begin{cases} 3^{n/3} & \text{if } n \equiv 0 \pmod{3}, \\ 4 \times 3^{\lfloor n/3 \rfloor - 1} & \text{if } n \equiv 1 \pmod{3}, \\ 2 \times 3^{\lfloor n/3 \rfloor} & \text{if } n \equiv 2 \pmod{3}. \end{cases}$$

There are several natural generalizations of this problem to hypergraphs, especially 3-uniform graphs. Cory Palmer [11] explicitly asked the following question and inspired our research.

Problem Determine the maximum number of maximal independent sets in an n -vertex 3-uniform hypergraph.

A generalization of Theorem 1.1 to hypergraphs was studied by Tomescu [12] and by Lonc and Truszczyński [6]. They showed in 3-uniform n -vertex hypergraphs the largest number of maximal independent sets is between roughly 1.5849^n and 1.6702^n .

As it is often the case with graph theoretical notions, independent sets have multiple generalizations to hypergraphs. A set S of vertices in a hypergraph is *strongly independent*, if every hyperedge shares at most one vertex with S . We denote by $MSIS(\mathcal{H})$ the number of maximal strongly independent sets in a hypergraph \mathcal{H} . We show the following analogue of Theorem 1.1 for 3-graphs.

Theorem 1.2. *If \mathcal{H} is a 3-uniform hypergraph on n vertices, then $MSIS(\mathcal{H}) \leq g(n)$, where*

$$g(n) = \begin{cases} 3^{n/3} & \text{if } n \equiv 0 \pmod{3}, \\ 4 \times 3^{\frac{n-4}{3}} & \text{if } n \equiv 1 \pmod{3}, \\ 16 \times 3^{\frac{n-8}{3}} & \text{if } n \equiv 2 \pmod{3}. \end{cases}$$

Moreover, for all $n \geq 6$, there is an n -vertex 3-uniform hypergraph \mathcal{H} with $MSIS(\mathcal{H}) = g(n)$.

One can consider intermediate independence notions, when a set shares a bounded number of vertices with any hyperedge. Instead of that, we consider the following generalization. Given an r -uniform hypergraph \mathcal{H} and a non-empty set A of non-negative integers, we say that a set S is an A -*transversal* of \mathcal{H} if for any hyperedge H of \mathcal{H} , we have $|H \cap S| \in A$. Independent sets are $\{0, 1, \dots, r-1\}$ -transversals, while strongly independent sets are $\{0, 1\}$ -transversals. Note that for some sets A , there may exist hypergraphs without

any A -transversals. We study the maximum number of A -transversals for every A , but we focus on the more natural sets, e.g. $A = \{a\}$, $A = \{0, 1, \dots, a\}$ or A being the set of odd or the set of even numbers.

Observe that the point of studying maximal independent sets, rather than arbitrary independent sets, is that in the empty graph or empty hypergraph all the 2^n subsets are independent. We have the same problem for every A . However, it turns out, in some cases a more natural way to avoid this degenerate example is assuming that each vertex is contained in at least one edge/hyperedge, i.e. there are no isolated vertices. Note that this is only possible if $n \geq r$; we omit this extra assumption later. Galvin [5] studied the maximum number of independent sets in graphs with minimum degree at least δ and showed that for sufficiently large n , $K_{\delta, n-\delta}$ contains the most independent sets. The threshold on n was shown to be 2δ in [3].

Let us denote by $g_A^{(r)}(n)$ the largest number of A -transversals in an n -vertex r -uniform hypergraph without isolated vertices. We denote by $h_A^{(r)}(n)$ the largest number of maximal A -transversals in an n -vertex r -uniform hypergraph. We start with completely resolving the graph case.

Theorem 1.3. (i) $g_{\{0\}}^{(2)}(n) = h_{\{0\}}^{(2)}(n) = 1$.

(ii) $g_{\{1\}}^{(2)}(n) = h_{\{1\}}^{(2)}(n) = 2^{\lfloor n/2 \rfloor}$.

(iii) $g_{\{2\}}^{(2)}(n) = h_{\{2\}}^{(2)}(n) = 1$.

(iv)

$$h_{\{0,1\}}^{(2)}(n) = \begin{cases} 3^{n/3} & \text{if } n \equiv 0 \pmod{3}, \\ 4 \times 3^{\lfloor n/3 \rfloor - 1} & \text{if } n \equiv 1 \pmod{3} \\ 2 \times 3^{\lfloor n/3 \rfloor} & \text{if } n \equiv 2 \pmod{3}. \end{cases}$$

(v) $g_{\{0,1\}}^{(2)}(n) = g_{\{1,2\}}^{(2)}(n) = 1 + 2^{n-1}$.

(vi) $h_{\{0,2\}}^{(2)}(n) = h_{\{0,1,2\}}^{(2)}(n) = 1$.

(vii) $g_{\{0,1,2\}}^{(2)}(n) = 2^n$.

(viii) $g_{\{0,2\}}^{(2)}(n) = 2^{\lfloor n/2 \rfloor}$.

We remark that (iv) of the above theorem is Theorem 1.1, and (v) is the only other non-trivial statement. The above mentioned theorem of Galvin [5] implies (v) for $n \geq 14$, and the stronger result from [3] implies (v) for $n \geq 2$. We will present a simple proof.

Now we list our general results. We collect the simple observations in the following proposition.

Proposition 1.4. (i) $g_{\{0\}}^{(r)}(n) = h_{\{0\}}^{(r)}(n) = 1$ for any r .

(ii) If $r \in A$, then $h_A^{(r)}(n) = 1$ for any r .

(iii) $g_{\{r\}}^{(r)}(n) = 1$.

(iv) $g_{\{0,1,\dots,r\}}^{(r)}(n) = 2^n$.

(v) $g_{\{0,r\}}^{(r)}(n) = 2^{\lfloor n/r \rfloor}$.

- (vi) If $A \subset \{0, \dots, r\}$ and $B = \{r - a : a \in A\}$, then $g_A^{(r)}(n) = g_B^{(r)}(n)$.
(vii) $g_{\{a\}}^{(r)}(n) = h_{\{a\}}^{(r)}(n)$ for every $n \geq r$.

Finally, we list our more substantial statements for general r . Among other results, we determine the order of magnitude of the function $g_A^{(r)}(n)$ in every instance.

Theorem 1.5. (i) Let $A(i)$ denote the non-negative integers of the form $a - i$ with $a \in A$, i.e., $A(i) = \{a - i : a \in A\} \cap \mathbb{N}$. Let $f(q, i, A, r) = \sum_{b \in A(i)} \binom{q}{b}$ and consider the largest $f(q, i, A, r)^{1/q}$ with $q \leq r$ and $0 \leq i \leq r - q$. If the largest value is obtained at $p = q$, then $g_A^{(r)}(n) = \Theta(f(p, i, A, r)^{n/p})$.

(ii) Let A be the set of even integers and B the set of odd integers. Then $g_A^{(r)}(n) = g_B^{(r)}(n) = 2^{\lfloor \frac{r-1}{r} n \rfloor}$.

(iii) Let $a < k < r$. Then $g_{\{0,1,\dots,a\}}^{(k)}(n) \geq g_{\{0,1,\dots,a\}}^{(r)}(n) \geq g_{\{0,1,\dots,a\}}^{(k)}(n - r + k)$ and $h_{\{0,1,\dots,a\}}^{(k)}(n) \geq h_{\{0,1,\dots,a\}}^{(r)}(n) \geq h_{\{0,1,\dots,a\}}^{(k)}(n - r + k)$.

Using the Θ notation, we emphasize that the function may depend on A and r but not on n .

2 Proofs

Let us start with the proof of Theorem 1.2. We adapt a simple inductive proof of Theorem 1.1 due to Chang and Jou [1] and independently Wood [13].

Proof of Theorem 1.2. We can quickly verify that the theorem holds for 3-uniform hypergraphs on 0, 1, or 2 vertices. Now, let $n \geq 3$, and assume that the theorem holds for all positive integers less than n . Let \mathcal{H} be a 3-uniform hypergraph on n vertices. Let $N[v]$ denote the closed neighborhood of v , i.e., the set of vertices u such that either $u = v$ or there is a hyperedge containing both v and u . Let us choose $x \in V(\mathcal{H})$ with $|N[x]|$ minimal, and define $d = |N[x]|$. Observe, any maximal strongly independent set X of \mathcal{H} intersects $N[x]$ at least once, since otherwise $X \cup \{x\}$ remains strongly independent and properly contains X . Also note that if $v \in N[x] \cap X$, then $X - \{v\}$ is a maximal strongly independent set in the subhypergraph $\mathcal{H} - N[v]$ of \mathcal{H} which is induced on $V(\mathcal{H}) \setminus N[v]$. These observations together imply that

$$\text{MSIS}(\mathcal{H}) \leq \sum_{v \in N[x]} \text{MSIS}(\mathcal{H} - N[v]),$$

since a maximal strongly independent set X of \mathcal{H} is counted $|X \cap N[x]|$ times by the right hand sum.

Now, for any $v \in V(\mathcal{H})$, we have $\text{MSIS}(\mathcal{H} - N[v]) \leq g(n - |N[v]|) \leq g(n - d)$, since g is clearly a non-decreasing function and d is the minimum size of a closed neighborhood in \mathcal{H} . So we have

$$\text{MSIS}(\mathcal{H}) \leq d \cdot g(n - d).$$

Now, we only need to show $d \cdot g(n-d) \leq g(n)$ regardless of the value of d . This is clearly true if $d = 1$; note also that, since d represents the size of a closed neighborhood and \mathcal{H} is 3-uniform, we cannot have $d = 0$ or $d = 2$.

When $d = 3$, note that we have $3g(n-3) = g(n)$ regardless of the value of n , so indeed, $3 \cdot g(n-3) \leq g(n)$.

When $d = 4$, we have

$$4g(n-4) = \begin{cases} \frac{64}{81} \times 3^{n/3} & \text{if } n \equiv 0 \pmod{3}, \\ 4 \times 3^{\frac{n-4}{3}} & \text{if } n \equiv 1 \pmod{3}, \\ 16 \times 3^{\frac{n-8}{3}} & \text{if } n \equiv 2 \pmod{3}. \end{cases}$$

We observe that in each case, $4g(n-4) \leq g(n)$.

When $d = 5$, we have

$$5g(n-5) = \begin{cases} \frac{20}{27} \times 3^{n/3} & \text{if } n \equiv 0 \pmod{3}, \\ \frac{80}{27} \times 3^{\frac{n-4}{3}} & \text{if } n \equiv 1 \pmod{3}, \\ 15 \times 3^{\frac{n-8}{3}} & \text{if } n \equiv 2 \pmod{3}. \end{cases}$$

Again, we observe in each case that $5g(n-5) \leq g(n)$.

Observe also that for any $d \geq 5$, we have

$$(d-3)g(n-d+3) = 3(d-3)g(n-d) > d \cdot g(n-d).$$

Thus, if $d > 5$, we may repeatedly apply the above inequality to show that

$$dg(n-d) \leq \begin{cases} 3g(n-3) & \text{if } d \equiv 0 \pmod{3}, \\ 4g(n-4) & \text{if } d \equiv 1 \pmod{3}, \\ 5g(n-5) & \text{if } d \equiv 2 \pmod{3}. \end{cases}$$

We conclude that for any 3-graph \mathcal{H} on n -vertices, we have $MSIS(\mathcal{H}) \leq g(n)$. It remains to describe n -vertex constructions achieving $g(n)$ maximal strongly independent sets. These are as follows.

When $n \equiv 0 \pmod{3}$, we take $\frac{n}{3}$ independent 3-edges. When $n \equiv 1 \pmod{3}$ (and $n \geq 4$), we take $\frac{n-4}{3}$ independent 3-edges and a copy of K_4^3 . When $n \equiv 2 \pmod{3}$ (and $n \geq 8$), we take $\frac{n-8}{3}$ independent 3-edges and two disjoint copies of K_4^3 . ■

We continue with the proof of Theorem 1.3, which deals with the graph case.

Proof of Theorem 1.3. Observe that (i) follows from (i) of Proposition 1.4.

To prove (ii), observe that for any $\{1\}$ -transversal S , we have that S and $V(G) \setminus S$ is a proper 2-coloring of G . In particular, there are no $\{1\}$ -transversals in non-bipartite graphs. There are two $\{1\}$ -transversals in any connected bipartite component. If there are no isolated vertices, then there are at most $\lfloor n/2 \rfloor$ connected components. If there are isolated vertices, then each of them is contained in any maximal $\{1\}$ -transversal, thus they do not change $h_{\{1\}}^{(2)}(n)$ and there are at most $\lfloor n/2 \rfloor$ connected components that are not isolated vertices.

Observe that **(iii)** follows from **(ii)** and **(iii)** of Proposition 1.4. Recall that **(iv)** is Theorem 1.1.

To prove **(v)**, observe first that the complement of a $\{0, 1\}$ -transversal is a $\{1, 2\}$ -transversal, which proves that $g_{\{0,1\}}^{(2)}(n) = g_{\{1,2\}}^{(2)}(n)$. The lower bound is given by the star $K_{1,n-1}$: the center is a $\{0, 1\}$ -transversal, and any subset of leaves is also a $\{0, 1\}$ -transversal.

To prove the upper bound, we apply induction on n ; the base step $n = 2$ is trivial. Assume that the statement holds for $n - 1$ and consider an n -vertex graph G without isolated vertices. Let v be a vertex of G , and let u be a neighbor of v with minimum degree. If u has another neighbor in G , then we can apply the induction hypothesis on the graph G' we obtain by deleting v from G . Thus, there are at most $1 + 2^{n-2}$ independent sets in G' . This is the number of independent sets in G that do not contain v . The independent sets containing v must avoid u , and thus there are at most 2^{n-2} of them. Hence there are at most $1 + 2^{n-2} + 2^{n-2} = 1 + 2^{n-1}$ independent sets in G .

Observe that we can pick an arbitrary vertex v , thus we are done unless each vertex has only neighbors with degree 1, i.e., G is a matching. In this case n is even and there are at most $2^{n/2} \leq 1 + 2^{n-1}$ independent sets in G , completing the proof.

Observe that **(vi)** follows from **(ii)** of Proposition 1.4, **(vii)** follows from **(iv)** of Proposition 1.4, and **(viii)** follows from **(v)** of Proposition 1.4. ■

We continue with proving Proposition 1.4. Recall that it contains several simple observations.

Proof of Proposition 1.4. To prove **(i)**, observe that a $\{0\}$ -transversal is a set of isolated vertices. There is only one such maximal set in any hypergraph, and in the case that there are no isolated vertices, \emptyset is the only $\{0\}$ -transversal.

To prove **(ii)**, observe that the whole vertex set is an A -transversal, thus it is the only maximal one.

To prove **(iii)**, observe that an $\{r\}$ -transversal contains all the hyperedges, i.e., all the non-isolated vertices, thus the whole vertex set in our case.

To prove **(iv)**, observe that every subset of vertices is a $\{0, 1, \dots, r\}$ -transversal.

To prove **(v)**, observe that for any connected component C in a hypergraph \mathcal{H} , a $\{0, r\}$ -transversal either contains C or avoids C . Therefore, the only choice to make is to pick some of the at most $\lfloor n/r \rfloor$ components. Clearly, there are hypergraphs with $\lfloor n/r \rfloor$ components and no isolated vertices.

To prove **(vi)**, observe that the complement of an A -transversal is a B -transversal, thus the graph with $g_A^{(r)}(n)$ A -transversals contains at least that many B -transversals, showing $g_A^{(r)}(n) \leq g_B^{(r)}(n)$. By symmetry, we also have $g_B^{(r)}(n) \leq g_A^{(r)}(n)$, completing the proof.

To prove **(vii)**, observe first that if S is an $\{a\}$ -transversal and $v \notin S$ such that v is contained in a hyperedge, then $S \cup \{v\}$ is not an $\{a\}$ -transversal. This means that in a hypergraph without isolated vertices, every $\{a\}$ -transversal is maximal, showing that $g_{\{a\}}^{(r)}(n) \leq h_{\{a\}}^{(r)}(n)$.

Let \mathcal{H} be an n -vertex r -uniform hypergraph with $h_{\{a\}}^{(r)}(n)$ maximal $\{a\}$ -transversals. We pick one with the smallest number i of isolated vertices among those. Let I be the set of isolated vertices. If $i = 0$, we are done. If $i \geq r$, then we can add a hyperedge on r isolated vertices. Clearly, the number of $\{a\}$ -transversals increases, completing the proof.

If $0 < i < r$, we pick $r - i$ vertices from an arbitrary hyperedge and add the hyperedge containing them and I , to obtain the hypergraph \mathcal{H}' . Then we know the number j of vertices that an $\{a\}$ -transversal needs to contain from I . Observe that every maximal $\{a\}$ -transversal S of \mathcal{H} contains I . Now this is replaced by $\{a\}$ -transversals consisting of $S \setminus I$ and an arbitrary j -subset of I . This means that \mathcal{H}' contains at least as many $\{a\}$ -transversals as the number of maximal $\{a\}$ -transversals in \mathcal{H} and \mathcal{H}' contains no isolated vertices, completing the proof. \blacksquare

Finally, we prove Theorem 1.5. Recall that **(i)** determines the order of magnitude of $g_A^r(n)$, **(ii)** deals with A being the set of even (or odd) integers, and **(iii)** deals with $A = \{0, 1, \dots, a\}$.

Proof of Theorem 1.5. The lower bound in **(i)** is given by the following construction. We consider a set U of $r - p = i$ vertices, and we select $\lfloor (n - i - p)/p \rfloor$ vertex-disjoint p -sets of the remaining vertices. We take the hyperedges containing U and any of the p -sets. We form one or two r -sets containing the at most $2p - 1$ remaining vertices. The resulting r -uniform hypergraph \mathcal{H} has no isolated vertices and at least $\lfloor (n - i - p)/p \rfloor$ hyperedges that pairwise share exactly U . Let \mathcal{H}' denote the subhypergraph having the p -sets as hyperedges. Consider the A -transversals of \mathcal{H} that contain U . Those intersect any member of \mathcal{H}' exactly in an element of $A(i)$. For each hyperedge of \mathcal{H}' , there are $f(p, i, A, r)$ ways to pick such an intersection. As we can pick the vertices from distinct hyperedges of \mathcal{H}' separately, there are at least $f(p, i, A, r)^{\lfloor (n - i - p)/p \rfloor}$ A -transversals in \mathcal{H} .

Let us continue with proving the upper bound $g_A^{(r)}(n) \leq cf(p, i, A, r)^{n/p}$ for some c that depends on A and r but not on n . We apply induction on n ; the base cases are covered by the option of picking c large enough. Let \mathcal{H} be an n -vertex r -uniform hypergraph without isolated vertices. Let v be a vertex of minimum degree $d \geq 1$ and let H_1, \dots, H_d be the hyperedges containing v . Let Q denote the set of vertices that are contained in $H_1 \cup \dots \cup H_d$ and no other hyperedges. Thus $v \in Q$.

Now let us delete Q and H_1, \dots, H_d , and let \mathcal{H}' be the resulting hypergraph. Observe that there are no isolated vertices in \mathcal{H}' , as each vertex not in Q is contained in another hyperedge (here we also use that d is the minimum degree). Thus we can apply the induction hypothesis to obtain that there are at most $cf(p, i, A, r)^{(n - |Q|)/p}$ A -transversals in \mathcal{H}' .

Observe that if S is an A -transversal in \mathcal{H} , then $S \setminus Q$ is an A -transversal in \mathcal{H}' . Given $S \setminus Q$, we know the cardinality i of $(S \setminus Q) \cap H_1$. Then we know that S is an A -transversal in \mathcal{H} if and only if the number of vertices in $S \cap Q$ is an element of $A(i)$. This means that for each A -transversal of \mathcal{H}' , there are at most $f(|Q|, i, A, r)$ ways to extend it to an A -transversal in \mathcal{H} (note that it can be less if $d > 1$ and some H_j satisfies $|(S \setminus Q) \cap H_j| \neq i$). This implies that there are at most $cf(|Q|, i, A, r)f(p, i, A, r)^{(n - |Q|)/p} =$

$cf(|Q|, i, A, r)f(p, i, A, r)^{n/p}/f(p, i, A, r)^{|Q|/p}$ A -transversals in \mathcal{H} . We have $f(|Q|, i, A, r) \leq f(p, i, A, r)^{|Q|/p}$ by the definition of p , completing the proof.

To prove (ii), let $n = ar + b$, so the stated bound becomes $2^{a(r-1)+b-1}$. Let us show the lower bound. If $b = 0$, then n/r independent hyperedges give the lower bound, since there are 2^{r-1} ways to pick a subset of even/odd cardinality from any hyperedge. If $b > 0$, we pick $(a - 1)$ independent edges, and two edges H and H' covering together $a + b$ vertices, thus sharing $a - b$ vertices. There are 2^{r-1} ways to pick a subset of even/odd cardinality from H , and it determines whether odd or even many vertices are needed to be selected from $H' \setminus H$ so that the resulting set intersects H' in an element of A (or B). There are 2^{b-1} ways to pick such a set from $H' \setminus H$, completing the proof of the lower bound.

To prove the upper bound, we apply induction on n . The base case $n = r$ is trivial. Now we proceed as in the proof of the upper bound in (i), defining Q and \mathcal{H}' the same way. The inductive hypothesis implies there are at most $2^{\lfloor \frac{r-1}{r}(n-|Q|) \rfloor}$ A -transversals and B -transversals in \mathcal{H}' . As in (i), we know the number of vertices we add from Q is a member of $A(i)$ or $B(i)$; thus we know whether we need to select an odd or even number of vertices from Q to obtain an A -transversal (or B -transversal) in \mathcal{H} .

In both cases, there are $2^{|Q|-1}$ ways to pick them, thus we have that there are at most $2^{\lfloor \frac{r-1}{r}(n-|Q|) \rfloor + |Q| - 1}$ A -transversals (B -transversals) in \mathcal{H} . If $|Q| < b$, then this upper bound is $2^{a(r-1)+b-|Q|-1+|Q|-1} < 2^{a(r-1)+b-1}$. If $|Q| = b$, then this upper bound is $2^{a(r-1)+b-1}$. If $|Q| > b$, then (using that $|Q| \leq r$) this upper bound is $2^{(a-1)(r-1)+r+b-|Q|-1+|Q|-1} = 2^{a(r-1)+b-1}$.

To prove (iii), let \mathcal{H}_1 be an n -vertex r -uniform hypergraph and let \mathcal{H}'_1 be its k -shadow, i.e., the k -uniform hypergraph where a k -set H is a hyperedge if and only if H is contained in a hyperedge of \mathcal{H}_1 . Let S be a $\{0, 1, \dots, a\}$ -transversal in \mathcal{H}_1 . We claim that S is also a $\{0, 1, \dots, a\}$ -transversal in \mathcal{H}'_1 . Indeed, if a hyperedge H of \mathcal{H}'_1 shares more than a elements with S , then the hyperedge of \mathcal{H}_1 containing H also shares more than a elements with S . This shows that $g_{\{0,1,\dots,a\}}^{(r)}(n) \leq g_{\{0,1,\dots,a\}}^{(k)}(n)$.

If S is a maximal $\{0, 1, \dots, a\}$ -transversal in \mathcal{H}_1 , then we claim that S is also a maximal $\{0, 1, \dots, a\}$ -transversal in \mathcal{H}'_1 . Indeed, for any $s \notin S$, $S \cup \{s\}$ is not a $\{0, 1, \dots, a\}$ -transversal in \mathcal{H}_1 , thus it shares at least $a + 1$ vertices with a hyperedge of \mathcal{H}_1 . Then a hyperedge of \mathcal{H}'_1 contains these $a + 1$ vertices, a contradiction. This shows that $h_{\{0,1,\dots,a\}}^{(r)}(n) \leq h_{\{0,1,\dots,a\}}^{(k)}(n)$.

Let \mathcal{H}_2 be an $(n - r + k)$ -vertex k -uniform hypergraph and let \mathcal{H}'_2 be the n -vertex r -uniform hypergraph we obtain by adding the same $r - k$ new vertices to each hyperedge. If S is a $\{0, 1, \dots, a\}$ -transversal in \mathcal{H}_2 , then clearly S is also a $\{0, 1, \dots, a\}$ -transversal in \mathcal{H}'_2 . This shows that $g_{\{0,1,\dots,a\}}^{(r)}(n) \geq g_{\{0,1,\dots,a\}}^{(k)}(n - r + k)$.

If S is a maximal $\{0, 1, \dots, a\}$ -transversal in \mathcal{H}_2 , then it is not necessarily maximal in \mathcal{H}'_2 , as some of the $r - k$ new elements may be added to S . However, let S_1 and S_2 be two distinct maximal $\{0, 1, \dots, a\}$ -transversals in \mathcal{H}_2 , and let S'_i be a maximal $\{0, 1, \dots, a\}$ -transversal in \mathcal{H}'_2 that contains S_i . We claim that $S'_1 \neq S'_2$. Indeed, there exists $s_1 \in S_1 \setminus S_2$ and $s_2 \in S_2 \setminus S_1$ by the maximality of these two transversals. Then $s_1 \in S'_1 \setminus S'_2$ and $s_2 \in S'_2 \setminus S'_1$, thus $S'_1 \neq S'_2$. This shows that $h_{\{0,1,\dots,a\}}^{(r)}(n) \geq h_{\{0,1,\dots,a\}}^{(k)}(n - r + k)$. \blacksquare

3 Concluding remarks

One could avoid the degenerate example of the empty (hyper)graph by different assumptions instead of having no isolated vertices. We have mentioned that in the graph case, the maximum number of independent sets in graphs with minimum degree at least $\delta > 1$ was studied. Another possible assumption is that the host hypergraph is connected. The order of magnitude is determined by **(i)** of Theorem 1.5 in several cases. In the remaining cases, the construction is mostly a matching. It is easy to see from the proof that a similar statement holds in these cases; we only need to strengthen the assumption $q \leq r$ to $q < r$ when defining $f(q, i, A, r)$. Note that the largest number of *maximal* independent sets in connected graphs has been determined in [2] and [4].

One can consider these problems for non-uniform hypergraphs as well. In the case of the function g and $1 \in A$, one should assume that the hyperedges contain more than 1 vertex, as there are 2^n independent sets in the hypergraph consisting of n 1-element hyperedges. An even better assumption is that each hyperedge contains more than $\max A$ vertices, as there are $2^n \{0, 1, \dots, a\}$ -transversals among all the k -sets for each $k \leq a$. We remark that the proof of Theorem 1.5 **(iii)** extends to the non-uniform setting, and thus it is enough to determine the number of (maximum) $\{0, 1, \dots, a\}$ -transversals in $(a + 1)$ -uniform families in order to determine the order of magnitude of the analogous quantity in non-uniform families.

Finite projective planes provide prototype examples of (intersecting) r -uniform hypergraphs. In that context, a much studied concept is the k -sets of (m, n) -type [9]. It means A consists of two non-zero elements m and n and the ground-set has k elements. Some values lead to beautiful constructions, some constructions lead to applications even in coding theory. It shows already the existence problem for two-element A has useful applications.

References

- [1] G.J. Chang, M.J. Jou. Survey on counting maximal independent sets. In Proceedings of the Second Asian Mathematical Conference (1995) 265–275.
- [2] Z. Füredi. The number of maximal independent sets in connected graphs, J. Graph Theory 11(4) (1987) 463–470.
- [3] W. Gan, P.-S. Loh, and B. Sudakov. Maximizing the number of independent sets of a fixed size. Combinatorics, Probability and Computing, 24(3) (2015) 521–527.
- [4] J.M. Griggs, C.M. Grinstead, D.R. Guichard. The number of maximal independent sets in a connected graph, Discrete Mathematics 68(2-3) (1988) 211–220.
- [5] D. Galvin. (2011). Two problems on independent sets in graphs. Discrete Mathematics, 311(20), 2105–2112.
- [6] Z. Lonc, M. Truszczyński. On the number of minimal transversals in 3-uniform hypergraphs, Discrete Mathematics, 308(16) (2008) 3668–3687.

- [7] R.E. Miller, D.E. Muller. A problem of maximum consistent subsets, 1960. IBM Research Report RC-240, J.T. Watson Research Center, New York, USA.
- [8] J.W. Moon, L. Moser. On cliques in graphs, *Israel J. Math.* 3 (1965) 23–28.
- [9] V. Napolitano. Two-character sets in finite linear spaces. *Contributions to Discrete Mathematics* 3(1) (2007) 123–133.
- [10] J.M. Nielsen. On the number of maximal independent sets in a graph. *BRICS Report Series*, 9(15) (2002).
- [11] C. Palmer. Problem booklet of the 11th Emléktábla Workshop held in 2022.
<https://www.renyi.hu/~emlektab/emlektabla11.pdf>
- [12] I. Tomescu. Le nombre maximum de cliques et de recouvrements par cliques des hypergraphes chromatiques complets, *Discrete Mathematics* 37(2-3) (1981) 263–277.
- [13] D.R. Wood. On the number of maximal independent sets in a graph. *Discrete Maths. & Theoretical Computer Science* 13(3) (2011) 17–20.