

JORDAN PRODUCT AND ANALYTIC CORE PRESERVERS

S. Elouazzani

*Departement of Mathematics, Labo LIABM,
Faculty of Sciences, 60000 Oujda, Morocco
elouazzani.soufiane@ump.ac.ma*

M. Elhodaibi

*Departement of Mathematics, Labo LIABM,
Faculty of Sciences, 60000 Oujda, Morocco
m.elhodaibi@ump.ac.ma*

Abstract Let $\mathcal{B}(X)$ be the algebra of all bounded linear operators on an infinite-dimensional complex Banach space X . For an operator $T \in \mathcal{B}(X)$, $K(T)$ denotes as usual the analytic core of T . We determine the form of surjective maps ϕ on $\mathcal{B}(X)$ satisfying

$$K(\phi(T)\phi(S) + \phi(S)\phi(T)) = K(TS + ST)$$

for all $T, S \in \mathcal{B}(X)$.

2020 Mathematics Subject Classification. 47B49, 47A10, 47A11

Keywords— Analytic core; Inner local spectral radius; Jordan product; Pre-server.

1 Introduction

Throughout this paper, $\mathcal{B}(X)$ denotes the algebra of all bounded linear operators on an infinite-dimensional complex Banach space X . Recall that the local resolvent set $\rho_T(x)$ of an operator $T \in \mathcal{B}(X)$ at a point $x \in X$ is the set of all λ in \mathbb{C} for which there exist an open neighborhood U_λ of λ in \mathbb{C} and an X -valued analytic function $f : U_\lambda \rightarrow X$ such that $(T - \mu I)f(\mu) = x$ for all $\mu \in U_\lambda$. As it is known,

$\sigma_T(x)$ denotes the local spectrum of T at x , it is the complement in \mathbb{C} of $\rho_T(x)$ and is a compact subset of the spectrum $\sigma(T)$ of T , possibly empty. An operator $T \in \mathcal{B}(X)$ is said to have *the single – valued extension property* (SVEP) if for every open subset U of \mathbb{C} , the equation $(T - \lambda I)f(\lambda) = 0$ for all $\lambda \in U$, has no nontrivial analytic solution f on U . If T has the SVEP, then $\sigma_T(x) \neq \emptyset$ for all nonzero vector $x \in X$. Let r be a positive scalar, we denote by $D(0, r)$ and $\overline{D}(0, r)$, respectively, the open and the closed disc centered at the origin with radius r . For every closed subset F of \mathbb{C} , the global spectral of an operator $T \in \mathcal{B}(X)$ is defined by $\mathcal{X}_T(F) := \{x \in X : \text{there exists an analytic function } f : \mathbb{C} \setminus F \rightarrow X \text{ such that } (T - \lambda)f(\lambda) = x \text{ for all } \lambda \in \mathbb{C} \setminus F\}$. The local spectral radius of T at $x \in X$ is defined by

$$r_T(x) = \limsup_{n \rightarrow \infty} \|T^n(x)\|^{\frac{1}{n}},$$

it coincides with $r_T(x) = \inf\{r \geq 0 : x \in \mathcal{X}_T(\overline{D}(0, r))\}$, and also with the maximum modulus of $\sigma_T(x)$ if T has the SVEP, see for instance [19]. Recall that for an operator $T \in \mathcal{B}(X)$, the analytical core $K(T)$ of T is the set of all $x \in X$ for which there exist $\delta > 0$ and a sequence $(x_n) \subset X$ such that $x_0 = x, Tx_{n+1} = x_n$ and $\|x_n\| \leq \delta^n \|x\|$ for all $n \geq 0$, for more information, see [3, 19]. Recall also that the inner local spectral radius $i_T(x)$ of T at $x \in X$ is defined by $i_T(x) := \sup\{r \geq 0 : x \in \mathcal{X}_T(\mathbb{C} \setminus D(0, r))\}$; see [20].

The study of local spectra preserver problems was initiated by A. Bourhim and T. Ransford [12], they characterized additive maps on $\mathcal{B}(X)$ which preserve the local spectrum of all operators $T \in \mathcal{B}(X)$ at each vector $x \in X$. After that, the maps preserving local spectrum and local spectral radius have been studied by many authors; see [4, 7, 9, 13, 14]. In [16], M.E. El Kettani and H. Benbouziane showed that a surjective additive map ϕ on $\mathcal{B}(X)$ satisfies

$$i_T(x) = 0 \quad \text{if and only if} \quad i_{\phi(T)}(x) = 0$$

for all $x \in X$ and $T \in \mathcal{B}(X)$ if and only if there exists a nonzero scalar $c \in \mathbb{C}$ such that $\phi(T) = cT$ for all $T \in \mathcal{B}(X)$. M. Elhodaibi and A. Jaatit, in [15], proved the result of T. Jari [17] for all maps ϕ on $\mathcal{B}(X)$ (not necessarily surjective) satisfying

$$i_{\phi(T) - \phi(S)}(x) = 0 \iff i_{T-S}(x) = 0$$

for every $x \in X$ and $T, S \in \mathcal{B}(X)$ if and only if there exists a nonzero scalar $c \in \mathbb{C}$ such that $\phi(T) = cT + \phi(0)$. For the inner local spectral radius preservers of generalized product, A. Achchi [2], showed that if a surjective map ϕ on $\mathcal{B}(X)$ satisfies

$$i_{\phi(T_1) * \dots * \phi(T_k)}(x) = 0 \iff i_{T_k * \dots * T_k}(x) = 0$$

for all $x \in X$ and all $T_1, \dots, T_k \in \mathcal{B}(X)$, then there exists a map $\gamma : \mathcal{B}(X) \rightarrow \mathbb{C} \setminus \{0\}$ such that $\phi(T) = \gamma(T)T$. Let us mention other papers that characterize

nonlinear maps preserving certain spectral and local spectral quantities of product and Jordan product of operators or matrices, see for instance [5, 6, 10, 11]. In the present paper, we characterize surjective maps ϕ (not necessarily linear) on $\mathcal{B}(X)$ that satisfying

$$K(TS + ST) = K(\phi(T)\phi(S) + \phi(S)\phi(T))$$

for all $T, S \in \mathcal{B}(X)$. This gives immediately the characterization of surjective maps $\phi : \mathcal{B}(X) \rightarrow \mathcal{B}(X)$ satisfying

$$i_{TS+ST}(x) = 0 \quad \text{if and only if} \quad i_{\phi(T)\phi(S)+\phi(S)\phi(T)}(x) = 0$$

for all $x \in X$ and $T, S \in \mathcal{B}(X)$.

2 Preliminaries and Notations

For every vector $x \in X$ and every linear functional $f \in X^*$, where X^* is the topological dual space of X . Denote by $x \otimes f$ the operator of rank at most one, with $(x \otimes f)(z) = f(z)x$ for all $z \in X$. Recall that $x \otimes f$ is nilpotent if and only if $f(x) = 0$, and it is idempotent if and only if $f(x) = 1$. Let $\mathcal{F}_1(X)$ denotes the set of all rank at most one operators on X . For a subspace A of X , the subspace spanned by A is denoted by $span(A)$. Observe that, if $f(x) = 0$ then $K(x \otimes f) = \{0\}$, if not then $K(x \otimes f) = span\{x\}$. Let $\dim(Y)$ be the dimension of a subspace Y of X , and let for every operator $T \in \mathcal{B}(X)$, $N(T)$ be the kernel of T and $R(T)$ be its range.

In The next lemma, we give some basic properties of $K(T)$ and $i_T(x)$, see for instance [3, 19, 20].

Lemma 1. *Let $T \in \mathcal{B}(X)$, then the following statements hold.*

- (i) $K(T) \subset R(T)$.
- (ii) $K(\lambda T) = K(T)$ for all nonzero scalar $\lambda \in \mathbb{C}$.
- (iii) If M is a closed subspace of X and $TM = M$ then $M \subset K(T)$.
- (iv) If T is quasi-nilpotent then $K(T) = \{0\}$.
- (v) $N(T - \lambda) \subset K(T)$ for all nonzero scalar $\lambda \in \mathbb{C}$.
- (vi) $K(T) = \{x \in X : 0 \notin \sigma_T(x)\}$ for all $T \in \mathcal{B}(X)$.
- (vii) $i_T(x) = 0$ if and only if $0 \in \sigma_T(x)$.

The next lemma and its proof are quoted in [18].

Lemma 2. *Let $T, S \in \mathcal{B}(X)$. Assume that for every $x \in X$ the vector Tx belongs to the linear span of x and Sx . Then $T = \lambda I + \mu S$ for some $\lambda, \mu \in \mathbb{C}$.*

Proof. See lemma 2.4 in [18]. □

The following lemma was introduced in [2] for generalized product of operators, but in this paper we use another techniques to characterize rank one operators by the dimension of analytical core of Jordan product of operators.

Lemma 3. *For a nonzero operator $A \in \mathcal{B}(X)$, the following statements are equivalent.*

- (i) *A is a rank one operator.*
- (ii) *$\dim(K(TA + AT)) \leq 2$ for all $T \in \mathcal{B}(X)$.*

Proof. For $A \in \mathcal{B}(X)$, we consider the following operator $R = AT + TA$ with $T \in \mathcal{B}(X)$ is an arbitrary operator. It is clear that we have (i) \implies (ii). For (ii) \implies (i) suppose that $\text{Rank}(A) \geq 2$, and let us show that there exists $T \in \mathcal{B}(X)$ such that $\dim(K(R)) \geq 3$. We will distinguish two cases.

Case 1. If $\text{Rank}(A) \geq 3$.

Let $y_1, y_2, y_3 \in X$ be linearly independent vectors such that $y_1 = Ax_1, y_2 = Ax_2$ and $y_3 = Ax_3$ where $x_1, x_2, x_3 \in X$ are obviously linearly independent. We will distinguish four cases.

- (1) If $\text{span}\{x_1, x_2, x_3\} = \text{span}\{y_1, y_2, y_3\}$.
Let $T \in \mathcal{B}(X)$ be an operator satisfying $Ty_1 = x_1, Ty_2 = x_2$ and $Ty_3 = x_3$. As $AT = TA = I$ on $\text{span}\{x_1, x_2, x_3\}$, then $Rx_1 = 2x_1, Rx_2 = 2x_2$ and $Rx_3 = 2x_3$. Hence $\text{span}\{x_1, x_2, x_3\} \subset N(R - 2I) \subset K(R)$, and so $\dim(K(R)) \geq 3$.
- (2) If $\dim(\text{span}\{x_1, x_2, x_3, y_1, y_2, y_3\}) = 4$.
Let for example $\text{span}\{x_1, x_2, x_3, y_1, y_2, y_3\} = \text{span}\{x_1, y_1, y_2, y_3\}$. Consider an operator $T \in \mathcal{B}(X)$ satisfying $Ty_1 = x_1, Ty_2 = x_2, Ty_3 = x_3$ and $Tx_1 = 0$. For $x_2 = \alpha x_1 + \beta y_1 + \gamma y_2 + \delta y_3$ and $x_3 = \alpha' x_1 + \beta' y_1 + \gamma' y_2 + \delta' y_3$ where $\alpha, \beta, \gamma, \delta, \alpha', \beta', \gamma', \delta' \in \mathbb{C}$, we get that $Rx_2 = 2x_2 - \alpha x_1$ and $Rx_3 = 2x_3 - \alpha' x_1$. Since $Rx_1 = x_1$, then $RM = M$ with $M = \text{span}\{x_1, x_2, x_3\}$. Hence $M \subset K(R)$, therefore $\dim(K(R)) \geq 3$.

(3) If $\dim(\text{span}\{x_1, x_2, x_3, y_1, y_2, y_3\}) = 5$.

Let for example $\text{span}\{x_1, x_2, x_3, y_1, y_2, y_3\} = \text{span}\{x_1, x_2, y_1, y_2, y_3\}$. Take an operator $T \in \mathcal{B}(X)$ satisfying $Ty_1 = x_1, Ty_2 = x_2, Ty_3 = x_3$ and $Tx_1 = Tx_2 = 0$. For $x_3 = \alpha x_1 + \beta x_2 + y$ where $y \in \text{Span}\{y_1, y_2, y_3\}$ and $\alpha, \beta \in \mathbb{C}$, we get that $Rx_1 = x_1, Rx_2 = x_2$ and $Rx_3 = 2x_3 - \alpha x_1 - \beta x_2$. It follows that $RM = M$ with $M = \text{span}\{x_1, x_2, x_3\}$. Then $M \subset K(R)$, thus $\dim(K(R)) \geq 3$.

(4) If $\dim(\text{span}\{x_1, x_2, x_3, y_1, y_2, y_3\}) = 6$.

Let $T \in \mathcal{B}(X)$ be an operator satisfying $Ty_1 = x_1, Ty_2 = x_2, Ty_3 = x_3$ and $Tx_1 = Tx_2 = Tx_3 = 0$. Since $Rx_1 = x_1, Rx_2 = x_2$ and $Rx_3 = x_3$. It follows that $\text{span}\{x_1, x_2, x_3\} \subset N(R - I) \subset K(R)$, and so $\dim(K(R)) \geq 3$.

Case 2. If $\text{Rank}(A) = 2$.

Let y_1 and y_2 two linearly independent vectors such that $y_1 = Ax_1, y_2 = Ax_2$ and $Ay_1 = ay_1 + by_2$ where $a, b \in \mathbb{C}$.

If $x_1, y_1, y_2 \in X$ are linearly dependent, we can choose $u \in N(A)$ such that $x_1 + u, y_1, y_2 \in X$ are linearly dependent. Now, if $x_1 + u, x_2, y_1 \in X$ are linearly dependent. As $x_1 + u$ and y_1 are linearly independent, and $N(A)$ is an infinite-dimensional subspace of X . Then we can take $v \in N(A)$ such that $x_1 + u, x_2 + v, y_1 \in X$ are linearly independent. Thus without loss of generality, we may assume that (x_1, y_1, y_2) and (x_1, x_2, y_1) are linearly independent. Let us distinguish two cases.

(1) If $x_2 \in \text{span}\{x_1, y_1, y_2\}$.

Let $x_2 = \alpha x_1 + \beta y_1 + \gamma y_2$ where $\alpha, \beta, \gamma \in \mathbb{C}$. For an operator $T \in \mathcal{B}(X)$ satisfying $Ty_1 = x_1, Ty_2 = x_2$ and $Tx_1 = 0$, we get that

$$\begin{cases} Rx_1 = x_1 \\ Rx_2 = 2x_2 - \alpha x_1 \\ Ry_1 = y_1 + ax_1 + bx_2. \end{cases}$$

This yields that $RM = M$ with $M = \text{span}\{x_1, x_2, y_1\}$. Hence $M \subset K(R)$, thus $\dim(K(R)) \geq 3$.

(2) If (x_1, x_2, y_1, y_2) are linearly independent.

Consider an operator $T \in \mathcal{B}(X)$ satisfying $Ty_1 = x_1, Ty_2 = x_2$ and $Tx_1 =$

$Tx_2 = 0$. Hence

$$\begin{cases} Rx_1 = x_1 \\ Rx_2 = x_2 \\ Ry_1 = y_1 + ax_1 + bx_2. \end{cases}$$

It follows that $RM = M$ with $M = \text{span}\{x_1, x_2, y_1\}$. Thus $M \subset K(R)$, and so $\dim(K(R)) \geq 3$. This establishes the lemma. \square

For Jordan product of operators, we give a characterization for two operators to be linearly dependent.

Lemma 4. *For A and B two operators in $\mathcal{B}(X)$, the following statements are equivalent.*

(i) $K(AT + TA) = K(BT + TB)$ for every $T \in \mathcal{B}(X)$.

(ii) $K(AF + FA) = K(BF + FB)$ for every $F \in \mathcal{F}_1(X)$.

(iii) $B = \lambda A$ for some nonzero scalar $\lambda \in \mathbb{C}$.

Proof. For $A, B \in \mathcal{B}(X)$, we consider the following operators $R = AF + FA$ and $S = BF + FB$ with $F \in \mathcal{F}_1(X)$ is an arbitrary operator. The implications (i) \implies (ii) and (iii) \implies (i) are straightforward. We only need to show the implication (ii) \implies (iii). Let us distinguish two cases.

Case 1. (x, Ax, Bx) are linearly independent for some nonzero vector $x \in X$. Let f be a linear functional in X^* such that $f(x) = f(Ax) = 0$ and $f(Bx) = 1$. For $F = x \otimes f$, we obtain that

$$\begin{cases} Rx = 0 \\ RAx = f(A^2x)x \end{cases} \quad \text{and} \quad \begin{cases} Sx = x \\ SBx = f(B^2x)x + Bx. \end{cases}$$

This yields that $R^2 = 0$, thus $K(R) = \{0\}$ but $x \in K(S)$, which is a contradiction.

Case 2. (x, Ax, Bx) are linearly dependent for all $x \in X$.

Let x be a vector in X . If x and Ax are linearly independent, then we get that $Bx \in \text{span}\{x, Ax\}$. If not, then $Ax = \mu x$ where $\mu \in \mathbb{C}$. Suppose that Bx and x are linearly independent. It follows that there exists a linear functional $f \in X^*$ such that $f(x) = 0$ and $f(Bx) = 1$. For $T = x \otimes f$ we obtain that $Rx = 0$ and $Sx = x$. Thus $x \in K(S) = K(R) = \{0\}$. This contradiction shows that $Bx \in \text{span}\{x\} \subset \text{span}\{x, Ax\}$. We conclude that $Bx \in \text{span}\{x, Ax\}$ for all

$x \in X$. Lemma 2 implies that $B = \lambda A + \alpha I$ where $\alpha, \lambda \in \mathbb{C}$. It is clear that we have

$$A = 0 \text{ if and only if } B = 0$$

and also

$$A \in \mathbb{C}I \text{ if and only if } B \in \mathbb{C}I.$$

Now, if $A \notin \mathbb{C}I$, assume that $\alpha \neq 0$. For that we distinguish two cases.

- (1) If (x, Ax, A^2x) are linearly independent for some $x \in X$.

Take $F = x \otimes f$ such that $f(x) = 1$ and $f(Ax) = f(A^2x) = 0$. Hence $R^2 = 0$, thus $K(R) = \{0\}$. On the other hand, we have $K(S) = \text{span}\{\lambda Ax + 2\alpha x\}$, this is a contradiction.

- (2) If (x, Ax, A^2x) are linearly dependent for all $x \in X$.

It is obvious that we have $A^2x \in \text{span}\{x, Ax\}$ for all $x \in X$. It follows by Lemma 2 that $A^2 = aA + bI$ where $a, b \in \mathbb{C}$. Without loss of generality, we may assume that $\lambda = 1$. Since $A \notin \mathbb{C}I$. Then there exists a nonzero vector $x \in X$ such that Ax and x are linearly independent. Consider an operator $F = x \otimes f$ such that $f(x) \neq 0$ and $f(Ax)^2 = f(x)f(A^2x)$, then we obtain

$$\begin{cases} Rx = f(Ax)x + f(x)Ax \\ RAx = f(A^2x)x + f(Ax)Ax \end{cases} \quad \text{and} \quad \begin{cases} Sx = f(Bx)x + f(x)Bx \\ SAx = f(BAx)x + f(Ax)Bx. \end{cases}$$

Hence

$$\begin{cases} Rx = f(Ax)x + f(x)Ax \\ RAx = \frac{f(Ax)}{f(x)}Rx \end{cases} \quad \text{and} \quad \begin{cases} Sx = f(Ax)x + f(x)Ax + 2\alpha f(x)x \\ SAx = (f(A^2x) + \alpha f(Ax))x + f(Ax)(Ax + \alpha x). \end{cases}$$

Thus

$$\begin{cases} Sx = Rx + 2\alpha f(x)x \\ SAx = f(A^2x)x + f(Ax)Ax + 2\alpha f(Ax)x, \end{cases}$$

and so

$$\begin{cases} Sx = Rx + 2\alpha f(x)x \\ SAx = \frac{f(Ax)}{f(x)}Sx. \end{cases}$$

- If $f(Ax) = 0$, we have $R^2 = 0$ and $K(R) = \{0\}$. Since $Sx = f(x)Ax + 2\alpha f(x)x$ and $SAx = 0$, then $S^2x = 2\alpha f(x)Sx$. This yields that $Sx \in K(S) = K(R) = \{0\}$, contradiction.

- Now, if $f(Ax) \neq 0$, we have $K(R) = \text{span}\{Rx\}$ and $K(S) = \text{span}\{Sx\}$. Hence there exists a nonzero scalar $\delta \in \mathbb{C}$ such that $Sx = \delta Rx$. This implies that $2\alpha f(x)x = (\delta - 1)Rx$. As x and Ax are linearly independent, then $\delta = 1$, contradiction.

Finally, we get that $\alpha = 0$ and the proof is complete. \square

3 Main Result

We start this section with the following theorem that gives a characterization of maps preserving the analytic core of the Jordan product of operators.

Theorem 1. *Let $\phi : \mathcal{B}(X) \longrightarrow \mathcal{B}(X)$ be a surjective map. Then the following assertions are equivalent:*

(i) *For every $T, S \in \mathcal{B}(X)$, we have*

$$K(\phi(T)\phi(S) + \phi(S)\phi(T)) = K(TS + ST)$$

(ii) *There is a map $\gamma : \mathcal{B}(X) \longrightarrow \mathbb{C} \setminus \{0\}$ such that $\phi(T) = \gamma(T)T$ for all $T \in \mathcal{B}(X)$.*

Proof. Clearly, we have (ii) \implies (i). It remains to show that (i) \implies (ii). We divide the proof into several steps.

Step 1. For any $R \in \mathcal{B}(X)$, $\phi(R) = 0$ if and only if $R = 0$.

Let us show that $\phi(0) = 0$. We have

$$\begin{aligned} K(\phi(T)\phi(0) + \phi(0)\phi(T)) &= K(T0 + 0T) \\ &= \{0\} \\ &= K(\phi(T)0 + 0\phi(T)). \end{aligned}$$

As ϕ is surjective, then Lemma 4 implies that $\phi(0) = 0$.

Now, assume that $\phi(R) = 0$ for an operator $R \in \mathcal{B}(X)$. Hence

$$\begin{aligned} K(RT + TR) &= K(\phi(R)\phi(T) + \phi(T)\phi(R)) \\ &= \{0\} \\ &= K(0T + T0). \end{aligned}$$

From Lemma 4, we infer that $R = 0$.

Step 2. For any operator $F \in \mathcal{B}(X)$, $\phi(F) \in \mathcal{F}_1(X)$ if and only if $F \in \mathcal{F}_1(X)$.

Let $F \in \mathcal{B}(X)$ such that $\phi(F)$ is a rank one operator. Step 1 implies that $F \neq 0$. By Lemma 3, we obtain that

$$\dim(K(S\phi(F) + \phi(F)S)) \leq 2$$

for all $S \in \mathcal{B}(X)$. As ϕ is surjective, then

$$\dim(K(TF + FT)) = \dim(K(\phi(T)\phi(F) + \phi(F)\phi(T))) \leq 2$$

for all $T \in \mathcal{B}(X)$. It follows from Lemma 3 that F is a rank one operator. Just as before, we obtain in the same way that if $F \in \mathcal{F}_1(X)$ then $\phi(F) \in \mathcal{F}_1(X)$.

Step 3. There is a nonzero scalar $\lambda_F \in \mathbb{C}$ such that $\phi(F) = \lambda_F F$ for all $F \in \mathcal{F}_1(X)$. For that we discuss two cases, if F is nilpotent or not.

- (1) There is a nonzero scalar λ_P such that $\phi(P) = \lambda_P P$, for every rank one idempotent operator $P \in \mathcal{F}_1(X)$.

Let $x \in X$ and $f \in X^*$ such that $f(x) = 1$. Step 2 implies that there exist a nonzero vector $y \in X$ and a linear functional $g \in X^*$ such that $\phi(x \otimes f) = y \otimes g$. Since $x \otimes f$ is a rank one idempotent operator, then

$$\begin{aligned} \text{span}\{x\} &= K(x \otimes f) \\ &= K(x \otimes fx \otimes f + x \otimes fx \otimes f) \\ &= K(y \otimes gy \otimes g + y \otimes gx \otimes g) \\ &= K(2g(y)y \otimes g). \end{aligned}$$

This yields that $g(y) \neq 0$ and $\text{span}\{y\} = K(2g(y)y \otimes g) = \text{span}\{x\}$. Therefore $y = \alpha x$ for a nonzero scalar $\alpha \in \mathbb{C}$. Without loss of generality, we may and shall assume that $\alpha = 1$, then we get that $\phi(x \otimes f) = x \otimes g_{x,f}$ for certain linear functional $g_{x,f} \in X^*$. We claim that f and $g_{x,f}$ are linearly dependent. If not, take a nonzero vector $z \in X$ such that x and z are linearly independent in X , with $f(z) = 1$ and $g_{x,f}(z) = 0$. Just as before, one shows that there exists a linear functional $g_{z,f} \in X^*$ such that $\phi(z \otimes f) = z \otimes g_{z,f}$. Observe that

$$(x \otimes fz \otimes f + z \otimes fx \otimes f)(x + z) = 2(x + z).$$

Then

$$(x + z) \in K(x \otimes fz \otimes f + z \otimes fx \otimes f).$$

On the other hand, we have

$$\begin{aligned}
\{0\} &= K(g_{z,f}(x)z \otimes g_{x,f}) \\
&= K(x \otimes g_{x,f}z \otimes g_{z,f} + z \otimes g_{z,f}x \otimes g_{x,f}) \\
&= K(\phi(x \otimes f)\phi(z \otimes f) + \phi(z \otimes f)\phi(x \otimes f)) \\
&= K(x \otimes fz \otimes f + z \otimes fx \otimes f),
\end{aligned}$$

which is a contradiction. Hence $g_{x,f}$ and f are linearly dependent, thus $\phi(x \otimes f) = \alpha x \otimes f$ for some nonzero scalar $\alpha \in \mathbb{C}$.

- (2) There is a nonzero scalar λ_N such that $\phi(N) = \lambda_N N$, for every rank one nilpotent operator $N \in \mathcal{F}_1(X)$.

Let $x \in X$ and $f \in X^*$ such that $f(x) = 0$. Set $T = x \otimes f$, then $\phi(T) = y \otimes g$ with $g(y) = 0$. Assume that x and y are linearly independent, and so let $z \in X$ and $h \in X^*$ such that $f(z) = h(x) = 1$, $h(z) \neq 0$ and $h(y) = 0$. It follows that

$$\begin{aligned}
\{0\} &= K(g(z)y \otimes h) \\
&= K(y \otimes gz \otimes h + z \otimes hy \otimes g) \\
&= K(\phi(x \otimes f)\phi(z \otimes h) + \phi(z \otimes h)\phi(x \otimes f)) \\
&= K(x \otimes fz \otimes h + z \otimes hx \otimes f).
\end{aligned}$$

Observe that

$$(x \otimes fz \otimes h + z \otimes hx \otimes f)x = x,$$

hence

$$\text{span}\{x\} \subset K(x \otimes fz \otimes h + z \otimes hx \otimes f).$$

This contradiction shows that x and y are linearly dependent. As before, without loss of generality, we may assume that $\phi(x \otimes f) = x \otimes g_{x,f}$ for certain linear functional $g_{x,f} \in X^*$.

Suppose that f and g are linearly independent, and let $z \in X$ and $h \in X^*$ such that $f(z) = 1$, $g(z) = 0$, $h(x) = 1$ and $h(z) \neq 0$. Since

$$(x \otimes fz \otimes h + z \otimes hx \otimes f)x = x,$$

then

$$\text{span}\{x\} \subset K(x \otimes fz \otimes h + z \otimes hx \otimes f).$$

On the other hand, we get that

$$\begin{aligned}
\{0\} &= K(z \otimes g) \\
&= K(x \otimes gz \otimes h + z \otimes hx \otimes g) \\
&= K(\phi(x \otimes f)\phi(z \otimes h) + \phi(z \otimes h)\phi(x \otimes f)) \\
&= K(x \otimes fz \otimes h + z \otimes hx \otimes f).
\end{aligned}$$

Then, this contradiction shows also that f and g are linearly dependent. Thus $\phi(x \otimes f) = \beta x \otimes f$ for some nonzero scalar $\beta \in \mathbb{C}$. Now, if $f(x) \in \mathbb{C} \setminus \{0, 1\}$, obviously this can be obtained as in (1) of Step 3. Finally, there is a nonzero scalar $\lambda_F \in \mathbb{C}$ such that $\phi(F) = \lambda_F F$ for all rank one operator $F \in \mathcal{F}_1(X)$.

Step 4. There exists a map $\gamma : \mathcal{B}(X) \longrightarrow \mathbb{C} \setminus \{0\}$ such that $\phi(T) = \gamma(T)T$ for all $T \in \mathcal{B}(X)$.

For every $F \in \mathcal{F}_1(X)$ and every $T \in \mathcal{B}(X)$, we have

$$K(TF + FT) = K(\phi(T)\phi(F) + \phi(F)\phi(T)).$$

By using step 3, we get that

$$\begin{aligned} K(\phi(T)\phi(F) + \phi(F)\phi(T)) &= K(\lambda_F(\phi(T)F + F\phi(T))) \\ &= K(\phi(T)F + F\phi(T)). \end{aligned}$$

Then $K(TF + FT) = K(\phi(T)F + F\phi(T))$, for all operator $F \in \mathcal{F}_1(X)$. Lemma 4 implies that $\phi(T)$ and T are linearly dependent. Hence, there exists a map $\gamma : \mathcal{B}(X) \longrightarrow \mathbb{C} \setminus \{0\}$ such that $\phi(T) = \gamma(T)T$ for all $T \in \mathcal{B}(X)$. The proof is then complete. \square

As a consequence of theorem 1, we get the following corollary.

Corollary 1. *Let $\phi : \mathcal{B}(X) \longrightarrow \mathcal{B}(X)$ be a surjective map. Then the following assertions are equivalent.*

- (i) $i_{\phi(T)\phi(S)+\phi(S)\phi(T)}(x) = 0 \iff i_{TS+ST}(x) = 0$ for every $T, S \in \mathcal{B}(X)$ and $x \in X$.
- (ii) There exists a map $\gamma : \mathcal{B}(X) \longrightarrow \mathbb{C} \setminus \{0\}$ such that $\phi(T) = \gamma(T)T$ for all $T \in \mathcal{B}(X)$.

Proof. The implication (ii) \implies (i) is straightforward.

(i) \implies (ii) From theorem 1, and the assertions (vi) and (vii) in lemma 1, we get the desired form of ϕ . \square

References

- [1] Z. ABDELALI, A. ACHCHI AND R. MARZOUKI, *Maps preserving the local spectrum of skew-product of operators*, Linear Algebra and its Applications **485** (2015), 58-71.

- [2] A. ACHCHI, *Maps preserving the inner local spectral radius zero of generalized product of operators*, Rendiconti del Circolo Matematico di Palermo Series 2 **68** (2018), 355-362.
- [3] P. AIENA, *Abstract Fredholm theory*, Fredholm and Local Spectral Theory with Applications to Multipliers (2007), 239-308.
- [4] A. BOURHIM, *Surjective linear maps preserving local spectra*, Linear Algebra and its Applications **432** (2010), 383-393.
- [5] A. BOURHIM AND M. MABROUK, *Maps preserving the local spectrum of Jordan product of matrices*, Linear Algebra and its Applications **484** (2015), 379-395.
- [6] A. BOURHIM AND M. MABROUK, *Maps preserving the local spectrum of Jordan product*, Studia mathematica **234** (2016), 97-120.
- [7] A. BOURHIM AND J. MASHREGHI, *Local Spectral Radius Preservers*, Integral Equations and Operator Theory **76** (2013), 95-104.
- [8] A. BOURHIM AND J. MASHREGHI, *Maps preserving the local spectrum of triple product of operators*, Linear and Multilinear Algebra **63** (2014), 765-773.
- [9] A. BOURHIM AND J. MASHREGHI, *A survey on preservers of spectra and local spectra*, Invariant Subspaces of the Shift Operator Contemporary Mathematics (2015), 45-98.
- [10] A. BOURHIM AND J. MASHREGHI, *Maps Preserving The Local Spectrum Of Product Of Operators*, Glasgow Mathematical Journal **57** (2014), 709-718.
- [11] A. BOURHIM AND V. G. MILLER, *Linear maps on $M_n(C)$ preserving the local spectral radius*, Studia Mathematica **188** (2008), 67-75.
- [12] A. BOURHIM AND T. RANSFORD, *Additive Maps Preserving Local Spectrum*, Integral Equations and Operator Theory **55** (2005), 377-385.
- [13] C. COSTARA, *Continuous maps preserving local spectra of matrices*, Linear Algebra and its Applications **492** (2016), 1-8.
- [14] C. COSTARA, *Local spectrum linear preservers at non-fixed vectors*, Linear Algebra and its Applications **457** (2014), 154-161.
- [15] M. ELHODAIBI AND A. JAATIT, *Inner local spectral radius preservers*, Rendiconti del Circolo Matematico di Palermo Series 2 **67** (2017), 215-225.

- [16] M. E.-C. E. KETTANI AND H. BENBOUZIANE, *Additive maps preserving operators of inner local spectral radius zero*, Rendiconti del Circolo Matematico di Palermo **63** (2014), 311-316.
- [17] T. JARI, *Nonlinear maps preserving the inner local spectral radius*, Rendiconti del Circolo Matematico di Palermo **64** (2015), 67-76.
- [18] C.-K. LI, P. ŠEMRL, AND N.-S. SZE, *Maps preserving the nilpotency of products of operators*, Linear Algebra and its Applications **424** (2007), 222-239.
- [19] K. B. LAURSEN AND M. NEUMANN, *An Introduction to Local Spectral Theory*, Oxford University Press (2000).
- [20] T. MILLER, V. MILLER AND M. NEUMANN, *Local spectral properties of weighted shifts*, Journal of Operator Theory **51** (2004), 71-88.