

A natural pseudometric on homotopy groups of metric spaces

Jeremy Brazas and Paul Fabel

November 28, 2022

Abstract

For a path-connected metric space (X, d) , the n -th homotopy group $\pi_n(X)$ inherits a natural pseudometric from the n -th iterated loop space with the uniform metric. This pseudometric gives $\pi_n(X)$ the structure of a topological group and when X is compact, the induced pseudometric topology is independent of the metric d . In this paper, we study the properties of this pseudometric and how it relates to previously studied structures on $\pi_n(X)$. Our main result is that the pseudometric topology agrees with the shape topology on $\pi_n(X)$ if X is compact and UV^{n-1} or if X is an inverse limit of finite polyhedra with retraction bonding maps.

1 Introduction

There are many ways to enrich the n -th homotopy group $\pi_n(X, x_0)$ of a based topological space (X, x_0) with geometric or topological structure that remembers local features of the space X , which are “unseen” by the usual group-theoretic structure. For example, the natural quotient topology [11], the τ -topology [9], the Spanier topology [1, 3], the whisker topology [2], and variations on these (e.g. coreflections in a Convenient category). Of particular interest is the initial topology on $\pi_n(X, x_0)$ with respect to the canonical homomorphism $\Psi_n : \pi_n(X, x_0) \rightarrow \tilde{\pi}_n(X, x_0)$ to the n -th shape homotopy group. This topology is often referred to as the *shape topology* and we denote the resulting topological group as $\pi_n^{\text{sh}}(X, x_0)$. The shape topology gives $\pi_n^{\text{sh}}(X, x_0)$ the structure of a pro-discrete group and is closely related to Dugundji’s pre-shape-theory approach in [16]. While all of these topologies are well-suited for characterizing various properties in the topological setting, they often forget geometric features determined by a choice of metric.

In this paper¹, we show that a metric space (X, d) with basepoint $x_0 \in X$ naturally induces a pseudometric ρ on $\pi_n(X, x_0)$. While the vast majority of papers on topologized homotopy groups focus on the fundamental group,

¹In 2014, the authors wrote a version of this manuscript which remained unpublished for several years. The authors were motivated update it in 2022 following relevant advances in higher wild homotopy theory.

our results hold in arbitrary dimensions. The resulting pseudometric group $(\pi_n(X, x_0), \rho)$ gives the n -th homotopy group the structure of a topological group, which we denote as $\pi_n^{\text{met}}(X, x_0)$. While ρ depends entirely on d , the topology of $\pi_n^{\text{met}}(X, x_0)$ is independent of the choice of metric d when X is compact. Indeed, examples such as punctured real space $\mathbb{R}^{n+1} \setminus \mathbf{0}$ show that for non-compact metrizable spaces, the topology of $\pi_n^{\text{met}}(X, x_0)$ can vary with d . We establish these results among the basic theory of $\pi_n^{\text{met}}(X, x_0)$ developed in Section 4.

In Section 5, we prove the following comparison with the shape topology, which we view as our main result. Recall that a space X is UV^n if for every open neighborhood U of a point $x \in X$, there is an open neighborhood V of x such that $V \subseteq U$ and such that every map $f : S^k \rightarrow V$ where $0 \leq k \leq n$ is null-homotopic in U . The UV^0 property is equivalent to local path connectedness for metrizable spaces.

Theorem 1.1. *Let (X, d) be a path-connected compact metric space and $n \geq 1$. The topology induced by the natural pseudometric ρ on $\pi_n(X, x_0)$ is finer than the shape topology. Moreover, these two topologies agree in the following two cases:*

1. *if X is UV^{n-1} ,*
2. *if $X = \varprojlim_{j \in \mathbb{N}} (X_j, r_{j+1, j})$ is an inverse limit of finite polyhedra where the bonding maps $r_{j+1, j} : X_{j+1} \rightarrow X_j$ are retractions.*

To prove this result, we show that for a compact metric space (X, d) the pseudometric topology on $\pi_n(X, x_0)$ lies between the shape topology and a third topology called the *Spanier topology* (Definition 5.5). For Case (1), we apply a result from [3] to conclude that, in the UV^{n-1} case, the Spanier and shape topologies (and thus all three topologies) agree. We prove Case (2) separately using the fact that we may alter the metric on X without affecting the topologies of $\pi_n^{\text{met}}(X, x_0)$ or $\pi_n^{\text{sh}}(X, x_0)$.

While Case (2) may appear structurally restrictive, it provides insight to some prominent examples. For example, if X is an infinite shrinking wedge of finite polyhedra X_j , then X may be identified with $\varprojlim_{k \in \mathbb{N}} \bigvee_{j=1}^k X_j$, which has retraction bonding maps (see Example 5.18). While the higher homotopy groups of spaces of the form X are only known in some special cases, Theorem 5.19 implies that $\pi_n^{\text{met}}(X, x_0) \cong \pi_n^{\text{sh}}(X, x_0)$ for all $n \geq 1$.

The upshot of Case (1) of Theorem 1.1 is that in the presence of local $(n-1)$ -connectedness conditions, one can often characterize the pseudometric topology by appealing to shape-theoretic methods, which are likely to be simpler than an analysis of the uniform metric on the n -th loop space. We can modify an example from [15] to show Theorem 1.1 does not hold for $n \geq 1$ without the UV^0 condition (see Example 5.17). The authors do not know of a Peano continuum for which the pseudometric topology is strictly finer than the shape topology. This leaves the following problem.

Problem 1.2. Give an example of a based Peano continuum (X, x_0) for which the topology of $\pi_2^{\text{met}}(X, x_0)$ is strictly finer than that of $\pi_2^{\text{sh}}(X, x_0)$.

2 Preliminaries

Throughout this paper, (X, d) will denote a path-connected metric space with basepoint $x_0 \in X$. The unit interval is denoted $[0, 1]$, the unit n -disk is denoted $D^n = \{\mathbf{x} \in \mathbb{R}^n \mid \|\mathbf{x}\| \leq 1\}$ and $S^{n-1} = \partial D^n$ is the unit $(n-1)$ -sphere. The latter two spaces have basepoint $d_0 = (1, 0, \dots, 0)$. The n -th homotopy group of (X, x_0) is denoted $\pi_n(X, x_0)$. If $f : (X, x_0) \rightarrow (Y, y_0)$ is a based map, then $f_\# : \pi_n(X, x_0) \rightarrow \pi_n(Y, y_0)$ is the induced homomorphism.

Let $\Omega^n(X, x_0)$ be the space of maps $\alpha : ([0, 1]^n, \partial[0, 1]^n) \rightarrow (X, x_0)$ based at x_0 (which we call n -loops) with the metric of uniform convergence. It is well-known that the uniform metric topology agrees with the usual compact-open topology on $\Omega^n(X, x_0)$ (when convenient, we may identify $\Omega^n(X, x_0)$ with the space of based maps $(S^n, d_0) \rightarrow (X, x_0)$ also with the uniform metric). Let $\pi : \Omega^n(X, x_0) \rightarrow \pi_n(X, x_0)$, $\pi(\alpha) = [\alpha]$ denote the canonical surjection taking a map α to its homotopy class.

Given $\alpha \in \Omega^n(X, x_0)$, let $\alpha^-(t_1, t_2, \dots, t_n) = \alpha(1 - t_1, t_2, \dots, t_n)$ denote the reverse of α and if $\alpha_1, \alpha_2, \dots, \alpha_n$ is a sequence of n -loops, then $\alpha_1 \cdot \alpha_2 \cdots \alpha_n$ is the usual n -fold concatenation defined as α_i on $[\frac{i-1}{n}, \frac{i}{n}] \times [0, 1]^{n-1}$. Generally, $c_{x_0} \in \Omega^n(X, x_0)$ will denote the constant map at x_0 so that $e = [c_{x_0}]$ serves as the identity element of $\pi_n(X, x_0)$.

3 Topologies on homotopy groups

Here, we briefly recall some previously studied topologies on the homotopy groups.

3.1 The quotient topology

Let $\pi_n^{\text{qtop}}(X, x_0)$ denote the n -th homotopy group with the quotient topology with respect to the canonical map $\pi : \Omega^n(X, x_0) \rightarrow \pi_n(X, x_0)$, that is, the finest topology on $\pi_n(X, x_0)$ such that π is continuous. In particular $A \subseteq \pi_n^{\text{qtop}}(X, x_0)$ is open (closed) if and only if $\pi^{-1}(A)$ is open (closed) in $\Omega^n(X, x_0)$. It is known that this topology gives $\pi_n(X, x_0)$ the structure of a *quasitopological group* [5] (in the sense that inversion is continuous and multiplication is continuous in each variable) which can fail to be a topological group [8, 19, 20] even when X is a compact metric space. For a general study of π_1 with the quotient topology, we refer the reader to [11]. It is known that if X is locally $(n-1)$ -connected and semilocally n -connected, then $\Omega^n(X, x_0)$ is locally path-connected [25]. In such a situation, $\pi_n^{\text{qtop}}(X, x_0)$ is guaranteed to be discrete (see also [14] in dimension $n = 1$). In particular, if X is a polyhedron, manifold, or CW-complex, then $\pi_n^{\text{qtop}}(X, x_0)$ is a discrete group for all $n \geq 1$.

3.2 The “tau topology”

In [9], it was observed that for any quasitopological group G , there is a finest group topology on the group G , which is coarser than that of G . The resulting topological group is denoted $\tau(G)$. In other words, the category of topological groups is a reflective subcategory of the category of quasitopological groups, where τ is the reflection functor. In the case of fundamental groups, the τ -reflection $\pi_n^\tau(X, x_0) = \tau(\pi_n^{\text{qtop}}(X, x_0))$ is a topological group. The topology of $\pi_n^\tau(X, x_0)$ is coarser than that of $\pi_n^{\text{qtop}}(X, x_0)$ and agrees with that of $\pi_n^{\text{qtop}}(X, x_0)$ if and only if $\pi_n^{\text{qtop}}(X, x_0)$ is a topological group. The τ -topology is the finest *group* topology on $\pi_n(X, x_0)$ for which $\pi : \Omega^n(X, x_0) \rightarrow \pi_n(X, x_0)$ is continuous. In [24], an analogous construction is given for the “universal path space,” i.e. the set of path-homotopy classes of paths in X starting at x_0 .

3.3 The shape topology

We give a few details regarding the construction of the n -th shape homotopy group and refer the reader to [21] for a detailed treatment of shape theory. Let $\text{cov}(X)$ be the directed set of pairs (\mathcal{U}, U_0) where \mathcal{U} is a locally finite open cover of X and U_0 is a distinguished element of \mathcal{U} containing x_0 . Here, $\text{cov}(X)$ is directed by refinement. Given $(\mathcal{U}, U_0) \in \text{cov}(X)$ let $N(\mathcal{U})$ be the abstract simplicial complex which is the nerve of \mathcal{U} . In particular, \mathcal{U} is the vertex set of $N(\mathcal{U})$ and the n vertices U_1, \dots, U_n span an n -simplex $\Leftrightarrow \bigcap_{i=1}^n U_i \neq \emptyset$. The geometric realization $|N(\mathcal{U})|$ is a polyhedron and thus $\pi_n(|N(\mathcal{U})|, U_0)$ may be regarded naturally as a discrete group, e.g. if it is given the quotient topology.

Given a pair (\mathcal{V}, V_0) which refines (\mathcal{U}, U_0) , a simplicial map $p_{\mathcal{U}\mathcal{V}} : |N(\mathcal{V})| \rightarrow |N(\mathcal{U})|$ is constructed by sending a vertex $V \in \mathcal{V}$ to some $U \in \mathcal{U}$ for which $V \subseteq U$ (in particular, V_0 is mapped to U_0) and extending linearly. The map $p_{\mathcal{U}\mathcal{V}}$ is unique up to homotopy and thus induces a unique homomorphism $p_{\mathcal{U}\mathcal{V}\#} : \pi_n(|N(\mathcal{V})|, V_0) \rightarrow \pi_n(|N(\mathcal{U})|, U_0)$. The inverse system

$$(\pi_n(|N(\mathcal{U})|, U_0), p_{\mathcal{U}\mathcal{V}\#}, \text{cov}(X))$$

of discrete groups is the *fundamental pro-group* and the limit $\tilde{\pi}_n(X, x_0)$ (topologized with the usual inverse limit topology) is the *n -th shape homotopy group*.

Given a partition of unity $\{\phi_U\}_{U \in \mathcal{U}}$ subordinated to \mathcal{U} and such that $\phi_{U_0}(x_0) = 1$, a map $p_{\mathcal{U}} : X \rightarrow |N(\mathcal{U})|$ is constructed by taking $\phi_U(x)$ (for $x \in U$, $U \in \mathcal{U}$) to be the barycentric coordinate of $p_{\mathcal{U}}(x)$ corresponding to the vertex U . The induced continuous homomorphism $p_{\mathcal{U}\#} : \pi_n(X, x_0) \rightarrow \pi_n(|N(\mathcal{U})|, U_0)$ satisfies $p_{\mathcal{U}\#} \circ p_{\mathcal{U}\mathcal{V}\#} = p_{\mathcal{V}\#}$ whenever (\mathcal{V}, V_0) refines (\mathcal{U}, U_0) . Thus there is a canonical, continuous homomorphism $\Psi_n : \pi_n(X, x_0) \rightarrow \tilde{\pi}_n(X, x_0)$ to the n -th shape homotopy group, given by $\Psi_n([\alpha]) = ([p_{\mathcal{U}} \circ \alpha]_{\mathcal{U}})$.

Definition 3.1. The *shape topology* on $\pi_n(X, x_0)$ is the initial topology with respect to the first shape homomorphism $\Psi_n : \pi_n(X, x_0) \rightarrow \tilde{\pi}_n(X, x_0)$. Let $\pi_n^{\text{sh}}(X, x_0)$ denote $\pi_n(X, x_0)$ equipped with the shape topology.

The shape topology is characterized as follows: $A \subset \pi_n^{\text{sh}}(X, x_0)$ is open (resp. closed) $\Leftrightarrow A = \Psi_n^{-1}(B)$ for an open (resp. closed) set $B \subset \tilde{\pi}_n(X, x_0)$. Equivalently, a set $A \subset \pi_n^{\text{sh}}(X, x_0)$ containing the identity element is open \Leftrightarrow there exists $(\mathcal{U}, U_0) \in \text{cov}(X)$ such that $\ker(p_{\mathcal{U}\#} : \pi_n(X, x_0) \rightarrow \pi_n(|N(\mathcal{U})|, U_0))$ lies in A . Since $\tilde{\pi}_n(X, x_0)$ is a topological group, it follows immediately that $\pi_n^{\text{sh}}(X, x_0)$ is a topological group.

Proposition 3.2. [9, 3.24] For any space X , the shape topology of $\pi_n^{\text{sh}}(X, x_0)$ is coarser than that of $\pi_n^\tau(X, x_0)$.

Definition 3.3. We say a space X is π_n -shape injective if $\Psi_n : \pi_n(X, x_0) \rightarrow \tilde{\pi}_n(X, x_0)$ is a monomorphism.

By construction, $\pi_n^{\text{sh}}(X, x_0)$ is Hausdorff $\Leftrightarrow X$ is π_n -shape injective.

4 A pseudometric on homotopy groups

Let (X, d) be a path-connected metric space and consider the uniform metric

$$\mu(\alpha, \beta) = \sup_{t \in [0, 1]^n} \{d(\alpha(t), \beta(t))\}$$

on $\Omega^n(X, x_0)$. Observe that

$$\mu(\alpha \cdot \alpha', \beta \cdot \beta') = \max\{\mu(\alpha, \beta), \mu(\alpha', \beta')\}$$

and $\mu(\alpha, \beta) = \mu(\alpha^-, \beta^-)$. We consider the following function $\rho : \pi_n(X, x_0) \times \pi_n(X, x_0) \rightarrow [0, \infty)$ on the homotopy group $\pi_n(X, x_0)$:

$$\rho(a, b) = \inf\{\mu(\alpha, \beta) \mid \alpha \in a, \beta \in b\}.$$

We claim that ρ is a pseudometric on $\pi_n(X, x_0)$. Certainly ρ is symmetric and $\rho(a, a) = 0$; however, a little more work is required to verify the triangle inequality.

Remark 4.1. In general, if (X, d) is a metric space, \sim is an equivalence relation, and $Y = X/\sim$, then the definition $\rho : Y \times Y \rightarrow [0, \infty)$, $\rho(a, b) = \inf\{d(\alpha, \beta) \mid \alpha \in a, \beta \in b\}$ need not satisfy the triangle inequality. Thus in our situation, we must make use of the group structure of $\pi_n(X, x_0)$ and the nature of the uniform metric μ in order to verify the triangle inequality.

Lemma 4.2 (Isometric Inversion). For all $a, b \in \pi_n(X, x_0)$, $\rho(a, b) = \rho(a^{-1}, b^{-1})$.

Proof. Since $\mu(\alpha, \beta) = \mu(\alpha^-, \beta^-)$ for all $\alpha \in a, \beta \in b$, it is clear that

$$\rho(a, b) = \rho([\alpha^-], [\beta^-]) = \rho(a^{-1}, b^{-1}).$$

□

Lemma 4.3 (Isometric translations). Fix $c \in \pi_n(X, x_0)$. Then for all $a, b \in \pi_n(X, x_0)$, we have $\rho(a, b) = \rho(ac, bc) = \rho(ca, cb)$.

Proof. Fix $c = [\gamma] \in \pi_n(X, x_0)$. Let $a, b \in \pi_n(X, x_0)$. For $\alpha \in a$ and $\beta \in b$, we have $\mu(\alpha \cdot \gamma, \beta \cdot \gamma) = \mu(\alpha, \beta)$. It follows that

$$\rho(a, b) = \rho([\alpha \cdot \gamma], [\beta \cdot \gamma]) = \rho(ac, bc).$$

The symmetric argument gives $\rho(a, b) = \rho(ca, cb)$. \square

Lemma 4.4. For all $a, b \in \pi_n(X, x_0)$, we have $\rho(ab, e) \leq \max\{\rho(a, e), \rho(b, e)\}$

Proof. Suppose $a, b \in \pi_n(X, x_0)$ and $\epsilon > 0$. Find $\alpha \in a$, $\beta \in b$, and $\gamma_1, \gamma_2 \in e = [c_{x_0}]$ such that $\mu(\alpha, \gamma_1) < \rho(a, e) + \frac{\epsilon}{2}$ and $\mu(\beta, \gamma_2) < \rho(b, e) + \frac{\epsilon}{2}$. Note that

$$\begin{aligned} \mu(\alpha \cdot \beta, \gamma_1 \cdot \gamma_2) &= \max\{\mu(\alpha, \gamma_1), \mu(\beta, \gamma_2)\} \\ &< \max\left\{\rho(a, e) + \frac{\epsilon}{2}, \rho(b, e) + \frac{\epsilon}{2}\right\} \\ &< \max\{\rho(a, e), \rho(b, e)\} + \epsilon \end{aligned}$$

Thus $\rho(ab, e) \leq \max\{\rho(a, e), \rho(b, e)\}$. \square

Theorem 4.5. ρ is a pseudometric on $\pi_n(X, x_0)$.

Proof. As noted above, it suffices to verify the triangle inequality. Let $a, b, c \in \pi_n(X, x_0)$. Using the previous three lemmas, we have:

$$\begin{aligned} \rho(a, c) &= \rho(ac^{-1}, e) \\ &= \rho(ab^{-1}bc, e) \\ &\leq \max\{\rho(ab^{-1}, e), \rho(bc^{-1}, e)\} \\ &= \max\{\rho(ab^{-1}, e), \rho(cb^{-1}, e)\} \\ &\leq \rho(ab^{-1}, e) + \rho(cb^{-1}, e) \\ &= \rho(a, b) + \rho(c, b) \\ &= \rho(a, b) + \rho(b, c) \end{aligned}$$

\square

Proposition 4.6. Equipped with the topology induced by the pseudometric ρ , $\pi_n(X, x_0)$ is a topological group whose open balls $B_\rho(e, r)$, $r > 0$ are open normal subgroups.

Proof. Since the open balls $B_\rho(e, r)$ form a neighborhood base at e and translations are homeomorphisms (Lemma 4.3), it will follow that $\pi_n(X, x_0)$ is a topological group once we show that $B_\rho(e, r)$ is an open normal subgroup.

Since $\rho(a, e) = \rho(a^{-1}, e)$, $B_\rho(e, r)$ is closed under inversion. Additionally, by Lemma 4.3, we have

$$\rho(cac^{-1}, e) = \rho(a, c^{-1}c) = \rho(a, e)$$

for any $c \in \pi_n(X, x_0)$. Thus $B_\rho(e, r)$ is closed under conjugation (particularly when $n = 1$). Finally, if $\rho(a, e) < r$ and $\rho(b, e) < r$, then $\rho(ab, e) \leq \max\{\rho(a, e), \rho(b, e)\} < r$ by Lemma 4.4 and it follows that $B_\rho(e, r)$ is closed under multiplication. \square

Definition 4.7. For a metric space (X, d) and $x_0 \in X$, let $\pi_n^{\text{met}}(X, x_0)$ denote the n -th homotopy group equipped with the topology induced by the pseudometric ρ . We call this topology the *pseudometric topology (induced by d)*.

Since $\pi_n^{\text{met}}(X, x_0)$ is a topological group, open subgroups of $\pi_n^{\text{met}}(X, x_0)$ are also closed. Therefore, $\pi_n^{\text{met}}(X, x_0)$ is zero-dimensional. On the other hand, $\pi_n^{\text{met}}(X, x_0)$ need not be Hausdorff, since the closed normal subgroup $\bigcap_{r>0} B_\rho(e, r) = \{[\alpha] \in \pi_1(X, x_0) \mid \rho([\alpha], e) = 0\}$, is equal to $\overline{\{e\}}$, the closure of the identity element, which may be non-trivial. In particular, $\overline{\{e\}}$ is non-empty if and only if there exist sequences $\{\alpha_k\}_{k \in \mathbb{N}}$ and $\{\beta_k\}_{k \in \mathbb{N}}$ in $\Omega^n(X, x_0)$ such that $[\alpha_k] = [\alpha_{k+1}]$, $[\beta_k] = [\beta_{k+1}]$ and $\lim_{k \rightarrow \infty} \mu(\alpha_k, \beta_k) = 0$.

Proposition 4.8. If $f : (X, x_0) \rightarrow (Y, y_0)$ is a uniformly continuous map, then the induced homomorphism $f_\# : \pi_n^{\text{met}}(X, x_0) \rightarrow \pi_n^{\text{met}}(Y, y_0)$ is uniformly continuous.

Proof. Suppose $\epsilon > 0$. There is a $\delta > 0$ such that $\mu(\alpha, \beta) < \delta \Rightarrow \mu(f \circ \alpha, f \circ \beta) < \epsilon/2$. Suppose $\rho(g, h) < \delta$ for $g, h \in \pi_n^{\text{met}}(X, x_0)$. There are $\alpha \in g$, $\beta \in h$ such that $\mu(\alpha, \beta) < \delta$. Thus $\mu(f \circ \alpha, f \circ \beta) < \epsilon/2$. It follows that $\rho(f_\#(g), f_\#(h)) = \rho([f \circ \alpha], [f \circ \beta]) < \epsilon$. \square

Corollary 4.9. If X is compact, then the topology of $\pi_n^{\text{met}}(X, x_0)$ induced by ρ is independent of the choice of metric on X .

Proof. Suppose metrics d_1 and d_2 on X induce the same topology. Let ρ_1 and ρ_2 be the respective pseudometrics on $\pi_1^{\text{top}}(X, x_0)$. Since X is compact, the identity maps $id : (X, d_1) \rightarrow (X, d_2)$ and $id : (X, d_2) \rightarrow (X, d_1)$ are uniformly continuous. By Proposition 4.8, the induced homomorphisms on $\pi_n(X, x_0)$ (with the respective pseudometrics) are inverse isomorphisms. \square

Next, we observe that, in general, the isomorphism class of the topological group $\pi_n^{\text{met}}(X, x_0)$ does not depend on the choice of basepoint. Fix a retraction $r : S^n \times [0, 1] \rightarrow S^n \times \{0\} \cup \{d_0\} \times [0, 1]$. For any path $\gamma : [0, 1] \rightarrow X$ and map $\alpha : (S^n, d_0) \rightarrow (X, \gamma(1))$, we define the *path-conjugate* $\gamma * \alpha : (S^n, d_0) \rightarrow (X, \gamma(0))$ to be the composition of $r(d, 0) : S^n \rightarrow S^n \times \{0\} \cup \{d_0\} \times [0, 1]$ followed by applying γ to $\{d_0\} \times [0, 1]$ and α to $S^n \times \{0\}$. This defines a natural map $\Omega^n(X, \gamma(1)) \rightarrow \Omega^n(X, \gamma(0))$, $\beta \mapsto \gamma * \beta$ which induces a change-of-basepoint isomorphism $\Gamma : \pi_n(X, \gamma(1)) \rightarrow \pi_n(X, \gamma(0))$. Moreover, when γ is a loop, the action of loops on n -loops defines a jointly continuous map $\Omega(X, x_0) \times \Omega^n(X, x_0) \rightarrow \Omega^n(X, x_0)$, $(\gamma, \alpha) \mapsto \gamma * \alpha$. This map induces the π_1 -action $[\gamma] * [\alpha] = [\gamma * \alpha]$ on $\pi_n(X, x_0)$.

Proposition 4.10. For any path $\gamma : [0, 1] \rightarrow X$, the group isomorphism

$$\Gamma : \pi_n^{\text{met}}(X, \gamma(1)) \rightarrow \pi_n^{\text{met}}(X, \gamma(0)),$$

$\Gamma([\beta]) = [\gamma * \beta]$ is an isometry.

Proof. For all $\alpha, \beta \in \Omega^n(X, \gamma(1))$, we have $\mu(\alpha, \beta) = \mu(\gamma * \alpha, \gamma * \beta)$ and thus

$$\rho([\gamma * \alpha], [\gamma * \beta]) \leq \rho([\alpha], [\beta]).$$

Thus Γ is non-expansive. The inverse $\Gamma^{-1} : \pi_1^{\text{met}}(X, \gamma(1)) \rightarrow \pi_1^{\text{met}}(X, \gamma(0))$, $\Gamma^{-1}([\beta]) = [\gamma^{-1} * \beta]$ is non-expansive for the same reason (replacing γ with γ^{-1}). Thus Γ is an isometry. \square

Theorem 4.11. *The π_1 -action $\pi_1^{\text{met}}(X, x_0) \times \pi_n^{\text{met}}(X, x_0) \rightarrow \pi_n^{\text{met}}(X, x_0)$, $([\gamma], [\alpha]) \mapsto [\gamma * \alpha]$ is jointly continuous. Moreover, $\pi_1^{\text{met}}(X, x_0)$ acts on $\pi_n^{\text{met}}(X, x_0)$ by isometry.*

Proof. That the action is by isometry follows immediately from Proposition 4.10. For continuity, suppose $\{g_k\}_{k \in \mathbb{N}} \rightarrow g$ in $\pi_1^{\text{met}}(X, x_0)$ and $\{a_k\}_{k \in \mathbb{N}} \rightarrow a$ in $\pi_n^{\text{met}}(X, x_0)$. Let $\epsilon > 0$. Find $K \in \mathbb{N}$ such that $\max\{\rho(g_k, g), \rho(a_k, a)\} < \epsilon/2$ for all $k \geq K$. Thus for $k \geq K$, there exist $\gamma_k \in g_k$ and $\delta_k \in g$ with $\mu(\gamma_k, \delta_k) < \epsilon/2$ and $\alpha_k \in a_k$ and $\beta_k \in a$ with $\mu(\alpha_k, \beta_k) < \epsilon/2$. Our definition of the action $*$ of loops on n -loops using a fixed retraction r ensures that $\mu(\gamma_k * \alpha_k, \delta_k * \beta_k) < \epsilon/2$. Since $g_k * a_k = [\gamma_k * \alpha_k]$ and $g * a = [\delta_k * \beta_k]$ for all $k \geq K$, we have $\rho(g_k * a_k, g * a) < \epsilon$. We conclude that $\{g_k * a_k\}_{k \in \mathbb{N}} \rightarrow g * a$ in $\pi_n^{\text{met}}(X, x_0)$. \square

Finally, we compare the pseudometric topology to the quotient and τ -topologies.

Proposition 4.12. The function $\pi : \Omega^n(X, x_0) \rightarrow \pi_n^{\text{met}}(X, x_0)$, $\pi(\alpha) = [\alpha]$ is continuous. Thus the topology of both $\pi_n^{\text{qtop}}(X, x_0)$ and $\pi_n^\tau(X, x_0)$ is finer than that of $\pi_n^{\text{met}}(X, x_0)$.

Proof. Suppose $\{\alpha_k\}_{k \in \mathbb{N}} \rightarrow \alpha$ in $\Omega^n(X, x_0)$ and $\epsilon > 0$. There is an $K \in \mathbb{N}$ such that $\mu(\alpha, \alpha_k) < \epsilon/2$ for $k \geq K$. Thus $\rho([\alpha], [\alpha_k]) < \epsilon$ for $k \geq K$ showing that $\{[\alpha_k]\}_{k \in \mathbb{N}} \rightarrow [\alpha]$ in $\pi_n^{\text{met}}(X, x_0)$. Thus π is continuous. The second statement follows directly from the characterizations of the quotient and τ -topologies in Section 3. \square

Remark 4.13. Although basepoint-change isomorphisms are continuous when π_n is given the quotient and τ -topologies, the π_1 -action $\pi_1(X, x) \times \pi_n(X, x) \rightarrow \pi_n(X, x)$ is discontinuous when π_1 and π_n are given the quotient topology [10]. Apparently, it is unknown if the π_1 -action $\pi_1(X, x) \times \pi_n(X, x) \rightarrow \pi_n(X, x)$ is continuous in the τ -topology.

The following example illustrates that when X is non-compact the pseudometric on $\pi_n^{\text{met}}(X, x_0)$ may vary with our choice of metric on X .

Example 4.14. Consider the cylinder $X = \mathbb{R} \times S^n$ with basepoint $x_0 = (1, b)$. It is easy to see that with the natural product metric d_1 , the resulting pseudometric ρ_1 on $\pi_n(\mathbb{R} \times S^n, x_0)$ is discrete. For instance, $\pi_n^{\text{met}}(S^n, (1, 0))$ is discrete and we may then apply Proposition 4.8 to the projection (and homotopy equivalence) $\mathbb{R} \times S^n \rightarrow S^n$.

On the other hand, we may give $\mathbb{R} \times S^n$ the metric d_2 of the homeomorphic punctured plane $\mathbb{R}^{n+1} \setminus \{\mathbf{0}\}$. Let $x_0 = (1, 0, 0, \dots, 0) \in \mathbb{R}^{n+1} \setminus \{\mathbf{0}\}$ be

the basepoint and ρ_2 denote the resulting pseudometric on $\pi_n(X, x_0)$. Take $\gamma_n : [0, 1] \rightarrow X$ to be the linear path from x_0 to $(1/n, 0, \dots, 0)$ and let α_n be the embedding of the n -sphere of radius $1/n$ centered at the origin. Now consider the path-conjugates $\gamma_n * \alpha_n$, all of which represent a generator g of $\pi_n(X, x_0) \cong \mathbb{Z}$. We also consider the null-homotopic maps $\gamma_n * c_n$ where $c_n \in \Omega^n(X, \gamma_n(1))$ is the constant map. But

$$\lim_{n \rightarrow \infty} \mu(\gamma_n * c_n, \gamma_n * \alpha_n) = 0.$$

Thus $\rho_2(g, 1) = 0$ and it follows that the resulting pseudometric group $\pi_n^{\text{met}}(X, x_0)$ is indiscrete.

5 Comparison with the Shape Topology

Lemma 5.1. If K is a finite polyhedron and $n \geq 1$, then $\pi_n^{\text{sh}}(K, x_0)$ is discrete.

Proof. If the compact metric space (K, d) is the underlying space of a finite simplicial complex, then K admits a topologically compatible $CAT(1)$ metric d' [13, Corollary 5.19]. Consequently there exists $\epsilon > 0$ so that for any space Y , if the maps $f : Y \rightarrow K$ and $g : Y \rightarrow K$ are uniformly close with $d(f(y), g(y)) < \epsilon$ for all $y \in Y$, then f and g are canonically homotopic via a homomotopy H . The homotopy H maps $f(y)$ to $g(y)$ linearly with time, along the unique geodesic $[f(y), g(y)]$ in (K, d') . The existence and continuity of H follows from Proposition 1.4 and Corollary 3.13 of [13]. Thus $\pi_n^{\text{met}}(K, x_0)$ is discrete with the induced pseudometric generated by d' . Since K is compact, Corollary 4.9 gives that the topology of $\pi_n^{\text{met}}(K, x_0)$ is independent of the choice of metric on K . Hence, $\pi_n^{\text{met}}(K, x_0)$ is discrete with the induced pseudometric generated by d . \square

Proposition 5.2. If X is a compact metric space, then the topology of $\pi_n^{\text{met}}(X, x_0)$ is finer than that of $\pi_n^{\text{sh}}(X, x_0)$.

Proof. Since X is compact, we may construct $\tilde{\pi}_n(X, x_0)$ by replacing $\text{cov}(X)$ with a cofinal sequence of finite covers. Whenever $\mathcal{U} \in \text{cov}(X)$ is finite, $|N(\mathcal{U})|$ is a finite polyhedron and thus $\pi_n^{\text{met}}(|N(\mathcal{U})|, U_0)$ is discrete by Lemma 5.1. Since X is compact, any canonical map $p_{\mathcal{U}} : X \rightarrow |N(\mathcal{U})|$ will be uniformly continuous and thus the induced homomorphism $p_{\mathcal{U}\#} : \pi_n^{\text{met}}(X, x_0) \rightarrow \pi_n^{\text{met}}(|N(\mathcal{U})|, U_0)$ will be continuous by Proposition 4.8. Since $\pi_n^{\text{sh}}(X, x_0)$ has the initial topology with respect to homomorphisms $p_{\mathcal{U}\#}$ with discrete codomains, it follows that the topology of $\pi_n^{\text{met}}(X, x_0)$ is finer than that of $\pi_n^{\text{sh}}(X, x_0)$. \square

Remark 5.3. The previous proposition fails when X is no longer required to be compact. If $X = \mathbb{R}^{n+1} \setminus \{\mathbf{0}\}$ is punctured real $(n+1)$ -space as in Example 4.14, then $\pi_n^{\text{met}}(X, x_0)$ is isomorphic to \mathbb{Z} with the indiscrete topology whereas $\pi_n^{\text{sh}}(X, x_0)$ is isomorphic to \mathbb{Z} with the discrete topology.

Definition 5.4. Let $n \geq 1$ and \mathcal{U} be an open cover of X . The n -th Spanier group of (X, x_0) with respect to \mathcal{U} is the subgroup $\pi_n^{\text{SP}}(\mathcal{U}, x_0)$ of $\pi_n(X, x_0)$ generated by path-conjugates $[\gamma * f] \in \pi_n(X, x_0)$ where $\gamma : ([0, 1], 0) \rightarrow (X, x_0)$ is a path and $\alpha : (S^n, d_0) \rightarrow (X, \gamma(1))$ has image in U for some $U \in \mathcal{U}$. The n -th Spanier group of (X, x_0) is the intersection $\pi_n^{\text{SP}}(X, x_0) = \bigcap_{\mathcal{U} \in \text{cov}(X)} \pi_n^{\text{SP}}(\mathcal{U}, x_0)$.

Spanier groups (in dimension $n = 1$) first appeared in [23] and have been used frequently in the past decade. Higher Spanier groups were more recently introduced [4]. Note that since Spanier groups $\pi_n^{\text{SP}}(\mathcal{U}, x_0)$, $\text{cov}(X)$ are always normal subgroups of $\pi_n(X, x_0)$, the sets of left cosets $\{[g]\pi_n^{\text{SP}}(\mathcal{U}, x_0) \mid \mathcal{U} \in \text{cov}(X), [g] \in \pi_n(X, x_0)\}$ forms a basis for a topology on $\pi_n(X, x_0)$.

Definition 5.5. We refer to the topology on $\pi_n(Y, y_0)$ generated by cosets of the Spanier groups $\pi_n^{\text{SP}}(\mathcal{U}, y_0)$ as the *Spanier topology*.

In the current paper, the Spanier topology will not be studied in detail but will only serve as a convenient bound for the pseudometric topology. In order to make these comparisons, we recall known results that compare Spanier groups with homomorphisms $p_{\mathcal{U}\#} : \pi_n(X, x_0) \rightarrow \pi_n(|N(\mathcal{U})|, U_0)$ induced by canonical maps. The following has a straightforward proof. See [4, Prop 4.13] or [3, Lemma 3.10].

Proposition 5.6. For every $\mathcal{U} \in \text{cov}(X)$ and choice of canonical map $p_{\mathcal{U}} : X \rightarrow |N(\mathcal{U})|$, there exists a $\mathcal{V} \in \text{cov}(X)$ such that $\pi_n^{\text{SP}}(\mathcal{V}, x_0) \leq \ker(p_{\mathcal{U}\#} : \pi_n(X, x_0) \rightarrow \pi_n(|N(\mathcal{U})|, U_0))$.

Since the shape topology on $\pi_n(X, x_0)$ is generated by left cosets of subgroups of the form $\ker(p_{\mathcal{U}\#})$, we have the following.

Corollary 5.7. [3, Remark 5.5] For any metrizable space X , Spanier topology on $\pi_n(X, x_0)$ is finer than the shape topology.

Determining when the Spanier topology agrees with the shape topology is a more technical matter that has been addressed in [3]. To state this properly, we recall the following definition.

Definition 5.8. [22] Let $n \geq 0$. A topological space Y is UV^n at $y \in Y$ if for every neighborhood U of y , there exists a neighborhood V of y such that $V \subseteq U$ and such that for all $0 \leq k \leq n$, every map $f : S^k \rightarrow V$ extends to a map $g : D^{k+1} \rightarrow U$. We say Y is UV^n if Y is UV^n at all of its points.

Remark 5.9. In some of the Shape Theory literature, the UV^n property is called the “ LC^n property.” For example, see [21, p. 40]. This is easily confused with local n -connectedness in which one has a basis of n -connected open sets. Since the two are not equivalent even for Peano continua, we prefer the “ UV^n ” terminology. Note that if Y is locally n -connected at $y \in Y$, then Y is UV^n at y .

The following lemma was proved in dimension $n = 1$ in [12]. For $n \geq 2$, the proof requires other techniques from shape theory.

Lemma 5.10. [3, Lemma 5.1] Suppose X is UV^{n-1} . Then for every $\mathcal{U} \in \text{cov}(X)$, there exists $\mathcal{V} \in \text{cov}(X)$ such that for any canonical map $p_{\mathcal{V}} : X \rightarrow |N(\mathcal{V})|$, we have $\ker(p_{\mathcal{V}\#} : \pi_n(X, x_0) \rightarrow \pi_n(N(\mathcal{V}), V_0)) \leq \pi_n^{\text{Sp}}(\mathcal{U}, x_0)$.

Corollary 5.7 and Lemma 5.10 now combine to give the following.

Corollary 5.11. If X is a UV^{n-1} metrizable space, then the Spanier and shape topologies on $\pi_n(X, x_0)$ agree.

Next, we compare the Spanier topology and pseudometric topologies by comparing neighborhood bases at the identity element.

Proposition 5.12. Let (X, d) be a path-connected metric space. Then for every $r > 0$, there exists $\mathcal{U} \in \text{cov}(X)$ such that $\pi_n^{\text{Sp}}(\mathcal{U}, x_0) \leq B_{\rho}(e, r)$.

Proof. Given $r > 0$, let $\mathcal{U} = \{B_d(x, r/2) \mid x \in X\}$ be the cover of X by $r/2$ -balls. Consider a generator $[\alpha * f] \in \pi_n^{\text{Sp}}(\mathcal{U}, x_0)$ where $\alpha : ([0, 1], 0) \rightarrow (X, x_0)$ is a path and $f : (S^n, d_0) \rightarrow (X, \alpha(1))$ is a map with image in $B_d(x, r/2)$ for some $x \in X$. Let $g : S^n \rightarrow X$ be the constant map at $\alpha(1)$. Since $e = [\alpha * g]$ and $\mu(\alpha * f, \alpha * g) < r$, we have $\rho([\alpha * f], e) < r$, giving $[\alpha * f] \in B_{\rho}(e, r)$. \square

Corollary 5.13. For a metric space (X, d) , the Spanier topology on $\pi_n(X, x_0)$ is finer than the pseudometric topology induced by d .

At this point, we have sufficient results to prove Case (1) of Theorem 1.1. The following lemma addresses Case (2).

Lemma 5.14. If (X, d) is homeomorphic to an inverse limit $\varprojlim_{j \in \mathbb{N}} (X_j, r_{j+1, j})$ of finite polyhedra X_j where the bonding maps $r_{j+1, j} : X_{j+1} \rightarrow X_j$ are retractions, then for all $n \geq 1$, the shape topology on $\pi_n(X, x_0)$ is finer than the pseudometric topology.

Proof. Identify $X = \varprojlim_{j \in \mathbb{N}} (X_j, r_{j+1, j})$ and let $r_j : X \rightarrow X_j$, $j \in \mathbb{N}$ denote the projection maps. Let $s_{j, j+1} : X_j \rightarrow X_{j+1}$ be a section to $r_{j+1, j}$. When $i > j$, $s_{j, i} = s_{i-1, i} \circ s_{i-2, i-1} \circ \cdots \circ s_{j, j+1}$ is a section to $r_{i, j} = r_{j, j+1} \circ \cdots \circ r_{i-2, i-2} \circ r_{i-1, i-1}$. For fixed all pairs $i, j \in \mathbb{N}$, let $t_{j, i} : X_j \rightarrow X_i$ be $r_{j, i}$ if $j > i$, id_{X_j} if $i = j$, and $s_{j, i}$ if $j < i$. Then the maps $\{t_{j, i}\}_{i \in \mathbb{N}}$ induce a map $t_j : X_j \rightarrow X$ which is a section to r_j . Thus we may identify $X_1 \subseteq X_2 \subseteq X_3 \subseteq \cdots$ as a nested sequence of closed subspaces of X where $\overline{\bigcup_{j \in \mathbb{N}} X_j} = X$. In particular X_k consists of the sequences $(x_j)_{j \in \mathbb{N}} \in X$ where $x_j = x_k$ for all $j \geq k$ and $r_k((x_j)_{j \in \mathbb{N}}) = (x_1, x_2, \dots, x_{k-1}, x_k, x_k, x_k, \dots)$ for all $(x_j)_{j \in \mathbb{N}} \in X$.

Pick basepoint $x_0 \in X_1$ and fix $n \geq 1$. Let $\delta > 0$ so that $B_{\rho}(e, \delta)$ is a basic neighborhood of the identity in $\pi_n^{\text{met}}(X, x_0)$. Since X is compact, the sequence $\{X_j\}_{j \in \mathbb{N}}$ gives an **HPol**-expansion of X and therefore, the subgroups $H_j = \ker(r_{j\#} : \pi_n(X, x_0) \rightarrow \pi_n(X_j, x_0))$ form a neighborhood base at the identity in $\pi_n^{\text{sh}}(X, x_0)$. To show that the shape topology is finer than the pseudometric topology it suffices to show that there exists $k \in \mathbb{N}$ such that $H_k \subseteq B_{\rho}(e, \delta)$.

First, we claim that there exists $k \in \mathbb{N}$ such that $d(x, r_k(x)) < \frac{\delta}{2}$ for all $x \in X$. By Corollary 4.9, we may alter the metric d on X without changing the

topology of $\pi_n^{\text{met}}(X, x_0)$. Since $\pi_n^{\text{sh}}(X, x_0)$ is a topological invariant, doing so will not change the topology of $\pi_n^{\text{sh}}(X, x_0)$ either. In particular, let d_j be a metric inducing the topology of X_j that is bounded by 1. For elements $x = (x_1, x_2, \dots)$ and $y = (y_1, y_2, \dots)$ of $\prod_{j=1}^{\infty} X_j$, the formula $d(x, y) = \sum_{j=1}^{\infty} \frac{d_j(x_j, y_j)}{2^j}$ defines a metric that induces the topology of $\prod_{j=1}^{\infty} X_j$ with the product topology [18, 4.2.2]. Since X is topologized a subspace of $\prod_{j=1}^{\infty} X_j$, $d(x, y)$ restricts to a metric that induces the topology of X . Find $k \in \mathbb{N}$ such that $\sum_{j=k+1}^{\infty} \frac{1}{2^j} < \frac{\delta}{2}$. Then, for any $x = (x_1, x_2, \dots) \in X$, we have

$$d(x, r_k(x)) = \sum_{j=k+1}^{\infty} \frac{d_j(x_j, x_k)}{2^j} \leq \sum_{j=k+1}^{\infty} \frac{1}{2^j} < \frac{\delta}{2}.$$

This proves the desired claim.

Finally, we check that $H_k \subseteq B_\rho(e, \delta)$. Given $a \in H_k$, find an n -loop $\alpha \in a$. Then $\beta = r_k \circ \alpha$ is null-homotopic in X_k and since $X_k \subseteq X$, β is null-homotopic in X . Moreover, our choice of k ensures that $d(\alpha(\mathbf{t}), \beta(\mathbf{t})) = d(\alpha(\mathbf{t}), r_k(\alpha(\mathbf{t}))) < \frac{\delta}{2}$ for all $\mathbf{t} \in [0, 1]^n$. Thus $\mu(\alpha, \beta) < \frac{\delta}{2}$. It follows that $\rho(a, e) < \delta$, giving $a \in B_\rho(e, \delta)$. The inclusion $H_J \subseteq B_\rho(e, \delta)$ follows. \square

At this point, we have established all results needed to prove Theorem 1.1.

Proof of Theorem 1.1. Suppose (X, d) is a compact metric space. By Proposition 5.2, the pseudometric topology is finer than the shape topology (note that this is where compactness is required). To prove equivalence of the two topologies in Case (1), we assume X is UV^{n-1} . By Corollary 5.13, the Spanier topology is finer than the pseudometric topology. Thus, we have the following situation:

$$\text{shape topology} \subseteq \text{pseudometric topology} \subseteq \text{Spanier topology}.$$

Finally, Corollary 5.11 implies that the Spanier topology agrees with the shape topology. Thus the pseudometric topology agrees with the shape topology. Equivalence in Case (2) follows directly from Lemma 5.14. \square

Remark 5.15. The UV^{n-1} condition in Case (1) of Theorem 1.1 implies the well-known n -movability condition from shape theory [21, II§8.1, Theorem 6]. Moreover, the hypothesis in Case (2) of Theorem 1.1 directly implies that X is n -movable. However, Theorem 1.1 does not appear to generalize readily to n -movable spaces. Indeed, the main structure of interest is the homotopy group $\pi_n(X, x_0)$ itself, rather than an $H\text{Pol}_*$ -expansion or the shape-type of X . In general, movability conditions allow one to lift maps inductively along an inverse system. However, for the standard shape category these lifts are only lifts up to homotopy. One must be able to obtain strict lifts in order to induce a map to the limit.

Next, we consider an example where the equivalence of the shape topology and pseudometric topology allows one to characterize the pseudometric topology by appealing to the simpler shape-theoretic setting.

Example 5.16 (Earring mapping tori). Fix $n \geq 2$. The n -dimensional infinite earring space is $\mathbb{E}_n = \bigcup_{k \in \mathbb{N}} C_k$ where $C_k \subseteq \mathbb{R}^{n+1}$ is the n -sphere of radius $1/k$ centered at $(1/k, 0, 0, \dots, 0)$ [17]. Sometimes this space is referred to as the *Barratt-Milnor Spheres* [6]. Let $\ell_k : S^k \rightarrow C_k$ denote the canonical homeomorphism based at the origin $x_0 = (0, 0, \dots, 0)$. Let $f : \mathbb{E}_n \rightarrow \mathbb{E}_n$ be the shift map given by $f \circ \ell_k = \ell_{k+1}$ for each $k \in \mathbb{N}$ and $T_f = \mathbb{E}_n \times [0, 1] / \sim$, $(x, 0) \sim (f(x), 1)$ be the mapping torus of f . We identify \mathbb{E}_n with the image of $\mathbb{E}_n \times \{0\}$ in T_f . Let $\alpha : [0, 1] \rightarrow T_f$ be the path where $\alpha(t)$ is the image of (x_0, t) . In [3], it is shown that \mathbb{E}_n is UV^{n-1} and that every element of $\pi_n(T_f, x_0)$ is homotopic to a map of the form $\alpha^m * \gamma$ for some $\gamma : S^n \rightarrow \mathbb{E}_n$ and $m \in \mathbb{Z}$. It is clear that T_f is shape equivalent to the aspherical space S^1 and thus $\tilde{\pi}_n(T_f, x_0) = 0$. Hence, $\pi_n^{\text{sh}}(T_f, x_0)$ is an uncountable indiscrete group. It follows from Case (1) of Theorem 1.1 that $\pi_n^{\text{met}}(T_f, x_0)$ is also indiscrete.

We also consider an example showing that Theorem 1.1 fails to hold in any dimension $n \geq 1$ without the local path-connectedness hypothesis, i.e the assumption that X is UV^0 .

Example 5.17. Fix $n \geq 1$. Let $A_0 = \{0\} \times [0, 1] \subseteq \mathbb{R}^2$ and $A_1 = \{(x, \sin(1/x)) \in \mathbb{R}^2 \mid 0 < x < 1/\pi\}$ so that $A = A_0 \cup A_1$ is the closed topologists sine curve. Let $a_0 = (0, 0)$ and $a_1 = (1/\pi, 0)$ and $B = A \times S^n / \{a_0\} \times S^n$ and with quotient map $q : A \times S^n \rightarrow B$. Set $x_0 = q(a_0, t)$ and, for each $a \in A$, let $\ell_a : S^n \rightarrow B$ be the embedding $\ell_a(t) = q(a, t)$. Note that ℓ_a is null-homotopic if and only if $a \in A_1$. Let X be obtained by attaching an arc $[0, 1]$ to B by $0 \sim a_0$ and $1 \sim a_1$ where we metrize X with the quotient metric. Let $\alpha : [0, 1] \rightarrow X$ be a reparameterization of this arc from a_1 to a_0 and note that $[\alpha * \ell_{a_0}]$ is non-trivial in $\pi_n(X, a_1)$. It is not difficult to show that $\pi_n(X, a_1)$ is infinite cyclic generated by $[\alpha * \ell_{a_0}]$.

The space X is a compact metric space, which may be embedded in \mathbb{R}^{n+2} . However, X is not locally path-connected at any point in $q(A_0 \times [0, 1])$. Moreover, X is shape equivalent to the aspherical space S^1 and therefore $\pi_m^{\text{sh}}(X, a_1)$ is indiscrete for all $m \geq 2$. Even though $\tilde{\pi}_1(X, a_1) \cong \mathbb{Z}$, the canonical homomorphism $\Psi_1 : \pi_1(X, a_1) \rightarrow \tilde{\pi}_1(X, a_1)$ is trivial since A is not path connected. Hence, $\pi_1^{\text{sh}}(X, a_1)$ is also indiscrete.

On the other hand, let $Y \subseteq X$ be the compact CW-complex consisting of $q(A_0 \times S^n)$ and the arc. Since $\pi_n^{\text{met}}(Y, a_1)$ is discrete by Lemma 5.1, there exists $\epsilon > 0$ such that whenever $\mu(\beta, \gamma) < \epsilon$, $[\beta] = [\gamma]$ in $\pi_n(Y, a_1)$. The image of any loop in X homotopic to $[\alpha * \ell_{a_0}]$ must contain the attached arc and the projection $A_0 \times S^n \rightarrow S^n$ must map it onto S^n . Based on these observations, we may choose ϵ small enough so that whenever $\beta \in \Omega^n(X, a_1)$, $\gamma \simeq \alpha * \ell_{a_0}$ with $\mu(\beta, \gamma) < \epsilon$, then we must have $\beta \simeq \gamma$. It follows that $\pi_n^{\text{met}}(X, a_1)$ is discrete.

In the next example, we illustrate an application of Case (2) of Theorem 1.1 as an important case where Case (1) does not apply.

Example 5.18. Let K_1, K_2, K_3, \dots be a sequence of based, compact polyhedra and $X_k = \bigvee_{j=1}^k K_j$ for $k \in \mathbb{N}$. We take $r_{k+1, k} : X_{k+1} \rightarrow X_k$ to be the retraction

which collapses K_{k+1} to the basepoint and $X = \varprojlim_{k \in \mathbb{N}} (X_k, r_{k+1, k})$. Then $X = \widetilde{\bigvee}_{j \in \mathbb{N}} K_j$ is a Peano continuum, which we refer to as the *shrinking wedge* of the sequence $\{K_j\}_{j \in \mathbb{N}}$. Since X is an inverse limit of a nested sequence of polyhedral retracts, Case (2) of Theorem 1.1 implies that for any choice of metric d on X , $\pi_n^{\text{met}}(X, x_0) = \pi_n^{\text{sh}}(X, x_0)$ for all $n \geq 1$. For example, the m -dimensional earring space \mathbb{E}_m is homeomorphic to $\widetilde{\bigvee}_{j \in \mathbb{N}} S^m$ and the conclusion $\pi_n^{\text{met}}(\mathbb{E}_m, x_0) = \pi_n^{\text{sh}}(\mathbb{E}_m, x_0)$ only follows from Case (1) in dimension $n = m$. Interestingly, this equivalence of topologies holds even when infinitely many K_j fail to be m -connected, in which case X is not UV^m and the algebraic structure of $\pi_n(X, x_0)$ is complicated significantly by Whitehead products.

We conclude by remarking that in dimension $n = 1$, Theorem 1.1 provides an avenue for relating the topology of $\pi_1^{\text{met}}(X, x_0)$ and the covering spaces of X .

Corollary 5.19. For a Peano continuum X and $x_0 \in X$, the following are equivalent:

1. ρ is a metric on $\pi_1(X, x_0)$,
2. X is π_1 -shape injective,
3. for every $[\alpha] \in \pi_1(X, x_0)$, there exists a covering map $p : (E, e) \rightarrow (X, x_0)$ such that $[\alpha] \notin p_{\#}(\pi_1(E, e))$.

Proof. The equivalence (2) \Leftrightarrow (3) was proved in [12]. (1) \Rightarrow (2) if ρ is a metric, then $\pi_1^{\text{met}}(X, x_0)$ is Hausdorff. Since $\pi_1^{\text{met}}(X, x_0) = \pi_1^{\text{sh}}(X, x_0)$ by Theorem 1.1, $\pi_1^{\text{sh}}(X, x_0)$ is Hausdorff and it follows that $\Psi_1 : \pi_1(X, x_0) \rightarrow \tilde{\pi}_1(X, x_0)$ is injective. (2) \Rightarrow (1) If $\Psi_1 : \pi_1(X, x_0) \rightarrow \tilde{\pi}_1(X, x_0)$ is injective, then $\pi_1^{\text{met}}(X, x_0) = \pi_1^{\text{sh}}(X, x_0)$ is Hausdorff. Since the topology of $\pi_1^{\text{met}}(X, x_0)$ is generated by the pseudometric ρ , ρ is a metric. \square

References

- [1] M. Abdullahi Rashid, N. Jamali, B. Mashayekhy, S.Z. Pashaei, H. Torabi, *On subgroup topologies on the fundamental group*. Hacettepe Journal of Mathematics & Statistics **49** (2020), no. 3, 935-949.
- [2] M. Abdullahi Rashid, S.Z. Pashaei, B. Mashayekhy, H. Torabi, *On the Whisker Topology on Fundamental Group*. Conference Paper from 46th Annual Iranian Mathematics Conference **46** (2015).
- [3] J. Aceti, J. Brazas, *Elements of homotopy groups undetectable by polyhedral approximation*, Preprint, 2022. arXiv:2208.06645.
- [4] A. Akbar Bahredar, N. Kouhestani, H. Passandideh, *The n -dimensional Spanier group*, Filomat **35** (2021), no. 9, 3169-3182.

- [5] A. Arhangel'skii, M. Tkachenko, *Topological Groups and Related Structures*, Series in Pure and Applied Mathematics, Atlantis Studies in Mathematics, 2008.
- [6] M.G. Barratt, J. Milnor, *An Example of Anomalous Singular Homology*, Proc. Amer. Math. Soc. **13** (1962), no. 2, 293–297.
- [7] K. Borsuk, *On the n -movability*, Bull. Acad. Polon. Sci. Ser. Sci. Math. Astr. Phys. **20** (1972), 859-864.
- [8] J. Brazas, *The topological fundamental group and free topological groups*, Topology Appl. **158** (2011) 779-802.
- [9] J. Brazas, *The fundamental group as a topological group*, Topology Appl. **160** (2013) 170-188.
- [10] J. Brazas, *On the discontinuity of the π_1 -action*, Topology Appl. **247** (2018) 29-40.
- [11] J. Brazas, P. Fabel, *On fundamental groups with the quotient topology*, J. Homotopy and Related Structures **10** (2015) 71-91.
- [12] J. Brazas, P. Fabel, *Thick Spanier groups and the first shape group*, Rocky Mountain J. Math. **44** (2014) 1415-1444.
- [13] M.R. Bridson, A. Haefliger, *Metric spaces of non-positive curvature*, Grundle Math. Wissen. 319, Springer, 1999.
- [14] J.S. Calcut; J.D. McCarthy, *Discreteness and homogeneity of the topological fundamental group*, Topology Proc. **34** (2009) 339-349.
- [15] G. Conner, M. Meilstrup, D. Repovš, A. Zastrow, M. Željko, *On small homotopies of loops*, Topology Appl. **155** (2008) 1089-1097.
- [16] J. Dugundji, *A topologized fundamental group*, Proc. Natl. Acad. Sci. **36** (1950) 141–143.
- [17] K. Eda, K. Kawamura, *Homotopy and homology groups of the n -dimensional Hawaiian earring*, Fund. Math. **165** (2000) 17-28.
- [18] R. Engelking, *General topology*, Heldermann Verlag Berlin, 1989.
- [19] P. Fabel, *Multiplication is discontinuous in the Hawaiian earring group*, Bull. Pol. Acad. Sci. Math. **59** (2011) 77-83.
- [20] P. Fabel, *Compactly generated quasitopological homotopy groups with discontinuous multiplication*, Topology Proc. **40** (2012) 303-309.
- [21] S. Mardešić and J. Segal, *Shape theory*, North-Holland Publishing Company, 1982.

- [22] K. Sakai, *Geometric aspects of general topology*, Springer Japan, Tokyo, 2013.
- [23] E. Spanier, *Algebraic Topology*, McGraw-Hill, 1966.
- [24] Z. Virk, A. Zastrow *A new topology on the universal path space*, *Topology Appl.* **231** (2017), no. 1, 186-196.
- [25] H. Wada, *Local connectivity of mapping spaces*, *Duke Math. J.* **22** (1955), 419-425.