

Progress towards the two-thirds conjecture on locating-total dominating sets*

Dipayan Chakraborty[†] Florent Foucaud[†] Anni Hakanen^{†‡}
 Michael A. Henning[§] Annegret K. Wagler[†]

November 28, 2022

Abstract

We study upper bounds on the size of optimum locating-total dominating sets in graphs. A set S of vertices of a graph G is a locating-total dominating set if every vertex of G has a neighbor in S , and if any two vertices outside S have distinct neighborhoods within S . The smallest size of such a set is denoted by $\gamma_t^L(G)$. It has been conjectured that $\gamma_t^L(G) \leq \frac{2n}{3}$ holds for every twin-free graph G of order n without isolated vertices. We prove that the conjecture holds for cobipartite graphs, split graphs, block graphs, subcubic graphs and outerplanar graphs.

1 Introduction

Our aim is to study upper bounds on the smallest size of locating-total dominating sets in graphs. This notion is part of the extended research area of *identification problems* in graphs and, more generally, discrete structures like hypergraphs. In this type of problems, one seeks to select a small solution set, generally vertices of a graph, in order to uniquely identify each vertex of the graph by its relationship with the selected vertices. More precisely, given a set D of vertices of a graph G , we say that two vertices v and w of G are *located* by D if they have distinct sets of neighbors in D . Various notions based on this location property have been studied, such as locating-dominating sets [33], identifying codes [27] or separating sets [8], to name a few. We refer to the online bibliography on these topics maintained by Lobstein [30] (almost 500 references by the time of writing, in 2022). This type of problems have a wide range of applications, such as fault-detection in sensor or computer networks [27, 34], biological testing [31], machine learning [10], or canonical representations of graphs [3, 29], to name a few.

In this paper, we study the notion of a locating-total dominating set. A set D of vertices is a *total dominating set*, abbreviated TD-set, of a graph G if every vertex in G has a neighbor in D . Total dominating sets are a natural and widely studied variant of the domination problem in graphs. We refer to the book [26] for an overview on the topic. A *locating-total dominating set* of G is a TD-set $D \subset V(G)$ such that any two vertices of G not in D are located by D . The smallest size of such a locating-total dominating set of a graph G is called the *locating-total domination number* of G and is denoted by $\gamma_t^L(G)$. A graph admits a (locating-)total dominating set if and only if it has no isolated vertex. We abbreviate a locating-total dominating set by LTD-set, and we say that a γ_t^L -set of G is a LTD-set of minimum cardinality, $\gamma_t^L(G)$, in G .

The concept of a locating-total dominating set was first considered in [23], based on the similar concept of a locating-dominating set (where total domination is replaced with usual domination) introduced by Slater in the 1980s [33]. It was studied for example in [1, 2, 5, 6, 7, 13, 14, 15, 18, 19, 24, 25]. The

*This work was sponsored by a public grant overseen by the French National Research Agency as part of the “Investissements d’Avenir” through the IMobS3 Laboratory of Excellence (ANR-10-LABX-0016) and the IDEX-ISITE initiative CAP 20-25 (ANR-16-IDEX-0001). We also acknowledge support of the ANR project GRALMECO (ANR-21-CE48-0004).

[†]Université Clermont-Auvergne, CNRS, Mines de Saint-Étienne, Clermont-Auvergne-INP, LIMOS, 63000 Clermont-Ferrand, France

[‡]Research supported by Jenny and Antti Wihuri Foundation

[§]Department of Mathematics and Applied Mathematics, University of Johannesburg

associated decision problem is NP-hard [28, 32]. The related concept where *all* vertices (not just the ones outside of D) must be located was studied in [13, 21, 23].

It is known that any graph of order n with every component of order at least 3 has a TD-set size at most $\frac{2}{3}n$ [16], and this bound is tight only for the triangle, the 6-cycle and the family of 2-coronas of graphs [11]. (The 2-*corona* $H \circ P_2$ of a connected graph H is the graph of order $3|V(H)|$ obtained from H by attaching a path of length 2 to each vertex of H so that the resulting paths are vertex-disjoint.)

However, such a bound does not hold for locating-total dominating sets. Two vertices of a graph are *twins* if they either have the same open neighborhood (*open twins*) or the same closed neighborhood (*closed twins*). Consider a set S of vertices that are pairwise twins of size at least 2 in a graph G (and S forms either a clique or an independent set). Then, any locating-total dominating set D needs to contain all vertices of S except possibly one. Indeed, any two such vertices not in D would otherwise not be located. For example, any complete graph of order at least 2 has only twins, and thus has its locating-total domination number equal to its order minus one (while any two vertices form a total dominating set). Other families of (twin-free) graphs with a total dominating set of size 2 and arbitrarily large locating-total domination number have been described in [18].

Nevertheless, it seems that in the absence of twins, the locating-total domination number cannot be as close to the graph's order as in the general case. Towards such a fact, inspired by a similar problem for (non-total) locating-dominating sets [17, 20, 22], two of the authors posed the following conjecture (a graph is called *twin-free* if it contains no set of twins). A graph G is *isolate-free* if it contains no isolated vertex.

Conjecture 1 ([18]). Every twin-free isolate-free graph G of order n satisfies $\gamma_t^L(G) \leq \frac{2}{3}n$.

Conjecture 1 was proved for graphs with no 4-cycle as a subgraph in [18]. It was also proved for line graphs in [19]. It was also proved in [18] to hold for all graphs with minimum degree at least 26 for which another related conjecture [20, 22] holds (which is the case for example for bipartite graphs and cubic graphs). It was proved in a stronger form for claw-free cubic graphs in [24] (there, the $\frac{2}{3}$ factor in the upper bound is in fact replaced with $\frac{1}{2}$, and the authors conjectured that $\frac{1}{2}$ holds for all connected cubic graphs, except K_4 and $K_{3,3}$). An approximation of the conjecture was proved to hold for all twin-free graphs in [18], where the $\frac{2}{3}$ factor in the upper bound is replaced with $\frac{3}{4}$.

Note that, if true, the bound of Conjecture 1 is tight for the 6-cycle and 2-coronas, by the following.

Observation 2. If a graph G of order n is a triangle, a 6-cycle or a 2-corona of any graph, then $\gamma_t^L(G) = \frac{2}{3}n$.

A graph is *cobipartite* if its vertex set can be partitioned into two cliques, and *split* if it can be partitioned into a stable set (also called an independent set in the literature) and a clique. A graph is a *block graph* if every 2-connected component forms a clique. A graph is *subcubic* if each vertex has degree at most 3. A graph is *outerplanar* if it can be embedded in the plane without any edge-crossing, so that all vertices lie on the same face of the embedding.

In this paper, we give further evidence towards Conjecture 1, by showing that it holds for cobipartite graphs (Section 2), split graphs (Section 3), block graphs (Section 4), subcubic graphs (Section 5) and outerplanar graphs (Section 6). We conclude in Section 7.

Some of our results are actually slightly stronger. Indeed, the proved upper bound for cobipartite graphs is in fact $\lceil \frac{n}{2} \rceil$ (which is tight). For twin-free split graphs, we show that the $\frac{2n}{3}$ bound of the conjecture can never be reached. However, we construct infinitely many connected split graphs that come very close to the bound; this is interesting in its own right, showing that not only 2-coronas have such large locating-total domination numbers. Moreover, the bound for subcubic graphs is proved to hold for all subcubic graphs, even if they have twins.

We now introduce some of the notations used in the paper. The open and closed neighborhoods of a vertex v in a graph G are denoted $N_G(v)$ and $N_G[v]$, respectively (or $N(v)$ and $N[v]$ if G is clear from the context). We denote by $\deg_G(v)$ the *degree* of the vertex v in the graph G , that is, $\deg_G(v) = |N_G(v)|$. The *distance* in G between two vertices u, v is denoted $d_G(u, v)$. If two graphs G and H are isomorphic, we note $G \cong H$. For a graph G with a vertex or edge x , we denote by $G - x$ the subgraph of G obtained by removing x , and by $G + x$ the supergraph of G obtained by adding x . Similarly, if X is a set of vertices and edges, we use the notations $G - X$ and $G + X$ for the subgraph and supergraph of G obtained by

deleting or adding all the elements of X . A *leaf* is a vertex of degree 1, and its unique neighbor is called a *support vertex*. We denote by $\delta(G)$ and $\Delta(G)$ the minimum and maximum degree, respectively, in the graph G . We denote a *path*, a *cycle*, and a *complete* graph on n vertices by P_n , C_n , and K_n , respectively.

2 Cobipartite graphs

We now prove a stronger variant of the bound of Conjecture 1, whose proof is a refinement of a similar proof for the (non-total) locating-domination number from [20]. The bound is tight for *complements of half-graphs*, which are graphs made from two cliques of the same size [20, Definition 5] and whose locating(-total) domination number is equal to $\frac{n}{2}$.

Theorem 3. For any twin-free cobipartite graph G of order n , we have $\gamma_t^L(G) \leq \lceil \frac{n}{2} \rceil$.

Proof. Note that G is connected (as a disconnected cobipartite graph is the disjoint union of two cliques, and thus not twin-free: it either has closed twins if one of the cliques has order at least 2, or is a pair of open twins if both cliques have order 1). Let C_1 and C_2 be two cliques of G that partition its vertex set. Since G is twin-free, both C_1 and C_2 have size at least 2 (assume that $|C_1| \leq |C_2|$). Moreover, no two vertices of C_1 have the same neighborhood in C_2 , and vice-versa. Thus, if any of C_1, C_2 is a TD-set, then it is also an LTD-set. Thus, if C_1 is a TD-set, we are done, as $|C_1| \leq \lfloor \frac{n}{2} \rfloor$. If however C_1 is not a TD-set, it means that some vertex v of C_2 has no neighbor in C_1 ; this vertex is unique since G is twin-free.

If moreover, there is a vertex w in C_1 with no neighbor in C_2 , we select $D = (C_1 \setminus \{w\}) \cup \{x\}$ as a solution set, where $x \neq v$ is any vertex of C_2 other than v . This set is clearly a TD-set. Any two vertices from C_2 are located, as v is only dominated by x , and any two other vertices from $C_2 \setminus \{x\}$ have a distinct and nonempty neighborhood within C_1 (and thus, within $C_1 \setminus \{w\}$). Moreover, w is the only vertex not in D not dominated by x . Hence, D is an LTD-set. Since $|D| = |C_1| \leq \lfloor \frac{n}{2} \rfloor$, we are done.

Otherwise, every vertex of C_1 has a neighbor in C_2 , and so C_2 is a TD-set and, in fact, as seen previously, an LTD-set. Thus, if $|C_2| \leq \lceil \frac{n}{2} \rceil$, we are done. Otherwise, we have $|C_1| < \lfloor \frac{n}{2} \rfloor$, and thus $|C_1| + 1 \leq \lceil \frac{n}{2} \rceil$. Moreover, by similar arguments as in the previous paragraph, the set C_1 together with any vertex of C_2 other than v produces a LTD-set of size $|C_1| + 1$, and we are done. \square

3 Split graphs

Consider a split graph $G = (Q \cup S, E)$ where Q induces a clique and S a stable set. We suppose that G is isolate-free to ensure the existence of an LTD-set in G , which further implies that G is connected and Q non-empty (as every component not containing the clique Q needs to be an isolated vertex from S).

Theorem 4. For any twin-free isolate-free split graph $G = (Q \cup S, E)$ of order n , we have $\gamma_t^L(G) < \frac{2}{3}n$.

Proof. First, note that we have $|Q|, |S| \geq 2$ as otherwise G is a single vertex or not twin-free.

Observe next that Q is an LTD-set of G since Q is a TD-set and no two vertices in S have the same neighbors in Q (as G is twin-free) showing that Q is also locating. Hence, the assertion is true if $|Q| < \frac{2}{3}n$.

Consider now a set D consisting of all vertices in S and, for each $s \in S$, some arbitrary neighbor $q_s \in Q$ (which exists since G is connected). The set D is an LTD-set of G since D is a TD-set and no two vertices in $Q \setminus D$ have the same neighbors in S (as G is twin-free) implying that D is also locating. Hence, the assertion is true if $|Q| > \frac{2}{3}n$ and, thus, $|S| < \frac{1}{3}n$ and $|D| = 2|S| < \frac{2}{3}n$.

It is left to consider the case $|Q| = \frac{2}{3}n$ and $|S| = \frac{1}{3}n$. We observe that any set D constructed as above is an LTD-set of G , and we are going to show that it is not minimum. In fact, as G is twin-free, there are two vertices $s, s' \in S$ so that $N(s) \cap N(s')$ is non-empty. To see this, observe that otherwise all sets $N(s_i)$ need to be pairwise disjoint, which would imply that

- either $N(s_i)$ is composed of twins when $|N(s_i)| \geq 2$ holds, or
- each set $N(s_i)$ contains exactly one vertex and the $\frac{1}{3}n$ vertices in Q without a neighbor in S are twins.

As G is twin-free, none of these cases can happen. Now, the set D' obtained from D by replacing q_s and $q_{s'}$ by a common neighbor $q_{s,s'}$ of s and s' in Q is an LTD-set of G , indeed:

- each vertex from S has a neighbor in D' , so does every vertex from Q (as all vertices in $Q \setminus \{q_{s,s'}\}$ have $q_{s,s'}$ as a neighbor in D' and $q_{s,s'}$ has s and s' as neighbors in D') which shows that D' is a TD-set;
- no two vertices in $Q \setminus D'$ have the same neighbors in S (as G is twin-free) which implies that D' is also locating.

Thus, D' is an LTD-set of G of size

$$|D'| = |D| - 1 = 2 \cdot |S| - 1 = 2 \cdot \frac{1}{3}n - 1 < \frac{2}{3}n,$$

which finally proves the assertion. \square

We next show that the bound of Theorem 4 cannot be improved.

Proposition 5. For each integer $k \geq 3$, there is a connected twin-free split graph G_k of order $n = 3k$ and $\gamma_t^L(G_k) \geq 2k - 1$.

Proof. Let $Q = \{q_1, \dots, q_k\} \cup \{q'_1, \dots, q'_k\}$ be a clique and $S = \{s_1, \dots, s_k\}$ a stable set, so that $N(s_i) = \{q_i, q'_i\}$ for $1 \leq i < k$ and $N(s_k) = \{q_1, \dots, q_k\}$. Note that q'_k has no neighbor in S and that the sets $N(s_i)$ are disjoint for $1 \leq i < k$. See Figure 1 for an illustration.




Figure 1: The construction of graph G_k in the proof of Proposition 5, with an optimal LTD-set (black vertices).

Let C be an LTD-set of G_k . Consider the $k - 1$ closed neighborhoods $N[s_i]$ for $1 \leq i < k$. If we have $|N[s_i] \cap C| \geq 2$ for all i with $1 \leq i < k$, then $|\bigcup_{1 \leq i < k} (N[s_i] \cap C)| \geq 2k - 2$, and at least one of the remaining vertices s_k, q_k, q'_k must belong to C , as otherwise $N(q_k) \cap C = N(q'_k) \cap C$ would follow, a contradiction. This implies $|C| \geq 2k - 1$.

If, however, for some i with $1 \leq i < k$, we have $|N[s_i] \cap C| = 1$, then $s_i \notin C$ since otherwise s_i is not totally dominated by the set C . If $N[s_i] \cap C = \{q_i\}$, then $N(q'_i) \cap C = Q \cap C$. If $N[s_i] \cap C = \{q'_i\}$, then $N(q_i) \cap C = (Q \cup \{s_k\}) \cap C$. The two possibilities can occur at most once each. Assume that they both occur once each, with $N[s_a] \cap C = \{q'_a\}$ and $N[s_b] \cap C = \{q_b\}$ (with $1 \leq a < b < k$). Note that $s_k \in C$, otherwise q_a and q'_b are not located. Moreover, C must contain q'_k (otherwise q'_b and q'_k are not located) and q_k (otherwise q_a and q_k are not located), and so $|C| \geq 2k - 1$, as claimed.

Similarly, if we have $|N[s_i] \cap C| \geq 2$ for all i with $1 \leq i < k$ except that $N[s_a] \cap C = \{q'_a\}$, if $s_k \in C$, then $q_k \in C$, otherwise q_a and q_k are not located. If $s_k \notin C$, then both q_k, q'_k are in C to locate the vertices q_a, q_k, q'_k . Thus, again $|C| \geq 2k - 1$.

Finally, if we have $|N[s_i] \cap C| \geq 2$ for all i with $1 \leq i < k$ except that $N[s_b] \cap C = \{q_b\}$, if $s_k \in C$, then $q'_k \in C$, otherwise q'_b and q'_k are not located. If $s_k \notin C$, then both q_k, q'_k are in C to locate the vertices q'_b, q_k, q'_k , and again $|C| \geq 2k - 1$.

Thus, in all the above cases, we have $|C| \geq 2k - 1$ and, together with the upper bound $\gamma_t^L(G_k) < \frac{2}{3} \times 3k = 2k$ from Theorem 4, we finally obtain $\gamma^{LTD}(G_k) = 2k - 1$. \square

4 Block graphs

A block graph is a graph in which every maximal 2-connected subgraph (henceforth referred to as a *block*) is complete. Equivalently, block graphs are diamond-free chordal graphs [4]. A *cut-vertex* v of a graph

G is one such that the graph $G - v$ has more components than G . For any block graph G , a *leaf block* of G is a block that contains only a single cut-vertex of G . In this section, we show that our $\frac{2}{3}$ -conjecture (namely, Conjecture 1), holds for block graphs. Trees are a subclass of block graphs in which every block is of order 2. There are some concepts of trees which we use quite often in proving our result for block graphs and we, therefore, define these concepts formally here.

A *root* of a tree is a fixed vertex of the tree to which the name is designated. Having fixed a root r of a tree T , for any vertex u of T ,

- (1) a *child* of u is a vertex v of T such that uv is an edge of T and $d_T(v, r) = d_T(u, r) + 1$;
- (2) a *grandchild* of u is a vertex w of T such that uv and vw are edges of T and $d_T(w, r) = d_T(u, r) + 2$; and
- (3) a *great-grandchild* of u is a vertex x of T such that uv , vw and wx are edges of T and $d_T(x, r) = d_T(u, r) + 3$.

Conversely, the vertex u of T is called the *parent*, the *grandparent* and the *great-grandparent* of v , w and x , respectively. Given any two vertices u and v of a tree T , we say that u is *above* v in T (or v is *below* u in T) if there exists a sequence of vertices x_1, x_2, \dots, x_m of T such that, for each $i = 1, 2, \dots, m - 1$, x_{i+1} is a child of x_i in T , where $m \geq 2$, $x_1 = u$ and $x_m = v$.

Theorem 6. If $G \cong P_3$ or if G is a twin-free isolate-free block graph of order $n \geq 4$, then $\gamma_t^L(G) \leq \frac{2}{3}n$.

Proof. Since the locating-total domination number of a graph is the sum of the locating-total domination numbers of each of the components of the graph, it is therefore enough to prove the theorem for a connected twin-free block graph. Thus, let us assume that G is either isomorphic to a 3-path or is a connected twin-free block graph of order $n \geq 4$. The proof is by induction on $n \geq 3$. The base case of the induction hypothesis is when $n = 3$, in which case $G \cong P_3$ and $\gamma_t^L(G) = 2 = \frac{2}{3}n$. Clearly, any two consecutive vertices of P_3 constitute a minimum LTD-set of P_3 and hence, the result holds for the base case of the induction hypothesis. We now assume, therefore, that $n \geq 4$ and that the induction hypothesis is true for all connected twin-free block graphs of order at least 3 and at most $n - 1$. Next, we construct a new graph T_G from G in the following way (see Figure 2 for an example of the construction).

For every block B of G , introduce a vertex $u_B \in V(T_G)$ and for every cut-vertex $c \in V(G)$, introduce a vertex $v_c \in V(T_G)$. Next, introduce edges $u_B v_c \in E(T_G)$ if and only if the cut-vertex c belongs to the block B of G . By construction, therefore, T_G is a tree. Thus, the vertices of the tree T_G are of two types:

- (1) *u-type*: u_B introduced in a one-to-one association with a block B of G ; and
- (2) *v-type*: v_c introduced in a one-to-one association with a cut-vertex c of G .

Notice that any pair of vertices w, z of the tree T_G such that w is the grandparent/grandchild of z in T_G are of the same vertex type. For a fixed cut-vertex $r \in V(G)$, designate $v_r \in V(T_G)$ as the root of T_G (indeed, such a cut-vertex exists as $n \geq 4$ and the twin-free property of G implies that G has at least two blocks). Notice that any leaf of the tree T_G is a vertex of the type u_B for some leaf block B of G . By the twin-free nature of G every leaf block B of G has order exactly 2. Now, fix a leaf u_F of T_G that is at the farthest distance, in T_G , from the root v_r of T_G . We now look at the great-grandparent of the leaf u_F in the tree T_G (indeed, the great-grandparent of u_F in T_G exists because, on account of G being twin-free, at least one of the blocks of G containing the vertex r has a cut-vertex other than r). Notice that the great-grandparent of u_F in T_G must be a vertex of the type v_p for some cut-vertex p of G . We next define the following.

$$\begin{aligned} \mathcal{B}_p &= \{B : B \text{ is a block of } G \text{ and } u_B \text{ is either a child or a great-grandchild of } v_p \text{ in } T_G\}; \\ U &= \cup_{B \in \mathcal{B}_p} V(B); \text{ and} \\ A &= \{x \in U : x \text{ is a cut-vertex of } G\}. \end{aligned}$$

We now establish the following two claims related to the sets defined above.

Claim A. The set A is an LTD-set of $G[U]$.




Figure 2: Figure (a) represents a twin-free block graph G and Figure (b) represents T_G . The vertices underneath the dashed curve represent those deleted from G to obtain G' . The black vertices represent vertices in the set A . (All notations are as in the proof of Theorem 6.)

Proof of claim. That A is a TD-set of $G[U]$ is clear from the structure of G . We show that A is also a locating set of $G[U]$. So, let us assume that vertices $w, z \in U \setminus A$. Since G is twin-free, notice that w and z belong to distinct blocks of \mathcal{B}_p , say B and B' , respectively.

First, assume that u_B and $u_{B'}$ are both children of v_p in T_G . Then, at least one of u_B and $u_{B'}$ must have a child in T_G (or equivalently, at least one of B and B' must contain a cut-vertex of G). This is because if none of u_B and $u_{B'}$ had a child in T_G , it would mean that B and B' are leaf blocks of G and so, $|B| = |B'| = 2$ (since G is twin-free). That, in turn, would mean that G has open twins (with the leaves of G in the blocks B and B' sharing the common support vertex v_p), contrary to our assumption. Thus, if q is a cut-vertex of G in $V(B) \cup V(B')$, then $q \in A$ locates w and z . Next, we assume that u_B and $u_{B'}$ are both great-grandchildren of v_p in T_G (or equivalently, B and B' are both leaf blocks of G). Then, u_B and $u_{B'}$ have parents v_q and $v_{q'}$, say, respectively, in T_G , where $q \neq q'$ due to the fact that G is twin-free. This implies that w and z are located in G by $\{q, q'\}$. Finally, we assume that u_B is a child of v_p and that $u_{B'}$ is a great-grandchild of v_p in T_G . Then, w and z are located in G by p . This proves the claim. \square

Claim B. $|U| \geq 2|A| - 1$.

Proof of claim. Let $u_{B_1}, u_{B_2}, \dots, u_{B_m}$ be $m \geq 1$ children of v_p in T_G and let each block B_i of G be of order n_i . Now, due to the twin-free nature of G , each vertex u_{B_i} of T_G has at least $n_i - 2$ and at most $n_i - 1$ children (and hence at least $n_i - 2$ and at most $n_i - 1$ grandchildren as well). To be more precise, assume that, for $0 \leq s \leq m$, the vertices $u_{B_1}, u_{B_2}, \dots, u_{B_s}$ have exactly $n_1 - 2, n_2 - 2, \dots, n_s - 2$ children, respectively, in T_G ; and that the vertices $u_{B_{s+1}}, u_{B_{s+2}}, \dots, u_{B_m}$ have exactly $n_{s+1} - 1, n_{s+2} - 1, \dots, n_m - 1$ children, respectively, in T_G . This implies that we have the following.

$$|U| = 1 - s + 2 \sum_{1 \leq i \leq m} (n_i - 1) = 1 - 2m - s + 2 \sum_{1 \leq i \leq m} n_i.$$

Moreover, we have

$$|A| = 1 - s + \sum_{1 \leq i \leq m} (n_i - 1) = 1 - m - s + \sum_{1 \leq i \leq m} n_i.$$

Combining the above two equations, therefore, we have

$$|U| - (2|A| - 1) = s \geq 0$$

which proves the claim. \square

Now, let $G' = G - U$, that is, G' is the graph obtained by deleting from G all vertices (and edges incident with them) in the blocks $B \in \mathcal{B}_p$. Notice that G' is still a connected block graph; and assume that the order of G' is n' (which is strictly less than n). We next divide the proof according to whether G' is twin-free, has twins or is isomorphic to a 3-path.

Case 1 (G' is either twin-free or is isomorphic to a 3-path). We further subdivide this case into the following.

Subcase 1.1 ($n' \leq 2$). In this subcase, if $n' = 2$, then the two vertices of G' form an edge of G' (since G is connected). Hence, the two vertices of G' are closed twins of degree 1, contrary to our initial assumption in this case. Therefore, let us assume that $n' \leq 1$. If $n' = 1$, then v_p has no grandparent in T_G . In other words, there is no vertex of v -type above v_p in T_G . This implies that v_p must itself be the root vertex of T_G . However, this, in turn, implies that G' is an empty graph which contradicts the fact that $n' = 1$. Thus, we must have $n' = 0$. In this case too, v_p must itself be the root vertex of T_G and so, $G = G[U]$ and $|U| = n$. Therefore, by Claim A, the set A is an LTD-set of $G[U] = G$. Moreover, by Claim B, we have

$$|A| \leq \frac{1}{2}(|U| + 1) = \frac{1}{2}(n + 1) \leq \frac{2}{3}n,$$

where the last inequality is true since $n \geq 4$.

Subcase 1.2 ($n' \geq 3$). In this subcase, by the induction hypothesis, we have $\gamma_t^L(G') \leq \frac{2}{3}n'$. Suppose now that $S' \subset V(G')$ is a minimum LTD-set of G' , that is with $|S'| = \gamma_t^L(G')$. We then claim that the set $S = S' \cup A$ is an LTD-set of G . To prove so, we first see that the set S is a TD-set of G , since S' is a TD-set of G' and A is a TD-set of $G[U]$. Moreover, S is also a locating set of G due to the following two reasons.

- (1) Any two distinct vertices $w \in V(G') \setminus S'$ and $z \in V(G) \setminus S$ are located by S' .
- (2) By Claim A, the set A is a locating set of $G[U]$.

Using Claim B, therefore, the two-thirds bound on $\gamma_t^L(G)$ in this subcase is established by the following inequality.


$$\begin{aligned} \gamma_t^L(G) &\leq |S| = |S'| + |A| \leq \gamma_t^L(G') + \frac{2}{3}(2|A| - 1) \quad [\text{since } |A| \geq 2] \\ &\leq \frac{2}{3}(n' + 2|A| - 1) \leq \frac{2}{3}(n' + |U|) = \frac{2}{3}n. \end{aligned}$$

We next turn to the case that G' has twins.

Case 2 (G' is neither twin-free nor is isomorphic to a 3-path). Assume that x and y are two vertices of G' which are twins in G' . Then, without loss of generality, there exists an edge in G between the vertices p and x and there is no edge in G between p and y . This implies that x and p belong to the same block X , say, of G to which y does not belong. Let the block of G to which y belongs be called Y . Next, we prove the following claim.

Claim C. The vertex x also belongs to the block Y of G .

Proof of claim. Toward a contradiction, let us assume that $x \notin V(Y)$. If the blocks X and Y of G had no common vertex in G' , by the connectedness of G' , it would mean that the symmetric difference of the sets $N_{G'}(x)$ and $N_{G'}(y)$ is non-empty, and so, x and y would not be twins in G' , a contradiction. So, let $V(X) \cap V(Y) = \{v\}$ (note that two blocks of a block graph can intersect at not more than a single vertex). Now, if either of x and y were cut-vertices of G , or if $V(X) \cup V(Y)$ contained any vertex



(a) The vertices to the left of the dashed curve represent those deleted from G to obtain G'' . $G'' \cong P_3$. The black vertices constitute the set A and the grey vertices constitute an LTD-set S'' of G'' .

(b) The vertices to the left of the dashed curve represent those deleted from G to obtain G^* . G'' has twins x'' and y ; and G^* is a twin-free block graph. The black vertices constitute the set $A \cup \{x\}$ and the grey vertices constitute an LTD-set S^* of G^* .

Figure 3: Twin-free block graph G . The dotted boxes mark the blocks X , X'' and Y of G as in the proof of Theorem 6.

other than p, x, v and y , then again, x and y would not be twins in G' , the same contradiction as before. This implies that $V(G') = \{x, v, y\}$ and that G' is isomorphic to a 3-path, again a contradiction to our assumption in this case. Hence, the claim holds. \square

Now, clearly, $X \neq Y$, or else, py would be an edge in G , contradicting our earlier observation. Thus, x is a cut-vertex of G belonging to the distinct blocks X and Y of G . Moreover, $|X| = 2$, or else, again, x and y would not be twins in G' , a contradiction. More precisely, $V(X) = \{x, p\}$. We also observe here that y cannot be a cut-vertex of G , or else, x and y would not be twins in G' , again the contradiction as before. Therefore, since G is twin-free, every vertex other than y of the block Y must be a cut-vertex of G .

Now, look at the block graph $G'' = G' - x$ on, say, n'' vertices (the graph induced by the vertices on the right of the dashed curve in Figure 3a). Notice that, in the tree T_G , the vertex v_x cannot have any children other than u_X , or else, x and y cannot be twins in G' , contrary to our assumption for this case. This implies that the block graph G'' is also connected. We also have $y \in V(G'')$. Thus, $n'' \geq 1$. However, we claim that $n'' \neq 2$.

Claim D. The order n'' of the block graph G'' cannot be 2.

Proof of claim. Toward a contradiction, let us assume $n'' = 2$ such that z is the vertex of G'' other than y . That is, $V(G'') = \{y, z\}$ and that yz is an edge of G , by the connectedness of G'' . Therefore, y and z belong to the same block Z , say, of G . If $Z = Y$, then by our observation preceding this claim, the vertex z , being different from y , must be a cut-vertex of G making $n'' \geq 3$, contradicting our assumption. So, Z and Y are distinct blocks of G to both of which the vertex y belongs. This makes y a cut-vertex of G , again contradicting our earlier observation that y cannot be a cut-vertex of G . This proves the claim. \square

Thus, we next divide this case into the following two subcases according to the order of G'' .

Subcase 2.1 ($n'' = 1$). In this subcase, we claim that $S = A \cup \{x\}$ is an LTD-set of G . It is clear that S is a TD-set of G ; by Claim A, A is an LTD-set of $G[U]$. The vertex y is located from any vertex in

$U \setminus A$ by the vertex x , and thus the set S is also locating. Therefore, in this case, we have

$$\begin{aligned}\gamma_t^L(G) &\leq |S| = |A| + 1 < \frac{2}{3}(2|A| - 1) + \frac{4}{3} \quad [\text{since } |A| \geq 2] \\ &\leq \frac{2}{3}(|U| + 2) = \frac{2}{3}n. \quad [\text{by Claim B}]\end{aligned}$$

Subcase 2.2 ($n'' \geq 3$ and G'' is either twin-free or is isomorphic to a 3-path). Since n'' is at least 3 and is strictly less than n , by the induction hypothesis, we have $\gamma_t^L(G'') \leq \frac{2}{3}n''$. Moreover, let S'' be a minimum LTD-set of G'' , that is with $|S''| = \gamma_t^L(G'')$. We next claim the following.

Claim E. The set $S = S'' \cup A$ is an LTD-set of G .

Proof of claim. Since S'' is a TD-set of G'' and A is a TD-set of $G[U \cup \{x\}]$, S is therefore a TD-set of G . Next we show that S is also a locating set of G . To begin with, we note that $Y'' = Y - x$ is a block of G'' containing the vertex y . Now, since y is not a cut-vertex of G , we have $S'' \cap Y'' \neq \emptyset$ (or else, y is not dominated by S''). This implies that x is located by S'' from all vertices in $U \setminus A$. Moreover, x is also located by p from all vertices in $V(G'') \setminus S''$. Next, any pair w, z of distinct vertices with $w \in V(G'') \setminus S''$ and $z \in V(G) \setminus (S \cup \{x\})$ are located by S'' . Finally, any distinct pair of vertices $w, z \in U \setminus A$ are located by A , since the latter is an LTD-set of $G[U]$ by Claim A. \square

Therefore, in this subcase, using Claim B again, the theorem follows from the following inequality.

$$\begin{aligned}\gamma_t^L(G) &\leq |S| = |S''| + |A| \leq \gamma_t^L(G'') + \frac{2}{3}(2|A| - 1) \quad [\text{since } |A| \geq 2] \\ &\leq \frac{2}{3}(n'' + 2|A| - 1) \leq \frac{2}{3}(n'' + |U|) < \frac{2}{3}n.\end{aligned}$$

Subcase 2.3 ($n'' \geq 3$ and G'' is neither twin-free nor is isomorphic to a 3-path). Assume that x'' and y'' are a pair of twins of G'' . Moreover, for x'' and y'' to be twins in G'' , at least one of them must be in the block Y . Let us, without loss of generality, assume that $y'' \in V(Y)$.

We next observe that the vertices y and y'' are the same. To prove so, by contradiction, let us assume that $y'' \neq y$. Then y'' is a cut-vertex of G and so, for x'' and y'' to be twins in G'' , x'' must *not* belong to the block Y of G . However, this, in turn, implies that y is a neighbor of y'' but not of x'' and so, x'' and y'' are not twins in G'' , a contradiction all the same. This, therefore, proves the observation.

Again, the vertex $x'' \notin Y$, since otherwise, $x'' \neq y'' = y$ implies that x'' is a cut-vertex of G , thus forcing x'' and y'' to not be twins, contrary to our assumption. Let x'' belong to the block $X'' (\neq Y)$ of G'' (and of G). We now try to establish the structure of the block Y of G . Notice that, by the structure of a block graph, the twins x'' and y in G'' must have a *single* common neighbor z , say, in G'' such that z is a cut-vertex of G belonging to both the blocks Y and X'' of G . Furthermore, if the block Y contains any vertex of G other than the vertices x, y and z , then x'' and y are not twins in G'' , a contradiction. Thus, we have $V(Y) = \{x, y, z\}$.

Next, to understand the structure of the block X'' of G'' , we see that neither can X'' contain any vertex other than z and x'' , nor can x'' be a cut-vertex of G ; or else, we again have the contradiction that x'' and y are not twins in G'' . Therefore, this implies that $V(X'') = \{x'', z\}$, that is, X'' is a leaf block of G'' (and of G). See Figure 3 for the structure of the blocks X'' and Y .

With that, we look at the block graph $G^* = G'' - y$ (the graph induced by the vertices on the right of the dashed curve in Figure 3b). Then, G^* is again a connected graph, since y is not a cut-vertex of G . Moreover, the order n^* of G^* is at least 2 (since $x'', z \in V(G^*)$). If, however, $n^* = 2$, then we have $V(G'') = \{x'', y, z\}$ and thus, G'' is isomorphic to a 3-path, contrary to our assumption in this subcase. Therefore, we have $n^* \geq 3$. We next show the following claim.

Claim F. The graph G^* is twin-free.

Proof of claim. Toward a contradiction, let us assume that the block graph G^* has a pair of twins. Then one of them must be the cut-vertex z of G . Let x^* be the other vertex of G^* such that x^* and z are twins in G^* . Since x'' is a neighbor of z alone in G^* , therefore z cannot be a twin in G^* of any vertex other

than x'' . In other words, $x^* = x''$. However, since $\deg_{G^*}(x'') = 1$, we have $\deg_{G^*}(z) = 1$ and, hence, the graph G^* is simply the edge $x''z$ of G . This however, contradicts the fact that $n^* \geq 3$. Hence, this proves that G^* is twin-free. (\square)

Since n^* is at least 3 and is strictly less than n , by the induction hypothesis, we have $\gamma_t^L(G^*) \leq \frac{2}{3}n^*$. Moreover, let S^* be a minimum LTD-set of G^* , that is with $|S^*| = \gamma_t^L(G^*)$. We next claim the following.

Claim G. The set $S = S^* \cup A \cup \{x\}$ is an LTD-set of G .

Proof of claim. Since S^* is a TD-set of G^* and $A \cup \{x\}$ is a TD-set of $G[U \cup \{x, y\}]$, S is therefore a TD-set of G . Next we show that S is also a locating set of G . To begin with, we note that, since x'' is a leaf in G^* , its support vertex z must be in the LTD-set S^* of G^* . Thus, the vertex y is located from every other vertex in $V(G) \setminus S$ by the set $\{x, z\}$. Next, any pair w_1, w_2 of distinct vertices with $w_1 \in V(G^*) \setminus S^*$ and $w_2 \in V(G) \setminus S$, respectively, are located by S^* . Finally, by Claim A, any pair of distinct vertices $w_1, w_2 \in U \setminus A$ are located by the set A . (\square)

Therefore, again using Claim B, in this subcase, the theorem follows from the following inequality.

$$\begin{aligned} \gamma_t^L(G) &\leq |S| = |S^*| + |A| + 1 < \gamma_t^L(G^*) + \frac{2}{3}(2|A| + 1) \quad [\text{since } |A| \geq 2] \\ &\leq \frac{2}{3}(n^* + 2|A| + 1) \leq \frac{2}{3}(n^* + |U| + 2) = \frac{2}{3}n. \end{aligned}$$

This completes the proof. \square

For any block graph H of order $k \geq 2$, the 2-corona $G = H \circ P_2$ is a twin-free block graph of order $n = 3k$ and by Observation 2, it has locating-total domination number equal to its total domination number, that is, $\gamma_t^L(G) = \gamma_t(G) = 2k = \frac{2}{3}n$. See Figure 4 for an illustration with H a complete graph. Thus, we obtain the following.

Proposition 7. There are infinitely many connected twin-free block graphs G of order n with $\gamma_t^L(G) = \frac{2}{3}n$.




Figure 4: The 2-corona $K_6 \circ P_2$ of a complete graph of order 6.

5 Subcubic graphs

In this section, we establish a tight upper bound on the locating-total domination number of a subcubic graph, where a subcubic graph is a graph with maximum degree at most 3. For this purpose, let $\mathcal{F}_{\text{tdom}}$ be the family consisting of the three complete graphs K_1 , K_2 , and K_4 , and a star $K_{1,3}$, that is,

$$\mathcal{F}_{\text{tdom}} = \{K_1, K_2, K_4, K_{1,3}\}.$$

We denote a *path*, a *cycle*, and a *complete* graph on n vertices by P_n , C_n , and K_n , respectively. A *diamond* is the graph $K_4 - e$ where e is an arbitrary edge of the K_4 . A *paw* is the graph obtained from a

triangle K_3 by adding a new vertex and joining it with an edge to one vertex of the triangle. Equivalently, a paw is obtained from $K_{1,3}$ by adding an edge between two leaves.

For $k \geq 2$, we say a graph G contains a (d_1, d_2, \dots, d_k) -sequence if there exists a path $v_1 v_2 \dots v_k$ such that $\deg_G(v_i) = d_i$ for all $i \in [k]$. We are now in a position to prove the following upper bound on the locating-total domination number of a subcubic graph.

Theorem 8. If $G \notin \mathcal{F}_{\text{tdom}}$ is a connected subcubic graph of order $n \geq 3$, then $\gamma_t^L(G) \leq \frac{2}{3}n$.

Proof. Suppose, to the contrary, that the theorem is false. Among all counterexamples, let G be one of minimum order n . If $n = 3$, then $G \cong P_3$ or $G \cong K_3$, and in both cases $\gamma_t^L(G) = 2 = \frac{2}{3}n$, a contradiction. Hence, $n \geq 4$. Suppose $n = 4$. By assumption, $G \notin \{K_4, K_{1,3}\}$. If G is a diamond or a paw, then let S consist of one vertex of degree 2 and one vertex of degree 3, and if G is a path or a cycle, then let S consist of two adjacent vertices of degree 2. In all cases, S is a LTD-set of G of cardinality 2, and so $\gamma_t^L(G) \leq 2 < \frac{2}{3}n$, a contradiction. Hence, $n \geq 5$.

Suppose that $n = 5$. If G is a path P_5 or a cycle C_5 , then $\gamma_t^L(G) = 3 < \frac{2}{3}n$ (choose three consecutive vertices of degree 2), a contradiction. Hence, $\Delta(G) = 3$. Let v be a vertex of degree 3 in G with neighbors v_1, v_2, v_3 . Let v_4 be the remaining vertex of G . Since G is connected, we may assume, renaming vertices if necessary, that $v_1 v_4$ is an edge. The set $\{v, v_1, v_2\}$ is a LTD-set of G , and so $\gamma_t^L(G) \leq 3 < \frac{2}{3}n$, a contradiction. Hence, $n \geq 6$.

Suppose that $n = 6$. If G is a path P_6 or a cycle C_6 , then $\gamma_t^L(G) = 4 = \frac{2}{3}n$ (choose four consecutive vertices of degree 2), a contradiction. Hence, $\Delta(G) = 3$. Let v be a vertex of degree 3 in G with neighbors v_1, v_2, v_3 , and let v_4 and v_5 be the two remaining vertices of G . Since G is connected, we may assume, renaming vertices if necessary, that $v_1 v_4$ is an edge. One of the sets $\{v, v_1, v_2, v_4\}$ and $\{v, v_1, v_3, v_4\}$ is a LTD-set of G , and so $\gamma_t^L(G) \leq 4 = \frac{2}{3}n$, a contradiction. Hence, $n \geq 7$.

In what follows, we adopt the notation that if there is a (d_1, d_2, \dots, d_k) -sequence in G , then $P: v_1 v_2 \dots v_k$ denotes a path in G associated with such a sequence, where $\deg_G(v_i) = d_i$ for all $i \in [k]$. Further, we let $G' = G - V(P)$ and let G' have order n' , and so $n' = n - k$. Recall that $n \geq 7$.

We show firstly that there is no vertex of degree 1.

Claim H. $\delta(G) \geq 2$.

Proof of claim. Suppose, to the contrary, that $\delta(G) = 1$. We proceed further with a series of structural properties of the graph G that show that certain (d_1, d_2, \dots, d_k) -sequences are forbidden.

Subclaim H.1. The following properties hold in the graph G .

- (a) There is no $(1, 3, 1)$ -sequence.
- (b) There is no $(1, 2, 2)$ -sequence.
- (c) There is no $(1, 2, 3, 1)$ -sequence.
- (d) There is no $(1, 2, 3, 2, 1)$ -sequence.
- (e) There is no $(1, 2, 3)$ -sequence.
- (f) There is no $(1, 2)$ -sequence.
- (g) There is no $(1, 3, 2)$ -sequence.
- (h) There is no $(1, 3, 3, 1)$ -sequence.

Proof of subclaim. (a) Suppose that there is a $(1, 3, 1)$ -sequence in G . In this case, $n' = n - 3 \geq 4$. Since G is connected, so too is the graph G' . Thus, G' is not a counterexample to our theorem, except possibly when $n = 4$ and $G' \in \mathcal{F}_{\text{tdom}}$. Let v' be the third neighbor of v_2 in G not on the path P . Suppose $G' \in \mathcal{F}_{\text{tdom}}$, implying that $G' \cong K_{1,3}$ with v' as a leaf in G' . The graph G is therefore determined, and has order $n = 7$. In this case, choosing S to consist of the two support vertices (of degree 3) and a leaf neighbor of each support vertex produces a LTD-set of G of cardinality 4, and so $\gamma_t^L(G) \leq 4 < \frac{2}{3}n$. Hence, $G' \notin \mathcal{F}_{\text{tdom}}$. Since G' is not a counterexample, it holds that $\gamma_t^L(G') \leq \frac{2}{3}n' = \frac{2}{3}n - 2$. Every

γ_t^L -set of G' can be extended to a LTD-set of G by adding to it the vertices v_2 and v_3 , implying that $\gamma_t^L(G) \leq \gamma_t^L(G') + 2 \leq \frac{2}{3}n$, a contradiction.

(b) Suppose that there is a $(1, 2, 2)$ -sequence in G . Let v' be the second neighbor of v_3 . As in the previous case, $n' = n - 3 \geq 4$ and G' is connected. By part (a), there is no $(1, 3, 1)$ -sequence, implying that $G' \notin \mathcal{F}_{\text{tdom}}$ and $\gamma_t^L(G') \leq \frac{2}{3}n' = \frac{2}{3}n - 2$. As before every γ_t^L -set of G' can be extended to a LTD-set of G by adding to it the vertices v_2 and v_3 , implying that $\gamma_t^L(G) \leq \frac{2}{3}n$, a contradiction.

(c) Suppose that there is a $(1, 2, 3, 1)$ -sequence in G . In this case, $n' = n - 4 \geq 3$ and G' is connected. By part (a), there is no $(1, 3, 1)$ -sequence, implying that $G' \notin \mathcal{F}_{\text{tdom}}$ and $\gamma_t^L(G') \leq \frac{2}{3}n' = \frac{2}{3}(n-4) < \frac{2}{3}n - 2$. Every γ_t^L -set of G' can be extended to a LTD-set of G by adding to it the vertices v_2 and v_3 , implying that $\gamma_t^L(G) \leq \frac{2}{3}n$, a contradiction.

(d) Suppose that there is a $(1, 2, 3, 2, 1)$ -sequence in G . In this case, G' is connected and $n' = n - 5 \geq 2$. If $G' \in \mathcal{F}_{\text{tdom}}$, then $G' \cong K_2$ by the fact that there is no $(1, 3, 1)$ -sequence in G by part (a). The graph G is therefore determined, and is obtained from a star $K_{1,3}$ by subdividing every edge once. We note that G has order $n = 7$ and the set $N[v_3]$ (of non-leaves of G) is a LTD-set of G , implying that $\gamma_t^L(G) \leq 4 < \frac{2}{3}n$, a contradiction. Hence, $G' \notin \mathcal{F}_{\text{tdom}}$. Thus, $\gamma_t^L(G') \leq \frac{2}{3}n' = \frac{2}{3}(n-5) < \frac{2}{3}n - 3$. Every γ_t^L -set of G' can be extended to a LTD-set of G by adding to it the vertices v_2, v_3 , and v_4 , implying that $\gamma_t^L(G) < \frac{2}{3}n$, a contradiction.

(e) Suppose that there is a $(1, 2, 3)$ -sequence in G . In this case, $n' = n - 3 \geq 4$ and G' contains at most two components. Let v_4 and v'_4 be the two neighbors of v_3 different from v_2 . By our earlier observations, each of v_4 and v'_4 has degree at least 2 in G , and therefore degree at least 1 in G' .

Suppose that G' is disconnected. In this case, since there is no $(1, 3, 1)$ -sequence, no $(1, 2, 3, 1)$ -sequence, and no $(1, 2, 3, 2, 1)$ -sequence in G , neither component of G' belongs to $\mathcal{F}_{\text{tdom}}$. By linearity, we therefore have that $\gamma_t^L(G') \leq \frac{2}{3}n' = \frac{2}{3}n - 2$. Every γ_t^L -set of G' can be extended to a LTD-set of G by adding to it the vertices v_2 and v_3 , implying that $\gamma_t^L(G) \leq \frac{2}{3}n$, a contradiction. Hence, G' is connected. Recall that $n' \geq 4$.

Suppose now that G' is connected. If $G' \in \mathcal{F}_{\text{tdom}}$, then $G' \cong K_{1,3}$. Let v_5 be the central vertex of G' , and so each of v_4 and v'_4 is a leaf neighbor of v_5 in G' . The graph G is therefore determined and $n = 7$. The set $\{v_2, v_3, v_4, v_5\}$ is a LTD-set of G , implying that $\gamma_t^L(G) \leq 4 < \frac{2}{3}n$, a contradiction. Hence, $G' \notin \mathcal{F}_{\text{tdom}}$. Thus, $\gamma_t^L(G') \leq \frac{2}{3}n' = \frac{2}{3}n - 2$. Every γ_t^L -set of G' can be extended to a LTD-set of G by adding to it the vertices v_2 and v_3 , implying that $\gamma_t^L(G) \leq \frac{2}{3}n$, a contradiction.

(f) Since there is no $(1, 2, 1)$ -sequence (since $n \geq 7$), no $(1, 2, 2)$ -sequence by (b) and no $(1, 2, 3)$ -sequence by (e), there can be no $(1, 2)$ -sequence in G . Hence, part (f) follows immediately from parts (b) and (e).

(g) Suppose that there is a $(1, 3, 2)$ -sequence in G . In this case, $n' = n - 3 \geq 4$. If G' is disconnected, then by parts (a)–(f), neither component of G' belongs to $\mathcal{F}_{\text{tdom}}$. By linearity, we therefore have that $\gamma_t^L(G') \leq \frac{2}{3}n' = \frac{2}{3}n - 2$. Every γ_t^L -set of G' can be extended to a LTD-set of G by adding to it the vertices v_2 and v_3 , implying that $\gamma_t^L(G) \leq \frac{2}{3}n$, a contradiction. Hence, G' is connected. If $G' \in \mathcal{F}_{\text{tdom}}$, then $G' \cong K_{1,3}$. In this case, the graph G has order $n = 7$ and is obtained from a 5-cycle by selecting two non-adjacent vertices on the cycle and adding a pendant edge to these two vertices. In this case, the set consisting of the two vertices of degree 3 and any two vertices of degree 2 is a LTD-set of G , implying that $\gamma_t^L(G) \leq 4 < \frac{2}{3}n$, a contradiction. Hence, $G' \notin \mathcal{F}_{\text{tdom}}$. Thus, $\gamma_t^L(G') \leq \frac{2}{3}n' = \frac{2}{3}n - 2$. Every γ_t^L -set of G' can be extended to a LTD-set of G by adding to it the vertices v_2 and v_3 , implying that $\gamma_t^L(G) \leq \frac{2}{3}n$, a contradiction.

(h) Suppose that there is a $(1, 3, 3, 1)$ -sequence in G . In this case, $n' = n - 4 \geq 3$ and G' contains at most two components. Let u_i be the neighbor of v_i not on P for $i \in \{2, 3\}$. Possibly, $u_2 = u_3$. By parts (a) and (g), the vertex u_i has degree 3 in G for $i \in \{2, 3\}$. Suppose that G' is disconnected. In this case, by parts (a)–(g), neither component of G' belongs to $\mathcal{F}_{\text{tdom}}$. By linearity, we therefore have that $\gamma_t^L(G') \leq \frac{2}{3}n' = \frac{2}{3}(n-4) < \frac{2}{3}n - 2$. Every γ_t^L -set of G' can be extended to a LTD-set of G by adding to it the vertices v_2 and v_3 , implying that $\gamma_t^L(G) < \frac{2}{3}n$, a contradiction. Hence, G' is connected. Recall that $n' \geq 3$. By parts (a)–(g), we note that $G' \notin \mathcal{F}_{\text{tdom}}$, implying that $\gamma_t^L(G') \leq \frac{2}{3}n' < \frac{2}{3}n - 2$. Every γ_t^L -set of G' can be extended to a LTD-set of G by adding to it the vertices v_2 and v_3 , implying that $\gamma_t^L(G) < \frac{2}{3}n$, a contradiction.

Thus, the proof of the subclaim is complete. \diamond

We now return to the proof of Claim H. By Subclaim H.1(f), the neighbor of every vertex of degree 1 has degree 3 in G . Further by Subclaim H.1(g), such a vertex of degree 3 has both its other two neighbors of degree 3. Therefore the existence of a vertex of degree 1 implies that there is a $(1, 3, 3)$ -sequence in G . In this case, $n' = n - 3 \geq 4$. Let u_2 be the neighbor of v_2 not on P , and let u_3 and w_3 be the two neighbors of v_3 not on P . By our earlier observations, the vertex u_2 has degree 3 in G , and, by Subclaim H.1(h), both vertices u_3 and w_3 have degree at least 2 in G .

Suppose that G' contains a component that belongs to $\mathcal{F}_{\text{tdom}}$. By Subclaim H.1, this is only possible if u_3 and w_3 are adjacent and both vertices have degree 2 in G . In this case, $G[\{v_3, u_3, w_3\}]$ is a triangle in G . We now consider the connected graph $G^* = G - \{v_1, v_2, v_3, u_3, w_3\}$ of order $n^* = n - 5$. Since u_2 has degree 2 in G^* , we note that $n^* \geq 3$ and $G^* \notin \mathcal{F}_{\text{tdom}}$. Hence, $\gamma_t^L(G^*) \leq \frac{2}{3}n^* = \frac{2}{3}(n - 5) < \frac{2}{3}n - 3$. Every γ_t^L -set of G^* can be extended to a LTD-set of G by adding to it the vertices v_2, v_3 , and u_3 , implying that $\gamma_t^L(G) \leq \gamma_t^L(G^*) + 3 < \frac{2}{3}n$, a contradiction. Hence, no component of G' belongs to the family $\mathcal{F}_{\text{tdom}}$. By linearity, we therefore have that $\gamma_t^L(G') \leq \frac{2}{3}n' = \frac{2}{3}n - 2$. Every γ_t^L -set of G' can be extended to a LTD-set of G by adding to it the vertices v_2 and v_3 , implying that $\gamma_t^L(G) < \frac{2}{3}n$, a contradiction. This completes the proof of Claim H. \square

By Claim H, every vertex in G has degree 2 or 3.

Claim I. The graph G is triangle-free.

Proof of claim. Suppose that G contains a triangle K_3 . Among all triangles in G , let T contain the maximum number of vertices of degree 2 in G . Let $V(T) = \{v_1, v_2, v_3\}$, where $2 \leq \deg_G(v_1) \leq \deg_G(v_2) \leq \deg_G(v_3) \leq 3$. Since $n \geq 7$, the triangle T contains at most two vertices of degree 2, and so $\deg_G(v_3) = 3$. Let $G' = G - V(T)$ and let G' have order n' , and so $n' = n - 3 \geq 4$.

Suppose that $\deg_G(v_1) = 2$. We note that $\deg_G(v_2) = 2$ or $\deg_G(v_2) = 3$. Since every vertex in G has degree 2 or 3, no component of G' belongs to $\mathcal{F}_{\text{tdom}}$. Hence by linearity, $\gamma_t^L(G') \leq \frac{2}{3}n' = \frac{2}{3}n - 2$. Every γ_t^L -set of G' can be extended to a LTD-set of G by adding to it the vertices v_2 and v_3 , implying that $\gamma_t^L(G) < \frac{2}{3}n$, a contradiction. Hence, $\deg_G(v_1) = 3$, implying that every vertex in T has degree 3 in G . Hence by our choice of the triangle T , no vertex of degree 2 in G belongs to a triangle.

Let u_i be the neighbor of v_i not in the triangle T for $i \in [3]$. We note that the vertices u_1, u_2 , and u_3 are not necessarily distinct. Suppose that G' contains no component that belongs to $\mathcal{F}_{\text{tdom}}$. By linearity, this yields $\gamma_t^L(G') \leq \frac{2}{3}n' = \frac{2}{3}n - 2$. Every γ_t^L -set of G' can be extended to a LTD-set of G by adding to it the vertices v_2 and v_3 , implying that $\gamma_t^L(G) \leq \frac{2}{3}n$, a contradiction. Hence, G' contains a component that belongs to $\mathcal{F}_{\text{tdom}}$. Since $n \geq 7$ and no vertex of degree 2 in G belongs to a triangle, this is only possible if either $G' \cong K_{1,3}$ or if G' contains a K_2 -component.

On the one hand, if $G' \cong K_{1,3}$, then the three vertices u_1, u_2 , and u_3 are leaves in G' that are adjacent to a common neighbor (of degree 3) in G' . In this case, the graph G is determined and $n = 7$, and the set $V(T) \cup \{u_1\}$ is a LTD-set of G , implying that $\gamma_t^L(G) \leq 4 < \frac{2}{3}n$, a contradiction.

On the other hand, if G' contains a K_2 -component, then renaming vertices if necessary, we may assume that u_1 and u_2 belong to such a component. We note that u_1 and u_2 both have degree 2 in G , and $u_1v_1v_2u_2u_1$ is a 4-cycle in G . Further we note that in this case, G' contains two components, where the second component contains the vertex u_3 . We now consider the graph $G^* = G - \{v_1, v_2, v_3, u_1, u_2\}$. Let G^* have order $n^* = n - 5$. By the fact that $\delta(G) \geq 2$ by Claim H, the graph $G^* \notin \mathcal{F}_{\text{tdom}}$, implying that $\gamma_t^L(G^*) \leq \frac{2}{3}n^* = \frac{2}{3}(n - 5) < \frac{2}{3}n - 3$. Every γ_t^L -set of G^* can be extended to a LTD-set of G by adding to it, for example, the vertices u_2, v_2 and v_3 , implying that $\gamma_t^L(G) < \frac{2}{3}n$, a contradiction. \square

By Claim I, the graph G is triangle-free. We show next that there is no vertex of degree 2.

Claim J. The graph G is a cubic graph.

Proof of claim. Suppose, to the contrary, that $\delta(G) = 2$. As before, we obtain a series of structural properties of the graph G that show that certain (d_1, d_2, \dots, d_k) -sequences are forbidden. These forbidden sequences will enable us to deduce the desired result of the claim that G must be a cubic graph.

Subclaim J.1. The following properties hold in the graph G .

- (a) There is no $(2, 2, 2)$ -sequence.
- (b) There is no $(2, 3, 2)$ -sequence.
- (c) There is no $(2, 2, 3)$ -sequence.
- (d) There is no $(2, 2)$ -sequence.
- (e) There is no $(2, 3, 3)$ -sequence.

Proof of subclaim. (a) Suppose that there is a $(2, 2, 2)$ -sequence in G . In this case, $n' = n - 3 \geq 4$. Since $n \geq 7$, $\delta(G) = 2$, and G contains no triangle, no component of G' belongs to $\mathcal{F}_{\text{tdom}}$. Hence by linearity, $\gamma_t^L(G') \leq \frac{2}{3}n' = \frac{2}{3}n - 2$. Every γ_t^L -set of G' can be extended to a LTD-set of G by adding to it the vertices v_2 and v_3 , implying that $\gamma_t^L(G) \leq \frac{2}{3}n$, a contradiction.

(b) Suppose that there is a $(2, 3, 2)$ -sequence in G . As before, $n' = n - 3 \geq 4$. Suppose that G' contains a component that belongs to $\mathcal{F}_{\text{tdom}}$. Since there is no $(2, 2, 2)$ -sequence and $n \geq 7$, and since $\delta(G) \geq 2$ and G contains no triangle, this is only possible if $G' \cong K_{1,3}$. But then the graph G is determined and $n = 7$, and $\gamma_t^L(G') = 4 < \frac{2}{3}n$ (by considering the set $N[v_2]$), a contradiction. Hence, no component of G' belongs to $\mathcal{F}_{\text{tdom}}$. By linearity, this yields $\gamma_t^L(G') \leq \frac{2}{3}n' = \frac{2}{3}n - 2$. Every γ_t^L -set of G' can be extended to a LTD-set of G by adding to it the vertices v_2 and v_3 , implying that $\gamma_t^L(G) \leq \gamma_t^L(G') + 2 \leq \frac{2}{3}n$, a contradiction.

(c) Suppose that there is a $(2, 2, 3)$ -sequence in G . Since there is no $(2, 2, 2)$ -sequence and no $(2, 3, 2)$ -sequence in G , every vertex different from v_2 that is adjacent to v_1 or v_3 has degree 3 in G . Together with our earlier observations, the graph G' therefore cannot contain a component that belongs to $\mathcal{F}_{\text{tdom}}$. By linearity, we have $\gamma_t^L(G') \leq \frac{2}{3}n' = \frac{2}{3}n - 2$. As before this yields $\gamma_t^L(G) \leq \gamma_t^L(G') + 2 \leq \frac{2}{3}n$, a contradiction.

(d) Since there is no $(2, 2, 2)$ -sequence and no $(2, 2, 3)$ -sequence, there can be no $(2, 2)$ -sequence in G noting that every vertex has degree 2 or 3.

(e) Suppose that there is a $(2, 3, 3)$ -sequence in G . Since there is no $(2, 2, 2)$ -sequence, no $(2, 3, 2)$ -sequence, and no $(2, 2, 3)$ -sequence in G , the graph G' cannot contain a component that belongs to $\mathcal{F}_{\text{tdom}}$. By linearity, this yields $\gamma_t^L(G') \leq \frac{2}{3}n' = \frac{2}{3}n - 2$. Every γ_t^L -set of G' can be extended to a LTD-set of G by adding to it the vertices v_2 and v_3 , implying that $\gamma_t^L(G) \leq \gamma_t^L(G') + 2 \leq \frac{2}{3}n$, a contradiction.

Thus, the proof of the subclaim is complete. (\diamond)

By Subclaim J.1(d), there is no $(2, 2)$ -sequence. Hence every vertex of degree 2 has both its neighbors of degree 3. Moreover since there is no $(2, 3, 2)$ -sequence, every vertex of degree 3 has at most one neighbor of degree 2. But this would imply the existence of a $(2, 3, 3)$ -sequence, contradicting Subclaim J.1(e). Therefore, there can be no vertex of degree 2 in G , that is, G is a cubic graph. This completes the proof of Claim J. (\square)

By Claim J, the graph G is a cubic graph. Recall that G is triangle-free. We now consider a $(3, 3, 3)$ -sequence. The graph G' cannot contain a component that belongs to $\mathcal{F}_{\text{tdom}}$. By linearity, this yields $\gamma_t^L(G') \leq \frac{2}{3}n' = \frac{2}{3}n - 2$. Every γ_t^L -set of G' can be extended to a LTD-set of G by adding to it the vertices v_2 and v_3 , implying that $\gamma_t^L(G) \leq \gamma_t^L(G') + 2 \leq \frac{2}{3}n$, a contradiction. This completes the proof of Theorem 8. (\square)

For $k \geq 3$, the 2-corona $G = C_k \circ P_2$ of a cycle C_k has order $n = 3k$ and by Observation 2, it has locating total domination number equal to its total domination number, that is, $\gamma_t^L(G) = \gamma_t(G) = 2k = \frac{2}{3}n$. See Figure 5 for an illustration. Moreover for $k \geq 1$, the 2-corona $G = P_k \circ P_2$ of a path P_k has order $n = 3k$ and also satisfies $\gamma_t^L(G) = \gamma_t(G) = 2k = \frac{2}{3}n$. Thus, we obtain the following.

Proposition 9. There are infinitely many connected twin-free subcubic graphs G of order n with $\gamma_t^L(G) = \frac{2}{3}n$.




Figure 5: The 2-corona $C_6 \circ P_2$ of a 6-cycle.

6 Outerplanar graphs

In this section, we prove that Conjecture 1 holds for twin-free outerplanar graphs. The conjecture was shown to hold for trees in [18] (as trees are C_4 -free). Thus, we focus on outerplanar graphs that are not trees. We use the following well-known characterisation of outerplanar graphs in our proofs.

Theorem 10 ([12]). A graph G is outerplanar if and only if it does not contain K_4 or $K_{2,3}$ as a minor.

A leg $l_0 \cdots l_k$ at l_0 of length $k \geq 1$ is an induced subgraph of G such that it is isomorphic to a path of length k , and $\deg_G(l_0) \geq 3$, $\deg_G(l_k) = 1$, and $\deg_G(l_i) = 2$ for all $i \in \{1, \dots, k-1\}$. When there exists a leg $l_0 \cdots l_k$, we say that vertex l_0 has a leg.

Lemma 11. Let G be a connected twin-free outerplanar graph of order n that is not a tree. If the smallest component of $G - e$ has at most three vertices for every bridge $e \in E(G)$, then $\gamma_t^L(G) \leq \frac{2}{3}n$.

Proof. Since G is twin-free, every bridge of G is an edge of a leg of length at most 3. The structure of G can be described as follows. Denote by G' the graph we obtain from G by removing all bridges and resulting isolated vertices. The graph G' is a connected outerplanar graph with $\delta(G') \geq 2$. We denote the maximal 2-connected components of G' by G'_i , where $i \in \{1, \dots, k\}$. As the 2-connected components form a tree-like structure, we choose the indexing of the components G'_i so that

$$|V(G'_i) \cap \bigcup_{j < i} V(G'_j)| \leq 1$$

for all $i \in \{1, \dots, k\}$ (such an indexing can be found by choosing a leaf-component as a root and indexing the components in a breadth-first ordering starting from the root). We denote by G_i the induced subgraph of G obtained by adding the necessary legs to G'_i for each $i \in \{1, \dots, k\}$. If a cut-vertex v separating some 2-connected components has legs, then those legs are in the subgraph G_i where $i = \min\{j \mid v \in V(G'_j)\}$ and no others. Now, we have $|V(G_i) \cap V(G_j)| \leq 1$ when $i \neq j$, and $V(G) = \bigcup_{i=1}^k V(G_i)$.

In what follows, we construct a LTD-set $S \subseteq V(G)$ of G by considering the subgraphs G_i in the order of their indexing. Denote by c_i the unique element of $V(G'_i) \cap \bigcup_{j < i} V(G'_j)$ if such a vertex exists (i.e. the vertex c_i is the cut-vertex separating G_i from previous subgraphs G_j). If c_i exists, then whether c_i is contained in S or not has already been decided when considering some previous subgraph G_j , and this decision will not change in later steps. Recall that any 2-connected outerplanar graph is Hamiltonian [12]. Let $v_0^i v_1^i \cdots v_{n_i}^i v_0^i$ be a Hamiltonian cycle of G'_i such that $v_0^i = c_i$ if c_i exists, otherwise v_0^i can be chosen arbitrarily. We include vertices from G_i to the set S as follows:

1. For each leg $l_0 \cdots l_k$, let

$$\begin{cases} l_0 \in S & \text{if } k = 1, \\ l_0, l_1 \in S & \text{if } k = 2, \\ l_1, l_2 \in S & \text{if } k = 3, \end{cases}$$

(notice that $l_0 = v_j^i$ for some j).




Figure 6: Illustrations for step 3 of the construction in the proof of Lemma 11. The black vertices are in S by steps 1 and 3. The dotted lines depict a possible partition of the vertices to obtain an upper bound on $|S|$.

2. Assume that v_j^i has a leg of length 1, and v_{j-1}^i has at least one leg. If $v_{j+1}^i \neq c_i$ has no legs, then let $v_{j+1}^i \in S$. If $v_{j+1}^i = c_i$, then let $v_{j-1}^i \in S$.
Similarly, assume that v_j^i has a leg of length 1, and v_{j+1}^i has at least one leg. If $v_{j-1}^i \neq c_i$ has no legs, then let $v_{j-1}^i \in S$. If $v_{j-1}^i = c_i$, then let $v_{j+1}^i \in S$.
3. If v_j^i has a leg of length 1, and v_{j-1}^i and v_{j+1}^i have no legs, then let

$$\begin{cases} v_0^i \in S & \text{if } j \in \{1, n_i\} \text{ and } v_0^i \neq c_i, \\ v_{n_i-1}^i \in S & \text{if } j = n_i \text{ and } v_0^i = c_i, \\ v_1^i \in S & \text{if } j = 2, \\ v_{j+1}^i \in S & \text{otherwise.} \end{cases}$$

See Figure 6 for illustrations of the different cases.

4. Relabel the vertices v_j^i that have not been included in S in the previous steps as $u_0^i, \dots, u_{n_i'}^i$ such that $u_0^i = v_m^i$, where $m = \min\{j \mid v_j^i \neq c_i \text{ and } v_j^i \notin S\}$, and if $j < l$, $u_j^i = v_j^i$, and $u_l^i = v_{l'}^i$, then $j' < l'$ (for some $l \in [n_i']$ and $l' \in [n_i]$). Let $u_j^i \in S$ if and only if $j \equiv 1, 2 \pmod{3}$, except let $u_{n_i'}^i \notin S$ and $u_{n_i'-1}^i \in S$ when $n_i' \equiv 1 \pmod{3}$.

See Figure 7 for an example of the set S in one subgraph G_i . Notice that the edges of G_i' that are not in the Hamiltonian cycle do not affect the construction. The vertices outside the dotted outlines (other than c_i) are the vertices u_j^i considered in step 4. Notice that in this example we have the exceptional case of step 4.

Claim K. The set S is a total dominating set of G .

Proof of claim. Let $v \in V(G_i) \setminus V(G_i')$ for some i . Now $v = l_j$ for some leg $l_0 \cdots l_k$ and $j \neq 0$. Due to step 1 of the construction of S , we have $l_{j-1} \in S$ or $l_{j+1} \in S$.

Let us then consider the vertices of G_i' for some i . Consider a vertex $v_j^i \neq c_i$. Due to step 1, if the vertex v_j^i has a leg of length 2 or 3, then $l_1 \in N(v_j^i) \cap S$ for such a leg.

Assume that the vertex v_j^i has a leg of length 1. If at least one of v_{j-1}^i and v_{j+1}^i has no legs, then one of them is in $N(v_j^i) \cap S$ by step 2 or 3. Assume that both v_{j-1}^i and v_{j+1}^i have legs. Suppose to the contrary that $v_{j-1}^i \notin S$ and $v_{j+1}^i \notin S$. Since $v_j^i \in S$ by step 1, we have $v_{j-1}^i = u_{j'}^i$, and $v_{j+1}^i = u_{j'+1}^i$ for some $j' \in \{0, \dots, n_i' - 1\}$. However, according to step 4, we have $u_{j'}^i \in S$ or $u_{j'+1}^i \in S$, a contradiction.




Figure 7: An example of a subgraph G_i . The black vertices are included in the set S constructed in the proof of Lemma 11. The cycle with the solid edges is a Hamiltonian cycle of G_i' , and the dashed lines within the cycle represent edges of G_i' not in the Hamiltonian cycle. The dotted lines depict a possible partition of the vertices considered in steps 1-3 to obtain an upper bound on $|S|$.

Assume then that the vertex v_j^i has no legs. If v_{j-1}^i or v_{j+1}^i has a leg of length 1 or 2, then v_{j-1}^i or v_{j+1}^i is in $N(v_j^i) \cap S$ due to step 1. Assume that v_{j-1}^i and v_{j+1}^i have no legs of length 1 or 2. Suppose to the contrary that $v_{j-1}^i \notin S$ and $v_{j+1}^i \notin S$. Now $v_j^i = u_{j'}^i$, $v_{j-1}^i = u_{j'-1}^i$, and $v_{j+1}^i = u_{j'+1}^i$ for some $j' \in \{0, \dots, n_i'\}$, and $N(v_j^i) \cap S$ is nonempty due to step 4.

The vertex c_i also has a neighbor in S , however, this neighbor is not necessarily in $V(G_i)$. Let G_j be the subgraph with the smallest index that contains c_i . Now, $c_i \neq c_j$ and c_i has a neighbor in S by the arguments above. (\square)

The following claim is used repeatedly in the final part of the proof where we prove that the set S is locating.

Claim L. Let $v_j^i \notin S$ such that $v_j^i \neq c_i$. If $j \neq n_i$, then $v_{j+1}^i \in S$, and if $j \notin \{0, 1\}$, then $v_{j-1}^i \in S$.

Proof of claim. Since $v_j^i \notin S$ and $v_j^i \neq c_i$, we have $v_j^i = u_{j'}^i$ for some $j' \in \{0, \dots, n_i'\}$.

Assume that $j \neq n_i$. Suppose, to the contrary, that $v_{j+1}^i \notin S$. Since $j \neq n_i$, we have $v_{j+1}^i \neq c_i$. Now $v_{j+1}^i = u_{j'+1}^i$, and due to step 4, either $v_j^i \in S$ or $v_{j+1}^i \in S$, a contradiction.

Assume then that $j \notin \{0, 1\}$, and suppose, to the contrary, that $v_{j-1}^i \notin S$. Now $j-1 \geq 1$, and thus $v_{j-1}^i \neq c_i$. As in the previous case, we have $v_{j-1}^i = u_{j'-1}^i$, and either $v_j^i \in S$ or $v_{j-1}^i \in S$, a contradiction. (\square)

Claim M. The set S is a LTD-set of G .

Proof of claim. By Claim K, the set S is a TD-set of G . What remains to be shown is that for all distinct $v, w \notin S$, we have $N(v) \cap S \neq N(w) \cap S$.

Consider a vertex v that is a leaf. Suppose, to the contrary, that there exists a vertex $w \in V(G) \setminus S$ such that $w \neq v$ and $N(v) \cap S = N(w) \cap S$. Since the vertex v is a leaf, its support vertex is in S due to step 1 of the construction of S . The vertex w is not a leaf, since otherwise the graph G would have

twins. Due to step 1, the vertex w cannot be in the same leg as v . Thus, $w = v_j^i$ for some i and j , and v is a leaf of a leg of length 1. Without loss of generality, we may assume that i is the minimum index such that $w \in V(G'_i)$. This implies that $w \neq c_i$. Since $w \notin S$, due to step 1 the vertex w either has no legs or all the legs of w are of length 3. If w has a leg of length 3, then that leg contains an element of $N(w) \cap S$ that is not in $N(v)$. Thus, the vertex w has no legs. Since the vertex v is a leaf of a leg of length 1, we have $N(v) \cap S = \{v_l^{i'}\}$ for some l and i' . If $j \notin \{0, 1, n_i\}$, then $|N(w) \cap S| \geq 2$ due to Claim L as $w \neq c_i$. Assume that $j \in \{0, 1, n_i\}$.

- If $j = 1$, then $v_2^i \in N(w) \cap S$ due to Claim L. Consequently, $v_l^{i'} = v_2^i$. Since $w = v_1^i$ has no legs and v_2^i has a leg of length 1, we have $w \in S$ due to step 2 or 3, a contradiction.
- If $j = n_i$, then $v_{n_i-1}^i \in N(w) \cap S$ due to Claim L. Consequently, $v_l^{i'} = v_{n_i-1}^i$. Since $w = v_{n_i}^i$ has no legs and $v_{n_i-1}^i$ has a leg of length 1, we have $w \in S$ due to step 2 or 3, a contradiction.
- Assume that $j = 0$. According to (the proof of) Claim K and the minimality of i , the vertex w has a neighbor in $V(G_i) \cap S$. Now $N(v) \cap S = \{v_l^{i'}\} = N(w) \cap S$. Since $w \neq c_i$ and $w \notin S$, we have $w = v_0^i$ and $v_1^i \in S$, which implies that $l = 1$ and $i = i'$. However, now $w \in S$ due to step 2 or 3, a contradiction.

Thus, all leaves are located by S .

Consider then the case where v and w are in the same 2-connected component of G , that is, $v = v_j^i$ and $w = v_l^i$ for some j, l and i . Assume without loss of generality that $j < l$.

- Assume that $v_j^i \neq c_i$. Since $j < l \leq n_i$, we have $v_{j+1}^i \in S$ according to Claim L. This implies that $l \geq 2$, and thus $v_{l-1}^i \in S$. If $j+1 = l-1$, then having both $v_j^i \notin S$ and $v_l^i \notin S$ contradicts step 4 of the construction of S . Thus, $j+1 \neq l-1$. However, now $N(v_j^i) \cap S = N(v_l^i) \cap S$ implies that the graph G contains K_4 as a minor. Indeed, the vertices v_{j+1}^i and v_{l-1}^i are both neighbors of v_j^i and v_l^i . Since $v_0^i \cdots v_j^i v_{j+1}^i \cdots v_{l-1}^i v_l^i \cdots v_{n_i}^i v_0^i$ is a Hamiltonian cycle of G'_i , there exist vertex disjoint paths $v_{j+1}^i - v_{l-1}^i$ and $v_l^i - v_j^i$, whose edges can be contracted to obtain the minor K_4 . Now the graph G is not outerplanar according to Theorem 10, a contradiction.
- Assume that $j = 0$ and $v = v_j^i = c_i$. Now for some $i' < i$, we have $v \in V(G_{i'})$ and $w \notin V(G_{i'})$. Due to (the proof of) Claim K, the vertex v has a neighbor in $V(G_{i'}) \cap S$. This neighbor cannot be a neighbor of w , since the only vertex of G_i that can have neighbors in $G_{i'}$ is the cut-vertex v .

Consider finally the case where v and w belong to two different 2-connected components of G , that is, $v = v_j^i$ and $w = v_l^{i'}$ for some j, l, i and i' with $i \neq i'$. If $V(G_i) \cap V(G_{i'}) = \emptyset$, then v and w are clearly located, since S is a total dominating set by Claim K. Assume that $V(G_i) \cap V(G_{i'}) \neq \emptyset$. The only vertex G_i and $G_{i'}$ share is the cut-vertex separating them. If either of v or w is this cut-vertex, then $v, w \in V(G_i)$ or $v, w \in V(G_{i'})$, and we can handle this case as in the previous case. So assume that neither v nor w is the cut-vertex separating G_i and $G_{i'}$. Assume without loss of generality that $i' < i$. Now the cut-vertex separating G_i and $G_{i'}$ is c_i . Since $v = v_j^i \neq c_i$, we have $v_{j-1}^i \in S$ and $v_{j-1}^i \neq c_i$, or $v_{j+1}^i \in S$ and $v_{j+1}^i \neq c_i$ due to Claim L. Thus, the vertex v has a neighbor in S that is not a neighbor of w . (□)

To conclude the proof, we note that $|S| \leq \frac{2}{3}n$. This is easy to see when each $V(G_i) \setminus \{c_i\}$ is partitioned into small sets according to the steps of the construction as illustrated in Figures 6 and 7. The dotted outlines group vertices considered in steps 1-3. Each set of three vertices contains two elements of S , and each set of two elements contains one element of S . The vertices outside the dotted lines are the vertices u_j^i considered in step 4, of which at most two thirds are in S . □

Theorem 12. If G is a connected twin-free isolate-free outerplanar graph of order n , then $\gamma_t^L(G) \leq \frac{2}{3}n$.

Proof. We prove the claim by induction on n . If the graph G is a tree, the claim holds by [18] as trees do not contain 4-cycles. If G is not a tree and does not contain a bridge $e \in E(G)$ such that both components of $G - e$ have at least four vertices, then the claim holds by Lemma 11.




Figure 8: Illustrations for Case 1, where (a) $|N_{G_1}(v) \setminus \{u\}| = 1$, and (b) $|N_{G_1}(v) \setminus \{u\}| = 2$.

In particular, for sufficiently small n , the claim holds by Lemma 11, or the graph G is a tree and the claim holds due to [18].

Thus, assume that G contains a bridge $e \in E(G)$ such that both components of $G - e$ have at least four vertices.

For the inductive step, assume that when G' is a connected twin-free outerplanar graph of order $n' < n$, then there exists a LTD-set $S \subset V(G')$ such that $|S| \leq \frac{2}{3}n'$.

The graph $G - e$ consists of two components G_1 and G_2 . Assume that both G_1 and G_2 are twin-free. By the induction hypothesis, there exist LTD-sets S_1 and S_2 of G_1 and G_2 , respectively, such that $|S_1| \leq \frac{2}{3}n_1$ and $|S_2| \leq \frac{2}{3}n_2$, where $n_i = |V(G_i)|$. The set $S = S_1 \cup S_2$ is clearly a LTD-set of G . Thus, $\gamma_t^L(G) \leq \frac{2}{3}n_1 + \frac{2}{3}n_2 = \frac{2}{3}n$.

Assume then that G_1 is not twin-free. Let $v \in V(G_1)$ be the end-point of e in G_1 . There exists exactly one pair of twins, and v is one of the twins (otherwise the graph G is not twin-free). Let u be the twin of v in G_1 . If $|N_{G_1}(v) \setminus \{u\}| \geq 3$, then v and u together with their shared neighborhood contains $K_{2,3}$, and the graph G is not outerplanar according to Theorem 10. Thus, $|N_{G_1}(v) \setminus \{u\}| \in \{1, 2\}$. We denote $N_{G_1}(v) \setminus \{u\} = \{s\}$ and $N_{G_1}(v) \setminus \{u\} = \{s, t\}$ when $|N_{G_1}(v) \setminus \{u\}|$ equals 1 and 2, respectively. In what follows, we have combined the case analyses of both of these cases. Thus, the vertex t is included in the case analysis, when in fact t may not exist. The places where the existence of t truly matters are clearly indicated. In other places, keep in mind that t may or may not exist.

Instead of $G - e$, let us now consider the graph $G - \{vs, vt, vu\}$ (note that the edge vu may or may not be present). The graph $G - \{vs, vt, vu\}$ has two components, which are clearly outerplanar, and we denote them by G'_1 and G'_2 so that $V(G'_1) = V(G_1) \setminus \{v\}$ and $V(G'_2) = V(G_2) \cup \{v\}$ (see Figure 8).

Case 1. Assume that G'_1 is twin-free. By the induction hypothesis, there exists a LTD-set S_1 of G'_1 such that $|S_1| \leq \frac{2}{3}n'_1$. Notice that since the set S_1 totally dominates the vertex u , we have $\{s, t\} \cap S_1 \neq \emptyset$.

Let us then consider G'_2 . Notice that the vertex v is a leaf in G'_2 . Denote the support vertex of v by s' . Assume that G'_2 is twin-free. By the induction hypothesis, there exists a LTD-set S_2 of G'_2 such that $|S_2| \leq \frac{2}{3}n'_2$. Notice that $s' \in S_2$, since s' is a support vertex. Assume that G'_2 is not twin-free. Then, there exists a leaf $v' \neq v$ attached to s' . Now the graph $G'_2 - v$ is connected, outerplanar, and twin-free. By the induction hypothesis, there exists a LTD-set S'_2 of $G'_2 - v$ such that $|S'_2| \leq \frac{2}{3}(n'_2 - 1)$. Moreover, we have $s' \in S'_2$, since s' is a support vertex in $G'_2 - v$. Notice that the set S'_2 is a TD-set of G'_2 . However, it might not be locating, since $N_{G'_2}(v) \cap S'_2 = N_{G'_2}(v') \cap S'_2 = \{s'\}$.

Now, either the set $S = S_1 \cup S_2$ or $S' = S_1 \cup S'_2$ is a LTD-set of G depending on whether the graph G'_2 has twins. Indeed, the vertex v is the unique vertex that is dominated by a vertex of G'_1 and a vertex of G'_2 . Since $|S| \leq \frac{2}{3}n'_1 + \frac{2}{3}n'_2 = \frac{2}{3}n$ and $|S'| \leq \frac{2}{3}n'_1 + \frac{2}{3}(n'_2 - 1) < \frac{2}{3}n$, the claim holds when G'_1 is twin-free.

Case 2. Assume that G'_1 contains twins. Then, one of the twins must be a neighbor of v in G . If $|N_{G_1}(v) \setminus \{u\}| = 1$, then one of the twins is u , since u is a leaf in G'_1 , and thus s cannot have a twin in G'_1 . The other twin is a leaf $u' \neq u$ attached to s (see Figure 9a). Notice that the edge vu is present in G , since otherwise u and u' would be twins also in G .




Figure 9: Illustrations for Case 2, where (a) $|N_{G_1}(v) \setminus \{u\}| = 1$, and (b) $|N_{G_1}(v) \setminus \{u\}| = 2$.

If $|N_{G_1}(v) \setminus \{u\}| = 2$, then one of the twins must be u, s or t (possibly, the two twins are among these three vertices). If u is a twin with some vertex other than s, t , then we would have a $K_{2,3}$ subgraph in G , contradicting Theorem 10. Thus, either s or t is one of the twins. However, the vertices s and t are not twins with each other in G'_1 , since if they were, they would be twins also in G , and G would not be twin-free. Thus, s or t has a neighbor that is not a neighbor of the other, say, $x \in N(s) \setminus N(t)$, $x \neq t$. Since $N_{G'_1}(u) = \{s, t\}$, if s or t has a twin, then it must be u . However, the vertex u cannot be a twin of s , because the vertex x is not a neighbor of u . Thus, if G'_1 has twins, then these twins are u and t , $N_{G'_1}(t) = \{u, s\}$ and $N_{G'_1}(u) = \{s, t\}$ (see Figure 9b).

Let us return to the original components G_1 and G_2 of $G - e$.

Claim N. There is a LTD-set of G_1 of size at most $\frac{2}{3}n_1$ which contains v and which is also a LTD-set of $G_1 + \{e, s'\}$.

Proof of claim. Denote $G_1^* = G_1 - \{v, u\}$ when $|N_{G_1}(v) \setminus \{u\}| = 1$, and $G_1^* = G_1 - \{v, t\}$ when $|N_{G_1}(v) \setminus \{u\}| = 2$ (see Figure 9).

If $G_1^* = K_2$, then the set $S_1 = \{v, s\}$ is clearly a LTD-set of both G_1 and $G_1 + \{e, s'\}$ with $|S_1| < \frac{2}{3}n_1$.

Assume that $G_1^* \neq K_2$, i.e., $|V(G_1^*)| \geq 3$. If G_1^* is twin-free, then by the induction hypothesis, there exists a LTD-set S_1^1 of G_1^* such that $|S_1^1| \leq \frac{2}{3}(n_1 - 2)$, and we have $s \in S_1^1$, since s is the support vertex of u' or u . Now, the set $S_1 = S_1^1 \cup \{v\}$ is a LTD-set of G_1 . Indeed, S_1 is clearly a TD-set, and it is a locating set as v dominates u (resp. t) and the only neighbor of v in G_1^* is s . Since $v \in S_1$, the set S_1 is also a TD-set of $G_1 + \{e, s'\}$. We have $N(s') \cap S_1 = \{v\}$. Since u and t are dominated by $s \in S_1$, the set S_1 is also locating in $G_1 + \{e, s'\}$. We have $|S_1| \leq \frac{2}{3}(n_1 - 2) + 1 = \frac{2}{3}(n_1 - \frac{1}{2})$.

Assume then that G_1^* has twins. Then, $|N_{G_1}(v) \setminus \{u\}| = 2$ and there exists a leaf $w \neq u$ attached to s . Now, the graph $G_1^* - u$ is either twin-free or K_2 . If $G_1^* - u = K_2$, then the set $S_1' = \{v, s, u\}$ fulfils the claim. Indeed, it is clearly a TD-set of both G_1 and $G_1 + \{e, s'\}$. Since the neighbors of t, w , and s' in S_1' are $\{v, s, u\}, s$, and v , respectively, the set S_1' is a locating set of both G_1 and $G_1 + \{e, s'\}$. Additionally, we have $|S_1'| < \frac{2}{3}n_1$.

Assume finally that $G_1^* - u$ is twin-free. Then, by the induction hypothesis, there exists a LTD-set S_1^2 of $G_1^* - u$ such that $|S_1^2| \leq \frac{2}{3}(n_1 - 3)$, and we have $s \in S_1^2$. Let $S_1' = S_1^2 \cup \{v, u\}$. As before, the set S_1' is a LTD-set of both G_1 and $G_1 + \{e, s'\}$, and we have $|S_1'| \leq \frac{2}{3}(n_1 - 3) + 2 = \frac{2}{3}n_1$. \square

If there exists a LTD-set of G_2 of cardinality at most $\frac{2}{3}n_2$, then it forms a LTD-set of G together with the set of Claim N. If G_2 is twin-free, then such a LTD-set of G_2 exists by the induction hypothesis. If G_2 is not twin-free, then we can repeat the same case analysis as for G_1 and obtain a LTD-set of G_2 of cardinality at most $\frac{2}{3}n_2$. \square

The bound in Theorem 12 is tight due to Observation 2. Indeed, if G is an outerplanar graph, then so is its 2-corona $G \circ P_2$, and we have $\gamma_t^L(G \circ P_2) = 2|V(G)| = \frac{2}{3}|V(G \circ P_2)|$.

Proposition 13. There are infinitely many connected twin-free outerplanar graphs G of order n with $\gamma_t^L(G) = \frac{2}{3}n$.

7 Conclusion

We have proved Conjecture 1 for several important graph classes: cobipartite graphs, split graphs, block graphs, subcubic graphs and outerplanar graphs.

It would be interesting to extend these results to larger classes, for example chordal graphs (which include split graphs and block graphs) or K_4 -minor-free graphs (which include outerplanar graphs). Another interesting subclass of chordal graphs to consider is the class of interval graphs.

It would also be interesting to prove that the bound $\gamma_t^L(G) \leq \frac{n}{2}$ holds for sufficiently large (twin-free) cubic graphs, as conjectured in [24].

References

- [1] G. R. Argiroffo, S. M. Bianchi, Y. Lucarini and A. Wagler. The Identifying Code, the Locating-dominating, the Open Locating-dominating and the Locating Total-dominating problems under some graph operations. Proceedings of the X Latin and American Algorithms, Graphs and Optimization Symposium (LAGOS 2019), *Electron. Notes Theor. Comput. Sci.* **346** (2019), 135–145.
- [2] G. R. Argiroffo, S. M. Bianchi, Y. Lucarini and A. Wagler. Polyhedra associated with open locating-dominating and locating total-dominating sets in graphs. Proceedings of the 6th International Symposium on Combinatorial Optimization (ISCO 2020), *Lecture Notes in Computer Science* **12176** (2020), 3–14.
- [3] L. Babai. On the complexity of canonical labeling of strongly regular graphs. *SIAM J. Comput.* **9** (1980), 212–216.
- [4] H.-J. Bandelt and H. M. Mulder. Distance-hereditary graphs. *J. Comb. Theory, Ser. B* **41** (1986), 182–208.
- [5] M. Blidia, M. Chellali, F. Maffray, J. Moncel, and A. Semri. Locating-domination and identifying codes in trees, *Australas. J. Combin.* **39** (2007), 219–232.
- [6] M. Blidia and W. Dali. A characterization of locating-total domination edge critical graphs, *Discuss. Math. Graph Theory* **31**(1) (2011), 197–202.
- [7] M. Blidia, O. Favaron, and R. Lounes. Locating-domination, 2-domination and independence in trees, *Australas. J. Combin.* **42** (2008), 309–319.
- [8] B. Bollobás, and A. D. Scott. On separating systems. *Eur. J. Comb.* **28** (2007), 1068–1071.
- [9] J. A. Bondy. Induced subsets. *J. Comb. Theory, Ser. B* **12** (1972), 201–202.
- [10] É. Bonnet, P. Giannopoulos, and M. Lampis. On the parameterized complexity of red-blue points separation. *J. Comput. Geom.* **10** (2019), 181–206.
- [11] R. C. Brigham, J. R. Carrington, and R. P. Vitray. Connected graphs with maximum total domination number. *J. Combin. Comput. Combin. Math.* **34** (2000), 81–96.
- [12] G. Chartrand and F. Harary. Planar permutation graphs. *Annales de l’Institut Henri Poincaré B* **3**(4) (1967), 433–438.
- [13] M. Chellali. On locating and differentiating-total domination in trees, *Discuss. Math. Graph Theory* **28**(3) (2008), 383–392.
- [14] M. Chellali and N. Jafari Rad. Locating-total domination critical graphs, *Australas. J. Combin.* **45** (2009), 227–234.
- [15] X. G. Chen and M. Y. Sohn. Bounds on the locating-total domination number of a tree, *Discrete Appl. Math.* **159** (2011), 769–773.

- [16] E. J. Cockayne, R. M. Dawes, and S. T. Hedetniemi. Total domination in graphs. *Networks* **10** (1980), 211–219.
- [17] F. Foucaud and M. A. Henning. Location-domination and matching in cubic graphs. *Discrete Math.* **339** (2016), 1221–1231.
- [18] F. Foucaud and M. A. Henning. Locating-total dominating sets in twin-free graphs: a conjecture. *The Electronic Journal of Combinatorics* **23** (2016), #P3.9.
- [19] F. Foucaud and M. A. Henning. Location-domination in line graphs. *Discrete Math.* **340** (2017), 3140–3153.
- [20] F. Foucaud, M. A. Henning, C. Löwenstein, and T. Sasse. Locating-dominating sets in twin-free graphs. *Discrete Appl. Math.* **200** (2016), 52–58.
- [21] F. Foucaud and T. Lehtilä. Bounds and extremal graphs for total dominating identifying codes. Manuscript, 2022. <https://arxiv.org/abs/2206.09481>
- [22] D. Garijo, A. González, and A. Márquez. The difference between the metric dimension and the determining number of a graph. *Appl. Math. Comput.* **249** (2014), 487–501.
- [23] T. W. Haynes, M. A. Henning, and J. Howard. Locating and total dominating sets in trees. *Discrete Appl. Math.* **154** (2006), 1293–1300.
- [24] M. A. Henning and C. Löwenstein. Locating-total domination in claw-free cubic graphs. *Discrete Math.* **312** (2012), 3107–3116.
- [25] M. A. Henning and N. J. Rad. Locating-total domination in graphs, *Discrete Appl. Math.* **160** (2012), 1986–1993.
- [26] M. A. Henning and A. Yeo. *Total domination in graphs*, Springer-Verlag, 2013.
- [27] M. G. Karpovsky, K. Chakrabarty, and L. B. Levitin. On a new class of codes for identifying vertices in graphs. *IEEE Trans. Inf. Theory* **44** (1998), 599–611.
- [28] M. Miller, R.S. Rajan, R. Jayagopal, I. Rajasingh and P. Manuel. A note on locating-total domination in graphs. *Discuss. Math. Graph Theory* **37** (2017), 745–754.
- [29] J. H. Kim, O. Pikhurko, J. Spencer and O. Verbitsky. How complex are random graphs in First Order logic? *Random Struct. Algorithms* **26** (2005), 119–145.
- [30] A. Lobstein. Watching systems, identifying, locating-dominating and discriminating codes in graphs: a bibliography. <https://www.lri.fr/~lobstein/debutBIBidetlocdom.pdf>
- [31] B. M. E. Moret, and H. D. Shapiro. On minimizing a set of tests. *SIAM J. Sci. Stat. Comput.* **6** (1985), 983–1003.
- [32] H. Rahbani, N. J. Rad and M.-R. Sadeghi. A note on the complexity of locating-total domination in graphs. *Theor. Comput. Sci.* **799** (2019), 32–39.
- [33] P. J. Slater. Dominating and reference sets in graphs. *J. Math. Phys. Sci.* **22** (1988), 445–455.
- [34] R. Ungrangsi, A. Trachtenberg, and D. Starobinski. An implementation of indoor location detection systems based on identifying codes. Proceedings of the IFIP International Conference on Intelligence in Communication Systems, INTELLICOMM 2004. *Lecture Notes in Computer Science* 3283:175–189, 2004.