

A Rigorous Proof of the Index Theorem for Economists

Yuhki Hosoya*

Faculty of Economics, Chuo University†

Abstract

This paper provides a rigorous and gap-free proof of the index theorem used in the theory of regular economy. In the index theorem that is the subject of this paper, the assumptions for the excess demand function are only several usual assumptions and continuous differentiability around any equilibrium price, and thus it has a form that is applicable to many economies. However, the textbooks on this theme contain only abbreviated proofs and there is no known monograph that contains a rigorous proof of this theorem. Hence, the purpose of this paper is to make this theorem available to more economists by constructing a readable proof.

Keywords: index theorem, regular economy, transversality.

JEL codes: C65, D41, D51

MSC2020 codes: 91B24, 57R25

1 Introduction

How many equilibrium prices exist? Arrow and Debreu (1954) showed that at least one equilibrium price exists. It seems that economists of this period believed that the fewer the number of equilibria, the better. Therefore, Arrow et al. (1959) proposed a sufficient condition for the existence of a unique equilibrium price. However, their condition is very complicated and does not seem to be valid in many economies.

*TEL: +81-90-5525-5142, E-mail: ukki(at)gs.econ.keio.ac.jp

†742-1, Higashinakano, Hachioji-shi, Tokyo, 192-0393, Japan.

Debreu (1970) proved that the number of equilibrium prices is at least finite in “almost all” economies. In his view, it was difficult to prove that there is exactly one equilibrium price, and thus he proved that at least there are only a finite number of equilibria. This result had been refined and reformulated by many economists. This research area is now known as the theory of regular economy.

The most central result in the theory of regular economy is called the index theorem. This theorem states that the sum of “indices” computed by a certain method for equilibrium prices is equal to 1. This result is powerful. Because “indices” must be able to be summed, the number of equilibrium prices must be a finite odd number. We can immediately reproduce the result of Debreu (1970) using this theorem.

The problem is that no straightforward proof of this index theorem is known. The most famous index theorem currently published in a form applicable to many economies appears to be Proposition 5.6.1 of Mas-Colell (1985). However, the proof of this theorem contains many gaps that ordinary readers will find impossible to fill.

Therefore, in this paper, we focus on the proof of this result and provide a rigorous and gap-free proof of this theorem in the shortest manner. The proof presented in this paper requires basic knowledge of differential topology supplied by Guillemin and Pollack (1974). Therefore, in this paper, we construct a proof under the assumption that the contents of the above book is known. However, since Guillemin-Pollack’s formulation of differential topology is a bit peculiar, we have chosen to extract only the claims necessary for this paper and offer them to readers.

The proof in this paper requires a classification theorem for one-dimensional manifolds. However, the problem is that a well-written proof of the classification theorem contained in a book is missing. Therefore, in the appendix of this paper, we present a proof of this classification theorem.

The structure of this paper is as follows. In Section 2, we explain a part of the knowledge of differential topology contained in Guillemin-Pollack’s book that is necessary to understand the proof. We do not prove any of the results that appear in this section, but the proofs are either in Guillemin-Pollack or can be easily reconstructed from what is written in Guillemin-Pollack. Section 3 contains the assertion of the main result and its proof. In Section 4, we explain the relationships between Proposition 5.6.1 of Mas-Colell (1985) and our main result, and the proof and its gaps in Mas-Colell’s book. In the appendix, we prove the classification theorem for one-dimensional manifolds.

2 Preliminaries

In this paper, we assume that readers are sufficiently familiar with the results in Guillemin and Pollack (1974). In particular, we assume that readers know the definition of manifolds, tangent spaces and the tangent bundle, the derivatives of smooth functions on manifolds, transversality, and oriented manifolds. However, the definition of manifolds in Guillemin and Pollack (1974) is not standard in differential topology, and with very few exceptions such as Milnor (1965), another definition has been used. Thus, readers may not be familiar with the manifolds treated in this paper, even though they have appropriately deep knowledge on manifolds. Therefore, in this section, we present the definitions of the terms in differential topology treated in this paper and the assertion of some theorems. All of the results in this section have been proved or can be easily verified from the results proved in Guillemin and Pollack (1974).

Note that, although Guillemin and Pollack (1974) treated only C^∞ manifolds and functions, in most cases, their proofs work well for C^k manifolds and functions, even if $k \neq \infty$. However, several techniques depend on the assumption $k = +\infty$. For example, Sard's theorem does not necessarily hold if the function is not C^∞ . The concept of Morse function cannot be appropriately defined in C^1 manifolds. The tangent bundle $T(X)$ is not necessarily a C^1 manifold when X is a C^1 manifold. We avoid these technical problems in this paper. First, we use Sard's theorem only for a C^∞ function defined on a C^∞ manifold. Second, we consider the tangent bundle only for a C^∞ manifold. Third, we avoid the use of Morse functions.

2.1 Manifolds

First, suppose that $X \subset \mathbb{R}^N$ and let $f : X \rightarrow \mathbb{R}^M$ be a continuous function. Choose any $x \in X$ and let $k \in \mathbb{N} \cup \{\infty\}$. We say that f is C^k around x if and only if there exists a C^k function $F : U \rightarrow \mathbb{R}^M$ such that U is an open neighborhood of x in \mathbb{R}^N and $F(y) = f(y)$ whenever $y \in U \cap X$. As usual, a function that is C^k everywhere is called a C^k function.

A function $f : X \rightarrow Y$ is called a **homeomorphism** if and only if f is bijective and both f and f^{-1} are continuous. Suppose that $f : X \rightarrow Y$ is a homeomorphism. We say that f is a C^k **diffeomorphism** if and only if both f and f^{-1} are C^k . A nonempty set $X \subset \mathbb{R}^N$ is called an **n -dimensional C^k manifold** if and only if for every $x \in X$, there exists a C^k diffeomorphism $\phi : U \rightarrow V$ such that U is an open set in \mathbb{R}^n and V is an open neighborhood of x in X . Such a ϕ is called a **local parametrization around x** . The number n is called the **dimension of X** and denoted by $\dim X$. It is known

that if X is a C^k manifold, then $\dim X$ is uniquely determined.

Sometimes, C^0 manifolds are considered. A function is said to be C^0 if and only if it is continuous, and a set $X \subset \mathbb{R}^N$ is called an n -dimensional C^0 manifold if and only if every $x \in X$ has an open neighborhood that is homeomorphic to some open set in \mathbb{R}^n . A C^0 manifold is sometimes called a **topological manifold**. However, in C^0 manifolds, we cannot define the tangent space, which is explained later. This is very restrictive for our argument, and thus we do not use a C^0 manifold in this paper, except in the appendix.

We define $\mathbb{R}^0 = \{0\}$. Hence, 0-dimensional manifolds can be defined. A set X is a 0-dimensional C^k manifold if and only if it is a discrete set.

Let X be an n -dimensional C^k manifold, and choose any $x \in X$. Choose any local parametrization ϕ around x . Let $\phi(z) = x$ and suppose that $D\phi(z)$ denotes the Fréchet derivative of ϕ at z . Let $T_x(X)$ denote the range of $D\phi(z)$. Then, it is known that $T_x(X)$ is independent of the choice of ϕ . We call this linear space $T_x(X)$ the **tangent space** of X at x . It is known that the dimension of $T_x(X)$ is n , and for every $v \in \mathbb{R}^N$, $v \in T_x(X)$ if and only if there exist $\varepsilon > 0$ and a C^1 curve $c : (-\varepsilon, \varepsilon) \rightarrow X$ such that $c(0) = x$ and $c'(0) = v$.

Next, let X, Y be C^k manifolds, and let $f : X \rightarrow Y$ be a C^k function. Choose $x \in X$, and a local parametrization ϕ around x . Let $\phi(z) = x$. Then, $g = f \circ \phi$ is a C^k function defined on an open set $U \subset \mathbb{R}^n$, where n is the dimension of X . Hence, the Fréchet derivative $Dg(z)$ can be defined. Choose any $v \in T_x(X)$, and let $D\phi(z)w = v$. Then, we can define $df_x(v) = Dg(z)w$, and it can be shown that the left-hand side is independent of the choice of ϕ . Moreover, $df_x(v) \in T_{f(x)}(Y)$ for all $x \in X$ and $v \in T_x(X)$. This linear function df_x is called the **derivative** of f at x . Suppose that $v \in T_x(X)$ and $c : (-\varepsilon, \varepsilon) \rightarrow X$ is a C^1 function that satisfies $c(0) = x$ and $c'(0) = v$. For a C^k function $f : X \rightarrow Y$, if we define $d = f \circ c$, then we can check that $df_x(v) = d'(0)$.

Suppose that X, Y are C^k manifolds whose dimensions are the same, and $f : X \rightarrow Y$ is a C^k function. Choose $x \in X$ and suppose also that df_x is a bijection from $T_x(X)$ onto $T_{f(x)}(Y)$. Then, there exists an open neighborhood U of x such that $F(U)$ is an open neighborhood of $f(x)$ and the restriction of f is a C^k diffeomorphism between U and $f(U)$. This result is known as the **inverse function theorem**.

2.2 Submanifolds and Transversality

Let X be an n -dimensional C^k manifold, and $Y \subset X$. We call Y a **submanifold** of X if Y itself is a C^k manifold. It is known that if Y is a

submanifold of X , then $T_x(Y) \subset T_x(X)$. If $\dim Y = m$, then $n - m$ is called the **codimension** of Y , and denoted by $\text{codim } Y$.

Suppose that X, Y are C^k manifolds, Z is a submanifold of Y , and $f : X \rightarrow Y$ is C^k . We say that f is **transversal to Z** if and only if for every $x \in f^{-1}(Z)$,

$$\text{Im}(df_x) + T_{f(x)}(Z) = T_{f(x)}(Y).$$

Then, the following fact is known.

Fact 1. *Suppose that X, Y are C^k manifolds, Z is a C^k submanifold of Y , and $f : X \rightarrow Y$ is a C^k function that is transversal to Z . If $S = f^{-1}(Z)$ is nonempty, then S is also a submanifold of X , and $\text{codim } S = \text{codim } Z$.*

2.3 Tangent Bundle

Suppose that X is an n -dimensional C^∞ manifold. The following set

$$T(X) = \{(x, v) | x \in X, v \in T_x(X)\}$$

is called the **tangent bundle** of X . It is known that $T(X)$ is a $2n$ -dimensional C^∞ manifold, and if $(x, v) \in T(X)$ and ϕ is a local parametrization around x in X , then

$$(z, w) \mapsto (\phi(z), d\phi_z(w))$$

is a local parametrization around (x, v) in $T(X)$.

By construction, it is easy to show that $T_{(x,v)}(T(X)) = (T_x(X))^2$.

2.4 Manifolds with Boundary

We need the definition of manifolds with boundary. First, let $H^n = \{x \in \mathbb{R}^n | x_n \geq 0\}$. A set $X \subset \mathbb{R}^n$ is called an **n -dimensional C^k manifold with boundary** if and only if for every $x \in X$, there exists a C^k diffeomorphism $\phi : U \rightarrow V$ such that U is an open set in H^n and V is an open neighborhood of x in X . The mapping ϕ is called a local parametrization around x as usual. Tangent space $T_x(X)$ can be defined in the usual manner. Note that, $T_x(X)$ is an n -dimensional linear space, even when X is not a usual manifold but a manifold with boundary.

Suppose that X is an n -dimensional C^k manifold with boundary, and define the set ∂X as the set of all $x \in X$ such that there is no local parametrization around x whose domain is an open set in \mathbb{R}^n . It can be shown that if ∂X is nonempty, then it is an $(n - 1)$ -dimensional C^k manifold, and $T_x(\partial X)$ is a subspace of $T_x(X)$.

Note that, $X \times Y$ is not necessarily a manifold with boundary, even if both X, Y are manifolds with boundary. For example, it is known that $[0, 1]^2$ is not a manifold with boundary. However, if X is an n -dimensional C^k manifold with boundary and Y is an m -dimensional C^k manifold without boundary, then $X \times Y$ is an $(n+m)$ -dimensional C^k manifold with boundary, and $\partial(X \times Y) = \partial X \times Y$ and $T_{(x,y)}(X \times Y) = T_x(X) \times T_y(Y)$. In particular, because the unit interval $[0, 1]$ is a 1-dimensional C^∞ manifold with boundary, if X is an n -dimensional C^k manifold, then $Y = [0, 1] \times X$ is an $(n+1)$ -dimensional C^k manifold with boundary, and $\partial Y = \{0, 1\} \times X$. If we write $X_t = \{t\} \times X$, then $\partial Y = X_0 \cup X_1$ in this sense.

Suppose that X, Y are C^k manifolds with boundary, and $f : X \rightarrow Y$ is C^k . Then, we can define the derivative $df_x : T_x(X) \rightarrow T_{f(x)}(Y)$ in the same manner as in Subsection 2.1. Usually, ∂f denotes the restriction of f into ∂X . If Y has no boundary and Z is a submanifold of Y , we can define transversality as usual: that is, f is transversal to Z if and only if for any $x \in f^{-1}(Z)$,

$$\text{Im}(df_x) + T_{f(x)}(Z) = T_{f(x)}(Y).$$

Then, the following fact holds.

Fact 2. *Suppose that X is a C^k manifold with boundary, Y is a C^k manifold without boundary, and Z is a submanifold of Y without boundary. Let $f : X \rightarrow Y$ be a C^k function such that both f and ∂f are transversal to Z . If $S = f^{-1}(Z)$ is nonempty, then S is a C^k manifold with boundary whose codimension is the same as that of Z , and $\partial S = \partial X \cap S$.*

2.5 Orientation

Suppose that V is an n -dimensional linear space. Choose any two bases $v = (v_1, \dots, v_n)$ and $w = (w_1, \dots, w_n)$. Then, there uniquely exists an $(n \times n)$ matrix A such that

$$w_i = a_{i1}v_1 + \dots + a_{in}v_n$$

for all $i \in \{1, \dots, n\}$. We say that v is **equivalent** to w if and only if $\det A > 0$.¹

Suppose that $X \subset \mathbb{R}^N$ is an n -dimensional C^k manifold with/without boundary. Let

$$T^n(X) = \{(x, v_1, \dots, v_n) | x \in X, v_i \in T_x(X)\}.$$

Choose any function $\text{sign}_X : T^n(X) \rightarrow \{+1, -1\}$. We say that sign_X is **sign-preserving** if for any two equivalent bases $v = (v_1, \dots, v_n)$ and $w =$

¹We use both expressions $\det A$ and $|A|$ to denote the determinant of A .

(w_1, \dots, w_n) of $T_x(X)$, $\text{sign}_X(x, v) = \text{sign}_X(x, w)$. Suppose that for every $x \in X$, there exists a local parametrization ϕ around x such that for $v_i(z) = D\phi(z)e_i$, $\text{sign}_X(\phi(z), v_1(z), \dots, v_n(z))$ is constant, where $e_i \in \mathbb{R}^n$ is the i -th unit vector. Then, we call this sign-preserving function sign_X an **orientation** of X .

We must note an exception. Suppose that X is a 0-dimensional manifold. By definition, X is a discrete set, and $T_x(X) = \{0\}$. In this case, any function $\text{sign}_X : X \rightarrow \{-1, +1\}$ is called an orientation of X .

Note that, there is a manifold with no orientation: the Möbius strip is a typical example. If X is a manifold and there is an orientation, then we say that X is **orientable**. If we think that an orientation of X is given, then X is said to be **oriented**. It is known that if X is a connected and orientable manifold, then there are exactly two possible orientations.

If X, Y are manifolds and either $\partial X = \emptyset$ or $\partial Y = \emptyset$, then $X \times Y$ is also a manifold. If, in addition, both X, Y are oriented, then $X \times Y$ is also oriented, and the orientation is made in the following manner. Suppose that $\text{sign}_X, \text{sign}_Y$ denote orientations. Then, there uniquely exists a sign-preserving function $\text{sign}_{X \times Y}$ such that

$$\begin{aligned} & \text{sign}_{X \times Y}((x, y), (v_1, 0), \dots, (v_n, 0), (0, w_1), \dots, (0, w_m)) \\ &= \text{sign}_X(x, v_1, \dots, v_n) \text{sign}_Y(y, w_1, \dots, w_m) \end{aligned}$$

for all $(x, y) \in X \times Y$ and bases $v = (v_1, \dots, v_n)$ of $T_x(X)$ and $w = (w_1, \dots, w_m)$ of $T_y(Y)$. We can easily check that this function $\text{sign}_{X \times Y}$ is an orientation of $X \times Y$, and thus, $X \times Y$ is oriented by this orientation.

Suppose that $X \subset \mathbb{R}^N$ is an n -dimensional C^k oriented manifold, and $Y \subset X$ is an $(n-1)$ -dimensional submanifold. Choose any orientation sign_X of X . Suppose also that there exists a C^{k-1} function $n : Y \rightarrow \mathbb{R}^N$ such that $n(x) \in T_x(X)$, $\|n(x)\| = 1$, and $n(x) \cdot v = 0$ for all $v \in T_x(Y)$. Then, there exists an orientation sign_Y of Y such that

$$\text{sign}_Y(x, v_1, \dots, v_{n-1}) = \text{sign}_X(x, n(x), v_1, \dots, v_{n-1}).$$

Hence, Y is an oriented manifold.

Next, suppose that $X \subset \mathbb{R}^N$ is a C^k manifold with boundary. Choose any $x \in \partial X$. Then, there exists $p \in T_x(X)$ such that $\|p\| = 1$ and $p \cdot v = 0$ for all $v \in T_x(\partial X)$. Choose any local parametrization ϕ around x , and suppose that $p = D\phi_x(q)$. Then, $q_n \neq 0$. Define $n(x) = p$ if $q_n < 0$ and $n(x) = -p$ if $q_n > 0$. This function $n : \partial X \rightarrow \mathbb{R}^N$ is called the **outward normal vector field** of ∂X . We can show that $n(x)$ is C^{k-1} , and thus, we can conclude that ∂X is oriented by the following sign-preserving function:

$$\text{sign}_{\partial X}(x, v_1, \dots, v_{n-1}) = \text{sign}_X(x, n(x), v_1, \dots, v_{n-1}).$$

Suppose that X is a C^k oriented manifold with boundary, Y is a C^k oriented manifold without boundary, Z is a C^k oriented submanifold of Y without boundary, $f : X \rightarrow Y$ is a C^k function such that both f and ∂f are transversal to Z , and $S = f^{-1}(Z)$ is nonempty. Then, S is a manifold with boundary. Let $\ell = \text{codim } Z$. Choose any $x \in f^{-1}(Z)$ and suppose that H is an ℓ -dimensional linear subspace of $T_x(X)$ such that

$$df_x(H) \oplus T_{f(x)}(Z) = T_{f(x)}(Y).$$

Then, we can show that

$$H \oplus T_x(S) = T_x(X).$$

Choose any basis (v_1, \dots, v_ℓ) of H , and define $w_i = df_x(v_i)$. Choose any basis $(w_{\ell+1}, \dots, w_m)$ such that $\text{sign}_Z(f(x), w_{\ell+1}, \dots, w_m) = +1$. Define

$$h(v_1, \dots, v_\ell) = \text{sign}_Y(f(x), w_1, \dots, w_m),$$

and

$$\text{sign}_S(x, v_{\ell+1}, \dots, v_n) = \text{sign}_X(x, v_1, \dots, v_n)h(v_1, \dots, v_\ell).$$

Then, we can show that sign_S is an orientation. We call this orientation the **preimage orientation**.

2.6 Classification of One-dimensional Manifolds

Let

$$S^1 = \{x \in \mathbb{R}^2 \mid \|x\| = 1\}.$$

Suppose that X is a compact and connected 1-dimensional C^k manifold with/without boundary. Then, it is known that X is C^k diffeomorphic to either S^1 or $[0, 1]$. In Guillemin and Pollack (1974), this classification theorem was proved using a Morse function. However, Morse function can be defined only if $k \geq 2$, and thus their proof cannot be applied if $k = 1$. And actually, we need this result for $k = 1$. Hence, we prove this theorem in appendix.

It is known that every connected component of a manifold is open. Suppose that X is a compact 1-dimensional C^k manifold with/without boundary. Because X is compact, the number of connected components of X is finite. Because each connected component X_i is diffeomorphic to either S^1 or $[0, 1]$, either ∂X_i is empty or it contains 2 points. In particular, ∂X is a finite set that includes even number of points.

Moreover, suppose that X is oriented. Then, for each connected component X_i that is diffeomorphic to $[0, 1]$, if $\partial X_i = \{a, b\}$, then $\text{sign}_{\partial X}(a) +$

$\text{sign}_{\partial X}(b) = 0$. This implies that

$$\sum_{x \in \partial X} \text{sign}_{\partial X}(x) = 0.$$

This result is crucial for our main result.

3 Result

In this section, we fix $n \geq 2$. We define $\mathbb{R}_+^n = \{x \in \mathbb{R}^n | x_i \geq 0 \text{ for all } i\}$ and $\mathbb{R}_{++}^n = \{x \in \mathbb{R}^n | x_i > 0 \text{ for all } i\}$. Let $S = \{p \in \mathbb{R}_{++}^n | \|p\| = 1\}$. It is known that S is an $(n - 1)$ -dimensional C^∞ manifold. Moreover, because $g(p) = p$ is a natural normal vector field on S , we can assume that S is oriented. Our main result is as follows.

Theorem 1. *Let $f : \mathbb{R}_{++}^n \rightarrow \mathbb{R}^n$ be a continuous mapping such that the following five conditions hold.*

- (i) $p \cdot f(p) = 0$ for all $p \in S$, and $f(ap) = f(p)$ for all $a > 0$ and $p \in S$.
- (ii) There exists $s > 0$ such that $f_i(p) > -s$ for all $p \in S$ and $i \in \{1, \dots, n\}$.
- (iii) If (p^k) is a sequence on S such that $p^k \rightarrow p \notin S$ as $k \rightarrow \infty$, then $\|f(p^k)\| \rightarrow \infty$ as $k \rightarrow \infty$.
- (iv) If $f(p^*) = 0$, then f is continuously differentiable around p^* .
- (v) If $f(p^*) = 0$, then

$$g(p^*) = \begin{vmatrix} Df(p^*) & p^* \\ (p^*)^T & 0 \end{vmatrix} \neq 0.$$

Let $E^* = \{p \in S | f(p) = 0\}$, and for each $p^* \in E^*$, define

$$\text{index}(p) = \begin{cases} +1 & \text{if } (-1)^n g(p^*) > 0, \\ -1 & \text{if } (-1)^n g(p^*) < 0. \end{cases}$$

Then, E^* is a finite set and

$$\sum_{p^* \in E^*} \text{index}(p^*) = +1.$$

This result is actually the same as Proposition 5.6.1 of Mas-Colell (1985). However, the proof of this proposition includes big gaps, and thus many readers cannot judge whether this proposition is correct. Hence, we provide a rigorous and gap-free proof for this theorem. This is the purpose of this paper.

Proof. We separate the proof into nine steps.

Step 1. *The set E^* is finite.*

Proof of Step 1. Choose any $p^* \in E^*$, and let $A = Df(p^*)$ and $V = \{v \in \mathbb{R}^n \mid v \cdot p^* = 0\}$. If $v \in V$, $v \neq 0$ and $Av = 0$, then

$$\begin{pmatrix} Df(p^*) & p^* \\ (p^*)^T & 0 \end{pmatrix} \begin{pmatrix} v \\ 0 \end{pmatrix} = 0,$$

which contradicts (v). Therefore, we have that $v \in V$ and $v \neq 0$ imply $Av \neq 0$. Hence,

$$m = \min\{\|Av\|/\|v\| \mid v \in V, v \neq 0\} > 0.$$

Because of the definition of the Fréchet derivative, there exists $\varepsilon > 0$ such that if $v \in V$ and $0 < \|v\| < \varepsilon$, then

$$\|f(p^* + v) - f(p^*) - Av\| < \frac{\|v\|m}{2}.$$

Because $f(p^*) = 0$,

$$\begin{aligned} \|v\|m &\leq \|Av\| \leq \|f(p^* + v) - f(p^*) - Av\| + \|f(p^* + v)\| \\ &< \frac{\|v\|m}{2} + \|f(p^* + v)\|, \end{aligned}$$

which implies that $f(p^* + v) \neq 0$. Now, for any $q \in (p^* + V) \cap \mathbb{R}_{++}^n$, define $P(q) = q/\|q\|$. Then, P is a C^∞ mapping from $(p^* + V)$ into S , and it is easy to show that dP_{p^*} is the identity mapping. By the inverse function theorem, there exists $\delta > 0$ such that P^{-1} is defined, injective on $\{p \in S \mid \|p - p^*\| < \delta\}$, and if $p \in S$ and $\|p - p^*\| < \delta$, then $\|P^{-1}(p) - p^*\| < \varepsilon$. By (i), $p \in S, 0 < \|p - p^*\| < \delta$ implies that $f(p) \neq 0$, and thus E^* is a discrete set.

By (iii), there exists $\varepsilon > 0$ such that if $p \in S$ and $p_i \geq \varepsilon$ for all $i \in \{1, \dots, n\}$, then $f(p) \neq 0$. This implies that E^* is compact, and thus finite. This completes the proof of Step 1. \square

Step 2. *Let (p^k) be a sequence on S such that p^k converges to $p \notin S$ and $\frac{1}{\|f(p^k)\|} f(p^k)$ converges to $z \in \mathbb{R}^n$ as $k \rightarrow \infty$. Then, $z \geq 0$, $z \neq 0$ and $p \cdot z = 0$.*

Proof of Step 2. By (ii), $f_i(p^k) \rightarrow -\infty$ is impossible, and thus by (iii), we have that $z_i \geq 0$. Because $\|z\| = 1$, $z \neq 0$. Finally, suppose that $p_i > 0$. By (i),

$$f_i(p^k) = -\frac{1}{p_i^k} \left(\sum_{j \neq i} p_j^k f_j(p^k) \right).$$

Because the right-hand side is bounded from above, we have that the sequence $(f_i(p^k))$ is bounded, and thus $z_i = 0$, which implies that $p \cdot z = 0$. This completes the proof of Step 2. \square

Step 3. Choose any $q \neq 0$. Suppose that A is an $n \times n$ matrix such that $v^T A v < 0$ for all $v \in \mathbb{R}^n$ such that $v \cdot q = 0$ and $v \neq 0$. Then,

$$(-1)^n \begin{vmatrix} A & q \\ q^T & 0 \end{vmatrix} > 0.$$

Proof of Step 3. Let $A_t = (1-t)A + tA^T$. By assumption, we have that $v^T A_t v < 0$ for all $t \in [0, 1]$ and $v \in \mathbb{R}^n$ such that $v \cdot q = 0$ and $v \neq 0$. Suppose that there exists $t \in [0, 1]$ such that

$$\begin{vmatrix} A_t & q \\ q^T & 0 \end{vmatrix} = 0,$$

then, there exist $v \in \mathbb{R}^n$ and $w \in \mathbb{R}$ such that $(v, w) \neq 0$ and

$$\begin{pmatrix} A_t & q \\ q^T & 0 \end{pmatrix} \begin{pmatrix} v \\ w \end{pmatrix} = 0.$$

This implies that $q^T v = 0$, and thus $v \cdot q = 0$. Therefore,

$$0 = v^T 0 = v^T A_t v + v^T q w = v^T A_t v,$$

which implies that $v = 0$. However, this implies that $w q = 0$, and thus $w = 0$, which is a contradiction. By the intermediate value theorem, we have that

$$c(t) \equiv \begin{vmatrix} A_t & q \\ q^T & 0 \end{vmatrix}$$

has a constant sign on $[0, 1]$. If we set $t = \frac{1}{2}$, then A_t is symmetric, and thus by Theorem 5 of Debreu (1952), $(-1)^n c(1/2) > 0$. Therefore, $(-1)^n c(0) > 0$, as desired. This completes the proof of Step 3. \square

Step 4. Suppose that $p \in \mathbb{R}^n$ and $\|p\| = 1$, and A is an $n \times n$ matrix such that $A p = 0$ and

$$\begin{vmatrix} A & p \\ p^T & 0 \end{vmatrix} \neq 0.$$

Let $x_1, \dots, x_{n-1} \in \mathbb{R}^n$ satisfy $p \cdot x_i = 0$ and $\det(p, x_1, \dots, x_{n-1}) > 0$, and define $y_i = A x_i$. Then,

$$\text{sign} \begin{vmatrix} A & p \\ p^T & 0 \end{vmatrix} = -\text{sign}(\det(p, y_1, \dots, y_{n-1})).$$

Proof of Step 4. Let

$$B = \begin{pmatrix} A & p \\ p^T & 0 \end{pmatrix}.$$

Then,

$$B \begin{pmatrix} x_i \\ 0 \end{pmatrix} = \begin{pmatrix} y_i \\ 0 \end{pmatrix}, \quad B \begin{pmatrix} p \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad B \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} p \\ 0 \end{pmatrix}.$$

Define

$$C = \begin{pmatrix} 0 & x_{11} & \cdots & x_{n-1,1} & p_1 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & x_{1n} & \cdots & x_{n-1,n} & p_n \\ 1 & 0 & \cdots & 0 & 0 \end{pmatrix}.$$

Then, $\det(C) < 0$ and

$$BC = \begin{pmatrix} p_1 & y_{11} & \cdots & y_{n-1,1} & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ p_n & y_{1n} & \cdots & y_{n-1,n} & 0 \\ 0 & 0 & \cdots & 0 & 1 \end{pmatrix},$$

and thus

$$\det(B) = \det(p, y_1, \dots, y_{n-1})(\det(C))^{-1},$$

which implies that our claim is correct. This completes the proof of Step 4. \square

Choose any $q \in S$, and define $f^q(p) = \frac{\|p\|}{p \cdot q} q - \frac{1}{\|p\|} p$. Then, f^q is defined on $\mathbb{R}_+^n \setminus \{0\}$.

Step 5. f^q satisfies (i), (ii), (iv), (v) of this theorem.

Proof of Step 5. Clearly, $p \cdot f^q(p) = 0$ and $f^q(p) = f^q(ap)$ for all $a > 0$, and thus (i) holds. Let \bar{S} be the closure of S in \mathbb{R}^n . Then, f^q is defined on \bar{S} and continuous, which implies that (ii) holds. Clearly, f^q is C^∞ and thus (iv) holds. It is obvious that $f^q(p) = 0$ if and only if $p = aq$ for some $a > 0$. Let $v \cdot q = 0$ and $v \neq 0$, and define $p(t) = q + tv$. Then, $p(t) \cdot q = \|q\|^2 = 1$, and thus

$$\frac{d}{dt} f^q(p(t)) = \frac{t\|v\|^2}{\|p(t)\|} q + \frac{t\|v\|^2}{2\|p(t)\|^3} p(t) - \frac{1}{\|p(t)\|} v,$$

which implies that

$$v^T Df^q(q)v = v \cdot \left(\frac{d}{dt} f^q(p(t)) \Big|_{t=0} \right) = -\|v\|^2 < 0.$$

Therefore, $v \cdot q = 0$ and $v \neq 0$ imply that $v^T Df^q(q)v < 0$, and by Step 3,

$$(-1)^n \begin{vmatrix} Df^q(q) & q \\ q^T & 0 \end{vmatrix} > 0. \quad (1)$$

Because $f^q(ap) = f^q(p)$ for all $a > 0$, we have that $Df^q(aq) = a^{-1}Df^q(q)$, and thus (v) holds. This completes the proof of Step 5. \square

Define

$$\begin{aligned} S_\varepsilon &= \{p \in S \mid p_i \geq \varepsilon \text{ for all } i \in \{1, \dots, n\}\}, \\ \tilde{S}_\varepsilon &= \{p \in S \mid p_i > \varepsilon \text{ for all } i \in \{1, \dots, n\}\}, \\ \partial S_\varepsilon &= S_\varepsilon \setminus \tilde{S}_\varepsilon. \end{aligned}$$

Note that, S_ε and ∂S_ε are not manifolds. Choose any $\varepsilon' > 0$ such that $\tilde{S}_{\varepsilon'} \neq \emptyset$.

Step 6. *If $\varepsilon > 0$ is sufficiently small, then $\tilde{S}_\varepsilon \neq \emptyset$ and*

$$(1-t)f(p) + tf^q(p) \neq 0$$

for all $t \in [0, 1]$, $q \in S_{\varepsilon'}$, and $p \in \partial S_\varepsilon$.

Proof of Step 6. Suppose not. Then, there exist sequences (t^k) on $[0, 1]$, (p^k) on S , and (q^k) on $S_{\varepsilon'}$ such that $p^k \rightarrow p \notin S$ as $k \rightarrow \infty$ and

$$(1-t^k)f(p^k) + t^k f^{q^k}(p^k) = 0$$

for all k . Let $z^k = \frac{1}{\|f(p^k)\|} f(p^k)$. By taking subsequences, we can assume that $z^k \rightarrow z$ as $k \rightarrow \infty$. By Step 2, $z \geq 0$, $z \neq 0$ and $p \cdot z = 0$. Choose any i such that $z_i > 0$. Then, $p_i = 0$, and $\min_{q \in S_{\varepsilon'}} f_i^q(p) > 0$. This implies that $\min_{q \in S_{\varepsilon'}} f_i^q(p^k) > 0$ for sufficiently large k . Moreover, because $z_i > 0$, $f_i(p^k) > 0$ for sufficiently large k . Thus, for such a k ,

$$(1-t^k)f_i(p^k) + t^k f_i^{q^k}(p^k) > 0,$$

which is a contradiction. This completes the proof of Step 6. \square

Hereafter, we fix $\varepsilon > 0$ that satisfies the requirements in Step 6, $E^* \subset S_\varepsilon$, and $\varepsilon < \varepsilon'$.

Choose any $\eta > 0$ such that if $p^* \in E^*$, then f is C^1 on $\{p \in \mathbb{R}_{++}^n \mid \|p - p^*\| < 4\eta\}$, and $\{p \in \mathbb{R}_{++}^n \mid f(p) = 0, \|p - p^*\| < 4\eta\} \subset \{ap^* \mid a > 0\}$. By Step 1, E^* is finite, and thus such an η exists. Define

$$\hat{S}_{\varepsilon, a} = \{p \in S_\varepsilon \mid \forall p^* \in E^*, \|p - p^*\| \geq a\}.$$

Then, $\hat{S}_{\varepsilon, a}$ is compact. We assume without loss of generality that $\hat{S}_{\varepsilon, 2\eta}$ is nonempty. Define

$$\varphi(p) = \begin{cases} Ce^{-\frac{1}{1-\|p\|^2}} & \text{if } \|p\| < 1, \\ 0 & \text{if } \|p\| \geq 1, \end{cases}$$

where $C > 0$ is chosen as

$$\int_{\mathbb{R}^n} \varphi(p) dp = 1.$$

For $\delta > 0$, define

$$\varphi_\delta(p) = \delta^n \varphi(p/\delta),$$

and

$$f^\delta(p) = \int_{\mathbb{R}^n} f(p-q) \varphi_\delta(q) dq.$$

It is known that f^δ uniformly converges to f on the compact set S_ε as $\delta \downarrow 0$. Moreover, f^δ is C^∞ .² Choose any C^∞ function $\xi : \mathbb{R} \rightarrow [0, 1]$ such that $\xi^{-1}(1) =]-\infty, 1]$ and $\xi^{-1}(0) = [2, +\infty[$, and for any $p^* \in E^*$, let

$$t_{p^*}(p) = \xi(\eta^{-1}\|p - p^*\|).$$

For every $p \in S_\varepsilon$, define³

$$\begin{aligned} \hat{f}^\delta(p) &= f^\delta(p) - (f^\delta(p) \cdot p)p, \\ \tilde{f}^\delta(p) &= \left(1 - \sum_{p^* \in E^*} t_{p^*}(p)\right) \hat{f}^\delta(p) + \sum_{p^* \in E^*} t_{p^*}(p) f(p). \end{aligned}$$

Because f^δ converges to f uniformly on S_ε as $\delta \downarrow 0$, there exists $\delta > 0$ such that

- 1) For any $p \in S_\varepsilon$, $\tilde{f}^\delta(p) = 0$ if and only if $p \in E^*$.
- 2) If $\tilde{f}^\delta(p) = 0$, then $\tilde{f}^\delta = f$ on some neighborhood of p .
- 3) $p \cdot \tilde{f}^\delta(p) = 0$ for all $p \in S_\varepsilon$.
- 4) For all $p \in \partial S_\varepsilon$, $q \in S_{\varepsilon'}$, and $t \in [0, 1]$, $(1-t)\tilde{f}^\delta(p) + tf^q(p) \neq 0$.
- 5) \tilde{f}^δ is C^1 on S_ε .

²See Appendix C.5 of Evans (2010).

³If necessary, we can define $\tilde{f}^\delta(p) = \tilde{f}^\delta(p/\|p\|)$, and extend the domain of \tilde{f}^δ to a cone generated by S_ε , which includes an open neighborhood of E^* .

Clearly, if \tilde{f}^δ satisfies the claim of this theorem, then f also satisfies the same claim. Hence, hereafter we implicitly use \tilde{f}^δ instead of f , and thus assume that f is C^1 on S_ε .

Recall the definition of the tangent bundle:

$$T(\tilde{S}_\varepsilon) = \{(p, v) | p \in \tilde{S}_\varepsilon, v \in T_p(\tilde{S}_\varepsilon)\}.$$

Define a function $F : [0, 1] \times S_\varepsilon \times \tilde{S}_{\varepsilon'} \rightarrow T(\tilde{S}_\varepsilon)$ as

$$F(t, p, q) = (p, (1-t)f(p) + tf^q(p)),$$

and \tilde{F} as the restriction of F into $[0, 1] \times \tilde{S}_\varepsilon \times \tilde{S}_{\varepsilon'}$. Moreover, we define

$$G_q(t, p) = F(t, p, q), \quad \tilde{G}_q(t, p) = \tilde{F}(t, p, q).$$

Finally, for any $(t, p, q) \in [0, 1] \times \tilde{S}_\varepsilon \times \tilde{S}_{\varepsilon'}$, define $\pi(t, p, q) = q$. Because $[0, 1] \times \tilde{S}_\varepsilon \times \tilde{S}_{\varepsilon'}$ is a C^∞ manifold and π is C^∞ , we can apply Sard's theorem to π and $\partial\pi$, and thus there exists $q \in \tilde{S}_{\varepsilon'}$ such that it is a regular value of π and $\partial\pi$.

Step 7. *If q is a regular value of both π and $\partial\pi$, then both \tilde{G}_q and $\partial\tilde{G}_q$ are transversal to $\tilde{S}_\varepsilon \times \{0\}$.*

Proof of Step 7. First, we verify that \tilde{F} and $\partial\tilde{F}$ are transversal to $\tilde{S}_\varepsilon \times \{0\}$. Note that

$$\begin{aligned} T_p(\tilde{S}_\varepsilon) &= \{v \in \mathbb{R}^n | v \cdot p = 0\}, \\ T_{(p,0)}(\tilde{S}_\varepsilon \times \{0\}) &= T_p(\tilde{S}_\varepsilon) \times \{0\}, \\ T_{(p,0)}(T(\tilde{S}_\varepsilon)) &= T_p(\tilde{S}_\varepsilon) \times T_p(\tilde{S}_\varepsilon). \end{aligned}$$

Suppose that $\tilde{F}(t, p, q) = (p, 0)$. Let $c :]-\varepsilon'', \varepsilon''[\rightarrow \tilde{S}_\varepsilon \times \tilde{S}_{\varepsilon'}$ be a C^1 function such that $c(0) = (p, q)$ and $c'(0) = (v, w)$, and define $d(s) = \tilde{F}(t, c(s))$. Then, $v \cdot p = w \cdot q = 0$, and thus

$$d'(0) = (v, (1-t)df_p(v) + tdf_p^q(v) + (t/p \cdot q)w),$$

which implies that both $d\tilde{F}_{(t,p,q)}$ and $d(\partial\tilde{F})_{(t,p,q)}$ are surjective when $t \neq 0$. Suppose that $t = 0$. By the same argument as in the proof of Step 1, we have that $df_p(v) = 0$ if and only if $v = 0$, and thus for every $w \in T_p(\tilde{S}_\varepsilon)$, there exists $v \in T_p(\tilde{S}_\varepsilon)$ such that $df_p(v) = w$. Therefore, in this case,

$$\begin{aligned} \text{Im}(d\tilde{F}_{(0,p,q)}) + T_{(p,0)}(\tilde{S}_\varepsilon \times \{0\}) &= T_{(p,0)}(T(\tilde{S}_\varepsilon)), \\ \text{Im}(d(\partial\tilde{F})_{(0,p,q)}) + T_{(p,0)}(\tilde{S}_\varepsilon \times \{0\}) &= T_{(p,0)}(T(\tilde{S}_\varepsilon)), \end{aligned}$$

which implies that \tilde{F} and $\partial\tilde{F}$ are transversal to $\tilde{S}_\varepsilon \times \{0\}$. As in the proof of the transversality theorem in Section 2.3 of Guillemin and Pollack (1974), if both π and $\partial\pi$ have a common regular value q , then \tilde{G}_q and $\partial\tilde{G}_q$ are transversal to $\tilde{S}_\varepsilon \times \{0\}$. This completes the proof of Step 7. \square

Step 8. Define $X \equiv \tilde{G}_q^{-1}(\tilde{S}_\varepsilon \times \{0\})$. Then, X is a compact 1-dimensional C^1 manifold with boundary, where $\partial X = \{(0, p^*) | p^* \in E^*\} \cup \{(1, q)\}$.⁴

Proof of Step 8. Because of Step 6,⁵ we have that $X = G_q^{-1}(S_\varepsilon \times \{0\})$. Because G_q is continuous, X is closed. Moreover, because $X \subset [0, 1] \times S_\varepsilon$, X is compact. On the other hand, because \tilde{G}_q and $\partial\tilde{G}_q$ are transversal to $\tilde{S}_\varepsilon \times \{0\}$, we have that X is 1-dimensional C^1 manifold with boundary, where $\partial X = X \cap (\{0, 1\} \times \tilde{S}_\varepsilon)$. Then,

$$X \cap (\{0, 1\} \times \tilde{S}_\varepsilon) = \{(0, p^*) | p^* \in E^*\} \cup \{(1, q)\},$$

and our claim is correct. This completes the proof of Step 8. \square

Step 9. $\text{sign}_{\partial X}(1, q) = (-1)^{n-1}$, and for all $p^* \in E^*$,

$$\text{sign}_{\partial X}(0, p^*) = (-1)^n \text{index}(p^*).$$

Proof of Step 9. Let $n(p) = p$. Then, $n(p)$ is a normal vector field on \tilde{S}_ε . Therefore, using this vector field, we can obtain an orientation $\text{sign}_{\tilde{S}_\varepsilon}$: that is, for a linearly independent family $v_1, \dots, v_{n-1} \in T_p(\tilde{S}_\varepsilon)$,

$$\text{sign}_{\tilde{S}_\varepsilon}(p, v_1, \dots, v_{n-1}) = \text{sign}(\det(p, v_1, \dots, v_{n-1})).$$

Note that $[0, 1]$ can be oriented by the usual function

$$\text{sign}_{[0,1]}(t, a) = \text{sign}(a).$$

Therefore, $[0, 1] \times \tilde{S}_\varepsilon$ is oriented. The tangent bundle $T(\tilde{S}_\varepsilon)$ is oriented by the following function $\text{sign}_{T(\tilde{S}_\varepsilon)}$:

$$\begin{aligned} & \text{sign}_{T(\tilde{S}_\varepsilon)}((p, v), v_1, \dots, v_{n-1}, w_1, \dots, w_{n-1}) \\ &= \text{sign}(\det((p, 0), v_1, \dots, v_{n-1}, (0, p), w_1, \dots, w_{n-1})). \end{aligned}$$

Therefore, X is also oriented by the preimage orientation. Suppose that $t \in \{0, 1\}$, $(t, p) \in X$, and $(1, v) \in T_{(t,p)}(X)$. Choose $v_1, \dots, v_{n-1} \in T_p(\tilde{S}_\varepsilon)$ such that $\det(p, v_1, \dots, v_{n-1}) > 0$. Because $\partial\tilde{G}_{(t,p)}$ is transversal to $\tilde{S}_\varepsilon \times \{0\}$, there exists $v'_1, \dots, v'_{n-1} \in T_p(\tilde{S}_\varepsilon)$ such that $\det(p, v'_1, \dots, v'_{n-1}) > 0$ and for $(v'_i, w'_i) = d\tilde{G}_{(1,q)}(0, v'_i)$, $x = ((v'_1, w'_1), \dots, (v'_{n-1}, w'_{n-1}), (v_1, 0), \dots, (v_{n-1}, 0))$ consists of a basis of $T_{G(t,p)}(T(\tilde{S}_\varepsilon))$.⁶ Then,

$$\begin{aligned} & \text{sign}_X((t, p), (1, v)) \\ &= \text{sign}_{T(\tilde{S}_\varepsilon)}((p, 0), x) \times \text{sign}_{[0,1] \times \tilde{S}_\varepsilon}((t, p), (0, v'_1), \dots, (0, v'_{n-1}), (1, v)). \end{aligned}$$

⁴Note that, because $\varepsilon < \varepsilon'$, we have that $q \in \tilde{S}_\varepsilon$.

⁵Actually, we use property 4) of \tilde{f}^δ .

⁶Note that $d\tilde{G}_{(t,p)}(0, v') = (v', w')$ by the definition of \tilde{G} .

Note that, for $x_1, \dots, x_n \in T_{(t,p)}([0, 1] \times \tilde{S}_\varepsilon)$,

$$\text{sign}_{[0,1] \times \tilde{S}_\varepsilon}((t, p), x_1, \dots, x_n) = \text{sign}(\det(x_1, (0, p), x_2, \dots, x_n)),$$

and thus, we have that

$$\begin{aligned} & \text{sign}_{[0,1] \times \tilde{S}_\varepsilon}((t, p), (0, v'_1), \dots, (0, v'_{n-1}), (1, v)) \\ &= \text{sign}((-1)^{n+1} \det(p, v'_1, \dots, v'_{n-1})) = (-1)^{n+1}. \end{aligned}$$

On the other hand,

$$\begin{aligned} \text{sign}_{T(\tilde{S}_\varepsilon)}((p, 0), x) &= \text{sign}((-1)^{n^2+1} \det(p, w'_1, \dots, w'_{n-1}) \det(p, v_1, \dots, v_{n-1})) \\ &= (-1)^{n+1} \text{sign}(\det(p, w'_1, \dots, w'_{n-1})). \end{aligned}$$

Hence, we obtain that

$$\text{sign}_X((t, p), (1, v)) = \text{sign}(\det(p, w'_1, \dots, w'_{n-1})).$$

Suppose that $t = 1$. Then, $p = q$. Because $w'_i = Df^q(q)v'_i$, by Step 4 and (1), we have that $\text{sign}_X((1, q), (1, v)) = (-1)^{n-1}$, and thus $\text{sign}_{\partial X}(1, q) = (-1)^{n-1}$.

Next, suppose that $t = 0$. Then, $p = p^*$ for some $p^* \in E^*$. Because $w'_i = Df(p)v'_i$, by Step 4, we have that $\text{sign}_X((0, p^*), (1, v)) = (-1)^{n-1} \text{index}(p^*)$. Because $(1, v)$ is an inward vector at $T_{(0,p^*)}(X)$, we have that

$$\text{sign}_{\partial X}(0, p^*) = \text{sign}_X((0, p^*), (-1, -v)) = (-1)^n \text{index}(p^*).$$

This completes the proof of Step 9. \square

Because any compact 1-dimensional C^1 manifold is a finite union of connected components, and each of them is diffeomorphic to either S^1 or $[0, 1]$, we have that

$$\sum_{(t,p) \in \partial X} \text{sign}_{\partial X}(t, p) = 0.$$

By Step 9,

$$\sum_{p^* \in E^*} \text{index}(p^*) = 1,$$

as desired. This completes the proof of Theorem 1. \square

4 Remarks

Our Theorem 1 is, at least formally, different from Proposition 5.6.1 of Mas-Colell (1985). That is, Mas-Colell treated a function f whose domain is S , whereas in our Theorem 1, we treat a function f defined on \mathbb{R}_{++}^n that is homogeneous of degree zero. Moreover, the definitions of $\text{index}(p^*)$ are a little different. However, readers can check that these differences are superficial. First, the restriction of a function treated in Theorem 1 into S satisfies all the assumptions of Mas-Colell's proposition. Conversely, a function treated in Mas-Colell's proposition can be extended to \mathbb{R}_{++}^n by the following formula $f(p) = f(p/\|p\|)$, and this extended function satisfies all requirements of our Theorem 1. Furthermore, Mas-Colell mentioned in Section 5.3 of his book that his definition of $\text{index}(p^*)$ is equivalent to our definition. Therefore, actually our Theorem 1 is equivalent to Mas-Colell's proposition.

This result is closely related to the Poincaré-Hopf index theorem. This theorem claims for every smooth vector field in a compact smooth manifold, the sum of the "indices" of zeroes are the same as the Euler characteristic of this manifold. This theorem is often explained as the existence theorem of "hair whorl." Let us look at the human head as a manifold and consider the direction of hair growth as a vector field. The human head can be seen as the upper hemisphere in \mathbb{R}^3 , and its Euler characteristic is 1. Therefore, there must be a zero with a positive index. This is the hair whorl. From this explanation, this theorem is also called the "hairy ball theorem."

However, upon closer examination, we find something strange. What would happen to the head of an alopecia areata patient? In this case, if we remove the part of the head that is bald from the upper hemisphere, a new manifold is created. This manifold is diffeomorphic to the compact domain of a large disc from which a small disc is removed. It is easy to see that the Euler characteristic of this manifold is 0, because the rotating vector field has no zero. Hence, there must either be no hair whorl or an even number of hair whorls, although we have never heard of a hair whorl disappearing caused by alopecia areata. In other words, if a region is removed by alopecia areata, the Poincaré-Hopf index theorem does not appear to hold.

The reason for this lies in a hidden assumption in the Poincaré-Hopf index theorem. If the manifold has a boundary, then this theorem requires that the vector field is outward on that boundary. Normal human head hair satisfies this condition, and thus this theorem is applicable. However, this condition could not be satisfied on the boundary of the hair lost area caused by alopecia areata, and thus this theorem is not applicable.

With this in mind, let us look at our Theorem 1. Unfortunately, S is not compact, and thus the index theorem is not applicable. On the other

hand, S_ε is compact. Although there is a problem that S_ε is not a manifold, we ignore this problem and check whether the index theorem is applicable. Then, we encounter the problem that we indicated above. That is, our vector field f is not necessarily outward on ∂S_ε . Therefore, the index theorem is still inapplicable.

To solve this problem, Mas-Colell prepared a function f^q , to which the conclusion of the index theorem holds. It is known that the sum of indices is a homotopy invariant. Because f^q is homotopic to f , the sum of indices of f is equal to that of f^q . However, f^q has a unique zero q , whose index is $+1$. Hence, the sum of indices of f is also $+1$. This is the ‘first’ proof of this theorem provided by Mas-Colell. Curiously, Mas-Colell made the ‘second’ proof of this theorem, but this proof is profound and unfamiliar.

There are a number of immediately visible problems with this proof. First, S_ε is not a manifold. Although \tilde{S}_ε is a manifold, it is not compact. If we change the definition of S_ε , then it may possibly be a manifold with boundary. However, even in this case, $[0, 1] \times S_\varepsilon$ is not a manifold with boundary. As we have already seen in the proof, to show that the sum of indices is homotopy invariant, we need to use the transversality of $G_q(t, p)$ to prove that X is a one-dimensional manifold and that the sum of orientations of its boundary is 0. However, for the reasons mentioned above, the domain of G_q is not a manifold, and thus it is unclear whether the result of transversality is applicable. Step 6 is essential for solving this problem. In fact, Mas-Colell himself proved Step 6, but he did not explain why it is needed at all.

Second, to apply the transversality result to G_q , the range must be a manifold. However, because f is a vector field, the range of this function is not a manifold. Therefore, it is necessary to make the range of G_q a manifold artificially by introducing the notion of tangent bundle, and then apply the transversality theorem to F and show that there exists q such that G_q is transversal. However, Mas-Colell did not mention the tangent bundle in his proof.

Hence, Mas-Colell’s proof has a number of big gaps. We think that our proof fulfills these gaps completely, and thus it is worthwhile to many economists that want to use the index theorem.

A Proof of the Classification Theorem

The classification theorem for one-dimensional manifolds can be found in Guillemin and Pollack (1974). However, as already mentioned, their proof uses the Morse function. Since the definition of the Morse function involves second-order derivatives, this argument cannot be applied to C^1 manifolds.

Milnor (1965) provided another proof using arc length. However, this proof is too difficult to understand. On the other hand, Exercise 1.4.9 in Hirsch (1976) asks us to prove this theorem. However, no answer is provided there. Moreover, Hirsch stated in the preface to his book that the exercises contain open problems. Therefore, the existence of this exercise does not solve this problem in any way. Gale (1987) treated this classification theorem. However, he only proved that there is a function that is not a diffeomorphism, but only a homeomorphism. Thus, this cannot be used for our purpose.

In fact, as of November 2022, using Google search, we found a document that proved this theorem rigorously on the server of the University of Lisbon. Upon investigation, we found that the document was written by Gustavo Granja, who provided it for his student. However, there is no signature of the author inside the document, and thus this monograph is not in a position to be added to the reference list. In addition, since it is a lecture document, it could disappear at any time because of Granja's transfer or other reasons. Therefore, in this section, we reconstruct Granja's proof only as we need it.

Recall the definition of S^1 :

$$S^1 = \{x \in \mathbb{R}^2 \mid \|x\| = 1\}.$$

Suppose that $k \in \mathbb{N} \cup \{0, \infty\}$ is given, and suppose that X is a 1-dimensional compact and connected C^k manifold. Our goal is to show the following theorem.

Theorem 2. *X is C^k diffeomorphic to either S^1 or $[0, 1]$.*

Proof. In this proof, we call a function $f : W \rightarrow X$ a **local parametrization of X** if and only if it is a C^k diffeomorphism from W onto $f(W)$, where either $W = (a, b)$ or $W = [a, b)$ for some $a, b \in \mathbb{R}$ with $a < b$ and $f(W)$ is open in X . By definition, we can easily check that $W = [a, b)$ means that $f(a) \in \partial X$.

Let $\varphi : U \rightarrow X$ and $\psi : V \rightarrow X$ be given C^k functions such that φ^{-1} and ψ^{-1} are local parametrizations whose domains are I and J , respectively, and $U, V \subset X$ satisfies

$$U \cap V \neq \emptyset, U \setminus V \neq \emptyset, V \setminus U \neq \emptyset. \quad (2)$$

If a pair (φ, ψ) of functions satisfies the above requirements, then we say that (φ, ψ) satisfies **condition A**.

We separate the proof into nine steps. In Steps 1-6, we assume that (φ, ψ) satisfies condition A. By definition, $I, J \subset \mathbb{R}$ are bounded intervals.⁷ Define

⁷We call a subset of \mathbb{R} an **interval** if this set is a convex set that contains at least two points. If I is a bounded interval whose closure is $[a, b]$, then a, b are called **endpoints** of I .

$\alpha : \varphi(U \cap V) \rightarrow \psi(U \cap V)$ as $\alpha = \psi \circ \varphi^{-1}$. It is easy to show that $\varphi(U \cap V)$ is a (possibly infinitely many) disjoint union of relatively open separated intervals of I .

Step 1. *Suppose that K is a connected component of $\varphi(U \cap V)$ and a is an endpoint of K . Then, $a \notin K$ and if $a \in I$, then*

$$c \equiv \lim_{t \rightarrow a, t \in K} \alpha(t) \notin J,$$

and c is an endpoint of J .

Proof of Step 1. Because J is bounded, $\alpha(t)$ is bounded on K . Moreover, because α is a homeomorphism defined on K , it is monotone, which implies that c is defined and a real number.

Because K is relatively open in I , if a is not an endpoint of I , then $a \notin K$. Suppose that a is an endpoint of I and $a \in K$. Then, $a \in I$ and thus $I = [a, b)$ for some $b > a$, which implies that $\varphi^{-1}(a) \in \partial X$ and $\alpha(a)$ is also an endpoint of J . Let $\sup K = d$. Suppose that $d = b$. Then, $U \subset V$, which contradicts (2). Hence, $d < b$ and $a < d < b$. Let $\delta = \lim_{t \rightarrow d, t \in K} \alpha(t)$. If $\delta \notin J$, then $J = [\alpha(a), \delta)$, and thus $V \subset U$, which contradicts (2). Therefore, $\delta \in J$. Because φ^{-1} and ψ^{-1} are continuous, $\varphi^{-1}(d) = \psi^{-1}(\delta)$. Since ψ is a homeomorphism, $U \cap V$ must include a neighborhood of $\psi^{-1}(\delta)$, and thus K includes $[a, d + \varepsilon)$ for some $\varepsilon > 0$, which is a contradiction. Thus, we conclude that $a \notin K$.

Next, suppose that $c \in J$. If $\varphi^{-1}(a) = \psi^{-1}(c)$, then $a \in K$, which is a contradiction. Therefore, $\varphi^{-1}(a) \neq \psi^{-1}(c)$, which contradicts the continuity of φ^{-1} and ψ^{-1} . Hence, we have that $c \notin J$. This completes the proof. \square

Step 2. *$\varphi(U \cap V)$ consists of either one or two open intervals, and if K is a connected component of $\varphi(U \cap V)$, then K has at least one endpoint in common with I .*

Proof of Step 2. Suppose that K is a connected component of $\varphi(U \cap V)$. By Step 1, we have that K is an open interval, and thus $K = (a, b)$ for some a, b such that $a < b$. Let c, d be endpoints of I such that $c < d$. Then, $c \leq a < b \leq d$. If $c < a$ and $b < d$, then $V = \varphi^{-1}(K) \subset U$, which contradicts (2). Therefore, we have that either $c = a$ or $b = d$, which implies that K has at least one endpoint in common with I . Therefore, there are at most two such K , which completes the proof of Step 2. \square

Step 3. *If $\varphi(U \cap V)$ consists of two connected components, then I and J are open and $U \cup V$ is homeomorphic to S^1 .*

Proof of Step 3. Suppose that a, b are endpoints of I such that $a < b$. Then, there are two connected components $K_1 = (a, c)$ and $K_2 = (d, b)$ of $\varphi(U \cap V)$ such that $c \leq d$. Define

$$e = \lim_{t \uparrow c} \alpha(t), \quad f = \lim_{t \downarrow d} \alpha(t).$$

By Step 1, we have that $e, f \notin J$, and e, f are endpoints of J . If $e = f$, then α is monotone in $K_1 \cup K_2$, and because J is convex, there exists $r \in V \setminus U$ such that $\psi(r) = e = f$. However, this implies that $e, f \in J$, which is a contradiction. Therefore, $e \neq f$, which implies that J is an open interval. By symmetrical arguments, we can show that I is an open interval, and thus $I = (a, b)$.

Thus, either $e < f$ or $e > f$. Because arguments can be done symmetrically, we assume that $e > f$, and thus $J = (f, e)$. Then, α is increasing on both K_i , and $\alpha(K_1) = (g, e)$ and $\alpha(K_2) = (f, h)$ for

$$g = \lim_{t \downarrow a} \alpha(t), \quad h = \lim_{t \uparrow b} \alpha(t).$$

Because $e > f$, we have that $g \geq h$. If $g = h$, then $(U \cup V) \setminus U$ is a singleton, and thus $U \cup V$ can be seen as a one-point compactification of U , which is homeomorphic to S^1 . If $g > h$, then choose an increasing homeomorphism $\lambda : (0, \pi) \rightarrow (a, b)$ and $\mu : [\pi, 2\pi] \rightarrow [h, g]$, and define

$$f(e^{i\theta}) = \begin{cases} \varphi^{-1}(\lambda(\theta)) & \text{if } 0 < \theta < \pi, \\ \psi^{-1}(\mu(\theta)) & \text{if } \pi \leq \theta \leq 2\pi. \end{cases}$$

We can easily check that f is a homeomorphism between S^1 and $U \cup V$. This completes the proof of Step 3. \square

Step 4. Suppose that $\varphi(U \cap V) = K$ for an open interval K whose endpoints are in common with I . Then, $I = [a, b)$ and $K = (a, b)$ for some a, b with $a < b$, and $U \cap \partial X \neq \emptyset$. If $V \cap \partial X = \emptyset$, then $U \cup V$ is homeomorphic to $[0, 1)$. If $V \cap \partial X \neq \emptyset$, then $U \cup V$ is homeomorphic to $[0, 1]$.

Proof of Step 4. Suppose that $K = (a, b)$. Because of (2), we have that $I \neq K$, and thus $I = [a, b)$. Therefore, $\varphi^{-1}(a) \in \partial X$, and thus $U \cap \partial X = \{\varphi^{-1}(a)\}$. Let c, d be endpoints of $\psi(U \cap V)$, and

$$c = \lim_{t \downarrow a} \alpha(t).$$

By Step 1, $c \notin J$ and c is an endpoint of J . Because of (2), we have that $d \in J$, and thus $\psi^{-1}(d) \in V \setminus U$. Suppose that e is another endpoint of J . Then,

J is (e, c) , (c, e) , or $[e, c]$. If $J = (e, c)$, then $J \cap \partial X = \emptyset$ and $e < d$. In this case, choose homeomorphisms $\lambda : [0, 1/2) \rightarrow [a, b)$ and $\mu : [1/2, 1) \rightarrow (e, d]$, and define

$$f(x) = \begin{cases} \varphi^{-1}(\lambda(x)) & \text{if } 0 \leq x < 1/2, \\ \psi^{-1}(\mu(x)) & \text{if } 1/2 \leq x < 1. \end{cases}$$

Then, we can easily check that f is a homeomorphism between $[0, 1)$ and $U \cup V$. The cases $J = (c, e)$ and $J = [e, c]$ can be similarly treated. This completes the proof. \square

Step 5. Suppose that $\varphi(U \cap V) = K$ for an open interval K and there exists an endpoint b of K such that it is not an endpoint of I . Then, $U \cup V$ is homeomorphic to Y , where Y is $(0, 1)$, $[0, 1)$ or $[0, 1]$. Suppose that $f : Y \rightarrow (U \cup V)$ is the diffeomorphism. If $Y = (0, 1)$, then $(U \cup V) \cap \partial X = \emptyset$. If $Y = [0, 1)$, then $(U \cup V) \cap \partial X = \{f(0)\}$. If $Y = [0, 1]$, then $(U \cup V) \cap \partial X = \{f(0), f(1)\}$.

Proof of Step 5. This step can be proved in almost the same manner as the proof of Step 4. Hence, we omit the proof. \square

Step 6. If $U \cup V$ is homeomorphic to S^1 or $[0, 1]$, then $U \cup V = X$.

Proof of Step 6. In this case, $U \cup V$ becomes a clopen set. Therefore, by connectedness of X , we have that $U \cup V = X$. This completes the proof of Step 6. \square

Choose any pair (φ, ψ) that satisfies condition A. Suppose that there exist an open set U' of X and an extension $\varphi' : U' \rightarrow \mathbb{R}$ of φ such that the closure \bar{U} of U is included in U' and (φ', ψ) satisfies condition A. Then, we say that (φ, ψ) satisfies **condition B**.

In Steps 7-8, we suppose that (φ, ψ) satisfies condition B and $\varphi' : U' \rightarrow I'$ is fixed.

Step 7. Suppose that $k \geq 1$, and $\varphi(U \cap V)$ has two connected components. Then, $U \cup V$ is C^k diffeomorphic to S^1 .

Proof of Step 7. As in the proof of Step 3, we define a, b, c, d, e, f, g, h . Then, $I = (a, b)$, $\varphi(U \cap V) = (a, c) \cup (d, b)$,

$$\lim_{t \uparrow c} \alpha(t) = e, \quad \lim_{t \downarrow d} \alpha(t) = f,$$

$$\lim_{t \downarrow a} \alpha(t) = g, \quad \lim_{t \uparrow b} \alpha(t) = h,$$

and $e \neq f$. Because arguments can be done symmetrically, we assume that $e > f$, and thus $J = (f, e)$. Then, α is increasing on both K_i , and $\alpha(K_1) =$

(g, e) and $\alpha(K_2) = (f, h)$. If $g = h$, then $U \cup V = U \cup \{\psi(g)\}$. By Step 3, in this case $U \cup V$ is homeomorphic to S^1 , and thus by Step 6, $U \cup V = X$, which implies that $V \subset X = U'$. This contradicts that U', V satisfies (2). Therefore, we must have that $h < g$. Choose an increasing linear function $\lambda : [-\delta, \pi + \delta] \rightarrow [a - \varepsilon, b + \varepsilon]$ such that $\delta < \pi/2$, $\lambda(0) = a$, $\lambda(\pi) = b$, and $[a - \varepsilon, c) \cup (d, b + \varepsilon] \subset \varphi'(U' \cap V)$, and thus the domain of α can be extended to $[a - \varepsilon, c) \cup (d, b + \varepsilon]$. Define $\beta = \alpha \circ \lambda$. Choose $p, q \in (h, g)$ such that

$$\beta(\pi + \delta/2) < p < q < \beta(-\delta/2).$$

Because $\beta(\pi + \delta) \leq \beta(-\delta)$, we have that such p, q exist. Let $\mu(\theta) = ((2\pi - \theta)p + (\theta - \pi)q)/\pi$ on $[\pi, 2\pi]$, and choose any C^∞ function $\nu : [\pi, 2\pi] \rightarrow [0, 1]$ such that $\nu^{-1}(0) = [\pi, \pi + \delta/2] \cup [2\pi - \delta/2, 2\pi]$, $\nu^{-1}(1) = [\pi + \delta, 2\pi - \delta]$ and $\nu'(\theta) \neq 0$ if $\theta \in (\pi + \delta/2, \pi + \delta) \cup (2\pi - \delta/2, 2\pi)$, and define

$$g(\theta) = (1 - \nu(\theta))(\beta(\theta) + \beta(\theta - 2\pi)) + \nu(\theta)\mu(\theta),$$

where we define $\beta(\theta) = 0$ if $\theta \notin [-\delta, \pi + \delta]$. Then, we can check that g is C^k on $[\pi, 2\pi]$, $g'(\theta) > 0$ for all $\theta \in [\pi, 2\pi]$, and $g \equiv \beta$ on some neighborhood of $\{\pi, 2\pi\}$. Define

$$f(e^{i\theta}) = \begin{cases} \varphi^{-1}(\lambda(\theta)) & \text{if } 0 < \theta < \pi, \\ \psi^{-1}(g(\theta)) & \text{if } \pi \leq \theta \leq 2\pi. \end{cases}$$

Then, f is a C^k diffeomorphism between S^1 and $U \cup V$. This completes the proof of Step 7. \square

Step 8. *Suppose that $k \geq 1$, and $\varphi(U \cap V)$ is an open interval. Then, "homeomorphic" that appears in Steps 4-5 can be replaced with " C^k diffeomorphic."*

Proof of Step 8. We treat only the case in which I, J are open intervals and $\varphi(U \cap V) = K$ for an open interval K , because the remaining cases can be treated in almost the same manner. It is easy to show that there exists a C^∞ diffeomorphism $g_1 : I \rightarrow (0, 1)$ and $g_2 : J \rightarrow (1, 2)$ such that $g_1(\varphi(U \cap V)) = (1/2, 1)$ and $g_2(\psi(U \cap V)) = (1, 3/2)$. Therefore, without loss of generality, we can assume that $I = (0, 1)$, $J = (1, 2)$, and $\varphi(U \cap V) = (1/2, 1)$, $\psi(U \cap V) = (1, 3/2)$. This implies that α is increasing. Because (φ, ψ) satisfies condition B, α can be extended to $(1/2, 1 + \varepsilon]$ for some $\varepsilon \in (0, 1/2)$ such that $\alpha(1 + \varepsilon) < 2$. Choose any C^∞ function $\lambda : (0, 3/2) \rightarrow [0, 1]$ such that $\lambda(x) = 1$ if $x \leq 1$, $\lambda(x) = 0$ if $x \geq 1 + \varepsilon$, and $\lambda'(x) < 0$ for all $x \in (1, 1 + \varepsilon)$. Define

$$g(x) = \alpha(1 + \varepsilon) + 2(x - 1)(2 - \alpha(1 + \varepsilon)),$$

and

$$f(x) = \begin{cases} \varphi^{-1}(x) & \text{if } x < 1, \\ \psi^{-1}(\lambda(x)\alpha(x) + (1 - \lambda(x))g(x)) & \text{if } x \geq 1. \end{cases}$$

Then, f is a C^k diffeomorphism between $(0, 3/2) \rightarrow U \cup V$. This completes the proof of Step 8. \square

Step 9. X is C^k diffeomorphic to S^1 or $[0, 1]$.

Proof of Step 9. In this proof, let \bar{W} denote the closure of W . Choose any open covering (U_i) of X such that there exists a C^k diffeomorphism $\varphi_i : U_i \rightarrow A_i$ such that A_i is either (a, b) or $[a, b]$. Shrinking U_i if needed, we can assume that φ_i can be extended to some open set U'_i of X that includes \bar{U}_i , and φ_i is a C^k diffeomorphism defined on U'_i . Because X is compact, there exists a finite subcovering $(U_{i_j})_{j=1}^n$. Define

$$V_j = U_{i_1} \cup \dots \cup U_{i_j}.$$

Renumbering and removing some U_{i_j} if needed, we can assume that

$$U_{i_j} \cap V_{j-1} \neq \emptyset, U_{i_j} \setminus V_{j-1} \neq \emptyset, V_{j-1} \setminus U_{i_j} \neq \emptyset.$$

Shrinking $U'_{i_1}, \dots, U'_{i_n}$ if needed, we can assume without loss of generality that

$$V_{j-1} \setminus U'_{i_j} \neq \emptyset.$$

If $n = 1$, then $X = U_{i_1}$ is C^k diffeomorphic to either $[0, 1)$ or $(0, 1)$. However, because X is compact, it is impossible. Hence, we have that $n \geq 2$. By Steps 7-8, there exists a C^k diffeomorphism $\psi_2 : V_2 \rightarrow I_2$ such that I_2 is S^1 , $[0, 1]$, $(0, 1)$, or $[0, 1)$. Because X is compact, $V_2 = X$ implies that I_2 is either S^1 or $[0, 1]$, and by Step 6, the converse is also true. Therefore, I_2 is S^1 or $[0, 1]$ if and only if $n = 2$, and if $n > 2$, then I_2 is either $(0, 1)$ or $[0, 1)$. Moreover, if $I_2 = [0, 1)$, then $\psi_2^{-1}(0) \in \partial X$. Hence, (φ_3, ψ_2) satisfies condition B. Inductively, we can assume that for each $j \in \{2, \dots, n\}$, there is a C^k diffeomorphism $\psi_j : V_j \rightarrow I_j$ that satisfies the same condition as ψ_2 , and because $V_n = X$, I_n is either S^1 or $[0, 1]$. This completes the proof of Step 9. \square

Step 9 says that our initial claim is correct. This completes the proof. \square

Reference

- [1] Arrow, K. J., Block, H. D., and Hurwicz, L. (1959) “On the Stability of the Competitive Equilibrium, II.” *Econometrica* 27, pp.82-109.
- [2] Arrow, K. J. and Debreu, G. (1954) “Existence of an Equilibrium for a Competitive Economy”. *Econometrica* 22, pp.265-290.
- [3] Debreu, G. (1952) “Definite and Semi-Definite Quadratic Forms.” *Econometrica* 20, pp.295-300.
- [4] Debreu, G. (1970) “Economies with a Finite Set of Equilibria.” *Econometrica* 38, pp.387-392.
- [5] Evans, L. C. (2010) *Partial Differential Equation: Second Edition*. American Mathematical Society, Providence.
- [6] Gale, D. (1987) “The Classification of 1-Manifolds: A Take-Home Exam.” *American Mathematical Monthly* 94, pp.170-175.
- [7] Guillemin, V. and Pollack, A. (1974) *Differential Topology*. Prentice Hall, New Jersey.
- [8] Hirsch, M. W. (1976) *Differential Topology*. Springer, New York.
- [9] Mas-Colell, A. (1985) *The Theory of General Economic Equilibrium: A Differentiable Approach*. Cambridge University Press, New York.
- [10] Milnor, J. W. (1965) *Topology from the Differential Viewpoint*. The University Press of Virginia, Charlottesville.