

Implications of the Weak Gravity Conjecture for Tidal Love Numbers of Black Holes

Valerio De Luca^{*}, Justin Khoury[†] and Sam S. C. Wong[‡]

*Center for Particle Cosmology, Department of Physics and Astronomy, University of Pennsylvania,
Philadelphia, PA 19104*

Abstract

The Weak Gravity Conjecture indicates that extremal black holes in the low energy effective field theory should be able to decay. This criterion gives rise to non-trivial constraints on the coefficients of higher-order derivative corrections to gravity. In this paper, we investigate the tidal deformability of neutral black holes due to higher-order derivative corrections. As a case in point, we consider a correction of cubic order in the Riemann curvature tensor. The tidal Love numbers of neutral black holes receive leading-order corrections from higher-order derivative terms, since black holes in pure General Relativity have vanishing tidal Love number. We conclude that the tidal deformability of neutral black holes is constrained by the Weak Gravity Conjecture, and therefore provides a test for quantum gravity.

1 Introduction

The recent discovery of gravitational waves has provided a powerful new tool to dig deeper into open questions regarding the interaction of fundamental theory and observation, in particular in probing gravity in its most extreme regime and in searching for signatures of new physics. One of the most important prediction of General Relativity (GR) are black holes. Within this theory, the black hole mass, angular momentum, and electric charge uniquely define its entire multipolar structure as well as its quasinormal modes spectrum, whose observation can be used to probe the so-called no-hair theorems and to test GR [1].

The multipolar structure of a compact object may be modified under the presence of external fields, whose gravitational interaction can result into tidal deformations. Such effects are usually captured in terms of the tidal Love numbers (TLNs) [2]. These strongly depend on the internal properties and structure of the deformed body. TLNs are also found to affect the dynamics of the inspiral of a binary system of compact objects and impact the consequent gravitational waves emission at the fifth post-Newtonian order [3].

A powerful result of GR is that the TLNs of non-rotating and spinning black holes are precisely zero [4–13]. This result has generated an issue of “naturalness” in the gravitational theory [10], and it has been connected with special symmetries of the perturbation fields around black holes [14–21]. This property is, however, broken in higher dimensions [14, 22, 23] and especially in the context of

^{*}vdeluca@sas.upenn.edu

[†]jkhoury@sas.upenn.edu

[‡]scswong@sas.upenn.edu

modified gravity [24–26]. In this paper, as an example case, we will be interested in a theory with a $(R^{\mu\nu}{}_{\rho\sigma})^3$ correction. Such an operator naturally appears when one includes six derivative terms in higher-derivative gravity [27–29] and may also be generated by integrating out massive fields [30]. We will argue that this operator represents the leading higher-order curvature correction that gives rise to non-vanishing TLNs for neutral black holes.

Aside from the TLNs or quasinormal modes, black holes also provide essential information about quantum gravity. The strong gravity environment created by black holes is sensitive to corrections to GR. Especially small black holes are important as space-time curvature gets strong at the horizon. Hawking radiation is one example. Effective field theory (EFT) provides a convenient framework to systematically study higher-derivative corrections to GR. Most of the complicated microscopic degrees of freedom are effectively captured by effective operators at low energies.

However, it is well known that not every bottom-up EFT of gravity can be consistently UV completed. Theories that do not meet this requirement are said to be living in the Swampland [31]. Of many UV consistency conditions, the Weak Gravity Conjecture (WGC) [32] is a well-studied criterion for consistent quantum gravity theories. It is conjectured that there must exist a state with charge to mass ratio (in proper units) larger than unity. There has been evidence for the WGC from the consideration of holography [33–36], black hole entropy [37–39], cosmic censorship [37, 40–42], dimensional reduction [43–47] and importantly, IR consistency [48–52]. The form of WGC we utilize deals with extremal black holes in the EFT [39, 48, 50, 53, 54]. It simply demands that black holes are able to decay and, as a consequence, the small extremal black holes are the states that satisfy the charge to mass ratio bound.

We compute the spectrum of extremal black holes in the theory of interest, and derive a bound on the coefficient of the higher curvature corrections. This constraint carries over to TLNs of neutral black holes in the theory. Broadly speaking, we believe that there is a set of bounds on TLNs of black holes coming from the UV consistency of quantum gravity. Hence, measuring tidal deformations of black holes can potentially provide a test for quantum gravity.

In this work we first extract the WGC constraint on coefficient of the higher-derivative operator $(R^{\mu\nu}{}_{\rho\sigma})^3$ by considering the spectrum of extremal black holes. Then we compute TLNs in the same theory for neutral black holes. We will conclude that the WGC imposes a type of positivity bound on the tidal deformability of black holes.

2 Weak Gravity Conjecture constraints on extremal black holes

In this Section we review the basic idea of the WGC and then apply the conjecture to derive constraints on the theory of interest. The original conjecture [32] states that a consistent theory of quantum gravity should contain a state of charge to mass ratio (in proper units) greater than unity. There are different aspects and different versions of the WGC. We will focus on the infrared consistency of low energy effective theories that stems from the WGC [39, 48, 50, 53, 54], which requires that extremal black holes in the effective field theory be able to decay into sub-extremal black holes.

Typically, in an effective field theory of gravity, the black hole geometry is corrected in the presence of higher-derivative operators. Corrections to the geometry coming from curvature invariants




Figure 1: The curves are indicating the charge $|Q|$ and mass M of extremal black holes in different effective theories. The dotted line is the GR extremality relation. Sub-extremal black holes are under the curves. Black hole configurations above the curves are forbidden as they contain naked singularities. The green curve represents a healthy effective field theory, since extremal black holes can decay into sub-extremal black holes. The red curve represents an effective field theory in which decay products of an extremal black hole must contain states above the curves, and therefore violates the weak gravity conjecture.

such as $R^{\mu\nu}{}_{\rho\sigma}R^{\rho\sigma}{}_{\mu\nu}$, are proportional to some negative power of the size of the black hole. Therefore higher-derivative corrections are more important as the size of the black hole decreases. This is also consistent with the intuition that smaller black holes are more sensitive to UV degrees of freedom.

When it comes to extremal black holes, the charge to mass ratio $|Q|/M$ is corrected in a way that the decay of extremal black holes becomes a nontrivial story, as depicted in Fig. 1. If the corrected extremal relation $|Q|/M$ is below the GR relation, illustrated by the red curve in the figure, then any decay of an extremal black hole into macroscopic black holes must contain a super-extremal black hole with naked singularity. Therefore, this is forbidden by the WGC. On the contrary, if the corrected extremal relation lies above the GR relation (illustrated by the green curve), then extremal black holes can decay into sub-extremal black holes. Consequently, the WGC effectively imposes constraints on the effective operators. It turns out that this constraint is consistent with the relation between corrections to entropy and extremality derived from the thermodynamic point of view [55].

We now study extremal black holes in the specific theory of interest.

2.1 Charged black holes in $(R^{\mu\nu}{}_{\rho\sigma})^3$

To demonstrate the idea that WGC constrains tidal deformations of black holes, we consider a simple theory, namely Einstein-Maxwell gravity corrected by a term cubic in Riemann:

$$S = - \int d^4x \sqrt{-g} \left[\frac{R}{2\kappa^2} - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \alpha R^{\mu\nu}{}_{\rho\sigma} R^{\rho\sigma}{}_{\kappa\lambda} R^{\kappa\lambda}{}_{\mu\nu} \right], \quad (1)$$

where $\kappa^2 = 8\pi G = 1/M_{\text{Pl}}^2$, and α is a coefficient of dimension mass^{-2} . One could also include four-derivative operators, such as R^2 , $R_{\mu\nu}R^{\mu\nu}$ and $R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}$. However, we are interested in the tidal

deformability of neutral black holes in later sections. Since R and $R_{\mu\nu}$ both vanish for a neutral black hole, while $R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}$ can be expressed in terms of the first two terms using the Gauss-Bonnet theorem in four dimensions, they do not modify the Schwarzschild geometry. Moreover, one can also verify that they do not contribute to the tidal Love numbers perturbatively for neutral black holes.¹ Therefore, $(R^{\mu\nu}{}_{\rho\sigma})^3$ is the first non-trivial term that modifies a neutral black hole.²

We consider spherically symmetric, charged black hole solutions of the form

$$\begin{aligned} ds^2 &= -f_t(r)dt^2 + \frac{1}{f_r(r)}dr^2 + r^2d\Omega^2; \\ F &= E(r)dt \wedge dr. \end{aligned} \quad (2)$$

The generic solution at $\mathcal{O}(\alpha)$ is given by,

$$\begin{aligned} f_t(r) &= \frac{(r-r_-)(r-r_+)}{r^2} \left[1 + \frac{\alpha\kappa^2}{r_+^3(r-r_-)} \left\{ 98x^5 - 10\frac{x^5-1}{x-1} \right. \right. \\ &\quad + \frac{6r_-}{7r_+} \left(922x^6 - 317x^5 - 65\frac{x^5-x}{x-1} - 37 \right) \\ &\quad + \frac{6}{7} \left(\frac{r_-}{r_+} \right)^2 \left(-2198x^7 + 1214x^6 - 25x^5 + 101\frac{x^5-1}{x-1} \right) \\ &\quad \left. \left. + \frac{2}{21} \left(\frac{r_-}{r_+} \right)^3 \left(12068x^8 - 7714x^7 + 584x^6 - 445\frac{x^6-1}{x-1} \right) \right\} \right]; \\ f_r(r) &= \frac{(r-r_-)(r-r_+)}{r^2} \left[1 + \frac{\alpha\kappa^2}{r_+^3(r-r_-)} \left\{ -10\frac{x^6-1}{x-1} + \frac{6r_-}{7r_+} \left(36x^6 + 65\frac{x^6-x}{x-1} + 37 \right) \right. \right. \\ &\quad + \frac{6}{7} \left(\frac{r_-}{r_+} \right)^2 \left(-62x^7 - 130x^6 - 101\frac{x^6-1}{x-1} \right) \\ &\quad \left. \left. + \frac{2}{21} \left(\frac{r_-}{r_+} \right)^3 \left(784x^8 + 226x^7 + 550x^6 + 445\frac{x^6-1}{x-1} \right) \right\} \right]; \\ E(r) &= \frac{q}{r^2} \left[1 + 12\alpha\kappa^2 \left(\frac{1}{r_+^4} - \frac{51r_-^2r_+^2}{r^8} + \frac{208r_-r_+(r_-+r_+)}{7r^7} - \frac{9(r_-+r_+)^2}{2r^6} \right) \right], \end{aligned} \quad (3)$$

where r_+ is the outer horizon, and

$$x = \frac{r_+}{r}; \quad \kappa^2 q^2 = 2r_-r_+. \quad (4)$$

Note that r_+ is always the outer horizon, but the inner horizon receives correction from α , *i.e.*, $r_- + \mathcal{O}(\alpha)$ is the inner horizon. A neutral black hole corresponds to $r_- = 0$. In the extremal limit,

¹This is due to the essential fact that the linearized equation of motion receives corrections that are either proportional to the background $\{\bar{R}, \bar{R}_{\mu\nu}\}$ or to the linearized GR equation of motion $\{\delta\bar{R}, \delta\bar{R}_{\mu\nu}\}$.

²There is another independent operator involving six derivatives, that is the parity violating term $\tilde{R}^{\mu\nu}{}_{\rho\sigma}R^{\rho\sigma}{}_{\kappa\lambda}R^{\kappa\lambda}{}_{\mu\nu}$ [27, 28]. Such a term would also modify a neutral black hole. However, it mixes parity even and odd perturbations and therefore introduces an ambiguity in the definition of tidal Love numbers [25].

$r_+ = r_- = r_h$, and the above solution reduces to

$$\begin{aligned}
f_t(r) &= \frac{(r - r_h)^2}{r^2} \left[1 + \frac{\alpha\kappa^2}{r_h^4} \left\{ -\frac{472}{21} \frac{r_h}{r} - \frac{440}{21} \left(\frac{r_h}{r}\right)^2 - \frac{136}{7} \left(\frac{r_h}{r}\right)^3 - \frac{376}{21} \left(\frac{r_h}{r}\right)^4 \right. \right. \\
&\quad \left. \left. - \frac{344}{21} \left(\frac{r_h}{r}\right)^5 + \frac{2920}{7} \left(\frac{r_h}{r}\right)^6 - \frac{4408}{3} \left(\frac{r_h}{r}\right)^7 + \frac{3448}{3} \left(\frac{r_h}{r}\right)^8 \right\} \right]; \\
f_r(r) &= \frac{(r - r_h)^2}{r^2} \left[1 + \frac{\alpha\kappa^2}{r_h^4} \left\{ -\frac{472}{21} \frac{r_h}{r} - \frac{440}{21} \left(\frac{r_h}{r}\right)^2 - \frac{136}{7} \left(\frac{r_h}{r}\right)^3 - \frac{376}{21} \left(\frac{r_h}{r}\right)^4 \right. \right. \\
&\quad \left. \left. - \frac{344}{21} \left(\frac{r_h}{r}\right)^5 + \frac{104}{7} \left(\frac{r_h}{r}\right)^6 - \frac{904}{21} \left(\frac{r_h}{r}\right)^7 + \frac{224}{3} \left(\frac{r_h}{r}\right)^8 \right\} \right]; \\
E(r) &= \frac{q}{r^2} \left[1 + \frac{12\alpha\kappa^2}{r_h^4} \left\{ 1 - 51 \left(\frac{r_h}{r}\right)^8 + \frac{416}{7} \left(\frac{r_h}{r}\right)^7 - 18 \left(\frac{r_h}{r}\right)^6 \right\} \right]. \tag{5}
\end{aligned}$$

Obviously, the charge of the extremal black hole is corrected at $\mathcal{O}(\alpha)$:

$$Q = \frac{1}{4\pi} \int \star F = \pm q \left(1 + 12 \frac{\alpha\kappa^2}{r_h^4} \right) = \pm \frac{\sqrt{2}r_h}{\kappa} \left(1 + 12 \frac{\alpha\kappa^2}{r_h^4} \right), \tag{6}$$

where we have used $r_{\pm} = r_h$. There is also an $\mathcal{O}(\alpha)$ correction to the relation between the ADM mass and the horizon r_h ,

$$M = r_h \left(1 + \frac{236}{21} \frac{\alpha\kappa^2}{r_h^4} \right). \tag{7}$$

It follows that the charge to mass ratio of an extremal black hole is

$$\frac{|Q|}{M} = \frac{\sqrt{2}}{\kappa} \left(1 + \frac{16}{21} \frac{\alpha\kappa^2}{r_h^4} \right). \tag{8}$$

Since the weak gravity conjecture demands that $\frac{\kappa|Q|}{\sqrt{2}M} \geq 1$, we have

$$\boxed{\alpha \geq 0}. \tag{9}$$

This is a non-trivial constraint on the effective theory in Eq. (1). It must be emphasized that from the EFT point of view, it is natural to include all four-derivative operators composed of $R_{\mu\nu\rho\sigma}$ and $F_{\mu\nu}$ such that the WGC bounds a certain linear combination of the coefficients with proper factors of κ . In such case the $\mathcal{O}(\alpha)$ contribution to the extremal condition becomes subleading but is still bounded from below. The simple theory we considered above is just for the purpose of demonstrating the connection between the WGC and TLNs in this work. A detailed analysis will be carried out in a following work.

Let us now switch to study tidal Love numbers of neutral black holes in the same effective theory, and find the implications of the above constraint on the tidal deformability of black holes.

3 Tidal Love numbers

In this Section we review the basics of the computation of tidal Love numbers. We start from their definition in the Newtonian regime, and then move to their relativistic computation for massless spin-1 and spin-2 tidal perturbations in full GR, assuming for simplicity a Schwarzschild black hole background. The interested reader can find a more comprehensive discussion in Refs. [14, 16, 18].

3.1 Newtonian limit

Tidal Love numbers are defined as the response coefficients of a spherically-symmetric body under the action of external tidal perturbations. Consider a spherical body of mass M placed at the origin of a Cartesian coordinate frame. One can adiabatically apply a static external gravitational field U_{ext} perturbing the body. In the multipole expansion, this field can be written as

$$U_{\text{ext}} = - \sum_{\ell=2}^{\infty} \frac{(\ell-2)!}{\ell!} \mathcal{E}_L r^L, \quad (10)$$

in terms of the distance from the origin r , the multi-index $L \equiv i_1 \cdots i_\ell$, and the symmetric trace-free multipole moments \mathcal{E}_L . In response to the external perturbation, the body will deform and develop internal multipole moments given by

$$I_L = \int d^3x \delta\rho(\vec{x}) x^{(L)}, \quad (11)$$

as a function of the body's mass density perturbation $\delta\rho$ and $x^{(L)} \equiv x^{i_1} \cdots x^{i_\ell}$.

Adopting spherical coordinates, the external source and induced response can be expanded in terms of spherical harmonics Y_ℓ^m :

$$\mathcal{E}_{\ell m} \equiv \mathcal{E}_L \int_{\mathbb{S}^2} d\Omega n^L Y_\ell^{m*}; \quad I_{\ell m} \equiv I_L \int_{\mathbb{S}^2} d\Omega n^L Y_\ell^{m*}, \quad (12)$$

where $d\Omega \equiv \sin\theta d\theta d\phi$, and $n^i \equiv x^i/|\vec{x}|$. One can then write the total potential of the system as

$$U_{\text{tot}} = -\frac{GM}{r} - \sum_{\ell=2}^{\infty} \sum_{m=-\ell}^{\ell} Y_{\ell m} \left[\frac{(\ell-2)!}{\ell!} \mathcal{E}_{\ell m} r^\ell - \frac{(2\ell-1)!!}{\ell!} \frac{I_{\ell m}}{r^{\ell+1}} \right]. \quad (13)$$

Assuming an adiabatic and weak external tidal perturbation, linear response theory dictates that the response multipoles should be proportional to the perturbing multipole moments as

$$GI_{\ell m}(\omega) = -\frac{(\ell-2)!}{(2\ell-1)!!} \lambda_{\ell m}(\omega) r_h^{2\ell+1} \mathcal{E}_{\ell m}(\omega), \quad (14)$$

in terms of the characteristic size of the object r_h . The dimensionless coefficients $\lambda_{\ell m}$ describe the response. They are given in terms of the perturbation's frequency ω in the external inertial frame as

$$\lambda_{\ell m} \simeq k_{\ell m} + i\nu_{\ell m}(\omega - m\Omega) + \dots, \quad (15)$$

as a function of the azimuthal harmonic number m and the body's angular velocity Ω . The real term in Eq. (15) describes the conservative response, and the corresponding coefficients $k_{\ell m}$ are called tidal Love numbers, while the imaginary contribution $i\nu_{\ell m}$ describes dissipation effects.

So far our discussion has been entirely based on the Newtonian regime, which is only a long-distance approximation to the full general relativistic picture. In the following we will therefore move to the computation of the tidal Love numbers for a massless spin-1 and spin-2 tidal perturbations, generalising the results found above to a fully relativistic theory. For the purposes of our discussion we will focus on static perturbations ($\omega = 0$), and we will outline the TLN computation for a Schwarzschild black hole.

3.2 GR: vector TLN

The vector TLN of a black hole is equivalent to its electric polarizability and magnetic susceptibility. It can be studied by considering a massless spin-1 vector field in the background of a Schwarzschild black hole, which is described by the Maxwell action

$$S = - \int d^4x \sqrt{-g} \frac{1}{4} F_{\mu\nu}^2. \quad (16)$$

Because of the rotational invariance of the background, one can expand the gauge potential A_μ into perturbations and express them in spherical harmonics as

$$A_\mu = \sum_{\ell, m} \begin{pmatrix} a_0 Y_\ell^m \\ a_r Y_\ell^m \\ a^{(L)} \nabla_i Y_\ell^m + a^{(T)} \epsilon_i^j \nabla_j Y_\ell^m \end{pmatrix}, \quad (17)$$

where the index “ i ” runs over the coordinates on the two sphere, ∇_i and ϵ_i^j are the covariant derivative and Levi-Civita tensor with respect to γ_{ij} on the 2-sphere, respectively. The variables $a^{(L)}$ and $a^{(T)}$ denote the longitudinal and transverse perturbations. On the 2-sphere, the variables a_0 , a_r and $a^{(L)}$ are parity-even, while $a^{(T)}$ is parity odd. Thus they decouple. They are related to each other by electromagnetic duality in four dimensions, and therefore have equal TLN. For simplicity, we focus on $a^{(T)}$ for the computation of the TLN.

Introducing the variable $\Psi_v = a^{(T)}$ and going to tortoise coordinates $dr_\star = dr/f$, where $f = 1 - r_h/r$, one can write down the action for this perturbation as [14]

$$S = \int dt dr_\star \left(\frac{1}{2} \dot{\Psi}_v^2 - \frac{1}{2} \left(\frac{\partial \Psi_v}{\partial r_\star} \right)^2 - \frac{\ell(\ell+1)}{2r^2} f \Psi_v^2 \right). \quad (18)$$

The corresponding equation of motion then takes the form

$$\left(-\frac{\partial^2}{\partial t^2} + \frac{\partial^2}{\partial r_\star^2} - f(r) \frac{\ell(\ell+1)}{r^2} \right) \Psi_v = 0. \quad (19)$$

Focusing on the static limit $\omega = 0$ and assuming that the background is asymptotically flat at spatial infinity ($f \rightarrow 1$ as $r \rightarrow \infty$), one finds that Ψ_v can be expanded asymptotically as

$$\Psi_v \simeq c_1 r^{\ell+1} \left[1 + \dots + k_v^{(\ell)} \left(\frac{r}{r_h} \right)^{-2\ell-1} + \dots \right]. \quad (20)$$

The first term $\sim r^{\ell+1}$ denotes the external tidal field applied at spatial infinity, while the second term $\sim r^{-\ell}$ encodes the quadrupolar response. The coefficient $k_v^{(\ell)}$ denotes the axial (magnetic) vector tidal Love number, and its value depends on the assumed background geometry.

An obvious problem with the above definition is a possible ambiguity due to an overlap between the source series generated by the gravitational non-linearity and the response contribution [11, 16, 22], where subleading corrections to the source appear to have the same power in r as the response in the physical case $\ell \in \mathbb{N}$. In order to get around this ambiguity and properly define the Love numbers through a matching procedure, one needs to compute the graviton corrections to the source term and subtract them from the full GR solution. An alternative approach consists of an analytic continuation to the unphysical region $\ell \in \mathbb{R}$ [11], where the source and response series do not overlap. However obtaining such a solution can be challenging, and for simplicity we will ignore it in the rest of the paper.

Furthermore, we stress that the functional form of the field profile can be more complicated than Eq. (20) due to the presence of logarithmic corrections multiplying the response contribution, coming from EFT loop integrals and resulting into a running Love number for $\ell \gtrsim 3$. Such logarithms are, however, found to cancel out when summing over all loop diagrams for Schwarzschild black holes [56].

To conclude this discussion, let us comment on the relation between the static response coefficient calculated above and the coefficients which appear in the point-particle effective field theory approach. This connection is important in order to provide a gauge invariant definition of the TLN.

The worldline effective field theory approach is based on the fact that, at very large distance, a black hole behaves as a point particle, and corrections due to the object's finite size and internal structure are encoded in higher-derivative operators in the effective theory. Focusing on couplings to a tidal magnetic field, the worldline operator can be built starting from the magnetic field $B_{ab} \equiv F_{ab} = \partial_a A_b - \partial_b A_a$, such that the effective field theory action is given by

$$S = \frac{1}{2} \int d\tau \left(e^{-1} g_{\mu\nu} \frac{dx^\mu}{d\tau} \frac{dx^\nu}{d\tau} - em^2 \right) - \frac{1}{4} \int d^4x F_{\mu\nu} F^{\mu\nu} + \sum_{\ell=1}^{\infty} \frac{1}{2\ell!} \int d\tau e \left[\frac{\lambda_\ell^{(B)}}{2} (\partial_{(a_1} \cdots \partial_{a_{\ell-1}} B_{a_\ell)Tb})^2 \right]. \quad (21)$$

The first term describes the free point particle action, expressed in terms of the worldline vielbein e , while the last operator encodes the magnetic susceptibility of the black hole, proportional to the coefficient $\lambda_\ell^{(B)}$, with $(\cdots)_T$ indicating the trace-free symmetrized part of the enclosed indices. Following Ref. [14], one can relate this Wilson coefficient with the tidal response coefficient $k_v^{(\ell)}$ by matching the gauge-invariant magnetic field as $B_{ij} = \Psi_v \nabla_{[i} \epsilon_{j]}^k \nabla_k Y_\ell^m$, such that

$$\lambda_\ell^{(B)} = k_v^{(\ell)} (-1)^\ell \frac{\pi^{\frac{3}{2}}}{2^{\ell-2}} \frac{\Gamma(\frac{1}{2} - \ell)}{\Gamma(\frac{1}{2})^2} r_h^{2\ell+1}. \quad (22)$$

This result shows that the coupling to B^2 operators in the worldline EFT is absent when the magnetic polarizability vanishes, and that they have the same sign for any $\ell \in \mathbb{N}$.

3.3 GR: tensor TLN

The spherical symmetry of the background geometry allows us to decompose metric perturbations into scalar, vector and tensor spherical harmonics:

$$\begin{aligned}
h_{tt} &= \sum_{\ell,m} f_t(r) H_0(t,r) Y_\ell^m; \\
h_{tr} &= \sum_{\ell,m} H_1(t,r) Y_\ell^m; \\
h_{rr} &= \sum_{\ell,m} f_r(r)^{-1} H_2(t,r) Y_\ell^m; \\
h_{ti} &= \sum_{\ell,m} \left[\mathcal{H}_0(t,r) \nabla_i Y_\ell^m + h_0(t,r) \epsilon_i^j \nabla_j Y_\ell^m \right]; \\
h_{ri} &= \sum_{\ell,m} \left[\mathcal{H}_1(t,r) \nabla_i Y_\ell^m + h_1(t,r) \epsilon_i^j \nabla_j Y_\ell^m \right]; \\
h_{ij} &= \sum_{\ell,m} r^2 \left[\mathcal{K}(t,r) \gamma_{ij} Y_\ell^m + G(t,r) \nabla_{(i} \nabla_{j)T} Y_\ell^m + h_2(t,r) \epsilon_{(i}^k \nabla_{j)} \nabla_k Y_\ell^m \right]. \tag{23}
\end{aligned}$$

The perturbations $H_0, H_1, H_2, \mathcal{H}_0, \mathcal{H}_1, \mathcal{K}, G$ belong to the parity-even sector, while h_0, h_1, h_2 belong to the parity-odd sector on the 2-sphere. At the linearized level, the two types of perturbations decouple in any theory that is parity invariant (such as (1)).

For proof of principle, we focus on parity-odd perturbations. Since there is only one physical combination of these degrees of freedom, we will work in Regge–Wheeler gauge [57], defined by the condition $h_2 = 0$. Following Ref. [58], one can introduce an additional auxiliary field χ , such that the action becomes

$$S_{\text{RW}} = \int dt dr \left[2\chi \left(\dot{h}_1 + \frac{2}{r} h_0 - h_0' \right) - \chi^2 + \frac{2f + 2rf' - \ell(\ell+1)}{r^2} f h_1^2 + \frac{\ell(\ell+1) - 2f - 2rf'}{r^2 f} h_0^2 \right]. \tag{24}$$

We can now integrate out h_0 and h_1 to obtain an action for χ only. Their equations of motion then set

$$h_0 = -\frac{rf}{(\ell-1)(\ell+2)} (2\chi + r\chi'); \quad h_1 = -\frac{r^2}{(\ell-1)(\ell+2)f} \dot{\chi}. \tag{25}$$

Lastly, introducing the Regge–Wheeler perturbation variable $\Psi_{\text{RW}} \equiv \sqrt{\frac{2r^2}{(\ell-1)(\ell+2)}} \chi$, the action reduces to

$$S_{\text{RW}} = \int dt dr_\star \left(\frac{1}{2} \dot{\Psi}_{\text{RW}}^2 - \frac{1}{2} \left(\frac{\partial \Psi_{\text{RW}}}{\partial r_\star} \right)^2 - \frac{1}{2} V_{\text{RW}}(r) \Psi_{\text{RW}}^2 \right); \quad V_{\text{RW}}(r) = f \frac{\ell(\ell+1)}{r^2} - \frac{3ff'}{r}. \tag{26}$$

The equation of motion of Ψ_{RW} takes the form

$$\left(-\frac{\partial^2}{\partial t^2} + \frac{\partial^2}{\partial r_\star^2} - V_{\text{RW}}(r) \right) \Psi_{\text{RW}} = 0. \tag{27}$$

As before, focusing on the static limit and assuming asymptotic flatness at spatial infinity, one finds that Ψ_{RW} can be expanded asymptotically as

$$\Psi_{\text{RW}} = c_1 r^{\ell+1} \left[1 + \dots + k_{\text{RW}}^{(\ell)} \left(\frac{r}{r_h} \right)^{-2\ell-1} + \dots \right]. \quad (28)$$

The coefficient $k_{\text{RW}}^{(\ell)}$ denotes the axial (magnetic) tensor tidal Love number.

In the worldline effective field theory approach, we consider coupling the black hole point particle to gravity. To determine the tidal response of the point particle and match it to the Love numbers found in the full theory, one has to include higher-derivative worldline couplings to the graviton, usually built from the Weyl tensor $C_{\mu\nu\rho\sigma}$. The magnetic part can then be constructed as $B_{ab|c}^{(2)} \equiv C_{0abc}$, such that the corresponding worldline effective action at quadratic order in the graviton fluctuation $g_{\mu\nu} = \eta_{\mu\nu} + 2h_{\mu\nu}/M_{\text{Pl}}$ is given by

$$S = \frac{1}{2} \int d\tau \left(e^{-1} g_{\mu\nu} \frac{dx^\mu}{d\tau} \frac{dx^\nu}{d\tau} - em^2 \right) + \int d^4x \sqrt{-g} \left(\frac{R}{2\kappa^2} + \dots \right) + \sum_{\ell=1}^{\infty} \frac{1}{2\ell!} \int d\tau \frac{\lambda_\ell^{(CB)}}{2} \left(\partial_{(a_1} \dots \partial_{a_{\ell-2}} B_{a_{\ell-1}a_\ell)T|b}^{(2)} \right)^2. \quad (29)$$

The last operator encodes the magnetic susceptibility of the black hole. The Wilson coefficient $\lambda_\ell^{(CB)}$ can be related to the tidal response coefficient k_{RW} using the matching $C_{0rij} \xrightarrow{r \rightarrow \infty} \nabla_{[i} \epsilon_{j]}^k \nabla_k Y_\ell^m r^{-1} \Psi_{\text{RW}}$, to obtain [14]

$$\lambda_\ell^{(CB)} = -k_{\text{RW}}^{(\ell)} (-1)^\ell \frac{\ell + 1}{\ell} \frac{\pi^{\frac{3}{2}}}{2^{\ell-2}} \frac{\Gamma(\frac{1}{2} - \ell)}{\Gamma(\frac{1}{2})^2} r_h^{2\ell+1}. \quad (30)$$

This equation shows therefore how to relate the response coefficients in the worldline EFT to the tidal Love number computed within GR, and it highlights that they have opposite sign for any $\ell \in \mathbb{N}$.

4 Neutral black holes in $(R^{\mu\nu}_{\rho\sigma})^3$

In this Section we describe the calculation of the tidal Love numbers for vector and tensor perturbations in the case of neutral black holes, including a $(R^{\mu\nu}_{\rho\sigma})^3$ correction in the Lagrangian. Recall that the corrected black hole background metric in this case is given by Eq. (3) with $r_- = 0$:

$$f_t = \frac{(r - r_h)}{r} \left[1 + \frac{\alpha\kappa^2}{rr_h^3} \left(98 \left(\frac{r_h}{r} \right)^5 - \frac{10}{r^4} \frac{r_h^5 - r^5}{r_h - r} \right) \right];$$

$$f_r = \frac{(r - r_h)}{r} \left[1 + \frac{\alpha\kappa^2}{rr_h^3} \left(-\frac{10}{r^5} \frac{r_h^6 - r^6}{r_h - r} \right) \right]. \quad (31)$$

4.1 Vector TLN

The presence of a term $(R^{\mu\nu}_{\rho\sigma})^3$ in the action for a spin-1 tidal field has the effect of modifying the background metric of a neutral black hole, as shown above, but otherwise leaves the vector field

equation of motion invariant. The latter is given by

$$\nabla_\nu F^{\mu\nu} = 0, \quad (32)$$

in terms of the vector field strength $F^{\mu\nu}$. The corresponding axial perturbation satisfies then the equation of motion in the static limit

$$\frac{d^2\Psi_\ell}{dr_\star^2} - f_t(r)\frac{\ell(\ell+1)}{r^2}\Psi_\ell = 0, \quad (33)$$

where the tortoise coordinate is now defined as $dr_\star = dr/\sqrt{f_r f_t}$. This equation can be solved perturbatively in the coupling strength α . Considering the multipole moments $\ell = 1$ and $\ell = 2$, and imposing regularity of the solution at the black hole horizon r_h , the solution at zeroth order in α has the following form

$$\begin{aligned} \Psi_{\ell=1}^{(0)} &= c_1 r^2; \\ \Psi_{\ell=2}^{(0)} &= c_1 \left(r^3 - \frac{3}{4} r^2 r_h \right), \end{aligned} \quad (34)$$

in terms of a dimensional constant c_1 . We stress that we have made use of the fact that the metric perturbations f_r and f_t coincide at zeroth-order in α . The absence of an induced dipolar or quadrupolar term implies that the dipole and axial vector TLNs of an unperturbed neutral black hole are both zero.

At first order in α , the metric perturbations f_r and f_t receive a correction, as shown in Eq. (31). Expanding the solution as $\Psi = \Psi^{(0)} + \alpha\Psi^{(1)}$, one finds that the first-order solutions are given by

$$\begin{aligned} \Psi_{\ell=1}^{(1)} &= -c_1 \frac{\kappa^2}{r_h^2} \left[4 \left(\frac{r_h}{r} \right) + 3 \left(\frac{r_h}{r} \right)^2 + \frac{12}{5} \left(\frac{r_h}{r} \right)^3 - 22 \left(\frac{r_h}{r} \right)^4 \right]; \\ \Psi_{\ell=2}^{(1)} &= -c_1 \frac{\kappa^2}{r_h} \left[\frac{15}{2} \left(\frac{r}{r_h} \right)^2 + \frac{219}{20} \left(\frac{r_h}{r} \right)^2 - \frac{197}{5} \left(\frac{r_h}{r} \right)^3 + \frac{33}{2} \left(\frac{r_h}{r} \right)^4 \right]. \end{aligned} \quad (35)$$

Combined with (34), the full solutions in the asymptotic regime $r \rightarrow \infty$ thus take the form

$$\begin{aligned} \Psi_{\ell=1} &= \Psi_{\ell=1}^{(0)} + \alpha\Psi_{\ell=1}^{(1)} \simeq c_1 r^2 \left[1 + k_V^{(\ell=1)} \left(\frac{r_h}{r} \right)^3 \right]; \\ \Psi_{\ell=2} &= \Psi_{\ell=2}^{(0)} + \alpha\Psi_{\ell=2}^{(1)} \simeq c_1 r^3 \left[1 + k_V^{(\ell=2)} \left(\frac{r_h}{r} \right)^5 \right]. \end{aligned} \quad (36)$$

From these one can extract the tidal response coefficients as

$$\begin{aligned} k_V^{(\ell=1)} &= -4 \frac{\alpha\kappa^2}{r_h^4}; \\ k_V^{(\ell=2)} &= -\frac{219}{20} \frac{\alpha\kappa^2}{r_h^4}. \end{aligned} \quad (37)$$

The corresponding Wilson coefficients in the worldline effective field theory approach are then

$$\begin{aligned}\lambda_{\ell=1}^{(B)} &= -16\pi \frac{\alpha\kappa^2}{r_h}; \\ \lambda_{\ell=2}^{(B)} &= -\frac{73}{5}\pi r_h \alpha\kappa^2.\end{aligned}\tag{38}$$

This result shows that a positive value for the coupling α , as dictated by the WGC, would result in a negative axial vector tidal Love number.

Proceeding to the computation for higher ℓ , we know that the extraction of the Love number is not well defined, as discussed in the previous Section. In particular, the perturbative solution at $\ell = 3$ is given by

$$\begin{aligned}\Psi_{\ell=3} &= c_1 \left(r^4 - \frac{4r^3 r_h}{3} + \frac{2r^2 r_h^2}{5} \right) + c_1 \alpha\kappa^2 \left\{ -\frac{40}{3} \left(\frac{r}{r_h} \right)^3 + 8 \left(\frac{r_h}{r} \right)^2 \right. \\ &\quad \left. - \left[\frac{184616}{1225} - \frac{704}{7} \ln \left(\frac{r}{r_h} \right) \right] \left(\frac{r_h}{r} \right)^3 + \mathcal{O} \left[\left(\frac{r_h}{r} \right)^4, \left(\frac{r_h}{r} \right)^4 \ln \left(\frac{r}{r_h} \right) \right] \right\}.\end{aligned}\tag{39}$$

By identifying the coefficient of $r^{-\ell}$, one finds

$${}^{\text{“}k_V^{\ell=3}\text{”}} = \frac{\alpha\kappa^2}{r_h^4} \left[-\frac{184616}{1225} + \frac{704}{7} \ln \left(\frac{r}{r_h} \right) \right],\tag{40}$$

where a running behaviour appears.

Let us also stress that, in principle, a non-vanishing vector TLN may arise if one considers higher-order derivative terms in the effective action containing the field strength $F_{\mu\nu}$, *e.g.*, $R^{\mu\nu\rho\sigma} F_{\mu\nu} F_{\rho\sigma}$ studied in [59]. However, since such terms would not generate any tensor response for neutral black holes, we have neglected them in our analysis.

4.2 Tensor TLN

In this Section we compute the TLN in the parity-odd sector of tensor perturbations, where only h_0 , h_1 and h_2 in Eq. (23) are relevant. We further choose Regge-Wheeler gauge $h_2 = 0$. The quadratic action, in terms of h_0 , h_1 in a neutral black hole background, of the theory (1) is

$$\begin{aligned}S_{(2)} &= \int dt dr \left[\frac{1}{4\kappa^2} \left(h_0' - \frac{2}{r} h_0 + \dot{h}_1 \right)^2 + \frac{j^2 + 2 - 2f - 6rf' - r^2 f''}{4\kappa^2 r^2 f} h_0^2 + f \frac{4f + 6rf' + r^2 f''}{4\kappa^2 r^2} h_1^2 \right. \\ &\quad \left. + \alpha \tilde{\mathcal{L}}[h_0, h_1] \right],\end{aligned}\tag{41}$$

where $j^2 = \ell(\ell + 1)$, and $f = 1 - r_h/r$. We have written $f_t = f + \alpha\delta f_t$, $f_r = f + \alpha\delta f_r$ and collected all $\mathcal{O}(\alpha)$ terms in $\alpha\tilde{\mathcal{L}}$. The explicit form of the latter is given in Appendix A. The first line in (41) is the same as in GR, while the second line includes terms coming from the $\mathcal{O}(\alpha)$ corrections to the background metric as well as the quadratic action from the $\alpha(R^{\mu\nu}{}_{\rho\sigma})^3$ operator.

We now extract the gauge-invariant variable by introducing once again an auxiliary field χ as in [58],

$$\begin{aligned}
& S_{(2)}[h_0, h_1, \chi] \\
&= \int dt dr \left[\frac{1}{4\kappa^2} \left(h_0' - \frac{2}{r} h_0 + \dot{h}_1 \right)^2 + \frac{j^2 + 2 - 2f - 6rf' - r^2 f''}{4\kappa^2 r^2 f} h_0^2 + f \frac{4f + 6rf' + r^2 f''}{4\kappa^2 r^2} h_1^2 \right. \\
&\quad \left. - \frac{1}{4\kappa^2} \left(h_0' - \frac{2}{r} h_0 + \dot{h}_1 - \chi \right)^2 + \alpha \tilde{\mathcal{L}}[h_0, h_1] \right]. \tag{42}
\end{aligned}$$

It is obvious that χ is a gauge invariant variable [60], since its equation of motion gives

$$\chi = h_0' - \frac{2}{r} h_0 + \dot{h}_1. \tag{43}$$

Since we are treating α perturbatively, one can introduce

$$\begin{aligned}
h_0 &= h_0^{(0)} + \alpha h_0^{(1)}; \\
h_1 &= h_1^{(0)} + \alpha h_1^{(1)}; \\
\chi &= \chi^{(0)} + \alpha \chi^{(1)}. \tag{44}
\end{aligned}$$

Keeping terms up to first order in α , the quadratic action becomes

$$\begin{aligned}
& S_{(2)}[h_0^{(0)}, h_1^{(0)}, \chi^{(0)}, h_0^{(1)}, h_1^{(1)}, \chi^{(1)}] \\
&= S_{(2)}[h_0^{(0)}, h_1^{(0)}, \chi^{(0)}] + \alpha \int dt dr \left[\frac{1}{2\kappa^2} \left(h_0^{(0)'} - \frac{2}{r} h_0^{(0)} + \dot{h}_1^{(0)} \right) \left(h_0^{(1)'} - \frac{2}{r} h_0^{(1)} + \dot{h}_1^{(1)} \right) \right. \\
&\quad + \frac{j^2 + 2 - 2f - 6rf' - r^2 f''}{2\kappa^2 r^2 f} h_0^{(0)} h_0^{(1)} + f \frac{4f + 6rf' + r^2 f''}{2\kappa^2 r^2} h_1^{(0)} h_1^{(1)} \\
&\quad \left. - \frac{1}{2\kappa^2} \left(h_0^{(0)'} - \frac{2}{r} h_0^{(0)} + \dot{h}_1^{(0)} - \chi^{(0)} \right) \left(h_0^{(1)'} - \frac{2}{r} h_0^{(1)} + \dot{h}_1^{(1)} - \chi^{(1)} \right) \right]. \tag{45}
\end{aligned}$$

It is clear that $h_0^{(1)}$ and $h_1^{(1)}$ are now Lagrange multipliers. Their equations of motion give

$$\begin{aligned}
h_0^{(0)} &= \frac{r_h - r}{j^2 - 2} \left(2\chi^{(0)} + r\chi^{(0)'} \right); \\
h_1^{(0)} &= \frac{r^3}{(j^2 - 2)(r_h - r)} \dot{\chi}^{(0)}. \tag{46}
\end{aligned}$$

After substituting these back to the action (45), we are left with an action for $\chi^{(0)}$ and $\chi^{(1)}$ only.³ The equation of motion for χ from this action can be recast in the Regge-Wheeler form after rescaling χ by

$$\chi^{(0)} + \alpha \chi^{(1)} = \frac{\sqrt{2(j^2 - 2)}}{r} \left(\Psi^{(0)} + \alpha \Psi^{(1)} \right). \tag{47}$$

³Of course, even though $h_0^{(1)}, h_1^{(1)}$ are gone in the Lagrangian, there are still equations of motion relating them to $\chi^{(0)}, \chi^{(1)}$.




Figure 2: Pictorial representation of the tidal deformability of a black hole with a $(R^{\mu\nu}_{\rho\sigma})^3$ term and corresponding sign of the tensor tidal Love number for the multipole $\ell = 2$.

The resulting equations are

$$\begin{aligned} \left(-\partial_t^2 + \partial_{r^*}^2 + \left(1 - \frac{r_h}{r}\right) \frac{3r_h - j^2 r}{r^3}\right) \Psi^{(0)} &= 0; \\ \left(-\partial_t^2 + \partial_{r^*}^2 + \left(1 - \frac{r_h}{r}\right) \frac{3r_h - j^2 r}{r^3}\right) \Psi^{(1)} &= \mathcal{S}[\Psi^{(0)}], \end{aligned} \quad (48)$$

where $\partial_{r^*} = (1 - r_h/r) \partial_r$. The explicit form of the source term $\mathcal{S}[\Psi^{(0)}]$ is given in Appendix A.

We are now ready to study the static limit where Ψ only depends on r . It should be emphasized that the static limit should be taken in the gauge invariant combination Ψ instead of h_0 and h_1 . Then the solution for h_0, h_1 are calculated from Ψ . Following the procedure in the previous sections, the regular solution for $\Psi_{\ell=2}$ is

$$\Psi_{\ell=2} = c_1 r^3 + \frac{\alpha \kappa^2}{r_h} c_1 \left(-184 \left(\frac{r_h}{r}\right)^2 + 80 \left(\frac{r_h}{r}\right)^3 + 184 \left(\frac{r_h}{r}\right)^4 \right). \quad (49)$$

Therefore the tidal Love number in the Regge-Wheeler variable for $\ell = 2$ is⁴

$$k_{\text{RW}}^{(\ell=2)} = -184 \frac{\alpha \kappa^2}{r_h^4}. \quad (50)$$

Correspondingly, the coupling in the worldline effective action is

$$\boxed{\lambda_{\ell=2}^{(CB)} = 368\pi r_h \alpha \kappa^2}. \quad (51)$$

This result shows that a positive value for the coupling α , as dictated by the WGC, would result in a positive parity-odd tidal Love number. See Fig. 2 for a pictorial representation.

When one continues the analysis to higher ℓ , the story becomes subtle, as mentioned earlier, since the identification of the Love number is unclear. For instance, the perturbative solution

⁴A similar computation has been performed in Ref. [61] for the metric tensor perturbations (h, H) , without however estimating the TLN for the more appropriate gauge invariant variable. The same functional dependence on the coupling α and black hole Schwarzschild radius is found.

at $\ell = 3$ is

$$\Psi_{\ell=3} = c_1 \left(r^4 - \frac{5}{6} r_h r^3 \right) + c_1 \alpha \kappa^2 \left\{ -\frac{25}{3} \left(\frac{r}{r_h} \right)^3 - 372 \left(\frac{r_h}{r} \right)^2 - \left[\frac{190418}{441} - \frac{4730}{7} \ln \left(\frac{r}{r_h} \right) \right] \left(\frac{r_h}{r} \right)^3 + \mathcal{O} \left[\left(\frac{r_h}{r} \right)^4, \left(\frac{r_h}{r} \right)^4 \ln \left(\frac{r}{r_h} \right) \right] \right\}. \quad (52)$$

One can still identify the coefficient of $1/r^\ell$, which is

$$“k_{\text{RW}}^{(\ell=3)}” = \frac{\alpha \kappa^2}{r_h^4} \left[-\frac{190418}{441} + \frac{4730}{7} \ln \left(\frac{r}{r_h} \right) \right]. \quad (53)$$

However, it is unclear whether this is still the Love number as it is no longer the dominating tail. The same issue appears when one consider a tidal scalar field in the same background [56].

We conclude this section with a final remark on the assumed response function. In particular, we have computed only the “linear” response of the black hole to the external tidal perturbation. In principle a $\alpha (R^{\mu\nu}{}_{\rho\sigma})^3$ term would generate also a nonlinear response when the non-linear source term in the equation of motion is taken into account. At each order of the perturbative series in α , the non-linear source are simply the product of lower-order terms in the series. For instance, at $\mathcal{O}(\alpha)$ and $\ell = 4$, a source proportional to $(C^{\ell=2})^2$ should appear, where $C^{\ell=2}$ is the magnitude of the source tidal field. However, it mixes parity-odd and -even modes, and requires the use of Clebsch-Gordan coefficients between vector and tensor spherical harmonics. The computation of this response would be quite challenging, and the extraction of the corresponding TLN is still not well defined. We have therefore neglected this contribution for the sake of our discussion.

5 Conclusions

Within the low energy effective field theory, extremal black holes should be able to decay according to the WGC. The latter sets non-trivial bounds on the sign of the coefficients of higher-order derivative corrections to GR. On the other hand, the presence of these operators can have an impact on the multipolar structure of black holes. Indeed, even though neutral black holes in pure GR are characterized by a vanishing tidal Love number, which measures the static deformability of compact objects under external tidal perturbations, the presence of these higher-order derivative corrections can modify this result and give rise to a nonzero tidal Love number.

Adopting the example case of a $(R^{\mu\nu}{}_{\rho\sigma})^3$ term in the effective gravity action with coupling α , we showed that the WGC dictates only positive values for the coupling α . By computing the tidal deformability of neutral black holes, which is found to be proportional to α , we concluded that only positive values for the tidal Love numbers (in the point particle effective action) are allowed, therefore providing a test for quantum gravity. Given the dimensionful nature of this coupling, we stress that integrating out massive fields with a lower cut-off scale would eventually enhance the tidal Love numbers.

Constraints on the sign and size of the couplings of higher-derivative terms in the effective action of gravity can also be set from other observables. For example, the presence of terms like Gauss-Bonnet operators are known to affect also the graviton 3-point interaction, leading to violations of

causality unless the value of the coupling α is very small, independently of its sign [62]. A term like $(R^{\mu\nu}{}_{\rho\sigma})^3$ would therefore impact as well the scattering amplitude, probably resulting into a similar constraints on its coupling coefficient [27, 63]. Furthermore, a complete unitarity analysis of all derivative-six operators, composed of $R^{\mu\nu}{}_{\rho\sigma}$ and $F_{\mu\nu}$, along the line of [50–52] would be valuable.

Finally, we stress that higher-derivative operators like $(R^{\mu\nu}{}_{\rho\sigma})^3$ may impact as well on the black hole entropy, which would be corrected by a term proportional to α , with the entropy increasing only for positive values of α [61].

Acknowledgments

We thank A. Kehagias, D. Lombardo, T. Noumi, R. Penco, A. Riotto, I. Rothstein and L. Santoni for interesting comments and discussions. V.DL. is supported by funds provided by the Center for Particle Cosmology at the University of Pennsylvania. The work of J.K. and S.W. is supported in part by the DOE (HEP) Award DE-SC0013528.

A Explicit form of the action and equation of motion for Ψ

For completeness, we report the explicit form of the action $\tilde{\mathcal{L}}[h_0, h_1]$ in Eq. (41) and the source term for the equation of motion of the perturbation field Ψ in the tensor sector of Eq. (48). The former is given by

$$\begin{aligned}
& \tilde{\mathcal{L}}[h_0, h_1] = \\
& \frac{3h_1r_h\dot{h}_0(j^2r(7r-5r_h)+r_h(21r_h-25r))}{r^7(r-r_h)} - \frac{3h_0r_h\dot{h}_1'((8j^2-4)r-9r_h)}{2r^6} - \frac{3h_1r_h\dot{h}_0'(4(2j^2-3)r-9r_h)}{2r^6} \\
& + \frac{3h_0r_h\dot{h}_1((13j^2+10)r^2-3(5j^2+24)rr_h+66r_h^2)}{r^7(r-r_h)} + \frac{6h_0r_h\dot{h}_1''(3r_h-2r)}{r^5} + \frac{3h_1r_h\dot{h}_0''(6r_h-5r)}{r^5} \\
& + \frac{3h_0r_h\dot{h}_1'''(r-r_h)}{r^4} + \frac{3h_1r_h\dot{h}_0'''(r-r_h)}{r^4} + \frac{6h_1r_h\ddot{h}_0}{r^3(r-r_h)} + \frac{3h_0r_h\ddot{h}_1}{r^2(r-r_h)^2} - \frac{3h_0r_h\ddot{h}_1'}{r^2(r-r_h)} - \frac{3h_1r_h\ddot{h}_0'}{r^2(r-r_h)} \\
& + \frac{3h_0r_h\dot{h}_0'(39r_h-10j^2r)}{r^7} + \frac{3h_0r_h\dot{h}_0''(4(j^2-6)r+27r_h)}{2r^6} + \frac{6h_0r_h\ddot{h}_0(j^2r-r_h)}{r^4(r-r_h)^2} + \frac{6h_0r_h\dot{h}_0'''(3r-4r_h)}{r^5} \\
& - \frac{3h_0r_h\dot{h}_0''''(r-r_h)}{r^4} + \frac{h_0\ddot{h}_0'(6r_h^2-9rr_h)}{r^3(r-r_h)^2} + \frac{3h_0r_h\ddot{h}_0''}{r^2(r-r_h)} + \frac{6h_1(j^2-2)r_h\dot{h}_1''(r-r_h)^2}{r^7} \\
& + \frac{3h_1r_h\ddot{h}_1(4(j^2+4)r-33r_h)}{2r^6} - \frac{6h_1(j^2-2)r_h\dot{h}_1'(5r^2-12rr_h+7r_h^2)}{r^8} + \frac{3h_1r_h\ddot{h}_1'(3r-4r_h)}{r^5} \\
& - \frac{3h_1r_h\ddot{h}_1''(r-r_h)}{r^4} + \frac{3h_1r_h\ddot{h}_1'''}{r^2(r-r_h)} + \frac{h_0^2((-6j^4+185j^2-10)r^2r_h^4+5(j^2-2)r^6+5(j^2-2)r^5r_h)}{2r^8r_h^3(r-r_h)} \\
& + \frac{h_0^2(5(j^2-2)r^4r_h^2+5(j^2-2)r^3r_h^3-2(113j^2+176)rr_h^5+354r_h^6)}{2r^8r_h^3(r-r_h)} \\
& + \frac{h_1^2(-3(2j^4+13j^2+200)r^2r_h^5+5(j^2-2)r^7+6(j^2-2)^2r^3r_h^4+2(29j^2+701)rr_h^6-816r_h^7)}{2r^{10}r_h^3}.
\end{aligned} \tag{54}$$

We stress that in principle the expression can be further simplified. The source term is instead given by

$$\begin{aligned}
& \frac{1}{\kappa^2}\mathcal{S}[\Psi^{(0)}] = \\
& (r-r_h)^2 \left(\frac{1080(r-r_h)r_h^2}{(j^2-2)r^9} - \frac{4(194r_h^5+r[-211r_h^4+5r(r_h+r)(r_h^2+r^2)])}{r^8r_h^3} \right) \partial_r^2\Psi^{(0)} \\
& + \left(\frac{24(81r-116r_h)(r-r_h)^2r_h^2}{(j^2-2)r^{10}} + \frac{2(5r^7-10r^6r_h-198r^2r_h^5+415rr_h^6-212r_h^7)}{r^9r_h^3} \right) \partial_r\Psi^{(0)} \\
& + \left(-\frac{24r_h^2(r-r_h)^2(9r+26r_h)}{(j^2-2)r^{11}} + \frac{2j^2(5r^6-432r^2r_h^4+900rr_h^5-473r_h^6)}{r^9r_h^3} \right. \\
& \left. + \frac{6(5r^7-10r^6r_h+198r^2r_h^5-409rr_h^6+216r_h^7)}{r^{10}r_h^3} \right) \Psi^{(0)},
\end{aligned} \tag{55}$$

where we have repeatedly used the equation of motion (48) for $\Psi^{(0)}$ to reduce the higher-order derivative terms from $(R^{\mu\nu}{}_{\rho\sigma})^3$.

References

- [1] LIGO SCIENTIFIC, VIRGO, KAGRA collaboration, *Tests of General Relativity with GWTC-3*, [2112.06861](#).
- [2] E. Poisson and C.M. Will, *Gravity: Newtonian, Post-Newtonian, Relativistic*, Cambridge University Press (2014), [10.1017/CBO9781139507486](#).
- [3] E.E. Flanagan and T. Hinderer, *Constraining neutron star tidal Love numbers with gravitational wave detectors*, *Phys. Rev. D* **77** (2008) 021502 [[0709.1915](#)].
- [4] T. Binnington and E. Poisson, *Relativistic theory of tidal Love numbers*, *Phys. Rev. D* **80** (2009) 084018 [[0906.1366](#)].
- [5] T. Damour and A. Nagar, *Relativistic tidal properties of neutron stars*, *Phys. Rev. D* **80** (2009) 084035 [[0906.0096](#)].
- [6] T. Damour and O.M. Lecian, *On the gravitational polarizability of black holes*, *Phys. Rev. D* **80** (2009) 044017 [[0906.3003](#)].
- [7] P. Pani, L. Gualtieri, A. Maselli and V. Ferrari, *Tidal deformations of a spinning compact object*, *Phys. Rev. D* **92** (2015) 024010 [[1503.07365](#)].
- [8] P. Pani, L. Gualtieri and V. Ferrari, *Tidal Love numbers of a slowly spinning neutron star*, *Phys. Rev. D* **92** (2015) 124003 [[1509.02171](#)].
- [9] N. Gürlebeck, *No-hair theorem for Black Holes in Astrophysical Environments*, *Phys. Rev. Lett.* **114** (2015) 151102 [[1503.03240](#)].
- [10] R.A. Porto, *The Tune of Love and the Nature(ness) of Spacetime*, *Fortsch. Phys.* **64** (2016) 723 [[1606.08895](#)].
- [11] A. Le Tiec and M. Casals, *Spinning Black Holes Fall in Love*, *Phys. Rev. Lett.* **126** (2021) 131102 [[2007.00214](#)].
- [12] H.S. Chia, *Tidal deformation and dissipation of rotating black holes*, *Phys. Rev. D* **104** (2021) 024013 [[2010.07300](#)].
- [13] A. Le Tiec, M. Casals and E. Franzin, *Tidal Love Numbers of Kerr Black Holes*, *Phys. Rev. D* **103** (2021) 084021 [[2010.15795](#)].
- [14] L. Hui, A. Joyce, R. Penco, L. Santoni and A.R. Solomon, *Static response and Love numbers of Schwarzschild black holes*, *JCAP* **04** (2021) 052 [[2010.00593](#)].

- [15] P. Charalambous, S. Dubovsky and M.M. Ivanov, *Hidden Symmetry of Vanishing Love Numbers*, *Phys. Rev. Lett.* **127** (2021) 101101 [[2103.01234](#)].
- [16] P. Charalambous, S. Dubovsky and M.M. Ivanov, *On the Vanishing of Love Numbers for Kerr Black Holes*, *JHEP* **05** (2021) 038 [[2102.08917](#)].
- [17] L. Hui, A. Joyce, R. Penco, L. Santoni and A.R. Solomon, *Near-zone symmetries of Kerr black holes*, *JHEP* **09** (2022) 049 [[2203.08832](#)].
- [18] P. Charalambous, S. Dubovsky and M.M. Ivanov, *Love symmetry*, [2209.02091](#).
- [19] M.M. Ivanov and Z. Zhou, *Vanishing of black hole tidal Love numbers from scattering amplitudes*, [2209.14324](#).
- [20] T. Katagiri, M. Kimura, H. Nakano and K. Omukai, *Vanishing Love of Black Holes in General Relativity: From Spacetime Conformal Symmetry of a Two-dimensional Reduced Geometry*, [2209.10469](#).
- [21] A. Kehagias, D. Perrone and A. Riotto, *Quasinormal Modes and Love Numbers of Kerr Black Holes from AdS₂ Black Holes*, [2211.02384](#).
- [22] B. Kol and M. Smolkin, *Black hole stereotyping: Induced gravito-static polarization*, *JHEP* **02** (2012) 010 [[1110.3764](#)].
- [23] V. Cardoso, L. Gualtieri and C.J. Moore, *Gravitational waves and higher dimensions: Love numbers and Kaluza-Klein excitations*, *Phys. Rev. D* **100** (2019) 124037 [[1910.09557](#)].
- [24] V. Cardoso, E. Franzin, A. Maselli, P. Pani and G. Raposo, *Testing strong-field gravity with tidal Love numbers*, *Phys. Rev. D* **95** (2017) 084014 [[1701.01116](#)].
- [25] V. Cardoso, M. Kimura, A. Maselli and L. Senatore, *Black Holes in an Effective Field Theory Extension of General Relativity*, *Phys. Rev. Lett.* **121** (2018) 251105 [[1808.08962](#)].
- [26] M. Cvetič, G.W. Gibbons, C.N. Pope and B.F. Whiting, *Supergravity black holes, Love numbers, and harmonic coordinates*, *Phys. Rev. D* **105** (2022) 084035 [[2109.03254](#)].
- [27] R.R. Metsaev and A.A. Tseytlin, *Curvature Cubed Terms in String Theory Effective Actions*, *Phys. Lett. B* **185** (1987) 52.
- [28] S. Endlich, V. Gorbenko, J. Huang and L. Senatore, *An effective formalism for testing extensions to General Relativity with gravitational waves*, *JHEP* **09** (2017) 122 [[1704.01590](#)].
- [29] M. Ruhdorfer, J. Serra and A. Weiler, *Effective Field Theory of Gravity to All Orders*, *JHEP* **05** (2020) 083 [[1908.08050](#)].
- [30] C. de Rham and A.J. Tolley, *Speed of gravity*, *Phys. Rev. D* **101** (2020) 063518 [[1909.00881](#)].
- [31] C. Vafa, *The String landscape and the swampland*, [hep-th/0509212](#).

- [32] N. Arkani-Hamed, L. Motl, A. Nicolis and C. Vafa, *The String landscape, black holes and gravity as the weakest force*, *JHEP* **06** (2007) 060 [[hep-th/0601001](#)].
- [33] Y. Nakayama and Y. Nomura, *Weak gravity conjecture in the AdS/CFT correspondence*, *Phys. Rev. D* **92** (2015) 126006 [[1509.01647](#)].
- [34] D. Harlow, *Wormholes, Emergent Gauge Fields, and the Weak Gravity Conjecture*, *JHEP* **01** (2016) 122 [[1510.07911](#)].
- [35] N. Benjamin, E. Dyer, A.L. Fitzpatrick and S. Kachru, *Universal Bounds on Charged States in 2d CFT and 3d Gravity*, *JHEP* **08** (2016) 041 [[1603.09745](#)].
- [36] M. Montero, G. Shiu and P. Soler, *The Weak Gravity Conjecture in three dimensions*, *JHEP* **10** (2016) 159 [[1606.08438](#)].
- [37] G. Shiu, P. Soler and W. Cottrell, *Weak Gravity Conjecture and extremal black holes*, *Sci. China Phys. Mech. Astron.* **62** (2019) 110412 [[1611.06270](#)].
- [38] A. Hebecker and P. Soler, *The Weak Gravity Conjecture and the Axionic Black Hole Paradox*, *JHEP* **09** (2017) 036 [[1702.06130](#)].
- [39] C. Cheung, J. Liu and G.N. Remmen, *Proof of the Weak Gravity Conjecture from Black Hole Entropy*, *JHEP* **10** (2018) 004 [[1801.08546](#)].
- [40] G.T. Horowitz, J.E. Santos and B. Way, *Evidence for an Electrifying Violation of Cosmic Censorship*, *Class. Quant. Grav.* **33** (2016) 195007 [[1604.06465](#)].
- [41] T. Crisford, G.T. Horowitz and J.E. Santos, *Testing the Weak Gravity - Cosmic Censorship Connection*, *Phys. Rev. D* **97** (2018) 066005 [[1709.07880](#)].
- [42] T.-Y. Yu and W.-Y. Wen, *Cosmic censorship and Weak Gravity Conjecture in the Einstein–Maxwell–dilaton theory*, *Phys. Lett. B* **781** (2018) 713 [[1803.07916](#)].
- [43] J. Brown, W. Cottrell, G. Shiu and P. Soler, *Fencing in the Swamp: Quantum Gravity Constraints on Large Field Inflation*, *JHEP* **10** (2015) 023 [[1503.04783](#)].
- [44] J. Brown, W. Cottrell, G. Shiu and P. Soler, *On Axionic Field Ranges, Loopholes and the Weak Gravity Conjecture*, *JHEP* **04** (2016) 017 [[1504.00659](#)].
- [45] B. Heidenreich, M. Reece and T. Rudelius, *Sharpening the Weak Gravity Conjecture with Dimensional Reduction*, *JHEP* **02** (2016) 140 [[1509.06374](#)].
- [46] B. Heidenreich, M. Reece and T. Rudelius, *Evidence for a sublattice weak gravity conjecture*, *JHEP* **08** (2017) 025 [[1606.08437](#)].
- [47] S.-J. Lee, W. Lerche and T. Weigand, *Tensionless Strings and the Weak Gravity Conjecture*, *JHEP* **10** (2018) 164 [[1808.05958](#)].
- [48] C. Cheung and G.N. Remmen, *Infrared Consistency and the Weak Gravity Conjecture*, *JHEP* **12** (2014) 087 [[1407.7865](#)].

- [49] S. Andriolo, D. Junghans, T. Noumi and G. Shiu, *A Tower Weak Gravity Conjecture from Infrared Consistency*, *Fortsch. Phys.* **66** (2018) 1800020 [[1802.04287](#)].
- [50] Y. Hamada, T. Noumi and G. Shiu, *Weak Gravity Conjecture from Unitarity and Causality*, *Phys. Rev. Lett.* **123** (2019) 051601 [[1810.03637](#)].
- [51] B. Bellazzini, M. Lewandowski and J. Serra, *Positivity of Amplitudes, Weak Gravity Conjecture, and Modified Gravity*, *Phys. Rev. Lett.* **123** (2019) 251103 [[1902.03250](#)].
- [52] N. Arkani-Hamed, Y.-t. Huang, J.-Y. Liu and G.N. Remmen, *Causality, unitarity, and the weak gravity conjecture*, *JHEP* **03** (2022) 083 [[2109.13937](#)].
- [53] Y. Kats, L. Motl and M. Padi, *Higher-order corrections to mass-charge relation of extremal black holes*, *JHEP* **12** (2007) 068 [[hep-th/0606100](#)].
- [54] T. Noumi and H. Satake, *Higher derivative corrections to black brane thermodynamics and the weak gravity conjecture*, [2210.02894](#).
- [55] G. Goon and R. Penco, *Universal Relation between Corrections to Entropy and Extremality*, *Phys. Rev. Lett.* **124** (2020) 101103 [[1909.05254](#)].
- [56] M.M. Ivanov and Z. Zhou, *Black Hole Tidal Love Numbers and Dissipation Numbers in Worldline Effective Field Theory*, [2208.08459](#).
- [57] T. Regge and J.A. Wheeler, *Stability of a Schwarzschild singularity*, *Phys. Rev.* **108** (1957) 1063.
- [58] A. De Felice, T. Suyama and T. Tanaka, *Stability of Schwarzschild-like solutions in $f(R,G)$ gravity models*, *Phys. Rev. D* **83** (2011) 104035 [[1102.1521](#)].
- [59] S. Garcia-Saenz, A. Held and J. Zhang, *Schwarzschild quasi-normal modes of non-minimally coupled vector fields*, *JHEP* **05** (2022) 139 [[2202.07131](#)].
- [60] K. Martel and E. Poisson, *Gravitational perturbations of the Schwarzschild spacetime: A Practical covariant and gauge-invariant formalism*, *Phys. Rev. D* **71** (2005) 104003 [[gr-qc/0502028](#)].
- [61] S. Cai and K.-D. Wang, *Non-vanishing of tidal Love numbers*, [1906.06850](#).
- [62] X.O. Camanho, J.D. Edelstein, J. Maldacena and A. Zhiboedov, *Causality Constraints on Corrections to the Graviton Three-Point Coupling*, *JHEP* **02** (2016) 020 [[1407.5597](#)].
- [63] A. Gruzinov and M. Kleban, *Causality Constrains Higher Curvature Corrections to Gravity*, *Class. Quant. Grav.* **24** (2007) 3521 [[hep-th/0612015](#)].