

ON INCLUSION RELATIONS BETWEEN WEIGHTED SPACES OF ENTIRE FUNCTIONS

GERHARD SCHINDL

ABSTRACT. We characterize the inclusions of weighted classes of entire functions in terms of the defining weights resp. weight systems. First we treat weights defined in terms of a so-called associated weight function where the weight(system) is based on a given sequence. The abstract weight function case is then reduced to the weight sequence setting by using the so-called associated weight sequence. As an application of the main statements we characterize closedness under point-wise multiplication of these classes.

1. INTRODUCTION

Weighted spaces of entire functions are defined as follows:

$$(1.1) \quad H_v^\infty(\mathbb{C}) := \{f \in H(\mathbb{C}) : \|f\|_v := \sup_{z \in \mathbb{C}} |f(z)|v(|z|) < +\infty\}.$$

Here $H(\mathbb{C})$ denotes the class of entire functions and $v : [0, +\infty) \rightarrow (0, +\infty)$ is the (radial and strictly positive) *weight function*. From now on, except stated explicitly otherwise, we assume that v is

- (*) continuous,
- (*) non-increasing and
- (*) rapidly decreasing, i.e. $\lim_{t \rightarrow +\infty} t^k v(t) = 0$ for all $k \geq 0$.

This is the same notion to be a weight (function) as it has been considered in [7], [5], [22] and in [3]. (In [3, Sect. 2] such weighted classes are even introduced for weights which are not necessarily non-increasing and rapidly decreasing; such v 's are called *general weights*.) It is known that $H_v^\infty(\mathbb{C})$ is a Banach space w.r.t. the norm $\|\cdot\|_v$ and similarly such weighted classes can be defined when \mathbb{C} is replaced by any other (non-empty) open subset $G \subseteq \mathbb{C}$ and $v : G \rightarrow (0, +\infty)$. (Another very prominent example in the literature is the unit disc $G = \mathbb{D}$; in this article we will focus on $G = \mathbb{C}$ except in Section 3.1.)

Such weighted classes (and operators on them) have been studied during the last years from different aspects; very recently in [3] a detailed survey of available results on this topic is given, see also the numerous literature citations therein.

The aim of this paper is to study and compare growth relations between weights and inclusion relations of the corresponding weighted spaces. To be more precise, let v and w be two weights and

Date: November 29, 2022.

2020 Mathematics Subject Classification. 30D15, 30D60, 46E05, 46E15.

Key words and phrases. Weighted classes of entire functions, inclusion relation, weight sequences and weight functions, associated weight function.

G. Schindl is supported by FWF-Project P33417-N.

write $v \preceq w$ if

$$(1.2) \quad w(t) = O(v(t)) \text{ as } t \rightarrow +\infty.$$

Let us call v and w *equivalent*, written $v \sim w$, if

$$v \preceq w \text{ and } w \preceq v.$$

Note that relation \sim is denoted by the symbol \approx in [3] and by \asymp in [10] and [11]. Obviously, by definition one has

$$(1.3) \quad v \preceq w \implies H_v^\infty(\mathbb{C}) \subseteq H_w^\infty(\mathbb{C}),$$

with continuous inclusion (and similarly if \mathbb{C} is replaced by any other subset $G \subseteq \mathbb{C}$). So equivalent weights yield equivalent weighted classes (as locally convex vector spaces) and the main question is: What can be said about the implication \Leftarrow in (1.3), i.e., can the inclusion (resp. equality) of weighted spaces be characterized in terms of a growth relation between two given weights?

To treat this problem the idea is to focus first on the case when $v = v_M : t \mapsto \exp(-\omega_M(t))$ with $M \in \mathbb{R}_{>0}^{\mathbb{N}}$ a given (weight) sequence satisfying mild standard growth assumptions and ω_M denotes the so-called *associated weight function*, see Section 2.4. Such particular weights have been considered in [22] where solid hulls and cores of the corresponding weighted classes have been studied. The advantage in this setting is that the weights v_M are based on a given sequence M ; therefore more structure and information is involved and available. In particular, this fact is illustrated by Theorem 3.2. Desired properties for v_M can be expressed in terms of growth and regularity assumptions (known and used in the ultradifferentiable and ultraholomorphic setting) for M .

Then, when an abstract weight v is given, the aim is to reduce this case to the weight sequence setting. This can be achieved by considering the so-called *associated weight sequence* M^v , see Section 2.7. Note that in view of Remarks 2.13, 2.14 in order to involve M^v our standard assumptions on v to be a weight are convenient.

By combining a recent result from [11] and the additional information for v_M (see the crucial Lemma 2.16) in the first main result we prove in Theorem 3.2 that any v_M is *essential*. More generally, if given v is *essential*, then by using a consequence of the recent works [10] and [11] we show that in this case in (1.3) equivalence holds true (for $G = \mathbb{D}$ and $G = \mathbb{C}$), see Theorem 3.3.

Combining both statements, in Theorem 3.4 (see also Corollary 3.15) the equivalence $v_M \preceq v_N \iff H_{v_M}^\infty(\mathbb{C}) \subseteq H_{v_N}^\infty(\mathbb{C})$ is characterized by the following growth relation:

$$(1.4) \quad \exists A \geq 1 \forall j \in \mathbb{N} : N_j \leq AM_j.$$

All this can be found in Section 3.1. It turns out that in order to use the associated weight sequence M^v it suffices to assume that v is (log)-convex which is in general weaker than essentiality for v in the weighted entire case, see Remark 3.5. The comments summarized in this Remark together with Theorem 3.6 give also a comparison between essential and (log)-convex weights which are satisfying the crucial technical assumption *moderate growth*, see Definition 2.18.

Moreover, (1.4) is much more restrictive than the (known) comparison relation

$$(1.5) \quad \sup_{j \in \mathbb{N}_{>0}} \left(\frac{M_j}{N_j} \right)^{1/j} < +\infty,$$

which is crucial in the ultradifferentiable setting. In (1.5) instead resp. in addition of an uniform constant A the geometrically growing factor $j \mapsto c^j$ is appearing with a parameter $c > 0$. This

difference is connected to a dilatation in the function argument in the weight v_M , see (2.17) and (2.18). It turns out (see Proposition 3.8) that (1.5) is becoming relevant for weighted spaces given by *non-increasing* resp. *non-decreasing systems of weights*, see e.g. [1, Sect. 2]. On the other hand, when treating such spaces we have to replace \preceq by a more general growth relation involving a parameter, see (2.6).

Moreover, in view of this observation two kinds of weight systems can be considered "naturally" also for abstractly given weights. We can either consider $t \mapsto v(ct)$ (dilatation-type, motivated by the aforementioned explanations) or $t \mapsto v(t)^c$ (exponential-type). Here $c > 0$ is the parameter and it means, for v_M , that c appears "inside" the function argument of ω_M as a dilatation/stretching parameter resp. "outside the function" as a multiplicative parameter. The latter case requires another generalization of \preceq , see (2.7). All precise definitions can be found in detail in Section 2.2.

Summarizing, we are able to prove characterizations for the inclusion (resp. equality) of weighted spaces purely in terms of given sequences M and N both for dilatation- and exponential-type systems (when the system is based on a weight sequence) and to reduce then the general weight function case to the weight sequence situation by involving the associated weight sequence.

Based on these results it is then possible to study and characterize closedness under the point-wise multiplication operator $\mathfrak{m} : (f, g) \mapsto f \cdot g$ acting on such weighted classes. This is mainly due to the fact that in the weight sequence setting point-wise multiplication yields naturally the appearance of the so-called *convolved sequence*, see Section 7.2. However, our characterization for weight systems is in contrast to the fact that a weighted class defined by a single weight function v (see (1.1)) is not closed under \mathfrak{m} , see Proposition 7.2 and Corollary 7.15. Moreover, note that concerning the behaviour of \mathfrak{m} acting on $H_v^\infty(G)$ there is a difference between $G = \mathbb{C}$ and $G = \mathbb{D}$; in [4] the case $H_v^\infty(\mathbb{D})$ has been studied in detail.

The paper is structured as follows: In Section 2 all relevant information on weights (functions, sequences and associated functions and sequences), weight systems of both types and their corresponding classes is gathered. Then, in Section 3 we state and prove the main characterizing results; apart from the first statement Theorem 3.4 see Theorems 3.14, 3.21, 3.30 and 3.34.

In Section 4 we comment on the corresponding results for classes of σ -growth restriction (see e.g. [1, Sect. 1.B]) and in Section 5 we compare the dilatation- and exponential-type structures and prove a characterization when both notions coincide, see Theorems 5.4 and 5.6.

In Section 6 we treat the situation when relation (1.2) resp. the weaker growth relations (2.6), (2.7) for weight systems are replaced by their corresponding σ -growth counterparts. We obtain analogue results, see Theorems 6.6, 6.9, 6.13 and 6.16. Finally, in Section 7 we characterize closedness under \mathfrak{m} , see Theorems 7.8 and 7.13. In particular, in the dilatation-type weight sequence case the characterizing condition for M is precisely the classical *moderate growth* (i.e. (M.2) in [17]) resp. the corresponding condition (2.28) in the weight function approach.

Acknowledgements. The author thanks Prof. José Bonet Solves from the Universitat Politècnica de València for helpful and clarifying discussions and explanations during the preparation of this work. In particular, for forwarding and explaining (the proof of) Proposition 7.2 and the recent article [3].

2. WEIGHTS, WEIGHTED SPACES AND GROWTH CONDITIONS

2.1. Basic convention and comments. We use the notation $\mathbb{N} = \{0, 1, \dots\}$ and $\mathbb{N}_{>0} = \{1, 2, \dots\}$.

We call a weight v *normalized* when $v(t) = 1$ for all $t \in [0, 1]$. A normalized v satisfies $v(t) \leq 1$ for all $t \geq 0$.

When both v, w are (normalized) weight functions, then the product $v \cdot w$ is so, too.

For defining the class in (1.1) normalization can be assumed w.l.o.g.: Otherwise replace v by a normalized weight, say v^n , and such that $v(t) = v^n(t)$ for all large $t > 1$. This gives $H_v^\infty(\mathbb{C}) = H_{v^n}^\infty(\mathbb{C})$ as l.c.v.s.

Since $w(t), v(t) \neq 0$ for all t , (1.2) precisely means that

$$\exists C \geq 1 \forall t \geq 0 : w(t) \leq Cv(t).$$

2.2. Weight systems and corresponding weighted classes. Let $\underline{v} = (v_n)_{n \in \mathbb{N}_{>0}}$ be a non-increasing sequence of weights, i.e. $v_n \geq v_{n+1}$ for all n . Then we define the (LB)-space $H_{\underline{v}}^\infty(\mathbb{C})$ of the Banach spaces $H_{v_n}^\infty(\mathbb{C})$, so

$$(2.1) \quad H_{\underline{v}}^\infty(\mathbb{C}) := \varprojlim_{n \rightarrow \infty} H_{v_n}^\infty(\mathbb{C}).$$

Analogously, if $\overline{v} = (v_n)_{n \in \mathbb{N}_{>0}}$ is a non-decreasing sequence of weights, i.e. $v_n \leq v_{n+1}$ for all n , then we define the Fréchet-space $H_{\overline{v}}^\infty(\mathbb{C})$ to be the projective limit of the Banach spaces $H_{v_n}^\infty(\mathbb{C})$, so

$$(2.2) \quad H_{\overline{v}}^\infty(\mathbb{C}) := \varprojlim_{n \rightarrow \infty} H_{v_n}^\infty(\mathbb{C}).$$

Let v be a weight function and $c > 0$, then set

$$(2.3) \quad v_c : t \mapsto v(ct), \quad v^c : t \mapsto v(t)^c,$$

and write again v instead of v_1 resp. v^1 . We summarize:

- (i) v is a weight function if and only if some/each weight v_c resp. if and only if some/each v^c is so.
- (ii) For any $d \geq c > 0$ we have the continuous inclusions

$$H_{v_c}^\infty(\mathbb{C}) \subseteq H_{v_d}^\infty(\mathbb{C}), \quad H_{v^c}^\infty(\mathbb{C}) \subseteq H_{v^d}^\infty(\mathbb{C}).$$

Note that $v^d(t) \leq v^c(t)$ for all large t since eventually $v(t) \leq 1$ (and for all $t \geq 0$ if v is normalized).

- (iii) $v \preceq w$ if and only if $v_c \preceq w_c$ if and only if $v^c \preceq w^c$ for some/each $c > 0$.

Consider the corresponding weight systems

$$(2.4) \quad \underline{v}_c := (v_c)_{c \in \mathbb{N}_{>0}}, \quad \overline{v}_c := (v_{\frac{1}{c}})_{c \in \mathbb{N}_{>0}}.$$

Thus \underline{v}_c is a non-increasing sequence of weights, whereas \overline{v}_c is a non-decreasing sequence. Similarly, if $v \leq 1$, then consider

$$(2.5) \quad \underline{v}^c := (v^c)_{c \in \mathbb{N}_{>0}}, \quad \overline{v}^c := (v^{\frac{1}{c}})_{c \in \mathbb{N}_{>0}}.$$

Since $v \leq 1$ the set \underline{v}^c is a non-increasing sequence of weights, whereas \overline{v}^c is a non-decreasing sequence. The assumption $v \leq 1$ holds if v is normalized.

Definition 2.1. *The weight systems in (2.4) are called to be of dilatation-type and the systems in (2.5) of exponential-type.*

Let us write $v \preceq_c w$ if

$$(2.6) \quad \exists c \geq 1 : v \preceq_w c,$$

and $v \sim_c w$, if

$$v \preceq_c w \quad \text{and} \quad w \preceq_c v.$$

Similarly, write $v \preceq^c w$ if

$$(2.7) \quad \exists c \geq 1 : v \preceq_w^c,$$

and $v \sim^c w$, if

$$v \preceq^c w \quad \text{and} \quad w \preceq^c v.$$

Note:

(*) Both $v \preceq_c w$ and $v \preceq^c w$ are weaker than $v \preceq w$.

(*) $v \preceq_c w$ resp. $v \preceq^c w$ if and only if

$$\exists c \geq 1 \forall a > 0 : v_a \preceq_w a c \quad \text{resp.} \quad v^a \preceq_w^c a c.$$

These observations imply the following:

Proposition 2.2. *Let v and w be (normalized) weight functions. If $v \preceq_c w$, then*

$$H_{\underline{v}_c}^\infty(\mathbb{C}) \subseteq H_{\underline{w}_c}^\infty(\mathbb{C}), \quad H_{\overline{v}_c}^\infty(\mathbb{C}) \subseteq H_{\overline{w}_c}^\infty(\mathbb{C}),$$

and if $v \preceq^c w$, then

$$H_{\underline{v}^c}^\infty(\mathbb{C}) \subseteq H_{\underline{w}^c}^\infty(\mathbb{C}), \quad H_{\overline{v}^c}^\infty(\mathbb{C}) \subseteq H_{\overline{w}^c}^\infty(\mathbb{C}),$$

with continuous inclusions.

We close with the following observation:

Remark 2.3. *Let v be a (normalized) weight function, then*

$$(2.8) \quad \forall d > c > 0 : \lim_{t \rightarrow +\infty} \frac{v^c(t)}{v^d(t)} = \lim_{t \rightarrow +\infty} v^{c-d}(t) = +\infty.$$

However, in general $\lim_{t \rightarrow +\infty} \frac{v_c(t)}{v_d(t)} = +\infty$ is not clear.

2.3. Weight sequences and conditions. Let $M = (M_j)_j \in \mathbb{R}_{>0}^{\mathbb{N}}$. The corresponding sequence of quotients is denoted by $\mu_j := \frac{M_j}{M_{j-1}}$, $j \geq 1$, and $\mu_0 := 1$. M is called *normalized* if $1 = M_0 \leq M_1$ and M is called *log-convex* if

$$\forall j \in \mathbb{N}_{>0} : M_j^2 \leq M_{j-1} M_{j+1},$$

equivalently if the sequence $(\mu_j)_j$ is non-decreasing. If $M_0 = 1$ and M is log-convex, then $(M_j)^{1/j} \leq \mu_j$ for all $j \in \mathbb{N}_{>0}$.

Definition 2.4. *M is called a weight sequence if*

$$1 = M_0 \quad \text{and} \quad \lim_{j \rightarrow +\infty} (M_j)^{1/j} = +\infty.$$

Moreover consider the set

$$\mathcal{LC} := \{M \in \mathbb{R}_{>0}^{\mathbb{N}} : M \text{ is normalized, log-convex, } \lim_{j \rightarrow +\infty} (M_j)^{1/j} = +\infty\},$$

i.e. $M \in \mathcal{LC}$ if and only if M is a log-convex weight sequence with $M_1 \geq 1$. Note that each $M \in \mathcal{LC}$ is non-decreasing since $\mu_j \geq 1$ for all j and $j \mapsto \mu_j$ is non-decreasing.

M (with $M_0 = 1$) has condition *moderate growth*, denoted by (mg), if

$$\exists C \geq 1 \forall j, k \in \mathbb{N} : M_{j+k} \leq C^{j+k} M_j M_k.$$

In [17] this is denoted by (M.2) and also known under the name *stability under ultradifferential operators*.

Let $M, N \in \mathbb{R}_{>0}^{\mathbb{N}}$, we write $M \leq N$ if $M_j \leq N_j$ for all j and $M \preceq N$ if (1.5), i.e.

$$\sup_{j \in \mathbb{N}_{>0}} \left(\frac{M_j}{N_j} \right)^{1/j} < +\infty.$$

M and N are called *equivalent*, written $M \approx N$, if

$$M \preceq N \quad \text{and} \quad N \preceq M.$$

Property (mg) is clearly preserved under \approx .

Remark 2.5. Let M be a log-convex weight sequence then there exists an equivalent $N \in \mathcal{LC}$ such that $\mu_j = \nu_j$ for all j sufficiently large:

Since $\lim_{j \rightarrow +\infty} \mu_j = +\infty$ and $(\mu_j)_j$ is non-decreasing, there exists $j_0 \in \mathbb{N}_{>0}$ such that $\mu_j \geq \mu_{j_0} > 1$ for all $j \geq j_0$. Then set

$$\nu_j := 1, \quad j < j_0, \quad \text{and} \quad \nu_j := \mu_j, \quad j \geq j_0,$$

and consequently

$$(2.9) \quad \exists C \geq 1 \forall j \in \mathbb{N} : \frac{1}{C} N_j \leq M_j \leq C N_j.$$

2.4. Associated weight function. Let $M \in \mathbb{R}_{>0}^{\mathbb{N}}$ (with $M_0 = 1$), then the *associated weight function* $\omega_M : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R} \cup \{+\infty\}$ is defined by

$$\omega_M(t) := \sup_{j \in \mathbb{N}} \log \left(\frac{t^j}{M_j} \right) \quad \text{for } t \in \mathbb{R}_{>0}, \quad \omega_M(0) := 0.$$

For an abstract introduction of the associated function we refer to [18, Chapitre I], see also [17, Definition 3.1].

If $\liminf_{j \rightarrow +\infty} (M_j)^{1/j} > 0$, then $\omega_M(t) = 0$ for sufficiently small t , since $\log \left(\frac{t^j}{M_j} \right) < 0 \Leftrightarrow t < (M_j)^{1/j}$ holds for all $j \in \mathbb{N}_{>0}$. (In particular, if $M_j \geq 1$ for all $j \in \mathbb{N}$, then ω_M is vanishing on $[0, 1]$.) Moreover, under this assumption $t \mapsto \omega_M(t)$ is a continuous non-decreasing function, which is convex in the variable $\log(t)$ and tends faster to infinity than any $\log(t^j)$, $j \geq 1$, as $t \rightarrow +\infty$. $\lim_{j \rightarrow +\infty} (M_j)^{1/j} = +\infty$ implies that $\omega_M(t) < +\infty$ for each finite t which shall be considered as a basic assumption for defining ω_M . In particular, this holds true provided M is a weight sequence.

If M is a log-convex weight sequence, then let us introduce the counting function

$$(2.10) \quad \Sigma_M(t) := |\{j \geq 1 : \mu_j \leq t\}|.$$

In this case we can compute M by involving ω_M as follows, see [18, Chapitre I, 1.4, 1.8] and also [17, Prop. 3.2]:

$$(2.11) \quad M_j = \sup_{t \geq 0} \frac{t^j}{\exp(\omega_M(t))}, \quad j \in \mathbb{N}.$$

If M is a weight sequence which is not log-convex, then the right-hand side of (2.11) yields M_j^{lc} with M^{lc} denoting the *log-convex minorant* of M and M^{lc} is a log-convex weight sequence.

Moreover, one has

$$\forall t \in [0, \mu_1] : \omega_M(t) = 0,$$

which follows by the known integral representation formula (see [18, 1.8. III] and also [17, (3.11)])

$$(2.12) \quad \omega_M(t) = \int_0^t \frac{\Sigma_M(u)}{u} du = \int_{\mu_1}^t \frac{\Sigma_M(u)}{u} du.$$

In particular, if $M \in \mathcal{LC}$, then ω_M vanishes on $[0, 1]$.

Finally, for any weight sequence M we have

$$(2.13) \quad \omega_{M^{lc}} = \omega_M.$$

2.5. Weighted spaces of entire functions defined in terms of ω_M . For any weight sequence M (in the sense of Definition 2.4) and for any parameter $c > 0$ we set

$$(2.14) \quad v_{M,c}(t) := \exp(-\omega_M(ct)), \quad v_M^c(t) := \exp(-c\omega_M(t)), \quad t \geq 0.$$

Note that each $v_{M,c}$ and v_M^c is a weight by the properties of the associated weight function. Then define

$$\begin{aligned} H_{v_{M,c}}^\infty(\mathbb{C}) &:= \{f \in H(\mathbb{C}) : \|f\|_{v_{M,c}} := \sup_{z \in \mathbb{C}} |f(z)| v_{M,c}(|z|) < +\infty\} \\ &= \{f \in H(\mathbb{C}) : \|f\|_{v_{M,c}} := \sup_{z \in \mathbb{C}} |f(z)| \exp(-\omega_M(c|z|)) < +\infty\}, \end{aligned}$$

and the class $H_{v_M^c}^\infty(\mathbb{C})$ is introduced accordingly. $M \in \mathcal{LC}$ has been the standard assumption in [22].

Remark 2.6. *We comment on the requirements for M :*

- (*) *If M is a weight sequence, then in view of (2.13) there is no difference between the weighted classes defined in terms of M or M^{lc} and thus one can assume w.l.o.g. that the defining sequence is log-convex.*
- (*) *Let M be a log-convex weight sequence. By Remark 2.5 there exists $N \in \mathcal{LC}$ satisfying (2.9) and so by definition*

$$\exists D \geq 0 \forall t \geq 0 : \omega_M(t) - D \leq \omega_N(t) \leq \omega_M(t) + D,$$

which yields as l.c.v.s.

$$\forall c > 0 : H_{v_{M,c}}^\infty(\mathbb{C}) = H_{v_{N,c}}^\infty(\mathbb{C}), \quad H_{v_M^c}^\infty(\mathbb{C}) = H_{v_N^c}^\infty(\mathbb{C}).$$

Remark 2.7. *We summarize more consequences for weight sequences M and N :*

- (i) *For $c = 1$ we write v_M instead of $v_{M,1}$ resp. v_M^1 . v_M is normalized if ω_M vanishes on $[0, 1]$, in particular this holds for any $M \in \mathcal{LC}$.*
- (ii) *One has that*

$$\forall d \geq c > 0 : H_{v_{M,c}}^\infty(\mathbb{C}) \subseteq H_{v_{M,d}}^\infty(\mathbb{C}), \quad H_{v_M^c}^\infty(\mathbb{C}) \subseteq H_{v_M^d}^\infty(\mathbb{C}),$$

with continuous inclusions.

- (iii) *For the weight systems in (2.4) and (2.5) we set (M is fixed)*

$$(2.15) \quad \underline{\mathcal{M}}_c := (v_{M,c})_{c \in \mathbb{N}_{>0}}, \quad \overline{\mathcal{M}}_c := (v_{M, \frac{1}{c}})_{c \in \mathbb{N}_{>0}}, \quad \underline{\mathcal{M}}^c := (v_M^c)_{c \in \mathbb{N}_{>0}}, \quad \overline{\mathcal{M}}^c := (v_M^{\frac{1}{c}})_{c \in \mathbb{N}_{>0}},$$

and the (LB)-spaces from (2.1) resp. the Fréchet-spaces from (2.2) are defined accordingly.

- (iv) *By Remark 2.3*

$$\forall d > c > 0 : \lim_{t \rightarrow +\infty} \frac{v_M^c(t)}{v_M^d(t)} = +\infty.$$

- (v) The following are equivalent by definition:
- (*) $\omega_M(t) = O(\omega_N(t))$, i.e. $\omega_N \preceq \omega_M$,
 - (*) for some $c \geq 1$ we get $v_M \preceq v_N^c$, i.e. $v_M \preceq^c v_N$.

We prove now a first characterization in terms of (1.4).

Proposition 2.8. *Let M and N be weight sequences and assume that N is log-convex. Then the following are equivalent:*

- (i) The sequences are satisfying (1.4), i.e.

$$\exists A \geq 1 \forall j \in \mathbb{N} : N_j \leq AM_j.$$

- (ii) The weights are satisfying

$$v_M \preceq v_N.$$

For (i) \Rightarrow (ii) also log-convexity for N is not required necessarily and recall that $v_M \preceq v_N$ implies $H_{v_M}^\infty(\mathbb{C}) \subseteq H_{v_N}^\infty(\mathbb{C})$ (with continuous inclusion).

Proof. (i) \Rightarrow (ii) By assumption and definition $\omega_M(t) \leq \omega_N(t) + \log(A)$ for all $t \geq 0$ follows and so (ii) is immediate.

(ii) \Rightarrow (i) By assumption there exists some $C \geq 1$ such that $\exp(-\omega_N(t)) = v_N(t) \leq Cv_M(t) = C \exp(-\omega_M(t))$ for all $t \geq 0$. Then (2.11) applied to both weights yields $N_j \leq CM_j^{1c} \leq CM_j$ for all $j \in \mathbb{N}$, i.e. (1.4) with $A = C$. \square

The next statement should be compared with Remark 2.3.

Lemma 2.9. *Let M be a log-convex weight sequence. Then*

$$(2.16) \quad \forall d > c > 0 : \lim_{t \rightarrow +\infty} \frac{v_{M,c}(t)}{v_{M,d}(t)} = +\infty.$$

Proof. Let $d > c > 0$, then by the definition of Σ_M in (2.10) we get

$$\omega_M(dt) - \omega_M(ct) = \int_{ct}^{dt} \frac{\Sigma_M(u)}{u} du \geq \Sigma_M(ct) \int_{ct}^{dt} \frac{1}{u} du = \Sigma_M(ct) \log(d/c) \rightarrow +\infty.$$

\square

The appearing parameter $c > 0$ in (2.14) in the dilatation-type structure is closely connected to and naturally appearing in the weight sequence approach.

Remark 2.10. *Let M be a given weight sequence.*

- (a) For any $c > 0$ we set $M^c := (c^j M_j)_{j \in \mathbb{N}}$ with corresponding quotient sequence $\mu_j^c = c\mu_j$ for $j \geq 1$ (and again $\mu_0^c := 1$). Then $1 = M_0^c$ and $\lim_{j \rightarrow +\infty} (M_j^c)^{1/j} = +\infty$, i.e. M^c is a weight sequence as well and M is log-convex if and only if some/each M^c is so.

We also have

$$(2.17) \quad \forall c > 0 \forall t \geq 0 : M^c \approx M, \quad \omega_{M^c}(t) = \omega_M\left(\frac{t}{c}\right),$$

which gives

$$(2.18) \quad \forall c > 0 \forall t \geq 0 : v_{M,c}(t) = v_{M^{\frac{1}{c}}}(t).$$

If $c < 1$, then even if M is normalized in general normalization is not clear for M^c .

(b) For arbitrary $c > 0$ the spaces $H_{v_{M,c}}^\infty(\mathbb{C})$ and $H_{v_M}^\infty(\mathbb{C})$ are isometrically isomorphic, written $H_{v_{M,c}}^\infty(\mathbb{C}) \cong H_{v_M}^\infty(\mathbb{C})$: This follows by taking into account the mappings $f(z) \mapsto f(zc)$ and $f(z) \mapsto f(z/c)$, see also [22, Cor. 4.5, Rem. 7.1].

Analogously, this comment holds true for an abstractly given weight v .

Let $f_k(z) := z^k$, then for all $k \in \mathbb{N}$ and any weight sequence M we have $f_k \in H_{\mathcal{M}_c}^\infty(\mathbb{C}), H_{\mathcal{M}^c}^\infty(\mathbb{C})$, because $|f_k(z)|v_{M,c}(|z|) = |z|^k v_M(c|z|) \rightarrow 0$ and $|f_k(z)|v_M^c(|z|) = |z|^k v_M^c(|z|) \rightarrow 0$ as $|z| \rightarrow +\infty$ for all $c > 0$. For this recall that v_M is rapidly decreasing. The same statement holds true if v_M is replaced by any abstractly given weight v : In fact $H_v^\infty(\mathbb{C})$ contains all polynomials if and only if v is rapidly decreasing, see [3, p. 3].

2.6. On the class of (log-)convex weights v . We recall some statements from [22, Sect. 6]. Let $v : [0, +\infty) \rightarrow (0, +\infty)$ be a normalized (radial) weight function, then set

$$(2.19) \quad \omega^v(t) := -\log(v(t)), \quad t \in [0, +\infty),$$

and we get (cf. [22, Lemma 6.1]): $\omega^v : [0, +\infty) \rightarrow [0, +\infty)$ is continuous, non-decreasing, $\lim_{t \rightarrow +\infty} \omega^v(t) = +\infty$ and $\omega^v(t) = 0$ for $t \in [0, 1]$ (normalization). Moreover, the fact that v is rapidly decreasing is equivalent to

$$(\omega_3) : \quad \log(t) = o(\omega^v(t)) \quad t \rightarrow +\infty,$$

and

$$(2.20) \quad t \mapsto \omega^v(e^t) (= -\log(v(e^t))) \quad \text{is convex on } \mathbb{R},$$

if and only if $\varphi_{\omega^v} : t \mapsto \omega^v(e^t)$ is convex. (ω_3) and (2.20) are standard assumptions in the theory of ultradifferentiable functions defined by so-called *Braun-Meise-Taylor weight functions* ω , see [8]. (ω_3) is named after [21] and there (2.20) is abbreviated with (ω_4) . Note that v has (2.20) if and only if v_c resp. v^c for some/each $c > 0$ does so and $\omega^{v^c}(t) = -\log(v_c(t)) = -\log(v(tc))$, whereas $\omega^{v^c}(t) = -\log(v^c(t)) = -c \log(v(t)) = -c\omega^v(t)$.

Conversely, we write that $\omega : [0, +\infty) \rightarrow [0, +\infty)$ has (ω_0) if ω is continuous, non-decreasing, $\omega(t) = 0$ for all $t \in [0, 1]$ (normalization) and $\lim_{t \rightarrow +\infty} \omega(t) = +\infty$. We put

$$\mathcal{W}_0 := \{\omega : [0, \infty) \rightarrow [0, \infty) : \omega \text{ has } (\omega_0), (\omega_3), (2.20)\},$$

and for any $\omega \in \mathcal{W}_0$ we set

$$(2.21) \quad v^\omega(t) := \exp(-\omega(t)).$$

Then v^ω is a normalized weight function satisfying (2.20). Summarizing, we have shown:

Lemma 2.11. *The set of normalized weight functions v satisfying (2.20) is isomorphic to \mathcal{W}_0 via the mappings*

$$v \mapsto \omega^v, \quad \omega \mapsto v^\omega.$$

In particular, Lemma 2.11 can be applied to $\omega \equiv \omega_M$ provided $M \in \mathcal{LC}$ since in this case $\omega_M \in \mathcal{W}_0$, see [16, Lemma 3.1] and the citations there. In this situation $v^{\omega_M} \equiv v_M$ and $\omega^{v_M} \equiv \omega_M$.

Definition 2.12. *We call a weight v convex if it satisfies (2.20).*

Note that in [3, p. 5, (4)] a weight satisfying (2.20) has been called *log-convex*; this has also been done in [10] and [11].

2.7. From weight functions to weight sequences. The following is also taken from [22, Sect. 6]. Let v be a normalized and convex weight and let ω^v be given by (2.19). In [22, (6.3)] we have introduced and studied the sequence $M^v = (M_j^v)_{j \in \mathbb{N}}$ defined by

$$(2.22) \quad M_j^v := \sup_{t>0} \frac{t^j}{\exp(\omega^v(t))} = \sup_{t>0} t^j v(t), \quad j \in \mathbb{N}.$$

Note that this expression has already been introduced on [1, p. 157].

We are also interested in the weight $t \mapsto \exp(-\omega_{M^v}(t)) (= v_{M^v}(t))$. This notation is suggested by (2.14) but confusing. Thus when explicitly using this function we denote the abstractly given weight with a different symbol than v , say w or u , e.g. $v_{M^u}(t) = \exp(-\omega_{M^u}(t))$.

When $M \equiv M^u$, then in (2.15) replace \mathcal{M} by \mathcal{M}^u .

We gather several properties, see [22, Prop. 6.2]:

- (i) $M^u \in \mathcal{LC}$,
- (ii) $\omega_{M^u} \sim \omega^u$, more precisely $\exists A \geq 1 \forall t \geq 0$:

$$(2.23) \quad \frac{1}{A} (v_{M^u}(t))^2 = \frac{1}{A} \exp(-2\omega_{M^u}(t)) \leq u(t) = \exp(-\omega^u(t)) \leq \exp(-\omega_{M^u}(t)) = v_{M^u}(t),$$

consequently

$$(2.24) \quad \exists A \geq 1 \forall c > 0 \forall t \geq 0: \quad \frac{1}{A} v_{M^u, c}^2(t) \leq u_c(t) \leq v_{M^u, c}(t), \quad \frac{1}{A} v_{M^u}^{2c}(t) \leq u^c(t) \leq v_{M^u}^c(t),$$

which implies

$$(2.25) \quad \forall c > 0: \quad H_{v_{M^u, c}}^\infty(\mathbb{C}) \subseteq H_{u, c}^\infty(\mathbb{C}) \subseteq H_{v_{M^u}^{2c}}^\infty(\mathbb{C}), \quad H_{v_{M^u}^c}^\infty(\mathbb{C}) \subseteq H_{u^c}^\infty(\mathbb{C}) \subseteq H_{v_{M^u}^{2c}}^\infty(\mathbb{C}),$$

with continuous inclusions. Recall that the second part in (2.24) precisely means $v_{M^u} \sim^c u$.

- (iii) If $u \equiv v_M$ with $M \in \mathcal{LC}$, and so $\omega^u \equiv \omega_M$, then $M^u \equiv M$. Moreover, for any $c > 0$ by definition $(M^u)^{\frac{1}{c}} \equiv (c^j M_j^u)_{j \in \mathbb{N}}$.

We comment on the consequences of the standard requirements for a given weight.

Remark 2.13. *A careful inspection of the proof of [22, Prop. 6.2] (and of the used citations [21, Thm. 4.0.3], [20, Lemma 5.7]) gives that convexity for u is in the whole approach only required in order to get the first estimate in (2.23), i.e. the second inclusions in (2.25).*

Crucially we remark that

$$\omega_{M^u} \sim \omega^u \not\Rightarrow v_{M^u} \sim u,$$

which is due to the appearance of the quadratic power in the first estimate in (2.23). This fact requires a technical condition studied in Section 2.8.

Remark 2.14. *If v is a function satisfying all properties to be a normalized weight except that v is rapidly decreasing, then (2.22) implies that $M_j^v = +\infty$ for all sufficiently large j and so the technique of associating a weight sequence is not available anymore.*

Next let us see that $\sup_{j \in \mathbb{N}_{>0}} (M^v)^{1/j} < +\infty$ if and only if $v(t) = 0$ for all t sufficiently large (i.e. v is violating strict positivity):

If the supremum is finite, then by (2.22) there exists $D \geq 1$ such that for all $t \geq 0$ and $j \in \mathbb{N}_{>0}$ we get $t^j v(t) \leq D^j \Leftrightarrow v(t) \leq (D/t)^j$. Thus $v(t) = 0$ for all $t > D$ follows.

Conversely, if there exists t_0 with $v(t) = 0$ for all $t \geq t_0$, then $M_j^v = \sup_{t>0} t^j v(t) = \max_{0 \leq t \leq t_0} t^j v(t) \leq v(0)t_0^j = t_0^j$ for all $j \in \mathbb{N}$ since v is non-decreasing and $v(0) = 1$.

Note that for weights v satisfying $v(t) = 0$ for all sufficiently large t we get that $H_v^\infty(\mathbb{C}) = H(\mathbb{C})$ (as sets!).

The second part of (2.25) immediately gives:

Proposition 2.15. *Let u be a normalized and convex weight function, M^u the associated weight sequence and consider the corresponding weight systems \underline{U}^c resp. \overline{U}^c (see (2.5)) and $\underline{\mathcal{M}}^{u^c}$ resp. $\overline{\mathcal{M}}^{u^c}$ (see (2.15)). Then as locally convex vector spaces we get*

$$H_{\underline{\mathcal{M}}^{u^c}}^\infty(\mathbb{C}) = H_{\underline{U}^c}^\infty(\mathbb{C}), \quad H_{\overline{\mathcal{M}}^{u^c}}^\infty(\mathbb{C}) = H_{\overline{U}^c}^\infty(\mathbb{C}).$$

If u is not convex, then in view of Remark 2.13 we only get the (continuous) inclusions \subseteq .

We close with the following important observation concerning the norm of the monomials $f_j(z) = z^j$, $j \in \mathbb{N}$. For a given weight v we set (see [3, p. 5]):

$$(2.26) \quad P_v(t) := \sup_{j \in \mathbb{N}} \frac{t^j}{\|f_j\|_v}, \quad t \geq 0.$$

Lemma 2.16. *Let u be a given normalized weight. Then*

$$\forall j \in \mathbb{N}: \quad \|f_j\|_u = M_j^u,$$

and

$$(2.27) \quad \forall t \geq 0: \quad P_u(t) = \exp(\omega_{M^u}(t)) = \frac{1}{v_{M^u}(t)} \leq \frac{1}{u(t)}.$$

If $u = v_M$ for a given log-convex weight sequence M , then

$$\forall j \in \mathbb{N}: \quad \|f_j\|_{v_M} = M_j, \quad \forall t \geq 0: \quad P_{v_M}(t) = \exp(\omega_M(t)) = \frac{1}{v_M(t)}.$$

The inequality in (2.27) has already been observed in [11, Sect. 2.1]. This follows immediately by definition and also by the second half in (2.23).

Proof. By definition and (2.22) we have for all j that

$$\|f_j\|_u = \sup_{z \in \mathbb{C}} |z^j| u(|z|) = \sup_{t \geq 0} t^j u(t) = M_j^u,$$

and so $P_u(t) = \sup_{j \in \mathbb{N}} \frac{t^j}{\|f_j\|_u} = \sup_{j \in \mathbb{N}} \frac{t^j}{M_j^u} = \exp(\omega_{M^u}(t)) = \frac{1}{v_{M^u}(t)}$ is immediate.

In view of (2.11) the case $u = v_M$ is then clear. Note that, if M is not log-convex, then we get the equality $\|f_j\|_{v_M} = M_j^{lc}$ but by Remark 2.6 log-convexity is not restricting. \square

2.8. Technical growth conditions on weight functions. From $\omega_{M^u} \sim \omega^u$ it does not follow that $u \sim v_{M^u}$ is valid, recall Remark 2.13. In order to get more information we recall the following property appearing in the theory of ultradifferentiable function classes:

$$(\omega_6): \quad \exists H \geq 1 \forall t \geq 0: \quad 2\omega(t) \leq \omega(Ht) + H.$$

This condition is named after [20] and [21]. For abstractly given weight functions ω (in the sense of Braun-Meise-Taylor) (ω_6) has appeared in [6] and for a log-convex weight sequence M the function ω_M has (ω_6) if and only if M has (mg), see [17, Prop. 3.6].

Lemma 2.17. *Let u be normalized and convex, then the following are equivalent:*

(a) u satisfies

$$(2.28) \quad \exists H \geq 1 \forall t \geq 0: \quad u_H(t) = u(Ht) \leq e^H u^2(t).$$

(b) The function ω^u (see (2.19)) satisfies (ω_6) .

(c) The associated weight function ω_{M^u} satisfies (ω_6) .

(d) The sequence M^u satisfies (mg).

Proof. (a) \Leftrightarrow (b) follows with the same H directly by definition (see (2.19)); (b) \Leftrightarrow (c) holds by $\omega_{M^u} \sim \omega^u$ (for which convexity is required!) and since (ω_6) is preserved under \sim ; (c) \Leftrightarrow (d) holds by [17, Prop. 3.6], see also (iii) in Remark 7.4. \square

Note:

(*) (2.28) holds if and only if there exist $H_1, H_2 \geq 1$ such that $u_{H_1}(t) \leq e^{H_2} u^2(t)$ for all $t \geq 0$ (take $H := \max\{H_1, H_2\}$).

(*) (2.28) holds for u if and only if for some/each u_c even with the same choice for H and it is clearly preserved under relation \sim .

(*) The choice $H > 1$ is required since for $0 < H \leq 1$ one has $u(t) \leq u(Ht) \leq e^H u^2(t)$ and so $1 \leq e^H u(t)$ for all $t \geq 0$, a contradiction.

Definition 2.18. A weight v is called to be of moderate growth if v satisfies (2.28).

Combining the first part of (2.25) and (2.28) we get:

Proposition 2.19. Let u be a normalized and convex weight function which is of moderate growth. Let M^u be the associated weight sequence and consider the corresponding weight systems \underline{U}_c resp. \overline{U}_c (see (2.4)) and \underline{M}_c^u resp. \overline{M}_c^u (see (2.15)). Then as locally convex vector spaces we get

$$H_{\underline{U}_c}^\infty(\mathbb{C}) = H_{\underline{M}_c^u}^\infty(\mathbb{C}), \quad H_{\overline{U}_c}^\infty(\mathbb{C}) = H_{\overline{M}_c^u}^\infty(\mathbb{C}).$$

Lemma 2.17 can be used to transfer (2.16) to the weight function setting.

Lemma 2.20. We have the following:

(i) Let u be normalized and convex. Assume that u is also of moderate growth, then

$$(2.29) \quad \exists H \geq 1 \forall c > 0 \forall d > Hc: \quad \lim_{t \rightarrow +\infty} \frac{u_c(t)}{u_d(t)} = +\infty.$$

Here H can be chosen to be the same constant as the one appearing in (2.28).

(ii) Let v and u be normalized weight functions satisfying $v \preceq_c u$. If either v or u is in addition convex and of moderate growth, then

$$(2.30) \quad \exists c \geq 1 \exists H \geq 1 \forall a > Hc: \quad u_a(t) = o(v(t)), \quad t \rightarrow +\infty.$$

Here c denotes the value from $v \preceq_c u$ and H can be taken as the constant appearing in (2.28) for the particular weight.

Proof. (i) Let $d > Hc$ with H the constant appearing in (2.28), then by combining (2.23) and (2.28) we get

$$\frac{u_c(t)}{u_d(t)} \geq \frac{1}{A} \frac{v_{M^u, c}(t)^2}{v_{M^u, d}(t)} \geq \frac{1}{Ae^H} \frac{v_{M^u, Hc}(t)}{v_{M^u, d}(t)},$$

hence (2.16) applied to $M \equiv M^u$ yields the conclusion.

(ii) Let $c > 0$ be the value from $v \preceq_c u$ and $H \geq 1$ the constant appearing in (2.28). If u has in addition (2.20) and (2.28), then write

$$\frac{u_a(t)}{v(t)} = \frac{u(at) u(ct)}{u(ct) v(t)}.$$

So, by (2.29) applied to u we get that $\lim_{t \rightarrow +\infty} \frac{u(at)}{v(t)} = 0$ for all $a > Hc$.

Similarly, if v has in addition (2.20) and (2.28), then write

$$\frac{u(t)}{v(t/a)} = \frac{u(t) v(t/c)}{v(t/c) v(t/a)},$$

and (2.29) applied to v yields $\lim_{t \rightarrow +\infty} \frac{u(at)}{v(t)} = 0$ for all a satisfying $c^{-1} > Ha^{-1} \Leftrightarrow a > Hc$. \square

For weight functions ω in the sense of Braun-Meise-Taylor also the following assumption is standard, see [8]:

$$(\omega_1) : \exists L \geq 1 \forall t \geq 0 : \omega(2t) \leq L(\omega(t) + 1).$$

This condition is named again after [20] and [21]. Recently (ω_1) for ω_M has been characterized in terms of M in [23, Thm. 3.1]: If $M \in \mathcal{LC}$ then (ω_1) holds true for ω_M if and only if

$$(2.31) \quad \exists L \in \mathbb{N}_{>0} : \liminf_{j \rightarrow +\infty} \frac{(M_{Lj})^{1/(Lj)}}{(M_j)^{1/j}} > 1.$$

By definition (2.19) it is immediate that ω^v has (ω_1) if and only if

$$(2.32) \quad \exists L \geq 1 \forall t \geq 0 : v^L(t) \leq e^L v_2(t).$$

Summarizing, we get the following characterization (analogously to Lemma 2.17):

Lemma 2.21. *Let u be a normalized and convex weight function. Then the following are equivalent:*

- (a) u satisfies (2.32).
- (b) ω^u has (ω_1) .
- (c) ω_{M^u} has (ω_1) .
- (d) M^u satisfies (2.31).

3. CHARACTERIZATION OF INCLUSIONS AND EQUALITIES OF WEIGHTED CLASSES

In this section we give a complete characterization of the inclusion (and equality) of weighted classes of entire functions in terms of the growth of the corresponding weights. We treat both dilatation- and exponential-type weight systems. But before we focus on (single) essential weights and show that in this case one has an equivalence in (1.3).

3.1. Characterization for essential weights. We recall the notions from [1, Def. 1.1, Sect. 1.B], see also [3, Sect. 3]. Let a weight v be given, then consider the function (radial growth condition) $w_v : \mathbb{C} \rightarrow (0, +\infty)$ given by

$$(3.1) \quad w_v := \frac{1}{v}$$

and set

$$B_{w_v}(\mathbb{C}) := \{f \in H(\mathbb{C}) : |f(z)| \leq w_v(|z|), \forall z \in \mathbb{C}\}.$$

The function $\widetilde{w}_v : \mathbb{C} \rightarrow [0, +\infty)$ associated with w_v is defined by

$$(3.2) \quad \widetilde{w}_v(z) := \sup\{|f(z)| : f \in B_{w_v}(\mathbb{C})\}, \quad z \in \mathbb{C},$$

and [1, Observation 1.5] gives that \widetilde{w}_v is again radial. Set

$$(3.3) \quad \widetilde{v} := \frac{1}{\widetilde{w}_v}.$$

If $v = v_M$, then we write \widetilde{v}_M . Similarly these notions can be introduced for arbitrary (open and connected) sets G .

Definition 3.1. *A weight v is called essential if $v \sim \widetilde{v}$ (e.g. see [3, Sect. 3]).*

The following result illustrates that the notion of essentiality and the weight sequence setting are very closely connected via involving the associated weight sequence.

Theorem 3.2. *We get the following:*

- (i) *Let M be a log-convex weight sequence. Then v_M is essential (for the weighted entire case) and in fact we even get $v_M = \widetilde{v}_M$.*
- (ii) *The following are equivalent:*
 - (a) *u is an essential weight.*
 - (b) *$u \sim v_{M^u}$ is valid.*

By recalling (2.17) and (2.18) we thus have that $v_{M,c}$ is essential and $\widetilde{v}_{M,c} = v_{M,c}$ for each $c > 0$. The difference between $u \sim v_{M^u}$ and the weaker relation (2.23) (due to the appearing quadratic power) illustrates that (log)-convexity is in general a weaker requirement than essentiality, see also Remark 3.5 for more explanations.

Proof. (i) For any weight u in [11, Lemma 1], by using a statement from [12], it has been shown that (see also [3, p.5]):

$$(3.4) \quad \forall t \geq 0: \quad P_u(t) \leq \frac{1}{\widetilde{u}(t)} \leq 6P_u(t).$$

For this recall the notation $w = \frac{1}{u}$, $\widetilde{w} = \frac{1}{\widetilde{u}}$ used in [11]. Therefore, by combining (3.4) with Lemma 2.16 applied to $u = v_M$ we get

$$\forall t \geq 0: \quad \frac{1}{v_M(t)} \leq \frac{1}{\widetilde{v}_M(t)} \leq \frac{6}{v_M(t)},$$

i.e. $v_M \sim \widetilde{v}_M$ is verified and so v_M is essential.

Moreover, for any growth condition w we get $0 \leq \widetilde{w} \leq w$, see [1, Prop. 1.2]. Thus by definitions (3.1), (3.3) and the first part in (3.4)

$$v_M = \frac{1}{w_{v_M}} \leq \frac{1}{\widetilde{w}_{v_M}} = \widetilde{v}_M \leq v_M$$

holds and we are done.

(ii)(a) \Rightarrow (b) Since u is essential the non-trivial estimate $\widetilde{u}(t) \leq Cu(t)$ holds true for some $C \geq 1$ and all $t \geq 0$. Then the second part in (3.4) and (2.27) yield

$$\forall t \geq 0: \quad \frac{1}{v_{M^u}(t)} = P_u(t) \leq \frac{1}{u(t)} \leq \frac{C}{\widetilde{u}(t)} \leq 6CP_u(t) = \frac{6C}{v_{M^u}(t)},$$

and so $v_{M^u} \sim u$ is verified.

(ii)(b) \Rightarrow (a) The assumption and the first part in (3.4) give

$$\exists C \geq 1 \forall t \geq 0: \quad \frac{1}{Cu(t)} \leq \frac{1}{v_{M^u}(t)} = P_u(t) \leq \frac{1}{\widetilde{u}(t)},$$

hence $\tilde{u}(t) \leq Cu(t)$. Since always $u(t) \leq \tilde{u}(t)$ (see [1, Prop. 1.2]) relation $u \sim \tilde{u}$ is verified and we are done. \square

The main result in this context is the following:

Theorem 3.3. *Let u and v be weights and assume that u is essential. Then the following are equivalent:*

- (i) *The weights are related by $u \preceq v$.*
- (ii) *We have $H_u^\infty(G) \subseteq H_v^\infty(G)$ (with continuous inclusion) for $G = \mathbb{D}$ and/or $G = \mathbb{C}$.*

When $G = \mathbb{D}$ then the relation $u \preceq v$ means $v(t) \leq Cu(t)$ for all $0 \leq t < 1$ and some $C \geq 1$.

Proof. It remains to prove (ii) \Rightarrow (i). For $G = \mathbb{C}$ we use [11, Thm. 2] (see also [3, Thm. 3]) and get that u is essential if and only if u is *approximable by the maximum of an analytic function modulus*, i.e.

$$(3.5) \quad \exists f_u \in H(\mathbb{C}) : \quad \frac{1}{u} \sim t \mapsto M(f_u, t),$$

with $M(f, t) := \sup\{|f(z)| : |z| = t\}$, see [11, Def. 3], [3, p. 3, p. 5]. Recall that $w = \frac{1}{u}$ denotes the weight used in the notation in [11]. (3.5) gives that $f_u \in H_u^\infty(\mathbb{C})$ and more precisely this relation ensures the existence of *optimal functions* $f_u \in H_u^\infty(\mathbb{C})$ admitting sharp estimates from below. Then (3.5) and the inclusion $H_u^\infty(\mathbb{C}) \subseteq H_v^\infty(\mathbb{C})$ (as sets) give

$$\exists C, D \geq 1 \forall t \geq 0 : \quad \frac{1}{u(t)} \leq CM(f_u, t) \leq CD \frac{1}{v(t)},$$

i.e. $v(t) = O(u(t))$ and so $u \preceq v$ is verified.

When $G = \mathbb{D}$, then by [10, Thm. 1.2], see also [11, Thm. 1, Rem. 1, Def. 1, Thm. 5] and [3, Thm. 2] again u is essential if and only if (3.5) is valid. The rest follows as before.

Note that for this case in order to apply the cited results we do not require necessarily that u is rapidly decreasing. \square

Finally, when combining Theorems 3.2 and 3.3 with Proposition 2.8 we arrive at the following characterization:

Theorem 3.4. *Let M and N be log-convex weight sequences. Then both v_M and v_N are essential and the following are equivalent:*

- (i) *The sequences are satisfying (1.4), i.e.*

$$\exists A \geq 1 \forall j \in \mathbb{N} : \quad N_j \leq AM_j.$$

- (ii) *The weights are satisfying*

$$v_M \preceq v_N.$$

- (iii) *We have*

$$H_{v_M}^\infty(\mathbb{C}) \subseteq H_{v_N}^\infty(\mathbb{C}),$$

with continuous inclusion.

Thus (2.9) if and only if $v_M \sim v_N$ if and only if $H_{v_M}^\infty(\mathbb{C}) = H_{v_N}^\infty(\mathbb{C})$ (as Banach spaces).

Remark 3.5. *We comment on the known relation between essentiality and convexity and its consequences.*

(A) Let u and r be given weights. Of course, Theorem 3.4 applies to the associated sequences M^u and M^r and the proof of Theorem 3.2 yields $v_{M^u} \sim \tilde{u}$ and $v_{M^r} \sim \tilde{r}$. Moreover the first part in (2.25), for which even convexity is not required, gives the following: Assume that the inclusion $H_u^\infty(\mathbb{C}) \subseteq H_r^\infty(\mathbb{C})$ is valid, then by Theorem 3.2 the weight v_{M^u} is essential and hence $v_{M^u} \preceq r$ follows as in the proof of Theorem 3.3.

However, from this the desired relation $u \preceq r$ does not follow automatically which is due to the quadratic power appearing in (2.23). In view of (ii) in Theorem 3.2 we can conclude provided u is essential.

(B) By [10], [11] for the weighted entire case each essential weight u is equivalent to a convex one. Since v_{M^u} is convex, (ii) in Theorem 3.2 proves this implication and provides even more precise information.

For $G = \mathbb{D}$ one even has that u is essential, equivalently (3.5) holds true, if and only if it is equivalent to a convex weight.

(C) In the discussion after Thm. 1 in [11] resp. after Thm. 2 in [3] it is explained that if u is rapidly decreasing, in particular for weight functions in our setting, in general the characterization [3, Thm. 2] fails for $G = \mathbb{C}$. More precisely, there exist (convex!) weights u on \mathbb{C} such that (3.5) is violated. In particular, by [11, Thm. 2], [3, Thm. 3], such weights cannot be essential and by (ii) in Theorem 3.2 they cannot be equivalent to v_{M^u} . In any case, for non-essential weights u defined on \mathbb{C} property (3.5) fails.

(D) Hence by the previous comments we see that for the weighted entire case when dealing with general convex weights the proof of \Leftarrow in (1.3) does not follow from (the proof of) Theorem 3.3 since (3.5) fails in general. On the other hand convexity of u is sufficient for our techniques (for involving the sequence M^u , see Theorem 3.30). Note that in view of Remark 2.14 the assumption that u is rapidly decreasing should be considered as standard when involving M^v .

(E) In this context also note that (3.5) implies (3.40) (with $\phi = -\log \circ u$ there), see Remark 3.31 for more explanations.

(F) In [1, Prop. 3.1] sufficient conditions on (log)-convex weights v are given to ensure essentiality.

We close this section by showing a partial converse to (B).

Theorem 3.6. *Let u be a normalized and convex weight function.*

(i) *We get*

$$(3.6) \quad \exists B \geq 1 \forall t \geq 0: \quad u(t) \leq \tilde{u}(t) \leq Bu^{1/2}(t),$$

i.e. $\tilde{u} \preceq u \preceq \tilde{u}$. Thus $u \sim \tilde{u}$ holds true.

(ii) *If in addition u is of moderate growth, then*

$$(3.7) \quad \exists B, H \geq 1 \forall t \geq 0: \quad u(t) \leq \tilde{u}(t) \leq Bu_{H^{-1}}(t) = Bu(t/H),$$

i.e. $\tilde{u} \preceq u \preceq \tilde{u}$. Thus $u \sim \tilde{u}$ holds true.

Proof. Let us write w_u and $w_{M,c} := w_{v_{M,c}}$ for the corresponding growth conditions, see (3.1). We will also use the following immediate properties, see e.g. [1, Prop. 1.2]: For any growth condition w we get $0 \leq \tilde{w} \leq w$, write (\star) for this, and given weights u, v with $u \leq v \Leftrightarrow w_v \leq w_u$, then $\tilde{w}_v \leq \tilde{w}_u$ (written as relation $(\star\star)$).

(i) First, (2.24) applied to the weight u transfers into

$$(3.8) \quad \exists A \geq 1 \forall c > 0 \forall t \geq 0 : w_{M^u, c}(t) \leq w_{u_c}(t) \leq Aw_{M^u, c}(t)^2.$$

Take $c = 1$ and from this we get

$$(3.9) \quad \exists A \geq 1 \forall t \geq 0 : \frac{1}{\sqrt{A}} \sqrt{w_u(t)} \leq w_{M^u}(t) = \widetilde{w_{M^u}}(t) \leq \widetilde{w_u}(t).$$

More precisely, the first estimate holds by the second half in (3.8), the second estimate since v_{M^u} is essential and coincides with $\widetilde{v_{M^u}}$ by Theorem 3.2. The third one holds by the first estimate in (3.8) and $(\star\star)$.

So (3.9) is verified and thus the second half of (3.6) follows by (3.1), (3.3) (the first half is precisely (\star)).

(ii) In view of (3.1) and since for any weight u and $c > 0$ we get

$$(3.10) \quad w_{u_c}(t) = \frac{1}{u_c(t)} = \frac{1}{u(ct)} = w_u(ct) = (w_u)_c(t),$$

a weight u is of moderate growth if and only if the corresponding growth condition satisfies (with the same H)

$$(3.11) \quad \exists H \geq 1 \forall t \geq 0 : w_u(t)^2 \leq e^H w_{u_H}(t) = e^H w_u(Ht).$$

Next let us verify

$$(3.12) \quad \exists A \geq 1 \forall c > 0 \forall t \geq 0 : w_{u_c}(t) \leq Aw_{M^u, c}(t)^2 = A\widetilde{w_{M^u, c}}(t)^2 \leq A\widetilde{w_{u_c}}(t)^2.$$

The first estimate holds by (3.8) (second part), the second one since each $v_{M^u, c}$ is essential by Theorem 3.2 and finally the third estimate in (3.12) holds by combining the first part of (3.8) with $(\star\star)$.

Now we continue (3.12) (with $c = H^{-1}$) and prove

$$(3.13) \quad A\widetilde{w_{u_{H^{-1}}}}(t)^2 \leq Ae^H \widetilde{w_u}(t).$$

By (3.11) one has $e^H w_u(t) \geq w_{u_{H^{-1}}}(t)^2$ and then note that for any $f \in B_w(\mathbb{C})$ and $c \in \mathbb{N}_{>0}$ we have $f^c \in B_{w^c}(\mathbb{C})$. So

$$\widetilde{w^c}(z) = \sup\{|f(z)|^c : f \in B_w(\mathbb{C})\} = \sup\{|f^c(z)| : f \in B_w(\mathbb{C})\} \leq \sup\{|g(z)| : g \in B_{w^c}(\mathbb{C})\} = \widetilde{w^c}(z)$$

follows and by combining both estimates (take $c = 2$) with $(\star\star)$ and the fact that $\widetilde{Cw} = C\widetilde{w}$ for any $C > 0$ and any growth condition w (see again [1, Prop. 1.2]) the estimate (3.13) holds.

Summarizing, by combining (3.12) and (3.13) we have verified

$$\exists A, H \geq 1 \forall t \geq 0 : w_{u_{H^{-1}}}(t) \leq Ae^H \widetilde{w_u}(t),$$

and so by taking into account (3.1), (3.3) we get the second half of (3.7) with $B := Ae^H$ and H being the constant appearing in (2.28) for u . \square

(*) For the sake of completeness let us mention that for any growth condition w and any $c > 0$ one gets $f \in B_{w^c}(\mathbb{C})$ if and only if $f_c \in B_w(\mathbb{C})$ with $f_c(z) := f(z/c)$. Thus for any $z \in \mathbb{C}$ and $c > 0$ we have

$$\widetilde{w^c}(z) = \sup\{|f(z)| : f \in B_{w^c}(\mathbb{C})\} = \sup\{|f_c(cz)| : f_c \in B_w(\mathbb{C})\} = \widetilde{w}(cz) = \widetilde{w^c}(z).$$

(*) The proof shows that in (3.7) we have to take H to be the constant appearing in (2.28) for u .

(*) Thus, since $H > 1$, from the previous result we cannot conclude that u is essential. In view of Proposition 2.2 the corresponding dilatation- resp. exponential-type class defined in terms of u and \tilde{u} coincide as l.c.v.s. when having (i) resp. (ii). However, as stated on [3, p. 4] even $H_u^\infty(\mathbb{C}) = H_{\tilde{u}}^\infty(\mathbb{C})$ coincide isometrically (which holds for general weights).

But note that, in order to ensure the existence of optimal functions satisfying (3.5), essentiality for u is unavoidable in any case and the shown relations are too weak.

(*) Summarizing, by (ii) in Theorem 3.2, (B) in Remark 3.5 and Theorem 3.6 we get in the weighted entire case: A weight u is essential if and only if it is equivalent to v_{M^u} , so u is equivalent to a convex weight. If u is convex, then $u \sim_\epsilon \tilde{u}$ and if in addition u is of moderate growth, then also $u \sim_\epsilon \tilde{u}$.

3.2. The dilatation-type weight sequence setting. We start with the following:

Lemma 3.7. *Let M and N be weight sequences and assume that N is log-convex. Consider the following assertions:*

(i) *We have (1.4), i.e.*

$$\exists A \geq 1 \forall j \in \mathbb{N}: N_j \leq AM_j.$$

(ii) *The weights are satisfying*

$$(3.14) \quad \forall c > 0: v_{M,c} \preceq_c v_{N,c},$$

which implies $v_M \preceq_c v_N$.

(iii) *We have that*

$$(3.15) \quad \forall c > 0: H_{v_{M,c}}^\infty(\mathbb{C}) \subseteq H_{v_{N,c}}^\infty(\mathbb{C}),$$

with continuous inclusion.

Then (i) \Leftrightarrow (ii) \Rightarrow (iii) holds true. Alternatively, in (3.14) and (3.15) we can assume $\exists c > 0$ instead and (iii) implies both $H_{\underline{M}_c}^\infty(\mathbb{C}) \subseteq H_{\underline{N}_c}^\infty(\mathbb{C})$ and $H_{\overline{M}_c}^\infty(\mathbb{C}) \subseteq H_{\overline{N}_c}^\infty(\mathbb{C})$ with continuous inclusions.

If log-convexity for N fails, then only (i) \Rightarrow (ii) \Rightarrow (iii) is valid.

Proof. (i) \Leftrightarrow (ii) follows from resp. as in Proposition 2.8 and (ii) \Rightarrow (iii) is clear by definition (1.1). \square

In the next result we replace (1.4) by the weaker requirement $N \preceq M$.

Proposition 3.8. *Let M and N be weight sequences. Consider the following assertions:*

(i) $N \preceq M$,

(ii) $v_M \preceq_c v_N$,

(iii)

$$H_{\underline{M}_c}^\infty(\mathbb{C}) \subseteq H_{\underline{N}_c}^\infty(\mathbb{C}), \quad H_{\overline{M}_c}^\infty(\mathbb{C}) \subseteq H_{\overline{N}_c}^\infty(\mathbb{C}),$$

with continuous inclusions.

Then (i) \Rightarrow (ii) \Rightarrow (iii) is valid.

Consequently, $M \approx N$ implies $v_M \sim_\epsilon v_N$ and this relation yields

$$H_{\underline{M}_c}^\infty(\mathbb{C}) = H_{\underline{N}_c}^\infty(\mathbb{C}), \quad H_{\overline{M}_c}^\infty(\mathbb{C}) = H_{\overline{N}_c}^\infty(\mathbb{C}),$$

as l.c.v.s.

Proof. (i) \Rightarrow (ii) By $N \preceq M$ there exist $C, h \geq 1$ such that $N_j \leq Ch^j M_j$ for all $j \in \mathbb{N}$ and by definition of associated weight functions we get

$$(3.16) \quad \exists C, h \geq 1 \forall t \geq 0 : \omega_M(t) \leq \omega_N(ht) + \log(C),$$

hence in view of (2.14)

$$(3.17) \quad \exists h \geq 1 \forall c > 0 : v_{N,hc}(t) = O(v_{M,c}(t)), \quad t \rightarrow +\infty.$$

This is precisely $v_M \preceq_c v_N$.

(ii) \Rightarrow (iii) This is clear by definition (recall Proposition 2.2). \square

Now let us see to prove a complete characterization. In order to do so, in the inductive structure, the idea is to use an optimal function. For the weight sequence approach by the previous section we can use $f_{v_{M,c}}$ from (3.5). We present here a different, weaker but sufficient construction involving a dilatation parameter. Let M be a weight sequence and $c > 0$, then consider the entire function

$$(3.18) \quad \theta_{M,c}(z) := \sum_{j \geq 0} \frac{1}{2^j M_j} (cz)^j, \quad z \in \mathbb{C}.$$

The definition of this function is inspired by the approach in [9]; more precisely compare $\theta_{M,c}$ with the function F in [9], see also Remark 3.31 for more explanations.

Lemma 3.9. *Let M be a weight sequence. Then*

$$\forall c > 0 : \theta_{M,c} \in H_{v_{M,c}}^\infty(\mathbb{C}) \subseteq H_{\underline{M}_c}^\infty(\mathbb{C}),$$

and

$$(3.19) \quad \forall c > 0 \forall t \geq 0 : \exp(\omega_M(ct/2)) \leq |\theta_{M,c}(t)| = \theta_{M,c}(t).$$

Proof. For any $z \in \mathbb{C}$ we estimate by

$$|\theta_{M,c}(z)| \leq \sum_{j \geq 0} \frac{1}{2^j M_j} (c|z|)^j \leq \sup_{k \in \mathbb{N}} \frac{(c|z|)^k}{M_k} \sum_{j \geq 0} \frac{1}{2^j} = 2 \exp(\omega_M(c|z|)),$$

showing $\theta_{M,c} \in H_{v_{M,c}}^\infty(\mathbb{C})$. If $t \geq 0$, then $\theta_{M,c}(t) \in \mathbb{R}_{\geq 0}$ and we get

$$\theta_{M,c}(t) = \sum_{j \geq 0} \frac{1}{2^j M_j} (ct)^j \geq \sup_{k \in \mathbb{N}} \frac{(ct)^k}{2^k M_k} = \exp(\omega_M(ct/2)),$$

since each summand is non-negative. \square

Remark 3.10. *The difference between the estimate (3.19) for $\theta_{M,c}$ and $f_{v_{M,c}}$ in (3.5) is that $|\theta_{M,c}| \sim \frac{1}{v_{M,c}}$ is not clear in general since an additional dilatation parameter $\frac{1}{2}$ appears on the left-hand side in (3.19). However, it turns out that this fact is not disturbing for dilatation-type systems. On the other hand in the previous Lemma formally log-convexity for M is not required but in view of Remark 2.6 this difference is negligible.*

Let M be a log-convex weight sequence. Any function satisfying the estimate from (3.19) for some arbitrary but fixed parameter $c > 0$ cannot be contained in the space $H_{\underline{M}_c}^\infty(\mathbb{C})$: Assume that there exists $\psi \in H_{\underline{M}_c}^\infty(\mathbb{C})$ satisfying (3.19) for some $c > 0$, then

$$\forall d > 0 \exists D \geq 1 \forall t \geq 0 : \exp(\omega_M(ct/2)) \leq |\psi(t)| \leq D \exp(\omega_M(dt)),$$

which implies $\omega_M(ct/2) - \omega_M(dt) \leq \log(D)$. But this is contradicting (2.16) as $t \rightarrow +\infty$ when $d < c/2$.

In particular this comment also applies to $f_{v_{M,c}}$ from (3.5) and hence for projective structures a different technique is required.

Remark 3.11. Similarly, for any $c \in \mathbb{N}_{>0}$ we can put

$$(3.20) \quad \theta_M^c(z) := \sum_{j \geq 0} \frac{1}{2^j (M_j)^c} z^{cj}, \quad z \in \mathbb{C},$$

and show analogously as before that $\theta_M^c \in H_{v_M^c}^\infty(\mathbb{C}) \subseteq H_{\mathcal{M}^c}^\infty(\mathbb{C})$ and

$$(3.21) \quad \forall c \in \mathbb{N}_{>0} \forall t \geq 0: \quad \exp(c\omega_M(t/2^{1/c})) \leq |\theta_M^c(t)| = \theta_M^c(t).$$

However, it turns out that for exponential-type structures this construction is less optimal than applying the functions from (3.5) since in (3.21) the appearing dilatation parameter $\frac{1}{2^{1/c}}$ requires additional technical assumptions on the weights, see Section 3.3 for more details.

Using $\theta_{M,c}$ we obtain the following result:

Proposition 3.12. Let M, N be weight sequences. Assume that N is log-convex and that (as sets)

$$(3.22) \quad \exists c > 0 \exists d > 0: \quad H_{v_{M,c}}^\infty(\mathbb{C}) \subseteq H_{v_{N,d}}^\infty(\mathbb{C}).$$

Then $N \preceq M$ follows.

Alternatively, in order to conclude we can replace in (3.22) one or both \exists by \forall (giving stronger assumptions).

For the proof only log-convexity for N is required but indispensable (by using (2.11)); otherwise we only get $N^{\text{lc}} \preceq M^{\text{lc}} (\leq M)$, i.e. the desired relation for the log-convex minorants. Note also that for the proof in (3.22) we only require the inclusion as sets.

Proof. Let the parameters c and d be given. The inclusion $H_{v_{M,c}}^\infty(\mathbb{C}) \subseteq H_{v_{N,d}}^\infty(\mathbb{C})$ and Lemma 3.9 yield $\theta_{M,c} \in H_{v_{N,d}}^\infty(\mathbb{C})$ and so

$$\exists C \geq 1 \forall t \geq 0: \quad \exp(\omega_M(ct/2)) \leq \theta_{M,c}(t) = |\theta_{M,c}(t)| \leq C \exp(\omega_N(dt)).$$

Hence, by applying (2.11) we obtain for all $j \in \mathbb{N}$ that

$$\begin{aligned} M_j &\geq M_j^{\text{lc}} = \sup_{t \geq 0} \frac{t^j}{\exp(\omega_M(t))} = \sup_{s \geq 0} \frac{(cs/2)^j}{\exp(\omega_M(cs/2))} \geq \frac{(c/2)^j}{C} \sup_{s \geq 0} \frac{s^j}{\exp(\omega_N(ds))} \\ &= \frac{(c/(2d))^j}{C} \sup_{u \geq 0} \frac{u^j}{\exp(\omega_N(u))} = \frac{(c/(2d))^j}{C} N_j, \end{aligned}$$

so

$$\exists C \geq 1 \forall j \in \mathbb{N}: \quad N_j \leq C \left(\frac{2d}{c} \right)^j M_j,$$

i.e. $N \preceq M$ is verified.

A different technique is to apply Theorem 3.4 to $v_{M^{\frac{1}{c}}}$ and $v_{N^{\frac{1}{d}}}$ (see (2.18)) and get $\frac{1}{d^j} N_j = N_j^{\frac{1}{d}} \leq A M_j^{\frac{1}{c}} = A \frac{1}{c^j} M_j$. Thus the same estimate as before is obtained apart from the fact that the constant 2 disappears. But this does not effect the conclusion (in general $d > c$). Note that in order to apply Theorem 3.4 formally we require that *both* sequences are log-convex. \square

For the characterization of $H_{\mathcal{M}^c}^\infty(\mathbb{C}) \subseteq H_{\mathcal{N}^c}^\infty(\mathbb{C})$ by Remark 3.10 we have to come up with a different technique; we use the family of monomials $(f_j)_{j \in \mathbb{N}}$.

Proposition 3.13. *Let M, N be weight sequences. Assume that N is log-convex and (with continuous inclusion)*

$$(3.23) \quad H_{\mathcal{M}_c}^\infty(\mathbb{C}) \subseteq H_{\mathcal{N}_c}^\infty(\mathbb{C}).$$

Then $N \preceq M$ holds true.

Proof. Both spaces $H_{\mathcal{M}_c}^\infty(\mathbb{C})$ and $H_{\mathcal{N}_c}^\infty(\mathbb{C})$ are Fréchet and by the closed graph theorem the inclusion in (3.23) is continuous since convergence in $H_{\mathcal{M}_c}^\infty(\mathbb{C})$ implies point-wise convergence, see the proof of [20, Prop. 4.6 (1)] and more generally [14, Prop. 4.5, Rem. 4.6]. This argument is analogously used in all forthcoming results dealing with non-decreasing sequences of weights (projective structures). So one has

$$\forall d > 0 \exists c > 0 \exists C \geq 1 \forall f \in H_{\mathcal{M}_c}^\infty(\mathbb{C}) : \|f\|_{v_{N,d}} \leq C \|f\|_{v_{M,c}}.$$

We apply this estimate to the family of monomials $(f_k)_{k \in \mathbb{N}}$ and get

$$\forall d > 0 \exists c > 0 \exists C \geq 1 \forall k \in \mathbb{N} \forall z \in \mathbb{C} : \frac{|z|^k}{\exp(\omega_N(d|z|))} \leq C \frac{|z|^k}{\exp(\omega_M(c|z|))}.$$

Thus, by using again (2.11), we obtain for all $j \in \mathbb{N}$ that

$$\begin{aligned} N_j &= \sup_{t \geq 0} \frac{t^j}{\exp(\omega_N(t))} = \sup_{s \geq 0} \frac{(ds)^j}{\exp(\omega_N(ds))} \leq Cd^j \sup_{s \geq 0} \frac{s^j}{\exp(\omega_M(cs))} \\ &= C \left(\frac{d}{c}\right)^j \sup_{u \geq 0} \frac{u^j}{\exp(\omega_M(u))} = C \left(\frac{d}{c}\right)^j M_j^{\text{lc}} \leq C \left(\frac{d}{c}\right)^j M_j, \end{aligned}$$

i.e. $N \preceq M$ is verified. \square

Combining now Propositions 3.8, 3.12 and 3.13 we arrive at the following characterization.

Theorem 3.14. *Let M, N be weight sequences and assume that N is log-convex. Then the following are equivalent:*

- (a) *The sequences are satisfying $N \preceq M$.*
- (b) *The weights v_M and v_N are satisfying $v_M \preceq_c v_N$.*
- (c) *We have*

$$H_{\mathcal{M}_c}^\infty(\mathbb{C}) \subseteq H_{\mathcal{N}_c}^\infty(\mathbb{C}),$$

with continuous inclusion.

- (d) *We have*

$$H_{\mathcal{M}_c}^\infty(\mathbb{C}) \subseteq H_{\mathcal{N}_c}^\infty(\mathbb{C}),$$

with continuous inclusion.

Thus for log-convex weight sequences M and N we get the following equivalences:

- (*) $M \approx N$,
- (*) $v_M \sim_c v_N$,
- (*) as l.c.v.s.

$$H_{\mathcal{M}_c}^\infty(\mathbb{C}) = H_{\mathcal{N}_c}^\infty(\mathbb{C}),$$

- (*) as l.c.v.s.

$$H_{\mathcal{M}_c}^\infty(\mathbb{C}) = H_{\mathcal{N}_c}^\infty(\mathbb{C}).$$

On the other hand, when assuming instead of (3.23) the (continuous) inclusion $H_{v_{M,c}}^\infty(\mathbb{C}) \subseteq H_{v_{N,c}}^\infty(\mathbb{C})$, where $c > 0$ is arbitrary, and following then the proof in Proposition 3.13 with $c = d$, then we get (iii) \Rightarrow (i) in Lemma 3.7. Thus the following characterization is shown:

Corollary 3.15. *Let M, N be weight sequences and assume that N is log-convex. Then the following are equivalent:*

(a) *We have (1.4), i.e.*

$$\exists A \geq 1 \forall j \in \mathbb{N}: N_j \leq AM_j.$$

(b) *The weights are satisfying*

$$\forall c > 0: v_{M,c} \preceq v_{N,c}.$$

(c) *We have that*

$$\forall c > 0: H_{v_{M,c}}^\infty(\mathbb{C}) \subseteq H_{v_{N,c}}^\infty(\mathbb{C}),$$

with continuous inclusion.

Alternatively, we can replace in (b), (c) also \forall by \exists . Thus, for log-convex sequences M and N the following are equivalent:

(*) (2.9) holds,

(*) $v_{M,c} \sim v_{N,c}$ for some/any $c > 0$,

(*) $H_{v_{M,c}}^\infty(\mathbb{C}) = H_{v_{N,c}}^\infty(\mathbb{C})$ (as Banach spaces) for some/any $c > 0$.

This gives an independent proof of the equivalences listed in Theorem 3.4 (involving a dilatation parameter $c > 0$, see (2.18)); indeed Corollary 3.15 is formally slightly more general since we only require log-convexity for N whereas the techniques in the previous section require log-convexity for both weights (see Theorem 3.2, Lemma 2.16). In view of Remark 2.6 this difference is negligible.

3.3. The exponential-type weight sequence setting. In this setting the relation $N \preceq M$ is, in general, not the proper condition for the characterization of the inclusions. It turns out that we have to deal with auxiliary sequences. First let us gather some immediate and known facts:

(*) Since $\omega_M(t) \geq 0$ for all $t \geq 0$, we have $v_M \leq 1$ and so $v_M^d \leq v_M^c$ for all $0 < c \leq d$.

(*) Let M, N be weight sequences satisfying $N \leq AM$ (i.e. (1.4)). Then

$$\forall c > 0 \forall t \geq 0: c\omega_M(t) \leq c\omega_N(t) + c \log(A) \Leftrightarrow v_N^c(t) \leq A^c v_M^c(t),$$

hence

$$\forall c > 0: H_{v_M^c}^\infty(\mathbb{C}) \subseteq H_{v_N^c}^\infty(\mathbb{C}),$$

with continuous inclusion.

(*) However, the natural relation in this setting is expected to be \preceq^c since $v_M \preceq^c v_N$ implies

$$H_{\underline{M}^c}^\infty(\mathbb{C}) \subseteq H_{\underline{N}^c}^\infty(\mathbb{C}), \quad H_{\overline{M}^c}^\infty(\mathbb{C}) \subseteq H_{\overline{N}^c}^\infty(\mathbb{C}),$$

with continuous inclusions, see Proposition 2.2.

(*) By (v) in Remark 2.7 we know that $\omega_M(t) = O(\omega_N(t))$ if and only if $v_M \preceq^c v_N$.

(*) In [14, Lemma 6.5] we have shown: Let $M, N \in \mathcal{LC}$, then $\omega_M(t) = O(\omega_N(t))$ if and only if

$$(3.24) \quad \exists c \in \mathbb{N}_{>0} \exists A \geq 1 \forall j \in \mathbb{N}: N_j \leq A(M_{cj})^{1/c}.$$

We summarize:

Proposition 3.16. *Let $M, N \in \mathcal{LC}$ be given. Consider the following assertions:*

(i) *M and N are related by (3.24).*

(ii) *The weights are satisfying $v_M \preceq^c v_N$.*

(iii) *We have*

$$H_{\underline{M}^c}^\infty(\mathbb{C}) \subseteq H_{\underline{N}^c}^\infty(\mathbb{C}), \quad H_{\overline{M}^c}^\infty(\mathbb{C}) \subseteq H_{\overline{N}^c}^\infty(\mathbb{C}),$$

with continuous inclusions.

Then (i) \Leftrightarrow (ii) \Rightarrow (iii) holds true.

Moreover, we get the following characterization (even under formally more general assumptions on M and N):

Theorem 3.17. *Let M and N be weight sequences and assume that N is log-convex. Then the following are equivalent:*

(a) *The sequences are satisfying (1.4), so*

$$\exists A \geq 1 \forall j \in \mathbb{N}: N_j \leq AM_j.$$

(b) *The weights are satisfying*

$$\forall c > 0: v_M^c \preceq v_N^c.$$

(c) *We have*

$$\forall c > 0: H_{v_M^c}^\infty(\mathbb{C}) \subseteq H_{v_N^c}^\infty(\mathbb{C}),$$

with continuous inclusion.

Proof. (a) \Rightarrow (b) \Rightarrow (c) follows from the previous comments and (c) \Rightarrow (a) follows by applying (c) \Rightarrow (a) in Corollary 3.15 to $c = 1$ (for which log-convexity of N is required). \square

In order to proceed we have to recall and introduce some notation: First, for given $M \in \mathbb{R}_{>0}^{\mathbb{N}}$ and $c \in \mathbb{N}_{>0}$ let us set (see [23, (2.5), (2.6)] and [24, Sect.3])

$$(3.25) \quad \widetilde{M}_j^c := (M_{cj})^{1/c}.$$

Hence $\widetilde{M}^1 \equiv M$ is clear. If $M_0 = 1$ and M is log-convex, then each \widetilde{M}^c is log-convex as well, $\widetilde{M}_0^c = 1$ and $\widetilde{M}^c \geq M$. If $M \in \mathcal{LC}$, then $\widetilde{M}^c \in \mathcal{LC}$ (for some/any $c \in \mathbb{N}_{>0}$). Recall also the technical estimate [14, Lemma 6.5, (6.7)]:

Lemma 3.18. *Let $M \in \mathcal{LC}$ be given, then*

(3.26)

$$\forall c \in \mathbb{N}_{>0} \exists D \geq 1 \forall t \geq 0: c\omega_{\widetilde{M}^c}(t) \leq \omega_M(t) \leq 2c\omega_{\widetilde{M}^c}(t) + Dc \Leftrightarrow \frac{1}{e^{Dc}}v_{\widetilde{M}^c}^{2c}(t) \leq v_M(t) \leq v_{\widetilde{M}^c}^c(t).$$

Moreover, for any given $M \in \mathcal{LC}$ and $c \in \mathbb{N}_{>0}$ we introduce

$$(3.27) \quad \underline{M}_j^c := \sup_{t \geq 0} \frac{t^j}{\exp(c\omega_M(t))} = \exp(\varphi_{c\omega_M}^*(j)),$$

and the second equality holds as follows (analogously as in the proof of [14, Lemma 6.5] for \widetilde{M}^c): Recall that by $M \in \mathcal{LC}$ we get $\omega_M(t) = 0$ for $t \in [0, 1]$ (i.e. normalization) and so

$$\begin{aligned} \underline{M}_j^c &:= \sup_{t \geq 0} \frac{t^j}{\exp(c\omega_M(t))} = \sup_{t \geq 1} \frac{t^j}{\exp(c\omega_M(t))} = \exp(\sup_{t \geq 1} j \log(t) - c\omega_M(t)) \\ &= \exp(\sup_{s \geq 0} js - c\omega_M(e^s)) =: \exp(\varphi_{c\omega_M}^*(j)), \end{aligned}$$

where φ_ω^* denotes the so-called Young-conjugate of $\varphi_\omega : t \mapsto \omega(e^t)$. Since $\omega_M \in \mathcal{W}_0$ we have $c\omega_M \in \mathcal{W}_0$. By this fact and (3.27), in particular, we know that $\underline{M}^c \in \mathcal{LC}$. By applying [21, Theorem 4.0.3, Lemma 5.1.3] (see also [15, Lemma 2.5]) to $\omega \equiv c\omega_M$ we have $c\omega_M \sim \omega_{\underline{M}^c}$, more precisely (by setting the weight matrix parameter $x = 1$)

$$(3.28) \quad \forall c \in \mathbb{N}_{>0} \exists D > 0 \forall t \geq 0: \omega_{\underline{M}^c}(t) \leq c\omega_M(t) \leq 2\omega_{\underline{M}^c}(t) + D.$$

Proposition 3.19. *Let $M, N \in \mathcal{LC}$ be given.*

(i) If we have as sets

$$(3.29) \quad \exists c, d \in \mathbb{N}_{>0} : H_{v_M^c}^\infty(\mathbb{C}) \subseteq H_{v_N^d}^\infty(\mathbb{C}),$$

then (with the same c and $c_1 = 2d$)

$$(3.30) \quad \exists c, c_1 \in \mathbb{N}_{>0} \exists C \geq 1 \forall j \in \mathbb{N} : \tilde{N}_j^c \leq C \tilde{M}_j^{c_1}.$$

(ii) If we have the (continuous) inclusion

$$(3.31) \quad H_{\mathcal{M}^c}^\infty(\mathbb{C}) \subseteq H_{\mathcal{N}^c}^\infty(\mathbb{C}),$$

then

$$(3.32) \quad \forall d \in \mathbb{N}_{>0} \exists c \in \mathbb{N}_{>0} \exists C \geq 1 \forall j \in \mathbb{N} : \tilde{N}_j^d = (N_{jd})^{1/d} \leq C(M_{jc})^{1/c} = C \tilde{M}_j^c.$$

Proof. (i) Let the parameters c and d be given and w.l.o.g. assume that $d = kc$ for some $k \in \mathbb{N}_{>0}$. We apply Theorem 3.2 to \underline{M}^c and so (3.5) holds for the corresponding class. Thus there exists $f_{v_{\underline{M}^c}} \in H_{v_{\underline{M}^c}}^\infty(\mathbb{C})$ such that $\frac{1}{v_{\underline{M}^c}} \sim t \mapsto M(f_{v_{\underline{M}^c}}, t)$. By (3.28) and the inclusion (3.29) one has $H_{v_{\underline{M}^c}}^\infty(\mathbb{C}) \subseteq H_{v_M^c}^\infty(\mathbb{C}) \subseteq H_{v_N^d}^\infty(\mathbb{C})$. Consider the function $f_{v_{\underline{M}^c}}^2$ and so $M(f_{v_{\underline{M}^c}}^2, t) = M(f_{v_{\underline{M}^c}}, t)^2$. Gathering this information we arrive at

$$\exists A, D, C \geq 1 \forall t \geq 0 : \frac{1}{AD} \exp(c\omega_M(t)) \leq \frac{1}{A} \exp(2\omega_{\underline{M}^c}(t)) \leq M(f_{v_{\underline{M}^c}}^2, t) \leq C \exp(2d\omega_N(t)),$$

and for the first estimate (3.28) has been used again. By applying this and (2.11) we obtain for all $j \in \mathbb{N}$ that

$$\begin{aligned} M_{2dj} &= \sup_{t \geq 0} \frac{t^{2dj}}{\exp(\omega_M(t))} = \left(\sup_{t \geq 0} \frac{t^{2cdj}}{\exp(c\omega_M(t))} \right)^{1/c} \geq \frac{1}{(ADC)^{1/c}} \left(\sup_{t \geq 0} \frac{t^{2cdj}}{\exp(2d\omega_N(t))} \right)^{1/c} \\ &= \frac{1}{(ADC)^{1/c}} \left(\sup_{t \geq 0} \frac{t^{cj}}{\exp(\omega_N(t))} \right)^{2d/c} = \frac{1}{(ADC)^{1/c}} (N_{cj})^{2d/c}. \end{aligned}$$

Thus

$$\exists c, d \in \mathbb{N}_{>0} \exists B \geq 1 \forall j \in \mathbb{N} : \tilde{N}_j^c = (N_{cj})^{1/c} \leq B(M_{2dj})^{1/(2d)} = B \tilde{M}_j^{2d},$$

and so (3.30) is verified.

(ii) Both spaces $H_{\mathcal{M}^c}^\infty(\mathbb{C})$ and $H_{\mathcal{N}^c}^\infty(\mathbb{C})$ are Fréchet and by the continuity of the inclusion in (3.31) we get

$$\forall d > 0 \exists c > 0 \exists C \geq 1 \forall f \in H_{\mathcal{M}^c}^\infty(\mathbb{C}) : \|f\|_{v_N^d} \leq C \|f\|_{v_M^c}.$$

We apply this estimate to the family of monomials $(f_k)_{k \in \mathbb{N}}$ and so

$$\forall d > 0 \exists c > 0 \exists C \geq 1 \forall k \in \mathbb{N} \forall z \in \mathbb{C} : \frac{|z|^k}{\exp(d\omega_N(|z|))} \leq C \frac{|z|^k}{\exp(c\omega_M(|z|))}.$$

W.l.o.g. we can assume $c_1 := \frac{1}{c} \in \mathbb{N}_{>0}$, $d_1 := \frac{1}{d} \in \mathbb{N}_{>0}$ (with $c_1 \geq d_1$) and thus, by using again (2.11) we obtain for all $j \in \mathbb{N}$ that

$$\begin{aligned} N_{j/d} &= \sup_{t \geq 0} \frac{t^{j/d}}{\exp(\omega_N(t))} = \left(\sup_{t \geq 0} \frac{t^j}{\exp(d\omega_N(t))} \right)^{1/d} \leq C^{1/d} \left(\sup_{t \geq 0} \frac{t^j}{\exp(c\omega_M(t))} \right)^{1/d} \\ &= C^{1/d} \left(\sup_{t \geq 0} \frac{t^{j/c}}{\exp(\omega_M(t))} \right)^{c/d} = C^{1/d} (M_{j/c})^{c/d}. \end{aligned}$$

So we have verified

$$\forall d_1 \in \mathbb{N}_{>0} \exists c_1 \in \mathbb{N}_{>0} \exists C \geq 1 \forall j \in \mathbb{N} : (N_{jd_1})^{1/d_1} \leq C(M_{jc_1})^{1/c_1},$$

i.e. (3.32). \square

Lemma 3.20. *Let $M, N \in \mathcal{LC}$ be given. Then the following are equivalent:*

- (i) (3.24) holds true,
- (ii) (3.30) holds true,
- (iii) (3.32) holds true.

Note: If in these assertions the symbol \leq is replaced by relation \preceq , then the analogous statement holds true as well.

Proof. Clearly, (3.32) implies (3.30) and (3.24) (for the latter set $c := 1$ and recall that $\tilde{N}^1 = N$). If (3.30) is valid, then (3.24) holds true because $\tilde{N}^c \geq N (= \tilde{N}^1)$.

Finally, if (3.24) holds, then take an arbitrary $d \in \mathbb{N}_{>0}$ be arbitrary and then $N_j \leq A(M_{cj})^{1/c}$ for some $A \geq 1$ and all $j \in \mathbb{N}$ implies $(N_{dj})^{1/d} \leq A^{1/d}(M_{cdj})^{1/(cd)}$. This verifies (3.32) (and hence (3.30)). \square

By combining Proposition 3.16, Proposition 3.19 and Lemma 3.20 we get:

Theorem 3.21. *Let $M, N \in \mathcal{LC}$ be given. Then the following are equivalent:*

- (i) M and N are related by (3.24), i.e.

$$\exists c \in \mathbb{N}_{>0} \exists A \geq 1 \forall j \in \mathbb{N} : N_j \leq A(M_{cj})^{1/c} = A\tilde{M}_j^c.$$

- (ii) The weights are related by $v_M \preceq^c v_N$.
- (iii) We have the continuous inclusion

$$H_{\tilde{M}^c}^\infty(\mathbb{C}) \subseteq H_{\tilde{N}^c}^\infty(\mathbb{C}).$$

- (iv) We have the continuous inclusion

$$H_{\tilde{M}^c}^\infty(\mathbb{C}) \subseteq H_{\tilde{N}^c}^\infty(\mathbb{C}).$$

Remark 3.22. *Recall Remark 3.11: When using in the proof of (i) in Proposition 3.19 alternatively the function θ_M^c (see (3.20)), then due to the appearance of the dilatation parameter we get that the inclusion (3.29) implies (with the same c, d):*

$$(3.33) \quad \forall c \in \mathbb{N}_{>0} \exists d \in \mathbb{N}_{>0} : \tilde{N}^c \preceq \tilde{M}^d,$$

equivalently

$$(3.34) \quad \exists c \in \mathbb{N}_{>0} : N \preceq \tilde{M}^c.$$

In order to conclude from this weaker relation we have to assume in addition that either M or N has (2.31): (3.34) gives by definition of the associated functions

$$\exists c \in \mathbb{N}_{>0} \exists h \geq 1 \exists C \geq 0 \forall t \geq 0 : \omega_{\tilde{M}^c}(t) \leq \omega_N(ht) + C,$$

hence by the technical estimate (3.26) we get

$$\frac{1}{2c}\omega_M(t) - \frac{D}{2} \leq \omega_{\tilde{M}^c}(t) \leq \omega_N(ht) + C,$$

and so

$$\exists D \geq 1 \exists c \in \mathbb{N}_{>0} \exists h \geq 1 \exists C \geq 0 \forall t \geq 0 : \omega_M(t) \leq 2c\omega_N(ht) + 2Cc + Dc.$$

(2.31) gives (ω_1) for either ω_M or ω_N . An iterated application of this property yields (see the proof of (i) in Proposition 5.1) that the previous estimate turns into

$$\omega_M(t) = O(\omega_N(t)) \Rightarrow v_M \preceq^c v_N.$$

Based on this observation the next result involves relation $N \preceq M$.

Lemma 3.23. *Let $M, N \in \mathcal{LC}$. Assume that either M or N has in addition (2.31). Consider the following assertions:*

- (i) *The sequences satisfy $N \preceq M$.*
- (ii) *The weights are related by $v_M \preceq^c v_N$,*
- (iii) *We have the continuous inclusions*

$$H_{\widetilde{M}}^\infty(\mathbb{C}) \subseteq H_{\widetilde{N}}^\infty(\mathbb{C}), \quad H_{\widetilde{M}}^\infty(\mathbb{C}) \subseteq H_{\widetilde{N}}^\infty(\mathbb{C}).$$

Then (i) \Rightarrow (ii) \Rightarrow (iii) holds true.

Proof. (i) \Rightarrow (ii) By Proposition 3.8, $N \preceq M$ implies $v_M \preceq^c v_N$. (2.31) gives (ω_1) for either ω_M or ω_N . Then the conclusion follows by an iterated application of (2.32), see the proof of (i) in Proposition 5.1.

(ii) \Rightarrow (iii) is clear. \square

We continue with a more detailed study concerning the sequences \widetilde{M}^c and their relation to M .

Lemma 3.24. *Let $M, N \in \mathcal{LC}$ be given.*

- (i) *$M \leq \widetilde{M}^c \leq \widetilde{M}^d$ for all $c, d \in \mathbb{N}_{>0}$ with $c \leq d$.*
- (ii) *$\widetilde{M}^c \preceq M$ holds true for some/any $c \in \mathbb{N}_{\geq 2}$ if and only if M has (mg). Consequently, $M \approx \widetilde{M}^c$ for some/any $c \in \mathbb{N}_{\geq 2}$ if and only if M has (mg).*
- (iii) *If M has (2.31), then each \widetilde{M}^c , too.*
- (iv) *If $M \preceq N$, then $\widetilde{M}^c \preceq \widetilde{N}^c$ for each $c \in \mathbb{N}_{>0}$.*
- (v) *If at least one of the sequences M or N has (mg), then the following are equivalent:*
 - (a) *we have $N \preceq M$;*
 - (b) *we have (3.34), i.e. $\exists c \in \mathbb{N}_{>0} : N \preceq \widetilde{M}^c$.*

Proof. (i) and (ii) have been shown in [23, Lemma 2.2].

(iii) is immediate by definition: If M has (2.31) with some L , then for \widetilde{M}^c we choose Lc .

(iv) is immediate by definition.

(v) (a) \Rightarrow (b) is clear by (i) (even without (mg)).

(v) (b) \Rightarrow (a) If M has in addition (mg), then (a) holds by (ii). If N has in addition (mg), then we argue as follows:

By assumption $N_j \leq AB^j(M_{cj})^{1/c}$ holds for some $A, B \geq 1, c \in \mathbb{N}_{>0}$ and all $j \in \mathbb{N}$. First let $k = cj$ and so $N_j \leq AB^j(M_k)^{1/c}$. Since N has (mg) we get $N_k = N_{cj} \leq B_1^{cj} N_j^c$ for some $B_1 \geq 1$ and all $j \in \mathbb{N}$. Combining both estimates gives

$$N_k \leq A^c (BB_1)^{cj} M_k, \quad \forall j, k \in \mathbb{N}, k = cj.$$

For the remaining cases let k with $cj < k < cj + c$ for some $j \in \mathbb{N}$ and then, since both M and N are also non-decreasing, we get

$$N_k \leq N_{cj+c} \leq B_2^{cj+c} N_c N_{cj} \leq N_c (AB_2)^c (BB_1 B_2)^{cj} M_{cj} \leq N_c (AB_2)^c (BB_1 B_2)^k M_k.$$

\square

Using this additional information we prove the following variant of Theorem 3.21.

Theorem 3.25. *Let $M, N \in \mathcal{LC}$. Assume that either M or N has in addition (2.31) and that either M or N has in addition (mg). Then the following are equivalent:*

- (i) *The sequences satisfy $N \preceq M$.*
- (ii) *We have the continuous inclusion*

$$H_{\underline{M}^\epsilon}^\infty(\mathbb{C}) \subseteq H_{\underline{N}^\epsilon}^\infty(\mathbb{C}).$$

- (iii) *We have the continuous inclusion*

$$H_{\overline{M}^\epsilon}^\infty(\mathbb{C}) \subseteq H_{\overline{N}^\epsilon}^\infty(\mathbb{C}).$$

Proof. (i) \Rightarrow (ii), (iii) holds by Lemma 3.23.

(ii), (iii) \Rightarrow (i) By Theorem 3.21 we get relation (3.24) and so (3.34), too. Then (v) in Lemma 3.24 yields the conclusion. \square

We close this section by showing that, when considering exponential-type systems, then the weighted spaces defined in terms of M and \widetilde{M}^d coincide.

Theorem 3.26. *Let $M \in \mathcal{LC}$ be given. Then as l.c.v.s. we get*

$$\forall d \in \mathbb{N}_{>0} : H_{\underline{M}^\epsilon}^\infty(\mathbb{C}) = H_{\underline{\widetilde{M}^d}^\epsilon}^\infty(\mathbb{C}), \quad H_{\overline{M}^\epsilon}^\infty(\mathbb{C}) = H_{\overline{\widetilde{M}^d}^\epsilon}^\infty(\mathbb{C}),$$

with $\widetilde{M}^{d^\epsilon} := (v_{M^d}^\epsilon)_{c \in \mathbb{N}_{>0}}$ and $\overline{\widetilde{M}^d}^\epsilon := (v_{\widetilde{M}^d}^{\frac{1}{\epsilon}})_{c \in \mathbb{N}_{>0}}$.

Proof. (3.26) precisely means

$$\forall d \in \mathbb{N}_{>0} : v_{\widetilde{M}^d}^d \preceq v_M \preceq v_{\widetilde{M}^d}^{2d},$$

and thus, since $v_{\widetilde{M}^d}^d \leq v_{\widetilde{M}^d}$ for all $d \in \mathbb{N}_{>0}$,

$$\forall d \in \mathbb{N}_{>0} : v_{\widetilde{M}^d} \preceq v_M \preceq v_{\widetilde{M}^d}^{2d}.$$

With other words, one obtains $v_{\widetilde{M}^d} \sim^\epsilon v_M$ for any given $d \in \mathbb{N}_{>0}$ which finishes the proof. \square

3.4. The dilatation-type weight function setting. The strategy is to reduce the abstract weight function setting to the weight sequence setting by applying the results from Section 3.2 to M^u (see (2.22)).

Proposition 3.27. *Let u and w be normalized weight functions. Assume that u is convex and that $H_u^\infty(\mathbb{C}) \subseteq H_w^\infty(\mathbb{C})$ (as sets).*

- (i) *Then the weights are related by*

$$(3.35) \quad \exists a \geq 1 \forall t \geq 0 : w^2(t) \leq au(t),$$

and so $u \preceq^\epsilon w$.

- (ii) *If in addition one of the weights is of moderate growth, then*

$$(3.36) \quad \exists a, b \geq 1 \forall t \geq 0 : w(at) \leq bu(t),$$

i.e. $u \preceq_c w$. The proof shows that we can take $a := H$ with H the constant appearing in (2.28) for the weight being of moderate growth.

Proof. (i) By assumption and (2.25) we get (as sets)

$$H_{v_{M^u}}^\infty(\mathbb{C}) \subseteq H_u^\infty(\mathbb{C}) \subseteq H_w^\infty(\mathbb{C}).$$

We apply Theorem 3.2 to M^u and get the existence of an optimal $f_{v_{M^u}}$ satisfying (3.5). The previous inclusion yields now

$$(3.37) \quad \exists C, D \geq 1 \forall t \geq 0: \quad \omega_{M^u}(t) - \log(D) \leq \log(C) - \log(w(t)) = \log(C) + \omega^w(t),$$

and recall (2.19) for the last equality.

There exist now two arguments to conclude; both make only use of the convexity for u (see Remark 2.13).

1) By combining the first part of (2.23) applied to u and (3.37) we get

$$\exists C_1 \geq 1 \exists A \geq 1 \forall t \geq 0: \quad -\log(A) + \frac{1}{2}\omega^u(t) \leq \omega_{M^u}(t) \leq \log(C_1) + \omega^w(t).$$

In view of (2.19) estimate (3.35) is shown.

2) The second method uses again (3.37) but follows then the proof of Proposition 3.12: By taking into account (2.22) we get for all $j \in \mathbb{N}$

$$M_j^w = \sup_{t>0} \frac{t^j}{\exp(\omega^w(t))} \leq CD \sup_{t>0} \frac{t^j}{\exp(\omega_{M^u}(t))} = CDM_j^u,$$

i.e. M^w and M^u are related by (1.4). Thus we have shown

$$\exists C \geq 1 \forall t \geq 0: \quad \omega_{M^u}(t) \leq \omega_{M^w}(t) + \log(C) \Leftrightarrow v_{M^w}(t) \leq Cv_{M^u}(t).$$

The second estimate in (2.23) applied to w and the first one applied to u give

$$\exists B \geq 1 \exists C \geq 1 \forall t \geq 0: \quad w(t) \leq v_{M^w}(t) \leq Cv_{M^u}(t) \leq BC\sqrt{u(t)},$$

which is verifying again (3.35).

(ii) If now w satisfies in addition (2.28), then by combining this with (3.35) we get

$$w(Ht) \leq e^H w^2(t) \leq e^H au(t),$$

hence (3.36) is shown. If u satisfies (2.28), then similarly $\sqrt{u(Ht)} \leq e^{H/2}u(t)$ for some $H \geq 1$ and all $t \geq 0$ and so

$$w(Ht) \leq \sqrt{a}\sqrt{u(Ht)} \leq \sqrt{a}e^{H/2}u(t).$$

□

Remark 3.28. If in (i) in Proposition 3.27 we use instead of $f_{v_{M^u}}$ the function $\theta_{M^u,1}$ from Lemma 3.9, then a dilatation parameter $\frac{1}{2}$ appears and instead of (3.35) we get the weaker estimate

$$(3.38) \quad \exists a \geq 1 \forall t \geq 0: \quad (w(2t))^2 \leq au(t).$$

Consequently, (3.36) holds then with $a := 2H$ instead of H . (However, in view of (ii) this technical difference is negligible since there in any case a dilatation appears.)

To treat the classes $H_{\mathbb{V}_c}^\infty(\mathbb{C})$ we follow Proposition 3.13.

Proposition 3.29. *Let u and w be normalized weights. Assume that u is convex and that*

$$(3.39) \quad H_{\underline{u}_c}^\infty(\mathbb{C}) \subseteq H_{\underline{w}_c}^\infty(\mathbb{C})$$

holds (with continuous inclusion). Then the conclusion of (ii) in Proposition 3.27 holds true; concerning (i) the parameter 2 in (3.38) has to be replaced by some $a' \geq 1$ and therefore $a := a'H$ in (3.36).

Proof. The spaces $H_{\underline{u}_c}^\infty(\mathbb{C})$ and $H_{\underline{w}_c}^\infty(\mathbb{C})$ are Fréchet and by the continuity of the inclusion in (3.39) we get

$$\forall d > 0 \exists c > 0 \exists C \geq 1 \forall f \in H_{\underline{u}_c}^\infty(\mathbb{C}) : \|f\|_{w_d} \leq C \|f\|_{u_c}.$$

We apply this estimate again to the family of monomials $(f_k)_{k \in \mathbb{N}}$ and so

$$\forall d > 0 \exists c > 0 \exists C \geq 1 \forall k \in \mathbb{N} \forall z \in \mathbb{C} : |z|^k w(d|z|) \leq C |z|^k u(c|z|).$$

Thus, by taking into account (2.19), this estimate turns into

$$\forall d > 0 \exists c > 0 \exists C \geq 1 \forall k \in \mathbb{N} \forall z \in \mathbb{C} : \frac{|z|^k}{\exp(\omega^w(d|z|))} \leq C \frac{|z|^k}{\exp(\omega^u(c|z|))}.$$

Then, by using (2.22), we get for all $j \in \mathbb{N}$ that

$$\begin{aligned} M_j^w &= \sup_{t>0} \frac{t^j}{\exp(\omega^w(t))} = \sup_{s>0} \frac{(ds)^j}{\exp(\omega^w(ds))} \leq C d^j \sup_{s>0} \frac{s^j}{\exp(\omega^u(cs))} \\ &= C \left(\frac{d}{c}\right)^j \sup_{s>0} \frac{s^j}{\exp(\omega^u(s))} = C \left(\frac{d}{c}\right)^j M_j^u, \end{aligned}$$

i.e. $M^w \preceq M^u$ is verified. Then take $d := 1$ and the rest follows analogously as the second method in (i) in Proposition 3.27 and so (3.38) holds with $a' := \frac{1}{c}$ instead of 2. \square

Gathering all this information we arrive at the following result:

Theorem 3.30. *Let u and w be normalized weight functions. Assume that u is convex and that either u or w is of moderate growth. Then the following are equivalent:*

- (i) *The weights satisfy $u \preceq_c w$.*
- (ii) *We have*

$$H_{\underline{u}_c}^\infty(\mathbb{C}) \subseteq H_{\underline{w}_c}^\infty(\mathbb{C}),$$

with continuous inclusion.

- (iii) *We have*

$$H_{\underline{u}_c}^\infty(\mathbb{C}) \subseteq H_{\underline{w}_c}^\infty(\mathbb{C}),$$

with continuous inclusion.

Consequently, if both weights are normalized and convex and at least one of them is of moderate growth, then as l.c.v.s.

$$H_{\underline{u}_c}^\infty(\mathbb{C}) = H_{\underline{w}_c}^\infty(\mathbb{C}), \quad \text{and/or} \quad H_{\underline{u}_c}^\infty(\mathbb{C}) = H_{\underline{w}_c}^\infty(\mathbb{C}),$$

if and only if $u \sim_c w$.

Moreover, under this additional assumption the previous list of equivalences can be extended by

- (iv) *The associated sequences satisfy $M^w \preceq M^u$.*
- (v) *The weights are satisfying $v_{M^u} \preceq_c v_{M^w}$.*

Proof. (i) \Leftrightarrow (ii) \Leftrightarrow (iii) follows by Propositions 2.2, 3.27 and 3.29.

(iv) \Leftrightarrow (v) holds by Theorem 3.14 (even without requiring moderate growth) and (i) \Leftrightarrow (v) follows by combining (2.24) with (2.28). Note that for this equivalence we have to use convexity for *both* weights whereas it is enough to assume moderate growth for one of the weights, see the proof of (ii) in Proposition 3.27. \square

Remark 3.31. We have reduced the proof of Theorem 3.30 to the weight sequence setting by involving the associated weight sequence M^w . For this procedure the appearing technical growth assumption (2.28) seems to be unavoidable in order to get control on the quadratic power in (2.23). Similarly, in the ultradifferentiable setting moderate growth is needed when transferring techniques from the weight function to the weight sequence setting.

In [9] the author has also started with an abstractly given increasing function ϕ satisfying $\phi(t) \neq O(\log(t))$ and such that $t \mapsto \phi(e^t)$ is convex. Note that the first requirement is weaker than $\phi(t) = o(\log(t))$ which always holds true for ω^v when v is a weight function, and the second basic assumption holds true for ω^v if and only if v is convex (see (2.20)). Summarizing, we have the relation $\phi \equiv -\log \circ v$, with v being a weight function.

Moreover, the representation $\phi(t) = \int_1^t \frac{\psi(s)}{s} ds$ is used which should be compared with (2.12), i.e. $\psi \equiv \Sigma_M$ with Σ_M denoting the counting function in (2.10). By using ψ the sequence $r = (r_i)_{i \geq 1}$ is defined via $\psi(i) = r_i$ and finally $a_n := \frac{1}{r_1 \cdots r_n}$, i.e. r corresponds to the quotient sequence μ^v and $a_n = (M_n^v)^{-1}$. Note that $\psi(1) = 0$ precisely corresponds to normalization of given v .

Then, in [9, Sect. 2] it is shown that there exists an entire function f satisfying

$$\lim_{t \rightarrow +\infty} \frac{\log(M(f, t))}{\phi(t)} = 1.$$

In fact, in [9, Lemma 2] it is shown that $\tilde{f} := f|_{\mathbb{R}}$ satisfies

$$(3.40) \quad \lim_{t \rightarrow +\infty} \frac{\log(M(f, t))}{\phi(t)} = \lim_{t \rightarrow +\infty} \frac{\log(\tilde{f}(t))}{\phi(t)} = 1.$$

(3.40) means that for all $1 > \epsilon > 0$ we have $\tilde{f}(t) \geq \exp((1 - \epsilon)\phi(t))$ for all $t \geq 0$ large enough and so the inclusion $H_v^\infty(\mathbb{C}) \subseteq H_w^\infty(\mathbb{C})$ implies (by the correspondence $\phi_v \equiv -\log \circ v$, $\phi_w \equiv -\log \circ w$)

$$\exists C \geq 1 \forall \epsilon > 0 \exists D \geq 1 \forall t \geq 0: \quad \frac{1}{D}(v(t))^{\epsilon-1} = \frac{1}{D} \exp((1 - \epsilon)\phi_v(t)) \leq C \exp(\phi_w(t)) = C \frac{1}{w(t)}.$$

But, since $1 - \epsilon < 1$, in order to verify (3.36), property (2.28) has to be applied again (to one of the weights).

3.5. The exponential-type weight function setting. Recall that by Proposition 2.2 for given weights v and w satisfying $v \leq^c w$ we get

$$H_{\mathcal{V}^c}^\infty(\mathbb{C}) \subseteq H_{\mathcal{W}^c}^\infty(\mathbb{C}), \quad H_{\mathcal{V}^c}^\infty(\mathbb{C}) \subseteq H_{\mathcal{W}^c}^\infty(\mathbb{C}),$$

with continuous inclusions.

On the other hand, by combining already shown statements we obtain the following characterization:

Theorem 3.32. Let u and w be normalized and convex weights with corresponding associated sequences M^u and M^w . Then the following are equivalent:

(i) M^u and M^w are related by (3.24), i.e.

$$\exists c \in \mathbb{N}_{>0} \exists A \geq 1 \forall j \in \mathbb{N}: \quad M_j^w \leq A(M_{c_j}^u)^{1/c}.$$

- (ii) The weights are related by $v_{M^u} \preceq^c v_{M^w}$.
 (iii) We have that $u \preceq^c w$.
 (iv) We have

$$H_{\underline{U}}^\infty(\mathbb{C}) \subseteq H_{\underline{W}}^\infty(\mathbb{C}),$$

with continuous inclusion.

- (v) We have

$$H_{\overline{U}}^\infty(\mathbb{C}) \subseteq H_{\overline{W}}^\infty(\mathbb{C}),$$

with continuous inclusion.

Proof. (i) \Leftrightarrow (ii) is precisely (i) \Leftrightarrow (ii) in Proposition 3.16 and (ii) \Leftrightarrow (iii) holds by (2.24), i.e. $v_{M^u} \sim^c u$, $v_{M^w} \sim^c w$. For this convexity is required for *both* weights; more precisely convexity for u is needed for (ii) \Rightarrow (iii) and convexity for w for (iii) \Rightarrow (ii).

Since (iii) \Rightarrow (iv), (v) holds by Proposition 2.2 it suffices to show (iv), (v) \Rightarrow (ii) in order to conclude.

By Proposition 2.15 applied to both weights and assumption (iv) resp. (v) we have (with continuous inclusion)

$$H_{\underline{M^u}}^\infty(\mathbb{C}) = H_{\underline{U}}^\infty(\mathbb{C}) \subseteq H_{\underline{W}}^\infty(\mathbb{C}) = H_{\underline{M^w}}^\infty(\mathbb{C}), \text{ resp. } H_{\overline{M^u}}^\infty(\mathbb{C}) = H_{\overline{U}}^\infty(\mathbb{C}) \subseteq H_{\overline{W}}^\infty(\mathbb{C}) = H_{\overline{M^w}}^\infty(\mathbb{C}).$$

Then (iii) \Rightarrow (ii) resp. (iv) \Rightarrow (ii) in Theorem 3.21 applied to $M \equiv M^u$ and $N \equiv M^w$ gives (ii) and hence we are done.

Note: In order to have the above relation again we need convexity for both weights. In fact for (iv), (v) \Rightarrow (ii) only convexity for w is needed (see Proposition 2.15) but for closing the circle convexity of u is required in any case by the comment in (ii) \Leftrightarrow (iii). \square

However, this results seems to be not optimal and we want to get rid of assumption convexity for w . We continue with the following statement:

Proposition 3.33. *Let u and w be normalized weights. Assume that u is convex and that either*

- (i) *we have (as sets)*

$$(3.41) \quad \exists c, d \in \mathbb{N}_{>0} : H_{u^c}^\infty(\mathbb{C}) \subseteq H_{w^d}^\infty(\mathbb{C}),$$

or that

- (ii)

$$(3.42) \quad H_{\underline{U}}^\infty(\mathbb{C}) \subseteq H_{\overline{W}}^\infty(\mathbb{C})$$

holds (with continuous inclusion).

Then

$$(3.43) \quad \exists a \geq 1 \exists b \geq 1 \forall t \geq 0 : w^a(t) \leq bu(t)$$

is valid, i.e. $u \preceq^c w$.

Proof. (i) We follow the proof of (i) in Proposition 3.19. Let the parameters c and d be given (w.l.o.g. $d \geq c \geq 1$). Then apply Theorem 3.2 to $\underline{M^u}^c$ (recall (3.27)) and so (3.5) holds for the corresponding class. Thus there exists $f_{v_{\underline{M^u}^c}} \in H_{v_{\underline{M^u}^c}}^\infty(\mathbb{C})$ such that $\frac{1}{v_{\underline{M^u}^c}} \sim t \mapsto M(f_{v_{\underline{M^u}^c}}, t)$. The inclusion (3.41) together with (2.25) and (3.28) give $H_{v_{\underline{M^u}^c}}^\infty(\mathbb{C}) \subseteq H_{v_{M^u}^c}^\infty(\mathbb{C}) \subseteq H_{u^c}^\infty(\mathbb{C}) \subseteq H_{w^d}^\infty(\mathbb{C})$. Consider the function $f_{v_{\underline{M^u}^c}}^2$ and so gathering all this information we arrive at

$$\exists A, D, C \geq 1 \forall t \geq 0 : \frac{1}{AD} \exp(c\omega_{M^u}(t)) \leq \frac{1}{A} \exp(2\omega_{\underline{M^u}^c}(t)) \leq M(f_{v_{\underline{M^u}^c}}^2, t) \leq C(w^d(t))^{-2},$$

where for the first estimate (3.28) applied to M^u has been used again. Thus, by the first part of (2.23) applied to u and for which *convexity* is needed, we have

$$\exists A, B, C, D \geq 1 \forall t \geq 0: \quad \frac{1}{ADB^c} (u^{c/2}(t))^{-1} \leq \frac{1}{AD} \exp(c\omega_{M^u}(t)) \leq C(w^d(t))^{-2},$$

hence $(w^{4d/c}(t)) \leq (AB^cCD)^{2/c}u(t)$ follows, i.e. (3.43).

Alternatively, the above estimate yields

$$\begin{aligned} \exists A, C, D \geq 1 \forall t \geq 0: \\ c\omega_{M^u}(t) - \log(AD) \leq \log(C) - 2d \log(w(t)) = \log(C) + 2d\omega^w(t), \end{aligned}$$

and then in view of (2.22) we can follow part (i) in Proposition 3.19 and get $\widetilde{M}^w_j \leq C_1 \widetilde{M}^u_j^{2d}$, i.e. (3.30) for M^w and M^u . By Lemma 3.20 the sequences M^w and M^u are related by (3.24) and so (i) \Leftrightarrow (ii) in Proposition 3.16 yields $v_{M^u} \preceq^c v_{M^w}$. Finally (2.24) implies (3.43). Note that for this last step only convexity for u is required.

(ii) Both spaces $H_{\mathcal{U}}^\infty(\mathbb{C})$ and $H_{\mathcal{W}}^\infty(\mathbb{C})$ are Fréchet and by the continuity of the inclusion in (3.42) we get

$$\forall d > 0 \exists c > 0 \exists C \geq 1 \forall f \in H_{\mathcal{U}}^\infty(\mathbb{C}): \quad \|f\|_{w^d} \leq C \|f\|_{u^c}.$$

We apply this estimate to the family of monomials $(f_k)_{k \in \mathbb{N}}$ and so

$$\begin{aligned} \forall d > 0 \exists c > 0 \exists C \geq 1 \forall k \in \mathbb{N} \forall z \in \mathbb{C}: \\ \frac{|z|^k}{\exp(d\omega^w(|z|))} = |z|^k w^d(|z|) \leq C |z|^k u^c(|z|) = C \frac{|z|^k}{\exp(c\omega^u(|z|))}. \end{aligned}$$

W.l.o.g. we can assume $c_1 := \frac{1}{c} \in \mathbb{N}_{>0}$, $d_1 := \frac{1}{d} \in \mathbb{N}_{>0}$ (with $c_1 \geq d_1$) and thus, by taking into account (2.22) we follow the proof of (ii) in Proposition 3.19 and get (3.32) between M^w and M^u . The rest follows now as in the second method in case (i). \square

Note: Alternatively, in the proof of (i) we can use the function $\theta_{M^u}^c$ from (3.20). But then the appearance of a dilatation parameter requires to involve (2.32) (for one of the weights) in order to obtain the conclusion. For this note that (2.32) for either u or w is equivalent to having (2.31) for either M^u or M^w , see Lemma 2.21.

Gathering Propositions 2.2, 3.33 and Theorem 3.32 give the following characterization:

Theorem 3.34. *Let u and w be normalized weights and assume that u is convex. Then the following are equivalent:*

- (i) *The weights satisfy $u \preceq^c w$.*
- (ii) *We have*

$$H_{\mathcal{U}}^\infty(\mathbb{C}) \subseteq H_{\mathcal{W}}^\infty(\mathbb{C}),$$

with continuous inclusion.

- (iii) *We have*

$$H_{\mathcal{U}}^\infty(\mathbb{C}) \subseteq H_{\mathcal{W}}^\infty(\mathbb{C}),$$

with continuous inclusion.

Consequently, when u and w both are convex, then we have

- (*) *as l.c.v.s. the equalities*

$$H_{\mathcal{U}}^\infty(\mathbb{C}) = H_{\mathcal{W}}^\infty(\mathbb{C}), \quad \text{and/or} \quad H_{\mathcal{U}}^\infty(\mathbb{C}) = H_{\mathcal{W}}^\infty(\mathbb{C}),$$

(*) if and only if $u \sim^\epsilon w$.

Moreover, under this additional assumption the previous list of equivalences can be extended by

(iv) M^u and M^w are related by (3.24), i.e.

$$\exists c \in \mathbb{N}_{>0} \exists A \geq 1 \forall j \in \mathbb{N} : M_j^w \leq A(M_{cj}^u)^{1/c}.$$

(v) The weights are related by $v_{M^u} \preceq^\epsilon v_{M^w}$.

This result should be compared with Theorem 3.32: There the proof for the equivalences (i) \Leftrightarrow (ii) \Leftrightarrow (iii) also requires the convexity for *both* weights.

4. WEIGHTED CLASSES OF ENTIRE FUNCTIONS WITH o -GROWTH RESTRICTION

Let $v : [0, +\infty) \rightarrow (0, +\infty)$ be a (radial and strictly positive) weight. We define weighted classes of entire functions with o -growth restriction as follows, see e.g. [1, Sect. 1.B]:

$$H_v^o(\mathbb{C}) := \{f \in H(\mathbb{C}) : |f(z)|v(|z|) \text{ vanishes at } \infty\},$$

i.e. for all $\epsilon > 0$ we can find a (radial symmetric) compact set $K \subseteq \mathbb{C}$ such that $|f(z)|v(|z|) < \epsilon$ for all $z \in \mathbb{C} \setminus K$. In [3] these classes are denoted by $H_v^0(\mathbb{C})$.

The spaces $H_{\underline{v}_c}^o(\mathbb{C})$, $H_{\underline{v}_c}^o(\mathbb{C})$, $H_{\overline{v}_c}^o(\mathbb{C})$ and $H_{\overline{v}_c}^o(\mathbb{C})$ are defined accordingly (and similarly for the weight sequence setting). Concerning the growth relations we summarize:

- (a) Given two weights v and w , then again the relation $v \preceq w$ implies $H_v^o(\mathbb{C}) \subseteq H_w^o(\mathbb{C})$ with continuous inclusion and Proposition 2.2 clearly transfers to this setting.
- (b) We remark that the crucial function f_v from (3.5) cannot belong to the class $H_v^o(\mathbb{C})$ and so the proof of Theorem 3.3 cannot be transferred to this setting.
- (c) On the other hand, when assuming the (continuous) inclusion $H_{v_M}^o(\mathbb{C}) \subseteq H_{v_N}^o(\mathbb{C})$ then we can follow the proof in Proposition 3.13 (again with $c = d$) and so the o -analogue of Corollary 3.15 holds true. For this note that the monomials f_k are elements of $H_v^o(\mathbb{C})$ since v is rapidly decreasing.
- (d) Moreover, $H_{v_c}^o(\mathbb{C}) \subseteq H_{v_c}^\infty(\mathbb{C})$, $H_{\overline{v}_c}^o(\mathbb{C}) \subseteq H_{\overline{v}_c}^\infty(\mathbb{C})$ for any (normalized) weight function v and any $c > 0$ and so with continuous inclusions

$$H_{\underline{v}_c}^o(\mathbb{C}) \subseteq H_{\underline{v}_c}^\infty(\mathbb{C}), \quad H_{\overline{v}_c}^o(\mathbb{C}) \subseteq H_{\overline{v}_c}^\infty(\mathbb{C}),$$

and

$$H_{\underline{v}_c}^o(\mathbb{C}) \subseteq H_{\underline{v}_c}^\infty(\mathbb{C}), \quad H_{\overline{v}_c}^o(\mathbb{C}) \subseteq H_{\overline{v}_c}^\infty(\mathbb{C}).$$

For weight systems we find the following result which shows that an immediate reduction to the already shown statements in the previous sections is possible. This is due to the fact that the appearing parameter allows for having "enough space" and so the corresponding spaces of o - and O -growth restriction coincide.

Theorem 4.1. *Let v be a (normalized) weight function.*

(i) *We get*

$$\forall d > c \geq c' > 0 : H_{v_{c'}}^\infty(\mathbb{C}) \subseteq H_{v_c}^\infty(\mathbb{C}) \subseteq H_{v_d}^o(\mathbb{C}) \subseteq H_{v_d}^\infty(\mathbb{C})$$

with continuous inclusions. Consequently,

$$H_{\underline{v}_c}^o(\mathbb{C}) = H_{\underline{v}_c}^\infty(\mathbb{C}), \quad H_{\overline{v}_c}^o(\mathbb{C}) = H_{\overline{v}_c}^\infty(\mathbb{C}),$$

as locally convex vector spaces.

(ii) If v has in addition

$$(4.1) \quad \exists H \geq 1 \forall c > 0 \forall d > Hc > 0: \quad \lim_{t \rightarrow +\infty} \frac{v_c(t)}{v_d(t)} = +\infty,$$

then

$$\exists H \geq 1 \forall d > Hc \geq Hc' > 0: \quad H_{v_{c'}}^\infty(\mathbb{C}) \subseteq H_{v_c}^\infty(\mathbb{C}) \subseteq H_{v_d}^o(\mathbb{C}) \subseteq H_{v_d}^\infty(\mathbb{C}),$$

with continuous inclusions. This implies

$$H_{\underline{v}_c}^o(\mathbb{C}) = H_{\underline{v}_c}^\infty(\mathbb{C}), \quad H_{\overline{v}_c}^o(\mathbb{C}) = H_{\overline{v}_c}^\infty(\mathbb{C}),$$

as locally convex vector spaces.

Proof. (i) The first and the third inclusion is trivial, the second one follows from (2.8).

(ii) The first and the third inclusion is trivial, the second one follows from (4.1) (with the same H). \square

Recall that (4.1) is not clear for abstractly given weights v . However:

- (*) By Lemma 2.9, see (2.16), this requirement is satisfied with $H = 1$ in the weight sequence setting when M is a given log-convex weight sequence.
- (*) By Lemma 2.20 this holds true provided v is a normalized and convex weight function of moderate growth and we can choose $H > 1$ to be the constant appearing in (2.28), see (2.29).

By using these comments and Theorem 4.1 we can use the main results concerning weight systems from the previous Section 3 and transfer them directly to the corresponding analogues for classes with o -growth restriction. Note that only in the analogue of Theorem 3.30 we require now an extra assumption, i.e. that *both* weights are of moderate growth.

5. COMPARISON BETWEEN DILATATION-TYPE AND EXPONENTIAL-TYPE SYSTEMS

The goal of this section is to compare the inductive and projective structures defined by (2.4) and (2.5) and to characterize equality as l.c.v.s. in terms of the given weights.

For this both technical growth restrictions (2.28) resp. (2.32), which can equivalently be expressed in terms of (ω_6) resp. (ω_1) for the function ω^v (see (2.19)), are becoming crucial.

5.1. On the sufficiency of conditions (ω_6) and (ω_1) .

Proposition 5.1. *Let v be a (normalized) weight function and let ω^v be the function from (2.19).*

(i) *If v has (2.32), equivalently if ω^v has (ω_1) , then*

$$H_{\underline{v}_c}^\infty(\mathbb{C}) \subseteq H_{\underline{v}_c}^\infty(\mathbb{C}), \quad H_{\overline{v}_c}^\infty(\mathbb{C}) \subseteq H_{\overline{v}_c}^\infty(\mathbb{C}),$$

with continuous inclusions.

(ii) *If v has (2.28), equivalently if ω^v has (ω_6) , then*

$$H_{\underline{v}_c}^\infty(\mathbb{C}) \subseteq H_{\underline{v}_c}^\infty(\mathbb{C}), \quad H_{\overline{v}_c}^\infty(\mathbb{C}) \subseteq H_{\overline{v}_c}^\infty(\mathbb{C}),$$

with continuous inclusions.

Summarizing, if v has both (2.32) and (2.28), then as l.c.v.s.

$$H_{\underline{v}_c}^\infty(\mathbb{C}) = H_{\underline{v}_c}^\infty(\mathbb{C}), \quad H_{\overline{v}_c}^\infty(\mathbb{C}) = H_{\overline{v}_c}^\infty(\mathbb{C}).$$

Proof. (i) If ω^v has (ω_1) , then (by iteration)

$$\forall C \geq 1 \exists C' \geq 1 \forall t \geq 0: \quad \omega^v(Ct) \leq C' \omega^v(t) + C' \Leftrightarrow v^{C'}(t) \leq e^{C'} v_C(t),$$

yielding the first part. Similarly,

$$\forall 0 < c \leq 1 \exists 0 < c' \leq 1 \forall t \geq 0: \quad \omega^v(t/c) \leq \frac{1}{c'} \omega^v(t) + \frac{1}{c'} \Leftrightarrow v_c(t) \leq e^{c'} v^{c'}(t),$$

showing the second part.

(ii) If ω^v has (ω_6) , then (by iteration)

$$\forall C \geq 1 \exists C' \geq 1 \forall t \geq 0: \quad C \omega^v(t) \leq \omega^v(C't) + C' \Leftrightarrow v_{C'}(t) \leq e^{C'} v^C(t),$$

yielding the first part. Similarly,

$$\forall 0 < c \leq 1 \exists 0 < c' \leq 1 \forall t \geq 0: \quad \frac{1}{c} \omega^v(t) \leq \omega^v(t/c') + \frac{1}{c'} \Leftrightarrow v^c(t) \leq e^{c'/c'} v_{c'}(t),$$

showing the second part. □

We apply Proposition 5.1 to $v = v_M$, $\omega^v = \omega_M$, and by taking into account the characterizations [23, Thm. 3.1] and [17, Prop. 3.6] we have:

Corollary 5.2. *Let $M \in \mathcal{LC}$ be given such that (mg) and (2.31) hold true. Then as l.c.v.s. we get*

$$H_{\underline{M}_c}^\infty(\mathbb{C}) = H_{\underline{M}^c}^\infty(\mathbb{C}), \quad H_{\overline{M}_c}^\infty(\mathbb{C}) = H_{\overline{M}^c}^\infty(\mathbb{C}).$$

Note: By [23, Prop. 3.4] when assuming (mg) then condition (2.31) is equivalent to

$$\exists Q \in \mathbb{N}_{>0}: \quad \liminf_{j \rightarrow +\infty} \frac{\mu_{Qj}}{\mu_j} > 1.$$

5.2. On the necessity of conditions (ω_6) and (ω_1) for the weight sequence setting. Using the developed techniques from Section 3 we are able to prove the converse statement of Proposition 5.1 and Corollary 5.2.

Proposition 5.3. *Let $M \in \mathcal{LC}$ be given.*

- (i) *The inclusion (as sets) $H_{\underline{M}_c}^\infty(\mathbb{C}) \subseteq H_{\underline{M}^c}^\infty(\mathbb{C})$ implies (ω_1) for ω_M , whereas $H_{\underline{M}^c}^\infty(\mathbb{C}) \subseteq H_{\underline{M}_c}^\infty(\mathbb{C})$ yields (ω_6) .*
- (ii) *The (continuous) inclusion $H_{\overline{M}_c}^\infty(\mathbb{C}) \subseteq H_{\overline{M}^c}^\infty(\mathbb{C})$ implies (ω_1) for ω_M , whereas the (continuous) inclusion $H_{\overline{M}^c}^\infty(\mathbb{C}) \subseteq H_{\overline{M}_c}^\infty(\mathbb{C})$ yields (ω_6) .*

For the proof it is convenient resp. sufficient to use $\theta_{M,c}$ from Lemma 3.9 resp. θ_M^c from Remark 3.11.

Proof. (i) $H_{\underline{M}_c}^\infty(\mathbb{C}) \subseteq H_{\underline{M}^c}^\infty(\mathbb{C})$ and Lemma 3.9 yield

$$\forall c > 0 \exists c' > 0 \exists D \geq 1 \forall t \geq 0: \quad \exp(\omega_M(ct/2)) \leq \theta_{M,c}(t) = |\theta_{M,c}(t)| \leq D \exp(c' \omega_M(t)).$$

When taking $c = 4$ we immediately get (ω_1) for ω_M with $L := \max\{c', \log(D)\}$.

Similarly, the inclusion $H_{\underline{M}^c}^\infty(\mathbb{C}) \subseteq H_{\underline{M}_c}^\infty(\mathbb{C})$ and Remark 3.11 imply

$$\forall c \in \mathbb{N}_{>0} \exists c' > 0 \exists D \geq 1 \forall t \geq 0: \quad \exp(c \omega_M(t/2^{1/c})) \leq \theta_M^c(t) = |\theta_M^c(t)| \leq D \exp(\omega_M(c't)).$$

We take $c = 2$ and get $2\omega_M(t) \leq \omega_M(c'\sqrt{2}t) + \log(D)$, i.e. (ω_6) with $H := \max\{c'\sqrt{2}, \log(D)\}$. Note that (formally) we only require the inclusion as sets.

(ii) Assume now that $H_{\overline{\mathcal{M}}^c}^\infty(\mathbb{C}) \subseteq H_{\overline{\mathcal{M}}_c}^\infty(\mathbb{C})$. Both spaces are Fréchet and by the continuity of this inclusion we get

$$\forall d > 0 \exists c > 0 \exists C \geq 1 \forall f \in H_{\overline{\mathcal{M}}^c}^\infty(\mathbb{C}) : \|f\|_{v_{M,d}} \leq C \|f\|_{v_M^c}.$$

We apply this estimate again to the family of monomials $(f_k)_{k \in \mathbb{N}}$, so

$$\forall d > 0 \exists c > 0 \exists C \geq 1 \forall k \in \mathbb{N} \forall z \in \mathbb{C} : \frac{|z|^k}{\exp(\omega_M(d|z|))} \leq C \frac{|z|^k}{\exp(c\omega_M(|z|))}.$$

Now fix $d < 1$ and w.l.o.g. $c' := \frac{1}{c} \in \mathbb{N}_{>0}$ (since $c \mapsto \|\cdot\|_{v_M^c}$ is non-increasing). Thus, by using again (2.11) we obtain for all $j \in \mathbb{N}$ that

$$\begin{aligned} M_{c'j} &= \sup_{t \geq 0} \frac{t^{c'j}}{\exp(\omega_M(t))} = \left(\sup_{t \geq 0} \frac{t^j}{\exp(c\omega_M(t))} \right)^{c'} \geq \frac{1}{C^{c'}} \left(\sup_{t \geq 0} \frac{t^j}{\exp(\omega_M(dt))} \right)^{c'} \\ &= \frac{1}{C^{c'}} \frac{1}{d^{c'j}} \left(\sup_{s \geq 0} \frac{s^j}{\exp(\omega_M(s))} \right)^{c'} = \frac{1}{C^{c'}} \frac{1}{d^{c'j}} (M_j)^{c'}. \end{aligned}$$

Consequently, we have shown

$$(5.1) \quad \exists c' \in \mathbb{N}_{>0} \exists C \geq 1 \forall j \in \mathbb{N}_{>0} : \frac{(M_{c'j})^{1/(c'j)}}{(M_j)^{1/j}} \geq \frac{1}{C^{1/j}} \frac{1}{d},$$

i.e. (2.31) with $c' = L$. Thus (ω_1) for ω_M follows by [23, Thm. 3.1].

Finally, if $H_{\overline{\mathcal{M}}_c}^\infty(\mathbb{C}) \subseteq H_{\overline{\mathcal{M}}^c}^\infty(\mathbb{C})$ with continuous inclusion, then

$$\forall d > 0 \exists c > 0 \exists C \geq 1 \forall f \in H_{\overline{\mathcal{M}}_c}^\infty(\mathbb{C}) : \|f\|_{v_M^d} \leq C \|f\|_{v_{M,c}},$$

and applied to the monomials we get

$$\forall d > 0 \exists c > 0 \exists C \geq 1 \forall k \in \mathbb{N} \forall z \in \mathbb{C} : \frac{|z|^k}{\exp(d\omega_M(|z|))} \leq C \frac{|z|^k}{\exp(\omega_M(c|z|))}.$$

Take $d < 1$ and assume w.l.o.g. $\frac{1}{d} \in \mathbb{N}_{>0}$, so for all $j \in \mathbb{N}$

$$\begin{aligned} M_j &= \sup_{t \geq 0} \frac{t^j}{\exp(\omega_M(t))} = c^j \sup_{t \geq 0} \frac{t^j}{\exp(\omega_M(ct))} \geq \frac{1}{C} c^j \sup_{t \geq 0} \frac{t^j}{\exp(d\omega_M(t))} \\ &= \frac{1}{C} c^j \left(\sup_{s \geq 0} \frac{s^{j/d}}{\exp(\omega_M(s))} \right)^d = \frac{1}{C} c^j M_{j/d}. \end{aligned}$$

In particular, when choosing $d := \frac{1}{2}$ we have shown

$$(5.2) \quad \exists c > 0 \exists C \geq 1 \forall j \in \mathbb{N} : M_{2j} \leq C c^{-j} M_j.$$

By [21, Thm. 9.5.1] applied to the (constant) matrix $\{M\}$ (see also [19, Thm. 1]) the estimate (5.2) is equivalent to (mg) for M and so [17, Prop. 3.6] yields (ω_6) for ω_M . \square

By taking into account Lemmas 2.17 and 2.21 and by combining Corollary 5.2 and Proposition 5.3 we get the following characterization:

Theorem 5.4. *Let $M \in \mathcal{LC}$ be given. Then the following are equivalent:*

- (i) M has (2.31) and (mg).
- (ii) ω_M has (ω_1) and (ω_6) .
- (iii) v_M satisfies (2.32) and (2.28).

(iv) As l.c.v.s. we get

$$H_{\underline{M}_c}^\infty(\mathbb{C}) = H_{\underline{M}^c}^\infty(\mathbb{C}).$$

(v) As l.c.v.s. we get

$$H_{\overline{M}_c}^\infty(\mathbb{C}) = H_{\overline{M}^c}^\infty(\mathbb{C}).$$

5.3. On the necessity of conditions (ω_6) and (ω_1) for the weight function setting. We study the comparison between the structures from (2.4) and (2.5) in the weight function setting.

Proposition 5.5. *Let u be a normalized and convex weight.*

- (i) *The inclusion (as sets) $H_{\underline{U}_c}^\infty(\mathbb{C}) \subseteq H_{\underline{U}^c}^\infty(\mathbb{C})$ implies (2.32) for u , whereas $H_{\underline{U}^c}^\infty(\mathbb{C}) \subseteq H_{\underline{U}_c}^\infty(\mathbb{C})$ implies (2.28).*
- (ii) *The (continuous) inclusion $H_{\overline{U}^c}^\infty(\mathbb{C}) \subseteq H_{\overline{U}_c}^\infty(\mathbb{C})$ implies (2.32) for u , whereas the (continuous) inclusion $H_{\overline{U}_c}^\infty(\mathbb{C}) \subseteq H_{\overline{U}^c}^\infty(\mathbb{C})$ implies (2.28).*

Note: For the proof convexity for u is indispensable.

Proof. (i) By assumption and (2.25) we get

$$\forall c > 0: H_{v_{M^u, c}}^\infty(\mathbb{C}) \subseteq H_{\underline{U}_c}^\infty(\mathbb{C}) \subseteq H_{\underline{U}^c}^\infty(\mathbb{C}).$$

These inclusions, Lemma 3.9 applied to M^u and (2.24) yield

$$\begin{aligned} \exists A \geq 1 \forall c > 0 \exists d > 0 \exists D \geq 1 \forall t \geq 0: \\ \frac{1}{A} u^{-1/2}(ct/2) \leq \exp(\omega_{M^u}(ct/2)) \leq \theta_{M^u, c}(t) = |\theta_{M^u, c}(t)| \leq D(u^d(t))^{-1}. \end{aligned}$$

When taking $c = 4$ then we get $u^{2d}(t) \leq (AD)^2 u(2t) = (AD)^2 u_2(t)$ and thus (2.32) is verified with $L := \max\{2d, 2 \log(AD)\}$.

Similarly, by assumption and (2.25) we get

$$\forall c > 0: H_{v_{M^u}^c}^\infty(\mathbb{C}) \subseteq H_{\overline{U}^c}^\infty(\mathbb{C}) \subseteq H_{\overline{U}_c}^\infty(\mathbb{C}).$$

These inclusions, Remark 3.11 applied to M^u and (2.24) yield

$$\begin{aligned} \exists A \geq 1 \forall c \in \mathbb{N}_{>0} \exists d > 0 \exists D \geq 1 \forall t \geq 0: \\ \frac{1}{A^c} u^{-c/2}(t/2^{1/c}) \leq \exp(c\omega_{M^u}(t/2^{1/c})) \leq \theta_{M^u}^c(t) = |\theta_{M^u}^c(t)| \leq D(u(dt))^{-1}. \end{aligned}$$

We choose $c = 4$ and get $u_{d2^{1/4}}(t) = u_d(t2^{1/4}) \leq A^4 D u^2(t)$. So (2.28) is verified with $H := \max\{d2^{1/4}, \log(A^4 D)\}$.

(ii) Assume now that $H_{\overline{U}^c}^\infty(\mathbb{C}) \subseteq H_{\overline{U}_c}^\infty(\mathbb{C})$. Both spaces are Fréchet and by the continuity of this inclusion we get

$$\forall d > 0 \exists c > 0 \exists C \geq 1 \forall f \in H_{\overline{U}^c}^\infty(\mathbb{C}): \|f\|_{u_d} \leq C \|f\|_{u^c}.$$

We apply this estimate again to the family of monomials $(f_k)_{k \in \mathbb{N}}$ and get

$$\forall d > 0 \exists c > 0 \exists C \geq 1 \forall k \in \mathbb{N} \forall z \in \mathbb{C}: \frac{|z|^k}{\exp(\omega^u(d|z|))} \leq C \frac{|z|^k}{\exp(c\omega^u(|z|))}.$$

Now fix $d < 1$ and w.l.o.g. $c' := \frac{1}{c} \in \mathbb{N}_{>0}$ ($c \mapsto \|\cdot\|_{u^c}$ is non-increasing). Then, by using (2.22) and following the estimation in (ii) in Proposition 5.3 we obtain (5.1) for M^u , i.e. (2.31) for this sequence. Thus (ω_1) for ω_{M^u} follows by [23, Thm. 3.1] and so Lemma 2.21 yields the conclusion.

Finally, if $H_{\underline{u}_c}^\infty(\mathbb{C}) \subseteq H_{\underline{u}}^\infty(\mathbb{C})$ with continuous inclusion, then

$$\forall d > 0 \exists c > 0 \exists C \geq 1 \forall f \in H_{\underline{u}_c}^\infty(\mathbb{C}) : \|f\|_{u^d} \leq C \|f\|_{u_c},$$

and applied to the monomials we get

$$\forall d > 0 \exists c > 0 \exists C \geq 1 \forall k \in \mathbb{N} \forall z \in \mathbb{C} : \frac{|z|^k}{\exp(d\omega^u(|z|))} \leq C \frac{|z|^k}{\exp(\omega^u(c|z|))}.$$

Take $d < 1$ such that $\frac{1}{d} \in \mathbb{N}_{>0}$ and then, by (2.22) and following the estimation in (ii) in Proposition 5.3 we obtain (with $d := \frac{1}{2}$) that M^u satisfies (5.2). Thus (mg) for M^u is verified and by Lemma 2.17 we are done. \square

Propositions 5.1 and 5.5 and Lemmas 2.17 and 2.21 yield the following characterization:

Theorem 5.6. *Let u be a normalized and convex weight. Then the following are equivalent:*

- (i) u satisfies (2.32) and (2.28).
- (ii) ω^u has (ω_1) and (ω_6) .
- (iii) ω_{M^u} has (ω_1) and (ω_6) .
- (iv) M^u satisfies (2.31) and (mg).
- (v) As l.c.v.s. we get

$$H_{\underline{u}}^\infty(\mathbb{C}) = H_{\underline{u}_c}^\infty(\mathbb{C}).$$

- (vi) As l.c.v.s. we get

$$H_{\underline{u}_c}^\infty(\mathbb{C}) = H_{\underline{u}}^\infty(\mathbb{C}).$$

6. STRONG INCLUSION RELATIONS FOR WEIGHTED SPACES OF ENTIRE FUNCTIONS

The aim of this section is to replace the introduced growth relations by their stronger o -growth counterparts and to study these conditions and their consequences.

6.1. Strong growth relations for weight functions and weight sequences. Let v and w be (normalized) weight functions. We write $v \triangleleft w$ if

$$w(t) = o(v(t)) \quad \text{as } t \rightarrow +\infty.$$

Obviously $v \triangleleft w$ implies $v \preceq w$ and this relation is neither reflexive nor symmetric. Next we introduce

$$v \triangleleft_c w \quad \text{resp.} \quad v \triangleleft^c w,$$

defined by

$$\forall c > 0 : v \triangleleft_c w \quad \text{resp.} \quad v \triangleleft^c w.$$

Remark 6.1. *We gather some facts:*

- (i) In both \triangleleft_c and \triangleleft^c we are interested in small $c (\leq 1)$.
- (ii) Clearly, $v \triangleleft_c w$ resp. $v \triangleleft^c w$ is equivalent to

$$(6.1) \quad \forall c, d > 0 : v_d \triangleleft_c w \quad \text{resp.} \quad v^d \triangleleft^c w.$$

- (iii) Since by Remark 2.3 we have $w^c \triangleleft w^d$ for all $0 < c < d$, condition $v \triangleleft^c w$ is equivalent to

$$(6.2) \quad \forall c > 0 : v \preceq w^c.$$

On the other hand, let $M, N \in \mathbb{R}_{>0}^{\mathbb{N}}$ and write

$$M \triangleleft N \Leftrightarrow \forall h > 0 \exists C_h \geq 1 \forall j \in \mathbb{N} : M_j \leq C_h h^j N_j \Leftrightarrow \lim_{j \rightarrow +\infty} \left(\frac{M_j}{N_j} \right)^{1/j} = 0.$$

Obviously $M \triangleleft N$ implies $M \preceq N$ and this relation is neither reflexive nor symmetric. We gather consequences of these relations; compare this also with Lemma 2.20.

Lemma 6.2. *Let M, N be weight sequences and assume that M is log-convex. Then the following are equivalent:*

- (a) $M \triangleleft N$ is valid,
- (b) $v_N \triangleleft_c v_M$ is valid,
- (c) $v_N \preceq_{v_M, c}$ is valid for all $c > 0$.

Proof. (a) \Rightarrow (b) $M \triangleleft N$ implies by definition

$$(6.3) \quad \forall h > 0 \exists C_h \geq 1 \forall t \geq 0 : \omega_N(t) \leq \omega_M(ht) + \log(C_h).$$

Let now $c > c' > 0$ be given and fixed (the case $c' \geq c$ follows then) and choose $d > 0$ and $h > 0$ such that $c' > d$ and $hc \leq d$. So $\omega_N(ct) \leq \omega_M(dt) + \log(C_h)$ for all $t \geq 0$ follows and (2.12) yields

$$\omega_M(c't) - \omega_M(dt) = \int_{dt}^{c't} \frac{\Sigma_M(u)}{u} du \geq \Sigma_M(dt) \int_{dt}^{c't} \frac{1}{u} du = \Sigma_M(dt) \log(c'/d) \rightarrow +\infty \quad \text{as } t \rightarrow +\infty,$$

by the definition of Σ_M in (2.10), recall also (2.16). Thus

$$\forall \epsilon > 0 \exists t_\epsilon > 0 \forall t \geq t_\epsilon : \omega_M(dt) + \frac{1}{\epsilon} \leq \omega_M(c't),$$

and gathering everything we get

$$\forall \epsilon > 0 \exists t_\epsilon > 0 \forall t \geq t_\epsilon : \omega_N(ct) \leq \omega_M(dt) + \log(C_h) \leq \log(C_h) - \frac{1}{\epsilon} + \omega_M(c't).$$

This verifies (the first half of) (6.1) since C_h is only depending on given (fixed) choices c, c' but not on $\epsilon \rightarrow 0$.

(b) \Rightarrow (c) This is trivial.

(c) \Rightarrow (a) If $v_N \preceq_{v_M, c}$ is valid for all $c > 0$, then

$$\forall c > 0 \exists C \geq 1 \forall t \geq 0 : v_{M,c}(t) \leq C v_N(t) \Leftrightarrow \frac{1}{\exp(\omega_M(ct))} \leq C \frac{1}{\exp(\omega_N(t))}.$$

Thus (2.11) yields for all $j \in \mathbb{N}$

$$M_j = \sup_{t \geq 0} \frac{t^j}{\exp(\omega_M(t))} = c^j \sup_{t \geq 0} \frac{t^j}{\exp(\omega_M(ct))} \leq C c^j \sup_{t \geq 0} \frac{t^j}{\exp(\omega_N(t))} = C c^j N_j^{1c} \leq C c^j N_j,$$

which gives $M \triangleleft N$ (as $c \rightarrow 0$). □

Lemma 6.3. *Let $M, N \in \mathcal{LC}$.*

- (a) *If either M or N has in addition (mg), then $v_N \triangleleft_c v_M$ implies $v_N \triangleleft^{c'} v_M$.*
- (b) *If either M or N has in addition (2.31), then $v_N \triangleleft^{c'} v_M$ implies $v_N \triangleleft_c v_M$.*
- (c) *The following are equivalent:*
 - (i) $v_N \triangleleft^{c'} v_M$,
 - (ii) $\omega_N(t) = o(\omega_M(t))$ as $t \rightarrow +\infty$,

(iii) we have that

$$(6.4) \quad \forall c \in \mathbb{N}_{>0} \exists A \geq 1 : \widetilde{M}_j^c = (M_{cj})^{1/c} \leq AN_j.$$

Proof. The proof follows similarly like in Proposition 5.1.

(a) If M has (mg) then we get (ω_6) for ω_M . Let $1 > c > 0$ be given, then iterating (ω_6) yields

$$\exists H \geq 1 \forall t \geq 0 : \omega_M(t) \leq c\omega_M(Ht) + cH.$$

By Lemma 6.2 relation $v_N \triangleleft_c v_M$ implies $M \triangleleft N$ and so (6.3). We apply this to $h := H^{-1}$ and get

$$\begin{aligned} \exists H \geq 1 \exists C_{H^{-1}} \geq 1 \forall t \geq 0 : \omega_N(t) &\leq \omega_M(H^{-1}t) + \log(C_{H^{-1}}) \leq c\omega_M(t) + cH + \log(C_{H^{-1}}) \\ &\implies v_M^c(t) \leq e^{cH} C_{H^{-1}} v_N(t). \end{aligned}$$

Thus (6.2) is shown and in view of Remark 6.1 we are done.

On the other hand, if N has (mg), then similarly by iterating (ω_6) for ω_N and (6.3) we have

$$\omega_N(t) \leq c\omega_N(Ht) + cH \leq c\omega_M(Hht) + c \log(C_h) + cH.$$

(6.3) again applied to $h := H^{-1}$ yields the conclusion.

(b) If M satisfies (2.31), then ω_M has (ω_1) . Let $1 > h > 0$ be given, then iterating (ω_1) gives

$$\exists L \geq 1 \forall t \geq 0 : \omega_M(t) \leq L\omega_M(th) + L.$$

$v_N \triangleleft_c v_M$ is equivalent to (6.2), see Remark 6.1. We apply this relation to $c := L^{-1}$ and get

$$\exists C \geq 1 \forall t \geq 0 : \omega_N(t) \leq c\omega_M(t) + C \leq \omega_M(th) + 1 + C \implies v_{M,h}(t) \leq e^{1+C} v_N(t).$$

Since $h > 0$ was arbitrary we have verified $v_N \preceq v_{M,h}$ for all $h > 0$ (small) and thus by Lemma 6.2 we are done.

On the other hand, if N satisfies (2.31), then similarly $\omega_N(t) \leq L\omega_N(th) + L$ and so again (6.2) applied to $c := L^{-1}$ gives

$$\omega_N(t) \leq L\omega_N(th) + L \leq Lc\omega_M(th) + LC + L \implies v_{M,h}(t) \leq e^{L(C+1)} v_N(t).$$

(c)(i) \Leftrightarrow (ii) First (compare with (v) in Remark 2.7) relation $\omega_N(t) = o(\omega_M(t))$ as $t \rightarrow +\infty$, i.e. $\omega_M \triangleleft \omega_N$, holds if and only if for all $0 < c \leq 1$ we get $v_N \preceq v_M^c$, i.e. (6.2). By (ii), (iii) in Remark 6.1 this is equivalent to $v_N \triangleleft_c v_M^c$ for all $0 < c \leq 1$, i.e. $v_N \triangleleft_c v_M$.

(c)(ii) \Leftrightarrow (iii) This is precisely the second part of [14, Lemma 6.5]. \square

6.2. The dilatation-type weight sequence case. We start with the following result.

Proposition 6.4. *Let M, N be weight sequences and assume that M is log-convex. Consider the following assertions:*

- (i) *The weights are satisfying $M \triangleleft N$.*
- (ii) *The weights are satisfying $v_N \triangleleft_c v_M$.*
- (iii) *We have that*

$$(6.5) \quad \forall c, c' > 0 : H_{v_{N,c}}^\infty(\mathbb{C}) \subseteq H_{v_{M,c'}}^\infty(\mathbb{C}),$$

with continuous inclusion.

- (iv) *We have that*

$$(6.6) \quad H_{\underline{N}_c}^\infty(\mathbb{C}) \subseteq H_{\underline{M}_c}^\infty(\mathbb{C}),$$

with continuous inclusion.

Then (i) \Leftrightarrow (ii) \Rightarrow (iii) \Rightarrow (iv) holds true.

Proof. (i) \Leftrightarrow (ii) is (a) \Leftrightarrow (b) in Lemma 6.2.

(ii) \Rightarrow (iii) \Rightarrow (iv) is clear. \square

Now we treat the converse statement.

Proposition 6.5. *Let M, N be weight sequences and assume that M is log-convex. If we have (as sets)*

$$(6.7) \quad \exists c > 0 \forall c' > 0 : H_{v_{N,c}}^\infty(\mathbb{C}) \subseteq H_{v_{M,c'}}^\infty(\mathbb{C}),$$

then $M \triangleleft N$ holds true.

Note that (6.5) is stronger than (6.7).

Proof. Let $c > 0$ be the parameter and by assumption $\theta_{N,c} \in H_{v_{M,c'}}^\infty(\mathbb{C})$ for any $c' > 0$, hence

$$\forall c' > 0 \exists C \geq 1 \forall t \geq 0 : \exp(\omega_N(ct/2)) \leq |\theta_{N,c}(t)| \leq C \exp(\omega_M(c't)).$$

By using (2.11) we obtain for all $j \in \mathbb{N}$ that

$$\begin{aligned} M_j &= \sup_{t \geq 0} \frac{t^j}{\exp(\omega_M(t))} = \sup_{s \geq 0} \frac{(c's)^j}{\exp(\omega_M(c's))} \leq C(c')^j \sup_{s \geq 0} \frac{s^j}{\exp(\omega_N(cs/2))} \\ &= C(2c'/c)^j \sup_{u \geq 0} \frac{u^j}{\exp(\omega_N(u))} = C(2c'/c)^j N_j^{\text{lc}} \leq C(2c'/c)^j N_j. \end{aligned}$$

Then let $c' \rightarrow 0$ and so $M \triangleleft N$ is verified (note that c is given and fixed). \square

An independent argument is to apply Theorem 3.4 to $v_{N \frac{1}{c}}$ and $v_{M \frac{1}{c'}}$ (see (2.18)) and get

$$\exists c > 0 \forall c' > 0 \exists A \geq 1 \forall j \in \mathbb{N} : \frac{1}{(c')^j} M_j = M_j^{\frac{1}{c'}} \leq A N_j^{\frac{1}{c}} = A \frac{1}{c^j} N_j.$$

As $c' \rightarrow 0$ this verifies again $M \triangleleft N$ and thus the same estimate as before (apart from the fact that the constant 2 disappears but which does not effect the conclusion). Note that for this method we also require log-convexity for N .

Gathering Lemma 6.2 and Propositions 6.4 and 6.5 we arrive at the following characterization:

Theorem 6.6. *Let M, N be weight sequences and assume that M is log-convex. Then the following are equivalent:*

- (i) *The sequences are satisfying $M \triangleleft N$.*
- (ii) *The corresponding weights are satisfying $v_N \triangleleft_c v_M$.*
- (iii) *The corresponding weights are satisfying $v_N \preceq_c v_{M,c}$ for all $c > 0$.*
- (iv) *(6.5) holds true, i.e.*

$$\forall c, c' > 0 : H_{v_{N,c}}^\infty(\mathbb{C}) \subseteq H_{v_{M,c'}}^\infty(\mathbb{C}),$$

with continuous inclusion.

(v) (6.7) holds true, i.e.

$$\exists c > 0 \forall c' > 0 : H_{v_{N,c}}^\infty(\mathbb{C}) \subseteq H_{v_{M,c'}}^\infty(\mathbb{C}),$$

with continuous inclusion.

(vi) (6.6) holds true, i.e.

$$H_{\underline{N}_c}^\infty(\mathbb{C}) \subseteq H_{\underline{M}_c}^\infty(\mathbb{C}),$$

with continuous inclusion.

6.3. The exponential-type weight sequence case. We start with the following result.

Proposition 6.7. *Let $M, N \in \mathcal{LC}$ be given and consider the following assertions:*

- (i) *We have $M \triangleleft N$ and such that either M or N has in addition (mg) resp. such that (6.4) holds true.*
- (ii) *The weights are satisfying $v_N \triangleleft^c v_M$.*
- (iii) *We have*

$$(6.8) \quad \forall c, c' > 0: \quad H_{v_N^c}^\infty(\mathbb{C}) \subseteq H_{v_M^{c'}}^\infty(\mathbb{C}),$$

with continuous inclusion.

- (iv) *We have*

$$(6.9) \quad H_{\underline{N}^c}^\infty(\mathbb{C}) \subseteq H_{\widetilde{M}^c}^\infty(\mathbb{C}),$$

with continuous inclusion.

Then (i) \Rightarrow (ii) \Rightarrow (iii) \Rightarrow (iv) holds true.

Proof. (i) \Rightarrow (ii) follows by combining Lemma 6.2 and (a) resp. (c) in Lemma 6.3. (ii) \Rightarrow (iii) follows immediately by definition of relation $v_N \triangleleft^c v_M$ and (iii) \Rightarrow (iv) is clear. \square

Proposition 6.8. *Let $M, N \in \mathcal{LC}$ be given. Then the following are equivalent:*

- (i) *We have (as sets)*

$$\forall c \in \mathbb{N}_{>0} \forall c' > 0: \quad H_{v_N^c}^\infty(\mathbb{C}) \subseteq H_{v_M^{c'}}^\infty(\mathbb{C}).$$

- (ii) *We have (as sets)*

$$(6.10) \quad \exists c \in \mathbb{N}_{>0} \forall c' > 0: \quad H_{v_N^c}^\infty(\mathbb{C}) \subseteq H_{v_M^{c'}}^\infty(\mathbb{C}).$$

- (iii) *The weight sequences satisfy*

$$(6.11) \quad \exists c \in \mathbb{N}_{>0} \forall d \in \mathbb{N}_{>0} \exists C \geq 1 \forall j \in \mathbb{N}: \quad \widetilde{M}_j^{cd} = (M_{cdj})^{1/(cd)} \leq CN_j.$$

- (iv) *The weight sequences satisfy (6.4), i.e.*

$$\forall d \in \mathbb{N}_{>0} \exists C \geq 1 \forall j \in \mathbb{N}: \quad \widetilde{M}_j^d \leq CN_j.$$

- (v) *The weights satisfy $v_N \triangleleft^c v_M$.*

Proof. (i) \Rightarrow (ii) and (v) \Rightarrow (i) are clear. (iv) \Leftrightarrow (v) holds by (c) in Lemma 6.3, (iv) \Rightarrow (iii) is clear and (iii) \Rightarrow (iv) holds as follows: Let $d \in \mathbb{N}_{>0}$ be given and take $d_1 \in \mathbb{N}_{>0}$ such that $d \leq cd_1$ (with c appearing in (6.11)). By (i) in Lemma 3.24 we get $\widetilde{M}^d \leq \widetilde{M}^{cd_1}$ and then (6.11) yields the conclusion.

It remains to show (ii) \Rightarrow (iii) and we follow (i) in Proposition 3.19. For the given and fixed c we apply Theorem 3.2 to \underline{N}^c (recall (3.27)) and so there exists an optimal $f_{v_{\underline{N}^c}} \in H_{v_{\underline{N}^c}}^\infty(\mathbb{C})$ such that $\frac{1}{v_{\underline{N}^c}} \sim t \mapsto M(f_{v_{\underline{N}^c}}, t)$. By (3.28) and the inclusion (6.10) one has $H_{v_{\underline{N}^c}}^\infty(\mathbb{C}) \subseteq H_{v_N^c}^\infty(\mathbb{C}) \subseteq H_{v_M^{c'}}^\infty(\mathbb{C})$ for all $c' > 0$. Gathering everything we arrive at

$$\exists c \in \mathbb{N}_{>0} \exists A, D \geq 1 \forall c' > 0 \exists C \geq 1 \forall t \geq 0:$$

$$\frac{1}{AD} \exp(c\omega_N(t)) \leq \frac{1}{A} \exp(2\omega_{\underline{N}^c}(t)) \leq M(f_{v_{\underline{N}^c}}^2, t) \leq C \exp(2c'\omega_M(t)),$$

and for the first estimate (3.28) has been used again.

Let now $c' = \frac{1}{2d}$ for any given $d \in \mathbb{N}_{>0}$. Then we use the previous estimate and (2.11) and obtain for all $j \in \mathbb{N}$ that

$$\begin{aligned} M_{cdj} &= M_{cj/(2c')} = \sup_{t \geq 0} \frac{t^{cj/(2c')}}{\exp(\omega_M(t))} = \left(\sup_{t \geq 0} \frac{t^{cj}}{\exp(2c'\omega_M(t))} \right)^d \leq (ADC)^d \left(\sup_{t \geq 0} \frac{t^{cj}}{\exp(c\omega_N(t))} \right)^d \\ &= (ADC)^d \left(\sup_{t \geq 0} \frac{t^j}{\exp(\omega_N(t))} \right)^{cd} = (ADC)^d (N_j)^{cd}. \end{aligned}$$

Thus (6.11) is verified. \square

When combining Propositions 6.7 and 6.8 and Lemma 6.3 we get the following final statement:

Theorem 6.9. *Let $M, N \in \mathcal{LC}$ be given. Then the following are equivalent:*

(i) *The weight sequences satisfy (6.4), i.e.*

$$\forall d \in \mathbb{N}_{>0} \exists C \geq 1 \forall j \in \mathbb{N}: (M_{dj})^{1/d} = \widetilde{M}_j^d \leq CN_j.$$

(ii) *The corresponding weights are satisfying $v_N \triangleleft_c v_M$.*

(iii) (6.8) holds true, i.e.

$$\forall c, c' > 0: H_{v_N^c}^\infty(\mathbb{C}) \subseteq H_{v_M^{c'}}^\infty(\mathbb{C}),$$

with continuous inclusion.

(iv) (6.10) holds true, i.e.

$$\exists c > 0 \forall c' > 0: H_{v_N^c}^\infty(\mathbb{C}) \subseteq H_{v_M^{c'}}^\infty(\mathbb{C}),$$

with continuous inclusion.

(v) (6.9) holds true, i.e.

$$H_{\underline{N}^c}^\infty(\mathbb{C}) \subseteq H_{\overline{M}^c}^\infty(\mathbb{C}),$$

with continuous inclusion.

If in addition either M or N has (mg) and either M or N has (2.31), then the list of equivalences can be extended by

(vi) *The sequences are satisfying $M \triangleleft N$.*

(vii) *The weights are satisfying $v_N \triangleleft_c v_M$.*

6.4. The dilatation-type weight function setting. We start with the following characterization.

Lemma 6.10. *Let u and w be normalized weight functions and assume that u is convex. Consider the following assertions:*

(a) *The associated sequences satisfy $M^w \triangleleft M^u$.*

(b) *The weights are satisfying*

$$(6.12) \quad \exists A \geq 1 \forall h > 0 \exists C_h \geq 1 \forall t \geq 0: w(Aht) \leq AC_h \sqrt{u(t)}.$$

(c) *The weights are satisfying*

$$(6.13) \quad \exists A \geq 1 \forall h > 0 \exists C_h \geq 1 \forall t \geq 0: w(Aht) \leq AC_h u(t).$$

(d) *The weights are satisfying $u \triangleleft_c w$.*

If u is also of moderate growth, then (a) \Rightarrow (b) \Rightarrow (c) \Rightarrow (d).

If w is convex and either w or u is of moderate growth, then (d) \Rightarrow (a).

Note: For $(a) \Rightarrow (b)$ moderate growth is not needed, for $(b) \Rightarrow (c)$ it is sufficient to require that either u or w has moderate growth, $(c) \Rightarrow (d)$ holds for arbitrary u if w is convex and of moderate growth and for $(d) \Rightarrow (a)$ convexity for u is also not required.

Proof. $(a) \Rightarrow (b)$ By definition we get

$$\forall h > 0 \exists C_h \geq 1 \forall t \geq 0: \quad \omega_{M^u}(t) \leq \omega_{M^w}(ht) + \log(C_h) \Leftrightarrow v_{M^w, h}(t) \leq C_h v_{M^u}(t).$$

Now use (2.23): We apply the second estimate there to w and the first one to u for which we require convexity. Thus we arrive at

$$\exists A \geq 1 \forall h > 0 \exists C_h \geq 1 \forall t \geq 0: \quad w(ht) \leq v_{M^w}(ht) \leq C_h v_{M^u}(t) \leq AC_h \sqrt{u(t)},$$

which yields (6.12).

$(b) \Rightarrow (c)$ If u satisfies (2.28), then $\sqrt{u(Ht)} \leq e^{H/2}u(t)$ for some $H \geq 1$ and all $t \geq 0$ and so

$$w(AHht) \leq AC_h \sqrt{u(Ht)} \leq AC_h e^{H/2}u(t),$$

i.e. (6.13).

Similarly, if w satisfies in addition (2.28), then by combining this with (6.12) we get

$$\exists A \geq 1 \exists H \geq 1 \forall h > 0 \exists C_h \geq 1 \forall t \geq 0: \quad w(HAht) \leq e^H (w(Aht))^2 \leq e^H (AC_h)^2 u(t).$$

$(c) \Rightarrow (d)$ If u satisfies in addition (2.28), then we write

$$\frac{w(t)}{u(ct/d)} = \frac{w(t)}{u((Ah)^{-1}t)} \frac{u((Ah)^{-1}t)}{u(ct/d)},$$

and apply (2.29) to u : For given $d, c > 0$ we have to choose $h > 0$ small enough to ensure $(Ah)^{-1} > Hc/d \Leftrightarrow \frac{d}{AHc} > h$ with H denoting the constant appearing in (2.28) and A the one appearing in (6.13). Thus $w(t) = o(u(ct/d))$ follows and we have shown $w_d(t) = o(u_c(t))$ for any $d, c > 0$, i.e. $u \triangleleft_c w$. Note that in order to apply (2.29) also convexity for u is used.

Similarly, if u is arbitrary but w satisfies in addition (2.28) and is convex, then for all $h, d, c > 0$ we write

$$\frac{w_d(t)}{u_c(t)} = \frac{w(dt)}{u(ct)} = \frac{w(dt)}{w(Ahct)} \frac{w(Ahct)}{u(ct)}.$$

By (6.13) and (2.29) applied to w we get $w(dt) = o(u(ct))$ for all $d > HAhc$. So, when given $d, c > 0$, we are choosing (again) $h < \frac{d}{AHc}$ in order to conclude.

$(d) \Rightarrow (a)$ $u \triangleleft_c w$ means $w(ct) = o(u(t))$ as $t \rightarrow +\infty$ for all $c > 0$. Then by the first estimate in (2.23) applied to w , for which we require convexity, and the second one applied to u we get $v_{M^w}^2(ct) = o(v_{M^u}(t))$ for all $c > 0$, i.e. $v_{M^u} \triangleleft_c v_{M^w}^2$. Analogously as in $(b) \Rightarrow (c)$, if either u or w has property (2.28), then $v_{M^u} \triangleleft_c v_{M^w}$ follows. Finally $(b) \Rightarrow (a)$ in Lemma 6.2 finishes the conclusion. \square

The following statement is analogous to Proposition 6.4.

Proposition 6.11. *Let u and w be weights. Consider the following assertions:*

- (i) *The weights are satisfying $u \triangleleft_c w$.*
- (ii) *We have that*

$$(6.14) \quad \forall c, c' > 0: \quad H_{u_c}^\infty(\mathbb{C}) \subseteq H_{w_{c'}}^\infty(\mathbb{C}),$$

with continuous inclusion.

(iii) We have that

$$H_{\underline{u}}^\infty(\mathbb{C}) \subseteq H_{\overline{w}_c}^\infty(\mathbb{C}),$$

with continuous inclusion.

Then (i) \Rightarrow (ii) \Rightarrow (iii) holds true.

Proof. (i) \Rightarrow (ii) Since $u \triangleleft_c w$ is equivalent to $u_d \triangleleft w_c$ for all $c, d > 0$ (i.e. the first part of (6.1)) the assertion follows immediately.

(ii) \Rightarrow (iii) is clear. \square

Next we deal with the converse statement which is analogous to Proposition 6.5; we make again a reduction to the weight sequence setting.

Proposition 6.12. *Let u and w be normalized weights. Assume that u is convex and that (as sets)*

$$(6.15) \quad \exists c > 0 \forall c' > 0 : H_{u,c}^\infty(\mathbb{C}) \subseteq H_{w,c'}^\infty(\mathbb{C}).$$

(i) Then we get

$$\exists A \geq 1 \forall h > 0 \exists C_h \geq 1 \forall t \geq 0 : w(ht) \leq AC_h \sqrt{u(t)},$$

i.e. (6.12).

(ii) If u is in addition of moderate growth or if in addition w is convex and of moderate growth, then

$$(6.16) \quad \exists A \geq 1 \exists H \geq 1 \forall h > 0 \exists C_h \geq 1 \forall t \geq 0 : w(AHht) \leq AC_h u(t),$$

with H the constant appearing in (2.28) for the particular weight.

Finally, in this case (6.16) implies $u \triangleleft_c w$.

Proof. (i) We follow the proof given in Proposition 3.27.

Let $c > 0$ be the fixed parameter. Then, by assumption and the second part of (2.23) applied to u , we get

$$\exists c > 0 \forall c' > 0 : H_{v_{M^u,c}}^\infty(\mathbb{C}) \subseteq H_{u,c}^\infty(\mathbb{C}) \subseteq H_{w,c'}^\infty(\mathbb{C}).$$

Hence, by applying this to the function $\theta_{M^u,c}$, we have

$$\exists c > 0 \forall c' > 0 \exists C \geq 1 \forall t \geq 0 : \omega_{M^u}(ct/2) \leq \log(C) - \log(w(c't)) = \log(C) + \omega^w(c't),$$

which gives for all $j \in \mathbb{N}$:

$$\begin{aligned} M_j^w &= \sup_{t>0} \frac{t^j}{\exp(\omega^w(t))} = \sup_{t>0} \frac{(c't)^j}{\exp(\omega^w(c't))} \leq C(c')^j \sup_{t>0} \frac{t^j}{\exp(\omega_{M^u}(ct/2))} \\ &= C(2c'/c)^j \sup_{s>0} \frac{s^j}{\exp(\omega_{M^u}(s))} = C(2c'/c)^j M_j^u. \end{aligned}$$

Now let $c' \rightarrow 0$ and since $c > 0$ is fixed we have verified $M^w \triangleleft M^u$. Then (a) \Rightarrow (b) in Lemma 6.10, for which we require convexity for u , yields the conclusion. (When working with the optimal $f_{v_{M^u,c}}$ instead of $\theta_{M^u,c}$ we can only get rid of the constant 2 above.)

(ii) This follows by (b) \Rightarrow (c) \Rightarrow (d) in Lemma 6.10. \square

Gathering Lemmas 6.2, 6.10 and Propositions 6.11 and 6.12 we get the following characterization:

Theorem 6.13. *Let u and w be normalized weights and assume that u is convex. Consider the following assertions:*

(i) The associated weight sequences satisfy $M^w \triangleleft M^u$.

- (ii) The weights satisfy $v_{M^u} \triangleleft_c v_{M^w}$.
- (iii) The weights satisfy $u \triangleleft_c w$.
- (iv) We have (6.14), i.e.

$$\forall c, c' > 0 : H_{u_c}^\infty(\mathbb{C}) \subseteq H_{w_{c'}}^\infty(\mathbb{C}),$$

with continuous inclusion.

- (v) We have (6.15), i.e.

$$\exists c > 0 \forall c' > 0 : H_{u,c}^\infty(\mathbb{C}) \subseteq H_{w,c'}^\infty(\mathbb{C}),$$

with continuous inclusion.

- (vi) We have

$$H_{\underline{u}_c}^\infty(\mathbb{C}) \subseteq H_{\overline{w}_c}^\infty(\mathbb{C}),$$

with continuous inclusion.

If u is of moderate growth then (i) \Leftrightarrow (ii) \Rightarrow (iii) \Leftrightarrow (iv) \Leftrightarrow (v) \Leftrightarrow (vi); if w is also convex (and either v or w is of moderate growth), then all listed assertions are equivalent.

6.5. The exponential-type weight function setting.

Proposition 6.14. *Let u and w be normalized weights. Consider the following assertions:*

- (i) The weights satisfy $u \triangleleft^c w$.
- (ii) We have that

$$(6.17) \quad \forall c, d > 0 : H_{u^c}^\infty(\mathbb{C}) \subseteq H_{w^d}^\infty(\mathbb{C}),$$

with continuous inclusion.

- (iii) We have that

$$H_{\underline{u}^c}^\infty(\mathbb{C}) \subseteq H_{\overline{w}^c}^\infty(\mathbb{C}),$$

with continuous inclusion.

Then (i) \Rightarrow (ii) \Rightarrow (iii) holds true.

Proof. (i) \Rightarrow (ii) Since $u \triangleleft^c w$ is equivalent to $u^d \triangleleft w^c$ for all $c, d > 0$ (i.e. the second part of (6.1)) the assertion follows immediately.

(ii) \Rightarrow (iii) is clear. \square

Proposition 6.15. *Let u and w be normalized weights and assume that u is convex. If (as sets)*

$$(6.18) \quad \exists c \in \mathbb{N}_{>0} \forall d > 0 : H_{u^c}^\infty(\mathbb{C}) \subseteq H_{w^d}^\infty(\mathbb{C}),$$

then $u \triangleleft^c w$ holds true.

Proof. A word-by-word repetition of the proof given in (i) in Proposition 3.33 gives

$$\exists c \in \mathbb{N}_{>0} \exists A, B, D \geq 1 \forall d > 0 \exists C \geq 1 \forall t \geq 0 :$$

$$\frac{1}{ADB^c} (u^{c/2}(t))^{-1} \leq \frac{1}{AD} \exp(c\omega_{M^u}(t)) \leq C(w^d(t))^{-2},$$

equivalently $(w^{4d/c}(t)) \leq (AB^c CD)^2 u(t)$. As $d \rightarrow 0$ we have shown $u \preceq w^{d_1}$ for all $d_1 > 0$ which is precisely (6.2). By (iii) in Remark 6.1 this is equivalent to $u \triangleleft^c w$. \square

We summarize:

Theorem 6.16. *Let u and w be normalized weights and assume that u is convex. Then the following are equivalent:*

- (i) The weights are satisfying $u \triangleleft^\epsilon w$.
- (ii) (6.17) holds true, i.e.

$$\forall c, d > 0 : H_{u^c}^\infty(\mathbb{C}) \subseteq H_{w^d}^\infty(\mathbb{C}),$$

with continuous inclusion.

- (iii) (6.18) holds true, i.e.

$$\exists c \in \mathbb{N}_{>0} \forall d > 0 : H_{u^c}^\infty(\mathbb{C}) \subseteq H_{w^d}^\infty(\mathbb{C}),$$

with continuous inclusion.

- (iv) We have

$$H_{\mathcal{U}^c}^\infty(\mathbb{C}) \subseteq H_{\mathcal{W}^c}^\infty(\mathbb{C}),$$

with continuous inclusion.

If w is also convex, then the list of equivalences can be extended by:

- (v) The weights are satisfying $v_{M^u} \triangleleft^\epsilon v_{M^w}$.
- (vi) The associated weight sequences are satisfying (6.4), i.e.

$$\forall d \in \mathbb{N}_{>0} \exists C \geq 1 \forall j \in \mathbb{N} : \widetilde{M}_j^w \leq CM_j^u.$$

Proof. The equivalences of the first four assertions hold by Propositions 6.14 and 6.15.

(i) \Leftrightarrow (v) is valid by the second part in (2.24), i.e. $v_{M^u} \sim^\epsilon u$, $v_{M^w} \sim^\epsilon w$. (For this convexity is required for *both* weights, see the proof of Theorem 3.32.)

(v) \Leftrightarrow (vi) is (iv) \Leftrightarrow (v) in Proposition 6.8. \square

7. STABILITY UNDER POINT-WISE MULTIPLICATION

The aim of this section is to characterize the stability of weighted classes of entire functions under the point-wise multiplication of functions. In the following consider the (bilinear) multiplication operator

$$\mathfrak{m} : (f, g) \mapsto f \cdot g.$$

7.1. Preliminaries. We start with the following observation:

Proposition 7.1. *Let v be a (normalized) weight function. Then both classes $H_{\underline{v}^c}^\infty(\mathbb{C})$ and $H_{\overline{v}^c}^\infty(\mathbb{C})$ defined in terms of exponential-type weight systems (see (2.5)) are automatically closed under the action of \mathfrak{m} and*

$$\mathfrak{m} : H_{\underline{v}^c}^\infty(\mathbb{C}) \times H_{\underline{v}^c}^\infty(\mathbb{C}) \rightarrow H_{\underline{v}^c}^\infty(\mathbb{C}), \quad \text{resp.} \quad \mathfrak{m} : H_{\overline{v}^c}^\infty(\mathbb{C}) \times H_{\overline{v}^c}^\infty(\mathbb{C}) \rightarrow H_{\overline{v}^c}^\infty(\mathbb{C}),$$

is continuous.

The analogous statement is valid for classes with o -growth condition.

Proof. Let $f \in H_{v^c}^\infty(\mathbb{C})$, $g \in H_{v^d}^\infty(\mathbb{C})$ for some $c, d > 0$. Then $f \cdot g \in H_{v^{c+d}}^\infty(\mathbb{C})$. \square

On the other hand we have the following statement which has been sent to the author in a private communication by Prof. José Bonet.

Proposition 7.2. *Let v be an essential weight. Then $H_v^\infty(\mathbb{C})$ is not closed under point-wise multiplication.*

Proof. Suppose that $\mathbf{m} : H_v^\infty(\mathbb{C}) \rightarrow H_v^\infty(\mathbb{C})$ is continuous, so

$$\exists C > 0 \forall f \in H_v^\infty(\mathbb{C}) : \|f^2\|_v \leq C\|f\|_v.$$

Since v is essential we have the following: Given an arbitrary $x \in \mathbb{C}$ we find $g \in H_v^\infty(\mathbb{C})$ such that $|g(z)| \leq \frac{1}{v(z)}$ for all $z \in \mathbb{C}$ and $g(x) = \frac{1}{v(x)}$, i.e. $\|g\|_v = 1$.

Then

$$\frac{1}{v(x)} \leq v(x)|g(x)|^2 \leq \|g^2\|_v \leq C\|g\|_v = C,$$

and because x is arbitrary we conclude $\frac{1}{C} \leq v(x)$ for all $x \in \mathbb{C}$.

Let $f \in H_v^\infty(\mathbb{C})$ be arbitrary and so there exists $M > 0$ such that $v(z)|f(z)| \leq M$ for all $z \in \mathbb{C}$. By the previous estimate $|f(z)| \leq MC$ for all $z \in \mathbb{C}$ follows and since f is entire this boundedness implies that f is constant, a contradiction. (Recall that since v is rapidly decreasing at least all polynomials are contained in $H_v^\infty(\mathbb{C})$.) \square

The last argument in the proof before shall be compared with the characterization [4, Prop. 2.1] for the unit disc case.

7.2. Convolved weight sequences. Let $M, N \in \mathbb{R}_{>0}^{\mathbb{N}}$ be given, then define the *convolved sequence* $M \star N$ by

$$(7.1) \quad M \star N_j := \min_{0 \leq k \leq j} M_k N_{j-k}, \quad j \in \mathbb{N},$$

see [17, (3.15)]. Hence, obviously $M \star N = N \star M$ and $M \star N_0 = N_0 = 1$ provided that $M_0 = N_0 = 1$.

Remark 7.3. In [17, Lemma 3.5] the following is shown for log-convex weight sequences M, N (resp. $M, N \in \mathcal{LC}$):

- (*) $M \star N$ is also a log-convex weight sequence (resp. $M \star N \in \mathcal{LC}$).
- (*) The corresponding quotient sequence $\mu \star \nu$ is obtained when rearranging resp. ordering the sequences μ and ν in the order of growth.
- (*) This identity yields by definition

$$\forall t \geq 0 : \Sigma_{M \star N}(t) = \Sigma_M(t) + \Sigma_N(t),$$

and so by (2.12)

$$\forall t \geq 0 : \omega_{M \star N}(t) = \omega_M(t) + \omega_N(t).$$

- (*) Thus, by recalling (2.14), we have

$$(7.2) \quad \forall c > 0 \forall t \geq 0 : v_{M \star N, c}(t) = \exp(-\omega_{M \star N}(ct)) = \exp(-\omega_M(ct) - \omega_N(ct)) = v_{M, c}(t)v_{N, c}(t).$$

Remark 7.4. Let $M, N \in \mathbb{R}_{>0}^{\mathbb{N}}$, we summarize more properties for $M \star N$.

- (i) For any $c > 0$ by definition we have $M^c \star N^c = (M \star N)^c$, see (a) in Remark 2.10.
- (ii) For all $j \in \mathbb{N}$ we have $M \star N_j \leq \min\{M_0 N_j, N_0 M_j\}$. So if in addition $M_0 = N_0 = 1$, then we get $M \star N \leq \min\{M, N\}$.

On the other hand $M \preceq M \star M$ amounts to requiring (mg) (and $N \preceq M \star M$ yields the mixed version of (mg) between N and M).

Summarizing, for any $M \in \mathbb{R}_{>0}^{\mathbb{N}}$ we have that M and $M \star M$ are equivalent if and only if M satisfies (mg).

(iii) If M is a log-convex weight sequence, then

$$(7.3) \quad \forall t \geq 0: \quad \omega_M(t) \leq 2\omega_M(t) = \omega_{M \star M}(t),$$

i.e.

$$\forall c > 0 \forall t \geq 0: \quad v_{M,c}(t)^2 = v_{M \star M,c}(t) \leq v_{M,c}(t).$$

And M satisfies (mg), equivalently $M \preceq M \star M$, if and only if

$$(7.4) \quad \exists H \geq 1 \forall t \geq 0: \quad \omega_{M \star M}(t) = 2\omega_M(t) \leq \omega_M(Ht) + H,$$

i.e. (ω_6) for ω_M , see [17, Prop. 3.6]. In this situation, (7.2) turns into

$$(7.5) \quad \exists H \geq 1 \forall c > 0 \forall t \geq 0: \quad v_{M,cH}(t) \leq e^H v_{M \star M,c}(t) = e^H v_{M,c}(t)^2 \leq e^H v_{M,c}(t).$$

7.3. The weight sequence setting. Let M, N be log-convex weight sequences and $f \in H_{v_{M,c_1}}^\infty(\mathbb{C})$, $g \in H_{v_{N,c_2}}^\infty(\mathbb{C})$. Then by Remark 7.3 we get

$$\begin{aligned} \sup_{z \in \mathbb{C}} |(f \cdot g)(z)| &= \sup_{z \in \mathbb{C}} |f(z)| \cdot |g(z)| \leq AB \exp(\omega_M(c_1|z|) + \omega_N(c_2|z|)) \\ &\leq C \exp(\omega_M(d|z|) + \omega_N(d|z|)) = C \exp(\omega_{M \star N}(d|z|)), \end{aligned}$$

when taking $C := AB$ and $d := \max\{c_1, c_2\}$. Thus, the bilinear operator

$$(7.6) \quad \mathfrak{m} : H_{v_{M,c_1}}^\infty(\mathbb{C}) \times H_{v_{N,c_2}}^\infty(\mathbb{C}) \longrightarrow H_{v_{M \star N, \max\{c_1, c_2\}}}^\infty(\mathbb{C}), \quad (f, g) \mapsto f \cdot g,$$

is well-defined and continuous. In particular, take $M = N$ and then by (7.5) we get the following result:

Proposition 7.5. *Let M be a log-convex weight sequence satisfying (mg). Then*

$$\exists H \geq 1 \forall c_1, c_2 > 0 \forall f \in H_{v_{M,c_1}}^\infty(\mathbb{C}), g \in H_{v_{M,c_2}}^\infty(\mathbb{C}): \quad f \cdot g \in H_{v_{M,H \max\{c_1, c_2\}}}^\infty(\mathbb{C}),$$

with H being the constant appearing in (ω_6) for ω_M .

Consequently, both spaces $H_{\underline{M}_c}^\infty(\mathbb{C})$ and $H_{\overline{M}_c}^\infty(\mathbb{C})$ are closed under point-wise multiplication and the multiplication operator \mathfrak{m} is continuously acting on these spaces.

The next result deals with the converse implication.

Proposition 7.6. *Let M, N, L be log-convex weight sequences.*

(i) *Assume that*

$$\exists c_1, c_2, d > 0 \forall f \in H_{v_{M,c_1}}^\infty(\mathbb{C}), g \in H_{v_{N,c_2}}^\infty(\mathbb{C}): \quad f \cdot g \in H_{v_{L,d}}^\infty(\mathbb{C}).$$

Then the sequences satisfy

$$L \preceq M \star N.$$

(ii) *If (i) holds for $M \equiv N \equiv L$, then M has to satisfy (mg).*

Alternatively, in order to conclude in the assumption we can replace some or all \exists by \forall .

Proof. (i) Let $c_1, c_2, d > 0$ be given and by assumption for any $f \in H_{v_{M,c_1}}^\infty(\mathbb{C})$, $g \in H_{v_{N,c_2}}^\infty(\mathbb{C})$ we get

$$\exists C \geq 1 \forall z \in \mathbb{C}: \quad |f(z) \cdot g(z)| \leq C \exp(\omega_L(d|z|)).$$

We apply this estimate to $f \equiv \theta_{M,c_1}$ and $g \equiv \theta_{N,c_2}$ from Lemma 3.9 and so

$$\forall t \geq 0: \quad \exp(\omega_M(c_1 t/2)) \exp(\omega_N(c_2 t/2)) \leq |\theta_{M,c_1}(t) \cdot \theta_{N,c_2}(t)| \leq C \exp(\omega_L(dt)).$$

Take $c := \min\{c_1, c_2\}$ and then Remark 7.3 gives

$$\omega_{M \star N}(ct/2) = \omega_M(ct/2) + \omega_N(ct/2) \leq \omega_M(c_1 t/2) + \omega_N(c_2 t/2) \leq \log(C) + \omega_L(dt).$$

Since $M \star N$ is also a log-convex weight sequence (see again Remark 7.3) we can apply (2.11) and get for all $j \in \mathbb{N}$

$$\begin{aligned} L_j &= \sup_{t \geq 0} \frac{t^j}{\exp(\omega_L(t))} = \sup_{s \geq 0} \frac{(ds)^j}{\exp(\omega_L(ds))} \leq Cd^j \sup_{s \geq 0} \frac{s^j}{\exp(\omega_{M \star N}(cs/2))} \\ &= C \left(\frac{2d}{c} \right)^j \sup_{u \geq 0} \frac{u^j}{\exp(\omega_{M \star N}(u))} = C \left(\frac{2d}{c} \right)^j M \star N_j, \end{aligned}$$

i.e. $L \preceq M \star N$.

(ii) If we take $M \equiv N \equiv L$ in (i), then $M \preceq M \star M$ and this precisely means (mg) for M , see (ii) in Remark 7.4. \square

For the class $H_{\mathcal{M}_c}^\infty(\mathbb{C})$ we cannot make use of the optimal functions $\theta_{M,c}$.

Proposition 7.7. *Let M be a log-convex weight sequence. Assume that $H_{\mathcal{M}_c}^\infty(\mathbb{C})$ is closed under point-wise multiplication and that \mathfrak{m} is continuously acting on this space. Then M has to satisfy (mg).*

Proof. The proof is inspired by [13, Prop. 4.7]. By assumption

$$\mathfrak{m}_\Delta : H_{\mathcal{M}_c}^\infty(\mathbb{C}) \rightarrow H_{\mathcal{M}_c}^\infty(\mathbb{C}), \quad f \mapsto f^2,$$

is continuous. This amounts to $H_{\mathcal{M}_c}^\infty(\mathbb{C}) \subseteq H_{\mathcal{M}_c}^{2,\infty}(\mathbb{C})$ for the Fréchet space

$$H_{\mathcal{M}_c}^{2,\infty}(\mathbb{C}) := \{f \in H(\mathbb{C}) : \|f^2\|_{v_{M,c}} < +\infty, \forall c > 0\}.$$

By the closed graph theorem the inclusion $H_{\mathcal{M}_c}^\infty(\mathbb{C}) \subseteq H_{\mathcal{M}_c}^{2,\infty}(\mathbb{C})$ is continuous and hence

$$(7.7) \quad \forall d > 0 \exists c > 0 \exists C \geq 1 \forall f \in H_{\mathcal{M}_c}^\infty(\mathbb{C}) : \|f^2\|_{v_{M,d}} \leq C \|f\|_{v_{M,c}}.$$

We apply this estimate to the monomials $f_k = z^k$ and get:

$$\forall d > 0 \exists c > 0 \exists C \geq 1 \forall k \in \mathbb{N} \forall z \in \mathbb{C} : \frac{|z|^{2k}}{\exp(\omega_M(d|z|))} \leq C \frac{|z|^k}{\exp(\omega_M(c|z|))}.$$

(2.11) gives for all $j \in \mathbb{N}$ that

$$\begin{aligned} M_{2j} &= \sup_{t \geq 0} \frac{t^{2j}}{\exp(\omega_M(t))} = \sup_{s \geq 0} \frac{(ds)^{2j}}{\exp(\omega_M(ds))} \leq Cd^{2j} \sup_{s \geq 0} \frac{s^j}{\exp(\omega_M(cs))} \\ &= C \left(\frac{d^2}{c} \right)^j \sup_{u \geq 0} \frac{u^j}{\exp(\omega_M(u))} = C \left(\frac{d^2}{c} \right)^j M_j. \end{aligned}$$

This verifies (5.2) and so M has to satisfy (mg). \square

Thus, by combining Propositions 7.5, 7.6 and 7.7 we arrive at the following characterization:

Theorem 7.8. *Let M be a log-convex weight sequence. Then the following are equivalent:*

- (i) $H_{\mathcal{M}_c}^\infty(\mathbb{C})$ is closed under point-wise multiplication and \mathfrak{m} is continuously acting on this space.
- (ii) $H_{\mathcal{M}_c}^\infty(\mathbb{C})$ is closed under point-wise multiplication and \mathfrak{m} is continuously acting on this space.
- (iii) M has (mg).

We close this Section with the following observation.

Remark 7.9. If M is violating (mg), then one is able to infer a mixed version of Theorem 7.8: When taking $M = N$ in (7.6) we see that both $\mathfrak{m} : H_{\underline{M}_c}^\infty(\mathbb{C}) \times H_{\underline{M}_c}^\infty(\mathbb{C}) \rightarrow H_{\underline{L}_c}^\infty(\mathbb{C})$ and $\mathfrak{m} : H_{\overline{M}_c}^\infty(\mathbb{C}) \times H_{\overline{M}_c}^\infty(\mathbb{C}) \rightarrow H_{\overline{L}_c}^\infty(\mathbb{C})$ are continuous with $L := M \star M$.

On the other hand, the choice $M = N$ in (i) in Proposition 7.6 gives $L \preceq M \star M$, i.e. the mixed version of (mg) between M and L . Similarly, when assuming that $\mathfrak{m} : H_{\overline{M}_c}^\infty(\mathbb{C}) \times H_{\overline{M}_c}^\infty(\mathbb{C}) \rightarrow H_{\overline{L}_c}^\infty(\mathbb{C})$ is continuous, then the proof of Proposition 7.7 gives the mixed version of (5.2) and [21, Thm. 9.5.1] implies again $L \preceq M \star M$.

In both cases $L \preceq M \star M$ yields $H_{\underline{M \star M}_c}^\infty(\mathbb{C}) \subseteq H_{\underline{L}_c}^\infty(\mathbb{C})$ and $H_{\overline{M \star M}_c}^\infty(\mathbb{C}) \subseteq H_{\overline{L}_c}^\infty(\mathbb{C})$, recall Proposition 3.8.

7.4. The weight function setting. Let u and v be weight functions and $f \in H_{u_{c_1}}^\infty(\mathbb{C})$, $g \in H_{v_{c_2}}^\infty(\mathbb{C})$. Then we get $f \cdot g \in H_{w_d}^\infty(\mathbb{C})$ with $d := \max\{c_1, c_2\}$ and $w := u \cdot v$ by the same estimate as in Section 7.3 above. Thus, the bilinear operator

$$\mathfrak{m} : H_{u_{c_1}}^\infty(\mathbb{C}) \times H_{v_{c_2}}^\infty(\mathbb{C}) \longrightarrow H_{w_{\max\{c_1, c_2\}}}^\infty(\mathbb{C}), \quad (f, g) \mapsto f \cdot g,$$

is well-defined and continuous.

When taking $u \equiv v$, then we get the following result which is analogous to Proposition 7.5.

Proposition 7.10. *Let v be a (normalized) weight function of moderate growth. Then*

$$\exists H \geq 1 \forall c_1, c_2 > 0 \forall f \in H_{v_{c_1}}^\infty(\mathbb{C}), g \in H_{v_{c_2}}^\infty(\mathbb{C}) : f \cdot g \in H_{v_{H \max\{c_1, c_2\}}}^\infty(\mathbb{C}),$$

and H can be chosen to be the constant appearing in (2.28).

Proof. Let $f \in H_{v_{c_1}}^\infty(\mathbb{C})$, $g \in H_{v_{c_2}}^\infty(\mathbb{C})$ be given and so $f \cdot g \in H_{v_c \cdot v_c}^\infty(\mathbb{C})$ with $c := \max\{c_1, c_2\}$. By (2.28) we find $v(Ht) \leq e^H v(t)^2$ for some $H \geq 1$ and all $t \geq 0$, hence $f \cdot g \in H_{v_{Hc}}^\infty(\mathbb{C})$. \square

The next result deals with the converse implication and is analogous to Proposition 7.6; we reduce its proof to the weight sequence setting.

Proposition 7.11. *Let r, u, w be normalized weights and let r and u be convex.*

(i) *Assume that*

$$\exists c_1, c_2, d > 0 \forall f \in H_{r_{c_1}}^\infty(\mathbb{C}), g \in H_{u_{c_2}}^\infty(\mathbb{C}) : f \cdot g \in H_{w_d}^\infty(\mathbb{C}).$$

Then the weights are satisfying

$$\exists A, B \geq 1 \forall t \geq 0 : (w(At))^2 \leq Br(t)u(t).$$

(ii) *If (i) holds for $r \equiv u \equiv w$, then u has to be of moderate growth.*

Alternatively, in order to conclude in (i) we can replace some or all \exists by \forall .

Proof. (i) Let $c_1, c_2, d > 0$ be the parameters. The second estimate in (2.23) and the assumption imply that for any $f \in H_{v_{M^r, c_1}}^\infty(\mathbb{C})$, $g \in H_{v_{M^u, c_2}}^\infty(\mathbb{C})$ we get $f \cdot g \in H_{w_d}^\infty(\mathbb{C})$.

We apply this to $f \equiv \theta_{M^r, c_1}$ and $g \equiv \theta_{M^u, c_2}$ from Lemma 3.9 and restrict to $t \geq 0$ in order to get

$$\exists C \geq 1 \forall t \geq 0 : \omega_{M^r}(c_1 t/2) + \omega_{M^u}(c_2 t/2) \leq \log(C) - \log(w(dt)) = \log(C) + \omega^w(dt),$$

recall (2.19) for the last equality. Set $c := \min\{c_1, c_2\}$ and thus we have

$$\omega_{M^r}(c_1 t/2) + \omega_{M^u}(c_2 t/2) \geq \omega_{M^r}(ct/2) + \omega_{M^u}(ct/2) = \omega_{M^r \star M^u}(ct/2),$$

see again Remark 7.3. Similarly like in the proof of Proposition 7.6 and by taking into account (2.22) (for M^w) we get for all $j \in \mathbb{N}$:

$$\begin{aligned} M_j^w &= \sup_{t>0} \frac{t^j}{\exp(\omega^w(t))} = \sup_{s>0} \frac{(ds)^j}{\exp(\omega^w(ds))} \leq Cd^j \sup_{s>0} \frac{s^j}{\exp(\omega_{M^r \star M^u}(cs/2))} \\ &= C \left(\frac{2d}{c} \right)^j M^r \star M_j^u. \end{aligned}$$

Hence $M^w \preceq M^r \star M^u$ is verified and by Remarks 7.3 and 7.4 this relation yields

$$\exists H \geq 1 \forall t \geq 0: \quad \omega_{M^r}(t) + \omega_{M^u}(t) \leq \omega_{M^w}(Ht) + H \Leftrightarrow v_{M^w, H}(t) \leq e^H v_{M^r}(t) v_{M^u}(t).$$

We use this, the first estimate in (2.23) applied to r and u (for which convexity is required) and the second one applied to w and get

$$\exists A \geq 1 \exists H \geq 1 \forall t \geq 0: \quad w(Ht) \leq v_{M^w}(Ht) \leq e^H v_{M^r}(t) v_{M^u}(t) \leq Ae^H \sqrt{r(t)u(t)}.$$

(ii) When $r \equiv u \equiv w$, then following the proof in (i) we get $M^u \preceq M^u \star M^u$ and so M^u satisfies (mg) by Remark 7.4. Thus, by Lemma 2.17 we get (2.28) for u . \square

Proposition 7.12. *Let u be a normalized and convex weight function. Assume that $H_{\mathcal{U}_c}^\infty(\mathbb{C})$ is closed under point-wise multiplication and that \mathfrak{m} is continuously acting on this space. Then u satisfies (2.28), i.e. u is of moderate growth.*

Proof. Analogously, as in the proof of Proposition 7.7 we get $H_{\mathcal{U}_c}^\infty(\mathbb{C}) \subseteq H_{\mathcal{U}_c}^{2,\infty}(\mathbb{C})$ for the Fréchet space

$$H_{\mathcal{U}_c}^{2,\infty}(\mathbb{C}) := \{f \in H(\mathbb{C}) : \|f^2\|_{u_c} < +\infty, \forall c > 0\}.$$

By the closed graph theorem the inclusion $H_{\mathcal{U}_c}^\infty(\mathbb{C}) \subseteq H_{\mathcal{U}_c}^{2,\infty}(\mathbb{C})$ is continuous and hence

$$\forall d > 0 \exists c > 0 \exists C \geq 1 \forall f \in H_{\mathcal{U}_c}^\infty(\mathbb{C}): \quad \|f^2\|_{u_d} \leq C \|f\|_{u_c}.$$

We apply this estimate to the monomials $f_k = z^k$ and get (see (2.19)):

$$\forall d > 0 \exists c > 0 \exists C \geq 1 \forall k \in \mathbb{N} \forall z \in \mathbb{C}:$$

$$\frac{|z|^{2k}}{\exp(\omega^u(d|z|))} = |z|^{2k} u(d|z|) \leq C |z|^{2k} u(c|z|) = C \frac{|z|^{2k}}{\exp(\omega^u(c|z|))}.$$

Then, we use (2.22) and follow the proof of Proposition 7.7 to get that M^u satisfies (5.2) and thus (mg). Consequently, by Lemma 2.17 we get (2.28) for u . \square

Combining Lemma 2.17 and Propositions 7.10, 7.11 and 7.12 we arrive at the following characterization.

Theorem 7.13. *Let u be a normalized and convex weight function. Then the following are equivalent:*

- (i) $H_{\mathcal{U}_c}^\infty(\mathbb{C})$ is closed under point-wise multiplication and \mathfrak{m} is continuously acting on this space.
- (ii) $H_{\mathcal{U}_c}^{2,\infty}(\mathbb{C})$ is closed under point-wise multiplication and \mathfrak{m} is continuously acting on this space.
- (iii) u is of moderate growth, i.e. it satisfies (2.28).
- (iv) The associated sequence M^u satisfies (mg).

Note that the characterization Theorem 7.8 resp. Theorem 7.13 is consistent with Proposition 7.1 and Theorem 5.4 resp. Theorem 5.6.

Remark 7.14. *By Theorem 4.1 (and the comments after this statement) we get: Theorem 7.8 transfers directly to the corresponding o -growth setting and if the characterization established in Theorem 7.13 is valid, then the analogue of (i) and (ii) holds, i.e. $H_{\underline{u}_c}^o(\mathbb{C})$ and $H_{\overline{u}_c}^o(\mathbb{C})$ are closed under point-wise multiplication and \mathfrak{m} is continuously acting on these spaces.*

We close by showing that the proofs of Proposition 7.7 and 7.12 give an independent argument for the conclusion of Proposition 7.2; even under more general assumptions on the weight.

Corollary 7.15. *Let $M \in \mathcal{LC}$ resp. a normalized weight u be given. Then $H_{v_M}^\infty(\mathbb{C})$ resp. $H_u^\infty(\mathbb{C})$ is not closed under point-wise multiplication.*

Proof. For the weight sequence case we present two proofs: First, by Theorem 3.2 we know that v_M is essential and thus Proposition 7.2 yields the conclusion.

Second, given $M \in \mathcal{LC}$ let us assume that $H_{v_M}^\infty(\mathbb{C})$ is closed under point-wise multiplication. Then follow the arguments in the proof of Proposition 7.7 with $d = c = 1$ and thus get

$$(7.8) \quad \exists C \geq 1 \forall j \in \mathbb{N}: M_{2j} \leq CM_j.$$

But this violates $\lim_{j \rightarrow +\infty} (M_j)^{1/j} = +\infty$: Let $k \in \mathbb{N}$ be such that $2^n \leq k < 2^{n+1}$ for some $n \in \mathbb{N}_{>0}$, then $M_k \leq M_{2^{n+1}} \leq C^{n+1} M_1$ and so $(M_k)^{1/k} \leq M_1^{1/k} C^{(n+1)/k} \leq M_1^{1/k} C^{(n+1)/2^n} \leq M_1 C < +\infty$. Now given a normalized weight u and assuming that $H_u^\infty(\mathbb{C})$ is closed under point-wise multiplication, then the proof of Proposition 7.12 with $d = c = 1$ gives (7.8) for M^u . However, since $M^u \in \mathcal{LC}$ this is impossible by the arguments just given before. \square

Note:

- (*) For the conclusion convexity for u is not required since in Proposition 7.12 convexity was only needed in order to apply Lemma 2.17 in the very last step which is not needed in Corollary 7.15.
- (*) For the contradictory argument we have used that $\lim_{j \rightarrow +\infty} (M_j^u)^{1/j} = +\infty$. In view of Remark 2.14 we know that this argument would break down if $u(t) = 0$ for all t sufficiently large. But then as sets $H_u^\infty(\mathbb{C}) = H(\mathbb{C})$ and $H(\mathbb{C})$ is clearly closed under point-wise multiplication.
- (*) The difference between the failure in the single weight case and the characterization in the weight system case is that the latter situation allows for having "enough place" when changing the parameter, see the proof of Proposition 7.1 and note that $H > 1$ in (2.28).
- (*) The characterization [4, Prop. 2.1] suggests that Corollary 7.15 fails for classes $H_{v_M}^\infty(\mathbb{D})$ resp. $H_u^\infty(\mathbb{D})$. Note that by [22, Sect. 7], in the weight sequence setting, for $G = \mathbb{D}$ the weights are given by $t \mapsto \exp(-\omega_M(\frac{1}{1-t})) =: v_{M,\mathbb{D}}(t)$ and hence the corresponding norms are expressed in terms of $v_{M,\mathbb{D}}$. Consequently, when following the proof of Proposition 7.7 we arrive in (7.7) (again with $c = d = 1$) at expressions which are not well-behaved w.r.t. the crucial formula (2.11).

7.5. On the point-wise product of weight sequences. Let M, N be log-convex weight sequences ($M, N \in \mathcal{LC}$) and set $M \cdot N := (M_j \cdot N_j)_{j \in \mathbb{N}}$. Thus $M \cdot M$ is again a log-convex weight sequence ($M \cdot N \in \mathcal{LC}$) and one may ask how this operation is related to weighted classes of entire functions.

In [17, Prop. 4.5] the following result has been shown which characterizes the growth of the Taylor-coefficients of entire functions (belonging to $H_{\underline{M}_c}^\infty(\mathbb{C})$, $H_{\overline{M}_c}^\infty(\mathbb{C})$).

Proposition 7.16. *Let M be a log-convex weight sequence and let $P(\xi) = \sum_{j=0}^{+\infty} a_j \xi^j$ be an entire function.*

- (i) *The following are equivalent ("Roumieu-type variant"):*
- (a) $\exists L, C > 0 \forall j \in \mathbb{N}: |a_j| \leq \frac{CL^j}{M_j}$,
 - (b) $\exists L, C > 0 \forall \xi \in \mathbb{C}: |P(\xi)| \leq C \exp(\omega_M(L|\xi|))$, i.e. $P \in H_{\underline{M}_c}^\infty(\mathbb{C})$.
- (ii) *The following are equivalent ("Beurling-type variant"):*
- (a) $\forall L > 0 \exists C > 0 \forall j \in \mathbb{N}: |a_j| \leq \frac{CL^j}{M_j}$,
 - (b) $\forall L > 0 \exists C > 0 \forall \xi \in \mathbb{C}: |P(\xi)| \leq C \exp(\omega_M(L|\xi|))$, i.e. $P \in H_{\overline{M}_c}^\infty(\mathbb{C})$.

The proof shows that in (a) \Rightarrow (b) we have to replace L by $2L$, in (b) \Rightarrow (a) we can take the same constant L (since we are considering the case dimension $d = 1$).

Let $f(z) = \sum_{j \geq 0} a_j z^j$, $g(z) = \sum_{j \geq 0} b_j z^j$ be two entire functions, then set

$$f \odot g(z) := \sum_{j \geq 0} a_j b_j z^j.$$

Theorem 7.17. *Let M, N be log-convex weight sequences, then*

$$\forall c, d, h > 0 \forall f \in H_{v_{M,c}}^\infty(\mathbb{C}), g \in H_{v_{N,d}}^\infty(\mathbb{C}): f \odot g \in H_{v_{M \cdot N, 2cd}}^\infty(\mathbb{C}) \subseteq H_{v_{M,h}}^\infty(\mathbb{C}), H_{v_{N,h}}^\infty(\mathbb{C}).$$

Consequently, for any log-convex M and each $c > 0$ the class $H_{v_{M,c}}^\infty(\mathbb{C})$ is closed under \odot which is acting as a continuous operator on this space.

In particular, both $H_{\underline{M}_c}^\infty(\mathbb{C})$ and $H_{\overline{M}_c}^\infty(\mathbb{C})$ are closed under \odot which is acting as a continuous operator on these classes.

Proof. Let $c, d, h > 0$ be arbitrary. If $f \in H_{v_{M,c}}^\infty(\mathbb{C})$, $g \in H_{v_{N,d}}^\infty(\mathbb{C})$, then (b) \Rightarrow (a) in Proposition 7.16 implies

$$(7.9) \quad \exists C_1, C_2 \geq 1 \forall j \in \mathbb{N}: |a_j b_j| \leq C_1 C_2 \frac{(cd)^j}{M_j N_j},$$

hence (a) \Rightarrow (b) in Proposition 7.16 yields $f \odot g \in H_{v_{M \cdot N, 2cd}}^\infty(\mathbb{C})$. Thus

$$\odot: H_{v_{M,c}}^\infty(\mathbb{C}) \times H_{v_{N,d}}^\infty(\mathbb{C}) \rightarrow H_{v_{M \cdot N, 2cd}}^\infty(\mathbb{C}),$$

is a continuous operator. But moreover, since $\lim_{j \rightarrow +\infty} (M_j)^{1/j} = \lim_{j \rightarrow +\infty} (N_j)^{1/j} = +\infty$ we have

$$\forall c, d, h > 0 \exists A \geq 1 \forall j \in \mathbb{N}: \frac{(cd)^j}{M_j N_j} \leq A \min \left\{ \frac{(h/2)^j}{M_j}, \frac{(h/2)^j}{N_j} \right\}.$$

Combining this with (7.9) and (a) \Rightarrow (b) in Proposition 7.16 we get $f \odot g \in H_{v_{M,h}}^\infty(\mathbb{C}), H_{v_{N,h}}^\infty(\mathbb{C})$. \square

Corollary 7.18. *Let u be a normalized and convex weight of moderate growth. Then both $H_{\underline{u}_c}^\infty(\mathbb{C})$ and $H_{\overline{u}_c}^\infty(\mathbb{C})$ are closed under \odot which is acting as a continuous operator on these spaces.*

Proof. We use Proposition 2.19 (see the first part in (2.25)) and apply Theorem 7.17 to the associated sequence M^u . \square

For the sake of completeness we mention that in [2, Lemma 4] it has been shown that

$$\exp(\omega_{M \cdot N}(t)) = \inf_{s > 0} \{ \exp(\omega_M(s)) \cdot \exp(\omega_N(t/s)) \} = \inf_{s > 0} \{ \exp(\omega_M(t/s)) \cdot \exp(\omega_N(s)) \}.$$

REFERENCES

- [1] K.-D. Bierstedt, J. Bonet, and J. Taskinen. Associated weights and spaces of holomorphic functions. *Studia Math.*, 127(2):137–168, 1998.
- [2] J. Boman. Uniqueness and non-uniqueness for microanalytic continuation of ultradistributions. *Contemporary Mathematics*, 251:61–82, 1998.
- [3] J. Bonet. Weighted Banach spaces of analytic functions with sup-norms and operators between them: a survey. *Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM*, (116):art. no. 184, 2022.
- [4] J. Bonet, P. Domański, and M. Lindström. Pointwise multiplication operators on weighted Banach spaces of analytic functions. *Studia Math.*, 137(2):177–194, 1999.
- [5] J. Bonet, W. Lusky, and J. Taskinen. Solid hulls and cores of weighted H^∞ -spaces. *Rev. Mat. Complut.*, 31(3):781–804, 2019.
- [6] J. Bonet, R. Meise, and S. N. Melikhov. A comparison of two different ways to define classes of ultradifferentiable functions. *Bull. Belg. Math. Soc. Simon Stevin*, 14:424–444, 2007.
- [7] J. Bonet and J. Taskinen. Solid hulls of weighted Banach spaces of entire functions. *Rev. Mat. Iberoam.*, 34(2):593–608, 2018.
- [8] R. W. Braun, R. Meise, and B. A. Taylor. Ultradifferentiable functions and Fourier analysis. *Results Math.*, 17(3-4):206–237, 1990.
- [9] J. Clunie. On integral functions having prescribed asymptotic growth. *Canad. J. Math.*, 17:396–404, 1965.
- [10] Abakumov E. and E. Doubtsov. Moduli of holomorphic functions and logarithmically convex radial weights. *Bull. Lond. Math. Soc.*, 47(3):519–532, 2015.
- [11] Abakumov E. and E. Doubtsov. Approximation by proper holomorphic maps and tropical power series. *Constr. Approx.*, 47(2):321–338, 2018.
- [12] P. Erdős and T. Kövári. On the maximum modulus of entire functions. *Acta Math. Acad. Sci. Hungar.*, (7):305–317, 1956.
- [13] C. Esser and G. Schindl. On the construction of large algebras not contained in the image of the Borel map. *Res. Math.*, 75(22), 2020.
- [14] J. Jiménez-Garrido, D. N. Nenning, and G. Schindl. On generalized definitions of ultradifferentiable classes. 2022, available online at <https://arxiv.org/pdf/2211.08090.pdf>.
- [15] J. Jiménez-Garrido, J. Sanz, and G. Schindl. Sectorial extensions, via Laplace transforms, in ultraholomorphic classes defined by weight functions. *Results Math.*, 74(27), 2019.
- [16] J. Jiménez-Garrido, J. Sanz, and G. Schindl. Sectorial extensions for ultraholomorphic classes defined by weight functions. *Math. Nachr.*, 293(11):2140–2174, 2020.
- [17] H. Komatsu. Ultradistributions. I. Structure theorems and a characterization. *J. Fac. Sci. Univ. Tokyo Sect. IA Math.*, 20:25–105, 1973.
- [18] S. Mandelbrojt. *Séries adhérentes, Régularisation des suites, Applications*. Gauthier-Villars, Paris, 1952.
- [19] W. Matsumoto. Characterization of the separativity of ultradifferentiable classes. *J. Math. Kyoto Univ.*, 24(4):667–678, 1984.
- [20] A. Rainer and G. Schindl. Composition in ultradifferentiable classes. *Studia Math.*, 224(2):97–131, 2014.
- [21] G. Schindl. Exponential laws for classes of Denjoy-Carleman-differentiable mappings, 2014. PhD Thesis, Universität Wien, available online at http://othes.univie.ac.at/32755/1/2014-01-26_0304518.pdf.
- [22] G. Schindl. Solid hulls and cores of classes of weighted entire functions defined in terms of associated weight functions. *Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM*, 114(176), 2020.
- [23] G. Schindl. On subadditivity-like conditions for associated weight functions. *Bull. Belg. Math. Soc. Simon Stevin*, 28(3):399–427, 2022.
- [24] G. Schindl. On the equivalence between moderate growth-type conditions in the weight matrix setting. *Note di Mat.*, 42(1):1–35, 2022.

G. SCHINDL: FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT WIEN, OSKAR-MORGENSTERN-PLATZ 1, A-1090 WIEN, AUSTRIA.

Email address: gerhard.schindl@univie.ac.at