

# ISOCLINISM OF SKEW BRACES

T. LETOURMY AND L. VENDRAMIN

ABSTRACT. We define isoclinism of skew braces and present several applications. We study some properties of skew braces that are invariant under isoclinism. For example, we prove that right nilpotency is an isoclinism invariant. This result has application in the theory of set-theoretic solutions to the Yang–Baxter equation. We define isoclinic solutions and study multipermutation solutions under isoclinism.

## 1. INTRODUCTION

The fundamental problem of constructing combinatorial solutions of the Yang–Baxter equation (YBE) is nowadays based on the use of specific (associative and non-associative) algebraic structures acting on the solution. One particular algebraic structure stands out: skew braces. Such structures were introduced in [13] and [10]. The theory originates in Jacobson radical rings, and it is now considered a hot topic, as skew braces appear in several different areas of mathematics, see for example [7, 15, 20].

Skew braces classify solutions [1, 2]. This justifies the search for classification results on skew braces. In this vein, several results are known; see for example [4, 10]. However, the classification of skew braces of order  $p^n$  ( $p$  prime) is still hard to achieve.

To classify  $p$ -groups, in [11] Hall introduced a specific equivalence relation that simplifies the problem. Without technicalities, one defines isoclinic as groups that have “essentially the same” commutator functions. Recall that the commutator function for the group  $G$  is the map

$$G/Z(G) \times G/Z(G) \rightarrow G, \quad (xZ(G), yZ(G)) \mapsto [x, y] = xyx^{-1}y^{-1}.$$

Isoclinism is an equivalence relation that generalizes isomorphisms.

There are several different motivations to consider this notion in the context of skew braces. With isoclinism of skew braces, we will have a new tool to classify finite skew braces of prime-power order, something that ultimately will have applications in other branches of mathematics, for example in the theory of pre-Lie algebras [17, 18, 19].

In this work, we define isoclinism in the context of skew braces. We present several properties of this notion (e.g. Theorems 2.8, 2.18, 3.10, 3.18 and 4.1). As an application, we define isoclinism of set-theoretic solutions to the YBE. This new equivalence relation defined on the space of solutions suggests a way of attacking the classification problem of set-theoretic solutions to the YBE. Isoclinism classes of solutions will be relevant in the study of multipermutation solutions [3, 5, 8, 9, 12].

---

2020 *Mathematics Subject Classification.* Primary:16T25, Secondary:17D99.

*Key words and phrases.* Skew brace, isoclinism, Yang–Baxter, multipermutation.

We prove in Section 4 that if  $(X, r)$  and  $(Y, s)$  are isoclinic set-theoretic solutions and  $(X, r)$  is multipermutation, then  $(Y, s)$  is multipermutation (see Theorem 4.4).

## 2. ISOCLINISM OF SKEW BRACES

Recall that a *skew brace* is a triple  $(A, +, \circ)$ , where  $(A, +)$  and  $(A, \circ)$  are groups such that the compatibility condition  $a \circ (b + c) = a \circ b - a + a \circ c$  holds for all  $a, b, c \in A$ . The inverse of an element  $a \in A$  with respect to the circle operation  $\circ$  will be denoted by  $a'$ . We will denote the groups  $(A, +)$  and  $(A, \circ)$  respectively by  $A_+$  and  $A_\circ$ .

Let  $A$  be a skew brace. There are two canonical actions by automorphism

$$\begin{aligned} \lambda: A_\circ &\rightarrow \text{Aut}(A_+), & \lambda_a(b) &= -a + a \circ b, \\ \rho: A_\circ &\rightarrow \text{Aut}(A_+), & \rho_a(b) &= a \circ b - a. \end{aligned}$$

By definition, a sub skew brace  $B$  of  $A$  is an ideal of  $A$  if  $B_+$  is normal in  $A_+$ ,  $B_\circ$  is normal in  $A_\circ$  and  $\lambda_a(B) \subseteq B$  for all  $a \in A$ .

We write  $A^{(2)} = A * A$  to denote the additive subgroup of  $A$  generated by  $\{a * b : a, b \in A\}$ , where  $a * b = -a + a \circ b - b$  for all  $a, b \in A$ . One proves that  $A^{(2)}$  is an ideal of  $A$ . If  $a, b \in A$ , we write  $[a, b]_+ = a + b - a - b$  and  $[a, b]_\circ = a \circ b \circ a' \circ b'$ . We write  $[A, A]_+$  to denote the commutator subgroup of the additive group of  $A$ .

**Definition 2.1.** Let  $A$  be a skew brace. The *commutator*  $A'$  of  $A$  is the additive subgroup of  $A$  generated by  $[A, A]_+$  and  $A^{(2)}$ .

**Proposition 2.2.** *Let  $A$  be a skew brace. Then  $A'$  is an ideal of  $A$ .*

*Proof.* Let  $a \in A$  and  $x \in A'$ . Since  $a + x - a = x + [-x, a]_+ \in A'$ , it follows that  $(A', +)$  is a normal subgroup of  $(A, +)$ . Since  $\lambda_a(x) \in A'$  and  $x * a \in A'$ , the claim follows from [6, Lemma 1.9].  $\square$

**Example 2.3.** In the case of a trivial brace  $A$ ,  $A'$  is the commutator of the underlying group.

**Notation 2.4.** *In general, if  $A/B$  is a quotient of a skew brace, we will denote the equivalence class of  $a \in A$  in  $A/B$  by  $\bar{a}$ .*

**Remark 2.5.**  $A'$  is the smallest ideal of  $A$  such that  $A^{\text{ab}} = A/A'$  is an abelian group.

**Lemma 2.6.** *We define two collections of maps  $\phi_+$  and  $\phi_*$  that associate respectively to every skew brace  $B$  the maps*

$$(2.1) \quad \phi_+^B: (B/\text{Ann } B)^2 \rightarrow B', \quad (\bar{a}, \bar{b}) \mapsto [a, b]_+,$$

$$(2.2) \quad \phi_*^B: (B/\text{Ann } B)^2 \rightarrow B', \quad (\bar{a}, \bar{b}) \mapsto a * b.$$

*The maps (2.1) and (2.2) are well-defined.*

*Proof.* A direct calculation shows that (2.1) is well-defined. To prove that (2.2) is well defined, let  $(\bar{a}, \bar{b}) = (\bar{c}, \bar{d}) \in (B/\text{Ann } B)^2$ .

$$-a + a \circ b - b - (-c + c \circ d - d) = -a + a \circ d - c \circ d + c.$$

After conjugating by  $c$ ,

$$\begin{aligned} c - a + a \circ d - c \circ d &= (c - a) \circ a \circ d - c \circ d \\ &= ((c - a) + a) \circ d - c \circ d \\ &= c \circ d - c \circ d \\ &= 0. \end{aligned}$$

Since the conjugation by  $c$  is an automorphism, we have  $a * b = c * d$ .  $\square$

**Definition 2.7.** We say that the braces  $A$  and  $B$  are *isoclinic* if there are two isomorphisms  $\xi: A/\text{Ann } A \rightarrow B/\text{Ann } B$  and  $\theta: A' \rightarrow B'$  such that

$$(2.3) \quad \begin{array}{ccccc} A' & \xleftarrow{\phi_+^A} & (A/\text{Ann } A)^2 & \xrightarrow{\phi_*^A} & A' \\ \downarrow \theta & & \downarrow \xi \times \xi & & \downarrow \theta \\ B' & \xleftarrow{\phi_+^B} & (B/\text{Ann } B)^2 & \xrightarrow{\phi_*^B} & B' \end{array}$$

commutes. We call the pair  $(\xi, \theta)$  a skew brace *isoclinism*.

**Theorem 2.8.** *Let  $A$  and  $B$  be skew braces. If  $A$  and  $B$  are isoclinic then  $A$  is trivial if and only if  $B$  is trivial.*

*Proof.* It is a direct consequence of the commutativity of the right part of the diagram (2.3).  $\square$

**Remark 2.9.** Two trivial skew braces are isoclinic if and only if their underlying groups are isoclinic. Therefore, we will not distinguish between the two notions when dealing with trivial skew braces.

**Remark 2.10.** Let  $A$  and  $B$  be isoclinic skew braces. The quotients  $A/\text{Ann}(A)$  and  $B/\text{Ann}(B)$  are isomorphic. It follows that  $A$  is annihilator nilpotent if and only if  $B$  is annihilator nilpotent.

We now present the notions of terms and term functions from universal algebra in the context of skew braces. A more general approach can be found in §10, and §11 of the book [16].

Let  $X$  be a set of objects called variables. The set of *skew brace terms* over  $X$  is the smallest set  $T(X)$  such that

- (1)  $X \cup \{0\} \subseteq T(X)$ , where  $0$  is a formal element, and
- (2) if  $p_1, p_2 \in T(X)$ , then the ‘‘strings’’  $p_1 \circ p_2, p_1 + p_2, -p_1, p_2' \in T(X)$ .

For  $p \in T(X)$  we write  $p$  as  $p(x_1, \dots, x_n)$  to indicate that the variables occurring in  $p$  are among  $x_1, \dots, x_n$ . We say that a skew brace term  $p$  is  $n$ -ary if the number of variables appearing explicitly in  $p$  is  $\leq n$ .

Given a skew brace term  $p(x_1, \dots, x_n)$  over some set of variables  $X$  and a skew brace  $A$ , we define a map  $p^A: A^n \rightarrow A$  as follows:

- (1) if  $p$  is a variable  $x_i \in X$ , then  $p^A(a_1, \dots, a_n) = a_i$  for all  $a_1, \dots, a_n \in A$ ;
- (2) if  $p$  is of the form  $p_1(x_1, \dots, x_n) \star p_2(x_1, \dots, x_n)$ , then

$$p^A(a_1, \dots, a_n) = p_1^A(a_1, \dots, a_n) \star p_2^A(a_1, \dots, a_n),$$

where  $\star$  denotes either  $\circ$  or  $+$ ;

- (3) if  $p$  is of the form  $p_1(x_1, \dots, x_n)'$ , then  $p^A(a_1, \dots, a_n) = p_1^A(a_1, \dots, a_n)'$ ;
- (4) if  $p$  is of the form  $-p_1(x_1, \dots, x_n)$ , then  $p^A(a_1, \dots, a_n) = -p_1^A(a_1, \dots, a_n)$ .

**Example 2.11.** Let  $X = \{x, y\}$ , then  $p(x, y) = -x + x \circ y - y$  is a skew brace term over  $X$ . Given a skew brace  $A$ , its term function over  $A$  is the map  $p^A: A^2 \rightarrow A$ ,  $(a, b) \mapsto a * b$ .

**Notation 2.12.** Let  $A$  be a skew brace. From now on, we denote by  $\overline{A}$  the quotient  $A / \text{Ann } A$ .

**Lemma 2.13.** Let  $n, m$  be integers,  $\eta_1, \dots, \eta_{2m}$  be  $n$ -ary skew brace terms and  $p$  be an  $m$ -ary skew brace term. Let  $\phi_1, \dots, \phi_m$  be collections of maps where each  $\phi_i$  is either  $\phi_+$  or  $\phi_*$ . Then one can construct a collection of well-defined maps  $\phi$  that associates to every skew brace  $B$  a map  $\phi^B: \overline{B}^n \rightarrow B'$  such that

$$\phi^B(\overline{b}_1, \dots, \overline{b}_n) = p^B(a_1(\overline{b}_1, \dots, \overline{b}_n), \dots, a_m(\overline{b}_1, \dots, \overline{b}_n)),$$

where  $a_i: \overline{B}^n \mapsto B$  is the map

$$(\overline{b}_1, \dots, \overline{b}_n) \mapsto \phi_i^B(\eta_{2i-1}^{\overline{B}}(\overline{b}_1, \dots, \overline{b}_n), \eta_{2i}^{\overline{B}}(\overline{b}_1, \dots, \overline{b}_n))$$

for all  $1 \leq i \leq m$ . In addition, if  $A$  and  $B$  are two isoclinic skew braces with isoclinism  $(\xi, \theta)$ , then the following diagram

$$(2.4) \quad \begin{array}{ccc} \overline{A}^n & \xrightarrow{\phi^A} & A' \\ \xi^n \downarrow & & \downarrow \theta \\ \overline{B}^n & \xrightarrow{\phi^B} & B' \end{array}$$

commutes.

*Proof.* Let  $B$  be a skew brace. One has that  $\phi^B$  is the composition of the maps

$$\overline{B}^n \xrightarrow{\theta} \overline{B}^{2m} \xrightarrow{\gamma} B^m \xrightarrow{p^B} B$$

where

$$\begin{aligned} \theta(\overline{b}_1, \dots, \overline{b}_n) &= (\eta_1^{\overline{B}}(\overline{b}_1, \dots, \overline{b}_n), \dots, \eta_{2m}^{\overline{B}}(\overline{b}_1, \dots, \overline{b}_n)), \\ \gamma(\overline{f}_1, \dots, \overline{f}_{2m}) &= (\phi_1^B(\overline{f}_1, \overline{f}_2), \dots, \phi_m^B(\overline{f}_{2m-1}, \overline{f}_{2m})). \end{aligned}$$

Suppose that  $A$  and  $B$  are isoclinic skew braces with isoclinism  $(\xi, \theta)$ . Since  $(\xi, \theta)$  is an isoclinism, [16, Theorem 10.3] implies that for all  $a_1, \dots, a_n \in A$  and  $1 \leq i \leq m$ ,

$$\begin{aligned} \phi_i^B(\eta_{2i-1}^{\overline{B}}(\xi(\overline{a}_1), \dots, \xi(\overline{a}_n)), \eta_{2i}^{\overline{B}}(\xi(\overline{a}_1), \dots, \xi(\overline{a}_n))) \\ = \phi_i^B(\xi(\eta_{2i-1}^{\overline{A}}(\overline{a}_1, \dots, \overline{a}_n)), \xi(\eta_{2i}^{\overline{A}}(\overline{a}_1, \dots, \overline{a}_n))) \\ = \theta(\phi_i^A(\eta_{2i-1}^{\overline{A}}(\overline{a}_1, \dots, \overline{a}_n), \eta_{2i}^{\overline{A}}(\overline{a}_1, \dots, \overline{a}_n))). \end{aligned}$$

Therefore

$$\phi^B(\xi^n(\overline{a}_1, \dots, \overline{a}_n)) = \theta(\phi^A(\overline{a}_1, \dots, \overline{a}_n)).$$

This means that the diagram (2.5) is commutative.  $\square$

**Notation 2.14.** Let  $X$  be a skew brace and  $x, y \in X$ . Then

$$\begin{aligned} r(x, y) &= -y - x + x \circ y = [-y, -x + x \circ y]_+ + x * y, \\ l(x, y) &= x \circ y - y - x = [x \circ y - y, -x]_+ + x * y. \end{aligned}$$

**Proposition 2.15.** *For all skew braces  $X$ , the map*

$$\phi_{\circ}^X : \overline{X}^2 \rightarrow X', \quad (\bar{a}, \bar{b}) \mapsto [a, b]_{\circ},$$

*is well-defined. In addition, if  $A$  and  $B$  are isoclinic skew braces with isoclinism  $(\xi, \theta)$ , then*

$$(2.5) \quad \begin{array}{ccc} \overline{A}^2 & \xrightarrow{\phi_{\circ}^A} & A' \\ \xi \times \xi \downarrow & & \downarrow \theta \\ \overline{B}^2 & \xrightarrow{\phi_{\circ}^B} & B' \end{array}$$

*commutes.*

*Proof.* Let  $X$  be a skew brace. For all  $a, b \in X$ ,

$$[a, b]_{\circ} = [a, b]_{+} - l(b, a) - r(b \circ a, a' \circ b') + r(b, a' \circ b') + r(a, b \circ a' \circ b').$$

Lemma 2.13 concludes the proof.  $\square$

**Proposition 2.16.** *Let  $A$  and  $B$  be two skew braces. If  $A$  and  $B$  are isoclinic, then  $A_{+}$  is isoclinic to  $B_{+}$  and  $A_{\circ}$  is isoclinic to  $B_{\circ}$ .*

*Proof.* Assume  $A$  is isoclinic to  $B$  with isoclinism  $(\xi, \theta)$ . Because of the commutativity of (2.3), the map  $\theta$  restricts to an isomorphism  $\theta_1 : [A, A]_{+} \rightarrow [B, B]_{+}$  and  $\xi(Z(A_{+})/\text{Ann } A) = Z(B_{+})/\text{Ann } B$ . Therefore  $\xi$  induces a group isomorphism  $\xi_1 : A_{+}/Z(A_{+}) \rightarrow B_{+}/Z(B_{+})$  such that

$$\begin{array}{ccc} (A_{+}/Z(A_{+}))^2 & \longrightarrow & A'_{+} \\ \xi \times \xi \downarrow & & \downarrow \theta_1 \\ (B_{+}/Z(B_{+}))^2 & \longrightarrow & B'_{+} \end{array}$$

commutes, where the horizontal maps are the classical commutator maps for groups.

Proposition 2.15 and a similar argument shows that  $A_{\circ}$  is isoclinic to  $B_{\circ}$ .  $\square$

**Remark 2.17.** If  $A$  and  $B$  are isoclinic skew braces, then  $A$  is of nilpotent type if and only if  $B$  is of nilpotent type.

Recall that a skew brace  $A$  is said to be *two-sided* if  $(a+b) \circ c = a \circ c - c + b \circ c$  holds for all  $a, b, c \in A$ . In [13], Rump proved that radical rings are exactly two-sided skew braces with an abelian additive group.

**Theorem 2.18.** *Let  $A$  and  $B$  be skew braces. If  $A$  and  $B$  are isoclinic, then  $A$  is two-sided if and only if  $B$  is two-sided.*

*Proof.* We claim that for all skew braces  $X$

$$\phi^X : (\overline{X})^3 \rightarrow X', \quad (\bar{a}, \bar{b}, \bar{c}) \mapsto (a+b) \circ c - b \circ c + c - a \circ c,$$

is well-defined. In addition, if  $A$  and  $B$  be isoclinic skew braces with isoclinism  $(\xi, \theta)$ . the following diagram

$$(2.6) \quad \begin{array}{ccc} \overline{A}^3 & \xrightarrow{\phi^A} & A' \\ \xi^3 \downarrow & & \downarrow \theta \\ \overline{B}^3 & \xrightarrow{\phi^B} & B' \end{array}$$

commutes. This is a direct consequence of the Lemma 2.13 and the fact that for any elements  $a, b, c$  of any brace, the following equation holds

$$(a + b) \circ c - b \circ c + c - a \circ c = l(a + b, c) - l(b, c) - [a, b \circ c - c - b]_+ - l(a, c)$$

This concludes the proof.  $\square$

**Example 2.19.** Computer calculations using the database of [10] show that among the 47 skew braces of size eight, there are 20 isoclinism classes. Moreover, there are eight isoclinism classes of radical rings of size eight, 12 isoclinism classes of skew braces of abelian type of size eight and 16 isoclinism classes of two-sided skew braces of size eight. See Table 2.1 for other numbers.

size	radical rings	abelian type	two-sided	all
8	8	12	16	20
27	10	13	25	38

TABLE 2.1. Number of isoclinism classes of skew braces.

### 3. ISOCLINISM INVARIANTS

**Notation 3.1.** Let  $A$  be a skew brace and  $B$  and  $C$  be sub skew braces. Then

$$B + C = \{b + c : b \in B, c \in C\}.$$

Note that, in general, if  $B$  and  $C$  are sub skew braces of  $A$ , then  $B + C$  is not a skew brace. However,  $B + C$  is always a skew brace if  $C$  is an ideal.

**Proposition 3.2.** Let  $A$  and  $B$  be isoclinic skew braces. Then sub skew braces of  $A$  containing  $\text{Ann}(A)$  are in bijective correspondence with sub skew braces of  $B$  containing  $\text{Ann}(B)$ . Furthermore, these corresponding sub skew braces are isoclinic.

*Proof.* Let  $(\xi, \theta)$  be an isoclinism between  $A$  and  $B$ . The correspondence is given by  $A_1 \mapsto \pi^{-1}(\xi(A_1/\text{Ann}(A)))$  for all skew braces  $\text{Ann}(A) \subseteq A_1 \subseteq A$ , where the map  $\pi: B \rightarrow B/\text{Ann}(B)$  is the canonical map. Let  $\text{Ann}(A) \subseteq A_1 \subseteq A$  and  $\text{Ann}(B) \subseteq B_1 \subseteq B$  be corresponding skew braces. Then, by the commutativity of the diagram (2.3),  $\theta$  restricts to an isomorphism  $A_1' \rightarrow B_1'$  and  $\xi$  factors through an isomorphism  $\bar{A}_1 \rightarrow \bar{B}_1$ . These two maps form an isoclinism from  $A_1$  to  $B_1$ .  $\square$

**Proposition 3.3.** Let  $A$  be a skew brace and  $K$  be a sub skew brace of  $A$ . Then  $K$  is isoclinic to  $K + \text{Ann}(A)$ .

*Proof.* It is a direct consequence of the fact that  $K' = (K + \text{Ann}(A))'$  and that  $K/\text{Ann}(K) \simeq (K + \text{Ann}(A))/\text{Ann}(K + \text{Ann}(A))$ .  $\square$

**Proposition 3.4.** Let  $A$  be a skew brace and  $K$  be a sub skew brace of  $A$ . If  $A/\text{Ann}(A)$  is finite, then  $A$  is isoclinic to  $K$  if and only if  $K + \text{Ann}(A) = A$ .

*Proof.* Assume  $A$  is isoclinic to  $K$ . Then

$$|K/\text{Ann} K| = |A/\text{Ann} A| \geq |K/K \cap \text{Ann}(A)| \geq |K/\text{Ann} K|.$$

Therefore,  $(K + \text{Ann}(A))/\text{Ann}(A) = A/\text{Ann}(A)$ . Thus,  $K + \text{Ann}(A) = A$ .  $\square$

**Proposition 3.5.** Let  $A$  and  $B$  be isoclinic skew braces with isoclinism  $(\xi, \theta)$ . If  $K \subseteq A'$  is an ideal of  $A$ , then  $\theta(K) \subseteq B'$  is an ideal of  $B$ . In addition,  $A/K$  is isoclinic to  $B/\theta(K)$ .

*Proof.* First notice that for all elements  $x \in A'$  we have  $\xi(\bar{x}) = \overline{\theta(x)}$ . It is enough to verify it on the generators of  $A'$ . The commutativity of (2.3) implies that for all  $a, a_1 \in A$ ,

$$\overline{\theta([a, a_1]_+)} = [\xi(\bar{a}), \xi(\bar{a}_1)]_+, \quad \overline{\theta(a * a_1)} = \xi(\bar{a}) * \xi(\bar{a}_1).$$

We show that  $\theta(K)$  is an ideal of  $B$ . Clearly  $\theta(K)_+$  is a subgroup of  $B_+$  and  $\theta(K)_\circ$  is a subgroup of  $B_\circ$ . Let  $k \in K$ . Let  $b \in B$  and  $a \in A$  be such that  $\xi(\bar{a}) = \bar{b}$ . The commutativity of the diagram (2.3) implies that  $\theta([a, k]_+) = [b, \theta(k)]_+$  and  $\theta(a * k) = b * \theta(k)$ . Thus,  $b + \theta(k) - b = \theta(a + k - a) \in \theta(K)$  and  $\lambda_b(\theta(k)) = \theta(\lambda_a(k))$ . Therefore  $\theta(K)$  is a strong left ideal of  $B$ . Similarly, one shows that  $K_\circ$  is a normal subgroup of  $B_\circ$  using the commutativity of the diagram (2.5). It is left to see that  $A/K$  is isoclinic to  $B/\theta(K)$ . Firstly,  $(A/K)' = A'/K$  and  $(B/\theta(K))' = B'/\theta(K)$ . Thus  $\theta$  factors through an isomorphism  $\bar{\theta}: (A/K)' \rightarrow (B/\theta(K))'$ . The annihilator of  $A/K$  corresponds by the canonical homomorphism to an ideal  $Q$  of  $A$  such that

$$\text{Ann}(A) + K \subseteq Q \subseteq A.$$

Similarly, the annihilator of  $B/\theta(K)$  corresponds to an ideal  $Q_1$  of  $B$  such that

$$\text{Ann}(B) + \theta(K) \subseteq Q_1 \subseteq B.$$

Thus

$$\overline{A/K} \simeq A/Q \simeq \overline{A}/(Q/\text{Ann}(A)), \quad \overline{B/\theta(K)} \simeq B/Q_1 \simeq \overline{B}/(Q_1/\text{Ann}(B)).$$

It is left to see that  $\xi(Q/\text{Ann}(A)) = Q_1/\text{Ann}(B)$ . Let  $q \in Q$ , then  $[q, x]_+ \in K$ ,  $q * x \in K$  and  $[q, x]_\circ \in K$  for all  $x \in A$ . Let  $q_1 \in B$  be a representative of  $\xi(\bar{q})$ . By the commutativity of the diagrams (2.3) and (2.5),  $[q_1, x]_+ \in \theta(K)$ ,  $q_1 * x \in \theta(K)$  and  $[q_1, x]_\circ \in \theta(K)$  for all  $x \in B$ . By a symmetric argument,  $Q_1/\text{Ann}(B) \subseteq \xi(Q/\text{Ann}(A))$ . The commutativity of (2.3) is straightforward.  $\square$

**Proposition 3.6.** *Let  $A$  be a skew brace and  $K$  an ideal of  $A$ . Then  $A/K$  is isoclinic to  $A/(K \cap A')$ .*

*Proof.* We have that  $(A/K)' = (A' + K)/K \cong A'/(K \cap A') = (A/(K \cap A'))'$ . The annihilator of  $A/K$  corresponds to an ideal  $Q$  of  $A$  such that  $\text{Ann}(A) + K \subseteq Q \subseteq A$ . Similarly, the annihilator of  $A/(K \cap A')$  corresponds to an ideal  $Q_1$  of  $A$  such that  $\text{Ann}(A) + K \cap A' \subseteq Q_1 \subseteq A$ . Thus  $\overline{A/K} \simeq A/Q$  and  $\overline{A/(K \cap A')} \simeq A/Q_1$ . It is left to see that  $Q = Q_1$ . It is clear that  $Q_1 \subseteq Q$ . Let  $q \in Q$ . Then  $[q, x]_+ \in K$ ,  $q * x \in K$  and  $[q, x]_\circ \in K$  for all  $x \in A$ . In addition, by the definition of  $A'$ ,  $[q, x]_+ \in A'$ ,  $q * x \in A'$  and  $[q, x]_\circ \in A'$  for all  $x \in A$ . Thus  $q \in Q_1$ . The commutativity of (2.3) is straightforward.  $\square$

**Corollary 3.7.** *Let  $A$  be a finite skew brace and  $K$  an ideal of  $A$ . Then  $A$  is isoclinic to  $A/K$  if and only if  $K \cap A' = \{0\}$ .*

*Proof.* If  $K \cap A' \neq \{0\}$ , then

$$|(A/K)'| = |A'/(K \cap A')| = |A'|/|K \cap A'| < |A'|.$$

Thus  $A$  cannot be isoclinic to  $A/K$ . The other implication is a direct consequence of 3.6.  $\square$

**Remark 3.8.** Let  $A, A_1, B$  and  $B_1$  be skew braces. If  $A$  is isoclinic to  $A_1$  and  $B$  is isoclinic to  $B_1$ , then  $A \times B$  is isoclinic to  $A_1 \times B_1$ .

**Definition 3.9.** A skew brace  $A$  such that  $\text{Ann}(A) \subseteq A'$  will be called a *stem skew brace*.

**Theorem 3.10.** *Every skew brace is isoclinic to a stem skew brace.*

*Proof.* Let  $A$  be a skew brace and  $I$  a set of indices (possibly uncountable). Let  $\{\xi_i : i \in I\}$  be a set of generators of  $A$  (as a skew brace). Let  $G$  be the free abelian group generated by  $\{\eta_i : i \in I\}$ . Then  $A$  is isoclinic to  $A \times G$ . Let  $A_1 \subseteq A \times G$  be the skew brace generated by  $\{(\xi_i, \eta_i) : i \in I\}$ . Then clearly  $A_1 + G$  coincides with  $A \times G$ . Since  $G$  is contained in the annihilator of  $A \times G$ , it follows that  $A_1$  is isoclinic to  $A \times G$ . Thus  $A_1$  is isoclinic to  $A$ . Let  $\pi : A_1 \rightarrow G$  denote the projection on the second coordinate. Then  $\pi$  factors through a morphism of abelian groups  $\bar{\pi} : A_1/A'_1 \rightarrow G$ . By the universal property of the free abelian group  $G$ , there is a homomorphism  $\phi : G \rightarrow A_1/A'_1$  that maps  $\eta_i$  to  $(\xi_i, \eta_i)$  for all  $i \in I$ . In fact,  $\phi$  is the inverse map of  $\bar{\pi}$ . Therefore  $A_1/A'_1 \simeq G$ . Moreover,

$$\text{Ann}(A_1)/(\text{Ann}(A_1) \cap A'_1) \simeq (\text{Ann}(A_1) + A'_1)/A'_1,$$

that is  $\text{Ann}(A_1)/(\text{Ann}(A_1) \cap A'_1)$  is isomorphic to a subgroup of a free abelian group. Thus  $\text{Ann}(A_1)$  is the direct product of  $\text{Ann}(A_1) \cap A'_1$  with another group  $K$ . Since  $K \cap A'_1 = \{0\}$ , the skew brace  $A_1/K$  is isoclinic to  $A_1$ . The annihilator of  $A_1/K$  is  $\text{Ann}(A_1)/K$  and is contained in the commutator, which is  $(A'_1 + K)/K$ .  $\square$

For  $n \geq 2$ , let  $C_n$  be the cyclic group of order  $n$ .

**Notation 3.11.** *Let  $n \geq 2$  be an integer. Let  $d$  be an integer such that  $d|n$  and every prime divisor of  $n$  divides  $d$ . We will denote by  $C(n, d)$  the skew brace  $C_n$  with multiplication defined as  $x \circ y = x + dxy + y$ .*

The skew braces  $C(n, d)$  appear in the work of Rump [14].

**Example 3.12.** Since  $C(p^2, p)' = \text{Ann}(C(p^2, p))$ ,  $C(p^2, p)$  is a stem skew brace. It is straightforward to see that  $C(p^2, p)$  is isoclinic to  $C(p^n, p^{n-1})$  for all  $n > 1$ .

**Notation 3.13.** *Let  $B$  be a skew brace. There is a canonical group homomorphism*

$$B_+ \rtimes_{\rho} B_{\circ} \rightarrow \text{Aut}(B_+), \quad (a, b) \mapsto (c \mapsto a + \rho_b(c) - a),$$

*We write  ${}^{(a,b)}c = {}^a\rho_b(c)$ .*

**Notation 3.14.** *Let  $A$  be a skew brace, denote respectively by  $\overline{A_+}$  and  $\overline{A_{\circ}}$  the groups  $A_+/\text{Ann}(A)$  and  $A_{\circ}/\text{Ann}(A)$ .*

**Remark 3.15.** The group homomorphism  $\rho : A_{\circ} \rightarrow \text{Aut}(A_+)$  induces a group homomorphism  $\bar{\rho} : \overline{A_{\circ}} \rightarrow \text{Aut}(\overline{A_+})$ . One can check that the group  $\overline{A_+} \rtimes_{\bar{\rho}} \overline{A_{\circ}}$  is isomorphic to  $(A_+ \rtimes_{\rho} A_{\circ})/(\text{Ann}(A) \times \text{Ann}(A))$ . Thus  $\overline{A_+} \rtimes_{\bar{\rho}} \overline{A_{\circ}}$  acts canonically on  $A_+$ .

**Remark 3.16.** If  $A$  and  $B$  are isoclinic skew braces, then  $\overline{A_+} \rtimes_{\bar{\rho}} \overline{A_{\circ}} \simeq \overline{B_+} \rtimes_{\bar{\rho}} \overline{B_{\circ}}$ .

**Notation 3.17.** *Let  $A$  be a skew brace. Let  $H$  be a subgroup of  $\overline{A_+} \rtimes_{\bar{\rho}} \overline{A_{\circ}}$ . We call the orbit of an element  $a \in A$  under the induced action of  $H$  an  $H$ -orbit.*

**Theorem 3.18.** *Let  $A$  and  $B$  be isoclinic skew braces. Let  $H$  be a subgroup of  $\overline{A_+} \rtimes_{\bar{\rho}} \overline{A_{\circ}}$  and  $K$  be the corresponding subgroup of  $\overline{B_+} \rtimes_{\bar{\rho}} \overline{B_{\circ}}$ . For  $c \in \mathbb{Z}_{\geq 1}$ , let  $m_1$  (resp.  $m_2$ ) be the number of  $H$ -orbits (resp.  $K$ -orbits) of size  $c$ . Then*

$$m_1 = m_2|A|/|B|.$$



*Proof.* Lemma 2.13 and the fact that

$${}^{(a,b)}c - c = [a, b \circ c - b]_+ + [b \circ c, -b]_+ + b * c$$

imply that the map

$$\phi^X: \overline{X}^3 \rightarrow X', \quad (\overline{a}, \overline{b}, \overline{c}) \mapsto {}^{(a,b)}c - c$$

is well-defined for all skew brace  $X$ . In addition, the diagram

$$(3.1) \quad \begin{array}{ccc} \overline{A}^3 & \xrightarrow{\phi^A} & A' \\ \xi^3 \downarrow & & \downarrow \theta \\ \overline{B}^3 & \xrightarrow{\phi^B} & B' \end{array}$$

commutes.

An element  $a \in A$  has an  $H$ -orbit of size  $c$  if and only if the index of the subgroup  $C(\overline{a}) = \{w \in H : \phi^A(w, \overline{a}) = 0\}$  in  $H$  is  $c$ . Let  $S$  be the subset of  $\overline{A}$  that consists of elements  $w$  such that  $C(w)$  has index  $c$  in  $H$ . If  $\pi: A \rightarrow \overline{A}$  is the canonical homomorphism,  $\pi^{-1}(S)$  is the set of elements of  $A$  that have an  $H$ -orbit of size  $c$ . Hence  $m_1c = |\text{Ann } A||S|$ . Because of the commutativity of (3.1), one also has that  $m_2c = |\text{Ann } B||S|$ . Hence the claim follows.  $\square$

**Remark 3.19.** We use the notations of Theorem 3.18.

(1) Let

$$\begin{aligned} H &= \{(-\overline{a}, \overline{a}) : a \in A\} \subseteq \overline{A}_+ \rtimes_{\overline{\rho}} \overline{A}_\circ, \\ K &= \{(-\overline{b}, \overline{b}) : b \in B\} \subseteq \overline{B}_+ \rtimes_{\overline{\rho}} \overline{B}_\circ. \end{aligned}$$

Note that  $K$  is the subgroup of  $\overline{B}_+ \rtimes_{\overline{\rho}} \overline{B}_\circ$  corresponding to  $H$  by the isomorphism induced by isoclinism. Then the  $H$ -orbits and the  $K$ -orbits are, respectively, the orbits of the canonical actions  $\lambda: A_\circ \rightarrow \text{Aut}(A_+)$  and  $\lambda: B_\circ \rightarrow \text{Aut}(B_+)$ .

(2) Similarly, Theorem 3.18 applies to the pair  $H = \{(\overline{0}, \overline{a}) : a \in A\}$  and  $K = \{(\overline{0}, \overline{b}) : b \in B\}$ . In this case, the  $H$ -orbits and the  $K$ -orbits are, respectively, the orbits of the canonical actions  $\rho: A_\circ \rightarrow \text{Aut}(A_+)$  and  $\rho: B_\circ \rightarrow \text{Aut}(B_+)$ .

**Example 3.20.** Let  $A$  be the skew brace  $C_2 \times C_4$  with multiplication given by

$$(x_1, x_2) \circ (y_1, y_2) = (x_1 + x_2, y_1 + y_2 + 2x_1y_2)$$

and  $B$  the skew brace  $C_2 \times C_4$  with multiplication given by

$$(x_1, x_2) \circ (y_1, y_2) = (x_1 + x_2, y_1 + y_2 + 2(x_1 + y_2)x_2 + 2y_1y_2).$$

Both  $A$  and  $B$  have commutator  $C_2$  and annihilator quotient  $C_2 \times C_2$ . In addition,  $A_+$  is isoclinic to  $B_+$  and  $A_\circ$  is isoclinic to  $B_\circ$ . However,  $A$  and  $B$  are not isoclinic as  $A$  has four  $\lambda$ -orbits of size one and two of size two and  $B$  has two  $\lambda$ -orbits of size one and three of size two.

In group theory, the notion of isoclinism is very convenient in the study of finite  $p$ -groups as these groups have non-trivial center and their commutator is a proper subgroup. This implies that isoclinism only depends on relations between groups of smaller order. However, there exist skew braces of prime-power size that have trivial annihilator. This is not the case for two-sided skew braces. One can extend

the action defined earlier (see Notation 3.13) for two-sided skew braces. Let  $B$  be a two-sided skew brace. Since the underlying multiplicative group acts on itself by conjugation, we can consider the semidirect product  $B_\circ \rtimes B_\circ$ . Straightforward calculations show that the map

$$B_\circ \rtimes B_\circ \rightarrow \text{Aut}(B_+), \quad (a, b) \mapsto (c \mapsto a \circ \rho_b(c) \circ a'),$$

defines an action by automorphisms of  $B_\circ \rtimes B_\circ$  over  $B_+$ . Thus we can consider the semi-direct product  $B_+ \rtimes (B_\circ \rtimes B_\circ)$ . Finally, straightforward calculations show that the map

$$B_+ \rtimes (B_\circ \rtimes B_\circ) \rightarrow \text{Aut}(B_+), \quad (a, b, c) \mapsto (d \mapsto a + b \circ \rho_c(d) \circ b' - a),$$

defines an action by automorphisms of  $B_+ \rtimes (B_\circ \rtimes B_\circ)$  over  $B_+$ . It is straightforward to see that the elements of  $B$  whose orbits have size one are exactly the elements of the annihilator. Moreover, if  $B$  is a two-sided skew brace of size  $p^n$ , the group  $B_+ \rtimes (B_\circ \rtimes B_\circ)$  has size  $p^{3n}$ , thus the non-trivial orbits of the action have size a power of  $p$ . Let  $n_1, \dots, n_m$  denote the sizes of the  $m$  non-trivial orbits of  $B$ , then we have the following class equation

$$p^n = |\text{Ann}(B)| + \sum_{i=1}^m n_i.$$

Therefore  $p$  divides  $|\text{Ann}(B)|$ . We have proved the following result:

**Proposition 3.21.** *Let  $p$  be a prime number and  $B$  be a two-sided skew brace of size  $p^n$  for some integer  $n \geq 1$ . Then  $\text{Ann}(B)$  is non-trivial.*

**Proposition 3.22.** *Let  $p$  be a prime number and  $B$  be a two-sided skew brace of size  $p^n$  for some integer  $n \geq 1$ . Then  $B'$  is a proper ideal of  $B$ .*

*Proof.* We proceed by induction on  $n$ . If  $n = 1$ , then  $B$  is the trivial skew brace  $C_p$ . Assume now that  $n \geq 2$ . Since  $|B/\text{Ann}(B)| < p^n$ , it follows by the induction hypothesis that  $\overline{B'} = (B' + \text{Ann}(B))/\text{Ann}(B)$  is a proper sub skew brace of  $\overline{B}$ . Thus  $B' + \text{Ann}(B) \subsetneq B$ .  $\square$

#### 4. AN APPLICATION TO THE YANG-BAXTER EQUATION

**Theorem 4.1.** *Let  $A$  and  $B$  be isoclinic skew braces. Then  $A$  is right nilpotent if and only if  $B$  is right nilpotent.*

*Proof.* Let  $(\xi, \theta)$  be an isoclinism between  $A$  and  $B$ . Assume that  $A$  is right nilpotent. By [6, Lemma 2.5],  $A/\text{Soc}(A)$  is right nilpotent.

We claim that

$$(4.1) \quad \xi(\text{Soc}(A)/\text{Ann}(A)) = \text{Soc}(B)/\text{Ann}(B).$$

Let  $a \in \text{Soc}(A)$  be such that  $\xi(a + \text{Ann}(A)) = b + \text{Ann}(B)$  for some  $b \in B$ . Let  $a_1 \in A$  and  $b_1 \in B$  such that  $\xi(a_1 + \text{Ann}(A)) = b_1 + \text{Ann}(B)$ . By the commutativity of (2.3),

$$\begin{aligned} b * b_1 &= \phi_*(b + \text{Ann}(B), b_1 + \text{Ann}(B)) \\ &= \phi_*(a + \text{Ann}(A), a_1 + \text{Ann}(A)) \\ &= a * a_1 = 0. \end{aligned}$$

Similarly,  $b$  is central in  $(B, +)$ . Now (4.1) follows.

It follows from the first isomorphism theorem that

$$(A/\text{Ann}(A))/(\text{Soc}(A)/\text{Ann}(A)) \simeq (B/\text{Ann}(B))/(\text{Soc}(B)/\text{Ann}(B)).$$

Hence  $A/\text{Soc}(A) \simeq B/\text{Soc}(B)$ . Thus  $B/\text{Soc}(B)$  is right nilpotent. By [6, Proposition 2.17],  $B$  is right nilpotent.  $\square$

A *set-theoretic solution* to the Yang–Baxter equation (YBE) is a pair  $(X, r)$ , where  $X$  is a set and  $r: X \times X \rightarrow X \times X$  is a bijective map such that

$$(r \times \text{id})(\text{id} \times r)(r \times \text{id}) = (\text{id} \times r)(r \times \text{id})(\text{id} \times r).$$

By convention, we will consider finite *non-degenerate* solutions, that is solutions  $(X, r)$ , where  $X$  is a finite set and

$$r(x, y) = (\sigma_x(y), \tau_y(x))$$

where the maps  $\sigma_x: X \rightarrow X$  and  $\tau_x: X \rightarrow X$  are bijective for every  $x \in X$ .

If  $(X, r)$  is a solution, there is an equivalence relation on  $X$  given by

$$x \sim y \iff \sigma_x = \sigma_y \text{ and } \tau_x = \tau_y.$$

This equivalence relation induces a solution  $\text{Ret}(X, r)$  on the set of equivalence classes. The solution  $\text{Ret}(X, r)$  is called the *retraction* of  $(X, r)$ .

A solution  $(X, r)$  is said to be *multipermutation* if there exists an integer  $m \geq 1$  such that  $|\text{Ret}^m(X, r)| = 1$ , where  $\text{Ret}^1(X, r) = \text{Ret}(X, r)$  and

$$\text{Ret}^k(X, r) = \text{Ret}(\text{Ret}^{k-1}(X, r))$$

for  $k \geq 2$ .

**Definition 4.2.** Let  $(X, r)$  and  $(Y, s)$  be solutions to the YBE. We say that  $(X, r)$  and  $(Y, s)$  are *isoclinic* if the skew braces  $\mathcal{G}(X, r)$  and  $\mathcal{G}(Y, s)$  are isoclinic.

The following result follows from [6, Proposition 2.17], [6, Theorem 2.20] and [5, Theorem 4.13].

**Lemma 4.3.** *Let  $(X, r)$  be a finite non-degenerate solution to the YBE. The following statements are equivalent:*

- (1)  $(X, r)$  is multipermutation.
- (2)  $\mathcal{G}(X, r)$  is right nilpotent of nilpotent type.
- (3)  $\mathcal{G}(X, r)$  is right nilpotent of nilpotent type.

**Theorem 4.4.** *Let  $(X, r)$  and  $(Y, s)$  be isoclinic solutions to the YBE. Then  $(X, r)$  is multipermutation if and only if  $(Y, s)$  is multipermutation.*

*Proof.* If  $(X, r)$  is multipermutation, then  $\mathcal{G}(X, r)$  is right nilpotent. Thus  $\mathcal{G}(Y, s)$  is right nilpotent and hence  $(Y, s)$  is multipermutation.  $\square$

We conclude the paper with concrete examples of involutive solutions up to isoclinism. Recall that solution  $(X, r)$  is said to be *involutive* if  $r^2 = \text{id}$ . If  $(X, r)$  is involutive, then

$$\tau_y(x) = \sigma_{\sigma_x(y)}^{-1}(x)$$

for all  $x, y \in X$ .

**Example 4.5.** There are four isoclinism classes of involutive solutions of size five. Let  $X = \{1, 2, 3, 4\}$ . The following list provides a complete set of representatives over the set  $X$ :

- (1) The flip  $(x, y) \mapsto (y, x)$ .
- (2)  $\sigma_1 = \sigma_2 = \text{id}$ ,  $\sigma_3 = (12)$  and  $\sigma_4 = (12)(34)$ .
- (3)  $\sigma_1 = (34)$ ,  $\sigma_2 = (1324)$ ,  $\sigma_3 = (1423)$  and  $\sigma_4 = (1, 2)$ .
- (4)  $\sigma_1 = (12)$ ,  $\sigma_2 = (1324)$ ,  $\sigma_3 = (34)$  and  $\sigma_4 = (1423)$ .

**Remark 4.6.** Isoclinism of solutions does not preserve indecomposability. For example, let  $X = \{1, 2, 3, 4\}$  and  $\sigma = (1234)$ . Then  $(X, r)$  has multipermutation level two. Moreover,  $(X, r)$  is isoclinic to the flip.

**Remark 4.7.** Isoclinism of solutions does not preserve the multipermutation level. For example, let  $X = \{1, 2, 3, 4\}$  and  $\sigma_1 = \sigma_2 = \sigma_3 = \text{id}$ ,  $\sigma_4 = (23)$ . Then  $(X, r)$ , where  $r(x, y) = (\sigma(y), \sigma^{-1}(x))$ , is an indecomposable solution isoclinic to the flip.

**Example 4.8.** There are six isoclinism classes of involutive solutions of size five. Let  $X = \{1, 2, 3, 4, 5\}$ . The following list provides a complete set of representatives over  $X$ :

- (1) The flip  $(x, y) \mapsto (y, x)$ .
- (2)  $\sigma_1 = \sigma_2 = \sigma_3 = \text{id}$ ,  $\sigma_4 = (45)$ ,  $\sigma_5 = (23)(45)$ .
- (3)  $\sigma_1 = \sigma_2 = \sigma_3 = \text{id}$ ,  $\sigma_4 = (23)(45)$ ,  $\sigma_5 = (12)(45)$ .
- (4)  $\sigma_1 = \text{id}$ ,  $\sigma_2 = (45)$ ,  $\sigma_3 = (2435)$ ,  $\sigma_4 = (2534)$  and  $\sigma_5 = (23)$ .
- (5)  $\sigma_1 = \text{id}$ ,  $\sigma_2 = (23)$ ,  $\sigma_3 = (2435)$ ,  $\sigma_4 = (45)$  and  $\sigma_5 = (2534)$ .
- (6)  $\sigma_1 = \sigma_2 = (45)$ ,  $\sigma_3 = (14)(25)$  and  $\sigma_4 = \sigma_5 = (12)$ .

Note that the fourth solution is isoclinic to the third solution of Example 4.5. The fifth solution is isoclinic to the fourth solution of Example 4.5.

#### ACKNOWLEDGEMENTS

Vendramin is supported in part by OZR3762 of Vrije Universiteit Brussel.

#### REFERENCES

- [1] D. Bachiller. Solutions of the Yang-Baxter equation associated to skew left braces, with applications to racks. *J. Knot Theory Ramifications*, 27(8):1850055, 36, 2018.
- [2] D. Bachiller, F. Cedó, and E. Jespers. Solutions of the Yang-Baxter equation associated with a left brace. *J. Algebra*, 463:80–102, 2016.
- [3] D. Bachiller, F. Cedó, and L. Vendramin. A characterization of finite multipermutation solutions of the Yang-Baxter equation. *Publ. Mat.*, 62(2):641–649, 2018.
- [4] V. G. Bardakov, M. V. Neshchadim, and M. K. Yadav. Computing skew left braces of small orders. *Internat. J. Algebra Comput.*, 30(4):839–851, 2020.
- [5] F. Cedó, E. Jespers, L. Kubat, A. Van Antwerpen, and C. Verwimp. On various types of nilpotency of the structure monoid and group of a set-theoretic solution of the Yang-Baxter equation. *J. Pure Appl. Algebra*, 227(2):Paper No. 107194, 38, 2023.
- [6] F. Cedó, A. Smoktunowicz, and L. Vendramin. Skew left braces of nilpotent type. *Proc. Lond. Math. Soc. (3)*, 118(6):1367–1392, 2019.
- [7] L. N. Childs, C. Greither, K. P. Keating, A. Koch, T. Kohl, P. J. Truman, and R. G. Underwood. *Hopf algebras and Galois module theory*, volume 260 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, [2021] ©2021.
- [8] P. Etingof, T. Schedler, and A. Soloviev. Set-theoretical solutions to the quantum Yang-Baxter equation. *Duke Math. J.*, 100(2):169–209, 1999.
- [9] T. Gateva-Ivanova and P. Cameron. Multipermutation solutions of the Yang-Baxter equation. *Comm. Math. Phys.*, 309(3):583–621, 2012.
- [10] L. Guarnieri and L. Vendramin. Skew braces and the Yang-Baxter equation. *Math. Comp.*, 86(307):2519–2534, 2017.
- [11] P. Hall. The classification of prime-power groups. *J. Reine Angew. Math.*, 182:130–141, 1940.
- [12] V. Lebed and L. Vendramin. On structure groups of set-theoretic solutions to the Yang-Baxter equation. *Proc. Edinb. Math. Soc. (2)*, 62(3):683–717, 2019.

- [13] W. Rump. Braces, radical rings, and the quantum Yang-Baxter equation. *J. Algebra*, 307(1):153–170, 2007.
- [14] W. Rump. Classification of cyclic braces. *J. Pure Appl. Algebra*, 209(3):671–685, 2007.
- [15] W. Rump. The brace of a classical group. *Note Mat.*, 34(1):115–144, 2014.
- [16] H. P. Sankappanavar and S. Burris. A course in universal algebra. *Graduate Texts Math*, 78:56, 1981.
- [17] A. Smoktunowicz. Algebraic approach to Rump’s results on relations between braces and pre-Lie algebras. *J. Algebra Appl.*, 21(3):Paper No. 2250054, 13, 2022.
- [18] A. Smoktunowicz. A new formula for Lazard’s correspondence for finite braces and pre-Lie algebras. *J. Algebra*, 594:202–229, 2022.
- [19] A. Smoktunowicz. On the passage from finite braces to pre-Lie rings. *Adv. Math.*, 409:Paper No. 108683, 2022.
- [20] A. Smoktunowicz and L. Vendramin. On skew braces (with an appendix by N. Byott and L. Vendramin). *J. Comb. Algebra*, 2(1):47–86, 2018.

(L. Vendramin) DEPARTMENT OF MATHEMATICS, VRIJE UNIVERSITEIT BRUSSEL, PLEINLAAN 2, 1050 BRUSSEL, BELGIUM

*Email address:* `Leandro.Vendramin@vub.be`

(T. Letourmy) DÉPARTEMENT DE MATHÉMATIQUE, UNIVERSITÉ LIBRE DE BRUXELLES, BOULEVARD DU TRIOMPHE, B-1050 BRUSSELS, BELGIUM

*Email address:* `thomas.letourmy@ulb.be`