

MODULUS OF CONTINUITY OF WEAK SOLUTIONS TO A CLASS OF SINGULAR ELLIPTIC EQUATIONS

XIANGSHENG XU

ABSTRACT. In this paper we study the modulus of continuity of weak solutions to a singular elliptic equation in the plane under very weak assumption on the integrability of the elliptic coefficients. Our investigation reveals that the modulus of continuity can be described by the reciprocal of the logarithmic function raised to a power. However, the power can be arbitrarily large. This is in sharp contrast with a result by J. Onninen and X. Zhong (*Ann. Scuola Norm. Sup. Pisa Cl. Sci.*, **6**(2007), 103–116) for a degenerate elliptic equation in the plane, in which the power must be suitably small.

1. INTRODUCTION

In this paper we investigate the modulus of continuity of a weak solution to the Dirichlet boundary value problem

$$(1.1) \quad -\operatorname{div} [(I + \mathbf{m} \otimes \mathbf{m})\nabla p] = S \text{ in } \Omega,$$

$$(1.2) \quad p = 0 \text{ on } \partial\Omega$$

under the following assumptions:

- (H1) Ω is a bounded domain in \mathbb{R}^2 with Lipschitz boundary $\partial\Omega$;
- (H2) $S \in L^q(\Omega)$ for some $q > 1$;
- (H3) $\mathbf{m} = (m_1, m_2)$ is a vector-valued function in $(L^4(\Omega))^2$.

Recall that the out product $\mathbf{m} \otimes \mathbf{m}$ is the matrix given by

$$\mathbf{m} \otimes \mathbf{m} = \mathbf{m}^T \mathbf{m}.$$

Thus,

$$\mathbf{m} \otimes \mathbf{m} \nabla p = (\mathbf{m} \cdot \nabla p) \mathbf{m}.$$

We say that p is a weak solution to (1.1)-(1.2) if:

- (D1) $p \in W_0^{1,2}(\Omega)$, $\mathbf{m} \cdot \nabla p \in L^2(\Omega)$;
- (D2) for each $\zeta \in W_0^{1,2}(\Omega)$ with $\mathbf{m} \cdot \nabla \zeta \in L^2(\Omega)$ one has

$$(1.3) \quad \int_{\Omega} [\nabla p \nabla \zeta + (\mathbf{m} \cdot \nabla p)(\mathbf{m} \cdot \nabla \zeta)] dx = \int_{\Omega} S(x) \zeta dx.$$

Note from Theorem 7.15 in [8] that the test function ζ in (D2) satisfies

$$\int_{\Omega} e^{c_0|\zeta|^2} dx < \infty \text{ for some positive number } c_0.$$

Thus, each integral in (1.3) is well-defined.

2020 *Mathematics Subject Classification.* 35D30, 35B65, 35J15, 35J75, 35J47 .

Key words and phrases. Modulus of continuity of weak solutions, singular elliptic equations, the Stummel-Kato class of functions, the De Giorgi iteration scheme .

Department of Mathematics and Statistics, Mississippi State University, Mississippi State, MS 39762. *Email:* xxu@math.msstate.edu.

The out product of two vectors appears in many mathematical models. Our situation here is directly related to the analysis of biological transport networks [9, 11, 12]. We summarize known results concerning (1.1)-(1.2) in the following

Proposition 1.1. *Let (H1)-(H3) Hold. Then there is a unique weak solution p to (1.1)-(1.2). Moreover,*

(C1) *there is a positive number $c = c(\Omega)$ such that*

$$\|p\|_{\infty, \Omega} \leq c \|S\|_{q, \Omega}, \quad \text{and}$$

(C2) *for each $x_0 \in \Omega$ and each $R \in (0, \min\{1, \text{dist}(x_0, \partial\Omega)\})$ we can find a positive constant c with the property*

$$\text{osc}_{B_r(x_0)} p \equiv \text{ess sup}_{B_r(x_0)} p - \text{ess inf}_{B_r(x_0)} p \leq \frac{c}{\ln \frac{1}{2} \frac{R}{r}} \quad \text{for all } r \in (0, R],$$

where $B_r(x_0)$ is the ball centered at x_0 with radius r .

The existence of a unique weak solution can be inferred from a result in [18]. The conclusion (C1) is true even for the space dimension $N > 2$ [14]. A suitable modification of the proof in [14] can be applied to the case $N = 2$ (see Claim 2.4 below). The continuity property (C2) was established in [19].

To describe our results here, we introduce some notations first. We say that $f \in K_2(\Omega)$, the Stummel-Kato class of functions [1, 13], if f is a measurable function on Ω and

$$(1.4) \quad \eta(f; \Omega; r) \equiv \sup_{y \in \Omega} \int_{B_r(y)} |f(x)| \chi_{\Omega} |\ln |x - y|| dx \rightarrow 0 \quad \text{as } r \rightarrow 0^+,$$

while $f \in K_2^{\text{loc}}(\Omega)$ means that $\lim_{r \rightarrow 0} \eta(f; \Omega_1; r) = 0$ for each bounded subdomain Ω_1 of Ω with $\overline{\Omega_1} \subset \Omega$. Note that in (1.4) we have used $y \in \Omega$ instead of $y \in \mathbb{R}^2$ as was done in [13]. Our definition here seems to be more suitable for PDE applications. As usual, the letter c or $c_i, i = 0, 1, \dots$, will be used to represent a generic positive constant.

Our main result is:

Theorem 1.2. *Let (H1)-(H3) hold and p be the weak solution to (1.1)-(1.2).*

(C3) *Then for each $y \in \Omega$, $\ell > 0$, and $R \in (0, \min\{1, \text{dist}(y, \partial\Omega)\})$ there is a positive constant c such that*

$$(1.5) \quad \text{osc}_{B_r(y)} p \leq \frac{c}{\ln^{\ell} \frac{R}{r}} + cr^{\frac{2(q-1)}{q}} \quad \text{for all } r \in (0, R];$$

(C4) *If we further assume that*

$$(H4) \quad \mathbf{m} \in (\text{BMO}(\Omega))^2 \quad [5],$$

then

$$(1.6) \quad F \equiv |\nabla p|^2 + (\mathbf{m} \cdot \nabla p)^2 \in K_2^{\text{loc}}(\Omega).$$

To put our results into some mathematical context, we recall a conjecture by De Giorgi [2, 3]. In 1995, De Giorgi gave a lecture in Lecce, Italy on the continuity of weak solutions to second-order elliptic equations of the form

$$(1.7) \quad \text{div}(A \nabla u) = 0 \quad \text{in } \Omega \subset \mathbb{R}^N, \quad N \geq 2.$$

Here the entries of the coefficient matrix $A = A(x)$ are measurable functions, satisfying

$$\lambda(x) |\xi|^2 \leq A(x) \xi \cdot \xi \leq \Lambda(x) |\xi|^2$$

for some non-negative measurable functions $\lambda(x), \Lambda(x)$, a.e $x \in \Omega$, and each $\xi \in \mathbb{R}^N$. Obviously, if $\Lambda(x)$ is bounded above, we can replace it with the number 1, and we can do the same with $\lambda(x)$ if it is bounded away from 0 below. With this in mind, we say that (1.7) is singular if $\lambda(x) = 1$, and

degenerate if $\Lambda(x) = 1$. De Giorgi proposed several open problems [3] about the equation, one of which was stated as follows:

Conjecture 1.3 (De Giorgi). *If (1.7) is singular, the space dimension $N \geq 3$, and*

$$(1.8) \quad \int_{\Omega} e^{\Lambda(x)} dx < \infty,$$

then weak solutions of (1.7) are continuous.

This conjecture remains open. When $A = I + \mathbf{m} \otimes \mathbf{m}$ for some $\mathbf{m} \in (L^2(\Omega))^N$ then we have

$$|\xi|^2 \leq A(x)\xi \cdot \xi = |\xi|^2 + (\mathbf{m} \cdot \xi)^2 \leq (1 + |\mathbf{m}|^2)|\xi|^2 \quad \text{for a.e } x \in \Omega \text{ and each } \xi \in \mathbb{R}^N.$$

One way of gaining a condition like (1.8) is to assume that $N = 2$ and $\mathbf{m} \in (W_0^{1,2}(\Omega))^2$. Then (1.8) is a consequence of Theorem 7.15 in [8]. However, the case $N = 2$ is not included in the above conjecture even though our integrability condition on $\Lambda(x) = 1 + |\mathbf{m}|^2$ is much weaker.

It is well known that when $\Lambda(x) \leq c\lambda(x)$ for some $c > 0$ and $\lambda(x)$ is an A_2 weight weak solutions of (1.7) are Hölder continuous no matter what the space dimensions are [10]. The maximum Hölder exponent was investigated in [16] under the assumptions that $N=2$ and A is bounded and uniformly elliptic. If (1.7) is degenerate, $N = 2$, and

$$K \equiv \int_{\Omega} e^{\frac{\gamma}{\sqrt{\lambda(x)}}} dx < \infty \quad \text{for some } \gamma > 1,$$

a result of [15] asserts that for each $\beta \in (0, \gamma - 1)$ there is a c such that

$$\text{osc}_{B_r(x_0)} u \leq \frac{c \left(\int_{\Omega} A \nabla u \cdot \nabla u dx \right)^{\frac{1}{2}}}{\ln^{\frac{\beta}{2}} \frac{K}{\pi r^2}} \quad \text{for } r \text{ sufficiently small.}$$

In view of this, our result (C3) is really surprising because the power ℓ in (1.5) can be arbitrarily large and the entries of A are only square-integrable under (H3). It is natural for us to make the following

Conjecture 1.4. *Let (H1)-(H4) hold. Then the weak solution p to (1.1)-(1.2) is Hölder continuous.*

We point out that (H4) is not enough to guarantee (1.8) because $|\mathbf{m}|^2$ may not belong to $\text{BMO}(\Omega)$. If the preceding conjecture were true, we would immediately be able to find applications for it in the mathematical analysis of the biological network formation model [20]. Unfortunately, we have not been able to prove or disprove the conjecture. However, we do have

Theorem 1.5. *Let (H1)-(H4) hold. Then whenever $x_0 \in \Omega$ is such that*

$$(1.9) \quad \sup_{r>0} |\mathbf{m}_{x_0,r}| < \infty,$$

where

$$\mathbf{m}_{x_0,r} = \frac{1}{|B_r(x_0)|} \int_{B_r(x_0)} \mathbf{m} dx = \int_{B_r(x_0)} \mathbf{m} dx.$$

we can find two positive numbers c, α with

$$\text{osc}_{B_r(x_0)} p \leq cr^{\alpha} \quad \text{for } r \text{ sufficiently small.}$$

This is in essence a partial regularity result [20].

In [13] the relationship between the continuity of p and $|\nabla p|^2 \in K_2^{\text{loc}}(\Omega)$ was investigated. It turns out that the former implies the latter when A is bounded and uniformly elliptic. Here, due to the singularity in our problem, the continuity of p is not enough to guarantee (1.6). As we shall see below, we must have $\ell > 3$ in (1.5) to ensure that (1.6) holds. Of course, the most interesting result associated with the space $K_2^{\text{loc}}(\Omega)$ is the following

Lemma 1.6. *There is a constant c such that*

$$\int_{B_r(x_0)} |F|v^2 dx \leq c\eta(F; B_r(x_0); r) \int_{B_r(x_0)} |\nabla v|^2 dx \quad \text{for each } v \in W_0^{1,2}(B_r(x_0)).$$

This lemma is not really new, and it is only slightly different from Lemma 1.1 in [6], which has had many applications in the study of partial differential equations [6, 13]. We include the lemma here in the hope that this version of the inequality may be more suitable for applications in certain cases [1].

The rest of the paper is organized as follows. The proof of Theorem 1.2 is given in Section 2. It relies on a decomposition of the logarithmic function. Theorem 1.5 and Lemma 1.6 will be established in Section 3.

2. PROOF OF THEOREM 1.2

In this section we offer the proof of Theorem 1.2. The proof is divided into several lemmas.

Lemma 2.1. *Let (H1)-(H3) hold and p be the weak solution to (1.1)-(1.2). Then there is a positive constant c such that*

$$(2.1) \quad \int_{B_r(y)} F |\ln |x - y|| dx \leq (\ln 2 + |\ln r|) \int_{B_r(y)} F dx + c \int_{B_r(y)} |\nabla p|^2 dx \\ + c \left(\int_{B_r(y)} |\mathbf{m}|^4 dx \right)^{\frac{1}{2}} \int_{B_r(y)} |\nabla p|^2 dx + cr^{\frac{2(q-1)}{q}} \|S(x)\|_{q, B_r(y)}$$

for each $y \in \Omega$ and $r \in (0, \text{dist}(y, \partial\Omega))$.

Proof. Let y, r be given as in the lemma. Pick a smooth function $\zeta(x)$ on \mathbb{R}^N such that $\zeta(x) = 1$ on $B_1(0)$, $0 \leq \zeta(x) \leq 1$ on \mathbb{R}^N , and $\zeta(x) = 0$ when $|x| \geq 2$. Set

$$(2.2) \quad l(x) = - \sum_{k=1}^{\infty} \zeta^2 \left(\frac{2^k(y-x)}{r} \right).$$

Then $\ln \frac{|y-x|}{r} - l(x) \ln 2 \in L^\infty(B_r(y))$. Indeed, for each $x \in B_r(y)$ there must exist a $j \in \{1, 2, \dots\}$ such that

$$\frac{1}{2^j} \leq \frac{|y-x|}{r} < \frac{1}{2^{j-1}}.$$

We easily verify from the properties of ζ that

$$\zeta^2 \left(\frac{2^k(y-x)}{r} \right) = \begin{cases} 0 & \text{if } k > j, \\ 1 & \text{if } k \leq j-1. \end{cases}$$

That is, the series in (2.2) is actually a finite sum and

$$-l(x) = j - 1 + \zeta^2 \left(\frac{2^j(y-x)}{r} \right).$$

It immediately follows that

$$j - 1 \leq -l(x) \leq j, \quad -j \ln 2 \leq \ln \frac{|y-x|}{r} \leq -(j-1) \ln 2.$$

Hence,

$$(2.3) \quad \ln \frac{|y-x|}{r} - l(x) \ln 2 \leq \ln \frac{|y-x|}{r} + j \ln 2 \leq \ln 2, \\ \ln \frac{|y-x|}{r} - l(x) \ln 2 \geq \ln \frac{|y-x|}{r} + (j-1) \ln 2 \geq -\ln 2.$$

That is,

$$(2.4) \quad \left| \ln \frac{|y-x|}{r} - l(x) \ln 2 \right| \leq \ln 2.$$

We would like to remark that this type of decomposition of the logarithmic function is often found in the study of real variable Hardy spaces [17].

Let

$$A_j = B_{\frac{r}{2^{j-1}}}(y) \setminus B_{\frac{r}{2^j}}(y).$$

Then

$$B_r(y) = \cup_{j=1}^{\infty} A_j.$$

Denote by p_{A_j} the average of p over A_j , i.e.,

$$p_{A_j} = \fint_{A_j} p dx = \frac{1}{|A_j|} \int_{A_j} p dx.$$

Subsequently, we can infer from the Sobolev-Poincaré inequality ([8], p.174) that for each $s \geq 2$ there is a positive number c such that

$$(2.5) \quad \left(\fint_{A_j} |p - p_{A_j}|^s dx \right)^{\frac{1}{s}} \leq \frac{cr}{2^j} \left(\fint_{A_j} |\nabla p|^{\frac{2s}{s+2}} dx \right)^{\frac{s+2}{2s}} \leq \frac{cr}{2^j} \left(\fint_{A_j} |\nabla p|^2 dx \right)^{\frac{1}{2}}.$$

With this in mind, we use $(p - p_{A_j})\zeta^2 \left(\frac{2^j(y-x)}{r} \right)$ as a test function in (1.1) to derive

$$(2.6) \quad \begin{aligned} & \int_{B_{\frac{r}{2^{j-1}}}(y)} (|\nabla p|^2 + (\mathbf{m} \cdot \nabla p)^2) \zeta^2 \left(\frac{2^j(y-x)}{r} \right) dx \\ &= \frac{2^{j+1}}{r} \int_{A_j} \zeta \left(\frac{2^j(y-x)}{r} \right) \nabla p \cdot \nabla \zeta \left(\frac{2^j(y-x)}{r} \right) (p - p_{A_j}) dx \\ & \quad + \frac{2^{j+1}}{r} \int_{A_j} \zeta \left(\frac{2^j(y-x)}{r} \right) (\mathbf{m} \cdot \nabla p) \mathbf{m} \cdot \nabla \zeta \left(\frac{2^j(y-x)}{r} \right) (p - p_{A_j}) dx \\ & \quad + \int_{B_{\frac{r}{2^{j-1}}}(y)} S(y)(p - p_{A_j})\zeta^2 \left(\frac{2^j(y-x)}{r} \right) dx. \end{aligned}$$

We proceed to estimate each term on the right hand side. First, by (2.5), we have

$$\begin{aligned} & \frac{2^{j+1}}{r} \int_{A_j} \zeta \left(\frac{2^j(y-x)}{r} \right) \nabla p \cdot \nabla \zeta \left(\frac{2^j(y-x)}{r} \right) (p - p_{A_j}) dx \\ & \leq \frac{\|\nabla \zeta\|_{\infty, B_2(0)} 2^{j+1}}{r} \left(\int_{A_j} |\nabla p|^2 dx \right)^{\frac{1}{2}} \left(\int_{A_j} |p - p_{A_j}|^2 dx \right)^{\frac{1}{2}} \leq c \int_{A_j} |\nabla p|^2 dx. \end{aligned}$$

Similarly, for each $\varepsilon > 0$ we calculate

$$\begin{aligned}
& \frac{2^{j+1}}{r} \int_{A_j} \zeta \left(\frac{2^j(y-x)}{r} \right) (\mathbf{m} \cdot \nabla p) \mathbf{m} \cdot \nabla \zeta \left(\frac{2^j(y-x)}{r} \right) (p - p_{A_j}) dx \\
& \leq \varepsilon \int_{B_{\frac{r}{2^{j-1}}}(y)} (\mathbf{m} \cdot \nabla p)^2 \zeta^2 \left(\frac{2^j(y-x)}{r} \right) dx + \frac{\|\nabla \zeta\|_{\infty, B_2(0)}^2 4^{j+1}}{\varepsilon r^2} \int_{A_j} |\mathbf{m}|^2 |p - p_{A_j}|^2 dx \\
& \leq \varepsilon \int_{B_{\frac{r}{2^{j-1}}}(y)} (\mathbf{m} \cdot \nabla p)^2 \zeta^2 \left(\frac{2^j(y-x)}{r} \right) dx + \frac{c 4^j}{\varepsilon r^2} \left(\int_{A_j} |\mathbf{m}|^4 dx \right)^{\frac{1}{2}} \left(\int_{A_j} |p - p_{A_j}|^4 dx \right)^{\frac{1}{2}} \\
& \leq \varepsilon \int_{B_{\frac{r}{2^{j-1}}}(y)} (\mathbf{m} \cdot \nabla p)^2 \zeta^2 \left(\frac{2^j(y-x)}{r} \right) dx + \frac{c}{\varepsilon} \left(\int_{B_r(y)} |\mathbf{m}|^4 dx \right)^{\frac{1}{2}} \int_{A_j} |\nabla p|^2 dx.
\end{aligned}$$

Finally,

$$\int_{B_{\frac{r}{2^{j-1}}}(y)} S(x) (p - p_{A_j}) \zeta^2 \left(\frac{2^j(y-x)}{r} \right) dx \leq c \|S(x)\|_{q, B_r(y)} \left(\frac{r}{2^{j-1}} \right)^{\frac{2(q-1)}{q}}.$$

Substitute the preceding three estimates into (2.6) and choose ε suitably small in the resulting inequality to obtain

$$\begin{aligned}
\int_{B_{\frac{r}{2^{j-1}}}(y)} F \zeta^2 \left(\frac{2^j(y-x)}{r} \right) dx &= \int_{B_{\frac{r}{2^{j-1}}}(y)} (|\nabla p|^2 + (\mathbf{m} \cdot \nabla p)^2) \zeta^2 \left(\frac{2^j(y-x)}{r} \right) dx \\
&\leq c \int_{A_j} |\nabla p|^2 dx + c \left(\int_{B_r(y)} |\mathbf{m}|^4 dx \right)^{\frac{1}{2}} \int_{A_j} |\nabla p|^2 dx \\
(2.7) \quad &+ c \|S(x)\|_{q, B_r(y)} \left(\frac{r}{2^{j-1}} \right)^{\frac{2(q-1)}{q}}.
\end{aligned}$$

With this and (2.4) in mind, we derive

$$\begin{aligned}
\int_{B_r(y)} F |\ln |x-y|| dx &\leq \int_{B_r(y)} F |\ln |x-y| - \ln r - l(x) \ln 2| dx \\
&+ |\ln r| \int_{B_r(y)} F dx + \ln 2 \int_{B_r(y)} F |l(x)| dx \\
&\leq (\ln 2 + |\ln r|) \int_{B_r(y)} F dx + \ln 2 \int_{B_r(y)} F \sum_{j=1}^{\infty} \zeta^2 \left(\frac{2^j(y-x)}{r} \right) dx \\
&\leq (\ln 2 + |\ln r|) \int_{B_r(y)} F dx + c \int_{B_r(y)} |\nabla p|^2 dx \\
&+ c \left(\int_{B_r(y)} |\mathbf{m}|^4 dx \right)^{\frac{1}{2}} \int_{B_r(y)} |\nabla p|^2 dx + cr^{\frac{2(q-1)}{q}} \|S(x)\|_{q, B_r(y)}.
\end{aligned}$$

The lemma follows. \square

Lemma 2.2. *Let the assumptions of the preceding lemma hold. Then there is a positive constant c such that*

$$\begin{aligned}
 & \int_{B_r(y)} F \ln^2 |x - y| dx \\
 & \leq c(1 + \ln^2 r) \int_{B_r(y)} F dx + c \int_{B_r(y)} |\ln |x - y|| |\nabla p|^2 dx \\
 & \quad + c \left(\int_{B_r(y)} |\mathbf{m}|^4 dx \right)^{\frac{1}{2}} \int_{B_r(y)} |\ln |x - y|| |\nabla p|^2 dx \\
 & \quad + c(1 + |\ln r|) \left(\int_{B_r(y)} |\mathbf{m}|^4 dx \right)^{\frac{1}{2}} \int_{B_r(y)} |\nabla p|^2 dx \\
 (2.8) \quad & \quad + c(1 + |\ln r|) r^{\frac{(2-\sigma)(q-1)}{q}} \|S(x)\|_{q, B_r(y)}.
 \end{aligned}$$

for each $y \in \Omega$ and $r \in (0, \text{dist}(y, \partial\Omega))$.

Proof. We see from (2.4) that

$$\begin{aligned}
 \int_{B_r(y)} F |\ln |x - y||^2 dx & \leq 2 \int_{B_r(y)} F |\ln |x - y| - \ln r - l(x) \ln 2|^2 dx \\
 & \quad + 4 \ln^2 r \int_{B_r(y)} F dx + 4 \ln^2 2 \int_{B_r(y)} F |l(x)|^2 dx \\
 (2.9) \quad & \leq (2 \ln^2 2 + 4 \ln^2 r) \int_{B_r(y)} F dx + 4 \ln^2 2 \int_{B_r(y)} F |l(x)|^2 dx.
 \end{aligned}$$

It is easy to verify that

$$\zeta^2 \left(\frac{2^j(y-x)}{r} \right) \zeta^2 \left(\frac{2^i(y-x)}{r} \right) = \zeta^2 \left(\frac{2^i(y-x)}{r} \right) \quad \text{whenever } i < j.$$

Consequently,

$$\begin{aligned}
 |l(x)|^2 & = \sum_{i=1}^{\infty} \zeta^2 \left(\frac{2^i(y-x)}{r} \right) \sum_{j=1}^{\infty} \zeta^2 \left(\frac{2^j(y-x)}{r} \right) \\
 (2.10) \quad & = \sum_{j=1}^{\infty} \zeta^4 \left(\frac{2^j(y-x)}{r} \right) + 2 \sum_{j=1}^{\infty} (j-1) \zeta^2 \left(\frac{2^j(y-x)}{r} \right).
 \end{aligned}$$

We can infer from the proof of (2.7) that

$$\begin{aligned}
 & \int_{B_{\frac{r}{2^{j-1}}}(y)} (|\nabla p|^2 + (\mathbf{m} \cdot \nabla p)^2) \zeta^4 \left(\frac{2^j(y-x)}{r} \right) dx \\
 (2.11) \quad & \leq c \int_{A_j} |\nabla p|^2 dx + c \left(\int_{B_r(y)} |\mathbf{m}|^4 dx \right)^{\frac{1}{2}} \int_{A_j} |\nabla p|^2 dx + c \|S(x)\|_{q, B_r(y)} \left(\frac{r}{2^{j-1}} \right)^{\frac{2(q-1)}{q}}.
 \end{aligned}$$

Use $(p - p_{A_j})(j - 1)\zeta^2 \left(\frac{2^j(y-x)}{r} \right)$ as a test function in (1.1) to derive

$$\begin{aligned}
& \int_{B_{\frac{r}{2^{j-1}}}(y)} (|\nabla p|^2 + (\mathbf{m} \cdot \nabla p)^2) (j - 1)\zeta^2 \left(\frac{2^j(y-x)}{r} \right) dx \\
&= \frac{2^{j+1}}{r} \int_{A_j} (j - 1)\zeta \left(\frac{2^j(y-x)}{r} \right) \nabla p \cdot \nabla \zeta \left(\frac{2^j(y-x)}{r} \right) (p - p_{A_j}) dx \\
&\quad + \frac{2^{j+1}}{r} \int_{A_j} (j - 1)\zeta \left(\frac{2^j(y-x)}{r} \right) (\mathbf{m} \cdot \nabla p) \mathbf{m} \cdot \nabla \zeta \left(\frac{2^j(y-x)}{r} \right) (p - p_{A_j}) dx \\
(2.12) \quad &+ \int_{B_{\frac{r}{2^{j-1}}}(y)} S(y)(p - p_{A_j})(j - 1)\zeta^2 \left(\frac{2^j(y-x)}{r} \right) dx.
\end{aligned}$$

Recall from (2.3) that

$$j - 1 \leq -\frac{1}{\ln 2} \ln \frac{|x - y|}{r}.$$

Keep this in mind to deduce for each $\varepsilon > 0$ that

$$\begin{aligned}
& \frac{2^{j+1}}{r} \int_{A_j} (j - 1)\zeta \left(\frac{2^j(y-x)}{r} \right) \nabla p \cdot \nabla \zeta \left(\frac{2^j(y-x)}{r} \right) (p - p_{A_j}) dx \\
&\leq \varepsilon \int_{B_{\frac{r}{2^{j-1}}}(y)} |\nabla p|^2 (j - 1)\zeta^2 \left(\frac{2^j(y-x)}{r} \right) dx + \frac{\|\nabla \zeta\|_{\infty, B_2(0)}^2 4^{j+1} (j - 1)}{r^2 \varepsilon} \int_{A_j} |p - p_{A_j}|^2 dx \\
&\leq \varepsilon \int_{B_{\frac{r}{2^{j-1}}}(y)} |\nabla p|^2 (j - 1)\zeta^2 \left(\frac{2^j(y-x)}{r} \right) dx + \frac{c}{\varepsilon} \int_{A_j} (j - 1) |\nabla p|^2 dx \\
&\leq \varepsilon \int_{B_{\frac{r}{2^{j-1}}}(y)} |\nabla p|^2 (j - 1)\zeta^2 \left(\frac{2^j(y-x)}{r} \right) dx + \frac{c}{\varepsilon} \int_{A_j} |\ln |x - y| - \ln r| |\nabla p|^2 dx.
\end{aligned}$$

By the same token,

$$\begin{aligned}
& \frac{2^{j+1}}{r} \int_{A_j} (j - 1)\zeta \left(\frac{2^j(y-x)}{r} \right) (\mathbf{m} \cdot \nabla p) \mathbf{m} \cdot \nabla \zeta \left(\frac{2^j(y-x)}{r} \right) (p - p_{A_j}) dx \\
&\leq \varepsilon \int_{B_{\frac{r}{2^{j-1}}}(y)} (\mathbf{m} \cdot \nabla p)^2 (j - 1)\zeta^2 \left(\frac{2^j(y-x)}{r} \right) dx \\
&\quad + \frac{c 4^j (j - 1)}{\varepsilon r^2} \left(\int_{A_j} |\mathbf{m}|^4 dx \right)^{\frac{1}{2}} \left(\int_{A_j} |p - p_{A_j}|^4 dx \right)^{\frac{1}{2}} \\
&\leq \varepsilon \int_{B_{\frac{r}{2^{j-1}}}(y)} (\mathbf{m} \cdot \nabla p)^2 (j - 1)\zeta^2 \left(\frac{2^j(y-x)}{r} \right) dx \\
&\quad + \frac{c}{\varepsilon} \left(\int_{B_r(y)} |\mathbf{m}|^4 dx \right)^{\frac{1}{2}} \int_{A_j} |\ln |x - y| - \ln r| |\nabla p|^2 dx.
\end{aligned}$$

The last term in (2.12) can be estimated as follows:

$$\begin{aligned}
 & \int_{B_{\frac{r}{2^{j-1}}}(y)} S(x)(p - p_{A_j})(j-1)\zeta^2 \left(\frac{2^j(y-x)}{r} \right) dx \\
 & \leq c \int_{B_{\frac{r}{2^{j-1}}}(x)} |S(x)| |\ln|x-y| - \ln r| dx \\
 & \leq c |\ln r| \|S(x)\|_{q, B_r(y)} \left(\frac{r}{2^{j-1}} \right)^{\frac{2(q-1)}{q}} + c \|S(x)\|_{q, B_r(y)} \left(\frac{r}{2^{j-1}} \right)^{\frac{(2-\sigma)(q-1)}{q}}, \quad \sigma \in (0, 2).
 \end{aligned}$$

Collect the preceding three estimates in (2.12) to deduce

$$\begin{aligned}
 & \int_{B_{\frac{r}{2^{j-1}}}(y)} (|\nabla p|^2 + (\mathbf{m} \cdot \nabla p)^2) (j-1)\zeta^2 \left(\frac{2^j(y-x)}{r} \right) dx \\
 & \leq c \int_{A_j} |\ln|x-y||\nabla p|^2 dx + c \left(\int_{B_r(y)} |\mathbf{m}|^4 dx \right)^{\frac{1}{2}} \int_{A_j} |\ln|x-y||\nabla p|^2 dx \\
 & \quad + c |\ln r| \int_{A_j} |\nabla p|^2 dx + c |\ln r| \left(\int_{B_r(y)} |\mathbf{m}|^4 dx \right)^{\frac{1}{2}} \int_{A_j} |\nabla p|^2 dx \\
 & \quad + c(1 + |\ln r|) \|S(x)\|_{q, B_r(y)} \left(\frac{r}{2^{j-1}} \right)^{\frac{(2-\sigma)(q-1)}{q}}.
 \end{aligned}$$

Combining this with (2.11) and (2.10) yields

$$\begin{aligned}
 & \int_{B_r(y)} F |l^2(x)| dx \\
 & \leq \int_{B_r(y)} F \sum_{j=1}^{\infty} \zeta^4 \left(\frac{2^j(y-x)}{r} \right) dx + 2 \int_{B_r(y)} F \sum_{j=1}^{\infty} (j-1)\zeta^2 \left(\frac{2^j(y-x)}{r} \right) dx \\
 & \leq c \int_{B_r(y)} |\ln|x-y||\nabla p|^2 dx + c \left(\int_{B_r(y)} |\mathbf{m}|^4 dx \right)^{\frac{1}{2}} \int_{B_r(y)} |\ln|x-y||\nabla p|^2 dx \\
 & \quad + c(1 + |\ln r|) \int_{B_r(y)} |\nabla p|^2 dx + c(1 + |\ln r|) \left(\int_{B_r(y)} |\mathbf{m}|^4 dx \right)^{\frac{1}{2}} \int_{B_r(y)} |\nabla p|^2 dx \\
 & \quad + c(1 + |\ln r|) r^{\frac{(2-\sigma)(q-1)}{q}} \|S(x)\|_{q, B_r(y)}.
 \end{aligned}$$

Use this in (2.9) to obtain (2.8). The proof is complete. \square

We can also verify that

$$\begin{aligned}
 |l(x)|^3 & = \sum_{j=1}^{\infty} \zeta^6 \left(\frac{2^j(y-x)}{r} \right) + 2 \sum_{j=1}^{\infty} (j-1)\zeta^4 \left(\frac{2^j(y-x)}{r} \right) + \sum_{j=1}^{\infty} (j-1)\zeta^2 \left(\frac{2^j(y-x)}{r} \right) \\
 & \quad + \sum_{j=1}^{\infty} (j-1)^2 \zeta^2 \left(\frac{2^j(y-x)}{r} \right) + \sum_{j=2}^{\infty} \frac{(j-1)(j-2)}{2} \zeta^2 \left(\frac{2^j(y-x)}{r} \right).
 \end{aligned}$$

It is not difficult to deduce that for each positive integer n we have

$$|l(x)|^n = \sum_{j=1}^{\infty} \zeta^{2n} \left(\frac{2^j(y-x)}{r} \right) + \text{lower order terms.}$$

Our earlier proof indicates that

$$\int_{B_r(y)} F |\ln |x - y||^{n-1} dx < \infty \implies \int_{B_r(y)} F |\ln |x - y||^n dx < \infty.$$

In summary, we have:

Lemma 2.3. *Let the assumptions of the preceding lemma hold. For each positive integer $n, y \in \Omega, r \in (0, \text{dist}(y, \partial\Omega))$ there is a positive number c such that*

$$(2.13) \quad \int_{B_r(y)} F |\ln |x - y||^n dx \leq c.$$

We are ready to prove (C3).

Proof of (C3). Let $y \in \Omega, R \in (0, \min\{1, \text{dist}(y, \partial\Omega)\})$ be given. For each $r \in (0, R]$, we consider the boundary value problem

$$(2.14) \quad -\text{div} [(I + \mathbf{m} \otimes \mathbf{m}) \nabla p_1] = S(x) \quad \text{in } B_r(y),$$

$$(2.15) \quad p_1 = 0 \quad \text{on } \partial B_r(y).$$

Even though the elliptic coefficients in (2.14) may not be bounded, we can easily infer from the proof of Lemma 2.3 in [18] that this problem has a unique solution $p_1 = p_1(x, t)$ in the sense of (D1)-(D2).

Claim 2.4. *For each $q > 1$ there is a positive number $c = c(N, q)$ such that*

$$(2.16) \quad \text{ess sup}_{B_r(y)} |p_1| \leq cr^{\frac{2(q-1)}{q}} \left(\int_{B_r(y)} |S(x)|^q dx \right)^{\frac{1}{q}}.$$

Proof. This result was established in [14] for $N > 2$. The proof is similar for $N = 2$. For completeness, we offer a proof here. We employ the classical Moser-Nash-De Giorgi type of arguments. Without loss of generality, we assume

$$\text{ess sup}_{B_r(y)} p_1 = \text{ess sup}_{B_r(y)} |p_1|.$$

Let κ be a positive number to be determined. Write

$$\kappa_n = \kappa - \frac{\kappa}{2^n}, \quad A_n = \{x \in B_r(y) : p_1(x, t) \geq \kappa_n\}, \quad n = 0, 1, 2, \dots$$

Use $(p_1 - \kappa_n)^+$ as a test function in (2.14) to deduce

$$(2.17) \quad \int_{B_r(y)} |\nabla(p_1 - \kappa_n)^+|^2 dx + \int_{B_r(y)} (\mathbf{m} \cdot \nabla(p_1 - \kappa_n)^+)^2 dx = \int_{B_r(y)} S(x)(p_1 - \kappa_n)^+ dx.$$

Here we have used the fact that

$$\nabla p_1 \chi_{A_n} = \nabla(p_1 - \kappa_n)^+.$$

By Poincaré's inequality,

$$\begin{aligned} \int_{B_r(y)} S(x)(p_1 - \kappa_n)^+ dx &\leq \left(\int_{B_r(y)} |S(x)|^q dx \right)^{\frac{1}{q}} \left(\int_{B_r(y)} [(p_1 - \kappa_n)^+]^{\frac{q}{q-1}} dx \right)^{\frac{q-1}{q}} \\ &\leq c \|S(x)\|_{q, B_r(y)} \left(\int_{B_r(y)} |\nabla(p_1 - \kappa_n)^+|^{\frac{2q}{3q-2}} dx \right)^{\frac{3q-2}{2q}} \\ &\leq c \|S(x)\|_{q, B_r(y)} \left(\int_{B_r(y)} |\nabla(p_1 - \kappa_n)^+|^2 dx \right)^{\frac{1}{2}} |A_n|^{\frac{q-1}{q}}. \end{aligned}$$

Use this in (2.17) to obtain

$$\left(\int_{B_r(y)} |\nabla(p_1 - \kappa_n)^+|^2 dx \right)^{\frac{1}{2}} \leq c \|S(x)\|_{q, B_r(y)} |A_n|^{\frac{q-1}{q}}.$$

For each $s > 1$ we have

$$\begin{aligned} \frac{\kappa}{2^{n+1}} |A_{n+1}|^{\frac{1}{s}} &\leq \left(\int_{B_r(y)} [(p_1 - \kappa_n)^+]^s dx \right)^{\frac{1}{s}} \\ &\leq c \left(\int_{B_r(y)} |\nabla(p_1 - \kappa_n)^+|^{\frac{2s}{s+2}} dx \right)^{\frac{2+s}{2s}} \\ &\leq c \left(\int_{B_r(y)} |\nabla(p_1 - \kappa_n)^+|^2 dx \right)^{\frac{1}{2}} r^{\frac{2}{s}} \\ &\leq cr^{\frac{2}{s}} \|S(x)\|_{q, B_r(y)} |A_n|^{\frac{q-1}{q}}, \end{aligned}$$

from whence follows

$$|A_{n+1}| \leq cr^2 \|S(x)\|_{q, B_r(y)}^s \frac{2^{(n+1)s}}{\kappa^s} |A_n|^{\frac{(q-1)s}{q}}.$$

Now we pick s so large that

$$\alpha \equiv \frac{(q-1)s}{q} - 1 = \frac{s(q-1) - q}{q} > 0.$$

We are in a position to apply Lemma 4.1 in ([4], p. 12). To this end, we choose κ so large that

$$A_0 \leq |B_r(y)| \leq c \left(\frac{\kappa^s}{r^2 \|S(x)\|_{q, B_r(y)}^s} \right)^{\frac{q}{s(q-1)-q}}.$$

Then

$$p_1 \leq \kappa \quad \text{on } B_r(y).$$

It is enough for us to take

$$\kappa = cr^{\frac{2(q-1)}{q}} \|S(x)\|_{q, B_r(y)}.$$

The lemma follows. The proof is complete. \square

Obviously, $p_0 \equiv p - p_1$ satisfies

$$\begin{aligned} -\operatorname{div} [(I + \mathbf{m} \otimes \mathbf{m}) \nabla p_0] &= 0 \quad \text{in } B_r(y), \\ p_0(x, t) &= p(x, t) \quad \text{on } \partial B_r(y) \end{aligned}$$

in the sense of (D1)-(D2) with an obvious modification to the boundary condition. Obviously, the weak maximum principle still holds, from which it follows that

$$\operatorname{osc}_{B_r(y)} p_0 = \operatorname{osc}_{\partial B_r(y)} p_0 = \operatorname{osc}_{\partial B_r(y)} p \leq \int_{\partial B_r(y)} |\nabla p| ds.$$

(Here we may assume that p is a smooth function because p can be viewed as the limit of a sequence of smooth functions.) For each $\ell > 1$ and $\tau \in (0, R)$ we multiply through the above inequality by

$\frac{|\ln r|^\ell}{r}$ and then integrate the resulting one over $[\tau, R]$ to derive

$$\begin{aligned}
\int_\tau^R \frac{|\ln r|^\ell}{r} \operatorname{osc}_{B_r(y)} p_0 dr &\leq \int_\tau^R \frac{|\ln r|^\ell}{r} \int_{\partial B_r(y)} |\nabla p| ds dr \\
&\leq \int_{B_R(y) \setminus B_\tau(y)} \frac{|\nabla p| |\ln |x-y||^\ell}{|x-y|} dx \\
&\leq \left(\int_{B_R(y)} |\nabla p|^2 |\ln |x-y||^{3\ell} dx \right)^{\frac{1}{2}} \left(\int_{B_R(y) \setminus B_\tau(y)} \frac{1}{|\ln |x-y||^\ell |x-y|^2} dx \right)^{\frac{1}{2}} \\
&\leq c(R, \ell) \left(\int_0^R \frac{1}{|\ln r|^\ell r} dr \right)^{\frac{1}{2}} \leq c(R, \ell).
\end{aligned}$$

Here we have employed (2.13). Observe that $\operatorname{osc}_{B_r(y)} p_0$ is an increasing function of r . With this in mind, we obtain

$$\begin{aligned}
\operatorname{osc}_{B_\tau(y)} p_0 &\leq \frac{\int_\tau^R \frac{|\ln r|^\ell}{r} \operatorname{osc}_{B_r(y)} p_0 dr}{\int_\tau^R \frac{|\ln r|^\ell}{r} dr} \\
&\leq \frac{c(R, \ell)}{\int_0^{\ln \frac{R}{\tau}} (s - \ln R)^\ell ds} \leq \frac{c(R, \ell)}{\ln^{\ell+1} \frac{R}{\tau}}.
\end{aligned}$$

Here we have used the fact that $\ln R < 0$. The above inequality together with (2.16) implies

$$\begin{aligned}
\operatorname{osc}_{B_\tau(y)} p &\leq \operatorname{osc}_{B_\tau(y)} p_0 + \operatorname{osc}_{B_\tau(y)} p_1 \\
&\leq \frac{c(R, \ell)}{\ln^{\ell+1} \frac{R}{\tau}} + c\tau^{\frac{2(q-1)}{q}} \quad \text{for } \tau \in (0, R).
\end{aligned}$$

This finishes the proof of (C3). □

Lemma 2.5. *Let $x_0 \in \Omega$ be given and define*

$$d(x_0) = \operatorname{dist}(x_0, \partial\Omega).$$

If $\mathbf{m} \in (\operatorname{BMO}(\Omega))^N$, i.e.,

$$\|\mathbf{m}\|_{\operatorname{BMO}(\Omega)} \equiv \sup_{B_r(y) \subset \Omega} \int |\mathbf{m} - \mathbf{m}_{y,r}| dx < \infty,$$

then there exists a positive number $c = c(\|\mathbf{m}\|_{\operatorname{BMO}(\Omega)}, d(x_0))$ such that

$$|\mathbf{m}_{x_0, \rho}| \leq c \ln \left(\frac{d(x_0)}{\rho} \right) + c \quad \text{for each } \rho \in (0, d(x_0)).$$

Proof. We follow the proof of Proposition 1.2 in ([7], p. 68). For each $0 < \rho \leq r \leq d(x_0)$, we have

$$|\mathbf{m}_{x_0, r} - \mathbf{m}_{x_0, \rho}| \leq |\mathbf{m}_{x_0, r} - \mathbf{m}(x)| + |\mathbf{m}(x) - \mathbf{m}_{x_0, \rho}|.$$

Integrate the above inequality over $B_\rho(x_0)$ to derive

$$|\mathbf{m}_{x_0, r} - \mathbf{m}_{x_0, \rho}| \leq \left(\frac{r}{\rho} \right)^2 \int_{B_r(x_0)} |\mathbf{m}_{x_0, r} - \mathbf{m}| dx + \int_{B_\rho(x_0)} |\mathbf{m} - \mathbf{m}_{x_0, \rho}| dx \leq c \left(\frac{r}{\rho} \right)^2.$$

In particular, take $r = \frac{d(x_0)}{2^i}$, $\rho = \frac{d(x_0)}{2^{i+1}}$, where $i \in \{0, 1, \dots\}$. Then we have

$$\left| \mathbf{m}_{x_0, \frac{d(x_0)}{2^i}} - \mathbf{m}_{x_0, \frac{d(x_0)}{2^{i+1}}} \right| \leq c.$$

For each $\rho \in (0, d(x_0))$ there is an $i \in \{0, 1, \dots\}$ such that

$$(2.18) \quad \frac{d(x_0)}{2^{i+1}} \leq \rho < \frac{d(x_0)}{2^i}.$$

Subsequently,

$$(2.19) \quad \begin{aligned} |\mathbf{m}_{x_0, \rho}| &\leq \left| \mathbf{m}_{x_0, \frac{d(x_0)}{2^i}} \right| + \left| \mathbf{m}_{x_0, \rho} - \mathbf{m}_{x_0, \frac{d(x_0)}{2^i}} \right| \\ &\leq \left| \mathbf{m}_{x_0, \frac{d(x_0)}{2^i}} - \mathbf{m}_{x_0, d(x_0)} \right| + |\mathbf{m}_{x_0, d(x_0)}| + c \\ &\leq \sum_{j=1}^i \left| \mathbf{m}_{x_0, \frac{d(x_0)}{2^j}} - \mathbf{m}_{x_0, \frac{d(x_0)}{2^{j-1}}} \right| + |\mathbf{m}_{x_0, d(x_0)}| + c \\ &\leq c(i+1) + |\mathbf{m}_{x_0, d(x_0)}|. \end{aligned}$$

We easily see from (2.18) that

$$i \leq \frac{1}{\ln 2} \ln \left(\frac{d(x_0)}{\rho} \right).$$

Using this in (2.19) yields the desired result. \square

We are in a position to prove (C4).

Proof of (C4). Let $y \in \Omega, r > 0$ be such that $B_{2r}(y) \subset \Omega$. Then for each $x_0 \in B_r(y)$ and $\rho \in (0, \frac{r}{2})$, we have $B_{2\rho}(x_0) \subset \Omega$. Let ρ be so chosen. Select a cutoff function $\zeta \in C_0^\infty(B_{2\rho}(x_0))$ the properties

$$\zeta = 1 \text{ on } B_\rho(x_0), 0 \leq \zeta \leq 1 \text{ on } B_{2\rho}(x_0), \text{ and } |\nabla \zeta| \leq \frac{c}{\rho} \text{ on } B_{2\rho}(x_0).$$

Use $(p - p_{x_0, 2\rho})\zeta^2$ as a test function in (1.1) to obtain

$$\begin{aligned} &\int_{B_{2\rho}(x_0)} |\nabla p|^2 \zeta^2 dx + \int_{B_{2\rho}(x_0)} (\mathbf{m} \cdot \nabla p)^2 \zeta^2 dx \\ &= - \int_{B_{2\rho}(x_0)} \nabla p \cdot \nabla \zeta \zeta (p - p_{x_0, 2\rho}) dx - \int_{B_{2\rho}(x_0)} (\mathbf{m} \cdot \nabla p) \mathbf{m} \cdot \nabla \zeta \zeta (p - p_{x_0, 2\rho}) dx \\ &\quad + \int_{B_{2\rho}(x_0)} S(x) (p - p_{x_0, 2\rho}) \zeta^2 dx \\ &\leq \frac{1}{2} \int_{B_{2\rho}(x_0)} (|\nabla p|^2 + (\mathbf{m} \cdot \nabla p)^2) \zeta^2 dx + \frac{c}{\rho^2} \int_{B_{2\rho}(x_0)} (p - p_{x_0, 2\rho})^2 dx \\ &\quad + \frac{c}{\rho^2} \int_{B_{2\rho}(x_0)} |\mathbf{m}|^2 (p - p_{x_0, 2\rho})^2 dx + \int_{B_{2\rho}(x_0)} S(x) (p - p_{x_0, 2\rho}) \zeta^2 dx, \end{aligned}$$

from whence follows

$$(2.20) \quad \begin{aligned} \int_{B_\rho(x_0)} F dx &= \int_{B_\rho(x_0)} |\nabla p|^2 dx + \int_{B_\rho(x_0)} (\mathbf{m} \cdot \nabla p)^2 dx \\ &\leq c \left(\text{osc}_{B_{2\rho}(x_0)} p \right)^2 + \frac{c}{\rho^2} \int_{B_{2\rho}(x_0)} |\mathbf{m} - \mathbf{m}_{x_0, 2\rho}|^2 (p - p_{x_0, 2\rho})^2 dx \\ &\quad + c |\mathbf{m}_{x_0, 2\rho}|^2 \left(\text{osc}_{B_{2\rho}(x_0)} p \right)^2 + c \|S\|_{q, B_\rho(x_0)} \rho^{\frac{2(q-1)}{q}} \text{osc}_{B_{2\rho}(x_0)} p \\ &\leq c (1 + (|\ln \rho| + 1)^2) \left(\text{osc}_{B_{2\rho}(x_0)} p \right)^2 + c \rho^{\frac{2(q-1)}{q}}. \end{aligned}$$

Here we have applied the John-Nirenberg Theorem [5], which asserts that for each $s > 1$ and subdomain Ω_0 of Ω with $\overline{\Omega_0} \subset \Omega$ there is a constant c such that

$$(2.21) \quad \sup_{B_r(y) \subset \Omega_0} \left(\int_{B_r(y)} |\mathbf{m} - \mathbf{m}_{y,r}|^s dx \right)^{\frac{1}{s}} \leq c \|\mathbf{m}\|_{\text{BMO}(\Omega)}.$$

We derive from (2.1), (2.20), and (C3) that

$$(2.22) \quad \begin{aligned} & \int_{B_\rho(x_0)} F \chi_{B_r(y)} |\ln|x - x_0|| dx \\ & \leq \int_{B_\rho(x_0)} F |\ln|x - x_0|| dx \\ & \leq (\ln 2 + |\ln \rho|) \int_{B_\rho(x_0)} F dx + c \int_{B_\rho(x_0)} |\nabla p|^2 dx \\ & \quad + c \left(\int_{B_\rho(x_0)} |\mathbf{m}|^4 dx \right)^{\frac{1}{2}} \int_{B_\rho(x_0)} |\nabla p|^2 dx + c \rho^{\frac{2(q-1)}{q}} \|S(x)\|_{q, B_\rho(x_0)} \\ & \leq (c + |\ln \rho|) \int_{B_\rho(x_0)} F dx + c \rho^{\frac{2(q-1)}{q}} \\ & \leq c(1 + (|\ln \rho| + 1)^2) (c + |\ln \rho|) \left(\text{osc}_{B_{2\rho}(x_0)} p \right)^2 + c(c + |\ln \rho|) \rho^{\frac{2(q-1)}{q}} \\ & \leq \frac{c(1 + (|\ln \rho| + 1)^2) (c + |\ln \rho|)}{\ln^{2\ell} \frac{R}{\rho}} + c(c + |\ln \rho|) \rho^{\frac{2(q-1)}{q}} \rightarrow 0 \text{ as } \rho \rightarrow 0, \end{aligned}$$

provided that $\ell > \frac{3}{2}$. The proof is complete. \square

3. PROOF OF THEOREM 1.5

In this section we prove the last two results in the introduction.

Proof of Theorem 1.5. For each $r \in (0, 1)$ and $x_0 \in \Omega$ such that $B_{2r}(x_0) \subset \Omega$, we define

$$M_{x_0,r} = \text{ess sup}_{B_r(x_0)} p, \quad m_{x_0,r} = \text{ess inf}_{B_r(x_0)} p, \quad \omega_{x_0,r} = \text{osc}_{B_r(x_0)} p = M_{x_0,r} - m_{x_0,r}.$$

Let q be given as in (H1). Then we can pick a number

$$(3.1) \quad \delta \in \left(0, \frac{2q-2}{q} \right).$$

Subsequently, set

$$\bar{k}(r) = r^\delta \|S\|_{q, B_r(x_0)}.$$

Either

$$(3.2) \quad \left| B_r(x_0) \cap \left\{ p(x) \leq m_{x_0,r} + \frac{\omega_{x_0,r}}{2} \right\} \right| \geq \frac{1}{2} |B_r(x_0)|, \quad \text{or}$$

$$(3.3) \quad \left| B_r(x_0) \cap \left\{ p(x) > m_{x_0,r} + \frac{\omega_{x_0,r}}{2} \right\} \right| \geq \frac{1}{2} |B_r(x_0)|.$$

If (3.2) is true, we consider the function

$$w = \ln \left(\frac{\omega_{x_0,r} + \bar{k}(r)}{2(M_{x_0,r} - p) + \bar{k}(r)} \right).$$

If (3.3) holds, we take

$$w = \ln \left(\frac{\omega_{x_0,r} + \bar{k}(r)}{2(p - m_{x_0,r}) + \bar{k}(r)} \right).$$

For definiteness, we assume the first case (3.2). Then we easily verify that w satisfies the equation

$$(3.4) \quad \begin{aligned} & -\operatorname{div} [(I + \mathbf{m} \otimes \mathbf{m}) \nabla w] + (I + \mathbf{m} \otimes \mathbf{m}) \nabla w \cdot \nabla w \\ & = \frac{2S}{2(M_{x_0,r} - p) + \bar{k}(r)} \text{ in } B_r(x_0). \end{aligned}$$

The rest of the proof is divided into several claims.

Claim 3.1. *There is a $\delta_0 \in (0, 1)$ such that*

$$\int_{B_{(1-\delta_0)r}(x_0)} (w^+)^2 dx \leq c \int_{B_r(x_0)} |\mathbf{m}|^2 dx + cr^{2-\frac{2}{q}-\delta} + c.$$

Proof. Select $\delta_0 \in (0, 1)$ as below. Let $\theta(\eta)$ be a smooth decreasing function on $[0, r]$ with the properties

$$\theta(\eta) = 1 \text{ for } \eta \leq (1 - \delta_0)r, \theta(r) = 0, \text{ and } |\theta'(\eta)| \leq \frac{c}{r\delta_0} \text{ on } [0, r].$$

We easily see that

$$|B_r(x_0) \setminus B_{(1-\delta_0)r}(x_0)| \leq N\delta_0 |B_r(x_0)|.$$

We can pick δ_0 so that

$$(3.5) \quad |\{\theta(|x - x_0|) = 1\} \cap \{x \in B_r, w^+(x, t) = 0\}| \geq c_0 |B_r(x_0)| \text{ some } c_0 > 0.$$

To see this, take

$$N\delta_0 = \frac{1}{16}.$$

Note that

$$w^+(x, t) = 0 \Leftrightarrow x \in B_r(x_0) \cap \left\{ w(x, t) \leq m_{x_0,r} + \frac{\omega_{x_0,r}}{2} \right\}.$$

Combining this with (3.2) yields (3.5) with $c_0 = \frac{7}{16}$. Now we are in a position to apply Proposition 2.1 in ([4], p.5). Upon doing so, we obtain

$$(3.6) \quad \int_{B_r(x_0)} \theta^2(|x - x_0|) (w^+)^2 dx \leq cr^2 \int_{B_r(x_0)} \theta^2(|x - x_0|) |\nabla w|^2 dx.$$

Use θ^2 as a test function in (3.4) to derive

$$(3.7) \quad \begin{aligned} & \int_{B_r(x_0)} (|\nabla w|^2 + (\mathbf{m} \cdot \nabla w)^2) \theta^2 dx \\ & = -2 \int_{B_r(x_0)} \nabla w \cdot \nabla \theta \theta dx - 2 \int_{B_r(x_0)} (\mathbf{m} \cdot \nabla w) \mathbf{m} \cdot \nabla \theta \theta dx \\ & + \int_{B_r(x_0)} \frac{2S\theta^2}{2(M_{x_0,r} - p) + \bar{k}(r)} dx. \end{aligned}$$

We easily see that

$$\frac{1}{2(M_{x_0,r} - p) + r^\delta \|S\|_{q, B_r(x_0)}} \leq \frac{1}{r^\delta \|S\|_{q, B_r(x_0)}}.$$

We shall always use this inequality when dealing with the last term in (3.4). Indeed, with this in mind, we can derive from (3.7) that

$$\int_{B_r(x_0)} |\nabla w|^2 \theta^2 dx \leq c + \frac{c}{r^2} \int_{B_r(x_0)} |\mathbf{m}|^2 dx + cr^{2-\frac{2}{q}-\delta}.$$

This together with (3.6) yields

$$\begin{aligned}
\int_{B_{(1-\delta_0)r}(x_0)} (w^+)^2 dx &\leq \int_{B_r(x_0)} \theta^2 (w^+)^2 dx \\
&\leq cr^2 \int_{B_r(x_0)} \theta^2 |\nabla w|^2 dx \\
&\leq cr^2 + cr^2 \int_{B_r(x_0)} |\mathbf{m}|^2 dx + cr^{4-\frac{2}{q}-\delta}.
\end{aligned}$$

The lemma follows. □

Claim 3.2. *We have*

$$\sup_{B_{\frac{r}{2}}(x_0)} w \leq c \left(\int_{B_r(x_0)} (w^+)^2 dx \right)^{\frac{1}{2}} + \frac{1}{r} \|\mathbf{m}\|_{4, B_r(x_0)}^2 + r^{2-\delta-\frac{2}{q}}.$$

Proof. We employ the classical De Giorgi iteration scheme. For this purpose, we define

$$r_n = \frac{r}{2} + \frac{r}{2^{1+n}}, \quad n = 0, 1, \dots.$$

Select a sequence of smooth functions $\zeta_n(x)$ so that

$$\zeta_n(x) = 1 \text{ on } B_{r_n}(x_0), \zeta_n(x) = 0 \text{ outside } B_{r_{n-1}}(x_0), |\nabla \zeta_n| \leq \frac{c2^n}{r}, \text{ and } 0 \leq \zeta_n \leq 1.$$

Choose $k > 0$ as below. Set

$$k_n = k - \frac{k}{2^{n+1}} \text{ for } n = 0, 1, \dots.$$

Use $(w - k_{n+1})^+ \zeta_{n+1}^2$ as a test function in (3.4) and make use of the fact that

$$\nabla w = \nabla(w - k_{n+1})^+ \text{ on } \{w \geq k_{n+1}\}$$

to derive

$$\begin{aligned}
&\int_{B_{r_n}(x_0)} \left(|\nabla(w - k_{n+1})^+|^2 + [\mathbf{m} \cdot \nabla(w - k_{n+1})^+]^2 \right) \zeta_{n+1}^2 dx \\
&\quad + \int_{B_{r_n}(x_0)} (I + \mathbf{m} \otimes \mathbf{m}) \nabla(w - k_{n+1})^+ \cdot \nabla(w - k_{n+1})^+ (w - k_{n+1})^+ \zeta_{n+1}^2 dx \\
(3.8) \quad &= -2 \int_{B_{r_n}(x_0)} \nabla(w - k_{n+1})^+ \cdot \nabla \zeta_{n+1} \zeta_{n+1} (w - k_{n+1})^+ dx \\
&\quad - 2 \int_{B_{r_n}(x_0)} \mathbf{m} \cdot \nabla(w - k_{n+1})^+ \mathbf{m} \cdot \nabla \zeta_{n+1} \zeta_{n+1} (w - k_{n+1})^+ dx \\
&\quad + \int_{B_{r_n}(x_0)} \frac{2S(w - k_{n+1})^+ \zeta_{n+1}^2}{2(M_{x_0, r} - p) + r^\delta \|S\|_{q, B_r(x_0)}} dx.
\end{aligned}$$

Fix $s \geq \max\{2, \frac{q}{q-1}\}$ so that

$$\frac{s}{s-1} \leq q \text{ and } \frac{2s}{s+2} \geq 1.$$

We can use the Sobolev inequality to estimate the last integral in (3.8) as follows:

$$\begin{aligned}
 & \int_{B_{r_n}(x_0)} \frac{2S(w - k_{n+1})^+ \zeta_{n+1}^2}{2(M_{x_0, r} - p) + r^\delta \|S\|_{q, B_r(x_0)}} dx \\
 & \leq \frac{2}{r^\delta \|S\|_{q, B_r(x_0)}} \|S\|_{\frac{s}{s-1}, B_{r_n}(x_0) \cap \{w \geq k_{n+1}\}} \|(w - k_{n+1})^+ \zeta_{n+1}\|_{s, B_{r_n}(x_0)} \\
 & \leq \frac{2}{r^\delta} |B_{r_n}(x_0) \cap \{w \geq k_{n+1}\}|^{\frac{s-1}{s} - \frac{1}{q}} \|\nabla [(w - k_{n+1})^+ \zeta_{n+1}]\|_{\frac{2s}{s+2}, B_{r_n}(x_0)} \\
 & \leq \frac{2}{r^\delta} |B_{r_n}(x_0) \cap \{w \geq k_{n+1}\}|^{1 - \frac{1}{q}} \|\nabla [(w - k_{n+1})^+ \zeta_{n+1}]\|_{2, B_{r_n}(x_0)} \\
 & \leq \varepsilon \|\nabla [(w - k_{n+1})^+ \zeta_{n+1}]\|_{2, B_{r_n}(x_0)}^2 + \frac{c}{\varepsilon} |B_{r_n}(x_0) \cap \{w \geq k_{n+1}\}|^{2 - \delta - \frac{2}{q}}, \quad \varepsilon > 0.
 \end{aligned}$$

Here we have used the fact that $|B_{r_n}(x_0) \cap \{w \geq k_{n+1}\}| \leq cr^2$. Substitute this into (3.8), choose ε suitably small in the resulting equation, and thereby obtain

$$\begin{aligned}
 & \int_{B_{r_n}(x_0)} |\nabla [(w - k_{n+1})^+ \zeta_{n+1}]|^2 dx \\
 & \leq \frac{cA^n}{r} y_n + \frac{cA^n}{r^2} \int_{B_{r_n}(x_0)} |\mathbf{m}|^2 [(w - k_{n+1})^+]^2 dx \\
 & \quad + c |B_{r_n}(x_0) \cap \{w \geq k_{n+1}\}|^{2 - \delta - \frac{2}{q}},
 \end{aligned}$$

where

$$y_n = \left(\int_{B_{r_n}(x_0)} [(w - k_n)^+]^4 dx \right)^{\frac{1}{2}}.$$

It is easy to see that

$$(3.9) \quad y_n \geq \left(\int_{B_{r_n}(x_0) \cap \{w > k_{n+1}\}} [(w - k_n)^+]^4 dx \right)^{\frac{1}{2}} \geq \frac{k^2}{4^{n+2}} |B_{r_n}(x_0) \cap \{w > k_{n+1}\}|^{\frac{1}{2}}.$$

Let s be given as before. We derive from the Sobolev inequality that

$$\begin{aligned}
 (3.10) \quad y_{n+1} & \leq \int_{B_{r_n}(x_0)} [(w - k_{n+1})^+ \zeta_{n+1}]^2 dx \\
 & \leq \left(\int_{B_{r_n}(x_0)} [(w - k_{n+1})^+ \zeta_{n+1}]^{2s} dx \right)^{\frac{1}{s}} |B_{r_n}(x_0) \cap \{w > k_{n+1}\}|^{\frac{s-1}{s}} \\
 & \leq c \left(\int_{B_{r_n}(x_0)} |\nabla [(w - k_{n+1})^+ \zeta_{n+1}]|^{\frac{2s}{1+s}} dx \right)^{\frac{1+s}{s}} |B_{r_n}(x_0) \cap \{w > k_{n+1}\}|^{\frac{s-1}{s}} \\
 & \leq c \int_{B_{r_n}(x_0)} |\nabla [(w - k_{n+1})^+ \zeta_{n+1}]|^2 dx |B_{r_n}(x_0) \cap \{w > k_{n+1}\}| \\
 & \leq \frac{cA^n}{r^2} y_n |B_{r_n}(x_0) \cap \{w > k_{n+1}\}| \\
 & \quad + \frac{cA^n}{r^2} \int_{B_{r_n}(x_0)} |\mathbf{m}|^2 [(w - k_{n+1})^+]^2 dx |B_{r_n}(x_0) \cap \{w > k_{n+1}\}| \\
 & \quad + c |B_{r_n}(x_0) \cap \{w \geq k_{n+1}\}|^{3 - \delta - \frac{2}{q}}.
 \end{aligned}$$

Set

$$\sigma = \min \left\{ 2 - \delta - \frac{2}{q}, \frac{1}{2} \right\} > 0 \quad \text{due to (3.1).}$$

Then we obtain from (3.9) that

$$\begin{aligned} y_n |B_{r_n}(x_0) \cap \{w > k_{n+1}\}| &= y_n |B_{r_n}(x_0) \cap \{w > k_{n+1}\}|^{1-\sigma+\sigma} \\ &\leq cr^{2(1-\sigma)} \left(\frac{4^{n+2}y_n}{k^2} \right)^\sigma y_n \leq \frac{cA^{\sigma n}r^2}{r^{2\sigma}k^{2\sigma}} y_n^{1+\sigma}. \end{aligned}$$

Similarly,

$$\begin{aligned} &\int_{B_{r_n}(x_0)} |\mathbf{m}|^2 [(w - k_{n+1})^+]^2 dx |B_{r_n}(x_0) \cap \{w > k_{n+1}\}| \\ &\leq \|\mathbf{m}\|_{4, B_r(x_0)}^2 \left(\int_{B_{r_n}(x_0)} [(w - k_{n+1})^+]^4 dx \right)^{\frac{1}{2}} |B_{r_n}(x_0) \cap \{w > k_{n+1}\}|^{\frac{1}{2}+\sigma+\frac{1}{2}-\sigma} \\ &\leq \operatorname{ess\,sup}_{B_r(y)} w^+ \|\mathbf{m}\|_{4, B_r(x_0)}^2 y_n^{\frac{1}{2}} r^{1-2\sigma} \left(\frac{4^{n+2}y_n}{k^2} \right)^{\frac{1}{2}+\sigma} \\ &\leq \frac{cA^{(\frac{1}{2}+\sigma)n}r^2}{r^{2\sigma}k^{2\sigma}} y_n^{1+\sigma}. \end{aligned}$$

Here we have assumed that

$$(3.11) \quad k \geq \frac{1}{r} \|\mathbf{m}\|_{4, B_r(x_0)}^2.$$

As for the last term in (3.10), we have

$$\begin{aligned} &|B_{r_n}(x_0) \cap \{w \geq k_{n+1}\}|^{1+\sigma+2-\delta-\frac{2}{q}-\sigma} \\ &\leq cr^{4-2\delta-\frac{4}{q}-2\sigma} \left(\frac{4^{n+2}y_n}{k^2} \right)^{1+\sigma} \leq \frac{cA^{(1+\sigma)n}}{r^{2\sigma}k^{2\sigma}} y_n^{1+\sigma}. \end{aligned}$$

In this case, we have taken

$$(3.12) \quad k \geq r^{2-\delta-\frac{2}{q}}.$$

Collecting these estimates in (3.10) yields

$$y_{n+1} \leq \frac{cA^{(\frac{3}{2}+\sigma)n}}{r^{2\sigma}k^{2\sigma}} y_n^{1+\sigma}.$$

According to Lemma 4.1 in ([4], p.12), if we pick k so large that

$$y_0 \leq \frac{r^2k^2}{c^{\frac{1}{\sigma}}4^{\frac{3}{2}+\sigma}},$$

then

$$\sup_{B_{\frac{r}{2}}(x_0)} w \leq k.$$

Recall that

$$y_0 = \int_{B_r(x_0)} \left[\left(w - \frac{k}{2} \right)^+ \right]^2 dx \leq \int_{B_r(x_0)} (w^+)^2 dx.$$

This together with (3.11) and (3.12) implies that it is enough for us to take

$$k = c \left(\int_{B_r(x_0)} (w^+)^2 dx \right)^{\frac{1}{2}} + \frac{1}{r} \|\mathbf{m}\|_{4, B_r(x_0)}^2 + r^{2-\delta-\frac{2}{q}}.$$

This gives the desired result. \square

We are ready to complete the proof of Theorem 1.5. We estimate from (2.21) that

$$\begin{aligned} \frac{1}{r} \|\mathbf{m}\|_{4, B_r(x_0)}^2 &= c \left(\int_{B_r(x_0)} |\mathbf{m}|^4 dx \right)^{\frac{1}{2}} \\ &\leq c \left(\int_{B_r(x_0)} |\mathbf{m} - \mathbf{m}_{x_0, r}|^4 dx \right)^{\frac{1}{2}} + c |\mathbf{m}_{x_0, r}|^2 \\ &\leq c |\mathbf{m}_{x_0, r}|^2 + c. \end{aligned}$$

Similarly,

$$\int_{B_r(x_0)} |\mathbf{m}|^2 dx \leq c |\mathbf{m}_{x_0, r}|^2 + c.$$

We conclude from Claims 3.1 and 3.2 that

$$\begin{aligned} \sup_{B_{\frac{(1-\delta_0)r}{2}}(x_0)} w &\leq c \left(\int_{B_{(1-\delta_0)r}(x_0)} w^+ dx \right)^{\frac{1}{2}} + \frac{1}{r} \|\mathbf{m}\|_{4, B_r(x_0)}^2 + cr^{2-\delta-\frac{2}{q}} \\ &\leq c \left(\int_{B_r(x_0)} |\mathbf{m}|^2 dx \right)^{\frac{1}{2}} + cr^{1-\frac{1}{q}-\frac{\delta}{2}} + c + \frac{1}{r} \|\mathbf{m}\|_{4, B_r(x_0)}^2 + cr^{2-\delta-\frac{2}{q}} \\ &\leq c |\mathbf{m}_{x_0, r}|^2 + c \leq c. \end{aligned}$$

The last step is due to (1.9). By the definition of w ,

$$\omega_{x_0, r} + \bar{k}(r) \leq e^c [2(M_{x_0, r} - p) + \bar{k}(r)] \quad \text{on } B_{\frac{(1-\delta_0)r}{2}}(x_0),$$

from whence follows

$$\begin{aligned} 2e^c M_{x_0, \frac{(1-\delta_0)r}{2}} &\leq (2e^c - 1)\omega_{x_0, r} + 2e^c m_{x_0, r} \\ &\quad + (e^c - 1)\bar{k}(r). \end{aligned}$$

Subtract $2e^c m_{x_0, \frac{(1-\delta_0)r}{2}}$ from both sides of the above inequality and make use of the fact that $m_{x_0, r}$ is a decreasing function of r to deduce.

$$\omega_{x_0, \frac{(1-\delta_0)r}{2}} \leq \frac{2e^c - 1}{2e^c} \omega_{x_0, r} + \frac{e^c - 1}{2e^c} \bar{k}(r).$$

Obviously, recall the definition of $\bar{k}(r)$ to get

$$\omega_{x_0, \frac{(1-\delta_0)r}{2}} \leq \frac{2e^c - 1}{2e^c} \omega_{x_0, r} + r^\delta \|S\|_{q, B_r(x_0)} \leq \frac{2e^c - 1}{2e^c} \omega_{x_0, r} + cr^\delta.$$

It is very important to note that the two coefficients on the right-hand side of the above inequality can be made independent of r for r small. This enables us to invoke Lemma 8.23 in ([8], p.201) to get

$$\omega_{x_0, \rho} \leq c\rho^\alpha \quad \text{for some } c > 0, \alpha > 0 \text{ and } \rho \text{ sufficiently small.}$$

The proof is complete. \square

Finally, we give the proof of Lemma 1.6.

Proof of Lemma 1.6. Take

$$0 < r \leq \frac{1}{2}$$

so that

$$\ln|x-y| \leq 0 \quad \text{whenever } x, y \in B_r(x_0).$$

Consider the problem

$$(3.13) \quad -\Delta\psi = F \quad \text{in } B_r(x_0),$$

$$(3.14) \quad \psi = 0 \quad \text{on } \partial B_r(x_0).$$

Claim 3.3. *There exists a constant c such that*

$$(3.15) \quad \sup_{B_r(x_0)} |\psi| \leq c\eta(F; B_r(x_0); r),$$

where $\eta(F; B_r(x_0); r)$ is given as in (1.4).

Proof. We can write

$$F = F^+ - F^-.$$

Then solve the following two problems

$$(3.16) \quad \begin{aligned} -\Delta\psi_1 &= F^+ \quad \text{in } B_r(x_0), \\ \psi_1 &= 0 \quad \text{on } \partial B_r(x_0) \end{aligned}$$

and

$$\begin{aligned} -\Delta\psi_2 &= F^- \quad \text{in } B_r(x_0), \\ \psi_2 &= 0 \quad \text{on } \partial B_r(x_0). \end{aligned}$$

Clearly,

$$\psi = \psi_1 - \psi_2.$$

The non-negative function

$$-\frac{1}{2\pi} \int_{B_r(x_0)} F^+ \ln|x-y| dy$$

satisfies the equation (3.16). The comparison principle asserts

$$0 \leq \psi_1 \leq -\frac{1}{2\pi} \int_{B_r(x_0)} F^+ \ln|x-y| dy \quad \text{on } B_r(x_0).$$

Similarly,

$$0 \leq \psi_2 \leq -\frac{1}{2\pi} \int_{B_r(x_0)} F^- \ln|x-y| dy \quad \text{on } B_r(x_0).$$

The preceding inequalities yield the claim. □

The rest of the proof here largely follows [6]. Without loss of generality, assume

$$F \geq 0.$$

Let ψ be given as in (3.13)-(3.14). For $u \in W_0^{1,2}(B_r(x_0)) \cap L^\infty(B_r(x_0))$, we calculate

$$(3.17) \quad \begin{aligned} \int_{B_r(x_0)} Fu^2 dx &= - \int_{B_r(x_0)} \Delta\psi u^2 dx \\ &= 2 \int_{B_r(x_0)} u \nabla\psi \cdot \nabla u dx \\ &\leq 2 \left(\int_{B_r(x_0)} |\nabla u|^2 dx \right)^{\frac{1}{2}} \left(\int_{B_r(x_0)} u^2 |\nabla\psi|^2 dx \right)^{\frac{1}{2}}. \end{aligned}$$

It is easy to verify

$$|\nabla\psi|^2 = \frac{1}{2}\Delta\psi^2 - \psi\Delta\psi = \frac{1}{2}\Delta\psi^2 - \psi F.$$

This together with (3.15) implies

$$\begin{aligned} \int_{B_r(x_0)} u^2 |\nabla\psi|^2 dx &= \frac{1}{2} \int_{B_r(x_0)} u^2 \Delta\psi^2 dx - \int_{B_r(x_0)} \psi F u^2 dx \\ &\leq -\frac{1}{2} \int_{B_r(x_0)} \nabla u^2 \cdot \nabla\psi^2 dx + \|\psi\|_{\infty, B_r(x_0)} \int_{B_r(x_0)} F u^2 dx \\ &\leq \frac{1}{2} \int_{B_r(x_0)} u^2 |\nabla\psi|^2 dx + 2 \int_{B_r(x_0)} \psi^2 |\nabla u|^2 dx \\ &\quad + c\eta(F; B_r(x_0); r) \int_{B_r(x_0)} F u^2 dx, \end{aligned}$$

from whence follows

$$\int_{B_r(x_0)} u^2 |\nabla\psi|^2 dx \leq c\eta^2(r) \int_{B_r(x_0)} |\nabla u|^2 dx + c\eta(F; B_r(x_0); r) \int_{B_r(x_0)} F u^2 dx.$$

Plug this into (3.17) to derive

$$\begin{aligned} \int_{B_r(x_0)} |F|u^2 dx &\leq c\eta(F; B_r(x_0); r) \int_{B_r(x_0)} |\nabla u|^2 dx \\ &\quad + c \left(\eta(F; B_r(x_0); r) \int_{B_r(x_0)} F u^2 dx \right)^{\frac{1}{2}} \left(\int_{B_r(x_0)} |\nabla u|^2 dx \right)^{\frac{1}{2}} \\ &\leq \frac{1}{2} \int_{B_r(x_0)} |F|u^2 dx + c\eta(F; B_r(x_0); r) \int_{B_r(x_0)} |\nabla u|^2 dx. \end{aligned}$$

The lemma follows. □

REFERENCES

- [1] F. Chiarenza, E. Fabes, and N. Garofalo, Harnack's inequality for Schrödinger operators and the continuity of solutions, *Proc. Amer. Math. Soc.*, **98**(1986), 415-425.
- [2] S. Cho, A certain example for a De Giorgi conjecture, *J. Chungcheong Math. Soc.*, **27**(2014), 763-769.
- [3] E. De Giorgi, Congettura sulla continuità delle soluzioni di equazioni lineari ellittiche autoaggiunte a coefficienti illimitati, Unpublished (1995)
- [4] E. DiBenedetto, *Degenerate Parabolic Equations*, Springer-Verlag, New York, 1993.
- [5] G. Di Fazio and M.S. Fanciullo, BMO regularity for elliptic systems in Carnot-Carathéodory spaces, *Comm. Appl. Nonlinear Anal.*, **10**(2003), 81-95.
- [6] E.B. Fabes, N. Garofalo, and Fang-Hua Lin, A partial answer to a conjecture of B. Simon concerning unique continuation, *J. Func. Anal.*, **88** (1990), 194-210.
- [7] M. Gianquinta, *Multiple Integrals in the Calculus of Variations and Nonlinear Elliptic Systems*, Princeton University Press, Princeton, NJ, 1983.
- [8] D. Gilbarg and N. S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Springer-Verlag, Berlin, 1983.
- [9] J. Haskovec, P. Markowich, and B. Perthame, Mathematical analysis of a PDE system for biological network formulation, *Comm. Partial Differential Equations*, **40** (2015), 918-956.
- [10] J. Heinonen, T. Kilpeläinen, and O. Martio, *Nonlinear Potential Theory of Degenerate Equations*, Clarendon Press, Oxford, New York, Tokyo, 1993.
- [11] D. Hu, *Optimization, adaptation, and initialization of biological transport networks*, Workshop on multi scale problems from physics, biology, and material sciences, May 28-31, 2014, Shanghai.
- [12] D. Hu and D. Cai, Adaptation and optimization of biological transport networks, *Phys. Rev. Lett.*, **111**(2013), 138701.
- [13] K. Kurara, Continuity and Harnack's inequality for solutions of elliptic partial differential equations of second order, *Indiana Univ. Math. J.*, **43**(1994), 411-440.

- [14] J.-G. Liu and X. Xu, Partial regularity of weak solutions to a PDE system with cubic nonlinearity, *J. Differential Equations*, **264**(2018), 5489-5526.
- [15] J. Onninen and X. Zhong, Continuity of solutions of linear, degenerate elliptic equations. *Ann. Scuola Norm. Sup. Pisa Cl. Sci.*, **6**(2007), 103–116.
- [16] L.C. Piccinini and S. Spagnolo, On the Hölder continuity of solutions of second order elliptic equations in two variables, *Ann. Scuola Norm. Sup. Pisa Cl. Sci.*, **26** (1972), 391–402.
- [17] S. Semmes, A primer on Hardy spaces, and some remarks on a theorem of Evans and Müller, *Comm. Partial Differential Equations*, **19**(1994), 277-319.
- [18] X. Xu, Existence theorems for the quantum drift-diffusion system with mixed boundary conditions, *Commun. Contemp. Math.*, **18**(2016), no. 4, 1550048, 21 pp. arXiv:1911.01970v2 [math.AP], 2019.
- [19] X. Xu, Regularity theorems for a biological network formulation model in two space dimensions, *Kinetic and Related Models*, **11**(2018), 397-408.
- [20] X. Xu, Partial regularity of weak solutions and life-span of smooth solutions to a biological network formulation model, *SN Partial Differ. Equ. Appl.*, **1**, 18 (2020). <https://doi.org/10.1007/s42985-020-00021-3>