

# Multi-indexed poly-Bernoulli numbers

Yuna Baba, Maki Nakasuji\* and Mika Sakata

## Abstract

As properties of poly-Bernoulli numbers, a number of formulas such as the duality formula, explicit formula using the Stirling numbers of the second kind and periodicity for negative upper-index have been established. For the multi-indexed poly-Bernoulli numbers generalized by Kaneko-Tsumura, among such properties only the duality formula was obtained. In this paper, we restrict the double-indexed poly-Bernoulli numbers and show the explicit formula using the Stirling numbers of the second kind and periodicity for negative upper-index for them. Further, we define the variant of multiple-indexed poly-Bernoulli numbers using the star-version of multiple-indexed logarithms and obtain the relation between this kind of double and triple-indexed poly-Bernoulli numbers with multi-indexed poly-Bernoulli numbers ahead.

Keywords: Bernoulli numbers

AMS classification: 11B68.

## 1 Introduction

The Bernoulli numbers  $B_k$  which was defined by

$$\sum_{i=0}^n \binom{n+1}{i} B_i = n+1 \quad (n = 0, 1, 2, \dots),$$

were introduced by Jakob Bernoulli in his book. One of the most famous properties of them which was conceived by Euler is the following power series expansions:

$$\frac{t}{1 - e^{-t}} = \sum_{n=0}^{\infty} B_n \frac{t^n}{n!}. \quad (1.1)$$

---

\*The second author is supported by Grants-in-Aid for Scientific Research (C) 22K03274.

The parameter  $t$  can be real or complex. Since their introduction, Bernoulli numbers have been extended in a various way. poly-Bernoulli numbers by using polylogarithm series  $\text{Li}_k(z) = \sum_{m=1}^{\infty} z^m/m^k$  by Kaneko [K] are the first extension:

$$\frac{\text{Li}_k(1 - e^{-t})}{1 - e^{-t}} = \sum_{n=0}^{\infty} B_n^{(k)} \frac{t^n}{n!}$$

for  $k \in \mathbb{Z}$  and  $n \in \mathbb{Z}_{\geq 0}$  (see also Arakawa-Kaneko [AK1] and Arakawa-Ibukiyama-Kaneko [AIK]). When  $k = 1$ ,  $B_n^{(1)}$  is the usual Bernoulli number  $B_n$ . He found the explicit formula for  $B_n^{(k)}$  in terms of Stirling numbers of the second kind as a generalization of that for classical Bernoulli numbers.

**Theorem 1.1** ([K, Theorem 1]) *For a non negative integer  $n$  and an integer  $k$ , we have*

$$B_n^{(k)} = (-1)^n \sum_{m=0}^n \frac{(-1)^m m! \left\{ \begin{matrix} n \\ m \end{matrix} \right\}}{(m+1)^k}.$$

Here,  $\left\{ \begin{matrix} n \\ m \end{matrix} \right\}$  is a Stirling number of the second kind, which is defined by the following recurrence formula:

$$\left\{ \begin{matrix} 0 \\ 0 \end{matrix} \right\} = 1, \left\{ \begin{matrix} n \\ 0 \end{matrix} \right\} = \left\{ \begin{matrix} 0 \\ m \end{matrix} \right\} = 0 \ (n, m \neq 0), \left\{ \begin{matrix} n+1 \\ m \end{matrix} \right\} = \left\{ \begin{matrix} n \\ m-1 \end{matrix} \right\} + m \left\{ \begin{matrix} n \\ m \end{matrix} \right\}.$$

Ohno-Sakata considered the poly-Bernoulli numbers with negative upper-index and found the following properties for their periodicity ([OS], [S]).

**Theorem 1.2** ([OS, Theorem 4.2], [S, Theorem 6.1]) *Let  $k, N$  be positive integers. For a prime number  $p$  and positive integers  $n, m \geq N$  with  $n \equiv m \pmod{p^{N-1}(p-1)}$ , we have*

$$B_n^{(-k)} \equiv B_m^{(-k)} \pmod{p^N}.$$

**Theorem 1.3** ([S, Theorem 6.10]) *Let  $k, N$  be positive integers. For a prime number  $p$  and any integer  $n \geq N$ , we have*

$$\sum_{i=0}^{\varphi(p^N)-1} B_{n+i}^{(-k)} \equiv 0 \pmod{p^N},$$

where  $\varphi(x)$  is a Euler's  $\varphi$ -function.

Arakawa-Kaneko [AK1] introduced the multi-logarithmic function defined by

$$\text{Li}_{k_1, k_2, \dots, k_r}(z) = \sum_{0 < m_1 < m_2 < \dots < m_r} \frac{z^{m_r}}{m_1^{k_1} m_2^{k_2} \dots m_r^{k_r}}$$

for  $k_i \geq 1$  and  $|z| < 1$ . And they defined a further generalization of  $B_n^{(k)}$ , which is known as the multiple poly-Bernoulli numbers  $\mathbb{B}_n^{(k_1, \dots, k_r)}$ , as

$$\frac{\text{Li}_{k_1, \dots, k_r}(1 - e^{-t})}{(1 - e^{-t})^r} = \sum_{n=0}^{\infty} \mathbb{B}_n^{(k_1, \dots, k_r)} \frac{t^n}{n!}$$

for  $k_1, \dots, k_r \in \mathbb{Z}$ . When  $r = 1$ ,  $\mathbb{B}_n^{(k_1)} = B_n^{(k_1)}$  being the poly-Bernoulli numbers mentioned as above. Hamahata-Masubuchi and Bayad-Hamataha investigated some properties of  $\mathbb{B}_n^{(k_1, \dots, k_r)}$  ([HM], [BH]). Kaneko-Tsumura ([KT]) introduced further variant using the multiple polylogarithms of  $*$ -type and of  $\text{III}$ -type defined by Goncharov ([G]):

$$\text{Li}_{s_1, \dots, s_r}^*(z_1, \dots, z_r) = \sum_{1 \leq m_1 < \dots < m_r} \frac{z_1^{m_1} z_2^{m_2} \dots z_r^{m_r}}{m_1^{s_1} m_2^{s_2} \dots m_r^{s_r}}, \quad (1.2)$$

$$\begin{aligned} \text{Li}_{s_1, \dots, s_r}^{\text{III}}(z_1, \dots, z_r) &= \sum_{1 \leq m_1 < \dots < m_r} \frac{z_1^{m_1} z_2^{m_2 - m_1} \dots z_r^{m_r - m_{r-1}}}{m_1^{s_1} m_2^{s_2} \dots m_r^{s_r}} \\ &= \sum_{\ell_1, \dots, \ell_r=1}^{\infty} \frac{z_1^{\ell_1} z_2^{\ell_2} \dots z_r^{\ell_r}}{\ell_1^{s_1} (\ell_1 + \ell_2)^{s_2} \dots (\ell_1 + \dots + \ell_r)^{s_r}} \end{aligned} \quad (1.3)$$

for  $s_1, \dots, s_r \in \mathbb{C}$  and  $z_1, \dots, z_r \in \mathbb{C}$ . They defined the multi-indexed poly-Bernoulli numbers  $B_{m_1, \dots, m_r}^{(s_1, \dots, s_r), (d)}$  in Theorem 5.1 in [KT] by using multiple polylogarithms of  $\text{III}$ -type as

$$\begin{aligned} F(x_1, \dots, x_r; s_1, \dots, s_r; d) &= \frac{\text{Li}_{s_1, \dots, s_r}^{\text{III}}(1 - e^{-\sum_{\nu=1}^r x_\nu}, \dots, 1 - e^{-x_{r-1} - x_r}, 1 - e^{-x_r})}{\prod_{j=1}^d (1 - e^{-\sum_{\nu=j}^r x_\nu})} \\ &= \left( \sum_{\ell_1, \dots, \ell_r=1}^{\infty} \frac{\prod_{j=1}^r (1 - e^{-\sum_{\nu=j}^r x_\nu})^{\ell_j - \delta_j(d)}}{\prod_{j=1}^r (\sum_{\nu=1}^j \ell_\nu)^{s_j}} \right) \\ &= \sum_{m_1, \dots, m_r=0}^{\infty} B_{m_1, \dots, m_r}^{(s_1, \dots, s_r), (d)} \frac{x_1^{m_1} \dots x_r^{m_r}}{m_1! \dots m_r!} \end{aligned} \quad (1.4)$$

for  $x_1, \dots, x_r \in \mathbb{C}$  and  $d \in \mathbb{Z}$  ( $1 \leq d \leq r$ ). When  $d = r$ , they denote  $B_{m_1, \dots, m_r}^{(s_1, \dots, s_r), (r)}$  by  $\mathbb{B}_{m_1, \dots, m_r}^{(s_1, \dots, s_r)}$  and proved the beautiful duality formula.

**Theorem 1.4** ([KT, Theorem 5.4]) For  $m_1, \dots, m_r, k_1, \dots, k_r \in \mathbb{Z}_{\geq 0}$ , we have

$$\mathbb{B}_{m_1, \dots, m_r}^{(-k_1, \dots, -k_r)} = \mathbb{B}_{k_1, \dots, k_r}^{(-m_1, \dots, -m_r)}.$$

Note that for  $r = 1$  and  $s_1 = k_1 \in \mathbb{Z}$ ,  $\mathbb{B}_{m_1}^{(s_1)} = \mathbb{B}_{m_1}^{(k_1)} = B_{m_1}^{(k_1)}$ . Theorem 1.4 is known as the generalization of the following duality theorem by Kaneko.

**Theorem 1.5** ([K, Theorem 2]) For any  $m, k \geq 0$ , we have

$$B_m^{(-k)} = B_k^{(-m)}.$$

Therefore, it is natural to ask other properties such as the generalization of Theorems 1.1, 1.2 and 1.3.

On the other hand, as we can see in (1.2) and (1.3),  $\text{Li}_{s_1, \dots, s_r}^*(z_1, \dots, z_r)$  and  $\text{Li}_{s_1, \dots, s_r}^{\text{III}}(z_1, \dots, z_r)$  are defined by the sum over  $m_i$  with the order without equal sign,  $1 \leq m_1 < \dots < m_r$ , so it is a natural question what happens in the case of equal ordering in the sum. Therefore, in this article we introduce the multiple-indexed polylogarithms.

**Definition 1.6** Let

$$\begin{aligned} \text{Li}_{s_1, \dots, s_r}^{*,*}(z_1, \dots, z_r) &= \sum_{1 \leq m_1 \leq \dots \leq m_r} \frac{z_1^{m_1} z_2^{m_2} \dots z_r^{m_r}}{m_1^{s_1} m_2^{s_2} \dots m_r^{s_r}}, \\ \text{Li}_{s_1, \dots, s_r}^{\text{III},*}(z_1, \dots, z_r) &= \sum_{1 \leq m_1 \leq \dots \leq m_r} \frac{z_1^{m_1} z_2^{m_2 - m_1} \dots z_r^{m_r - m_{r-1}}}{m_1^{s_1} m_2^{s_2} \dots m_r^{s_r}} \\ &= \sum_{\ell_1=1, \ell_2, \dots, \ell_r=0}^{\infty} \frac{z_1^{\ell_1} z_2^{\ell_2} \dots z_r^{\ell_r}}{\ell_1^{s_1} (\ell_1 + \ell_2)^{s_2} \dots (\ell_1 + \dots + \ell_r)^{s_r}}, \end{aligned}$$

for  $s_1, \dots, s_r \in \mathbb{C}$  and  $z_1, \dots, z_r \in \mathbb{C}$  with  $|z_j| < 1$  ( $1 \leq j \leq r$ ).

Similarly with (1.4), we define multiple-indexed poly-Bernoulli numbers  $\mathbb{B}_{m_1, \dots, m_r}^{*,(s_1, s_2, \dots, s_r)}$  by using multiple polylogarithms of  $\text{III}$ -type as

$$\begin{aligned} F^*(x_1, \dots, x_r; s_1, \dots, s_r; r) &= \frac{\text{Li}_{s_1, \dots, s_r}^{\text{III},*}(1 - e^{-\sum_{\nu=1}^r x_\nu}, \dots, 1 - e^{-x_{r-1} - x_r}, 1 - e^{-x_r})}{\prod_{j=1}^r (1 - e^{-\sum_{\nu=j}^r x_\nu})} \\ &= \sum_{m_1, \dots, m_r=0}^{\infty} \mathbb{B}_{m_1, \dots, m_r}^{*,(s_1, \dots, s_r)} \frac{x_1^{m_1} \dots x_r^{m_r}}{m_1! \dots m_r!}. \end{aligned} \quad (1.5)$$

Hereafter, we may write  $F^*(x_1, \dots, x_r; s_1, \dots, s_r)$  for  $F^*(x_1, \dots, x_r; s_1, \dots, s_r; r)$ . Then for

the double-indexed and triple-indexed poly-Bernoulli numbers, we obtain the following

**Theorem 1.7** *For  $m_1, m_2 \in \mathbb{Z}_{\geq 0}$ ,  $s_1, s_2 \in \mathbb{C}$ , the double-indexed poly-Bernoulli numbers, we have*

$$m_2 \mathbb{B}_{m_1, m_2-1}^{*,(s_1, s_2)} = m_2 \mathbb{B}_{m_1, m_2-1}^{(s_1, s_2)} + B_{m_2} \mathbb{B}_{m_1}^{(s_1+s_2)}.$$

**Theorem 1.8** *For  $m_1, m_2, m_3 \in \mathbb{Z}_{\geq 0}$ ,  $s_1, s_2, s_3 \in \mathbb{C}$ , the triple-indexed poly-Bernoulli numbers, we have*

$$\begin{aligned} & m_2 m_3 \mathbb{B}_{m_1, m_2-1, m_3-1}^{*,(s_1, s_2, s_3)} + m_3(m_3 - 1) \mathbb{B}_{m_1, m_2, m_3-2}^{*,(s_1, s_2, s_3)} \\ &= m_2 m_3 \mathbb{B}_{m_1, m_2-1, m_3-1}^{(s_1, s_2, s_3)} + m_3(m_3 - 1) \mathbb{B}_{m_1, m_2, m_3-2}^{(s_1, s_2, s_3)} + m_3 B_{m_2} \mathbb{B}_{m_1, m_3-1}^{(s_1+s_2, s_3)} \\ & \quad + m_2 B_{m_3} \mathbb{B}_{m_1, m_2-1}^{(s_1, s_2+s_3)} + m_3 B_{m_3-1} \mathbb{B}_{m_1, m_2}^{(s_1, s_2+s_3)} + B_{m_2} B_{m_3} \mathbb{B}_{m_1}^{(s_1+s_2+s_3)}. \end{aligned}$$

In this article, in Section 2, we discuss the generalization of Theorems 1.1, 1.2 and 1.3 to the double-indexed poly-Bernoulli numbers  $\mathbb{B}_{m_1, m_2}^{(s_1, s_2)}$ . And in Sections 3 and 4, we give the proof of Theorem 1.7 and 1.8, respectively.

## 2 Double-indexed poly-Bernoulli numbers

In this section, we restrict double-indexed poly-Bernoulli numbers. We first extend Theorem 1.1 to double-indexed poly-Bernoulli numbers.

**Theorem 2.1** *For non negative integers  $\ell_1, \ell_2$  and some integers  $k_1, k_2$ , we have*

$$\mathbb{B}_{\ell_1, \ell_2}^{(k_1, k_2)} = \sum_{0 \leq m_1, m_2}^{\ell_1 + \ell_2} \frac{(-1)^{m_1 + m_2 + \ell_1 + \ell_2} m_1! m_2!}{(m_1 + 1)^{k_1} (m_1 + m_2 + 2)^{k_2}} \sum_{n=0}^{\ell_1 + \ell_2} \begin{Bmatrix} n \\ m_1 \end{Bmatrix} \begin{Bmatrix} \ell_1 + \ell_2 - n \\ m_2 \end{Bmatrix} \binom{\ell_2}{n - \ell_1},$$

where  $\begin{Bmatrix} n \\ m \end{Bmatrix}$  is a Stirling number and  $\binom{\ell}{n}$  is a binomial coefficient.

**Proof** This can be proved by faithful calculation. First, by the definition of  $\text{Li}_{k_1, k_2}^{\text{III}}$ , we have

$$\frac{\text{Li}_{k_1, k_2}^{\text{III}}(1 - e^{-t_1 - t_2}, 1 - e^{-t_2})}{(1 - e^{-t_1 - t_2})(1 - e^{-t_2})} = \sum_{1 \leq m_1 < m_2} \frac{(1 - e^{-t_1 - t_2})^{m_1 - 1} (1 - e^{-t_2})^{m_2 - m_1 - 1}}{m_1^{k_1} m_2^{k_2}}.$$

Putting  $m_3 = m_2 - m_1$ , then we obtain  $1 \leq m_3$  and

$$\begin{aligned}
(RHS) &= \sum_{1 \leq m_1, m_3} \frac{(1 - e^{-t_1 - t_2})^{m_1 - 1} (1 - e^{-t_2})^{m_3 - 1}}{m_1^{k_1} (m_1 + m_3)^{k_2}} \\
&= \sum_{0 \leq m_1, m_3} \frac{(1 - e^{-t_1 - t_2})^{m_1} (1 - e^{-t_2})^{m_3}}{(m_1 + 1)^{k_1} (m_1 + m_3 + 2)^{k_2}} \\
&= \sum_{0 \leq m_1, m_3} \frac{(-1)^{m_1 + m_3} (e^{-t_1 - t_2} - 1)^{m_1} (e^{-t_2} - 1)^{m_3} m_1! m_3!}{(m_1 + 1)^{k_1} (m_1 + m_3 + 2)^{k_2} m_1! m_3!}.
\end{aligned}$$

Using the properties of the Stirling numbers, we can proceed the calculation:

$$\begin{aligned}
&= \sum_{0 \leq m_1, m_3} \frac{(-1)^{m_1 + m_3} m_1! m_3!}{(m_1 + 1)^{k_1} (m_1 + m_3 + 2)^{k_2}} \sum_{n_1=0}^{\infty} \binom{n_1}{m_1} \frac{(-t_1 - t_2)^{n_1}}{n_1!} \sum_{n_2=0}^{\infty} \binom{n_2}{m_3} \frac{(-t_2)^{n_2}}{n_2!} \\
&= \sum_{0 \leq m_1, m_3} \frac{(-1)^{m_1 + m_3} m_1! m_3!}{(m_1 + 1)^{k_1} (m_1 + m_3 + 2)^{k_2}} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{n_1} \binom{n_2}{m_1} \binom{n_1 - n_2}{m_3} \frac{(-t_1 - t_2)^{n_2}}{n_2!} \frac{(-t_2)^{n_1 - n_2}}{(n_1 - n_2)!} \\
&= \sum_{0 \leq m_1, m_3} \frac{(-1)^{m_1 + m_3} m_1! m_3!}{(m_1 + 1)^{k_1} (m_1 + m_3 + 2)^{k_2}} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{n_1} \binom{n_2}{m_1} \binom{n_1 - n_2}{m_3} \sum_{\ell_1=0}^{n_2} \binom{n_2}{\ell_1} \frac{(-t_1)^{\ell_1} (-t_2)^{n_1 - \ell_1}}{n_2! (n_1 - n_2)!} \\
&= \sum_{\ell_1=0}^{\infty} \sum_{0 \leq m_1, m_3} \frac{(-1)^{m_1 + m_3} m_1! m_3!}{(m_1 + 1)^{k_1} (m_1 + m_3 + 2)^{k_2}} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{n_1} \binom{n_2}{m_1} \binom{n_1 - n_2}{m_3} \frac{(-t_1)^{\ell_1} (-t_2)^{n_1 - \ell_1}}{\ell_1! (n_1 - n_2)! (n_2 - \ell_1)!} \\
&= \sum_{\ell_1=0}^{\infty} \sum_{0 \leq m_1, m_3} \frac{(-1)^{m_1 + m_3} m_1! m_3!}{(m_1 + 1)^{k_1} (m_1 + m_3 + 2)^{k_2}} \\
&\quad \times \sum_{\ell_2=0}^{\infty} \sum_{n_2=0}^{\ell_1 + \ell_2} \binom{n_2}{m_1} \binom{\ell_1 + \ell_2 - n_2}{m_3} \frac{(-t_1)^{\ell_1} (-t_2)^{\ell_2}}{\ell_1! (\ell_1 + \ell_2 - n_2)! (n_2 - \ell_1)!} \\
&= \sum_{\ell_1=0}^{\infty} \sum_{0 \leq m_1, m_3} \frac{(-1)^{m_1 + m_3} m_1! m_3!}{(m_1 + 1)^{k_1} (m_1 + m_3 + 2)^{k_2}} \\
&\quad \times \sum_{\ell_2=0}^{\infty} \sum_{n_2=0}^{\ell_1 + \ell_2} \binom{n_2}{m_1} \binom{\ell_1 + \ell_2 - n_2}{m_3} \binom{\ell_2}{n_2 - \ell_1} \frac{(-t_1)^{\ell_1} (-t_2)^{\ell_2}}{\ell_1! \ell_2!} \\
&= \sum_{\ell_1=0}^{\infty} \sum_{\ell_2=0}^{\infty} \sum_{0 \leq m_1, m_3}^{\ell_1 + \ell_2} \frac{(-1)^{m_1 + m_3 + \ell_1 + \ell_2} m_1! m_3!}{(m_1 + 1)^{k_1} (m_1 + m_3 + 2)^{k_2}} \sum_{n_2=0}^{\ell_1 + \ell_2} \binom{n_2}{m_1} \binom{\ell_1 + \ell_2 - n_2}{m_3} \binom{\ell_2}{n_2 - \ell_1} \frac{t_1^{\ell_1} t_2^{\ell_2}}{\ell_1! \ell_2!}.
\end{aligned}$$

Comparing the coefficients of both sides, we obtain

$$\mathbb{B}_{\ell_1, \ell_2}^{(k_1, k_2)} = \sum_{0 \leq m_1, m_3}^{\ell_1 + \ell_2} \frac{(-1)^{m_1 + m_3 + \ell_1 + \ell_2} m_1! m_3!}{(m_1 + 1)^{k_1} (m_1 + m_3 + 2)^{k_2}} \sum_{n_2=0}^{\ell_1 + \ell_2} \begin{Bmatrix} n_2 \\ m_1 \end{Bmatrix} \begin{Bmatrix} \ell_1 + \ell_2 - n_2 \\ m_3 \end{Bmatrix} \binom{\ell_2}{n_2 - \ell_1}.$$

□

Second, we consider the periodicity to extend Theorem 1.2.

**Theorem 2.2** *Let  $k_1, k_2, N$  be positive integers. Then for a prime number  $p$ , and positive integers  $n_1, n_2, m_1, m_2 \geq N$  with  $n_1 \equiv m_1 \pmod{p^{N-1}(p-1)}$  and  $n_2 \equiv m_2 \pmod{p^{N-1}(p-1)}$ , we have*

$$\mathbb{B}_{n_1, n_2}^{(-k_1, -k_2)} \equiv \mathbb{B}_{n_1, m_2}^{(-k_1, -k_2)} \pmod{p^N}, \quad (2.1)$$

$$\mathbb{B}_{n_1, n_2}^{(-k_1, -k_2)} \equiv \mathbb{B}_{m_1, n_2}^{(-k_1, -k_2)} \pmod{p^N}. \quad (2.2)$$

**Proof** Let us begin with the first assertion. Assume  $n_2 \geq m_2$ . From Theorem 2.1,

$$\begin{aligned} \mathbb{B}_{k_1, k_2}^{(-n_1, -n_2)} &= \sum_{0 \leq \ell_1}^{k_1 + k_2} \sum_{0 \leq \ell_2}^{k_1 + k_2} (-1)^{\ell_1 + \ell_2 + k_1 + k_2} \ell_1! \ell_2! (\ell_1 + 1)^{n_1} (\ell_1 + \ell_2 + 2)^{n_2} \\ &\quad \times \sum_{h_2=0}^{k_1 + k_2} \begin{Bmatrix} h_2 \\ \ell_1 \end{Bmatrix} \begin{Bmatrix} k_1 + k_2 - h_2 \\ \ell_2 \end{Bmatrix} \binom{k_2}{h_2 - k_1}. \end{aligned} \quad (2.3)$$

Dividing  $\sum_{0 \leq \ell_2}^{k_1 + k_2}$  into two parts  $\sum_{\substack{\ell_2=0 \\ p|\ell_1 + \ell_2 + 2}}^{k_1 + k_2} + \sum_{\substack{\ell_2=0 \\ p \nmid \ell_1 + \ell_2 + 2}}^{k_1 + k_2}$ , we obtain

$$\begin{aligned} \mathbb{B}_{k_1, k_2}^{(-n_1, -n_2)} &\equiv \sum_{0 \leq \ell_1}^{k_1 + k_2} \sum_{\substack{\ell_2=0 \\ p|\ell_1 + \ell_2 + 2}}^{k_1 + k_2} (-1)^{\ell_1 + \ell_2 + k_1 + k_2} \ell_1! \ell_2! (\ell_1 + 1)^{n_1} (\ell_1 + \ell_2 + 2)^{n_2} \\ &\quad \times \sum_{h_2=0}^{k_1 + k_2} \begin{Bmatrix} h_2 \\ \ell_1 \end{Bmatrix} \begin{Bmatrix} k_1 + k_2 - h_2 \\ \ell_2 \end{Bmatrix} \binom{k_2}{h_2 - k_1} \pmod{p^N}. \end{aligned}$$

The Euler theorem leads to

$$\begin{aligned} \mathbb{B}_{k_1, k_2}^{(-n_1, -n_2)} &\equiv \sum_{0 \leq \ell_1}^{k_1+k_2} \sum_{\substack{\ell_2=0 \\ p \nmid \ell_1 + \ell_2 + 2}}^{k_1+k_2} (-1)^{\ell_1 + \ell_2 + k_1 + k_2} \ell_1! \ell_2! (\ell_1 + 1)^{n_1} (\ell_1 + \ell_2 + 2)^{m_2} \\ &\times \sum_{h_2=0}^{k_1+k_2} \begin{Bmatrix} h_2 \\ \ell_1 \end{Bmatrix} \begin{Bmatrix} k_1 + k_2 - h_2 \\ \ell_2 \end{Bmatrix} \binom{k_2}{h_2 - k_1} = \mathbb{B}_{k_1, k_2}^{(-n_1, -m_2)} \pmod{p^N}. \end{aligned}$$

Applying Theorem 1.4, we obtain

$$\mathbb{B}_{n_1, n_2}^{(-k_1, -k_2)} \equiv \mathbb{B}_{n_1, m_2}^{(-k_1, -k_2)} \pmod{p^N}.$$

The second assertion is obtained in a similar way. Assume  $n_1 \geq m_1$ . We consider

$$\begin{aligned} \mathbb{B}_{k_1, k_2}^{(-n_1, -n_2)} &= \sum_{0 \leq \ell_2}^{k_1+k_2} \sum_{0 \leq \ell_1}^{k_1+k_2} (-1)^{\ell_1 + \ell_2 + k_1 + k_2} \ell_1! \ell_2! (\ell_1 + 1)^{n_1} (\ell_1 + \ell_2 + 2)^{n_2} \\ &\times \sum_{h_2=0}^{k_1+k_2} \begin{Bmatrix} h_2 \\ \ell_1 \end{Bmatrix} \begin{Bmatrix} k_1 + k_2 - h_2 \\ \ell_2 \end{Bmatrix} \binom{k_2}{h_2 - k_1}. \end{aligned}$$

Dividing  $\sum_{0 \leq \ell_1}^{k_1+k_2}$  into two parts  $\sum_{\substack{\ell_1=0 \\ p \nmid \ell_1 + 1}}^{k_1+k_2} + \sum_{\substack{\ell_1=0 \\ p \mid \ell_1 + 1}}^{k_1+k_2}$ , we obtain

$$\begin{aligned} \mathbb{B}_{k_1, k_2}^{(-n_1, -n_2)} &\equiv \sum_{0 \leq \ell_2}^{k_1+k_2} \sum_{\substack{\ell_1=0 \\ p \nmid \ell_1 + 1}}^{k_1+k_2} (-1)^{\ell_1 + \ell_2 + k_1 + k_2} \ell_1! \ell_2! (\ell_1 + 1)^{n_1} (\ell_1 + \ell_2 + 2)^{n_2} \\ &\times \sum_{h_2=0}^{k_1+k_2} \begin{Bmatrix} h_2 \\ \ell_1 \end{Bmatrix} \begin{Bmatrix} k_1 + k_2 - h_2 \\ \ell_2 \end{Bmatrix} \binom{k_2}{h_2 - k_1} \pmod{p^N}. \end{aligned}$$

The Euler theorem leads to

$$\begin{aligned} \mathbb{B}_{k_1, k_2}^{(-n_1, -n_2)} &\equiv \sum_{0 \leq \ell_2}^{k_1+k_2} \sum_{\substack{\ell_1=0 \\ p \nmid \ell_1 + 1}}^{k_1+k_2} (-1)^{\ell_1 + \ell_2 + k_1 + k_2} \ell_1! \ell_2! (\ell_1 + 1)^{m_1} (\ell_1 + \ell_2 + 2)^{n_2} \\ &\times \sum_{h_2=0}^{k_1+k_2} \begin{Bmatrix} h_2 \\ \ell_1 \end{Bmatrix} \begin{Bmatrix} k_1 + k_2 - h_2 \\ \ell_2 \end{Bmatrix} \binom{k_2}{h_2 - k_1} = \mathbb{B}_{k_1, k_2}^{(-m_1, -n_2)} \pmod{p^N}. \end{aligned}$$

Applying Theorem 1.4, we obtain the second assertion:

$$\mathbb{B}_{n_1, n_2}^{(-k_1, -k_2)} \equiv \mathbb{B}_{m_1, n_2}^{(-k_1, -k_2)} \pmod{p^N}.$$

□

Next, we consider the periodicity to extend Theorem 1.3.

**Theorem 2.3** *Let  $k_1, k_2, N$  be positive integers. For a prime number  $p$ , and integers  $n_1, n_2 \geq N$ , we have*

$$\sum_{i=0}^{\varphi(p^N)-1} \mathbb{B}_{n_1+i, n_2}^{(-k_1, -k_2)} \equiv 0 \pmod{p^N}, \quad (2.4)$$

$$\sum_{i=0}^{\varphi(p^N)-1} \mathbb{B}_{n_1, n_2+i}^{(-k_1, -k_2)} \equiv 0 \pmod{p^N}. \quad (2.5)$$

**Proof** For (2.4), we first consider the case of  $n_1 = N$ . From Theorems 1.4 and 2.1,

$$\begin{aligned} \sum_{i=0}^{\varphi(p^N)-1} \mathbb{B}_{N+i, n_2}^{(-k_1, -k_2)} &= \sum_{i=0}^{\varphi(p^N)-1} \mathbb{B}_{k_1, k_2}^{(-N-i, -n_2)} \\ &= \sum_{i=0}^{\varphi(p^N)-1} \sum_{\ell_1, \ell_2=0}^{k_1+k_2} (-1)^{\ell_1+\ell_2+k_1+k_2} \ell_1! \ell_2! (\ell_1+1)^{N+i} (\ell_1+\ell_2+2)^{n_2} \\ &\quad \times \sum_{h_2=0}^{k_1+k_2} \begin{Bmatrix} h_2 \\ \ell_1 \end{Bmatrix} \begin{Bmatrix} k_1+k_2-h_2 \\ \ell_2 \end{Bmatrix} \binom{k_2}{h_2-k_1} \\ &= \sum_{\ell_1, \ell_2=0}^{k_1+k_2} (-1)^{\ell_1+\ell_2+k_1+k_2} \ell_1! \ell_2! (\ell_1+\ell_2+2)^{n_2} \\ &\quad \times \sum_{h_2=0}^{k_1+k_2} \begin{Bmatrix} h_2 \\ \ell_1 \end{Bmatrix} \begin{Bmatrix} k_1+k_2-h_2 \\ \ell_2 \end{Bmatrix} \binom{k_2}{h_2-k_1} \frac{(\ell_1+1)^N \{(\ell_1+1)^{\varphi(p^N)} - 1\}}{\ell_1}. \end{aligned}$$

The right-hand side above identically 0 modulo  $p^N$  from the Euler theorem and we get the assertion. In case of  $n_1 > N$ , applying the case of  $n_1 = N$  and Theorem 2.2 leads to the desired result. For (2.5), we similarly consider the case of  $n_2 = N$  first. From Theorems 1.4

and 2.1,

$$\begin{aligned}
\sum_{i=0}^{\varphi(p^N)-1} \mathbb{B}_{n_1 N+i}^{(-k_1, -k_2)} &= \sum_{i=0}^{\varphi(p^N)-1} \mathbb{B}_{k_1, k_2}^{(-n_1, -N-i)} \\
&= \sum_{i=0}^{\varphi(p^N)-1} \sum_{\ell_1, \ell_2=0}^{k_1+k_2} (-1)^{\ell_1+\ell_2+k_1+k_2} \ell_1! \ell_2! (\ell_1+1)^{n_1} (\ell_1+\ell_2+2)^{N+i} \\
&\quad \times \sum_{h_2=0}^{k_1+k_2} \begin{Bmatrix} h_2 \\ \ell_1 \end{Bmatrix} \begin{Bmatrix} k_1+k_2-h_2 \\ \ell_2 \end{Bmatrix} \binom{k_2}{h_2-k_1} \\
&= \sum_{\ell_1, \ell_2=0}^{k_1+k_2} (-1)^{\ell_1+\ell_2+k_1+k_2} \ell_1! \ell_2! (\ell_1+1)^{n_1} \\
&\quad \times \sum_{h_2=0}^{k_1+k_2} \begin{Bmatrix} h_2 \\ \ell_1 \end{Bmatrix} \begin{Bmatrix} k_1+k_2-h_2 \\ \ell_2 \end{Bmatrix} \binom{k_2}{h_2-k_1} \frac{(\ell_1+\ell_2+2)^N \{(\ell_1+\ell_2+2)^{\varphi(p^N)} - 1\}}{\ell_1+\ell_2+1}.
\end{aligned}$$

The right-hand side above also identically 0 modulo  $p^N$  from the Euler theorem and we get the assertion. The case of  $n_2 > N$  is similarly derived.  $\square$

**Corollary 2.4** *Let  $k_1, k_2, M$  be positive integers. For  $M = p_1^{e_1} p_2^{e_2} \cdots p_m^{e_m}$ , where  $p_i$ 's are relatively prime, and any integer  $n \geq e_1, e_2, \dots, e_m$ , we have*

$$\sum_{i=0}^{\varphi(M)-1} \mathbb{B}_{n_1+i, n_2}^{(-k_1, -k_2)} \equiv 0 \pmod{M}, \tag{2.6}$$

$$\sum_{i=0}^{\varphi(M)-1} \mathbb{B}_{n_1, n_2+i}^{(-k_1, -k_2)} \equiv 0 \pmod{M}. \tag{2.7}$$

**Proof** From Theorem 2.2, for  $1 \leq r \leq m$ , we have

$$\begin{aligned}
\sum_{i=0}^{\varphi(M)-1} \mathbb{B}_{n_1+i, n_2}^{(-k_1, -k_2)} &= \sum_{i=0}^{\varphi(p_1^{e_1}) \cdots \varphi(p_m^{e_m})-1} \mathbb{B}_{n_1+i, n_2}^{(-k_1, -k_2)} \\
&\equiv \prod_{\substack{j=1 \\ j \neq r}}^m \varphi(p_j^{e_j}) \sum_{i=0}^{\varphi(p_r^{e_r})-1} \mathbb{B}_{n_1+i, n_2}^{(-k_1, -k_2)} \pmod{p_r^{e_r}}.
\end{aligned}$$

The right-hand side above also identically 0 modulo  $p_r^{e_r}$  from Theorem 2.3, we can get (2.6). We similarly consider the case of  $n_2$  and obtain (2.7).  $\square$

**Remark 1** *As we can see, Theorem 2.1 is the key theorem for Theorems 2.2 and 2.3. Therefore, if Theorem 2.1 can be generalized for  $n$ -indexed poly-Bernoulli numbers with  $n \geq 3$ , the related periodicity theorems may also be derived in a similar way. However, we have not pursued here because the binomial coefficients in these cases are complicated.*

**Remark 2** *According to our calculations (by using Mathematica), there seems to have finer periodicities in (2.1) and (2.2) sometimes, but not always. This means that we may be able to obtain better results under suitable conditions.*

### 3 Double-indexed star-poly-Bernoulli numbers

This section is devoted to the proof of Theorem 1.7, where we restrict double-indexed polylogarithms. By the harmonic calculations, we have

$$\begin{aligned} \text{Li}_{k_1, k_2}^{\text{III}, \star}(z_1, z_2) &= \sum_{1 \leq m_1 \leq m_2} \frac{z_1^{m_1} z_2^{m_2 - m_1}}{m_1^{k_1} m_2^{k_2}} \\ &= \sum_{1 \leq m_1 < m_2} \frac{z_1^{m_1} z_2^{m_2 - m_1}}{m_1^{k_1} m_2^{k_2}} + \sum_{1 \leq m_1} \frac{z_1^{m_1}}{m_1^{k_1 + k_2}} \\ &= \text{Li}_{k_1, k_2}^{\text{III}}(z_1, z_2) + \text{Li}_{k_1 + k_2}^{\text{III}}(z_1). \end{aligned}$$

So,  $F^*(x_1, x_2; s_1, s_2)$  defined in (1.5) is written as

$$\begin{aligned} F^*(x_1, x_2; s_1, s_2) &= \frac{\text{Li}_{s_1, s_2}^{\text{III}, \star}(1 - e^{-x_1 - x_2}, 1 - e^{-x_2})}{(1 - e^{-x_1 - x_2})(1 - e^{-x_2})} \\ &= \frac{\text{Li}_{s_1, s_2}^{\text{III}}(1 - e^{-x_1 - x_2}, 1 - e^{-x_2})}{(1 - e^{-x_1 - x_2})(1 - e^{-x_2})} + \frac{1}{(1 - e^{-x_2})} \frac{\text{Li}_{s_1 + s_2}^{\text{III}}(1 - e^{-x_1 - x_2})}{(1 - e^{-x_1 - x_2})}. \end{aligned} \tag{3.1}$$

Using (1.4), this can be expressed in terms of double-indexed poly-Bernoulli numbers,

$$F^*(x_1, x_2; s_1, s_2) = \sum_{m_{11}, m_{12}=0}^{\infty} \mathbb{B}_{m_{11}, m_{12}}^{(s_1, s_2)} \frac{x_1^{m_{11}} x_2^{m_{12}}}{m_{11}! m_{12}!} + \frac{1}{(1 - e^{-x_2})} \sum_{m_{21}=0}^{\infty} \mathbb{B}_{m_{21}}^{(s_1 + s_2)} \frac{(x_1 + x_2)^{m_{21}}}{m_{21}!}. \tag{3.2}$$

Further, by (1.1), we have

$$\begin{aligned}
& F^*(x_1, x_2; s_1, s_2) \\
&= \sum_{m_{11}, m_{12}=0}^{\infty} \mathbb{B}_{m_{11}, m_{12}}^{(s_1, s_2)} \frac{x_1^{m_{11}} x_2^{m_{12}}}{m_{11}! m_{12}!} + \frac{1}{x_2} \sum_{n_2=0}^{\infty} B_{n_2} \frac{x_2^{n_2}}{n_2!} \sum_{m_{21}=0}^{\infty} \mathbb{B}_{m_{21}}^{(s_1+s_2)} \frac{(x_1+x_2)^{m_{21}}}{m_{21}!} \\
&= \sum_{m_{11}, m_{12}=0}^{\infty} \mathbb{B}_{m_{11}, m_{12}}^{(s_1, s_2)} \frac{x_1^{m_{11}} x_2^{m_{12}}}{m_{11}! m_{12}!} + \frac{1}{x_2} \sum_{n_2=0}^{\infty} B_{n_2} \frac{x_2^{n_2}}{n_2!} \sum_{m_{21}=0}^{\infty} \mathbb{B}_{m_{21}}^{(s_1+s_2)} \frac{\sum_{i_2=0}^{m_{21}} \binom{m_{21}}{i_2} x_1^{i_2} x_2^{m_{21}-i_2}}{m_{21}!} \\
&= \sum_{m_{11}, m_{12}=0}^{\infty} \mathbb{B}_{m_{11}, m_{12}}^{(s_1, s_2)} \frac{x_1^{m_{11}} x_2^{m_{12}}}{m_{11}! m_{12}!} + \frac{1}{x_2} \sum_{n_2, m_{21}=0}^{\infty} B_{n_2} \mathbb{B}_{m_{21}}^{(s_1+s_2)} \sum_{i_2=0}^{m_{21}} \binom{m_{21}}{i_2} \frac{x_1^{i_2} x_2^{m_{21}-i_2+n_2}}{n_2! m_{21}!} \\
&= \sum_{m_1, m_2=0}^{\infty} \mathbb{B}_{m_1, m_2}^{(s_1, s_2)} \frac{x_1^{m_1} x_2^{m_2}}{m_1! m_2!} + \frac{1}{x_2} \sum_{m_1, m_2=0}^{\infty} B_{m_2} \mathbb{B}_{m_1}^{(s_1+s_2)} \sum_{i_2=0}^{m_1} \binom{m_1}{i_2} \frac{x_1^{i_2} x_2^{m_1-i_2+m_2}}{m_1! m_2!}
\end{aligned}$$

Therefore, from the definition of double-indexed poly-Bernoulli numbers, we have

$$\begin{aligned}
& x_2 \sum_{m_1, m_2=0}^{\infty} \mathbb{B}_{m_1, m_2}^{*,(s_1, s_2)} \frac{x_1^{m_1} x_2^{m_2}}{m_1! m_2!} = x_2 F^*(x_1, x_2; s_1, s_2) \\
&= x_2 \sum_{m_1, m_2=0}^{\infty} \mathbb{B}_{m_1, m_2}^{(s_1, s_2)} \frac{x_1^{m_1} x_2^{m_2}}{m_1! m_2!} + \sum_{m_1, m_2=0}^{\infty} B_{m_2} \mathbb{B}_{m_1}^{(s_1+s_2)} \sum_{i_2=0}^{m_1} \binom{m_1}{i_2} \frac{x_1^{i_2} x_2^{m_1-i_2+m_2}}{m_1! m_2!} \\
&= \sum_{m_1, m_2=0}^{\infty} m_2 \mathbb{B}_{m_1, m_2-1}^{(s_1, s_2)} \frac{x_1^{m_1} x_2^{m_2}}{m_1! m_2!} + \sum_{m_1, m_2=0}^{\infty} B_{m_2} \mathbb{B}_{m_1}^{(s_1+s_2)} \sum_{i_2=0}^{m_1} \binom{m_1}{i_2} \frac{x_1^{i_2} x_2^{m_1-i_2+m_2}}{m_1! m_2!}.
\end{aligned} \tag{3.3}$$

Theorem 1.7 is obtained by comparing the coefficients of this equation.

Since for an odd number  $n_2$  where  $n_2 > 1$   $B_{n_2} = 0$ , Theorem 1.7 becomes

$$\mathbb{B}_{n_1, n_2-1}^{*,(s_1, s_2)} = \mathbb{B}_{n_1, n_2-1}^{(s_1, s_2)}.$$

This gives the following corollary.

**Corollary 3.1** *Let  $k_1, k_2, N$  be positive integers and  $n_2$  be an odd number where  $n_2 > 1$ . Then for a prime number  $p$ , and integers  $n_1, n_2, m_1, m_2 \geq N$  with  $n_1 \equiv m_1 \pmod{p^{N-1}(p-1)}$  and  $n_2 \equiv m_2 \pmod{p^{N-1}(p-1)}$ , we have*

$$\mathbb{B}_{n_1, n_2-1}^{*,(-k_1, -k_2)} \equiv \mathbb{B}_{n_1, m_2-1}^{(-k_1, -k_2)} \pmod{p^N}, \quad \mathbb{B}_{n_1, n_2-1}^{*,(-k_1, -k_2)} \equiv \mathbb{B}_{m_1, n_2-1}^{(-k_1, -k_2)} \pmod{p^N},$$

$$\sum_{i=0}^{\varphi(p^N)-1} \mathbb{B}_{n_1+i, n_2-1}^{*,(-k_1, -k_2)} \equiv 0 \pmod{p^N}, \quad \sum_{i=0}^{\varphi(p^N)-1} \mathbb{B}_{n_1, n_2-1+i}^{*,(-k_1, -k_2)} \equiv 0 \pmod{p^N}.$$

Let  $M$  be positive integers. For  $M = p_1^{e_1} p_2^{e_2} \cdots p_m^{e_m}$ , where  $p_i$ 's are relatively prime, and any integer  $n \geq e_1, e_2, \dots, e_m$ , we have

$$\sum_{i=0}^{\varphi(M)-1} \mathbb{B}_{n_1+i, n_2-1}^{*,(-k_1, -k_2)} \equiv 0 \pmod{M}, \quad \sum_{i=0}^{\varphi(M)-1} \mathbb{B}_{n_1, n_2-1+i}^{*,(-k_1, -k_2)} \equiv 0 \pmod{M}.$$

## 4 Triple-indexed poly-Bernoulli numbers

In this section, we restrict triple-indexed polylogarithms and give similar discussion in the previous section to lead the proof of Theorem 1.8. By the harmonic calculations, we have

$$\begin{aligned} & \text{Li}_{k_1, k_2, k_3}^{\text{III},*}(z_1, z_2, z_3) \\ &= \sum_{1 \leq m_1 \leq m_2 \leq m_3} \frac{z_1^{m_1} z_2^{m_2-m_1} z_3^{m_3-m_2}}{m_1^{k_1} m_2^{k_2} m_3^{k_3}} \\ &= \sum_{1 \leq m_1 < m_2 < m_3} \frac{z_1^{m_1} z_2^{m_2-m_1} z_3^{m_3-m_2}}{m_1^{k_1} m_2^{k_2} m_3^{k_3}} + \sum_{1 \leq m_1 < m_3} \frac{z_1^{m_1} z_3^{m_3-m_1}}{m_1^{k_1+k_2} m_3^{k_3}} \\ &+ \sum_{1 \leq m_1 < m_2} \frac{z_1^{m_1} z_2^{m_2-m_1}}{m_1^{k_1} m_2^{k_2+k_3}} + \sum_{1 \leq m_1} \frac{z_1^{m_1}}{m_1^{k_1+k_2+k_3}} \\ &= \text{Li}_{k_1, k_2, k_3}^{\text{III}}(z_1, z_2, z_3) + \text{Li}_{k_1+k_2, k_3}^{\text{III}}(z_1, z_3) + \text{Li}_{k_1, k_2+k_3}^{\text{III}}(z_1, z_2) + \text{Li}_{k_1+k_2+k_3}^{\text{III}}(z_1). \end{aligned}$$

$F^*(x_1, x_2, x_3; s_1, s_2, s_3)$  is written as

$$\begin{aligned} F^*(x_1, x_2, x_3; s_1, s_2, s_3) &= \frac{\text{Li}_{s_1, s_2, s_3}^{\text{III},*}(1 - e^{-x_1-x_2-x_3}, 1 - e^{-x_2-x_3}, 1 - e^{-x_3})}{(1 - e^{-x_1-x_2-x_3})(1 - e^{-x_2-x_3})(1 - e^{-x_3})} \\ &= \frac{\text{Li}_{s_1, s_2, s_3}^{\text{III}}(1 - e^{-x_1-x_2-x_3}, 1 - e^{-x_2-x_3}, 1 - e^{-x_3})}{(1 - e^{-x_1-x_2-x_3})(1 - e^{-x_2-x_3})(1 - e^{-x_3})} \\ &+ \frac{1}{(1 - e^{-x_2-x_3})} \frac{\text{Li}_{s_1+s_2, s_3}^{\text{III}}(1 - e^{-x_1-x_2-x_3}, 1 - e^{-x_3})}{(1 - e^{-x_1-x_2-x_3})(1 - e^{-x_3})} \\ &+ \frac{1}{(1 - e^{-x_3})} \frac{\text{Li}_{s_1, s_2+s_3}^{\text{III}}(1 - e^{-x_1-x_2-x_3}, 1 - e^{-x_2-x_3})}{(1 - e^{-x_1-x_2-x_3})(1 - e^{-x_2-x_3})} \\ &+ \frac{1}{(1 - e^{-x_2-x_3})(1 - e^{-x_3})} \frac{\text{Li}_{s_1+s_2+s_3}^{\text{III}}(1 - e^{-x_1-x_2-x_3})}{(1 - e^{-x_1-x_2-x_3})}. \end{aligned}$$

In terms of triple-indexed poly-Bernoulli numbers and classical Bernoulli numbers, it is

$$\begin{aligned}
& F^*(x_1, x_2, x_3; s_1, s_2, s_3) \\
&= \sum_{m_{11}, m_{12}, m_{13}=0}^{\infty} \mathbb{B}_{m_{11}, m_{12}, m_{13}}^{(s_1, s_2, s_3)} \frac{x_1^{m_{11}} x_2^{m_{12}} x_3^{m_{13}}}{m_{11}! m_{12}! m_{13}!} + \frac{1}{(1 - e^{-x_2 - x_3})} \sum_{m_{21}, m_{22}=0}^{\infty} \mathbb{B}_{m_{21}, m_{22}}^{(s_1 + s_2, s_3)} \frac{(x_1 + x_2)^{m_{21}} x_3^{m_{22}}}{m_{21}! m_{22}!} \\
&\quad + \frac{1}{(1 - e^{-x_3})} \sum_{m_{31}, m_{32}=0}^{\infty} \mathbb{B}_{m_{31}, m_{32}}^{(s_1, s_2 + s_3)} \frac{x_1^{m_{31}} (x_2 + x_3)^{m_{32}}}{m_{31}! m_{32}!} \\
&\quad + \frac{1}{(1 - e^{-x_2 - x_3})(1 - e^{-x_3})} \sum_{m_{41}=0}^{\infty} \mathbb{B}_{m_{41}}^{(s_1 + s_2 + s_3)} \frac{(x_1 + x_2 + x_3)^{m_{41}}}{m_{41}!} \\
&= \sum_{m_{11}, m_{12}, m_{13}=0}^{\infty} \mathbb{B}_{m_{11}, m_{12}, m_{13}}^{(s_1, s_2, s_3)} \frac{x_1^{m_{11}} x_2^{m_{12}} x_3^{m_{13}}}{m_{11}! m_{12}! m_{13}!} \\
&\quad + \frac{1}{(x_2 + x_3)} \sum_{n_2=0}^{\infty} B_{n_2} \frac{(x_2 + x_3)^{n_2}}{n_2!} \sum_{m_{21}, m_{22}=0}^{\infty} \mathbb{B}_{m_{21}, m_{22}}^{(s_1 + s_2, s_3)} \frac{(x_1 + x_2)^{m_{21}} x_3^{m_{22}}}{m_{21}! m_{22}!} \\
&\quad + \frac{1}{x_3} \sum_{n_3=0}^{\infty} B_{n_3} \frac{x_3^{n_3}}{n_3!} \sum_{m_{31}, m_{32}=0}^{\infty} \mathbb{B}_{m_{31}, m_{32}}^{(s_1, s_2 + s_3)} \frac{x_1^{m_{31}} (x_2 + x_3)^{m_{32}}}{m_{31}! m_{32}!} \\
&\quad + \frac{1}{(x_2 + x_3)x_3} \sum_{n_{41}=0}^{\infty} B_{n_{41}} \frac{(x_2 + x_3)^{n_{41}}}{n_{41}!} \sum_{n_{42}=0}^{\infty} B_{n_{42}} \frac{x_3^{n_{42}}}{n_{42}!} \sum_{m_{41}=0}^{\infty} \mathbb{B}_{m_{41}}^{(s_1 + s_2 + s_3)} \frac{(x_1 + x_2 + x_3)^{m_{41}}}{m_{41}!} \\
&= \sum_{m_1, m_2, m_3=0}^{\infty} \mathbb{B}_{m_1, m_2, m_3}^{(s_1, s_2, s_3)} \frac{x_1^{m_1} x_2^{m_2} x_3^{m_3}}{m_1! m_2! m_3!} \\
&\quad + \frac{1}{(x_2 + x_3)} \sum_{m_2, m_1, m_3=0}^{\infty} B_{m_2} \mathbb{B}_{m_1, m_3}^{(s_1 + s_2, s_3)} \sum_{i_2=0}^{m_2} \binom{m_2}{i_2} \sum_{j_{21}=0}^{m_1} \binom{m_1}{j_{21}} \frac{x_1^{j_{21}} x_2^{i_2 + m_1 - j_{21}} x_3^{m_2 - i_2 + m_3}}{m_1! m_2! m_3!} \\
&\quad + \frac{1}{x_3} \sum_{m_3, m_1, m_2=0}^{\infty} B_{m_3} \mathbb{B}_{m_1, m_2}^{(s_1, s_2 + s_3)} \sum_{j_{32}=0}^{m_2} \binom{m_2}{j_{32}} \frac{x_1^{m_1} x_2^{j_{32}} x_3^{m_3 + m_2 - j_{32}}}{m_1! m_2! m_3!} \\
&\quad + \frac{1}{(x_2 + x_3)x_3} \sum_{m_2, m_3, m_1=0}^{\infty} B_{m_2} B_{m_3} \mathbb{B}_{m_1}^{(s_1 + s_2 + s_3)} \\
&\quad \quad \times \sum_{i_{41}=0}^{m_2} \binom{m_2}{i_{41}} \sum_{j_{41}=0}^{m_1} \sum_{j_{411}=0}^{j_{41}} \binom{m_1}{j_{41}} \binom{j_{41}}{j_{411}} \frac{x_1^{j_{411}} x_2^{i_{41} + j_{41} - j_{411}} x_3^{m_2 - i_{41} + m_3 + m_1 - j_{41}}}{m_1! m_2! m_3!}.
\end{aligned}$$

Therefore, from the definition of triple-indexed poly-Bernoulli numbers, we have

$$\begin{aligned}
& \sum_{m_1, m_2, m_3=0}^{\infty} \mathbb{B}_{m_1, m_2, m_3}^{*, (s_1, s_2, s_3)} \frac{x_1^{m_1} x_2^{m_2+1} x_3^{m_3+1}}{m_1! m_2! m_3!} + \sum_{m_1, m_2, m_3=0}^{\infty} \mathbb{B}_{m_1, m_2, m_3}^{*, (s_1, s_2, s_3)} \frac{x_1^{m_1} x_2^{m_2} x_3^{m_3+2}}{m_1! m_2! m_3!} \\
= & \sum_{m_1, m_2, m_3=0}^{\infty} \mathbb{B}_{m_1, m_2, m_3}^{(s_1, s_2, s_3)} \frac{x_1^{m_1} x_2^{m_2+1} x_3^{m_3+1}}{m_1! m_2! m_3!} + \sum_{m_1, m_2, m_3=0}^{\infty} \mathbb{B}_{m_1, m_2, m_3}^{(s_1, s_2, s_3)} \frac{x_1^{m_1} x_2^{m_2} x_3^{m_3+2}}{m_1! m_2! m_3!} \\
& + \sum_{m_2, m_1, m_3=0}^{\infty} B_{m_2} \mathbb{B}_{m_1, m_3}^{(s_1+s_2, s_3)} \sum_{i_2=0}^{m_2} \binom{m_2}{i_2} \sum_{j_{21}=0}^{m_1} \binom{m_1}{j_{21}} \frac{x_1^{j_{21}} x_2^{i_2+m_1-j_{21}} x_3^{m_2-i_2+m_3+1}}{m_1! m_2! m_3!} \\
& + \sum_{m_3, m_1, m_2=0}^{\infty} B_{m_3} \mathbb{B}_{m_1, m_2}^{(s_1, s_2+s_3)} \sum_{j_{32}=0}^{m_2} \binom{m_2}{j_{32}} \frac{x_1^{m_1} x_2^{j_{32}+1} x_3^{m_3+m_2-j_{32}}}{m_1! m_2! m_3!} \\
& + \sum_{m_3, m_1, m_2=0}^{\infty} B_{m_3} \mathbb{B}_{m_1, m_2}^{(s_1, s_2+s_3)} \sum_{j_{32}=0}^{m_2} \binom{m_2}{j_{32}} \frac{x_1^{m_1} x_2^{j_{32}} x_3^{m_3+m_2-j_{32}+1}}{m_1! m_2! m_3!} \\
& + \sum_{m_2, m_3, m_1=0}^{\infty} B_{m_2} B_{m_3} \mathbb{B}_{m_1}^{(s_1+s_2+s_3)} \sum_{i_{41}=0}^{m_2} \binom{m_2}{i_{41}} \sum_{j_{41}=0}^{m_1} \sum_{j_{411}=0}^{j_{41}} \binom{m_1}{j_{41}} \binom{j_{41}}{j_{411}} \\
& \times \frac{x_1^{j_{411}} x_2^{i_{41}+j_{41}-j_{411}} x_3^{m_2-i_{41}+m_3+m_1-j_{41}}}{m_1! m_2! m_3!} \\
= & \sum_{m_1, m_2, m_3=0}^{\infty} m_2 m_3 \mathbb{B}_{m_1, m_2-1, m_3-1}^{(s_1, s_2, s_3)} \frac{x_1^{m_1} x_2^{m_2} x_3^{m_3}}{m_1! m_2! m_3!} + \sum_{m_1, m_2, m_3=0}^{\infty} m_3 (m_3 - 1) \mathbb{B}_{m_1, m_2, m_3-2}^{(s_1, s_2, s_3)} \frac{x_1^{m_1} x_2^{m_2} x_3^{m_3}}{m_1! m_2! m_3!} \\
& + \sum_{m_2, m_1, m_3=0}^{\infty} m_3 B_{m_2} B_{m_1, m_3-1}^{(s_1+s_2, s_3)} \sum_{i_2=0}^{m_2} \binom{m_2}{i_2} \sum_{j_{21}=0}^{m_1} \binom{m_1}{j_{21}} \frac{x_1^{j_{21}} x_2^{i_2+m_1-j_{21}} x_3^{m_2-i_2+m_3}}{m_1! m_2! m_3!} \\
& + \sum_{m_3, m_1, m_2=0}^{\infty} m_2 B_{m_3} \mathbb{B}_{m_1, m_2-1}^{(s_1, s_2+s_3)} \sum_{j_{32}=0}^{m_2} \binom{m_2}{j_{32}} \frac{x_1^{m_1} x_2^{j_{32}} x_3^{m_3+m_2-j_{32}}}{m_1! m_2! m_3!} \\
& + \sum_{m_3, m_1, m_2=0}^{\infty} m_3 B_{m_3-1} \mathbb{B}_{m_1, m_2}^{(s_1, s_2+s_3)} \sum_{j_{32}=0}^{m_2} \binom{m_2}{j_{32}} \frac{x_1^{m_1} x_2^{j_{32}} x_3^{m_3+m_2-j_{32}}}{m_1! m_2! m_3!} \\
& + \sum_{m_2, m_3, m_1=0}^{\infty} B_{m_2} B_{m_3} \mathbb{B}_{m_1}^{(s_1+s_2+s_3)} \sum_{i_{41}=0}^{m_2} \binom{m_2}{i_{41}} \sum_{j_{41}=0}^{m_1} \sum_{j_{411}=0}^{j_{41}} \binom{m_1}{j_{41}} \binom{j_{41}}{j_{411}} \\
& \times \frac{x_1^{j_{411}} x_2^{i_{41}+j_{41}-j_{411}} x_3^{m_2-i_{41}+m_3+m_1-j_{41}}}{m_1! m_2! m_3!}.
\end{aligned}$$

Comparing the coefficients of this equation leads to Theorem 1.8.

**Remark 3** Similar discussion can be made for  $n$ -indexed star-poly-Bernoulli numbers with  $n \geq 4$ . For example, in the case of  $n = 4$ , it is expected that single, double, triple and

*quadruple-indexed poly-Bernoulli numbers will appear. Next stage is to find the regularity there and obtain the general formula.*

## References

- [AIK] T. Arakawa, T. Ibukiyama and M. Kaneko, *Bernoulli Numbers and Zeta Functions*, Springer, Tokyo, 2014.
- [AK1] T. Arakawa and M. Kaneko, Multiple zeta values, poly-Bernoulli numbers, and related zeta functions, *Nagoya Math.J.*, **153** (1999), 189–209.
- [AK2] T. Arakawa and M. Kaneko, On multiple L-values, *J. Math. Soc. Japan*, **56** (2004), 967–991.
- [BH] A. Bayad and Y. Hamahata, Multiple polylogarithms and multi-poly-Bernoulli poly-nomials, *Funct. Approx. Comment. Math.* **46** (2012), 45–61.
- [G] A.B. Goncharov, Multiple polylogarithms, cyclotomy, and modular complexes, *Math.Res.Lett.*, **5** (1998), 497–516.
- [HM] Y. Hamahata and H. Masubuchi, Special multi-poly-Bernoulli numbers, *J. Integer Seq.* **10** (2007), Article 07.4.1.
- [K] M. Kaneko, poly-Bernoulli numbers, *Jour. Th. Nombre Bordeaux*, **9** (1997), 199–206.
- [KT] M. Kaneko and H. Tsumura, Multi-poly-Bernoulli numbers and related zeta functions, *Nagoya Math.J.*, **232** (2018), 19–54.
- [OS] Y. Ohno and M. Sakata, On certain properties of poly-Bernoulli numbers with negative index, *J. School sci. Eng.*, **49** (2013), 5–7.
- [S] M. Sakata, On p-adic properties of poly-Bernoulli numbers (in Japanese), *Kindai University*, master thesis (2012).

YUNA BABA  
Faculty of Science and Technology,  
Sophia University,

7-1 Kio-cho, Chiyoda-ku, Tokyo, 102-8554, Japan

MAKI NAKASUJI

Department of Information and Communication Science, Faculty of Science and Technology,  
Sophia University,

7-1 Kio-cho, Chiyoda-ku, Tokyo, 102-8554, Japan

`nakasuji@sophia.ac.jp`

and

Mathematical Institute,

Tohoku University,

Sendai 980-8578, Japan

MIKA SAKATA

Department of Health and Sport management, School of Physical Education,  
Osaka University of Health and Sport Sciences,

1-1 Asashirodai, Kumatori-cho, Sennan-gun, Osaka, 590-0496, Japan

`m.sakata@ouhs.ac.jp`