

# DISTRIBUTION ESTIMATION AND CHANGE-POINT DETECTION FOR TIME SERIES VIA DNN-BASED GANS

JIANYA LU, YINGJUN MO, ZHIJIE XIAO, LIHU XU, AND QIURAN YAO

**ABSTRACT.** The generative adversarial networks (GANs) have recently been applied to estimating the distribution of independent and identically distributed data, and got excellent performances. In this paper, we use the blocking technique to demonstrate the effectiveness of GANs for estimating the distribution of stationary time series. Theoretically, we obtain a non-asymptotic error bound for the Deep Neural Network (DNN)-based GANs estimator for the stationary distribution of the time series. Based on our theoretical analysis, we put forward an algorithm for detecting the change-point in time series. We simulate in our first experiment a stationary time series by the multivariate autoregressive model to test our GAN estimator, while the second experiment is to use our proposed algorithm to detect the change-point in a time series sequence. Both perform very well. The third experiment is to use our GAN estimator to learn the distribution of a real financial time series data, which is not stationary, we can see from the experiment results that our estimator cannot match the distribution of the time series very well but give the right changing tendency.

**Key words:** time series; blocking technique; s-dependence form; GAN; Wasserstein distance; nonasymptotic error bounds; change-point detection

## 1. INTRODUCTION

Let  $\{\mathbf{X}_n\}_{n \in \mathbb{N}}$  be a stationary time series with the stationary measure  $\pi$  satisfying appropriate regularity assumptions, see the details in Section 2.2. We aim in this paper to investigate the validity of DNN-based generative adversarial networks (GANs) for estimating the distribution of this time series, and propose an algorithm based on our theoretical result to detect the change-point in time series.

GANs are a class of deep learning methods designed to estimate the distribution of data [Goodfellow et al. (2014)]. The approach is usually implemented by training two neural networks with conflicting objectives, one generator (g), and one discriminator (d), forcing each other to improve. The generator is designed to produce samples that look real, while the discriminator measures the distance between generated samples and the real data.

In the research of GANs, extensive theoretical and empirical analysis have been established for independent and identically distributed (i.i.d.) data, however limited work has exploited them for estimating the distribution of time-series data, which is important in many applications. In addition, the evaluation of GANs remains a largely-unsolved problem, researchers often rely on visual evaluation of generated examples, an approach that is both impractical and inappropriate for multi-dimensional time series. There have been several results related to the GANs for time series, but most of them only showed the successes in experiments without a rigorous justification, see more details in the literature review below.

Our first main result contributes to the theory of the GAN method for estimating the distribution of stationary time-dependent time series. We establish the convergence rate of the proposed estimator measured by integral probability metric under the Hölder evaluation function class. Our Theorem 3.1 below provides a theoretical guarantee for the application of GANs in time series under certain conditions.

We also notice that the time series sometimes may not admit the stationary distribution. There might have been abrupt variations in time series data. The detection of change-points gains popularity in modeling and prediction of time series, for example, in finance, biology, engineering, etc. Most of the traditional change-point detection methods conduct the hypothesis test for the density function to monitor the change-point [Harchaoui et al. (2009)]. However, these methods may have low detection efficiency when encountering high-dimensional time series data. Some change-point detection methods for high-dimensional time series indirectly use the model selection or testing to build the parametric detection model [Cho and Fryzlewicz (2015), Chen and Zhang (2015)]. The algorithms of parametric models are hard to extend to the massive data case due to the computational cost and some strong assumptions [Chen and Zhang (2015)]. Motivated by the property that GANs can directly learn the distribution of high-dimensional complex time series data, we propose a simple algorithm without any constraint to detect the change-point directly by tracking the variations in the distribution of the time series based on GANs.

**1.1. Literature review.** The research of the GANs architecture for sequential data has received more and more attention. In recent years, many methods of modeling time series with GANs have been proposed, such as continuous recurrent neural networks with adversarial training (C-RNN-GAN) [Mogren (2016)], Recurrent Conditional GAN (RCGAN) [Esteban et al. (2017)], time series-GAN [Yoon et al. (2019)], Conditional Sig-Wasserstein GANs (SigCW-GAN) [Ni et al. (2020)]. However, they only showed the success of the GAN-based methods for realistic sequence generation in experiments but there is no theory to support the validity of this model. We refer to the survey [Brophy et al. (2021)] for more details about GANs in time series.

Compared with the vast number of theoretical results for neural networks used to approximate functions, the use of neural networks to express distributions is much less theoretically understood. Some literature focus on how well GAN and its variants can express probability distributions. Lu and Lu (2020) studied the universal approximation property of deep neural networks for representing probability distributions, Zhu et al. (2020) studied a fundamental trade-off in the approximation error and statistical error in GANs, Huang et al. (2022) studied the convergence rates of GAN estimators under the integral probability metric with the Hölder function class. Liang (2021) established the optimal minimax rates for distribution estimation under GANs in view of nonparametric density estimation. Zhang et al. (2017) explored the generalization capacity of GANs and extended the generalization bounds for neural net distance.

For the theoretical analysis of GANs, many scholars consider the probability distribution estimating of i.i.d. data by GANs. Gao et al. (2022) estimated the error bound of the approximation of Wasserstein GANs using GroupSort neural networks as discriminators. Bai et al. (2019) used GANs with some special discriminators to learn distributions of i.i.d. samples in Wasserstein distance and KL-divergence. Liu et al. (2021) introduced a nearly sharp bound for the bidirectional GAN estimation error under the Dudley distance. We refer the reader to [Chae et al. (2021); Taghvaei and Jalali (2019)] for more details. Although a lot of work on the theoretical basis of GAN has appeared in academia, there seems very few of theoretical analysis for generating time-dependent data.

The traditional high-dimensional change-point detections are around two aspects: model selection and testing. Tibshirani and Wang (2008), Li et al. (2016), Harchaoui and Lévy-Leduc (2010) utilized the Lasso-type regularization model to estimate the change points and improve

the computational efficiency. For the aspect of testing, [Harchaoui et al. \(2009\)](#) used a regularized kernel-based test statistic to test the consistency of distributions between time-series samples. [Enikeeva and Harchaoui \(2019\)](#) considered the hypothesis test to detect the change point in a set of high-dimensional Gaussian vectors. [Chen and Zhang \(2015\)](#) proposed a nonparametric approach based on the graphs to test the similarity between observations. The aggregation technique testing whether there exists a gap between two time regions is also a popular way for change-point analysis. For example, [Jirak \(2015\)](#) showed asymptotic properties by using the coordinate-wise test statistic to detect the single change point. [Bai \(2010\)](#) theoretically analyzed the consistency of the single-change point in panel data. [Chen et al. \(2021\)](#) improved the aggregation way of the data and explored to detect the multiple change-points in high-dimensional time series. For the change-point detection of complex high-dimensional data, [Romanenkova et al. \(2022\)](#) considered the change-point detection of the complex sequential stream, video data, by representation learning for the deep model. [Chang et al. \(2019\)](#) combined the kernel two-sample test with recurrent neural networks (RNNs) to detect the different types of change-points. They introduced an auxiliary generative model to generate an approximate distribution and use the maximum mean discrepancy to conduct the two-sample test, but there is no information about the location of the change-point. Although there are various change-point detection techniques, fewer works directly focus on tracking the distribution of time series.

**1.2. Our contributions.** Our main contributions are summarized as the following two aspects. Our first contribution is theoretically confirming that GANs are effective for estimating the distribution of stationary time series. More precisely, for the stationary time series whose dependence satisfies appropriate regularity assumptions, we show that DNN-based GANs can be applied to learn its stationary distribution, i.e., like the i.i.d. samples, the time series can be fed into the device of the neural networks of the GANs to obtain a stable generator  $\hat{g}$  such that  $\hat{g}(z)$  has a distribution very close to  $\pi$  ( $z$  is a Gaussian random variable). We prove a non-asymptotic error bound between the estimator of the measure  $\pi$  and itself, which clearly demonstrates the estimator is consistent and thus effective. Our proof of this theoretical result is based on a typical block technique, see [[Berkes et al. \(2014\)](#); [Gouëzel \(2010\)](#); [Liu and Lin \(2009\)](#); [Lu et al. \(2022\)](#)].

By the validity of GANs for estimating the distribution of stationary times series, we put forward an algorithm based on GANs to detect the single change-point for time series. The idea is straightforward. Since the observed data on both sides of the change point follow two different stationary distributions, there exists an inconsistency of distributions of the observed data around the change point. This inspires us to learn the initial stationary distribution by GAN and take it as a benchmark to detect the change in the distributions of the subsequent data. Specifically, we compute the loss value of GAN using the trained discriminator and generator as the measure. When we use GAN’s loss to evaluate a sequence of observed data following the same distribution as the initial stationary distribution, the value would be consistently small. On the contrary, the GAN’s loss value would be consistently large for the data from another distribution. If the loss value between the adjacent sequences is inconsistent, this alerts us that there exists a change in the distributions. Following this idea, we design an effective algorithm to accurately locate the change point. We also implement three experiments to validate our main results. In the first experiment, we show some generated examples by GAN learning a set of stationary sequences from the multivariate autoregressive model to validate our theoretical results. The results show that GANs can generate higher-quality simulated time series data. The second experiment demonstrates an example of our proposed single change-point detection algorithm, in which the change point can be accurately detected. In the third experiment, we

also test the performance of GANs on a set of nonstationary financial time series, the generated samples can not better overlap with the real data, but they reveal the general tendency of the distribution of time series.

**1.3. Organization of the paper and some notations.** The paper is organized as follows. In Section 2, we provide some preliminary knowledge and assumptions for our main result. In Section 3, we give the convergence rate of GANs estimator for time series and the algorithm to detect the change-point in a time series sequence. Three examples of simulation and real data analysis are given in Section 4. The proof of our main result is deferred to Section 5.

We finish this section by introducing some notations which will be frequently used in sequel. Denote  $[n] := \{1, \dots, n\}$ . Given a vector  $\mathbf{w} \in \mathbb{R}^p$ ,  $\|\mathbf{w}\|$  will refer to the Euclidean norm, and for  $h \geq 1$ ,  $\|\mathbf{w}\|_h = (\sum_{i=1}^p |\mathbf{w}_i|^h)^{1/h}$  will refer to the  $\ell_h$  norm,  $\|\mathbf{w}\|_\infty$  refer to the largest elements of  $\mathbf{w}$ . Given a matrix  $\mathbf{W}$  and reals  $h, l \geq 1$ , we let  $\|\mathbf{W}\|_{h,l} := (\sum_k (\sum_j |\mathbf{W}_{j,k}|^h)^{l/h})^{1/l}$  denote the  $l$ -norm of the  $h$ -norms of the columns of  $\mathbf{W}$ .

For a real-valued function  $f : \mathcal{X} \rightarrow \mathbb{R}$ , we define  $\|f\|_\infty := \mathbf{P}_{x \in \mathcal{X}} |f(x)|$ . Let  $\mathfrak{L}(\Omega, c)$  be all the real-valued Lipschitz functions from  $\Omega$  to  $\mathbb{R}$  with Lipschitz constant  $c$ , that is,

$$\mathfrak{L}(\Omega, c) := \{f : \Omega \rightarrow \mathbb{R} : |f(z_1) - f(z_2)| \leq c \|z_1 - z_2\| \text{ for any } z_1, z_2 \in \Omega\}.$$

For  $b > 0$  with  $b = r + q$ , where  $r \in \mathbb{N}_0 := \mathbb{N} \cup \{0\}$ ,  $q \in (0, 1]$ , denote the Hölder class with smoothness index  $b$  as

$$\mathcal{H}^b(\mathbb{R}^p) := \left\{ f : \mathbb{R}^p \rightarrow \mathbb{R}, \mathbf{m}_{\|\alpha\|_1 \leq r} \|\partial^\alpha f\|_\infty \leq 1, \mathbf{m}_{\|\alpha\|_1 = r} \mathbf{P}_{x \neq y} \frac{|\partial^\alpha f(x) - \partial^\alpha f(y)|}{\|x - y\|^q} \leq 1 \right\},$$

and we denote  $\mathcal{H} = \mathcal{H}^b(\mathbb{R}^p)$  for the simplicity. For any subset  $\mathcal{X} \subseteq \mathbb{R}^p$ , we denote  $\mathcal{H}^b(\mathcal{X}) := \{f : \mathcal{X} \rightarrow \mathbb{R}, f \in \mathcal{H}^b(\mathbb{R}^p)\}$ .

Let  $\mu$  and  $\nu$  be a pair of probability measures on a space  $\mathcal{X}$ , and  $\mathcal{F}$  be a class of functions  $f : \mathcal{X} \rightarrow \mathbb{R}$  that are integrable with respect to  $\mu$  and  $\nu$ . Define integral probability metric as

$$d_{\mathcal{F}}(\mu, \nu) := \mathbf{P}_{f \in \mathcal{F}} \left| \int f(d\mu - d\nu) \right| = \mathbf{P}_{f \in \mathcal{F}} |\mathbb{E}_\mu[f(X)] - \mathbb{E}_\nu[f(Z)]|.$$

Wasserstein distance between two probability measures  $\mu$  and  $\nu$  is defined as

$$\mathcal{W}_1(\mu, \nu) := \mathbf{h}_{\gamma \in \Gamma(\mu, \nu)} \int_{\mathcal{X} \times \mathcal{X}} \|x - y\| d\gamma(x, y),$$

where  $\Gamma(\mu, \nu)$  denotes the collection of all measures on  $\mathcal{X} \times \mathcal{X}$  with marginals  $\mu$  and  $\nu$  on the first and second factors respectively.

By the duality theorem of Kantorovich and Rubinstein (1958), we have

$$\mathcal{W}_1(\mu, \nu) = d_{\text{Lip}(\mathcal{X}, 1)}(\mu, \nu) = \mathbf{P}_{f \in \text{Lip}(\mathcal{X}, 1)} \left\{ \int_{\mathcal{X}} f(x) (\mu - \nu)(x) \mid \text{continuous } f : \mathcal{X} \rightarrow \mathbb{R} \right\}.$$

## 2. PRELIMINARIES

In this section, we give the preliminaries about the definition of GANs, the assumptions on the time series, and the architectures of DNNs that we will use to model the generator and discriminator.

**2.1. GANs.** Let  $\mathcal{G}$  and  $\mathcal{D}$  be the collections of neural networks. Let  $g \in \mathcal{G}$  be the generator function from  $\mathbb{R}$  to  $\mathbb{R}^p$  and  $d \in \mathcal{D}$  be the discriminator function from  $\mathbb{R}^p$  to  $\mathbb{R}$ . GANs can be formulated as the following optimal problem,

$$(2.1) \quad g^* \in \underset{g \in \mathcal{G}}{\text{gm}} \underset{d \in \mathcal{D}}{\text{m}} \{ \mathbb{E}_{x \sim \pi} [d(x)] - \mathbb{E}_{z \sim \nu} [d(g(z))] \},$$

where  $\nu$  is the Gaussian distribution defined on the one-dimensional space  $\mathbb{R}$  to generate fake signal. For convenience,  $\mathcal{D}$  is assumed in symmetric class, which means if  $d \in \mathcal{D}$ , then  $-d \in \mathcal{D}$ . For any subset  $\mathcal{X} \subseteq \mathbb{R}^p$ , we denote  $\mathcal{D}(\mathcal{X}) := \{f : \mathcal{X} \rightarrow \mathbb{R}, f \in \mathcal{D}(\mathbb{R}^p)\}$ . The empirical version of (2.1) is considered as

$$(2.2) \quad \hat{g} \in \underset{g \in \mathcal{G}}{\text{gm}} \underset{d \in \mathcal{D}}{\text{m}} \left\{ \frac{1}{n} \sum_{i=1}^n d(\mathbf{X}_i) - \frac{1}{m} \sum_{i=1}^m d(g(z_i)) \right\},$$

where  $\{z_i\}_{i=1, \dots, m}$  are i.i.d. random variables with distribution  $\nu$ .

**2.2. Time Series Assumptions.** Wu (2005) proposed predictive dependence measures to provide another look at the fundamental issue of dependence. Using this tool, he established a theory for high-dimensional inferences under dependence. It is different from most of the current research on a high-dimensional inference that assumes the underlying observations are independent.

The  $p$ -dimensional stationary time series in Wu (2005) is in the form of

$$(2.3) \quad \mathbf{X}_n := \mathbf{G}(\dots, \varepsilon_{n-1}, \varepsilon_n),$$

where  $\varepsilon_n, n \in \mathbb{Z}$  are i.i.d. random variables and  $\mathbf{G}$  is a measurable function such that  $\mathbf{X}_n$  is well-defined.

Let  $(\varepsilon'_i)$  be an i.i.d copy of  $(\varepsilon_i)$ ,  $\xi_i = (\dots, \varepsilon_{i-1}, \varepsilon_i)$  and  $\xi'_i = (\dots, \varepsilon'_{i-1}, \varepsilon'_i)$ . For any set  $I \subset \mathbb{Z}$ ,  $\varepsilon_{j,I} = \varepsilon'_j$  if  $j \in I$  and  $\varepsilon_{j,I} = \varepsilon_j$  if  $j \notin I$ ;  $\xi_{i,I} = (\dots, \varepsilon_{i-1,I}, \varepsilon_{i,I})$ .

Define the predictive dependence measure

$$(2.4) \quad \omega_\beta(I, n) = [\mathbb{E}(\|\mathbb{E}(\mathbf{X}_n | \xi_0) - \mathbb{E}(\mathbf{X}_n | \xi_{0,I})\|^\beta)]^{1/\beta},$$

where  $n \geq 0, \beta \geq 1$ .

Under the assumption that  $\{\mathbf{X}_n\}_{n \in \mathbb{N}}$  satisfies geometric moment contraction,

$$(2.5) \quad \mathbb{E}\|\mathbf{X}_n - \mathbf{G}(\xi'_0, \varepsilon_1, \dots, \varepsilon_n)\|^\beta \leq C_1 e^{-C_2 n},$$

Wu (2007) extended KMT approximation from independent random variables to a large class of dependent stationary processes. Berkes et al. (2014) extended this result to optimal rate.

**Assumption 2.1.** We assume that the time series  $\{\mathbf{X}_n\}_{n \in \mathbb{N}}$  in the form of (2.3) satisfies geometric moment contraction, that is,

$$(2.6) \quad \mathbb{E}\|\mathbf{X}_n - \mathbf{G}(\xi'_0, \varepsilon_1, \dots, \varepsilon_n)\|^\beta \leq C_1 e^{-C_2 n},$$

where  $C_1$  and  $C_2$  are positive constants,  $n \geq 0, \beta \geq 1, \xi_0 = (\dots, \varepsilon_{-1}, \varepsilon_0), \xi'_0 = (\dots, \varepsilon'_{-1}, \varepsilon'_0)$  and  $\{\varepsilon'_n\}_{n \in \mathbb{Z}}$  is an i.i.d. copy of  $\{\varepsilon_n\}_{n \in \mathbb{Z}}$ .

**Assumption 2.2.** The stationary measure  $\pi$  of time series  $\{\mathbf{X}_n\}_{n \in \mathbb{N}}$  is sub-Gaussian, that is, there exist  $\mathbf{m} \in \mathbb{R}^p$  and  $v > 0$  such that

$$\mathbb{E}_{x \sim \pi} \left[ \exp\left\{ \frac{1}{2} \alpha^T (x - \mathbf{m}) \right\} \right] \leq e^{\|\alpha\|^2 v^2 / 2},$$

for all  $\alpha \in \mathbb{R}^p$ , where  $T$  is the transpose operator. Without loss of generality, we assume  $\mathbf{m} = \mathbf{0}$ .

**2.3. DNN Model Assumptions.** We consider the feed-forward DNN as the L-compositional function class indexed by parameter

$$(2.7) \quad \Theta = (\mathbf{W}_0, \mathbf{W}_1, \dots, \mathbf{W}_L),$$

where  $\mathbf{W}_l \in \mathbb{R}^{D_{l+1} \times D_l}$ , for  $l = 0, 1, \dots, L$ ,  $D_0, \dots, D_{L+1} \in \mathbb{N}$ .  $L$  and  $N := \mathfrak{n} \{D_1, \dots, D_L\}$  are depth and width of the DNNs respectively.

We denote the DNN with an input  $\mathbf{x} \in \mathbb{R}^p$  by

$$(2.8) \quad \mathcal{NN}_p(N, L) := \{f(\mathbf{x}; \Theta) = \mathbf{W}_L \sigma_L(\dots \mathbf{W}_1 \sigma(\mathbf{W}_0 \mathbf{x}) \dots) \in \mathbb{R}^{D_{L+1}} | \Theta\}.$$

When the input dimension  $p$  is clear from contexts, we simply denote it by  $\mathcal{NN}(N, L)$ .

We consider that the neural networks  $\mathcal{NN}(N, L)$  in (2.8) has ReLU activation functions  $\left\{ \sigma_j(\mathbf{z}) = (z_1 \vee 0, \dots, z_{D_j} \vee 0) \text{ for } \mathbf{z} := (z_1, \dots, z_{D_j})^T \in \mathbb{R}^{D_j} \right\}_{j=1}^L$ .

**Assumption 2.3.** We further denote  $\mathcal{NN}(N, L, K)$  by the function of  $\mathcal{NN}(N, L)$  with

$$\prod_{l=0}^L \|\mathbf{W}_l\| \leq K,$$

where  $\|\cdot\|$  is the operator norm of a matrix.

*Remark 2.4.* Since ReLU function is a Lipschitz function with Lipschitz constant 1, it is easy to see that for any function  $f \in \mathcal{NN}(N, L, K)$ , its Lipschitz constant is smaller than  $\prod_{l=0}^L \|\mathbf{W}_l\|$  which is bounded by  $K$ , that is,

$$\|f(\mathbf{x}) - f(\mathbf{y})\| \leq K \|\mathbf{x} - \mathbf{y}\|.$$

For more details, see [Jiao et al. \(2022, Section 3\)](#).

### 3. MAIN RESULTS

In this chapter, two main results are introduced, the first result is the first result is a non-asymptotic error bound between the stationary measure  $\pi$  of the time series and its estimator  $\hat{\mathbf{g}}_{\#}\nu$  which is defined below. Based on the theoretical result, the second main result for detecting the single change-point in time series sequence is also introduced.

**3.1. Non-asymptotic bound for the estimator.** The following theorem is our first main result, whose proof will be postponed to the last section.

Let the stationary distribution of time series  $\mathbf{X}_n := \mathbf{G}(\dots, \varepsilon_{n-1}, \varepsilon_n)$  in the form of (2.3) be  $\pi$ . Here  $\pi$  can be a high-dimensional distribution. Our goal is to establish a non-asymptotic error bound of the GANs estimator for  $\pi$  based on integral probability metric,

$$d_{\mathcal{H}}(\pi, \hat{\mathbf{g}}_{\#}\nu) = \inf_{h \in \mathcal{H}} \left\{ \mathbb{E}_{\mathbf{x} \sim \pi}[h(\mathbf{x})] - \mathbb{E}_{\mathbf{z} \sim \nu}[h(\hat{\mathbf{g}}(\mathbf{z}))] \right\},$$

where  $\hat{\mathbf{g}}_{\#}\nu$  is the push-forward measure defined as  $\hat{\mathbf{g}}_{\#}\nu(A) = \nu(\hat{\mathbf{g}}^{-1}(A))$  for measurable set  $A \subset \mathbb{R}^p$ . For ease of notation, we use  $\mathbb{E}_{\gamma}[f]$  instead of  $\mathbb{E}_{\mathbf{x} \sim \gamma}[f(\mathbf{x})]$  for any distribution  $\gamma$  and function  $f$ .

**Theorem 3.1** (Non-asymptotic Bound). *Assume measure  $\nu$  is absolutely continuous on  $\mathbb{R}$  and  $\mathbf{g}_{\#}\nu$  is supported on  $[-a_n, a_n]^p$  for all  $\mathbf{g}$ ,  $a_n = \mathfrak{g} \quad pn$ .*

*Suppose  $\mathbb{E} \mathfrak{n} \quad i \in [m] \|z_i\| < \infty$ ,  $z_i \sim \nu$  and Assumptions 2.1-2.3 are satisfied. The evaluation class is  $\mathcal{H} = \mathcal{H}^b(\mathbb{R}^p)$  with  $b = r + q$ ,  $r \in \mathbb{N}_0$  and  $q \in (0, 1]$ .*

*Then, for the given sample size  $n$ , there exists a generator*

$$\mathcal{G} = \{\mathbf{g} \in \mathcal{NN}(N_{\mathcal{G}}, L_{\mathcal{G}}, K_{\mathcal{G}}) : \mathbf{g}(\mathbb{R}) \subseteq [-\mathfrak{g} \quad pn), \mathfrak{g} \quad pn)^p\},$$

with  $N_G \geq 7p + 1$ ,  $L_G \geq 2$  and a discriminator

$$\mathcal{D} = \{d \in \mathcal{NN}(N_D, L_D, K_D) : d(\mathbb{R}^p) \subseteq [-B, B]\} \cap \mathbb{K}(\mathbb{R}^p, C), B, C < \infty,$$

with  $N_D \geq c(K_D/\mathfrak{h}^{-\gamma} K_D)^{(2p+b)/(2p+2)}$ ,  $L_D \geq 4\gamma + 2$ ,  $\gamma := \lceil \mathfrak{h}^{-2}(p+r) \rceil$ , such that

$$n \leq \frac{N_G - p - 1}{2} \left\lfloor \frac{N_G - p - 1}{6p} \right\rfloor \left\lfloor \frac{L_G}{2} \right\rfloor + 2,$$

such that GAN estimator  $\hat{g}$  satisfies

$$\begin{aligned} d_{\mathcal{H}}(\pi, \hat{g}_{\#}\nu) &:= \sup_{h \in \mathcal{H}} \left\{ \mathbb{E}_{\mathbf{x} \sim \pi} [h(\mathbf{x})] - \mathbb{E}_{\mathbf{z} \sim \nu} [h(\hat{g}(\mathbf{z}))] \right\} \\ &\leq O\left(\left(\mathfrak{h}^{-pn}\right)^b (K_D/\mathfrak{h}^{-\gamma} K_D)^{-\frac{b}{p+1}}\right) + O(K_D e^{-C_2 n^\alpha}) + O\left(\sqrt{\frac{t_1}{n^{1-\alpha}}}\right) \\ &\quad + O\left(\sqrt{\frac{1}{n}} K_D \sqrt{L_D + \mathfrak{h}^{-p}}\right) + O\left(\frac{1}{m} K_D K_G \sqrt{L_D + L_G} + \sqrt{\frac{t_2}{m}}\right), \end{aligned}$$

with probability at least  $1 - e^{-C_2 n^\alpha/2\beta} - 2e^{-t_1} - e^{-t_2}$ , for  $t_1, t_2 > 0$ ,  $0 < \alpha < 1/2$ .

*Remark 3.2.* The error of estimating the distribution with i.i.d. sample data by GANs is given as  $n^{-b/p} \vee n^{-1/2} \mathfrak{h}^{-c(b,p)} n$ , where  $c(b, p)$  is a positive constant depending on the index  $b$  of Hölder class and the dimension  $p$ , see [Huang et al. (2022)]. Considering the time series case, we further assume that  $K_D = Cn^{\frac{1}{2} - \frac{b}{2(p+1+b)}}$  and  $m \geq CnK_G\sqrt{L_G}$ , and get for large enough sample size  $n$ ,

$$d_{\mathcal{H}}(\pi, \hat{g}_{\#}\nu) \leq O\left(\left(\mathfrak{h}^{-pn}\right)^b n^{-\frac{b}{2(p+1+b)}}\right) + O\left(n^{-\frac{1}{2}(1-\alpha)}\right),$$

with high probability. Here  $\alpha \in (0, \frac{1}{2})$  is the parameter in the blocking technique to handle the dependence issue of time series. For any  $p$  and  $b$ , one can choose  $\alpha < \frac{p+1}{p+1+b}$  such that

$$d_{\mathcal{H}}(\pi, \hat{g}_{\#}\nu) \leq O\left(\left(\mathfrak{h}^{-pn}\right)^b n^{-\frac{b}{2(p+1+b)}}\right),$$

which is close to the error of of i.i.d. case as  $p \geq 2b$ , i.e.  $n^{-b/p}$ .

**3.2. Single change-point detection for time series.** Let  $\mathbf{X} = \{\mathbf{X}_1, \dots, \mathbf{X}_n\}$  be a  $p$ -dimensional time series sequence such that there exists a  $\tau \in \mathbb{N}$  such that  $\{\mathbf{X}_1, \dots, \mathbf{X}_\tau\}$  is stationary with a stationary distribution  $\pi$  and that  $\{\mathbf{X}_{\tau+1}, \dots, \mathbf{X}_n\}$  is also stationary but with a distribution  $\pi'$ . This form of time series is called the time series with a single change-point [Chen and Zhang (2015)]. Detecting the change-point  $\tau$  is an important research topic in Statistics and time series.

Based on the theoretical result in the previous subsection, we put forward an algorithm for detecting the single change-point in time series using GANs, see Algorithm 1 below. It is reasonable to assume that  $\tau$  is sufficiently large and that we can use the first  $t$  sample points, with  $t$  being sufficiently large but smaller than  $\tau$ , in  $\mathbf{X}$  to train a stable GAN model. In order to use the following  $K$ -block detection procedure, with  $K$  some number satisfying  $1 \ll K \ll n$ , we need to assume  $\tau < n - K$ .

The process for detecting  $\tau$  contains three stages. The **first** stage is to train a stable GAN model by the first  $t$  sample points in  $\mathbf{X}$ , i.e., we obtain a generator  $g$  and a discriminator  $d$ . At the **second** stage, we divide the remaining time series  $\{\mathbf{X}_{t+1}, \dots, \mathbf{X}_n\}$  into  $\lceil (n -$

$t)/K\lceil$  blocks, each having  $K$  samples, as the following:  $\mathbf{B}_1, \dots, \mathbf{B}_{\lceil(n-t)/K\rceil}$ , where  $\mathbf{B}_i = \{\mathbf{X}_{t+(i-1)K+1}, \dots, \mathbf{X}_{t+iK}\}$  for each  $i$ . For every  $\mathbf{B}_i$ , we compute

$$l_i = \frac{1}{K} \sum_{\mathbf{X}_j \in \mathbf{B}_i} d(\mathbf{X}_j) - \frac{1}{K} \sum_{j=1}^K d(g(\mathbf{z}_j)), \quad \text{for } i = 1, \dots, \lceil(n-t)/K\rceil,$$

where  $\mathbf{z}_1, \dots, \mathbf{z}_K$  are i.i.d. standard normal distributed random variables. Obviously, for the blocks that contain the samples following the same distribution as the ones used to train GAN, the loss  $l_k$  would be much smaller than the ones containing the samples with different distributions. The  $l_k$  of the block in which the change point is located will be very different from that of its neighboring blocks. Motivated by these properties, we can compute the difference,  $|l_{i+1} - l_i|$ , between the neighboring two blocks and find some  $\mathbf{B}_k$  and  $\mathbf{B}_{k+1}$  that have the largest difference on the loss values. After determining the two candidate blocks  $\mathbf{B}_k, \mathbf{B}_{k+1}$ , we merge them into one group,  $\mathbf{G} = \{\mathbf{B}_k, \mathbf{B}_{k+1}\} = \{\dot{\mathbf{X}}_1, \dots, \dot{\mathbf{X}}_{2K}\}$ . This operation can help us narrow down the time series data set and further facilitate the detection of change-points. In the **third** stage, for each sample  $\dot{\mathbf{X}}_j \in \mathbf{G}$  for  $j = 1, \dots, 2K$ , we build a rolling window  $\mathbf{H}_j = \{\dot{\mathbf{X}}_{j-w}, \dots, \dot{\mathbf{X}}_j, \dots, \dot{\mathbf{X}}_{j+w}\}$  centered on the sample  $\dot{\mathbf{X}}_j$  and with the radius  $w$  to do the detection. In each loop, all the samples in  $\mathbf{H}_j$  are used to compute

$$\pi_j = \frac{1}{2w+1} \sum_{\dot{\mathbf{X}}_i \in \mathbf{H}_j} d(\dot{\mathbf{X}}_i) - \frac{1}{2w+1} \sum_{i=1}^{2w+1} d(g(\mathbf{z}_i)) \quad \text{for } j = 1, \dots, 2K.$$

where  $\mathbf{z}_1, \dots, \mathbf{z}_{2w+1}$  are i.i.d. standard normal distributed random variables. At the change-point (e.g.,  $\dot{\mathbf{X}}_{j^*}$ ), there exists an obvious difference between the loss of two adjacent windows,  $\pi_{j^*}$  and  $\pi_{j^*-1}$ , because the window  $\mathbf{H}_{j^*}$  contains more samples following different distribution than  $\mathbf{H}_{j^*-1}$ . Thus, we compute the difference  $\pi_j - \pi_{j-1}$  for  $j = 2, \dots, 2K$  between two adjacent loss values to locate the change-point  $\dot{\mathbf{X}}_{j^*}$  where the index satisfies

$$j^* = \arg \max_{j \in \{2, \dots, 2K\}} [\pi_j - \pi_{j-1}].$$

## 4. EXPERIMENTS

**4.1. Sequences from the multivariate autoregressive model.** In this section, we illustrate the effectiveness of GANs for modeling the stationary time series data from the multivariate autoregressive model as follows:

$$\mathbf{x}_t = \mathbf{x}_{t-1} - 0.01\mathbf{x}_{t-2} - 0.5\mathbf{x}_{t-3} + \mathbf{v}_t,$$

where  $\mathbf{x}_t \in \mathbb{R}^{20}$  and  $\mathbf{v}_t \sim \mathcal{N}(\mathbf{0}, 0.2\mathbf{I}_{20})$  for  $t = 1, \dots, 10500$ . The initial values are set as  $\mathbf{x}_1 = \mathbf{x}_2 = \mathbf{0}$ , and each element of  $\mathbf{x}_3$  is independently sampled from uniform distribution  $U(0, 0.01)$ , the seed is set 3 (Python language, `random.seed(3)`). Considering that the sequences at initial times are nonstationary, we remove the first 500 observations to guarantee the stationarity of the training data and retain the remaining 10000 observations to train GAN. We adopt the general training method of GAN (Algorithm 1 in [Arjovsky et al. \(2017\)](#)) to train the discriminator and generator. The neural network of the discriminator for GAN is composed of nine-linear layers with the LeakyReLU activation function. The structure of the generator contains two convolution layers and five deconvolution layers, in which each layer is also followed by a LeakyReLU activation function. Specifically, the two convolutional layers first transform the Gaussian input noises into latent variables, and then the subsequent deconvolution layers upsample the latent variables to produce fake samples with the same dimension as the real ones. In the training process, we first train the discriminator 100 times to enhance its ability to distinguish the real and generated samples. Then, we train the generator one time once the



---

**Algorithm 1:** Single change-point detection

---

**Input:**  $n$ : sample size;  $K$ : the number of samples in each block;  $t$ : the number of samples to train a stable GAN model;  $L$ : a  $\lceil (n-t)/K \rceil$ -dimensional zero vector used for storing the loss values in the second stage;  $w$ : is the radius of the rolling window,  $\Pi$ : a  $2K$ -dimensional zero vector used for storing the loss values in the third stage;

- 1 **Stage 1: Training a stable discriminator and generator**
  - 2 Sample a  $p$ -dimensional time series sequence  $\mathbf{X} = \{\mathbf{X}_1, \dots, \mathbf{X}_n\}$ ;
  - 3 **Initialize**  $\theta_d, \theta_g \leftarrow$  Gaussian.
  - 4 Use the first  $t$  sample points in  $\mathbf{X}$ ,  $\{\mathbf{X}_1, \dots, \mathbf{X}_t\}$ , to train a discriminator  $d$ ;
  - 5 Sample i.i.d noise  $\{z_1, \dots, z_t\}$  from the standard normal distribution to train a generator  $g$ ;
  - 6 **Stage 2: Select the blocks possibly including the change-point**
  - 7 Divide the remaining samples  $\{\mathbf{X}_{t+1}, \dots, \mathbf{X}_n\}$  into  $\lceil (n-t)/K \rceil$  blocks,  $\mathbf{B}_1, \dots, \mathbf{B}_{\lceil (n-t)/K \rceil}$  with  $\mathbf{B}_i = \{\mathbf{X}_{t+(i-1)K+1}, \dots, \mathbf{X}_{t+iK}\}$ , for each  $i$ ;
  - 8 **for**  $i = 1, \dots, \lceil (n-t)/K \rceil$  **do**
  - 9     Input  $\mathbf{X}_j \in \mathbf{B}_i$  into  $d$ ;
  - 10     Sample  $K$  i.i.d noises  $\{z_1, \dots, z_K\}$  from the the standard normal distribution;
  - 11     Input  $\{z_1, \dots, z_K\}$  into  $g$ ;
  - 12     Compute  $l_i = \frac{1}{K} \sum_{\mathbf{X}_j \in \mathbf{B}_i} d(\mathbf{X}_j) - \frac{1}{K} \sum_{j=1}^K d(g(z_j))$ ;
  - 13     Store  $l_i$  as the  $i$ th element of  $L$ .
  - 14 **end**
  - 15 Compute the distances  $|l_{i+1} - l_i|$  for  $i = 1, \dots, \lceil (n-t)/K \rceil - 1$ , and find the  $\mathbf{B}_k$  and  $\mathbf{B}_{k+1}$  blocks having the largest distance. Then merge  $\mathbf{B}_k$  and  $\mathbf{B}_{k+1}$  into one group  $\mathbf{G} = \{\mathbf{B}_k, \mathbf{B}_{k+1}\} = \{\dot{\mathbf{X}}_1, \dots, \dot{\mathbf{X}}_{2K}\}$ ;
  - 16 **Stage 3: Detect the location of change-point by rolling windows**
  - 17 **for**  $j = 1, \dots, 2K$  **do**
  - 18     Input samples  $\mathbf{H}_j = \{\dot{\mathbf{X}}_{j-w}, \dots, \dot{\mathbf{X}}_j, \dots, \dot{\mathbf{X}}_{j+w}\}$  in  $\mathbf{G}$  into  $d$ ;
  - 19     Sample  $2w+1$  i.i.d noises  $\{z_1, \dots, z_{2w+1}\}$  from the standard normal distribution;
  - 20     Input  $\{z_1, \dots, z_{2w+1}\}$  into  $g$ ;
  - 21     Compute  $\pi_j = \frac{1}{2w+1} \sum_{\dot{\mathbf{X}}_i \in \mathbf{H}_j} d(\dot{\mathbf{X}}_i) - \frac{1}{2w+1} \sum_{i=1}^{2w+1} d(g(z_i))$ ;
  - 22     Store  $\pi_j$  as  $j$ th element of  $\Pi$ .
  - 23 **end**
  - 24 Compute the distances  $\pi_j - \pi_{j-1}$  for  $j = 2, \dots, 2K$ , and record the  $j^*$ th index having the largest distance.
- 

discriminator is trained 25 times. The batch size is set as 64. After training a stable generator, we feed a set of random Gaussian noises into the generator and then obtain 20-dimensional generated samples on the time cross section. Figure 1 compares two sets of generated samples (blue curves: Fake) with the real samples (red curves: Real) at two different times. It can be seen that the synthetic samples generated by GAN show substantially good overlap with the real data. The results reveal that GAN can effectively estimate the distribution of stationary time series data.

**4.2. An example of the Algorithm 1.** We now present an example of single change-point detection for time series so as to illustrate the usefulness of our proposed detection algorithm. Specifically, we generate a sequence of 23-dimensional 10500 time-series samples  $\{\mathbf{y}_k\}_{k=1}^{10500}$




FIGURE 1. GAN’s generated samples of the multivariate autoregressive data on the time cross section.




FIGURE 2. Loss values of 10000 stationary samples  $\{\tilde{\mathbf{y}}_t\}_{t=1}^{10000}$ . The samples located in the interval  $[4800, 5200]$  are as the elements of  $\mathbf{G}$  (in Stage 2).

with a single change-point, where  $\mathbf{y}_k$  is generated by

$$\mathbf{y}_k = 0.6\mathbf{y}_{k-1} - 0.5\mathbf{y}_{k-2} + \epsilon_k,$$




FIGURE 3. Loss values of each samples in  $\mathcal{G}$  corresponding to the interval  $[4800, 5200]$  in Figure 2. The change point detected by our algorithm is accurately the same as the real one.

and  $\epsilon_k$  is a Gaussian vector from  $\mathcal{N}(\mathbf{0}, \mathbf{I}_{23})$  for  $k < 5500$ ,  $\mathcal{N}(\mathbf{2}, \mathbf{I}_{23})$  for  $k \geq 5500$ . The initial values  $\mathbf{y}_1 = \mathbf{0}$  and  $\mathbf{y}_2 = \mathbf{0}$ . We remove the first 500 non-stationary observations to obtain a sequence of stationary time series  $\{\tilde{\mathbf{y}}_t\}_{t=1}^{10000}$ . So the change-point exists at  $t = 5000$  in the new time series. We use our proposed single-point detection algorithm to detect the change point. We utilize the first 2000 samples to train a stable generator and discriminator (in **Stage 1**). Figure 2 shows the loss values of the 10000 stationary samples. Then, following the **stage 2**, we set the sample size in each block as  $K = 200$  and find the blocks possibly including the change-point, i.e., the 24th and 25th blocks (in the time interval  $[4800, 5200]$ ). Next, we merge the 24th and 25th blocks into group  $\mathcal{G}$  shown in Figure 2. Then, we use the rolling window strategy with radius  $w = 6$  (in **Stage 3**) to finally search the change-point on  $\mathcal{G}$ . As shown in Figure 3, the change-point in  $\mathcal{G}$  can be accurately detected.

**4.3. Real data applications.** In previous experiments, we demonstrated the outperformance of GAN in estimating a set of stationary sequences. In this section, we also test the performance of GAN for learning a set of nonstationary financial time series data, the monthly returns of forty-eight Fama French (FF-48) portfolios<sup>1</sup>. The FF-48 portfolios contain the monthly returns of 48 stocks from December 1992 to December 2016. This data is not stationary since the asset returns of the 48 stocks gradually increase over time. We use the same training methods and network configurations in the experiments in section 4.1. We input a set of random Gaussian noises into the trained generator and obtain 48-dimensional generated samples corresponding to different times. Figure 4 shows two sets of generated samples and their corresponding real monthly returns of the 48 stocks in two different months. The results show that the generated samples can not completely recover the real data, but their general tendencies are consistent with the real ones. We can conclude that GAN also can estimate the basic changing tendency of the nonstationary sequences.

<sup>1</sup>[http://mba.tuck.dartmouth.edu/pages/faculty/ken.french/data\\_library.html](http://mba.tuck.dartmouth.edu/pages/faculty/ken.french/data_library.html)




FIGURE 4. Generated samples of the monthly returns of the 48 stocks.

## 5. PROOF OF THEOREM 3.1

In this section, we prove Theorem 3.1, based on a decomposition for  $d_{\mathcal{H}}(\pi, \hat{\mathbf{g}}_{\#}\nu)$  and the estimates for each term in the decomposition.

**5.1. Auxiliary lemmas.** In order to prove Theorem 3.1, we need the following auxiliary lemmas. The first lemma is an error decomposition for  $d_{\mathcal{H}}(\pi, \hat{\mathbf{g}}_{\#}\nu)$ , while the others are estimates for each term in this decomposition.

**5.1.1. Error decomposition of  $d_{\mathcal{H}}(\pi, \hat{\mathbf{g}}_{\#}\nu)$ .**

**Lemma 5.1** (Error Decomposition). *Suppose  $\mathbf{g}_{\#}\nu$  is supported on  $[-a_n, a_n]^p$  for all  $\mathbf{g} \in \mathcal{G}$  and assume  $\mathcal{D}$  is in symmetric class. Let  $\hat{\mathbf{g}}$  be defined in (2.2). Suppose Assumption 2.2 is satisfied, for any function class  $\mathcal{H}$  defined on  $\mathbb{R}^p$ ,  $h \in \mathcal{H}^b$ ,  $b = r + q$ , there exists  $c > 0$  such that for any*

$$\mathcal{D} = \{d \in \mathcal{NN}(N_{\mathcal{D}}, L_{\mathcal{D}}, K_{\mathcal{D}}) : d(\mathbb{R}^p) \subseteq [-B, B]\}, B < \infty,$$

with  $N_{\mathcal{D}} \geq c(K_{\mathcal{D}}/\mathfrak{b}^{-\gamma} K_{\mathcal{D}})^{(2p+b)/(2p+2)}$ ,  $L_{\mathcal{D}} \geq 4\gamma + 4$ ,  $\gamma := \lceil \mathfrak{b}^{-2}(p+r) \rceil$ , one has

$$\begin{aligned} d_{\mathcal{H}}(\pi, \hat{\mathbf{g}}_{\#}\nu) &\leq 2(2a_n)^b (K_{\mathcal{D}}/\mathfrak{b}^{-\gamma} K_{\mathcal{D}})^{-\frac{b}{p+1}} + 2(1+B)pe^{-a_n+v^2/2} \\ &\quad + \mathfrak{P}_{d \in \mathcal{D}} \left\{ \mathbb{E}_{\pi}[d] - \frac{1}{n} \sum_{i=1}^n d(\mathbf{X}_i) \right\} + \mathfrak{f}_{\mathbf{g} \in \mathcal{G}} \mathfrak{P}_{d \in \mathcal{D}} \left\{ \frac{1}{n} \sum_{i=1}^n d(\mathbf{X}_i) - \mathbb{E}_{\mathbf{g}_{\#}\nu}[d] \right\} \\ &\quad + 2 \mathfrak{P}_{d \circ \mathbf{g} \in \mathcal{D} \circ \mathcal{G}} \left\{ \mathbb{E}_{\hat{\nu}_m}[d \circ \mathbf{g}] - \mathbb{E}_{\nu}[d \circ \mathbf{g}] \right\}. \end{aligned}$$

where  $\hat{\nu}_m := \frac{1}{m} \sum_{i=1}^m \delta_{\mathbf{z}_i}$ .

*Proof.* For any  $\varepsilon > 0$ , it is easy to know that there exist an  $h_{\varepsilon} \in \mathcal{H}$  and a  $d_{\varepsilon} \in \mathcal{D}$  such that

$$\mathfrak{P}_{h \in \mathcal{H}} \left\{ \mathbb{E}_{\pi}[h] - \mathbb{E}_{\hat{\mathbf{g}}_{\#}\nu}[h] \right\} \leq \mathbb{E}_{\pi}[h_{\varepsilon}] - \mathbb{E}_{\hat{\mathbf{g}}_{\#}\nu}[h_{\varepsilon}] + \varepsilon,$$

and

$$\|h_{\varepsilon} - d_{\varepsilon}\|_{\infty} \leq \mathfrak{f}_{d \in \mathcal{D}} \|h_{\varepsilon} - d\|_{\infty} + \varepsilon.$$

Then, we have

$$\begin{aligned}
& \mathbb{P}_{h \in \mathcal{H}} \{ \mathbb{E}_\pi[h] - \mathbb{E}_{\hat{g}_{\#\nu}}[h] \} \\
& \leq \mathbb{E}_\pi[h_\varepsilon] - \mathbb{E}_{\hat{g}_{\#\nu}}[h_\varepsilon] + \varepsilon \\
& = \mathbb{E}_\pi[h_\varepsilon] - \mathbb{E}_\pi[d_\varepsilon] + \mathbb{E}_\pi[d_\varepsilon] - \mathbb{E}_{\hat{g}_{\#\nu}}[d_\varepsilon] + \mathbb{E}_{\hat{g}_{\#\nu}}[d_\varepsilon] - \mathbb{E}_{\hat{g}_{\#\nu}}[h_\varepsilon] + \varepsilon \\
(5.1) \quad & = \{ \mathbb{E}_\pi[h_\varepsilon - d_\varepsilon] + \mathbb{E}_{\hat{g}_{\#\nu}}[d_\varepsilon - h_\varepsilon] \} + \{ \mathbb{E}_\pi[d_\varepsilon] - \mathbb{E}_{\hat{g}_{\#\nu}}[d_\varepsilon] \} + \varepsilon.
\end{aligned}$$

Step 1: We first deal with the first terms of (5.1). Let  $A_n = [-a_n, a_n]^p$ , it is easy to see that

$$\begin{aligned}
(5.2) \quad & \mathbb{E}_\pi[ h_\varepsilon - d_\varepsilon ] + \mathbb{E}_{\hat{g}_{\#\nu}}[d_\varepsilon - h_\varepsilon] \\
& = \mathbb{E}_\pi[ h_\varepsilon - d_\varepsilon ] 1_{A_n^c} + \mathbb{E}_\pi[ h_\varepsilon - d_\varepsilon ] 1_{A_n} + \mathbb{E}_{\hat{g}_{\#\nu}}[ d_\varepsilon - h_\varepsilon ].
\end{aligned}$$

For the first term of (5.2), the boundedness of  $h_\varepsilon, d_\varepsilon$  and sub-Gaussian assumption yield

$$\begin{aligned}
(5.3) \quad & \mathbb{E}_\pi[ h_\varepsilon - d_\varepsilon ] 1_{A_n^c} \leq \|h_\varepsilon - d_\varepsilon\|_\infty \mathbb{E}_\pi[ \mathbb{1}_{A_n^c} ] \\
& \leq (1 + \|d\|_\infty + \varepsilon) \mathbb{P}_{\mathbf{x} \sim \pi} (\|\mathbf{x}\|_\infty \geq a_n) \\
& \leq (1 + \|d\|_\infty + \varepsilon) \sum_{i=1}^p \mathbb{P}_{\mathbf{x} \sim \pi} (|x_i| \geq a_n) \\
& \leq 2(1 + B + \varepsilon) p e^{-a_n + v^2/2}.
\end{aligned}$$

Since  $\mathbf{g}(\mathbb{R}) \subset A_n$ , one can get the estimate of last two terms of (5.2)

$$\mathbb{E}_\pi[ h_\varepsilon - d_\varepsilon ] 1_{A_n} + \mathbb{E}_{\hat{g}_{\#\nu}}[ d_\varepsilon - h_\varepsilon ] \leq 2 \mathbb{P}_{h \in \mathcal{H}(A_n)} \mathbb{f}_{d \in \mathcal{D}(A_n)} \|h - d\|_\infty + 2\varepsilon.$$

Jiao et al. (2022, Theorem 3.2) guarantees that, for any  $\bar{h} \in \mathcal{H}^b(\bar{\mathfrak{b}}, 1]^p$ ,  $b = r + q$ , there exists  $c > 0$  such that for any  $\bar{d} \in \mathcal{NN}(N, L, K)$ ,  $N \geq c(K/\bar{\mathfrak{b}}^\gamma K)^{(2p+b)/(2p+2)}$  and  $L \geq 4\gamma + 2$ ,  $\gamma := \lceil \bar{\mathfrak{b}}^{-2}(p+r) \rceil$ ,

$$\mathbb{P}_{\bar{h} \in \mathcal{H}([0,1]^p)} \mathbb{f}_{\bar{d} \in \mathcal{NN}(N,L,K)} \|\bar{h} - \bar{d}\|_\infty \leq (K/\bar{\mathfrak{b}}^\gamma K)^{-\frac{b}{p+1}}.$$

Let  $h(x)1_{A_n} = (2a_n)^b \bar{h}(\frac{x}{2a_n} + \frac{1}{2})$ ,  $d_0(x)1_{A_n} = (2a_n)^b \bar{d}(\frac{x}{2a_n} + \frac{1}{2})$ , and

$$d(x)1_{A_n} = (d_0(x) \wedge B) \vee (-B) = \sigma(d_0(x) + B) - \sigma(d_0(x) - B) - B.$$

Then, we have

$$\begin{aligned}
\mathbb{P}_{h \in \mathcal{H}(A_n)} \mathbb{f}_{d \in \mathcal{D}(A_n)} \|h - d\|_\infty & = \mathbb{P}_{\bar{h} \in \mathcal{H}([0,1]^p)} \mathbb{f}_{d \in \mathcal{D}(A_n)} \left\| (2a_n)^b \bar{h}\left(\frac{x}{2a_n} + \frac{1}{2}\right) - d(x) \right\|_\infty \\
& \leq (2a_n)^b \mathbb{P}_{\bar{h} \in \mathcal{H}([0,1]^p)} \mathbb{f}_{\bar{d} \in \mathcal{D}([0,1]^p)} \|\bar{h} - \bar{d}\|_\infty \leq (2a_n)^b (K_{\mathcal{D}}/\bar{\mathfrak{b}}^\gamma K_{\mathcal{D}})^{-\frac{b}{p+1}},
\end{aligned}$$

where  $d \in \mathcal{NN}(N_{\mathcal{D}}, L_{\mathcal{D}}, K_{\mathcal{D}})$ , with  $N_{\mathcal{D}} \geq c(K_{\mathcal{D}}/\bar{\mathfrak{b}}^\gamma K_{\mathcal{D}})^{(2p+b)/(2p+2)}$  and  $L_{\mathcal{D}} \geq 4\gamma + 4$ ,  $\gamma := \lceil \bar{\mathfrak{b}}^{-2}(p+r) \rceil$ . Thus, we obtain

$$\mathbb{E}_\pi[h_\varepsilon - d_\varepsilon] + \mathbb{E}_{\hat{g}_{\#\nu}}[d_\varepsilon - h_\varepsilon] \leq 2(2a_n)^b (K_{\mathcal{D}}/\bar{\mathfrak{b}}^\gamma K_{\mathcal{D}})^{-\frac{b}{p+1}} + 2(1 + B + \varepsilon) p e^{-a_n + v^2/2} + 2\varepsilon.$$

Step 2: For the rest terms of (5.1), that is  $\mathbb{E}_\pi[d_\varepsilon] - \mathbb{E}_{\hat{g}_{\#\nu}}[d_\varepsilon]$ , one considers the empirical process and gets

$$\begin{aligned}
\mathbb{E}_\pi[d_\varepsilon] - \mathbb{E}_{\hat{g}_{\#\nu}}[d_\varepsilon] &= \mathbb{E}_\pi[d_\varepsilon] - \frac{1}{n} \sum_{i=1}^n d_\varepsilon(\mathbf{X}_i) + \frac{1}{n} \sum_{i=1}^n d_\varepsilon(\mathbf{X}_i) - \frac{1}{m} \sum_{i=1}^m d_\varepsilon(\hat{\mathbf{g}}(\mathbf{z}_i)) \\
&\quad + \frac{1}{m} \sum_{i=1}^m d_\varepsilon(\hat{\mathbf{g}}(\mathbf{z}_i)) - \mathbb{E}_\nu[d_\varepsilon(\hat{\mathbf{g}}(\mathbf{z}))] \\
&\leq \mathbf{P}_{d \in \mathcal{D}} \left\{ \mathbb{E}_\pi[d] - \frac{1}{n} \sum_{i=1}^n d(\mathbf{X}_i) \right\} + \mathbf{f}_{\mathbf{g} \in \mathcal{G}} \mathbf{P}_{d \in \mathcal{D}} \left\{ \frac{1}{n} \sum_{i=1}^n d(\mathbf{X}_i) - \frac{1}{m} \sum_{i=1}^m d(\mathbf{g}(\mathbf{z}_i)) \right\} \\
&\quad + \mathbf{P}_{d \circ \mathbf{g} \in \mathcal{D} \circ \mathcal{G}} \left\{ \mathbb{E}_{\hat{\nu}_m}[d \circ \mathbf{g}] - \mathbb{E}_\nu[d \circ \mathbf{g}] \right\} \\
&\leq \mathbf{P}_{d \in \mathcal{D}} \left\{ \mathbb{E}_\pi[d] - \frac{1}{n} \sum_{i=1}^n d(\mathbf{X}_i) \right\} + \mathbf{f}_{\mathbf{g} \in \mathcal{G}} \mathbf{P}_{d \in \mathcal{D}} \left\{ \frac{1}{n} \sum_{i=1}^n d(\mathbf{X}_i) - \mathbb{E}_{\mathbf{g}_{\#\nu}}[d] \right\} \\
&\quad + \mathbf{f}_{\mathbf{g} \in \mathcal{G}} \mathbf{P}_{d \in \mathcal{D}} \left\{ \mathbb{E}_{\mathbf{g}_{\#\nu}}[d] - \frac{1}{m} \sum_{i=1}^m d(\mathbf{g}(\mathbf{z}_i)) \right\} + \mathbf{P}_{d \circ \mathbf{g} \in \mathcal{D} \circ \mathcal{G}} \left\{ \mathbb{E}_{\hat{\nu}_m}[d \circ \mathbf{g}] - \mathbb{E}_\nu[d \circ \mathbf{g}] \right\},
\end{aligned}$$

where the first inequality follows the definition of  $\hat{\mathbf{g}}$  and  $\hat{\nu}_m := \frac{1}{m} \sum_{i=1}^m \delta_{\mathbf{z}_i}$  is the empirical measure. Combining the inequality above with (5.1) and a straight calculation implies

$$\begin{aligned}
d_{\mathcal{H}}(\pi, \hat{\mathbf{g}}_{\#\nu}) &\leq 2(2a_n)^b (K_{\mathcal{D}}/\mathfrak{g}^\gamma K_{\mathcal{D}})^{-\frac{b}{p+1}} + 2(1+B+\varepsilon)pe^{-a_n+v^2/2} + 3\varepsilon \\
&\quad + \mathbf{P}_{d \in \mathcal{D}} \left\{ \mathbb{E}_\pi[d] - \frac{1}{n} \sum_{i=1}^n d(\mathbf{X}_i) \right\} + \mathbf{f}_{\mathbf{g} \in \mathcal{G}} \mathbf{P}_{d \in \mathcal{D}} \left\{ \frac{1}{n} \sum_{i=1}^n d(\mathbf{X}_i) - \mathbb{E}_{\mathbf{g}_{\#\nu}}[d] \right\} \\
&\quad + 2 \mathbf{P}_{d \circ \mathbf{g} \in \mathcal{D} \circ \mathcal{G}} \left\{ \mathbb{E}_{\hat{\nu}_m}[d \circ \mathbf{g}] - \mathbb{E}_\nu[d \circ \mathbf{g}] \right\}.
\end{aligned}$$

Let  $\varepsilon \rightarrow 0$  and the proof is complete.  $\square$

**5.1.2. Statistical error of the GAN estimator.** Recall that  $\mathbf{X}_i := \mathbf{G}(\dots, \varepsilon_i)$ , which means for any  $i, j \in \mathbb{Z}$ ,  $\mathbf{X}_i, \mathbf{X}_j$  is not independent. We denote the  $s$ -dependence form of  $\mathbf{X}_i$  by  $\tilde{\mathbf{X}}_i = \mathbb{E}[\mathbf{X}_i | \mathcal{F}_{i-s}^i]$  and  $\mathcal{F}_{i-s}^i = \sigma(\varepsilon_{i-s}, \dots, \varepsilon_i)$ . Thus  $\tilde{\mathbf{X}}_i$  only depends on  $\{\varepsilon_{i-s}, \dots, \varepsilon_i\}$ . We denote the distribution of  $\tilde{\mathbf{X}}_i$  as  $\tilde{\pi}$ . In general, we choose  $n$  large enough that for some  $s \in \mathbb{N}_+$ , such that  $n/2s \in \mathbb{N}_+$ .

Statistical error of the GAN estimator,  $\mathbf{P}_{d \in \mathcal{D}} \left\{ \mathbb{E}_\pi[d] - \frac{1}{n} \sum_{i=1}^n d(\mathbf{X}_i) \right\}$ , can be controlled by the concentration for suprema of the GAN estimator as in following Lemma.

**Lemma 5.2** (Concentration for Supremum of GAN Empirical Processes). *Suppose Assumption 2.1 is satisfied and*

$$\mathcal{D} = \{d \in \mathcal{NN}(N_{\mathcal{D}}, L_{\mathcal{D}}, K_{\mathcal{D}}) : d(\mathbb{R}^p) \subseteq [-B, B]\}, B < \infty,$$

then with probability at least  $1 - 2e^{-t_1} - e^{-C_2s/2\beta}$

$$\begin{aligned}
&\mathbf{P}_{d \in \mathcal{D}} \left| \frac{1}{n} \sum_{i=1}^n (d(\mathbf{X}_i) - \mathbb{E}_\pi[d(\mathbf{X}_i)]) \right| \\
&\leq C_1^{1/\beta} K_{\mathcal{D}} e^{-C_2(s+1)/\beta} / (1 - e^{-C_2/\beta}) + \frac{1}{1 - e^{-C_2/\beta}} C_1^{1/\beta} K_{\mathcal{D}} e^{-C_2s/2\beta - C_2/\beta} \\
&\quad + 4B\sqrt{st_1/n} + 4\sqrt{\frac{2C_3}{n}} K_{\mathcal{D}} \cdot \sqrt{L_{\mathcal{D}} + 1 + \mathfrak{g}^{-p}}.
\end{aligned}$$

In order to prove Lemma 5.2, we need to prove following lemma first,

**Lemma 5.3.** *Suppose Assumption 2.1 is satisfied and*

$$\mathcal{D} = \{d \in \mathcal{NN}(N_{\mathcal{D}}, L_{\mathcal{D}}, K_{\mathcal{D}}) : d(\mathbb{R}^p) \subseteq [-B, B]\}, B < \infty.$$

(i) We have

$$\mathbb{P}_{d \in \mathcal{D}} \left| \sum_{1 \leq i \leq n} \left( \mathbb{E}_{\pi} [d(\mathbf{X}_i)] - \mathbb{E}_{\tilde{\pi}} [d(\tilde{\mathbf{X}}_i)] \right) \right| \leq n C_1^{1/\beta} K_{\mathcal{D}} e^{-C_2(s+1)/\beta} / (1 - e^{-C_2/\beta}).$$

(ii) With probability at least  $1 - e^{-C_2s/2\beta}$ , we have

$$\mathbb{P}_{d \in \mathcal{D}} \left\{ \sum_{i=1}^n d(\mathbf{X}_i) - \sum_{i=1}^n d(\tilde{\mathbf{X}}_i) \right\} \leq \frac{1}{1 - e^{-C_2/\beta}} n C_1^{1/\beta} K_{\mathcal{D}} e^{-C_2s/2\beta - C_2/\beta}.$$

*Proof.* (i) A straight calculation yields that

$$\begin{aligned} d(\mathbf{X}_i) - d(\tilde{\mathbf{X}}_i) &= d(\mathbb{E}[\mathbf{X}_i | \mathcal{F}_{-\infty}^i]) - d(\mathbb{E}[\mathbf{X}_i | \mathcal{F}_{i-s}^i]) \\ &= \sum_{j=s-i+1}^{\infty} (d(\mathbb{E}[\mathbf{X}_i | \mathcal{F}_{-j}^i]) - d(\mathbb{E}[\mathbf{X}_i | \mathcal{F}_{-j+1}^i])) \\ &= \sum_{j=s-i+1}^{\infty} (d(\mathbb{E}[\mathbf{G}(\boldsymbol{\xi}_{-j}, \boldsymbol{\varepsilon}_{-j+1}, \dots, \boldsymbol{\varepsilon}_i) | \mathcal{F}_{-j}^i]) - d(\mathbb{E}[\mathbf{G}(\boldsymbol{\xi}'_{-j}, \boldsymbol{\varepsilon}_{-j+1}, \dots, \boldsymbol{\varepsilon}_i) | \mathcal{F}_{-j}^i])). \end{aligned}$$

Since  $\mathcal{D}$  is set of Lipschitz function with Lipschitz constant  $K_{\mathcal{D}}$ , combining equality above with condition (2.6), we can get

$$\begin{aligned} (5.4) \quad & \mathbb{E}_{\mathbb{P}} \mathbb{P}_{d \in \mathcal{D}} \left| \sum_{i=1}^n (d(\mathbf{X}_i) - d(\tilde{\mathbf{X}}_i)) \right| \\ & \leq \mathbb{E}_{\mathbb{P}} \mathbb{P}_{d \in \mathcal{D}} \sum_{i=1}^n \sum_{j=s-i+1}^{\infty} |d(\mathbb{E}[\mathbf{G}(\boldsymbol{\xi}_{-j}, \boldsymbol{\varepsilon}_{-j+1}, \dots, \boldsymbol{\varepsilon}_i) | \mathcal{F}_{-j}^i]) - d(\mathbb{E}[\mathbf{G}(\boldsymbol{\xi}'_{-j}, \boldsymbol{\varepsilon}_{-j+1}, \dots, \boldsymbol{\varepsilon}_i) | \mathcal{F}_{-j}^i])| \\ & \leq K_{\mathcal{D}} \mathbb{E} \left\{ \sum_{i=1}^n \sum_{j=s-i+1}^{\infty} (\mathbb{E}[\|\mathbf{G}(\boldsymbol{\xi}_{-j}, \boldsymbol{\varepsilon}_{-j+1}, \dots, \boldsymbol{\varepsilon}_i) - \mathbf{G}(\boldsymbol{\xi}'_{-j}, \boldsymbol{\varepsilon}_{-j+1}, \dots, \boldsymbol{\varepsilon}_i)\| | \mathcal{F}_{-j}^i]) \right\} \\ & = K_{\mathcal{D}} \sum_{i=1}^n \sum_{j=s-i+1}^{\infty} \mathbb{E} \|\mathbf{G}(\boldsymbol{\xi}_{-j}, \boldsymbol{\varepsilon}_{-j+1}, \dots, \boldsymbol{\varepsilon}_i) - \mathbf{G}(\boldsymbol{\xi}'_{-j}, \boldsymbol{\varepsilon}_{-j+1}, \dots, \boldsymbol{\varepsilon}_i)\| \\ & \leq K_{\mathcal{D}} \sum_{i=1}^n \sum_{j=s-i+1}^{\infty} C_1^{1/\beta} e^{-C_2(i+j)/\beta} = n C_1^{1/\beta} K_{\mathcal{D}} e^{-C_2(s+1)/\beta} / (1 - e^{-C_2/\beta}). \end{aligned}$$

Since

$$\mathbb{P}_{d \in \mathcal{D}} \left| \sum_{i=1}^n (\mathbb{E}_{\pi} [d(\mathbf{X}_i)] - \mathbb{E}_{\tilde{\pi}} [d(\tilde{\mathbf{X}}_i)]) \right| \leq \mathbb{E}_{\mathbb{P}} \mathbb{P}_{d \in \mathcal{D}} \left| \sum_{i=1}^n (d(\mathbf{X}_i) - d(\tilde{\mathbf{X}}_i)) \right|,$$

then the result in (5.4) yields that

$$\mathbb{P}_{d \in \mathcal{D}} \left| \sum_{i=1}^n (\mathbb{E}_{\pi} [d(\mathbf{X}_i)] - \mathbb{E}_{\tilde{\pi}} [d(\tilde{\mathbf{X}}_i)]) \right| \leq n C_1^{1/\beta} K_{\mathcal{D}} e^{-C_2(s+1)/\beta} / (1 - e^{-C_2/\beta}).$$

(ii) The Chebyshev inequality and the result in (i) imply

$$\begin{aligned} \mathbb{P}\left(\mathbb{P}_{d \in \mathcal{D}} \left\{ \sum_{i=1}^n d(\mathbf{X}_i) - \sum_{i=1}^n d(\tilde{\mathbf{X}}_i) \right\} \geq x\right) &\leq \frac{1}{x} \mathbb{E}_{\mathbb{P}_{d \in \mathcal{D}}} \left\{ \sum_{i=1}^n (d(\mathbf{X}_i) - d(\tilde{\mathbf{X}}_i)) \right\} \\ &\leq \frac{1}{x(1 - e^{-C_2/\beta})} n C_1^{1/\beta} K_{\mathcal{D}} e^{-C_2(s+1)/\beta}. \end{aligned}$$

We can obtain with probability at least  $1 - e^{-C_2 s/2\beta}$ ,

$$\mathbb{P}_{d \in \mathcal{D}} \left\{ \sum_{i=1}^n d(\mathbf{X}_i) - \sum_{i=1}^n d(\tilde{\mathbf{X}}_i) \right\} \leq \frac{1}{1 - e^{-C_2/\beta}} n C_1^{1/\beta} K_{\mathcal{D}} e^{-C_2 s/2\beta - C_2/\beta}.$$

□

Given a real-valued function class  $\mathcal{H}$  and some set of data points  $\mathbf{x}_1, \dots, \mathbf{x}_n \in \mathbb{R}^n$ , we define the (empirical) Rademacher complexity  $\hat{\mathcal{R}}_n(\mathcal{H})$  as

$$\hat{\mathcal{R}}_n(\mathcal{H}) := \mathbb{E}_{\boldsymbol{\eta}} \left[ \mathbb{P}_{h \in \mathcal{H}} \left[ \frac{1}{n} \sum_{i=1}^n \eta_i h(\mathbf{x}_i) \right] \right],$$

where  $\boldsymbol{\eta} = (\eta_1, \dots, \eta_n)$  is a vector uniformly distributed in  $\{-1, +1\}^n$ .

Theorem 2 in [Golowich et al. \(2018\)](#) shows that,

**Lemma 5.4** (Rademacher Complexity of Neural Networks). *Let  $\mathcal{H}_L$  be the class of real-valued networks of depth  $L$ , where  $\|W_j\|_{1,\infty} \leq M(j)$  for all  $j \in \{1, \dots, L\}$ , and  $\sigma$  is a 1-Lipschitz activation function with  $\sigma(0) = 0$ , applied element-wise. Then*

$$\hat{\mathcal{R}}_n(\mathcal{H}_L) \leq \frac{2}{n} \prod_{j=1}^L M(j) \cdot \sqrt{L + 1 + \frac{1}{p}} \cdot \sqrt{\mathbb{E}_{\mathbf{x} \in \{1, \dots, p\}} \sum_{i=1}^n \mathbf{x}_{i,j}^2},$$

where  $\mathbf{x}_{i,j}$  is the  $j$ -th coordinate of the vector  $\mathbf{x}_i$ .

*Proof of Lemma 5.2.* For  $j \in \{1, 2, \dots, n/(2s)\}$ , let

$$\begin{aligned} I_j &= \{i : 2s(j-1) + 1 \leq i \leq (2j-1)s\}, \\ J_j &= \{i : (2j-1)s + 1 \leq i \leq 2js\}, \end{aligned}$$

thus we decompose the interval  $\llbracket 1, n \rrbracket$  as a union of  $n/s$  blocks. Let the block sums be

$$\mathbf{Y}_{I_j} = \sum_{i \in I_j} d(\tilde{\mathbf{X}}_i), \quad \mathbf{Y}_{J_j} = \sum_{i \in J_j} d(\tilde{\mathbf{X}}_i).$$



By Lemma 5.3, it is easy to see that, with probability at least  $1 - e^{-C_2s/2\beta}$

$$\begin{aligned}
& \mathbb{P}_{d \in \mathcal{D}} \left| \frac{1}{n} \sum_{i=1}^n (d(\mathbf{X}_i) - \mathbb{E}_\pi [d(\mathbf{X}_i)]) \right| \\
& \leq \mathbb{P}_{d \in \mathcal{D}} \left| \frac{1}{n} \sum_{i=1}^n (d(\mathbf{X}_i) - \mathbb{E}_\pi [d(\mathbf{X}_i)]) - \frac{1}{n} \sum_{i=1}^n (d(\tilde{\mathbf{X}}_i) - \mathbb{E}_{\tilde{\pi}} [d(\tilde{\mathbf{X}}_i)]) \right| \\
& \quad + \mathbb{P}_{d \in \mathcal{D}} \left| \frac{1}{n} \sum_{i=1}^n (d(\tilde{\mathbf{X}}_i) - \mathbb{E}_{\tilde{\pi}} [d(\tilde{\mathbf{X}}_i)]) \right| \\
& \leq C_1^{1/\beta} K_{\mathcal{D}} e^{-C_2(s+1)/\beta} / (1 - e^{-C_2/\beta}) + \frac{1}{1 - e^{-C_2/\beta}} C_1^{1/\beta} K_{\mathcal{D}} e^{-C_2s/2\beta - C_2/\beta} \\
& \quad + \mathbb{P}_{d \in \mathcal{D}} \left| \frac{1}{n} \sum_{j=1}^{n/2s} (\mathbf{Y}_{I,j} - \mathbb{E}_{\tilde{\pi}} [\mathbf{Y}_{I,j}]) \right| + \mathbb{P}_{d \in \mathcal{D}} \left| \frac{1}{n} \sum_{j=1}^{n/2s} (\mathbf{Y}_{J,j} - \mathbb{E}_{\tilde{\pi}} [\mathbf{Y}_{J,j}]) \right|.
\end{aligned}$$

Denote  $Z_I := \mathbb{P}_{d \in \mathcal{D}} \left| \frac{1}{n} \sum_{j=1}^{n/2s} (\mathbf{Y}_{I,j} - \mathbb{E}_{\tilde{\pi}} [\mathbf{Y}_{I,j}]) \right|$ . Since  $\mathbb{E}(\mathbf{Y}_{I,j}) - \mathbb{E}_{\tilde{\pi}} [\mathbf{Y}_{I,j}] = 0$  and

$$|\mathbf{Y}_{I,j} - \mathbb{E}_{\tilde{\pi}} [\mathbf{Y}_{I,j}]| \leq \sum_{i=1}^s |d(\tilde{\mathbf{X}}_i) - \mathbb{E}_{\tilde{\pi}} [d(\tilde{\mathbf{X}}_i)]| \leq \sum_{i=1}^s (|d(\tilde{\mathbf{X}}_i)| + \mathbb{E}_{\tilde{\pi}} |d(\tilde{\mathbf{X}}_i)|) \leq 2sB,$$

for all  $1 \leq j \leq n/2s$ , by Massart's concentration inequality (cf. Massart (2000, Theorem 9)) for suprema of the bounded empirical processes, we have for all  $t_1 > 0$

$$\mathbb{P} \left\{ Z_I \geq \mathbb{E} Z_I + 2B \sqrt{\frac{st_1}{n}} \right\} \leq e^{-t_1}.$$

For the expectation, we consider the Rademacher random variables  $\{\eta_i^j\}_{i=1, \dots, s, 1 \leq j \leq n/2s}$ , that is  $\eta_i^j$  equals to 1 and  $-1$  with propability  $1/2$ . Then by using symmetrization theorem for empirical processes (see Lemma 2.3.1 in van der Vaart and Wellner, 1996) and Lemma 5.4, we can get

$$\begin{aligned}
\mathbb{E} Z_I &= \mathbb{E} \mathbb{P}_{d \in \mathcal{D}} \left| \frac{1}{n} \sum_{j=1}^{n/2s} (\mathbf{Y}_{I,j} - \mathbb{E}_{\tilde{\pi}} [\mathbf{Y}_{I,j}]) \right| \\
&= \mathbb{E} \mathbb{P}_{d \in \mathcal{D}} \left| \frac{1}{n} \sum_{j=1}^{n/2s} \left( \sum_{i \in I_j} d(\tilde{\mathbf{X}}_i) - \mathbb{E}_{\tilde{\pi}} \left[ \sum_{i \in I_j} d(\tilde{\mathbf{X}}_i) \right] \right) \right| \\
&= \mathbb{E} \mathbb{P}_{d \in \mathcal{D}} \left| \frac{1}{n} \sum_{j=1}^{n/2s} \sum_{i \in I_j} d(\tilde{\mathbf{X}}_i) - \mathbb{E}_{\tilde{\pi}} [d] \right| \\
&\leq \frac{2}{n} \mathbb{E} \mathbb{P}_{d \in \mathcal{D}} \left| \sum_{j=1}^{n/2s} \sum_{i \in I_j} \eta_i^j d(\tilde{\mathbf{X}}_i) \right| \\
&\leq \frac{4}{n} K_{\mathcal{D}} \cdot \sqrt{L_{\mathcal{D}} + 1 + \mathfrak{g}(p)} \cdot \mathbb{E} \sqrt{\sum_{k \in \{1, \dots, p\}} \sum_{j=1}^{n/2s} \sum_{i \in I_j} \tilde{\mathbf{X}}_{i,k}^2},
\end{aligned}$$

where  $\tilde{\mathbf{X}}_{i,k}$  denote the  $k$  th coordinate of  $\tilde{\mathbf{X}}_i$ .

Assumption 2.2 implies that  $\pi$  has second order moment bound and thus

$$\mathbb{E}_{\mathbf{m}} \sum_{k \in \{1, \dots, p\}} \tilde{\mathbf{X}}_{i,k}^2 \leq \mathbb{E} \sum_{k=1}^p \tilde{\mathbf{X}}_{i,k}^2 = \sum_{k=1}^p \mathbb{E} [\mathbb{E}(\mathbf{X}_{i,k} | \mathcal{F}_{i-s}^i)^2] \leq \sum_{k=1}^p \mathbb{E} \mathbf{X}_{i,k}^2 \leq C_3.$$

Jensen's inequality implies

$$\begin{aligned} \mathbb{E} \sqrt{\sum_{k \in \{1, \dots, p\}} \sum_{r=1}^{n/2s} \tilde{\mathbf{X}}_{i,k}^2} &\leq \left\{ \mathbb{E}_{\mathbf{m}} \sum_{r=1}^{n/2s} \sum_{i \in I_j} \tilde{\mathbf{X}}_{i,k}^2 \right\}^{1/2} \\ &\leq \left\{ \sum_{r=1}^{n/2s} \sum_{i \in I_j} \mathbb{E}_{\mathbf{m}} \tilde{\mathbf{X}}_{i,k}^2 \right\}^{1/2} \leq \sqrt{C_3 n/2}. \end{aligned}$$

Hence, it is easy to know that,

$$\mathbb{P} \left\{ Z_I \geq 4 \sqrt{\frac{C_3}{2n}} K_{\mathcal{D}} \cdot \sqrt{L_{\mathcal{D}} + 1 + \lfloor p \rfloor} + 2B \sqrt{\frac{st_1}{n}} \right\} \leq e^{-t_1}.$$

Let  $Z_J := \{\mathbf{Y}_{J,j}\}_{1 \leq j \leq n/2s}$ , one can get similarly

$$\mathbb{P} \left\{ Z_J \geq 4 \sqrt{\frac{C_3}{2n}} K_{\mathcal{D}} \cdot \sqrt{L_{\mathcal{D}} + 1 + \lfloor p \rfloor} + 2B \sqrt{\frac{st_1}{n}} \right\} \leq e^{-t_1}.$$

Using above result, it is easy to know that, with probability at least  $1 - 2e^{-t_1} - e^{-C_2 s/2\beta}$ ,

$$\begin{aligned} &\mathbb{P}_{d \in \mathcal{D}} \left| \frac{1}{n} \sum_{i=1}^n (d(\mathbf{X}_i) - \mathbb{E}_{\pi} [d(\mathbf{X}_i)]) \right| \\ &\leq C_1^{1/\beta} K_{\mathcal{D}} e^{-C_2(s+1)/\beta} / (1 - e^{-C_2/\beta}) + \frac{1}{1 - e^{-C_2/\beta}} C_1^{1/\beta} K_{\mathcal{D}} e^{-C_2 s/2\beta - C_2/\beta} \\ &\quad + 4B \sqrt{st_1/n} + 4 \sqrt{\frac{2C_3}{n}} K_{\mathcal{D}} \cdot \sqrt{L_{\mathcal{D}} + 1 + \lfloor p \rfloor}. \end{aligned}$$

□

### 5.1.3. Generator approximation error.

**Lemma 5.5.** Suppose that  $\mathcal{G} = \mathcal{NN}(N_{\mathcal{G}}, L_{\mathcal{G}})$  satisfies  $N_{\mathcal{G}} \geq 7p + 1$ ,  $L_{\mathcal{G}} \geq 2$  and

$$n \leq \frac{N_{\mathcal{G}} - p - 1}{2} \left\lfloor \frac{N_{\mathcal{G}} - p - 1}{6p} \right\rfloor \left\lfloor \frac{L_{\mathcal{G}}}{2} \right\rfloor + 2.$$

For any  $\mathcal{D} \subset \{d \in \mathcal{NN}(N_{\mathcal{D}}, L_{\mathcal{D}}, K_{\mathcal{D}}) : d(\mathbb{R}^p) \subseteq [-B, B]\} \cap \mathbb{R}^p$ ,  $B, C < \infty$ , we have

$$\mathbb{E}_{\mathbf{g} \in \mathcal{G}} \mathbb{P}_{d \in \mathcal{D}} \left\{ \frac{1}{n} \sum_{i=1}^n d(\mathbf{X}_i) - \mathbb{E}_{\mathbf{g}_{\# \nu}} [d] \right\} = 0.$$

*Proof.* Suppose that  $\mathcal{G} = \mathcal{NN}(N_{\mathcal{G}}, L_{\mathcal{G}})$  satisfies  $N_{\mathcal{G}} \geq 7p + 1$ ,  $L_{\mathcal{G}} \geq 2$  and

$$n \leq \frac{N_{\mathcal{G}} - p - 1}{2} \left\lfloor \frac{N_{\mathcal{G}} - p - 1}{6p} \right\rfloor \left\lfloor \frac{L_{\mathcal{G}}}{2} \right\rfloor + 2.$$

For any  $\mathcal{D} \subset \{d \in \mathcal{NN}(N_{\mathcal{D}}, L_{\mathcal{D}}, K_{\mathcal{D}}) : d(\mathbb{R}^p) \subseteq [-B, B]\} \cap \mathfrak{H}(\mathbb{R}^p, C)$ ,  $B, C < \infty$ , by similar calculate of (5.3), we have

$$\begin{aligned} \mathfrak{h}_{g \in \mathcal{G}} \mathfrak{P}_{d \in \mathcal{D}} \left\{ \frac{1}{n} \sum_{i=1}^n d(\mathbf{X}_i) - \mathbb{E}_{g_{\#}\nu}[d] \right\} &\leq \mathfrak{h}_{g \in \mathcal{G}} \mathfrak{P}_{f \in \text{Lip}(\mathbb{R}^p, C)} \int f(d\pi_n - dg_{\#}\nu) \\ &\leq \mathfrak{h}_{g \in \mathcal{G}} C \mathfrak{P}_{\|f\|_{\text{Lip}} \leq 1} \int f(d\pi_n - dg_{\#}\nu) \\ &= C \mathfrak{h}_{g \in \mathcal{G}} \mathcal{W}_1(\hat{\pi}_n, g_{\#}\nu), \end{aligned}$$

where  $\hat{\pi}_n := \frac{1}{n} \sum_{i=1}^n \delta_{x_i}$  is the empirical distribution of  $\{X_i\}_{i=1, \dots, n}$ .

Since  $\mathcal{W}_1(\hat{\pi}_n, g_{\#}\nu) \rightarrow 0$  for some  $g_{\epsilon}$  as  $\epsilon \rightarrow 0$  by Lemma 3.2 of Yang et al. (2022).

$$\mathfrak{h}_{g \in \mathcal{G}} \mathfrak{P}_{d \in \mathcal{D}} \left\{ \frac{1}{n} \sum_{i=1}^n d(\mathbf{X}_i) - \mathbb{E}_{g_{\#}\nu}[d] \right\} = 0.$$

□

#### 5.1.4. Statistical error for simulated sample.

**Lemma 5.6.** Suppose  $\mathcal{G} = \mathcal{NN}(N_{\mathcal{G}}, L_{\mathcal{G}}, K_{\mathcal{G}})$ , and

$$\mathcal{D} = \{d \in \mathcal{NN}(N_{\mathcal{D}}, L_{\mathcal{D}}, K_{\mathcal{D}}) : d(\mathbb{R}^p) \subseteq [-B, B]\}, B < \infty,$$

then with probability at least  $1 - e^{-t_2}$ ,

$$d_{\mathcal{D} \circ \mathcal{G}}(\hat{\nu}_m, \nu) \leq \frac{2}{m} K^{L_{\mathcal{D}} + L_{\mathcal{G}} + 1} \sqrt{L_{\mathcal{D}} + L_{\mathcal{G}} + 2} \cdot \mathbb{E}_{\mathfrak{m}} \mathfrak{P}_{i \in [m]} \|\mathbf{z}_i\|_2 + 2B \sqrt{\frac{8t_2}{m}}.$$

*Proof.* Statistical error for simulated sample is defined as

$$d_{\mathcal{D} \circ \mathcal{G}}(\hat{\nu}_m, \nu) = \mathfrak{P}_{d \circ g \in \mathcal{D} \circ \mathcal{G}} \left| \frac{1}{m} \sum_{i=1}^m ((d \circ g)(\mathbf{z}_i) - \mathbb{E}_{\mathbf{z}_i \sim \nu}[(d \circ g)(\mathbf{z}_i)]) \right|.$$

Let  $\mathcal{R}_m(\mathcal{D} \circ \mathcal{G}; \mathbf{z}) := \mathbb{E} \mathfrak{P}_{d \circ g \in \mathcal{D} \circ \mathcal{G}} \left| \frac{1}{m} \sum_{i=1}^m \eta_i d \circ g(\mathbf{z}_i) \right|$  is the Rademacher complexity of functional class  $\mathcal{D} \circ \mathcal{G}$ , and  $\eta_1, \dots, \eta_m$  are independent Rademacher random variables.

We can obtain  $\mathbb{E} d_{\mathcal{D} \circ \mathcal{G}}(\hat{\nu}_m, \nu) \leq 2\mathcal{R}_m(\mathcal{D} \circ \mathcal{G}; \mathbf{z})$  by the symmetrization theorem. Using assumption  $\mathfrak{P}_{d \in \mathcal{D}} \|d\|_{\infty} \leq B$ , Massart's concentration inequality for suprema of the bounded empirical processes shows that

$$\begin{aligned} &\mathbb{P} \left\{ d_{\mathcal{D} \circ \mathcal{G}}(\hat{\nu}_m, \nu) \geq 2\mathcal{R}_m(\mathcal{D} \circ \mathcal{G}) + 2B \sqrt{\frac{8t_2}{m}} \right\} \\ &\leq \mathbb{P} \left\{ d_{\mathcal{D} \circ \mathcal{G}}(\hat{\nu}_m, \nu) \geq \mathbb{E} d_{\mathcal{D} \circ \mathcal{G}}(\hat{\nu}_m, \nu) + 2B \sqrt{\frac{8t_2}{m}} \right\} \leq e^{-t_2}. \end{aligned}$$

Next, we need to calculate  $\mathcal{R}_m(\mathcal{D} \circ \mathcal{G})$ .

Since the composition of two neural networks  $d \in \mathcal{D} = \mathcal{NN}(N_{\mathcal{D}}, L_{\mathcal{D}}, K_{\mathcal{D}})$  and  $g \in \mathcal{G} = \mathcal{NN}(N_{\mathcal{G}}, L_{\mathcal{G}}, K_{\mathcal{G}})$  is still a neural network with width  $\mathfrak{m} = \{N_{\mathcal{D}}, N_{\mathcal{G}}\}$  and depth  $L_{\mathcal{D}} + L_{\mathcal{G}} + 1$ . Lemma 5.4 implies

$$\mathbb{E} d_{\mathcal{D} \circ \mathcal{G}}(\hat{\nu}_m, \nu) \leq 2\mathcal{R}_m(\mathcal{D} \circ \mathcal{G}; \mathbf{z}) \leq \frac{2}{m} K_{\mathcal{D}} K_{\mathcal{G}} \sqrt{L_{\mathcal{D}} + L_{\mathcal{G}} + 2} \cdot \mathbb{E}_{\mathfrak{m}} \mathfrak{P}_{i \in [m]} \|\mathbf{z}_i\|_2.$$

Thus, with probability at least  $1 - e^{-t_2}$ ,

$$d_{\mathcal{D} \circ \mathcal{G}}(\widehat{\nu}_m, \nu) \leq \frac{2}{m} K_{\mathcal{D}} K_{\mathcal{G}} \sqrt{L_{\mathcal{D}} + L_{\mathcal{G}} + 2} \cdot \mathbb{E}_{\mathbf{z}_i} \|\mathbf{z}_i\|_2 + 2B \sqrt{\frac{8t_2}{m}}.$$

□

## 5.2. Proof of Theorem 3.1.

*Proof of Theorem 3.1.* By Lemmas 5.1, 5.2, 5.5 and 5.6, it is easy to see that, with probability at least  $1 - e^{-C_2 n^\alpha/2} - 2e^{-t_1} - e^{-t_2}$ ,

$$\begin{aligned} d_{\mathcal{H}}(\pi, \widehat{\mathbf{g}}_{\#} \nu) &\leq 2(2a_n)^b (K_{\mathcal{D}} / \mathfrak{g}^\gamma K_{\mathcal{D}})^{-\frac{b}{p+1}} + 2(1+B) p e^{-a_n + v^2/2} + \mathfrak{P}_{d \in \mathcal{D}} \left\{ \mathbb{E}_{\pi}[d] - \frac{1}{n} \sum_{i=1}^n d(\mathbf{X}_i) \right\} \\ &\quad + \mathfrak{f}_{\mathbf{g} \in \mathcal{G}} \mathfrak{P}_{d \in \mathcal{D}} \left\{ \frac{1}{n} \sum_{i=1}^n d(\mathbf{X}_i) - \mathbb{E}_{\mathbf{g}_{\#} \nu}[d] \right\} + 2 \mathfrak{P}_{d \circ \mathbf{g} \in \mathcal{D} \circ \mathcal{G}} \left\{ \mathbb{E}_{\widehat{\nu}_m}[d \circ \mathbf{g}] - \mathbb{E}_{\nu}[d \circ \mathbf{g}] \right\} \\ &\leq 2(2a_n)^b (K_{\mathcal{D}} / \mathfrak{g}^\gamma K_{\mathcal{D}})^{-\frac{b}{p+1}} + 2(1+B) p e^{-a_n + v^2/2} + \frac{1}{1 - e^{-C_2/\beta}} C_1^{1/\beta} K_{\mathcal{D}} e^{-C_2(s+1)/\beta} \\ &\quad + \frac{1}{1 - e^{-C_2/\beta}} C_1^{1/\beta} K_{\mathcal{D}} e^{-C_2 s/2\beta - C_2/\beta} + 4B \sqrt{\frac{st_1}{n}} + 4 \sqrt{\frac{2C_3}{n}} K_{\mathcal{D}} \cdot \sqrt{L_{\mathcal{D}} + 1 + \mathfrak{g}^\gamma p} \\ &\quad + \frac{4}{m} K_{\mathcal{D}} K_{\mathcal{G}} \sqrt{L_{\mathcal{D}} + L_{\mathcal{G}} + 2} \cdot \mathbb{E}_{\mathbf{z}_i} \|\mathbf{z}_i\|_2 + 4B \sqrt{\frac{8t_2}{m}} \\ &\leq 2(\mathfrak{g}^\gamma p n)^b (K_{\mathcal{D}} / \mathfrak{g}^\gamma K_{\mathcal{D}})^{-\frac{b}{p+1}} + 2(1+B) p e^{-\log p n + v^2/2} + \frac{C_1^{1/\beta} K_{\mathcal{D}}}{e^{C_2(n^\alpha+1)/\beta} (1 - e^{-C_2/\beta})} \\ &\quad + \frac{C_1^{1/\beta} K_{\mathcal{D}}}{(1 - e^{-C_2/\beta}) e^{C_2 n^\alpha/2 + C_2/\beta}} + 4B \sqrt{\frac{t_1}{n^{1-\alpha}}} + 4 \sqrt{\frac{2C_3}{n}} K_{\mathcal{D}} \cdot \sqrt{L_{\mathcal{D}} + 1 + \mathfrak{g}^\gamma p} \\ &\quad + \frac{4}{m} K_{\mathcal{D}} K_{\mathcal{G}} \sqrt{L_{\mathcal{D}} + L_{\mathcal{G}} + 2} \cdot \mathbb{E}_{\mathbf{z}_i} \|\mathbf{z}_i\|_2 + 4B \sqrt{\frac{8t_2}{m}} \\ &= O\left((\mathfrak{g}^\gamma p n)^b (K_{\mathcal{D}} / \mathfrak{g}^\gamma K_{\mathcal{D}})^{-\frac{b}{p+1}}\right) + O\left(\frac{1}{n}\right) + O\left(K_{\mathcal{D}} e^{-C_2 n^\alpha}\right) + O\left(K_{\mathcal{D}} e^{-C_2 n^\alpha/2}\right) \\ &\quad + O\left(\sqrt{\frac{t_1}{n^{1-\alpha}}}\right) + O\left(\frac{1}{\sqrt{n}} K_{\mathcal{D}} \sqrt{L_{\mathcal{D}} + \mathfrak{g}^\gamma p}\right) + O\left(\frac{1}{m} K_{\mathcal{D}} K_{\mathcal{G}} \sqrt{L_{\mathcal{D}} + L_{\mathcal{G}}} + \sqrt{\frac{t_2}{m}}\right) \\ &= O\left((\mathfrak{g}^\gamma p n)^b (K_{\mathcal{D}} / \mathfrak{g}^\gamma K_{\mathcal{D}})^{-\frac{b}{p+1}}\right) + O\left(K_{\mathcal{D}} e^{-C_2 n^\alpha}\right) + O\left(\sqrt{\frac{t_1}{n^{1-\alpha}}}\right) \\ &\quad + O\left(\sqrt{\frac{1}{n}} K_{\mathcal{D}} \sqrt{L_{\mathcal{D}} + \mathfrak{g}^\gamma p}\right) + O\left(\frac{1}{m} K_{\mathcal{D}} K_{\mathcal{G}} \sqrt{L_{\mathcal{D}} + L_{\mathcal{G}}} + \sqrt{\frac{t_2}{m}}\right). \end{aligned}$$

□

## REFERENCES

- Martin Arjovsky, Soumith Chintala, and Léon Bottou. Wasserstein generative adversarial networks. In *International conference on machine learning*, pages 214–223. PMLR, 2017.
- Jushan Bai. Common breaks in means and variances for panel data. *Journal of Econometrics*, 157(1):78–92, 2010.

- Yu Bai, Tengyu Ma, and Andrej Risteski. Approximability of discriminators implies diversity in gans. *International Conference on Learning Representations*, 2019.
- István Berkes, Weidong Liu, and Wei Biao Wu. Komlós–major–tusnády approximation under dependence. *The Annals of Probability*, 42(2):794–817, 2014.
- Eoin Brophy, Zhengwei Wang, Qi She, and Tomas Ward. Generative adversarial networks in time series: A survey and taxonomy. *arXiv preprint arXiv:2107.11098*, 2021.
- Minwoo Chae, Dongha Kim, Yongdai Kim, and Lizhen Lin. A likelihood approach to non-parametric estimation of a singular distribution using deep generative models. *arXiv preprint arXiv:2105.04046*, 2021.
- Wei-Cheng Chang, Chun-Liang Li, Yiming Yang, and Barnabás Póczos. Kernel change-point detection with auxiliary deep generative models. *arXiv preprint arXiv:1901.06077*, 2019.
- Hao Chen and Nancy Zhang. Graph-based change-point detection. *The Annals of Statistics*, 43(1):139–176, 2015.
- Likai Chen, Weining Wang, and Wei Biao Wu. Inference of breakpoints in high-dimensional time series. *Journal of the American Statistical Association*, pages 1–13, 2021.
- Haeran Cho and Piotr Fryzlewicz. Multiple-change-point detection for high dimensional time series via sparsified binary segmentation. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 77(2):475–507, 2015.
- Farida Enikeeva and Zaid Harchaoui. High-dimensional change-point detection under sparse alternatives. *The Annals of Statistics*, 47(4):2051–2079, 2019.
- Cristóbal Esteban, Stephanie L Hyland, and Gunnar Rätsch. Real-valued (medical) time series generation with recurrent conditional gans. *arXiv preprint arXiv:1706.02633*, 2017.
- Yihang Gao, Michael K. Ng, and Mingjie Zhou. Approximation for probability distributions by wasserstein gan. *arXiv preprint arXiv:2103.10060*, 2022.
- Noah Golowich, Alexander Rakhlin, and Ohad Shamir. Size-independent sample complexity of neural networks. In *Conference On Learning Theory*, pages 297–299. PMLR, 2018.
- Ian J Goodfellow, Jean Pouget-Abadie, Mehdi Mirza, Bing Xu, David Warde-Farley, Sherjil Ozair, Aaron Courville, and Yoshua Bengio. Generative adversarial nets. *stat*, 1050:10, 2014.
- Sébastien Gouëzel. Almost sure invariance principle for dynamical systems by spectral methods. *The Annals of Probability*, 38(4):1639–1671, 2010.
- Zaid Harchaoui and Céline Lévy-Leduc. Multiple change-point estimation with a total variation penalty. *Journal of the American Statistical Association*, 105(492):1480–1493, 2010.
- Zaid Harchaoui, Félicien Vallet, Alexandre Lung-Yut-Fong, and Olivier Cappé. A regularized kernel-based approach to unsupervised audio segmentation. In *2009 IEEE international conference on acoustics, speech and signal processing*, pages 1665–1668. IEEE, 2009.
- Jian Huang, Yuling Jiao, Zhen Li, Shiao Liu, Yang Wang, and Yunfei Yang. An error analysis of generative adversarial networks for learning distributions. *Journal of Machine Learning Research*, 23(116):1–43, 2022.
- Yuling Jiao, Yang Wang, and Yunfei Yang. Approximation bounds for norm constrained neural networks with applications to regression and gans. *arXiv preprint arXiv:2201.09418*, 2022.
- Moritz Jirak. Uniform change point tests in high dimension. *The Annals of Statistics*, 43(6):2451–2483, 2015.
- Degui Li, Junhui Qian, and Liangjun Su. Panel data models with interactive fixed effects and multiple structural breaks. *Journal of the American Statistical Association*, 111(516):1804–1819, 2016.
- Tengyuan Liang. How well generative adversarial networks learn distributions. *The Journal of Machine Learning Research*, 22(1):10366–10406, 2021.

- Shiao Liu, Yunfei Yang, Jian Huang, Yuling Jiao, and Yang Wang. Non-asymptotic error bounds for bidirectional gans. *Advances in Neural Information Processing Systems*, 34: 12328–12339, 2021.
- Weidong Liu and Zhengyan Lin. Strong approximation for a class of stationary processes. *Stochastic Processes and their Applications*, 119(1):249–280, 2009.
- Jianya Lu, Wei Biao Wu, Zhijie Xiao, and Lihu Xu. Almost sure invariance principle of  $\beta$ -mixing time series in hilbert space. *arXiv preprint arXiv:2209.12535*, 2022.
- Yulong Lu and Jianfeng Lu. A universal approximation theorem of deep neural networks for expressing probability distributions. *Advances in neural information processing systems*, 33: 3094–3105, 2020.
- Pascal Massart. About the constants in talagrand’s concentration inequalities for empirical processes. *The Annals of Probability*, 28(2):863–884, 2000.
- Olof Mogren. C-rnn-gan: Continuous recurrent neural networks with adversarial training. *arXiv preprint arXiv:1611.09904*, 2016.
- Hao Ni, Lukasz Szpruch, Magnus Wiese, Shujian Liao, and Baoren Xiao. Conditional sig-wasserstein gans for time series generation. *arXiv preprint arXiv:2006.05421*, 2020.
- Evgenia Romanenkova, Alexander Stepikin, Matvey Morozov, and Alexey Zaytsev. Indid: Instant disorder detection via a principled neural network. In *Proceedings of the 30th ACM International Conference on Multimedia*, pages 3152–3162, 2022.
- Amirhossein Taghvaei and Amin Jalali. 2-wasserstein approximation via restricted convex potentials with application to improved training for gans. *arXiv preprint arXiv:1902.07197*, 2019.
- Robert Tibshirani and Pei Wang. Spatial smoothing and hot spot detection for cgh data using the fused lasso. *Biostatistics*, 9(1):18–29, 2008.
- Wei Biao Wu. Nonlinear system theory: Another look at dependence. *Proceedings of the National Academy of Sciences*, 102(40):14150–14154, 2005.
- Wei Biao Wu. Strong invariance principles for dependent random variables. *The Annals of Probability*, 35(6):2294–2320, 2007.
- Yunfei Yang, Zhen Li, and Yang Wang. On the capacity of deep generative networks for approximating distributions. *Neural Networks*, 145:144–154, 2022.
- Jinsung Yoon, Daniel Jarrett, and Mihaela Van der Schaar. Time-series generative adversarial networks. *Advances in neural information processing systems*, 32, 2019.
- Pengchuan Zhang, Qiang Liu, Dengyong Zhou, Tao Xu, and Xiaodong He. On the discrimination-generalization tradeoff in gans. *arXiv preprint arXiv:1711.02771*, 2017.
- Banghua Zhu, Jiantao Jiao, and David Tse. Deconstructing generative adversarial networks. *IEEE Transactions on Information Theory*, 66(11):7155–7179, 2020.

(J. Lu) DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF ESSEX  
*Email address:* jianya.lu@essex.ac.uk

(Y. Mo) DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND TECHNOLOGY, UNIVERSITY OF  
MACAU, MACAU S.A.R., CHINA  
*Email address:* yc27477@um.edu.mo

(Z. Xiao) DEPARTMENT OF ECONOMICS, BOSTON COLLEGE  
*Email address:* zhijie.xiao@bc.edu

(L. Xu) DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND TECHNOLOGY, UNIVERSITY OF  
MACAU, MACAU S.A.R., CHINA  
*Email address:* lihuxu@um.edu.mo

(Q. Yao) DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND TECHNOLOGY, UNIVERSITY OF  
MACAU, MACAU S.A.R., CHINA  
*Email address:* yb97478@um.edu.mo