

A MOLECULAR RECONSTRUCTION THEOREM FOR $H_{\omega}^{p(\cdot)}(\mathbb{R}^n)$

PABLO ROCHA

Abstract

In this article we give a molecular reconstruction theorem for $H_{\omega}^{p(\cdot)}(\mathbb{R}^n)$. As an application of this result and the atomic decomposition developed in [5] we show that classical singular integrals can be extended to bounded operators on $H_{\omega}^{p(\cdot)}(\mathbb{R}^n)$. We also prove, for certain exponents $q(\cdot)$ and certain weights ω , that Riesz potential I_{α} , with $0 < \alpha < n$, can be extended to a bounded operator from $H_{\omega}^{p(\cdot)}(\mathbb{R}^n)$ into $H_{\omega}^{q(\cdot)}(\mathbb{R}^n)$, for $\frac{1}{p(\cdot)} := \frac{1}{q(\cdot)} + \frac{\alpha}{n}$.

1 Introduction

In the celebrated paper [4], Fefferman and Stein defined the Hardy space $H^p(\mathbb{R}^n)$. Since then, the study of Hardy spaces has received the attention of a substantial number of researchers. One of the most remarkable results for the study of Hardy spaces is the atomic characterization of $H^p(\mathbb{R}^n)$ (see [1, 7]). Roughly speaking, every distribution $f \in H^p$ can be expressed of the form

$$f = \sum_j \lambda_j a_j,$$

where the a_j 's are p -atoms, $\{\lambda_j\} \in \ell^p$ and $\|f\|_{H^p}^p \approx \sum_j |\lambda_j|^p$. For $0 < p \leq 1$, an p -atom is a function $a(\cdot)$ supported on a cube Q such that

$$\|a\|_{\infty} \leq |Q|^{-1/p} \quad \text{and} \quad \int x^{\alpha} a(x) dx = 0 \quad \text{for all } |\alpha| \leq n \left(\frac{1}{p} - 1 \right).$$

2020 *Mathematics Subject Classification*. Primary 42B30; Secondary 42B20, 42B25, 42B35.

Key words and phrases: weighted variable Hardy spaces, molecular decomposition, singular integrals, Riesz potential.

Such decomposition allows to study the behavior of certain operators on H^p by focusing one's attention on individual atoms. In principle, the continuity of an operator T on H^p can often be proved by estimating Ta when $a(\cdot)$ is an atom. In [2] was observed that, in general, the atoms are not mapped into atoms. However, for many convolution operators (e.g.: singular integrals or Riesz potentials), $m = Ta$ behaves like an atom. These functions m were called *molecules*, their properties as well as the molecular characterization of $H^p(\mathbb{R}^n)$ were established in [15]. Then, in essence, the continuity $H^p \rightarrow H^p$ of an operator reduces to showing that it maps atoms into molecules.

The theory of variable Hardy spaces $H^{p(\cdot)}(\mathbb{R}^n)$ was first developed by Nakai and Sawano in [8], where they proved the infinite atomic decomposition and molecular decomposition for $H^{p(\cdot)}$. As a corollary of these decompositions, they obtained the boundedness of singular integral operators on $H^{p(\cdot)}$. Later, Cruz-Uribe and Wang in [3] independently considered the same problem under a weaker condition for variable exponents $p(\cdot)$. They gave a finite atomic decomposition for $H^{p(\cdot)}$ and also proved the boundedness of singular integral operators on $H^{p(\cdot)}$. Both theories prove equivalent definitions in terms of maximal operators using different approaches. In [12], the author jointly with Urciuolo proved the $H^{p(\cdot)} - L^{q(\cdot)}$ boundedness of certain generalized Riesz potentials and the $H^{p(\cdot)} - H^{q(\cdot)}$ boundedness of Riesz potential via the infinite atomic and molecular decomposition developed in [8]. In [10], the author gave another proof of the results obtained in [12], but by using the finite atomic decomposition given in [3].

Recently, Kwok-Pun Ho in [5] developed the weighted theory for variable Hardy spaces on \mathbb{R}^n . He established the atomic decompositions for the weighted Hardy spaces with variable exponents $H_\omega^{p(\cdot)}(\mathbb{R}^n)$ and also revealed some intrinsic structures of atomic decomposition for Hardy type spaces. His results generalize the infinite atomic decomposition obtained in [8]. By means of the atomic decomposition given in [5], the author in [11] proved, for certain exponents $q(\cdot)$ and certain weights ω , the $H_\omega^{p(\cdot)} - L_\omega^{q(\cdot)}$ boundedness for Riesz potential.

The purpose of this article is to generalize the molecular decomposition obtained in [8, Theorem 5.2] to weighted variable Hardy spaces by using the framework developed in [5]. As an application of our molecular decomposition, we prove two theorems concerning the $H_\omega^{p(\cdot)} - H_\omega^{p(\cdot)}$ boundedness of a class of singular integral operators and the $H_\omega^{p(\cdot)} - H_\omega^{q(\cdot)}$ boundedness of Riesz potential. These results are established in Theorems 4.2 and 5.2 below.

This paper is organized as follows. Section 2 gives weighted variable Hardy spaces $H_\omega^{p(\cdot)}$ and some of their preliminary results. Section 3 contains a molecular reconstruction theorem for $H_\omega^{p(\cdot)}$. The boundedness of singular integrals on $H_\omega^{p(\cdot)}$ is established in Section 4. The $H_\omega^{p(\cdot)} - H_\omega^{q(\cdot)}$ boundedness of Riesz potential is proved in Section 5.

Notation. The symbol $A \lesssim B$ stands for the inequality $A \leq cB$ for some positive constant c . The symbol $A \approx B$ stands for $B \lesssim A \lesssim B$. We denote by $Q(z, r)$ the cube centered at $z \in \mathbb{R}^n$ with side length r . Given a cube $Q = Q(z, r)$, we set $kQ = Q(z, kr)$ and $\ell(Q) = r$. For a measurable subset $E \subseteq \mathbb{R}^n$ we denote by $|E|$ and χ_E the Lebesgue measure of E and the characteristic function of E respectively. Given a real number $s \geq 0$, we write $[s]$ for the integer part of s . As usual we denote with $\mathcal{S}(\mathbb{R}^n)$ the space of smooth and rapidly decreasing functions and with $\mathcal{S}'(\mathbb{R}^n)$ the dual space. If β is the multiindex $\beta = (\beta_1, \dots, \beta_n)$, then $|\beta| = \beta_1 + \dots + \beta_n$.

2 Preliminaries

We begin with the definition of weighted variable Lebesgue spaces $L_\omega^{p(\cdot)}(\mathbb{R}^n)$.

Let $p(\cdot) : \mathbb{R}^n \rightarrow (0, \infty)$ be a measurable function. Given a measurable set E , let

$$p_-(E) = \operatorname{ess\,inf}_{x \in E} p(x), \quad \text{and} \quad p_+(E) = \operatorname{ess\,sup}_{x \in E} p(x).$$

When $E = \mathbb{R}^n$, we will simply write $p_- := p_-(\mathbb{R}^n)$ and $p_+ := p_+(\mathbb{R}^n)$.

Given a measurable function f on \mathbb{R}^n , define the modular ρ associated with $p(\cdot)$ by

$$\rho(f) = \int_{\mathbb{R}^n} |f(x)|^{p(x)} dx.$$

We define the variable Lebesgue space $L^{p(\cdot)} = L^{p(\cdot)}(\mathbb{R}^n)$ to be the set of all measurable functions f such that, for some $\lambda > 0$, $\rho(f/\lambda) < \infty$. This becomes a quasi normed space when equipped with the Luxemburg norm

$$\|f\|_{L^{p(\cdot)}} = \inf \{ \lambda > 0 : \rho(f/\lambda) \leq 1 \}.$$

Given a weight ω , i.e.: a locally integrable function on \mathbb{R}^n such that $0 < \omega(x) < \infty$ almost everywhere, we define the weighted variable Lebesgue space $L_\omega^{p(\cdot)}$ as the set of all measurable functions $f : \mathbb{R}^n \rightarrow \mathbb{C}$ such that $\|f\omega\|_{L^{p(\cdot)}} < \infty$. If $f \in L_\omega^{p(\cdot)}$, we define its "norm" by

$$\|f\|_{L_\omega^{p(\cdot)}} := \|f\omega\|_{L^{p(\cdot)}}.$$

The following result follows from the definition of the $L_\omega^{p(\cdot)}$ -norm.

LEMMA 2.1. *Given a measurable function $p(\cdot) : \mathbb{R}^n \rightarrow (0, \infty)$ with $0 < p_- \leq p_+ < \infty$ and a weight ω , then for every $s > 0$,*

$$\|f\|_{L_\omega^{p(\cdot)}}^s = \| |f|^s \|_{L_{\omega^s}^{p(\cdot)/s}}.$$

Next, we introduce the weights used in [5] to define weighted Hardy spaces with variable exponents.

DEFINITION 2.2. (See [11, Remark 2.5]) Let $p(\cdot) : \mathbb{R}^n \rightarrow (0, \infty)$ be a measurable function with $0 < p_- \leq p_+ < \infty$. We define $\mathcal{W}_{p(\cdot)}$ as the set of all weights ω such that

(i) there exists $0 < p_* < \min\{1, p_-\}$ such that $\|\chi_Q\|_{L_{\omega}^{p(\cdot)/p_*}} < \infty$, and

$$\|\chi_Q\|_{L_{\omega^{-p_*}}^{(p(\cdot)/p_*)'}} < \infty, \text{ for all cube } Q;$$

(ii) there exist $\kappa > 1$ and $s > 1$ such that Hardy-Littlewood maximal

operator M is bounded on $L_{\omega^{-\kappa/s}}^{(sp(\cdot))'/\kappa}$.

For a measurable function $p(\cdot) : \mathbb{R}^n \rightarrow (0, \infty)$ such that $0 < p_- \leq p_+ < \infty$ and $\omega \in \mathcal{W}_{p(\cdot)}$, in [5] the author give a variety of distinct approaches, based on differing definitions, all lead to the same notion of weighted variable Hardy space $H_{\omega}^{p(\cdot)}$.

We recall some terminologies and notations from the study of maximal functions. Given $N \in \mathbb{N}$, define

$$\mathcal{F}_N = \left\{ \varphi \in \mathcal{S}(\mathbb{R}^n) : \sum_{|\beta| \leq N} \sup_{x \in \mathbb{R}^n} (1 + |x|)^N |\partial^{\beta} \varphi(x)| \leq 1 \right\}.$$

For any $f \in \mathcal{S}'(\mathbb{R}^n)$, the grand maximal function of f is given by

$$\mathcal{M}_N f(x) = \sup_{t > 0} \sup_{\varphi \in \mathcal{F}_N} |(\varphi_t * f)(x)|,$$

where $\varphi_t(x) = t^{-n} \varphi(t^{-1}x)$. It is common to denote the grand maximal by \mathcal{M} .

DEFINITION 2.3. Let $p(\cdot) : \mathbb{R}^n \rightarrow (0, \infty)$, $0 < p_- \leq p_+ < \infty$, and $\omega \in \mathcal{W}_{p(\cdot)}$. The weighted variable Hardy space $H_{\omega}^{p(\cdot)}(\mathbb{R}^n)$ is the set of all $f \in \mathcal{S}'(\mathbb{R}^n)$ for which $\|\mathcal{M}f\|_{L_{\omega}^{p(\cdot)}} < \infty$. In this case we define $\|f\|_{H_{\omega}^{p(\cdot)}} := \|\mathcal{M}f\|_{L_{\omega}^{p(\cdot)}}$.

DEFINITION 2.4. Let $p(\cdot) : \mathbb{R}^n \rightarrow (0, \infty)$, $0 < p_- \leq p_+ < \infty$, $p_0 > 1$, and $\omega \in \mathcal{W}_{p(\cdot)}$. Fix an integer $N \geq 1$. A function $a(\cdot)$ on \mathbb{R}^n is called a $\omega - (p(\cdot), p_0, N)$ atom if there exists a cube Q such that

$$a_1) \text{ supp}(a) \subset Q,$$

$$a_2) \|a\|_{L^{p_0}} \leq \frac{|Q|^{\frac{1}{p_0}}}{\|\chi_Q\|_{L_{\omega}^{p(\cdot)}}},$$

$$a_3) \int x^{\beta} a(x) dx = 0 \text{ for all } |\beta| \leq N.$$

Now, we introduce two indices, which are related to the intrinsic structure of the atomic decomposition of $H_\omega^{p(\cdot)}$. Given $\omega \in \mathcal{W}_{p(\cdot)}$, we write

$$s_{\omega, p(\cdot)} := \inf \left\{ s \geq 1 : M \text{ is bounded on } L_{\omega^{-1/s}}^{(sp(\cdot))'} \right\}$$

and

$$\mathbb{S}_{\omega, p(\cdot)} := \left\{ s \geq 1 : M \text{ is bounded on } L_{\omega^{-\kappa/s}}^{(sp(\cdot))'/\kappa} \text{ for some } \kappa > 1 \right\}.$$

Then, for every fixed $s \in \mathbb{S}_{\omega, p(\cdot)}$, we define

$$\kappa_{\omega, p(\cdot)}^s := \sup \left\{ \kappa > 1 : M \text{ is bounded on } L_{\omega^{-\kappa/s}}^{(sp(\cdot))'/\kappa} \right\}.$$

The index $\kappa_{\omega, p(\cdot)}^s$ is used to measure the left-openness of the boundedness of M on the family $\left\{ L_{\omega^{-\kappa/s}}^{(sp(\cdot))'/\kappa} \right\}_{\kappa > 1}$. The index $s_{\omega, p(\cdot)}$ is related to the vanishing moment condition and the index $\kappa_{\omega, p(\cdot)}^s$ is related to the size condition of the atoms (see [5, Theorems 5.3 and 6.3]). These indices are also related to the structure of our molecular decomposition (see Definition 3.1 and Theorem 3.4 below).

DEFINITION 2.5. Let $p(\cdot) : \mathbb{R}^n \rightarrow (0, \infty)$ be a measurable function such that $0 < p_- \leq p_+ < \infty$, and let $\omega \in \mathcal{W}_{p(\cdot)}$. Given a sequence of scalars $\{\lambda_j\}_{j=1}^\infty$, a family of cubes $\{Q_j\}_{j=1}^\infty$ and $0 < \theta < \infty$, we define

$$\mathcal{A}(\{\lambda_j\}_{j=1}^\infty, \{Q_j\}_{j=1}^\infty, p(\cdot), \omega, \theta) := \left\| \sum_{j=1}^\infty \left(\frac{|\lambda_j|}{\|\chi_{Q_j}\|_{L_\omega^{p(\cdot)}}} \right)^\theta \chi_{Q_j} \right\|_{L_{\omega^\theta}^{p(\cdot)/\theta}}^{1/\theta}.$$

The following theorem is a version of the atomic decomposition for $H_\omega^{p(\cdot)}$ obtained in [5].

THEOREM 2.6. *Let $1 < p_0 < \infty$, $p(\cdot) : \mathbb{R}^n \rightarrow (0, \infty)$ be a measurable function with $0 < p_- \leq p_+ < \infty$ and $\omega \in \mathcal{W}_{p(\cdot)}$. Then, for every $f \in H_\omega^{p(\cdot)}(\mathbb{R}^n) \cap L^{p_0}(\mathbb{R}^n)$ and every integer $N \geq \lceil ns_{\omega, p(\cdot)} - n \rceil$ fixed, there exist a sequence of scalars $\{\lambda_j\}_{j=1}^\infty$, a sequence of cubes $\{Q_j\}_{j=1}^\infty$ and $\omega - (p(\cdot), p_0, N)$ atoms a_j supported on Q_j such that $f = \sum_{j=1}^\infty \lambda_j a_j$ converges in $L^{p_0}(\mathbb{R}^n)$ and*

$$(1) \quad \mathcal{A}(\{\lambda_j\}_{j=1}^\infty, \{Q_j\}_{j=1}^\infty, p(\cdot), \omega, \theta) \lesssim \|f\|_{H_\omega^{p(\cdot)}}, \text{ for all } 0 < \theta < \infty,$$

where the implicit constant in (1) is independent of $\{\lambda_j\}_{j=1}^\infty$, $\{Q_j\}_{j=1}^\infty$, and f .

PROOF. The existence of a such atomic decomposition is guaranteed by [5, Theorem 6.2]. Its construction is analogous to that given for classical Hardy spaces (see [14, Chapter III]). So, following the proof in [9, Theorem 3.1], we obtain the convergence of the atomic series to f in $L^{p_0}(\mathbb{R}^n)$. \square

The following three results will be useful in what follows.

LEMMA 2.7. ([11, Lemma 3.4]) Let $p(\cdot) : \mathbb{R}^n \rightarrow (0, \infty)$ be a measurable function with $0 < p_- \leq p_+ < \infty$. If $\omega \in \mathcal{W}_{p(\cdot)}$, then, for every cube $Q \subset \mathbb{R}^n$,

$$\|\chi_{2\sqrt{n}Q}\|_{L_\omega^{p(\cdot)}} \approx \|\chi_Q\|_{L_\omega^{p(\cdot)}}.$$

We say that an exponent function $p(\cdot) : \mathbb{R}^n \rightarrow (0, \infty)$ such that $0 < p_- \leq p_+ < \infty$ belongs to $\mathcal{P}^{\log}(\mathbb{R}^n)$, if there exist two positive constants C and C_∞ such that $p(\cdot)$ satisfies the local log-Hölder continuity condition, i.e.:

$$|p(x) - p(y)| \leq \frac{C}{-\log(|x - y|)}, \quad |x - y| \leq \frac{1}{2},$$

and is log-Hölder continuous at infinity, i.e.:

$$|p(x) - p_\infty| \leq \frac{C_\infty}{\log(e + |x|)}, \quad x \in \mathbb{R}^n,$$

for some $p_- \leq p_\infty \leq p_+$.

We define the set $\mathcal{S}_0(\mathbb{R}^n)$ by

$$\mathcal{S}_0(\mathbb{R}^n) = \left\{ \varphi \in \mathcal{S}(\mathbb{R}^n) : \int x^\beta \varphi(x) dx = 0, \text{ for all } \beta \in \mathbb{N}_0^n \right\}.$$

PROPOSITION 2.8. ([6, Proposition 2.1]) Let $p(\cdot) \in \mathcal{P}^{\log}(\mathbb{R}^n)$ with $0 < p_- \leq p_+ < \infty$. If $\omega \in \mathcal{W}_{p(\cdot)}$, then $\mathcal{S}_0(\mathbb{R}^n) \subset H_\omega^{p(\cdot)}(\mathbb{R}^n)$ densely.

We conclude this preliminaries with a supporting result, which will allow us to study the behavior of Riesz potential on $H_\omega^{p(\cdot)}$.

PROPOSITION 2.9. ([11, Corollary 4.2]) Let $0 < \alpha < n$, $q(\cdot) : \mathbb{R}^n \rightarrow (0, \infty)$ be a measurable function with $0 < q_- \leq q_+ < \infty$ and $\omega \in \mathcal{W}_{q(\cdot)}$. If $\frac{1}{p(\cdot)} := \frac{1}{q(\cdot)} + \frac{\alpha}{n}$ and $\|\chi_Q\|_{L_\omega^{q(\cdot)}} \approx |Q|^{-\alpha/n} \|\chi_Q\|_{L_\omega^{p(\cdot)}}$ for every cube Q , then for any sequence of scalars $\{\lambda_j\}_{j=1}^\infty$, any family of cubes $\{Q_j\}_{j=1}^\infty$, and any $\theta \in (0, \infty)$ fixed we have

$$\mathcal{A}(\{\lambda_j\}_{j=1}^\infty, \{Q_j\}_{j=1}^\infty, q(\cdot), \omega, \theta) \lesssim \mathcal{A}(\{\lambda_j\}_{j=1}^\infty, \{Q_j\}_{j=1}^\infty, p(\cdot), \omega, \theta).$$

3 Molecular reconstruction for $H_\omega^{p(\cdot)}(\mathbb{R}^n)$

Our definition of ω -molecule is a slight modification from that given in [8] by Nakai and Sawano for variable Hardy spaces. Namely, we replace the amount $\|\chi_Q\|_{L^{p(\cdot)}}$ by $\|\chi_Q\|_{L_\omega^{p(\cdot)}}$.

DEFINITION 3.1. Let $p(\cdot) : \mathbb{R}^n \rightarrow (0, \infty)$ be a measurable function with $0 < p_- \leq p_+ < \infty$ and let $\omega \in \mathcal{W}_{p(\cdot)}$. Let $p_0 > 1$ and $d_{p(\cdot)} := \lfloor ns_{\omega, p(\cdot)} - n \rfloor$. A function $m(\cdot)$ on \mathbb{R}^n is called a $\omega - (p(\cdot), p_0, d_{p(\cdot)})$ molecule centered at a cube $Q = Q(z, r)$ (with side length $\ell(Q) = r$) if

$$\begin{aligned} m_1) \quad & \|m\|_{L^{p_0}(2\sqrt{n}Q)} \leq \frac{|Q|^{\frac{1}{p_0}}}{\|\chi_Q\|_{L_\omega^{p(\cdot)}}}, \\ m_2) \quad & |m(x)| \leq \|\chi_Q\|_{L_\omega^{p(\cdot)}}^{-1} \left(1 + \frac{|x-z|}{\ell(Q)}\right)^{-2n-2d_{p(\cdot)}-3} \quad \text{for all } x \in \mathbb{R}^n \setminus Q(z, 2\sqrt{n}r), \\ m_3) \quad & \int x^\beta m(x) dx = 0 \text{ for all multi-index } \beta \text{ with } |\beta| \leq d_{p(\cdot)}. \end{aligned}$$

REMARK 3.2. The conditions $m_1)$ and $m_2)$ imply that $\|m\|_{L^{p_0}(\mathbb{R}^n)} \leq C \frac{|Q|^{\frac{1}{p_0}}}{\|\chi_Q\|_{L_\omega^{p(\cdot)}}}$, where C is a positive constant independent of the molecule m .

Next, we give a molecular reconstruction theorem for $H_\omega^{p(\cdot)}(\mathbb{R}^n)$. For them, we need to introduce the following discrete maximal: given $\phi \in \mathcal{S}(\mathbb{R}^n)$ and $f \in \mathcal{S}'(\mathbb{R}^n)$, we define

$$M_\phi^{dis} f(x) = \sup_{j \in \mathbb{Z}} |(\phi^j * f)(x)|,$$

where $\phi^j(x) = 2^{jn}\phi(2^jx)$. From [8, Lemma 3.2 and Proof of Theorem 3.3], it follows that for all $f \in \mathcal{S}'(\mathbb{R}^n)$ and all $0 < \theta < 1$

$$(2) \quad \mathcal{M}_N f(x) \leq C \left[M \left((M_\phi^{dis} f)^\theta \right) (x) \right]^{\frac{1}{\theta}}, \quad \text{for all } x \in \mathbb{R}^n,$$

if N is sufficiently large. This inequality gives the following result.

LEMMA 3.3. *Let $\phi \in \mathcal{S}(\mathbb{R}^n)$ and $f \in \mathcal{S}'(\mathbb{R}^n)$. If $\omega \in \mathcal{W}_{p(\cdot)}$, then $\|f\|_{H_\omega^{p(\cdot)}} \leq C \|M_\phi^{dis} f\|_{L_\omega^{p(\cdot)}}$, where C is a positive constant which does not depend on f .*

PROOF. From (2) above and Lemma 2.1, we have

$$\|f\|_{H_\omega^{p(\cdot)}} = \|\mathcal{M}_N f\|_{L_\omega^{p(\cdot)}} \lesssim \left\| \left[M \left((M_\phi^{dis} f)^\theta \right) (\cdot) \right]^{\frac{1}{\theta}} \right\|_{L_\omega^{p(\cdot)}} = \left\| M \left((M_\phi^{dis} f)^\theta \right) \right\|_{L_\omega^{p(\cdot)/\theta}}^{1/\theta}.$$

By taking $\frac{1}{\theta} > s_{\omega, p(\cdot)}$, by [5, Theorem 3.1] and Lemma 2.1, we get

$$\|f\|_{H_\omega^{p(\cdot)}} \lesssim \left\| M \left((M_\phi^{dis} f)^\theta \right) \right\|_{L_\omega^{p(\cdot)/\theta}}^{1/\theta} \lesssim \left\| (M_\phi^{dis} f)^\theta \right\|_{L_\omega^{p(\cdot)/\theta}}^{1/\theta} = \|M_\phi^{dis} f\|_{L_\omega^{p(\cdot)}}.$$

Consequently, we obtain the desired result. \square

THEOREM 3.4. *Let $p(\cdot) : \mathbb{R}^n \rightarrow (0, \infty)$ be a measurable function with $0 < p_- \leq p_+ < \infty$, and $\omega \in \mathcal{W}_{p(\cdot)}$. Suppose that $0 < \theta < 1$ satisfies $\frac{1}{\theta} \in \mathbb{S}_{\omega, p(\cdot)}$. Let $\{\lambda_j\}_{j=1}^\infty$ be a sequence of scalars and let $\{Q_j\}_{j=1}^\infty$ be a family of cubes such that*

$$\mathcal{A}(\{\lambda_j\}_{j=1}^\infty, \{Q_j\}_{j=1}^\infty, p(\cdot), \omega, \theta) := \left\| \sum_{j=1}^\infty \left(\frac{|\lambda_j|}{\|\chi_{Q_j}\|_{L_\omega^{p(\cdot)}}} \right)^\theta \chi_{Q_j} \right\|_{L_\omega^{p(\cdot)/\theta}}^{1/\theta} < \infty.$$

If $\{m_j\}_{j=1}^\infty$ is a sequence of $\omega - (p(\cdot), p_0, [ns_{\omega, p(\cdot)} - n])$ molecules, with $p_0 > \theta \left(\kappa_{\omega, p(\cdot)}^{1/\theta} \right)'$, such that m_j is centered at Q_j for every $j \in \mathbb{N}$ and the series

$$g := \sum_{j=1}^\infty \lambda_j m_j$$

converges in $\mathcal{S}'(\mathbb{R}^n)$, then $g \in H_\omega^{p(\cdot)}(\mathbb{R}^n)$ with

$$(3) \quad \|g\|_{H_\omega^{p(\cdot)}} \lesssim \mathcal{A}(\{\lambda_j\}_{j=1}^\infty, \{Q_j\}_{j=1}^\infty, p(\cdot), \omega, \theta),$$

where the implicit constant in (3) does not depend on g , $\{\lambda_j\}_{j=1}^\infty$, and $\{Q_j\}_{j=1}^\infty$.

PROOF. Let $\phi \in C_c^\infty(\mathbb{R}^n)$ such that $\chi_{B(0,1)} \leq \phi \leq \chi_{B(0,2)}$, we set $\phi_{2^k}(x) = 2^{kn} \phi(2^k x)$ where $k \in \mathbb{Z}$. Since $g = \sum_j \lambda_j m_j$ in $\mathcal{S}'(\mathbb{R}^n)$, it follows that

$$|(\phi_{2^k} * g)(x)| \leq \sum_j |\lambda_j| |(\phi_{2^k} * m_j)(x)|, \quad \text{for all } x \in \mathbb{R}^n, \text{ and all } k \in \mathbb{Z}.$$

We observe that the argument used in [8, Proof of Theorem 5.2] to obtain the pointwise inequality (5.2) works in this setting, but considering now the conditions m_1 , m_2 and m_3 in Definition 3.1. Therefore we get

$$(4) \quad M_\phi^{dis} g(x) \lesssim \sum_j |\lambda_j| \chi_{2\sqrt{n}Q_j}(x) M(m_j)(x) + \sum_j |\lambda_j| \frac{[M(\chi_{Q_j})(x)]^{\frac{n+d_{p(\cdot)}+1}{n}}}{\|\chi_{Q_j}\|_{L_\omega^{p(\cdot)}}}, \quad (x \in \mathbb{R}^n),$$

where M is the Hardy-Littlewood maximal operator and $d_{p(\cdot)} = [ns_{\omega, p(\cdot)} - n]$.

Then, from (4) above and Lemma 3.3, we have

$$\begin{aligned} & \|g\|_{H_\omega^{p(\cdot)}} \lesssim \|M_\phi^{dis} g\|_{L_\omega^{p(\cdot)}} \\ & \lesssim \left\| \sum_j |\lambda_j| \chi_{2\sqrt{n}Q_j} \cdot M(m_j) \right\|_{L_\omega^{p(\cdot)}} + \left\| \sum_j |\lambda_j| \frac{[M(\chi_{Q_j})(\cdot)]^{\frac{n+d_{p(\cdot)}+1}{n}}}{\|\chi_{Q_j}\|_{L_\omega^{p(\cdot)}}} \right\|_{L_\omega^{p(\cdot)}} = J_1 + J_2. \end{aligned}$$

Now, we consider J_1 . The boundedness of the Hardy-Littlewood maximal operator M on L^{p_0} , Remark 3.2 and Lemma 2.7 yield

$$(5) \quad \begin{aligned} \|[M(m_j)]^\theta\|_{L^{p_0/\theta}(2\sqrt{n}Q_j)} &= \|M(m_j)\|_{L^{p_0}(2\sqrt{n}Q_j)}^\theta \lesssim \|m_j\|_{L^{p_0}(\mathbb{R}^n)}^\theta \\ &\lesssim \frac{|Q_j|^{\frac{\theta}{p_0}}}{\|\chi_{Q_j}\|_{L_\omega^{p(\cdot)/\theta}}^\theta} \lesssim \frac{|2\sqrt{n}Q_j|^{\frac{\theta}{p_0}}}{\|\chi_{2\sqrt{n}Q_j}\|_{L_\omega^{p(\cdot)/\theta}}^\theta}, \end{aligned}$$

where $0 < \theta < 1$ and $\frac{1}{\theta} \in \mathbb{S}_{\omega, p(\cdot)}$, and $p_0 > \theta \left(\kappa_{\omega, p(\cdot)}^{1/\theta}\right)'$. By applying the θ -inequality and [5, Lemma 5.4] with $b_j = (\chi_{2\sqrt{n}Q_j} \cdot M(m_j))^\theta$, (5) and $A_j = \|\chi_{2\sqrt{n}Q_j}\|_{L_\omega^{p(\cdot)/\theta}}^{-1}$, we get

$$J_1 \lesssim \left\| \sum_j (|\lambda_j| \chi_{2\sqrt{n}Q_j} \cdot M(m_j))^\theta \right\|_{L_\omega^{p(\cdot)/\theta}}^{1/\theta} \lesssim \left\| \sum_j \left(\frac{|\lambda_j|}{\|\chi_{2\sqrt{n}Q_j}\|_{L_\omega^{p(\cdot)}}} \right)^\theta \chi_{2\sqrt{n}Q_j} \right\|_{L_\omega^{p(\cdot)/\theta}}^{1/\theta}.$$

Being $\chi_{2\sqrt{n}Q_j} \leq M(\chi_{Q_j})^2$, by Lemma 2.7 and [5, Theorem 3.1], we have

$$J_1 \lesssim \left\| \left\{ \sum_j \left(\frac{|\lambda_j|^{\theta/2}}{\|\chi_{Q_j}\|_{L_\omega^{p(\cdot)}}^{\theta/2}} M(\chi_{Q_j}) \right)^2 \right\}^{1/2} \right\|_{L_\omega^{2p(\cdot)/\theta}}^{2/\theta} \lesssim \left\| \sum_j \left(\frac{|\lambda_j|}{\|\chi_{Q_j}\|_{L_\omega^{p(\cdot)}}} \right)^\theta \chi_{Q_j} \right\|_{L_\omega^{p(\cdot)/\theta}}^{1/\theta}.$$

So,

$$(6) \quad J_1 \lesssim \mathcal{A}(\{\lambda_j\}_{j=1}^\infty, \{Q_j\}_{j=1}^\infty, p(\cdot), \omega, \theta) < \infty.$$

To estimate J_2 , we write $r = \frac{n + d_{p(\cdot)} + 1}{n}$. Thus

$$J_2 \lesssim \left\| \left(\sum_j |\lambda_j| \frac{[M(\chi_{Q_j})(\cdot)]^r}{\|\chi_{Q_j}\|_{L_\omega^{p(\cdot)}}^r} \right)^{1/r} \right\|_{L_\omega^{rp(\cdot)/r}}^r.$$

Since

$$r = \frac{n + d_{p(\cdot)} + 1}{n} = \frac{n + \lfloor ns_{\omega, p(\cdot)} - n \rfloor + 1}{n} > s_{\omega, p(\cdot)},$$

to apply again [5, Theorem 3.1] followed by the θ -inequality we obtain

$$\begin{aligned} J_2 &\lesssim \left\| \sum_j \left(\frac{|\lambda_j|}{\|\chi_{Q_j}\|_{L_\omega^{p(\cdot)}}} \chi_{Q_j} \right)^{1/r} \right\|_{L_\omega^{rp(\cdot)/r}}^r = \left\| \sum_j \frac{|\lambda_j|}{\|\chi_{Q_j}\|_{L_\omega^{p(\cdot)}}} \chi_{Q_j} \right\|_{L_\omega^{p(\cdot)}} \\ &\lesssim \left\| \sum_j \left(\frac{|\lambda_j|}{\|\chi_{Q_j}\|_{L_\omega^{p(\cdot)}}} \right)^\theta \chi_{Q_j} \right\|_{L_\omega^{p(\cdot)/\theta}}^{1/\theta}. \end{aligned}$$

Hence,

$$(7) \quad J_2 \lesssim \mathcal{A}(\{\lambda_j\}_{j=1}^\infty, \{Q_j\}_{j=1}^\infty, p(\cdot), \omega, \theta) < \infty.$$

Finally, (6) and (7) give (3) and, with that, $g \in H_\omega^{p(\cdot)}(\mathbb{R}^n)$. \square

4 Weighted variable estimates for singular integrals

Let $\Omega \in C^\infty(S^{n-1})$ with $\int_{S^{n-1}} \Omega(u) d\sigma(u) = 0$. We define the operator T by

$$(8) \quad Tf(x) = \lim_{\epsilon \rightarrow 0^+} \int_{|y| > \epsilon} \frac{\Omega(y/|y|)}{|y|^n} f(x-y) dy, \quad x \in \mathbb{R}^n.$$

It is well known that the operator T is bounded on $L^{p_0}(\mathbb{R}^n)$ for all $1 < p_0 < +\infty$ and of weak-type $(1, 1)$ (see e.g. [13]).

We start studying the behavior of the operator T on atoms. Then, we prove the boundedness of T on $H_\omega^{p(\cdot)}$.

PROPOSITION 4.1. *Let T be the operator given by (8) and let $p(\cdot) : \mathbb{R}^n \rightarrow (0, \infty)$ be a measurable function with $0 < p_- \leq p_+ < \infty$, and $\omega \in \mathcal{W}_{p(\cdot)}$. Then, for some universal constant $C > 0$, $C(Ta)(\cdot)$ is a $\omega - (p(\cdot), p_0, [ns_{\omega, p(\cdot)} - n])$ molecule for each $\omega - (p(\cdot), p_0, n + 2[ns_{\omega, p(\cdot)} - n] + 2)$ atom $a(\cdot)$.*

PROOF. It is well known that $\widehat{Tf}(\xi) = m(\xi)\widehat{f}(\xi)$, where the multiplier m is homogeneous of degree 0 and is indefinitely differentiable on $\mathbb{R}^n \setminus \{0\}$ (see e.g. [13]). Moreover, for $K(y) = \frac{\Omega(y/|y|)}{|y|^n}$ we have

$$(9) \quad |\partial_y^\alpha K(y)| \leq C|y|^{-n-|\alpha|}, \quad \text{for all } y \neq 0 \text{ and all multi-index } \alpha.$$

Let $p_0 > 1$ and $d_{p(\cdot)} = [ns_{\omega, p(\cdot)} - n]$. Given a $w - (p(\cdot), p_0, n + 2d_{p(\cdot)} + 2)$ atom $a(\cdot)$ with support in the cube $Q = Q(z, r)$ (with side length $\ell(Q) = r$), we have that

$$(10) \quad \|Ta\|_{L^{p_0}(2\sqrt{n}Q)} \leq C\|a\|_{L^{p_0}} \leq C \frac{|Q|^{1/p_0}}{\|\chi_Q\|_{L_\omega^{p(\cdot)}}},$$

since T is bounded on $L^{p_0}(\mathbb{R}^n)$. In view of the moment condition of $a(\cdot)$ we obtain

$$Ta(x) = \int_Q K(x-y)a(y)dy = \int_Q [K(x-y) - q_{n+2d+2}(x, y)]a(y)dy, \quad x \notin 2\sqrt{n}Q$$

where $y \rightarrow q_{n+2d_{p(\cdot)}+2}(x, y)$ is the degree $n + 2d_{p(\cdot)} + 2$ Taylor polynomial of the function $y \rightarrow K(x-y)$ expanded around z . From the estimate in (9), and the standard estimate of the remainder term of the Taylor expansion, there exists ξ between y and z such that

$$(11) \quad |Ta(x)| \leq C\|a\|_1 \frac{\ell(Q)^{n+2d_{p(\cdot)}+3}}{|x-\xi|^{2n+2d_{p(\cdot)}+3}} \leq C \frac{\ell(Q)^{2n+2d_{p(\cdot)}+3}}{\|\chi_Q\|_{L_\omega^{p(\cdot)}}} |x-z|^{-2n-2d_{p(\cdot)}-3}, \quad x \notin 2\sqrt{n}Q,$$

this inequality and a simple computation allow us to obtain

$$(12) \quad |Ta(x)| \leq C\|\chi_Q\|_{L_\omega^{p(\cdot)}}^{-1} \left(1 + \frac{|x-z|}{\ell(Q)}\right)^{-2n-2d_{p(\cdot)}-3}, \quad \text{for all } x \notin 2\sqrt{n}Q.$$

From the estimate in (11) we obtain that the function $x \rightarrow x^\alpha Ta(x)$ belongs to $L^1(\mathbb{R}^n)$ for each $|\alpha| \leq d_{p(\cdot)}$, so

$$\begin{aligned} |((-2\pi ix)^\alpha Ta)^\wedge(\xi)| &= |\partial_\xi^\alpha(m(\xi)\widehat{a}(\xi))| = \left| \sum_{\beta \leq \alpha} c_{\alpha,\beta} (\partial_\xi^{\alpha-\beta} m)(\xi) (\partial_\xi^\beta \widehat{a})(\xi) \right| \\ &= \left| \sum_{\beta \leq \alpha} c_{\alpha,\beta} (\partial_\xi^{\alpha-\beta} m)(\xi) ((-2\pi ix)^\beta a)^\wedge(\xi) \right|, \end{aligned}$$

from the homogeneity of the function $\partial_\xi^{\alpha-\beta} m$ we obtain that

$$(13) \quad |((-2\pi ix)^\alpha Ta)^\wedge(\xi)| \leq C \sum_{\beta \leq \alpha} |c_{\alpha,\beta}| \frac{|((-2\pi ix)^\beta a)^\wedge(\xi)|}{|\xi|^{|\alpha|-\beta}}, \quad \xi \neq 0.$$

Since $\lim_{\xi \rightarrow 0} \frac{|((-2\pi ix)^\beta a)^\wedge(\xi)|}{|\xi|^{|\alpha|-\beta}} = 0$ for each $\beta \leq \alpha$ (see 5.4, pp. 128, in [14]), taking the limit as $\xi \rightarrow 0$ at (13), we get

$$(14) \quad \int_{\mathbb{R}^n} (-2\pi ix)^\alpha Ta(x) dx = ((-2\pi ix)^\alpha Ta)^\wedge(0) = 0, \quad \text{for all } |\alpha| \leq d_{p(\cdot)}.$$

From (10), (12) and (14) it follows that there exists an universal constant $C > 0$ such that $C(Ta)(\cdot)$ is a $w - (p(\cdot), p_0, d_{p(\cdot)})$ molecule if $a(\cdot)$ is a $w - (p(\cdot), p_0, n + 2d_{p(\cdot)} + 2)$ atom. \square

THEOREM 4.2. *Let $p(\cdot) \in \mathcal{P}^{\log}(\mathbb{R}^n)$, $\omega \in \mathcal{W}_{p(\cdot)}$ and let T be the singular integral operator given by (8). Then T can be extended to a bounded operator on $H_\omega^{p(\cdot)}(\mathbb{R}^n)$.*

PROOF. Given $\omega \in \mathcal{W}_{p(\cdot)}$, by Definition 2.2, there exists $0 < \theta < 1$ such that $\frac{1}{\theta} \in \mathbb{S}_{\omega, p(\cdot)}$. Let $p_0 > \max \left\{ \theta \left(\kappa_{\omega, p(\cdot)}^{1/\theta} \right)', 1 \right\}$. Given $f \in \mathcal{S}_0(\mathbb{R}^n)$, by Theorem 2.6, there exist a sequence of real numbers $\{\lambda_j\}_{j=1}^\infty$, a sequence of cubes $\{Q_j\}_{j=1}^\infty$, and $\omega - (p(\cdot), p_0, n + 2d_{p(\cdot)} + 2)$ atoms a_j supported on Q_j , such that $f = \sum_{j=1}^\infty \lambda_j a_j$ converges in $L^{p_0}(\mathbb{R}^n)$ and

$$(15) \quad \mathcal{A}(\{\lambda_j\}_{j=1}^\infty, \{Q_j\}_{j=1}^\infty, p(\cdot), \omega, \theta) \lesssim \|f\|_{H_\omega^{p(\cdot)}} < \infty.$$

Since $f = \sum_{j=1}^\infty \lambda_j a_j$ converges in $L^{p_0}(\mathbb{R}^n)$ and T is bounded on L^{p_0} , we have that

$$(16) \quad Tf = \sum_{j=1}^\infty \lambda_j Ta_j \quad \text{in } \mathcal{S}'(\mathbb{R}^n).$$

Now, by (16) and (15) above and Proposition 4.1, we can apply Theorem 3.4 to obtain

$$\|Tf\|_{H_\omega^{p(\cdot)}} \lesssim \mathcal{A}(\{\lambda_j\}_{j=1}^\infty, \{Q_j\}_{j=1}^\infty, p(\cdot), \omega, \theta) \lesssim \|f\|_{H_\omega^{p(\cdot)}},$$

for all $f \in \mathcal{S}_0(\mathbb{R}^n)$. Finally, the theorem follows from the density of $\mathcal{S}_0(\mathbb{R}^n)$ in $H_\omega^{p(\cdot)}(\mathbb{R}^n)$ (see Proposition 2.8). \square

In particular, if $p(\cdot) \in \mathcal{P}^{\log}$ and $\omega \in \mathcal{W}_{p(\cdot)}$, then Hilbert transform and Riesz transforms admit a continuous extension on $H_\omega^{p(\cdot)}(\mathbb{R})$ and on $H_\omega^{p(\cdot)}(\mathbb{R}^n)$ respectively.

5 Weighted variable estimates for Riesz potential

Let $0 < \alpha < n$. The *Riesz potential* of order α is the operator I_α defined, say on $\mathcal{S}(\mathbb{R}^n)$, by

$$(17) \quad I_\alpha f(x) = \int_{\mathbb{R}^n} f(y) |x - y|^{\alpha-n} dy, \quad x \in \mathbb{R}^n.$$

In [11], the author proved that the operator I_α extends to a bounded operator $H_\omega^{p(\cdot)}(\mathbb{R}^n) \rightarrow L_\omega^{q(\cdot)}(\mathbb{R}^n)$, for $\frac{1}{p(\cdot)} := \frac{1}{q(\cdot)} + \frac{\alpha}{n}$; under the following assumptions:

- 1) $q(\cdot) \in \mathcal{P}^{\log}(\mathbb{R}^n)$ and $\omega \in \mathcal{W}_{q(\cdot)}$;
- 2) for every cube $Q \subset \mathbb{R}^n$, $\|\chi_Q\|_{L_\omega^{q(\cdot)}} \approx |Q|^{-\alpha/n} \|\chi_Q\|_{L^{p(\cdot)}}$.

We observe that if $q(\cdot) \in \mathcal{P}^{\log}(\mathbb{R}^n)$ and $\omega \equiv 1$, then the condition 2) holds. This was proved in [12]. In [11], the author gave non trivial examples of power weights satisfying 2). So, the condition 2) is an admissible hypothesis.

Next, assuming the conditions 1) and 2) above, we will prove that Riesz potential I_α extends to a bounded operator $H_\omega^{p(\cdot)}(\mathbb{R}^n) \rightarrow H_\omega^{q(\cdot)}(\mathbb{R}^n)$. For them, we will first show that I_α maps atoms into molecules.

PROPOSITION 5.1. *Let $0 < \alpha < n$, $q(\cdot) : \mathbb{R}^n \rightarrow (0, \infty)$ be a measurable function with $0 < q_- \leq q_+ < \infty$, and $\omega \in \mathcal{W}_{q(\cdot)}$. If $\frac{1}{p(\cdot)} := \frac{1}{q(\cdot)} + \frac{\alpha}{n}$ and $\|\chi_Q\|_{L_\omega^{q(\cdot)}} \approx |Q|^{-\alpha/n} \|\chi_Q\|_{L^{p(\cdot)}}$ for every cube Q , then, for some universal constant $C > 0$, $C(I_\alpha a)(\cdot)$ is a $\omega - (q(\cdot), q_0, [ns_{\omega, q(\cdot)} - n])$ molecule for each $\omega - (p(\cdot), p_0, 2[ns_{\omega, q(\cdot)} - n] + [\alpha] + 3 + n)$ atom $a(\cdot)$, where $q_0 > \frac{n}{n-\alpha}$ and $\frac{1}{p_0} := \frac{1}{q_0} + \frac{\alpha}{n}$.*

PROOF. By [11, Proposition 3.1], we have that $\mathcal{W}_{q(\cdot)} \subset \mathcal{W}_{p(\cdot)}$ and $s_{\omega, p(\cdot)} \leq s_{\omega, q(\cdot)} + \frac{\alpha}{n}$. Let $d_{q(\cdot)} := [ns_{\omega, q(\cdot)} - n]$ and $N = 2d_{q(\cdot)} + [\alpha] + 3 + n$, we observe that $N \geq d_{p(\cdot)}$. Fix $q_0 > \frac{n}{n-\alpha}$ and we put $\frac{1}{p_0} := \frac{1}{q_0} + \frac{\alpha}{n}$. Given a $\omega - (p(\cdot), p_0, N)$ atom $a(\cdot)$ supported on a cube $Q = Q(z, r)$, we will show that there exists an universal constant $C > 0$ such that $CI_\alpha a(\cdot)$ is a $\omega - (q(\cdot), q_0, d_{q(\cdot)})$ molecule centered at Q . Indeed,

m_1) by Sobolev's theorem, the condition a_1) of the atom $a(\cdot)$, and since $\|\chi_Q\|_{L_\omega^{q(\cdot)}} \approx |Q|^{-\alpha/n} \|\chi_Q\|_{L^{p(\cdot)}}$ for every cube Q , we have

$$\|I_\alpha a\|_{L^{q_0}(2\sqrt{n}Q)} \lesssim \|a\|_{L^{p_0}} \leq \frac{|Q|^{\frac{1}{p_0}}}{\|\chi_Q\|_{L_\omega^{p(\cdot)}}} \lesssim \frac{|Q|^{\frac{1}{q_0}}}{\|\chi_Q\|_{L_\omega^{q(\cdot)}}};$$

m_2) by doing use of the conditions a_3) and a_2) of the atom $a(\cdot)$, as in the estimate

(24) obtained in [11, Theorem 5.1], we get, for x outside $2\sqrt{n}Q$, that

$$\begin{aligned}
|(I_\alpha a)(x)| &\lesssim \frac{\ell(Q)^{N+1}}{|x-z|^{n-\alpha+N+1}} \|a\|_{L^1} \\
&\lesssim \frac{1}{\|\chi_Q\|_{L_\omega^{q(\cdot)}}} \left(\frac{\ell(Q)}{|x-z|} \right)^{n-\alpha+N+1} \\
&\lesssim \frac{1}{\|\chi_Q\|_{L_\omega^{q(\cdot)}}} \left(\frac{\ell(Q)}{\ell(Q)+|x-z|} \right)^{2n+2d_{q(\cdot)}+3} \\
&= \frac{1}{\|\chi_Q\|_{L_\omega^{q(\cdot)}}} \left(1 + \frac{|x-z|}{\ell(Q)} \right)^{-2n-2d_{q(\cdot)}-3},
\end{aligned}$$

where the third inequality follows since x outside $2\sqrt{n}Q$ implies $\ell(Q)+|x-z| < 2|x-z|$;

m_3) the moment condition was proved by Taibleson and Weiss in [15].

Thus $I_\alpha a$ satisfies Definition 3.1 with an universal implicit constant. This completes the proof. \square

THEOREM 5.2. *Let $0 < \alpha < n$, $q(\cdot) \in \mathcal{P}^{\log}(\mathbb{R}^n)$ with $0 < q_- \leq q_+ < \infty$ and $\omega \in \mathcal{W}_{q(\cdot)}$. If $\frac{1}{p(\cdot)} := \frac{1}{q(\cdot)} + \frac{\alpha}{n}$ and $\|\chi_Q\|_{L_\omega^{q(\cdot)}} \approx |Q|^{-\alpha/n} \|\chi_Q\|_{L_\omega^{p(\cdot)}}$ for every cube Q , then the Riesz potential I_α given by (17) can be extended to a bounded operator $H_\omega^{p(\cdot)}(\mathbb{R}^n) \rightarrow H_\omega^{q(\cdot)}(\mathbb{R}^n)$.*

PROOF. Let $\omega \in \mathcal{W}_{q(\cdot)}$, by Definition 2.2, there exists $0 < \theta < 1$ such that $\frac{1}{\theta} \in \mathbb{S}_{\omega, q(\cdot)}$. Now, we take $q_0 > \max \left\{ \theta \left(\kappa_{\omega, q(\cdot)}^{1/\theta} \right)', \frac{n}{n-\alpha} \right\}$, and define $\frac{1}{p_0} := \frac{1}{q_0} + \frac{\alpha}{n}$. By [11, Proposition 3.1], we have that $\mathcal{W}_{q(\cdot)} \subset \mathcal{W}_{p(\cdot)}$ and $s_{\omega, p(\cdot)} \leq s_{\omega, q(\cdot)} + \frac{\alpha}{n}$. So, given $f \in \mathcal{S}_0(\mathbb{R}^n)$, by Theorem 2.6 there exist a sequence of real numbers $\{\lambda_j\}_{j=1}^\infty$, a sequence of cubes $\{Q_j\}_{j=1}^\infty$, and $\omega - (p(\cdot), p_0, 2[n s_{\omega, q(\cdot)} - n] + [\alpha] + 3 + n)$ atoms a_j supported on Q_j , satisfying

$$(18) \quad \mathcal{A}(\{\lambda_j\}_{j=1}^\infty, \{Q_j\}_{j=1}^\infty, p(\cdot), \omega, \theta) \lesssim \|f\|_{H_\omega^{p(\cdot)}} < \infty,$$

and $f = \sum_{j=1}^\infty \lambda_j a_j$ converges in $L^{p_0}(\mathbb{R}^n)$. Then, by Sobolev's Theorem, $I_\alpha f = \sum_{j=1}^\infty \lambda_j I_\alpha(a_j)$ in $L^{\frac{np_0}{n-\alpha p_0}}(\mathbb{R}^n)$ thus

$$(19) \quad I_\alpha f = \sum_{j=1}^\infty \lambda_j I_\alpha(a_j) \text{ in } \mathcal{S}'(\mathbb{R}^n).$$

Now, by (19), (18) and Proposition 2.9, and Proposition 5.1, we can apply Theorem 3.4 to obtain

$$\begin{aligned}
\|I_\alpha f\|_{H_\omega^{q(\cdot)}} &\lesssim \mathcal{A}(\{\lambda_j\}_{j=1}^\infty, \{Q_j\}_{j=1}^\infty, q(\cdot), \omega, \theta) \\
&\lesssim \mathcal{A}(\{\lambda_j\}_{j=1}^\infty, \{Q_j\}_{j=1}^\infty, p(\cdot), \omega, \theta) \\
&\lesssim \|f\|_{H_\omega^{p(\cdot)}},
\end{aligned}$$

for all $f \in \mathcal{S}_0(\mathbb{R}^n)$. Finally, since $p(\cdot) \in \mathcal{P}^{\log}(\mathbb{R}^n)$, $0 < p_- \leq p_+ < \infty$ and $\omega \in \mathcal{W}_{p(\cdot)}$, the theorem follows from the density of $\mathcal{S}_0(\mathbb{R}^n)$ in $H_\omega^{p(\cdot)}(\mathbb{R}^n)$ (see Proposition 2.8). \square

References

- [1] R. COIFMAN, A real characterization of H^p , *Studia Math.* 51 (1974), 269-274.
- [2] R. COIFMAN and G. WEISS, Extensions of Hardy spaces and their use in analysis, *Bull. Amer. Math. Soc.* 83 (1977), 569-645.
- [3] D. CRUZ-URIBE and D. WANG, Variable Hardy Spaces, *Indiana Univ. Math. J.* 63 (2) (2014), 447-493.
- [4] C. FEFFERMAN and E. STEIN, H^p spaces of several variables, *Acta Math.* 129 (1972), 137-193.
- [5] K.-P. HO, Atomic decompositions of weighted Hardy spaces with variable exponents, *Tohoku Math. J.* 69 (2) (2017), no. 3, 383-413.
- [6] K.-P. HO, Sublinear operators on weighted Hardy spaces with variable exponents, *Forum Math.* 31 (3) (2019), 607-617.
- [7] R. H. LATTER, A characterization of $H^p(\mathbb{R}^n)$ in terms of atoms, *Studia Math.* 62 (1978), 93-101.
- [8] E. NAKAI and Y. SAWANO, Hardy spaces with variable exponents and generalized Campanato spaces, *J. Funct. Anal.* 262 (2012), 3665-3748.
- [9] P. ROCHA, A note on Hardy spaces and bounded linear operators, *Georgian Math. J.* 25 (1) (2018), 73-76.
- [10] P. ROCHA, Boundedness of generalized Riesz potentials on the variable Hardy spaces, *J. Aust. Math. Soc.* 104 (2018), 255-273.
- [11] P. ROCHA, Inequalities for weighted spaces with variable exponents, (2022), Preprint: arXiv:2211.12218
- [12] P. ROCHA and M. URUIUOLO, Fractional type integral operators on variable Hardy spaces, *Acta Math. Hung.* 143 (2) (2014), 502-514.
- [13] E. STEIN, *Singular Integrals and Differentiability Properties of Functions*, Princeton Univ. Press, Princeton, NJ, 1970.
- [14] E. STEIN, *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals*, Princeton University Press, 1993.

- [15] M. H. TAIBLESON and G. WEISS, The molecular characterization of certain Hardy spaces, *Astérisque* 77 (1980), 67-149.

DEPARTAMENTO DE MATEMÁTICA
UNIVERSIDAD NACIONAL DEL SUR
BAHÍA BLANCA, 8000 BUENOS AIRES
ARGENTINA

E-mail address: pablo.rocha@uns.edu.ar