

# ENTROPY SOLUTIONS TO THE DIRICHLET PROBLEM FOR NONLINEAR DIFFUSION EQUATIONS WITH CONSERVATIVE NOISE

KAI DU, RUOYANG LIU, AND YUXING WANG

ABSTRACT. Motivated by porous medium equations with randomly perturbed velocity field, this paper considers a class of nonlinear degenerate diffusion equations with nonlinear conservative noise in bounded domains. The existence, uniqueness and  $L_1$ -stability of non-negative entropy solutions under the homogeneous Dirichlet boundary condition are proved. The approach combines Kruzhkov's doubling variables technique with a revised strong entropy condition that is automatically satisfied by the solutions of approximate equations.

## 1. INTRODUCTION

This paper is concerned with the Dirichlet problem for nonlinear diffusion equation

$$(1.1) \quad \begin{aligned} du(t, x) &= (\Delta\Phi(u) + \nabla \cdot G(x, u) + F(x, u))dt \\ &\quad + (\nabla \cdot \sigma^k(x, u)) \circ dW^k(t), \quad (t, x) \in (0, T) \times D; \\ u(0, x) &= \xi(x), \quad x \in D; \\ u(t, x) &= 0, \quad (t, x) \in [0, T] \times \partial D, \end{aligned}$$

where  $D$  is a bounded domain in  $\mathbb{R}^d$  with smooth boundary, and  $\Phi : \mathbb{R} \rightarrow \mathbb{R}$  is a monotone function. The noise  $\{W^k\}_{k \in \mathbb{N}}$  is a sequence of independent standard Brownian motions defined on a complete filtered probability space  $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$ , and the stochastic integral is in the Stratonovich sense. The Einstein summation convention is used throughout this paper.

A typical example of (1.1) and also our primary motivation of this paper is the porous medium equation in random environment, which describes the flow of an ideal gas in a homogeneous porous medium: let  $u$  be the gas density that satisfies

$$(1.2) \quad \varepsilon \partial_t u + \nabla \cdot (u\mathbf{V}) = 0,$$

where  $\varepsilon \in (0, 1)$  is the porosity of the medium, and  $\mathbf{V}$  is the randomly perturbed velocity field of the form

$$(1.3) \quad \mathbf{V} = \mathbf{V}_0 + \sigma^k \circ \dot{W}^k,$$

where  $\mathbf{V}_0$  is derived from Darcy's law

$$\mu \mathbf{V}_0 = -k \nabla p = -k \nabla (p_0 u^{m-1}).$$

Then, we can informally derive a stochastic porous medium equation

$$du(t, x) = c \Delta u^m dt + \varepsilon^{-1} \nabla \cdot (u \sigma^k) \circ dW^k(t).$$

For more details of the model, we refer to [Váz07] and references therein. Moreover, equations of type (1.1) also arises as limits of interacting particle systems driven by common noise from mean field models [LL06a, LL06b, LL07], and as simplified models of fluctuations in non-equilibrium

---

2010 *Mathematics Subject Classification.* 60H15; 35R60; 35K59.

*Key words and phrases.* nonlinear diffusion equation, entropy solution, Dirichlet problem, stochastic porous medium equation, conservative noise.

statistical physics [DSZ16]; for more applications we refer to [FG19, DG20] and the references therein.

The well-posedness of stochastic nonlinear diffusion equations has been investigated in various frameworks, for example, the variational approach in the space  $H^{-1}$  (see [BDPR08, BDPR16, Cio20] etc.), the kinetic formulation (cf. [DHV<sup>+</sup>16, GS17, GH18, FG19, FG21] etc.), and the entropy formulation (cf. [BVW15, DGG19, DG20, DGT20] etc.). In general, the entropy approach may require lower the regularity of the diffusion coefficients and treat a more general class of non-linearity  $\Phi$  than the kinetic formulation according to [FG19, DG20]. In this paper, we adopt the entropy approach to study (1.1).

The notion of entropy solution was proposed for nonlinear PDEs in 1970s to tackle the difficulty that the weak solution may not be unique (see, for example, [Kru70, Ser99, ABK00, DD05] for scalar conservation law and [Car99] for degenerate parabolic equations). The entropy solution discriminates the physical admissible solution and maintains the uniqueness theoretically.

The entropy formulation has been naturally extended to nonlinear SPDEs. Many works have been done for first order equations (see, for example, [Kim03] for additive noise and [FN08, VW09, CDK12, BVW12, BVW14] for multiplicative noise). For second order equations, most works considered the Cauchy problem in the whole space or a torus; for instance, [BVW15] studies the Cauchy problem of parabolic-hyperbolic SPDEs with the noise term  $\sigma(x, u)dW$  and Lipschitz function  $\Phi$ . Stochastic nonlinear diffusion equations in a torus is considered in [DGG19] for the noise term  $\sigma(x, u)dW$  and in [DG20] for the conservative noise  $\nabla \cdot \sigma(x, u)dW$ . The obstacle problem with the noise term  $\sigma(x, u)dW$  in a torus is introduced in [LT21].

The results on the Dirichlet problem for stochastic nonlinear diffusion equations are relatively few. The recent papers [BGV20, Hen21] adopts the variational approach from [RRW07] with the monotone condition and affine noise, thus not covering our setting; [DGT20] considers entropy solutions of stochastic porous medium equations with the noise term  $\sigma(x, u)dW$ . The preprint [Cli22] is quite relevant to our paper, studying kinetic solutions to the Dirichlet problem for porous medium equations with nonlinear gradient noise driven by rough path. Thanks to the entropy approach, our result is built on the same regularity conditions with the existing work on the Cauchy problem in torus (cf. [DG20]), and do not require extra technical assumptions like [Cli22, Condition (2.4)] that prevents the space characteristics from escaping the domain.

The strategy of the proof of our main result (Theorem 2.6) basically follows from [DGG19, DGT20, DG20], combining the method of strong entropy condition (called the  $(\star)$ -property in this paper) and Kruzhkov's doubling variables technique (cf. [Kru70]). The notion of strong entropy condition was introduced by [FN08] to tackle the uniqueness issue of stochastic scalar conservation laws. The strategy can be summarized to two steps:

- (1)  $L_1$ -estimates: to derive an estimate for  $\mathbb{E}\|u(t, \cdot) - \tilde{u}(t, \cdot)\|_{L_1(D)}$ , providing one of the entropy solutions  $u$  and  $\tilde{u}$  (with different initial data) satisfies the  $(\star)$ -property.
- (2) Approximation: to construct a sequence of non-degenerate equations whose solutions has the  $(\star)$ -property and converge to the entropy solution of (1.1).

Kruzhkov's doubling variables technique plays a key role in the proof of  $L^1$ -estimates.

The new difficulty arising in the Dirichlet problem, comparing to the Cauchy problem (cf. [DG20]), is how to deal with the boundary integral terms that may emerge when applying the divergence theorem in the proof of  $L_1$ -estimates. In the paper [DGT20] where the noise term is  $\sigma(x, u)dW$ , a weighted space with a weight function  $w \in H_0^1$  satisfying  $\Delta w = -1$  is introduced to eliminate the boundary terms. Unfortunately, in the case of gradient noise, there will be a new "trouble" term  $|(u - \tilde{u})\nabla w|$  appearing in the estimate, which cannot be dominated by any "good" terms like  $|u - \tilde{u}|w$ .

There are three key points in our approach to overcome the above difficulty: i) to expand the set of test functions in the definition of entropy solution (see Definition 2.2), ii) to refine the

strong entropy condition, and iii) to construct subtly a pair of the convex function  $\eta$  and the test function  $\phi$  to avoid the appearance of boundary terms. Specifically, when applying Kruzhkov's doubling variables technique to our problem, we have to estimate both  $(\tilde{u}(s, x) - u(t, y))^+$  and  $(u(t, x) - \tilde{u}(s, y))^+$  rather than  $|u(t, x) - \tilde{u}(s, y)|$  as in [DGG19, DGT20, DG20]. Inspired by [Car99, BVW14], we make use of a partition of unity and shifted mollifiers to keep the test function in  $C_c^\infty(D)$  with respect to the variable  $y$ ; and for the variable  $x$ , we choose a sequence of smooth convex functions to approach  $\eta(r) = r^+$ . Those carefully chosen functions along our modified definition of entropy solution and the refined strong entropy condition let the boundary terms vanish. More specific details are given in Remarks 3.5, 3.11, 3.14 and 5.1. It is worth noting that our approach avoids the involvement of weight functions and consider the  $L_1$ -estimates in standard Sobolev spaces, and may also apply to nonlinear diffusion equations with the noise term  $\sigma(x, u)dW$  to obtain an  $L_1$ -estimates without weight.

This paper is organized as follows. Section 2 describes the entropy formulation and presents the main theorem. Section 3, which is the main part of this paper, introduces a refined strong entropy condition and derives  $L_1$ -estimates for the difference of two entropy solutions. Section 4 constructs the approximate equations and proves the strong entropy condition of their solutions as well as their solvability. Section 5 completes the proof of the main theorem. Two auxiliary lemmas are proved in the final section.

We conclude the introduction with some notation. Fix  $T > 0$ . Define  $\Omega_T := \Omega \times [0, T]$  and  $D_T := [0, T] \times D$ . Define  $|D|$  and  $\overline{D}$  as the volume and closure of  $D$ , respectively.  $L_p$  and  $H_p^k$  are the usual Lebesgue spaces and Sobolev spaces. Denote by  $H_{p,0}^k$  the closure of  $C_c^\infty$  in  $H_p^k$ . When  $p = 2$ , we simplify the notation by  $H^k := H_2^k$  and  $H_0^k := H_{2,0}^k$ . Moreover, if a function space is given on  $\Omega$  or  $\Omega_T$ , we understand it to be defined with respect to  $\mathcal{F}_T$  and the predictable  $\sigma$ -field, respectively. Let  $E$  be a Banach space and  $U = D$  or  $\mathbb{R}$ . For a function  $f : U \rightarrow E$ , we define

$$[f]_{C^\alpha(U;E)} := \sup_{x,y \in U, x \neq y} \frac{\|f(x) - f(y)\|_E}{|x - y|^\alpha}, \quad \alpha \in (0, 1],$$

$$\|f\|_{C^\alpha(U;E)} := [f]_{C^\alpha(U;E)} + \sup_{x \in U} \|f(x)\|_E.$$

We define a non-negative smooth mollifier  $\rho : \mathbb{R} \rightarrow \mathbb{R}$ , such that  $\text{supp}(\rho) \subset (0, 1)$ ,  $\rho \leq 2$  and  $\int_{\mathbb{R}} \rho(r) dr = 1$ . For  $\delta > 0$ , we set  $\rho_\delta(r) := \delta^{-1} \rho(\delta^{-1} r)$  as a sequence of mollifiers.

## 2. ENTROPY FORMULATION AND MAIN RESULTS

First of all, we rewrite (1.1) into an Itô form (the notation follows from [DG20]):

$$(2.1) \quad \begin{aligned} du(t, x) &= [\Delta \Phi(u) + \partial_{x_i}(a^{ij}(x, u) \partial_{x_j} u + b^i(x, u) + f^i(x, u)) + F(x, u)] dt \\ &\quad + (\nabla \cdot \sigma^k(x, u)) dW^k(t), \quad (t, x) \in (0, T) \times D; \\ u(0, x) &= \xi(x), \quad x \in D; \\ u(t, x) &= 0, \quad (t, x) \in [0, T] \times \partial D, \end{aligned}$$

where  $i, j = 1, \dots, d$  and

$$\begin{aligned} a^{ij}(x, r) &= \frac{1}{2} \sigma_r^{ik}(x, r) \sigma_r^{jk}(x, r), \quad b^i(x, r) = \sigma_r^{ik}(x, r) \sigma_{x_j}^{jk}(x, r), \\ f^i(x, r) &= G^i(x, r) - \frac{1}{2} b^i(x, r). \end{aligned}$$

We denote by  $\Pi(\Phi, \xi)$  the Dirichlet problem (2.1) with given  $\Phi$  and  $\xi$ .

Throughout this paper, we denote

$$\mathbf{a}(r) := \sqrt{\Phi'(r)};$$

for a function  $g : D \times \mathbb{R} \rightarrow \mathbb{R}$ , we use the notation

$$\llbracket g \rrbracket(x, r) := \int_0^r g(x, s) ds,$$

and drop  $x$  in the above if  $g$  does not depend on  $x \in D$ . Our condition on the non-linearity  $\Phi$  is the same with [DGG19, DGT20, DG20].

**Assumption 2.1.**  $\Phi : \mathbb{R} \rightarrow \mathbb{R}$  is odd, differentiable, and strictly increasing. With  $\mathbf{a}(r) = \sqrt{\Phi'(r)}$ , there exist constants  $m > 1$  and  $K > 0$  such that

$$(2.2) \quad \begin{aligned} |\mathbf{a}(0)| &\leq K, \quad |\mathbf{a}'(r)| \leq K|r|^{\frac{m-3}{2}} \mathbf{1}_{r \neq 0}, \quad \mathbf{a}(r) \geq K^{-1} \mathbf{1}_{|r| \geq 1}, \\ \llbracket \mathbf{a} \rrbracket(r) - \llbracket \mathbf{a} \rrbracket(s) &\geq \begin{cases} K^{-1}|r-s|, & \text{if } |r| \vee |s| \geq 1, \\ K^{-1}|r-s|^{\frac{m+1}{2}}, & \text{if } |r| \vee |s| < 1. \end{cases} \end{aligned}$$

The following definition of entropy solution is based on the formulation in [DGT20, DG20] with a slight but significant modification inspired by [BVW14, Definition 1]. Define two sets

$$\begin{aligned} \mathcal{E} &:= \{\eta \in C^2(\mathbb{R}) : \eta'' \geq 0, \text{ supp } \eta'' \text{ is compact}\}, \\ \mathcal{E}_0 &:= \{\eta \in \mathcal{E} : \eta'(0) = 0\}. \end{aligned}$$

**Definition 2.2.** An entropy solution of (2.1) is a predictable stochastic process  $u : \Omega_T \rightarrow L_1(D)$  such that

- (i)  $u \in L_{m+1}(\Omega_T; L_{m+1}(D))$ ;
- (ii) For all  $f \in C_b(\mathbb{R})$ , we have  $\llbracket \mathbf{a}f \rrbracket(u) \in L_2(\Omega_T; H_0^1(D))$  and
$$\partial_{x_i} \llbracket \mathbf{a}f \rrbracket(u) = f(u) \partial_{x_i} \llbracket \mathbf{a} \rrbracket(u);$$

(iii) For all

$$(\eta, \varphi, \varrho) \in (\mathcal{E} \times C_c^\infty[0, T] \times C_c^\infty(D)) \cup (\mathcal{E}_0 \times C_c^\infty[0, T] \times C^\infty(\overline{D}))$$

such that  $\phi := \varphi \times \varrho \geq 0$ , we have almost surely

$$(2.3) \quad \begin{aligned} & - \int_0^T \int_D \eta(u) \partial_t \phi dx dt \\ & \leq \int_D \eta(\xi) \phi(0) dx + \int_0^T \int_D \left( \llbracket \mathbf{a}^2 \eta' \rrbracket(u) \Delta \phi + \llbracket a^{ij} \eta' \rrbracket(x, u) \partial_{x_i x_j} \phi \right) dx dt \\ & \quad + \int_0^T \int_D \left( \llbracket a_{x_j}^{ij} \eta' - f_r^i \eta' \rrbracket(x, u) - \eta'(u) b^j(x, u) \right) \partial_{x_i} \phi dx dt \\ & \quad + \int_0^T \int_D \left( \eta'(u) f_{x_i}^i(x, u) - \llbracket f_{rx_i}^i \eta' \rrbracket(x, u) + \eta'(u) F(x, u) \right) \phi dx dt \\ & \quad + \int_0^T \int_D \left( \frac{1}{2} \eta''(u) \sum_{k=1}^{\infty} |\sigma_{x_i}^{ik}(x, u)|^2 - \eta''(u) |\nabla \llbracket \mathbf{a} \rrbracket(u)|^2 \right) \phi dx dt \\ & \quad + \int_0^T \int_D \left( \eta'(u) \phi \sigma_{x_i}^{ik}(x, u) - \llbracket \sigma_{rx_i}^{ik} \eta' \rrbracket(x, u) \phi - \llbracket \sigma_r^{ik} \eta' \rrbracket(x, u) \partial_{x_i} \phi \right) dx dW^k(t). \end{aligned}$$

*Remark 2.3.* Comparing with [DGT20, Definition 2.4], we expand the set of test functions  $(\eta, \varphi, \varrho)$  in order to give a more precise characterization of the behavior of solutions near the boundary. This is a critical point in our proof of  $L^1$ - estimates. Moreover, the Dirichlet boundary condition is satisfied implicitly according to Definition 2.2 (ii) and (iii).

The regularity assumption on coefficients coincides with [DG20, Assumption 2.3] for the Cauchy problem in a torus.

**Assumption 2.4.**  $G^i : D \times \mathbb{R} \rightarrow \mathbb{R}$  and  $\sigma^i : D \times \mathbb{R} \rightarrow l^2$  for  $i \in \{1, \dots, d\}$  and  $F : D \times \mathbb{R} \rightarrow \mathbb{R}$  are all continuous. For all  $i, l \in \{1, \dots, d\}$ ,  $q \in \{1, 2\}$  and all multi-indices  $\gamma \in \mathbb{N}^d$  with  $q + |\gamma| \leq 3$ , the derivatives  $\partial_r G^i$ ,  $\partial_{x_l} G^i$ ,  $\partial_{rx_l} G^i$  and  $\partial_r^q \partial_x^\gamma \sigma^i$  exist and are continuous on  $D \times \mathbb{R}$ . Moreover, there exist  $\bar{\kappa} \in ((m \wedge 2)^{-1}, 1]$ ,  $\beta \in ((2\bar{\kappa})^{-1}, 1]$ ,  $N_0 > 0$  and  $\tilde{\beta} \in (0, 1)$  such that for all  $i, j, l \in \{1, \dots, d\}$  and  $r \in \mathbb{R}$ , we have:

$$(2.4) \quad \sup_r \|\sigma_r^i(\cdot, r)\|_{H_\infty^2(D; l^2)} + \|\sigma_{x_l}^i(\cdot, 0)\|_{C^{\bar{\kappa}}(D; l^2)} \leq N_0,$$

$$(2.5) \quad \sup_{x \in D} \left( [\sigma(x, \cdot)]_{C^\beta(\mathbb{R}; l^2)} + \|\sigma_{rx_l}^i(x, \cdot)\|_{H_\infty^1(\mathbb{R}; l^2)} \right) \leq N_0,$$

$$(2.6) \quad \|\partial_r(\sigma_{x_j}^{jk} \sigma_{rx_l}^{ik})\|_{L_\infty} \leq N_0,$$

$$(2.7) \quad \sup_{x \in D} \left( \|G_r^i(x, \cdot)\|_{C^\beta(\mathbb{R})} + \|\partial_r(\sigma_r^{ik} \sigma_{x_j}^{jk})(x, \cdot)\|_{C^\beta(\mathbb{R})} \right) \leq N_0,$$

$$(2.8) \quad [G_{x_l}^i(\cdot, r)]_{C^{\tilde{\beta}}(D)} + [\partial_{x_l}(\sigma_r^{ik}(\cdot, r) \sigma_{x_j}^{jk}(\cdot, r))]_{C^{\tilde{\beta}}(D)} \leq N_0(1 + |r|),$$

$$(2.9) \quad \|\partial_{x_l} \partial_r(\sigma_r^{ik} \sigma_{x_j}^{jk})\|_{L_\infty} + \|G_{x_l r}^i\|_{L_\infty} \leq N_0,$$

$$(2.10) \quad \sup_{x \in D} [F(x, \cdot)]_{C^1(\mathbb{R})} + \sup_r \|F(\cdot, r)\|_{C^{\tilde{\beta}}(D)} \leq N_0.$$

Due to the physical background, we focus on non-negative solutions to the concerned problem, for which we need the following natural condition.

**Assumption 2.5.** The coefficients satisfy

$$(2.11) \quad \nabla_x \cdot G(x, 0) + F(x, 0) \geq 0, \quad \nabla_x \cdot \sigma(x, 0) = \{0\}_{k=1}^\infty, \quad x \in D.$$

The main result of this paper is stated as follows.

**Theorem 2.6.** Under Assumptions 2.1, 2.4 and 2.5, for  $0 \leq \xi \in L_{m+1}(\Omega, \mathcal{F}_0; L_{m+1}(D))$ , we have that

- (i) the problem  $\Pi(\Phi, \xi)$  has an unique entropy solution  $u$ ;
- (ii)  $u \geq 0$  for almost all  $(\omega, t, x) \in \Omega \times D$ ;
- (iii) if  $\tilde{u}$  is the entropy solution to  $\Pi(\Phi, \tilde{\xi})$  with  $0 \leq \tilde{\xi} \in L_{m+1}(\Omega, \mathcal{F}_0; L_{m+1}(D))$ , then

$$(2.12) \quad \text{ess sup}_{t \in [0, T]} \mathbb{E} \|(u(t, \cdot) - \tilde{u}(t, \cdot))^+\|_{L_1(D)} \leq C \mathbb{E} \|(\xi - \tilde{\xi})^+\|_{L_1(D)},$$

where the constant  $C$  depends only on  $N_0, K, d, T$  and  $|D|$ .

*Remark 2.7.* In view of Theorem 2.6, it is natural to define the ‘‘entropy solution’’ of the problem  $\Pi(\Phi, \xi)$  with  $\xi \in L_1(\Omega, \mathcal{F}_0; L_1(D))$  to be the  $L_1$ -limit of the entropy solution  $u^k$  of  $\Pi(\Phi, \xi^k)$  with  $\xi^k \in L_{m+1}(\Omega, \mathcal{F}_0; L_{m+1}(D))$  that converges to  $\xi$  in  $L_1(\Omega \times D)$  as  $k \rightarrow \infty$ . Indeed, the  $L_1$ -estimates (2.12) ensures that the limit does exist and be independent of the choice of sequence  $\xi^k$ .

We conclude this section by proving non-negativity of entropy solutions under our assumption.

**Proposition 2.8.** Under the condition of Theorem 2.6, each entropy solution to  $\Pi(\Phi, \xi)$  with  $0 \leq \xi \in L_{m+1}(\Omega, \mathcal{F}_0; L_{m+1}(D))$  is non-negative for almost all  $(\omega, t, x) \in \Omega_T \times D$ .

*Proof.* For any small  $\delta > 0$ , we introduce a function  $\eta_\delta \in C^2(\mathbb{R})$  defined by

$$\eta_\delta(0) = \eta'_\delta(0) = 0, \quad \eta''_\delta(r) = \rho_\delta(r).$$

Applying the entropy formulation (2.3) with  $\eta(\cdot) = \eta_\delta(\cdot)$  and  $\phi$  independent of  $x$ , using the non-negativity of  $\xi$ , we have

$$(2.13) \quad -\mathbb{E} \int_0^T \int_D \eta_\delta(-u) \partial_t \phi dx dt$$

$$\begin{aligned}
&\leq \mathbb{E} \int_0^T \int_D \left( \llbracket f_{rx_i}^i \eta'_\delta(\cdot) \rrbracket(x, u) - \eta'_\delta(-u)(f_{x_i}^i(x, u) - G_{x_i}^i(x, 0)) \right) \phi dx dt \\
&\quad - \mathbb{E} \int_0^T \int_D \eta'_\delta(-u)(F(x, u) + G_{x_i}^i(x, 0)) \phi dx dt \\
&\quad + \mathbb{E} \int_0^T \int_D \left( \frac{1}{2} \eta''_\delta(-u) \sum_{k=1}^{\infty} |\sigma_{x_i}^{ik}(x, u)|^2 \phi - \eta''_\delta(-u) |\nabla \llbracket \mathbf{a} \rrbracket(u)|^2 \phi \right) dx dt.
\end{aligned}$$

With Assumption 2.5 and (2.10) in Assumption 2.4, we have

$$(2.14) \quad \sup_x (F(x, r) + G_{x_i}^i(x, 0)) \geq \sup_x (F(x, 0) + G_{x_i}^i(x, 0)) - C|r| \geq -C|r|$$

for a positive constant  $C$ . Moreover, using Assumption 2.5, the definition of  $f^i$  and (2.5) and (2.9) in Assumption 2.4, we have

$$(2.15) \quad \sup_x |\sigma_{x_i}^i(x, r)|_{l_2} + \sup_x |f_{x_i}^i(x, r) - G_{x_i}^i(x, 0)| \leq C|r|.$$

Combining (2.13)-(2.15) with (2.9) in Assumption 2.4, we have

$$-\mathbb{E} \int_0^T \int_D \eta_\delta(-u) \partial_t \phi dx dt \leq C \mathbb{E} \int_0^T \int_D (-u)^+ \phi dx dt + C\delta.$$

Since  $|\eta_\delta(r) - r^+| \leq \delta$ , taking  $\delta \rightarrow 0^+$ , we have

$$(2.16) \quad -\mathbb{E} \int_0^T \int_D (-u)^+ \partial_t \phi dx dt \leq C \mathbb{E} \int_0^T \int_D (-u)^+ \phi dx dt.$$

Let  $0 < s < \tau < T$  be the Lebesgue points of the function

$$t \mapsto \mathbb{E} \int_D (-u(t, x))^+ dx.$$

Fix a constant  $\gamma \in (0, (\tau - s) \vee (T - \tau))$ . We choose a sequence of functions  $\{\phi_n\}_{n \in \mathbb{N}}$  satisfying  $\phi_n \in C_c^\infty((0, T))$  and  $\|\phi_n\|_{L^\infty(0, T)} \vee \|\partial_t \phi_n\|_{L^1(0, T)} \leq 1$ , such that

$$\lim_{n \rightarrow \infty} \|\phi_n - V_{(\gamma)}\|_{H_0^1(0, T)} = 0,$$

where  $V_{(\gamma)} : [0, T] \rightarrow \mathbb{R}$  satisfies  $V_{(\gamma)}(0) = 0$  and  $V_{(\gamma)}' = \gamma^{-1} \mathbf{1}_{[s, s+\gamma]} - \gamma^{-1} \mathbf{1}_{[\tau, \tau+\gamma]}$ . Taking  $\phi = \phi_n$  in (2.16) and passing the limit  $n \rightarrow \infty$ , we have

$$\frac{1}{\gamma} \mathbb{E} \int_\tau^{\tau+\gamma} \int_D (-u)^+ dx dt \leq C \mathbb{E} \int_0^{\tau+\gamma} \int_D (-u)^+ dx dt + \frac{1}{\gamma} \mathbb{E} \int_s^{s+\gamma} \int_D (-u)^+ dx dt.$$

Let  $\gamma \rightarrow 0^+$ , we have

$$\mathbb{E} \int_D (-u(\tau, x))^+ dx \leq C \mathbb{E} \int_0^\tau \int_D (-u)^+ dx dt + \mathbb{E} \int_D (-u(s, x))^+ dx$$

holds for almost all  $s \in (0, \tau)$ . Then, for each  $\tilde{\gamma} \in (0, \tau)$ , by averaging over  $s \in (0, \tilde{\gamma})$ , we have

$$\mathbb{E} \int_D (-u(\tau, x))^+ dx \leq C \mathbb{E} \int_0^{\tau+\gamma} \int_D (-u)^+ dx dt + \frac{1}{\tilde{\gamma}} \mathbb{E} \int_0^{\tilde{\gamma}} \int_D (-u)^+ dx ds.$$

Taking the limit  $\tilde{\gamma} \rightarrow 0^+$  and using Lemma 6.1, the non-negativity of  $\xi$  and Gronwall's inequality, we have  $u \geq 0$  for almost all  $(\omega, t, x) \in \Omega_T \times D$ .  $\square$

3. STRONG ENTROPY CONDITION AND  $L_1$ -ESTIMATES

To deal with the Dirichlet problem, we need to split the bounded domain  $D$ . Choose an open covering of  $\overline{D}$  by  $B_0$  and balls  $\{B_i\}_{i=1}^K$  satisfying  $\text{dist}(B_0, \partial D) > 0$ , and for  $i > 0$ ,  $B_i \subset B'_i$  with  $B'_i \cap \partial D$  part of a Lipschitz graph and  $\text{dist}(B_i, \partial B'_i) > 0$ , where

$$\text{dist}(A_1, A_2) := \inf_{x_1 \in A_1, x_2 \in A_2} |x_1 - x_2|, \quad \forall A_1, A_2 \subset \mathbb{R}^d.$$

From [BB11, Lemma 9.3], we know that there exist functions  $\psi_0, \psi_1, \dots, \psi_K \in C^\infty(\mathbb{R}^d)$  such that  $0 \leq \psi_i \leq 1$ ,  $\text{supp} \psi_i \subseteq B_i$  for  $i = 0, 1, \dots, K$ , and

$$\sum_{i=0}^K \psi_i(x) \equiv 1, \quad x \in \overline{D}.$$

Without loss of generality, we assume that  $\text{dist}(\text{supp} \psi_i, \partial B_i) > 0$ . Otherwise, we can take a bigger open domain instead of  $B_i$  satisfying  $\text{dist}(B_0, \partial D) > 0$  and  $\text{dist}(B_i, \partial B'_i) > 0$ .

For  $x = (x_1, x_2, \dots, x_d) \in \mathbb{R}^d$ , define the function  $\tilde{\rho}(x) := \prod_{i=1}^d \rho(x_i - 1/2)$  and the mollifiers  $\tilde{\rho}_\varepsilon(\cdot) := \tilde{\rho}(\cdot/\varepsilon)/(\varepsilon^d)$ . Similar to [BVW14, Section 3.2.1], for each  $i = 1, \dots, K$ , there exist a constant  $\varepsilon_i$  and a vector  $\tilde{\eta}_i \in \mathbb{R}^d$ , such that the translated sequence of mollifiers  $\rho_{\varepsilon,i}(\cdot) := \tilde{\rho}_\varepsilon(\cdot - \tilde{\eta}_i)$  satisfying  $y \mapsto \rho_{\varepsilon,i}(x - \cdot) \in C_c^\infty(D)$  for all  $(x, \varepsilon) \in (B_i \cap \overline{D}) \times (0, \varepsilon_i)$ . The vector  $\tilde{\eta}_i$  depends only on the local representation of the boundary of  $D$  in  $B'_i$  as the graph of a Lipschitz function. We also define  $\rho_{\varepsilon,0}(\cdot) := \tilde{\rho}_\varepsilon(\cdot)$ .

*Remark 3.1.* From the construction of the function  $\rho_{\varepsilon,i}$ , there exists a constant  $N$  depending on the maximum norm of  $\eta_i$ ,  $i = 1, \dots, K$ , such that  $\text{supp} \rho_{\varepsilon,i} \subset \{x \in \mathbb{R}^d : |x| < N\varepsilon\}$  holds for all  $\varepsilon \in (0, \varepsilon_i)$  and  $i = 0, 1, \dots, K$ .

*Remark 3.2.* For each  $i = 1, \dots, K$ , since  $\text{dist}(\text{supp} \psi_i, \partial B_i) > 0$ , there exists a  $\bar{\varepsilon}_i \in (0, \varepsilon_i)$  such that  $\text{supp} \rho_{\varepsilon,i}(x - \cdot) \subset B_i \cap D$  for all  $(x, \varepsilon) \in (\text{supp} \psi_i \cap \overline{D}) \times (0, \bar{\varepsilon}_i)$ . Moreover, there exists a  $\varepsilon_0 \in (0, 1)$  such that  $\text{supp} \rho_{\varepsilon,0}(x - \cdot) \subset D$  for all  $x \in B_0$ . In the rest of this article, we define

$$\bar{\varepsilon} := \min\{\varepsilon_0, \varepsilon_1, \bar{\varepsilon}_1, \varepsilon_2, \bar{\varepsilon}_2, \dots, \varepsilon_K, \bar{\varepsilon}_K\}.$$

Now, we fix  $i \in \{0, 1, \dots, K\}$  and define  $B := B_i$ ,  $\psi := \psi_i$  and  $\rho_\varepsilon(x - y) := \rho_{\varepsilon,i}(x - y)$  for the sake of brevity.

**3.1. ( $\star$ )-property.** Define the sets

$$\mathcal{C}^- := \{f \in C^\infty(\mathbb{R}) : f' \in C_c(\mathbb{R}), \text{supp} f \subseteq (-\infty, 0]\},$$

$$\mathcal{C}^+ := \{f \in C^\infty(\mathbb{R}) : f' \in C_c(\mathbb{R}), \text{supp} f \subseteq (0, \infty)\},$$

$$\Gamma_B := \{f \in C^\infty(\overline{(B \cap D)} \times \overline{(B \cap D)}) : f(x, \cdot) \in C^\infty(D), \forall x \in B \cap \overline{D}\}.$$

Let  $(g, \varphi, u, h) \in \Gamma_B \times C_c^\infty((0, T)) \times L_{m+1}(\Omega_T; L_{m+1}(D)) \times (\mathcal{C}^- \cup \mathcal{C}^+)$ . For  $\theta > 0$ , we introduce

$$\phi_\theta(t, s, x, y) := g(x, y) \rho_\theta(s - t) \varphi\left(\frac{t + s}{2}\right), \quad \tilde{\phi}_\theta(t, s, x, y) := \phi_\theta(t, s, y, x),$$

$$\begin{aligned} \mathcal{H}_\theta(\phi_\theta(\cdot, s, x, \cdot), z) &:= \int_0^T \int_y \left( h(u(t, y) - z) \sigma_{y_i}^{ik}(y, u(t, y)) \phi_\theta(t, s, x, y) \right. \\ &\quad - \int_0^{u(t, y)} h(r - z) \sigma_{ry_i}^{ik}(y, r) dr \phi_\theta(t, s, x, y) \\ &\quad \left. - \int_0^{u(t, y)} h(r - z) \sigma_r^{ik}(y, r) dr \partial_{y_i} \phi_\theta(t, s, x, y) \right) dW^k(t), \end{aligned}$$

$$\begin{aligned}
\mathfrak{H}_\theta(u, w, \phi_\theta) &:= -\mathbb{E} \int_{t,s,x,y} \partial_{x_j y_i} \phi_\theta(t, s, x, y) \int_u^w \int_{\tilde{r}}^u h'(r - \tilde{r}) \sigma_r^{ik}(y, r) \sigma_r^{jk}(x, \tilde{r}) dr d\tilde{r} \\
&\quad - \mathbb{E} \int_{t,s,x,y} \partial_{y_i} \phi_\theta(t, s, x, y) \int_u^w \int_{\tilde{r}}^u h'(r - \tilde{r}) \sigma_r^{ik}(y, r) \sigma_{rx_j}^{jk}(x, \tilde{r}) dr d\tilde{r} \\
&\quad + \mathbb{E} \int_{t,s,x,y} \partial_{y_i} \phi_\theta(t, s, x, y) \int_w^u h'(r - w) \sigma_r^{ik}(y, r) \sigma_{x_j}^{jk}(x, w) dr \\
&\quad - \mathbb{E} \int_{t,s,x,y} \partial_{x_j} \phi_\theta(t, s, x, y) \int_u^w \int_{\tilde{r}}^u h'(r - \tilde{r}) \sigma_{ry_i}^{ik}(y, r) \sigma_r^{jk}(x, \tilde{r}) dr d\tilde{r} \\
&\quad - \mathbb{E} \int_{t,s,x,y} \phi_\theta(t, s, x, y) \int_u^w \int_{\tilde{r}}^u h'(r - \tilde{r}) \sigma_{ry_i}^{ik}(y, r) \sigma_{rx_j}^{jk}(x, \tilde{r}) dr d\tilde{r} \\
&\quad + \mathbb{E} \int_{t,s,x,y} \phi_\theta(t, s, x, y) \int_w^u h'(r - w) \sigma_{ry_i}^{ik}(y, r) \sigma_{x_j}^{jk}(x, w) dr \\
&\quad + \mathbb{E} \int_{t,s,x,y} \partial_{x_j} \phi_\theta(t, s, x, y) \int_u^w h'(u - \tilde{r}) \sigma_{y_i}^{ik}(y, u) \sigma_r^{jk}(x, \tilde{r}) d\tilde{r} \\
&\quad + \mathbb{E} \int_{t,s,x,y} \phi_\theta(t, s, x, y) \int_u^w h'(u - \tilde{r}) \sigma_{y_i}^{ik}(y, u) \sigma_{rx_j}^{jk}(x, \tilde{r}) d\tilde{r} \\
&\quad - \mathbb{E} \int_{t,s,x,y} \phi_\theta(t, s, x, y) h'(u - w) \sigma_{y_i}^{ik}(y, u) \sigma_{x_j}^{jk}(x, w),
\end{aligned}$$

where in the integrand  $u = u(t, y)$  and  $w = w(t, x)$ . For simplicity, here and below we write  $\int_t$  in the place of  $\int_0^T \cdot dt$  (and similarly for  $\int_s$ ),  $\int_x$  in place of  $\int_D \cdot dx$  (and similarly for  $\int_y$ ), and  $\int_z$  in place of  $\int_{\mathbb{R}} \cdot dz$ . However, to avoid confusion, we use the usual notation if the integral is taken on a different domain or is a stochastic integral. Then, we define

$$\begin{aligned}
H_\theta(s, x, z) &:= \mathcal{H}_\theta(\phi_\theta(\cdot, s, x, \cdot), z), & \tilde{H}_\theta(s, x, z) &:= \mathcal{H}_\theta(\tilde{\phi}_\theta(\cdot, s, x, \cdot), z), \\
\mathcal{E}(u, w, \theta) &:= \mathfrak{H}_\theta(u, w, \phi_\theta), & \tilde{\mathcal{E}}(u, w, \theta) &:= \mathfrak{H}_\theta(u, w, \tilde{\phi}_\theta).
\end{aligned}$$

From the different status of  $x$  and  $y$  in the function  $g \in \Gamma_B$ , the function  $H_\theta$  may not equal to  $\tilde{H}_\theta$ . For the sake of the consistence of the formulation of the test function  $\phi_\theta$ , we exchange  $x \leftrightarrow y$  in the integrand of the definition of  $\tilde{H}_\theta$  and  $\tilde{\mathcal{E}}(u, w, \theta)$ . Then, we can see that  $\tilde{H}_\theta(s, y, z)$  are the terms exchanging  $x$  and  $y$  in the integrand of  $H_\theta(s, x, z)$  except the test function  $\phi_\theta$ .

*Remark 3.3.* Since  $\text{supp} \varphi \subset (0, T)$ , for sufficiently small  $\theta$ , the function  $H_\theta$  has a similar form as [DGG19, (3.7)], and it is also true for  $\tilde{H}_\theta(s, y, z)$ . The definition of the set  $\Gamma_B$  also indicates  $\text{supp} H_\theta(s, \cdot, z), \text{supp} \tilde{H}_\theta(s, \cdot, z) \subset B \cap \bar{D}$  for all  $(s, z) \in [0, T] \times \mathbb{R}$ . Furthermore, there exists a modification of  $H_\theta$  which is smooth in  $(s, x, z)$  (see [Kun97, Exercise 3.15]). Throughout this paper, we will use this smooth version and still denote it by  $H_\theta$ .

Set  $\mu = (3m + 5)/(4m + 4)$ .

**Definition 3.4.** We say that a function  $w \in L_{m+1}(\Omega_T; L_{m+1}(D))$  has the  $(\star)$ -property, if there exists an open covering of  $\bar{D}$  by  $B_0$  and balls  $\{B_i\}_{i=1}^K$  such that

(i)  $B_0 \cap \partial D = \emptyset$ . and for  $i > 0$ , there exists a ball  $B'_i$  such that  $B_i \subset B'_i$  and  $B'_i \cap \partial D$  is part of a Lipschitz graph.

(ii) For each  $i \in \{0, 1, \dots, K\}$ , define  $B := B_i$ . For all  $(g, \varphi, u, h) \in \Gamma_B \times C_c^\infty((0, T)) \times L_{m+1}(\Omega_T; L_{m+1}(D)) \times C^-$  satisfying  $u \geq 0$  for almost all  $(\omega, t, x) \in \Omega_T \times D$ , and for all sufficiently small  $\theta \in (0, 1)$ , we have  $H_\theta(\cdot, \cdot, w(\cdot, \cdot)) \in L_1(\Omega_T \times D)$  and

$$\mathbb{E} \int_{s,x} H_\theta(s, x, w(s, x)) \leq C\theta^{1-\mu} + \mathcal{E}(u, w, \theta)$$



for a constant  $C$  independent of  $\theta$ .

(iii) For each  $i = 0, 1, \dots, K$  and all  $(g, \varphi, u, h) \in \Gamma_{B_i} \times C_c^\infty((0, T)) \times L_{m+1}(\Omega_T; L_{m+1}(D)) \times \mathcal{C}^+$  and sufficiently small  $\theta \in (0, 1)$ , we have  $\tilde{H}_\theta(\cdot, \cdot, w(\cdot, \cdot)) \in L_1(\Omega_T \times D)$  and

$$\mathbb{E} \int_{s,y} \tilde{H}_\theta(s, y, w(s, y)) \leq C\theta^{1-\mu} + \tilde{\mathcal{E}}(u, w, \theta)$$

for a constant  $C$  independent of  $\theta$ .

*Remark 3.5.* Differing from the  $(\star)$ -property in [DGG19, DGT20, DG20], we adjust the test functions  $g, u$  and  $h$  to apply the divergence theorem with the Dirichlet boundary condition (see the proof of Proposition 4.4). Remark 3.11 shows the reason we consider separately the cases (ii) and (iii) in Definition 3.4.

The following lemmas are introduced from [DGG19, DG20], and the proofs are similar no matter the space is  $D$  or  $\mathbb{T}^d$ . We omit the proofs here.

**Lemma 3.6.** *Under (2.4)-(2.5) in Assumption 2.4, for all  $\lambda \in ((m+3)/(2m+2), 1)$ ,  $\bar{k} \in \mathbb{N}$  and sufficiently small  $\theta \in (0, 1)$ , we have*

$$\begin{aligned} \mathbb{E} \|\partial_z H_\theta\|_{L_\infty([0, T]; H_{m+1}^{\bar{k}}(D \times \mathbb{R}))}^{m+1} &\leq C\theta^{-\lambda(m+1)} \mathcal{N}_m(u), \\ \mathbb{E} \|\partial_z \tilde{H}_\theta\|_{L_\infty([0, T]; H_{m+1}^{\bar{k}}(D \times \mathbb{R}))}^{m+1} &\leq C\theta^{-\lambda(m+1)} \mathcal{N}_m(u), \end{aligned}$$

where

$$\mathcal{N}_m(u) := \mathbb{E} \int_0^T \left( 1 + \|u(t)\|_{L_{\frac{m+1}{2}}(D)}^{m+1} + \|u(t)\|_{L_2(D)}^{m+1} \right) dt,$$

and the constant  $C = C(N_0, k, d, T, \lambda, |D|, m, h, g, \varphi)$  is independent of  $\theta$ . In particular, we have

$$\begin{aligned} \mathbb{E} \|\partial_z H_\theta\|_{L_\infty([0, T]; H_{m+1}^{\bar{k}}(D \times \mathbb{R}))}^{m+1} &\leq C\theta^{-\lambda(m+1)} \left( 1 + \|u\|_{L_{m+1}(D_T)}^{m+1} \right), \\ \mathbb{E} \|\partial_z \tilde{H}_\theta\|_{L_\infty([0, T]; H_{m+1}^{\bar{k}}(D \times \mathbb{R}))}^{m+1} &\leq C\theta^{-\lambda(m+1)} \left( 1 + \|u\|_{L_{m+1}(D_T)}^{m+1} \right). \end{aligned}$$

**Lemma 3.7.** *Let  $w \in L_2(\Omega_T \times D)$ . Then, for all sufficiently small  $\theta \in (0, 1)$ , we have*

$$\begin{aligned} \mathbb{E} \int_{s,x} H_\theta(s, x, w(s, x)) &= \lim_{\lambda \rightarrow 0} \mathbb{E} \int_{s,x,z} H_\theta(s, x, z) \rho_\lambda(w(s, x) - z), \\ \mathbb{E} \int_{s,y} \tilde{H}_\theta(s, y, w(s, y)) &= \lim_{\lambda \rightarrow 0} \mathbb{E} \int_{s,y,z} H_\theta(s, y, z) \rho_\lambda(w(s, y) - z). \end{aligned}$$

**Lemma 3.8.** *Let  $\{w_n\}_{n \in \mathbb{N}}$  be a sequence bounded in  $L_{m+1}(\Omega_T \times D)$  satisfying the  $(\star)$ -property uniformly in  $n$ , which means the constant  $C$  in Definition 3.4 (ii) and (iii) are independent of  $n$ . Suppose that  $w_n$  converges to a function  $w$  for almost all  $(\omega, t, x) \in \Omega_T \times D$ . Then,  $w$  has the  $(\star)$ -property.*

**3.2.  $L_1$ -estimates.** Now, we give the  $L_1$ -estimates of two entropy solutions  $u$  and  $\tilde{u}$  using the Kruzhkov's doubling variables technique (cf. [Kru70]), which can eliminate the coupling effect of  $u - \tilde{u}$ . One key point is to select the appropriate function  $\eta$  such that  $\eta(\cdot - z) \in \mathcal{E}_0$  for any  $z \geq 0$  to approximate the positive part function. Using the convolution, the non-negative  $z$  can be replaced by another entropy solution under suitable conditions.

For a non-negative  $\varphi \in C_c^\infty((0, T))$  satisfying  $\|\varphi\|_{L_\infty(0, T)} \vee \|\partial_t \varphi\|_{L_1(0, T)} \leq 1$ , we define the non-negative test functions

$$\phi_\varepsilon(t, x, y) := \varrho_\varepsilon(x - y) \varphi(t) \psi(x) \mathbf{1}_{\bar{D}}(x), \quad \phi_{\theta, \varepsilon}(t, x, s, y) := \rho_\theta(s - t) \phi_\varepsilon\left(\frac{s+t}{2}, x, y\right),$$

where  $\varrho_\varepsilon$  corresponding to  $B$  is defined in the beginning of Section 3.

*Remark 3.9.* Compare to the test functions  $\phi_\theta$  in Section 3. We want to take  $g(x, y) = \mathbf{1}_{\overline{D}}(x)\psi(x)\varrho_\varepsilon(x - y)$ . In this case, with  $\varepsilon \in (0, \bar{\varepsilon})$  (see Remark 3.2 for the definition of  $\bar{\varepsilon}$ ), we have  $g \in \Gamma_B$ .

To derived the  $L_1$ -estimates, we need the following lemma.

**Lemma 3.10.** *Let  $0 \leq \xi, \tilde{\xi} \in L_{m+1}(\Omega, \mathcal{F}_0; L_{m+1}(D))$ . Suppose that  $u$  and  $\tilde{u}$  are the entropy solutions to the Dirichlet problems  $\Pi(\Phi, \xi)$  and  $\Pi(\tilde{\Phi}, \tilde{\xi})$ , respectively. Let Assumptions 2.1, 2.4 and 2.5 hold for both  $\Phi$  and  $\tilde{\Phi}$ . For  $\delta \in (0, 1)$ ,  $\varepsilon \in (0, \bar{\varepsilon})$ ,  $\lambda \in [0, 1]$ ,  $\alpha \in (0, 1 \wedge (m/2))$  and every non-negative  $\varphi \in C_c^\infty((0, T))$  such that*

$$\|\varphi\|_{L^\infty(0, T)} \vee \|\partial_t \varphi\|_{L_1(0, T)} \leq 1,$$

define

$$\begin{aligned} R_\lambda &:= \sup \{R \in [0, \infty] : |\mathbf{a}(r) - \tilde{\mathbf{a}}(r)| \leq \lambda, \forall |r| < R\}, \\ \mathfrak{C}(\varepsilon, \delta, \lambda, \alpha) &:= \varepsilon^{-2}\delta^{2\beta} + \delta^\beta \varepsilon^{-1} + \varepsilon^{\bar{\kappa}} + \varepsilon^{\tilde{\beta}} + \varepsilon^{\frac{1}{m+1}} + \varepsilon^{2\bar{\kappa}}\delta^{-1} \\ &\quad + \varepsilon^{-2}\delta^{2\alpha} + \varepsilon^{-2}\lambda^2 + \varepsilon^{-1}\lambda, \\ \mathcal{Q}(\tilde{u}, u) &:= -\mathbb{E} \int_{t, x, y} (\tilde{u}(t, x) - u(t, y))^+ \varrho_\varepsilon(x - y) \partial_t \varphi(t) \psi(x), \\ \mathfrak{R}(\tilde{u}, \tilde{\Phi}, u, \Phi) &:= \mathbb{E} \int_{t, x} \varphi(t) \Delta_x \psi(x) (\tilde{\Phi}(\tilde{u}(t, x)) - \Phi(u(t, x)))^+ + C\varepsilon^{\frac{1}{m+1}} \mathbb{E} \|\nabla[\mathbf{a}](u)\|_{L_2(D_T)}^2 \\ &\quad + C\mathfrak{C}(\varepsilon, \delta, \lambda, \alpha) \mathbb{E}(1 + \|u\|_{L_{m+1}(D_T)}^{m+1} + \|\tilde{u}\|_{L_{m+1}(D_T)}^{m+1}) \\ &\quad + C\varepsilon^{-2} \mathbb{E} \left( \|\mathbf{1}_{|u| \geq R_\lambda} (1 + u)\|_{L_m(D_T)}^m + \|\mathbf{1}_{|\tilde{u}| \geq R_\lambda} (1 + \tilde{u})\|_{L_m(D_T)}^m \right) \\ &\quad + C\mathbb{E} \int_{t, x, y} \mathbf{1}_{B \cap \overline{D}}(x) \varphi(t) \\ &\quad \cdot \left( \varepsilon^2 \sum_{i, j} |\partial_{x_i y_j} \varrho_\varepsilon(x - y)| + \varepsilon \sum_i |\partial_{x_i} \varrho_\varepsilon(x - y)| + \varrho_\varepsilon(x - y) \right) (\tilde{u}(t, x) - u(t, y))^+. \end{aligned}$$

If  $\tilde{u}$  has the  $(\star)$ -property, we have

$$(3.1) \quad \mathcal{Q}(\tilde{u}, u) \leq \mathfrak{R}(\tilde{u}, \tilde{\Phi}, u, \Phi),$$

$$(3.2) \quad \mathcal{Q}(u, \tilde{u}) \leq \mathfrak{R}(u, \Phi, \tilde{u}, \tilde{\Phi}).$$

*Remark 3.11.* Estimates (3.1) and (3.2) are not symmetric for  $u$  and  $\tilde{u}$  (See Remark 3.14 for the reason we focus on such terms) due to the  $(\star)$ -property of  $\tilde{u}$  and the different status of  $x$  and  $y$  in test function  $\phi_\varepsilon(t, x, y)$ . Therefore, it is necessary to consider the  $(\star)$ -property separately as in Definition 3.4. This is a key point of our proof, which is different from [DGG19, DGT20, DG20].

*Proof.* We first prove (3.1). With Proposition 2.8, we have  $u, \tilde{u} \geq 0$  for almost all  $(\omega, t, x) \in \Omega_T \times D$ . For each  $\delta > 0$ , define the function  $\eta_\delta \in C^2(\mathbb{R})$  by

$$\eta_\delta(0) = \eta'_\delta(0) = 0, \quad \eta''_\delta(r) = \rho_\delta(r).$$

Thus, we have

$$|\eta_\delta(r) - r^+| \leq \delta, \quad \text{supp} \eta''_\delta \subset [0, \delta], \quad \int_{\mathbb{R}} |\eta''_\delta(r)| dr \leq 2, \quad |\eta''_\delta| \leq 2\delta^{-1}.$$

Fix  $(z, t, y) \in [0, \infty) \times D_T$ . Since  $\tilde{u}$  is the entropy solution to  $\Pi(\tilde{\Phi}, \tilde{\xi})$  and

$$\left( \eta_\delta(\cdot - z), \rho_\theta(\cdot - t) \varphi\left(\frac{\cdot + t}{2}\right), \varrho_\varepsilon(\cdot - y) \psi(\cdot) \mathbf{1}_{\overline{D}}(\cdot) \right) \in \mathcal{E}_0 \times C_c^\infty((0, T)) \times C^\infty(\overline{D})$$

for  $\varepsilon \in (0, \bar{\varepsilon})$  and small enough  $\theta$ , using the entropy inequality (2.3) of  $\tilde{u}$  with  $(\eta_\delta(r-z), \phi_{\theta,\varepsilon}(t, \cdot, \cdot, y))$  instead of  $(\eta(r), \phi)$ , we have

$$\begin{aligned}
& - \int_{s,x} \eta_\delta(\tilde{u} - z) \partial_s \phi_{\theta,\varepsilon} \\
& \leq \int_{s,x} [\tilde{\mathbf{a}}^2 \eta'_\delta(\cdot - z)](\tilde{u}) \Delta_x \phi_{\theta,\varepsilon} + \int_{s,x} [a^{ij} \eta'_\delta(\cdot - z)](x, \tilde{u}) \partial_{x_i x_j} \phi_{\theta,\varepsilon} \\
& \quad + \int_{s,x} [a_{x_j}^{ij} \eta'_\delta(\cdot - z) - f_r^i \eta'_\delta(\cdot - z)](x, \tilde{u}) \partial_{x_i} \phi_{\theta,\varepsilon} \\
& \quad - \int_{s,x} \eta'_\delta(\tilde{u} - z) b^i(x, \tilde{u}) \partial_{x_i} \phi_{\theta,\varepsilon} + \int_{s,x} \eta'_\delta(\tilde{u} - z) F(x, \tilde{u}) \phi_{\theta,\varepsilon} \\
& \quad + \int_{s,x} \eta'_\delta(\tilde{u} - z) f_{x_i}^i(x, \tilde{u}) \phi_{\theta,\varepsilon} - \int_{s,x} [f_{rx_i}^i \eta'_\delta(\cdot - z)](x, \tilde{u}) \phi_{\theta,\varepsilon} \\
& \quad + \int_{s,x} \frac{1}{2} \eta''_\delta(\tilde{u} - z) \sum_{k=1}^{\infty} |\sigma_{x_i}^{ik}(x, \tilde{u})|^2 \phi_{\theta,\varepsilon} - \int_{s,x} \eta''_\delta(\tilde{u} - z) |\nabla_x [\tilde{\mathbf{a}}](\tilde{u})|^2 \phi_{\theta,\varepsilon} \\
& \quad + \int_0^T \int_x \left( \eta'_\delta(\tilde{u} - z) \phi_{\theta,\varepsilon} \sigma_{x_i}^{ik}(x, \tilde{u}) - [\sigma_{rx_i}^{ik} \eta'_\delta(\cdot - z)](x, \tilde{u}) \phi_{\theta,\varepsilon} \right. \\
& \quad \left. - [\sigma_r^{ik} \eta'_\delta(\cdot - z)](x, \tilde{u}) \partial_{x_i} \phi_{\theta,\varepsilon} \right) dW^k(s),
\end{aligned}$$

where  $\tilde{u} = \tilde{u}(s, x)$ . Notice that all the expressions are continuous in  $(z, t, y)$ . We take  $z = u(t, y)$  by convolution and integrate over  $(t, y) \in D_T$ . By taking expectations, we have

$$\begin{aligned}
(3.3) \quad & - \mathbb{E} \int_{t,x,s,y} \eta_\delta(\tilde{u} - u) \partial_s \phi_{\theta,\varepsilon} \\
& \leq \mathbb{E} \int_{t,x,s,y} [\tilde{\mathbf{a}}^2 \eta'_\delta(\cdot - u)](\tilde{u}) \Delta_x \phi_{\theta,\varepsilon} + \mathbb{E} \int_{t,x,s,y} [a^{ij} \eta'_\delta(\cdot - u)](x, \tilde{u}) \partial_{x_i x_j} \phi_{\theta,\varepsilon} \\
& \quad + \mathbb{E} \int_{t,x,s,y} [a_{x_j}^{ij} \eta'_\delta(\cdot - u) - f_r^i \eta'_\delta(\cdot - u)](x, \tilde{u}) \partial_{x_i} \phi_{\theta,\varepsilon} \\
& \quad - \mathbb{E} \int_{t,x,s,y} \eta'_\delta(\tilde{u} - u) b^i(x, \tilde{u}) \partial_{x_i} \phi_{\theta,\varepsilon} + \mathbb{E} \int_{t,x,s,y} \eta'_\delta(\tilde{u} - u) F(x, \tilde{u}) \phi_{\theta,\varepsilon} \\
& \quad + \mathbb{E} \int_{t,x,s,y} \eta'_\delta(\tilde{u} - u) f_{x_i}^i(x, \tilde{u}) \phi_{\theta,\varepsilon} - \mathbb{E} \int_{t,x,s,y} [f_{rx_i}^i \eta'_\delta(\cdot - u)](x, \tilde{u}) \phi_{\theta,\varepsilon} \\
& \quad + \mathbb{E} \int_{t,x,s,y} \frac{1}{2} \eta''_\delta(\tilde{u} - u) \sum_{k=1}^{\infty} |\sigma_{x_i}^{ik}(x, \tilde{u})|^2 \phi_{\theta,\varepsilon} - \mathbb{E} \int_{t,x,s,y} \eta''_\delta(\tilde{u} - u) |\nabla_x [\tilde{\mathbf{a}}](\tilde{u})|^2 \phi_{\theta,\varepsilon} \\
& \quad + \int_{t,y} \mathbb{E} \left[ \int_0^T \int_x \left( \eta'_\delta(\tilde{u} - z) \phi_{\theta,\varepsilon} \sigma_{x_i}^{ik}(x, \tilde{u}) - [\sigma_{rx_i}^{ik} \eta'_\delta(\cdot - z)](x, \tilde{u}) \phi_{\theta,\varepsilon} \right. \right. \\
& \quad \left. \left. - [\sigma_r^{ik} \eta'_\delta(\cdot - z)](x, \tilde{u}) \partial_{x_i} \phi_{\theta,\varepsilon} \right) dW^k(s) \right]_{z=u},
\end{aligned}$$

where  $u = u(t, y)$  and  $\tilde{u} = \tilde{u}(s, x)$ .

Similarly, for each  $(z, s, x) \in [0, \infty) \times D_T$ , since

$$(\eta_\delta(z - \cdot), \rho_\theta(s - \cdot) \varphi(\frac{s + \cdot}{2}), \varrho_\varepsilon(x - \cdot) \psi(x) \mathbf{1}_{\bar{D}}(x)) \in \mathcal{E} \times C_c^\infty((0, T)) \times C_c^\infty(D)$$

for  $\varepsilon \in (0, \bar{\varepsilon})$  and small enough  $\theta$ , we apply the entropy inequality of  $u$  with  $\eta(r) := \eta_\delta(z - r)$  and  $\phi(t, y) := \phi_{\theta, \varepsilon}(t, x, s, y)$ . After substituting  $z = \tilde{u}(s, x)$  by convolution, integrating over  $(s, x) \in D_T$  and taking expectations, we have

$$\begin{aligned}
(3.4) \quad & - \mathbb{E} \int_{t,x,s,y} \eta_\delta(\tilde{u} - u) \partial_t \phi_{\theta, \varepsilon} \\
& \leq - \mathbb{E} \int_{t,x,s,y} [\mathbf{a}^2 \eta'_\delta(\tilde{u} - \cdot)](u) \Delta_y \phi_{\theta, \varepsilon} - \mathbb{E} \int_{t,x,s,y} [\mathbf{a}^{ij} \eta'_\delta(\tilde{u} - \cdot)](y, u) \partial_{y_i y_j} \phi_{\theta, \varepsilon} \\
& \quad - \mathbb{E} \int_{t,x,s,y} [\mathbf{a}^{ij} \eta'_\delta(\tilde{u} - \cdot) - f_r^i \eta'_\delta(\tilde{u} - \cdot)](y, u) \partial_{y_i} \phi_{\theta, \varepsilon} \\
& \quad + \mathbb{E} \int_{t,x,s,y} \eta'_\delta(\tilde{u} - u) b^i(y, u) \partial_{y_i} \phi_{\theta, \varepsilon} - \mathbb{E} \int_{t,x,s,y} \eta'_\delta(\tilde{u} - u) F(y, u) \phi_{\theta, \varepsilon} \\
& \quad - \mathbb{E} \int_{t,x,s,y} \eta'_\delta(\tilde{u} - u) f_{y_i}^i(y, u) \phi_{\theta, \varepsilon} + \mathbb{E} \int_{t,x,s,y} [f_{ry}^i \eta'_\delta(\tilde{u} - \cdot)](y, u) \phi_{\theta, \varepsilon} \\
& \quad + \mathbb{E} \int_{t,x,s,y} \frac{1}{2} \eta''_\delta(\tilde{u} - u) \sum_{k=1}^{\infty} |\sigma_{y_i}^{ik}(y, u)|^2 \phi_{\theta, \varepsilon} - \mathbb{E} \int_{t,x,s,y} \eta''_\delta(\tilde{u} - u) |\nabla_y [\mathbf{a}](u)|^2 \phi_{\theta, \varepsilon} \\
& \quad - \int_{s,x} \mathbb{E} \left[ \int_0^T \int_y \left( \eta'_\delta(z - u) \phi_{\theta, \varepsilon} \sigma_{y_i}^{ik}(y, u) - [\sigma_{ry}^{ik} \eta'_\delta(z - \cdot)](y, u) \phi_{\theta, \varepsilon} \right. \right. \\
& \quad \left. \left. - [\sigma_r^{ik} \eta'_\delta(z - \cdot)](y, u) \partial_{y_i} \phi_{\theta, \varepsilon} \right) dW^k(t) \right]_{z=\tilde{u}},
\end{aligned}$$

where  $u = u(t, y)$  and  $\tilde{u} = \tilde{u}(s, x)$ . Note that the stochastic integral term in (3.3) is zero due to the support of  $\rho_\theta$ . Meanwhile, we use the  $(\star)$ -property (ii) of  $\tilde{u}$  for the last term of (3.4) with  $h(r) = -\eta'_\delta(-r)$  (then  $h \in \mathcal{C}^-$ ) and  $g(x, y) = \mathbf{1}_{\bar{D}}(x) \psi(x) \varrho_\varepsilon(x - y)$ . Adding inequalities (3.3)-(3.4) and taking the limit  $\theta \rightarrow 0^+$ , we have

$$(3.5) \quad - \mathbb{E} \int_{t,x,y} \eta_\delta(\tilde{u} - u) \partial_t \phi_\varepsilon \leq \sum_{i=1}^2 I_i + \sum_{i=1}^2 D_i + \sum_{i=1}^2 J_i + \sum_{i=1}^9 \mathcal{E}_i,$$

where

$$\begin{aligned}
I_1 & := \mathbb{E} \int_{t,x,y} [\tilde{\mathbf{a}}^2 \eta'_\delta(\cdot - u)](\tilde{u}) \Delta_x \phi_\varepsilon - \mathbb{E} \int_{t,x,y} [\mathbf{a}^2 \eta'_\delta(\tilde{u} - \cdot)](u) \Delta_y \phi_\varepsilon, \\
I_2 & := - \mathbb{E} \int_{t,x,y} \eta''_\delta(\tilde{u} - u) (|\nabla_x [\tilde{\mathbf{a}}](\tilde{u})|^2 + |\nabla_y [\mathbf{a}](u)|^2) \phi_\varepsilon, \\
D_1 & := \mathbb{E} \int_{t,x,y} [\mathbf{a}^{ij} \eta'_\delta(\cdot - u)](x, \tilde{u}) \partial_{x_i x_j} \phi_\varepsilon - \mathbb{E} \int_{t,x,y} [\mathbf{a}^{ij} \eta'_\delta(\tilde{u} - \cdot)](y, u) \partial_{y_i y_j} \phi_\varepsilon \\
& \quad + \mathbb{E} \int_{t,x,y} [\mathbf{a}^{ij} \eta'_\delta(\cdot - u)](x, \tilde{u}) \partial_{x_i} \phi_\varepsilon - \mathbb{E} \int_{t,x,y} [\mathbf{a}^{ij} \eta'_\delta(\tilde{u} - \cdot)](y, u) \partial_{y_i} \phi_\varepsilon \\
& \quad - \mathbb{E} \int_{t,x,y} \eta'_\delta(\tilde{u} - u) b^i(x, \tilde{u}) \partial_{x_i} \phi_\varepsilon + \mathbb{E} \int_{t,x,y} \eta'_\delta(\tilde{u} - u) b^i(y, u) \partial_{y_i} \phi_\varepsilon, \\
D_2 & := \mathbb{E} \int_{t,x,y} \frac{1}{2} \eta''_\delta(\tilde{u} - u) \left( \sum_{k=1}^{\infty} |\sigma_{x_i}^{ik}(x, \tilde{u})|^2 + \sum_{k=1}^{\infty} |\sigma_{y_i}^{ik}(y, u)|^2 \right) \phi_\varepsilon, \\
J_1 & := - \mathbb{E} \int_{t,x,y} [f_r^i \eta'_\delta(\cdot - u)](x, \tilde{u}) \partial_{x_i} \phi_\varepsilon + \mathbb{E} \int_{t,x,y} [f_r^i \eta'_\delta(\tilde{u} - \cdot)](y, u) \partial_{y_i} \phi_\varepsilon
\end{aligned}$$

$$\begin{aligned}
& + \mathbb{E} \int_{t,x,y} \eta'_\delta(\tilde{u} - u)(f_{x_i}^i(x, \tilde{u}) - f_{y_i}^i(y, u)) \phi_\varepsilon \\
& - \mathbb{E} \int_{t,x,y} (\llbracket f_{rx_i}^i \eta'_\delta(\cdot - u) \rrbracket(x, \tilde{u}) - \llbracket f_{ry_i}^i \eta'_\delta(\tilde{u} - \cdot) \rrbracket(y, u)) \phi_\varepsilon, \\
J_2 & := \mathbb{E} \int_{t,x,y} \eta'_\delta(\tilde{u} - u)(F(x, \tilde{u}) - F(y, u)) \phi_\varepsilon,
\end{aligned}$$

and

$$\begin{aligned}
\mathcal{E}_1 & := -\mathbb{E} \int_{t,x,y} \partial_{x_j} \phi_\varepsilon \int_u^{\tilde{u}} \int_{\tilde{r}}^u \eta''_\delta(\tilde{r} - r) \sigma_r^{ik}(y, r) \sigma_r^{jk}(x, \tilde{r}) dr d\tilde{r}, \\
\mathcal{E}_2 & := -\mathbb{E} \int_{t,x,y} \partial_{y_i} \phi_\varepsilon \int_u^{\tilde{u}} \int_{\tilde{r}}^u \eta''_\delta(\tilde{r} - r) \sigma_r^{ik}(y, r) \sigma_{rx_j}^{jk}(x, \tilde{r}) dr d\tilde{r}, \\
\mathcal{E}_3 & := \mathbb{E} \int_{t,x,y} \partial_{y_i} \phi_\varepsilon \int_{\tilde{u}}^u \eta''_\delta(\tilde{u} - r) \sigma_r^{ik}(y, r) \sigma_{x_j}^{jk}(x, \tilde{u}) dr, \\
\mathcal{E}_4 & := -\mathbb{E} \int_{t,x,y} \partial_{x_j} \phi_\varepsilon \int_u^{\tilde{u}} \int_{\tilde{r}}^u \eta''_\delta(\tilde{r} - r) \sigma_{ry_i}^{ik}(y, r) \sigma_r^{jk}(x, \tilde{r}) dr d\tilde{r}, \\
\mathcal{E}_5 & := -\mathbb{E} \int_{t,x,y} \phi_\varepsilon \int_u^{\tilde{u}} \int_{\tilde{r}}^u \eta''_\delta(\tilde{r} - r) \sigma_{ry_i}^{ik}(y, r) \sigma_{rx_j}^{jk}(x, \tilde{r}) dr d\tilde{r}, \\
\mathcal{E}_6 & := \mathbb{E} \int_{t,x,y} \phi_\varepsilon \int_{\tilde{u}}^u \eta''_\delta(\tilde{u} - r) \sigma_{ry_i}^{ik}(y, r) \sigma_{x_j}^{jk}(x, \tilde{u}) dr, \\
\mathcal{E}_7 & := \mathbb{E} \int_{t,x,y} \partial_{x_j} \phi_\varepsilon \int_u^{\tilde{u}} \eta''_\delta(\tilde{r} - u) \sigma_{y_i}^{ik}(y, u) \sigma_r^{jk}(x, \tilde{r}) d\tilde{r}, \\
\mathcal{E}_8 & := \mathbb{E} \int_{t,x,y} \phi_\varepsilon \int_u^{\tilde{u}} \eta''_\delta(\tilde{r} - u) \sigma_{y_i}^{ik}(y, u) \sigma_{rx_j}^{jk}(x, \tilde{r}) d\tilde{r}, \\
\mathcal{E}_9 & := -\mathbb{E} \int_{t,x,y} \phi_\varepsilon \eta''_\delta(\tilde{u} - u) \sigma_{y_i}^{ik}(y, u) \sigma_{x_j}^{jk}(x, \tilde{u}),
\end{aligned}$$

and  $u = u(t, y)$  and  $\tilde{u} = \tilde{u}(t, x)$  in the integrand. Now, we estimate these terms. The following estimates are similar to the proof of [DGG19, Theorem 4.1] and [DG20, Theorem 4.1]. The differences are caused by introducing the function  $\psi$  and Dirichlet boundary condition. Therefore, we only focus on the application of the divergence theorem and the term about  $\psi$ . Firstly, using (2.4)-(2.5) in Assumption 2.4, we have

$$\begin{aligned}
(3.6) \quad \mathcal{E}_9 + D_2 & = \mathbb{E} \int_{t,x,y} \frac{1}{2} \eta''_\delta(\tilde{u} - u) \sum_{k=1}^{\infty} (\sigma_{x_j}^{jk}(x, \tilde{u}) - \sigma_{y_i}^{ik}(y, u))^2 \phi_\varepsilon \\
& \lesssim \varepsilon^{2\bar{\kappa}} \delta^{-1} \mathbb{E}(1 + \|u\|_{L^{m+1}(D_T)}^{m+1}) + \delta^2.
\end{aligned}$$

Next, we estimate the terms  $I_1 + I_2$ . From

$$\partial_{y_i} \int_0^{\tilde{u}} \mathbf{a}^2(r) \eta'_\delta(\tilde{u} - r) dr = 0,$$

the support of  $\eta'_\delta$  and  $\varrho_\varepsilon(x - \cdot) \in C_c^\infty(D)$  for all  $(x, \varepsilon) \in (B \cap \bar{D}) \times (0, \bar{\varepsilon})$ , we have

$$I_1 = -\mathbb{E} \int_{t,x,y} \mathbf{1}_{u \leq \tilde{u}} \partial_{x_i} \phi_\varepsilon \int_u^{\tilde{u}} \int_u^{\tilde{u}} \mathbf{1}_{r \leq \tilde{r}} \tilde{\mathbf{a}}^2(\tilde{r}) \eta''_\delta(\tilde{r} - r) d\tilde{r} dr$$

$$\begin{aligned}
& - \mathbb{E} \int_{t,x,y} \mathbf{1}_{u \leq \tilde{u}} \partial_{x_i y_i} \phi_\varepsilon \int_u^{\tilde{u}} \int_u^{\tilde{u}} \mathbf{1}_{r \leq \tilde{r}} \mathbf{a}^2(r) \eta_\delta''(\tilde{r} - r) d\tilde{r} dr \\
& + \mathbb{E} \int_{t,x,y} \varphi \partial_{x_i} (\varrho_\varepsilon \partial_{x_i} \psi) \int_u^{\tilde{u}} \tilde{\mathbf{a}}^2(\tilde{r}) \eta_\delta'(\tilde{r} - u) d\tilde{r} \\
& + \mathbb{E} \int_{t,x,y} \varphi \partial_{y_i} (\varrho_\varepsilon \partial_{x_i} \psi) \int_u^{\tilde{u}} \mathbf{a}^2(r) \eta_\delta'(\tilde{u} - r) dr \\
& =: \sum_{i=1}^4 I_{1,i}.
\end{aligned}$$

Terms  $I_{1,3}$  and  $I_{1,4}$  are arisen from introducing  $\psi$ . The term  $I_{1,3}$  can be written as

$$\begin{aligned}
I_{1,3} & = \mathbb{E} \int_{t,x,y} \varphi(t) \varrho_\varepsilon(x-y) \Delta_x \psi(x) \int_u^{\tilde{u}} \tilde{\mathbf{a}}^2(\tilde{r}) \operatorname{sgn}^+(\tilde{r} - u) d\tilde{r} \\
& + \mathbb{E} \int_{t,x,y} \varphi(t) \varrho_\varepsilon(x-y) \Delta_x \psi(x) \int_u^{\tilde{u}} \mathbf{1}_{0 \leq \tilde{r} - u \leq \delta} \tilde{\mathbf{a}}^2(\tilde{r}) (\eta_\delta'(\tilde{r} - u) - \operatorname{sgn}^+(\tilde{r} - u)) d\tilde{r} \\
& + \mathbb{E} \int_{t,x,y} \varphi(t) \partial_{x_i} \varrho_\varepsilon(x-y) \partial_{x_i} \psi(x) \int_u^{\tilde{u}} \tilde{\mathbf{a}}^2(\tilde{r}) \operatorname{sgn}^+(\tilde{r} - u) d\tilde{r} \\
& + \mathbb{E} \int_{t,x,y} \varphi(t) \partial_{x_i} \varrho_\varepsilon(x-y) \partial_{x_i} \psi(x) \int_u^{\tilde{u}} \mathbf{1}_{0 \leq \tilde{r} - u \leq \delta} \tilde{\mathbf{a}}^2(\tilde{r}) (\eta_\delta'(\tilde{r} - u) - \operatorname{sgn}^+(\tilde{r} - u)) d\tilde{r} \\
& =: \sum_{i=1}^4 I_{1,3,i},
\end{aligned}$$

where

$$\operatorname{sgn}^+(x) := \begin{cases} 1, & x > 0; \\ 0, & \text{otherwise.} \end{cases}$$

Combining the boundness of  $\Delta \psi$  and  $\varphi$ , the definition of  $\varrho_\varepsilon$  and Assumption 2.1, we obtain that

$$\begin{aligned}
|I_{1,3,2}| + |I_{1,3,4}| & \lesssim \delta \mathbb{E} \int_{t,x,y} (\varrho_\varepsilon(x-y) + |\partial_{x_i} \varrho_\varepsilon(x-y)|) \sup_{0 \leq \tilde{r} - u \leq \delta} \tilde{\mathbf{a}}^2(\tilde{r}) \\
& \lesssim \delta (1 + \varepsilon^{-1}) \mathbb{E} (1 + \|u\|_{L^m(D_T)}^m).
\end{aligned}$$

Therefore, noticing  $\varepsilon < 1$ , we have

$$\begin{aligned}
I_{1,3} & \leq \mathbb{E} \int_{t,x,y} \varphi(t) \varrho_\varepsilon(x-y) \Delta_x \psi(x) (\tilde{\Phi}(\tilde{u}) - \tilde{\Phi}(u))^+ \\
& + \mathbb{E} \int_{t,x,y} \varphi(t) \partial_{x_i} \varrho_\varepsilon(x-y) \partial_{x_i} \psi(x) (\tilde{\Phi}(\tilde{u}) - \tilde{\Phi}(u))^+ + \delta \varepsilon^{-1} \mathbb{E} (1 + \|u\|_{L^m(D_T)}^m).
\end{aligned}$$

With the same method, we also have

$$I_{1,4} = -\mathbb{E} \int_{t,x,y} \varphi(t) \partial_{x_i} \varrho_\varepsilon(x-y) \partial_{x_i} \psi(x) (\Phi(\tilde{u}) - \Phi(u))^+ + \delta \varepsilon^{-1} \mathbb{E} (1 + \|\tilde{u}\|_{L^m(D_T)}^m).$$

Using the triangle inequality, the definition of  $\varrho_\varepsilon$ , the boundness of  $\partial_{x_i} \psi$  and the fact

$$|\Phi(r) - \tilde{\Phi}(r)| \lesssim \lambda |r|^{\frac{m+1}{2}} + \mathbf{1}_{|r| \geq R_\lambda} |r|^m, \quad \forall r \in \mathbb{R},$$

we have

$$\mathbb{E} \int_{t,x,y} \varphi(t) \partial_{x_i} \varrho_\varepsilon(x-y) \partial_{x_i} \psi(x) ((\tilde{\Phi}(\tilde{u}) - \tilde{\Phi}(u))^+ - (\Phi(\tilde{u}) - \Phi(u))^+)$$

$$\begin{aligned} &\leq \mathbb{E} \int_{t,x,y} \varphi(t) \partial_{x_i} \varrho_\varepsilon(x-y) \partial_{x_i} \psi(x) ((\Phi(u) - \tilde{\Phi}(u))^+ + (\tilde{\Phi}(\tilde{u}) - \Phi(\tilde{u}))^+) \\ &\lesssim \varepsilon^{-1} \lambda \mathbb{E} (\|u\|_{L_{m+1}(D_T)}^{m+1} + \|\tilde{u}\|_{L_{m+1}(D_T)}^{m+1}) + \varepsilon^{-1} \mathbb{E} (\|\mathbf{1}_{|u| \geq R_\lambda} u\|_{L_m(D_T)}^m + \|\mathbf{1}_{|\tilde{u}| \geq R_\lambda} \tilde{u}\|_{L_m(D_T)}^m). \end{aligned}$$

Similarly, for the remaining term in  $I_{1,3}$ , we have

$$(3.7) \quad \begin{aligned} &\mathbb{E} \int_{t,x,y} \varphi(t) \varrho_\varepsilon(x-y) \Delta_x \psi(x) ((\tilde{\Phi}(\tilde{u}) - \tilde{\Phi}(u))^+ - (\tilde{\Phi}(\tilde{u}) - \Phi(u))^+) \\ &\lesssim \lambda \|u\|_{L_{m+1}(D_T)}^{m+1} + \mathbb{E} \|\mathbf{1}_{|u| \geq R_\lambda} u\|_{L_m(D_T)}^m. \end{aligned}$$

To estimate the new term in (3.7), using (6.8) in Lemma 6.2, we have

$$\begin{aligned} &\mathbb{E} \int_{t,x,y} \varphi(t) \varrho_\varepsilon(x-y) \Delta_x \psi(x) (\tilde{\Phi}(\tilde{u}(t,x)) - \Phi(u(t,y)))^+ \\ &\leq \mathbb{E} \int_{t,x} \varphi(t) \Delta_x \psi(x) (\tilde{\Phi}(\tilde{u}(t,x)) - \Phi(u(t,x)))^+ \\ &\quad + C \varepsilon^{\frac{1}{m+1}} \mathbb{E} (1 + \|u\|_{L_{m+1}(D_T)}^{m+1} + \|\nabla[\mathbf{a}](u)\|_{L_2(D_T)}^2). \end{aligned}$$

For  $I_2$ , using Definition 2.2 (ii) and applying [DGG19, Remark 3.1], we have

$$(3.8) \quad \begin{aligned} I_2 &\leq -2 \int_{t,x,y} \eta_\delta''(\tilde{u}-u) \nabla_x [\tilde{\mathbf{a}}](\tilde{u}) \cdot \nabla_y [\mathbf{a}](u) \phi_\varepsilon \\ &= 2 \int_{t,x,y} \phi_\varepsilon \partial_{x_i} [\tilde{\mathbf{a}}](\tilde{u}) \cdot \partial_{y_i} \int_u^{\tilde{u}} \eta_\delta''(\tilde{u}-r) \mathbf{a}(r) dr \\ &= 2 \int_{t,x,y} \partial_{x_i y_i} \phi_\varepsilon \int_0^{\tilde{u}} \int_u^{\tilde{r}} \eta_\delta''(\tilde{r}-r) \tilde{\mathbf{a}}(\tilde{r}) \mathbf{a}(r) d\tilde{r} dr \\ &= 2 \int_{t,x,y} \mathbf{1}_{u \leq \tilde{u}} \partial_{x_i y_i} \phi_\varepsilon \int_u^{\tilde{u}} \int_u^{\tilde{u}} \eta_\delta''(\tilde{r}-r) \tilde{\mathbf{a}}(\tilde{r}) \mathbf{a}(r) d\tilde{r} dr. \end{aligned}$$

Then, we have

$$I_{1,1} + I_{1,2} + I_2 \leq 2 \int_{t,x,y} \mathbf{1}_{u \leq \tilde{u}} |\partial_{x_i y_i} \phi_\varepsilon| \int_u^{\tilde{u}} \int_u^{\tilde{u}} \eta_\delta''(\tilde{r}-r) |\mathbf{a}(r) - \tilde{\mathbf{a}}(\tilde{r})|^2 d\tilde{r} dr.$$

Based on the estimates of  $|\mathbf{a}(r) - \tilde{\mathbf{a}}(\tilde{r})|^2$  in the proof of [DGG19, Theorem 4.1] which using Assumption 2.1, combining the proceeding estimates and noticing  $\varepsilon < 1$ , we have

$$(3.9) \quad \begin{aligned} \sum_{i=1}^2 I_i &\leq C(\delta \varepsilon^{-1} + \varepsilon^{\frac{1}{m+1}} + \varepsilon^{-2} \lambda^2 + \varepsilon^{-1} \lambda + \varepsilon^{-2} \delta^{2\alpha}) \mathbb{E} (1 + \|u\|_{L_{m+1}(D_T)}^{m+1} + \|\tilde{u}\|_{L_{m+1}(D_T)}^{m+1}) \\ &\quad + C \varepsilon^{-2} \mathbb{E} \left( \|\mathbf{1}_{|u| \geq R_\lambda} (1+u)\|_{L_m(D_T)}^m + \|\mathbf{1}_{|\tilde{u}| \geq R_\lambda} (1+\tilde{u})\|_{L_m(D_T)}^m \right) \\ &\quad + \mathbb{E} \int_{t,x} \varphi(t) \Delta_x \psi(x) (\tilde{\Phi}(\tilde{u}(t,x)) - \Phi(u(t,x)))^+ + C \varepsilon^{\frac{1}{m+1}} \mathbb{E} \|\nabla[\mathbf{a}](u)\|_{L_2(D_T)}^2, \end{aligned}$$

where  $\alpha \in (0, 1 \wedge (m/2))$ . Now, we focus on the terms  $D_1$ . With the fact that  $\phi_\varepsilon(t, x, \cdot) \in C_c^\infty(D)$  for any  $(t, x, \varepsilon) \in D_T \times (0, \bar{\varepsilon})$ , using the divergence theorem in  $y$ , we have

$$\mathbb{E} \int_{t,x,y} \partial_{y_i y_j} \phi_\varepsilon \int_0^{\tilde{u}} \eta_\delta'(\tilde{u}-r) a^{ij}(y, r) dr + \mathbb{E} \int_{t,x,y} \partial_{y_i} \phi_\varepsilon \int_0^{\tilde{u}} \eta_\delta'(\tilde{u}-r) a_{y_j}^{ij}(y, r) dr = 0.$$

Then, from the support of  $\eta'_\delta$ , we have

$$\begin{aligned}
D_1 &= -\mathbb{E} \int_{t,x,y} \mathbf{1}_{u \leq \tilde{u}} \partial_{x_i y_j} \phi_\varepsilon \int_u^{\tilde{u}} \int_u^{\tilde{u}} \eta''_\delta(\tilde{r} - r) (a^{ij}(y, r) + a^{ij}(x, \tilde{r})) d\tilde{r} dr \\
&\quad - \mathbb{E} \int_{t,x,y} \partial_{x_i} \phi_\varepsilon \eta'_\delta(\tilde{u} - u) b^i(x, \tilde{u}) + \mathbb{E} \int_{t,x,y} \partial_{y_i} \phi_\varepsilon \eta'_\delta(\tilde{u} - u) b^i(y, u) \\
&\quad + \mathbb{E} \int_{t,x,y} \partial_{x_i} \phi_\varepsilon \int_u^{\tilde{u}} a_{x_j}^{ij}(x, \tilde{r}) \eta'_\delta(\tilde{r} - u) d\tilde{r} + \mathbb{E} \int_{t,x,y} \partial_{y_i} \phi_\varepsilon \int_u^{\tilde{u}} a_{y_j}^{ij}(y, r) \eta'_\delta(\tilde{u} - r) dr \\
&\quad + \mathbb{E} \int_{t,x,y} \mathbf{1}_{u \leq \tilde{u}} \varphi(t) \partial_{x_i} (\varrho_\varepsilon(x - y) \partial_{x_j} \psi(x)) \int_u^{\tilde{u}} \eta'_\delta(\tilde{r} - u) a^{ij}(x, \tilde{r}) d\tilde{r} \\
&\quad + \mathbb{E} \int_{t,x,y} \mathbf{1}_{u \leq \tilde{u}} \varphi(t) \partial_{y_j} \varrho_\varepsilon(x - y) \partial_{x_i} \psi(x) \int_u^{\tilde{u}} \int_u^{\tilde{u}} \eta''_\delta(\tilde{r} - r) a^{ij}(y, r) d\tilde{r} dr \\
&=: \sum_{i=1}^5 D_{1,i}.
\end{aligned}$$

Firstly, with (2.4)-(2.5) in Assumption 2.4, we have

$$\begin{aligned}
D_{1,4} + D_{1,5} &= \mathbb{E} \int_{t,x,y} \mathbf{1}_{u \leq \tilde{u}} \varphi \varrho_\varepsilon \partial_{x_i x_j} \psi \int_u^{\tilde{u}} \eta'_\delta(\tilde{r} - u) a^{ij}(x, \tilde{r}) d\tilde{r} \\
&\quad + \mathbb{E} \int_{t,x,y} \mathbf{1}_{u \leq \tilde{u}} \varphi (\partial_{x_j} \varrho_\varepsilon \partial_{x_i} \psi) \int_u^{\tilde{u}} \int_u^{\tilde{u}} \eta''_\delta(\tilde{r} - r) (a^{ij}(x, \tilde{r}) - a^{ij}(y, r)) d\tilde{r} dr \\
&\lesssim \mathbb{E} \int_{t,x,y} \mathbf{1}_{B \cap \bar{D}}(x) \varphi(t) \left( \varepsilon \sum_i |\partial_{x_i} \varrho_\varepsilon(x - y)| + \varrho_\varepsilon(x - y) \right) \cdot (\tilde{u} - u)^+ \\
&\quad + \delta^\beta \varepsilon^{-1} \mathbb{E} \|\tilde{u}\|_{L_1(D_T)}.
\end{aligned}$$

Moreover, if  $|r - \tilde{r}| \leq \delta$ , from the definition of  $a^{ij}$  and (2.5) in Assumption 2.4, we have

$$\begin{aligned}
&\partial_{x_i y_j} \phi_\varepsilon (a^{ij}(x, \tilde{r}) + a^{ij}(y, r) - \sigma_r^{ik}(x, \tilde{r}) \sigma_r^{jk}(y, r)) \\
&= \frac{1}{2} \varphi \psi \partial_{x_i y_j} \varrho_\varepsilon (\sigma_r^{ik}(x, \tilde{r}) - \sigma_r^{ik}(y, r)) (\sigma_r^{jk}(x, \tilde{r}) - \sigma_r^{jk}(y, r)) \\
&\quad + \frac{1}{2} \varphi \partial_{y_j} \varrho_\varepsilon \cdot \partial_{x_i} \psi (\sigma_r^{ik}(x, \tilde{r}) (\sigma_r^{jk}(x, \tilde{r}) - \sigma_r^{jk}(y, r)) - \sigma_r^{jk}(y, r) (\sigma_r^{ik}(x, \tilde{r}) - \sigma_r^{ik}(y, r))) \\
&\lesssim (\varepsilon^2 + \delta^{2\beta}) \mathbf{1}_{B \cap \bar{D}}(x) \varphi(t) \sum_{i,j} |\partial_{x_i y_j} \varrho_\varepsilon(x - y)| + (\varepsilon + \delta^\beta) \mathbf{1}_{B \cap \bar{D}}(x) \varphi(t) \sum_j |\partial_{y_j} \varrho_\varepsilon(x - y)|.
\end{aligned}$$

Therefore,

$$\begin{aligned}
(3.10) \quad D_{1,1} + \mathcal{E}_1 &\lesssim \mathbb{E} \int_{t,x,y} \mathbf{1}_{B \cap \bar{D}}(x) \varphi(t) \left( \varepsilon^2 \sum_{i,j} |\partial_{x_i y_j} \varrho_\varepsilon(x - y)| + \varepsilon \sum_i |\partial_{x_i} \varrho_\varepsilon(x - y)| \right) (\tilde{u} - u)^+ \\
&\quad + (\delta^{2\beta} \varepsilon^{-2} + \delta^\beta \varepsilon^{-1}) \mathbb{E} \|\tilde{u}\|_{L_1(D_T)}.
\end{aligned}$$

Similarly, we have

$$\begin{aligned}
D_{1,3} + \mathcal{E}_2 + \mathcal{E}_4 &= \mathbb{E} \int_{t,x,y} \mathbf{1}_{u \leq \tilde{u}} \varphi \partial_{x_i} \varrho_\varepsilon \psi \int_u^{\tilde{u}} \int_u^{\tilde{u}} \eta''_\delta(\tilde{r} - r) (a_{x_j}^{ij}(x, \tilde{r}) - a_{y_j}^{ij}(y, r)) d\tilde{r} dr
\end{aligned}$$



$$\begin{aligned}
& + \mathbb{E} \int_{t,x,y} \varphi \varrho_\varepsilon \partial_{x_i} \psi \int_u^{\tilde{u}} a_{x_j}^{ij}(x, \tilde{r}) \eta'_\delta(\tilde{r} - u) d\tilde{r}, \\
& + \mathbb{E} \int_{t,x,y} \mathbf{1}_{u \leq \tilde{u}} \varphi \partial_{x_i} \varrho_\varepsilon \psi \int_u^{\tilde{u}} \int_u^{\tilde{u}} \eta''_\delta(\tilde{r} - r) (\sigma_r^{ik}(x, \tilde{r}) \sigma_{ry_j}^{jk}(y, r) - \sigma_r^{ik}(y, r) \sigma_{rx_j}^{jk}(x, \tilde{r})) dr d\tilde{r} \\
& + \mathbb{E} \int_{t,x,y} \mathbf{1}_{u \leq \tilde{u}} \varphi \varrho_\varepsilon \partial_{x_i} \psi \int_u^{\tilde{u}} \int_u^{\tilde{u}} \eta''_\delta(\tilde{r} - r) \sigma_{ry_j}^{jk}(y, r) \sigma_r^{ik}(x, \tilde{r}) dr d\tilde{r},
\end{aligned}$$

with relabeling  $i \leftrightarrow j$  in  $\mathcal{E}_4$ . From (2.4)-(2.5) in Assumption 2.4, we have

$$\begin{aligned}
(3.11) \quad & D_{1,3} + \mathcal{E}_2 + \mathcal{E}_4 \\
& \lesssim \delta^\beta \varepsilon^{-1} \mathbb{E}(\|u\|_{L_1(D_T)} + \|\tilde{u}\|_{L_1(D_T)}) \\
& + \mathbb{E} \int_{t,x,y} \mathbf{1}_{B \cap \overline{D}}(x) \varphi(t) \left( \varepsilon \sum_i |\partial_{x_i} \varrho_\varepsilon(x - y)| + \varrho_\varepsilon(x - y) \right) (\tilde{u} - u)^+.
\end{aligned}$$

For  $D_{1,2} + \mathcal{E}_3 + \mathcal{E}_7$ , define

$$\begin{aligned}
D_{1,2} = & -\mathbb{E} \int_{t,x,y} \partial_{x_i} \phi_\varepsilon \eta'_\delta(\tilde{u} - u) b^i(x, \tilde{u}) \\
& + \mathbb{E} \int_{t,x,y} \partial_{y_i} \phi_\varepsilon \eta'_\delta(\tilde{u} - u) b^i(y, u) =: D_{1,2,1} + D_{1,2,2}.
\end{aligned}$$

Using the definition of  $b^i$  and (2.4)-(2.5) in Assumption 2.4 and relabeling  $i \leftrightarrow j$  in  $\mathcal{E}_7$ , we have

$$\begin{aligned}
D_{1,2,2} + \mathcal{E}_7 = & \mathbb{E} \int_{t,x,y} \varphi \partial_{y_i} \varrho_\varepsilon \psi \int_u^{\tilde{u}} \eta''_\delta(\tilde{r} - u) \sigma_{y_j}^{jk}(y, u) (\sigma_r^{ik}(y, u) - \sigma_r^{ik}(x, u)) d\tilde{r} \\
& + \mathbb{E} \int_{t,x,y} \varphi \partial_{y_i} \varrho_\varepsilon \psi \int_u^{\tilde{u}} \eta''_\delta(\tilde{r} - u) \sigma_{y_j}^{jk}(y, u) (\sigma_r^{ik}(x, u) - \sigma_r^{ik}(x, \tilde{r})) d\tilde{r} \\
& + \mathbb{E} \int_{t,x,y} \varphi \varrho_\varepsilon \partial_{x_i} \psi \int_u^{\tilde{u}} \eta''_\delta(\tilde{r} - u) \sigma_{y_j}^{jk}(y, u) \sigma_r^{ik}(y, u) d\tilde{r} \\
& - \mathbb{E} \int_{t,x,y} \varphi \varrho_\varepsilon \partial_{x_i} \psi \int_u^{\tilde{u}} \eta''_\delta(\tilde{r} - u) \sigma_{y_j}^{jk}(y, u) (\sigma_r^{ik}(y, u) - \sigma_r^{ik}(x, \tilde{r})) d\tilde{r} \\
\leq & \mathbb{E} \int_{t,x,y} \varphi \partial_{y_i} \varrho_\varepsilon \psi \eta'_\delta(\tilde{u} - u) \sigma_{y_j}^{jk}(y, u) (y_l - x_l) \int_0^1 \sigma_{rx_l}^{ik}(x + \theta(y - x), u) d\theta \\
& + C \delta^\beta \varepsilon^{-1} (1 + \mathbb{E}\|u\|_{L_1(D_T)}) \\
& + \mathbb{E} \int_{t,x,y} \varphi \varrho_\varepsilon \partial_{x_i} \psi \eta'_\delta(\tilde{u} - u) b^i(y, u) + C(\delta^\beta + \varepsilon)(1 + \mathbb{E}\|u\|_{L_1(D_T)}).
\end{aligned}$$

Similarly, we have

$$\begin{aligned}
D_{1,2,1} + \mathcal{E}_3 = & -\mathbb{E} \int_{t,x,y} \varphi \partial_{y_i} \varrho_\varepsilon \psi \int_u^{\tilde{u}} \eta''_\delta(\tilde{u} - r) \sigma_{x_j}^{jk}(x, \tilde{u}) (\sigma_r^{ik}(y, r) - \sigma_r^{ik}(x, \tilde{u})) dr \\
& - \mathbb{E} \int_{t,x,y} \varphi \varrho_\varepsilon \partial_{x_i} \psi \int_u^{\tilde{u}} \eta''_\delta(\tilde{u} - r) \sigma_r^{ik}(x, \tilde{u}) \sigma_{x_j}^{jk}(x, \tilde{u}) dr \\
\leq & -\mathbb{E} \int_{t,x,y} \varphi \partial_{y_i} \varrho_\varepsilon \psi \eta'_\delta(\tilde{u} - u) \sigma_{x_j}^{jk}(x, \tilde{u}) (y_l - x_l) \int_0^1 \sigma_{rx_l}^{ik}(x + \theta(y - x), \tilde{u}) d\theta \\
& + C \delta^\beta \varepsilon^{-1} (1 + \mathbb{E}\|\tilde{u}\|_{L_1(D_T)}) - \mathbb{E} \int_{t,x,y} \varphi \varrho_\varepsilon \partial_{x_i} \psi \eta'_\delta(\tilde{u} - u) b^i(x, \tilde{u}).
\end{aligned}$$

From (2.4)-(2.6) in Assumption 2.4, we have

$$\begin{aligned} & |\sigma_{y_j}^{jk}(y, u) \int_0^1 \sigma_{rx_i}^{ik}(x + \theta(y-x), u) d\theta - \sigma_{x_j}^{jk}(x, \tilde{u}) \int_0^1 \sigma_{rx_i}^{ik}(x + \theta(y-x), \tilde{u}) d\theta| \\ & \lesssim |x-y|(1+|u|+|\tilde{u}|) + |\sigma_{y_j}^{jk}(y, u) \sigma_{ry_i}^{ik}(y, u) - \sigma_{x_j}^{jk}(x, \tilde{u}) \sigma_{rx_i}^{ik}(x, \tilde{u})| \\ & \lesssim (|x-y| + |x-y|^{\bar{\kappa}})(1+|u|+|\tilde{u}|) + |u-\tilde{u}|, \end{aligned}$$

and

$$|b^i(y, u) - b^i(x, \tilde{u})| \lesssim |u-\tilde{u}| + |x-y|^{\bar{\kappa}} + |x-y|(1+|u|).$$

Therefore, using the support of  $\eta'_\delta$ , we have

$$\begin{aligned} (3.12) \quad & D_{1,2} + \mathcal{E}_3 + \mathcal{E}_7 \\ & \lesssim (\delta^\beta \varepsilon^{-1} + \varepsilon^{\bar{\kappa}}) \mathbb{E}(1 + \|u\|_{L_1(D_T)} + \|\tilde{u}\|_{L_1(D_T)}) \\ & \quad + \mathbb{E} \int_{t,x,y} \mathbf{1}_{B \cap \bar{D}}(x) \left( \varepsilon \sum_i |\partial_{x_i} \varrho_\varepsilon(x-y)| + \varrho_\varepsilon(x-y) \right) \varphi(t) (\tilde{u}-u)^+. \end{aligned}$$

The remaining terms in (3.5) are  $J_1$ ,  $J_2$ ,  $\mathcal{E}_5$ ,  $\mathcal{E}_6$  and  $\mathcal{E}_8$ . Using (2.5) in Assumption 2.4 and the support of  $\eta'_\delta$ , we have

$$\mathcal{E}_5 \lesssim \mathbb{E} \int_{t,x,y} \mathbf{1}_{B \cap \bar{D}}(x) \varphi(t) \varrho_\varepsilon(x-y) (\tilde{u}-u)^+,$$

and

$$\begin{aligned} (3.13) \quad & \mathcal{E}_6 + \mathcal{E}_8 \\ & \leq \mathbb{E} \int_{t,x,y} \phi_\varepsilon \int_{\tilde{u}}^u \eta''_\delta(\tilde{u}-r) \sigma_{ry_i}^{ik}(y, \tilde{u}) \sigma_{x_j}^{jk}(x, \tilde{u}) dr + C\delta \mathbb{E}(1 + \|\tilde{u}\|_{L_1(D_T)}) \\ & \quad + \mathbb{E} \int_{t,x,y} \phi_\varepsilon \int_u^{\tilde{u}} \eta''_\delta(\tilde{r}-u) \sigma_{y_i}^{ik}(y, u) \sigma_{rx_j}^{jk}(x, u) d\tilde{r} + C\delta \mathbb{E}(1 + \|u\|_{L_1(D_T)}) \\ & \lesssim \mathbb{E} \int_{t,x,y} \phi_\varepsilon \eta'_\delta(\tilde{u}-u) |\sigma_{y_i}^{ik}(y, u) \sigma_{rx_j}^{jk}(x, u) - \sigma_{ry_i}^{ik}(y, \tilde{u}) \sigma_{x_j}^{jk}(x, \tilde{u})| \\ & \quad + \delta \mathbb{E}(1 + \|u\|_{L_1(D_T)} + \|\tilde{u}\|_{L_1(D_T)}) \\ & \lesssim (\delta + \varepsilon^{\bar{\kappa}}) \mathbb{E}(1 + \|u\|_{L_1(D_T)} + \|\tilde{u}\|_{L_1(D_T)}) + \mathbb{E} \int_{t,x,y} \mathbf{1}_{B \cap \bar{D}}(x) \varphi(t) \varrho_\varepsilon(x-y) (\tilde{u}-u)^+. \end{aligned}$$

From (2.7)-(2.9) in Assumption 2.4, we have

$$\begin{aligned} (3.14) \quad & J_1 = \mathbb{E} \int_{t,x,y} \mathbf{1}_{u \leq \tilde{u}} \varphi(t) \partial_{x_i} \varrho_\varepsilon(x-y) \psi(x) \int_u^{\tilde{u}} \int_u^{\tilde{u}} \eta''_\delta(\tilde{r}-r) (f_r^i(y, r) - f_r^i(x, \tilde{r})) d\tilde{r} dr \\ & \quad - \mathbb{E} \int_{t,x,y} \varphi(t) \varrho_\varepsilon(x-y) \partial_{x_i} \psi(x) \int_u^{\tilde{u}} f_r^i(x, \tilde{r}) \eta'_\delta(\tilde{r}-u) d\tilde{r} \\ & \quad + \mathbb{E} \int_{t,x,y} \eta'_\delta(\tilde{u}-u) (f_{x_i}^i(x, \tilde{u}) - f_{y_i}^i(y, u)) \phi_\varepsilon \\ & \quad - \mathbb{E} \int_{t,x,y} \phi_\varepsilon \left( \int_u^{\tilde{u}} f_{rx_i}^i(x, \tilde{r}) \eta'_\delta(\tilde{r}-u) d\tilde{r} + \int_u^{\tilde{u}} f_{ry_i}^i(y, r) \eta'_\delta(\tilde{u}-r) dr \right) \\ & \lesssim (\varepsilon^{\bar{\beta}} + \delta^\beta \varepsilon^{-1}) \mathbb{E}(1 + \|u\|_{L_1(D_T)} + \|\tilde{u}\|_{L_1(D_T)}) \\ & \quad + \mathbb{E} \int_{t,x,y} \mathbf{1}_{B \cap \bar{D}}(x) \varphi(t) \left( \varepsilon |\partial_{x_i} \varrho_\varepsilon(x-y)| + \varrho_\varepsilon(x-y) \right) (\tilde{u}-u)^+. \end{aligned}$$

With (2.10) in Assumption 2.4, we have

$$(3.15) \quad J_2 \lesssim \varepsilon^{\bar{\beta}} + \mathbb{E} \int_{t,x,y} \mathbf{1}_{B \cap \bar{D}}(x) \varphi(t) \varrho_\varepsilon(x-y) (\tilde{u} - u)^+.$$

Combining (3.5) with  $|\eta_\delta(r) - r^+| \leq \delta$ , (3.6) and (3.9)-(3.15), one has (3.1).

For (3.2), the proof is a similar procedure but with a different  $(\star)$ -property. Specifically, for each  $(z, t, x) \in [0, \infty) \times D_T$ , since

$$(\eta_\delta(z - \cdot), \rho_\theta(\cdot - t) \varphi(\frac{\cdot + t}{2}), \varrho_\varepsilon(x - \cdot) \psi(x) \mathbf{1}_{\bar{D}}(x)) \in \mathcal{E} \times C_c((0, T)) \times C_c^\infty(D)$$

for  $\varepsilon \in (0, \bar{\varepsilon})$  and small enough  $\theta$ , we apply the entropy formulation (2.3) of  $\tilde{u}(s, y)$  with  $(\eta_\delta(z - r), \phi_{\theta, \varepsilon}(t, x, \cdot, \cdot))$  instead of  $(\eta(r), \phi)$ . Taking  $z = u(t, x)$  by convolution, integrating over  $(t, x) \in D_T$  and taking expectations, we acquire an estimate of  $-\mathbb{E} \int_{t,x,s,y} \eta_\delta(u(t, x) - \tilde{u}(s, y)) \partial_t \phi_{\theta, \varepsilon}$ .

Similarly, fix  $(z, s, y) \in [0, \infty) \times D_T$ . Since

$$(\eta_\delta(\cdot - z), \rho_\theta(s - \cdot) \varphi(\frac{s + \cdot}{2}), \varrho_\varepsilon(\cdot - y) \psi(\cdot) \mathbf{1}_{\bar{D}}(\cdot)) \in \mathcal{E}_0 \times C_c((0, T)) \times C^\infty(\bar{D})$$

for small enough  $\theta$ , we use the entropy formulation (2.3) of  $u(t, x)$  with  $(\eta_\delta(r - z), \phi_{\theta, \varepsilon}(\cdot, \cdot, s, y))$  instead of  $(\eta(r), \phi)$ . Then, taking  $z = \tilde{u}(s, y)$  by convolution, integrating over  $(s, y) \in D_T$  and taking expectations, we also obtain an estimate of  $-\mathbb{E} \int_{t,x,s,y} \eta_\delta(u(t, x) - \tilde{u}(s, y)) \partial_s \phi_{\theta, \varepsilon}$ .

Note that  $\phi_{\theta, \varepsilon}$  only acts on the set  $\{s, t \in [0, T] : s > t\}$ , the stochastic integral term in the estimate of  $-\mathbb{E} \int_{t,x,s,y} \eta_\delta(u(t, x) - \tilde{u}(s, y)) \partial_t \phi_{\theta, \varepsilon}$  vanishes, while the one in the estimate of  $-\mathbb{E} \int_{t,x,s,y} \eta_\delta(u(t, x) - \tilde{u}(s, y)) \partial_s \phi_{\theta, \varepsilon}$  may not be zeros. Therefore, we apply the  $(\star)$ -property (iii) of  $\tilde{u}$  with  $h(r) = \eta'_\delta(r)$  (then  $h \in \mathcal{C}^+$ ) and  $g(x, y) = \mathbf{1}_{\bar{D}}(x) \psi(x) \varrho_\varepsilon(x - y)$ , then the stochastic integral terms is controlled by  $C\theta^{1-\mu} + \tilde{\mathcal{E}}(u, \tilde{u}, \theta)$ . After combining two estimates and taking the limit  $\theta \rightarrow 0^+$ , there is only one time variable in the integrand. The remaining terms can be estimated as in the proof of (3.1). Hence, this lemma is proved.  $\square$

*Remark 3.12.* In the whole proof of Lemma 3.10, the divergence theorem is applied mostly in  $y$  with  $\varrho_\varepsilon(x - \cdot) \in C_c^\infty(D)$  for all  $(x, \varepsilon) \in (B \cap \bar{D}) \times (0, \bar{\varepsilon})$ , except (3.8), in which the divergence theorem in  $x$  is used with the zero boundary condition of  $\tilde{u}$ .

**Theorem 3.13.** ( *$L_1$ -estimates*) Let  $0 \leq \xi, \tilde{\xi} \in L_{m+1}(\Omega, \mathcal{F}_0; L_{m+1}(D))$ . Suppose  $u$  and  $\tilde{u}$  are the entropy solutions to the Dirichlet problems  $\Pi(\Phi, \xi)$  and  $\Pi(\tilde{\Phi}, \tilde{\xi})$ , respectively. Let Assumptions 2.1, 2.4 and 2.5 hold for both  $\Phi$  and  $\tilde{\Phi}$ . If  $u$  or  $\tilde{u}$  has the  $(\star)$ -property. Then,

(1) if furthermore  $\Phi = \tilde{\Phi}$ , then

$$(3.16) \quad \operatorname{ess\,sup}_{t \in [0, T]} \mathbb{E} \|(\tilde{u}(t, \cdot) - u(t, \cdot))^+\|_{L_1(D)} \leq C \mathbb{E} \|(\tilde{\xi} - \xi)^+\|_{L_1(D)},$$

where the constant  $C$  depends only on  $N_0, K, d, T$  and  $|D|$ .

(2) for all  $\delta \in (0, 1)$ ,  $\varepsilon \in (0, \bar{\varepsilon})$ ,  $\lambda \in [0, 1]$  and  $\alpha \in (0, 1 \wedge (m/2))$ , we have that

$$(3.17) \quad \begin{aligned} & \mathbb{E} \int_0^T \int_x (\tilde{u}(\tau, x) - u(\tau, x))^+ d\tau \\ & \leq C \mathbb{E} \int_x (\tilde{\xi}(x) - \xi(x))^+ + C \sup_{|h| \leq 2\varepsilon} \mathbb{E} \|\xi(\cdot) - \tilde{\xi}(\cdot + h)\|_{L_1(D)} \\ & \quad + C \sup_{|h| \leq 2\varepsilon} \mathbb{E} \|\tilde{\xi}(\cdot) - \tilde{\xi}(\cdot + h)\|_{L_1(D)} \\ & \quad + C \varepsilon^{\frac{1}{m+1}} \mathbb{E} (\|\nabla[\mathbf{a}](u)\|_{L_2(D_T)}^2 + \|\nabla[\tilde{\mathbf{a}}](\tilde{u})\|_{L_2(D_T)}^2) \\ & \quad + C \mathfrak{C}(\varepsilon, \delta, \lambda, \alpha) \mathbb{E} (1 + \|u\|_{L_{m+1}(D_T)}^{m+1} + \|\tilde{u}\|_{L_{m+1}(D_T)}^{m+1}) \end{aligned}$$

$$+ C\varepsilon^{-2}\mathbb{E}\left(\|\mathbf{1}_{|u|\geq R_\lambda}(1+u)\|_{L_m(D_T)}^m + \|\mathbf{1}_{|\tilde{u}|\geq R_\lambda}(1+\tilde{u})\|_{L_m(D_T)}^m\right),$$

where  $\mathfrak{C}(\varepsilon, \delta, \lambda, \alpha)$  and  $R_\lambda$  are introduced in Lemma 3.10,

$$(3.18) \quad \bar{\xi}(x) := \begin{cases} \xi(x), & x \in D; \\ 0, & x \notin D, \end{cases}$$

and the constant  $C$  depends only on  $N_0, K, d, T$  and  $|D|$ .

*Remark 3.14.* Since the positive part function is not an even function, it requires to prove estimates with the  $(\star)$ -property of different entropy solution  $u$  or  $\tilde{u}$ . Using this fact, we obtain the  $L_1$ -estimates with only one of the entropy solutions has the  $(\star)$ -property, which is a key point in proving the uniqueness in the proof of Theorem 2.6. See Remark 5.1 for more detail.

*Proof.* We first assume that  $\tilde{u}$  has the  $(\star)$ -property. Let  $0 < s < \tau < T$  be Lebesgue points of the function

$$t \rightarrow \mathbb{E} \int_{x,y} (\tilde{u}(t,x) - u(t,y))^+ \psi(x) \varrho_\varepsilon(x-y),$$

and fix a constant  $\gamma \in (0, (\tau - s) \vee (T - \tau))$ . Choose a sequence of functions  $\{\phi_n\}_{n \in \mathbb{N}}$  and the limit  $V_{(\gamma)}$  as in the proof of Proposition 2.8, take  $\varphi = \phi_n$  in (3.1) and pass the limits  $n \rightarrow \infty$ . Then, taking  $\gamma \rightarrow 0^+$  and using (4.3), we have

$$\begin{aligned} & - \mathbb{E} \int_{x,y} (\tilde{u}(s,x) - u(s,y))^+ \varrho_\varepsilon(x-y) \psi(x) \\ & + \mathbb{E} \int_{x,y} (\tilde{u}(\tau,x) - u(\tau,y))^+ \varrho_\varepsilon(x-y) \psi(x) \\ & \leq \mathbb{E} \int_s^\tau \int_x \Delta_x \psi(x) (\tilde{\Phi}(\tilde{u}(t,x)) - \Phi(u(t,x)))^+ dt + M + C \mathbb{E} \int_0^\tau \int_{x,y} \mathbf{1}_{B \cap \bar{D}}(x) \\ & \quad \cdot \left( \varepsilon^2 \sum_{i,j} |\partial_{x_i y_j} \varrho_\varepsilon(x-y)| + \varepsilon \sum_i |\partial_{x_i} \varrho_\varepsilon(x-y)| + \varrho_\varepsilon(x-y) \right) (\tilde{u}(t,x) - u(t,y))^+ dt \end{aligned}$$

holds for almost all  $s \in (0, \tau)$ , where

$$\begin{aligned} M := & C\varepsilon^{\frac{1}{m+1}} \mathbb{E} \|\nabla[\mathbf{a}](u)\|_{L_2(D_T)}^2 + C\mathfrak{C}(\varepsilon, \delta, \lambda, \alpha) \mathbb{E}(1 + \|u\|_{L_{m+1}(D_T)}^{m+1} + \|\tilde{u}\|_{L_{m+1}(D_T)}^{m+1}) \\ & + C\varepsilon^{-2} \mathbb{E} \left( \|\mathbf{1}_{|u|\geq R_\lambda}(1+u)\|_{L_m(D_T)}^m + \|\mathbf{1}_{|\tilde{u}|\geq R_\lambda}(1+\tilde{u})\|_{L_m(D_T)}^m \right). \end{aligned}$$

Then, for  $\tilde{\gamma} \in (0, \tau)$ , by averaging over  $s \in (0, \tilde{\gamma})$ , setting  $\tilde{\gamma} \rightarrow 0^+$  and using Lemma 6.1, we have

$$(3.19) \quad \begin{aligned} & \mathbb{E} \int_{x,y} (\tilde{u}(\tau,x) - u(\tau,y))^+ \varrho_\varepsilon(x-y) \psi(x) \\ & \leq \mathbb{E} \int_{x,y} (\tilde{\xi}(x) - \xi(y))^+ \varrho_\varepsilon(x-y) \psi(x) \\ & \quad + \mathbb{E} \int_0^\tau \int_x \Delta_x \psi(x) (\tilde{\Phi}(\tilde{u}(t,x)) - \Phi(u(t,x)))^+ dt + M + C \mathbb{E} \int_0^\tau \int_{x,y} \mathbf{1}_{B \cap \bar{D}}(x) \\ & \quad \cdot \left( \varepsilon^2 \sum_{i,j} |\partial_{x_i y_j} \varrho_\varepsilon(x-y)| + \varepsilon \sum_i |\partial_{x_i} \varrho_\varepsilon(x-y)| + \varrho_\varepsilon(x-y) \right) (\tilde{u}(t,x) - u(t,y))^+ dt. \end{aligned}$$

To prove (3.16), taking  $\lambda = 0$  and  $R_\lambda = \infty$ , we have

$$\mathbb{E} \left( \|\mathbf{1}_{|u|\geq R_\lambda}(1+u)\|_{L_m(D_T)}^m + \|\mathbf{1}_{|\tilde{u}|\geq R_\lambda}(1+\tilde{u})\|_{L_m(D_T)}^m \right) = 0,$$

and  $\mathfrak{C}(\varepsilon, \delta, \lambda, \alpha)$  changes to

$$\mathfrak{C}(\varepsilon, \delta, \alpha) := \varepsilon^{-2} \delta^{2\beta} + \delta^\beta \varepsilon^{-1} + \varepsilon^{\bar{\kappa}} + \varepsilon^{\bar{\beta}} + \varepsilon^{1/(m+1)} + \varepsilon^{2\bar{\kappa}} \delta^{-1} + \varepsilon^{-2} \delta^{2\alpha}.$$

Since  $\beta \in ((2\bar{\kappa})^{-1}, 1]$ , we can choose  $\vartheta \in ((m \wedge 2)^{-1} \vee (2\beta)^{-1}, \bar{\kappa})$  and  $\alpha \in ((2\vartheta)^{-1}, 1 \wedge (m/2))$ . Let  $\delta = \varepsilon^{2\vartheta}$  and  $\varepsilon \rightarrow 0$ , we have  $\mathfrak{C}(\varepsilon, \delta, \alpha) \rightarrow 0$ . Notice that  $\varepsilon |\partial_{x_i} \varrho_\varepsilon|$  and  $\varepsilon^2 |\partial_{x_i x_j} \varrho_\varepsilon|$  are all approximations of the identity up to a constant. Adding over  $\psi_i$  from partition of unity, with the continuity of translations in  $L_1$ , we have

$$\mathbb{E} \int_x (\tilde{u}(\tau, x) - u(\tau, x))^+ \leq \mathbb{E} \int_x (\tilde{\xi}(x) - \xi(x))^+ + C \int_0^\tau \mathbb{E} \int_x (\tilde{u}(t, x) - u(t, x))^+ dt$$

holds for almost all  $\tau \in [0, T]$ . Hence, (3.16) follows from Gronwall's inequality.

Now, we prove (3.17). Notice that

$$\begin{aligned} \mathbb{E} \int_{x,y} (\xi(x) - \xi(y))^+ \varrho_\varepsilon(x-y) \psi(x) &\leq \int_{\mathbb{R}^d} \varrho_\varepsilon(h) \cdot \mathbb{E} \int_{B \cap D} (\xi(x) - \xi(x-h))^+ dx dh \\ &\leq \sup_{|h| \leq \varepsilon} \mathbb{E} \int_{B \cap D} (\bar{\xi}(x) - \bar{\xi}(x-h))^+ dx. \end{aligned}$$

Fixing  $s_1 \in (0, T]$  and integrating (3.19) over  $\tau \in (0, s_1)$ , we have

$$\begin{aligned} &\mathbb{E} \int_0^{s_1} \int_{x,y} (\tilde{u}(\tau, x) - u(\tau, y))^+ \varrho_\varepsilon(x-y) \psi(x) d\tau \\ &\leq T \mathbb{E} \int_x (\tilde{\xi}(x) - \xi(x))^+ \psi(x) + T \sup_{|h| \leq 2\varepsilon} \mathbb{E} \int_{B \cap D} (\bar{\xi}(x) - \bar{\xi}(x-h))^+ dx \\ &\quad + \mathbb{E} \int_0^{s_1} \int_0^\tau \int_x \Delta_x \psi(x) (\tilde{\Phi}(\tilde{u}(t, x)) - \Phi(u(t, x)))^+ dt d\tau + TM + C \mathbb{E} \int_0^{s_1} \int_0^\tau \int_{x,y} \mathbf{1}_{B \cap \bar{D}}(x) \\ &\quad \cdot \left( \varepsilon^2 \sum_{i,j} |\partial_{x_i y_j} \varrho_\varepsilon(x-y)| + \varepsilon \sum_i |\partial_{x_i} \varrho_\varepsilon(x-y)| + \varrho_\varepsilon(x-y) \right) (\tilde{u}(t, x) - u(t, y))^+ dt d\tau. \end{aligned}$$

Notice that  $\varepsilon |\partial_{x_i} \varrho_\varepsilon|$  and  $\varepsilon^2 |\partial_{x_i x_j} \varrho_\varepsilon|$  are approximations of the identity up to a constant. Taking  $\varepsilon \rightarrow 0$  and adding with different  $\psi_i$ , from the partition of unity, we have

$$\begin{aligned} \mathbb{E} \int_0^{s_1} \int_x (\tilde{u}(\tau, x) - u(\tau, x))^+ d\tau &\lesssim \mathbb{E} \int_x (\tilde{\xi}(x) - \xi(x))^+ + \sup_{|h| \leq 2\varepsilon} \mathbb{E} \int_D (\bar{\xi}(x) - \bar{\xi}(x-h))^+ dx \\ &\quad + M + \mathbb{E} \int_0^{s_1} \int_0^t \int_x (\tilde{u}(\tau, x) - u(\tau, x))^+ d\tau dt. \end{aligned}$$

Using Gronwall's inequality, we acquire (3.17).

For the case that  $u$  has the  $(\star)$ -property, we can relabel  $u \leftrightarrow \tilde{u}$  in (3.16)-(3.17). Using (3.2) and following the proceeding method, this theorem is proved.  $\square$

#### 4. APPROXIMATION

We approximate the function  $\Phi$  to make the approximate equations non-degenerate. The following proposition is taken from [DGG19, DG20] and we refer to [DGG19, Proposition 5.1] for the proof.

**Proposition 4.1.** *Let  $\Phi$  satisfy Assumption 2.1 with a constant  $K > 1$ . Then, for all  $n$  there exists an increasing function  $\Phi_n \in C^\infty(\mathbb{R})$  with bounded derivatives, satisfying Assumption 2.1 with constant  $3K$ , such that  $\mathbf{a}_n(r) \geq 2/n$ , and*

$$(4.1) \quad \sup_{|r| \leq n} |\mathbf{a}(r) - \mathbf{a}_n(r)| \leq 4/n.$$

Define  $\xi_n := \xi \wedge n$ . Denote by  $(\cdot, \cdot)$  the inner product in  $L_2(D)$ .

**Definition 4.2.** An  $L_2$ -solution to  $\Pi(\Phi_n, \xi_n)$  is a continuous  $L_2(D)$ -valued process  $u_n$ , such that  $u_n \in L_2(\Omega_T; H_0^1(D))$ ,  $\nabla \Phi_n(u_n) \in L_2(\Omega_T; L_2(D))$ , and the equality

$$\begin{aligned} (u_n(t, \cdot), \phi) &= (\xi_n, \phi) - \int_0^t \left( (\nabla \Phi_n(u_n), \nabla \phi) + (a^{ij}(\cdot, u_n) \partial_{x_j} u_n + b^i(\cdot, u_n) + f^i(\cdot, u_n), \partial_{x_i} \phi) \right. \\ &\quad \left. + (F(\cdot, u_n), \phi) \right) ds - \int_0^t (\sigma^k(\cdot, u_n), \nabla \phi) dW^k(s) \end{aligned}$$

holds for all  $\phi \in C_c^\infty(D)$ , almost surely for all  $t \in [0, T]$ .

Differing from Definition 2.2, with a strong regularity of the solution  $u_n$ , we can consider the Dirichlet boundary condition in the sense of trace. From (2.4), (2.8) and (2.10) in Assumption 2.4, we have the linear growth of  $\sigma_{x_i}^i$ ,  $f_{x_i}^i$  and  $F$  in  $r$ . Combining with (2.2) in Assumption 2.1, for all  $p \geq 2$ , the  $L_2$ -solution  $u_n$  to  $\Pi(\Phi_n, \xi_n)$  has the following a priori estimates

$$(4.2) \quad \mathbb{E} \sup_{t \leq T} \|u_n\|_{L_2(D)}^p + \mathbb{E} \|\nabla[\mathbf{a}_n](u_n)\|_{L_2(D_T)}^p \leq C(1 + \mathbb{E}\|\xi_n\|_{L_2(D)}^p),$$

$$(4.3) \quad \mathbb{E} \sup_{t \leq T} \|u_n\|_{L_{m+1}(D)}^{m+1} + \mathbb{E} \|\nabla \Phi_n(u_n)\|_{L_2(D_T)}^2 \leq C(1 + \mathbb{E}\|\xi_n\|_{L_{m+1}(D)}^{m+1}).$$

The proof of the above two estimates is almost the same as [DG20, Lemma A.1] with the Dirichlet boundary condition of  $u_n$ , and we omit it here. It is worth noting that the fact  $\mathbf{a}_n \geq 2/n > 0$  and (4.2) indicate

$$(4.4) \quad \mathbb{E} \|\nabla u_n\|_{L_2(D_T)}^p \leq C(n)(1 + \mathbb{E}\|\xi_n\|_{L_2(D)}^p).$$

*Remark 4.3.* Note that for all  $(\eta, \varphi, \varrho) \in \mathcal{E} \times C_c^\infty([0, T]) \times C_c^\infty(D)$  or  $(\eta, \varphi, \varrho) \in \mathcal{E}_0 \times C_c^\infty([0, T]) \times C^\infty(\overline{D})$  and  $\phi := \varphi \times \varrho \geq 0$ , applying Itô's formula (see e.g. [Kry13]) to  $u_n \mapsto \int_D \eta(u_n) \phi dx$ , with (4.2)-(4.3) and Assumption 2.1, we have that the  $L_2$ -solution  $u_n$  is also an entropy solution to  $\Pi(\Phi_n, \xi_n)$ . Using Proposition 2.8, when  $0 \leq \xi \in L_{m+1}(\Omega, \mathcal{F}_0; L_{m+1}(D))$ , we have  $u_n \geq 0$  for almost all  $(\omega, t, x) \in \Omega_T \times D$ .

**Proposition 4.4.** *Let  $0 \leq \xi \in L_{m+1}(\Omega, \mathcal{F}_0; L_{m+1}(D))$  and Assumptions 2.1, 2.4 and 2.5 hold. For each  $n \in \mathbb{N}$ , let  $u_n$  be an  $L_2$ -solution of  $\Pi(\Phi_n, \xi_n)$ . Then,  $u_n$  has the  $(\star)$ -property. If in addition  $\mathbb{E}\|\xi\|_{L_2(D)}^{2(m+1)/m} < \infty$ , the constants  $C$  in Definition 3.4 are independent of  $n$ .*

With the help of Remark 4.3, Proposition 4.4, and Theorem 3.13 (i), following almost the same argument as [DG20, Proposition 5.4], we have the existence and uniqueness of the  $L_2$ -solution  $u_n$ . Here, we omit the proof.

**Proposition 4.5.** *Let  $0 \leq \xi \in L_{m+1}(\Omega, \mathcal{F}_0; L_{m+1}(D))$  and Assumptions 2.1, 2.4 and 2.5 hold. Then, for each  $n \in \mathbb{N}$ ,  $\Pi(\Phi_n, \xi_n)$  admits a unique  $L_2$ -solution  $u_n$ .*

*Proof of Proposition 4.4.* Definition 3.4 (i) is obvious. We now prove (ii). Fix  $\gamma > 0$  small enough. Since  $y \mapsto \varrho_\gamma(x - y) \in C_c^\infty(D)$  for all  $(x, \gamma) \in (B \cap \overline{D}) \times (0, \bar{\varepsilon})$ , for a function  $f \in L_2(D)$ , let  $f^{(\gamma)}(x) := \int_D f(z) \varrho_\gamma(x - z) dz$ . Then, on  $B \cap D$ ,  $u_n^{(\gamma)}$  satisfies (pointwise) the equation

$$\begin{aligned} du_n^{(\gamma)} &= \Delta(\Phi_n(u_n))^{(\gamma)} + \partial_{x_i} (a^{ij}(\cdot, u_n) \partial_{x_j} u_n + b^i(\cdot, u_n) + f^i(\cdot, u_n))^{(\gamma)} \\ &\quad + (F(\cdot, u_n))^{(\gamma)} dt + \partial_{x_i} (\sigma^{ik}(\cdot, u_n))^{(\gamma)} dW^k(t). \end{aligned}$$

Note that

$$\mathbb{E} \int_{s,x} H_\theta(s, x, u_n(s, x)) = \lim_{\lambda \rightarrow 0} \mathbb{E} \int_{s,x,z} H_\theta(s, x, z) \mathbf{1}_{B \cap D}(x) \rho_\lambda(u_n(s, x) - z),$$

and

$$\begin{aligned} & \left| \mathbb{E} \int_{s,x,z} H_\theta(s,x,z) \mathbf{1}_{B \cap D}(x) \rho_\lambda(u_n(s,x) - z) - \mathbb{E} \int_{s,x,z} H_\theta(s,x,z) \mathbf{1}_{B \cap D}(x) \rho_\lambda(u_n^{(\gamma)}(s,x) - z) \right| \\ & \leq C \left( \mathbb{E} \|u_n - u_n^{(\gamma)}\|_{L^1([0,T] \times (B \cap \bar{D}))}^2 \right)^{\frac{1}{2}} \left( \mathbb{E} \|\partial_z H_\theta\|_{L^\infty(D_T \times \mathbb{R})}^2 \right)^{\frac{1}{2}} \xrightarrow{\gamma \rightarrow 0} 0. \end{aligned}$$

With Remark 3.3, we have  $\mathbb{E} H_\theta(s,x,z) X = 0$  for any  $\mathcal{F}_{s-\theta}$ -measurable bounded random variable  $X$ . Then,

$$\begin{aligned} & \mathbb{E} \int_{s,x,z} H_\theta(s,x,z) \mathbf{1}_{B \cap D}(x) \rho_\lambda(u_n^{(\gamma)}(s,x) - z) \\ & = \mathbb{E} \int_{s,x,z} H_\theta(s,x,z) \mathbf{1}_{B \cap D}(x) \left( \rho_\lambda(u_n^{(\gamma)}(s,x) - z) - \rho_\lambda(u_n^{(\gamma)}(s-\theta,x) - z) \right). \end{aligned}$$

Using Itô's formula, we have

$$\int_{s,x,z} H_\theta(s,x,z) \left( \rho_\lambda(u_n^{(\gamma)}(s,x) - z) - \rho_\lambda(u_n^{(\gamma)}(s-\theta,x) - z) \right) = \sum_{i=1}^5 N_{\lambda,\gamma}^{(i)},$$

where

$$\begin{aligned} N_{\lambda,\gamma}^{(1)} & := \int_{s,x,z} H_\theta(s,x,z) \int_{s-\theta}^s \rho'_\lambda(u_n^{(\gamma)}(t,x) - z) \Delta_x(\Phi_n(u_n))^{(\gamma)} dt, \\ N_{\lambda,\gamma}^{(2)} & := \int_{s,x,z} H_\theta(s,x,z) \int_{s-\theta}^s \rho'_\lambda(u_n^{(\gamma)}(t,x) - z) \partial_{x_i} (a^{ij}(\cdot, u_n) \partial_{x_j} u_n + b^i(\cdot, u_n))^{(\gamma)} dt, \\ N_{\lambda,\gamma}^{(3)} & := \int_{s,x,z} H_\theta(s,x,z) \int_{s-\theta}^s \rho'_\lambda(u_n^{(\gamma)}(t,x) - z) (\partial_{x_i} f^i(\cdot, u_n) + F(\cdot, u_n))^{(\gamma)} dt, \\ N_{\lambda,\gamma}^{(4)} & := \int_{s,x,z} H_\theta(s,x,z) \int_{s-\theta}^s \rho'_\lambda(u_n^{(\gamma)}(t,x) - z) \partial_{x_i} (\sigma^{ik}(\cdot, u_n))^{(\gamma)} dW^k(t), \\ N_{\lambda,\gamma}^{(5)} & := \frac{1}{2} \int_{s,x,z} H_\theta(s,x,z) \int_{s-\theta}^s \rho''_\lambda(u_n^{(\gamma)}(t,x) - z) \sum_{k=1}^{\infty} |\partial_{x_i} (\sigma^{ik}(\cdot, u_n))^{(\gamma)}|^2 dt. \end{aligned}$$

Using the divergence theorem in  $x$ , we have  $N_{\lambda,\gamma}^{(1)} = \sum_{i=1}^3 N_{\lambda,\gamma}^{(1,i)}$ , where

$$\begin{aligned} N_{\lambda,\gamma}^{(1,1)} & := - \int_{s,x,z} \mathbf{1}_{\{s>\theta\}} \nabla_x H_\theta(s,x,z) \int_{s-\theta}^s \rho'_\lambda(u_n^{(\gamma)}(t,x) - z) \nabla_x (\Phi_n(u_n))^{(\gamma)} dt, \\ N_{\lambda,\gamma}^{(1,2)} & := - \int_{s,x,z} \mathbf{1}_{\{s>\theta\}} H_\theta(s,x,z) \int_{s-\theta}^s \rho''_\lambda(u_n^{(\gamma)}(t,x) - z) \nabla_x u_n^{(\gamma)}(t,x) \nabla_x (\Phi_n(u_n))^{(\gamma)} dt, \\ N_{\lambda,\gamma}^{(1,3)} & := \int_{s,z} \int_{\partial D} \mathbf{1}_{\{s>\theta\}} H_\theta(s,x,z) \int_{s-\theta}^s \rho'_\lambda(u_n^{(\gamma)}(t,x) - z) \nabla_x (\Phi_n(u_n))^{(\gamma)} \cdot \nu dt dS, \end{aligned}$$

and  $\nu(x)$  is unit normal vector of  $\partial D$  at  $x$ . Using integration by parts in  $z$  and (4.3), we have

$$\begin{aligned} \mathbb{E} |N_{\lambda,\gamma}^{(1,1)}| & = \mathbb{E} \left| \int_{s,x,z} \mathbf{1}_{\{s>\theta\}} \nabla_x \partial_z H_\theta(s,x,z) \int_{s-\theta}^s \rho_\lambda(u_n^{(\gamma)}(t,x) - z) \nabla_x (\Phi_n(u_n))^{(\gamma)} dt \right| \\ & \leq C\theta (\mathbb{E} \|\nabla_x \partial_z H_\theta\|_{L^\infty(D_T \times \mathbb{R})}^2)^{\frac{1}{2}} (\mathbb{E} \|\nabla_x (\Phi_n(u_n))^{(\gamma)}\|_{L^1(D_T)}^2)^{\frac{1}{2}} \leq C\theta^{1-\mu}. \end{aligned}$$

Similarly, integrating by parts twice in  $z$  on  $N_{\lambda,\gamma}^{(1,2)}$ , we have

$$\lim_{\gamma \rightarrow 0} \mathbb{E} |N_{\lambda,\gamma}^{(1,2)}| \leq C\theta^{1-\mu} \lim_{\gamma \rightarrow 0} (\mathbb{E} \|\nabla_x u_n^{(\gamma)}(t,x) \nabla_x (\Phi_n(u_n))^{(\gamma)}\|_{L^1(D_T)}^{\frac{m+1}{m}})^{\frac{m}{m+1}}$$

$$\begin{aligned}
&\leq C\theta^{1-\mu}(\mathbb{E}\|\nabla_x u_n \nabla_x(\Phi_n(u_n))\|_{L^1_1(D_T)}^{\frac{m+1}{m}})^{\frac{m}{m+1}}. \\
&= C\theta^{1-\mu}(\mathbb{E}\|\nabla[\mathbf{a}_n](u_n)\|_{L^2(D_T)}^{\frac{2(m+1)}{m}})^{\frac{m}{m+1}} \leq C(n)\theta^{1-\mu}.
\end{aligned}$$

From the Dirichlet boundary condition of  $u_n$ , we have

$$\begin{aligned}
\lim_{\gamma \rightarrow 0} N_{\lambda, \gamma}^{(1,3)} &= \int_{s,z} \int_{\partial D} \mathbf{1}_{\{s>\theta\}} H_\theta(s, x, z) \int_{s-\theta}^s \rho'_\lambda(0-z) \nabla_x(\Phi_n(u_n))^{(\gamma)} \cdot \nu dt dS \\
&= \int_{s,z} \mathbf{1}_{\{z<0\}} \int_{\partial D} \mathbf{1}_{\{s>\theta\}} H_\theta(s, x, z) \int_{s-\theta}^s \rho'_\lambda(0-z) \nabla_x(\Phi_n(u_n))^{(\gamma)} \cdot \nu dt dS.
\end{aligned}$$

Since  $\text{supp} \rho'_\lambda \subset [0, \infty)$ , the integrand only acts when  $z \in (-\infty, 0]$ . However, based on the non-negativity of  $u$  and the definition of  $\mathcal{C}^-$ , we have for all  $(h, z) \in \mathcal{C}^- \times (-\infty, 0]$ ,

$$h(u(t, y) - z) = 0, \quad \text{a.s. } (t, \omega, y) \in \Omega_T \times D.$$

which indicates  $\lim_{\gamma \rightarrow 0} N_{\lambda, \gamma}^{(1,3)} = 0$ . Therefore,

$$\limsup_{\gamma \rightarrow 0} \mathbb{E}|N_{\lambda, \gamma}^{(1)}| \leq C(n)\theta^{1-\mu}.$$

Now we estimate  $N_{\lambda, \gamma}^{(2)} + N_{\lambda, \gamma}^{(5)}$ . As the estimate of  $N_{\lambda, \gamma}^{(1,3)}$ , using the divergence theorem in  $x$  and combining with the definition of  $a^{ij}$  and  $b^i$ , we have

$$\begin{aligned}
&\limsup_{\gamma \rightarrow 0} \mathbb{E}|N_{\lambda, \gamma}^{(2)} + N_{\lambda, \gamma}^{(5)}| \\
&\leq \mathbb{E} \left| \int_{s,x,z} \partial_{x_i} H_\theta(s, x, z) \int_{s-\theta}^s \rho'_\lambda(u_n(t, x) - z) (a^{ij}(\cdot, u_n) \partial_{x_j} u_n) dt \right| \\
&\quad + \mathbb{E} \left| \int_{s,x,z} \partial_{x_i} H_\theta(s, x, z) \int_{s-\theta}^s \rho'_\lambda(u_n(t, x) - z) (b^i(\cdot, u_n)) dt \right| \\
&\quad + \mathbb{E} \left| \frac{1}{2} \int_{s,x,z} H_\theta(s, x, z) \int_{s-\theta}^s \rho''_\lambda(u_n(t, x) - z) \sum_{k=1}^{\infty} |(\sigma_{x_i}^{ik}(\cdot, u_n))|^2 dt \right|.
\end{aligned}$$

Using the identity

$$\begin{aligned}
\rho'_\lambda(u_n(t, x) - z) a^{ij}(x, u_n) \partial_{x_j} u_n &= \partial_{x_j} \llbracket a^{ij} \rho'_\lambda(\cdot - z) \rrbracket(x, u_n(t, x)) \\
&\quad - \llbracket a^{ij} \rho'_\lambda(\cdot - z) \rrbracket(x, u_n(t, x)),
\end{aligned}$$

and

$$\begin{aligned}
&\mathbb{E} \left| \int_{s,x,z} \partial_{x_i} H_\theta(s, x, z) \int_{s-\theta}^s \partial_{x_j} \llbracket a^{ij} \rho'_\lambda(\cdot - z) \rrbracket(x, u_n(t, x)) dt \right| \\
&\leq C\theta (\mathbb{E} \|\partial_z \partial_{x_i} \partial_{x_j} H_\theta\|_{L^\infty(D_T \times \mathbb{R})}^2)^{\frac{1}{2}} (\mathbb{E} \|\llbracket a^{ij} \rho_\lambda(\cdot - z) \rrbracket(\cdot, u_n)\|_{L^1(D_T)}^2)^{\frac{1}{2}},
\end{aligned}$$

with the linear growth of  $\sigma_{x_i}^i$ ,  $b^i$  and and the boundness of  $a^{ij}_{x_j}$  and  $a^{ij}$  derived from (2.4)-(2.5) and (2.7) in Assumption 2.4, we have

$$\limsup_{\gamma \rightarrow 0} \mathbb{E}|N_{\lambda, \gamma}^{(2)} + N_{\lambda, \gamma}^{(5)}| \leq C\theta^{1-\mu} (1 + \mathbb{E}\|u_n\|_{L^2(D_T)}^{\frac{2(m+1)}{m}})^{\frac{m}{m+1}} \leq C(n)\theta^{1-\mu}.$$

For  $N_{\lambda, \gamma}^{(3)}$ , with the linear growth of  $f_{x_i}^i$ ,  $F$  and the boundness of  $f_r^i$  and  $f_{rx_i}^i$  derived from (2.7)-(2.10) in Assumption 2.4, we similarly obtain

$$\limsup_{\gamma \rightarrow 0} \mathbb{E}|N_{\lambda, \gamma}^{(3)}| \leq C\theta^{1-\mu} (1 + \mathbb{E}\|u_n\|_{L^2(D_T)}^2)^2 \leq C(n)\theta^{1-\mu}.$$



Using Itô's isometry and taking  $\gamma \rightarrow 0$ , since

$$(4.5) \quad \begin{aligned} & \int_{s,x,z,y} \int_{s-\theta}^s \int_0^z \sigma_{ry_i}^{ik}(y,r)h(r-z)dr\phi_\theta\rho'_\lambda(u_n(t,x)-z)\partial_{x_j}\sigma^{jk}(x,u_n(t,x))dt \\ &= - \int_{s,x,z,y} \int_{s-\theta}^s \int_0^z \sigma_r^{ik}(y,r)h(r-z)dr\partial_{y_i}\phi_\theta\rho'_\lambda(u_n(t,x)-z)\partial_{x_j}\sigma^{jk}(x,u_n(t,x))dt, \end{aligned}$$

we have

$$\lim_{\gamma \rightarrow 0} \mathbb{E}N_{\lambda,\gamma}^{(4)} = \sum_{i=1}^6 D_i,$$

where

$$\begin{aligned} D_1 &:= - \int_{s,x,z,y} \int_{s-\theta}^s \int_z^{u(t,y)} \sigma_{ry_i}^{ik}(y,r)h(r-z)dr\phi_\theta\rho'_\lambda(u_n(t,x)-z)\sigma_{x_j}^{jk}(x,u_n(t,x))dt, \\ D_2 &:= - \int_{s,x,z,y} \int_{s-\theta}^s \int_z^{u(t,y)} \sigma_{ry_i}^{ik}(y,r)h(r-z)dr\phi_\theta\rho'_\lambda(u_n(t,x)-z)\sigma_r^{jk}(x,u_n(t,x))\partial_{x_j}u_n(t,x)dt, \\ D_3 &:= - \int_{s,x,z,y} \int_{s-\theta}^s \int_z^{u(t,y)} \sigma_r^{ik}(y,r)h(r-z)dr\partial_{y_i}\phi_\theta\rho'_\lambda(u_n(t,x)-z)\sigma_{x_j}^{jk}(x,u_n(t,x))dt, \\ D_4 &:= - \int_{s,x,z,y} \int_{s-\theta}^s \int_z^{u(t,y)} \sigma_r^{ik}(y,r)h(r-z)dr\partial_{y_i}\phi_\theta\rho'_\lambda(u_n(t,x)-z)\sigma_r^{jk}(x,u_n(t,x))\partial_{x_j}u_n(t,x)dt, \\ D_5 &:= \int_{s,x,z,y} \int_{s-\theta}^s h(u(t,y)-z)\phi_\theta\sigma_{y_i}^{ik}(y,u(t,y))\rho'_\lambda(u_n(t,x)-z)\sigma_{x_j}^{jk}(x,u_n(t,x))dt, \\ D_6 &:= \int_{s,x,z,y} \int_{s-\theta}^s h(u(t,y)-z)\phi_\theta\sigma_{y_i}^{ik}(y,u(t,y))\rho'_\lambda(u_n(t,x)-z)\sigma_r^{jk}(x,u_n(t,x))\partial_{x_j}u_n(t,x)dt. \end{aligned}$$

For  $D_2 + D_4$ , notice that

$$\begin{aligned} \rho'_\lambda(u_n(t,x)-z)\sigma_r^{jk}(x,u_n(t,x))\partial_{x_j}u_n(t,x) &= \partial_{x_j} \int_0^{u_n(t,x)} \rho'_\lambda(\tilde{r}-z)\sigma_r^{jk}(x,\tilde{r})d\tilde{r} \\ &\quad - \int_0^{u_n(t,x)} \rho'_\lambda(\tilde{r}-z)\sigma_{rx_j}^{jk}(x,\tilde{r})d\tilde{r}. \end{aligned}$$

We can apply the divergence theorem in  $x$  and the Dirichlet boundary condition and integrate by parts in  $z$ . Moreover, since  $h \in \mathcal{C}^-$ , the integrand is non-zero only when  $u(t,y) < z$ . Using  $\text{supp}\rho_\lambda \in \mathbb{R}_+$  and the non-negativity of  $u$ , we have

$$(4.6) \quad \begin{aligned} & D_2 + D_4 \\ &= - \int_{s,x,z,y} \int_{s-\theta}^s \int_z^{u(t,y)} \sigma_{ry_i}^{ik}(y,r)h'(r-z)dr\partial_{x_j}\phi_\theta \int_{u(t,y)}^{u_n(t,x)} \rho_\lambda(\tilde{r}-z)\sigma_r^{jk}(x,\tilde{r})d\tilde{r}dt \\ &\quad - \int_{s,x,z,y} \int_{s-\theta}^s \int_z^{u(t,y)} \sigma_{ry_i}^{ik}(y,r)h'(r-z)dr\phi_\theta \int_{u(t,y)}^{u_n(t,x)} \rho_\lambda(\tilde{r}-z)\sigma_{rx_j}^{jk}(x,\tilde{r})d\tilde{r}dt \\ &\quad - \int_{s,x,z,y} \int_{s-\theta}^s \int_z^{u(t,y)} \sigma_r^{ik}(y,r)h'(r-z)dr\partial_{y_i x_j}\phi_\theta \int_{u(t,y)}^{u_n(t,x)} \rho_\lambda(\tilde{r}-z)\sigma_r^{jk}(x,\tilde{r})d\tilde{r}dt \\ &\quad - \int_{s,x,z,y} \int_{s-\theta}^s \int_z^{u(t,y)} \sigma_r^{ik}(y,r)h'(r-z)dr\partial_{y_i}\phi_\theta \int_{u(t,y)}^{u_n(t,x)} \rho_\lambda(\tilde{r}-z)\sigma_{rx_j}^{jk}(x,\tilde{r})d\tilde{r}dt. \end{aligned}$$

Similarly, we have

$$\begin{aligned}
D_6 &= - \int_{s,x,z,y} \int_{s-\theta}^s h'(u(t,y) - z) \phi_\theta \sigma_{y_i}^{ik}(y, u(t,y)) \partial_{x_j} \int_0^{u_n(t,x)} \rho_\lambda(\tilde{r} - z) \sigma_r^{jk}(x, \tilde{r}) d\tilde{r} dt \\
&\quad + \int_{s,x,z,y} \int_{s-\theta}^s h'(u(t,y) - z) \phi_\theta \sigma_{y_i}^{ik}(y, u(t,y)) \int_0^{u_n(t,x)} \rho_\lambda(\tilde{r} - z) \sigma_{rx_j}^{jk}(x, \tilde{r}) d\tilde{r} dt \\
&= \int_{s,x,z,y} \int_{s-\theta}^s h'(u(t,y) - z) \partial_{x_j} \phi_\theta \sigma_{y_i}^{ik}(y, u(t,y)) \int_{u(t,y)}^{u_n(t,x)} \rho_\lambda(\tilde{r} - z) \sigma_r^{jk}(x, \tilde{r}) d\tilde{r} dt \\
&\quad + \int_{s,x,z,y} \int_{s-\theta}^s h'(u(t,y) - z) \phi_\theta \sigma_{y_i}^{ik}(y, u(t,y)) \int_{u(t,y)}^{u_n(t,x)} \rho_\lambda(\tilde{r} - z) \sigma_{rx_j}^{jk}(x, \tilde{r}) d\tilde{r} dt.
\end{aligned}$$

For  $D_1$ ,  $D_3$  and  $D_5$ , we integrate by parts in  $z$ . Therefore, we have

$$\lim_{\lambda \rightarrow 0} \lim_{\gamma \rightarrow 0} \mathbb{E} N_{\lambda, \gamma}^{(4)} = \mathcal{E}(u, u_n, \theta).$$

Hence,

$$\mathbb{E} \int_{s,x} H_\theta(s, x, u_n(s, x)) \leq C(n) \theta^{1-\mu} + \mathcal{E}(u, u_n, \theta).$$

Furthermore, inspired by (4.2), if  $\mathbb{E} \|\xi\|_{L_2(D)}^{\frac{2(m+1)}{m}} < \infty$ , we can choose  $C$  independent of  $n$ .

For (iii) in the definition of the  $(\star)$ -property, note that  $g(x, \cdot) \in C_c(D)$  for all  $x \in B \cap \overline{D}$ . It is easy to prove following the proceeding proof. The reason is that the boundary terms vanish using the divergence theorem in  $y$ , and other differences in the proof are (4.5),  $D_2 + D_4$  and  $D_6$ . For (4.5), with  $h \in C^+$ , we have for all  $(x, z) \in D \times [0, \infty)$ ,

$$\int_0^z \sigma_{rx_i}^{ik}(x, r) h(r - z) dr = \int_0^z \sigma_r^{ik}(x, r) h(r - z) dr = 0.$$

When  $z < 0$ , using (2.4) in Assumption 2.4, (4.2), Remark 4.3 and the definition of  $C^+$ , the boundary term arose in the divergence theorem in  $x$  will disappear when  $\lambda \rightarrow 0$ . Therefore, we have (4.5) in a version exchanging  $x$  and  $y$  in the integrand except the test function  $\phi_\theta$ .

As for  $D_2 + D_4$  and  $D_6$ , we can apply

$$\partial_{y_j} \int_0^{u(t,x)} \rho_\lambda(\tilde{r} - z) \sigma_r^{jk}(y, \tilde{r}) d\tilde{r} dt - \int_0^{u(t,x)} \rho_\lambda(\tilde{r} - z) \sigma_{ry_j}^{jk}(y, \tilde{r}) d\tilde{r} dt = 0$$

to acquire (4.6) instead of using the support of  $h$ . Therefore, this proposition is proved.  $\square$

## 5. EXISTENCE AND UNIQUENESS

*Proof of Theorem 2.6.* In this part, we will use the  $L_1$ -estimates to prove that  $\{u_n\}_{n \in \mathbb{N}}$  constructed in Section 4 is a Cauchy sequence.

For  $n, n' \geq 1$ , let  $u_n$  and  $u_{n'}$  be the  $L_2$ -solution of  $\Pi(\Phi_n, \xi_n)$  and  $\Pi(\Phi_{n'}, \xi_{n'})$ , respectively. Remark 4.3 shows that  $u_n$  and  $u_{n'}$  are also entropy solution. Proposition 4.4 indicates that  $\{u_n\}_{n \in \mathbb{N}}$  has the  $(\star)$ -property uniformly in  $n$ . Without loss of generality, we can assume that  $n \leq n'$ . Since  $\beta \in ((2\bar{\kappa})^{-1}, 1]$ , we can choose  $\vartheta \in ((m \wedge 2)^{-1} \vee (2\beta)^{-1}, \bar{\kappa})$  and  $\alpha \in ((2\vartheta)^{-1}, 1 \wedge (m/2))$ . Let  $\delta = \varepsilon^{2\vartheta}$  and  $\lambda = 8/n$ . Using (4.1), we have  $R_\lambda \geq n$ . Applying Theorem 3.13 with  $u_n$  and  $u_{n'}$  and using (4.3) and the triangle inequality

$$\begin{aligned}
&\mathbb{E} \|\xi_n(\cdot) - \bar{\xi}_n(\cdot + h)\|_{L_1(D)} \\
&\leq \mathbb{E} \|\xi(\cdot) - \bar{\xi}(\cdot + h)\|_{L_1(D)} + 2\mathbb{E} \|\xi - \xi_n\|_{L_1(D)}, \quad \forall n \in \mathbb{N},
\end{aligned}$$

we have

$$\begin{aligned} & \mathbb{E} \int_0^T \int_D |u_{n'}(\tau, x) - u_n(\tau, x)| dx d\tau \\ & \leq M(\varepsilon) + C\mathbb{E}\|\xi - \xi_{n'}\|_{L_1(D)} + C\mathbb{E}\|\xi - \xi_n\|_{L_1(D)} + C\varepsilon^{-2}n^{-2} + C\varepsilon^{-1}n^{-1} \\ & \quad + C\varepsilon^{-2}\mathbb{E}\left(\|\mathbf{1}_{|u_n| \geq n}(1 + u_n)\|_{L_m(D_T)}^m + \|\mathbf{1}_{|u_{n'}| \geq n}(1 + u_{n'})\|_{L_m(D_T)}^m\right), \end{aligned}$$

where  $M(\varepsilon) \rightarrow 0$  as  $\varepsilon \rightarrow 0$ . For any  $\varepsilon_0 > 0$ , we select  $\varepsilon \in (0, \bar{\varepsilon})$  small enough such that  $M(\varepsilon) \leq \varepsilon_0$ . Then, using (4.3), we can choose  $n_0$  sufficiently large so that for  $n_0 \leq n \leq n'$ , we have

$$\begin{aligned} & C\mathbb{E}\|\xi_{n'} - \xi\|_{L_1(D)} + C\mathbb{E}\|\xi - \xi_n\|_{L_1(D)} + C\varepsilon^{-2}n^{-2} + C\varepsilon^{-1}n^{-1} \\ & \quad + C\varepsilon^{-2}\mathbb{E}\left(\|\mathbf{1}_{|u_n| \geq n}(1 + u_n)\|_{L_m(D_T)}^m + \|\mathbf{1}_{|u_{n'}| \geq n}(1 + u_{n'})\|_{L_m(D_T)}^m\right) \leq \varepsilon_0. \end{aligned}$$

Therefore, we have

$$\lim_{n, n' \rightarrow \infty} \mathbb{E} \int_0^T \int_D |u_{n'}(t, x) - u_n(t, x)| dx dt = 0.$$

Moreover, by taking a subsequence, we may assume

$$(5.1) \quad \lim_{n \rightarrow \infty} u_n = u, \quad \text{a.s. } (\omega, t, x) \in \Omega_T \times D.$$

In addition, the sequence  $\{|u_n(t, x)|^q\}_{n \in \mathbb{N}}$  is uniformly integrable on  $\Omega_T \times D$  for all  $q \in (0, m+1)$ . Now, we verify that  $u$  is an entropy solution to  $\Pi(\Phi, \xi)$  under Definition 2.2.

Firstly, with the definition of  $\xi_n$ , we have that  $\{\xi_n\}_{n \in \mathbb{N}}$  is weak convergence in the Banach space  $L_{m+1}(\Omega_T; L_{m+1}(D))$ . Applying the Banach-Saks Theorem, taking a subsequence and using (4.3), we have

$$\begin{aligned} \mathbb{E}\|u\|_{L_{m+1}(\Omega_T; L_{m+1}(D))} & \leq \liminf_{n \rightarrow \infty} \|u_n\|_{L_{m+1}(\Omega_T; L_{m+1}(D))} \\ & \leq C(1 + \liminf_{n \rightarrow \infty} \|\xi_n\|_{L_{m+1}(\Omega; L_{m+1}(D))}) \\ & \leq C(1 + \|\xi\|_{L_{m+1}(\Omega; L_{m+1}(D))}). \end{aligned}$$

To prove Definition 2.2 (ii) of  $u$ , let  $f \in C_b(\mathbb{R})$ . From Assumption 2.1 and (4.3), we have

$$\sup_n \mathbb{E} \int_0^T \int_D |[\mathbf{a}_n f](u_n)|^2 dx dt \lesssim \sup_n \mathbb{E} \int_0^T \int_D (|u_n| + |u_n|^{\frac{m+1}{2}})^2 dx dt < \infty.$$

Combining the estimate (4.2) with the fact that  $[\mathbf{a}_n f](u_n) \in L_2(\Omega_T; H_0^1(D))$  and  $\partial_{x_i} [\mathbf{a}_n f](u_n) = f(u_n) \partial_{x_i} [\mathbf{a}_n](u_n)$ , we have

$$\sup_n \mathbb{E} \int_t \|\partial_{x_i} [\mathbf{a}_n f](u_n)\|_{H^1(D)}^2 < \infty.$$

With the pointwise convergence and uniform integrability of  $u_n$  and Proposition 4.1, by taking a subsequence, we obtain the weak convergence of  $[\mathbf{a}_n f](u_n)$  and  $[\mathbf{a}_n](u_n)$  in  $L_2(\Omega_T; H_0^1(D))$  when  $n \rightarrow \infty$ , and the limits are  $[\mathbf{a} f](u)$  and  $[\mathbf{a}](u)$ , respectively. On the other hand, for all  $\phi \in C_c^\infty([0, T] \times D)$  and  $A \in \mathcal{F}$ , based on the strong convergence of  $f(u_n)\phi$  and weak convergence of  $[\mathbf{a}_n](u_n)$ , we have

$$\begin{aligned} \mathbb{E} \left[ \mathbf{1}_A \int_{t,x} \partial_{x_i} [\mathbf{a} f](u) \phi \right] & = \lim_{n \rightarrow \infty} \mathbb{E} \left[ \mathbf{1}_A \int_{t,x} \partial_{x_i} [\mathbf{a}_n f](u_n) \phi \right] \\ & = \lim_{n \rightarrow \infty} \mathbb{E} \left[ \mathbf{1}_A \int_{t,x} f(u_n) \partial_{x_i} [\mathbf{a}_n](u_n) \phi \right] \\ & = \mathbb{E} \left[ \mathbf{1}_A \int_{t,x} f(u) \partial_{x_i} [\mathbf{a}](u) \phi \right]. \end{aligned}$$

For Definition 2.2 (iii), let  $A \in \mathcal{F}$ . Combining Remark 4.3 with Itô's product rule, we have

$$\begin{aligned}
(5.2) \quad & - \mathbb{E} \left[ \mathbf{1}_B \int_0^T \int_D \eta(u_n) \partial_t \phi dx dt \right] \\
& = \mathbb{E} \mathbf{1}_B \left[ \int_D \eta(\xi_n) \phi(0) dx + \int_0^T \int_D (\llbracket \mathbf{a}_n^2 \eta' \rrbracket(u_n) \Delta \phi + \llbracket a^{ij} \eta' \rrbracket(x, u_n) \partial_{x_i x_j} \phi) dx dt \right. \\
& \quad + \int_0^T \int_D (\llbracket a_{x_j}^{ij} \eta' - f_r^i \eta' \rrbracket(x, u_n) - \eta'(u_n) b^i(x, u_n)) \partial_{x_i} \phi dx dt \\
& \quad + \int_0^T \int_D (- \llbracket f_{rx_i}^i \eta' \rrbracket(x, u_n) + \eta'(u_n) f_{x_i}^i(x, u_n) + \eta'(u_n) F(x, u_n)) \phi dx dt \\
& \quad + \int_0^T \int_D \left( \frac{1}{2} \eta''(u_n) \sum_{k=1}^{\infty} |\sigma_{x_i}^{ik}(x, u_n)|^2 \phi - \eta''(u_n) |\nabla \llbracket \mathbf{a}_n \rrbracket(u_n)|^2 \phi \right) dx dt \\
& \quad \left. + \int_0^T \int_D \left( \eta'(u_n) \phi \sigma_{x_i}^{ik}(x, u_n) - \llbracket \sigma_{rx_i}^{ik} \eta' \rrbracket(x, u_n) \phi - \llbracket \sigma_r^{ik} \eta' \rrbracket(x, u_n) \partial_{x_i} \phi \right) dx dW^k(t) \right].
\end{aligned}$$

Since  $(\eta'')^{1/2} \in C_b(\mathbb{R})$ . As the proof in checking (ii), we have

$$\partial_{x_i} \llbracket (\eta'')^{1/2} \mathbf{a}_n \rrbracket(u_n) = (\eta''(u_n))^{1/2} \partial_{x_i} \llbracket \mathbf{a}_n \rrbracket(u_n),$$

and can assume that  $\partial_{x_i} \llbracket (\eta'')^{1/2} \mathbf{a}_n \rrbracket(u_n)$  converges weakly to  $\partial_{x_i} \llbracket (\eta'')^{1/2} \mathbf{a} \rrbracket(u)$  in  $L_2(\Omega_T; L_2(D))$ . Then, we also have the weakly convergence in  $L_2(\Omega_T \times D, \bar{\mu})$ , where  $d\bar{\mu} := \mathbf{1}_B \phi d\mathbb{P} \otimes dt \otimes dx$ , which indicates

$$\mathbb{E} \mathbf{1}_B \int_{t,x} \phi \eta''(u) |\nabla \llbracket \mathbf{a} \rrbracket(u)|^2 \leq \liminf_{n \rightarrow \infty} \mathbb{E} \mathbf{1}_B \int_{t,x} \phi \eta''(u_n) |\nabla \llbracket \mathbf{a}_n \rrbracket(u_n)|^2.$$

Therefore, taking inferior limit on (5.2) and using Proposition 4.1 and Assumption 2.4, with the almost surely convergence and uniformly integrability of  $u_n$ , we have that  $u$  satisfies (2.3) almost surely. Therefore,  $u$  is actually an entropy solution.

Now, we focus on the uniqueness. For  $\bar{n} \in \mathbb{N}$ , define  $\xi_{\bar{n}} := \xi \wedge \bar{n}$  and denote by  $u_{\bar{n}}$  the entropy solution of  $\Pi(\Phi, \xi_{\bar{n}})$  constructed in the proof of existence. From the construction of  $u_{\bar{n}}$ , with Lemma 3.8 and Proposition 4.4, the entropy solution  $u_{\bar{n}}$  has the  $(\star)$ -property. Let  $\tilde{u}$  be an entropy solution of  $\Pi(\Phi, \tilde{\xi})$ . Using Theorem 3.13, we have

$$\operatorname{ess\,sup}_{t \in [0, T]} \mathbb{E} \|\tilde{u}(t, \cdot) - u_{\bar{n}}(t, \cdot)\|_{L_1(D)} \leq C \mathbb{E} \|\tilde{\xi} - \xi_{\bar{n}}\|_{L_1(D)},$$

where the constant  $C$  depends only on  $N_0, K, d, T$  and  $|D|$ . Taking the limit  $\bar{n} \rightarrow \infty$ , we obtain Theorem 2.6 (i). Following this method and using Theorem 3.13 (i), we complete the proof of Theorem 2.6 (iii).  $\square$

*Remark 5.1.* To obtain the uniqueness without the  $(\star)$ -property, we should prove the  $L_1$ -estimates with only one entropy solution satisfying the  $(\star)$ -property.

## 6. SOME AUXILIARY ESTIMATES

**Lemma 6.1.** *Suppose  $u$  is an entropy solution to the Dirichlet problem  $\Pi(\Phi, \xi)$ . Under Assumptions 2.1, 2.4 and 2.5, if  $\xi \in L_{m+1}(\Omega, \mathcal{F}_0; L_{m+1}(D))$ , we have*

$$(6.1) \quad \lim_{\gamma \rightarrow 0} \frac{1}{\gamma} \mathbb{E} \int_0^\gamma \int_x |u(t, x) - \xi(x)|^2 dt = 0.$$

*Proof.* From the partition of unity in Section 3.2, we can fix  $i \in \{0, 1, \dots, K\}$  and define  $B := B_i$ ,  $\psi := \psi_i$  and  $\varrho_\varepsilon := \varrho_{\varepsilon,i}$ . Notice that  $\text{dist}(\text{supp}\psi, \partial B) > 0$ . When  $\varepsilon$  is small enough, we have  $\text{supp}\varrho_\varepsilon(\cdot - x) \subset B$  for all  $x \in \text{supp}\psi$ . Then, from the definition of  $\varrho_\varepsilon$ , we have  $\psi(\cdot)\varrho_\varepsilon(y - \cdot) \in C_c^\infty(D)$  for all  $y \in \overline{D}$  and sufficient small  $\varepsilon > 0$ . Now, we only need to prove

$$\lim_{\gamma \rightarrow 0} \frac{1}{\gamma} \mathbb{E} \int_0^\gamma \int_x |u(t, x) - \xi(x)|^2 \psi(x) dt = 0.$$

We split the limitation into three parts

$$(6.2) \quad \begin{aligned} & \frac{1}{\gamma} \mathbb{E} \int_0^\gamma \int_x |u(t, x) - \xi(x)|^2 \psi(x) dt \\ & \leq 2\mathbb{E} \int_D \int_{\mathbb{R}^d} |\bar{\xi}(y) - \xi(x)|^2 \psi(x) \varrho_\varepsilon(y - x) dy dx \\ & \quad + \frac{2}{\gamma} \mathbb{E} \int_0^\gamma \int_D \int_D |u(t, x) - \bar{\xi}(y)|^2 \psi(x) \varrho_\varepsilon(y - x) dy dx dt \\ & \quad + \frac{2}{\gamma} \mathbb{E} \int_0^\gamma \int_D \int_{D_{y,\varepsilon}^1} |u(t, x)|^2 \psi(x) \varrho_\varepsilon(y - x) dy dx dt, \end{aligned}$$

where  $\gamma \in [0, T)$  and  $D_{y,\varepsilon}^1 := \{y \in \mathbb{R}^d : \varrho_\varepsilon(y - x) > 0, \exists x \in D\} \setminus D$ . See (3.18) for the definition of  $\bar{\xi}$ . Notice that  $|D_{y,\varepsilon}^1| = O(\varepsilon)$ . We first estimate the third term in the right hand side as

$$(6.3) \quad \begin{aligned} & \frac{2}{\gamma} \mathbb{E} \int_0^\gamma \int_D \int_{D_{y,\varepsilon}^1} |u(t, x)|^2 \psi(x) \varrho_\varepsilon(y - x) dy dx dt \\ & = \frac{2}{\gamma} \mathbb{E} \int_0^\gamma \int_{D_{x,\varepsilon}^2} \int_{D_{y,\varepsilon}^1} |u(t, x)|^2 \psi(x) \varrho_\varepsilon(y - x) dy dx dt \\ & \leq C \text{ess sup}_{t \in (0, \gamma)} \mathbb{E} \int_{D_{x,\varepsilon}^2} |u(t, x)|^2 dx, \end{aligned}$$

where  $D_{x,\varepsilon}^2 := \{x \in D : \varrho_\varepsilon(y - x) > 0, \exists y \in D_{y,\varepsilon}^1\}$ . We also have  $|D_{x,\varepsilon}^2| = O(\varepsilon)$ . Now, we choose a non-negative function  $w_\varepsilon \in C_c^\infty(\mathbb{R}^d)$  such that  $w_\varepsilon(x) = 1$  for all  $x \in D_{x,\varepsilon}^2$  and  $|\text{supp}w_\varepsilon| \leq C\varepsilon$ . Suppose that  $s \in (0, \gamma)$  be the Lebesgue point of the function

$$t \mapsto \mathbb{E} \int_x |u(t, x)|^2 w_\varepsilon(x),$$

and  $\theta \in (0, T - \gamma)$ . Define  $V_{(\theta)} : [0, T] \rightarrow \mathbb{R}^+$  by  $V_{(\theta)}(0) := 1$  and  $V_{(\theta)}' := -\theta^{-1} \mathbf{1}_{[s, s+\theta]}$ . We take a sequence of non-negative functions  $\varphi_{\theta,n} \in C_c^\infty([0, T])$  satisfying

$$\|\varphi_{\theta,n}\|_{L^\infty(0, T)} \vee \|\partial_t \varphi_{\theta,n}\|_{L^1(0, T)} \leq 1,$$

such that

$$\lim_{n \rightarrow \infty} \|\varphi_{\theta,n} - V_{(\theta)}\|_{H^1(0, T)} = 0.$$

For each  $\delta > 0$ , define  $\eta_\delta \in C^2(\mathbb{R})$  by

$$(6.4) \quad \eta_\delta(0) = \eta_\delta'(0) = 0, \quad \eta_\delta''(r) := 2 \cdot \mathbf{1}_{[0, \delta^{-1})}(|r|) + (-|r| + \delta^{-1} + 2) \cdot \mathbf{1}_{[\delta^{-1}, \delta^{-1}+2)}(|r|).$$

It is easy to find that  $\eta_\delta(r) \rightarrow r^2$  as  $\delta \rightarrow 0^+$ . Using the entropy inequality (2.3) for  $u$  with  $\eta = \eta_\delta \in \mathcal{E}_0$  and  $\phi_n(t, x) = \varphi_{\theta,n}(t)w_\varepsilon(x)$ , letting  $n \rightarrow \infty$  and taking expectations, with Assumption 2.1 and (2.4)-(2.10) in Assumption 2.4, we have

$$\frac{1}{\theta} \int_s^{s+\theta} \mathbb{E} \int_D \eta_\delta(u) w_\varepsilon(x) dx dt$$

$$\begin{aligned} &\leq \mathbb{E} \int_D \eta_\delta(\xi) w_\varepsilon(x) dx + \mathbb{E} \int_0^{s+\theta} \int_D \eta_\delta''(u) |\nabla[\mathbf{a}](u)|^2 w_\varepsilon(x) dx dt \\ &\quad + C \mathbb{E} \int_0^{s+\theta} \int_D (1 + |u|^{m+1}) \left( \sum_{i,j} |\partial_{x_i x_j} w_\varepsilon| + \sum_i |\partial_{x_i} w_\varepsilon| + w_\varepsilon \right) dx dt. \end{aligned}$$

Taking  $\delta \rightarrow 0^+$  and then letting  $\theta \rightarrow 0^+$ , we have

$$\begin{aligned} &\mathbb{E} \int_D |u(s, x)|^2 w_\varepsilon(x) dx dt \\ &\leq \mathbb{E} \int_D |\xi(x)|^2 w_\varepsilon(x) dx + 2 \mathbb{E} \int_0^s \int_D |\nabla[\mathbf{a}](u)|^2 w_\varepsilon(x) dx dt \\ &\quad + C \mathbb{E} \int_0^s \int_D (1 + |u|^{m+1}) \left( \sum_{i,j} |\partial_{x_i x_j} w_\varepsilon| + \sum_i |\partial_{x_i} w_\varepsilon| + w_\varepsilon \right) dx dt \end{aligned}$$

hold for almost all  $s \in (0, \gamma)$ . Combining (6.3) and taking  $\gamma \rightarrow 0$ , we have

$$(6.5) \quad \begin{aligned} &\limsup_{\gamma \rightarrow 0} \frac{2}{\gamma} \mathbb{E} \int_0^\gamma \int_D \int_{D_{y,\varepsilon}^1} |u(t, x) - \bar{\xi}(y)|^2 \psi(x) \varrho_\varepsilon(y - x) dy dx dt \\ &\leq C \limsup_{\gamma \rightarrow 0} \text{ess sup}_{t \in (0, \gamma)} \mathbb{E} \int_{D_{x,\varepsilon}^2} |u(t, x)|^2 dx \leq C \mathbb{E} \int_{D \cap \text{supp } w_\varepsilon} |\xi|^2 dx. \end{aligned}$$

Now, we estimate the second term on the right hand side of (6.2). For  $\gamma \in [0, T]$ , choose a decreasing, non-negative function  $\varpi \in C^\infty([0, T])$  such that

$$\varpi(0) = 2, \varpi \leq 2 \cdot \mathbf{1}_{[0, 2\gamma]}, \partial_t \varpi \leq -\frac{1}{\gamma} \cdot \mathbf{1}_{[0, \gamma]}.$$

Note that  $\psi(\cdot) \varrho_\varepsilon(y - \cdot) \in C_c^\infty(D)$  for all  $y \in \bar{D}$  and sufficient small  $\varepsilon > 0$ . For fixed  $(y, z) \in D \times \mathbb{R}$ , using the entropy inequality (2.3) with  $\phi(t, x) = \varpi(t) \psi(x) \varrho_\varepsilon(y - x)$  and  $\eta(r) = \eta_\delta(r - z)$  defined in (6.4), with Assumptions 2.1-2.4, we have

$$\begin{aligned} &-\int_{t,x} \eta_\delta(u - z) \partial_t \varpi(t) \psi(x) \varrho_\varepsilon(y - x) \\ &\leq 2 \int_x \eta_\delta(\xi - z) \psi(x) \varrho_\varepsilon(y - x) + \frac{C}{\varepsilon^2} \int_{t,x} (1 + |u|^{m+1} + |z|^{m+1}) \varpi(t) \\ &\quad - \int_{t,x} \eta_\delta''(u - z) |\nabla_x[\mathbf{a}](u)|^2 \phi + \int_0^T \int_x \left( \eta_\delta'(u - z) \phi \sigma_{x_i}^{ik}(x, u) \right. \\ &\quad \left. - [\sigma_{rx_i}^{ik} \eta_\delta'(\cdot - z)](x, u) \phi - [\sigma_r^{ik} \eta_\delta'(\cdot - z)](x, u) \partial_{x_i} \phi \right) dW^k(t). \end{aligned}$$

Notice that all the terms are continuous in  $z \in \mathbb{R}$ . Replacing  $z$  by  $\xi(y)$  by convolution, taking expectations, integrating over  $y \in D$  and taking  $\delta \rightarrow 0^+, \gamma \rightarrow 0$ , with the definition of  $\varpi$ , we have

$$(6.6) \quad \begin{aligned} &\limsup_{\gamma \rightarrow 0} \frac{1}{\gamma} \int_0^\gamma \mathbb{E} \int_{x,y} |u(t, x) - \xi(y)|^2 \psi(x) \varrho_\varepsilon(y - x) dt \\ &\leq 2 \mathbb{E} \int_{x,y} |\xi(x) - \xi(y)|^2 \psi(x) \varrho_\varepsilon(x - y). \end{aligned}$$

Combining (6.2) with (6.5)-(6.6), we have

$$\limsup_{\gamma \rightarrow 0} \frac{1}{\gamma} \mathbb{E} \int_0^\gamma \int_x |u(t, x) - \xi(x)|^2 \psi(x) dt$$

$$\begin{aligned} &\leq 2\mathbb{E} \int_D \int_{\mathbb{R}^d} |\bar{\xi}(y) - \xi(x)|^2 \psi(x) \varrho_\varepsilon(y-x) dy dx \\ &\quad + 4\mathbb{E} \int_{x,y} |\xi(x) - \xi(y)|^2 \psi(x) \varrho_\varepsilon(x-y) + C\mathbb{E} \int_{D \cap \text{supp} w_\varepsilon} |\xi|^2 dx. \end{aligned}$$

Since  $\xi \in L_{m+1}(\Omega \times D)$ , the right hand side goes to 0 as  $\varepsilon \rightarrow 0$ . The proof is complete.  $\square$

**Lemma 6.2.** *Let  $u \in L_{m+1}(\Omega \times D_T)$  and for some  $\varepsilon \in (0, 1)$ . Denote by  $N$  the constant in Remark 3.1. Let  $\varrho : \mathbb{R}^d \rightarrow \mathbb{R}$  be a non-negative function, integrating to 1 and supported on a ball of radius  $N\varepsilon$  centered at the origin. Under Assumption 2.1, one has*

$$(6.7) \quad \begin{aligned} &\mathbb{E} \int_{t,x,y} |u(t,x) - u(t,y)| \varrho(x-y) \\ &\leq C(\varepsilon^{\frac{m}{m+1}} + \varepsilon^{\frac{2}{m+1}}) \mathbb{E}(1 + \|u\|_{L_{m+1}(D_T)}^{m+1} + \|\nabla[\mathbf{a}](u)\|_{L_1(D_T)}), \end{aligned}$$

$$(6.8) \quad \begin{aligned} &\mathbb{E} \int_{t,x,y} |\Phi(u(t,x)) - \Phi(u(t,y))| \varrho(x-y) \\ &\leq C\varepsilon^{\frac{1}{m+1}} \mathbb{E}(1 + \|u\|_{L_{m+1}(D_T)}^{m+1} + \|\nabla[\mathbf{a}](u)\|_{L_2(D_T)}^2), \end{aligned}$$

where  $C$  depends on  $d, K$  and  $T$ .

*Proof.* We first estimate (6.8). Define a set

$$D_{-\varepsilon} := \{y \in D : y + z \in D, \text{ for all } z \in \mathbb{R}^d \text{ with } |z| \leq N\varepsilon\}.$$

Notice that  $|D \setminus D_{-\varepsilon}| = O(\varepsilon)$ , combining with Assumption 2.1 and Hölder's inequality, we have

$$\begin{aligned} &\mathbb{E} \int_{t,x,y} |\Phi(u(t,x)) - \Phi(u(t,y))| \varrho(x-y) \\ &\leq \mathbb{E} \int_t \int_{D_{-\varepsilon}} \int_{\mathbb{R}^d} |\Phi(u(t,y+z)) - \Phi(u(t,y))| \varrho(z) dz dy \\ &\quad + \mathbb{E} \int_{t,x} \int_{D \setminus D_{-\varepsilon}} |\Phi(u(t,x)) - \Phi(u(t,y))| \varrho(x-y) dy \\ &\lesssim \varepsilon \mathbb{E} \int_t \int_{D_{-\varepsilon}} \int_{\mathbb{R}^d} \varrho(z) \int_0^1 |\mathbf{a}(u) \nabla[\mathbf{a}](u)|(y+\lambda z) d\lambda dz dy \\ &\quad + \left( \mathbb{E} \int_{t,x} \int_{D \setminus D_{-\varepsilon}} |\Phi(u(t,x)) - \Phi(u(t,y))|^{\frac{m+1}{m}} \varrho(x-y) dy \right)^{\frac{m}{m+1}} \\ &\quad \cdot \left( \mathbb{E} \int_{t,x} \int_{D \setminus D_{-\varepsilon}} \varrho(x-y) dy \right)^{\frac{1}{m+1}} \\ &\leq C\varepsilon^{\frac{1}{m+1}} \mathbb{E}(1 + \|u\|_{L_{m+1}(D_T)}^{m+1} + \|\nabla[\mathbf{a}](u)\|_{L_2(D_T)}^2). \end{aligned}$$

Moreover, using the proceeding method and the ones in the proof of [DGG19, Lemma 3.1], (6.7) can be easily proved and we omit it here.  $\square$

*Acknowledgement.* The research of K. Du was partially supported by National Natural Science Foundation of China (12222103), by National Key R&D Program of China (2018YFA0703900), and by Natural Science Foundation of Shanghai (20ZR1403600).

## REFERENCES

- [ABK00] Boris P Andreianov, Philippe Benilan, and Stanislav N Kruzhkov,  *$L^1$ -theory of scalar conservation law with continuous flux function*, Journal of Functional Analysis **171** (2000), no. 1, 15–33.
- [BB11] Haim Brezis and Haim Brézis, *Functional analysis, sobolev spaces and partial differential equations*, vol. 2, Springer, 2011.
- [BCF88] Z Brzeźniak, M Capiński, and F Flandoli, *A convergence result for stochastic partial differential equations*, Stochastics: An International Journal of Probability and Stochastic Processes **24** (1988), no. 4, 423–445.
- [BDPR08] Viorel Barbu, Giuseppe Da Prato, and Michael Röckner, *Existence and uniqueness of nonnegative solutions to the stochastic porous media equation*, Indiana University mathematics journal (2008), 187–211.
- [BDPR16] ———, *Stochastic porous media equations*, vol. 2163, Springer, 2016.
- [BGV20] L’ubomír Bañas, Benjamin Gess, and Christian Vieth, *Numerical approximation of singular-degenerate parabolic stochastic PDEs*, arXiv preprint arXiv:2012.12150 (2020).
- [BVW12] Caroline Bauzet, Guy Vallet, and Petra Wittbold, *The Cauchy problem for conservation laws with a multiplicative stochastic perturbation*, Journal of Hyperbolic Differential Equations **9** (2012), no. 04, 661–709.
- [BVW14] ———, *The Dirichlet problem for a conservation law with a multiplicative stochastic perturbation*, Journal of Functional Analysis **266** (2014), no. 4, 2503–2545.
- [BVW15] ———, *A degenerate parabolic–hyperbolic Cauchy problem with a stochastic force*, Journal of Hyperbolic Differential Equations **12** (2015), no. 03, 501–533.
- [Car99] José Carrillo, *Entropy solutions for nonlinear degenerate problems*, Archive for rational mechanics and analysis **147** (1999), no. 4, 269–361.
- [CDK12] Gui-Qiang Chen, Qian Ding, and Kenneth H Karlsen, *On nonlinear stochastic balance laws*, Archive for Rational Mechanics and Analysis **204** (2012), no. 3, 707–743.
- [Cio20] Ioana Ciotir, *Stochastic porous media equations with divergence itô noise*, Evolution Equations & Control Theory **9** (2020), no. 2, 375.
- [Cli22] Andrea Clini, *Porous media equations with nonlinear gradient noise and Dirichlet boundary conditions*, arXiv preprint arXiv:2205.00836 (2022).
- [DD05] Constantine M Dafermos and Constantine M Dafermos, *Hyperbolic conservation laws in continuum physics*, vol. 3, Springer, 2005.
- [DG20] Konstantinos Dareiotis and Benjamin Gess, *Nonlinear diffusion equations with nonlinear gradient noise*, Electronic Journal of Probability **25** (2020).
- [DGG19] Konstantinos Dareiotis, Maté Gerencsér, and Benjamin Gess, *Entropy solutions for stochastic porous media equations*, Journal of Differential Equations **266** (2019), no. 6, 3732–3763.
- [DGT20] Konstantinos Dareiotis, Benjamin Gess, and Pavlos Tsatsoulis, *Ergodicity for stochastic porous media equations with multiplicative noise*, SIAM Journal on Mathematical Analysis **52** (2020), no. 5, 4524–4564.
- [DHV<sup>+</sup>16] Arnaud Debussche, Martina Hofmanová, Julien Vovelle, et al., *Degenerate parabolic stochastic partial differential equations: Quasilinear case*, The Annals of Probability **44** (2016), no. 3, 1916–1955.
- [DSZ16] Nicolas Dirr, Marios Stamatakis, and Johannes Zimmer, *Entropic and gradient flow formulations for nonlinear diffusion*, Journal of Mathematical Physics **57** (2016), no. 8, 081505.
- [FG19] Benjamin Fehrman and Benjamin Gess, *Well-posedness of nonlinear diffusion equations with nonlinear, conservative noise*, Archive for Rational Mechanics and Analysis **233** (2019), no. 1, 249–322.
- [FG21] ———, *Path-by-path well-posedness of nonlinear diffusion equations with multiplicative noise*, Journal de Mathématiques Pures et Appliquées **148** (2021), 221–266.
- [FN08] Jin Feng and David Nualart, *Stochastic scalar conservation laws*, Journal of Functional Analysis **255** (2008), no. 2, 313–373.
- [GH18] Benjamin Gess and Martina Hofmanová, *Well-posedness and regularity for quasilinear degenerate parabolic-hyperbolic SPDE*, The annals of probability **46** (2018), no. 5, 2495–2544.
- [GS17] Benjamin Gess and Panagiotis E Souganidis, *Stochastic non-isotropic degenerate parabolic-hyperbolic equations*, Stochastic Processes and their Applications **127** (2017), no. 9, 2961–3004.
- [Hen21] Sebastian Hensel, *Finite time extinction for the 1D stochastic porous medium equation with transport noise*, Stochastics and Partial Differential Equations: Analysis and Computations **9** (2021), no. 4, 892–939.
- [Kim03] Jong Uhn Kim, *On a stochastic scalar conservation law*, Indiana University mathematics journal (2003), 227–256.



- [Kru70] Stanislav N Kružkov, *First order quasilinear equations in several independent variables*, Mathematics of the USSR-Sbornik **10** (1970), no. 2, 217.
- [Kry13] Nikolai Vladimirovich Krylov, *A relatively short proof of Itô's formula for SPDEs and its applications*, Stochastic Partial Differential Equations: Analysis and Computations **1** (2013), no. 1, 152–174.
- [Kun97] Hiroshi Kunita, *Stochastic flows and stochastic differential equations*, vol. 24, Cambridge university press, 1997.
- [LL06a] Jean-Michel Lasry and Pierre-Louis Lions, *Jeux à champ moyen. i—le cas stationnaire*, Comptes Rendus Mathématique **343** (2006), no. 9, 619–625.
- [LL06b] ———, *Jeux à champ moyen. ii—horizon fini et contrôle optimal*, Comptes Rendus Mathématique **343** (2006), no. 10, 679–684.
- [LL07] ———, *Mean field games*, Japanese journal of mathematics **2** (2007), no. 1, 229–260.
- [LT21] Ruoyang Liu and Shanjian Tang, *The obstacle problem for stochastic porous media equations*, arXiv preprint arXiv:2111.10736 (2021).
- [MR18] Ionuț Munteanu and Michael Röckner, *Total variation flow perturbed by gradient linear multiplicative noise*, Infinite Dimensional Analysis, Quantum Probability and Related Topics **21** (2018), no. 01, 1850003.
- [RRW07] Jiagang Ren, Michael Röckner, and Feng-Yu Wang, *Stochastic generalized porous media and fast diffusion equations*, Journal of Differential Equations **238** (2007), no. 1, 118–152.
- [Ser99] Denis Serre, *Systems of conservation laws 1: Hyperbolicity, entropies, shock waves*, Cambridge University Press, 1999.
- [Váz07] Juan Luis Vázquez, *The porous medium equation: mathematical theory*, Oxford University Press on Demand, 2007.
- [VW09] Guy Vallet and Petra Wittbold, *On a stochastic first-order hyperbolic equation in a bounded domain*, Infinite Dimensional Analysis, Quantum Probability and Related Topics **12** (2009), no. 04, 613–651.

(K. Du) SHANGHAI CENTER FOR MATHEMATICAL SCIENCES, FUDAN UNIVERSITY, SHANGHAI 200438, CHINA.  
Email address: [kdu@fudan.edu.cn](mailto:kdu@fudan.edu.cn)

(R. Liu and Y. Wang) SCHOOL OF MATHEMATICAL SCIENCES, FUDAN UNIVERSITY, SHANGHAI 200433, CHINA.  
Email address: [ruoyangliu18@fudan.edu.cn](mailto:ruoyangliu18@fudan.edu.cn) (R. Liu), [wangyuxing@fudan.edu.cn](mailto:wangyuxing@fudan.edu.cn) (Y. Wang)