

THE JUCYS-MURPHY BASIS AND SEMISIMPLICITY CRITERIA FOR THE q -BRAUER ALGEBRA

HEBING RUI, MEI SI AND LINLIANG SONG

ABSTRACT. We construct the Jucys-Murphy elements and the Jucys-Murphy basis for the q -Brauer algebra in the sense of [11, Definition 2.4]. We also give a necessary and sufficient condition for the q -Brauer algebra being (split) semisimple over an arbitrary field.

1. INTRODUCTION

In his remarkable work [23], Schur classified the polynomial representations of the complex general linear group by decomposing tensor products of its natural representation. The symmetric group appears naturally in Schur's work and plays an important role. Later on Brauer [2] studied the related problem for the complex orthogonal group and the complex symplectic group and found a new family of complex associative algebras called the Brauer algebras or Brauer centralizer algebras. These algebras play roles similar to those of symmetric groups.

There are many ways to generalize results in [2, 23]. See [6, 25] where the complex classical groups are replaced by quantum enveloping algebras of classic types. Hecke algebras and Birman-Murakami-Wenzl algebras [1, 13] come into the picture. In [24], Wenzl considered tensor products of the natural representation of certain coideal algebras [7] and introduced a new associative algebra, called the q -Brauer algebra $\mathcal{B}_n(q, z)$ for any positive integer $n \geq 2$. See also [3] for the explicit Schur-Weyl duality between the q -quantum groups of types AI or AII and the q -Brauer algebras. The q -Brauer algebra is isomorphic to an endomorphism algebra of the q -Brauer category [20] and is isomorphic to a quotient of the associative algebra introduced by Molev [12].

The q -Brauer algebra has been used to construct new subfactors of type II_1 factors [27]. It has some application in module categories of fusion categories of type A corresponding to certain symmetric spaces [26]. In [24, Theorem 3.8], Wenzl proved that $\mathcal{B}_n(q, z)$ is free over a commutative ring with rank $(2n - 1)!!$. When the ground ring is a field, Wenzl also proved that $\mathcal{B}_n(q, z)$ is semisimple if $z^2 \neq q^{2k}$ for $k \in \mathbb{Z}$, $|k| \leq n$ and q^2 is not an ℓ -th root of unity, $\ell \leq n$ [24, Theorem 5.3]. Under the assumption $z^2 \neq 1$, Dung Tien Nguyen proved that $\mathcal{B}_n(q, z)$ is a cellular algebra in the sense of [8, Definition 1.1] and classified simple $\mathcal{B}_n(q, z)$ -modules over an arbitrary field [22, Theorem 3.2, Theorem 4.1]. He also gave an explicit condition for $\mathcal{B}_3(q, z)$ being semisimple over an arbitrary field [22, Proposition 5.1]. Dung Tien Nguyen's conditions on parameters [22, Example 5.7] are different from those for the corresponding Birman-Murakami-Weznl algebra in [17, Theorem 5.9]. Therefore the q -Brauer algebra is another q -deformation of the Brauer algebra.

We expect to generalize our previous results in [17, 18] to the q -Brauer algebra. More explicitly, we want to construct Jucys-Murphy basis and orthogonal basis of the q -Brauer algebra so as to compute Gram determinants, determine explicitly the semisimplicity criteria, classify blocks and compute decomposition matrices (when q is not a root of unity) etc. The first difficult problem that we faced is to find out a family of commutative elements in $\mathcal{B}_n(q, z)$ called the Jucys-Murphy elements or JM elements in the sense of [11, Definition 2.4]. By studying classical branching rule for the q -Brauer algebra in details, we are able to define the Jucys-Murphy basis of the q -Brauer algebra. Since the summation of Jucys-Murphy elements is not a central element, we have to make extra efforts to prove that Jucys-Murphy elements act on Jucys-Murphy basis as upper-triangular matrices. As a by-product one can use arguments in [11] to define an orthogonal basis of $\mathcal{B}_n(q, z)$ in generic case. In this paper, we do not give details since we have not found out an efficient way to compute Gram

Date: November 29, 2022.

2010 Mathematics Subject Classification. 17B10, 18D10, 33D80.

Key words and phrases. q -Brauer algebras, Jucys-Murphy basis, Semisimplicity.

H. Rui is supported partially by NSFC (grant No. 11971351). M. Si and L. Song are supported partially by NSFC (grant No. 12071346) and China Scholarship Council.

determinants associated to cell modules of $\mathcal{B}_n(q, z)$, and we could not use the method in [17] to give a semisimplicity criteria for $\mathcal{B}_n(q, z)$ over an arbitrary field. We describe explicitly the radicals of certain cell modules in section six and finally are able to prove the following result:

Theorem A. For any $n \geq 2$, let $\mathcal{B}_n(q, z)$ be defined over a field F which contains invertible elements $z, z - z^{-1}, q$ and $q - q^{-1}$. Let e be the quantum characteristic of q^2 . Then $\mathcal{B}_n(q, z)$ is (split) semisimple if and only if $e > n$ and $z^2 \neq q^{2a}$ where

$$a \in \{i \in \mathbb{Z} \mid 4 - 2n \leq i \leq n - 2\} \setminus \{i \in \mathbb{Z} \mid 4 - 2n < i \leq 3 - n, 2 \nmid i\}. \quad (1.1)$$

Here is a brief outline of the content of this paper. In the second section, we recall some basic results on the q -Brauer algebra. In section 3, we study the classical branching rule. In section 4, we construct the Jucys-Murphy basis and the Jucys-Murphy elements of the q -Brauer algebra. Restriction and induction functors are given in section 5. We describe explicitly the radical of certain cell modules in section 6. This enables us to prove Theorem A in section 7.

2. THE q -BRAUER ALGEBRA

In this section, we recall the definitions of the Hecke algebra and the q -Brauer algebra and give some basic results on them.

2.1. The Hecke algebra associated to the symmetric group. The symmetric group \mathfrak{S}_n is the Coxeter group with $\{s_i \mid 1 \leq i \leq n - 1\}$ as its distinguished generators subject to the relations:

$$s_i^2 = 1, \quad s_i s_j = s_j s_i, \quad s_k s_{k+1} s_k = s_{k+1} s_k s_{k+1} \quad (2.1)$$

where $1 < |i - j|$ and $1 \leq k \leq n - 2$. In this paper we assume $\mathcal{Z} = \mathbb{Z}[q, q^{-1}]$, the ring of Laurent polynomials in indeterminate q with coefficients in \mathbb{Z} . The Hecke algebra \mathcal{H}_n associated to \mathfrak{S}_n is the unital associative \mathcal{Z} -algebra generated by $\{T_i \mid 1 \leq i \leq n - 1\}$ subject to the relations

$$(T_i - q)(T_i + q^{-1}) = 0, \quad T_i T_j = T_j T_i, \quad T_k T_{k+1} T_k = T_{k+1} T_k T_{k+1} \quad (2.2)$$

where $1 < |i - j|$ and $1 \leq k \leq n - 2$.

Suppose $w \in \mathfrak{S}_n$. Then $s_{i_1} s_{i_2} \cdots s_{i_k}$ is a reduced expression of w if $w = s_{i_1} s_{i_2} \cdots s_{i_k}$ and k is minimal. In this case, k is the length of w and is denoted by $\ell(w)$. Let $T_w = T_{i_1} T_{i_2} \cdots T_{i_k}$ if $s_{i_1} s_{i_2} \cdots s_{i_k}$ is a reduced expression of w . It is known that T_w is independent of any reduced expression of w and $\{T_w \mid w \in \mathfrak{S}_n\}$ is a \mathcal{Z} -basis of \mathcal{H}_n . Let $\tau : \mathcal{H}_n \rightarrow \mathcal{Z}$ such that $\tau(\sum_w a_w T_w) = a_1$ where 1 is the identity element of \mathfrak{S}_n . Then τ is a trace function on \mathcal{H}_n such that

$$\tau(T_x T_y) = \delta_{x, y^{-1}} \quad (2.3)$$

for all $x, y \in \mathfrak{S}_n$ [9, Proposition 1.16].

2.2. The q -Brauer algebra. Let $\mathbb{k} = \mathbb{Z}[q, q^{-1}, z, z^{-1}, (q - q^{-1})^{-1}, (z - z^{-1})^{-1}]$, where z is another indeterminate. In this paper we always assume

$$\delta = \frac{z - z^{-1}}{q - q^{-1}}. \quad (2.4)$$

Then δ is invertible in \mathbb{k} . The original q -Brauer algebra is available over $\mathbb{Z}[q, q^{-1}, z, z^{-1}, (q - q^{-1})^{-1}]$. In this case, δ may not be invertible. In [21], Dung Tien Nguyen proved that q -Brauer algebra is a cellular algebra in the sense of [8, Definition 1.1] under the assumption that δ is invertible. Since our results depend on the cellular structure of the q -Brauer algebra, we need this additional assumption.

Definition 2.1. [24, Definition 3.1] [22, Definition 3.1] Suppose $n \geq 2$. The q -Brauer algebra $\mathcal{B}_n(q, z)$ or just \mathcal{B}_n is a unital associative \mathbb{k} -algebra generated by T_1, T_2, \dots, T_{n-1} and E_1 subject to the relations in (2.2) together with

- (1) $E_1^2 = \delta E_1$,
 - (2) $T_1 E_1 = q E_1 = E_1 T_1$,
 - (3) $E_1 T_2 E_1 = z E_1$,
 - (4) $T_i E_1 = E_1 T_i$ if $i > 2$,
 - (5) $T_2 T_3 T_1^{-1} T_2^{-1} E_{(2)} = E_{(2)} = E_{(2)} T_2 T_3 T_1^{-1} T_2^{-1}$,
- where $E_{(2)} = E_1 T_2 T_3 T_1^{-1} T_2^{-1} E_1$.

Remarks 2.2. Suppose $z = q^a$ for some $a \in \mathbb{Z} \setminus \{0\}$. It follows from [24, Remark 3.1] that the classical limit of $\mathcal{B}_n(q, q^a)$ is the Brauer algebra $B_n(a)$ over \mathbb{Z} . In this case, $\lim_{q \rightarrow 1} \delta = a$ and T_i becomes the simple reflection s_i and E_1 can be identified with the corresponding element in $B_n(a)$.

We write $\mathcal{B}_0 = \mathcal{B}_1 = \mathbb{k}$. Thanks to [24, Theorem 3.8], \mathcal{B}_n is free over \mathbb{k} with rank $(2n - 1)!!$ and the subalgebra of \mathcal{B}_n generated by T_1, T_2, \dots, T_{n-1} is isomorphic to the Hecke algebra \mathcal{H}_n . For this reason, one can use symbols T_i 's in both \mathcal{H}_n and \mathcal{B}_n . Moreover,

$$\mathcal{H}_n \cong \mathcal{B}_n / \langle E_1 \rangle \quad (2.5)$$

where $\langle E_1 \rangle$ is the two-sided ideal of \mathcal{B}_n generated by E_1 . Let F be a field containing units $z, q, q - q^{-1}, z - z^{-1}$. Throughout,

$$\mathcal{B}_{n,F} := \mathcal{B}_n \otimes_{\mathbb{k}} F,$$

where F can be considered as the left \mathbb{k} -module on which $z, q \in \mathbb{k}$ act via the corresponding z and q in F . If there is no confusion, we also denote $\mathcal{B}_{n,F}$ by \mathcal{B}_n .

Lemma 2.3. [21, Proposition 3.14] *There is a \mathbb{k} -linear anti-involution $\sigma : \mathcal{B}_n \rightarrow \mathcal{B}_n$ such that $\sigma(x) = x$ for any $x \in \{E_1, T_1, T_2, \dots, T_{n-1}\}$.*

Definition 2.4. Suppose $0 \leq k \leq \lfloor \frac{n}{2} \rfloor - 1$ and $1 \leq \ell \leq n - 1$. Define

$$(1) \ E^0 = 1 \text{ and } E^{k+1} = E_1 T_{2,2k+2} T_{2k+1,1}^{-1} E^k, \text{ where } T_{i,j} = T_{s_{i,j}}$$

$$s_{i,j} = \begin{cases} s_i s_{i+1, j} & \text{if } i < j, \\ 1 & \text{if } i = j, \\ s_{i,j+1} s_j & \text{if } i > j. \end{cases}$$

$$(2) \ E_\ell = T_{\ell,1} T_{2,\ell+1}^{-1} E_1 T_{2,\ell+1} T_{\ell,1}^{-1}.$$

Definition 2.4(1) was given in [24, (3.2)] where E^k was denoted by $E_{(k)}$.

Lemma 2.5. [24, (3.2), Lemma 3.3], [21, Lemma 3.4, Proposition 3.14] *Suppose $1 \leq k \leq \lfloor \frac{n}{2} \rfloor$ and $1 \leq \ell \leq \lfloor \frac{n}{2} \rfloor - 1$.*

$$\begin{aligned} (1) \ E^{\ell+1} &= E^\ell T_{1,2\ell+1}^{-1} T_{2\ell+2,2} E_1, & (4) \ E^k T_{2l+1,1} &= E^k T_{2,2l+2} \text{ if } l < k, \\ (2) \ \sigma(E^k) &= E^k, & (5) \ E^k T_{1,2l+1}^{-1} &= E^k T_{2l+2,2}^{-1} \text{ if } l < k, \\ (3) \ E_1 T_2^{-1} T_1^{-1} T_3 T_2 E_1 &= E^2 = E_1 T_2 T_1 T_3^{-1} T_2^{-1} E_1, & (6) \ E^k T_{2k-1} &= q E^k. \end{aligned}$$

Lemma 2.6. *For all admissible i and j , we have:*

$$\begin{aligned} (1) \ E_{i+1} &= T_i T_{i+1}^{-1} E_i T_{i+1} T_i^{-1}, & (5) \ E_i T_{i+1}^{\pm 1} E_i &= z^{\pm 1} E_i, \\ (2) \ T_i E_j &= E_j T_i \text{ if } |i - j| \geq 2, & (6) \ E_i^2 &= \delta E_i, \\ (3) \ E_i E_j &= E_j E_i \text{ if } |i - j| \geq 2, & (7) \ E_1 T_{2i,2}^{-1} T_{1,2i-1} E^{i-1} &= E^i \text{ [3, Lemma 3.3]}, \\ (4) \ E^j &= E_1 E_3 \cdots E_{2j-1}, & (8) \ E_{2i-1} T_{2i-1}^{-1} T_{2i}^{-1} E^i &= z^{-1} q^{-1} E^i. \end{aligned}$$

Proof. Obviously (1) follows immediately from Definition 2.4(2). If $i \geq j + 2$, then (2) follows from (2.2) and Definition 2.1(4). If $i \leq j - 2$, then $T_i E_j = T_{j,1} T_{2,j+1}^{-1} T_{i+2} E_1 T_{2,j+1} T_{j,1}^{-1} = E_j T_i$ and (2) follows. It is not difficult to verify

$$E_1 E_3 = E_{(2)} = E_3 E_1. \quad (2.6)$$

If $j > 3$, then $E_1 E_j = E_1 T_{j-1} T_j^{-1} E_{j-1} T_j T_{j-1}^{-1} = E_j E_1$, where the last equality follows from the induction assumption on $j - 1$ and Definition 2.1(4). Suppose $i > 1$. We can assume $j \geq i + 2$ without losing of any generality when we prove (3). By (2) and induction assumption on $i - 1$, $E_i E_j = T_{i-1} T_i^{-1} E_{i-1} T_i T_{i-1}^{-1} E_j = E_j E_i$. In any case, we have verified (3). Thanks to (2.6), $E^2 = E_1 E_3$. In general, using (3), induction assumption on $j - 1$, Lemma 2.5(5) and σ in Lemma 2.3 yields

$$\begin{aligned} E_1 E_3 \cdots E_{2j-1} &= T_{2j-1,1} T_{2,2j}^{-1} E_1 T_{2,2j} T_{2j-1,1}^{-1} E^{j-1} \\ &= T_{2j-1,1} T_{2,2j}^{-1} E^j \\ &= E^j, \end{aligned}$$

proving (4). $E_1 T_2 E_1 = z E_1$ is given in Definition 2.1(3). The required formula for $E_1 T_2^{-1} E_1$ follows from the quadratic relation in (2.2) and Definition 2.1(3). In general, by induction assumption on $i - 1$, we have

$$E_i T_{i+1}^{\pm 1} E_i = T_{i-1} T_i^{-1} E_{i-1} T_{i+1}^{-1} T_i^{\pm 1} T_{i+1} E_{i-1} T_i T_{i-1}^{-1} = z^{\pm 1} E_i,$$

proving (5). Obviously, (6) follows from Definition 2.1(1) and (7) was given in [3, Lemma 3.3]. Finally, (8) follows from the following computation:

$$\begin{aligned}
E_{2i-1}T_{2i-1}^{-1}T_{2i}^{-1}E^i &= T_{2i-1,1}T_{2,2i}^{-1}E_1T_{2i+1,1}^{-1}T_{1,2i-1}E^i \\
&= q^{-1}T_{2i-1,1}T_{2,2i}^{-1}E_1T_{2i+1,2}^{-1}T_{1,2i-1}E^i \\
&= q^{-1}T_{2i-1,1}T_{2,2i}^{-1}E_1T_{2i,2}^{-1}T_{1,2i-1}E^{i-1}T_{2i}^{-1}E_{2i-1} \\
&= q^{-1}T_{2i-1,1}T_{2,2i}^{-1}E^i T_{2i}^{-1}E_{2i-1} \\
&= z^{-1}q^{-1}T_{2i-1,1}T_{2,2i}^{-1}E^i \\
&= z^{-1}q^{-1}E^i
\end{aligned}$$

where the fourth (resp., fifth, resp., sixth) equation follows from (7) (resp., (5), resp., Lemma 2.5(2)(5)).
□

Proposition 2.7. *Suppose $n \geq 2$ and $\tilde{E}_1 = qz^{-1}T_1^{-1}T_2E_1$. Then $\tilde{E}_1^2 = \tilde{E}_1$. Moreover,*

- (1) *There is an algebra homomorphism $\phi : \mathcal{B}_{n-2} \rightarrow \tilde{E}_1\mathcal{B}_n\tilde{E}_1$ sending $1, E_1, T_i$ to $\tilde{E}_1, \tilde{E}_1E_3$ and $\tilde{E}_1T_{i+2}, 1 \leq i \leq n-3$, respectively.*
- (2) *$\phi(E^i) = qz^{-1}T_1^{-1}T_2E^{i+1}$ for any $0 \leq i \leq \lfloor \frac{n}{2} \rfloor - 1$.*

Proof. It is easy to verify that $\tilde{E}_1^2 = \tilde{E}_1$, $\tilde{E}_1E_3\tilde{E}_1 = \tilde{E}_1E_3$ and $\tilde{E}_1T_i\tilde{E}_1 = \tilde{E}_1T_i$ if $3 \leq i \leq n-1$. So $\phi(E_1), \phi(T_1), \dots, \phi(T_{n-3})$ satisfy (2.2) and Definition 2.1(1)-(4). We remark that the quadratic relation in (2.2) becomes $(\tilde{E}_1T_i - q\tilde{E}_1)(\tilde{E}_1T_i + q^{-1}\tilde{E}_1) = 0, 3 \leq i \leq n-1$. We have

$$\begin{aligned}
\phi(E^2) &= \tilde{E}_1E_3T_4T_5T_3^{-1}T_4^{-1}\tilde{E}_1E_3 \\
&= qz^{-1}T_1^{-1}T_2E_1T_2T_3T_1^{-1}T_2^{-1}E_1T_4T_5T_3^{-1}T_4^{-1}\tilde{E}_1E_3 \\
&= qz^{-1}T_1^{-1}T_2E_1T_2T_3T_4T_5T_1^{-1}T_2^{-1}T_3^{-1}T_4^{-1}E^2 \\
&= qz^{-1}T_1^{-1}T_2E^3,
\end{aligned}$$

where the last equality follows from Definition 2.4(1). So,

$$\phi(E^2T_2T_3T_1^{-1}T_2^{-1}) = qz^{-1}T_1^{-1}T_2E^3T_4T_5T_3^{-1}T_4^{-1} = qz^{-1}T_1^{-1}T_2E^3 = \phi(T_2T_3T_1^{-1}T_2^{-1}E^2).$$

This proves that the images of generators also satisfy Definition 2.1(5) and hence ϕ is an algebra homomorphism. When $i = 0, 1$, (2) follows from the definition of ϕ . In general, by induction assumption on $i-1$ we have

$$\begin{aligned}
\phi(E^i) &= \phi(E_1T_{2,2i}T_{2i-1,1}^{-1}E^{i-1}) \\
&= qz^{-1}\phi(E_1T_{2,2i}T_{2i-1,1}^{-1})T_1^{-1}T_2E^i \\
&= q^2z^{-2}T_1^{-1}T_2E_1T_2T_3T_1^{-1}T_2^{-1}E_1T_{4,2i+2}T_{2i+1,3}^{-1}T_1^{-1}T_2E^i \\
&= q^2z^{-2}T_1^{-1}T_2E_1T_2T_3T_1^{-1}T_2^{-1}T_{4,2i+2}T_{2i+1,3}^{-1}E_1T_1^{-1}T_2E^i \\
&= qz^{-1}T_1^{-1}T_2E_1T_{2,2i+2}T_{2i+1,1}^{-1}E^i \\
&= qz^{-1}T_1^{-1}T_2E^{i+1},
\end{aligned}$$

where the last equality follows from Definition 2.4(1). This proves (2). □

3. THE CLASSICAL BRANCHING RULE

In subsection 3.1, we recall some well-known results on Hecke algebras. In the remaining part of this section, we study the classical branching rule for the q -Brauer algebra.

3.1. Cell filtration of cell modules and permutation modules for Hecke algebras. For any $0 \leq f \leq \lfloor n/2 \rfloor$, let $\mathfrak{S}_{2f+1,n}$ be the symmetric group on letters $\{2f+1, 2f+2, \dots, n\}$. When n is even and $f = n/2$, we set $\mathfrak{S}_{2f+1,n} = 1$. Let $\mathcal{H}_{2f+1,n}$ be the Hecke algebra associated to $\mathfrak{S}_{2f+1,n}$. Then $\mathcal{H}_{n-2f} \cong \mathcal{H}_{2f+1,n}$. The corresponding isomorphism sends T_i to $T_{2f+i}, 1 \leq i \leq n-2f-1$.

Recall that a composition λ of a non-negative integer d is a sequence $(\lambda_1, \lambda_2, \dots)$ of non-negative integers such that $|\lambda| := \sum_{i \geq 1} \lambda_i = d$. If $\lambda_i \geq \lambda_{i+1}$ for all possible i , then λ is called a partition. Given a positive integer e , a partition λ of d is called e -restricted if $\lambda_i - \lambda_{i+1} < e$ for all possible i . When $e > d$, any partition of d is e -restricted. Let $\Lambda(d)$ (resp., $\Lambda^+(d)$, resp., $\Lambda_e^+(d)$) be the set of

all compositions (resp., partitions, resp., e -restricted partitions) of d , where e is always the quantum characteristic of q^2 . Then e is the minimal positive integer such that

$$1 + q^2 + \cdots + q^{2e-2} = 0.$$

If such a positive integer does not exist, then $e = \infty$. In this case, q^2 is not a root of unity. It is known that each of $\Lambda(d)$, $\Lambda^+(d)$ and $\Lambda_e^+(d)$ is a poset with dominance order \leq defined on it such that $\mu \leq \lambda$ if $\sum_{j=1}^i \mu_j \leq \sum_{j=1}^i \lambda_j$ for all possible i . If $\mu \leq \lambda$ and $\mu \neq \lambda$, we write $\mu < \lambda$.

Each composition λ of $n - 2f$ corresponds to the Young diagram $[\lambda]$ such that there are λ_i boxes in the i -th row of $[\lambda]$. A λ -tableau \mathfrak{t} is obtained from $[\lambda]$ by inserting $2f + 1, 2f + 2, \dots, n$ into $[\lambda]$ without repetition. In this case, we write $\text{shape}(\mathfrak{t}) = \lambda$ and call λ the shape of \mathfrak{t} . If the entries in \mathfrak{t} increase from left to right along row and down column, \mathfrak{t} is called standard. In this case, $\lambda \in \Lambda^+(n - 2f)$. Let $\mathcal{T}^{std}(\lambda)$ be the set of all standard λ -tableaux. Then $\mathfrak{t}^\lambda \in \mathcal{T}^{std}(\lambda)$, where \mathfrak{t}^λ is obtained from $[\lambda]$ by inserting $2f + 1, 2f + 2, \dots, n$ successively from left to right along the rows of $[\lambda]$. For any $\lambda \in \Lambda^+(n - 2f)$, let \mathfrak{S}_λ be the Young subgroup with respect to λ . Then \mathfrak{S}_λ is the subgroup of $\mathfrak{S}_{2f+1, n}$ which stabilizes the entries in each row of \mathfrak{t}^λ . For example, \mathfrak{S}_λ is the subgroup of $\mathfrak{S}_{3,9}$ generated by s_3, s_4, s_6 and s_8 if $n = 9, f = 1$ and $\lambda = (3, 2, 2)$. In this case,

$$\mathfrak{t}^\lambda = \begin{array}{|c|c|c|} \hline 3 & 4 & 5 \\ \hline 6 & 7 & \\ \hline 8 & 9 & \\ \hline \end{array}.$$

For each $\mathfrak{t} \in \mathcal{T}^{std}(\lambda)$, there is a distinguished right coset representative $d(\mathfrak{t})$ of \mathfrak{S}_λ in $\mathfrak{S}_{2f+1, n}$ such that $\mathfrak{t} = \mathfrak{t}^\lambda d(\mathfrak{t})$. Suppose $\mathfrak{s}, \mathfrak{t} \in \mathcal{T}^{std}(\lambda)$. Following [9], define

$$x_{\mathfrak{s}\mathfrak{t}} = T_{d(\mathfrak{s})}^* x_\lambda T_{d(\mathfrak{t})},$$

where $x_\lambda = \sum_{w \in \mathfrak{S}_\lambda} q^{\ell(w)} T_w$ and $*$ is the anti-involution on $\mathcal{H}_{2f+1, n}$ fixing all generators T_j 's.

Theorem 3.1. [9, Theorem 3.20] *The Hecke algebra $\mathcal{H}_{2f+1, n}$ is free over \mathcal{Z} with basis*

$$\{x_{\mathfrak{s}\mathfrak{t}} \mid \mathfrak{s}, \mathfrak{t} \in \mathcal{T}^{std}(\lambda), \lambda \in \Lambda^+(n - 2f)\}.$$

It is a cellular basis in the sense of [8, Definition 1.1]. The required anti-involution is $$ as above.*

Suppose $(\lambda, \mu) \in \Lambda^+(n - 2f) \times \Lambda^+(n - 2f - 1)$. The partition μ is obtained from λ by removing a removable node, say $p = (k, \lambda_k)$ of λ if $\mu_j = \lambda_j, j \neq k$ and $\mu_k = \lambda_k - 1$. In this case, λ is obtained from μ by adding the addable node p of μ . We write either $\lambda = \mu \cup p$ or $\mu = \lambda \setminus p$. Let \mathcal{R}_λ be the set of all partitions obtained from λ by removing a removable node and \mathcal{A}_λ the set of all partitions obtained from λ by adding an addable node.

For any $\lambda \in \Lambda^+(n - 2f)$, let S^λ be the cell module of $\mathcal{H}_{2f+1, n}$ with respect to the Jucys-Murphy basis in Theorem 3.1. By [8, Definition 2.1] S^λ can be identified with the free \mathcal{Z} -module with basis $\{x_{\mathfrak{t}} \mid \mathfrak{t} \in \mathcal{T}^{std}(\lambda)\}$ where

$$x_{\mathfrak{t}} := x_{\mathfrak{t}^\lambda \mathfrak{t}} + \mathcal{H}_{2f+1, n}^{\triangleright \lambda}$$

and $\mathcal{H}_{2f+1, n}^{\triangleright \lambda}$ is the free \mathcal{Z} -submodule of $\mathcal{H}_{2f+1, n}$ spanned by $\{x_{\mathfrak{u}\mathfrak{s}} \mid \mathfrak{u}, \mathfrak{s} \in \mathcal{T}^{std}(\mu), \mu \triangleright \lambda\}$. It is also a two-sided ideal of $\mathcal{H}_{2f+1, n}$. Suppose $n > 2f$. Write $\mathcal{R}_\lambda = \{\mu^{(i)} \mid 1 \leq i \leq a\}$ for some positive integer a such that

$$\mu^{(1)} \triangleright \mu^{(2)} \triangleright \cdots \triangleright \mu^{(a)}.$$

For any standard λ -tableau \mathfrak{t} , let $\mathfrak{t} \downarrow_{n-1}$ be obtained from \mathfrak{t} by removing the entry n . Then $\mathfrak{t} \downarrow_{n-1} \in \mathcal{T}^{std}(\mu)$ for some $\mu \in \mathcal{R}_\lambda$. Let $S_i^\lambda = \mathcal{Z}\text{-span}\{x_{\mathfrak{t}} \mid \mathfrak{t} \downarrow_{n-1} \in \mathcal{T}^{std}(\mu^{(j)}), 1 \leq j \leq i\}$. Then S_i^λ is a right $\mathcal{H}_{2f+1, n-1}$ -module such that

$$S^\lambda = S_a^\lambda \supset S_{a-1}^\lambda \supset \cdots \supset S_1^\lambda \supset S_0^\lambda = 0.$$

Theorem 3.2. [9, Proposition 6.1] *As $\mathcal{H}_{2f+1, n-1}$ -modules, $S_i^\lambda / S_{i-1}^\lambda \cong S^{\mu^{(i)}}, 1 \leq i \leq a$, where $S^{\mu^{(i)}}$ is the cell module with respect to the Jucys-Murphy basis of $\mathcal{H}_{2f+1, n-1}$ in Theorem 3.1.*

Suppose $(\lambda, \mu) \in \Lambda(n - 2f) \times \Lambda(n - 2f)$. Recall that a λ -tableau of type μ is obtained from $[\lambda]$ by inserting integers $2f + i$ into $[\lambda]$ such that $2f + i$ appears μ_i times. A λ -tableau S of type μ is called row semi-standard if the entries in each row of S are non-decreasing from left to right. A row semi-standard tableau S is called semi-standard if λ is a partition and the entries in each column of S are strictly increasing downward. Let $\mathcal{T}^{ss}(\lambda, \mu)$ be the set of all semi-standard λ -tableaux of type μ . When $\mu = (1, 1, \dots, 1) \in \Lambda^+(n - 2f)$, $\mathcal{T}^{ss}(\lambda, \mu)$ is $\mathcal{T}^{std}(\lambda)$.

For any $\mathfrak{t} \in \mathcal{T}^{std}(\lambda)$, let $\mu(\mathfrak{t})$ be the λ -tableau obtained from \mathfrak{t} by replacing each entry i in \mathfrak{t} with $2f+k$ if i appears in the k -th row of \mathfrak{t}^μ . By [9, Example 4.2(ii)], $\mu(\mathfrak{t})$ is a row semi-standard λ -tableau of type μ . Following [9, Chapter 4, §2], write

$$x_{S,\mathfrak{t}} = \sum_{\mathfrak{s} \in \mathcal{T}^{std}(\lambda), \mu(\mathfrak{s})=S} q^{\ell(d(\mathfrak{s}))} x_{\mathfrak{s},\mathfrak{t}} \quad (3.1)$$

where $(S, \mathfrak{t}) \in \mathcal{T}^{ss}(\lambda, \mu) \times \mathcal{T}^{std}(\lambda)$.

Lemma 3.3. [9, Corollary 4.10] *For any $\mu \in \Lambda(n-2f)$, the right $\mathcal{H}_{2f+1,n}$ -module $x_\mu \mathcal{H}_{2f+1,n}$ has basis $\{x_{S,\mathfrak{t}} \mid S \in \mathcal{T}^{ss}(\lambda, \mu), \mathfrak{t} \in \mathcal{T}^{std}(\lambda), \lambda \in \Lambda^+(n-2f)\}$. Arrange all such semi-standard tableaux as S_1, S_2, \dots, S_k such that $S_i \in \mathcal{T}^{ss}(\lambda^{(i)}, \mu)$ and $i > j$ whenever $\lambda^{(i)} \triangleright \lambda^{(j)}$. Let M_i be the \mathcal{Z} -submodule of $x_\mu \mathcal{H}_{2f+1,n}$ spanned by $\{x_{S_j,\mathfrak{t}} \mid \mathfrak{t} \in \mathcal{T}^{std}(\lambda^{(j)}), i \leq j\}$. Then $x_\mu \mathcal{H}_{2f+1,n}$ has a filtration*

$$x_\mu \mathcal{H}_{2f+1,n} = M_1 \supset M_2 \supset \dots \supset M_k \supset M_{k+1} = 0$$

such that $M_i/M_{i+1} \cong S^{\lambda^{(i)}}$. The required isomorphism sends $x_{S_i,\mathfrak{t}} + M_{i+1}$ to $x_{\lambda^{(i)}} T_{d(\mathfrak{t})} + \mathcal{H}_{2f+1,n}^{\triangleright \lambda^{(i)}}$.

3.2. A cellular basis of the q -Brauer algebra. In this subsection, we assume that $\mathcal{B}_n(q, z)$ is defined over \mathbb{k} . Let

$$\Lambda_n = \left\{ (f, \lambda) \mid \lambda \in \Lambda^+(n-2f), 0 \leq f \leq \lfloor \frac{n}{2} \rfloor \right\}.$$

There is a partial order \leq on Λ_n such that $(\ell, \mu) \leq (f, \lambda)$ if $\ell < f$ or $\ell = f$ and $\mu \leq \lambda$. Write $(\ell, \mu) \triangleleft (f, \lambda)$ if $(\ell, \mu) \leq (f, \lambda)$ and $(\ell, \mu) \neq (f, \lambda)$. Later on, we identify each (ℓ, μ) in Λ_n with μ . So $\mu \leq \lambda$ if $|\mu| > |\lambda|$ or $|\mu| = |\lambda|$ and $\mu \leq \lambda$.

Definition 3.4. [19, Lemma 4.3], [24, (1.5)] Suppose $0 \leq f \leq \lfloor \frac{n}{2} \rfloor$. Define

- (1) $\mathcal{D}_{f,n} = \{s_{2f,i_f} s_{2f-1,j_f} \cdots s_{2,i_1} s_{1,j_1} \mid i_1 < \dots < i_f, 2k-1 \leq j_k < i_k \leq n, \text{ for } 1 \leq k \leq f\}$,
- (2) $B_{f,n} = \{t_{n-1} t_{n-2} \cdots t_{2f} t_{2f-2} \cdots t_2 \mid t_j = 1 \text{ or } t_j = s_{i_j, j+1}, 1 \leq i_j \leq j+1\}$.

When $f = 0$, the RHS of the equality in Definition 3.4(1) is the empty set and we set $\mathcal{D}_{0,n} = \{1\}$.

Lemma 3.5. *Suppose $d^{-1} \in \mathcal{D}_{f,n-1}$ and $h \in \{s_{n-1}, s_{n-2,n}\}$. We have $dh = s_{1,n} d_1$ for some integer $1 \leq l \leq n$ and $d_1^{-1} \in \mathcal{D}_{f,n-2}$.*

Proof. Suppose $2f-1 \leq j_f < i_f \leq n-1$ and $2f < k \leq n-1$. We have

$$s_{j_f, 2f-1} s_{i_f, 2f} s_{k,n} = \begin{cases} s_{k,n} s_{j_f, 2f-1} s_{i_f, 2f} & \text{if } i_f < k, \\ s_{k-1,n} s_{j_f, 2f-1} s_{i_f-1, 2f} & \text{if } i_f \geq k, j_f < k-1, \\ s_{k-2,n} s_{j_f-1, 2f-1} s_{i_f-1, 2f} & \text{if } i_f \geq k, j_f \geq k-1. \end{cases} \quad (3.2)$$

Now the result follows if we use (3.2) f times to rewrite dh successively. \square

Lemma 3.6. *For any $0 \leq f \leq \lfloor \frac{n}{2} \rfloor$, $\mathcal{D}_{f,n}^{-1} \subset B_{f,n}$, where $\mathcal{D}_{f,n}^{-1} = \{d \mid d^{-1} \in \mathcal{D}_{f,n}\}$.*

Proof. The result for $n = 2$ is trivial. In general, write $d = d_{i_f, j_f} d_{i_{f-1}, j_{f-1}} \cdots d_{i_1, j_1}$ for any $d \in \mathcal{D}_{f,n}$, where

$$d_{i_f, j_f} = s_{2f, i_f} s_{2f-1, j_f}.$$

Suppose $n = 2f$. Thanks to Definition 3.4(1), $(i_f, j_f) = (n, n-1)$ and

$$d^{-1} \in \mathcal{D}_{f-1, n-1}^{-1} \subset B_{f-1, n-1} \subset B_{f,n},$$

where the first inclusion follows from induction assumption on $n-1$. Suppose $n > 2f$. If $i_f < n$ then $d^{-1} \in \mathcal{D}_{f, n-1}^{-1}$ and hence $d^{-1} \in B_{f,n}$ by induction assumption on $n-1$. If $i_f = n$, then there is a $d_1 \in \mathcal{D}_{f-1, n-1}$ such that

$$d^{-1} = d_1 s_{j_f, 2f-1} s_{i_f, 2f} = \begin{cases} d_1 s_{n-1} s_{j_f, 2f-1} s_{n-1, 2f} & \text{if } j_f < n-1, \\ d_1 s_{n-2} s_{n-1} s_{n-2, 2f-1} s_{n-1, 2f} & \text{if } j_f = n-1. \end{cases} \quad (3.3)$$

We use Lemma 3.5 to rewrite both $d_1 s_{n-1}$ and $d_1 s_{n-2} s_{n-1}$ in (3.3). So there is a $d'_1 \in \mathcal{D}_{f, n-1}^{-1}$ such that $d^{-1} = t_{n-1} d'_1$. Now, the result follows from induction assumption on $n-1$. \square

Recall that a Brauer diagram is a diagram with $2n$ vertices arranged in two rows and n edges such that each vertex belongs to a unique edge. The vertices in the top (resp., bottom) row are labeled as $n+1, n+2, \dots, 2n$ (resp., $1, 2, \dots, n$) from left to right. Then each Brauer diagram d corresponds to a unique partition of $1, 2, \dots, 2n$ into n pairs $\{(i_k, j_k) \mid 1 \leq k \leq n, i_k < j_k\}$. We denote $\{(i_k, j_k) \mid 1 \leq k \leq n\}$ by $\text{conn}(d)$ and call it the connector of d . An edge is called a horizontal edge if two vertices of it are at the same row. Otherwise, it is called a vertical edge. Let d_f be the Brauer diagram such that

$$\text{conn}(d_f) = \{(2i-1, 2i), (n+2i-1, n+2i) \mid 1 \leq i \leq f\} \cup \{(j, n+j) \mid 2f+1 \leq j \leq n\}.$$

Consider a Brauer diagram on which there are exactly f horizontal edges $(n+2i-1, n+2i)$, $1 \leq i \leq f$, at the top row and there is no crossing between any two vertical edges. Let $D_{f,n}$ be the set of all such Brauer diagrams. The symmetric group \mathfrak{S}_n acts transitively on the right of $D_{f,n}$. In [22, (2.11)], Dung Tien Nguyen defined $\mathcal{B}_{0,n} = \{1\}$ and

$$\mathcal{B}_{f,n} = \{w^{-1} \in B_{f,n} \mid d = d_f w \in D_{f,n}\} \quad (3.4)$$

for $f > 0$. There is a restriction on the length of d in [22, (2.11)]. However, this restriction is redundant since d automatically satisfies this condition if $w^{-1} \in B_{f,n}$ [24, Lemma 2.1].

Proposition 3.7. *For any $0 \leq f \leq \lfloor \frac{n}{2} \rfloor$, $\mathcal{D}_{f,n} = \mathcal{B}_{f,n}$.*

Proof. There is a well-known result for Brauer diagrams which says that $d_f w \in D_{f,n}$ if $w \in \mathcal{D}_{f,n}$. Thanks to Lemma 3.6 and (3.4), $\mathcal{D}_{f,n} \subseteq \mathcal{B}_{f,n}$. We have $\mathcal{D}_{f,n} = \mathcal{B}_{f,n}$ since the cardinalities of $\mathcal{D}_{f,n}$ and $\mathcal{B}_{f,n}$ are $\frac{n!}{2^f(n-2f)!}$ (see [19, Lemma 4.3] and [21, Remark 3.18]). \square

Thanks to Proposition 3.7, we can use $\mathcal{D}_{f,n}$ to replace $\mathcal{B}_{f,n}$ when we state the cellular basis of \mathcal{B}_n in [22, Theorem 3.2]. This has an advantage when we do some explicit computation later on.

Theorem 3.8. [22, Theorem 3.2] *The q -Brauer algebra \mathcal{B}_n is free over \mathbb{k} with cellular basis*

$$S = \{C_{(w,s),(v,t)}^\lambda \mid (w,s), (v,t) \in I(\lambda), (f,\lambda) \in \Lambda_n\}$$

in the sense of [8, Definition 1.1], where $I(\lambda) := \mathcal{D}_{f,n} \times \mathcal{T}^{std}(\lambda)$ and $C_{(w,s),(v,t)}^\lambda = \sigma(T_w)E^f x_{st} T_v$. The required anti-involution is σ in Lemma 2.3.

Lemma 3.9. [22, Corollary 3.1] *Suppose $0 \leq f \leq \lfloor \frac{n}{2} \rfloor$. Then $E^f \mathcal{B}_n + \mathcal{B}_n^{f+1}$ is a left $\mathcal{H}_{2f+1,n}$ -module spanned by $\{E^f T_d + \mathcal{B}_n^{f+1} \mid d \in \mathcal{D}_{f,n}\}$, where \mathcal{B}_n^{f+1} is the two-sided ideal of \mathcal{B}_n generated by E^{f+1} .*

Theorem 3.10. *Let $\phi : \mathcal{B}_{n-2} \rightarrow \tilde{E}_1 \mathcal{B}_n \tilde{E}_1$ be the algebra homomorphism in Proposition 2.7. Then ϕ is an algebra isomorphism.*

Proof. Thanks to Theorem 3.8, \mathcal{B}_{n-2} has basis

$$\{\sigma(T_{d_1})E^f T_w T_{d_2} \mid d_1, d_2 \in \mathcal{D}_{f,n-2}, w \in \mathcal{H}_{2f+1,n-2}, 0 \leq f \leq \lfloor \frac{n-2}{2} \rfloor\}.$$

By Proposition 2.7(2), $\phi(\sigma(T_{d_1})E^f T_w T_{d_2}) = qz^{-1}T_1^{-1}T_2\sigma(T_{e_1})E^{f+1}T_{w'}T_{e_2}$ where e_1, e_2, w' are obtained from d_1, d_2, w by replacing each factor s_i in d_1, d_2, w by s_{i+2} . This proves that $\phi(\sigma(T_{d_1})E^f T_w T_{d_2})$ is a basis element of \mathcal{B}_n , and hence ϕ is a monomorphism. On the other hand, there are some scalars $a, b \in \mathbb{k}$ and $w_1, w_2 \in \mathfrak{S}_{3,n}$ such that

$$E_1 T_{2,i_1} T_{1,j_1} E_1 = a E_1 E_3 T_{w_1} + b E_1 T_{w_2} \quad (3.5)$$

and $a = 0$ unless $i_1 \geq 4$ and $j_1 \geq 3$. In general, since $\mathcal{D}_{1,n}$ is a right coset representatives of $\mathfrak{S}_2 \times \mathfrak{S}_{3,n}$, by Definition 2.1(2)(3) and (3.5), $E_1 \mathcal{H}_n E_1 = E_1 E_3 \mathcal{H}_{3,n} + E_1 \mathcal{H}_{3,n}$. By Theorem 3.8 and (3.5), $E_1 \mathcal{B}_n E_1$ is generated by $E_1 E_3$, $E_1 T_j$, $3 \leq j \leq n-1$. Consequently, $\tilde{E}_1 \mathcal{B}_n \tilde{E}_1$ is generated by $\tilde{E}_1 E_3$, $\tilde{E}_1 T_j$, $3 \leq j \leq n-1$ and hence ϕ is an epimorphism. \square

Corollary 3.11. *Suppose $n \geq 2$.*

- (1) *There is an algebra isomorphism $\psi : \mathcal{B}_{n-2} \rightarrow E_1 \mathcal{B}_n E_1$ sending $1, E_1$ and T_i to $\delta^{-1} E_1, \delta^{-1} E_1 E_3$ and $\delta^{-1} E_1 T_{i+2}$ for all $1 \leq i \leq n-3$.*
- (2) *The right \mathcal{B}_n -module $E^f \mathcal{B}_n$ is a left $E^f \mathcal{B}_n E^f$ -module generated by $E^f T_d$, $d \in \mathcal{D}_{f,n}$. Further, $E^f \mathcal{B}_n E^f$ is generated by E^{f+1} and $E^f T_{2f+i}$, $1 \leq i \leq n-2f-1$.*

Proof. (1) follows from arguments in the proof of Theorem 3.10 and (2) follows from (3.5) and Lemma 3.9 successively for $\mathcal{B}_n, \mathcal{B}_{n-2}$ etc. \square

3.3. The classical branching rule. In this subsection, we go on assuming that \mathcal{B}_n is defined over \mathbb{k} . For any $(f, \lambda) \in \Lambda_n$, let $C(f, \lambda)$ be the (right) cell module of \mathcal{B}_n with respect to the cellular basis in Theorem 3.8. Up to an isomorphism, $C(f, \lambda)$ can be considered as the free \mathbb{k} -module with basis $\{E^f x_\lambda T_{d(\mathfrak{t})} T_v + \mathcal{B}_n^{\triangleright \lambda} \mid (v, \mathfrak{t}) \in I(\lambda)\}$, where

$$\mathcal{B}_n^{\triangleright \lambda} = \mathbb{k}\text{-span}\{C_{(w, \mathfrak{s}), (v, \mathfrak{t})}^\mu \mid (w, \mathfrak{s}), (v, \mathfrak{t}) \in I(\mu), (\ell, \mu) \in \Lambda_n, \mu \triangleright \lambda\}.$$

Definition 3.12. For any $(f, \lambda) \in \Lambda_n$, let $\mathcal{RA}(\lambda)$ be the set of all partitions $\mu \in \mathcal{A}_\lambda \cup \mathcal{R}_\lambda$ such that $(l, \mu) \in \Lambda_{n-1}$ for some $l \in \mathbb{N}$. Let

$$y_\mu^\lambda = \begin{cases} E^f x_\lambda T_{a_k, n} & \text{if } \mu = \lambda \setminus (k, \lambda_k), \\ E_{2f-1} T_{n, 2f}^{-1} T_{b_k, 2f-1}^{-1} E^{f-1} x_\mu & \text{if } \mu = \lambda \cup (k, \lambda_k + 1), \end{cases}$$

where $a_k = 2f + \sum_{i=1}^k \lambda_i$ and $b_k = 2f - 1 + \sum_{i=1}^k \lambda_i$.

Thanks to the defining relations for \mathcal{B}_n , we can rewrite y_μ^λ as follows:

$$y_\mu^\lambda = \begin{cases} \sum_{i=a_{k-1}+1}^{a_k} q^{a_k-i} T_{i, n} E^f x_\mu & \text{if } \mu = \lambda \setminus (k, \lambda_k), \\ E^f x_\lambda T_{n, 2f}^{-1} T_{b_k, 2f-1}^{-1} \sum_{i=b_{k-1}+1}^{b_k} q^{b_k-i} T_{b_k, i} & \text{if } \mu = \lambda \cup (k, \lambda_k + 1). \end{cases} \quad (3.6)$$

For any $(f, \lambda) \in \Lambda_n$, there is a pair $(a, m) \in \mathbb{N} \times \mathbb{N} \setminus \{0\}$ such that

$$\mathcal{RA}(\lambda) = \{\mu^{(1)}, \mu^{(2)}, \dots, \mu^{(a)}\} \cup \{\mu^{(a+1)}, \mu^{(a+2)}, \dots, \mu^{(m)}\}$$

and $(f, \mu^{(i)}) \in \Lambda_{n-1}$, $1 \leq i \leq a$ and $(f-1, \mu^{(i)}) \in \Lambda_{n-1}$, $a+1 \leq i \leq m$. Obviously $\mu^{(a+1)}$ occurs only if $f > 0$. We can arrange them so that

$$\mu^{(1)} \triangleright \mu^{(2)} \triangleright \dots \triangleright \mu^{(a)} \triangleright \mu^{(a+1)} \triangleright \mu^{(a+2)} \triangleright \dots \triangleright \mu^{(m)} \quad (3.7)$$

with respect the partial order \triangleright on Λ_{n-1} . Note that $\mu^{(m)} = \lambda \cup (k+1, 1)$ if $k = \max\{j \mid \lambda_j \neq 0\}$. So

$$y_{\mu^{(m)}}^\lambda = E^f x_\lambda T_{n, 2f}^{-1} T_{n-1, 2f-1}^{-1}. \quad (3.8)$$

Definition 3.13. For any $1 \leq k \leq m$, let $N_k = \sum_{j=1}^k y_{\mu^{(j)}}^\lambda \mathcal{B}_{n-1} + \mathcal{B}_n^{\triangleright \lambda}$.

It follows from Definition 3.13 that N_k , $1 \leq k \leq m$ are \mathcal{B}_{n-1} -submodules of $C(f, \lambda)$.

Lemma 3.14. Suppose $1 \leq k \leq a$.

(1) The \mathcal{B}_{n-1} -submodule N_k of $C(f, \lambda)$ is spanned by

$$\{y_{\mu^{(j)}}^\lambda T_{d(\mathfrak{u})} T_w + \mathcal{B}_n^{\triangleright \lambda} \mid \mathfrak{u} \in \mathcal{T}^{std}(\mu^{(j)}), w \in \mathcal{D}_{f, n-1}, 1 \leq j \leq k\}.$$

(2) As \mathcal{B}_{n-1} -modules, $N_k/N_{k-1} \cong C(f, \mu^{(k)})$,

(3) The submodule N_k has basis

$$\{E^f x_\lambda T_{d(\mathfrak{s})} T_v + \mathcal{B}_n^{\triangleright \lambda} \mid \mathfrak{s} \in \mathcal{T}^{std}(\lambda), \text{shape}(\mathfrak{s} \downarrow_{n-1}) \supseteq \mu^{(k)}, v \in \mathcal{D}_{f, n-1}\}.$$

Proof. Suppose $b \in \mathcal{B}_{n-1}$ and $\mu^{(k)} = \lambda \setminus (h, \lambda_h)$. Let $a_k = 2f + \sum_{i=1}^h \lambda_i$. Thanks to Definition 3.13, Lemma 3.9 and Theorem 3.2,

$$y_{\mu^{(k)}}^\lambda b = x_\lambda T_{a_k, n} \sum_{v \in \mathfrak{S}_{3, n}, w \in \mathcal{D}_{f, n-1}} a_{v, w} T_v E^f T_w \equiv \sum_{\mathfrak{u} \in \mathcal{T}^{std}(\mu^{(k)}), w \in \mathcal{D}_{f, n-1}} b_{\mathfrak{u}, w} y_{\mu^{(k)}}^\lambda T_{d(\mathfrak{u})} T_w \pmod{N_{k-1}}$$

where $a_{v, w}$ and $b_{\mathfrak{u}, w}$ are some scalars. This proves (1). Suppose $(\mathfrak{u}, w) \in \mathcal{T}^{std}(\mu^{(k)}) \times \mathcal{D}_{f, n-1}$. Then

$$y_{\mu^{(k)}}^\lambda T_{d(\mathfrak{u})} T_w + \mathcal{B}_n^{\triangleright \lambda} = E^f x_\lambda T_{d(\mathfrak{t})} T_w + \mathcal{B}_n^{\triangleright \lambda}$$

with $\mathfrak{t} \downarrow_{n-1} = \mathfrak{u}$. Since the RHS of the above equality is a basis element of $C(f, \lambda)$, by (1), N_k/N_{k-1} is free over \mathbb{k} with basis M_k , where

$$M_k = \{y_{\mu^{(k)}}^\lambda T_{d(\mathfrak{u})} T_w + N_{k-1} \mid \mathfrak{u} \in \mathcal{T}^{std}(\mu^{(k)}), w \in \mathcal{D}_{f, n-1}\}. \quad (3.9)$$

There is a well-defined \mathbb{k} -linear isomorphism $\phi : N_k/N_{k-1} \rightarrow C(f, \mu^{(k)})$ such that

$$\phi(y_{\mu^{(k)}}^\lambda T_{d(\mathfrak{u})} T_w + N_{k-1}) = E^f x_{\mu^{(k)}} T_{d(\mathfrak{u})} T_w + \mathcal{B}_{n-1}^{\triangleright \mu^{(k)}}, \quad (3.10)$$

for all $(\mathfrak{u}, w) \in \mathcal{T}^{std}(\mu^{(k)}) \times \mathcal{D}_{f, n-1}$. It follows immediately from Lemma 3.9 and Theorem 3.2 that ϕ is a right \mathcal{B}_{n-1} -homomorphism and hence ϕ is a \mathcal{B}_{n-1} -isomorphism. This completes the proof of (2). Finally, (3) immediately follows from (2), (3.10) and induction assumption on $k-1$. \square

From here to the end of this section, we assume $(f, \lambda) \in \Lambda_n$ and $f > 0$. We are going to deal with N_k/N_{k-1} , $a+1 \leq k \leq m$.

Lemma 3.15. [5] *Let $(f-1, \mu) \in \Lambda_{n-1}$ such that $\mu \triangleright \mu^{(m)}$.*

- (1) *Suppose $\mu \notin \{\mu^{(a+1)}, \dots, \mu^{(m)}\}$. If $(\mathfrak{s}, \mu^{(m)}(\mathfrak{s})) \in \mathcal{T}^{std}(\mu) \times \mathcal{T}(\mu, \mu^{(m)})$, then $\text{shape}(\mathfrak{s} \downarrow_{n-2}) \triangleright \lambda$.*
- (2) *Suppose $\mu = \mu^{(j)} = \lambda \cup (k, \lambda_k + 1)$ for some $a+1 \leq j \leq m$. Let $\mathfrak{s}_j = t^{\mu^{(j)}} s_{b_j, n-1}$, where $b_j = 2f-1 + \sum_{i=1}^k \lambda_i$. Then \mathfrak{s}_j is the unique standard $\mu^{(j)}$ -tableau \mathfrak{s} satisfying $\mu^{(m)}(\mathfrak{s}) \in \mathcal{T}^{ss}(\mu^{(j)}, \mu^{(m)})$. In this case, $\text{shape}(\mathfrak{s} \downarrow_{n-2}) = \lambda$.*

Proof. The results has already been given in the proof of [5, Corollary 5.4, Lemma 5.5]. \square

Lemma 3.16. *Suppose $(\mathfrak{s}, \mu^{(m)}(\mathfrak{s})) \in \mathcal{T}^{std}(\mu) \times \mathcal{T}^{ss}(\mu, \mu^{(m)})$ for some $(f-1, \mu) \in \Lambda_{n-1}$. If $\text{shape}(\mathfrak{s} \downarrow_{n-2}) \triangleright \lambda$, then $E^f T_{n,2f}^{-1} T_{n-1,2f-1}^{-1} \sigma(T_d(\mathfrak{s})) x_\mu \in \mathcal{B}_n^{\triangleright \lambda}$.*

Proof. The arguments in the proof of [5, Lemma 5.3] depends only on the braid relations for the Hecke algebras. Therefore, one can imitate arguments there to verify our current result. We leave details to the reader. \square

Lemma 3.17. *As \mathcal{B}_{n-1} -modules, $C(f, \lambda) = N_m$.*

Proof. Recall that the cell module $C(f, \lambda)$ has basis $\{E^f x_\lambda T_{d(t)} T_v + \mathcal{B}_n^{\triangleright \lambda} \mid (v, t) \in I(\lambda)\}$. If $v \in \mathcal{D}_{f, n-1}$, then

$$E^f x_\lambda T_{d(t)} T_v + \mathcal{B}_n^{\triangleright \lambda} \in E^f x_\lambda T_{d(t)} \mathcal{B}_{n-1} + \mathcal{B}_n^{\triangleright \lambda} \subset N_a. \quad (3.11)$$

The last inclusion follows from Lemma 3.14(3). Otherwise $T_v = T_{2f, n} b$ for some $b \in \mathcal{H}_{n-1}$. So,

$$E^f x_\lambda T_{d(t)} T_v + \mathcal{B}_n^{\triangleright \lambda} \in E^f x_\lambda T_{2f, n} \mathcal{B}_n + \mathcal{B}_n^{\triangleright \lambda} \subset \sum_{j=2f+1}^n E^f x_\lambda T_{j, n} \mathcal{B}_{n-1} + E^f x_\lambda T_{n, 2f}^{-1} \mathcal{B}_{n-1} + \mathcal{B}_n^{\triangleright \lambda}. \quad (3.12)$$

By Definitions 3.12–3.13, (3.8) and Lemma 3.14(3),

$$E^f x_\lambda T_{n, 2f}^{-1} \mathcal{B}_{n-1} + \mathcal{B}_n^{\triangleright \lambda} \in N_m \quad \text{and} \quad \sum_{j=2f+1}^n E^f x_\lambda T_{j, n} \mathcal{B}_{n-1} \in N_a \subset N_m.$$

This implies $C(f, \lambda) \subseteq N_m$ and hence $C(f, \lambda) = N_m$ as required. \square

Lemma 3.18. *If $b \in E^{f-1} \mathcal{B}_{n-1} \cap \mathcal{B}_{n-1}^f$, then there are $w \in \mathcal{D}_{f, n-1}$ and $h_w \in \mathcal{H}_{2f+1, n}$ such that*

$$E_{2f-1} T_{n, 2f}^{-1} T_{n-1, 2f-1}^{-1} b \equiv \sum_w h_w E^f T_w \pmod{\mathcal{B}_n^{f+1}}. \quad (3.13)$$

Proof. By Theorem 3.8, Lemma 3.9 and Corollary 3.11(2), any $b \in E^{f-1} \mathcal{B}_{n-1} \cap \mathcal{B}_{n-1}^f$ can be written as a linear combination of elements in S up to an element in \mathcal{B}_n^{f+1} , where

$$S = \{T_{j, 2f-1} T_{i, 2f} E^f T_u T_w \mid T_u \in \mathcal{H}_{2f+1, n-1}, w \in \mathcal{D}_{f, n-1}, 2f-1 \leq j < i \leq n-1\}. \quad (3.14)$$

So it is enough to assume $b = T_{j, 2f-1} T_{i, 2f} E^f T_u T_w$ when we verify (3.13).

Suppose $i = n-1$. Then $j < n-1$. Let $x = E_{2f-1} T_{n, 2f}^{-1} T_{n-1, 2f-1}^{-1} T_{j, 2f-1} T_{n-1, 2f} E^f$. By Lemma 2.6(2)(5) and Lemma 2.5(6),

$$\begin{aligned} x &= T_{j+2, 2f+1} E_{2f-1} T_{n-1, 2f}^{-1} T_{n-1, 2f-1}^{-1} T_{n-1, 2f} E^f \\ &= T_{j+2, 2f+1} E_{2f-1} T_{n, 2f}^{-1} T_{2f-1}^{-1} E^f \\ &= q^{-1} T_{j+2, 2f+1} E_{2f-1} T_{n, 2f}^{-1} E^f \\ &= q^{-1} T_{j+2, 2f+1} E_{2f-1} T_{2f}^{-1} E^f T_{n, 2f+1}^{-1} \\ &= z^{-1} q^{-1} E^f T_{j+2, 2f+1} T_{n, 2f+1}^{-1}. \end{aligned}$$

and (3.13) follows. Suppose $i < n-1$. By (2.2) and Lemma 2.6(2)

$$E_{2f-1} T_{n, 2f}^{-1} T_{n-1, 2f-1}^{-1} T_{j, 2f-1} T_{i, 2f} E^f = T_{j+2, 2f+1} T_{i+2, 2f+2} T_{n, 2f+2}^{-1} T_{n-1, 2f+1}^{-1}, \quad (3.15)$$

where $y = E_{2f-1} T_{2f}^{-1} T_{2f+1}^{-1} T_{2f-1}^{-1} T_{2f} E^f$. Thanks to (2.2), $T_i^{-1} = T_i - a$ for any $1 \leq i \leq n-1$, where $a = q - q^{-1}$. We use it to rewrite y as follows:

$$y = E_{2f-1} T_{2f} T_{2f+1} T_{2f-1}^{-1} T_{2f}^{-1} E^f - (aq^{-1} + aT_{2f+1}) E_{2f-1} T_{2f-1}^{-1} T_{2f}^{-1} E^f. \quad (3.16)$$

Since $E^{f+1} = E_1 T_{2,2f+2} T_{2f+1,1}^{-1} E^f$, the first term in the RHS of (3.16) is equal to

$$T_{2f-1,1} T_{2,2f}^{-1} E_1 T_{2,2f+2} T_{2f+1,1}^{-1} E^f = T_{2f-1,1} T_{2,2f}^{-1} E^{f+1} \in \mathcal{B}_n^{f+1}. \quad (3.17)$$

Using Lemma 2.6(8) to rewrite the second term in the RHS of (3.16), we see that (3.13) follows immediately from (3.15)–(3.17). \square

Lemma 3.19. *As \mathcal{B}_{n-1} -modules, $N_k/N_{k-1} \cong C(f-1, \mu^{(k)})$ for any $a+1 \leq k \leq m$.*

Proof. Fix k , $a+1 \leq k \leq m$. We want to compute $y_{\mu^{(k)}}^\lambda h$ for any $h \in \mathcal{B}_{n-1}$. Thanks to Theorem 3.8 and Lemma 3.9, $E^{f-1}h + \mathcal{B}_n^{f+1}$ can be written as a linear combination of elements $\bar{x} := x + \mathcal{B}_n^{f+1}$, where $x \in \{h_d E^{f-1} T_d, b\}$, and $b \in E^{f-1} \mathcal{B}_{n-1} \cap \mathcal{B}_{n-1}^f$, $d \in \mathcal{D}_{f-1, n-1}$ and $h_d \in \mathcal{H}_{2f-1, n-1}$. So

$$\begin{aligned} y_{\mu^{(k)}}^\lambda h &\equiv \sum_{d \in \mathcal{D}_{f-1, n-1}} y_{\mu^{(k)}}^\lambda h_d T_d + E_{2f-1} T_{n,2f}^{-1} T_{b_k, 2f-1}^{-1} x_{\mu^{(k)}} b \pmod{\mathcal{B}_n^{f+1}} \\ &\equiv \sum_{d \in \mathcal{D}_{f-1, n-1}} E^f T_{n,2f}^{-1} T_{n-1,2f-1}^{-1} x_{\mathfrak{s}_k, \mathfrak{t}^{(k)}} h_d T_d \pmod{N_a}, \end{aligned} \quad (3.18)$$

where \mathfrak{s}_k is given in Lemma 3.15(2). The second equivalence follows from (3.13) and Lemma 3.14(3). Further, $x_{\mathfrak{s}_k, \mathfrak{u}} \in x_{\mu^{(m)}} \mathcal{H}_{2f-1, n-1}$ for any $\mathfrak{u} \in \mathcal{T}^{std}(\mu^{(k)})$. Applying Lemma 3.3 on $x_{\mathfrak{s}_k, \mathfrak{t}^{(k)}} h_d$, we have

$$y_{\mu^{(k)}}^\lambda h \equiv \sum_{d \in \mathcal{D}_{f-1, n-1}} E^f T_{n,2f}^{-1} T_{n-1,2f-1}^{-1} \sum_{(\mathfrak{s}, S) \in \mathcal{T}^{std}(\mu) \times \mathcal{T}^{ss}(\mu, \mu^{(m)}), \mu \triangleright \mu^{(k)}} a_{S, \mathfrak{s}} x_{S, \mathfrak{s}} T_d \pmod{N_a}.$$

By Lemmas 3.15(1) and 3.16, the terms with respect to μ in the RHS of the above equality are in $\mathcal{B}_n^{\triangleright \lambda}$ if $\mu \notin \{\mu^{(a+1)}, \mu^{(a+2)}, \dots, \mu^{(k)}\}$. In the remaining case $S \in \mathcal{T}^{ss}(\mu^{(j)}, \mu^{(m)})$ for some $a+1 \leq j \leq m$. Thanks to Lemma 3.15(2), $\mu^{-1}(S)$ contains a unique element, say \mathfrak{s}_j in $\mathcal{T}^{std}(\mu^{(j)})$. So $x_{S, \mathfrak{s}} = q^{-\ell(d(\mathfrak{s}_j))} x_{\mathfrak{s}_j, \mathfrak{s}}$ and $d(\mathfrak{s}_j) = s_{b_j, n}$, where b_j is given in Lemma 3.15(2). Since

$$E^f T_{n,2f}^{-1} T_{n-1,2f-1}^{-1} x_{\mathfrak{s}_j, \mathfrak{s}} = E^f T_{n,2f}^{-1} T_{n-1,2f-1}^{-1} T_{b_j, n-1} x_{\mathfrak{s}} = y_{\mu^{(j)}}^\lambda T_{d(\mathfrak{s})},$$

N_k/N_{k-1} is the \mathbb{k} -module spanned by

$$M_k = \{y_{\mu^{(k)}}^\lambda T_{d(\mathfrak{s})} T_d + N_{k-1} \mid \mathfrak{s} \in \mathcal{T}^{std}(\mu^{(k)}), d \in \mathcal{D}_{f-1, n-1}\}. \quad (3.19)$$

By Lemma 3.19, $C(f, \lambda)$ can be spanned by a set whose cardinality is $\sum_{k=1}^m |M_k|$. Thanks to [25, Theorem 2.3], the rank of cell module for Birman-Murakami-Wenzl algebra with respect to (f, λ) is equal to $\sum_{k=1}^m |M_k|$. Since rank $C(f, \lambda)$ is equal to that for Birman-Murakami-Wenzl algebra, we have rank $C(f, \lambda) = \sum_{k=1}^m |M_k|$ and hence M_k is linear independent for any $1 \leq k \leq m$. Consequently, M_k is a basis of N_k/N_{k-1} for all $1 \leq k \leq m$. We have a well-defined \mathbb{k} -linear isomorphism $\phi: N_k/N_{k-1} \rightarrow C(f-1, \mu^{(k)})$ for any $a+1 \leq k \leq m$ such that

$$\phi(y_{\mu^{(k)}}^\lambda T_{d(\mathfrak{s})} T_d + N_{k-1}) = E^f x_{\mu^{(k)}} T_{d(\mathfrak{s})} T_d + \mathcal{B}_{n-1}^{\triangleright \mu^{(k)}} \quad (3.20)$$

for any $\mathfrak{s} \in \mathcal{T}^{std}(\mu^{(k)})$ and $d \in \mathcal{D}_{f-1, n-1}$. By arguments similar to those above (more explicitly, using Theorem 3.8, Lemmas 3.3 and 3.9) we see that ϕ is a right \mathcal{B}_{n-1} -homomorphism. \square

Theorem 3.20. *Suppose $(f, \lambda) \in \Lambda_n$. Then there is a filtration $0 = N_0 \subset N_1 \subset \dots \subset N_m = C(f, \lambda)$ of \mathcal{B}_{n-1} -modules such that*

$$N_k/N_{k-1} \cong \begin{cases} C(f, \mu^{(k)}) & \text{if } 1 \leq k \leq a, \\ C(f-1, \mu^{(k)}) & \text{if } a+1 \leq k \leq m. \end{cases}$$

Proof. The result follows immediately from Lemmas 3.14 and 3.19. \square

4. THE JUCYS-MURPHY ELEMENTS AND JUCYS-MURPHY BASIS

In this section, we construct the Jucys-Murphy elements and the Jucys-Murphy basis of \mathcal{B}_n . Unless otherwise stated, we assume that \mathcal{B}_n is defined over \mathbb{k} .

4.1. **Jucys-Murphy elements.** For any $1 \leq i \leq n$, define

$$L_i = \begin{cases} 0 & \text{if } i = 1, \\ \sum_{j=1}^{i-1} (j, i) - q^2 z^{-1} \sum_{j=1}^{i-1} E_{j,i}, & \text{if } 2 \leq i \leq n, \end{cases} \quad (4.1)$$

where

$$(j, i) = T_{j,i-1} T_{i-1} T_{i-1,j} \quad \text{and} \quad E_{j,i} = T_{1,j}^{-1} T_{i,2} E_1 T_{2,i} T_{j,1}^{-1} \quad (4.2)$$

for any $1 \leq j \leq i-1$. Note that $E_{i,i+1}$ is not the E_i in Definition 2.4 except the classical limit case.

Lemma 4.1. *If $n \geq 2$, then $L_n = T_{n-1} L_{n-1} T_{n-1} + T_{n-1} - q^2 z^{-1} E_{n-1,n}$.*

Proof. The result follows immediately from (4.1). \square

Lemma 4.2. *Suppose $n > 3$ and $x_m = E_{m-2,m} + E_{m-1,m}$ for $m \geq 3$. Then*

- (1) $E_1 E_{n-1,n} = E_{n-1,n} E_1$,
- (2) $T_{m-2} x_m = x_m T_{m-2}$,
- (3) $T_{n-2} E_{i,n} = E_{i,n} T_{n-2}$ if $1 \leq i \leq n-3$,
- (4) $T_j E_{n-1,n} = E_{n-1,n} T_j$ if $1 \leq j \leq n-3$.

Proof. Thanks to Definition 2.1(4)(5) and Lemma 2.5(3),

$$\begin{aligned} E_1 E_{n-1,n} &= T_{3,n-1}^{-1} T_{n,4} E_1 T_2^{-1} T_1^{-1} T_3 T_2 E_1 T_{2,n} T_{n-1,1}^{-1} \\ &= T_{3,n-1}^{-1} T_{n,4} E_1 T_2^{-1} T_1^{-1} T_3 T_2 E_1 T_{4,n} T_{n-1,3}^{-1} \\ &= E_{n-1,n} E_1. \end{aligned}$$

This proves (1). One can easily verify (2)-(4) by braid relations in (2.2) and Definition 2.4(4). \square

Lemma 4.3. *Suppose $n \geq 2$. For any $x \in \mathcal{B}_{n-1}$, $L_n x = x L_n$.*

Proof. We prove the result by induction on n . The result is trivial when $n = 2$. We have

$$L_3 = T_2 T_1 T_2 + T_2 - q^2 z^{-1} (E_{1,3} + E_{2,3}). \quad (4.3)$$

It is easy to verify $E_1 L_3 = L_3 E_1 = 0$. It is known that $T_2 + T_2 T_1 T_2$ is the corresponding Jucys-Murphy element of \mathcal{H}_3 (e.g., [9]) and hence commutes with T_1 . Then by Lemma 4.2(2) and (4.3), $L_3 T_1 = T_1 L_3$. This proves the result when $n = 3$. Suppose $n > 3$. It is enough to prove that $L_n x = x L_n$ for any $x \in \{E_1, T_1, \dots, T_{n-2}\}$. By induction assumption on $n-1$, Definition 2.1 and Lemma 4.2(1), E_1 commutes with both $T_{n-1} L_{n-1} T_{n-1} + T_{n-1}$ and $E_{n-1,n}$. By Lemma 4.1, $E_1 L_n = L_n E_1$. By induction assumption on $n-1$ and Lemma 4.2(4), T_1, T_2, \dots, T_{n-3} commute with both $T_{n-1} L_{n-1} T_{n-1} + T_{n-1}$ and $E_{n-1,n}$. Using Lemma 4.1 again yields $L_n x = x L_n$ for any $x \in \{T_1, T_2, \dots, T_{n-3}\}$. It is known that $T_{n-2} \sum_{i=1}^{n-1} (i, n) = \sum_{i=1}^{n-1} (i, n) T_{n-2}$ since $\sum_{i=1}^{n-1} (i, n)$ is the corresponding Jucys-Murphy element of \mathcal{H}_n . Then by Lemma 4.2(2)(3), $T_{n-2} L_n = L_n T_{n-2}$. \square

Lemma 4.4. *For any $1 \leq i \leq j \leq n$, $L_i L_j = L_j L_i$.*

Proof. Obviously, $L_i \in \mathcal{B}_{j-1}$ if $i < j$. Now, the result follows from Lemma 4.3. \square

Thanks to Lemma 4.4, the subalgebra L of \mathcal{B}_n generated by $\{L_1, L_2, \dots, L_n\}$ is a commutative subalgebra. By Remark 2.2, the Brauer algebra $B_n(a)$ can be considered as the classical limit of $\mathcal{B}_n(q, q^a)$. In this case, $\{L_i \mid 1 \leq i \leq m\}$ is the set of corresponding Jucys-Murphy elements in [5] and $\sum_{i=1}^n L_i$ is a central element. However, $\sum_{i=1}^n L_i$ may not be a central element in general. One can easily find a counterexample in \mathcal{B}_4 .

4.2. **A new basis of a cell module.** Suppose $(f, \lambda) \in \Lambda_n$. Write $\mu \rightarrow \lambda$ if $\mu \in \mathcal{RA}(\lambda)$. An up-down tableau \mathbf{t} of type λ is a sequence of partitions $\mathbf{t} = (\mathbf{t}_0, \mathbf{t}_1, \dots, \mathbf{t}_n)$ such that $\mathbf{t}_0 = \emptyset$, $\mathbf{t}_i \rightarrow \mathbf{t}_{i+1}$, $0 \leq i \leq n-1$ and $\mathbf{t}_n = \lambda$. Let $\mathcal{F}_n^{ud}(\lambda)$ be the set of all up-down tableaux of type λ . There is a partial order \preceq on $\mathcal{F}_n^{ud}(\lambda)$ such that $\mathbf{t} \prec \mathbf{s}$ if $\mathbf{t}_k \triangleleft \mathbf{s}_k$ for some k and $\mathbf{t}_j = \mathbf{s}_j$ for all $k < j \leq n$. In this case, we also write $\mathbf{t} \overset{k}{\prec} \mathbf{s}$.

Definition 4.5. Suppose $(f, \lambda) \in \Lambda_n$, $\mathbf{t} \in \mathcal{F}_n^{ud}(\lambda)$ and $\mathbf{t}_{n-1} = \mu$. Define $m_{\mathbf{t}} = E^f x_{\lambda} b_{\mathbf{t}}$, where $b_{\mathbf{t}} = b_{\mathbf{t}_n}$ which can be defined inductively as follows:

$$b_{\mathbf{t}_n} = \begin{cases} T_{a_k, n} b_{\mathbf{t}_{n-1}} & \text{if } \lambda = \mu \cup (k, \mu_k + 1), \\ T_{n, 2f}^{-1} \sum_{j=b_{k-1}+1}^{b_k} q^{b_k-j} T_{j, 2f-1}^{-1} b_{\mathbf{t}_{n-1}} & \text{if } \lambda = \mu \setminus (k, \mu_k), \end{cases}$$

and $a_k = 2f + \sum_{j=1}^k \lambda_j$ and $b_i = 2f - 1 + \sum_{j=1}^i \lambda_j$, for $i \in \{k-1, k\}$.

In fact one can easily check that

$$\mathbf{m}_{\mathbf{t}_n} = \begin{cases} \sum_{j=a_{k-1}+1}^{a_k} T_{j,n} \mathbf{m}_{\mathbf{t}_{n-1}} & \text{if } \lambda = \mu \cup (k, \mu_k + 1), \\ E_{2f-1} T_{n,2f}^{-1} T_{b_k,2f-1}^{-1} \mathbf{m}_{\mathbf{t}_{n-1}} & \text{if } \lambda = \mu \setminus (k, \mu_k), \end{cases} \quad (4.4)$$

and $\mathbf{m}_{\mathbf{t}_n} = y_\mu^\lambda b_{\mathbf{t}_{n-1}}$ in any case.

Theorem 4.6. *Suppose $(f, \lambda) \in \Lambda_n$.*

- (1) *The cell module $C(f, \lambda)$ has \mathbb{k} -basis $\{\mathbf{m}_{\mathbf{t}} + \mathcal{B}_n^{\triangleright\lambda} \mid \mathbf{t} \in \mathcal{F}_n^{ud}(\lambda)\}$.*
- (2) *For any $1 \leq j \leq m$, the \mathbb{k} -module M_j spanned by $\{\mathbf{m}_{\mathbf{t}} + \mathcal{B}_n^{\triangleright\lambda} \mid \mathbf{t} \in \mathcal{F}_n^{ud}(\lambda), \mathbf{t}_{n-1} \triangleright \mu^{(j)}\}$ is a \mathcal{B}_{n-1} -submodule of $C(f, \lambda)$, where $\mu^{(j)}$ is given in (3.7).*
- (3) *For any $1 \leq j \leq m$, $M_j = N_j$ where N_j 's are given in Definition 3.13.*
- (4) *As \mathcal{B}_{n-1} -modules, $M_j/M_{j-1} \cong C(\ell, \mu^{(j)})$, where $\ell = f$ if $1 \leq j \leq a$ and $\ell = f - 1$ if $a + 1 \leq j \leq m$. The required isomorphism sends $\mathbf{m}_{\mathbf{t}} + M_{j-1}$ to $\mathbf{m}_{\mathbf{t} \downarrow_{n-1}} + \mathcal{B}_{n-1}^{\triangleright\mu^{(j)}}$, where $\mathbf{t} \downarrow_{n-1} = (\mathbf{t}_0, \mathbf{t}_1, \dots, \mathbf{t}_{n-1}) \in \mathcal{F}_{n-1}^{ud}(\mu^{(j)})$.*
- (5) *The \mathbb{k} -algebra \mathcal{B}_n has cellular basis*

$$\{\mathbf{m}_{\mathbf{st}} \mid \mathbf{s}, \mathbf{t} \in \mathcal{F}_n^{ud}(\lambda), (f, \lambda) \in \Lambda_n\}$$

where $\mathbf{m}_{\mathbf{st}} = \sigma(b_{\mathbf{s}}) E^f x_\lambda b_{\mathbf{t}}$ and the required σ is given in Lemma 2.3.

Proof. Thanks to Lemmas 3.14, 3.19, the \mathbb{k} -linear map $\phi : C(\ell, \mu^{(j)}) \rightarrow N_j/N_{j-1}$ sending $E^\ell x_{\mu^{(j)}} T_{d(\mathbf{t})} T_d + \mathcal{B}_{n-1}^{\triangleright\mu^{(j)}}$ to $y_{\mu^{(j)}}^\lambda T_{d(\mathbf{t})} T_d + N_{j-1}$ is the required \mathcal{B}_{n-1} -isomorphism. In particular, $\phi(E^\ell x_{\mu^{(j)}} + \mathcal{B}_{n-1}^{\triangleright\mu^{(j)}}) = y_{\mu^{(j)}}^\lambda + N_{j-1}$. By induction assumption on $n - 1$, $C(\ell, \mu^{(j)})$ has basis

$$\{E^\ell x_{\mu^{(j)}} b_{\mathbf{u}} + \mathcal{B}_{n-1}^{\triangleright\mu^{(j)}} \mid \mathbf{u} \in \mathcal{F}_{n-1}^{ud}(\mu^{(j)})\}.$$

Note that $\phi^{-1}(\mathbf{m}_{\mathbf{u}} + \mathcal{B}_{n-1}^{\triangleright\mu^{(j)}}) = \mathbf{m}_{\mathbf{t}} + N_{j-1}$ where $\mathbf{m}_{\mathbf{t}} = y_{\mu^{(j)}}^\lambda b_{\mathbf{u}}$ and $\mathbf{t} \downarrow_{n-1} = \mathbf{u}$. Using the isomorphism ϕ and induction assumption on N_{j-1} , we see that N_j has basis $\{\mathbf{m}_{\mathbf{t}} + \mathcal{B}_n^{\triangleright\lambda} \mid \mathbf{t} \in \mathcal{F}_n^{ud}(\lambda), \mathbf{t}_{n-1} \triangleright \mu^{(j)}\}$. In particular, $M_j = N_j$ where M_j is defined in (2). This proves (1)–(4). Note that \mathcal{B}_n is a cellular algebra. Thanks to Lemma 2.5(2), E^f is fixed by the anti-involution σ . It is easy to verify $\sigma(x_\lambda) = x_\lambda$. By (1), we have the corresponding results for left cell modules. This implies (5) by using standard arguments on cellular algebras (see [16, Theorem 2.7] for Brauer algebras). \square

The basis in Theorem 4.6(1) (resp., (5)) is called the *Jucys-Murphy basis* of $C(f, \lambda)$ (resp., \mathcal{B}_n).

Lemma 4.7. *Suppose $0 \leq f \leq \lfloor \frac{n}{2} \rfloor$.*

- (1) *$E^f E_{i,n} \in \mathcal{B}_n^{f+1}$ if $2f + 1 \leq i \leq n - 1$.*
- (2) *$E^f((2j - 1, n) + (2j, n) - q^2 z^{-1} E_{2j-1,n} - q^2 z^{-1} E_{2j,n}) = 0$ if $1 \leq j \leq f$.*

Proof. Since $i \geq 2f + 1$, we have

$$\begin{aligned} E^f E_{i,n} &= E^f T_{1,i}^{-1} T_{2,n} E_1 T_{2,n} T_{i,1}^{-1} \\ &= T_{2f+1,i}^{-1} E^f T_{1,2f+1}^{-1} T_{2,2f+2} E_1 T_{2f+2,n} T_{2,n} T_{i,1}^{-1} \\ &\equiv 0 \pmod{\mathcal{B}_n^{f+1}} \end{aligned}$$

where the last equivalence (1) follows from Lemma 2.5(1). Obviously, (2) follows from (4.5) as follows:

$$E^f(2f - 1, n) = q E^f(2f, n) T_{2f-1}, \quad E^f E_{2f-1,n} = z E^f(2f, n), \quad E^f E_{2f,n} = z q^{-1} E^f(2f, n) T_{2f-1}^{-1}. \quad (4.5)$$

By Lemma 2.5(6),

$$\begin{aligned} E^f(2f - 1, n) &= E^f T_{2f-1}(2f, n) T_{2f-1} \\ &= q E^f(2f, n) T_{2f-1} \end{aligned}$$

proving the first equality in (4.5). By Lemma 2.5(5),

$$\begin{aligned} E^f E_{2f-1,n} &= E_3 E_5 \dots E_{2f-1} T_{n,3} E_1 T_2 E_1 T_{2,n} T_{2f,2}^{-1} T_{2f-1,1}^{-1} \\ &= z E^f(2f, n) \end{aligned}$$

where the last equality follows from Definition 2.1(3), (2.2) and Lemma 2.5(4). This proves the second equality in (4.5). By (4.2), Lemma 2.5(4) and the second equality in (4.5), we have

$$\begin{aligned} E^f E_{2f,n} &= E^f T_{2f-1}^{-1} E_{2f-1,n} T_{2f-1}^{-1} \\ &= zq^{-1} E^f(2f, n) T_{2f-1}^{-1}, \end{aligned}$$

proving the last equality in (4.5). \square

Proposition 4.8. *Suppose $(f, \lambda) \in \Lambda_n$ and $\mu^{(k)} = \lambda \setminus p_k$ in (3.7), $1 \leq k \leq a$. Then*

$$y_{\mu^{(k)}}^\lambda L_n \equiv \frac{q^{2c(p_k)} - 1}{q - q^{-1}} y_{\mu^{(k)}}^\lambda \pmod{N_{k-1}},$$

where $c(p_k) = j - i$ if p_k is in i -th row and j -th column of the Young diagram $[\lambda]$.

Proof. Thanks to Lemma 4.7, we immediately have

$$y_{\mu^{(k)}}^\lambda L_n \equiv y_{\mu^{(k)}}^\lambda \sum_{j=2f+1}^{n-1} (j, n) \pmod{\mathcal{B}_n^{f+1}}. \quad (4.6)$$

Now, the result follows from (4.6) and well-known results on the action of Jucys-Murphy elements for Hecke algebras in [9, Theorem 3.22]¹. \square

Lemma 4.9. *Suppose $(f, \lambda) \in \Lambda_n$. We have*

$$E^f x_\lambda T_{n,2f}^{-1} x \equiv 0 \pmod{N_a}$$

if $x \in \{(i, n) - q\delta_{2f-1,i}, E_{j,n} - z^{-1}(2f-1, j)^{-1}, E_{2f-1,n} - \delta\}$ and $2f-1 \leq i \leq n-1$, $2f-1 < j \leq n-1$.

Proof. Thanks to Lemma 2.5,

$$E^f x_\lambda T_{n,2f}^{-1}(i, n) = \begin{cases} x_\lambda E^f T_{i+1,n} T_{i,2f}^{-1} & \text{if } i \neq 2f-1, \\ qE^f x_\lambda T_{2f,n} & \text{if } i = 2f-1. \end{cases}$$

By Definition 3.12 and Lemma 3.14(3),

$$E^f x_\lambda T_{n,2f}^{-1}(i, n) \in N_a \text{ if } i \neq 2f-1. \quad (4.7)$$

In the remaining case, using the quadratic relation in (2.2) to rewrite $T_{2f,n}$ and using Lemma 3.14(3) again, we have

$$E^f x_\lambda T_{2f,n} - E^f x_\lambda T_{n,2f}^{-1} \in N_a. \quad (4.8)$$

When $x = (i, n) - q\delta_{2f-1,i}$, the result follows from (4.7)–(4.8). Similarly, by Lemma 2.5(4)(5), Lemma 2.6(7), and Definition 2.1(1)(2) we have

$$\begin{aligned} E^f x_\lambda T_{n,2f}^{-1} E_{2f-1,n} &= E^f x_\lambda T_{1,2f-1}^{-1} T_{2f,2} E_1 T_{2,n} T_{2f-1,1}^{-1} \\ &= \delta E^f x_\lambda T_{2,2f} T_{2f-1,1}^{-1} T_{2f,n} \\ &= \delta E^f x_\lambda T_{2f,n}, \end{aligned} \quad (4.9)$$

and

$$\begin{aligned} E^f T_{n,2f}^{-1} E_{2f,n} &= E^f T_{1,2f+1}^{-1} T_{2f+1,2} E_1 T_{2,n} T_{2f,1}^{-1} \\ &= E^f T_{2f,1} T_{1,2f+1}^{-1} E_1 T_{2,n} T_{2f,1}^{-1} \\ &= E^f T_{2f-1,1} T_{2,2f+1}^{-1} E_1 T_{2,n} T_{2f,1}^{-1} \\ &= E^f T_{2f}^{-1} T_{2f-1,1} T_{2,2f}^{-1} E_1 T_{2,n} T_{2f,1}^{-1} \\ &= E^f T_{2f}^{-1} E_{2f-1} T_{2f-1,1} T_{2,2f}^{-1} T_{2,n} T_{2f,1}^{-1} \\ &= z^{-1} E^f T_{2f-1,1} T_{2,2f}^{-1} T_{2,n} T_{2f,1}^{-1} \\ &= z^{-1} E^f T_{2f,n} T_{2f-1}^{-1}, \end{aligned} \quad (4.10)$$

¹The Hecke algebra \mathcal{H}_n is defined via $(T_i - q)(T_i + 1) = 0$ in [9]. Our $q^{-1}T_i$ (resp., q^2) is T_i (resp., q) in [9].

and

$$\begin{aligned}
E^f T_{n,2f}^{-1} E_{j,n} &= E^f T_{j+1,2f}^{-1} T_{1,j}^{-1} T_{n,j+1}^{-1} T_{n,2} E_1 T_{2,n} T_{j,1}^{-1} \\
&= E^f T_{j+1,2f}^{-1} T_{1,j}^{-1} T_{j+1,2} E_1 T_{2,n} T_{j,1}^{-1} \\
&= E^f T_{2f+1,j+1}^{-1} T_{j+1,2f}^{-1} T_{1,2f}^{-1} T_{j+1,2} E_1 T_{2,n} T_{j,1}^{-1} \\
&= E^f T_{2f+1,j+1}^{-1} T_{1,2f+1}^{-1} T_{2f+1,2} E_1 T_{2,n} T_{j,1}^{-1} \\
&= E^f T_{1,j+1}^{-1} T_{2f+1,2} E_1 T_{2,n} T_{j,1}^{-1} \\
&= E^f T_{2f,1} T_{1,j+1}^{-1} E_1 T_{2,n} T_{j,1}^{-1} \\
&= E^f T_{2f-1,1} T_{2,j+1}^{-1} E_1 T_{2,n} T_{j,1}^{-1} \\
&= E^f T_{2f,j+1}^{-1} T_{2f-1,1} T_{2,2f}^{-1} E_1 T_{2,n} T_{j,1}^{-1} \\
&= E^f T_{2f,j+1}^{-1} E_{2f-1} T_{2f-1,1} T_{2,2f}^{-1} T_{2,n} T_{j,1}^{-1} \\
&= z^{-1} E^f T_{2f-1,1} T_{2,2f}^{-1} T_{2f+1,j+1}^{-1} T_{2,n} T_{j,1}^{-1} \\
&= z^{-1} E^f T_{2,n} T_{2f,j}^{-1} T_{j,1}^{-1} \\
&= z^{-1} E^f T_{2,2f} T_{2f,n} T_{j,1}^{-1} T_{2f-1,j-1}^{-1} \\
&= z^{-1} E^f T_{2f-1,1} T_{2f,n} T_{j,1}^{-1} T_{2f-1,j-1}^{-1} \\
&= z^{-1} E^f T_{2f,n} T_{j,2f-1}^{-1} T_{2f-1,j-1}^{-1} \\
&= z^{-1} E^f T_{2f,n} (2f-1, j)^{-1}.
\end{aligned} \tag{4.11}$$

When $x = E_{j,n} - z^{-1}(2f-1, j)^{-1}$, the result follows from (4.8) and (4.10)-(4.11). In the remaining case, the result follows from (4.8)-(4.9). \square

Proposition 4.10. *Suppose $(f, \lambda) \in \Lambda_n$ and $\mu^{(k)} = \lambda \cup p_k$, $a < k \leq m$. Then*

$$y_{\mu^{(k)}}^\lambda L_n \equiv (q - q^{-1})^{-1} (z^{-2} q^{2-2c(p_k)} - 1) y_{\mu^{(k)}}^\lambda \pmod{N_{k-1}}.$$

Proof. Thanks to Lemma 4.7(2) and Lemma 4.9, we have

$$\begin{aligned}
E^f x_\lambda T_{n,2f}^{-1} L_n &= E^f x_\lambda T_{n,2f}^{-1} \left(\sum_{2f-1 \leq i \leq n-1} (i, n) - q^2 z^{-1} E_{i,n} \right) \\
&\equiv E^f x_\lambda T_{n,2f}^{-1} (q - q^2 z^{-1} \sum_{2f-1 \leq i \leq n-1} E_{i,n}) \\
&\equiv E^f x_\lambda T_{n,2f}^{-1} (q - q^2 z^{-1} \delta - q^2 z^{-2} \sum_{2f \leq j \leq n-1} (2f-1, j)^{-1}) \\
&\equiv -q^2 z^{-2} (E^f x_\lambda T_{n,2f}^{-1} x - E^f x_\lambda x' T_{n,2f}^{-1}) + (q - q^2 z^{-1} \delta) E^f x_\lambda T_{n,2f}^{-1} \pmod{N_a},
\end{aligned} \tag{4.12}$$

where

$$x' = \sum_{2f+1 \leq i < j \leq n} (i, j)^{-1} \text{ and } x = \sum_{2f-1 \leq i < j \leq n-1} (i, j)^{-1}.$$

Write $[k] = \frac{q^k - q^{-k}}{q - q^{-1}}$ for any $k \in \mathbb{Z}$. By [9, Theorem 3.22]²,

$$E^f x_\lambda x' - \sum_{p \in [\lambda]} q^{-c(p)} [c(p)] E^f x_\lambda \in \mathcal{B}_n^{\triangleright \lambda}, \quad y_{\mu^{(k)}}^\lambda x - \sum_{p \in [\mu^{(k)}]} q^{-c(p)} [c(p)] y_{\mu^{(k)}}^\lambda \in N_{k-1}. \tag{4.13}$$

Thanks to (4.12)-(4.13),

$$\begin{aligned}
y_{\mu^{(k)}}^\lambda L_n &\equiv (q - q^2 z^{-1} \delta - q^2 z^{-2} \frac{1 - q^{-2c(p_k)}}{q - q^{-1}}) E^f x_\lambda T_{n,2f}^{-1} h \\
&\equiv \frac{z^{-2} q^{2-c(p_k)} - 1}{q - q^{-1}} y_{\mu^{(k)}}^\lambda \pmod{N_{k-1}}
\end{aligned}$$

where $h \in \mathcal{B}_{n-1}$ such that $y_{\mu^{(k)}}^\lambda = E^f x_\lambda T_{n,2f}^{-1} h$ (see (3.6)).

²The Jucys-Murphy elements of the Hecke algebra in [9] were constructed via (i, j) . When we compute (4.13), the arguments in the proof of [9, Theorem 3.22] still work but the coefficients now are obtained from those in [9] by replacing q with q^{-1} .

□

Definition 4.11. Suppose $(f, \lambda) \in \Lambda_n$. For any $\mathbf{t} \in \mathcal{F}_n^{ud}(\lambda)$ and $1 \leq k \leq n$, define

$$c_{\mathbf{t}}(k) = \begin{cases} \frac{q^{2c(p)} - 1}{q - q^{-1}} & \text{if } \mathbf{t}_k = \mathbf{t}_{k-1} \cup p, \\ \frac{z^{-2} q^{2-c(p)} - 1}{q - q^{-1}} & \text{if } \mathbf{t}_k = \mathbf{t}_{k-1} \setminus p. \end{cases}$$

Theorem 4.12. Suppose $(f, \lambda) \in \Lambda_n$. For any $\mathbf{t} \in \mathcal{F}_n^{ud}(\lambda)$ and $1 \leq k \leq n$,

$$m_{\mathbf{t}} L_k = c_{\mathbf{t}}(k) m_{\mathbf{t}} + \sum_{\substack{\mathbf{s} \\ \mathbf{s} \triangleright \mathbf{t}}} a_{\mathbf{s}} m_{\mathbf{s}}$$

for some scalars $a_{\mathbf{s}} \in \mathbb{k}$.

Proof. We have $m_{\mathbf{t}} L_1 = 0$. Suppose $k \geq 2$. By Theorem 4.6(4) and induction assumption for $n - 1$, we have the required formulae on $m_{\mathbf{t}} L_k$, $1 \leq k \leq n - 1$. When $k = n$, the required formula follows from Propositions 4.8, 4.10 and Theorem 4.6(3)(4). □

By Theorem 4.12, $\{L_1, L_2, \dots, L_n\}$ are JM elements with respect to the Jucys-Murphy basis of any cell module $C(f, \lambda)$ in the sense of [11, Definition 2.4].

Now we assume that \mathcal{B}_n is defined over F . It follows from [8, Definition 2.3] that there is a canonical invariant form, say ψ on each cell module $C(f, \lambda)$. Let

$$D(f, \lambda) = C(f, \lambda) / \text{rad} C(f, \lambda),$$

where $\text{rad} C(f, \lambda)$ is the radical of $C(f, \lambda)$ with respect to ψ . Then all non-zero $D(f, \lambda)$ consist of all pair-wise non-isomorphic simple \mathcal{B}_n -modules [8, Theorem 3.4]. It follows from [22, Theorem 4.1] that $D(f, \lambda) \neq 0$ if and only if $\lambda \in \Lambda_e^+(n - 2f)$.

Corollary 4.13. Suppose $(f, \lambda) \in \Lambda_n$, $\mu \in \Lambda_e^+(n)$ and $[C(f, \lambda) : D(0, \mu)] \neq 0$. Then there are $(\mathbf{s}, \mathbf{t}) \in \mathcal{F}_n^{ud}(\mu) \times \mathcal{F}_n^{ud}(\lambda)$ such that $c_{\mathbf{s}}(k) = c_{\mathbf{t}}(k)$ for all $1 \leq k \leq n$.

Proof. Let L be the abelian subalgebra of \mathcal{B}_n generated by L_1, L_2, \dots, L_n . Restricting both $C(f, \lambda)$ and $D(0, \mu)$ to L and using Theorem 4.12 yields the result as required. □

5. RESTRICTION AND INDUCTION FUNCTORS

In this section, we consider the q -Brauer algebra over a field F . Recall the algebra homomorphism $\phi : \mathcal{B}_{n-2} \rightarrow \tilde{E}_1 \mathcal{B}_n \tilde{E}_1$ in Theorem 3.10. Identifying \mathcal{B}_{n-2} with $\tilde{E}_1 \mathcal{B}_n \tilde{E}_1$ via the algebra isomorphism ϕ , we have two functors $\mathcal{F} : \mathcal{B}_n\text{-mod} \rightarrow \mathcal{B}_{n-2}\text{-mod}$ and $\mathcal{G} : \mathcal{B}_{n-2}\text{-mod} \rightarrow \mathcal{B}_n\text{-mod}$ such that

$$\mathcal{F}(M) = M \tilde{E}_1, \text{ and } \mathcal{G}(N) = N \otimes_{\tilde{E}_1 \mathcal{B}_n \tilde{E}_1} \tilde{E}_1 \mathcal{B}_n \quad (5.1)$$

for any \mathcal{B}_n -module M and any \mathcal{B}_{n-2} -module N .

Proposition 5.1. Suppose $(f, \lambda) \in \Lambda_n$ and $(\ell, \mu) \in \Lambda_{n+2}$.

- (1) $\mathcal{F}\mathcal{G} = 1$,
- (2) $\mathcal{F}(C(f, \lambda)) \cong C(f - 1, \lambda)$ where $C(-1, \lambda)$ is defined to be 0,
- (3) $\mathcal{G}(C(f, \lambda)) \cong C(f + 1, \lambda)$,
- (4) $\text{Hom}_{\mathcal{B}_{n+2}}(\mathcal{G}(C(f, \lambda)), C(\ell, \mu)) \cong \text{Hom}_{\mathcal{B}_n}(C(f, \lambda), \mathcal{F}(C(\ell, \mu)))$.

Proof. Obviously, (1) follows from Theorem 3.10. Recall $\mathcal{D}_{f,n}$ in Definition 3.4(1). For any $d \in \mathcal{D}_{f,n-2}$, let \tilde{d} be obtained from d by replacing each factor s_i of d by s_{i+2} . Let $\mathcal{D}_{f,n}$ be the set of all such \tilde{d} . Then $\mathcal{D}_{f,n} \subset \mathcal{D}_{f+1,n}$. By Theorems 3.8, 3.10, $\tilde{E}_1 \mathcal{B}_n \tilde{E}_1$ has cellular basis

$$\{qz^{-1} T_1^{-1} T_2 \sigma(T_d) E^{f+1} x_{\mathbf{s}} T_e \mid d, e \in \mathcal{D}_{f,n}, \mathbf{s}, \mathbf{t} \in \mathcal{F}^{std}(\lambda), (f, \lambda) \in \Lambda_{n-2}\} \quad (5.2)$$

where x_{λ} corresponds to the index representation of $\mathcal{H}_{2f+3,n}$. If we denote by $\tilde{C}(f - 1, \lambda)$ the corresponding cell module of $\tilde{E}_1 \mathcal{B}_n \tilde{E}_1$ with respect to $(f - 1, \lambda)$, then

$$C(f - 1, \lambda) \cong \tilde{C}(f - 1, \lambda) \quad (5.3)$$

as \mathcal{B}_{n-2} -modules and the required isomorphism sends any basis element $E^{f-1} x_{\mathbf{s}} T_d + \mathcal{B}_{n-2}^{\triangleright \lambda}$ to $qz^{-1} T_1^{-1} T_2 E^f \phi(x_{\mathbf{s}}) \phi(T_d) + \tilde{B}^{\triangleright \lambda}$ where $\tilde{B}^{\triangleright \lambda}$ is the free \mathbb{k} -module spanned by

$$\bigcup_{(l, \mu) \in \Lambda_n} \{qz^{-1} T_1^{-1} T_2 \sigma(T_{e_1}) E^l x_{\mathbf{s}} T_{e_2} \mid \mathbf{s}, \mathbf{t} \in \mathcal{F}^{std}(\mu), e_1, e_2 \in \mathcal{D}_{l-1,n}, \mu \triangleright \lambda\}.$$

Now, we compute $\mathcal{F}(C(f, \lambda))$. Suppose $d \in \mathcal{D}_{f,n}$. Thanks to (3.5), $E^f T_d \tilde{E}_1$ can be written as a linear combination of elements $E^f T_e T_w$ with $w \in \mathfrak{S}_{3,n}$ and $e = s_{2f,if} s_{2f-1,jf} \cdots s_{4,i2} s_{3,j2} \in \mathcal{D}_{f-1,n}$. This shows $\mathcal{F}(C(f, \lambda))$ has basis

$$\{E^f x_t T_d + \mathcal{B}_n^{\triangleright \lambda} \mid t \in \mathcal{T}^{std}(\lambda), d \in \mathcal{D}_{f-1,n}\}.$$

So $\mathcal{F}_n(C(f, \lambda)) \cong \tilde{C}(f-1, \lambda)$ and the required right $\tilde{E}_1 \mathcal{B}_n \tilde{E}_1$ -isomorphism is the linear isomorphism sending $E^f x_t T_d + \mathcal{B}_n^{\triangleright \lambda}$ to $qz^{-1} T_1^{-1} T_2 E^f x_t T_d + \tilde{B}^{\triangleright \lambda}$. Now (2) follows from (5.3).

By definition, $\mathcal{G}(C(f, \lambda)) = \overline{E^f x_\lambda} \mathcal{B}_n \otimes_{\tilde{E}_1 \mathcal{B}_{n+2} \tilde{E}_1} \tilde{E}_1 \mathcal{B}_{n+2}$ where $\overline{E^f x_\lambda} = E^f x_\lambda + \mathcal{B}_n^{\triangleright \lambda}$. There is a right \mathcal{B}_{n+2} -homomorphism $\psi : \mathcal{G}(C(f, \lambda)) \rightarrow C(f+1, \lambda)$ sending $\overline{E^f x_\lambda} \otimes \tilde{E}_1 h$ to $\overline{E^{f+1} x_\lambda} h$ for any $h \in \mathcal{B}_{n+2}$. Since $C(f+1, \lambda)$ is a cyclic module generated by $\overline{E^{f+1} x_\lambda}$, ψ is surjective. By Lemma 3.9, any element in $\mathcal{G}(C(f, \lambda))$ can be written as a linear combination of elements $\overline{E^f x_\mathfrak{s}} \otimes \tilde{E}_1 T_d$, where $\mathfrak{s} \in \mathcal{T}^{std}(\lambda)$ and $d \in \mathcal{D}_{f+1, n+2}$. So the number of all such elements is equal to the rank of $C(f+1, \lambda)$. Since we are considering the q -Brauer algebra over a field F , $\dim_F \mathcal{G}(C(f, \lambda)) \leq \dim_F C(f+1, \lambda)$, forcing ψ to be an isomorphism. Finally, (4) follows from the adjoint associativity of tensor, hom functors and the well-known fact that $\mathcal{F}(C(\ell, \mu)) \cong \text{Hom}_{\mathcal{B}_n}(\tilde{E}_1 \mathcal{B}_n, C(\ell, \mu))$. \square

6. THE RADICAL OF $C(1, \mu)$

In this section we consider the q -Brauer algebra over the field F containing invertible q such that $e > n$ where e is the quantum characteristic of q^2 . We are going to describe explicitly the radical of cell modules $C(1, \mu)$ for all $(1, \mu) \in \Lambda_n$. This result can be considered as the counterpart of [4, Theorem 3.4] for the Brauer algebra. From Lemma 6.2 to the end of Remark 6.8 we assume that the q -Brauer algebra is defined over \mathbb{k} . The following result is motivated by Doran-Wales-Hanlon's work on Brauer algebra in [4].

Lemma 6.1. *Let V be the free \mathbb{k} -module with basis $\{E_1 T_w T_d + \mathcal{B}_n^2 \mid w \in \mathfrak{S}_{3,n}, d \in \mathcal{D}_{1,n}\}$. Then V is an $(\mathcal{H}_{3,n}, \mathcal{B}_n)$ -bimodule with the natural actions induced by the multiplication in \mathcal{B}_n .*

Proof. The result follows from Theorem 3.8 and Lemma 3.9. \square

In the remaining part of this section, we write $a = q - q^{-1}$.

Lemma 6.2. *Suppose $s_{2,j_1} s_{1,i_1}, s_{2,j_2} s_{1,i_2} \in \mathcal{D}_{1,n}$. There are $h \in \sum_{x \in \mathfrak{S}_{3,n}} q^{\mathbb{Z}} \mathbb{Z}[a] T_x$, $w \in \mathfrak{S}_{3,n}$ and $b \in \{0, 1, 2\}$ such that*

$$E_1 T_{2,j_1} T_{1,i_1} T_{i_2,1} T_{j_2,2} E_1 \equiv \delta_{(i_1,j_1),(i_2,j_2)} \delta E_1 + (1 - \delta_{(i_1,j_1),(i_2,j_2)}) z q^b T_w E_1 + a z h E_1 \pmod{\mathcal{B}_n^2}. \quad (6.1)$$

Proof. Suppose $k < l$. Thanks to the quadratic relation in (2.2), and Definition 2.1,

$$T_{k,l} T_{l,k} = 1 + a \sum_{k+1 \leq i \leq l} T_{i,k+1} T_k T_{k+1,i}, \text{ and } E_1 T_2 T_1 T_3 T_2 E_1 \equiv a z E_1 + a z q E_1 T_3 \pmod{\mathcal{B}_n^2}. \quad (6.2)$$

If $l \leq i_1$, then

$$E_1 T_{2,j_1} T_{l,2} T_{1,l} T_{j_1,2} E_1 = T_{l+1,3} E_1 T_{2,j_1} T_1 T_{j_1,2} E_1 T_{3,l+1}. \quad (6.3)$$

Let $x = E_1 T_{2,j_1} T_1 T_{j_1,2} E_1$. By braid relations in (2.2), Definition 2.1(3) and (6.2),

$$\begin{aligned} x &= E_1 T_2 T_1 T_{3,j_1} T_{j_1,3} T_2 E_1 \\ &= E_1 T_2 T_1 (1 + a \sum_{4 \leq m \leq j_1} T_{m,4} T_{j_1,3}) T_2 E_1 \\ &= q^2 z E_1 + a \sum_{4 \leq m \leq j_1} T_{m,4} E_1 T_2 T_1 T_3 T_2 E_1 T_{4,m} \\ &\equiv q^2 z E_1 + a \sum_{4 \leq m \leq j_1} T_{m,4} (a z E_1 + a z q E_1 T_3) T_{4,m} \pmod{\mathcal{B}_n^2}. \end{aligned} \quad (6.4)$$

Now, we are ready to verify (6.1). Suppose $(i_1, j_1) = (i_2, j_2)$. By (6.2) and Definition 2.1(3)

$$\begin{aligned} E_1 T_{2,j_1} T_{1,i_1} T_{i_1,1} T_{j_1,2} E_1 &= E_1 T_{2,j_1} T_{j_1,2} E_1 + a E_1 T_{2,j_1} \sum_{2 \leq l \leq i_1} T_{l,2} T_1 T_{2,l} T_{j_1,2} E_1 \\ &= \delta E_1 + a \sum_{3 \leq k \leq j_1} T_{k,3} E_1 T_2 E_1 T_{3,k} + a E_1 T_{2,j_1} \sum_{2 \leq l \leq i_1} T_{l,2} T_1 T_{2,l} T_{j_1,2} E_1 \\ &= \delta E_1 + a z E_1 \sum_{3 \leq k \leq j_1} T_{k,3} T_{3,k} + a E_1 T_{2,j_1} \sum_{2 \leq l \leq i_1} T_{l,2} T_1 T_{2,l} T_{j_1,2} E_1. \end{aligned}$$

By (6.3)-(6.4), $E_1 T_{2,j_1} T_{1,i_1} T_{i_1,1} T_{j_1,2} E_1$ can be written as the required form in (6.1).

Suppose $(i_1, j_1) \neq (i_2, j_2)$. Applying anti-involution σ on (6.1), we see that it is enough to prove (6.1) in two cases: (1) $i_1 < i_2 < j_2$, (2) $i_1 = i_2$ and $j_1 < j_2$. In the first case, we have

$$E_1 T_{2,j_1} T_{1,i_1} T_{i_2,1} T_{j_2,2} E_1 = y T_{3,i_1+2} \quad (6.5)$$

where $y = E_1 T_{2,j_1} T_{i_2,1} T_{j_2,2} E_1$. We have

$$y = \begin{cases} T_{i_2,3} T_{j_2,4} E_1 T_2 T_1 T_3 T_2 E_1 T_{4,j_1+2} & \text{if } j_1 < i_2, \\ qz E_1 T_{j_2,3} + a \sum_{3 \leq m \leq j_1} T_{m,3} T_{j_2,4} E_1 T_2 T_1 T_3 T_2 E_1 T_{4,m+1} & \text{if } j_1 = i_2, \\ T_{i_2+1,3} E_1 T_{2,j_1} T_1 T_{j_2,2} E_1 & \text{if } j_1 > i_2. \end{cases} \quad (6.6)$$

Applying anti-involution σ on $E_1 T_{2,j_1} T_1 T_{j_2,2} E_1$, we see that it is enough to assume $j_1 \leq j_2$ when we assume $j_1 > i_2$. If $j_1 < j_2$, then

$$E_1 T_{2,j_1} T_1 T_{j_2,2} E_1 = T_{j_2,4} E_1 T_2 T_1 T_3 T_2 E_1 T_{4,j_1+1}. \quad (6.7)$$

Under the assumption (1) except $j_1 = j_2 > i_2$, (6.1) follows immediately from (6.5)–(6.6) if we use (6.2) to rewrite $E_1 T_2 T_1 T_3 T_2 E_1$. In the remaining case, since $E_1 T_{2,j_1} T_1 T_{j_2,2} E_1$ has already been computed in (6.4) and (6.1) follows immediately. In the second case,

$$\begin{aligned} E_1 T_{2,j_1} T_{1,i_1} T_{i_1,1} T_{j_2,1} E_1 &= E_1 T_{2,j_1} T_{j_2,2} E_1 + a \sum_{2 \leq m \leq i_1} E_1 T_{2,j_1} T_{m,2} T_{1,m} T_{j_2,2} E_1 \\ &= z E_1 T_{j_2,3} T_{3,j_1+1} + a \sum_{2 \leq m \leq i_1} T_{m+1,3} E_1 T_2 T_1 T_{j_2,2} E_1 T_{4,j_1+1} T_{3,m+1} \\ &= z E_1 T_{j_2,3} T_{3,j_1+1} + a \sum_{2 \leq m \leq i_1} T_{m+1,3} T_{j_2,4} E_1 T_2 T_1 T_3 T_2 E_1 T_{4,j_1+1} T_{3,m+1}. \end{aligned}$$

Rewriting $E_1 T_2 T_1 T_3 T_2 E_1$ via (6.2) again, we have (6.1) immediately. \square

Definition 6.3. Let $\phi : V \times V \rightarrow \mathbb{k}$ be the bilinear form such that for any $w_1, w_2 \in \mathfrak{S}_{3,n}$ and $d_1, d_2 \in \mathcal{D}_{1,n}$,

$$\phi(E_1 T_{w_1} T_{d_1} + \mathcal{B}_n^2, E_1 T_{w_2} T_{d_2} + \mathcal{B}_n^2) = \tau(h)$$

where $h \in \mathcal{H}_{3,n}$ such that $E_1 T_{w_1} T_{d_1} \sigma(T_{d_2}) \sigma(T_{w_2}) E_1 \equiv E_1 h \pmod{\mathcal{B}_n^2}$ and $\tau : \mathcal{H}_{3,n} \rightarrow \mathcal{Z}$ is the trace function in (2.3).

Thanks to Theorem 3.8 and Corollary 3.11(1), the element h in Definition 6.3 is unique and the bilinear form ϕ is well-defined. The following result follows immediately from Definition 6.3.

Lemma 6.4. *Let $\phi : V \times V \rightarrow \mathbb{k}$ be the bilinear form in Definition 6.3. Then ϕ is a symmetric invariant form in the sense that $\phi(xa, y) = \phi(x, y\sigma(a))$ for all $x, y \in V$ and $a \in \mathcal{B}_n$.*

In order to simplify notation, define

$$v_{w,d_j,i} = E_1 T_w T_{d_j,i} + \mathcal{B}_n^2, \quad (6.8)$$

where $d_{j,i} = s_{2,j} s_{1,i} \in \mathcal{D}_{1,n}$ and $w \in \mathfrak{S}_{3,n}$. Then $\{v_{w,d_j,i} \mid 1 \leq i < j \leq n, w \in \mathfrak{S}_{3,n}\}$ is a basis of V .

Lemma 6.5. *For any $1 \leq k \leq n-1$, T_k acts on the right of V as a symmetric matrix.*

Proof. There are some $a_{(w',d_{j',i'}),(w,d_j,i)} \in \mathbb{k}$ and $(w',d_{j',i'}) \in \mathfrak{S}_{3,n} \times \mathcal{D}_{1,n}$ such that

$$v_{w,d_j,i} T_k = \sum_{(w',d_{j',i'})} a_{(w',d_{j',i'}),(w,d_j,i)} v_{w',d_{j',i'}}. \quad (6.9)$$

Let $A_k = (a_{(w', d_{j', i'}), (w, d_{j, i})})$ be the $m \times m$ -matrix where m is the rank of V . Then A_k is the matrix with respect to T_k . The LHS of (6.9) can be computed explicitly by Definition 2.1. There are five cases: (1) $i = k + 1$, (2) $i = k$, (3) $i > k + 1$, (4) $k > j$ (5) $i < k \leq j$.

Thanks to the braid relations in (2.2) and Definition 2.1(2), we have

$$v_{w, d_{j, i}} T_k = \begin{cases} (q - q^{-1})v_{w, d_{j, k+1}} + v_{w, d_{j, k}} & \text{if } i = k + 1, \\ v_{w, d_{j, k+1}} & \text{if } i = k, j > k + 1, \\ qv_{w, d_{j, k}} & \text{if } i = k, j = k + 1. \end{cases}$$

When $i > k + 1$,

$$v_{w, d_{j, i}} T_k = \begin{cases} v_{ws_{k+2}, d_{j, i}} & \text{if } \ell(ws_{k+2}) > \ell(w), \\ v_{ws_{k+2}, d_{j, i}} + (q - q^{-1})v_{w, d_{j, i}} & \text{if } \ell(ws_{k+2}) < \ell(w). \end{cases}$$

If $k > j$, then

$$v_{w, d_{j, i}} T_k = \begin{cases} v_{ws_k, d_{j, i}} & \text{if } \ell(ws_k) > \ell(w), \\ v_{ws_k, d_{j, i}} + (q - q^{-1})v_{w, d_{j, i}} & \text{if } \ell(ws_k) < \ell(w). \end{cases}$$

Finally, assume $i < k \leq j$. If $k \geq j - 1$, then

$$v_{w, d_{j, i}} T_k = \begin{cases} v_{w, d_{j+1, i}} & \text{if } k = j, \\ (q - q^{-1})v_{w, d_{j, i}} + v_{w, d_{j-1, i}} & \text{if } k = j - 1. \end{cases}$$

In the remaining case, $k < j - 1$. We have

$$v_{w, d_{j, i}} T_k = \begin{cases} v_{ws_{k+1}, d_{j, i}} & \text{if } \ell(ws_{k+1}) > \ell(w), \\ (q - q^{-1})v_{w, d_{j, i}} + v_{ws_{k+1}, d_{j, i}} & \text{if } \ell(ws_{k+1}) < \ell(w). \end{cases}$$

In summary, it follows from explicit computation on $v_{w, d_{j, i}} T_k$ that each diagonal entry of A_k is one of scalar in $\{q, q - q^{-1}, 1\}$ and each off diagonal entry of A_k is either 0 or 1. Further, for any $(w', d_{j', i'}) \neq (w, d_{j, i})$, $a_{(w', d_{j', i'}), (w, d_{j, i})} = 1$ if and only if $a_{(w, d_{j, i}), (w', d_{j', i'})} = 1$. In particular, A_k is a symmetric matrix, proving the result. \square

Lemma 6.6. *Let G be the Gram matrix with respect to the basis $\{v_{w, d_{j, i}} \mid 1 \leq i < j \leq n, w \in \mathfrak{S}_{3, n}\}$ of V and the symmetric invariant form ϕ in Definition 6.3. If $\psi : V \rightarrow V$ is the linear map given by G with respect to the basis above, then ψ is a $(\mathcal{H}_{3, n}, \mathcal{H}_n)$ -bimodule homomorphism.*

Proof. Thanks to Lemma 6.4, GA_k is a symmetric matrix. By Lemma 6.5, $GA_k = A_k^t G^t = A_k G$, which is equivalent to saying that ψ is a right \mathcal{H}_n -homomorphism. Suppose $3 \leq i \leq n$ and $(w, d), (w', d') \in \mathfrak{S}_{3, n} \times \mathcal{D}_{1, n}$. Thanks to Definition 6.3 there is an $h \in \mathcal{H}_{3, n}$ such that

$$\phi(T_i E_1 T_w T_d, E_1 T_{w'} T_{d'}) = \tau(T_i h) \quad \text{and} \quad \phi(E_1 T_w T_d, T_i E_1 T_{w'} T_{d'}) = \tau(h T_i).$$

Since τ is a trace function, we have $\phi(T_i E_1 T_w T_d, E_1 T_{w'} T_{d'}) = \phi(E_1 T_w T_d, T_i E_1 T_{w'} T_{d'})$, proving that ψ is a left $\mathcal{H}_{3, n}$ -homomorphism. Further, ψ is a $(\mathcal{H}_{3, n}, \mathcal{H}_n)$ -homomorphism since two actions are induced by the multiplication on \mathcal{B}_n . \square

Proposition 6.7. *For any two basis elements $v_{w, d}, v_{w', d'}$ of V in (6.8),*

$$\phi(v_{w, d}, v_{w', d'}) = \begin{cases} \delta + azf(q) & \text{if } (w, d) = (w', d'), \\ czq^b + azg(q) & \text{otherwise,} \end{cases}$$

for some $b \in \mathbb{Z}$, $c \in \{0, 1\}$ and $f(q), g(q) \in q^{\mathbb{Z}}\mathbb{Z}[a]$.

Proof. The result follows from (2.3) and (6.1). \square

Remark 6.8. Suppose $z = q^m$ for some $m \in \mathbb{Z}$. The classical limit of \mathcal{B}_n is the Brauer algebra $B_n(m)$ over \mathbb{Z} (see Remark 2.2). In this case, the above bilinear form becomes the corresponding bilinear form on V for $B_n(m)$ by setting $q \rightarrow 1$ and the corresponding Gram matrix is $G_1 = \lim_{q \rightarrow 1} G$. More explicitly, the diagonal (resp., off diagonal) entries of the Gram matrix G_1 are of form δ (resp., 0 or 1) and the linear map $\psi : V \rightarrow V$ induced by G_1 in Lemma 6.6 becomes an $(\mathfrak{S}_{3, n}, \mathfrak{S}_n)$ -bimodule homomorphism.

From here to the end of this section, we assume that \mathcal{B}_n is defined over the field F which contains a non-zero q such that $e > n$ where e is the quantum characteristic of q^2 .

Definition 6.9. Suppose $e > n$ and $(1, \mu), (0, \lambda) \in \Lambda_n$. We say that λ is $(1, \mu)$ -admissible if

- (1) $\mu \subset \lambda$ and $\lambda \setminus \mu = \{p_1, p_2\}$ such that p_1, p_2 are not at the same column,
- (2) $z^2 = q^{2-2(c(p_1)+c(p_2))}$.

Theorem 6.10. For any $(1, \mu), (0, \lambda) \in \Lambda_n$, $[C(1, \mu) : C(0, \lambda)] \neq 0$ if and only if λ is $(1, \mu)$ -admissible.

Proof. Since we are assuming $e > n$, \mathcal{H}_n is semisimple and any cell module S^λ of \mathcal{H}_n is irreducible. Recall the cell module $C(1, \mu)$ has a basis $\{E_1 x_\mu T_{d(t)} T_v + \mathcal{B}_n^{\triangleright \mu} \mid (v, t) \in I(\mu)\}$. So, it is easy to check that $C(1, \mu) \cong \text{Ind}_{\mathcal{H}_2 \otimes \mathcal{H}_{3,n}}^{\mathcal{H}_n} M \boxtimes N$ as \mathcal{H}_n -modules, where M (resp., N) is the \mathcal{H}_2 -module (resp., $\mathcal{H}_{3,n}$ -module) with basis $\{E_1\}$ (resp., $\{x_\mu T_{d(t)} + \mathcal{H}_{3,n}^{\triangleright \mu} \mid t \in \mathcal{T}^{std}(\mu)\}$). Since $M \cong S^\alpha$ with $\alpha = (2)$ (see Definition 2.1(2)) and $N \cong S^\mu$, we have

$$C(1, \mu) \cong \text{Ind}_{\mathcal{H}_2 \otimes \mathcal{H}_{3,n}}^{\mathcal{H}_n} S^\alpha \boxtimes S^\mu$$

as right \mathcal{H}_n -modules. Using Littlewood-Richardson rule for semisimple Hecke algebra \mathcal{H}_n ,

$$C(1, \mu) \cong \bigoplus_\gamma S^\gamma \tag{6.10}$$

as \mathcal{H}_n -modules, where $\gamma \in \Lambda^+(n)$ obtained from μ by adding two boxes which are not at the same column. Moreover, the multiplicity of S^γ is 1. So $[C(1, \mu) : C(0, \lambda)] \neq 0$ if and only if $[C(1, \mu) : C(0, \lambda)] = 1$. Further, by Corollary 4.13, $\lambda \setminus \mu = \{p_1, p_2\}$ such that

$$\frac{q^{2c(p_1)} - 1}{q - q^{-1}} = \frac{z^{-2} q^{2-c(p_2)} - 1}{q - q^{-1}}. \tag{6.11}$$

Therefore λ is $(1, \mu)$ -admissible.

We are going to prove $[C(1, \mu) : C(0, \lambda)] \neq 0$ if λ is $(1, \mu)$ -admissible. Consider the $(\mathcal{H}_{3,n}, \mathcal{H}_n)$ -bimodule homomorphism induced by $\psi : V \rightarrow V$ in Lemma 6.6 via base change. Since we are assuming that $e > n$, both \mathcal{H}_n and $\mathcal{H}_{3,n}$ are semisimple. This implies that V is direct sum of cell modules $C(1, \mu)$ with multiplicity $\dim S^\mu$. Since ψ is a left $\mathcal{H}_{3,n}$ -homomorphism, the restriction of ψ to $C(1, \mu)$ fixes $C(1, \mu)$. Let e_μ be the primitive idempotent with respect to S^μ . Then $S^\mu = e_\mu \mathcal{H}_n$ and $\tau(e_\mu) \neq 0$ (see [10, Lemma 1.6]). Since $\phi(E_1 e_\mu, E_1 e_\mu) = \delta \tau(e_\mu) \neq 0$, the restriction of ψ to the cell module $C(1, \mu)$ is non-trivial and hence

$$\text{Ker} \psi \cap C(1, \mu) \neq C(1, \mu) \tag{6.12}$$

Thanks to (6.10), $C(1, \mu) = \bigoplus C(1, \mu)_\gamma$ where $C(1, \mu)_\gamma$ is the right \mathcal{H}_n -submodule such that

$$C(1, \mu)_\gamma \cong S^\gamma.$$

Then $C(1, \mu) e_\gamma \mathcal{H}_n = C(1, \mu)_\gamma$. Since ψ is a right \mathcal{H}_n -homomorphism, ψ fixes each $C(1, \mu)_\gamma$. Pick up a basis of $C(1, \mu)_\gamma$ for each γ . The disjoint union of such bases forms a new basis of $C(1, \mu)$. Note that each $C(1, \mu)_\gamma$ is a right irreducible \mathcal{H}_n -module. With respect to this basis, the corresponding matrix with respect to the restriction of ψ to $C(1, \mu)$ is of form $\text{diag}(h_\gamma I_\gamma)_\gamma$ where I_γ is the $\dim S^\gamma \times \dim S^\gamma$ identity matrix and $h_\gamma \in F$.

Recall that $\{v_{w,d} \mid w \in \mathfrak{S}_{3,n}, d \in \mathcal{D}_{1,n}\}$ is a basis of V and the disjoint union of bases of $C(1, \mu)_\gamma$ form a new basis of V . Moreover, since our computation only involves Hecke algebra, each entry in the transition matrix between these two bases of V is of form $f(q) \otimes 1_F$ for some $f(q) = \frac{g(q)}{h(q)}$ where $g(q), h(q) \in \mathbb{Z}[q]$ such that $h(q) \otimes 1_F \neq 0$. Abusing of notation, we will denote $a \otimes_{\mathbb{k}} 1_F$ by a . Thanks to Proposition 6.7 each entry in the diagonal (resp., off diagonal) of the matrix G of ψ with respect to the basis is of form $\delta + zg(q)$ (resp., $zf(q)$) where $f(q), g(q) \in F$. Let $m = \dim V$. So, there exist $f_i(q) \in F$, $1 \leq i \leq m$ with $f_j(q) \neq 0$ for some j such that

$$G(f_1(q), \dots, f_m(q))^t = h_\gamma (f_1(q), \dots, f_m(q))^t$$

for each γ in (6.10). Then $(\delta + zg(q))f_j(q) + zf(q) = h_\gamma f_j(q)$ for some $f(q), g(q) \in F$ and hence

$$h_\gamma = \delta + zf_\gamma(q) \tag{6.13}$$

for some $f_\gamma(q) \in F$. We explain that there are z and q such that $h_\lambda = 0$. This can be seen as follows.

Let $z = q^b$ for some $b \in \mathbb{Z}$ and consider the classical limit of \mathcal{B}_n . It is the Brauer algebra $B_n(b)$ over \mathbb{Z} . The corresponding Brauer algebra $B_n(b)$ over \mathbb{C} is $B_n(b) \otimes_{\mathbb{Z}} \mathbb{C}$. Unless otherwise stated, we work on Brauer algebra $B_n(b)$ over \mathbb{C} at moment. Then

$$\left(\lim_{q \rightarrow 1} f_1(q), \dots, \lim_{q \rightarrow 1} f_m(q) \right),$$

is the corresponding eigenvector of G_1 in Remark 6.8 (In fact, it is the corresponding G_1 over \mathbb{C}). Further, $\lim_{q \rightarrow 1} \delta = b$ and $\lim_{q \rightarrow 1} h_\gamma = b + h$, where $\lim_{q \rightarrow 1} f_\gamma(q) = h$. We may choose b such that $b + h = 0$. This implies that $C(1, \mu)_\lambda$ is in the kernel of the ψ in classical limit case. Such a kernel is the corresponding radical of the invariant form ϕ . By (6.12), $C(1, \mu)_\lambda$ is a proper submodule of $C(1, \mu)$. If we denote $\text{rad}C(1, \mu)$ by the radical with respect to the canonical invariant form on $C(1, \mu)$, then $C(1, \mu)_\lambda \subset \text{rad}C(1, \mu)$. Standard arguments on cellular algebras in [15] shows that $\text{rad}C(1, \mu)$ is killed by E_1 and so is $C(1, \mu)_\lambda$. This proves that $C(1, \mu)_\lambda \cong C(0, \lambda)$. By [4, Theorem 3.2], $b = 1 - c(p_1) - c(p_2)$ and $h = -b$. Since $\delta \neq 0$, $\lim_{q \rightarrow 1} f_\lambda(q) = -b \neq 0$. So $f_\lambda(q) \neq 0$. Since $\lim_{q \rightarrow 1} f_\lambda(q)$ exist, $(q - q^{-1})f_\lambda(q) \neq 1$.

We may choose z and q such that $z^{-2} = 1 - (q - q^{-1})f_\lambda(q) \notin \{0, 1\}$. In other words, we can find z and q such that δ is invertible and $h_\lambda = 0$. By arguments similar to those for the Brauer algebra above, $C(1, \mu)_\lambda \cong C(0, \lambda)$ and hence $z^2 = q^{2-2(c(p_1)+c(p_2))}$ (see (6.11)). \square

7. PROOF OF THEOREM A

In this section, we give a proof of Theorem A. For any $(f, \lambda) \in \Lambda_n$, let $G_{f, \lambda}$ be the Gram matrix associated to the canonical invariant form (see [8, Definition 2.3]) on $C(f, \lambda)$ with respect to the cellular basis in Theorem 3.8.

Proposition 7.1. *For $n \geq 2$, let \mathcal{B}_n be defined over F containing non-zero $q, z, q - q^{-1}, z - z^{-1}$ such that the quantum characteristic e of q^2 is strictly bigger than n . Then \mathcal{B}_n is (split) semisimple if and only if $\det G_{1, \lambda} \neq 0$ for all $\lambda \in \Lambda^+(k - 2)$ and $2 \leq k \leq n$.*

Proof. Thanks to Theorem 3.8, \mathcal{B}_n is a cellular algebra over an arbitrary field. So any field is a split field of \mathcal{B}_n . By [8, Theorem 3.8], \mathcal{B}_n is (split) semisimple if and only if $\prod_{(f, \lambda) \in \Lambda_n} \det G_{f, \lambda} \neq 0$. Since we are assuming $e > n$, \mathcal{H}_n is semisimple and hence $\prod_{\lambda \in \Lambda^+(n)} \det G_{0, \lambda} \neq 0$. Consequently, \mathcal{B}_n is (split) semisimple if and only if $\prod_{(f, \lambda) \in \widetilde{\Lambda}_n} \det G_{f, \lambda} \neq 0$ where $\widetilde{\Lambda}_n = \{(f, \lambda) \in \Lambda_n \mid f > 0\}$.

We prove the result by induction on n . Since we are assuming that δ is invertible, $\det G_{1, \emptyset} = \delta \neq 0$. So \mathcal{B}_2 is always semisimple. In the following we assume $n > 2$ and the result is true for \mathcal{B}_r with $r \leq n - 1$.

“ \Leftarrow ”: If the result were false, there should be an (f, λ) with $f \geq 2$ such that $\det G_{f, \lambda} = 0$. In this case, there is a simple module $D(\ell, \mu) \subset \text{rad}C(f, \lambda)$. In particular, $(\ell, \mu) \triangleleft (f, \lambda)$ in the sense that either $\ell < f$ or $\ell = f$ and $\mu \triangleleft \lambda$. However, since $D(\ell, \mu) \hookrightarrow C(f, \lambda)$, there is a non zero homomorphism from $C(\ell, \mu)$ to $C(f, \lambda)$. Acting the exact functor \mathcal{F} and using Proposition 5.1(2)-(4) yields a non-zero homomorphism $C(0, \mu) \rightarrow C(f - \ell, \lambda)$. We claim $\ell < f$. Otherwise, both $C(0, \mu)$ and $C(0, \lambda)$ are simple modules of the semisimple Hecke algebra \mathcal{H}_{n-2f} . So $\lambda = \mu$ and hence $[C(f, \lambda) : D(f, \lambda)] \geq 2$, a contradiction. Hence $f - \ell > 0$.

If $\ell > 0$, then $\det G_{f-\ell, \lambda} = 0$, a contradiction since $\mathcal{B}_{n-2\ell}$ is semisimple by induction assumption. Suppose $\ell = 0$. Thanks to Theorem 3.20, there are two boxes p_1 and p_2 such that either $[C(f, \lambda \setminus p_1) : C(0, \mu \setminus p_2)] \neq 0$ or $[C(f - 1, \lambda \cup p_1) : C(0, \mu \setminus p_2)] \neq 0$. Since we are assuming $f \geq 2$, $f > f - 1 \geq 1$. Consequently, either $\det G_{f, \lambda \setminus p_1} \neq 0$ or $\det G_{f-1, \lambda \cup p_1} \neq 0$. This is a contradiction since \mathcal{B}_{n-1} is semisimple by induction assumption.

“ \Rightarrow ” Suppose $\det G_{1, \lambda} = 0$ for some $\lambda \in \Lambda^+(k), 0 \leq k \leq n - 2$. Then $k \neq 0$ and there is a monomorphism $C(0, \mu) \hookrightarrow C(1, \lambda)$ for some partition $|\mu| = k + 2$. Since \mathcal{B}_n is semisimple, we have $k < n - 2$. If $k \equiv n \pmod{2}$, then we apply the functor \mathcal{G} repeatedly so as to get a non-zero homomorphism from $C(f - 1, \mu)$ to $C(f, \lambda)$ where $f = \frac{1}{2}(n - k)$. So $\det G_{f, \lambda} = 0$ and \mathcal{B}_n is not semisimple, a contradiction. Suppose $k \equiv n + 1 \pmod{2}$. Since $D(0, \mu) \hookrightarrow C(1, \lambda)$, by Theorem 6.10, μ is $(1, \lambda)$ -admissible. By Definition 6.9, $z^2 = q^{2(1-c(p_1)-c(p_2))}$ where $\mu = \lambda \cup \{p_1, p_2\}$. The arguments in the proof of [14, Theorem 4.1] shows that there is a pair of partitions $\tilde{\lambda}$ and $\tilde{\mu}$ such that $(0, \tilde{\mu})$ is $(1, \tilde{\lambda})$ -admissible and $|\tilde{\lambda}| = |\lambda| \pm 1$. By Theorem 6.10, there is a non-zero homomorphism from $C(0, \tilde{\mu})$ to $C(1, \tilde{\lambda})$. Applying the functor \mathcal{G} repeatedly yields $\det G_{f, \tilde{\lambda}} = 0$ where $f = \frac{1}{2}(n - |\tilde{\lambda}|)$. So \mathcal{B}_n is not semisimple, a contradiction. \square

Proof of Theorem A: Thanks to (2.5), $\mathcal{H}_n \cong \mathcal{B}_n / \langle E_1 \rangle$, where $\langle E_1 \rangle$ is the two-sided ideal generated by E_1 . Therefore, \mathcal{B}_n is not semisimple if \mathcal{H}_n is not semisimple. The later is equivalent to the condition $e \leq n$. In summary, we can assume $e > n$ when we discuss semisimplicity of \mathcal{B}_n over an arbitrary field. In this case, Proposition 7.1 gives a necessary and sufficient condition for \mathcal{B}_n being semisimple

over an arbitrary field. Thanks to Definition 6.9 and Theorem 6.10, \mathcal{B}_n is semisimple if and only if $z^2 \neq q^{2a}$ where

$$a \in \bigcup_{k=2}^n \{1 - \sum_{p \in \lambda \setminus \mu} c(p) \mid \lambda \in \Lambda^+(k), \mu \subset \lambda, \lambda \setminus \mu = \{p_1, p_2\} \text{ and } p_1, p_2 \text{ are not at the same column}\}$$

By [15, Lemma 2.4], this is equivalent to saying that a is given in (1.1). \square

REFERENCES

- [1] J. BIRMAN, H. WENZL, “Braids, link polynomials and a new algebra”, *Trans. Amer. Math. Soc.*, **313** (1), (1989), 249–273.
- [2] R. BRAUER, “On algebras which are connected with the semisimple continuous groups”, *Ann. of Math.*, **38**, (1937), 857–872.
- [3] W. CUI, Y. SHEN, “Canonical basis of q -Brauer algebras and ι Schur dualities”, arXiv:2203.02082v1[math. QA].
- [4] W. DORAN, D. WALES, P. HANLON, “On the semisimplicity of the Brauer centralizer algebras”, *J. Algebra* **211** (1999), 647–685.
- [5] J. ENYANG, “Specht modules and semisimplicity criteria for Brauer and Birman-Murakami-Wenzl algebras”, *J. Algebr. Comb.*, **26**, (2007), 291–341.
- [6] M. JIMBO, “A q -difference analogue of $U(g)$ and the Yang-Baxter equation”, *Lett. Math. Phys.*, **10** (1985), no. 1, 63–69.
- [7] G. LETZTER, “Coideal subalgebras and quantum symmetric pairs, in: New Directions in Hopf Algebras”, *Math. Sci. Res. Inst. Publ.*, **43**, (2002), 117–165.
- [8] J. GRAHAM AND G. LEHRER, “Cellular algebras”, *Invent. Math.*, **123** (1996), 1–34.
- [9] A. MATHAS, “*Twahori-Hecke algebras and Schur algebras of the symmetric group*”, *University Lecture Series*, **15**, American Mathematical Society, Providence, RI, 1999.
- [10] A. MATHAS, “Matrix units and generic degrees for the Ariki-Koike algebras”, *J. Algebra*, **281** (2004), 695–730.
- [11] A. MATHAS “Seminormal forms and Gram determinants for cellular algebras”, With an appendix by Marcos Soriano, *J. Reine Angew. Math.* **619** (2008), 141–173.
- [12] A.I. MOLEV, “A new quantum analog of the Brauer algebra”, *Czechoslovak J. Phys.* **53** (2003), 1073–1078.
- [13] J. MURAKAMI, “The Kauffman polynomial of links and representation theory”, *Osaka Journal of Mathematics* **24** (1987) 745–758.
- [14] H. RUI, “A criterion on the semisimple Brauer algebras”, *J. Combin. Theory Ser. A* **111** (2005), no. 1, 78–88.
- [15] H. RUI AND M. SI, “A criterion on the semisimple Brauer algebras, II”, *J. Combin. Theory Ser. A* **113** (2006), 1199–1203.
- [16] H. RUI AND M. SI, “Discriminants of Brauer algebras”, *Math. Z.*, **258** (2008), no. 4, 925–944.
- [17] H. RUI AND M. SI, “Gram determinants and semisimplicity criteria for Birman-Wenzl algebras”, *J. Reine Angew. Math.* **631** (2009) 153–80.
- [18] H. RUI AND M. SI “Blocks of Birman-Murakami-Wenzl algebras”, *Int. Math. Res. Not. IMRN* 2011, no. 2, 452–486.
- [19] H. RUI, J. XU, “The representations of cyclotomic BMW algebras”, *J. Pure Appl. Algebra*, **213** (2009), no. 12, 2262–2288.
- [20] A. SARTORI, T. DANIEL, “Webs and q -Howe dualities in types BCD”, *Trans. Amer. Math. Soc.*, **371** (2019), no. 10, 7387–7431.
- [21] NGUYEN, DUNG TIEN, “Cellular structure of the q -Brauer algebras”, *Algebr. Represent. Theory*, **17** (2014), 1359–1400.
- [22] NGUYEN, DUNG TIEN, “A cellular basis of the q -Brauer algebra related with Murphy bases of Hecke algebras”, *J. Algebra Appl.*, **17** (2018), no. 5, 1850086, 26 pp.
- [23] I. SCHUR, Dissertation, Berlin, 1901. In I. Schur, “Gesammelte Abhandlungen I”, 1–70, Springer, Berlin, 1973.
- [24] H. WENZL, “A q -Brauer algebra”, *J. Algebra* **358** (2012), 102–127.
- [25] H. WENZL, “Quantum groups and subfactors of type B, C, and D”, *Comm. Math. Phys.*, **133** (1990), 383–432.
- [26] H. WENZL, “Quotients of representation rings”, *Represent. Theory*, **15** (2011), 385–406.
- [27] H. WENZL, “Fusion symmetric spaces and subfactors”, *Pacific J. Math.*, **259** (2012), no. 2, 483–510.

H.R. SCHOOL OF MATHEMATICAL SCIENCE, TONGJI UNIVERSITY, SHANGHAI, 200092, CHINA
Email address: hbrui@tongji.edu.cn

M.S. SCHOOL OF MATHEMATICS, SHANGHAI JIAOTONG UNIVERSITY, SHANGHAI, 200240, CHINA
Email address: simei@sjtu.edu.cn

L.S. SCHOOL OF MATHEMATICAL SCIENCE, TONGJI UNIVERSITY, SHANGHAI, 200092, CHINA
Email address: llsong@tongji.edu.cn