

p -LINEAR SCHEMES FOR SEQUENCES MODULO p^r

FRITS BEUKERS

ABSTRACT. We construct finite p -automata for the computation of interesting combinatorial sequences modulo p^r . They are presented in the form of so-called p -linear schemes.

1. INTRODUCTION

In this paper we study infinite sequences of integers that have special properties when they are reduced modulo a prime p or a prime power p^r . The most familiar sequences are those sequences $a(k)$ that have the so-called Lucas property modulo almost all primes p ,

$$a(k_m p^m + \dots + k_1 p + k_0) \equiv a(k_m) \cdots a(k_1) a(k_0) \pmod{p}$$

for all k_i with $0 \leq k_i \leq p - 1$. Examples are the exponential sequences like 2^k , the central binomial coefficients $\binom{2k}{k}$, the Apéry and Franel-type numbers given by

$$\sum_{m=0}^k \binom{k}{m}^2 \binom{k+m}{m}^2, \quad \sum_{m=0}^k \binom{k}{m}^\ell$$

and countless others. Our first result concerns constant term sequences of the form

$$g(k) = \text{ct}[g(\mathbf{x})^k] \quad k \geq 0,$$

where $g(\mathbf{x}) \in \mathbb{Z}[x_1^{\pm 1} \cdots x_n^{\pm 1}]$ is a Laurent polynomial in the n variables x_1, \dots, x_n and 'ct' is the operation of taking the constant term. In this paper we denote the Newton polytope of $g(\mathbf{x})$ by Δ . In [4, Thm 1.4] we find the following theorem.

Theorem 1.1. *Suppose that Δ has $\mathbf{0}$ as unique lattice point in its interior. Then, for any prime p the constant term sequence $g(k)$ has the Lucas property.*

In Section 3 we present a proof, which is basically the same as in [4] but which is amenable to the generalization studied in this paper. Here are some examples of interest. In [4, Example 2.3] we find

$$\binom{2k}{k} = \text{ct} \left[\left(x + 2 + \frac{1}{x} \right)^k \right].$$

In [4, Example 2.4] we find

$$\sum_{\substack{k_1, \dots, k_n \geq 0 \\ k_1 + \dots + k_n = k}} \left(\frac{k!}{k_1! \cdots k_n!} \right)^2 = \text{ct} \left[(1 + x_1 + \dots + x_n)^k \left(1 + \frac{1}{x_1} + \dots + \frac{1}{x_n} \right)^k \right].$$

In [4, Example 2.5] we find

$$\sum_{m=0}^k \binom{m+k}{m}^2 \binom{k}{m}^2 = \text{ct} \left[\left(\frac{(x+y)(z+1)(x+y+z)(y+z+1)}{xyz} \right)^k \right].$$

These are the famous Apéry numbers that occur in the irrationality proof of $\zeta(3)$. Finally we mention

$$\sum_{m=0}^k \binom{k}{m}^n = \text{ct} \left[(1 + x_1)^k (1 + x_2)^k \cdots (1 + x_n)^k \left(1 + \frac{1}{x_1 \cdots x_n} \right)^k \right]$$

which is not hard to show.

In practice one cannot expect the Lucas property to hold modulo p^r for $r > 1$. This can be circumvented by the introduction of matrix versions of the Lucas property.

Definition 1.2. A p -linear scheme modulo p^r for a set of sequences $\mathbf{a}(k) = (a_1(k), \dots, a_s(k))^t$, $k = 0, 1, \dots$ is a set of $s \times s$ -matrices $M(\ell)$ with entries in \mathbb{Z} such that

$$(1) \quad \mathbf{a}(kp + \ell) \equiv M(\ell)\mathbf{a}(k) \pmod{p^r}$$

for every $\ell \in \{0, 1, \dots, p-1\}$ and $k \geq 0$.

We say that any constant linear combination of the sequences $a_1(k), \dots, a_s(k)$ is a sequence that can be generated by a p -linear scheme modulo p^r . We call s the number of states of this p -linear scheme.

As explained in [8], p -linear schemes are special instances of finite p -automata. A simple corollary of Definition 1.2 is

$$\mathbf{a}(k_m p^m + \dots + k_1 p + k_0) \equiv M(k_0)M(k_1) \cdots M(k_m)\mathbf{a}(0) \pmod{p^r},$$

where $0 \leq k_i \leq p-1$ for all $i = 0, \dots, m$. So all one needs in order to compute $\mathbf{a}(k) \pmod{p^r}$ for any k are the matrices $M(\ell)$ for $\ell = 0, 1, \dots, p-1$, the vector $\mathbf{a}(0)$ and the base p expansion of k . Furthermore, if $r = 1$, $s = 1$ and $a(0) = 1$ then Definition 1.2 comes down to the Lucas property modulo p .

There is also a formulation of a p -linear scheme in terms of generating power series. Let $\mathbf{F}(t) = \sum_{k \geq 0} \mathbf{a}(k)t^k$, a vector of generating series, and $M(t) = \sum_{\ell=0}^{p-1} M(\ell)t^\ell$. Then (1) is equivalent to

$$(2) \quad \mathbf{F}(t) \equiv M(t)\mathbf{F}(t^p) \pmod{p^r}.$$

A very simple example of a 2-state p -linear scheme is given by the sequences $a(k) = 2^k, b(k) = k2^k$. It is easy to verify that for any odd prime p ,

$$\begin{aligned} a(kp + \ell) &\equiv 2^\ell a(k) + p\alpha 2^\ell b(k) \pmod{p^2} \\ b(kp + \ell) &\equiv \ell 2^\ell a(k) + p2^\ell(1 + \ell\alpha)b(k) \pmod{p^2}, \end{aligned}$$

where $\alpha \equiv \frac{2^{p-1}-1}{p} \pmod{p}$. So we can say that the sequence $2^k \pmod{p^2}$ is generated by a two state p -linear scheme. In general $2^k \pmod{p^r}$ can be generated by an r -state p -linear scheme.

An example of historical interest is Ira Gessel's mod p^2 congruence for the Apéry numbers $A(k) = \sum_{m=0}^k \binom{k}{m}^2 \binom{m+k}{m}^2$.

Theorem 1.3 (Gessel, 1982). Define for all integers $k \geq 0$,

$$A'(k) = \sum_{m=0}^k \binom{k}{m}^2 \binom{m+k}{m}^2 \left(\frac{1}{m+1} + \dots + \frac{1}{k} \right).$$

Then, for any prime p and any integers $k \geq 0$ and $\ell = 0, 1, \dots, p-1$ we have

$$A(kp + \ell) \equiv A(\ell)A(k) + pA'(\ell)kA(k) \pmod{p^2}.$$

Multiply this result with $kp + \ell$ to get

$$(kp + \ell)A(kp + \ell) \equiv \ell A(\ell)A(k) + p(A(\ell) + \ell A'(\ell))kA(k) \pmod{p^2}.$$

So the sequences $A(k), kA(k)$ form the two states of a p -linear scheme modulo p^2 .

Suppose again that $g(\mathbf{x})$ is a Laurent polynomial in x_1, \dots, x_n and coefficients in \mathbb{Z} . The following statement is an immediate consequence of Theorem 3.1 and Remark 3.2.

Corollary 1.4. Let $r \in \mathbb{N}$ and p a prime. Choose ρ such that $\rho - \lceil \rho/p \rceil \geq r - 1$. Then the numbers $g(k) \pmod{p^r}$ can be generated by a p -linear scheme with s states, where s is bounded by ρ times the number of lattice points in $\rho\Delta^\circ$. Here Δ° is the euclidean interior of the Newton polytope Δ of $g(\mathbf{x})$.

Note that if $r = 1$ we can take $\rho = 1$. Since $g(0) = 1$ we recover Theorem 1.1 if $\mathbf{0}$ is the unique lattice point in Δ° . Notice also that ρ can always be chosen such that $\rho \leq 2r$.

Our second result concerns coefficients of power series expansions for rational functions. Let $P(\mathbf{x}), Q(\mathbf{x}) \in \mathbb{Z}[x_1, \dots, x_n]$ and suppose that $P_0 := P(\mathbf{0}) \neq 0$. Then we can expand

$$\frac{Q(\mathbf{x})}{P(\mathbf{x})} = \sum_{\mathbf{k} \in \mathbb{Z}_{\geq 0}^n} a(\mathbf{k})\mathbf{x}^{\mathbf{k}}$$

with $a(\mathbf{k}) \in \mathbb{Z}[1/P_0]$. Of particular interest in the literature are the diagonal sequences $a(k, k, \dots, k)$. Some typical examples,

- The diagonal elements of $(1 - x_1 - \dots - x_n)^{-1}$ are $\frac{(kn)!}{(k!)^n}$.
- The diagonal elements of $((1 - x_1 - x_2)(1 - x_3 - x_4) - x_1x_2x_3x_4)^{-1}$ are the Apéry numbers introduced above. See [7].

In [9] we find a long list of further examples, with further references to Sloane's Online Encyclopedia.

In Theorem 4.1 we construct a p -linear scheme to compute the coefficients $a(\mathbf{k})$. This time the coefficients a have $\mathbb{Z}_{\geq 0}^n$ as domain, rather than $\mathbb{Z}_{\geq 0}$. However, it is straightforward to replace $\mathbf{a}(k)$ by $\mathbf{a}(\mathbf{k})$ in Definition 1.2. Here is a consequence of Corollary 4.2.

Corollary 1.5. *Let notations be as above and suppose that $Q(\mathbf{x}) = 1$, $P_0 = 1$ and the degree of $P(\mathbf{x})$ in each variable is 1. Then, for every prime p , $\mathbf{k} \in \mathbb{Z}_{\geq 0}^n$ and $\ell \in [0, 1, \dots, p-1]^n$ we have*

$$a(p\mathbf{k} + \ell) \equiv a(\ell)a(\mathbf{k}) \pmod{p}.$$

To prove this, apply Corollary 4.2 with $r = 1$ (hence $\rho = 1$) and $d_1 = \dots = d_n = 1$. Then we get a 1-state p -linear scheme to compute the numbers $a(\mathbf{k}) \pmod{p}$. Since $a(\mathbf{0}) = 1$ our corollary follows. Of course the restricted sequence $a(k, k, \dots, k)$ has now the Lucas property modulo every prime p .

In [5] and [6] the authors propose algorithms to construct finite p -automata and in particular p -linear schemes for the computation of sequences $a(k) \pmod{p^r}$. Unfortunately the algorithms seem rather inefficient (except in 'small' cases) and the resulting upper bounds for the number of states seem exceedingly large with a strong dependence on p (see [8, Thm 2.4]). These papers were the motivation for writing the present note. Its method is a simplified version of the techniques in [3], which studies the action of the so-called Cartier operator on spaces of rational functions. However, the present paper can be completely understood without having to read [3].

Acknowledgement: Many thanks Armin Straub for the discussions we had on the subject of the paper and his encouragement.

2. MAIN TOOL

The starting point in this paper is a Laurent polynomial f in the variables x_1, \dots, x_n and coefficients in a ring R which is either \mathbb{Z} or $\mathbb{Z}[t]$. Let p be any prime and let σ be the Frobius lift on R , which is the identity map on \mathbb{Z} and which maps t to t^p in the ring $\mathbb{Z}[t]$. Let Δ be the Newton polytope of f and $\mu \subset \Delta$ an "open" subset obtained by removing a finite number of closed subfaces from Δ . A common example is $\mu = \Delta^\circ$, the euclidean interior of Δ .

In [1],[2],[3] we study the action of the Cartier operator \mathcal{C}_p on the R -module $\Omega_f(\mu)$ of rational functions of the form $(r-1)!A(\mathbf{x})/f(\mathbf{x})^r$, where $A(\mathbf{x})$ is a Laurent polynomial with support in $r\mu$. Suppose we are given an infinite Laurent series $\sum_{\mathbf{k}} a_{\mathbf{k}}\mathbf{x}^{\mathbf{k}}$, $a_{\mathbf{k}} \in R$, with support in a positive cone with $\mathbf{0}$ as top. Then \mathcal{C}_p is defined by

$$\mathcal{C}_p \left(\sum_{\mathbf{k}} a_{\mathbf{k}}\mathbf{x}^{\mathbf{k}} \right) = \sum_{\mathbf{k}} a_{p\mathbf{k}}\mathbf{x}^{\mathbf{k}}.$$

The action of \mathcal{C}_p on a rational function $R(\mathbf{x}) \in \Omega_f(\mu)$ is obtained by expansion of $R(\mathbf{x})$ in an infinite Laurent series and then application of \mathcal{C}_p . Unfortunately the image is a Laurent series which usually does not come from an element in $\Omega_f(\mu)$. Fortunately, in [1] it is shown that the

image does lie in the p -adic completion of $\Omega_{f^\sigma}(\mu)$. Here f^σ is simply f with σ applied to its coefficients.

In the present paper we will only need a mod p^r version of this fact and for its derivation we need the following easy lemma.

Lemma 2.1. *Let $K(\mathbf{x}), L(\mathbf{x})$ be two infinite Laurent series in x_1, \dots, x_n supported in the same cone with top $\mathbf{0}$. Then,*

$$\mathcal{C}_p(K(\mathbf{x}^p)L(\mathbf{x})) = K(\mathbf{x})\mathcal{C}_p(L(\mathbf{x})).$$

Let r be a positive integer and consider the module of rational functions $\frac{A(\mathbf{x})}{f(\mathbf{x})^k}$ with support of $A(\mathbf{x})$ in $r\mu$. by $M(r)$. By $M^\sigma(r)$ we denote module generated by the elements of $M(r)$ where we have applied σ to all coefficients of the rational functions.

Proposition 2.2. *Let r be a positive integer and p a prime. Choose an integer ρ such that $\rho - \lceil \rho/p \rceil \geq r - 1$. Then the Cartier operator \mathcal{C}_p modulo p^r maps $M(\rho)$ to $M^\sigma(\rho)$.*

Proof. Rewrite $A(\mathbf{x})/f(\mathbf{x})^\rho$ as $A(\mathbf{x})f(\mathbf{x})^{p\lceil \rho/p \rceil - \rho}/f(\mathbf{x})^{p\lceil \rho/p \rceil}$. Then note that $f(\mathbf{x})^p = f^\sigma(\mathbf{x}^p) - pG(\mathbf{x})$ for some Laurent polynomial G with coefficients in R and support in $p\Delta$. Then we use the p -adic expansion

$$\frac{A(\mathbf{x})}{f(\mathbf{x})^\rho} = \frac{A(\mathbf{x})f(\mathbf{x})^{p\lceil \rho/p \rceil - \rho}}{(f^\sigma(\mathbf{x}^p) - pG(\mathbf{x}))^{\lceil \rho/p \rceil}} = \sum_{m \geq 0} p^m \binom{\lceil \rho/p \rceil + m - 1}{m} \frac{G(\mathbf{x})^m}{f^\sigma(\mathbf{x}^p)^{m + \lceil \rho/p \rceil}} A(\mathbf{x})f(\mathbf{x})^{p\lceil \rho/p \rceil - \rho}.$$

Apply \mathcal{C}_p . We find that

$$\mathcal{C}_p \left(\frac{A(\mathbf{x})}{f(\mathbf{x})^\rho} \right) = \sum_{m \geq 0} p^m \binom{\lceil \rho/p \rceil + m - 1}{m} \frac{Q_m(\mathbf{x})}{f^\sigma(\mathbf{x})^{m + \lceil \rho/p \rceil}},$$

where the $Q_m(\mathbf{x}) = \mathcal{C}_p(A(\mathbf{x})G(\mathbf{x})^m f(\mathbf{x})^{p\lceil \rho/p \rceil - \rho})$ are Laurent polynomials in x_1, \dots, x_n with support in $(m + \lceil \rho/p \rceil)\mu$ and coefficients in R . The terms with $m \geq r$ all vanish modulo p^r . Hence

$$(3) \quad \mathcal{C}_p \left(\frac{A(\mathbf{x})}{f(\mathbf{x})^\rho} \right) \equiv \sum_{m=0}^{r-1} p^m \binom{\lceil \rho/p \rceil + m - 1}{m} \frac{Q_m(\mathbf{x})}{f^\sigma(\mathbf{x})^{m + \lceil \rho/p \rceil}} \pmod{p^r}.$$

Since $m + \lceil \rho/p \rceil \leq r - 1 + \lceil \rho/p \rceil \leq \rho$ we see that modulo p^r the right hand side is in $M^\sigma(\rho)$. \square

3. CONSTANT TERMS OF POWERS OF LAURENT POLYNOMIALS

We apply Proposition 2.2 to construct p -linear schemes for sequences of the form

$$\text{ct}[q(\mathbf{x})g(\mathbf{x})^k] \pmod{p^r},$$

where 'ct' is the operation of selecting the constant term and $q(\mathbf{x}), g(\mathbf{x}) \in \mathbb{Z}[x_1^{\pm 1}, \dots, x_n^{\pm 1}]$ are Laurent polynomials. Let $f(\mathbf{x}) = 1 - tg(\mathbf{x})$ and $\mu = \Delta^\circ$, the euclidean interior of the Newton polytope Δ . Let p be a prime, r a positive integer and ρ an integer such that $r - 1 \leq \rho - \lceil \rho/p \rceil$. For any monomial $\mathbf{x}^{\mathbf{u}} \in \rho\Delta^\circ$ expand $\mathbf{x}^{\mathbf{u}}/f(\mathbf{x})^\rho$ as power series in t and take the constant term of each coefficient. We get a power series

$$F_{\mathbf{u}}(t) := \text{ct} \left[\frac{\mathbf{x}^{\mathbf{u}}}{(1 - tg(\mathbf{x}))^\rho} \right] = \sum_{k \geq 0} \binom{\rho + k - 1}{\rho} \text{ct}[\mathbf{x}^{\mathbf{u}}g(\mathbf{x})^k] t^k,$$

which is the generating series of the numbers $\binom{\rho + k - 1}{\rho} \text{ct}[\mathbf{x}^{\mathbf{u}}g(\mathbf{x})^k]$. The functions $t^\ell F_{\mathbf{u}}(t)$ with $\ell = 0, 1, \dots, \rho - 1$ and $\mathbf{u} \in \Delta^\circ$ will be the states of our p -linear scheme. Usually we are interested in sequences of the form $\text{ct}[g(\mathbf{x})^k]$, which correspond to the constant term of $1/f(\mathbf{x})$. However, we can use the observation

$$\frac{1}{f(\mathbf{x})} = \frac{f(\mathbf{x})^{\rho-1}}{f(\mathbf{x})^\rho}.$$

So the constant term of $1/f(\mathbf{x}) \pmod{p^r}$ is a \mathbb{Z} -linear combination of the states $t^\ell F_{\mathbf{v}}(t)$ with $\mathbf{v} \in \rho\Delta^\circ$ and $\ell = 0, 1, \dots, \rho - 1$.

Theorem 3.1. *Let p, r, ρ and $F_{\mathbf{u}t}$ be as before. Then there exist polynomials $\lambda_{\mathbf{u},\ell,\mathbf{v},m}(t)$ in t of degree $< p$ such that*

$$t^\ell F_{\mathbf{u}}(t) \equiv \sum_{\mathbf{v} \in \rho\Delta^\circ} \sum_{m=0}^{\rho-1} \lambda_{\mathbf{u},\ell,\mathbf{v},m}(t) (t^p)^m F_{\mathbf{v}}(t^p) \pmod{p^r}.$$

for all $\mathbf{u} \in (\rho\Delta^\circ)_{\mathbb{Z}}$ and $\ell = 0, 1, \dots, \rho - 1$.

Remark 3.2. *Note that we have found a p -linear scheme with states $t^\ell F_{\mathbf{u}} \pmod{p^r}$, where $\ell = 0, 1, \dots, \rho - 1$ and $\mathbf{u} \in \rho\Delta^\circ$. The cardinality of $(\rho\Delta^\circ)_{\mathbb{Z}}$ is bounded polynomially in ρ of degree n . For every prime p the choice $\rho = 2r$ suffices so the number of such states has order $O(r^{n+1})$, independent of p .*

Proof of Theorem 3.1. Consider (3) with $A(\mathbf{x}) = \mathbf{x}^{\mathbf{u}}$ and expand as

$$(4) \quad \mathcal{C}_p \left(\frac{\mathbf{x}^{\mathbf{u}}}{f(\mathbf{x})^\rho} \right) \equiv \frac{1}{f^\sigma(\mathbf{x})^\rho} \sum_{\mathbf{v} \in \rho\mu} q_{\mathbf{u},\mathbf{v}}(t) \mathbf{x}^{\mathbf{v}} \pmod{p^r}.$$

The coefficients $q_{\mathbf{u},\mathbf{v}}(t)$ are polynomials in t and one easily verifies that their degrees are at most $p(r-1) + p[\rho/p] - \rho$. Let us choose polynomials $\lambda_{\mathbf{u},\ell,\mathbf{v},m}$ of degree $< p$ such that

$$t^\ell q_{\mathbf{u},\mathbf{v}} = \sum_{m=0}^{\rho-1} \lambda_{\mathbf{u},\ell,\mathbf{v},m}(t) t^{pm}.$$

This is possible because the degree of the left hand side is at most

$$p(r-1) + p[\rho/p] - \rho + \rho - 1 = p(r-1 + [\rho/p]) - 1 \leq p\rho - 1.$$

The constant terms on both sides of (4) are equal modulo p^r and by a straightforward computation our assertion follows. \square

4. POWER SERIES COEFFICIENTS OF RATIONAL FUNCTIONS

Our second main application of Proposition 2.2 is to the coefficients of power series expansions of rational functions. Let P be a polynomial in x_1, \dots, x_n with integer coefficients and a constant term which is not divisible by the prime p . Let us call this constant P_0 . Let Δ be the Newton polytope of P . For any polynomial Q with integer coefficients we can expand Q/P as power series. Let us write

$$\frac{Q(\mathbf{x})}{P(\mathbf{x})} = \sum_{\mathbf{k}} a(\mathbf{k}) \mathbf{x}^{\mathbf{k}},$$

and note that $a(\mathbf{k}) \in \mathbb{Z}[1/P_0]$ for all \mathbf{k} . We will show that for every $r \geq 1$ there is a p -linear scheme which generates the numbers $a(\mathbf{k}) \pmod{p^r}$. In particular this holds for coefficients $a(k\mathbf{v})$ with $\mathbf{v} = (1, 1, \dots, 1)$ and $k \in \mathbb{Z}_{\geq 0}$. These are the diagonal elements of the power series. Starting with the Newton polytope Δ we consider its interior Δ° and its boxed closure. This is the union of all boxes $[0, x_1] \times \dots \times [0, x_n]$ for every point $(x_1, \dots, x_n) \in \Delta^\circ$. Let us denote this closure by $B(\Delta)$.

Theorem 4.1. *Let p be a prime not dividing P_0 . Choose ρ such that $r - 1 \leq \rho - [\rho/p]$. For every $\mathbf{u} \in (\rho B(\Delta))_{\mathbb{Z}}$ consider the power series*

$$\frac{\mathbf{x}^{\mathbf{u}}}{P(\mathbf{x})^\rho} = \sum_{\mathbf{k}} a_{\mathbf{u}}(\mathbf{k}) \mathbf{x}^{\mathbf{k}},$$

where $a_{\mathbf{u}}(\mathbf{k}) \in \mathbb{Z}[1/P_0]$. Then there exist $\lambda_{\mathbf{u},\mathbf{v}}(\ell) \in \mathbb{Z}$ with $\mathbf{u}, \mathbf{v} \in (\rho B(\Delta))_{\mathbb{Z}}$ and $\ell \in [0, 1, \dots, p-1]^n$ such that for every \mathbf{k} we have

$$a_{\mathbf{u}}(p\mathbf{k} + \ell) \equiv \sum_{\mathbf{v} \in (\rho B(\Delta))_{\mathbb{Z}}} \lambda_{\mathbf{u},\mathbf{v}}(\ell) a_{\mathbf{v}}(\mathbf{k}) \pmod{p^r}.$$

Proof. We apply (3) in Proposition 2.2 with $f = P$ and $\mu = B(\Delta)$. We see that for any ℓ and any \mathbf{u} we have

$$\mathcal{C}_p \left(\frac{\mathbf{x}^{\mathbf{u}-\ell}}{P(\mathbf{x})^\rho} \right) \equiv \sum_{m=0}^{r-1} p^m \binom{\lceil \rho/p \rceil + m - 1}{\rho} \frac{Q_m(\mathbf{x})}{P(\mathbf{x})^{m+\lceil \rho/p \rceil}} \pmod{p^r}.$$

with $Q_m(\mathbf{x}) = \mathcal{C}_p(\mathbf{x}^{\mathbf{u}-\ell} G(\mathbf{x})^m P(\mathbf{x})^{p\lceil \rho/p \rceil - \rho})$. A priori the factor $\mathbf{x}^{\mathbf{u}-\ell}$ may also contribute negative exponents, but since these are $\geq 1 - p$, they do not survive the Cartier operator and the terms in Q_m have their support in $B(\Delta)$. Hence there exist coefficients $\lambda_{\mathbf{u}\mathbf{v}}(\ell) \in \mathbb{Z}$ such that

$$\mathcal{C}_p \left(\frac{\mathbf{x}^{\mathbf{u}-\ell}}{P(\mathbf{x})^\rho} \right) \equiv \sum_{\mathbf{v} \in (\rho B(\Delta))_{\mathbb{Z}}} \lambda_{\mathbf{u}\mathbf{v}}(\ell) \frac{\mathbf{x}^{\mathbf{v}}}{P(\mathbf{x})^\rho} \pmod{p^r}.$$

The theorem follows by taking the power series expansion of both sides and then selection of the coefficient of $\mathbf{x}^{\mathbf{k}}$ on both sides. \square

Corollary 4.2. *Let $a(\mathbf{k})$ be as in the beginning of this section and suppose that the support of Q is inside $B(\Delta)$. Let d_i be the degree of P in the variable x_i for $i = 1, \dots, n$. Then there is p -linear scheme which generates the numbers $a(\mathbf{k}) \pmod{p^r}$. The number of states is bounded by $\rho^n d_1 \cdots d_n$, where ρ is chosen so that $r - 1 \leq \rho - \lceil \rho/p \rceil$.*

Proof. We apply Theorem 4.1. Note that the lattice points in $B(\Delta)$ are contained in the box $[0, \rho d_1 - 1] \times \cdots \times [0, \rho d_n - 1]$. Hence the number of lattice points is bounded above by $\rho^n d_1 \cdots d_n$. One easily checks that Q/P can be written as \mathbb{Z} -linear combination of $\mathbf{x}^{\mathbf{u}}/P^\rho$ with $\mathbf{u} \in (\rho B(\Delta))_{\mathbb{Z}}$. \square

REFERENCES

- [1] F.Beukers, M.Vlasenko, *Dwork crystals I*
- [2] F.Beukers, M.Vlasenko, *Dwork crystals II*
- [3] F.Beukers, M.Vlasenko, *Dwork crystals III*
- [4] J.A.Henningsen, A.Straub, *Generalized Lucas congruences and linear p -schemes*. Advances in Applied Mathematics 141, 2022, 1–20.
- [5] E. Rowland and R. Yassawi, *Automatic congruences for diagonals of rational functions*. Journal de Théorie des Nombres de Bordeaux, 27(1), 2015, 245–288.
- [6] E. Rowland and D. Zeilberger, *A case study in meta-automation: automatic generation of congruence automata for combinatorial sequences*. Journal of Difference Equations and Applications, 20(7), 2014, 973–988.
- [7] *Multivariate Apéry numbers and supercongruences of rational functions*. Algebra & Number Theory, 8(8), 2014, 1985–2008.
- [8] A.Straub, *On congruence schemes for constant terms and their applications*. Research in Number Theory 8(3), 2022, 1–21.
- [9] J.-A. Weil, *Diagonals of rational functions and selected differential Galois groups*, http://www.unilim.fr/pages_perso/jacques-arthur.weil/diagonals/