

CORRIGENDUM TO “FROM A_1 TO A_∞ : NEW MIXED INEQUALITIES FOR CERTAIN MAXIMAL OPERATORS”

FABIO BERRA

ABSTRACT. We devote this note to correct an estimate concerning mixed inequalities for the generalized maximal function M_Φ , when certain properties of the associated Young function Φ are assumed.

Although the obtained estimates turn out to be slightly different, they are good extensions of mixed inequalities for the classical Hardy-Littlewood maximal functions M_r , with $r \geq 1$. They also allow us to obtain mixed estimates for the generalized fractional maximal operator $M_{\gamma,\Phi}$, when $0 < \gamma < n$ and Φ is an $L \log L$ type function.

OVERVIEW

Throughout this note we shall consider a Young function Φ with the following properties. Given $r \geq 1$ and $\delta \geq 0$, we say that a Young function Φ belongs to the family $\mathfrak{F}_{r,\delta}$ if it is submultiplicative, has lower type r and satisfies the condition

$$\frac{\Phi(t)}{t^r} \leq C_0(\log t)^\delta, \quad \text{for } t \geq t^*,$$

for some constants $C_0 > 0$ and $t^* \geq 1$.

In [2] we obtained mixed estimates for the operator M_Φ , where Φ belongs to $\mathfrak{F}_{r,\delta}$. Concretely, we stated the inequality

$$uv^r \left(\left\{ x \in \mathbb{R}^n : \frac{M_\Phi(fv)(x)}{v(x)} > t \right\} \right) \leq C \int_{\mathbb{R}^n} \Phi \left(\frac{|f|}{t} \right) uv^r$$

where u and v^r are weights belonging to the A_1 -Muckenhoupt class.

Later, in [1], the same kind of estimate was obtained when v^r is only assumed to be an A_∞ weight.

In the the proofs of both results we used Claim 3.4 in [2], and Claims 1 and 3 in [1] as auxiliary tools. These claims have an error on a Hölder estimate, where a limiting argument was mistakenly used and it cannot be adapted to obtain the inequality given above.

The purpose of this note is give a proof of Theorem 1 in [1] that avoids this step on the claims and allows us to obtain a slightly different estimate, that will still be useful for our purposes. We shall only modify the results obtained in [1], since they are more general and the corresponding version of those in [2] will follow as an immediate consequence. The modified mixed estimate in [1] is the following.

2010 *Mathematics Subject Classification.* 26A33, 42B25.

Key words and phrases. Young functions, maximal operators, Muckenhoupt weights, fractional operators.

The author was supported by CONICET and UNL.

Theorem 1 (Corrected version of Theorem 1 in [1]). *Let $r \geq 1$, $\delta \geq 0$ and $\Phi \in \mathfrak{F}_{r,\delta}$. If $u \in A_1$ and $v^r \in A_\infty$ then there exists a positive constant ε_0 such that the inequality*

$$uv^r \left(\left\{ x \in \mathbb{R}^n : \frac{M_\Phi(fv)(x)}{v(x)} > t \right\} \right) \leq C \int_{\mathbb{R}^n} (\eta_\varepsilon \circ \Phi) \left(\frac{|f(x)|}{t} \right) u(x)v^r(x) dx$$

holds for every positive t and every $0 < \varepsilon < \varepsilon_0$, where $\eta_\varepsilon(z) = z(1 + \log^+ z)^{\delta/\varepsilon}$ and C depends on ε .

It is not difficult to see that $M_\Phi v \gtrsim v$ when Φ belongs to $\mathfrak{F}_{r,\delta}$. So we have the following result as an immediate consequence of the theorem above.

Corollary 2 (Corrected version of Corollary 2 in [1]). *Under the assumptions in Theorem 1, there exists a positive constant ε_0 such that*

$$uv^r \left(\left\{ x \in \mathbb{R}^n : \frac{M_\Phi(fv)(x)}{M_\Phi v(x)} > t \right\} \right) \leq C \int_{\mathbb{R}^n} (\eta_\varepsilon \circ \Phi) \left(\frac{|f(x)|}{t} \right) u(x)v^r(x) dx$$

holds for every positive t and every $0 < \varepsilon < \varepsilon_0$, where η_ε is as above and C depends on ε .

Throughout these notes, all references, lemmas and theorems will follow the label given in [1].

1. PROOF OF THEOREM 1

We shall first give some preliminaries in order to proceed with the proof. Recall that $w \in A_\infty$ if there exists a positive constant C such that

$$\left(\frac{1}{|Q|} \int_Q w \right) \exp \left(\frac{1}{|Q|} \int_Q \log w^{-1} \right) \leq C$$

for every cube Q in \mathbb{R}^n . The smallest constant for which the inequality above holds is denoted by $[w]_{A_\infty}$.

The following lemma will be useful in the sequel. It can be found in [3].

Lemma 3. *Let $w \in A_\infty$ and let $r_w = 1 + \frac{1}{\tau_n[w]_{A_\infty}}$. Then for any cube Q we have*

$$\left(\frac{1}{|Q|} \int_Q w^{r_w} \right)^{1/r_w} \leq \frac{2}{|Q|} \int_Q w.$$

As a consequence, given any cube Q and a measurable set $E \subseteq Q$ we have that

$$\frac{w(E)}{w(Q)} \leq 2 \left(\frac{|E|}{|Q|} \right)^{\varepsilon_w},$$

where $\varepsilon_w = 1/(1 + \tau_n[w]_{A_\infty})$. The constant τ_n is purely dimensional and can be chosen as 2^{11+n} .

Recall that we are dealing with a function $\Phi \in \mathfrak{F}_{r,\delta}$, where $r \geq 1$ and $\delta \geq 0$ are given. Since we are assuming $v^r \in A_\infty$, there exists $\varepsilon_1 > 0$ such that $v^{r+\varepsilon} \in A_\infty$ for every $0 < \varepsilon \leq \varepsilon_1$.

Fix $0 < \varepsilon < \min\{\varepsilon_1, \varepsilon_2\}$, where $\varepsilon_2 > 0$ will be chosen later. Then we have that $v^r \in \text{RH}_s$, where $s = 1 + \varepsilon/r$. We shall denote $\Psi_\varepsilon = \eta_\varepsilon \circ \Phi$. We shall follow the same sketch and steps as in [1], where the entire proof is included for the sake of clearness. Recall that it is enough to prove that

$$uv^r \left(\left\{ x \in \mathbb{R}^n : \frac{M_{\Phi, \mathcal{D}}(fv)(x)}{v(x)} > t \right\} \right) \leq C_\varepsilon \int_{\mathbb{R}^n} \Psi_\varepsilon \left(\frac{|f(x)|}{t} \right) u(x)v^r(x) dx,$$

where \mathcal{D} is a given dyadic grid. We can also assume that $t = 1$ and that $g = |f|v$ is a bounded function with compact support. Then, for a fixed number $a > 2^n$, we can write

$$\begin{aligned} uv^r \left(\left\{ x \in \mathbb{R}^n : \frac{M_{\Phi, \mathcal{D}}(fv)(x)}{v(x)} > 1 \right\} \right) &= \sum_{k \in \mathbb{Z}} uv^r \left(\left\{ x : \frac{M_{\Phi, \mathcal{D}}g(x)}{v(x)} > 1, a^k < v \leq a^{k+1} \right\} \right) \\ &=: \sum_{k \in \mathbb{Z}} uv^r(E_k). \end{aligned}$$

For every $k \in \mathbb{Z}$ we consider the set

$$\Omega_k = \left\{ x \in \mathbb{R}^n : M_{\Phi, \mathcal{D}}g(x) > a^k \right\},$$

and by virtue of the Calderón-Zygmund decomposition of the space (see [1, Lemma 6]) there exists a collection of disjoint dyadic cubes $\{Q_j^k\}_j$ that satisfies

$$\Omega_k = \bigcup_j Q_j^k,$$

and $\|g\|_{\Phi, Q_j^k} > a^k$ for each j . By maximality, we have

$$(1.1) \quad a^k < \|g\|_{\Phi, Q_j^k} \leq 2^n a^k, \quad \text{for every } j.$$

For every $k \in \mathbb{Z}$ we now proceed to split the obtained cubes in different classes, as in [4]. Given a nonnegative integer ℓ , we set

$$\Lambda_{\ell, k} = \left\{ Q_j^k : a^{(k+\ell)r} \leq \frac{1}{|Q_j^k|} \int_{Q_j^k} v^r < a^{(k+\ell+1)r} \right\},$$

and also

$$\Lambda_{-1, k} = \left\{ Q_j^k : \frac{1}{|Q_j^k|} \int_{Q_j^k} v^r < a^{kr} \right\}.$$

The next step is to split every cube in the family $\Lambda_{-1, k}$. Fixed $Q_j^k \in \Lambda_{-1, k}$, we perform the Calderón-Zygmund decomposition of the function $v^r \chi_{Q_j^k}$ at level a^{kr} . Then we obtain, for each k , a collection $\{Q_{j,i}^k\}_i$ of maximal cubes, contained in Q_j^k and which satisfy

$$(1.2) \quad a^{kr} < \frac{1}{|Q_{j,i}^k|} \int_{Q_{j,i}^k} v^r \leq 2^n a^{kr}, \quad \text{for every } i.$$

Also we define the sets

$$\Gamma_{\ell, k} = \left\{ Q_j^k \in \Lambda_{\ell, k} : \left| Q_j^k \cap \left\{ x : a^k < v \leq a^{k+1} \right\} \right| > 0 \right\},$$

and also

$$\Gamma_{-1, k} = \left\{ Q_{j,i}^k : Q_j^k \in \Lambda_{-1, k} \text{ and } \left| Q_{j,i}^k \cap \left\{ x : a^k < v \leq a^{k+1} \right\} \right| > 0 \right\}.$$

Since $E_k \subseteq \Omega_k$, we can estimate

$$\begin{aligned} \sum_{k \in \mathbb{Z}} uv^r(E_k) &= \sum_{k \in \mathbb{Z}} uv^r(E_k \cap \Omega_k) \\ &= \sum_{k \in \mathbb{Z}} \sum_j uv^r(E_k \cap Q_j^k) \end{aligned}$$

$$\leq \sum_{k \in \mathbb{Z}} \sum_{\ell \geq 0} \sum_{Q_j^k \in \Gamma_{\ell, k}} a^{(k+1)r} u(E_k \cap Q_j^k) + \sum_{k \in \mathbb{Z}} \sum_{i: Q_{j,i}^k \in \Gamma_{-1, k}} a^{(k+1)r} u(Q_{j,i}^k).$$

If we can prove that given a negative integer N , there exists a positive constant C_ε , independent of N for which the following estimate

$$(1.3) \quad \sum_{k \geq N} \sum_{\ell \geq 0} \sum_{Q_j^k \in \Gamma_{\ell, k}} a^{(k+1)r} u(E_k \cap Q_j^k) + \sum_{k \geq N} \sum_{i: Q_{j,i}^k \in \Gamma_{-1, k}} a^{(k+1)r} u(Q_{j,i}^k) \leq C_\varepsilon \int_{\mathbb{R}^n} \Psi_\varepsilon(|f|) uv^r$$

holds, then the proof would be completed by letting $N \rightarrow -\infty$.

We shall also need the following lemma from [4]. We include an adaptation of the proof involving our parameters for the sake of clearness.

Lemma 4. *Let $\ell \geq 0$ and $Q_j^k \in \Gamma_{\ell, k}$. If $u \in A_\infty$ and $v^r \in A_q$ for some $1 < q < \infty$, then there exists positive constants c_1 and c_2 depending on u and v^r such that*

$$u(E_k \cap Q_j^k) \leq c_1 e^{-c_2 r \ell} u(Q_j^k).$$

Furthermore, we can pick $c_1 = 2([v^r]_{A_q} a^r)^{1/((q-1)(1+\tau_n[u]_{A_\infty}))}$ and $c_2 = \ln a / ((q-1)(1+\tau_n[u]_{A_\infty}))$, where τ_n is the dimensional constant appearing in Lemma 3.

Proof. Since $v^r \in A_\infty$, there exists $q > 1$ such that $v^r \in A_q$. Since $Q_j^k \in \Gamma_{\ell, k}$, we have that

$$\left(\frac{|E_k \cap Q_j^k|}{|Q_j^k|} \right)^{q-1} \leq \left(\frac{1}{|Q_j^k|} \int_{Q_j^k} v^{r(1-q')} \right)^{q-1} a^{r(k+1)} \leq \frac{[v^r]_{A_q} |Q_j^k|}{v^r(Q_j^k)} a^{r(k+1)} \leq [v^r]_{A_q} a^{(1-\ell)r}.$$

Since $u \in A_1 \subseteq A_\infty$, by Lemma 3 and the estimate above we have that

$$\frac{u(E_k \cap Q_j^k)}{u(Q_j^k)} \leq 2 \left(\frac{|E_k \cap Q_j^k|}{|Q_j^k|} \right)^{1/(1+\tau_n[u]_{A_\infty})} \leq 2 \left([v^r]_{A_q} a^{(1-\ell)r} \right)^{1/((q-1)(1+\tau_n[u]_{A_\infty}))}.$$

From this last inequality we can obtain the thesis. \square

Proof of Theorem 1. Since $u \in A_1$, we have $u \in A_\infty$. Moreover, the assumption $v^r \in A_\infty$ implies that there exists $1 < q < \infty$ such that $v^r \in A_q$. We take $\varepsilon_2 = r/((q-1)(1+\tau_n[u]_{A_\infty}))$ and $\varepsilon_0 = \min\{\varepsilon_1, \varepsilon_2\}$. Fixed $0 < \varepsilon < \varepsilon_0$, recall that we have to estimate the two quantities

$$A_N := \sum_{k \geq N} \sum_{\ell \geq 0} \sum_{Q_j^k \in \Gamma_{\ell, k}} a^{(k+1)r} u(E_k \cap Q_j^k)$$

and

$$B_N := \sum_{k \geq N} \sum_{i: Q_{j,i}^k \in \Gamma_{-1, k}} a^{(k+1)r} u(Q_{j,i}^k)$$

by $C_\varepsilon \int_{\mathbb{R}^n} \Psi_\varepsilon(|f|) uv^r$, with C_ε independent of N .

We shall start with the estimate of A_N . Fix $\ell \geq 0$ and let $\Delta_\ell = \bigcup_{k \geq N} \Gamma_{\ell, k}$. We define recursively a sequence of sets as follows:

$$P_0^\ell = \{Q : Q \text{ is maximal in } \Delta_\ell \text{ in the sense of inclusion}\}$$

and for $m \geq 0$ given we say that $Q_j^k \in P_{m+1}^\ell$ if there exists a cube Q_s^t in P_m^ℓ which verifies

$$(1.4) \quad \frac{1}{|Q_j^k|} \int_{Q_j^k} u > \frac{2}{|Q_s^t|} \int_{Q_s^t} u$$

and it is maximal in this sense, that is,

$$(1.5) \quad \frac{1}{|Q_{j'}^k|} \int_{Q_{j'}^k} u \leq \frac{2}{|Q_s^t|} \int_{Q_s^t} u$$

for every $Q_j^k \subsetneq Q_{j'}^k \subsetneq Q_s^t$.

Let $P^\ell = \bigcup_{m \geq 0} P_m^\ell$, the set of principal cubes in Δ_ℓ . By applying Lemma 4 and the definition of $\Lambda_{\ell,k}$ we have that

$$\begin{aligned} \sum_{k \geq N} \sum_{\ell \geq 0} \sum_{Q_j^k \in \Gamma_{\ell,k}} a^{(k+1)r} u(E_k \cap Q_j^k) &\leq \sum_{k \geq N} \sum_{\ell \geq 0} \sum_{Q_j^k \in \Gamma_{\ell,k}} c_1 a^{(k+1)r} e^{-c_2 \ell r} u(Q_j^k) \\ &\leq \sum_{\ell \geq 0} c_1 e^{-c_2 \ell r} a^{r(1-\ell)} \sum_{k \geq N} \sum_{Q_j^k \in \Gamma_{\ell,k}} \frac{v^r(Q_j^k)}{|Q_j^k|} u(Q_j^k). \end{aligned}$$

Let us sort the inner double sum in a more convenient way. We define

$$\mathcal{A}_{(t,s)}^\ell = \left\{ Q_j^k \in \bigcup_{k \geq N} \Gamma_{\ell,k} : Q_j^k \subseteq Q_s^t \text{ and } Q_s^t \text{ is the smallest cube in } P^\ell \text{ that contains it} \right\}.$$

That is, every $Q_j^k \in \mathcal{A}_{(t,s)}^\ell$ is not a principal cube, unless $Q_j^k = Q_s^t$. Recall that $v^r \in A_\infty$ implies that there exist two positive constants C and θ verifying

$$(1.6) \quad \frac{v^r(E)}{v^r(Q)} \leq C \left(\frac{|E|}{|Q|} \right)^\theta,$$

for every cube Q and every measurable set E of Q .

By using (1.5) and Lemma 12 in [1] we have that

$$\begin{aligned} \sum_{k \geq N} \sum_{Q_j^k \in \Gamma_{\ell,k}} \frac{v^r(Q_j^k)}{|Q_j^k|} u(Q_j^k) &= \sum_{Q_s^t \in P^\ell} \sum_{(k,j): Q_j^k \in \mathcal{A}_{(t,s)}^\ell} \frac{u(Q_j^k)}{|Q_j^k|} v^r(Q_j^k) \\ &\leq 2 \sum_{Q_s^t \in P^\ell} \frac{u(Q_s^t)}{|Q_s^t|} \sum_{(k,j): Q_j^k \in \mathcal{A}_{(t,s)}^\ell} v^r(Q_j^k) \\ &\leq C \sum_{Q_s^t \in P^\ell} \frac{u(Q_s^t)}{|Q_s^t|} v^r(Q_s^t) \left(\frac{|\bigcup_{(k,j): Q_j^k \in \mathcal{A}_{(t,s)}^\ell} Q_j^k|}{|Q_s^t|} \right)^\theta \\ &\leq C \sum_{Q_s^t \in P^\ell} \frac{u(Q_s^t)}{|Q_s^t|} v^r(Q_s^t). \end{aligned}$$

Therefore,

$$\sum_{k \geq N} \sum_{\ell \geq 0} \sum_{Q_j^k \in \Gamma_{\ell,k}} a^{(k+1)r} u(E_k \cap Q_j^k) \leq C \sum_{\ell \geq 0} e^{-c_2 \ell r} a^{-\ell r} \sum_{Q_s^t \in P^\ell} \frac{v^r(Q_s^t)}{|Q_s^t|} u(Q_s^t)$$

$$\leq C \sum_{\ell \geq 0} e^{-c_2 \ell r} \sum_{Q_s^t \in P^\ell} a^{tr} u(Q_s^t).$$

Claim 1 (Corrected version of Claim 1 in [1]). Given $\ell \geq 0$ and $Q_j^k \in \bigcup_{k \geq N} \Gamma_{\ell, k}$, we have that

$$(1.7) \quad a^{kr} \leq C \frac{\rho^{\delta/\varepsilon} a^{\ell\varepsilon}}{|Q_j^k|} \int_{Q_j^k} \Psi_\varepsilon(|f(x)|) v^r(x) dx,$$

where C depends on ε .

By applying the estimate above we obtain

$$\begin{aligned} \sum_{k \geq N} \sum_{\ell \geq 0} \sum_{Q_j^k \in \Gamma_{\ell, k}} a^{(k+1)r} u(E_k \cap Q_j^k) &\leq C_\varepsilon \sum_{\ell \geq 0} e^{-c_2 \ell r} \rho^{\delta/\varepsilon} a^{\ell\varepsilon} \sum_{Q_s^t \in P^\ell} \frac{u(Q_s^t)}{|Q_s^t|} \int_{Q_s^t} \Psi_\varepsilon(|f|) v^r \\ &= C_\varepsilon \sum_{\ell \geq 0} e^{-c_2 \ell r} \rho^{\delta/\varepsilon} a^{\ell\varepsilon} \int_{\mathbb{R}^n} \Psi_\varepsilon(|f|) v^r \left(\sum_{Q_s^t \in P^\ell} \frac{u(Q_s^t)}{|Q_s^t|} \chi_{Q_s^t} \right) \\ &= C_\varepsilon \sum_{\ell \geq 0} e^{-c_2 \ell r} \rho^{\delta/\varepsilon} a^{\ell\varepsilon} \int_{\mathbb{R}^n} \Psi_\varepsilon(|f(x)|) v^r(x) h_1(x) dx \\ &\leq C_\varepsilon \int_{\mathbb{R}^n} \Psi_\varepsilon(|f(x)|) v^r(x) u(x) dx, \end{aligned}$$

by virtue of Claim 2 in [1]. Notice that the sum is finite since we are assuming $\varepsilon < \varepsilon_2$. Indeed, we have that

$$e^{-c_2 \ell r} a^{\ell\varepsilon} = e^{-c_2 \ell r + \ell\varepsilon \ln a} = e^{\ell(-c_2 r + \varepsilon \ln a)},$$

and this exponent is negative by the election of ε . This completes the estimate of A_N .

Let us center our attention on the estimate of B_N . Fix $0 < \beta < \theta$, where θ is the number appearing in (1.6). We shall build the set of principal cubes in $\Delta_{-1} = \bigcup_{k \geq N} \Gamma_{-1, k}$. Let

$$P_0^{-1} = \{Q : Q \text{ is a maximal cube in } \Delta_{-1} \text{ in the sense of inclusion}\}$$

and, recursively, we say that $Q_{j,i}^k \in P_{m+1}^{-1}$, $m \geq 0$, if there exists a cube $Q_{s,l}^t \in P_m^{-1}$ such that

$$(1.8) \quad \frac{1}{|Q_{j,i}^k|} \int_{Q_{j,i}^k} u > \frac{a^{(k-t)\beta r}}{|Q_{s,l}^t|} \int_{Q_{s,l}^t} u$$

and it is the biggest subcube of $Q_{s,l}^t$ that verifies this condition, that is

$$(1.9) \quad \frac{1}{|Q_{j',i'}^{k'}|} \int_{Q_{j',i'}^{k'}} u \leq \frac{a^{(k-t)\beta r}}{|Q_{s,l}^t|} \int_{Q_{s,l}^t} u$$

if $Q_{j,i}^k \subsetneq Q_{j',i'}^{k'} \subsetneq Q_{s,l}^t$. Let $P^{-1} = \bigcup_{m \geq 0} P_m^{-1}$, the set of principal cubes in Δ_{-1} . Similarly as before, we define the set

$$\mathcal{A}_{(t,s,l)}^{-1} = \left\{ Q_{j,i}^k \in \bigcup_{k \geq N} \Gamma_{-1, k} : Q_{j,i}^k \subseteq Q_{s,l}^t \text{ and } Q_{s,l}^t \text{ is the smallest cube in } P^{-1} \text{ that contains it} \right\}.$$

We can therefore estimate B_N as follows

$$\begin{aligned}
B_N &\leq a^r \sum_{k \geq N} \sum_{i: Q_{j,i}^k \in \Gamma_{-1,k}} \frac{v^r(Q_{j,i}^k)}{|Q_{j,i}^k|} u(Q_{j,i}^k) \\
&\leq a^r \sum_{Q_{s,l}^t \in P^{-1}} \sum_{k,j,i: Q_{j,i}^k \in \mathcal{A}_{(t,s,l)}^{-1}} \frac{u(Q_{j,i}^k)}{|Q_{j,i}^k|} v^r(Q_{j,i}^k) \\
&\leq a^r \sum_{Q_{s,l}^t \in P^{-1}} \frac{u(Q_{s,l}^t)}{|Q_{s,l}^t|} \sum_{k \geq t} a^{(k-t)\beta r} \sum_{j,i: Q_{j,i}^k \in \mathcal{A}_{(t,s,l)}^{-1}} v^r(Q_{j,i}^k).
\end{aligned}$$

Fixed $k \geq t$, observe that

$$\sum_{j,i: Q_{j,i}^k \in \mathcal{A}_{(t,s,l)}^{-1}} |Q_{j,i}^k| < \sum_{j,i: Q_{j,i}^k \in \mathcal{A}_{(t,s,l)}^{-1}} a^{-kr} v^r(Q_{j,i}^k) \leq a^{-kr} v^r(Q_{s,l}^t) \leq 2^n a^{(t-k)r} |Q_{s,l}^t|.$$

Combining this inequality with the A_∞ condition of v^r we have, for every $k \geq t$, that

$$\begin{aligned}
\sum_{j,i: Q_{j,i}^k \in \mathcal{A}_{(t,s,l)}^{-1}} a^{(k-t)\beta r} v^r(Q_{j,i}^k) &\leq C v^r(Q_{s,l}^t) \left(\frac{\sum_{j,i: Q_{j,i}^k \in \mathcal{A}_{(t,s,l)}^{-1}} |Q_{j,i}^k|}{|Q_{s,l}^t|} \right)^\theta \\
&\leq C a^{(t-k)r\theta}.
\end{aligned}$$

Thus,

$$\begin{aligned}
B_N &\leq C \sum_{Q_{s,l}^t \in P^{-1}} \frac{u(Q_{s,l}^t)}{|Q_{s,l}^t|} v^r(Q_{s,l}^t) \sum_{k \geq t} a^{(t-k)r(\theta-\beta)} \\
&= C \sum_{Q_{s,l}^t \in P^{-1}} \frac{v^r(Q_{s,l}^t)}{|Q_{s,l}^t|} u(Q_{s,l}^t) \\
&\leq C \sum_{Q_{s,l}^t \in P^{-1}} a^{tr} u(Q_{s,l}^t).
\end{aligned}$$

Claim 2 (Corrected version of Claim 3 in [1]). If $Q_j^k \in \Lambda_{-1,k}$ then there exists a positive constant C_ε such that

$$a^{kr} \leq \frac{C_\varepsilon}{|Q_j^k|} \int_{Q_j^k} \Psi_\varepsilon(|f(x)|) v^r(x) dx.$$

By using this estimate we can proceed as follows

$$\begin{aligned}
\sum_{k \geq N} \sum_{i: Q_{j,i}^k \in \Gamma_{-1,k}} a^{(k+1)r} u(Q_{j,i}^k) &\leq C \sum_{Q_{s,l}^t \in P^{-1}} a^{tr} u(Q_{s,l}^t) \\
&\leq C_\varepsilon \sum_{Q_{s,l}^t \in P^{-1}} \frac{u(Q_{s,l}^t)}{|Q_{s,l}^t|} \int_{Q_s^t} \Psi_\varepsilon(|f(x)|) v^r(x) dx \\
&\leq C_\varepsilon \int_{\mathbb{R}^n} \Psi_\varepsilon(|f(x)|) v^r(x) \left[\sum_{Q_{s,l}^t \in P^{-1}} \frac{u(Q_{s,l}^t)}{|Q_{s,l}^t|} \chi_{Q_s^t}(x) \right] dx \\
&= C_\varepsilon \int_{\mathbb{R}^n} \Psi_\varepsilon(|f(x)|) v^r(x) h_2(x) dx
\end{aligned}$$

$$\leq C_\varepsilon \int_{\mathbb{R}^n} \Psi_\varepsilon(|f(x)|) u(x) v^r(x) dx,$$

by virtue of Claim 4 in [1]. This concludes the proof. \square

We proceed with the proofs of the claims, in order to complete the argument above.

Proof of Claim 1. Fix $\ell \geq 0$ and a cube $Q_j^k \in \bigcup_{k \geq N} \Gamma_{\ell, k}$. We know that $\|g\|_{\Phi, Q_j^k} > a^k$ or, equivalently, $\|\frac{g}{a^k}\|_{\Phi, Q_j^k} > 1$. Denote with $A = \{x \in Q_j^k : v(x) \leq t^* a^k\}$ and $B = Q_j^k \setminus A$, where t^* is the number verifying that if $z \geq t^*$, then

$$\frac{\Phi(z)}{z^r} \leq C_0 (\log z)^\delta.$$

Then,

$$1 < \left\| \frac{g}{a^k} \right\|_{\Phi, Q_j^k} \leq \left\| \frac{g}{a^k} \chi_A \right\|_{\Phi, Q_j^k} + \left\| \frac{g}{a^k} \chi_B \right\|_{\Phi, Q_j^k} = I + II.$$

This inequality implies that either $I > 1/2$ or $II > 1/2$. Since $\Phi \in \mathfrak{F}_{r, \delta}$ we can easily see that $I > 1/2$ implies that

$$a^{kr} < \frac{2^r C_0 (\log(2t^*))^\delta}{|Q_j^k|} \int_{Q_j^k} \Phi(|f|) v^r \leq \frac{2^r C_0 (\log(2t^*))^\delta}{|Q_j^k|} \int_{Q_j^k} (\eta_\varepsilon \circ \Phi)(|f|) v^r,$$

because $\eta_\varepsilon(z) \geq z$.

On the other hand, if $II > 1/2$ then again

$$\begin{aligned} 1 &< \frac{1}{|Q_j^k|} \int_B \Phi\left(\frac{2|f|v}{a^k}\right) \\ &\leq \frac{\Phi(2)C_0}{|Q_j^k|} \int_B \Phi(|f|) \frac{v^r}{a^{kr}} \left(\log\left(\frac{v}{a^k}\right)\right)^\delta, \end{aligned}$$

since $\Phi \in \mathfrak{F}_{r, \delta}$. This implies that

$$a^{kr} \leq \frac{\Phi(2)C_0}{|Q_j^k|} \int_{Q_j^k} \Phi(|f|) v^r w_k,$$

where $w_k(x) = \left(\log\left(\frac{v(x)}{a^k}\right)\right)^\delta \chi_B(x)$. We shall now perform a generalized Hölder inequality with the Young functions

$$\eta_\varepsilon(z) = z(1 + \log^+ z)^{\delta/\varepsilon} \quad \text{and} \quad \tilde{\eta}_\varepsilon(z) \approx (e^{z^{\varepsilon/\delta}} - e) \chi_{(1, \infty)}(z),$$

with respect to the measure $d\mu(x) = v^r(x) dx$. Thus we have

$$(1.10) \quad \frac{1}{|Q_j^k|} \int_{Q_j^k} \Phi(|f|) w_k v^r \leq \frac{v^r(Q_j^k)}{|Q_j^k|} \|\Phi(|f|)\|_{\eta_\varepsilon, Q_j^k, v^r} \|w_k\|_{\tilde{\eta}_\varepsilon, Q_j^k, v^r}.$$

Let us first estimate the last factor. Since $e^{(\log z)^\varepsilon} \leq z^\varepsilon$ when $z \geq e^{1/(\varepsilon-1)}$, we proceed as follows

$$(1.11) \quad \frac{1}{v^r(Q_j^k)} \int_{Q_j^k} \tilde{\eta}_\varepsilon(w_k) v^r \leq \tilde{\eta}_\varepsilon\left(\varepsilon^{\delta/(\varepsilon-1)}\right) + \frac{1}{v^r(Q_j^k)} \int_{Q_j^k \cap \{v/a^k > e^{1/(\varepsilon-1)}\}} \frac{v^{r+\varepsilon}}{a^{k\varepsilon}}.$$

Since $\varepsilon^{-\varepsilon} \leq e$ we have that

$$\tilde{\eta}_\varepsilon\left(\varepsilon^{\delta/(\varepsilon-1)}\right) \leq e^{e^{1/(1-\varepsilon)}}.$$

On the other hand, our hypothesis on v implies that $v^r \in \text{RH}_s$, where $s = 1 + \varepsilon/r$. Since $v^r \in \Lambda_{\ell,k}$ we obtain

$$\begin{aligned} \frac{1}{v^r(Q_j^k)} \int_{Q_j^k \cap \{v/a^k > e^{\varepsilon^{1/(\varepsilon-1)}}\}} \frac{v^{r+\varepsilon}}{a^{k\varepsilon}} &\leq \frac{a^{-k\varepsilon}}{v^r(Q_j^k)} \int_{Q_j^k} v^{r+\varepsilon} \\ &\leq [v^r]_{\text{RH}_s}^s \frac{a^{-k\varepsilon} |Q_j^k|}{v^r(Q_j^k)} \left(\frac{1}{|Q_j^k|} \int_{Q_j^k} v^r \right)^s \\ &= [v^r]_{\text{RH}_s}^s a^{-k\varepsilon} \left(\frac{1}{|Q_j^k|} \int_{Q_j^k} v^r \right)^{s-1} \\ &\leq [v^r]_{\text{RH}_s}^s a^{-k\varepsilon} a^{(k+\ell+1)\varepsilon} \\ &= [v^r]_{\text{RH}_s}^s a^{(\ell+1)\varepsilon}. \end{aligned}$$

By using these two estimates in (1.11), we get

$$\|w_k\|_{\tilde{\eta}_\varepsilon, Q_j^k, v^r} \leq e^{\varepsilon^{1/(1-\varepsilon)}} + [v^r]_{\text{RH}_s}^s a^{(\ell+1)\varepsilon} \leq (e^{\varepsilon^{1/(1-\varepsilon)}} + [v^r]_{\text{RH}_s}^s) a^{(\ell+1)\varepsilon}.$$

We also observe that

$$(1.12) \quad \|\Phi(|f|)\|_{\eta_\varepsilon, Q_j^k, v^r} \approx \inf_{\tau > 0} \left\{ \tau + \frac{\tau}{v^r(Q_j^k)} \int_{Q_j^k} \eta_\varepsilon \left(\frac{\Phi(|f|)}{\tau} \right) v^r \right\}.$$

If we choose $\tau = (2a^{(\ell+1)(r+\varepsilon)}(e^{\varepsilon^{1/(1-\varepsilon)}} + [v^r]_{\text{RH}_s}^s))^{-1}$ then we can estimate the right-hand side of (1.10) as follows

$$\frac{v^r(Q_j^k)}{|Q_j^k|} \|\Phi(|f|)\|_{\eta_\varepsilon, Q_j^k, v^r} \|w_k\|_{\tilde{\eta}_\varepsilon, Q_j^k, v^r} \leq \frac{a^{kr}}{2} + (e^{\varepsilon^{1/(1-\varepsilon)}} + [v^r]_{\text{RH}_s}^s) a^{(\ell+1)\varepsilon} \tau \eta_\varepsilon \left(\frac{1}{\tau} \right) \frac{1}{|Q_j^k|} \int_{Q_j^k} \Psi_\varepsilon(|f|) v^r.$$

Notice that

$$\tau \eta_\varepsilon \left(\frac{1}{\tau} \right) = \left(1 + \log \left(\frac{1}{\tau} \right) \right)^{\delta/\varepsilon} \leq 2^{\delta/\varepsilon} \left(\log(2(e^{\varepsilon^{1/(1-\varepsilon)}} + [v^r]_{\text{RH}_s}^s)) + (\ell+1)(r+\varepsilon) \log a \right)^{\delta/\varepsilon} \leq C_\varepsilon \ell^{\delta/\varepsilon}.$$

By plugging these two estimates in (1.10) we arrive to

$$a^{kr} \leq \frac{C_\varepsilon \ell^{\delta/\varepsilon} a^{\ell\varepsilon}}{|Q_j^k|} \int_{Q_j^k} \Psi_\varepsilon(|f|) v^r,$$

and we are done. \square

Proof of Claim 2. The proof follows similar arguments as the previous one. By adopting the same notation, we have that $\|\frac{a}{a^k}\|_{\Phi, Q_j^k} > 1$, and this implies that either $I > 1/2$ or $II > 1/2$. If $I > 1/2$, we obtain

$$a^{kr} < \frac{C_0(\log(2t^*))^\delta}{|Q_j^k|} \int_{Q_j^k} \Phi(|f|) v^r \leq \frac{C_0(\log(2t^*))^\delta}{|Q_j^k|} \int_{Q_j^k} \Psi_\varepsilon(|f|) v^r.$$

We now assume that $II > 1/2$. By performing the same Hölder inequality as in Claim 1, we get

$$(1.13) \quad \frac{1}{|Q_j^k|} \int_{Q_j^k} \Phi(|f|) w_k v^r \leq \frac{v^r(Q_j^k)}{|Q_j^k|} \|\Phi(|f|)\|_{\eta_\varepsilon, Q_j^k, v^r} \|w_k\|_{\tilde{\eta}_\varepsilon, Q_j^k, v^r}.$$

In order to estimate the factor $\|w_k\|_{\tilde{\eta}_\varepsilon, Q_j^k, v^r}$ we proceed as before. Since

$$\begin{aligned} \frac{1}{v^r(Q_j^k)} \int_{Q_j^k \cap \{v/a^k > e^{\varepsilon-1/(1-\varepsilon)}\}} \frac{v^{r+\varepsilon}}{a^{k\varepsilon}} &\leq \frac{a^{-k\varepsilon}}{v^r(Q_j^k)} \int_{Q_j^k} v^{r+\varepsilon} \\ &\leq [v^r]_{\text{RH}_s}^s \frac{a^{-k\varepsilon}|Q_j^k|}{v^r(Q_j^k)} \left(\frac{1}{|Q_j^k|} \int_{Q_j^k} v^r \right)^s \\ &= [v^r]_{\text{RH}_s}^s a^{-k\varepsilon} \left(\frac{1}{|Q_j^k|} \int_{Q_j^k} v^r \right)^{s-1} \\ &\leq [v^r]_{\text{RH}_s}^s a^{-k\varepsilon} a^{k\varepsilon} \\ &= [v^r]_{\text{RH}_s}^s \end{aligned}$$

we obtain that

$$\|w_k\|_{\tilde{\eta}_\varepsilon, Q_j^k, v^r} \leq e^{e^{1/(1-\varepsilon)}} + [v^r]_{\text{RH}_s}^s = C_\varepsilon.$$

By using (1.12) and choosing $\tau = (2C_\varepsilon)^{-1}$, we can estimate the right-hand side in (1.13) as follows

$$\begin{aligned} \frac{v^r(Q_j^k)}{|Q_j^k|} \|\Phi(|f|)\|_{\eta_\varepsilon, Q_j^k, v^r} \|w_k\|_{\tilde{\eta}_\varepsilon, Q_j^k, v^r} &\leq \frac{a^{kr}}{2} + \tau \eta_\varepsilon \left(\frac{1}{\tau} \right) \frac{1}{|Q_j^k|} \int_{Q_j^k} \Psi_\varepsilon(|f|) v^r \\ &\leq \frac{a^{kr}}{2} + (1 + \log(2C_\varepsilon))^{\delta/\varepsilon} \frac{1}{|Q_j^k|} \int_{Q_j^k} \Psi_\varepsilon(|f|) v^r. \end{aligned}$$

This yields

$$a^{kr} \leq \frac{C_\varepsilon}{|Q_j^k|} \int_{Q_j^k} \Psi_\varepsilon(|f|) v^r.$$

This concludes the proof. \square

2. APPLICATIONS: MIXED ESTIMATES FOR THE GENERALIZED FRACTIONAL INTEGRAL OPERATOR

Mixed inequalities for the generalized fractional maximal operator $M_{\gamma, \Phi}$ were also given in [1]. One of the key properties in order to establish the following result was to define an auxiliary operator that is bounded in $L^\infty(wv^r)$ when $v^r \in A_\infty$. This operator is given by

$$\mathcal{T}_\Phi f(x) = \frac{M_\Phi(fv)(x)}{M_\Phi v(x)}.$$

It is not difficult to see that $M_\Phi v \approx v$ when $\Phi \in \mathfrak{F}_{r, \delta}$ and v^r is an A_1 -weight, so this operator is an extension of the Sawyer operator $S_\Phi f = M_\Phi(fv)/v$ considered in the main theorem.

Corollary 5 (Corrected version of Corollary 3 in [1]). *Let $r \geq 1$, $\delta \geq 0$ and $\Phi \in \mathfrak{F}_{r, \delta}$. Let $u \in A_1$, $v^r \in A_\infty$ and Ψ be a Young function that verifies $\Psi(t) \approx \Phi(t)$, for every $t \geq t_0 \geq 0$. Then, there exists $\varepsilon_0 > 0$ such that the inequality*

$$v^r \left(\left\{ x \in \mathbb{R}^n : \frac{M_\Psi(fv)(x)}{M_\Psi v(x)} > t \right\} \right) \leq C_1 \int_{\mathbb{R}^n} (\eta_\varepsilon \circ \Psi) \left(\frac{C_2 |f(x)|}{t} \right) u(x) v^r(x) dx$$

holds for every $t > 0$ and every $0 < \varepsilon < \varepsilon_0$, where C_1 depends on ε and $\eta_\varepsilon(z) = z(1 + \log^+ z)^{\delta/\varepsilon}$.

Proof. By combining the equivalence between Φ and Ψ and Proposition 8 in [1], we obtain that there exist positive constants A and B such that

$$A\|\cdot\|_{\Phi,Q} \leq \|\cdot\|_{\Psi,Q} \leq B\|\cdot\|_{\Phi,Q},$$

for every cube Q . By setting $c_1 = B/A$, we have that

$$\frac{M_{\Psi}(fv)(x)}{M_{\Psi}v(x)} \leq c_1 \frac{M_{\Phi}(fv)(x)}{M_{\Phi}v(x)}$$

for almost every x . By applying Corollary 2, there exists $\varepsilon_0 > 0$ such that for every $0 < \varepsilon < \varepsilon_0$ we have

$$\begin{aligned} uv^r \left(\left\{ x \in \mathbb{R}^n : \frac{M_{\Psi}(fv)(x)}{M_{\Psi}v(x)} > t \right\} \right) &\leq uv^r \left(\left\{ x \in \mathbb{R}^n : \frac{M_{\Phi}(fv)(x)}{M_{\Phi}v(x)} > \frac{t}{c_1} \right\} \right) \\ &\leq C \int_{\mathbb{R}^n} (\eta_{\varepsilon} \circ \Phi) \left(\frac{c_1|f|}{t} \right) uv^r. \end{aligned}$$

Observe that

$$\|\mathcal{T}_{\Psi}f\|_{L^{\infty}} = \left\| \frac{M_{\Psi}(fv)}{M_{\Psi}v} \right\|_{L^{\infty}} \leq \|f\|_{L^{\infty}},$$

which directly implies $\|\mathcal{T}_{\Psi}f\|_{L^{\infty}(uv^r)} \leq \|f\|_{L^{\infty}(uv^r)}$ since the measure given by $d\mu(x) = u(x)v^r(x) dx$ is absolutely continuous with respect to the Lebesgue measure. We now apply Lemma 13 in [1] to obtain

$$\begin{aligned} uv^r \left(\left\{ x \in \mathbb{R}^n : \frac{M_{\Psi}(fv)(x)}{M_{\Psi}v(x)} > t \right\} \right) &\leq C \int_{\{x:|f(x)|>t/2\}} (\eta_{\varepsilon} \circ \Phi) \left(\frac{2c_1|f(x)|}{t} \right) u(x)v^r(x) dx \\ &\leq C(\eta_{\varepsilon} \circ \Phi) \left(\frac{c_1}{t_0} \right) \int_{\{x:|f(x)|>t/2\}} (\eta_{\varepsilon} \circ \Phi) \left(\frac{2t_0|f(x)|}{t} \right) u(x)v^r(x) dx \\ &\leq C_1 \int_{\{x:|f(x)|>t/2\}} (\eta_{\varepsilon} \circ \Psi) \left(\frac{2t_0|f(x)|}{t} \right) u(x)v^r(x) dx \\ &\leq C_1 \int_{\mathbb{R}^n} (\eta_{\varepsilon} \circ \Psi) \left(\frac{C_2|f(x)|}{t} \right) u(x)v^r(x) dx. \quad \square \end{aligned}$$

The corollary above is key in order to obtain mixed inequalities for the generalized fractional maximal operator defined, for $0 < \gamma < n$ and a Young function Φ , by the expression

$$M_{\gamma,\Phi}f(x) = \sup_{Q \ni x} |Q|^{\gamma/n} \|f\|_{\Phi,Q}.$$

Mixed estimates for this operator are contained in the following theorems.

Theorem 6. *Let $\Phi(z) = z^r(1 + \log^+ z)^{\delta}$, with $r \geq 1$ and $\delta \geq 0$. Let $0 < \gamma < n/r$, $r < p < n/\gamma$ and $1/q = 1/p - \gamma/n$. If $u \in A_1$ and $v^{q(1/p+1/r')} \in A_{\infty}$, then the inequality*

$$uv^{q(1/p+1/r')} \left(\left\{ x \in \mathbb{R}^n : \frac{M_{\gamma,\Phi}(fv)(x)}{M_{\varphi}v(x)} > t \right\} \right)^{1/q} \leq C \left[\int_{\mathbb{R}^n} \left(\frac{|f(x)|}{t} \right)^p u^{p/q}(x)(v(x))^{1+p/r'} dx \right]^{1/p},$$

holds for every positive t , where $\varphi(z) = z^{q/p+q/r'}(1 + \log^+ z)^{n\delta/(n-r\gamma)}$.

Proof. We shall follow the scheme given in the proof of Theorem 4 in [1]. The difference lies when we apply Corollary 5, but the function controlling the right-hand side turns out to be auxiliary. We define

$$\sigma = \frac{nr}{n - r\gamma}, \quad \nu = \frac{n\delta}{n - r\gamma}, \quad \beta = \frac{q}{\sigma} \left(\frac{1}{p} + \frac{1}{r'} \right),$$

and let ξ be the auxiliary function given by

$$\xi(z) = \begin{cases} z^{q/\beta}, & \text{if } 0 \leq z \leq 1, \\ z^\sigma (1 + \log^+ z)^\nu, & \text{if } z > 1. \end{cases}$$

Observe that

$$\xi^{-1}(z)z^{\gamma/n} \approx \frac{z^{1/\sigma + \gamma/n}}{(1 + \log^+ z)^{\nu/\sigma}} = \frac{z^{1/r}}{(1 + \log^+ z)^{\delta/r}} \lesssim \Phi^{-1}(z),$$

for every $z \geq 1$. Observe that $\beta > 1$: indeed, since $p > r$ we have $q > \sigma$ and thus $q/(\sigma r') > 1/r'$. On the other hand, $q/(p\sigma) > 1/r$. By combining these two inequalities we have $\beta > 1$. Applying Proposition 10 and Lemma 9 with β from [1], we can conclude that

$$(2.1) \quad M_{\gamma, \Phi} \left(\frac{f_0}{w} \right) (x) \leq C \left[M_\xi \left(\frac{f_0^{p\beta/q}}{w^\beta} \right) (x) \right]^{1/\beta} \left(\int_{\mathbb{R}^n} f_0^p(y) dy \right)^{\gamma/n}.$$

Also observe that

$$(2.2) \quad \left(M_\xi v^\beta(x) \right)^{1/\beta} \lesssim M_\varphi v(x), \quad \text{a.e. } x.$$

Notice that ξ is equivalent to a Young function in $\mathfrak{F}_{\sigma, \nu}$ for $t \geq 1$. Since $q(1/p + 1/r') = \beta\sigma$, if we set $f_0 = |f|wv$, then we can use inequalities (2.1) and (2.2) to estimate

$$\begin{aligned} uv^{\frac{q}{p} + \frac{q}{r'}} \left(\left\{ x : \frac{M_{\gamma, \Phi}(fv)(x)}{M_\varphi v(x)} > t \right\} \right) &\lesssim uv^{\beta\sigma} \left(\left\{ x : \frac{M_{\gamma, \Phi}(fv)(x)}{(M_\xi v^\beta(x))^{1/\beta}} > t \right\} \right) \\ &\leq uv^{\beta\sigma} \left(\left\{ x : \frac{M_\xi \left(f_0^{p\beta/q} w^{-\beta} \right) (x)}{M_\xi v^\beta(x)} > \frac{t^\beta}{(f|f_0|^p)^{\beta\gamma/n}} \right\} \right). \end{aligned}$$

Since $v^{\beta\sigma} \in A_\infty$, by Corollary 5 there exists $\varepsilon_0 > 0$ such that the inequality

$$uv^{\beta\sigma} \left(\left\{ x : \frac{M_\xi \left(f_0^{p\beta/q} w^{-\beta} \right) (x)}{M_\xi v^\beta(x)} > t_0 \right\} \right) \leq C_\varepsilon \int_{\mathbb{R}^n} (\eta_\varepsilon \circ \xi) \left(c \frac{f_0^{p\beta/q} w^{-q}}{t_0} \right) uv^{\beta\sigma}$$

holds for every $0 < \varepsilon < \varepsilon_0$, with $t_0 = t^\beta \|f_0\|_p^{-p\beta\gamma/n}$. Notice that $\eta_\varepsilon(z) = z(1 + \log^+ z)^{\nu/\varepsilon}$ in this case.

Fixed ε , we write

$$\int_{\mathbb{R}^n} (\eta_\varepsilon \circ \xi) \left(c \frac{|f|^{p\beta/q} (wv)^{\beta(p/q-1)}}{t^\beta} \left[\int_{\mathbb{R}^n} |f|^p(wv)^p \right]^{\gamma/n\beta} \right) uv^{\beta\sigma} = \int_{\mathbb{R}^n} (\eta_\varepsilon \circ \xi)(\lambda) uv^{\beta\sigma},$$

where

$$\lambda = c \frac{|f|^{p\beta/q} (wv)^{\beta(p/q-1)}}{t^\beta} \left[\int_{\mathbb{R}^n} |f|^p(wv)^p \right]^{\gamma/n\beta}.$$

We further split \mathbb{R}^n into the sets $A = \{x \in \mathbb{R}^n : \lambda(x) \leq 1\}$ and $B = \mathbb{R}^n \setminus A$. Since $(\eta_\varepsilon \circ \xi)(z) = z^{q/\beta}$ for $0 \leq z \leq 1$, we have that

$$\int_A (\eta_\varepsilon \circ \xi)(\lambda(x)) u(x) [v(x)]^{\sigma\beta} dx = \int_A [\lambda(x)]^{q/\beta} u(x) [v(x)]^{\sigma\beta} dx.$$

If we set $w = u^{1/q} v^{1/p+1/r'-1}$, then

$$\begin{aligned} \lambda^{q/\beta} u v^{\sigma\beta} &= c^{q/\beta} \frac{|f|^p}{t^q} (wv)^{p-q} \left[\int_{\mathbb{R}^n} |f|^p (wv)^p \right]^{q\gamma/n} w v^{\sigma\beta} \\ &= c^{q/\beta} \frac{|f|^p}{t^q} \left[\int_{\mathbb{R}^n} |f|^p (wv)^p \right]^{q\gamma/n} u^{p/q} v^{\sigma\beta+(p-q)(1/p+1/r')}. \end{aligned}$$

Observe that

$$\sigma\beta + (p-q) \left(\frac{1}{p} + \frac{1}{r'} \right) = q \left(\frac{1}{p} + \frac{1}{r'} \right) + (p-q) \left(\frac{1}{p} + \frac{1}{r'} \right) = 1 + \frac{p}{r'}.$$

Also, notice that

$$(wv)^p = u^{p/q} v^{1+p/r'-p+p} = u^{p/q} v^{1+p/r'}.$$

Therefore,

$$\begin{aligned} \int_A (\eta_\varepsilon \circ \xi)(\lambda) u v^{\sigma\beta} &\leq \frac{c^{q/\beta}}{t^q} \left[\int_{\mathbb{R}^n} |f|^p u^{p/q} v^{1+p/r'} \right]^{q\gamma/n} \left[\int_{\mathbb{R}^n} |f|^p u^{p/q} v^{1+p/r'} \right] \\ &= \frac{c^{q/\beta}}{t^q} \left[\int_{\mathbb{R}^n} |f|^p u^{p/q} v^{1+p/r'} \right]^{1+q\gamma/n} \\ &= \frac{c^{q/\beta}}{t^q} \left[\int_{\mathbb{R}^n} |f|^p u^{p/q} v^{1+p/r'} \right]^{q/p}. \end{aligned}$$

On the other hand, $\lambda(x) > 1$ over B and

$$(\eta_\varepsilon \circ \xi)(z) \lesssim z^\sigma (1 + \log z)^{\nu(1+1/\varepsilon)},$$

and this function has an upper type q/β . Therefore we can estimate the integrand by $\lambda^{q/\beta} u v^{\sigma\beta}$ and proceed as we did with the set A . Thus, we obtain

$$u v^{q(1/p+1/r')} \left(\left\{ x \in \mathbb{R}^n : \frac{M_{\gamma, \Phi}(fv)(x)}{M_\varphi v(x)} > t \right\} \right)^{1/q} \leq C \left[\int_{\mathbb{R}^n} \left(\frac{|f|}{t} \right)^p u^{p/q} v^{1+p/r'} \right]^{1/p}. \quad \square$$

Theorem 7 (Corrected version of Theorem 5 in [1]). *Let $\Phi(z) = z^r (1 + \log^+ z)^\delta$, with $r \geq 1$ and $\delta \geq 0$. Let $0 < \gamma < n/r$ and $1/q = 1/r - \gamma/n$. If $u \in A_1$ and $v^q \in A_\infty$, then there exists a positive constant ε_0 such that the inequality*

$$u v^q \left(\left\{ x \in \mathbb{R}^n : \frac{M_{\gamma, \Phi}(fv)(x)}{v(x)} > t \right\} \right) \leq C \varphi_\varepsilon \left(\int_{\mathbb{R}^n} \Phi_{\gamma, \varepsilon} \left(\frac{|f(x)|}{t} \right) \Psi \left(u^{1/q}(x) v(x) \right) dx \right),$$

holds for $0 < \varepsilon < \varepsilon_0$, where $\varphi_\varepsilon(z) = [z(1 + \log^+ z)^{\delta(1+1/\varepsilon)}]^{q/r}$, $\Psi_\varepsilon(z) = z^r (1 + \log^+(z^{1-q/r}))^{q\delta(1+1/\varepsilon)/r}$, $\Phi_{\gamma, \varepsilon}(z) = \Phi(z) (1 + \log^+ z)^{\delta(1+1/\varepsilon)q\gamma/n + \delta/\varepsilon}$ and C depends on ε .

Proof. Set $\xi(z) = z^q (1 + \log^+ z)^\nu$, where $\nu = \delta q/r$. Thus $z^{\gamma/n} \xi^{-1}(z) \lesssim \Phi^{-1}(z)$. By applying Proposition 10 in [1] with $p = r$ we have that

$$M_{\gamma, \Phi} \left(\frac{f_0}{w} \right) (x) \leq C \left[M_\xi \left(\frac{f_0^{r/q}}{w} \right) \right] (x) \left(\int_{\mathbb{R}^n} f_0^r(y) dy \right)^{\gamma/n}.$$

By setting $f_0 = |f|wv$ we can write

$$\begin{aligned} uv^q \left(\left\{ x : \frac{M_{\gamma, \Phi}(fv)(x)}{v(x)} > t \right\} \right) &= uv^q \left(\left\{ x : \frac{M_{\gamma, \Phi}(f_0/w)(x)}{v(x)} > t \right\} \right) \\ &\leq uv^q \left(\left\{ x : \frac{M_{\xi}(f_0^{r/q}/w)(x)}{M_{\xi}v(x)} > \frac{t}{(\int f_0^r)^{\gamma/n}} \right\} \right). \end{aligned}$$

Since $\xi \in \mathfrak{F}_{q, \nu}$, by Corollary 2 there exists $\varepsilon_0 > 0$ such that

$$(2.3) \quad uv^q \left(\left\{ x : \frac{M_{\gamma, \Phi}(fv)(x)}{v(x)} > t \right\} \right) \leq C_{\varepsilon} \int_{\mathbb{R}^n} (\eta_{\varepsilon} \circ \xi) \left(\frac{f_0^{r/q} (\int f_0^r)^{\gamma/n}}{wvt} \right) uv^q,$$

for $0 < \varepsilon < \varepsilon_0$ and being $\eta_{\varepsilon}(z) = z(1 + \log^+ z)^{\nu/\varepsilon}$. Fixed ε , the argument of $\eta_{\varepsilon} \circ \xi$ above can be written as

$$\begin{aligned} \frac{f_0^{r/q} (\int f_0^r)^{\gamma/n}}{wvt} &= \left(\frac{|f|}{t} \right)^{r/q} (wv)^{r/q-1} \left(\int_{\mathbb{R}^n} \left(\frac{|f|}{t} \right)^r (wv)^r \right)^{\gamma/n} \\ &= \left[\left(\frac{|f|}{t} \right) (wv)^{1-q/r} \left(\int_{\mathbb{R}^n} \left(\frac{|f|}{t} \right)^r (wv)^r \right)^{\gamma q/(nr)} \right]^{r/q}. \end{aligned}$$

Observe that for $0 \leq z \leq 1$, $(\eta_{\varepsilon} \circ \xi)(z^{r/q}) = z^r$, and for $z > 1$ we have

$$(\eta_{\varepsilon} \circ \xi)(z^{r/q}) \lesssim z^r (1 + \log z)^{\nu(1+1/\varepsilon)},$$

which implies that $(\eta_{\varepsilon} \circ \xi)(z^{r/q}) \lesssim \Phi_{\gamma, \varepsilon}(z) = z^r (1 + \log^+ z)^{\nu(1+1/\varepsilon)}$, for every $z \geq 0$. Since $\Phi_{\gamma, \varepsilon}$ is submultiplicative, we can estimate as follows

$$\begin{aligned} (\eta_{\varepsilon} \circ \xi) \left(\frac{f_0^{r/q} (\int_{\mathbb{R}^n} f_0^r)^{\gamma/n}}{wvt} \right) &\leq \Phi_{\gamma, \varepsilon} \left(\left(\frac{|f|}{t} \right) (wv)^{1-q/r} \left(\int_{\mathbb{R}^n} \left(\frac{|f|}{t} \right)^r (wv)^r \right)^{\gamma q/(nr)} \right) \\ &\leq \Phi_{\gamma, \varepsilon} \left(\left[\int_{\mathbb{R}^n} \Phi_{\gamma, \varepsilon} \left(\frac{|f|}{t} \right) (wv)^r \right]^{\gamma q/(nr)} \right) \Phi_{\gamma, \varepsilon} \left(\frac{|f|}{t} (wv)^{1-q/r} \right) \end{aligned}$$

Returning to (2.3) and setting $w = u^{1/q}$, the right hand side is bounded by

$$\Phi_{\gamma, \varepsilon} \left(\left[\int_{\mathbb{R}^n} \Phi_{\gamma, \varepsilon} \left(\frac{|f|}{t} \right) (wv)^r \right]^{\gamma q/(nr)} \right) \int_{\mathbb{R}^n} \Phi_{\gamma, \varepsilon} \left(\frac{|f|}{t} (wv)^{1-q/r} \right) (wv)^q.$$

Notice that $\Phi_{\gamma, \varepsilon}(z^{1-q/r})z^q \leq \Psi_{\varepsilon}(z)$. Therefore, the expression above is bounded by

$$\Phi_{\gamma, \varepsilon} \left(\left[\int_{\mathbb{R}^n} \Phi_{\gamma, \varepsilon} \left(\frac{|f|}{t} \right) \Psi_{\varepsilon}(u^{1/q}v) \right]^{\gamma q/(nr)} \right) \int_{\mathbb{R}^n} \Phi_{\gamma, \varepsilon} \left(\frac{|f|}{t} \right) \Psi_{\varepsilon}(u^{1/q}v).$$

To finish, observe that

$$z \Phi_{\gamma, \varepsilon}(z^{\gamma q/(nr)}) \lesssim z^{1+\gamma q/n} (1 + \log^+ z)^{\nu(1+1/\varepsilon)} = z^{q/r} (1 + \log^+ z)^{\delta q(1+1/\varepsilon)/r} = \varphi_{\varepsilon}(z). \quad \square$$

REFERENCES

- [1] Fabio Berra, *From A_1 to A_∞ : new mixed inequalities for certain maximal operators*, Potential Anal. **57** (2022), no. 1, 1–27.
- [2] Fabio Berra, Marilina Carena, and Gladis Pradolini, *Improvements on Sawyer type estimates for generalized maximal functions*, Math. Nachr. **293** (2020), no. 10, 1911–1930.
- [3] Tuomas Hytönen and Carlos Pérez, *Sharp weighted bounds involving A_∞* , Anal. PDE **6** (2013), no. 4, 777–818.
- [4] K. Li, S. Ombrosi, and C. Pérez, *Proof of an extension of E. Sawyer’s conjecture about weighted mixed weak-type estimates*, Math. Ann. **374** (2019), no. 1-2, 907–929.

CONICET AND DEPARTAMENTO DE MATEMÁTICA (FIQ-UNL), SANTA FE, ARGENTINA.

Email address: `fberra@santafe-conicet.gov.ar`