

AN IMPROVEMENT OF THE FARRELL-JONES CONJECTURE FOR LOCALISING INVARIANTS

JOSÉ FRANCISCO REIS

ABSTRACT. The Farrell-Jones conjecture for lax monoidal finitary localising invariants was recently proved by Bunke-Kasprowski-Winges. In this short note, making use of the theory of noncommutative motives, we prove that the lax monoidal assumption is not necessary.

1. INTRODUCTION AND STATEMENT OF RESULTS

Let G be a group, $G\text{Orb}$ the orbit category¹ of G , \mathcal{F} a family of subgroups of G , $G_{\mathcal{F}}\text{Orb}$ the full subcategory of $G\text{Orb}$ of those left G -sets with stabilizers in \mathcal{F} , $\text{Cat}_{\infty,*}^{\text{Lex}}$ the (large) ∞ -category of small left-exact ∞ -categories² and finite limit preserving functors, and $H : \text{Cat}_{\infty,*}^{\text{Lex}} \rightarrow M$ a functor with values in a cocomplete stable ∞ -category. Also, let C be a small left-exact ∞ -category equipped with a G -action, i.e., an object of $\text{Fun}(BG, \text{Cat}_{\infty,*}^{\text{Lex}})$ where BG stands for the groupoid with one object and group of automorphism G . Note that we have a canonical inclusion $j^G : BG \rightarrow G\text{Orb}$ sending the unique object of BG to the transitive G -set G . The left Kan-extension of C along j^G will be denoted by C_G . Finally, note that for every subgroup H of G , we have an equivalence $C_G(G/H) \simeq \text{colim}_{BH} \text{Res}_H^G(C)$.

In a recent article [4], Bunke, Kasprowski and Wings proved the following beautiful result:

Theorem 1.1 ([4, Theorem 1.4]). *Assume that G is a Dress-Farrell-Hsiang-Jones (DFHJ) group relative to \mathcal{F} (consult [4, Definition 7.1]) and that H is a lax monoidal finitary localising invariant (consult [4, Definition 1.2]). Under these assumptions, the assembly map*

$$(1.1) \quad A_{\mathcal{F}, H \circ C_G} : \text{colim}_{G_{\mathcal{F}}\text{Orb}}(H \circ C_G) \longrightarrow H(\text{colim}_{BG} C)$$

is a phantom equivalence (consult [4, Definition 2.5]).

Theorem 1.1 proves the Farrell-Jones conjecture for a very large class of groups G , of functors H , and of coefficients C . The classical Farrell-Jones conjecture [5] corresponds to the particular case when \mathcal{F} is the family of virtually cyclic groups, H is the equivariant K -theory functor, and C is an additive category equipped with a G -action. In this short note, making use of the theory of noncommutative motives, we improve Theorem 1.1 as follows:

Theorem 1.2. *Theorem 1.1 holds ipso facto with H a finitary localising invariant (the lax monoidal assumption is not necessary).*

Remark 1.3. The proof of Theorem 1.2 makes essential use of Theorem 1.1.

2. PROOF OF THEOREM 1.2

Let $\text{Cat}_{\infty,*}^{\text{ex}}$ be the (large) ∞ -category of small stable ∞ -categories, $\text{stab} : \text{Cat}_{\infty,*}^{\text{Lex}} \rightarrow \text{Cat}_{\infty,*}^{\text{ex}}$ the stabilization functor and $\text{inc} : \text{Cat}_{\infty,*}^{\text{ex}} \rightarrow \text{Cat}_{\infty,*}^{\text{Lex}}$ the inclusion functor; consult [3, §7.4]. As proved in [3, Lemma 7.41], we have the following adjunction:

¹Recall that the objects of $G\text{Orb}$ are the left G -sets G/H , with H a subgroup of G , and the morphisms are the maps of G -sets $G/H \rightarrow G/H'$.

²Recall that a (small) ∞ -category is called *left-exact* if it is pointed and admits all finite limits.

$$\begin{array}{c} \text{Cat}_{\infty,*}^{\text{Lex}} \\ \text{stab} \downarrow \uparrow \text{inc} \\ \text{Cat}_{\infty,*}^{\text{ex}} \end{array}$$

Moreover, as explained in [3, §6.2], the composition $H \circ \text{inc}$ is a localising invariant in the sense of [1]. Furthermore, as proved in [3, Lemma 6.9(1)], the functors H and $H \circ \text{inc} \circ \text{stab}$ are naturally equivalent. Note that in [3, 4] a localising invariant is referred to as a stable finitary localising invariant; in what follows, we stick to this latter terminology.

Recall from [1, §8] the construction of the universal stable finitary localising invariant $\mathcal{U}_{\text{loc}} : \text{Cat}_{\infty,*}^{\text{ex}} \rightarrow \mathcal{M}_{\text{loc}}$; due to this universal property, \mathcal{M}_{loc} is called the (large) ∞ -category of noncommutative motives; consult [6]. Since the functor $H \circ \text{inc}$ is a stable finitary localising invariant, [1, Theorem 8.7] implies that it factors through \mathcal{U}_{loc} ; let us denote by $\overline{H \circ \text{inc}}$ the induced functor. Under these notations, we have the following commutative diagram (up to equivalence):

$$(2.1) \quad \begin{array}{ccccc} & & G\text{Orb} & \xrightarrow{H \circ C_G} & M \\ & & \uparrow j^G & \searrow^{C_G} & \\ BG & \xrightarrow{C} & \text{Cat}_{\infty,*}^{\text{Lex}} & \xrightarrow{H} & M \\ & & \text{stab} \downarrow \uparrow \text{inc} & \nearrow^{H \circ \text{inc}} & \\ & & \text{Cat}_{\infty,*}^{\text{ex}} & & \\ & & \mathcal{U}_{\text{loc}} \downarrow & \searrow^{\overline{H \circ \text{inc}}} & \\ & & \mathcal{M}_{\text{loc}} & & \end{array}$$

Since \mathcal{U}_{loc} is a stable finitary localising invariant, [3, Lemma 6.9(2)] implies that the composition $\mathcal{U}_{\text{loc}} \circ \text{stab}$ is a finitary localising. Moreover, as explained in [2, Theorem 5.8] and [4, §1], respectively, the functors \mathcal{U}_{loc} and stab are symmetric monoidal. This implies that the composition $\mathcal{U}_{\text{loc}} \circ \text{stab}$ is a lax monoidal finitary localising invariant. Consequently, Theorem 1.1 (with H replaced by $\mathcal{U}_{\text{loc}} \circ \text{stab}$) implies that the assembly map

$$(2.2) \quad A_{\mathcal{F},(\mathcal{U}_{\text{loc}} \circ \text{stab} \circ C_G)} : \text{colim}_{G\mathcal{F}\text{Orb}}(\mathcal{U}_{\text{loc}} \circ \text{stab} \circ C_G) \longrightarrow (\mathcal{U}_{\text{loc}} \circ \text{stab})(\text{colim}_{BG} C)$$

is a phantom equivalence. Let us denote by cof the cofiber of (2.2). As explained in the proofs of [4, Theorem 1.4 and Proposition 2.31], the diagonal map $\Delta : \text{cof} \rightarrow \Pi_{\mathbb{N}}\text{cof}$ factors through the canonical map $\oplus_{\mathbb{N}}\text{cof} \rightarrow \Pi_{\mathbb{N}}\text{cof}$. Therefore, since the functor $\overline{H \circ \text{inc}}$ is colimit preserving, it follows from Lemma 2.1 below that $\Delta : \overline{H \circ \text{inc}}(\text{cof}) \rightarrow \Pi_{\mathbb{N}}\overline{H \circ \text{inc}}(\text{cof})$ factors through the canonical morphism $\oplus_{\mathbb{N}}\overline{H \circ \text{inc}}(\text{cof}) \rightarrow \Pi_{\mathbb{N}}\overline{H \circ \text{inc}}(\text{cof})$. Following [4, Lemma 2.8], this implies, in particular, that $\overline{H \circ \text{inc}}(\text{cof})$ is a phantom object, consult [4, Definition 2.4].

Now, let us apply the functor $\overline{H \circ \text{inc}}$ to the above assembly map (2.2). Since the functor $\overline{H \circ \text{inc}}$ is colimit preserving, it follows from the above commutative diagram (2.1) that $\overline{H \circ \text{inc}}$ (2.2) identifies with the assembly map (1.1). Moreover, $\overline{H \circ \text{inc}}(\text{cof})$ identifies with the cofiber of (1.1). Consequently, since $\overline{H \circ \text{inc}}(\text{cof})$ is a phantom object, we conclude that the assembly map (1.1) is a phantom equivalence; consult [4, Remark 2.6]. \square

Lemma 2.1. *Let o be an object of an ∞ -category \mathcal{E} such that the diagonal map $\Delta : o \rightarrow \Pi_{\mathbb{N}}o$ factors through the canonical map $\oplus_{\mathbb{N}}o \rightarrow \Pi_{\mathbb{N}}o$. Given a colimit preserving functor $F : \mathcal{E} \rightarrow \mathcal{D}$, the diagonal map $\Delta : F(o) \rightarrow \Pi_{\mathbb{N}}F(o)$ factors through the canonical map $\oplus_{\mathbb{N}}F(o) \rightarrow \Pi_{\mathbb{N}}F(o)$.*

Proof. Apply the functor F to the factorization of the diagonal map $\Delta : o \rightarrow \Pi_{\mathbb{N}}o$ through the canonical map $\oplus_{\mathbb{N}}o \rightarrow \Pi_{\mathbb{N}}o$. Thanks to the universal property of the product and to the fact that F is colimit preserving, one hence obtains a factorization of the diagonal map $\Delta : F(o) \rightarrow \Pi_{\mathbb{N}}F(o)$ through the canonical map $\oplus_{\mathbb{N}}F(o) \rightarrow \Pi_{\mathbb{N}}F(o)$. \square

Acknowledgments. This work is funded by national funds through the FCT - Fundação para a Ciência e a Tecnologia, I.P., under the scope of the projects UIDB/00297/2020 and UIDP/00297/2020 (Center for Mathematics and Applications) and the PhD scholarship SFRH/BD/144305/2019.

REFERENCES

- [1] A. J. BLUMBERG, D. GEPNER, AND G. TABUADA, *A universal characterization of higher algebraic K-theory*, *Geom. Topol.*, **17** (2013), pp. 733–838.
- [2] ———, *Uniqueness of the multiplicative cyclotomic trace*, *Adv. Math.*, **260** (2014), pp. 191–232.
- [3] U. BUNKE, D.-C. CISINSKI, D. KASPROWSKI, AND C. WINGES, *Controlled objects in left-exact ∞ -categories and the Novikov conjecture*, (2019). Available at <https://arxiv.org/abs/1911.02338>.
- [4] U. BUNKE, D. KASPROWSKI, AND C. WINGES, *On the Farrell-Jones conjecture for localising invariants*, (2021). Available at <https://arxiv.org/abs/2111.02490>.
- [5] F. T. FARRELL AND L. E. JONES, *Isomorphism conjectures in algebraic K-theory*, *J. Am. Math. Soc.*, **6** (1993), pp. 249–297.
- [6] G. TABUADA, *Noncommutative motives* (with a preface by Yuri I. Manin), vol. **63** of University Lecture Series, Providence, RI: American Mathematical Society (AMS), 2015.

CENTER FOR MATHEMATICS AND APPLICATIONS (NOVAMATH), FCT NOVA AND DEPARTMENT OF MATHEMATICS, FCT NOVA
Email address: jfd.reis@campus.fct.unl.pt