

Evolution of the maximal hypersurface in a D-dimensional dynamical spacetime

Suraj Maurya,* Rahul Nigam,† and Sashideep Gutti‡
Birla Institute of Technology and Science Pilani (Hyderabad Campus)
Hyderabad 500078, India

(Dated: 28-Nov-2022)

In this article we set up a variational problem to arrive at the equation of a maximal hypersurface inside a spherically symmetric evolving trapped region. In the first part of the article, we present the Lagrangian and the corresponding Euler Lagrange equations that maximize the interior volume of a trapped region that is formed dynamically due to infalling matter in D-dimensions, with and without the cosmological constant. We explore the properties of special points in these maximal hypersurfaces at which the Kodama vector becomes tangential to the hypersurface. These points which we call steady state points, are shown to play a crucial role in approximating the maximal interior volume of a black hole. We derive a formula to locate these points on the maximal hypersurface in terms of coordinate invariants like area radius, principle values of energy momentum tensor, Misner Sharp mass and cosmological constant. Based on this formula, we estimate the location of these steady state points in various scenarios: (a) the case of static BTZ black holes in 2+1 dimensions and for the Schwarzschild, Schwarzschild-deSitter and Schwarzschild-Anti-deSitter black holes in D-dimensions. We plot the location of the steady state points in relation to the event horizon and cosmological horizon in a static D-dimensional scenario, (b) cosmological case: we prove that these steady state points do not exist for homogeneous evolving dust for the zero and negative cosmological constant but exist in the presence of positive cosmological constant when the scale factor is greater than a critical value.

I. INTRODUCTION

One of the most intriguing and mysterious objects in the Universe is the black hole. This area of research is now in the forefront ever since the black holes have been observationally identified using gravitational wave astronomy. The study of black holes is now considered mainstream and astrophysically relevant. The thermodynamical properties of the black holes, the singularities inside the black holes etc. are yet to be understood fully. The information loss of the black hole and its resolution are still sort after mysteries.

One of the intriguing aspects of a black hole is the volume in the interior of a black hole. This question has been addressed by various authors [6–8] using different approaches. Maulikh Parikh [6] discusses the definition of volume by constructing an invariant slice of the spacetime inside the black hole horizon. Gibbons et al.[7] have discussed the thermodynamical volume, V_{th} inside a black hole in the presence of a varying cosmological constant Λ . V_{th} is defined as the conjugate variable to Λ appearing in the first law of thermodynamics for black hole i.e., $dE = TdS + \Omega dJ + \Phi dQ + V_{th}\Lambda$, where E is the gravitational enthalpy of spacetime. Christodoulou and Rovelli [8] provided a somewhat different definition of the black hole volume, in which the volume grows indefinitely as a function of the advance time.

In the article, we focus on the approach due to

Christodoulou and Rovelli [8]. They find out the volume inside the black hole via a variational approach. From the event horizon to the singularity they define a space-like curve in general and from among them define an extremal curve or a maximal hypersurface that yields the maximum volume in the interior of the black hole. They set up a Lagrangian framework to solve the above question. A similar method is adopted in the following papers [10–17]

In the article, we work on two aspects concerning the volume of the black holes. The first part deals with the question about the evolution of the interior volume of the trapped region where the black hole is in the process of formation. The apparent horizon is the boundary of the trapped region. Due to the evolution of the trapped region, the maximal hypersurface evolves and so does the maximum interior volume. This evolution of the volume is not just due to the expanding trapped region, but also due to the evolution of the maximal hypersurface. We set up a variational problem where we write down a Lagrangian for the spacelike curve between the apparent horizon and the singularity that gives the maximal volume. We then obtain the equations for this maximal hypersurface from the Euler-Lagrange equations for the obtained Lagrangian. We first solve the problem for the simple case of 2+1 dimensions where the underlying equations and analysis is simpler. The features of 2+1 dimensional gravity do not generalize in a straightforward way for a general D-dimension. We then carry out the analysis for the case of D-dimensional spherically symmetric dust evolution that leads to the formation of black hole.

It is proved in [8] that the volume generated by maxi-

* p20200471@hyderabad.bits-pilani.ac.in (S. Maurya)

† rahul.nigam@hyderabad.bits-pilani.ac.in (R. Nigam)

‡ sashideep@hyderabad.bits-pilani.ac.in (S. Gutti)

mal hypersurface has maximum contribution from a certain region that we call in this article as R_{ss} (the subscript stands for 'steady state'). This region provides an excellent approximation for the interior volume of the black hole. For a Schwarzschild black hole with mass M , the event horizon is at $R = 2M$ while the value of $R_{ss} = 3M/2$. This steady state radius R_{ss} lies in the interior of the black hole. This region of the maximal hypersurface was first discovered by Reinhart [9] in 1973. Similar points of interest in the interior of a black hole have been found in the examples described below in various other black holes like BTZ, Kerr, Kerr-AdS blackholes. The volume of the interior of a black hole is shown to be dependent on this radius R_{ss} . The volume of a Schwarzschild black hole is shown to be equal to $V = 3\sqrt{3}M^2v$ where v is the advance time. The special feature of the surface is that the normal to the surface is divergence free, i.e the trace of the extrinsic curvature tensor vanishes. This unique point has played a pivotal role in the approximation of volume of a black hole in a more general setting. For instance, the asymptotic volume of a BTZ black hole is crucially dependent on a point $R = \sqrt{M/2|\Lambda|}$ in [16] with M being the ADM mass of the BTZ black hole and Λ is the cosmological constant. Even in the presence of rotation, the asymptotic formula for the interior volume of the BTZ black hole crucially depends on the point $R = \sqrt{M/2|\Lambda|}$ as shown in [17]. In the estimation of the volume of the Kerr family of black holes [11, 15], the asymptotic value of the volume is dependent on these special point.

When one explores the interesting aspects of a black hole interiors one thinks of the interior as a somewhat trivial region with the only interesting feature being the spacetime singularity (and the inner horizon in case of rotating or charged black holes). The presence of R_{ss} between the event horizon and singularity, therefore reveals yet another interesting region in the interior of the black hole. These points have not been explored in all its generality. In this article, we explore the features of this special point in the spacetime which plays a crucial role in the interior region. We point out the relation of this special point of the maximal hypersurface with the Kodama vector. We show that this point corresponds to the location in the maximal hypersurface which is tangential to the Kodama vector. We study the emergence and evolution aspects of this crucial piece of hypersurface during the evolution and formation of the black holes. The study is exhaustive where we explore the evolutionary aspects in D-dimensions with and without cosmological constant. We arrive at interesting results for the various cases discussed. We develop a formula for tracking the evolution of the R_{ss} from which we can deduce its location.

In section II we discuss the evolution of the maximal hypersurface for homogeneous dust model for 2+1 dimensional and D-dimensional case. In section III, we use the extrinsic curvature method to estimate the steady state radius in 2+1 dimensional case. In the subsections of section III, we discuss the vacuum case, static

black hole case and cosmological case in 2+1 dimensions. In section IV, we discuss the extrinsic curvature method to locate the steady state radius in D-dimensions. In the subsections of section IV, we discuss the vacuum scenario, D-dimensional Schwarzschild black hole case, D-dimensional Schwarzschild deSitter and anti-deSitter case and the cosmological case. In section V we discuss the conclusions for the work.

II. EVOLUTION OF THE MAXIMAL HYPERSURFACE FOR HOMOGENEOUS DUST COLLAPSE MODEL

In this section we track the evolution of the interior volume of the BTZ black hole that has zero angular momentum. We study a dynamical situation corresponding to the formation of the BTZ black hole so that we estimate the evolution of the maximal volume in the interior. The reason for the parameter J is kept to zero in order to obtain analytically tractable expressions. We moreover consider the simplest case of a homogeneous evolution of dust. The metric for a circularly symmetric evolving inhomogeneous dust model is given below [1, 2].

$$ds^2 = -dt^2 + \frac{(\cos(\sqrt{|\Lambda|}t) + B' \sin(\sqrt{|\Lambda|}t))^2 dr^2}{|\Lambda|r^2 + |\Lambda|B^2 - 2\kappa \int_0^r \rho_i(s) s ds + 1} + (r \cos(\sqrt{|\Lambda|}t) + B \sin(\sqrt{|\Lambda|}t))^2 d\phi^2 \quad (1)$$

Here t is the comoving time of dust cloud and Λ is the cosmological constant. As the cosmological constant $\Lambda = -1/l^2$ is negative, the background space is AdS. The coordinate r is the comoving radius that is fixed to a dust particle whose distribution is circularly symmetric. The metric is expressed in terms of two functions $B(r)$ and $\rho_i(r)$. Here $R(r, t) = r \cos(\sqrt{|\Lambda|}t) + B \sin(\sqrt{|\Lambda|}t)$ is the area radius defined geometrically using the Killing vector $\partial/\partial\phi$ such that the perimeter of the shell of comoving shell r is $2\pi R$. $B(r)$ decides the initial velocity of the dust cloud and $\rho_i(r)$ decides the initial density. As the cloud evolves, the density evolves and the area radius of each shell decreases in the black hole formation regime. In our present context, we focus on homogeneous dust interior since the model offers simpler equations without compromising the caveats involved.

We choose the boundary of the homogeneous dust to be at a comoving coordinate $r = r_0$. Outside this is comoving radius r_0 , we assume vacuum. We further choose the condition that $B(r) = 0$ and $\rho_i = |\Lambda|/\kappa$ so that the metric (1) is in its simplest form given by

$$ds^2 = -dt^2 + \cos(\sqrt{|\Lambda|}t)^2 dr^2 + r^2 \cos^2(\sqrt{|\Lambda|}t) d\phi^2 \quad (2)$$

or

$$ds^2 = -dt^2 + R'^2 dr^2 + R^2 d\phi^2 \quad (3)$$

where $R = R(t, r) = r \cos(\sqrt{|\Lambda|}t)$ and $R' = R'(t, r) = \cos(\sqrt{|\Lambda|}t)$, is the derivative of R w.r.t r . We define

$F(r)$ is given by the expression $F(r) = 2\kappa \int_0^r \rho_i(s) ds$. This is the Misner Sharp mass. For the parameters being considered in the article, it is equal to $|\Lambda|r^2$. We note that for the model under consideration the ADM mass M , if the upper limit is taken as the boundary r_0 , $M = \kappa\rho_i r_0^2 - 1 = |\Lambda|r_0^2 - 1$. It is easily shown that the metric (1) can be smoothly matched at the hypersurface $r = r_0$ by equating the first and second fundamental form to the exterior BTZ metric given by

$$ds^2 = -(|\Lambda|R^2 - M)dT^2 + \frac{1}{(|\Lambda|R^2 - M)}dR^2 + R^2 d\phi^2 \quad (4)$$

with T is the time coordinate corresponding to Killing time. It is more convenient to switch from Schwarzschild like coordinate system (t, R, ϕ) to Eddington-Finkelstein like coordinates (v, R, ϕ) to avoid the coordinate singularity at the event horizon. The Eddington-Finkelstein coordinates are defined as

$$v = t + \int^R \frac{dR}{N^2(R)} \quad (5)$$

where, lapse function $N^2(R) = (|\Lambda|R^2 - M)$. The metric (4) can be written as

$$ds^2 = -(|\Lambda|R^2 - M)dv^2 + 2dv dR + R^2 d\phi^2 \quad (6)$$

Let λ be the parameter on the hypersurface Σ , then the induced metric on the hypersurface Σ define as

$$ds_\Sigma^2 = [-(|\Lambda|R^2 - M)\dot{v}^2 + 2\dot{v}\dot{R}]d\lambda^2 + R^2 d\phi^2 \quad (7)$$

with $\dot{v} = dv/d\lambda$ and $\dot{R} = dR/d\lambda$ in the above equation. Similarly, the induced metric (2) in terms of induced parameter λ defined as

$$ds^2 = [-\dot{t}^2 + \dot{r}^2 \cos^2(\sqrt{|\Lambda|}t)]d\lambda^2 + r^2 \cos^2(\sqrt{|\Lambda|}t)d\phi^2 \quad (8)$$

Here \dot{t} and \dot{r} are derivatives w.r.t the parameter λ . To find out the volume in the interior of a trapped region, we track the position of apparent horizon given in [1, 2]. The condition for the expansion parameter for outgoing null geodesics to become zero for a general metric (1) is given by [2]

$$\frac{2\kappa \int_0^r \rho_i(s) ds - 1}{|\Lambda|R^2} = 1 \quad (9)$$

For the specific case we are considering here, we have

$$\frac{|\Lambda|r^2 - 1}{|\Lambda|r^2 \cos^2(\sqrt{|\Lambda|}t)} > 1 \quad (10)$$

This implies the curve of the apparent horizon is given by

$$r_a^2 = \frac{1}{|\Lambda| \sin^2(\sqrt{|\Lambda|}t)} \quad (11)$$

We consider the collapsing regime where t goes from 0 to $\pi/(2\sqrt{|\Lambda|})$. At time $t = \pi/(2\sqrt{|\Lambda|})$, we have the formation of singularity since the area radius R of each shell of label $r \leq r_0$ becomes zero. We note that in the time interval considered during the collapsing phase, the apparent horizon r_a is a decreasing function of time. This implies the shell r_0 becomes trapped first, and then the smaller values of r get trapped. This also implies that since there is no more mass collapsing beyond the shell r_0 , the apparent horizon is also an isolated horizon and hence becomes an event horizon. We note that the evolving apparent horizon at a comoving $r < r_0$ is an inner horizon [5]. The physical radius of this horizon is given by, $R_h = \sqrt{M/|\Lambda|}$ where $M = |\Lambda|r_0^2 - 1$. During the course of evolution, we therefore have two distinct regions inside the BTZ black hole. (A) is the region from the event horizon $R = R_h$ to the area radius of the outer shell $R_0 = r_0 \cos(\sqrt{|\Lambda|}t)$. Region (B) consists of interior region of the dust cloud. We note that the region (A) is vacuum whereas (B) is evolving dust that eventually becomes singular. The plan is to track the evolution of the maximal volume where we take into account both the regions (A) and (B). The maximal surface is then the union of both the regions. So the volume is expressed as

$$V_\Sigma[\gamma] = - \int_{\lambda_h}^{\lambda_0} d\lambda \int_{S^1} d\phi \sqrt{R^2 (-(|\Lambda|R^2 - M)\dot{v}^2 + 2\dot{v}\dot{R})} \\ - \int_{\lambda_0}^{\lambda_f} d\lambda \int_{S^1} d\phi \sqrt{r^2 \cos^2(\sqrt{|\Lambda|}t) (-\dot{t}^2 + \dot{r}^2 \cos^2(\sqrt{|\Lambda|}t))} \quad (12)$$

The minus sign is because the parameter λ is chosen to be monotonically decreasing with increasing radius. In the first term, $R(\lambda_h) = \sqrt{M/|\Lambda|}$, which is the event horizon. The above expression is obtained by connecting the hypersurface across the boundary $r = r_0$. The first term in the above equation is the volume inside BTZ black hole from the event horizon to an area radius $R_0 = r_0 \cos(\sqrt{|\Lambda|}t)$, which is the outer shell of the collapsing cloud. Therefore we have $R(\lambda_0) = r_0 \cos(\sqrt{|\Lambda|}t)$. The solution to this term where we seek the maximal volume in the interior of the vacuum part of BTZ black hole is solved completely in [16].

We now focus on the evolution of maximal surface in (B) as a function of comoving time t . We follow the same Lagrangian procedure outlined in [8] for the metric given by eq.(8). We define a spacelike surface with topology $R \times S^1$. (We note that though this topology of the trapped region is generic, but in the context of our cosmological dust model, the evolving trapped region is annular region extending from the event horizon to the marginally trapped radius R_a , since the outermost shell r_0 get trapped first followed by inner shells). We now address the following variations problem. We have the following boundary values $t(\lambda_0) = t$, $r(\lambda_0) = r_0$ and $t(\lambda_f) = (1/\sqrt{|\Lambda|}) \sin^{-1}(1/\sqrt{|\Lambda|}r_a)$, $r(\lambda_f) = r_a$ where λ_f ends at the marginally trapped region given by eq.(11). We note here that as far as the comoving coordinate chart

is concerned, the location of the apparent horizon or the event horizon do not appear as any coordinate singularities in the metric. So the Euler-Lagrange equations remain the same irrespective of the boundary limits we impose. This is one of the main advantages of using the comoving coordinates to describe dust evolution. We have to maximize the following functional after integrating over the angle variable

$$V_{\Sigma} = -2\pi \int_{\lambda_0}^{\lambda_f} d\lambda \sqrt{r^2 \cos^2(\sqrt{|\Lambda|}t) (-\dot{t}^2 + \dot{r}^2 \cos^2(\sqrt{|\Lambda|}t))} \quad (13)$$

We may choose the parameter to be r since this coordinate is monotonic in the domain under consideration. The above functional then has only one function $t(r)$ that needs to be determined.

$$V_{\Sigma} = -2\pi \int_{r_0}^0 dr \sqrt{r^2 \cos^2(\sqrt{|\Lambda|}t) (-\dot{t}^2 + \cos^2(\sqrt{|\Lambda|}t))} \quad (14)$$

where, $\dot{r} = 1$ and $\dot{t} = \partial t / \partial r$, is the derivative of t w.r.t to r . This can be viewed as an extremization problem and our goal is to find the equations of motion for the Lagrangian.

$$L = L(t, \dot{t}, r) = \sqrt{r^2 \cos^2(\sqrt{|\Lambda|}t) (-\dot{t}^2 + \cos^2(\sqrt{|\Lambda|}t))} \quad (15)$$

The Euler-Lagrange equations are

$$\frac{d}{dr} \left(\frac{\partial L}{\partial \dot{t}} \right) - \frac{\partial L}{\partial t} = 0 \quad (16)$$

for the Lagrangian, (16), we get

$$\ddot{t} - \left(\frac{\sec^2(\sqrt{|\Lambda|}t)}{r} \right) \dot{t}^3 + 3\sqrt{|\Lambda|} [\tan(\sqrt{|\Lambda|}t)] \dot{t}^2 + \frac{\dot{t}}{r} - \sqrt{|\Lambda|} \sin(2\sqrt{|\Lambda|}t) = 0 \quad (17)$$

eq.(17) is a non-linear differential equation with variable coefficients where, r is independent variable and $t(r)$ is a dependent variable. Solution of eq.(17) yields the comoving time as a function of r i.e $t(r)$, which corresponds to maximal hypersurface.

A. D-dimensions case

A convenient coordinate system for describing evolving pressureless matter is a comoving one. The metric that describes the evolving dust scenario in $D(= n+2)$ -dimensions can be written as

$$ds^2 = -dt^2 + R'^2 dr^2 + R^2(t, r) d\Omega_n^2 \quad (18)$$

Here the coordinates are $t, r, \theta_1, \theta_2, \dots, \theta_n$. The $d\Omega_n$ is the metric induced on the n -dimensional sphere at a constant "area" radius R . To define the volume let λ be the

induced parameter on the hypersurface Σ then induced metric defined as

$$ds_{\Sigma}^2 = (-\dot{t}^2 + \dot{r}^2 R'^2) d\lambda^2 + R^2(t, r) d\Omega_n^2 \quad (19)$$

Volume in $D(= n+2)$ -dimensions define as

$$\begin{aligned} V_{\Sigma}^{(D)} &= \int d\lambda d\Omega_n R^n \sqrt{-\dot{t}^2 + \dot{r}^2 R'^2} \\ &= \frac{2\pi^{\frac{n+1}{2}}}{\Gamma(\frac{n+1}{2})} \int d\lambda R^n \sqrt{-\dot{t}^2 + \dot{r}^2 R'^2} \end{aligned} \quad (20)$$

Hence, the Lagrangian is defined as

$$L(t, \dot{t}, \lambda) = R^n \sqrt{-\dot{t}^2 + \dot{r}^2 R'^2} \quad (21)$$

Now, let the parameter $\lambda = r$ then $\dot{r} = 1$, substitute $R(t, r) = ra(t)$ and $R'(t, r) = a(t)$ in the eq.(21), we get

$$L(t, \dot{t}, r) = [ra(t)]^n \sqrt{-\dot{t}^2 + [a(t)]^2} \quad (22)$$

Our next goal is to find the equation of motion for the above Lagrangian from eq.(15), we get

$$\ddot{t} - \left(\frac{n}{r[a(t)]^2} \right) \dot{t}^3 - \left(\frac{(n+2)a'(t)}{a(t)} \right) \dot{t}^2 + \left(\frac{n}{r} \right) \dot{t} + (n+1)a(t)a'(t) = 0 \quad (23)$$

where, $\dot{t} = \partial t / \partial r$ and $a'(t) = \partial a(t) / \partial t$. This is the differential equation in D -dimensions and we can easily recover the $2+1$ dimensional differential equation i.e eq.(17) by putting $n = 1$ and $a(t) = \cos(\sqrt{|\Lambda|}t)$ in the eq.(23).

For estimating the maximal hypersurface in the trapped region, we can choose the appropriate boundary values. We note that the condition for marginal trapping for the model under consideration is given by $r_a = -1/\dot{a}$ [5], where the \dot{a} is derivative of the scale factor w.r.t time t . In a collapsing scenario, we have \dot{a} is negative thereby giving a positive value for r_a .

III. EXTRINSIC CURVATURE METHOD FOR ESTIMATING STEADY STATE RADIUS IN 2+1 DIMENSIONS

We now track the existence of the 'steady state' hypersurface that is used to estimate the interior volume of black holes. The location and its evolution upon correlating with the evolution of the event horizon provides important clues towards estimating the volume of the evolving black hole. As mentioned in the introduction, special points on such a maximal hypersurface contribute significantly towards estimating the interior volume of the black hole. We evaluate the divergence of extrinsic curvature for the $R(t, r) = \text{const.}$ surface. To find the normal to the surface we differentiate $R(t, r) = \text{const.}$ and get

$$\dot{R}dt + R'dr = 0 \quad (24)$$

The covariant components of normal vector defined as $n_\alpha = (n_t, n_r)$, where $n_t = \dot{R}$ and $n_r = R'$.

We now show that the Kodama vector is tangential to the maximal hypersurface at these steady state points. The trace of the extrinsic curvature vanishes at all points on the maximal hypersurface by definition [9, 18]. Kodama vector in a spherically symmetric spacetime is defined as [19, 20]. We consider the 2-metric in the coordinate chart (t, r) . This is given by $-dt^2 + R'^2 dr^2$. The two dimensional volume form in (t, r) coordinates are of $\omega = R' dt \wedge dr$. Using the standard definition of Kodama vector, we evaluate the components to be: $K^t = 1$ and $K^r = -\dot{R}/R'$. Evaluating the dot product, we find that $K^\alpha n_\alpha = 0$. This shows that the Kodama vector is tangential to the maximal hypersurface at the steady state points. We note that the above result is independent of whether the Kodama vector is spacelike or timelike. Infact for the cases that were discovered in [8, 11–15], the Kodama vector is spacelike and is tangential to the maximal hypersurface at the steady state radius. Another interesting observation is that at the steady state radius, both the normal vector to the hypersurface and the Kodama vector have vanishing divergence.

We now analyze the case of 2 + 1 dimensional scenario separately here owing to the non trivial nature of gravity in 2 + 1 dimensions. We start with the area radius $R(t, r)$ of comoving shell as a function of the comoving time and shell label r . The normalized contravariant components of normal vector are $N^\alpha = (N^t, N^r) = \left(\frac{-\dot{R}}{\sqrt{\dot{R}^2 - 1}}, \frac{1}{R' \sqrt{\dot{R}^2 - 1}} \right)$. The condition for the vanishing trace of extrinsic curvature, which implies that normal vector is divergence free, is written as

$$N_{;\alpha}^\alpha = \frac{1}{\sqrt{-g}} \frac{\partial}{\partial x^\alpha} (\sqrt{-g} N^\alpha) = 0 \quad (25)$$

where, $\sqrt{-g} = R'R$, which is obtained from the determinant of metric (3). The eq.(25) can be written as

$$\frac{1}{R'R} \left[\frac{\partial}{\partial t} (R'RN^t) + \frac{\partial}{\partial r} (R'RN^r) \right] = 0 \quad (26)$$

or

$$-\frac{\partial}{\partial t} \left(\frac{\dot{R}R'R}{\sqrt{\dot{R}^2 - 1}} \right) + \frac{\partial}{\partial r} \left(\frac{R}{\sqrt{\dot{R}^2 - 1}} \right) = 0 \quad (27)$$

solution of eq.(27) gives,

$$(\dot{R}^2 - 1)^2 + \frac{R\dot{R}^3\dot{R}'}{R'} - R\ddot{R} = 0 \quad (28)$$

\dot{R}^2 can be written as

$$\dot{R}^2 = f(r) + \Lambda R^2 \quad (29)$$

We are taking $f(r) = 0$. Differentiating eq.(29) w.r.to t and r , we get expression for \ddot{R} and \dot{R}' as

$$\ddot{R} = \Lambda R \quad (30)$$

and

$$\dot{R}' = \frac{\Lambda R R'}{\dot{R}} + \frac{F'(r)}{2\dot{R}} \quad (31)$$

Substituting the value of \dot{R}^2 , \ddot{R} and \dot{R}' from eqs.(29), (30) and (31) in to eq.(28), we get

$$\frac{RF'(r)}{2R'} [\Lambda R^2 + F(r)] + [\Lambda R^2 + F(r) - 1][2\Lambda R^2 + F(r) - 1] = 0 \quad (32)$$

We now obtain a general formula connecting the comoving energy density at a point with the area radius, Misner-Sharp mass and cosmological constant. Let ϵ is the energy density of collapsing dust then we can define the energy density as $\frac{F'(r)}{2RR'} = \epsilon$. Therefore, from eq.(32), we can write ϵ as

$$\epsilon = \frac{F'(r)}{2RR'} = - \frac{[\Lambda R^2 + F(r) - 1][2\Lambda R^2 + F(r) - 1]}{R^2[\Lambda R^2 + F(r)]} \quad (33)$$

This is the condition for $R = const.$ hypersurface to have vanishing divergence of extrinsic curvature. Though this formula is true in general, we now examine few simple cases and gain an understanding of this special point in the hypersurface.

1. Vacuum Case

When $F(r) = 0$. This situation represents matter content is zero in the entire spacetime. We see that

$$[\Lambda R^2 - 1][2\Lambda R^2 - 1] = 0 \quad (34)$$

We have two values of area radius where the condition is met: $R_{CH} = 1/\sqrt{\Lambda}$ and $R_{ss} = 1/\sqrt{2\Lambda} = R_{CH}/\sqrt{2}$.

The former root is the deSitter cosmological horizon which is a null horizon. The normal to the horizon is also the null generator and has vanishing divergence. We look for timelike normals (timelike normals for $R = const$ occur in the interior of a black hole) and spacelike normals, so this root is not our answer. The other root is the answer and we observe that this root lies between the $R = 0$ and the cosmological horizon. This feature is also observed in the situations to come. We note that for negative cosmological constant, this case does not yield any solutions.

2. Static Black hole case

Suppose we consider the case when the Misner-Sharp mass is a constant, this is the situation of a black hole or a naked singularity where all the mass has already collapsed to a point. We can use the formula derived above to examine the situation of a black hole using the comoving chart and not static/stationary coordinates. So

we take the condition that $F(r) = \text{const.} = M + 1$ where, M is the ADM mass. Also we note that automatically $F'(r) = 0$. From eq.(32), we get

$$[\Lambda R^2 + M][2\Lambda R^2 + M] = 0 \quad (35)$$

there are again two roots of the eq.(35).

$$R = R_{CH} = \sqrt{\frac{-M}{\Lambda}} \text{ and } R = R_{ss} = \sqrt{\frac{-M}{2\Lambda}} \quad (36)$$

Here $R_{ss} = \frac{1}{\sqrt{2}}R_{CH}$. Here we have two sub cases. For a positive cosmological constant, we require that the mass at the naked singularity (which is a conical singularity) is not greater than 1 so that M is negative. In this case we have the cosmological horizon at R_{CH} and as can be seen in the above equation, $R_{ss} = R_{CH}/\sqrt{2}$. The fact that M has to be negative is observed in [1, 2]. This scenario is not present in the four and higher dimensions where the ADM mass is positive definite. Here $R_{ss} < R_{CH}$, which shows that the surface is not beyond the cosmological horizon. The second subcase involves negative cosmological constant. Here, in contrast, we require M to be positive for roots to be real. So $F > 1$ for black hole event horizon to exist. The total mass that collapses has to be greater than 1 so that an event horizon forms [1, 2]. We then have

$$R = R_e = \sqrt{\frac{M}{|\Lambda|}} \text{ and } R = R_{ss} = \sqrt{\frac{M}{2|\Lambda|}} \quad (37)$$

As is evident from the above equations, R_{ss} occurs in the interior of the BTZ black hole with zero angular momentum. The radius is $R_{ss} = R_e/\sqrt{2}$. This is similar to the situation of the case of four dimensional black holes as observed in the work of [8, 10, 14, 15].

3. Cosmological Case

We look at cosmological solutions in 2+1 dimensions with and without cosmological constant. The mass function for homogeneous collapsing dust in 2+1 dimensions define as $F(r) = gr^2$ and area radius of comoving shells is $R = R(t, r) = ra(t)$. Substituting these parameters in eq.(32), we get

$$2(\Lambda a^2 + g)^2 r^4 - (3\Lambda a^2 + 2g)r^2 + 1 = 0 \quad (38)$$

1. When $\Lambda = 0$. We set $\Lambda = 0$ in the above equation then we obtain, $g^2 r^4 + (gr^2 - 1)^2 = 0$. As can be seen readily, this is a sum of squares and is never zero. So we do not have R_{ss} for real values of the area radius. To analyze the case with non zero cosmological constant, we evaluate the roots of the eq.(38). This gives steady state radius r_{ss} as

$$r_{ss} = \left(\frac{(3\Lambda a^2 + 2g) \pm \sqrt{(3\Lambda a^2 + 2g)^2 - 8(\Lambda a^2 + g)^2}}{4(\Lambda a^2 + g)^2} \right)^{\frac{1}{2}} \quad (39)$$

2. When $\Lambda > 0$ i.e de-Sitter space: Here r_{ss} will be positive only when the terms inside the square root is positive i.e

$$\begin{aligned} & (3\Lambda a^2 + 2g)^2 - 8(\Lambda a^2 + g)^2 > 0 \\ \Rightarrow a^2 & > -\frac{2g(\sqrt{2} + 1)}{\Lambda(3 + 2\sqrt{2})} \text{ and } a^2 > \frac{2g(\sqrt{2} - 1)}{\Lambda(3 - 2\sqrt{2})} \quad (40) \\ \Rightarrow a^2 & > \frac{2g(\sqrt{2} - 1)}{\Lambda(3 - 2\sqrt{2})} \text{ or } a^2 > \frac{2g(\sqrt{2} + 1)}{\Lambda} \end{aligned}$$

This condition can be seen to be equal to the relation inside the square root has to be positive. This inequality implies that r_{ss} is only possible during the evolution if, the scale parameter is greater than a critical value given in the equation. This is a surprising fact that this point emerges during the course of evolution provided the scale parameter crosses a certain threshold value.

3. When $\Lambda < 0$ i.e Anti-deSitter space: Replacing Λ with $-\Lambda$ in eq.(39) we get

$$r_{ss} = \left(\frac{(-3|\Lambda|a^2 + 2g) \pm \sqrt{\Lambda^2 a^4 + 4|\Lambda|ga^2 - 4g^2}}{4(-|\Lambda|a^2 + g)^2} \right)^{\frac{1}{2}} \quad (41)$$

Let us define $A = -3|\Lambda|a^2 + 2g$ and $B = \sqrt{8}(-|\Lambda|a^2 + 2g)$, then the terms, $A^2 - B^2 = (-3|\Lambda|a^2 + 2g)^2 - 8(\Lambda a^2 + g)^2 = \Lambda^2 a^4 + 4|\Lambda|ga^2 - 4g^2$. There are two possible situation arises on A .

(i) Suppose A is negative i.e $A < 0$, then eq.(41) can be written as

$$r_{ss} = \left(\frac{-|(-3|\Lambda|a^2 + 2g)| \pm \sqrt{\Lambda^2 a^4 + 4|\Lambda|ga^2 - 4g^2}}{4(-|\Lambda|a^2 + g)^2} \right)^{\frac{1}{2}} \quad (42)$$

Now we check the existence of above steady state radius r_{ss} as follows

$$\begin{aligned} & (-3|\Lambda|a^2 + 2g)^2 > (-3|\Lambda|a^2 + 2g)^2 - 8(\Lambda a^2 + g)^2 \\ & \Rightarrow (-3|\Lambda|a^2 + 2g)^2 > \Lambda^2 a^4 + 4|\Lambda|ga^2 - 4g^2 \\ & \Rightarrow |(-3|\Lambda|a^2 + 2g)| > \sqrt{\Lambda^2 a^4 + 4|\Lambda|ga^2 - 4g^2} \\ & \Rightarrow -|(-3|\Lambda|a^2 + 2g)| < -\sqrt{\Lambda^2 a^4 + 4|\Lambda|ga^2 - 4g^2} \\ & \Rightarrow -|(-3|\Lambda|a^2 + 2g)| + \sqrt{\Lambda^2 a^4 + 4|\Lambda|ga^2 - 4g^2} < 0 \quad (43) \end{aligned}$$

and $-|(-3|\Lambda|a^2 + 2g)| - \sqrt{\Lambda^2 a^4 + 4|\Lambda|ga^2 - 4g^2} < 0$ also be possible. Hence, these conditions shows that the terms $-|(-3|\Lambda|a^2 + 2g)| \pm \sqrt{\Lambda^2 a^4 + 4|\Lambda|ga^2 - 4g^2} < 0$ in the eq.(42), which means r_{ss} becomes imaginary and hence there is no any r_{ss} exist.

(ii). Suppose A is positive i.e $A = (-3|\Lambda|a^2 + 2g) > 0$ then, it gives the condition on the scale parameter $a(t)$ as $a^2 < 2g/3|\Lambda|$ and using this condition inside the square root term $\sqrt{\Lambda^2 a^4 + 4|\Lambda|ga^2 - 4g^2}$ we get a imaginary value. Hence in both the scenarios there is no solution for r_{ss} and hence they do not exist for 2+1 dimensional AdS cosmological spacetimes.

IV. EXTRINSIC CURVATURE METHOD IN D-DIMENSIONS IN THE PRESENCE OF COSMOLOGICAL CONSTANT (Λ)

The general metric for an $D(= n + 2)$ -dimensional spherically symmetric spacetime given in [3] which is define as

$$ds^2 = -e^{\mu(t,r)} dt^2 + e^{\lambda(t,r)} dr^2 + R^2(t,r) d\Omega_n^2 \quad (44)$$

where $d\Omega_n^2 = d\theta_1^2 + \sin^2\theta_1(d\theta_2^2 + \sin^2\theta_2(d\theta_3^2 + \dots + \sin^2\theta_{n-1}d\theta_n^2))$, is the metric on unit n-dimensional sphere, t is the time coordinate and r is the comoving radial coordinate. It is easily shown in [3] that the g_{tt} and g_{rr} component of the metric can be define as $g_{tt} = -1$ and $g_{rr} = \frac{R'^2}{1+f(r)}$. For marginally bounded shells of dust where we can take $f(r) = 0$. The metric then has the following form

$$ds^2 = -dt^2 + R'^2 dr^2 + R^2(t,r) d\Omega_n^2 \quad (45)$$

Area radius $R(t,r)$ of comoving shell is constant at the maximal hypersurface i.e $R = R(t,r) = \text{constant}$.

$$\dot{R}dt + R'dr = 0 \quad (46)$$

The covariant components of normal vector defined as $n_\alpha = (n_t, n_r)$, where $n_t = \dot{R}$ and $n_r = R'$. Corresponding normalized contravariant components of normal vector is $N^\alpha = (N^t, N^r) = \left(\frac{-\dot{R}}{\sqrt{R'^2 - 1}}, \frac{1}{R'\sqrt{R'^2 - 1}} \right)$. The condition for the vanishing trace of extrinsic curvature is given by,

$$N_{;\alpha}^\alpha = \frac{1}{\sqrt{-g}} \frac{\partial}{\partial x^\alpha} (\sqrt{-g} N^\alpha) = 0 \quad (47)$$

where, $\sqrt{-g} = R' R^n \Theta_n$ which is obtained from the determinant of metric (45) where Θ_n contains the product of all the angular part of determinant of metric (45). The eq.(47) can be written as

$$\frac{1}{R' R^n \Theta_n} \left[\frac{\partial}{\partial t} (R' R^n \Theta_n N^t) + \frac{\partial}{\partial r} (R' R^n \Theta_n N^r) \right] = 0 \quad (48)$$

or

$$-\frac{\partial}{\partial t} \left(\frac{\dot{R} R' R^n}{\sqrt{R'^2 - 1}} \right) + \frac{\partial}{\partial r} \left(\frac{R^n}{\sqrt{R'^2 - 1}} \right) = 0 \quad (49)$$

solution of eq.(49) gives,

$$nR^{n-1}(\dot{R}^2 - 1)^2 + \frac{R^n \dot{R}^3 \dot{R}'}{R'} - R^n \ddot{R} = 0 \quad (50)$$

\dot{R}^2 can be written as

$$\dot{R}^2 = f(r) + \frac{2\Lambda R^2}{n(n+1)} + \frac{F(r)}{R^{n-1}} \quad (51)$$

We take the $f(r) = 0$ for marginally bounded shell of dust. Differentiating eq.(51) w.r.to t and r , we get expression for \ddot{R} and \dot{R}' as

$$\ddot{R} = \frac{2\Lambda R}{n(n+1)} - \frac{(n-1)F(r)}{2R^n} \quad (52)$$

and

$$\dot{R}' = \frac{2\Lambda R R'}{n(n+1)\dot{R}} + \frac{F'}{2R^{n-1}\dot{R}} - \frac{(n-1)F R'}{2R^n \dot{R}} \quad (53)$$

Substituting the value of \dot{R}^2 , \ddot{R} and \dot{R}' from eqs.(51), (52) and (53) in to eq.(50) and after simplifying we get

$$\begin{aligned} & \frac{R F'(r)}{2R'} \left(\frac{2\Lambda R^2}{n(n+1)} + \frac{F(r)}{R^{n-1}} \right) + \\ & \left(\frac{2\Lambda R^2}{n(n+1)} + \frac{F(r)}{R^{n-1}} - 1 \right) \left[\frac{2\Lambda}{n} R^{n+1} + \frac{(n+1)}{2} F(r) - nR^{n-1} \right] \\ & = 0 \quad (54) \end{aligned}$$

We will see below that the above formula simplifies to an expression involving coordinate invariant since the term containing F' gets related to the energy density. The formula therefore is an interesting relation between the principle value of the energy momentum tensor, the cosmological constant and the Misner-Sharp Mass.

In a general setting the Misner-Sharp mass is a monotonically increasing function of the comoving radius r . We take a non zero value for the mass function $F(r) \neq 0$ (and $F'(r) \neq 0$). Let ϵ is the energy density of collapsing dust then we can define the energy density as $\frac{nF'(r)}{2R^n R'} = \epsilon$. Therefore, from eq.(54), we can write ϵ as

$$\begin{aligned} \epsilon &= \frac{nF'(r)}{2R^n R'} = \\ & \frac{n \left(1 - \frac{2\Lambda R^2}{n(n+1)} - \frac{F(r)}{R^{n-1}} \right) \left[\frac{2\Lambda}{n} R^{n+1} + \frac{(n+1)}{2} F(r) - nR^{n-1} \right]}{R^{n+1} \left(\frac{2\Lambda R^2}{n(n+1)} + \frac{F(r)}{R^{n-1}} \right)} \quad (55) \end{aligned}$$

1. Vacuum scenario

When there is no matter in the spacetime we have $F(r) = 0$. In this case the above equation yields Minkowski, deSitter or Anti-deSitter spacetime. For the case of $F(r) = 0$, the formula above gives

$$\left(\frac{2\Lambda R^2}{n(n+1)} - 1 \right) \left[\frac{2\Lambda}{n} R^{n+1} - nR^{n-1} \right] = 0 \quad (56)$$

We solve for the R_{ss}

$$\frac{2\Lambda R^2}{n(n+1)} - 1 = 0 \quad (57)$$

and

$$\frac{2\Lambda}{n}R^{n+1} - nR^{n-1} = 0 \quad (58)$$

from eq.(57) and (58) we get

$$R_{CH} = \sqrt{\frac{n(n+1)}{2\Lambda}} \text{ and } R_{ss} = \sqrt{\frac{n^2}{2\Lambda}} \quad (59)$$

or

$$R_{ss} = \sqrt{\frac{n}{n+1}} R_{CH} \quad (60)$$

Here $n < n+1$ so $R_{ss} < R_{CH}$ which means steady state radius occurs before the cosmological horizon. The cosmological horizon is the boundary of anti-trapped region. So R_{ss} occurs in the accessible part of spacetime from the point of view of the interior of the cosmological horizon. We also note that this surface is possible only in deSitter (dS) space and not possible in Anti-deSitter (AdS) space.

2. D-dimensional Schwarzschild black hole

When $\Lambda = 0$ and $F'(r) = 0$. From the eq.(54) we get

$$\left(\frac{F(r)}{R^{n-1}} - 1\right) \left[\frac{n+1}{2}F(r) - nR^{n-1}\right] = 0 \quad (61)$$

or

$$R = R_e = [F(r)]^{\frac{1}{n-1}} \text{ and } R = R_{ss} = \left(\frac{n+1}{2n}F(r)\right)^{\frac{1}{n-1}} \quad (62)$$

From eq.(62), we can write

$$R_{ss} = \left(\frac{n+1}{2n}\right)^{\frac{1}{n-1}} R_e \quad (63)$$

For Schwarzschild case $n = 2$ (i. e 4-dimensions), the mass function $F(r) = 2M$, then from equation (62) we will get $R_e = 2M$ which is correspond to event horizon and $R_{ss} = \frac{3}{2}M$, which is steady state radius correspond to maximal hypersurface as identified in [8]. From eq.(63) $R_{ss} = \frac{3}{4}R_e$ i.e $R_{ss} < R_e$, which means R_{ss} lies inside the event horizon of a black hole. Since $2n > n+1$ for $n = 2, 3, \dots$ Which means R_{ss} always lies inside the event horizon for all $n > 1$. For the case $n = 1$ there is no black hole if there is no negative cosmological constant.

The mass of the dust cloud given in [3], which is define as

$$M = \int_0^r \epsilon(0, r) dV \quad (64)$$

where, dV is the volume element of a spherical shell lying between r and $r + dr$ in $n+1$ space dimensions. The volume element dV define as

$$dV = \frac{2\pi^{\frac{n+1}{2}}}{\Gamma(\frac{n+1}{2})} r^n dr \quad (65)$$

from eqs.(64) and (65) we get the mass of dust cloud M as

$$M = \frac{2\pi^{\frac{n+1}{2}}}{\Gamma(\frac{n+1}{2})} \int_0^r \epsilon(0, r) r^n dr \quad (66)$$

The mass function $F(r)$ for the distribution of dust cloud defined as

$$F(r) = \frac{2k}{n} \int_0^r \epsilon(0, r) r^n dr \quad (67)$$

Therefore, from eqs.(66) and (67), the relation between mass function $F(r)$ (i.e. Misner-Sharp mass) and the mass of dust cloud M in $n+1$ spatial dimensions define as

$$F(r) = \frac{2k}{n} \int_0^r \epsilon(0, r) r^n dr = \frac{2k}{n} \frac{\Gamma(\frac{n+1}{2})M}{2\pi^{\frac{n+1}{2}}} \quad (68)$$

where, $k = \frac{8\pi G}{c^4}$ is the Einstein's gravitation constant and G is the Newton's gravitation constant.

3. D-dimensional Schwarzschild deSitter/Anti-deSitter spacetime scenario

If the mass function $F(r) = \text{const.}$ then the change in mass function of dust vanishes i.e $F'(r) = 0$ and event horizon will be static. Let cosmological constant $\Lambda \neq 0$, then from eq.(54), we get

$$\left(\frac{2\Lambda R^2}{n(n+1)} + \frac{F(r)}{R^{n-1}} - 1\right) \left[\frac{2\Lambda R^{n+1}}{n} + \frac{(n+1)F(r)}{2} - nR^{n-1}\right] = 0 \quad (69)$$

or

$$P(R = R_e) = \frac{2\Lambda R_e^2}{n(n+1)} + \frac{F}{R_e^{n-1}} - 1 = 0 \quad (70)$$

and

$$P(R = R_{ss}) = \frac{2\Lambda R_{ss}^2}{n(n+1)} + \frac{F}{2R_{ss}^{n-1}} - \frac{n}{n+1} = 0 \quad (71)$$

Both of the eqs.(70) and (71) are $n+1$ dimensional polynomials and finding analytical roots are in general difficult. We rather obtain good insight into the relative locations of the event horizons and the R_{ss} by plotting the graphs of the polynomials explicitly. We observe that R_{ss} usually lies in the interior of a black hole. In the presence of a positive cosmological constant, we get another root for R_{ss} which is at a radius smaller than the cosmological horizon. These facts are illustrated in the following graphs on the next page.




FIG. 1. The graph (a) shows the location of all the event horizons corresponding to the polynomial $P(R = R_e)$ of eq.(70) and graph (b) show the location of all the steady state radius corresponding to the polynomial $P(R = R_{ss})$ of eq.(71) for $D = 3, 4, 5$ and 6 dimensions in AdS spacetime. Graph (c) shows that both the R_e and R_{ss} in 3-dimensions overlaps at zero and graphs (d),(e) and (f) shows that the steady state radius R_{ss} always lies inside the event horizon R_e . Here we take the cosmological constant $\Lambda = -0.05$ and mass function $F(r) = 1$ for all the graphs .




FIG. 2. The graph (a) shows the location of all the event horizons R_e and cosmological horizons R_{CH} corresponding to the polynomial $P(R = R_e)$ of eq.(70). The graph (b) show the location of all the inner and outer steady state radius R_{ss} corresponding to the polynomial $P(R = R_{ss})$ of eq.(71) for $D = 3, 4, 5$ and 6 dimensions in deSitter (dS) spacetime. The graph (c) shows that all three R_e, R_{ss} and R_{CH} overlaps at zero and graphs (d), (e) and (f) shows that the inner R_{ss} always lies inside the R_e and outer R_{ss} lies between R_e and R_{CH} . Here we take the cosmological constant $\Lambda = 0.05$ and mass function $F(r) = 1$ for all the graphs .

4. *D-dimensional Cosmological scenario*

We now explore the presence and evolution of the steady state radius for the cosmological solutions in D-dimensions. We can think of the cosmological solutions as being valid for the entire universe or we can

think of the solutions as a homogeneous interior of an evolving star. If we assume the latter, we need to define an outer comoving shell r_0 beyond which there is vacuum. The mass function for homogeneous dust in D(= n+2)-dimension define as $F(r) = \frac{2g}{n(n+1)}r^{n+1}$ where $g = k\epsilon(0, r) > 0$ and $R = R(t, r) = ra(t)$. The dynamics of various cases are given in the Appendix. Substituting these value in equation (54), we get

$$\left(\frac{4(\Lambda a^2 + ga^{1-n})^2}{n^3(n+1)}\right)r^4 - \left(\frac{2\Lambda a^2(2n+1) + g(3n+1)a^{1-n}}{n^2(n+1)}\right)r^2 + 1 = 0 \quad (72)$$

Case(1). For a general case (when $\Lambda \neq 0$), the solution

of equation (72) gives steady state radius r_{ss} as

$$r_{ss} = \frac{\sqrt{n}}{2} \left(\frac{[2\Lambda a^2(2n+1) + g(3n+1)a^{1-n}] \pm \sqrt{[2\Lambda a^2(2n+1) + g(3n+1)a^{1-n}]^2 - 16n(n+1)(\Lambda a^2 + ga^{1-n})^2}}{2(\Lambda a^2 + ga^{1-n})^2} \right)^{1/2} \quad (73)$$

Case (2). When $\Lambda = 0$ i.e without cosmological

constant: From the eq.(73), we get

$$r_{ss} = \frac{\sqrt{n}}{2} \left(\frac{(3n+1) \pm \sqrt{(3n+1)^2 - 16n(n+1)}}{2ga^{1-n}} \right)^{1/2} = \begin{cases} 0 & ; \text{if } n = 0 \text{ i.e (2 - dimensions)} \\ Im & ; \text{if } n \geq 1 \text{ i.e (3 and higher - dimensions)} \end{cases} \quad (74)$$

There are two possible situation arises from the eq.(73) for the validity of steady state radius and which we discuss as follows

(a). When $\Lambda < 0$ i.e Anti-deSitter spacetime: Replacing the cosmological constant Λ with $-\Lambda$ in eq.(73) we get the steady state radius r_{ss} as

$$r_{ss} = \frac{\sqrt{n}}{2} \left(\frac{[-2|\Lambda|a^2(2n+1) + g(3n+1)a^{1-n}] \pm \sqrt{[-2|\Lambda|a^2(2n+1) + g(3n+1)a^{1-n}]^2 - 16n(n+1)(|\Lambda|a^2 + ga^{1-n})^2}}{2(-|\Lambda|a^2 + ga^{1-n})^2} \right)^{1/2} \quad (75)$$

Now, let us define $x = [-2|\Lambda|a^2(2n+1) + g(3n+1)a^{1-n}]$, $y = 4\sqrt{n(n+1)}(-|\Lambda|a^2 + ga^{1-n})$ and $z = \sqrt{2}(-|\Lambda|a^2 + ga^{1-n})$. Now there are two possible situation arises on x . **(i).** Suppose x is negative i.e $x < 0$, then eq.(69) gives steady state radius r_{ss} as follows

Now we check the existence of above steady state radius r_{ss} as follows

$$\begin{aligned} x^2 &> x^2 - y^2 \\ \Rightarrow |x| &> \sqrt{x^2 - y^2} \\ \Rightarrow -|x| &< -\sqrt{x^2 - y^2} \\ \Rightarrow -|x| + \sqrt{x^2 - y^2} &< 0 \\ \text{and also, } -|x| - \sqrt{x^2 - y^2} &< 0 \end{aligned} \quad (77)$$

$$r_{ss} = \frac{\sqrt{n}}{2} \left(\frac{-|x| \pm \sqrt{x^2 - y^2}}{z^2} \right) \quad (76)$$

eq.(77) shows that the terms $-|x| \pm \sqrt{x^2 - y^2} < 0$, which means r_{ss} becomes imaginary and hence doesn't exist.

(ii). Suppose x is positive i.e $x > 0$, then eq.(69) gives steady state radius as

$$r_{ss} = \frac{\sqrt{n}}{2} \left(\frac{x \pm \sqrt{x^2 - y^2}}{z^2} \right) \quad (78)$$

Now, if $x > 0$ then, we get a condition on scale parameter $a(t)$ as $a^{n+1} < g(3n+1)/2|\Lambda|(2n+1)$. Using this condition for any value of n (i.e $n = 1, 2, 3, 4, \dots$) the term $\sqrt{x^2 - y^2}$ in the eq.(78) becomes imaginary. Hence, r_{ss} doesn't exist for positive x also. Therefore, these two cases prove that the steady state radius r_{ss} doesn't exist in D-dimensional cosmological AdS spacetime.

(b). When $\Lambda > 0$ i.e deSitter spacetime: From the eq.(73), the steady state radius r_{ss} is positive only when, the terms inside the square root is positive i.e

$$[2\Lambda a^2(2n+1) + g(3n+1)a^{1-n}]^2 - 16n(n+1)(\Lambda a^2 + ga^{1-n})^2 > 0 \quad (79)$$

the inequality in the eq.(79) gives condition on the scale parameter $a(t)$ as

$$a^{n+1} > -\frac{g}{2\Lambda} \left[\frac{4\sqrt{n(n+1)} + (3n+1)}{(2n+1) + 2\sqrt{n(n+1)}} \right] \quad (80)$$

and $a^{n+1} > \frac{g}{2\Lambda} \left[\frac{4\sqrt{n(n+1)} - (3n+1)}{(2n+1) - 2\sqrt{n(n+1)}} \right]$

from the inequalities in eq.(80), the general condition on scale parameter $a(t)$ in deSitter spacetime is

$$a^{n+1} > \frac{g}{2\Lambda} \left[\frac{4\sqrt{n(n+1)} - (3n+1)}{(2n+1) - 2\sqrt{n(n+1)}} \right] \quad (81)$$

where, $4\sqrt{n(n+1)} > 3n+1$ and $2n+1 > 2\sqrt{n(n+1)}$, which shows that the terms in the square bracket of eq.(81) is positive. Hence, in the above inequality the scale parameter is positive which means there exists r_{ss} in deSitter spacetime cosmological solution in D-dimensions.

V. CONCLUSIONS

In this work we address few aspects concerning maximal hypersurface of a black hole in a dynamically evolving scenario. We considered spherically symmetric collapse of dust, generalized to D-dimensions since the model is analytically tractable. We have carried out the analysis separately for 2+1 dimensions and grouped the other dimensions together. This is due to the fact that 2+1 dimensional gravity is fundamentally different from other dimensions owing to the topological nature of gravity in 2+1 dimensions. The dimensions $D > 3$ are qualitatively similar to each other. For the evolving setting, we choose Lemaitre-Tolman-Bondi model generalized to

D-dimensions since the model has simplicity in terms of analytical expressions while capturing the core essence of the problem.

We obtain the equation for the maximal hypersurface using the variational technique developed in [8]. We set up a Lagrangian whose solution to the Euler-Lagrange equation yields the maximal hypersurface in an evolving setting. By choosing the appropriate boundary values for the solutions one can arrive at the maximal volume inside a trapped region which is in the process of evolving. The same procedure is generalized to D-dimensions. We present the equations by considering a sub-class of Lemaitre-Tolman-Bondi model, the homogeneous dust evolution where the expressions greatly simplify.

We analyze an interesting region of the maximal hypersurfaces, which we denote as 'steady state radius' (R_{ss}). The reason for this nomenclature is due to the role these points play in the estimation of the maximal volume inside a black hole. Identified first by Reinhart [9] 1973, these steady state points were found in various other black holes. In this article, we explored the existence and evolution of these points during the course of formation of black holes. We have identified an interesting property of these points in relation with the maximal hypersurfaces. These points are located where the Kodama Vector becomes tangential to the maximal hypersurface. The geometrical meaning and consequence of this observation is left for future considerations. The Kodama vector works as a substitute for a timelike Killing vector in scenarios that do not have a timelike Killing vector. Kodama vector, when it is timelike, has been used to define surface gravity in a dynamically evolving setting. In this article, we find another role of the Kodama vector, viz, it is used to pinpoint the steady state regions of a maximal hypersurface. We note that inside the black holes, the Kodama vector is spacelike.

We develop a formula to find the location of R_{ss} in terms of coordinate invariants like area radius, cosmological constants, principle value of the energy momentum tensor and Misner Sharp mass. Using the formula one can locate the steady state radius in various situations. We have explicitly evaluated the location of R_{ss} for the vacuum case, black hole case with and without the cosmological constant. We have presented our analysis and compared the R_{ss} with the position of the event horizon and cosmological horizon. We showed that in the black hole scenario, the R_{ss} is located within the event horizon. If there is a positive cosmological constant, we showed that R_{ss} lies at a area radius smaller than the cosmological horizon. When we consider an evolving situation, we used the collapse of homogeneous dust. This can be viewed as a cosmological solution or the collapse of a star with homogeneous distribution of dust. We showed that for the case of Oppenheimer-Snyder scenario and the collapse with a negative cosmological horizon, there is no real solution for R_{ss} and therefore R_{ss} does not exist. For the dust evolution in the presence of a positive cosmological constant, we showed that R_{ss} exists provided

the evolving scale factor crosses a certain critical value.

The analysis raises many questions that are left for future considerations. Does the relation between Kodama vector and the maximal hypersurface continue to hold in a non spherically symmetric situation, like the Kerr family of solutions? A timelike Kodama vector has been used to define various quantities that have thermodynamic interpretation like surface gravity, etc in dynamical situations. Does a spacelike Kodama vector also have a geometric interpretation? The Lagrangian formulation for the maximal hypersurface in Kerr/Kerr-AdS/Kerr-Newman/Kerr-DeSitter has not been formulated though there are many interesting papers estimating the volume of the interior of Kerr family of black holes [11, 15]. The Lagrangian formulation of the Kerr family is a work in progress. At R_{ss} both the normal vector and its tangent has zero divergence. Is there a special geometric meaning associated with R_{ss} owing to the above property? These questions are left open.

ACKNOWLEDGMENTS

We would like to thank our institute BITS Pilani Hyderabad campus for providing the required infrastructure to carry out this research work. Suraj Maurya would further like to thank Government of India for providing CSIR (NET-JRF) fellowship to support this research work.

APPENDIX

A. Solution of scale parameter $a(t)$ for homogeneous dust evolution

As we know that the area radius $R(t, r)$ and mass function $F(r)$ of homogeneous dust defined as

$$R(t, r) = ra(t) \text{ and } F(r) = \frac{2g}{n(n+1)} r^{n+1} \quad (82)$$

also

$$\dot{R}^2 = \frac{2\Lambda}{n(n+1)} R^2 + \frac{2g}{n(n+1)} r^{n+1} \quad (83)$$

from eq.(82) and (83) we get

$$[\dot{a}(t)]^2 = \frac{2\Lambda}{n(n+1)} [a(t)]^2 + \frac{2g}{n(n+1)[a(t)]^{n-1}} \quad (84)$$

solution of eq.(84) gives the scale parameter $a(t)$ for different region of spacetime on the basis of cosmological constant (Λ).

1. For vacuum spacetime ($\Lambda = 0$).

For zero cosmological constant the eq.(84) becomes

$$[\dot{a}(t)]^2 = \frac{2g}{n(n+1)[a(t)]^{n-1}} \Rightarrow \frac{da(t)}{dt} = \pm \frac{1}{a^{\frac{n-1}{2}}} \sqrt{\frac{2g}{n(n+1)}} \quad (85)$$

solution of eq.(85) with initial scale parameter $a(0) = 1$ gives,

$$a(t) = \left(1 + \sqrt{\frac{g(n+1)}{2n}} t \right)^{\frac{n+1}{2}} \text{ and } \left(1 - \sqrt{\frac{g(n+1)}{2n}} t \right)^{\frac{n+1}{2}} \quad (86)$$

2. For deSitter (dS) spacetime ($\Lambda > 0$).

For positive cosmological constant the change in scale parameter defined as

$$\begin{aligned} [\dot{a}(t)]^2 &= \frac{2\Lambda}{n(n+1)} [a(t)]^2 + \frac{2g}{n(n+1)[a(t)]^{n-1}} \\ \Rightarrow \frac{da(t)}{dt} &= \pm \sqrt{\frac{2\Lambda}{n(n+1)} [a(t)]^2 + \frac{2g}{n(n+1)[a(t)]^{n-1}}} \end{aligned} \quad (87)$$

The solutions of eq.(87) with initial condition $a(0) = 1$ is

$$\begin{aligned} a(t) &= \left[\sqrt{\frac{g}{\Lambda}} \sinh \left(\sqrt{\frac{\Lambda(n+1)}{2n}} t + \operatorname{arcsinh} \sqrt{\frac{\Lambda}{g}} \right) \right]^{\frac{2}{n+1}} \\ \text{and } &\left[-\sqrt{\frac{g}{\Lambda}} \sinh \left(\sqrt{\frac{\Lambda(n+1)}{2n}} t - \operatorname{arcsinh} \sqrt{\frac{\Lambda}{g}} \right) \right]^{\frac{2}{n+1}} \end{aligned} \quad (88)$$

3. For Anti-deSitter (AdS) spacetime ($\Lambda < 0$).

For negative cosmological constant the change in scale parameter defined as

$$\begin{aligned} [\dot{a}(t)]^2 &= -\frac{2\Lambda}{n(n+1)} [a(t)]^2 + \frac{2g}{n(n+1)[a(t)]^{n-1}} \\ \Rightarrow \frac{da(t)}{dt} &= \pm \sqrt{-\frac{2\Lambda}{n(n+1)} [a(t)]^2 + \frac{2g}{n(n+1)[a(t)]^{n-1}}} \end{aligned} \quad (89)$$

The solutions of eq.(89) with initial condition $a(0) = 1$ is

$$\begin{aligned} a(t) &= \left[\sqrt{\frac{g}{\Lambda}} \sin \left(\sqrt{\frac{\Lambda(n+1)}{2n}} t + \operatorname{arcsin} \sqrt{\frac{\Lambda}{g}} \right) \right]^{\frac{2}{n+1}} \\ \text{and } &\left[-\sqrt{\frac{g}{\Lambda}} \sin \left(\sqrt{\frac{\Lambda(n+1)}{2n}} t - \operatorname{arcsin} \sqrt{\frac{\Lambda}{g}} \right) \right]^{\frac{2}{n+1}} \end{aligned} \quad (90)$$

-
- [1] R. B. Mann and S. F. Ross, "Gravitationally collapsing dust in 2+ 1 dimensions", *Phys. Rev. D.* 47.3319 (1993).
- [2] Sashideep Gutti, "Gravitational collapse of inhomogeneous dust in (2+1) dimensions", *Class. Quantum Grav.* 22 3223 (2005).
- [3] Rakesh Tibrewala, Sashideep Gutti, T. P. Singh and Cenalo Vaz, "Classical and quantum gravitational collapse in d-dimensional AdS spacetime: Classical solutions", *Phys. Rev. D.* 77, 064012 (2008).
- [4] Nilanjandev Bhaumik and Bibhas Ranjan Majhi, "Interior volume of (1+D)-dimensional Schwarzschild black hole", *International Journal of Modern Physics A* Vol. 33, No. 02, 1850011 (2018).
- [5] Konka Raviteja and Sashideep Gutti, "Aspects of marginally trapped and antitrapped surfaces in a D-dimensional evolving dust model", *Phys. Rev. D* 102, 024072 (2020).
- [6] M. Parikh, "Volume of Black holes", *Phys. Rev. D* 73,124021 (2006).
- [7] M. Cvetič, G. W. Gibbons, D. Kubiznak, and C. N. Pope, "Black hole enthalpy and an entropy inequality for the thermodynamic volume", *Phys. Rev. D* 84, 024037 (2011).
- [8] Marios Christodoulou and Carlo Rovelli, "How big is a black hole?", *Phys. Rev. D* 91, 064046 (2015).
- [9] Bruce L. Reinhart, "Maximal foliations of extended Schwarzschild space", *J. Math. Phys.* 14, 719 (1973).
- [10] Baocheng Zhang, "Entropy in the interior of a black hole and thermodynamics", *Phys. Rev. D* 92, 081501(R) (2015).
- [11] I. Bengtsson, E. Jakobsson, "Black holes: Their large interiors" *Mod. Phys. Lett. A* 30 (2015) 1550103.
- [12] Y. C. Ong, "Never judge a black hole by its area" *J. Cosmol. Astropart. Phys.* 04 (2015) 003.
- [13] Y.C. Ong, "The persistence of the large volumes in black holes", *Gen. Relativ. Gravit.* 47 (2015) 88.
- [14] M. Christodoulou, T. De Lorenzo, "Volume inside old black holes", *Phys. Rev. D* 94 (2016) 104002.
- [15] Xiao Yan Chew, Yen Chin Ong, "Interior volume of Kerr-AdS black holes", *Phys. Rev. D* 102 (6) (2020) 064055.
- [16] Ming Zhang, "Interior volume of Banados-Teitelboim-Zanelli black hole", *Physics Letter B* 790 (2019) 205-210.
- [17] Suraj Maurya, Sashideep Gutti and Rahul Nigam, "Volume of a rotating black hole in 2+1 dimensions", *Phys. Lett. B* Volume 833 j.physletb.137381(2022).
- [18] F. Estabrook, H. Wahlquist, S. Christensen, B. DeWitt, L. Smarr, and E. Tsiang, "Maximally slicing a black hole" *Phys. Rev. D* 7, 2814 (1973).
- [19] Hideo Kodama, "Conserved Energy Flux for the Spherically Symmetric System and the Backreaction Problem in the Black Hole Evaporation", *Progress of Theoretical Physics*, Volume 63, Pages 1217-1228 (1980).
- [20] Valerio Faraoni, "Cosmological and Black Hole Apparent Horizons", *Lect.Notes Phys.* 907, pp.1-199 (2015).