

ON SEELEY-TYPE UNIVERSAL EXTENSION OPERATORS FOR UPPER HALF SPACE

HAOWEN LU AND LIDING YAO

ABSTRACT. Modified from the standard half-space extension via reflection principle, we construct a linear extension operator for the upper half space \mathbb{R}_+^n that has the form $Ef(x) = \sum_{j=-\infty}^{\infty} a_j f(x', -b_j x_n)$ for $x_n < 0$. We prove that E is bounded in all C^k -spaces, Sobolev and Hölder spaces, Besov and Triebel-Lizorkin spaces, along with their Morrey generalizations. We also give an analogous construction on bounded smooth domains.

1. INTRODUCTION

Given a function f defined on the upper half space $\mathbb{R}_+^n = \{(x', x_n) : x_n > 0\}$, the *reflection principle* gives a well-known construction to extend f to the total space \mathbb{R}^n while preserving its regularity property: we can define the extension Ef that depends linearly on f by

$$(1) \quad Ef(x', x_n) = E^{a,b}f(x', x_n) := \sum_j a_j f(x', -b_j x_n), \quad \text{for } x_n < 0, \quad \text{where } b_j > 0.$$

Here $(a_j, b_j)_j$ are (finitely or infinitely many) real numbers that satisfy certain algebraic conditions (see below). This method traces back to Lichtenstein [Lic29] for C^1 -extensions and [Hes41] for C^k -extensions. See also [Nik53] and [Bab53] for the corresponding Sobolev extensions.

In the spaces of distributions, a linear extension operator E is viewed as a right inverse of the natural restriction map $\mathcal{D}'(\mathbb{R}^n) \rightarrow \mathcal{D}'(\mathbb{R}_+^n)$. Triebel [Tri95] and Franke [Fra86] showed that for every $\varepsilon > 0$ there is a $m = m(\varepsilon) > 0$, such that, if the finite collection $(a_j, b_j)_j$ satisfy $\sum_j a_j (-b_j)^k = 1$ for all integers $-m \leq k \leq m$, then for E in (1) have boundedness in Besov spaces $E : \mathcal{B}_{pq}^s(\mathbb{R}_+^n) \rightarrow \mathcal{B}_{pq}^s(\mathbb{R}^n)$ and in Triebel-Lizorkin spaces $E : \mathcal{F}_{pq}^s(\mathbb{R}_+^n) \rightarrow \mathcal{F}_{pq}^s(\mathbb{R}^n)$, for all $\varepsilon < p, q \leq \infty$ and $-\varepsilon^{-1} < s < \varepsilon^{-1}$ ($p < \infty$ in \mathcal{F} -cases). In [Tri06, Section 1.11.5] such E is called a *common extension operator* which depends on $\varepsilon (> 0)$.

In contrast, we call E a **universal extension operator**, if we have the boundedness $E : \mathcal{A}_{pq}^s(\mathbb{R}_+^n) \rightarrow \mathcal{A}_{pq}^s(\mathbb{R}^n)$ for all $\mathcal{A} \in \{\mathcal{B}, \mathcal{F}\}$, $0 < p, q \leq \infty$ and $s \in \mathbb{R}$ ($p < \infty$ in \mathcal{F} -cases), simultaneously (i.e. we can take $\varepsilon = 0$).

Note that the collections $(a_j, b_j)_j$ discussed above are all finite. In those cases, the range of spaces on which E is bounded would inevitably be finite and depends on $\varepsilon > 0$. Hence such E is never universal.

By taking some infinitely nonzero sequences $(a_j, b_j)_j$, Seeley [See64] constructed one such operator (1) such that $E : C^k(\mathbb{R}_+^n) \rightarrow C^k(\mathbb{R}^n)$ for all $k = 0, 1, 2, \dots$. In particular Seeley's extension preserve C^∞ -smoothness. However, to the best of authors' knowledge, there was no proof of the boundedness of Seeley's operator on general Besov and Triebel-Lizorkin spaces.

In this paper we generalize Seeley's construction by extending the boundedness to the spaces of negative index, and we show that such operator is a universal extension operator.

Theorem 1. (i) *There exists an extension operator $E : C^0(\overline{\mathbb{R}_+^n}) \rightarrow C^0(\mathbb{R}^n)$ that has the form*

$$(2) \quad Ef(x', x_n) = E^{a,b}f(x', x_n) := \begin{cases} \sum_{j=-\infty}^{\infty} a_j f(x', -b_j x_n) & x_n < 0 \\ f(x) & x_n > 0 \end{cases}, \quad \text{where } b_j > 0 \text{ such that,}$$

$$(3) \quad \text{for every } k \in \mathbb{Z}, \quad \sum_{j=-\infty}^{\infty} a_j (-b_j)^k = 1 \text{ and } \sum_{j \in \mathbb{Z}} 2^{\delta_k |j|} |a_j| \cdot b_j^k < \infty \text{ for some } \delta_k > 0.$$

Moreover, when (3) is satisfied, then (2) always defines an extension operator E for \mathbb{R}_+^n , that has the following boundedness (simultaneously):

(ii) (Sobolev, Hölder and C^k) $E : W^{k,p}(\mathbb{R}_+^n) \rightarrow W^{k,p}(\mathbb{R}^n)$ and $E : C^{k,s}(\mathbb{R}_+^n) \rightarrow C^{k,s}(\mathbb{R}^n)$ are defined and bounded for all $k \in \mathbb{Z}$, $0 < p \leq \infty$ and $0 < s < 1$.

$E : C^k(\mathbb{R}_+^n) \rightarrow C^k(\mathbb{R}^n)$ is bounded for all $k = 0, 1, 2, \dots$

(iii) (Besov and Triebel-Lizorkin) $E : \mathcal{S}'(\mathbb{R}_+^n) \rightarrow \mathcal{S}'(\mathbb{R}^n)$ is continuous. Moreover we have boundedness in Besov spaces $E : \mathcal{B}_{pq}^s(\mathbb{R}_+^n) \rightarrow \mathcal{B}_{pq}^s(\mathbb{R}^n)$ and in Triebel-Lizorkin spaces $E : \mathcal{F}_{pq}^s(\mathbb{R}_+^n) \rightarrow \mathcal{F}_{pq}^s(\mathbb{R}^n)$ for all $s \in \mathbb{R}$ and $0 < p, q \leq \infty$.

(iv) (Morrey-type) More generally E has boundedness on all Besov-type spaces $\mathcal{B}_{pq}^{s\tau}$, Triebel-Lizorkin-type spaces $\mathcal{F}_{pq}^{s\tau}$ and Besov-Morrey spaces $\mathcal{N}_{pq}^{s\tau}$. That is, $E : \mathcal{A}_{pq}^{s\tau}(\mathbb{R}_+^n) \rightarrow \mathcal{A}_{pq}^{s\tau}(\mathbb{R}^n)$ for $\mathcal{A} \in \{\mathcal{B}, \mathcal{F}, \mathcal{N}\}$ and all $0 < p, q \leq \infty$, $s \in \mathbb{R}$ and $0 \leq \tau \leq \frac{1}{p}$ ($p < \infty$ for the \mathcal{F} -cases).

See Definitions 3, 6 and 8 for the spaces in the theorem. Here the Sobolev spaces $W^{k,p}$ for $p < 1$ can be defined and are discussed in [Pee75], see also Remark 4.

The analogue of Theorem 1 on smooth domains is also true; see Theorem 26. The operator is given by (37).

Remark 2. Here we allow δ_k in (3) to tend to 0 as $|k| \rightarrow \infty$. In practice we can choose $(a_j, b_j)_j$ such that the sum $\sum_j 2^{\delta|j|} |a_j| b_j^k < \infty$ holds for all k and all arbitrarily large δ , see Proposition 14.

It is not known to the authors whether the results still hold if we remove $2^{\delta_k|j|}$ in (3), i.e. if we only assume $\sum_j |a_j| b_j^k < \infty$ for all $k \in \mathbb{Z}$. In the proof of Theorem 1 the terms $2^{\delta_k|j|}$ are used only when we consider $p < 1$ in (ii) and $\min(p, q) < 1$ in (iii) and (iv).

We should remark that the universal extension operator exists not only on smooth domains but also on Lipschitz domains. This is done by Rychkov [Ryc99] using Littlewood-Paley decompositions. Rychkov's extension operator E^R has the form $E^R f = \sum_{j=0}^{\infty} \psi_j * (\mathbf{1}_{\mathbb{R}_+^n} \cdot (\phi_j * f))$ where $\mathbf{1}_{\mathbb{R}_+^n}$ is the characteristic function of \mathbb{R}_+^n and $(\phi_j, \psi_j)_{j=0}^{\infty}$ is a carefully chosen family of Schwartz functions. The shape of E^R is totally different from (1) and the construction follows a different methodology.

Our result shows that the construction (1) can also be made to be universal extension. Cf. the comment below [JMHS15, Corollary 5.7].

In fact, our extension operator is “more universal” than the Rychkov extension E^R [Ryc99]. Although E^R is bounded in all Besov and Triebel-Lizorkin spaces (see [Ryc99, Theorem 4.1]), it is not known whether E^R is bounded on the endpoint Sobolev spaces $W^{k,1}$, $W^{k,\infty}$ and the C^k -spaces. On the other hand, our extension operator is also defined on the measurable function space $L^p(\mathbb{R}_+^n)$ for $0 < p < 1$, whose element may not be realized as a distribution on \mathbb{R}_+^n .

The range of the extension operator is important if we want to construct some operator that has the form

$$(4) \quad Tf(x) = \int_{\mathcal{U}} K(x, y) \mathcal{E}f(y) dy, \quad \text{where } \mathcal{E} \text{ is an extension operator for the domain } \Omega \subset \subset \mathcal{U}.$$

Here $K(x, y)$ can be some kernel of integral operator or singular integral operator.

This method has been used extensively in solving Cauchy-Riemann problem (the $\bar{\partial}$ -equation) on certain domains in \mathbb{C}^n . For example Wu [Wu98] used the Seeley's extension to construct a solution operator that is $W^{k,p}$ bounded for all $k \geq 0$. If \mathcal{E} is some common extension operator that is only bounded in a finite range of k , then T is only bounded in a finite range as well.

By replacing the Seeley's extension with our extension operator and doing more analysis on $K(x, y)$, it might be possible that one can show that T in (4) is bounded in $W^{k,p}$ for $k < 0$ as well.

The estimates on negative Sobolev spaces are discussed in [SY21, Yao22], where we took \mathcal{E} to be Rychkov's extension operator. In the case of smooth domains, if we replace Rychkov's extension by our Seeley-type extension, then it is possible not only to get some estimates related to L^1 -Sobolev spaces, but also to simplify some of the proofs in these papers (see for example [Yao22, Proposition 5.1 and Remark 5.2 (iii)]).

2. FUNCTION SPACES AND NOTATIONS

In the paper we use the following definitions for Sobolev spaces and Hölder spaces, including both positive and negative indices. We also include the case $p < 1$ for Sobolev spaces.

Definition 3. Let $U \subseteq \mathbb{R}^n$ be an arbitrary open subset. Let $k \in \mathbb{N}_0 = \{0, 1, 2, \dots\}$.

We use $C^k(U) = C^k(\bar{U})$ for the space of all continuous functions $f : U \rightarrow \mathbb{R}$ such that $\partial^\alpha f$ are bounded and uniformly continuous for all $|\alpha| \leq k$. We use $\|f\|_{C^k(U)} := \max_{|\alpha| \leq k} \|\partial^\alpha f\|_{C^0(U)}$.

We let $C^\infty(U) = C^\infty(\bar{U}) := \bigcap_{k=0}^{\infty} C^k(U)$ to be the space of bounded smooth functions.

For $0 < s < 1$, we define the Hölder space $C^{k,s}(U)$ to be the space of all functions $f \in C^k(U)$ such that $\|f\|_{C^{k,s}(U)} := \|f\|_{C^k(U)} + \max_{|\alpha| \leq k} \sup_{x,y \in U} |x-y|^{-s} |\partial^\alpha f(x) - \partial^\alpha f(y)| < \infty$.

For $0 < p \leq \infty$, we define the Sobolev space $W^{k,p}(U)$ by the following:

- (i) For $1 \leq p \leq \infty$, $W^{k,p}(U)$ consists of all $f \in L^p(U)$ whose distributional derivatives $\partial^\alpha f$ belong to $L^p(U)$ for all $|\alpha| \leq k$. We use the norm $\|f\|_{W^{k,p}(U)} := (\sum_{|\alpha| \leq k} \int_U |\partial^\alpha f|^p)^{\frac{1}{p}}$ for $1 \leq p < \infty$ and $\|f\|_{W^{k,\infty}(U)} := \sup_{|\alpha| \leq k} \|\partial^\alpha f\|_{L^\infty(U)}$.

- (ii) For $0 < p < 1$, the quasi-Banach space $W^{k,p}(U)$ is the abstract completion¹ of $C_c^\infty(\bar{U})$ under the quasi-norm $\|f\|_{W^{k,p}(U)} := (\sum_{|\alpha| \leq k} \int_U |\partial^\alpha f|^p)^{\frac{1}{p}}$.

For $k > 0$, $0 < s < 1$ and $1 \leq p \leq \infty$, we define

$$(5) \quad \|f\|_{C^{-k,s}(U)} := \inf \left\{ \max_{|\alpha| \leq k} \|g_\alpha\|_{C^{0,s}(U)} : \{g_\alpha\}_{|\alpha| \leq k} \subset C^{0,s}(U), \quad f = \sum_{|\alpha| \leq k} \partial^\alpha g_\alpha \text{ as distributions} \right\}.$$

$$(6) \quad \|f\|_{W^{-k,p}(U)} := \inf \left\{ \left(\sum_{|\alpha| \leq k} \|g_\alpha\|_{L^p(U)}^p \right)^{1/p} : \{g_\alpha\}_{|\alpha| \leq k} \subset L^p(U), \quad f = \sum_{|\alpha| \leq k} \partial^\alpha g_\alpha \text{ as distributions} \right\}.$$

- (iii) We define $C^{-k,s}(U)$ and $W^{-k,p}(U)$ ($1 \leq p \leq \infty$) to be the subset of distributions such that the above norms are finite, respectively.
- (iv) For $0 < p < 1$, we define $W^{-k,p}(U)$ to be the abstract completion of $C_c^\infty(\bar{U})$ under the quasi-norm (6) with $\{g_\alpha\}_\alpha \subset L^p(U)$ replacing by $\{g_\alpha\}_\alpha \subset C_c^\infty(\bar{U})$ (where we only consider $f \in C_c^\infty(\bar{U})$ on both sides).

Remark 4. When $0 < p < 1$ and $k \geq 1$, the natural (continuous) mapping $\iota : W^{k,p}(U) \rightarrow L^p(U)$ is **not** injective. The proof can be modified from [Pee75, Proposition 3.1]. If one use $W^{k,p}(U)/\ker \iota$ instead, then the elements can always be realized as measurable functions (in $L^p(U)$).

One can check that Theorem 1 (ii) implies the boundedness $E : W^{k,p}(\mathbb{R}_+^n)/\ker \iota_{\mathbb{R}_+^n} \rightarrow E : W^{k,p}(\mathbb{R}^n)/\ker \iota_{\mathbb{R}^n}$.

Remark 5. When U satisfies nice boundary condition, particularly when $U \in \{\mathbb{R}_+^n, \mathbb{R}^n\}$ or when U is a bounded smooth domain, we have the following:

- (a) Definition 3 (i) and (ii) coincide, in the sense that $C_c^\infty(\bar{U})$ is dense in $W^{k,p}(U)$ for $p \in [1, \infty)$. For the proof see [Eva10, Section 5.3] or [Ada75, Theorem 3.17]. Taking the distribution derivatives, we get the density of $C_c^\infty(\bar{U})$ in $W^{-k,p}(U)$ for $p \in [1, \infty)$ as well.
- (b) When $1 \leq p < \infty$, $W^{-k,p'}(U)$ is the dual of $W_0^{k,p}(U) := \overline{C_c^\infty(\bar{U})}^{\|\cdot\|_{W^{k,p}(U)}} (\subset W^{k,p}(U))$. See [Ada75, Theorem 3.9] for example.
- (c) For every $k \in \mathbb{Z}$, we have equivalent norms $\|f\|_{W^{k+1,p}(U)} \approx \|f\|_{W^{k,p}(U)} + \sum_{j=1}^n \|\partial_j f\|_{W^{k,p}(U)}$ for $1 < p < \infty$, and $\|f\|_{C^{k+1,s}(U)} \approx \|f\|_{C^{k,s}(U)} + \sum_{j=1}^n \|\partial_j f\|_{C^{k,s}(U)}$ for $0 < s < 1$.
However they fail for $p = 1, \infty$: in both cases we have $\|f\|_{L^p(U)} \gtrsim \|f\|_{W^{-1,p}(U)} + \sum_{j=1}^n \|\partial_j f\|_{W^{-1,p}(U)}$ but the converse inequalities are false. This can done by using (b) and [Pre97, Theorem 5].

We use the standard convention for spaces of tempered distributions.

Definition 6. We use $\mathcal{S}(\mathbb{R}^n)$ for the Schwartz space and $\mathcal{S}'(\mathbb{R}^n)$ for the space of tempered distributions.

For an arbitrary open subset $U \subseteq \mathbb{R}^n$, we define $\mathcal{S}'(U) := \{\tilde{f}|_U : \tilde{f} \in \mathcal{S}'(\mathbb{R}^n)\} (\subset \mathcal{D}'(U))$ to be the space of distributions in U that has tempered distributional extension.

We let $\mathcal{S}(U) := \{f \in \mathcal{S}(\mathbb{R}^n) : f|_{\bar{U}^c} \equiv 0\} (\subset \mathcal{S}(\mathbb{R}^n))$ to be the space of Schwartz functions supported in \bar{U} .

Remark 7. By definition $\mathcal{S}'(U) = \mathcal{S}'(\mathbb{R}^n)/\{\tilde{f} \in \mathcal{S}'(\mathbb{R}^n) : \tilde{f}|_U = 0\}$ is the quotient space. Since $\mathcal{S}(\mathbb{R}^n)$ and $\mathcal{S}'(\mathbb{R}^n)$ are dual to each other with respect to their standard topologies, we see that $\mathcal{S}'(U) = \mathcal{S}(U)'$. See also [Tri95, Theorem 2.10.5/1].

The Besov and Triebel-Lizorkin spaces, along with their Morrey analogies, can be defined using Littlewood-Paley decomposition. Note that we do not use these characterizations directly in the proof.

Definition 8. Let $\lambda = (\lambda_j)_{j=0}^\infty$ be a sequence of Schwartz functions satisfying:

- The Fourier transform $\hat{\lambda}_0(\xi) = \int_{\mathbb{R}^n} \lambda_0(x) 2^{-2\pi i x \xi} dx$ satisfies $\text{supp } \hat{\lambda}_0 \subset \{|\xi| < 2\}$, $\hat{\lambda}_0|_{\{|\xi| < 1\}} \equiv 1$.
- $\lambda_j(x) = 2^{jn} \lambda_0(2^j x) - 2^{(j-1)n} \lambda_0(2^{j-1} x)$ for $j \geq 1$.

¹ $C_c^\infty(\bar{U}) \subset C^\infty(\bar{U})$ is the set of all bounded smooth functions $f : U \rightarrow \mathbb{R}$ such that $f|_{U \cap \{|x| > R\}} \equiv 0$ for some R (depending on f).

Let $0 < p, q \leq \infty$, $s \in \mathbb{R}$ and $0 \leq \tau \leq \frac{1}{p}$, we define the **Besov-type** norm $\|\cdot\|_{\mathcal{B}_{pq}^{s\tau}(\lambda)}$, the **Triebel-Lizorkin-type** norm $\|\cdot\|_{\mathcal{F}_{pq}^{s\tau}(\lambda)}$ ($p < \infty$) and the **Besov-Morrey** norm $\|\cdot\|_{\mathcal{N}_{pq}^{s\tau}(\lambda)}$ as follow:

$$\begin{aligned} \|f\|_{\mathcal{B}_{pq}^{s\tau}(\lambda)} &:= \sup_{x \in \mathbb{R}^n, J \in \mathbb{Z}} 2^{nJ\tau} \left(\sum_{j=\max(0, J)}^{\infty} 2^{jsq} \left(\int_{B(x, 2^{-j})} |\lambda_j * f(x)|^p dx \right)^{\frac{q}{p}} \right)^{\frac{1}{q}}; \\ \|f\|_{\mathcal{F}_{pq}^{s\tau}(\lambda)} &:= \sup_{x \in \mathbb{R}^n, J \in \mathbb{Z}} 2^{nJ\tau} \left(\int_{B(x, 2^{-j})} \left(\sum_{j=\max(0, J)}^{\infty} |2^{js} \lambda_j * f(x)|^q \right)^{\frac{p}{q}} dx \right)^{\frac{1}{p}}, \quad \text{provided } p < \infty; \\ \|f\|_{\mathcal{N}_{pq}^{s\tau}(\lambda)} &:= \left(\sum_{j=0}^{\infty} \sup_{x \in \mathbb{R}^n, J \in \mathbb{Z}} 2^{nJ\tau q + jsq} \left(\int_{B(x, 2^{-j})} |\lambda_j * f(x)|^p dx \right)^{\frac{q}{p}} \right)^{\frac{1}{q}}. \end{aligned}$$

Here for $q = \infty$ we take natural modifications by replacing the ℓ^q -sums with the supremums over j .

We define the corresponding spaces $\mathcal{A}_{pq}^{s\tau}(\mathbb{R}^n) = \{f \in \mathcal{S}'(\mathbb{R}^n) : \|f\|_{\mathcal{A}_{pq}^s(\mathbb{R}^n)} := \|f\|_{\mathcal{A}_{pq}^s(\lambda)} < \infty\}$ where $\mathcal{A} \in \{\mathcal{B}, \mathcal{F}, \mathcal{N}\}$, for a fixed choice of λ ($p < \infty$ for the \mathcal{F} -cases).

For arbitrary open $U \subseteq \mathbb{R}^n$, we define $\mathcal{A}_{pq}^{s\tau}(U) := \{\tilde{f}|_U : \tilde{f} \in \mathcal{A}_{pq}^{s\tau}(\mathbb{R}^n)\}$ with $\|f\|_{\mathcal{A}_{pq}^{s\tau}(U)} := \inf_{\tilde{f}|_U=f} \|\tilde{f}\|_{\mathcal{A}_{pq}^{s\tau}(\mathbb{R}^n)}$.

For the classical **Besov** and **Triebel-Lizorkin** spaces, we use

$$\begin{aligned} \mathcal{B}_{pq}^s(U) &:= \mathcal{B}_{p,q}^{s,0}(U) (= \mathcal{N}_{p,q}^{s,0}(U)), \quad \mathcal{F}_{pq}^s(U) := \mathcal{F}_{p,q}^{s,0}(U), \quad \text{for } p, q \in (0, \infty] \text{ and } s \in \mathbb{R} \quad (p < \infty \text{ for } \mathcal{F}\text{-cases}); \\ \mathcal{F}_{\infty q}^s(U) &:= \mathcal{F}_{q,q}^{s,\frac{1}{q}}(U), \quad \text{for } q \in (0, \infty] \text{ and } s \in \mathbb{R}. \end{aligned}$$

We shall see that $\mathcal{A}_{pq}^{s\tau}(U)$ is always a (quasi-)Banach space, and different choices of $(\lambda_j)_{j=0}^{\infty}$ result in equivalent norms. See [Tri10, Proposition 2.3.2] and [Tri20, Propositions 1.3 and 1.8].

Remark 9. In the definition we only consider $\tau \leq \frac{1}{p}$. If we extend them to $\tau > \frac{1}{p}$ then by [YY13, Theorem 2] and [Sic12, Lemma 3.4] we have $\mathcal{B}_{pq}^{s\tau}(\mathbb{R}^n) = \mathcal{F}_{pq}^{s\tau}(\mathbb{R}^n) = \mathcal{B}_{\infty}^{s+n(\tau-\frac{1}{p})}$, for all $0 < p, q \leq \infty$ and $s \in \mathbb{R}$.

When $\tau = \frac{1}{p}$, by [YY13, Theorem 2] and [Sic12, Remark 11(ii)] we have

$$(7) \quad \mathcal{B}_{p,\infty}^{s,\frac{1}{p}}(\mathbb{R}^n) = \mathcal{F}_{p,\infty}^{s,\frac{1}{p}}(\mathbb{R}^n) = \mathcal{B}_{\infty,\infty}^s(\mathbb{R}^n), \quad \mathcal{N}_{p,q}^{s,\frac{1}{p}}(\mathbb{R}^n) = \mathcal{B}_{\infty,q}^s(\mathbb{R}^n), \quad \forall 0 < p, q \leq \infty, \quad s \in \mathbb{R}.$$

See also [Tri20, Proposition 1.18].

Our notation $\mathcal{N}_{pq}^{s\tau}$ corresponds to the $\mathcal{B}_{pq}^{s\tau}$ in [Sic12, Definition 5]. For the usual conventions of Besov-Morrey spaces and Triebel-Lizorkin-Morrey spaces: we have $\mathcal{N}_{u,q,p}^s(\mathbb{R}^n) = \mathcal{N}_{p,q}^{s,\frac{1}{p}-\frac{1}{u}}(\mathbb{R}^n)$ and $\mathcal{E}_{u,q,p}^s(\mathbb{R}^n) = \mathcal{F}_{p,q}^{s,\frac{1}{p}-\frac{1}{u}}(\mathbb{R}^n)$ see [Sic12, Remark 13(iii) and Proposition 3.3 (iii)].

For the notations in [Tri14, HT21], we have the correspondence $L^r \mathcal{A}_{pq}^s(\mathbb{R}^n) = \Lambda^{pr} \mathcal{A}_{pq}^s(\mathbb{R}^n) = \mathcal{A}_{p,q}^{s,\frac{1}{p}+\frac{r}{n}}(\mathbb{R}^n)$ for $\mathcal{A} \in \{\mathcal{B}, \mathcal{F}\}$ and $\Lambda_{pr} \mathcal{B}_{pq}^s(\mathbb{R}^n) = \mathcal{N}_{p,q}^{s,\frac{1}{p}+\frac{r}{n}}(\mathbb{R}^n)$, for $0 < p, q \leq \infty$, $s \in \mathbb{R}$ and $-\frac{n}{p} \leq r \leq 0$ ($p < \infty$ if $\mathcal{A} = \mathcal{F}$). See also [HT21, Section 2.1].

For more discussions, we refer [YSY10, Sic12, Tri14, HT21] to readers.

In the following, we will use the notation $x \lesssim y$ to mean $x \leq Cy$ where C is a constant independent of x, y , and $x \approx y$ for “ $x \lesssim y$ and $y \lesssim x$ ”. We use $x \lesssim_{\varepsilon} y$ to emphasize the dependence of C on the parameter ε .

For $r \neq 0$ we use ϑ^r as the dilation operator given by

$$(8) \quad \vartheta^r f(x', x_n) := f(x', rx_n)$$

We use M_{pq}^{ns} and \widehat{M}_{pq}^{ns} for some fixed positive constants in Proposition 22 and Theorem 23, both of which depend on $n \geq 1$, $0 < p, q \leq \infty$ and $s \in \mathbb{R}$.

We let $S = S_n$ be the zero extension operator of \mathbb{R}_+^n , that is

$$(9) \quad Sf(x) = S_n f(x) := \mathbf{1}_{\mathbb{R}_+^n}(x) \cdot f(x) = \begin{cases} f(x', x_n) & x_n > 0 \\ 0 & x_n < 0 \end{cases}.$$

Remark 10. We have the definedness and boundedness $S_n : \mathcal{A}_{pq}^s(\mathbb{R}_+^n) \rightarrow \mathcal{A}_{pq}^s(\mathbb{R}^n)$ for $\max(n(\frac{1}{p}-1), \frac{1}{p}-1) < s < \frac{1}{p}$, see [Tri20, Theorem 2.48]. We will use the case $n = 1$ and the case $p = \infty$ in the proof of Theorem 1 (iii).

3. THE CONSTRUCTION OF E : PROOF OF THEOREM 1 (I) AND (II)

First we need to fulfill condition (3). In [See64] Seeley took $b_j := 2^j$ for $j \geq 0$ and found $(a_j)_{j=0}^\infty$ such that $\sum_{j=0}^\infty a_j (-b_j)^k = 1$ for all $k \geq 0$. This can be done by constructing an entire function $F(z) = \sum_{j=0}^\infty a_j z^j$ such that $F(2^k) = (-1)^k$, we recall the construction below.

Lemma 11. (i) For $\beta > 1$, $W_\beta(z) := \prod_{j=0}^\infty (1 - z/\beta^j)$ ($z \in \mathbb{C}$) defines a nonzero entire function W_β , all of whose zeros are simple.

(ii) Let $u = (u_k)_{k=0}^\infty \in \ell^\infty$. The function F_u^β below is entire and satisfies $F_u^\beta(\beta^k) = u_k$, for all $k \geq 0$:

$$(10) \quad F_u^\beta(z) := \sum_{k=0}^\infty \frac{u_k}{W_\beta'(\beta^k)} \cdot \frac{W_\beta(z)}{z - \beta^k}.$$

(iii) When $\beta > 2$, we have the following quantitative estimate for $l = 1, 2, 3, \dots$:

$$(11) \quad \sum_{k=1}^\infty \left| \frac{W_\beta(\beta^{-l})}{W_\beta'(\beta^k) \cdot (\beta^{-l} - \beta^k)} \right| \leq \frac{\beta^3 \exp \frac{2}{\beta-1}}{(\beta^2 - 1)(\beta^2 - \beta - 1)}.$$

Proof. (i) is the result of the Weierstrass factorization theorem, see [Rud87, Theorem 15.9] for example.

For (ii), one can see that for $k \geq 0$,

$$(12) \quad W_\beta'(\beta^k) = \frac{W_\beta(z)}{z - \beta^k} \Big|_{z=\beta^k} = -\beta^{-k} \prod_{j \geq 0; j \neq k} (1 - \beta^k/\beta^j) = -\beta^{-k} W_\beta(\beta^{-1}) \cdot \prod_{l=1}^k (1 - \beta^l).$$

Since $\beta^l - 1 \geq \beta^{l-1}(\beta - 1)$ for $l \geq 1$, we see that for $k \geq 0$:

$$(13) \quad |W_\beta'(\beta^k)| \geq |W_\beta(\beta^{-1})| \beta^{-k} \prod_{l=1}^k \beta^{l-1}(\beta - 1) = |W_\beta(\beta^{-1})| (\beta - 1)^k \beta^{-k} \cdot \beta^{\frac{k^2-k}{2}}.$$

Since $\beta > 1$, we see that the sum $\sum_{k=0}^\infty u_k/W_\beta'(\beta^k)$ absolutely converges. Therefore the sum in (10) converges absolutely and locally uniformly in z . We conclude that F_u^β is indeed an entire function.

For each $j \geq 0$, we have $\frac{W_\beta(z)}{z - \beta^k} \Big|_{z=\beta^j} = W_\beta'(\beta^k)$ when $j = k$ and $\frac{W_\beta(z)}{z - \beta^k} \Big|_{z=\beta^j} = 0$ when $j \neq k$. Therefore $F_u^\beta(\beta^k) = u_k$, as for each $z = \beta^k$ there is only one nonzero term in the sum (ii).

For (iii), note that for each $l \geq 1$, $W_\beta(\beta^{-l}) = |W_\beta(\beta^{-l})| < 1$ because $|1 - \beta^{-l}/\beta^j| < 1$ for each $j \geq 0$. Therefore using (12) and (13) we see that for $k, l \geq 1$,

$$\left| \frac{W_\beta(\beta^{-l})}{W_\beta'(\beta^k)(\beta^{-l} - \beta^k)} \right| \leq \frac{W_\beta(\beta^{-1})^{-1} \cdot W_\beta(\beta^{-l}) \cdot \beta^k}{(\beta - 1)^k \beta^{\frac{k-1}{2}k} \cdot |\beta^{-l} - \beta^k|} \leq \frac{W_\beta(\beta^{-1})^{-1}}{(\beta - 1)^k \beta^{k-1}} \cdot \frac{1}{1 - \beta^{-k-l}} \leq \frac{W_\beta(\beta^{-1})^{-1}}{(\beta^2 - \beta)^k} \cdot \frac{\beta}{1 - \beta^{-2}}.$$

On the other hand

$$\begin{aligned} (W_\beta(\beta^{-1}))^{-1} &= \prod_{j=1}^\infty (1 - \beta^{-j})^{-1} = \exp \sum_{j=1}^\infty \log \frac{1}{1 - \beta^{-j}} \\ &< \exp \sum_{j=1}^\infty \log(1 + 2\beta^{-j}) && \left(\frac{1}{1-x} < 1 + 2x \text{ when } 0 < x < \frac{1}{2} \right) \\ &< \exp \sum_{j=1}^\infty 2\beta^{-j} = \exp \frac{2}{\beta - 1} && (\log(1 + x) < x \text{ when } 0 < x < 1). \end{aligned}$$

Therefore taking sum over $k \geq 1$ we obtain (11):

$$\sum_{k=1}^\infty \left| \frac{W_\beta(\beta^{-l})}{W_\beta'(\beta^k)(\beta^{-l} - \beta^k)} \right| \leq \frac{\beta}{1 - \beta^{-2}} \sum_{k=1}^\infty \frac{\exp \frac{2}{\beta-1}}{(\beta^2 - \beta)^k} = \frac{\beta^3 \exp \frac{2}{\beta-1}}{(\beta^2 - 1)(\beta^2 - \beta - 1)}. \quad \square$$

Remark 12. If one takes $u_k := (-1)^k$ in Lemma 11 (ii), then the corresponding sequence $a_j := \frac{1}{j!} (F_u^\beta)^{(j)}(0)$ ($j = 0, 1, 2, \dots$) satisfies $\sum_j a_j \beta^{jk} = (-1)^k$ for all $k \geq 0$. This is enough to prove the boundedness results in [See64] (C^k for $k \geq 0$), but in our case we also need the equality to be true for $k < 0$. However, one cannot construct an entire function F such that $F(2^{-k}) = (-1)^{-k}$ for all $k \geq 0$.

Remark 13. Alternatively, one can construct the $(a_j)_j$ using Vandermonde determinant, which is the method of [See64]. Let $M_1, M_2 \in \mathbb{N}_0$ and let $b_{-M_1}, \dots, b_{M_2} > 0$ be distinct numbers. By Cramer's rule we see that the solutions of the equation system $\sum_{j=-M_1}^{M_2} a_j b_j^k = (-1)^k$ for $-M_1 \leq k \leq M_2$ are (uniquely) determined by the following (see [Ada75, Page 149] for example):

$$(14) \quad a_j = (-b_j)^{M_1} \frac{V(b_{-M_1}, \dots, b_{j-1}, -1, b_{j+1}, \dots, b_{M_2})}{V(b_{-M_1}, \dots, b_{M_2})} = (-b_j)^{M_1} \prod_{\substack{-M_1 \leq k \leq M_2 \\ k \neq j}} \frac{b_k + 1}{b_k - b_j}, \quad -M_1 \leq j \leq M_2.$$

Here $V(x_{-M_1}, \dots, x_{M_2}) = \prod_{-M_1 \leq u < v \leq M_2} (x_v - x_u)$.

Now we set $b_j := \beta^j$ for some $\beta > 1$. If we let $M_1 = 0$ and let $M_2 \rightarrow +\infty$, we see that (14) converges and produces the same sequence $(a_j)_{j=0}^\infty$ to Remark 12.

We find $(a_j)_{j=-\infty}^\infty$ through the following:

Proposition 14. *There are real numbers $(\tilde{a}_j)_{j=0}^\infty$ such that $\sum_{j=0}^\infty \tilde{a}_j (4^{jk} + 4^{-jk}) = (-1)^k$ for all $k = 0, 1, 2, \dots$, with all the summations converging absolutely.*

Therefore the sequences $a_j := \tilde{a}_{|j|}$ ($j \neq 0$), $a_0 := \frac{1}{2}\tilde{a}_0$ and $b_j := 4^j$ ($j \in \mathbb{Z}$) fulfill condition (3).

The idea is to construct an iterated sequence $\tilde{a}^\nu = (\tilde{a}_0^\nu, \tilde{a}_1^\nu, \tilde{a}_2^\nu, \dots)$ such that

$$\sum_{j=0}^\infty \tilde{a}_j^\nu 4^{jk} = (-1)^k - \sum_{j=0}^\infty \tilde{a}_j^{\nu-1} 4^{-jk}, \quad \forall k \geq 0.$$

The limit $\lim_{\nu \rightarrow \infty} \tilde{a}_j^\nu$ will be our \tilde{a}_j , provided that it is convergent. In practice, it is more convenient to prove the convergence of $(u^\nu)_{\nu=0}^\infty \subset \ell^\infty$ via the correspondence $u_k^\nu := \sum_{j=0}^\infty \tilde{a}_j^\nu 4^{jk}$ and $\tilde{a}_j^\nu = \frac{1}{j!} \frac{d^j}{dz^j} F_{u^\nu}^4(z)|_{z=0}$, where $F_{u^\nu}^4(z)$ is the entire function in Lemma 11 (ii) with $\beta = 4$ and $u = u^\nu$.

Proof. Let $u^0 = (u_k^0)_{k=0}^\infty \in \ell^\infty$ be $u_0^0 := \frac{1}{2}$ and $u_k^0 := (-1)^k$ for $k \geq 1$. For each $\nu \geq 1$, we let $u_0^\nu := \frac{1}{2}$ and

$$u_k^\nu := (-1)^k - \sum_{j=0}^\infty \left(\frac{1}{j!} \frac{d^j}{dz^j} F_{u^{\nu-1}}^4(z) \Big|_{z=0} \right) 4^{-jk} = (-1)^k - F_{u^{\nu-1}}^4(4^{-k}), \quad \text{for } k \geq 1.$$

By Lemma 11 (ii), if $u^{\nu-1} \in \ell^\infty$ then $F_{u^{\nu-1}}^4(z)$ is entire in $|z| < 1$. Since $u^0 \in \ell^\infty$, by recursion we get $\|u^\nu\|_{\ell^\infty} \leq 1 + \|F_{u^{\nu-1}}\|_{L^\infty[0,1]} < \infty$ for every $\nu \geq 1$. Thus we have a sequence of bounded sequences $(u^\nu)_{\nu=0}^\infty \subset \ell^\infty$.

Moreover since $u_0^\nu \equiv \frac{1}{2}$, by (11),

$$\begin{aligned} \|u^{\nu+1} - u^\nu\|_{\ell^\infty} &= \sup_{l \geq 1} |u_l^{\nu+1} - u_l^\nu| \leq \sup_{l \geq 1} |F_{u^\nu}^4(4^{-l}) - F_{u^{\nu-1}}^4(4^{-l})| \\ &\leq \sup_{l \geq 1} \sum_{k=1}^\infty \left| \frac{(u_k^\nu - u_k^{\nu-1}) \cdot W_4(4^{-l})}{W_4^4(4^k) \cdot (4^{-l} - 4^k)} \right| \leq \|u^\nu - u^{\nu-1}\|_{\ell^\infty} \frac{4^3 \exp \frac{2}{4-1}}{(4^2 - 1)(4^2 - 4 - 1)} = \frac{64e^{\frac{2}{3}}}{165} \cdot \|u^\nu - u^{\nu-1}\|_{\ell^\infty}. \end{aligned}$$

Here $\frac{64e^{2/3}}{165} \approx 0.755 < 1$. We conclude that $(u^\nu)_{\nu=0}^\infty \subset \ell^\infty$ is a contraction sequence and must have a limit $u^\infty \in \ell^\infty$. Clearly $u_0^\infty = \frac{1}{2}$ and $u_k^\infty = (-1)^k - F_{u^\infty}^4(4^{-k})$ for $k \geq 1$.

By Lemma 11 (ii), the function $F_{u^\infty}^4(z)$ is entire. We define \tilde{a}_j in the way that $F_{u^\infty}^4(z) = \sum_{j=0}^\infty \tilde{a}_j z^j$, thus

$$\begin{aligned} \sum_{j=0}^\infty \tilde{a}_j &= u_0^\infty = 1 - u_0^\infty = (-1)^0 - \sum_{j=0}^\infty \tilde{a}_j, & k = 0, \\ \sum_{j=0}^\infty \tilde{a}_j 4^{jk} &= u_k^\infty = (-1)^k - F_{u^\infty}^4(4^{-k}) = (-1)^k - \sum_{j=0}^\infty \tilde{a}_j 4^{-jk}, & k \geq 1. \end{aligned}$$

Since $F_{u^\infty}^4$ is entire, the sum $\sum_{j=0}^\infty \tilde{a}_j 4^{jk} (= F_{u^\infty}^4(4^k))$ always converge absolutely. Therefore $(\tilde{a}_j)_{j=0}^\infty$ is as desired.

Let $a_j := \tilde{a}_{|j|}$ for $j \neq 0$ and $a_0 := \frac{1}{2}\tilde{a}_0$, we see that $\sum_{j=-\infty}^\infty a_j 4^{jk} = (-1)^k$ for all $k \in \mathbb{Z}$ with all summations converging absolutely. In particular $\sum_{j=-\infty}^\infty |a_j| 4^{|j|k} < \infty$ for all $k \geq 1$, which means that $\sum_{j=-\infty}^\infty |a_j| 2^{\delta|j|} 4^{jk} < \infty$ for all $\delta > 0$ and $k \in \mathbb{Z}$. Therefore $(a_j, 4^j)_{j=-\infty}^\infty$ satisfies the condition (3) and we prove Theorem 1 (i). \square

The direct computations yield the following. We omit the proof to reader.

Lemma 15. (i) Let $(a, b) = (a_j, b_j)_{j=-\infty}^{\infty}$ be the collection satisfying the condition (3). Then for every $k \in \mathbb{Z}$ the collections $(a(-b)^k, b) := (a_j(-b_j)^k, b_j)_{j=-\infty}^{\infty}$ and $(a(-b)^k, \frac{1}{b}) := (a_j(-b_j)^k, \frac{1}{b_j})_{j=-\infty}^{\infty}$ also satisfy (3).

(ii) For multi-index $\gamma \in \mathbb{N}_0^n$ we have $\partial^\gamma \circ E^{a,b} = E^{a(-b)^{\gamma_n}, b} \circ \partial^\gamma$ and $E^{a,b} \circ \partial^\gamma = \partial^\gamma \circ E^{a(-b)^{-\gamma_n}, b}$.

(iii) Let $E = E^{a,b}$ be the extension operator defined in (2). Then

$$(15) \quad E^* g(x', x_n) = g(x', x_n) - \sum_{j=-\infty}^{\infty} \frac{a_j}{-b_j} g(x', -\frac{1}{b_j} \cdot x_n), \quad x' \in \mathbb{R}^{n-1}, \quad x_n > 0.$$

Remark 16. In the notations of (8) and (9) we can write $E^{a,b} = S + \sum_j a_j \vartheta^{-b_j} \circ S$. For $r < 0$ we have

$$\partial_{x_n} (\vartheta^r S f)(x) = r(S \partial_{x_n} f)(x', r x_n) - \delta_0(x_n) f(x', r x_n),$$

where δ_0 is the Dirac measure at $0 \in \mathbb{R}$.

Therefore $\partial_n \circ E^{a,b} = E^{a(-b), b} \circ \partial_n$ (in the domain where either side is defined) if and only if $\sum_j a_j = 1$. Taking higher order derivatives we see that $\partial^\gamma \circ E^{a,b} = E^{a(-b)^{\gamma_n}, b} \circ \partial^\gamma$ still holds if we only assume the condition $\sum_j a_j (-b_j)^k = 1$ for $0 \leq k \leq \gamma_n$.

Lemma 17. Let $(X, \|\cdot\|)$ be a quasi-Banach space such that $(x, y) \mapsto \|x - y\|^q$ is a metric for some $0 < q \leq 1$, i.e. $\|x + y\|^q \leq \|x\|^q + \|y\|^q$ for all $x, y \in X$. Then for any $\delta > 0$ there is a constant $K_{q,\delta} > 0$ such that $\|\sum_{j=-\infty}^{\infty} x_j\| \leq K_{q,\delta} \sum_{j=-\infty}^{\infty} 2^{\delta|j|} \|x_j\|$ for all sequence $(x_j)_{j \in \mathbb{Z}} \subset X$ such that right hand summation converges. In this case the sum $\sum_{j=-\infty}^{\infty} x_j$ converges with respect to the quasi-norm topology of X .

Remark 18. The couple $(X, \|\cdot\|^q)$ is also call a q -Banach space.

In practice we use $X = W^{k,q}(U)$ and $X = \mathcal{S}'_{pr}(U)$ (see Definition 8) with $q = \min(p, r, 1)$.

Proof of Lemma 17. When $(x_j)_{j \in \mathbb{Z}}$ are finitely nonzero, by Hölder's inequality we have

$$\left\| \sum_j x_j \right\|_X \leq \left| \sum_j \|x_j\|_X^q \right|^{1/q} \leq \left(\left\| \sum_{j \in \mathbb{Z}} (2^{-\delta q |j|})_{j \in \mathbb{Z}} \left\| (2^{\delta q |j|} \|x_j\|_X^q)_{j \in \mathbb{Z}} \right\|_{\ell^{\frac{1}{1-q}}(\mathbb{Z})} \right)^{1/q} = K_{q,\delta} \sum_j 2^{\delta|j|} \|x_j\|_X.$$

Here $K_{q,\delta} := (\frac{2^{\delta/(1-q)} + 1}{2^{\delta/(1-q)} - 1})^{1/q-1}$ if $q < 1$ and $K_{q,\delta} := 1$ if $q = 1$.

When $\sum_{j=-\infty}^{\infty} 2^{\delta|j|} \|x_j\| < \infty$, we have $\lim_{\nu \rightarrow +\infty} \sup_{\nu \leq \nu' < \nu''} \left\| \sum_{\nu' < |j| \leq \nu''} x_j \right\| \leq K_{q,\delta} \lim_{\nu \rightarrow +\infty} \sum_{|j| \geq \nu} 2^{\delta|j|} \|x_j\| = 0$. Therefore $(\sum_{|j| \leq \nu} x_j)_{\nu=1}^{\infty}$ is a Cauchy sequence in X . Since X is complete, the sum must converge to a unique limit element and the same inequality $\|\sum_j x_j\| \leq K_{q,\delta} \sum_j 2^{\delta|j|} \|x_j\|$ holds. \square

The proof of boundedness on Sobolev spaces is standard:

Proof of Theorem 1 (ii). We only need to prove the Sobolev boundedness $W^{k,p}$ for $k \geq 0$, $0 < p \leq \infty$.

Indeed suppose we get $E : W^{k,\infty}(\mathbb{R}_+^n) \rightarrow W^{k,\infty}(\mathbb{R}^n)$, then $C^\infty(U) = \bigcap_{k=0}^{\infty} W^{k,\infty}(U)$ for $U \in \{\mathbb{R}^n, \mathbb{R}_+^n\}$ gives $E : C^\infty(\mathbb{R}_+^n) \rightarrow C^\infty(\mathbb{R}^n)$. Clearly $C^k(U) = \overline{C^\infty(U)}^{\|\cdot\|_{W^{k,\infty}(U)}}$ for $U \in \{\mathbb{R}^n, \mathbb{R}_+^n\}$, thus $E : C^k(\mathbb{R}_+^n) \rightarrow C^k(\mathbb{R}^n)$ holds for all $k \geq 0$.

The $C^{k,s}$ boundedness ($k \in \mathbb{Z}$, $0 < s < 1$) can follow either from the same argument for $W^{k,p}$ below, or the Besov correspondence $C^{k,s}(U) = \mathcal{B}_{\infty\infty}^{k+s}(U)$, for $U \in \{\mathbb{R}^n, \mathbb{R}_+^n\}$ (see Proposition 21 (ii)). When $k \geq 0$ one can also obtain the boundedness by the interpolation $C^{k,s}(U) = (C^k(U), C^{k+1}(U))_{s,\infty}$ (see for example [Tri95, Theorem 2.7.2/1] and the proof of [Tri95, Theorem 4.5.2/1]).

For the $W^{k,p}$ -boundedness, by Lemma 17, for every $f \in L^p(\mathbb{R}_+^n)$ if $1 \leq p \leq \infty$, and every $f \in C_c^\infty(\overline{\mathbb{R}_+^n})$ if $0 < p < 1$, we have (recall the notation ϑ^r in (8))

$$(16) \quad \begin{aligned} \|E^{a,b} f\|_{L^p(\mathbb{R}^n)} &\leq 2^{\max(0, \frac{1}{p}-1)} (\|f\|_{L^p(\mathbb{R}_+^n)} + \left\| \sum_{j \in \mathbb{Z}} a_j \cdot \vartheta^{-b_j} f \right\|_{L^p(\mathbb{R}_+^n)}) \\ &\lesssim_{p,\delta} \|f\|_{L^p(\mathbb{R}_+^n)} + \sum_{j \in \mathbb{Z}} 2^{\delta|j|} |a_j| \left(\int_{\mathbb{R}_+^n} |f(x', -b_j x_n)|^p dx' dx_n \right)^{\frac{1}{p}} \\ &= \|f\|_{L^p(\mathbb{R}_+^n)} + \sum_{j \in \mathbb{Z}} 2^{\delta|j|} |a_j| \left(b_j^{-1} \int_{\mathbb{R}_+^n} |f(x', y_n)|^p dx' dy_n \right)^{\frac{1}{p}} \\ &= \left(1 + \sum_{j \in \mathbb{Z}} 2^{\delta|j|} |a_j| b_j^{-\frac{1}{p}} \right) \|f\|_{L^p(\mathbb{R}_+^n)}. \end{aligned}$$

When $k \geq 0$, also for every $f \in L^p(\mathbb{R}_+^n)$ if $1 \leq p \leq \infty$, and every $f \in C_c^\infty(\overline{\mathbb{R}_+^n})$ if $0 < p < 1$, applying Lemma 15 (ii) to (16) we have

$$\begin{aligned} \|E^{a,b}f\|_{W^{k,p}(\mathbb{R}^n)} &\lesssim_{k,p} \sum_{|\gamma| \leq k} \|\partial^\gamma E^{a,b}f\|_{L^p(\mathbb{R}^n)} = \sum_{|\gamma| \leq k} \|E^{a(-b)^{\gamma_n}, b} \partial^\gamma f\|_{L^p(\mathbb{R}^n)} \\ &\lesssim_{p,\delta} \sum_{|\gamma| \leq k} \sum_{j \in \mathbb{Z}} 2^{\delta|j|} |a_j| b_j^{\gamma_n - \frac{1}{p}} \|\partial^\gamma f\|_{L^p(\mathbb{R}_+^n)} \lesssim_{k,p} \sum_{j \in \mathbb{Z}} 2^{\delta|j|} |a_j| (b_j^{-\frac{1}{p}} + b_j^{k-\frac{1}{p}}) \|f\|_{W^{k,p}(\mathbb{R}_+^n)}. \end{aligned}$$

By the assumption (3) we can take $\delta > 0$ small such that $\sum_{j \in \mathbb{Z}} 2^{\delta|j|} |a_j| b_j^{k-\lceil \frac{1}{p} \rceil} < \infty$ and $\sum_{j \in \mathbb{Z}} 2^{\delta|j|} |a_j| b_j^k < \infty$. We now get the boundedness of $W^{k,p}$ for $k \geq 0$.

When $k < 0$, applying Lemma 15 (ii) to (16) for

- every $f \in W^{k,p}(\mathbb{R}_+^n)$ and $\{g_\gamma\}_{|\gamma| \leq -k} \subset L^p(\mathbb{R}_+^n)$ such that $f = \sum_{|\gamma| \leq -k} \partial^\gamma g_\gamma$, if $1 \leq p \leq \infty$;
- every $f \in C_c^\infty(\overline{\mathbb{R}_+^n})$ and $\{g_\gamma\}_{|\gamma| \leq -k} \subset C_c^\infty(\overline{\mathbb{R}_+^n})$ such that $f = \sum_{|\gamma| \leq -k} \partial^\gamma g_\gamma$, if $0 < p < 1$,

we have the following

$$\begin{aligned} \|E^{a,b}f\|_{W^{k,p}(\mathbb{R}^n)} &\lesssim_{k,p} \sum_{|\gamma| \leq -k} \|E^{a,b} \partial^\gamma g_\gamma\|_{W^{k,p}(\mathbb{R}^n)} = \sum_{|\gamma| \leq -k} \|\partial^\gamma E^{a(-b)^{-\gamma_n}, b} g_\gamma\|_{W^{k,p}(\mathbb{R}^n)} \\ &\leq \sum_{|\gamma| \leq -k} \|E^{a(-b)^{-\gamma_n}, b} g_\gamma\|_{L^p(\mathbb{R}^n)} \lesssim_{p,\delta} \sum_{|\gamma| \leq -k} \sum_{j \in \mathbb{Z}} 2^{\delta|j|} |a_j| b_j^{-\gamma_n - \frac{1}{p}} \|g_\gamma\|_{L^p(\mathbb{R}_+^n)} \\ &\leq \sum_{j \in \mathbb{Z}} 2^{\delta|j|} |a_j| (b_j^{-\frac{1}{p}} + b_j^{k-\frac{1}{p}}) \sum_{|\gamma| \leq -k} \|g_\gamma\|_{L^p(\mathbb{R}_+^n)}. \end{aligned}$$

Taking the infimum over all $\{g_\gamma\}_{|\gamma| \leq -k}$ for f , and taking δ from (3) such that $\sum_{j \in \mathbb{Z}} 2^{\delta|j|} |a_j| (b_j^k + b_j^{k-\lceil \frac{1}{p} \rceil}) < \infty$ as well, we get the $W^{k,p}$ -boundedness for $k < 0$, finishing the proof. \square

Remark 19. The result can be extended to vector-valued functions. Indeed let $0 < q \leq 1$, $0 < p < \infty$, and let $(X, |\cdot|_X)$ be a quasi Banach space such that $|x+y|_X^q \leq |x|_X^q + |y|_X^q$. Then for every L^p (strongly measurable) function $f : \mathbb{R}_+^n \rightarrow X$, by the same calculation to (16),

$$\begin{aligned} &\left(\int_{\mathbb{R}^n} \left| \sum_{j \in \mathbb{Z}} a_j f(x', -b_j x_n) \right|_X^p dx' dx_n \right)^{\frac{1}{p}} \leq K_{q, \frac{\delta}{2}} \left(\int_{\mathbb{R}_+^n} \left(\sum_{j \in \mathbb{Z}} 2^{|j| \frac{\delta}{2}} |a_j| |f(x', -b_j x_n)|_X \right)^p dx' dx_n \right)^{\frac{1}{p}} \\ &\leq K_{q, \frac{\delta}{2}} K_{p, \frac{\delta}{2}} \sum_{j \in \mathbb{Z}} 2^{|j| (\frac{\delta}{2} + \frac{\delta}{2})} |a_j| \left(\int_{\mathbb{R}_+^n} b_j^{-1} |f(x', y_n)|_X^p dx' dy_n \right)^{\frac{1}{p}} = \left(K_{q, \frac{\delta}{2}} K_{p, \frac{\delta}{2}} \sum_{j \in \mathbb{Z}} 2^{|j| \delta} |a_j| b_j^{-\frac{1}{p}} \right) \|f\|_{L^p(\mathbb{R}_+^n; X)}. \end{aligned}$$

Here $K_{q, \delta/2}$ and $K_{p, \delta/2}$ are the constants in Lemma 17. Also by Lemma 17 the sum of functions $\sum_{j \in \mathbb{Z}} a_j f(x', -b_j x_n)$ converges in $L^p(\mathbb{R}^n; X)$.

Therefore $E^{a,b}$ defines a bounded linear map $E^{a,b} : L^p(\mathbb{R}_+^n; X) \rightarrow L^p(\mathbb{R}^n; X)$ with the control of operator norms $\|E^{a,b}\|_{L^p(\mathbb{R}_+^n; X) \rightarrow L^p(\mathbb{R}^n; X)} \lesssim_{p,q,\delta} \sum_{j \in \mathbb{Z}} 2^{|j| \delta} |a_j| b_j^{-1/p}$.

In practice we will take $X = \mathcal{D}_{pq}^s(\mathbb{R}^{n-1})$ in the proof of Theorem 23, Step A3.

4. BOUNDEDNESS ON BESOV AND TRIEBEL-LIZORKIN: PROOF OF THEOREM 1 (III) AND (IV)

Before we discuss the boundedness on Besov and Triebel-Lizorkin spaces, for completeness we prove that $E : \mathcal{S}' \rightarrow \mathcal{S}'$ is defined. Recall Definition 6 that $\mathcal{S}'(\mathbb{R}_+^n) = \mathcal{S}'(\mathbb{R}^n) / \{\tilde{f} : \tilde{f}|_{\mathbb{R}_+^n} = 0\}$.

Lemma 20. $E : \mathcal{S}'(\mathbb{R}_+^n) \rightarrow \mathcal{S}'(\mathbb{R}^n)$ is defined and bounded.

Proof. Since $\mathcal{S}'(\mathbb{R}_+^n) = \mathcal{S}(\mathbb{R}_+^n)'$ (see Remark 7), by duality it is equivalent to show that $E^* : \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}(\mathbb{R}_+^n)$ is continuous.

Let $g \in \mathcal{S}(\mathbb{R}^n)$, using (15) we see that for every $\alpha, \beta \in \mathbb{N}_0^n$ and $x \in \mathbb{R}_+^n$,

$$|x^\alpha \partial^\beta E^* g(x)| \leq |x^\alpha \partial^\beta g(x)| + \sum_{j \in \mathbb{Z}} |a_j (-b_j)^{\alpha_n - \beta_n - 1} (x', -\frac{1}{b_j} \cdot x_n)^\alpha (\partial^\beta g)(x', -\frac{x_n}{b_j})|.$$

By (3), $\sum_{j \in \mathbb{Z}} |a_j| \cdot b_j^{\alpha_n - \beta_n - 1} < \infty$. We conclude that $|x^\alpha \partial^\beta E^* g(x)| \lesssim_{a,b,\alpha,\beta} \sup_{y \in \mathbb{R}^n} |y^\alpha \partial^\beta g(y)|$ for all $x \in \mathbb{R}_+^n$.

On the other hand, by the condition $\sum_j a_j(-b_j)^{-k} = 1$ for $k = 1, 2, 3, \dots$ in (3), we have

$$\lim_{x_n \rightarrow 0^+} \partial^\beta E^* g(x) = \partial^\beta g(x', 0) - \sum_{j \in \mathbb{Z}} a_j(-b_j)^{-\beta_n-1} (\partial^\beta g)(x', 0) = 0, \quad \forall \beta \in \mathbb{N}_0^n.$$

Therefore along with the zero extension to \mathbb{R}_-^n , E^*g defines a Schwartz function in \mathbb{R}^n which has support $\overline{\mathbb{R}_+^n}$. Moreover $\sup_{x \in \mathbb{R}^n} |x^\alpha \partial^\beta E^* g(x)| \lesssim_{\alpha, \beta} \sup_{y \in \mathbb{R}^n} |y^\alpha \partial^\beta g(y)|$ with the implied constant being independent of g .

Since the seminorms $\{\sup_{x \in \mathbb{R}^n} |x^\alpha \partial^\beta E^* g(x)|\}_{\alpha, \beta \in \mathbb{N}_0^n}$ define the topology of $\mathcal{S}(\mathbb{R}^n)$, we conclude that $E^* : \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}(\mathbb{R}_+^n)$ is continuous. Therefore $E : \mathcal{S}'(\mathbb{R}_+^n) \rightarrow \mathcal{S}'(\mathbb{R}^n)$ is continuous as well. \square

We now turn to prove Theorem 1 (iii) and (iv).

To prove Theorem 1 (iii), it might be possible to use the mean oscillation approach in [Tri92, Chapter 4.5.5]. However [Tri92, (4.5.2/9)] fails when $(a_j)_j$ has infinite nonzero terms. This is critical if one wants to prove the Besov and Triebel-Lizorkin boundedness of Seeley's operator (see Remark 12) using such method. The authors do not know what the correct modification of [Tri92, (4.5.2/9)] is.

Instead, we adapt the arguments from [Tri10, Section 2.9]. The proof involves characteristic function multiplier [Tri10, Section 2.8.7], the Fubini decomposition [Tri10, Section 2.5.13] and interpolations. We recall these facts from literature.

Proposition 21. *Let $0 < p, q \leq \infty$, $s \in \mathbb{R}$ and $0 \leq \tau \leq \frac{1}{p}$. Let $U \subseteq \mathbb{R}^n$ be an arbitrary open subset.*

(i) *For $\mathcal{A} \in \{\mathcal{B}, \mathcal{F}\}$ and for every integer $m \geq 1$, $\mathcal{A}_{pq}^s(U) = \{\sum_{|\alpha| \leq m} \partial^\alpha g_\alpha : g_\alpha \in \mathcal{A}_{pq}^{s+m}(U)\}$ and has equivalent (quasi-)norm $\|f\|_{\mathcal{A}_{pq}^s(U)} \approx_{p,q,s,m} \inf \left\{ \sum_{|\alpha| \leq m} \|g_\alpha\|_{\mathcal{A}_{pq}^{s+m}(U)} : f = \sum_{|\alpha| \leq m} \partial^\alpha g_\alpha \text{ as distributions} \right\}$.*

(ii) *If $U \in \{\mathbb{R}^n, \mathbb{R}_+^n\}$ or U is a bounded smooth domain, then $C^{k,s}(U) = \mathcal{B}_{\infty\infty}^s(U)$ for all $k \in \mathbb{Z}$ and $0 < s < 1$, with equivalent norms.*

(iii) (Interpolations) *For $s_0, s_1 \in \mathbb{R}$ and $\theta \in (0, 1)$ such that $s_0 \neq s_1$ and $s = (1 - \theta)s_0 + \theta s_1$, we have*

$$(17) \quad \text{complex interpolation : } [\mathcal{F}_{pq}^{s_0}(\mathbb{R}^n), \mathcal{F}_{pq}^{s_1}(\mathbb{R}^n)]_\theta = \mathcal{F}_{pq}^s(\mathbb{R}^n);$$

$$(18) \quad \text{real interpolation : } (\mathcal{F}_{pp}^{s_0}(\mathbb{R}^n), \mathcal{F}_{pp}^{s_1}(\mathbb{R}^n))_{\theta, q} = \mathcal{B}_{pq}^s(\mathbb{R}^n);$$

$$(19) \quad (\mathcal{F}_{pp}^{s_0, \tau}(\mathbb{R}^n), \mathcal{F}_{pp}^{s_1, \tau}(\mathbb{R}^n))_{\theta, q} = \mathcal{N}_{pq}^{s, \tau}(\mathbb{R}^n), \quad \text{for } \tau \in [0, \frac{1}{p}] \quad (\text{provided } p < \infty).$$

(iv) (Fubini) *When $p < \infty$ or $p = q = \infty$, and when $s > n \max(0, \frac{1}{p} - 1, \frac{1}{q} - 1)$, we have decomposition*

$$\mathcal{F}_{pq}^s(\mathbb{R}^n) = L^p(\mathbb{R}_{x_n}; \mathcal{F}_{pq}^s(\mathbb{R}_{x'}^{n-1})) \cap L^p(\mathbb{R}_{x'}^{n-1}; \mathcal{F}_{pq}^s(\mathbb{R}_{x_n})), \text{ with equivalent (quasi-)norms.}$$

(v) (Local representations) *For $s < 0$, we have equivalent norms*

$$(20) \quad \|f\|_{\mathcal{B}_{pq}^{s, \tau}(\mathbb{R}^n)} \approx_{n,p,q,s,\tau} \sup_{x \in \mathbb{R}^n; R > 0} R^{-n\tau} \|f\|_{\mathcal{B}_{pq}^s(B(x,R))};$$

$$(21) \quad \|f\|_{\mathcal{F}_{pq}^{s, \tau}(\mathbb{R}^n)} \approx_{n,p,q,s,\tau} \sup_{x \in \mathbb{R}^n; R > 0} R^{-n\tau} \|f\|_{\mathcal{F}_{pq}^s(B(x,R))}, \quad \text{provided } p < \infty.$$

Proof. For (i), by [Tri20, Theorem 1.24] $\|\partial^\alpha \tilde{g}\|_{\mathcal{A}_{pq}^s(\mathbb{R}^n)} \lesssim \|\tilde{g}\|_{\mathcal{A}_{pq}^{s+m}(\mathbb{R}^n)}$ holds for $|\alpha| \leq m$. Note that $\tilde{g}|_U = g$ implies $\partial^\alpha \tilde{g}|_U = \partial^\alpha g$, thus taking restrictions we get $\|\sum_{|\alpha| \leq m} \partial^\alpha g_\alpha\|_{\mathcal{A}_{pq}^s(U)} \lesssim \sum_{|\alpha| \leq m} \|g_\alpha\|_{\mathcal{A}_{pq}^{s+m}(U)}$. Therefore $\|f\|_{\mathcal{A}_{pq}^s(U)} \lesssim \sum_{|\alpha| \leq m} \|g_\alpha\|_{\mathcal{A}_{pq}^{s+m}(U)}$ whenever $f = \sum_{|\alpha| \leq m} \partial^\alpha g_\alpha$.

Conversely for $f \in \mathcal{A}_{pq}^s(U)$, take an extension $\tilde{f} \in \mathcal{A}_{pq}^s(\mathbb{R}^n)$, we have $\tilde{f} = (I - \Delta)(I - \Delta)^{-1} \tilde{f} = \tilde{g}_0 + \sum_{j=1}^n \partial_j \tilde{g}_j$ where $\tilde{g}_0 := (I - \Delta)^{-1} \tilde{f}$ and $\tilde{g}_j := -\partial_j (I - \Delta)^{-1} \tilde{f}$ ($1 \leq j \leq n$). Thus by [Tri20, Theorems 1.22 and 1.24] we have $\sum_{j=0}^n \|\tilde{g}_j\|_{\mathcal{A}_{pq}^{s+1}(\mathbb{R}^n)} \lesssim \|\tilde{f}\|_{\mathcal{A}_{pq}^s(\mathbb{R}^n)}$. Taking infimum of all extensions \tilde{f} over f we get (i) for $m = 1$. Doing this recursively we get all $m \geq 1$.

For (ii), we have $C^{k,s}(\mathbb{R}^n) = \mathcal{B}_{\infty\infty}^{k+s}(\mathbb{R}^n)$ from [Tri10, Section 2.5.7] for $k \geq 0$ and $0 < s < 1$. Thus in this case $\mathcal{B}_{\infty\infty}^{k+s}(U) = \{\tilde{f}|_U : \tilde{f} \in C^{k,s}(\mathbb{R}^n)\} \subseteq C^{k,s}(U)$ for arbitrary open subset U . When $U = \mathbb{R}_+^n$ or U is bounded smooth, by the existence of extension operator $C^{k,s}(U) \rightarrow C^{k,s}(\mathbb{R}^n)$ (see [GT01, Lemma 6.37]) we get $C^{k,s}(U) \subseteq \mathcal{B}_{\infty\infty}^{k+s}(U)$. This proves the case $k \geq 0$.

Comparing the result (i) and (5) we see that $C^{k,s}(U) = \mathcal{B}_{\infty\infty}^{k+s}(U)$ for $k < 0$ as well.

See [Tri06, Corollary 1.111] and [HT21, (3.14)] for (iii).

See [Tri20, Theorem 3.25] and [Tri01, Theorem 4.4] for (iv).

See [Tri14, Theorem 3.64] for (v). In the reference the collection $\{B(x, 2^{-J}) : x \in \mathbb{R}^n, J \in \mathbb{Z}\}$ is replaced by $\{2Q_{J,v} := 2^{-J}(v + (-\frac{1}{2}, \frac{3}{2})^n) : J \in \mathbb{Z}, v \in \mathbb{Z}^n\}$, but there is no difference between the proofs. Note that one cannot

use $\{Q_{J,v} := 2^{-J}(v + (0, 1)^n) : J \in \mathbb{Z}, v \in \mathbb{Z}^n\}$ because for $J_0 \in \mathbb{Z}$, the set $\{x \in \mathbb{R}^n : x_k \in 2^{-J_0}\mathbb{Z}, \exists 1 \leq k \leq n\}$ is not contained in the union $\bigcup_{J \geq J_0; v \in \mathbb{Z}^n} Q_{J,v}$. \square

Recall the dilation operator $(\vartheta^r f)(x) := f(x', rx_n)$ for $r \in \mathbb{R} \setminus \{0\}$ from (8).

Proposition 22. *For any $0 < p, q \leq \infty$ and $s \in \mathbb{R}$ there is a $M = M_{p,q}^{n,s} > 0$ and $C = C_{n,p,q,s,M} > 0$ such that*

$$\|\vartheta^r f\|_{\mathcal{A}_{pq}^s(\mathbb{R}^n)} \leq C(|r|^{-M} + |r|^M) \|f\|_{\mathcal{A}_{pq}^s(\mathbb{R}^n)}, \quad \text{for } r \neq 0, \quad \mathcal{A} \in \{\mathcal{B}, \mathcal{F}\}, \quad f \in \mathcal{A}_{pq}^s(\mathbb{R}^n).$$

Proof. We only need to prove the Triebel-Lizorkin case, i.e. $\mathcal{A} = \mathcal{F}$. Indeed suppose we get the case $\mathcal{A} = \mathcal{F}$, then by real interpolation (18), for each $0 < p, q \leq \infty$ and $s \in \mathbb{R}$ there is a $C_{p,q,s} > 0$ such that (see [Tri95, Definition 1.2.2/2 and Theorem 1.3.3], in fact $C_{p,q,s} = C_q$)

$$(22) \quad \|T\|_{\mathcal{F}_{pq}^s(\mathbb{R}^n) \rightarrow \mathcal{F}_{pq}^s(\mathbb{R}^n)} \leq C_{p,q,s} (\|T\|_{\mathcal{F}_{pp}^{s-1}(\mathbb{R}^n) \rightarrow \mathcal{F}_{pp}^{s-1}(\mathbb{R}^n)} + \|T\|_{\mathcal{F}_{pp}^{s+1}(\mathbb{R}^n) \rightarrow \mathcal{F}_{pp}^{s+1}(\mathbb{R}^n)}),$$

whenever $T : \mathcal{F}_{pp}^{s-1}(\mathbb{R}^n) \rightarrow \mathcal{F}_{pp}^{s-1}(\mathbb{R}^n)$ is a bounded linear map such that $T : \mathcal{F}_{pp}^{s+1}(\mathbb{R}^n) \rightarrow \mathcal{F}_{pp}^{s+1}(\mathbb{R}^n)$ is also bounded. Taking $T = \vartheta^r$ and replacing $M_{p,q}^{n,s}$ (obtained from the \mathcal{F} -cases) with $\max(M_{p,q}^{n,s}, M_{p,p}^{n,s-1}, M_{p,p}^{n,s+1})$ we complete the proof.

Step 1: The case $n = 1$, $0 < p < \infty$, $0 < q \leq \infty$ and $s > \frac{1}{p} - 1$. We use [ET96, Section 2.3.1].

Recall $\lambda = (\lambda_j)_{j=0}^\infty$ from Definition 8. When $s > \max(0, 1 - \frac{1}{p})$, by [Tri92, Theorem 2.3.3], we have $\|f\|_{\mathcal{F}_{pq}^s(\mathbb{R})} \approx_{p,q,s} \|f\|_{L^p(\mathbb{R}^n)} + \|x \mapsto (\int_0^\infty t |(\lambda_1(\frac{\cdot}{t}) * f)(x)| \frac{dt}{t^{s+1}})^{1/q}\|_{L^p(\mathbb{R})}$. The same dilation argument from [ET96, (2.3.1/4) and (2.3.1/5)] shows that

$$(23) \quad \|\vartheta^r f\|_{\mathcal{F}_{pq}^s(\mathbb{R})} \approx |r|^{-\frac{1}{p}} \|f\|_{L^p} + |r|^{s-\frac{1}{p}} \|x \mapsto (\int_0^\infty |(\lambda_1(\frac{\cdot}{t}) * f)(x)| \frac{dt}{t^{s+1}})^{1/q}\|_{L^p} \lesssim_{p,q,s} (|r|^{-\frac{1}{p}} + |r|^{s-\frac{1}{p}}) \|f\|_{\mathcal{F}_{pq}^s(\mathbb{R})}.$$

In particular we can take $M_{p,q}^{1,s} := \max(\frac{1}{p}, s - \frac{1}{p}, \frac{1}{p} - s)$ in this case.

Step 2: The case $n \geq 2$, $0 < p < \infty$, $0 < q \leq \infty$ and $s > n \max(0, \frac{1}{p} - 1, \frac{1}{q} - 1)$. We use Proposition 21 (iv):

$$\begin{aligned} \|\vartheta^r f\|_{\mathcal{F}_{pq}^s(\mathbb{R}^n)} &\approx \left(\int_{\mathbb{R}^{n-1}} \|\vartheta^r(f(x', \cdot))\|_{\mathcal{F}_{pq}^s(\mathbb{R})}^p dx' \right)^{1/p} + \left(\int_{\mathbb{R}} \|f(\cdot, rx_n)\|_{\mathcal{F}_{pq}^s(\mathbb{R}^{n-1})}^p dx_n \right)^{1/p} \\ &\lesssim (|r|^{-\frac{1}{p}} + |r|^{s-\frac{1}{p}}) \left(\int_{\mathbb{R}^{n-1}} \|f(x', \cdot)\|_{\mathcal{F}_{pq}^s(\mathbb{R})}^p dx' \right)^{1/p} + |r|^{-\frac{1}{p}} \left(\int_{\mathbb{R}} \|f(\cdot, y_n)\|_{\mathcal{F}_{pq}^s(\mathbb{R}^{n-1})}^p dy_n \right)^{1/p} \\ &\lesssim (|r|^{-M_{p,q}^{1,s}} + |r|^{M_{p,q}^{1,s}}) \|f\|_{\mathcal{F}_{pq}^s(\mathbb{R}^n)}. \end{aligned}$$

In particular we can take $M_{p,q}^{n,s} := M_{p,q}^{1,s}$ in this case.

Step 3: The case $0 < p < \infty$, $0 < q \leq \infty$ and $s \in \mathbb{R}$. We use Proposition 21 (i).

Let $m \geq 0$ be such that $s + m > n \max(0, \frac{1}{p} - 1, \frac{1}{q} - 1)$. Note that $\vartheta^{\frac{1}{r}} \circ \partial^\alpha = r^\alpha \partial^\alpha \circ \vartheta^{\frac{1}{r}}$, thus

$$\begin{aligned} \|\vartheta^r f\|_{\mathcal{F}_{pq}^s(\mathbb{R}^n)} &\approx \inf \left\{ \sum_{|\alpha| \leq m} \|g_\alpha\|_{\mathcal{F}_{pq}^{s+m}(\mathbb{R}^n)} : \vartheta^r f = \sum_{|\alpha| \leq m} \partial^\alpha g_\alpha \right\} \\ &= \inf \left\{ \sum_{|\alpha| \leq m} \|g_\alpha\|_{\mathcal{F}_{pq}^{s+m}(\mathbb{R}^n)} : f = \sum_{|\alpha| \leq m} r^\alpha \partial^\alpha (\vartheta^{\frac{1}{r}} g_\alpha) \right\} = \inf \left\{ \sum_{|\alpha| \leq m} \frac{\|\vartheta^r h_\alpha\|_{\mathcal{F}_{pq}^{s+m}(\mathbb{R}^n)}}{|r|^{\alpha_n}} : f = \sum_{|\alpha| \leq m} \partial^\alpha h_\alpha \right\} \\ &\lesssim (|r|^{-m} + 1) (|r|^{-M_{p,q}^{n,s+m}} + |r|^{M_{p,q}^{n,s+m}}) \|f\|_{\mathcal{F}_{pq}^s(\mathbb{R}^n)}. \end{aligned}$$

In particular we can take $M_{p,q}^{n,s} := M_{p,q}^{n,s+m} + m$ in this case.

Step 4: The case $p = \infty$, $0 < q \leq \infty$ and $s \in \mathbb{R}$. We use Proposition 21 (v).

For $x \in \mathbb{R}^n$ and $R > 0$, we see that $f \in \mathcal{F}_{pq}^s(B((x', \frac{x_n}{r}), \max(R, \frac{R}{|r|})))$ implies $\vartheta^r f \in \mathcal{F}_{pq}^s(B(x, R))$. Therefore

$$\begin{aligned} \|\vartheta^r f\|_{\mathcal{F}_{\infty q}^s(\mathbb{R}^n)} &\approx \sup_{x \in \mathbb{R}^n, R > 0} R^{-\frac{n}{q}} \inf \{ \|\tilde{g}\|_{\mathcal{F}_{qq}^s(\mathbb{R}^n)} : \tilde{g}|_{B(x,R)} = (\vartheta^r f)|_{B(x,R)} \} \\ &\leq \sup_{x \in \mathbb{R}^n, R > 0} R^{-\frac{n}{q}} \inf \{ \|\vartheta^{\frac{1}{r}} \tilde{f}\|_{\mathcal{F}_{qq}^s(\mathbb{R}^n)} : \tilde{f}|_{B((x', \frac{x_n}{r}), \max(R, \frac{R}{|r|}))} = f|_{B((x', \frac{x_n}{r}), \max(R, \frac{R}{|r|}))} \} \\ &\lesssim (|r|^{-M_{q,q}^{n,s}} + |r|^{M_{q,q}^{n,s}}) \sup_{x \in \mathbb{R}^n, R > 0} R^{-\frac{n}{q}} \|f\|_{\mathcal{F}_{qq}^s(B((x', \frac{x_n}{r}), \max(R, \frac{R}{|r|})))} \\ &\lesssim (|r|^{-M_{q,q}^{n,s}} + |r|^{M_{q,q}^{n,s}}) (1 + |r|^{-\frac{n}{q}}) \sup_{x \in \mathbb{R}^n, R > 0} R^{-\frac{n}{q}} \|f\|_{\mathcal{F}_{qq}^s(B(x,R))} \lesssim (|r|^{-M_{q,q}^{n,s} - \frac{n}{q}} + |r|^{M_{q,q}^{n,s}}) \|f\|_{\mathcal{F}_{\infty q}^s(\mathbb{R}^n)}. \end{aligned}$$

Taking $M_{\infty,q}^{n,s} := M_{q,q}^{n,s} + n/q$, we complete the proof. \square

We now start to prove Theorem 1 (iii). In order to handle Theorem 1 (iv) later, we consider a more quantitative version of Theorem 1 (iii) that have controls of the operator norms in terms of $(a_j)_j$ and $(b_j)_j$.

Theorem 23 (Quantitative common extensions). *For every $0 < p, q \leq \infty$, $s \in \mathbb{R}$, there is a constant $\widehat{M} = \widehat{M}_{p,q}^{n,s}$, such that the following holds:*

Let $m_1, m_2 \in \mathbb{Z}_{\geq 0}$ be two non-negative integers such that $n \max(0, \frac{1}{p} - 1, \frac{1}{q} - 1) - m_1 < s < \frac{1}{p} + m_2$. Let $a = (a_j)_{j=-\infty}^{\infty} \subset \mathbb{R}$ and $b = (b_j)_{j=-\infty}^{\infty} \subset \mathbb{R}_+$ be two sequences satisfying

$$(24) \quad \sum_{j=-\infty}^{\infty} a_j (-b_j)^k = 1 \text{ for } -m_1 \leq k \leq m_2; \quad \sum_{j \in \mathbb{Z}} 2^{\delta|j|} |a_j| (b_j^{-\max(m_1, \widehat{M})} + b_j^{\max(m_2, \widehat{M})}) < \infty \text{ for some } \delta > 0.$$

Then $E = E^{a,b}$ given by (2) has the boundedness $E : \mathcal{A}_{pq}^s(\mathbb{R}_+^n) \rightarrow \mathcal{A}_{pq}^s(\mathbb{R}^n)$ for $\mathcal{A} \in \{\mathcal{B}, \mathcal{F}\}$. Moreover for the operator norms of E we have

$$(25) \quad \|Ef\|_{\mathcal{A}_{pq}^s(\mathbb{R}^n)} \leq C_{n,p,q,s,\delta,\widehat{M}} \sum_{j \in \mathbb{Z}} 2^{\delta|j|} |a_j| (b_j^{-\widehat{M}_{p,q}^{n,s}} + b_j^{\widehat{M}_{p,q}^{n,s}}) \|f\|_{\mathcal{A}_{pq}^s(\mathbb{R}_+^n)}.$$

Here $C_{n,p,q,s,\delta,\widehat{M}}$ does not depend on the choices of $(a_j)_j$ and $(b_j)_j$.

Remark 24. By Proposition 14, the sequences $(a_j)_j$ and $(b_j)_j$ can be chosen independently of p, q, n, s, m_1, m_2 . Note that δ in (24) is allowed to depend on $m_1, m_2, \widehat{M}_{pq}^{n,s}$ (cf. the δ_k in (3) and Remark 2).

Therefore, by letting $m_1, m_2 \rightarrow +\infty$ we obtain the all boundedness properties of E in Theorem 1 (iii).

Additionally, Theorem 23 also gives the following:

- If one take $m_1 = 0$ and let $m_2 \rightarrow +\infty$, we see that Seeley's extension operator (see Remark 12) has boundedness $E : \mathcal{A}_{pq}^s(\mathbb{R}_+^n) \rightarrow \mathcal{A}_{pq}^s(\mathbb{R}^n)$ for $\mathcal{A} \in \{\mathcal{B}, \mathcal{F}\}$, $p, q \in (0, \infty]$ and $s > n \max(0, \frac{1}{p} - 1, \frac{1}{q} - 1)$. It is possible that the range for s is not optimal.
- If one take $(a_j)_j$ to be finitely nonzero (depending on the upper bound of m_1, m_2), then the qualitative result $E : \mathcal{A}_{pq}^s(\mathbb{R}_+^n) \rightarrow \mathcal{A}_{pq}^s(\mathbb{R}^n)$ is well-known for *common extension operators*. See [Tri20, Remark 2.72].

Proof of Theorem 23 (hence Theorem 1 (iii)). Similar to the proof of Proposition 22, using (22) we only need to prove the case $\mathcal{A} = \mathcal{F}$.

To make the notation clear, we use $E_n^{a,b}$ for the extension operator of \mathbb{R}_+^n given in (2) associated with sequences $a = (a_j)_{j=-\infty}^{\infty}$ and $b = (b_j)_{j=-\infty}^{\infty}$. Recall the zero extension operator $S_n f(x) := \mathbf{1}_{\mathbb{R}_+^n}(x) f(x)$ from (9).

Case A: We consider $p < \infty$ and s suitably large. We sub-divide the discussion into three parts.

Step A1: The case $n = 1$, $0 < p < \infty$, $0 < q \leq \infty$ and $\frac{1}{p} - 1 < s < \frac{1}{p}$, with arbitrary $m_1, m_2 \geq 0$.

Indeed by [Tri20, Theorem 2.48] we have boundedness $S_1 : \mathcal{F}_{pq}^s(\mathbb{R}_+) \rightarrow \mathcal{F}_{pq}^s(\mathbb{R})$ at this range. Clearly $E_1^{a,b} = S_1 + \sum_{j=-\infty}^{\infty} a_j \cdot \vartheta^{-bj} \circ S_1$. Therefore by Lemma 17 and Proposition 22,

$$\begin{aligned} \|E_1^{a,b} f\|_{\mathcal{F}_{pq}^s(\mathbb{R})} &\lesssim_{\min(p,q,1),\delta} \|S_1 f\|_{\mathcal{F}_{pq}^s(\mathbb{R})} + \sum_{j \in \mathbb{Z}} 2^{\delta|j|} |a_j| \|\vartheta^{-bj} S_1 f\|_{\mathcal{F}_{pq}^s(\mathbb{R})} \\ &\lesssim_{p,q,s} \sum_{j \in \mathbb{Z}} 2^{\delta|j|} |a_j| (b_j^{-M_{pq}^{1s}} + b_j^{M_{pq}^{1s}}) \|S_1 f\|_{\mathcal{F}_{pq}^s(\mathbb{R})} \lesssim_{p,q,s} \sum_{j \in \mathbb{Z}} 2^{\delta|j|} |a_j| (b_j^{-M_{pq}^{1s}} + b_j^{M_{pq}^{1s}}) \|f\|_{\mathcal{F}_{pq}^s(\mathbb{R}_+)}. \end{aligned}$$

We conclude the case by taking $\widehat{M}_{pq}^{1s} := M_{pq}^{1s}$, the constant in Proposition 22.

Step A2: The case $n = 1$, $0 < p < \infty$, $0 < q \leq \infty$ and $s > \frac{1}{p} - 1$, with $m_1 \geq 0$ and $m_2 > s - \frac{1}{p}$.

By [Tri10, Theorem 2.3.8 and Remark 3.3.5/2] $\|f\|_{\mathcal{F}_{pq}^s(U)} \approx_{p,q,s,m_0} \sum_{k=0}^{m_0} \|\partial^k f\|_{\mathcal{F}_{pq}^{s-m_0}(U)}$ holds for every $m_0 \geq 1$ and $U \in \{\mathbb{R}^n, \mathbb{R}_+^n\}$. Recall from Remark 16 that $\partial^k \circ E_1^{a,b} = E_1^{a(-b)^k, b} \circ \partial^k$ holds for $0 \leq k \leq m_2$.

When $s - \frac{1}{p} \notin \mathbb{Z}$, we let $m_0 := \lceil s - \frac{1}{p} \rceil$. Therefore by Step A1 (since $m_2 \geq m_0$),

$$\begin{aligned} \|E_1^{a,b} f\|_{\mathcal{F}_{pq}^s(\mathbb{R})} &\approx_{p,q,s} \sum_{k=0}^{m_0} \|E_1^{a(-b)^k, b} \partial^k f\|_{\mathcal{F}_{pq}^{s-m_0}} \lesssim_{p,q,s,\delta} \sum_{k=0}^{m_0} \sum_{j \in \mathbb{Z}} 2^{\delta|j|} |a_j| (b_j^{k-\widehat{M}_{pq}^{1,s-m_0}} + b_j^{k+\widehat{M}_{pq}^{1,s-m_0}}) \|\partial^k f\|_{\mathcal{F}_{pq}^{s-m_0}} \\ &\lesssim_{m_0} \sum_{j \in \mathbb{Z}} 2^{\delta|j|} |a_j| (b_j^{-\widehat{M}_{pq}^{1,s-m_0}} + b_j^{m_0+\widehat{M}_{pq}^{1,s-m_0}}) \|f\|_{\mathcal{F}_{pq}^s(\mathbb{R}_+)}. \end{aligned}$$

In this case we can take $\widehat{M}_{pq}^{1s} := \widehat{M}_{pq}^{1,s-m_0} + m_0 = \widehat{M}_{pq}^{1,s-\lceil s-\frac{1}{p} \rceil} + \lceil s - \frac{1}{p} \rceil$.

The case $s - \frac{1}{p} \in \mathbb{Z}_+$ follows from the complex interpolation (17), where we can take $\widehat{M}_{pq}^{1s} := \max(\widehat{M}_{pq}^{1, s - \frac{1}{2}}, \widehat{M}_{pq}^{1, s + \frac{1}{2}})$.

Step A3: The case $n \geq 1$, $0 < p < \infty$, $0 < q \leq \infty$ and $s > n \max(0, \frac{1}{p} - 1, \frac{1}{q} - 1)$, with $m_1 \geq 0$ and $m_2 > s - \frac{1}{p}$.

Indeed we have $(E_n^{a,b} f)(x) = E_1^{a,b}(f(x', \cdot))(x_n)$. Therefore, for any extension $\tilde{f} \in \mathcal{F}_{pq}^s(\mathbb{R}^n)$ of $f \in \mathcal{F}_{pq}^s(\mathbb{R}_+^n)$,

$$\begin{aligned}
& \|E_n^{a,b} f\|_{\mathcal{F}_{pq}^s(\mathbb{R}^n)} \\
& \approx_{p,q,s} \left(\int_{\mathbb{R}^{n-1}} \|E_1^{a,b}(f(x', \cdot))\|_{\mathcal{F}_{pq}^s(\mathbb{R})}^p dx' \right)^{1/p} + \left(\int_{\mathbb{R}} \|(E_n^{a,b} f)(\cdot, x_n)\|_{\mathcal{F}_{pq}^s(\mathbb{R}^{n-1})}^p dx_n \right)^{1/p} \quad (\text{by Proposition 21 (iv)}) \\
& \leq \|E_1^{a,b}\|_{\mathcal{F}_{pq}^s} \left(\int_{\mathbb{R}^{n-1}} \|f(x', \cdot)\|_{\mathcal{F}_{pq}^s(\mathbb{R}_+)}^p dx' \right)^{1/p} + \|E_1^{a,b} f\|_{L^p(\mathbb{R}_{x_n}; \mathcal{F}_{pq}^s(\mathbb{R}_+^{n-1}))} \\
& \lesssim_{p,q,s} \|E_1^{a,b}\|_{\mathcal{F}_{pq}^s} \|f\|_{L^p(\mathbb{R}^{n-1}; \mathcal{F}_{pq}^s(\mathbb{R}_+))} + \|E_1^{a,b}\|_{L^p(\mathcal{F}_{pq}^s)} \|f\|_{L^p(\mathbb{R}_+; \mathcal{F}_{pq}^s(\mathbb{R}^{n-1}))} \\
& \leq_{p,q,s,\delta} \sum_{j \in \mathbb{Z}} 2^{\delta|j|} |a_j| (b_j^{-\widehat{M}_{pq}^{1s}} + b_j^{\widehat{M}_{pq}^{1s}}) \|\tilde{f}\|_{L^p(\mathbb{R}^{n-1}; \mathcal{F}_{pq}^s)} + \sum_{j \in \mathbb{Z}} 2^{\delta|j|} |a_j| b_j^{-\frac{1}{p}} \|\tilde{f}\|_{L^p(\mathbb{R}; \mathcal{F}_{pq}^s)} \quad (\text{by Step A2 and Remark 19}) \\
& \lesssim_{p,q,s} \sum_{j \in \mathbb{Z}} 2^{\delta|j|} |a_j| (b_j^{-\widehat{M}_{pq}^{1s} - \frac{1}{p}} + b_j^{\widehat{M}_{pq}^{1s}}) \|\tilde{f}\|_{\mathcal{F}_{pq}^s(\mathbb{R}^n)} \quad (\text{by Proposition 21 (iv)}).
\end{aligned}$$

Taking the infimum of the extensions \tilde{f} over f , we get the claim with $\widehat{M}_{pq}^{ns} := \widehat{M}_{pq}^{1s} + 1/p$.

Case B: We consider $p = \infty$, i.e. the $\mathcal{F}_{\infty q}^s$ -cases, for $s > -1$.

Step B1: The case $n \geq 1$, $p = \infty$, $0 < q \leq \infty$ and $-1 < s < 0$, with arbitrary $m_1, m_2 \geq 0$.

The proof is identical to **Step A1**, where we use $E_n^{a,b} = S_n + \sum_{j=-\infty}^{\infty} a_j \cdot \vartheta^{-b_j} \circ S_n$ and $S_n : \mathcal{F}_{pq}^s(\mathbb{R}_+^n) \rightarrow \mathcal{F}_{pq}^s(\mathbb{R}^n)$ (see [Tri20, Theorem 2.48]). We get the result with $\widehat{M}_{\infty q}^{ns} := M_{\infty q}^{ns}$, the constant in Proposition 22.

Step B2: The case $n \geq 1$, $p = \infty$, $0 < q \leq \infty$ and $s > -1$, with $m_1 \geq 0$ and $m_2 > s$.

It remains to prove the case $s \geq 0$. The proof is similar to **Step A2**, except we use (see [Tri20, Theorem 1.24]) $\|f\|_{\mathcal{F}_{\infty q}^s(\mathbb{R}^n)} \approx_{p,q,s,m_0} \sum_{|\gamma| \leq m_0} \|\partial^\gamma f\|_{\mathcal{F}_{pq}^{s-m_0}(\mathbb{R}^n)}$. When $s \notin \mathbb{Z}_+$, since $m_2 \geq \lceil s \rceil$ we get from **Step B1** that

$$\|E_n^{a,b} f\|_{\mathcal{F}_{pq}^s(\mathbb{R}^n)} \lesssim_{q,s,\delta} \sum_{j \in \mathbb{Z}} 2^{\delta|j|} |a_j| (b_j^{-\widehat{M}_{\infty q}^{n,s-\lceil s \rceil}} + b_j^{\lceil s \rceil + \widehat{M}_{\infty q}^{n,s-\lceil s \rceil}}) \|\tilde{f}\|_{\mathcal{F}_{\infty q}^s(\mathbb{R}^n)}, \quad \text{whenever } \tilde{f}|_{\mathbb{R}_+^n} = f.$$

Taking infimum of the extensions \tilde{f} over f we get the claim with $\widehat{M}_{\infty q}^{ns} := \widehat{M}_{\infty q}^{n,s-\lceil s \rceil} + \lceil s \rceil$.

The case $s \in \mathbb{Z}_+$ follows from the complex interpolation (17), where we can take $\widehat{M}_{\infty q}^{ns} := \max(\widehat{M}_{\infty q}^{n,s-\frac{1}{2}}, \widehat{M}_{\infty q}^{n,s+\frac{1}{2}})$.

Final step: The general case: $n \geq 1$, $0 < p, q \leq \infty$ and $s \in \mathbb{R}$, with $m_1 > n \max(0, \frac{1}{p} - 1, \frac{1}{q} - 1) - s$ and $m_2 > s - \frac{1}{p}$. Here we can take $m_1 := \lfloor n \max(0, \frac{1}{p} - 1, \frac{1}{q} - 1) - s \rfloor + 1$.

By Remark 16 again we have $E_n^{a,b} \circ \partial^\gamma = \partial^\gamma \circ E_n^{a(-b)^{-\gamma n}, b}$ for all $|\gamma| \leq m_1$, where $a(-b)^{-\gamma n} = (a_j(-b_j)^{-\gamma n})_{j=-\infty}^{\infty}$. Therefore when $f \in \mathcal{F}_{pq}^s(\mathbb{R}_+^n)$ and $g_\gamma \in \mathcal{F}_{pq}^{s+m_1}(\mathbb{R}_+^n)$ satisfy $f = \sum_{|\gamma| \leq m_1} \partial^\gamma g_\gamma$, by Steps A3 and B2 we have

$$\begin{aligned}
(26) \quad & \|E_n^{a,b} f\|_{\mathcal{F}_{pq}^s(\mathbb{R}^n)} = \left\| \sum_{|\gamma| \leq m_1} \partial^\gamma \circ E_n^{a(-b)^{-\gamma n}, b} g_\gamma \right\|_{\mathcal{F}_{pq}^s(\mathbb{R}^n)} \\
& \lesssim_{\min(p,q,1), m_1} \sum_{|\gamma| \leq m_1} \|\partial^\gamma \circ E_n^{a(-b)^{-\gamma n}, b} g_\gamma\|_{\mathcal{F}_{pq}^s(\mathbb{R}^n)} \\
& \lesssim_{p,q,s,m_1} \sum_{|\gamma| \leq m_1} \|E_n^{a(-b)^{-\gamma n}, b} g_\gamma\|_{\mathcal{F}_{pq}^{s+m_1}(\mathbb{R}^n)} \quad (\text{see [Tri10, (2.3.8/3)]}) \\
& \lesssim_{p,q,s,m_1,\delta} \sum_{|\gamma| \leq m_1} \sum_{j \in \mathbb{Z}} 2^{\delta|j|} |a_j| (b_j^{-\widehat{M}_{pq}^{n,s+m_1-\gamma n}} + b_j^{\widehat{M}_{pq}^{n,s+m_1-\gamma n}}) \|g_\gamma\|_{\mathcal{F}_{pq}^{s+m_1}(\mathbb{R}^n)} \quad (\text{by Steps A3 and B2}).
\end{aligned}$$

Taking the infimum over the decompositions $f = \sum_{|\alpha| \leq m} \partial^\alpha g_\alpha$, by Proposition 21 (i) we obtain the claim with $\widehat{M}_{pq}^{ns} := \widehat{M}_{pq}^{n,s+m_1} + m_1 = \widehat{M}_{pq}^{n,s+\lfloor n \max(0, \frac{1}{p} - 1, \frac{1}{q} - 1) - s \rfloor + 1} + \lfloor n \max(0, \frac{1}{p} - 1, \frac{1}{q} - 1) - s \rfloor + 1$. This completes the proof of $\mathcal{A} = \mathcal{F}$. The case $\mathcal{A} = \mathcal{B}$ is done by real interpolations, as discussed in the beginning of the proof. \square

We now prove Theorem 1 (iv), the boundedness on $\mathcal{B}_{pq}^{s\tau}$, $\mathcal{F}_{pq}^{s\tau}$ and $\mathcal{N}_{pq}^{s\tau}$, using Theorem 23.

In the following, for $m \in \mathbb{Z}_+$ and $r > 0$, we define a sequence $A^{mr} = (A_j^{mr})_{j=0}^{2m}$ to be the unique sequence such that $\sum_{j=0}^{2m} A_j^{mr} (-2^{-j}r)^k = 1$ for integers $-m \leq k \leq m$. Recall from Remark 13 we have

$$(27) \quad A_j^{mr} = (-r2^{-j})^m \prod_{0 \leq k \leq 2m; k \neq j} \frac{r2^{-k} + 1}{r2^{-k} - r2^{-j}} = (-1)^m \frac{2^{-jm}}{r^m} \prod_{0 \leq k \leq 2m; k \neq j} \frac{r2^{-k} + 1}{2^{-k} - 2^{-j}}, \quad 0 \leq j \leq 2m.$$

We see that $|A_j^{mr}| \lesssim_m r^m + r^{-m}$ for $0 \leq j \leq 2m$, where the implicit constant does not depend on $r > 0$.

We define

$$(28) \quad \mathbf{E}^{m,r} f(x', x_n) = \mathbf{E}_n^{m,r} f(x', x_n) := \begin{cases} \sum_{j=0}^{2m} A_j^{mr} f(x', -r2^{-j}x_n) & x_n < 0 \\ f(x) & x_n > 0 \end{cases}.$$

By Theorem 23, for every $0 < p, q \leq \infty$, $s \in \mathbb{R}$ and $m > \max(s - \frac{1}{p}, -s, n(\frac{1}{p} - 1) - s, n(\frac{1}{q} - 1) - s)$, there is a $C = C_{n,p,q,s,m} > 0$, such that $\mathbf{E}^{m,r} : \mathcal{A}_{pq}^s(\mathbb{R}_+^n) \rightarrow \mathcal{A}_{pq}^s(\mathbb{R}^n)$ is bounded for $\mathcal{A} \in \{\mathcal{B}, \mathcal{F}\}$ and for all $r > 0$, with

$$(29) \quad \|\mathbf{E}^{m,r}\|_{\mathcal{A}_{pq}^s(\mathbb{R}_+^n) \rightarrow \mathcal{A}_{pq}^s(\mathbb{R}^n)} \leq C_{n,p,q,s,m} (r^{-m} \widehat{M}_{pq}^{ns} + r^m \widehat{M}_{pq}^{ns}), \quad \forall r > 0.$$

Here $\widehat{M}_{pq}^{ns} > 0$ is the corresponding constant in Theorem 23.

Proof of Theorem 1 (iv). We only need to prove the case $\mathcal{A} \in \{\mathcal{B}, \mathcal{F}\}$. Indeed, when $\tau < \frac{1}{p}$ (hence $p < \infty$), the boundedness of $\mathcal{N}_{pq}^{s\tau}$ can be deduced from real interpolations (19). When $\tau = \frac{1}{p}$ we have coincidence $\mathcal{N}_{pq}^{s\frac{1}{p}} = \mathcal{B}_{\infty q}^s$ (see (7)), which is treated in Theorem 1 (iii).

We begin with the case $s < 0$.

Recall from Proposition 21 (v), $\|\tilde{f}\|_{\mathcal{A}_{pq}^{s\tau}(\mathbb{R}^n)} \approx_{n,p,q,s,\tau} \sup_{x \in \mathbb{R}^n; R > 0} R^{-n\tau} \|\tilde{f}\|_{\mathcal{A}_{pq}^s(B(x,R))}$ for $\mathcal{A} \in \{\mathcal{B}, \mathcal{F}\}$ ($p < \infty$ if $\mathcal{A} = \mathcal{F}$) under the assumption $s < 0$.

Clearly if $x_n \geq R$, then $\|Ef\|_{\mathcal{A}_{pq}^s(B(x,R))} = \|f\|_{\mathcal{A}_{pq}^s(B(x,R))}$.

If $x_n \leq -R$, then $(E^{a,b}f)|_{B(x,R)} = \sum_{j=-\infty}^{\infty} a_j \cdot \vartheta^{-bj} f$. Therefore, for every extension $\tilde{f} \in \mathcal{A}_{pq}^{s\tau}(\mathbb{R}^n)$ of f ,

$$(30) \quad \begin{aligned} R^{-n\tau} \|E^{a,b}f\|_{\mathcal{A}_{pq}^s(B(x,R))} &\lesssim_{p,q,\delta} R^{-n\tau} \sum_{j=-\infty}^{\infty} 2^{\delta|j|} |a_j| \|\vartheta^{-bj} f\|_{\mathcal{A}_{pq}^s(B(x,R))} && \text{(by Lemma 17)} \\ &\leq R^{-n\tau} \sum_{j=-\infty}^{\infty} 2^{\delta|j|} |a_j| (b_j^{-M_{pq}^{ns}} + b_j^{M_{pq}^{ns}}) \|f\|_{\mathcal{A}_{pq}^s(B((x', -\frac{x_n}{b_j}), \max(R, \frac{R}{b_j})) \cap \mathbb{R}_+^n)} && \text{(by Proposition 22)} \\ &\leq \sum_{j=-\infty}^{\infty} 2^{\delta|j|} |a_j| (b_j^{-M_{pq}^{ns}} + b_j^{M_{pq}^{ns}}) \max(R, \frac{R}{b_j})^{-n\tau} \sup_{y \in \mathbb{R}^n} \|\tilde{f}\|_{\mathcal{A}_{pq}^s(B(y, \max(R, R/b_j)))} \\ &\lesssim_{n,p,q,s,a,b} \sup_{y \in \mathbb{R}^n; \rho > 0} \rho^{-n\tau} \|\tilde{f}\|_{\mathcal{A}_{pq}^s(B(y,\rho))} \approx_{n,p,q,s,\tau} \|\tilde{f}\|_{\mathcal{A}_{pq}^{s\tau}(\mathbb{R}^n)} && \text{(by Proposition 21 (v)).} \end{aligned}$$

Here M_{pq}^{ns} is the constant in Proposition 22. Taking the infimum of the extension \tilde{f} over f we see that

$$R^{-n\tau} \|Ef\|_{\mathcal{A}_{pq}^s(B(x,R))} \lesssim_{n,p,q,s,\tau,a,b} \|f\|_{\mathcal{A}_{pq}^{s\tau}(\mathbb{R}_+^n)} \text{ uniformly for } R > 0 \text{ and } x_n \leq -R, \text{ and hence for all } |x_n| \geq R.$$

The difficult part is to prove the estimate for $|x_n| \leq R$. In this case $B(x, R) \subset B((x', 0), 2R)$, thus $R^{-n\tau} \|Ef\|_{\mathcal{A}_{pq}^s(B(x,R))} \leq 2^{n\tau} (2R)^{-n\tau} \|Ef\|_{\mathcal{A}_{pq}^s(B((x', 0), 2R))}$. Since E is translation invariant in x' , we can assume $x' = 0$ without loss of generality.

To summarize the discussion above, to prove the boundedness $E : \mathcal{A}_{pq}^{s\tau}(\mathbb{R}_+^n) \rightarrow \mathcal{A}_{pq}^s(\mathbb{R}^n)$ for $0 < p, q \leq \infty$, $s < 0$ and $0 \leq \tau \leq \frac{1}{p}$ ($p < \infty$ if $\mathcal{A} = \mathcal{F}$), it remains to show the existence of $C = C_{n,p,q,s,\tau,a,b} > 0$ such that

$$(31) \quad R^{-n\tau} \|Ef\|_{\mathcal{A}_{pq}^s(B(0,R))} \leq C \sup_{\rho > 0} \rho^{-n\tau} \|\tilde{f}\|_{\mathcal{A}_{pq}^s(B(0,\rho))}, \quad \forall R > 0, \quad \forall \tilde{f} \in \mathcal{A}_{pq}^s(\mathbb{R}^n) \text{ such that } \tilde{f}|_{\mathbb{R}_+^n} = f.$$

In the following we let

$$B_+(0, R) := B(0, R) \cap \mathbb{R}_+^n; \quad J_+ := \{j \in \mathbb{Z} : b_j > 1\}, \quad J_- := \{j \in \mathbb{Z} : b_j \leq 1\}, \quad \text{thus } \mathbb{Z} = J_+ \amalg J_-.$$

We fix a $R > 0$ temporally.

For each $j \in J_+$, take an extension $\tilde{f}_j = \tilde{f}_j^R \in \mathcal{A}_{pq}^s(\mathbb{R}^n)$ of $f|_{B_+(0, b_j R)}$ such that $\|\tilde{f}_j\|_{\mathcal{A}_{pq}^s(\mathbb{R}^n)} \leq 2\|f\|_{\mathcal{A}_{pq}^s(B_+(0, b_j R))}$. We let $\tilde{f}_- = \tilde{f}_-^R \in \mathcal{A}_{pq}^s(\mathbb{R}^n)$ be an extension of $f|_{B_+(0, R)}$ such that $\|\tilde{f}_-\|_{\mathcal{A}_{pq}^s(\mathbb{R}^n)} \leq 2\|f\|_{\mathcal{A}_{pq}^s(B_+(0, R))}$.

Clearly we have the following extension of $Ef|_{B(0,R)\cap\mathbb{R}_-^n}$:

$$\tilde{f}_{<0} = \tilde{f}_{<0}^R := \sum_{j \in J_+} a_j \cdot \vartheta^{-b_j} \tilde{f}_j + \sum_{j \in J_-} a_j \cdot \vartheta^{-b_j} \tilde{f}_-.$$

However $\tilde{f}_{<0}$ is not the extension of $f|_{B_+(0,R)} = Ef|_{B_+(0,R)}$. To modify it on the domain $B_+(0,R)$ we consider

$$\tilde{f}_{>0} = \tilde{f}_{>0}^R := \tilde{f}_- + \sum_{j \in J_+} \vartheta^{-1} \circ \mathbf{E}^{m, \frac{1}{b_j}} [\vartheta^{b_j} (\tilde{f}_j - \tilde{f}_-)] = \tilde{f}_- + \sum_{j \in J_+} \sum_{k=0}^{2m} A_k^{m, \frac{1}{b_j}} \cdot (\vartheta^{2^{-k}} \tilde{f}_j - \vartheta^{2^{-k}} \tilde{f}_-) \quad \text{in } \mathbb{R}_+^n,$$

where we have chosen $m > \max(s - \frac{1}{p}, -s, n(\frac{1}{p} - 1) - s, n(\frac{1}{q} - 1) - s)$, and $\mathbf{E}^{m, 1/b_j}$ is given by (28).

When $j \in J_+$ (i.e. $b_j > 1$) we have $(\tilde{f}_j - \tilde{f}_-)|_{B(0,R)} = 0$. Thus in this case $\mathbf{E}^{m, \frac{1}{b_j}} [\vartheta^{-b_j} (\tilde{f}_j - \tilde{f}_-)]|_{B(0,R)} \equiv 0$ (since $2^{-k}/b_j < 1$ for $j \in J_+$ and $k \geq 0$). We conclude that $\tilde{f}_{>0}|_{B_+(0,R)} = \tilde{f}_-|_{B_+(0,R)} = f|_{B_+(0,R)}$.

On the other hand we can rewrite $\tilde{f}_{<0}$ as

$$(32) \quad \tilde{f}_{<0} = \sum_{j=-\infty}^{\infty} a_j \cdot \vartheta^{-b_j} \tilde{f}_- + \sum_{j \in J_+} a_j \cdot \vartheta^{-b_j} (\tilde{f}_j - \tilde{f}_-) = E^{a,b} \tilde{f}_- + \sum_{j \in J_+} a_j \cdot \vartheta^{-1} \circ \vartheta^{b_j} (\tilde{f}_j - \tilde{f}_-) \quad \text{on } \mathbb{R}^n.$$

Here for global defined function \tilde{g} , we use the notation $E\tilde{g}$ for $E[(\tilde{g}|_{\mathbb{R}_+^n})]$.

Therefore, the function $\tilde{F} = \tilde{F}^R := (\tilde{f}_{>0}|_{\mathbb{R}_+^n}) \cup (\tilde{f}_{<0}|_{\mathbb{R}_-^n})$ is an extension of $Ef|_{B(0,R)}$ and has the expression

$$\tilde{F} = E^{a,b} [\tilde{f}_-] + \sum_{j \in J_+} a_j \cdot \vartheta^{-1} \circ \mathbf{E}^{m, \frac{1}{b_j}} [\vartheta^{b_j} (\tilde{f}_j - \tilde{f}_-)] \in \mathcal{S}'(\mathbb{R}^n).$$

We see that

$$\begin{aligned} \|\tilde{F}\|_{\mathcal{S}'_{pq}(\mathbb{R}^n)} &\lesssim_{p,q,\delta} \|E^{a,b} \tilde{f}_-\|_{\mathcal{S}'_{pq}(\mathbb{R}^n)} + \sum_{j \in J_+} 2^{\delta|j|} |a_j| \|\mathbf{E}^{m, \frac{1}{b_j}}\|_{\mathcal{S}'_{pq}(\mathbb{R}^n)} \|\vartheta^{b_j} (\tilde{f}_j - \tilde{f}_-)\|_{\mathcal{S}'_{pq}(\mathbb{R}^n)} && \text{(by Lemma 17)} \\ &\lesssim_{n,p,q,s,m} C'_{\delta,a,b,\widehat{M}} \|\tilde{f}_-\|_{\mathcal{S}'_{pq}(\mathbb{R}_+^n)} + \sum_{j \in J_+} 2^{\delta|j|} |a_j| b_j^{m \cdot \widehat{M}_{pq}^{ns}} \|\vartheta^{b_j} (\tilde{f}_j - \tilde{f}_-)\|_{\mathcal{S}'_{pq}(\mathbb{R}^n)} && \text{(by Theorem 23 and (29))} \\ &\lesssim_{n,p,q,s,a,b,\delta,\widehat{M}} \|\tilde{f}_-\|_{\mathcal{S}'_{pq}(\mathbb{R}^n)} + \sum_{j \in J_+} 2^{\delta|j|} |a_j| b_j^{m \cdot \widehat{M}_{pq}^{ns}} b_j^{M_{pq}^{ns}} \|\tilde{f}_j - \tilde{f}_-\|_{\mathcal{S}'_{pq}(\mathbb{R}^n)} && \text{(by Proposition 22)} \\ &\lesssim_{\min(1,p,q)} \|f\|_{\mathcal{S}'_{pq}(B_+(0,R))} + \sum_{j \in J_+} 2^{\delta|j|} |a_j| b_j^{m \cdot \widehat{M}_{pq}^{ns} + M_{pq}^{ns}} (\|f\|_{\mathcal{S}'_{pq}(B_+(0,R))} + \|f\|_{\mathcal{S}'_{pq}(B_+(0,b_j R))}). \end{aligned}$$

Here $C' = C'_{\delta,a,b,\widehat{M}} = \sum_{j \in \mathbb{Z}} 2^{\delta|j|} |a_j| (b_j^{\widehat{M}_{pq}^{ns}} + b_j^{-\widehat{M}_{pq}^{ns}})$. Notice that all implied constants in the above computation do not depend on the $R > 0$.

Therefore for general $R > 0$, applying (3) again with a small δ , we see that for every extension \tilde{f} of f ,

$$\begin{aligned} R^{-n\tau} \|Ef\|_{\mathcal{S}'_{pq}(B(0,R))} &\leq R^{-n\tau} \|\tilde{F}^R\|_{\mathcal{S}'_{pq}(\mathbb{R}^n)} \\ &\lesssim_{p,q,s,m,\delta} C'_{\delta,a,b,\widehat{M}} \cdot R^{-n\tau} \|f\|_{\mathcal{S}'_{pq}(B_+(0,R))} + \sum_{j \in J_+} 2^{\delta|j|} |a_j| b_j^{C''_{n,p,q,s,\tau,m}} (R^{-n\tau} \|f\|_{\mathcal{S}'_{pq}(B_+(0,R))} + (b_j R)^{-n\tau} \|f\|_{\mathcal{S}'_{pq}(B_+(0,b_j R))}) \\ &\lesssim_{a,b,n,\tau,m,\widehat{M},M,\delta} \sup_{\rho \in \{1, b_j; b_j > 1\}} \rho^{-n\tau} \|\tilde{f}\|_{\mathcal{S}'_{pq}(B(0,\rho))} \lesssim_{n,p,q,s,\tau} \|\tilde{f}\|_{\mathcal{S}'_{pq}(\mathbb{R}^n)}. \end{aligned}$$

Here $C'' = C''_{n,p,q,s,\tau,m} := m \cdot \widehat{M}_{pq}^{ns} + M_{pq}^{ns} + n\tau$ and we let $\delta > 0$ be such that $\sum_{j \in \mathbb{Z}} 2^{\delta|j|} |a_j| b_j^{[C'']} < \infty$. Taking the infimum over the extension \tilde{f} , we obtain (31) and conclude the proof of the case $s < 0$.

For $s \geq 0$, we use the following equivalent norm (see [WYY17, Theorem 1.6]):

$$(33) \quad \|\tilde{f}\|_{\mathcal{S}'_{pq}(\mathbb{R}^n)} \approx_{n,p,q,s,\tau,m} \sum_{|\gamma| \leq m} \|\partial^\gamma \tilde{f}\|_{\mathcal{S}'_{pq}^{-m,\tau}(\mathbb{R}^n)}, \quad \forall m \geq 1.$$

Using Lemma 15 (ii), and taking $m \geq s + 1$ we have

$$(34) \quad \|E^{a,b} f\|_{\mathcal{S}'_{pq}(\mathbb{R}^n)} \approx \sum_{|\gamma| \leq m} \|E^{a(-b)\gamma n, b} \partial^\gamma f\|_{\mathcal{S}'_{pq}^{-m,\tau}(\mathbb{R}^n)} \lesssim \sum_{|\gamma| \leq m} \|\partial^\gamma f\|_{\mathcal{S}'_{pq}^{-m,\tau}(\mathbb{R}_+^n)} \lesssim \|f\|_{\mathcal{S}'_{pq}(\mathbb{R}_+^n)}.$$

This completes the proof of $E : \mathcal{A}_{pq}^{s\tau}(\mathbb{R}_+^n) \rightarrow \mathcal{A}_{pq}^{s\tau}(\mathbb{R}^n)$ for $\mathcal{A} \in \{\mathcal{B}, \mathcal{F}\}$ and all $0 < p, q \leq \infty$, $s \in \mathbb{R}$ and $0 \leq \tau \leq \frac{1}{p}$ ($p < \infty$ if $\mathcal{A} = \mathcal{F}$). The case $\mathcal{A} = \mathcal{N}$ follows from real interpolations, as mentioned in the beginning of the proof. \square

Remark 25. By keeping track of the constants, and possibly enlarging the constant $\widehat{M} = \widehat{M}_{p,q}^{n,s}$ in Theorem 23, it is likely to prove that (25) is also true if we replace \mathcal{A}_{pq}^s by $\mathcal{A}_{pq}^{s\tau}$ for every $0 \leq \tau \leq \frac{1}{p}$ and $\mathcal{A} \in \{\mathcal{B}, \mathcal{F}, \mathcal{N}\}$ ($p < \infty$ for \mathcal{F} -cases).

5. THE ANALOGY ON BOUNDED SMOOTH DOMAINS

On bounded smooth domains the analogy of Theorem 1 is the following:

Theorem 26 (Universal Seeley's extension on smooth domains). *Let $\Omega \subset \mathbb{R}^n$ be a bounded smooth domain, and let $\mathcal{U} \subset \mathbb{R}^n$ be a bounded neighborhood of $\overline{\Omega}$. Then there exists an extension operator $\mathcal{E} : \mathcal{S}'(\Omega) \rightarrow \mathcal{E}'(\mathcal{U})$ that maps extendable distributions in Ω to compactly supported distributions in \mathcal{U} , such that*

(i) (One-parameter dependence) *There exists a smooth vector field $X \in C_c^\infty(\mathcal{U}; \mathbb{R}^n)$, such that for every $x \in \mathcal{U} \setminus \overline{\Omega}$ and $f \in C^0(\overline{\Omega})$, the value $\mathcal{E}f(x)$ depends only on the value of f on the (unique) integral curve for X passing through x that contains in Ω , which is, the set*

$$\{\gamma^x(t) : t \in \mathbb{R}, \gamma^x(t) \in \Omega\} \subset \mathcal{U}, \text{ where } \dot{\gamma}^x(t) = X(\gamma^x(t)) \forall t \in \mathbb{R} \text{ and } \gamma^x(0) = x.$$

(ii) (Universal boundedness) $\mathcal{E} : W^{k,p}(\Omega) \rightarrow W_c^{k,p}(\mathcal{U})$ and $\mathcal{E} : C^{k,s}(\Omega) \rightarrow C_c^{k,s}(\mathcal{U})$ are defined and bounded for all $k \in \mathbb{Z}$, $0 < p \leq \infty$ and $0 < s < 1$. In particular $\mathcal{E} : C^k(\Omega) \rightarrow C_c^k(\mathcal{U})$ for $k \geq 0$.

$\mathcal{E} : \mathcal{A}_{pq}^{s\tau}(\Omega) \rightarrow \mathcal{A}_{pq}^{s\tau}(\mathcal{U})$ is bounded for $\mathcal{A} \in \{\mathcal{B}, \mathcal{F}, \mathcal{N}\}$, $p, q \in (0, \infty]$, $s \in \mathbb{R}$ and $\tau \in [0, \frac{1}{p}]$ ($p < \infty$ for \mathcal{F} -cases). In particular $\mathcal{E} : \mathcal{A}_{pq}^s(\Omega) \rightarrow \mathcal{A}_{pq}^s(\mathcal{U})$ is bounded for $\mathcal{A} \in \{\mathcal{B}, \mathcal{F}\}$, $0 < p, q \leq \infty$ and $s \in \mathbb{R}$.

Note that (i) is false if one use the usual patching construction, for example [Tri10, (3.3.4/6)].

In the proof we need the standard partition of unity argument.

Lemma 27. *Let $\Phi : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a bounded diffeomorphism, i.e. Φ is bijective and*

$$(35) \quad \sup_{x \in \mathbb{R}^n} |\partial^\alpha \Phi(x)| + |\partial^\alpha (\Phi^{-1})(x)| < \infty, \quad \forall \alpha \in \mathbb{N}_0^n \setminus \{0\}.$$

Let $\chi \in C_c^\infty(\mathbb{R}^n)$ be a bounded smooth function. Define a linear map $Tf := (\chi \cdot f) \circ \Phi$.

(i) $T : W^{k,p}(\mathbb{R}^n) \rightarrow W^{k,p}(\mathbb{R}^n)$, $T : C^{k,s}(\mathbb{R}^n) \rightarrow C^{k,s}(\mathbb{R}^n)$ are both defined and bounded for all $k \in \mathbb{Z}$, $0 < p \leq \infty$ and $0 < s < 1$. In particular $T : C^k(\mathbb{R}^n) \rightarrow C^k(\mathbb{R}^n)$ for $k \in \mathbb{N}_0$.

(ii) $T : \mathcal{A}_{pq}^{s\tau}(\mathbb{R}^n) \rightarrow \mathcal{A}_{pq}^{s\tau}(\mathbb{R}^n)$ for $\mathcal{A} \in \{\mathcal{B}, \mathcal{F}, \mathcal{N}\}$, $p, q \in (0, \infty]$, $s \in \mathbb{R}$ and $\tau \in [0, \frac{1}{p}]$ ($p < \infty$ if $\mathcal{A} = \mathcal{F}$). In particular $T : \mathcal{A}_{pq}^s(\mathbb{R}^n) \rightarrow \mathcal{A}_{pq}^s(\mathbb{R}^n)$ is bounded for $\mathcal{A} \in \{\mathcal{B}, \mathcal{F}\}$, for $p, q \in (0, \infty]$ and $s \in \mathbb{R}$.

Proof. The result (i) follows from using chain rules, product rules and the formula of change of variables:

$$\int_{\mathbb{R}^n} |(\chi f)(\Phi(x))|^p dx = \int_{\mathbb{R}^n} |\chi(y)f(y)|^p |\det \nabla(\Phi^{-1})(y)| dy, \quad 0 < p < \infty, \quad f \in L_{\text{loc}}^p(\mathbb{R}^n).$$

Notice that $|\det \nabla(\Phi^{-1})|$ is a bounded smooth function, and all nonzero derivatives of Φ and χ are bounded. We leave the details to readers.

For (ii), see for example [YSY10, Sections 6.1.1 and 6.2] and [HT21, Sections 4.2 and 4.3]. \square

Proof of Theorem 26. In the following, we let \mathcal{X} be the function class that belongs to the following:

$$\begin{aligned} & \{W^{k,p}, C^{k,s} : k \in \mathbb{Z}, p \in (0, \infty], s \in (0, 1)\} \cup \{\mathcal{B}_{pq}^s, \mathcal{F}_{pq}^s, \mathcal{B}_{pq}^{s\tau}, \mathcal{N}_{pq}^{s\tau} : p, q \in (0, \infty], s \in \mathbb{R}, \tau \in [0, \frac{1}{p}]\} \\ & \cup \{C^k : k \in \mathbb{N}_0\} \cup \{\mathcal{F}_{pq}^{s\tau} : p \in (0, \infty), q \in (0, \infty], s \in \mathbb{R}, \tau \in [0, \frac{1}{p}]\}. \end{aligned}$$

Recall from Theorem 1 that $E^{a,b} : \mathcal{X}(\mathbb{R}_+^n) \rightarrow \mathcal{X}(\mathbb{R}^n)$ is bounded for all such \mathcal{X} .

For a function $g(\theta, t)$ defined on $\partial\Omega \times \mathbb{R}_+$, we define its extension $\mathcal{E}_0 g$ on $\partial\Omega \times \mathbb{R}$ by

$$(36) \quad \mathcal{E}_0 g(\theta, t) := \begin{cases} \sum_{j=-\infty}^{\infty} a_j \cdot g(\theta, -b_j t) & t < 0; \\ g(\theta, t) & t > 0 \end{cases}, \quad \text{where } (a_j, b_j)_{j=-\infty}^{\infty} \text{ satisfies (3).}$$

Recall from Proposition 14 that such $(a_j, b_j)_j$ exist.

We construct the following objects:

- A tubular neighborhood $U_0 \subset\subset \mathcal{U}$ of $\partial\Omega$ along with a diffeomorphism² $\Psi = (\psi', \psi_n) : U_0 \xrightarrow{\sim} \partial\Omega \times (-1, 1)$ such that $U_0 \cap \Omega = \{\psi_n > 0\}$.
- A coordinate cover $\{\phi'_\lambda : V_\lambda \subseteq \partial\Omega \rightarrow \mathbb{R}^{n-1}\}_{\lambda=1}^N$ of $\partial\Omega$, where V_λ are open in the submanifold $\partial\Omega$, along with the coordinate charts $\Phi_\lambda(\theta, t) := (\phi'_\lambda(\theta), t)$ defined on $V_\lambda \times (-1, 1) \subset \partial\Omega \times (-1, 1)$.

By choosing U_0 and each V_λ small enough, we can ensure that

- For every $1 \leq \lambda \leq N$, $\Phi_\lambda \circ \Psi : \Psi^{-1}(V_\lambda \times (-1, 1)) \subseteq \mathcal{U} \rightarrow \mathbb{R}^n$ can extend to a bounded diffeomorphism. In other words, there are $\Xi_\lambda : \mathbb{R}^n \rightarrow \mathbb{R}^n$ that satisfies (35) and $\Xi_\lambda|_{\Psi^{-1}(V_\lambda \times (-1, 1))} = \Phi_\lambda \circ \Psi$, for each λ .

For such Ψ and $\{\phi'_\lambda\}_\lambda$, we take the following cutoff functions:

$$\chi_0, \chi_1 \in C_c^\infty(-1, 1) \text{ such that } \chi_0|_{[-\frac{1}{2}, \frac{1}{2}]} \equiv 1, \chi_1|_{\text{supp } \chi_0} \equiv 1; \quad \text{and} \quad \tilde{\chi}_0 := \chi_0 \circ \psi_n, \tilde{\chi}_1 := \chi_1 \circ \psi_n \in C_c^\infty(U_0).$$

We define \mathcal{E} as

$$(37) \quad \mathcal{E}f := (1 - \tilde{\chi}_0)|_\Omega \cdot f + \tilde{\chi}_1 \cdot (\mathcal{E}_0[(\tilde{\chi}_0 \cdot f) \circ \Psi^{-1}] \circ \Psi).$$

Clearly $\mathcal{E}f|_\Omega = (1 - \tilde{\chi}_0) \cdot f + \tilde{\chi}_1 \cdot \tilde{\chi}_0 \cdot f = f$. Therefore \mathcal{E} is an extension operator.

Take $\tilde{\chi}_2 \in C_c^\infty(\mathcal{U})$ be such that $\tilde{\chi}_2|_{\text{supp } \tilde{\chi}_1} \equiv 1$. Immediately we see that the vector field $X := \tilde{\chi}_2 \cdot (\Psi^* \frac{\partial}{\partial t})$ is what we need for (i).

We postpone the proof of $\mathcal{E} : \mathcal{S}'(\Omega) \rightarrow \mathcal{E}'(\mathcal{U})$ to the end.

For (ii), to prove the boundedness for $\tilde{\chi}_1(\mathcal{E}_0[(\tilde{\chi}_0 f) \circ \Psi^{-1}] \circ \Psi)$, we take the following objects:

- $\{\mu_\lambda \in C_c^\infty(V_\lambda)\}_{\lambda=1}^N$ be a partition of unity of $\partial\Omega$, i.e. $\sum_{\lambda=1}^N \mu_\lambda \equiv 1$ on $\partial\Omega$.
- $\{\rho_\lambda \in C_c^\infty(V_\lambda)\}_{\lambda=1}^N$ be such that $\rho_\lambda|_{\text{supp } \mu_\lambda} \equiv 1$.
- For $1 \leq \lambda \leq N$, we let

$$\tilde{\mu}_\lambda := (\mu_\lambda \otimes \chi_0) \circ \Psi = (\mu_\lambda \circ \psi') \cdot (\chi_0 \circ \psi_n) \quad \text{and} \quad \tilde{\rho}_\lambda(x) := (\rho_\lambda \otimes \chi_1) \circ \Phi_\lambda^{-1}(x) = (\rho_\lambda \circ \phi'_\lambda)^{-1}(x') \cdot \chi_1(x_n).$$

Immediately we see that

- $\tilde{\mu}_\lambda \in C_c^\infty(\Psi^{-1}(V_\lambda \times (-1, 1)))$ and $\tilde{\rho}_\lambda \in C_c^\infty(\Phi_\lambda(V_\lambda \times (-1, 1)))$.
- When $\text{supp } g \subseteq (\text{supp } \mu_\lambda) \times [0, \infty)$, we have $\text{supp } \mathcal{E}_0 g \subseteq (\text{supp } \mu_\lambda) \times \mathbb{R}$.
In particular $(\mathbf{1}_{\partial\Omega} \otimes \chi_1) \cdot \mathcal{E}_0[(\mu_\lambda \otimes \chi_0) \cdot g] = (\rho_\lambda \otimes \chi_1) \cdot \mathcal{E}_0[(\mu_\lambda \otimes \chi_0) \cdot g]$ (since $\rho_\lambda|_{\text{supp } \mu_\lambda} \equiv 1$).
- When $\text{supp } g \subseteq V_\lambda \times [0, 1)$, we have $\mathcal{E}_0[g]|_{V_\lambda \times (-1, 1)} = E[g \circ \Phi_\lambda^{-1}] \circ \Phi_\lambda$, the right hand side of which is well-defined. Here E is the extension operator from (2).

Therefore

$$\begin{aligned} \tilde{\chi}_1 \cdot (\mathcal{E}_0[(\tilde{\chi}_0 \cdot f) \circ \Psi^{-1}] \circ \Psi) &= \sum_{\lambda=1}^N ((\mathbf{1}_{\partial\Omega} \otimes \chi_1) \cdot \mathcal{E}_0[(\tilde{\mu}_\lambda \cdot f) \circ \Psi^{-1}]) \circ \Psi \\ &= \sum_{\lambda=1}^N ((\rho_\lambda \otimes \chi_1) \cdot (E[(\tilde{\mu}_\lambda \cdot f) \circ \Psi^{-1} \circ \Phi_\lambda^{-1}] \circ \Phi_\lambda)) \circ \Psi = \sum_{\lambda=1}^N (\tilde{\rho}_\lambda \cdot E[(\tilde{\mu}_\lambda \cdot f) \circ (\Phi_\lambda \circ \Psi)^{-1}]) \circ (\Phi_\lambda \circ \Psi). \end{aligned}$$

Applying Lemma 27 and Theorem 1 (ii) - (iv), we have the boundedness of compositions

$$\mathcal{X}^c(\Omega) \xrightarrow{f \mapsto ((\tilde{\mu}_\lambda f) \circ (\Phi_\lambda \circ \Psi)^{-1})_{\lambda=1}^N} \mathcal{X}^c(\mathbb{R}_+^n)^N \xrightarrow{E} \mathcal{X}^c(\mathbb{R}^n)^N \xrightarrow{(g_\lambda)_{\lambda=1}^N \mapsto \sum_{\lambda=1}^N (\tilde{\rho}_\lambda g) \circ (\Phi_\lambda \circ \Psi)} \mathcal{X}^c(U_0) \subset \mathcal{X}^c(\mathcal{U}).$$

On the other hand, $\tilde{\chi}_0 = \chi_0 \circ \psi_n \in C_c^\infty(U_0)$ is identical to 1 near $\partial\Omega$, therefore $(1 - \tilde{\chi}_0)|_\Omega \in C_c^\infty(\Omega)$. By Lemma 27 $[f \mapsto (1 - \tilde{\chi}_0)|_\Omega \cdot f] : \mathcal{X}^c(\Omega) \rightarrow \mathcal{X}^c(\mathcal{U})$ is bounded. Altogether we conclude that $\mathcal{E} : \mathcal{X}^c(\Omega) \rightarrow \mathcal{X}^c(\mathcal{U})$ is bounded. This completes the proof of (ii).

Finally we illustrate the definedness of $\mathcal{E} : \mathcal{S}'(\Omega) \rightarrow \mathcal{E}'(\mathcal{U})$. Note that alternatively one can obtain this statement by using duality and prove the boundedness $\mathcal{E}^* : C_{\text{loc}}^\infty(\mathcal{U}) \rightarrow \mathcal{S}'(\Omega) (= \{g \in C^\infty(\mathbb{R}^n) : \text{supp } g \subseteq \bar{\Omega}\})$.

For a $f \in \mathcal{S}'(\Omega)$, let \tilde{f} be an extension of f . Since Ω is a bounded domain, we can assume \tilde{f} to have compact support, otherwise we replace \tilde{f} with $\kappa \tilde{f}$ where $\kappa \in C_c^\infty(\mathbb{R}^n)$ is such that $\kappa|_\Omega \equiv 1$.

By the structure theorem of distributions (see [Rud91, Theorem 6.27] for example), there are $M \geq 0$ and $\{g_\alpha\}_{|\alpha| \leq M} \subset C_c^0(\mathbb{R}^n)$ such that $\tilde{f} = \sum_{|\alpha| \leq M} \partial^\alpha g_\alpha$. By Proposition 21 (i) this is saying that $\tilde{f} \in \mathcal{B}_{\infty\infty}^{-M}(\mathbb{R}^n)$ for some $M > 0$, which means $f \in \mathcal{B}_{\infty\infty}^{-M}(\Omega)$ for the same M .

Now by (ii) $\mathcal{E}f \in \mathcal{B}_{\infty\infty}^{-M}(\mathcal{U}) \subset \mathcal{D}'(\mathcal{U})$ has compact support, we conclude that $\mathcal{E}f \in \mathcal{E}'(\mathcal{U})$. \square

²We do not say Ψ a coordinate chart of U_0 because its codomain is not a subset of \mathbb{R}^n .

REFERENCES

- [Ada75] Robert A. Adams, *Sobolev spaces*, Pure and Applied Mathematics, Vol. 65, Academic Press [Harcourt Brace Jovanovich, Publishers], New York-London, 1975. MR 0450957
- [Bab53] V. M. Babič, *On the extension of functions*, Uspehi Matem. Nauk (N.S.) **8** (1953), no. 2(54), 111–113. MR 0056675
- [ET96] D. E. Edmunds and H. Triebel, *Function spaces, entropy numbers, differential operators*, Cambridge Tracts in Mathematics, vol. 120, Cambridge University Press, Cambridge, 1996. MR 1410258
- [Eva10] Lawrence C. Evans, *Partial differential equations*, second ed., Graduate Studies in Mathematics, vol. 19, American Mathematical Society, Providence, RI, 2010. MR 2597943
- [Fra86] Jens Franke, *On the spaces \mathbf{F}_{pq}^s of Triebel-Lizorkin type: pointwise multipliers and spaces on domains*, Math. Nachr. **125** (1986), 29–68. MR 847350
- [GT01] David Gilbarg and Neil S. Trudinger, *Elliptic partial differential equations of second order*, Classics in Mathematics, Springer-Verlag, Berlin, 2001, Reprint of the 1998 edition. MR 1814364
- [Hes41] M. R. Hestenes, *Extension of the range of a differentiable function*, Duke Math. J. **8** (1941), 183–192. MR 3434
- [HT21] Dorothee D. Haroske and Hans Triebel, *Morrey smoothness spaces: a new approach*, arXiv:2110.10609v1, 2021.
- [JMHS15] J. Johnsen, S. Munch Hansen, and W. Sickel, *Anisotropic Lizorkin-Triebel spaces with mixed norms—traces on smooth boundaries*, Math. Nachr. **288** (2015), no. 11-12, 1327–1359. MR 3377120
- [Lic29] Leon Lichtenstein, *Eine elementare Bemerkung zur reellen Analysis*, Math. Z. **30** (1929), no. 1, 794–795. MR 1545093
- [Nik53] S. M. Nikol'skii, *Properties of certain classes of functions of several variables on differentiable manifolds*, Mat. Sb. N.S. **33(75)** (1953), 261–326. MR 0065605
- [Pee75] Jaak Peetre, *A remark on Sobolev spaces. The case $0 < p < 1$* , J. Approximation Theory **13** (1975), 218–228. MR 374900
- [Pre97] David Preiss, *Additional regularity for Lipschitz solutions of PDE*, J. Reine Angew. Math. **485** (1997), 197–207. MR 1442194
- [Rud87] Walter Rudin, *Real and complex analysis*, third ed., McGraw-Hill Book Co., New York, 1987. MR 924157
- [Rud91] ———, *Functional analysis*, second ed., International Series in Pure and Applied Mathematics, McGraw-Hill, Inc., New York, 1991. MR 1157815
- [Ryc99] Vyacheslav S. Rychkov, *On restrictions and extensions of the Besov and Triebel-Lizorkin spaces with respect to Lipschitz domains*, J. London Math. Soc. (2) **60** (1999), no. 1, 237–257. MR 1721827
- [See64] R. T. Seeley, *Extension of C^∞ functions defined in a half space*, Proc. Amer. Math. Soc. **15** (1964), 625–626. MR 165392
- [Sic12] W. Sickel, *Smoothness spaces related to Morrey spaces—a survey. I*, Eurasian Math. J. **3** (2012), no. 3, 110–149. MR 3024132
- [SY21] Ziming Shi and Liding Yao, *Existence of solutions for $\bar{\partial}$ -equations in Sobolev spaces of negative index*, arXiv:2111.09245, 2021.
- [Tri92] Hans Triebel, *Theory of function spaces. II*, Monographs in Mathematics, vol. 84, Birkhäuser Verlag, Basel, 1992. MR 1163193
- [Tri95] ———, *Interpolation theory, function spaces, differential operators*, second ed., Johann Ambrosius Barth, Heidelberg, 1995. MR 1328645
- [Tri01] ———, *The structure of functions*, Monographs in Mathematics, vol. 97, Birkhäuser Verlag, Basel, 2001. MR 1851996
- [Tri06] ———, *Theory of function spaces. III*, Monographs in Mathematics, vol. 100, Birkhäuser Verlag, Basel, 2006. MR 2250142
- [Tri10] ———, *Theory of function spaces*, Modern Birkhäuser Classics, Birkhäuser/Springer Basel AG, Basel, 2010, Reprint of 1983 edition [MR0730762], Also published in 1983 by Birkhäuser Verlag [MR0781540]. MR 3024598
- [Tri14] ———, *Hybrid function spaces, heat and Navier-Stokes equations*, EMS Tracts in Mathematics, vol. 24, European Mathematical Society (EMS), Zürich, 2014. MR 3308920
- [Tri20] ———, *Theory of function spaces IV*, Monographs in Mathematics, vol. 107, Birkhäuser/Springer, Cham, [2020] ©2020. MR 4298338
- [Wu98] Deyun Wu, *Sobolev and Hölder estimates for $\bar{\partial}$ on bounded convex domains in \mathbf{C}^2* , Illinois J. Math. **42** (1998), no. 3, 371–388. MR 1631224
- [WYY17] Suqing Wu, Dachun Yang, and Wen Yuan, *Equivalent quasi-norms of Besov-Triebel-Lizorkin-type spaces via derivatives*, Results Math. **72** (2017), no. 1-2, 813–841. MR 3684461
- [Yao22] Liding Yao, *Sobolev and Hölder estimates for homotopy operators of $\bar{\partial}$ -equations on convex domains of finite multitype*, arXiv:2210.15830v1, 2022.
- [YSY10] Wen Yuan, Winfried Sickel, and Dachun Yang, *Morrey and Campanato meet Besov, Lizorkin and Triebel*, Lecture Notes in Mathematics, vol. 2005, Springer-Verlag, Berlin, 2010. MR 2683024
- [YY13] Dachun Yang and Wen Yuan, *Relations among Besov-type spaces, Triebel-Lizorkin-type spaces and generalized Carleson measure spaces*, Appl. Anal. **92** (2013), no. 3, 549–561. MR 3021276

HAOWEN LU, SCHOOL OF MATHEMATICS, SUN YAT-SEN UNIVERSITY, GUANGZHOU, GUANGDONG, 510275, P. R. CHINA
Email address: sun7aris@gmail.com

LIDING YAO, DEPARTMENT OF MATHEMATICS, THE OHIO STATE UNIVERSITY, COLUMBUS, OH 43210, UNITED STATES
Email address: yao.1015@osu.edu