

# Spectrum of the $\bar{\partial}$ -Laplace operator on zero forms for the quantum quadric $\mathcal{O}_q(\mathbf{Q}_N)$

Fredy Díaz García

Mathematical Institute of Charles University, Prague, Czech Republic

E-mail address: *lenonndiaz@gmail.com*

## Abstract

We study the Laplacian operator  $\Delta_{\bar{\partial}}$  associated to a Kähler structure  $(\Omega^{(\bullet,\bullet)}, \kappa)$  for the Heckenberger–Kolb differential calculus of the quantum quadrics  $\mathcal{O}_q(\mathbf{Q}_N)$ , which is to say, the irreducible quantum flag manifolds of types  $B_n$  and  $D_n$ . We show that the eigenvalues of  $\Delta_{\bar{\partial}}$  on zero forms tend to infinity and have finite multiplicity.

## 1 Introduction

Since the discovery of quantum groups in the 1980's mathematicians have been trying to fit them into Connes' framework of noncommutative spin geometry. In fact, the construction of spectral triples for the Drinfeld–Jimbo quantum groups has been a very difficult problem in noncommutative geometry. One of the best-studied spectral triples is that of the Podleś sphere [6] which satisfies most of the Connes conditions for a noncommutative spin geometry, up to some slight modifications. Other constructions have been made in this direction, for example the equivariant isospectral Dirac operators for all the Podleś spheres [4], for the quantum projective spaces [2], for the quantum group  $\mathcal{O}_q(SU(2))$  [5] and for all the compact quantum groups [23]. In [7] a Dirac operator associated to Kähler structures is constructed for the quantum projective spaces  $\mathcal{O}_q(\mathbb{C}\mathbb{P}^n)$  generalizing the work of S. Majid in [20].

In recent years it has been shown that the quantum flag manifolds provide a large family of well behaved quantum homogeneous spaces whose noncommutative geometry is quite similar to the classical situation. It was shown by I. Heckenberger and S. Kolb that the subclass of irreducible quantum flag manifolds  $\mathcal{O}_q(G/L_S)$  admit a unique differential calculus which is a  $q$ -deformation of their classical de Rham complex [12], [13]. It is important to note that the existence of this differential calculus is an extremely especial property of these quantum spaces and it is not the case for all quantum spaces. Moreover the existence of this canonical deformation allows us to construct Dirac operators and investigate their spectrum.

The problem of the spectrum of the Dirac operator has been attacked for some cases. For example in [3] the spectrum for  $\mathcal{O}_q(\mathbb{C}\mathbb{P}^2)$  was calculated. In [2] the authors showed that the spectrum of the Dolbeault–Dirac operator for the quantum projective spaces has exponential growth by relating a Casimir operator. In [18] the authors gave a new approach for the construction of Dolbeault–Dirac operator and quantum Clifford algebras for the irreducible quantum

flag manifolds. Later, it was shown in [22] that this Dolbeault–Dirac operator gives a spectral triple for the quantum Lagrangian Grassmannian of rank 2 by giving a suitable version of the Parthasarathy formula.

Recently, a different approach to constructing Dolbeault–Dirac operators has appeared in [24] where the notion of noncommutative Kähler structure for a differential calculus was introduced. In the same paper the author showed the existence of a covariant Kähler structure for the Heckenberger–Kolb calculus for the quantum projective spaces. Later this result was generalized in [21] for all the irreducible quantum flag manifolds, for all but a finite number of values  $q$ . The existence of such a structure allows us to construct metrics, Hilbert space completions and to verify all the axioms of a spectral triple except the compact resolvent condition, see [8] for a more detailed discussion. This last axiom has turned out to be the most challenging one. This problem was solved in [7] for the quantum projective spaces  $\mathcal{O}_q(\mathbb{C}\mathbb{P}^n)$  by using the existence of the Kähler structure and the multiplicity-free property of the space of anti-holomorphic forms. Also the spectrum was calculated for the quantum quadric  $\mathcal{O}_q(\mathbf{Q}_5)$  in [10] by using the quantum version of the Gelfand–Bernstein–Gelfand resolution for quantized irreducible flag manifolds [14].

In this paper we investigate the asymptotic behaviour of the Laplace operator  $\Delta_{\bar{\partial}}$  on the space of zero forms  $\Omega^{(0,0)}$  for the irreducible quantum flag manifolds of quadric type. Besides of the existence of the Kähler form we make use of the FRT description of the quantum group  $\mathcal{O}_q(SO(N))$  as well as the fact that  $\Omega^{(0,0)}$  is multiplicity-free as a  $U_q(\mathfrak{g})$ -module to make explicit calculations of the eigenvalues. It has been conjectured in [7] that the Dolbeault–Dirac operator  $D_{\bar{\partial}}$  allows us to construct a spectral triple for all the irreducible quantum flag manifolds, here we take a small but significant step towards proving this conjecture.

The paper is organized as follows: in §2 we recall some necessary definitions and properties of the quantized enveloping algebras, differential calculi, Kähler structures as well as irreducible quantum flag manifolds and Takeuchi’s equivalence. We also recall the FRT description of the algebras  $\mathcal{O}_q(SO(N))$ .

In §3 we give the most important properties of the Heckenberger–Kolb calculus for the irreducible quantum flag manifolds and a proof of the existence of a Kähler structure for the case of quantum quadrics  $\mathcal{O}_q(\mathbf{Q}_N)$  by using tools of representation theory.

In §4 we present the main result of the paper. It is shown by explicit calculation that the eigenvalues of the Laplace operator  $\Delta_{\bar{\partial}}$  on zero forms tend to infinity, supporting the conjecture that the asymptotic behaviour of the Dolbeault–Dirac operator  $D_{\bar{\partial}}$  satisfies the compact resolvent condition for all the irreducible quantum flag manifolds.

## 2 Preliminaries

In this section we recall the necessary definitions of quantum enveloping algebras, covariant differential calculi as well as complex and Kähler structures. For a more detailed presentation see [17], [24], and [25]. We also introduce the definition of quantum flag manifolds, Takeuchi’s equivalence and the FRT presentation of the quantum coordinate algebra  $\mathcal{O}_q(SO(N))$ .

### 2.1 Drinfeld–Jimbo quantum groups

Let  $\mathfrak{g}$  a finite dimensional semisimple Lie algebra of rank  $r$ . Let  $\mathfrak{g}$  be a finite-dimensional complex semisimple Lie algebra of rank  $r$ . We fix a Cartan subalgebra  $\mathfrak{h}$  and choose a set of simple roots  $\Pi = \{\alpha_1, \dots, \alpha_r\}$  for the corresponding root system in the linear dual of  $\mathfrak{g}$ . We denote by  $(\cdot, \cdot)$  the

symmetric bilinear form induced on  $\mathfrak{h}^*$  by the Killing form of  $\mathfrak{g}$ , normalised so that any shortest simple root  $\alpha_i$  satisfies  $(\alpha_i, \alpha_i) = 2$ . The Cartan matrix  $(a_{ij})$  of  $\mathfrak{g}$  is defined by  $a_{ij} := (\alpha_i^\vee, \alpha_j)$ , where  $\alpha_i^\vee := 2\alpha_i/(\alpha_i, \alpha_i)$  is the *coroot* of  $\alpha_i$ .

Throughout the paper  $q$  stands for a real number in the interval  $(1, +\infty)$  and denote  $q_i := q^{(\alpha_i, \alpha_i)/2}$ . The *Drinfeld–Jimbo quantised enveloping algebra*  $U_q(\mathfrak{g})$  is the noncommutative associative algebra generated by the elements  $E_i, F_i, K_i$ , and  $K_i^{-1}$ , for  $i = 1, \dots, r$ , with the relations

$$\begin{aligned} K_i E_j &= q_i^{a_{ij}} E_j K_i, & K_i F_j &= q_i^{-a_{ij}} F_j K_i, & K_i K_j &= K_j K_i, & K_i K_i^{-1} &= K_i^{-1} K_i = 1, \\ E_i F_j - F_j E_i &= \delta_{ij} \frac{K_i - K_i^{-1}}{q_i - q_i^{-1}}, \end{aligned}$$

and the quantum Serre relations which we omit (see [17, §6.1.2] for details).

A Hopf algebra structure on  $U_q(\mathfrak{g})$  is defined by

$$\begin{aligned} \Delta(K_i) &= K_i \otimes K_i, & \Delta(E_i) &= E_i \otimes K_i + 1 \otimes E_i, & \Delta(F_i) &= F_i \otimes 1 + K_i^{-1} \otimes F_i, \\ S(K_i) &= K_i^{-1}, & S(E_i) &= -E_i K_i^{-1}, & S(F_i) &= -K_i F_i, & \varepsilon(K_i) &= 1, & \varepsilon(E_i) &= \varepsilon(F_i) = 0. \end{aligned}$$

A Hopf  $*$ -algebra structure is given by

$$K_i^* = K_i, \quad E_i^* = K_i F_i, \quad F_i^* = E_i K_i^{-1}.$$

Let  $\{\varpi_1, \dots, \varpi_r\}$  denote the corresponding set of *fundamental weights* of  $\mathfrak{g}$  which is the dual basis of the simple coroots  $\{\alpha_i^\vee, \dots, \alpha_r^\vee\}$ :

$$(\alpha_i^\vee, \varpi_j) = \delta_{ij}.$$

Let  $\mathcal{P}$  be the weight lattice of  $\mathfrak{g}$  and  $\mathcal{P}^+$  the set of *dominant integral weights* which are the  $\mathbb{Z}$ -span and  $\mathbb{Z}_{\geq 0}$ -span of the fundamental weights, respectively. For each  $\mu \in \mathcal{P}^+$  there exists an irreducible finite dimensional  $U_q(\mathfrak{g})$ -module  $V_\mu$  uniquely determined by the existence of a vector  $\nu_\mu$  which is called the *highest weight vector*, satisfying

$$K_i \triangleright \nu_\mu = q^{(\alpha_i, \mu)} \nu_\mu, \quad E_i \triangleright \nu_\mu = 0.$$

Moreover  $\nu_\mu$  is unique up to scalar multiple. A finite direct sum of such representations is called a *type-1 representation*. A vector  $\nu$  is called a *weight vector* of weight  $\text{wt}(\nu) \in \mathcal{P}$  if

$$K_i \triangleright \nu = q^{(\alpha_i, \text{wt}(\nu))} \nu.$$

In this paper we use the fact that  $U_q(\mathfrak{g})$  has invertible antipode and therefore we have an equivalence between  $U_q(\mathfrak{g})\text{Mod}$  and  $\text{Mod}_{U_q(\mathfrak{g})}$  induced by the antipode.

## 2.2 Differential calculi

A *differential calculus*  $(\Omega^\bullet \simeq \bigoplus_{k \in \mathbb{N}_0} \Omega^k, d)$  is a differential graded algebra (dg-algebra) which is generated in degree 0 as a dg-algebra, that is to say, it is generated as an algebra by the elements  $a, db$ , for  $a, b \in \Omega^0$ . We call an element  $\omega \in \Omega^\bullet$  a *form*, and if  $\omega \in \Omega^k$ , for some  $k \in \mathbb{N}_0$ , then  $\omega$  is said to be *homogeneous* of degree  $|\omega| := k$ . The product of two forms  $\omega, \nu \in \Omega^\bullet$  is usually

denoted by  $\omega \wedge \nu$ , unless one of the forms is of degree 0, where we just denote the product by juxtaposition.

For a given algebra  $B$ , a *differential calculus over  $B$*  is a differential calculus such that  $\Omega^0 = B$ . Note that for a differential calculus over  $B$ , each  $\Omega^k$  is a  $B$ -bimodule. A differential calculus is said to have *total degree*  $m \in \mathbb{N}_0$ , if  $\Omega^m \neq 0$ , and  $\Omega^k = 0$ , for every  $k > m$ .

A *differential \*-calculus* over a \*-algebra  $B$  is a differential calculus over  $B$  such that the \*-map of  $B$  extends to a (necessarily unique) conjugate-linear involutive map  $*$  :  $\Omega^\bullet \rightarrow \Omega^\bullet$  satisfying  $d(\omega^*) = (d\omega)^*$ , and

$$(\omega \wedge \nu)^* = (-1)^{kl} \nu^* \wedge \omega^*, \quad \text{for all } \omega \in \Omega^k, \nu \in \Omega^l.$$

We say that  $\omega \in \Omega^\bullet$  is *closed* if  $d\omega = 0$ , and *real* if  $\omega^* = \omega$ .

We now recall the definition of a complex structure as introduced in [25]. This abstracts the properties of the de Rham complex for a classical complex manifold [16].

**Definition 2.1.** An *almost complex structure*  $\Omega^{(\bullet, \bullet)}$  for a differential \*-calculus  $(\Omega^\bullet, d)$  is an  $\mathbb{N}_0^2$ -algebra grading  $\bigoplus_{(a,b) \in \mathbb{N}_0^2} \Omega^{(a,b)}$  for  $\Omega^\bullet$  such that, for all  $(a, b) \in \mathbb{N}_0^2$  the following hold:

- (i)  $\Omega^k = \bigoplus_{a+b=k} \Omega^{(a,b)}$ ,
- (ii)  $(\Omega^{(a,b)})^* = \Omega^{(b,a)}$ .

We call an element of  $\Omega^{(a,b)}$  an  $(a, b)$ -form. For the associated pair of projections  $\text{proj}_{\Omega^{(a+1,b)}} : \Omega^{a+b+1} \rightarrow \Omega^{(a+1,b)}$  and  $\text{proj}_{\Omega^{(a,b+1)}} : \Omega^{a+b+1} \rightarrow \Omega^{(a,b+1)}$ , we denote

$$\partial|_{\Omega^{(a,b)}} := \text{proj}_{\Omega^{(a+1,b)}} \circ d, \quad \bar{\partial}|_{\Omega^{(a,b)}} := \text{proj}_{\Omega^{(a,b+1)}} \circ d.$$

A *complex structure* is an almost complex structure which satisfies

$$d\Omega^{(a,b)} \subseteq \Omega^{(a+1,b)} \oplus \Omega^{(a,b+1)}, \quad \text{for all } (a, b) \in \mathbb{N}_0^2. \quad (1)$$

For a complex structure, (1) implies the identities

$$d = \partial + \bar{\partial}, \quad \bar{\partial} \circ \partial = -\partial \circ \bar{\partial}, \quad \partial^2 = \bar{\partial}^2 = 0.$$

In particular,  $(\bigoplus_{(a,b) \in \mathbb{N}_0^2} \Omega^{(a,b)}, \partial, \bar{\partial})$  is a double complex, which we call the *Dolbeault double complex* of  $\Omega^{(\bullet, \bullet)}$ . Moreover, it is easily seen that both  $\partial$  and  $\bar{\partial}$  satisfy the graded Leibniz rule, and

$$\partial(\omega^*) = (\bar{\partial}\omega)^*, \quad \bar{\partial}(\omega^*) = (\partial\omega)^*, \quad \text{for all } \omega \in \Omega^\bullet. \quad (2)$$

### 2.2.1 Hermitian and Kähler structures

We now present the definition of an Hermitian structure, as introduced in [24], as well as a Kähler structure, introduced in §7 of the same paper.

**Definition 2.2.** An *Hermitian structure*  $(\Omega^{(\bullet, \bullet)}, \sigma)$  for a differential  $*$ -calculus  $\Omega^\bullet$  over  $B$  of even total degree  $2n$  is a pair consisting of a complex structure  $\Omega^{(\bullet, \bullet)}$  and a central real  $(1, 1)$ -form  $\sigma$ , called the *Hermitian form*, such that, with respect to the *Lefschetz operator*

$$L : \Omega^\bullet \rightarrow \Omega^\bullet, \quad \omega \mapsto \sigma \wedge \omega,$$

isomorphisms are given by

$$L^{n-k} : \Omega^k \rightarrow \Omega^{2n-k}, \quad \text{for all } k = 0, \dots, n-1.$$

For  $L$  the Lefschetz operator of an Hermitian structure, we denote

$$P^{(a,b)} := \begin{cases} \{\alpha \in \Omega^{(a,b)} \mid L^{n-a-b+1}(\alpha) = 0\}, & \text{if } a+b \leq n, \\ 0, & \text{if } a+b > n. \end{cases}$$

Moreover, we denote  $P^k := \bigoplus_{a+b=k} P^{(a,b)}$ , and  $P^\bullet := \bigoplus_{k \in \mathbb{N}_0} P^k$ . An element of  $P^\bullet$  is called a *primitive form*.

An important consequence of the existence of the Lefschetz operator is the Lefschetz decomposition of the differential  $*$ -calculus, which we now recall below. For a proof see [24, §4.1].

**Proposition 2.3** (Lefschetz decomposition). *For  $L$  the Lefschetz operator of an Hermitian structure on a differential  $*$ -calculus  $\Omega^\bullet$ , a  $B$ -bimodule decomposition of  $\Omega^k$ , for all  $k \in \mathbb{N}_0$ , is given by*

$$\Omega^k \simeq \bigoplus_{j \geq 0} L^j (P^{k-2j}).$$

In classical Hermitian geometry, the Hodge map of an Hermitian metric is related to the associated Lefschetz decomposition through the well-known Weil formula [16, §1.2]. In the noncommutative setting, we take the direct generalisation of the Weil formula for our definition of the Hodge map, and build upon this to define an Hermitian metric.

**Definition 2.4.** The *Hodge map* associated to an Hermitian structure  $(\Omega^{(\bullet, \bullet)}, \sigma)$  is the  $B$ -bimodule map  $*_\sigma : \Omega^\bullet \rightarrow \Omega^\bullet$  satisfying, for any  $j \in \mathbb{N}_0$ ,

$$*_\sigma(L^j(\omega)) = (-1)^{\frac{k(k+1)}{2}} \mathbf{i}^{a-b} \frac{j!}{(n-j-k)!} L^{n-j-k}(\omega), \quad \omega \in P^{(a,b)} \subseteq P^{k=a+b},$$

where  $\mathbf{i}^2 = -1$ .

The metric associated to a Hermitian structure  $(\Omega^{(\bullet, \bullet)}, \sigma)$  is the map  $g_\sigma : \Omega^\bullet \times \Omega^\bullet \rightarrow B$ , for which  $g(\Omega^k, \Omega^l) = 0$  for  $k \neq l$  and

$$g_\sigma(\omega, \nu) := *_\sigma(*_\sigma(\omega^*) \wedge \nu), \quad \text{if } \omega, \nu \in \Omega^k.$$

**Definition 2.5.** We say that an Hermitian structure  $(\Omega^{(\bullet, \bullet)}, \sigma)$  is *positive definite* if the associated metric  $g_\sigma$  satisfies

$$g_\sigma(\omega, \omega) \in B_{>0} := \left\{ \sum_{i=1}^m \lambda_i b^* b_i : b_i \in B, \lambda_i \in \mathbb{R}_{>0}, m \in \mathbb{N} \right\}, \quad \text{for all non-zero } \omega \in \Omega^\bullet.$$

In this case we say that  $\sigma$  is a *positive definite Hermitian form*.

**Definition 2.6.** A *Kähler structure* for a differential  $*$ -calculus is an Hermitian structure  $(\Omega^{(\bullet, \bullet)}, \kappa)$  such that  $\kappa$  is closed, which is to say,  $d\kappa = 0$ . We call such a form  $\kappa$  a *Kähler form*.

### 2.3 Irreducible quantum flag manifolds

Let  $V$  be a finite-dimensional  $U_q(\mathfrak{g})$ -module,  $v \in V$ , and  $f \in V^*$ , the linear dual of  $V$ . Consider the function  $c_{f,v}^V : U_q(\mathfrak{g}) \rightarrow \mathbb{C}$  defined by  $c_{f,v}^V(X) := f(X(v))$ . The *coordinate ring* of  $V$  is the subspace

$$C(V) := \text{span}_{\mathbb{C}}\{c_{f,v}^V \mid v \in V, f \in V^*\} \subseteq U_q(\mathfrak{g})^*.$$

It is easy to check that  $C(V)$  is contained in  $U_q(\mathfrak{g})^\circ$ , the Hopf dual of  $U_q(\mathfrak{g})$ , and moreover that a Hopf subalgebra of  $U_q(\mathfrak{g})^\circ$  is given by

$$\mathcal{O}_q(G) := \bigoplus_{\lambda \in \mathcal{P}^+} C(V_\lambda). \quad (3)$$

We call  $\mathcal{O}_q(G)$  the *quantum coordinate algebra* of  $G$ , where  $G$  is the compact, simply-connected, simple Lie group having  $\mathfrak{g}$  as its complexified Lie algebra. Moreover, we call the decomposition of  $\mathcal{O}_q(G)$  given in (3) the *Peter–Weyl decomposition* of  $\mathcal{O}_q(G)$ .

For  $S$  a subset of simple roots, consider the Hopf subalgebra of  $U_q(\mathfrak{g})$  given by

$$U_q(\mathfrak{l}_S) := \langle K_i, E_j, F_j \mid i = 1, \dots, r; j \in S \rangle.$$

There are obvious notions of weight vectors, highest weight vectors as well as type-1 representations. It can be shown that every irreducible finite dimensional type-1  $U_q(\mathfrak{l}_S)$ -module admits a highest weight vector which is unique up to scalar multiple. Moreover, the weight of this highest weight vector determines the module up to isomorphism and there is a correspondence between irreducible type-1  $U_q(\mathfrak{l}_S)$  modules and weights in the sub-lattice

$$\mathcal{P}_S^+ + \mathcal{P}_{S^c} \subseteq \mathcal{P},$$

where  $S^c = \Pi \setminus S$  and

$$\begin{aligned} \mathcal{P}_S^+ &:= \left\{ \lambda \in \mathcal{P} : \lambda = \sum_{j \in S} \lambda_j \varpi_j, \lambda_j \in \mathbb{Z}_{\geq 0} \right\}, \\ \mathcal{P}_{S^c} &:= \left\{ \lambda \in \mathcal{P} : \lambda = \sum_{j \in S^c} \lambda_j \varpi_j, \lambda_j \in \mathbb{Z} \right\}. \end{aligned}$$

Just as for  $\mathcal{O}_q(G)$ , we can construct the type-1 dual of  $U_q(\mathfrak{l}_S)$  using matrix coefficients, this Hopf algebra will be denoted by  $\mathcal{O}_q(L_S)$ . Restriction of domains gives us a surjective Hopf  $*$ -algebra map  $\pi_S : \mathcal{O}_q(G) \rightarrow \mathcal{O}_q(L_S)$ , dual to the inclusion of  $U_q(\mathfrak{l}_S)$  in  $U_q(\mathfrak{g})$ . The *quantum flag manifold associated to  $S$*  is the quantum homogeneous space associated to  $\pi_S$

$$\begin{aligned} \mathcal{O}_q(G/L_S) &:= \mathcal{O}_q(G)^{co(\mathcal{O}_q(L_S))} \\ &= \{b \in \mathcal{O}_q(G) : b_{(1)}\langle X, b_{(2)} \rangle = \varepsilon(X)b \text{ for all } X \in U_q(\mathfrak{l}_S)\}. \end{aligned}$$

We say that the quantum flag manifold is irreducible if  $S = \Pi \setminus \{\alpha_x\}$  where  $\alpha_x$  has coefficient 1 in the expansion of the highest root of  $\mathfrak{g}$ . In the table 1 we give a diagrammatic presentation of the set of simple roots defining the irreducible quantum flag manifold, where the node corresponding to  $\alpha_x$  is the coloured one.

Table 1: Irreducible Quantum Flag Manifolds with nodes numbered according to [15, §11.4].

$A_n$		$\mathcal{O}_q(\mathrm{Gr}_{s,n+1})$	quantum Grassmannian
$B_n$		$\mathcal{O}_q(\mathbf{Q}_{2n+1})$	odd quantum quadric
$C_n$		$\mathcal{O}_q(\mathbf{L}_n)$	quantum Lagrangian Grassmannian
$D_n$		$\mathcal{O}_q(\mathbf{Q}_{2n})$	even quantum quadric
$D_n$		$\mathcal{O}_q(\mathbf{S}_n)$	quantum spinor variety
$E_6$		$\mathcal{O}_q(\mathbb{O}\mathbb{P}^2)$	quantum Cayley plane
$E_7$		$\mathcal{O}_q(\mathbf{F})$	quantum Freudenthal variety

### 2.3.1 Takeuchi's equivalence

In the following we denote  ${}_{\mathcal{O}_q(G/L_S)}^{\mathcal{O}_q(G)}\mathrm{Mod}_0$  the category whose objects are relative Hopf modules  $\mathcal{F}$  such that  $\mathcal{F}\mathcal{O}_q(G/L_S)^+ = \mathcal{O}_q(G/L_S)^+\mathcal{F}$  and morphisms are left  $\mathcal{O}_q(G)$ -comodule,  $\mathcal{O}(G/L_S)$ -bimodule maps. Let  ${}^{\mathcal{O}(L_S)}\mathrm{Mod}$  be the category whose objects are  $\mathcal{O}(L_S)$ -comodules and morphisms are  $\mathcal{O}_q(L_S)$ -comodule maps. By *Takeuchi's equivalence*, see for example [26], [13, §2.2], an equivalence of the categories  ${}_{\mathcal{O}_q(G/L_S)}^{\mathcal{O}_q(G)}\mathrm{Mod}_0$  and  ${}^{\mathcal{O}_q(L_S)}\mathrm{Mod}$  is given by the functors

$$\begin{aligned}\Phi : {}_{\mathcal{O}_q(G/L_S)}^{\mathcal{O}_q(G)}\mathrm{Mod}_0 &\rightarrow {}^{\mathcal{O}_q(L_S)}\mathrm{Mod}, & \Phi(\Gamma) &= \Gamma/\mathcal{O}_q(G/L_S)^+\Gamma, \\ \Psi : {}^{\mathcal{O}_q(L_S)}\mathrm{Mod} &\rightarrow {}_{\mathcal{O}(G/L_S)}^{\mathcal{O}_q(G)}\mathrm{Mod}_0, & \Psi(V) &= \mathcal{O}_q(G)\square_{\mathcal{O}_q(L_S)}V.\end{aligned}$$

Here for  $\Gamma \in {}_{\mathcal{O}_q(G/L_S)}^{\mathcal{O}_q(G)}\mathrm{Mod}_0$  the left  $\mathcal{O}(L_S)$ -comodule structure on  $\Gamma/\mathcal{O}_q(G/L_S)^+\Gamma$  is induced by the left  $\mathcal{O}_q(G)$ -comodule structure of  $\Gamma$ . The left  $\mathcal{O}_q(G)$ -comodule and  $\mathcal{O}(G/L_S)$ -bimodule structures on the cotensor  $\mathcal{O}_q(G)\square_{\mathcal{O}_q(L_S)}V$  are given by the coproduct and  $\mathcal{O}(G/L_S)$ -bimodule structure of  $\mathcal{O}_q(G)$ , respectively. Takeuchi's equivalence states that natural isomorphisms are given by

$$\begin{aligned}\mathbb{C} : \Phi \circ \Psi(V) &\rightarrow V, & \left[ \sum_i a^i \otimes v^i \right] &\mapsto \sum_i \varepsilon(a^i)v^i, \\ \mathbb{U} : \mathcal{F} &\rightarrow \Psi \circ \Phi(\mathcal{F}), & f &\mapsto f_{(-1)} \otimes [f_{(0)}].\end{aligned}$$

### 2.4 The Hopf algebras $\mathcal{O}_q(SO(N))$

In this subsection we give a brief description of the FRT presentation of the Hopf algebras  $\mathcal{O}_q(SO(N))$ , see [17] for details. We first give a bit of notation. We denote  $i' = N + 1 - i$ . Let  $N = 2n$  if  $N$  is even and  $N = 2n + 1$  if  $N$  is odd. We define

$$\rho_i = \frac{N}{2} - i \text{ if } i < i', \quad \rho'_i = -\rho_i \text{ if } i \leq i'.$$

Recall that the R-matrix associated to the quantum  $\mathcal{O}_q(SO(N))$  is given by

$$R_{mn}^{ij} = q^{\delta_{ij} - \delta_{ij'}} \delta_{im} \delta_{jn} + (q - q^{-1}) \theta(i - m) (\delta_{jm} \delta_{in} - \delta_{ji'} \delta_{mn'} q^{-\rho_j - \rho_m}). \quad (4)$$

Let  $U_q(\mathfrak{so}_N)^\circ$  denote the dual Hopf algebra of  $U_q(\mathfrak{so}_N)$ . Then  $\mathcal{O}_q(SO(N)) \subset U_q(\mathfrak{so}_N)^\circ$  is the Hopf subalgebra generated by matrix coefficients of the  $N$ -dimensional irreducible  $U_q(\mathfrak{so}_N)$ -representation of highest weight  $\varpi_1$ . We denote the matrix coefficients by  $u_j^i$ ,  $i, j = 1, \dots, N$ , they satisfy the relations

$$\sum_{k,l=1}^N R_{kl}^{ij} u_m^k u_n^l = \sum_{k,l=1}^N u_k^j u_l^i R_{mn}^{lk}, \quad i, j, m, n = 1, \dots, N, \quad (5)$$

$$\mathcal{D}_q = 1, \quad \sum_{i,j,k=1}^N C_j^i (C^{-1})_m^k u_i^n u_j^k = \sum_{i,j,k=1}^N C_i^n (C^{-1})_k^j u_i^j u_m^k = \delta_{mn}, \quad m, n = 1, \dots, N, \quad (6)$$

where the C-matrix coefficients and the quantum determinant are given as in [17, §9.3]. The Hopf  $*$ -algebra structure on  $\mathcal{O}_q(SO(N))$  is determined on generators by

$$\Delta(u_j^i) = \sum_k u_k^i \otimes u_j^k, \quad \varepsilon(u_j^i) = \delta_{ij}, \quad S(u_j^i) = q^{\rho_j - \rho_i} u_j^{i'}, \quad u_j^{i*} = S(u_i^j).$$

It follows from [17, §9.4] that there exists a dual pairing  $\langle \cdot, \cdot \rangle$  of the Hopf algebras  $U_q(\mathfrak{so}_N)$  and  $\mathcal{O}_q(SO(N))$  which in turn induces a left action  $U_q(\mathfrak{so}_N) \otimes \mathcal{O}_q(SO(N)) \ni X \otimes a \mapsto X \triangleright a \in \mathcal{O}_q(SO(N))$  and a right action  $\mathcal{O}_q(SO(N)) \otimes U_q(\mathfrak{so}_N) \ni a \otimes X \mapsto a \triangleleft X \in \mathcal{O}_q(SO(N))$  such that  $\mathcal{O}_q(SO(N))$  is a left and a right  $U_q(\mathfrak{so}_N)$ -module  $*$ -algebra. The actions on generators are given as follows:

$$E_j \triangleright u_j^i = u_{j+1}^i, \quad E_j \triangleright u_{(j+1)'}^i = -u_j^{i'}, \quad F_j \triangleright u_{j+1}^i = u_j^i, \quad F_j \triangleright u_{j'}^i = -u_{(j+1)'}^i, \quad j < n, \quad (7)$$

$$E_n \triangleright u_n^i = [2]_{q_2}^{1/2} u_{n+1}^i, \quad E_n \triangleright u_{n+1}^i = -q_2 [2]_{q_2}^{1/2} u_{n+2}^i, \quad (8)$$

$$F_n \triangleright u_{n+1}^i = [2]_{q_2}^{1/2} u_n^i, \quad F_n \triangleright u_{n+2}^i = -q_2^{-1} [2]_{q_2}^{1/2} u_{n+1}^i, \quad (9)$$

for  $i = 1, \dots, N$  and  $N = 2n + 1$ . For  $N = 2n$  we have

$$E_j \triangleright u_j^i = u_{j+1}^i, \quad E_j \triangleright u_{(j+1)'}^i = -u_j^{i'}, \quad F_j \triangleright u_{j+1}^i = u_j^i, \quad F_j \triangleright u_{j'}^i = u_{(j+1)'}^i, \quad i < n, \quad (10)$$

$$E_n \triangleright u_n^i = -u_{n+2}^i, \quad E_n \triangleright u_{n-1}^i = u_{n+1}^i, \quad (11)$$

$$F_n \triangleright u_{n+2}^i = -u_n^i, \quad F_n \triangleright u_{n+1}^i = u_{n-1}^i. \quad (12)$$

The corresponding right actions for  $N = 2n + 1$  are given by

$$u_j^i \triangleleft F_i = u_j^{i+1}, \quad u_j^{(i+1)'} \triangleleft F_i = -u_j^{i'}, \quad u_j^{i+1} \triangleleft E_i = u_j^i, \quad u_j^{i'} \triangleleft E_i = -u_j^{(i+1)'}, \quad i < n, \quad (13)$$

$$u_j^n \triangleleft F_n = [2]_{q_2}^{1/2} u_j^{n+1}, \quad u_j^{n+1} \triangleleft F_n = -q_2 [2]_{q_2}^{1/2} u_j^{n+2}, \quad (14)$$

$$u_j^{n+1} \triangleleft E_n = [2]_{q_2}^{1/2} u_j^n, \quad u_j^{n+2} \triangleleft E_n = -q_2^{-1} [2]_{q_2}^{1/2} u_j^{n+1}, \quad (15)$$

and for even  $N = 2n$

$$u_j^i \triangleleft F_i = u_j^{i+1}, \quad u_j^{(i+1)'} \triangleleft F_i = -u_j^{i'}, \quad u_j^{i+1} \triangleleft E_i = u_j^i, \quad u_j^{i'} \triangleleft E_i = u_j^{(i+1)'}, \quad i < n, \quad (16)$$

$$u_j^n \triangleleft F_n = -u_j^{n+2}, \quad u_j^{n-1} \triangleleft F_n = u_j^{n+1}, \quad (17)$$

$$u_j^{n+2} \triangleleft E_n = -u_j^n, \quad u_j^{n+1} \triangleleft E_n = u_j^{n-1}, \quad (18)$$



for  $i = 1, \dots, N$ . The values  $E_j \triangleright u_l^k$ ,  $F_j \triangleright u_l^k$ ,  $u_l^k \triangleleft E_j$  and  $u_l^k \triangleleft F_j$  for all other cases are zero.

For the convenience of the reader, the next lemma collects the commutation relations of the generators of  $\mathcal{O}_q(SO(N))$  which will be used in this paper.

**Lemma 2.7.** *If  $u_j^i, i, j = 1, \dots, N$  are the generators of  $\mathcal{O}_q(SO(N))$  then the following commutation relations hold*

$$u_1^i u_N^i = q^2 u_N^i u_1^i, \quad i \neq i', \quad (19)$$

$$u_l^i u_k^i = q u_k^i u_l^i, \quad l < k, \quad i \neq i', \quad (20)$$

$$u_l^j u_k^i = u_k^i u_l^j, \quad l < k, \quad i < j, \quad l \neq k', \quad i \neq j', \quad (21)$$

$$u_1^j u_N^i = q u_N^i u_1^j, \quad i < j, \quad i \neq j', \quad (22)$$

$$u_1^i u_N^j = q u_N^j u_1^i + (q^2 - 1) u_N^i u_1^j, \quad i < j, \quad i \neq j', \quad (23)$$

$$u_k^i u_l^j = u_l^j u_k^i - (q - q^{-1}) u_k^j u_l^i, \quad i < j, \quad k < l, \quad i \neq j', \quad k \neq l', \quad (24)$$

$$u_k^i u_k^j = q u_k^j u_k^i, \quad k \neq k', \quad i < j, \quad i \neq j', \quad (25)$$

$$(u_1^1 u_a^2 - q u_a^2 u_1^1)(u_1^1 u_N^2 - q u_1^2 u_N^1) = q^2 (u_1^1 u_N^2 - q u_1^2 u_N^1)(u_1^1 u_a^2 - q u_1^2 u_a^1), \quad (26)$$

$$(u_1^1 u_N^2 - q u_1^2 u_N^1)(u_a^1 u_N^2 - q u_a^2 u_N^1) = q^2 (u_a^1 u_N^2 - q u_a^2 u_N^1)(u_1^1 u_N^2 - q u_1^2 u_N^1), \quad (27)$$

where  $a = 2, \dots, N - 1$ .

*Proof.* Equations (19)-(24) follow directly from the R-matrix relations (5). For instance, if we take  $m = N$ ,  $n = 1$  and  $i > j$  with  $i \neq j'$  in relation (5) we obtain

$$u_N^i u_1^j + (q - q^{-1}) u_N^j u_1^i = q^{-1} u_1^j u_N^i,$$

which yields equation (23). Equation (25) follows by choosing  $k = l + 1$  in (21) and acting on both sides by the  $E_i$ 's or  $F_i$ 's from the left by using formulas (7)-(12). We only give the proof of (27) since the proof of (26) is analogous. By using (19)-(25) it can be show that  $u_N^1 (u_1^1 u_N^2 - q u_1^2 u_N^1) = q^{-1} (u_1^1 u_N^2 - q u_1^2 u_N^1) u_N^1$ . If we act on both sides by  $E_1$  from the left we get

$$u_N^1 (u_2^1 u_N^2 - q u_2^2 u_N^1) = q^{-1} (u_2^1 u_N^2 - q u_2^2 u_N^1) u_N^1.$$

Then the relation  $u_N^1 (u_a^1 u_N^2 - q u_a^2 u_N^1) = q^{-1} (u_a^1 u_N^2 - q u_a^2 u_N^1) u_N^1$  can be obtained by acting from the left by the  $E_i$ 's according to identities (7)-(12). The commutation relations

$$\begin{aligned} u_1^1 (u_a^1 u_N^2 - q u_a^2 u_N^1) &= q^2 (u_a^1 u_N^2 - q u_a^2 u_N^1) u_1^1, & u_N^2 (u_a^1 u_N^2 - q u_a^2 u_N^1) &= (u_a^1 u_N^2 - q u_a^2 u_N^1) u_N^2, \\ u_1^2 (u_a^1 u_N^2 - q u_a^2 u_N^1) &= (u_a^1 u_N^2 - q u_a^2 u_N^1) u_1^2 \end{aligned}$$

can be easily deduced from (19)-(25). Combining this results we have that (27) holds for  $a = 2, \dots, N - 1$ .  $\square$

### 3 The Heckenberger–Kolb calculi and the Kähler structure

In this section we recall important facts about Heckenberger–Kolb calculi for irreducible quantum flag manifolds as well as the existence of a Kähler structure for them. For more details we refer

the reader to the seminal papers [13] and [14] for the Heckenberger–Kolb calculus and [21], [24] for Kähler structures.

As mentioned before, the irreducible quantum flag manifolds are distinguished by existence of a unique  $q$ -deformed de Rham complex. The following theorem collects the main results of [12], [13], and [21].

**Theorem 3.1.** *For any irreducible quantum flag manifold  $\mathcal{O}_q(G/L_S)$ , there exists a unique finite dimensional left  $\mathcal{O}(G)$ -covariant differential  $*$ -calculus*

$$\Omega_q^\bullet(G/L_S) \in {}_{\mathcal{O}_q(G/L_S)}^{\mathcal{O}_q(G)}\text{Mod}_0,$$

of classical dimension

$$\dim \Phi(\Omega_q^k(G/L_S)) = \binom{2M}{k}, \quad \text{for all } k = 1, \dots, 2M,$$

where  $M$  is the complex dimension of the corresponding classical manifold. Moreover

(i)  $\Omega_q^\bullet(G/L_S)$  admits a unique left  $\mathcal{O}_q(G)$ -covariant complex structure

$$\Omega_q^\bullet(G/L_S) \simeq \bigoplus_{(a,b) \in \mathbb{N}_0} \Omega^{(a,b)} =: \Omega^{(\bullet, \bullet)},$$

(ii)  $\Omega^{(1,0)}$  and  $\Omega^{(0,1)}$  are irreducible objects in  ${}_{\mathcal{O}_q(G/L_S)}^{\mathcal{O}_q(G)}\text{Mod}_0$ .

We recall that the  $*$ -calculus and the covariant complex structure were given in [21]. The following theorem gives us the existence of the Kähler structure for the Heckenberger–Kolb calculus.

**Theorem 3.2.** ([21] Theorem 5.10). *Let  $\Omega_q^\bullet(G/L_S)$  be the Heckenberger–Kolb calculus of the irreducible quantum flag manifold  $\mathcal{O}_q(G/L_S)$ . Then there exists a form  $\kappa \in \Omega^{(1,1)}$  such that  $(\Omega^{(\bullet, \bullet)}, \kappa)$  is a covariant Kähler structure for all  $q \in \mathbb{R}_{>0} \setminus F$ , where  $F$  is a finite, possibly empty, subset of  $\mathbb{R}_{>0}$ . Moreover, any element of  $F$  is necessarily non-transcendental.*

It has been shown in [9, §5.7] that if  $\mathcal{O}_q(G)$  is a compact quantum group algebra and  $q$  belongs to a suitable open interval around 1 then an inner product on  $\Omega^{(\bullet, \bullet)}$  can be defined by

$$\langle \cdot, \cdot \rangle : \Omega^{(\bullet, \bullet)} \times \Omega^{(\bullet, \bullet)} \rightarrow \mathbb{C}, \quad (\omega, \nu) \mapsto \mathbf{h} \circ g_\sigma(\omega, \nu), \quad (28)$$

where  $\mathbf{h}$  is the Haar state associated to  $\mathcal{O}_q(G)$  [17, §11.3.2]. From now on we consider  $q \in (1, \epsilon)$  so that the inner product is well defined. With respect to this inner product we consider the adjoints  $d^\dagger$ ,  $\partial^\dagger$  and  $\bar{\partial}^\dagger$ . It is shown in [24, §5.4] that

$$d^\dagger = - *_\kappa \circ d \circ *_\kappa, \quad \partial^\dagger = - *_\kappa \circ \bar{\partial} \circ *_\kappa, \quad \bar{\partial}^\dagger = - *_\kappa \circ \partial \circ *_\kappa. \quad (29)$$

Then Theorem 3.1 and the inner product (28) allow us to define the  $\bar{\partial}$ -Dirac operator and the  $\bar{\partial}$ -Laplacian

$$D_{\bar{\partial}} := \bar{\partial} + \bar{\partial}^\dagger, \quad \Delta_{\bar{\partial}} := (\bar{\partial} + \bar{\partial}^\dagger)^2.$$

**Lemma 3.3.** *With respect to the notation in [13] we have*

- (i)  $u_j^i = c_{f_i, v_j}^{\varpi_1}$ ,  $S(u_j^i) = c_{v_j, f_i}^{-w_0(\varpi_1)}$  and  $z_{ij} := c_{f_i, v_N}^{\varpi_1} c_{v_j, f_N}^{-w_0 \varpi_1} = u_N^i S(u_j^N)$  are the generators of the algebra  $\mathcal{O}_q(G/L_S)$ ,
- (ii)  $\Phi(\Omega^{(0,1)}) = \Omega^{(0,1)}/\mathcal{O}_q(\mathbf{Q}_N)^+ \Omega^{(0,1)} = \text{Lin}_{\mathbb{C}}\{[\bar{\partial}z_{Ni}] : i = 2, \dots, N-1\}$  and  $\Phi(\Omega^{(1,0)}) = \text{Lin}_{\mathbb{C}}\{[\partial z_{iN}] : i = 2, \dots, N-1\}$
- (iii) if we consider  $\mathcal{O}_q(G/L_S)$  as a left  $U_q(\mathfrak{g})$ -module with action  $X \blacktriangleright z := z \triangleleft S(X)$  then the generators of  $\mathcal{O}_q(\mathbf{Q}_N)$  are given by  $z := u_1^1 u_N^1$ ,  $y := u_1^1 u_N^2 - q u_1^2 u_N^1$  with weights  $2\varpi_1$ ,  $\varpi_2$  respectively.

*Proof.* (i) follows from the notation in [13]. To prove (ii) we know from [13, §3.2] that  $I_{(1)} = \{i \in I : (\varpi_1, \varpi_1 - \alpha_1 - \text{wt}(v_i)) = 0\}$ . Since  $v_N$  is the highest weight vector  $(\varpi_1, \varpi_1 - \alpha_1 - \text{wt}(v_N)) = (\varpi_1, \varpi_1 - \alpha_1 - \varpi_1) = -(\varpi_1, \alpha_1) \neq 0$ , thus  $v_N \notin I_{(1)}$ . Similarly, we can consider  $v_1$  as the lowest weight vector, then  $\text{wt}(v_1) = w_0(\varpi_1) = -\text{id}(\varpi_1) = -\varpi_1$ . Then  $(\varpi_1, \varpi_1 - \alpha_1 - \text{wt}(v_1)) = (\varpi_1, \varpi_1 - \alpha_1 + \varpi_1) = 2(\varpi_1, \varpi_1) - (\varpi_1, \alpha_1) = (2r_1 - 1)(\varpi_1, \alpha_1)$  where  $r_1$  is the coefficient of  $\alpha_1$  in the expression of  $\varpi_1$  as linear combination of the  $\alpha_1, \dots, \alpha_n$ . From the form of the Cartan matrix of type  $B_n$  and  $D_n$  it can be shown that  $r_1 = 1$  and therefore  $v_1 \notin I_{(1)}$ . It also can be shown that  $M = N - 2$  for both cases  $\mathcal{O}_q(\mathbf{Q}_{2n+1})$  and  $\mathcal{O}_q(\mathbf{Q}_{2n})$ , see for example table 2 in [9]. Therefore  $I_{(1)} = \{2, \dots, N-1\}$ .

For the proof of (iii) we first note that the left and right actions on matrix coefficients are given by

$$X \triangleright c_{f_i, v_j}^{\varpi_1} \triangleleft Y = c_{f_i \triangleleft Y, X \triangleright v_j}^{\varpi_1}. \quad (30)$$

By (30) we have  $E_i \blacktriangleright (u_1^1 u_N^1) = (u_1^1 u_N^1) \triangleleft S(E_i) = -(c_{f_1 \triangleleft E_i, v_1}^{\varpi_1} c_{f_1 \triangleleft K_i, v_1}^{\varpi_1} + c_{f_1, v_1}^{\varpi_1} c_{f_1 \triangleleft E_i, v_N}^{\varpi_1}) \triangleleft K_i^{-1} = 0$  since  $f_1$  is the highest weight of the dual representation. On the other hand

$$K_i \blacktriangleright (u_1^1 u_N^1) = (u_1^1 u_N^1) \triangleleft K_i^{-1} = c_{f_1 \triangleleft K_i, v_j}^{\varpi_1} c_{f_1 \triangleleft K_i, v_N}^{\varpi_1} = q^{(2\varpi_1, \alpha_i)} u_1^1 u_N^1.$$

Similarly, since  $f_1$  is a highest weight vector we have for  $i \neq 1$

$$(u_1^1 u_N^2 - q u_1^2 u_N^1) \triangleleft E_i = c_{f_1, v_1}^{\varpi_1} c_{f_2 \triangleleft E_i, v_N}^{\varpi_1} - q c_{f_2 \triangleleft E_i, v_1}^{\varpi_1} c_{f_1 \triangleleft K_i, v_N}^{\varpi_1}. \quad (31)$$

However, since  $v_1$  is the lowest weight vector of  $V_{\varpi_1}$  with weight  $w_0 \varpi_1 = -\varpi_1$  then for  $i \neq 1$  the vector  $v_1$  is the lowest weight with weight zero of the  $U_q(\mathfrak{sl}_2)_i$ -module  $U_q(\mathfrak{sl}_2)_i v_N$  where  $U_q(\mathfrak{sl}_2)_i := \langle E_i, F_i, K_i \rangle$ . Therefore  $E_i v_1 = 0$  for  $i \neq 1$  and  $E_1 v_1 = v_2$ . This in turn implies that  $f_2 \triangleleft E_i = 0$  for  $i \neq 1$ . Then the right hand side of equation (31) is zero. Since  $f_2 \triangleleft E_1 = f_1$  and  $f_1 \triangleleft K_1 = f_1$  we have

$$\begin{aligned} E_1 \blacktriangleright (u_1^1 u_N^2 - q u_1^2 u_N^1) &= -(u_1^1 u_N^2 - q u_1^2 u_N^1) \triangleleft E_1 K_1^{-1} \\ &= -(c_{f_1, v_1}^{\varpi_1} c_{f_2 \triangleleft E_1, v_N}^{\varpi_1} - q c_{f_2 \triangleleft E_1, v_1}^{\varpi_1} c_{f_1 \triangleleft K_1, v_N}^{\varpi_1}) \triangleleft K_i^{-1} \\ &= 0. \end{aligned}$$

Finally, since  $K_i E_1 v_1 = q^{(\alpha_i, \alpha_1)} E_1 K_i v_1 = q^{(\alpha_i, \alpha_1) - (\varpi_1, \alpha_i)} E_1 v_1$  we have

$$\begin{aligned} K_i \blacktriangleright (u_1^1 u_N^2 - q u_1^2 u_N^1) &= (u_1^1 u_N^2 - q u_1^2 u_N^1) \triangleleft K_i^{-1} \\ &= c_{f_1 \triangleleft K_i^{-1}, v_1}^{\varpi_1} c_{f_2 \triangleleft K_i^{-1}, v_N}^{\varpi_1} - q c_{f_2 \triangleleft K_i^{-1}, v_1}^{\varpi_1} c_{f_1 \triangleleft K_i^{-1}, v_N}^{\varpi_1} \\ &= q^{(2\varpi_1 - \alpha_1, \alpha_i)} (c_{f_1, v_1}^{\varpi_1} c_{f_2, v_N}^{\varpi_1} - q c_{f_2, v_1}^{\varpi_1} c_{f_1, v_N}^{\varpi_1}) \\ &= q^{(\varpi_2, \alpha_i)} (u_1^1 u_N^2 - q u_1^2 u_N^1), \end{aligned}$$

where we used the fact that  $\alpha_i = \sum_j a_{ji} \varpi_j$  to derive the third equality.  $\square$

Part (iii) of the Lemma 3.3 says that the weights  $2\varpi_1, \varpi_2$  form a minimal set of generators for the weights of the highest weights vectors of  $\mathcal{O}_q(Q_N)$  as a monoid under addition which are called *spherical weights*. In table 2 below we present the spherical weights for each irreducible quantum flag manifold which was presented by Krämer [19] in the classical case. We note that for the case of  $\mathcal{O}_q(\mathbf{S}_{2n})$ , the weight  $2\varpi_{2n-1}$  or  $2\varpi_{2n}$  appears according to the defining crossed node, as presented in table 1.

Table 2: Spherical weights according to the numbering of the Dynkin nodes in [15, §11.4].

$\mathcal{O}_q(\mathrm{Gr}_{r,s})$	$\varpi_1 + \varpi_{r+s-1}, \varpi_2 + \varpi_{r+s-2}, \dots, \varpi_r + \varpi_s$
$\mathcal{O}_q(\mathbf{Q}_{2n+1})$	$2\varpi_1, \varpi_2$
$\mathcal{O}_q(\mathbf{L}_n)$	$2\varpi_1, 2\varpi_2, \dots, 2\varpi_n$
$\mathcal{O}_q(\mathbf{Q}_{2n})$	$2\varpi_1, \varpi_2$
$\mathcal{O}_q(\mathbf{S}_{2n})$	$\varpi_2, \varpi_4, \dots, \varpi_{2n-2}, 2\varpi_{2n-1}$ or $2\varpi_{2n}$
$\mathcal{O}_q(\mathbf{S}_{2n+1})$	$\varpi_2, \varpi_4, \dots, \varpi_{2n-2}, \varpi_{2n} + 2\varpi_{2n+1}$
$\mathcal{O}_q(\mathbb{O}\mathbb{P}^2)$	$\varpi_1 + \varpi_6, \varpi_2$
$\mathcal{O}_q(\mathbf{F})$	$2\varpi_7, \varpi_1, \varpi_6$

**Lemma 3.4.** *Let  $\partial, \bar{\partial}, z, y$  be as before. Then the following holds*

- (i)  $\bar{\partial}zz = q^2 z \bar{\partial}z$ ,
- (ii)  $\bar{\partial}yy = q^2 y \bar{\partial}y$ ,
- (iii)  $\partial zz = q^{-2} z \partial z$ ,
- (iv)  $\partial yy = q^{-2} y \partial y$ .

*Proof.* We only show (iii) and (iv), equations (i) and (ii) can be shown in a similar way by using Lemma 2.7 and Lemma 3.3. By Takeuchi's equivalence there exists an isomorphism in the category  ${}_{\mathcal{O}_q(G/L_S)}^{\mathcal{O}_q(G)} \mathrm{Mod}_0$  given by

$$U : \Omega^{(0,1)} \rightarrow \mathcal{O}(SO_q(5)) \square_{\mathcal{O}_q(L)} \Phi(\Omega^{(0,1)}), \quad \omega \mapsto \omega_{(-1)} \otimes [\omega_{(0)}].$$

Now we recall by Lemma 3.3 that  $\{[\bar{\partial}z_{Nj}] : j = 1, \dots, N-1\}$  and  $\{[\partial z_{jN}], j = 2, \dots, N-1\}$  are bases of the vector spaces  $\Phi(\Omega^{(0,1)})$  and  $\Phi(\Omega^{(1,0)})$ , respectively. Moreover, it was shown in [13, §3.2] that  $[\bar{\partial}z_{ij}] = [\partial z_{ji}] = 0$  if  $i \neq N$  and  $j \in \{1, N\}$ .

Then by definition of  $U$  and relations in Lemma 2.7 we have

$$\begin{aligned}
U((\bar{\partial}z_{1N})z_{1N}) &= U(\bar{\partial}z_{1N})z_{1N} \\
&= (z_{1N})_{(1)}z_{1N} \otimes [\bar{\partial}(z_{1N})_{(2)}] \\
&= (u_N^1 S(u_N^N))_{(1)}z_{1N} \otimes [\bar{\partial}(u_N^1 S(u_N^N))_{(2)}] \\
&= \sum_{j,k} u_j^1 S(u_N^k)z_{1N} \otimes [\bar{\partial}(u_N^j S(u_N^k))] \\
&= \sum_{j,k} u_j^1 S(u_N^k)z_{1N} \otimes [\bar{\partial}z_{jk}] \\
&= \sum_{k=2}^{N-1} u_N^1 S(u_N^k)z_{1N} \otimes [\bar{\partial}z_{Nk}] \\
&= \sum_{k=2}^{N-1} u_N^1 S(u_N^k)u_N^1 S(u_N^N) \otimes [\bar{\partial}z_{Nk}] \\
&= \sum_{k=2}^{N-1} q^{\rho_N - \rho_k} u_N^1 u_{k'}^1 u_N^1 u_1^1 \otimes [\bar{\partial}z_{Nk}] \\
&= q^{-2} \sum_{k=2}^{N-1} q^{\rho_N - \rho_k} u_N^1 u_1^1 u_N^1 u_{k'}^1 \otimes [\bar{\partial}z_{Nk}] \\
&= q^{-2} z_{1N} U(\bar{\partial}z_{1N}).
\end{aligned}$$

This yields equation (iii).

Since  $y = q(z_{2,N} - q^{-2}z_{1,N-1})$  then analogously by using relations of Lemma 2.7 we have

$$\begin{aligned}
U((\bar{\partial}y)y) &= U(\bar{\partial}y)y \\
&= U(\bar{\partial}(qz_{2,N} - q^{-1}z_{1,N-1}))(qz_{2,N} - q^{-1}z_{1,N-1}) \\
&= (qz_{2,N} - q^{-1}z_{1,N-1})_{(1)}(qz_{2,N} - q^{-1}z_{1,N-1}) \otimes [\bar{\partial}(qz_{2,N} - q^{-1}z_{1,N-1})_{(2)}] \\
&= \sum_{l=2}^{N-1} (qu_N^2 S(u_N^l) - q^{-1}u_N^1 S(u_{N-1}^l))(u_1^1 u_N^2 - qu_1^2 u_N^1) \otimes [\bar{\partial}(qz_{Nl} - q^{-1}z_{Nl})] \\
&= \sum_{l=2}^{N-1} q^{\rho_N - \rho_l} (qu_N^2 u_{l'}^1 - u_N^1 u_{l'}^2)(u_1^1 u_N^2 - qu_1^2 u_N^1) \otimes [\bar{\partial}(qz_{Nl} - q^{-1}z_{Nl})] \\
&= \sum_{l=2}^{N-1} q^{\rho_N - \rho_l + 1} (u_{l'}^1 u_N^2 - qu_{l'}^2 u_N^1)(u_1^1 u_N^2 - qu_1^2 u_N^1) \otimes [\bar{\partial}(qz_{Nl} - q^{-1}z_{Nl})] \\
&= q^{-2} \sum_{l=2}^{N-1} q^{\rho_N - \rho_l + 1} (u_1^1 u_N^2 - qu_1^2 u_N^1)(u_{l'}^1 u_N^2 - qu_{l'}^2 u_N^1) \otimes [\bar{\partial}(qz_{Nl} - q^{-1}z_{Nl})] \\
&= q^{-2} (u_1^1 u_N^2 - qu_1^2 u_N^1) \sum_{l=2}^{N-1} q^{\rho_N - \rho_l + 1} (u_{l'}^1 u_N^2 - qu_{l'}^2 u_N^1) \otimes [\bar{\partial}(qz_{Nl} - q^{-1}z_{Nl})] \\
&= q^{-2} y U(\bar{\partial}y).
\end{aligned}$$

This proves equation (iv). □

In the following lemma we give the existence of a Kähler structure for the Heckenberger–Kolb calculus over the quantum quadric  $\mathcal{O}_q(\mathbf{Q}_N)$ . Since the space of coinvariants  ${}^{co(L_S)}\Phi(\Omega^{(1,1)})$  is a 1-dimensional vector space (see for example [11, §3.3]) it can be shown that this Kähler form is up to scalar equal to the Kähler form given in [21, §5].

**Lemma 3.5.** *A Kähler form is given by*

$$\kappa = i \sum_{i,j=1}^N q^{-\rho_i} \partial z_{ij} \wedge \bar{\partial} z_{ji}.$$

For  $i = 1, \dots, N-2$  we denote  $e_i^+ := [\partial z_{i+1,N}]$  and  $e_i^- := q^{-\rho_{i+1}} [\bar{\partial} z_{N,i+1}]$ , then

$$U(\kappa) = i \sum_{i=1}^{N-2} 1 \otimes e_i^+ \wedge e_i^- \quad (32)$$

and there exist polynomials  $f_{I,J}(q) \in \mathbb{C}[q, q^{-1}]$  such that

$$U(\kappa^l) = i^l \sum_{I,J \in \Theta(l)} f_{I,I,J}(q) \cdot 1 \otimes e_I^+ \wedge e_J^-, \quad f_{I,I,I}(1) \neq 0, \quad \text{and} \quad f_{I,I,J}(1) = 0 \quad \text{for} \quad I \neq J, \quad (33)$$

where  $\Theta(l)$  denotes the set of all ordered subsets of  $\{1, \dots, N-2\}$  with  $l$  elements. Moreover,  $\text{sgn}(f_{I,I,I}(1)) = (-1)^{\frac{l(l-1)}{2}}$  for all  $I \in \Theta(l)$ .

*Proof.* That  $\kappa$  is real follows from identities (2) and the fact that  $z_{ij}^* = z_{ji}$ . Equation (32) follows from the calculation

$$\begin{aligned} U\left(\sum_{i,j=1}^N q^{-2\rho_i} \partial z_{ij} \wedge \bar{\partial} z_{ji}\right) &= \sum_{i,j=1}^N q^{-2\rho_i} (z_{ij})_{(1)} (z_{ji})_{(1)} \otimes [\partial((z_{ij})_{(2)})] \wedge [\bar{\partial}((z_{ji})_{(2)})] \\ &= \sum_{i,j=1}^N \sum_{a,b,c,d} q^{-2\rho_i} u_a^i S(u_j^b) u_c^j S(u_i^d) \otimes [\partial(u_N^a S(u_b^N))] \wedge [\bar{\partial}(u_N^c S(u_d^N))] \\ &= \sum_{i=1}^N \sum_{a,b,d} q^{-2\rho_i} u_a^i S(u_i^d) \otimes [\partial(u_N^a S(u_b^N))] \wedge [\bar{\partial}(u_N^b S(u_d^N))] \\ &= \sum_{i=1}^N \sum_{a,d} q^{-2\rho_i} u_a^i S(u_i^d) \otimes [\partial z_{aN}] \wedge [\bar{\partial} z_{Nd}] \\ &= \sum_{d=2}^{N-1} 1 \otimes [\partial z_{dN}] \wedge q^{-\rho_d} [\bar{\partial} z_{Nd}], \end{aligned}$$

where the identity  $\sum_{i=1}^N q^{-2\rho_i} u_a^i S(u_i^d) = q^{-2\rho_d} \delta_{da} \cdot 1$  was used in the penultimate line. This implies the left  $\mathcal{O}_q(G)$ -coinvariance of  $\kappa$  since

$$\left[ \sum_{i,j=1}^N q^{-2\rho_i} \partial z_{ij} \wedge \bar{\partial} z_{ji} \right] = \sum_{i,j=1}^N q^{-2\rho_i} [\partial z_{ij}] \wedge [\bar{\partial} z_{ji}] = \sum_{i=2}^{N-1} [\partial z_{iN}] \wedge q^{-2\rho_i} [\bar{\partial} z_{Ni}].$$

Now we show that  $\kappa$  is closed by a Lie theoretic argument: we set  $V^{(a,b)} := \Phi(\Omega^{(a,b)})$  and note that  $d\kappa = 0$  if and only if  $\partial\kappa = \bar{\partial}\kappa = 0$ , and both are coinvariant elements in  $\Omega^{(2,1)}$  and  $\Omega^{(1,2)}$ , respectively. But  ${}^{co(L_S)}V^{(1,2)} = {}^{co(L_S)}(V^{(1,0)} \otimes V^{(0,2)})$ , then  $\bar{\partial}\kappa = 0$  if  ${}^{co(L_S)}(V^{(1,0)} \otimes V^{(0,2)}) = 0$ . Since  $V^{(1,0)} \otimes W$  contains the trivial representation if and only if  $W \simeq (V^{(1,0)})^* \simeq V^{(0,1)}$  it is enough to show that  $V^{(0,2)}$  is not isomorphic to  $V^{(1,0)}$ . But  $V^{(0,2)}$  is an irreducible component of  $V^{(0,1)} \otimes V^{(0,1)}$ . The following formulas for the tensor product decomposition can be found in Table 5 of [27]: for odd  $N = 2n + 1$  we have

$$V^{(0,1)} \otimes V^{(0,1)} \simeq \begin{cases} V_{4\varpi_1} \oplus V_{2\varpi_1} \oplus \mathbb{C}, & n = 2 \quad (V^{(0,1)} \simeq V_{2\varpi_1} \text{ as } U_q(\mathfrak{sl}_2)\text{-modules}), \\ V_{2\varpi_1} \oplus V_{2\varpi_2} \oplus \mathbb{C}, & n = 3 \quad (V^{(0,1)} \simeq V_{\varpi_1} \text{ as } U_q(\mathfrak{so}_5)\text{-modules}), \\ V_{2\varpi_1} \oplus V_{\varpi_2} \oplus \mathbb{C}, & n \geq 4 \quad (V^{(0,1)} \simeq V_{\varpi_1} \text{ as } U_q(\mathfrak{so}_{2n-1})\text{-modules}). \end{cases}$$

For  $n = 2$ , it can be shown that the  $U_q(\mathfrak{sl}_2)$ -invariant element corresponding to  $\mathbb{C}$  is not  $U_q(\mathfrak{sl}_2)$ -invariant. For even  $N = 2n$  we have

$$V^{(0,1)} \otimes V^{(0,1)} \simeq \begin{cases} V_{2\varpi_2} \oplus V_{\varpi_3 + \varpi_1} \oplus \mathbb{C}, & n = 4 \quad (V^{(0,1)} \simeq V_{\varpi_2} \text{ as } U_q(\mathfrak{sl}_4)\text{-modules}), \\ V_{2\varpi_1} \oplus V_{\varpi_2} \oplus \mathbb{C}, & n \geq 5 \quad (V^{(0,1)} \simeq V_{\varpi_1} \text{ as } U_q(\mathfrak{so}_{2n-2})\text{-modules}). \end{cases}$$

This formulas show that  $V^{(1,0)}$  does not appear as an irreducible component of  $V^{(0,1)} \otimes V^{(0,1)}$  and therefore  $\bar{\partial}\kappa = 0$ . A similar argument shows that  $\partial\kappa = 0$ . By Lemma 4.6 in [24] we have that  $\kappa$  is central. The bijectivity of the Lefschetz operators  $L^{n-k}$  is shown in [21] for  $q$  belonging to a sufficiently small open interval around 1.

Now we proceed to calculate the powers of  $\kappa$  by using the relations in  $\Phi(\Omega^{(\bullet,0)})$  and  $\Phi(\Omega^{(0,\bullet)})$ . It follows from [13] that  $\Phi(\Omega^{(0,1)}) = \text{Lin}_{\mathbb{C}}\{e_1^-, \dots, e_{N-2}^-\}$  and the algebra  $\Phi(\Omega^{(0,\bullet)})$  is isomorphic to the quantum exterior algebra  $\Lambda(\mathcal{O}_q^{N-2})$  of the quantum Euclidean space  $\mathcal{O}_q^{N-2}$  in [17, §9.3]. Then the relations in [13] imply that the relations in  $\Phi(\Omega^{(0,\bullet)})$  for  $N = 2n + 1$  are given by

$$e_i^- \wedge e_{i'}^- = 0, \quad i \neq i', \quad e_i^- \wedge e_j^- = -q^{-1}e_j^- \wedge e_i^-, \quad i < j, \quad i \neq j', \quad (34)$$

$$e_{i'}^- \wedge e_i^- + e_i^- \wedge e_{i'}^- = (q - q^{-1}) \sum_{1 \leq j < i} \lambda_i^{-1} \lambda_j q^{j-i+1} e_j^- \wedge e_{i'}^-, \quad i < i', \quad (35)$$

$$e_n^- \wedge e_n^- = (q^{1/2} - q^{-1/2}) \sum_{1 \leq j \leq n-1} \lambda_n^{-1} \lambda_j q^{j-(n-1)} e_j^- \wedge e_{j'}^-, \quad (36)$$

where  $\lambda_i$  is a non-zero complex number. For even  $N = 2n$  the relations are given by (34) and (35) for possibly different constants  $\lambda_i$ . The relations for  $\Phi(\Omega^{(\bullet,0)})$  are giving by applying the involution to the relations (34)-(36). If we apply the involution to (35) we get

$$e_i^+ \wedge e_{i'}^+ + e_{i'}^+ \wedge e_i^+ = (q - q^{-1}) \sum_{1 \leq j < i} \bar{\lambda}_i^{-1} \bar{\lambda}_j q^{j-i+1} e_{j'}^+ \wedge e_j^+, \quad i < i'. \quad (37)$$

We claim that the relations in  $\Phi(\Omega^{(\bullet,0)})$  are given by

$$e_i^+ \wedge e_{i'}^+ = 0, \quad i \neq i', \quad e_i^+ \wedge e_j^+ = -qe_j^+ \wedge e_i^+, \quad i < j, \quad i \neq j', \quad (38)$$

$$e_{i'}^+ \wedge e_i^+ + e_i^+ \wedge e_{i'}^+ = -(q - q^{-1}) \sum_{1 \leq j < i} \bar{\lambda}_i^{-1} \bar{\lambda}_j q^{-(j-i+1)} e_j^+ \wedge e_{i'}^+, \quad i < i', \quad (39)$$

$$e_n^+ \wedge e_n^+ = -(q^{1/2} - q^{-1/2}) \sum_{1 \leq j \leq n-1} \bar{\lambda}_n^{-1} \bar{\lambda}_j q^{-(j-(n-1))} e_j^+ \wedge e_{j'}^+. \quad (40)$$

Relations (38) are shown by applying the involution to relations (34). For the rest of the paper we denote  $\nu_\tau := \tau - \tau^{-1}$  if  $\tau > 0$ . We use induction to prove (39):

$$\begin{aligned}
\bar{\lambda}_{i+1}(e_{i+1}^+ \wedge e_{(i+1)'}^+ + e_{(i+1)'}^+ \wedge e_{i+1}^+) &= \nu_q \sum_{1 \leq j < i+1} \bar{\lambda}_j q^{j-(i+1)+1} e_{j'}^+ \wedge e_j^+ && \text{by (37)} \\
&= q^{-i} \nu_q \sum_{1 \leq j < i+1} \bar{\lambda}_j q^j (-\nu_q \sum_{1 \leq k < j} \bar{\lambda}_j^{-1} \bar{\lambda}_k q^{-k+j-1} e_k^+ \wedge e_{k'}^+ - e_j^+ \wedge e_{j'}^+) && \text{(ind. hyp.)} \\
&= q^{-i} \nu_q \left( -\nu_q \sum_{1 \leq j < i+1} \sum_{1 \leq k < j} \bar{\lambda}_k q^{2j-k-1} e_k^+ \wedge e_{k'}^+ - \sum_{1 \leq j < i+1} \bar{\lambda}_j q^j e_j^+ \wedge e_{j'}^+ \right) \\
&= q^{-i} \nu_q \left( -\nu_q \sum_{1 \leq k < i} \sum_{k+1 \leq j \leq i} \bar{\lambda}_k q^{2j-k-1} e_k^+ \wedge e_{k'}^+ - \sum_{1 \leq j < i+1} \bar{\lambda}_j q^j e_j^+ \wedge e_{j'}^+ \right) \\
&= q^{-i} \nu_q \left( -\nu_q \sum_{1 \leq k < i} \bar{\lambda}_k q^{-k-1} \left( \frac{q^{2(k+1)} - q^{2i+2}}{1 - q^2} \right) e_k^+ \wedge e_{k'}^+ - \sum_{1 \leq j < i+1} \bar{\lambda}_j q^j e_j^+ \wedge e_{j'}^+ \right) \\
&= q^{-i} \nu_q \left( \sum_{1 \leq k < i} \bar{\lambda}_k (q^k - q^{2i-k}) e_k^+ \wedge e_{k'}^+ - \sum_{1 \leq j < i+1} \bar{\lambda}_j q^j e_j^+ \wedge e_{j'}^+ \right) \\
&= q^{-i} \nu_q \left( \sum_{1 \leq k < i} \bar{\lambda}_k q^k e_k^+ \wedge e_{k'}^+ - \sum_{1 \leq k < i} \bar{\lambda}_k q^{2i-k} e_k^+ \wedge e_{k'}^+ - \sum_{1 \leq j < i+1} \bar{\lambda}_j q^j e_j^+ \wedge e_{j'}^+ \right) \\
&= \nu_q \left( - \sum_{1 \leq k < i} \bar{\lambda}_k q^{i-k} e_k^+ \wedge e_{k'}^+ - \bar{\lambda}_i e_i^+ \wedge e_{i'}^+ \right).
\end{aligned}$$

Similar calculations show the identity (40) by using (39).

Now assuming that the equation (33) holds for  $l$ , we have

$$\begin{aligned}
U(\kappa^{l+1}) &= U(\kappa) \wedge i^l \sum_{I, J \in \Theta(l)} f_{I, J}(q) \cdot 1 \otimes e_I^+ \wedge e_J^- \\
&= i^{l+1} \sum_{I, J \in \Theta(l)} \sum_i f_{I, J}(q) \cdot 1 \otimes e_I^+ \wedge e_i^+ \wedge e_i^- \wedge e_J^-,
\end{aligned}$$

where we use the fact that  $\kappa$  is central. First we can focus on the case  $e_I^+ \wedge e_i^+ \wedge e_i^- \wedge e_J^-$  for  $I = \{i_1, \dots, i_l\}$  fixed. Then different cases arise:

**Case  $i \neq i'$  and  $i \in I$ :**  $e_I^+ \wedge e_i^+ \wedge e_i^- \wedge e_J^- = 0$  since we can move  $e_i^-$  to the right (resp. left) by using (34) (resp. (38)) if  $i < i'$  (resp.  $i > i'$ ).

**Case  $i, i' \notin I$ :** in this case we can move  $e_i^+$  to the left and  $e_i^-$  to the right by using (38) and (34), respectively and we get

$$e_I^+ \wedge e_i^+ \wedge e_i^- \wedge e_J^- = (-1)^l q^{l-2\alpha_i} e_{i_1}^+ \wedge \dots \wedge e_i^+ \wedge \dots \wedge e_i^- \wedge \dots \wedge e_{i_1}^- \wedge \dots \wedge e_l^-,$$

where  $\alpha_i \in \mathbb{N}_0$ .

**Case there is a  $k < l$  such that  $i_k = i' \geq i$  and  $i \notin I$ :** in this case we can move  $e_i^+$  to the



left by commuting with  $e_{i_1}^+, \dots, e_{i_{k+1}}^+$  by using relations (38) and we get

$$\begin{aligned} e_{i_1}^+ \wedge \dots \wedge e_{i_l}^+ \wedge e_i^- \wedge e_{i_1}^- \wedge \dots \wedge e_{i_l}^- \\ = (-q)^{-(l-k)} e_{i_1}^+ \wedge \dots \wedge e_{i_k}^+ \wedge e_i^+ \wedge \dots \wedge e_l^+ \wedge e_i^- \wedge e_{i_1}^- \wedge \dots \wedge e_{i_l}^- . \end{aligned}$$

Now if  $i_k = i' > i$  then we can use (39) with  $\gamma_{ij} := -\lambda_i^{-1} \lambda_j$  and we get

$$\begin{aligned} e_I^+ \wedge e_i^+ \wedge e_i^- \wedge e_I^- &= (-q)^{-(l-k)} e_{i_1}^+ \wedge \dots \wedge e_{i_k}^+ \wedge e_i^+ \wedge \dots \wedge e_{i_l}^+ \wedge e_i^- \wedge e_{i_1}^- \wedge \dots \wedge e_{i_l}^- \\ &= (-q)^{-(l-k)} \nu_q \sum_{1 \leq j < i} \bar{\gamma}_{ij} q^{-(j-i+1)} e_{i_1}^+ \wedge \dots \wedge e_{i_{k-1}}^+ \wedge e_j^+ \wedge e_{j'}^+ \wedge \dots \wedge e_{i_l}^+ \wedge e_i^- \wedge e_{i_1}^- \wedge \dots \wedge e_{i_l}^- \\ &\quad - (-q)^{-(l-k)} e_{i_1}^+ \wedge \dots \wedge e_{i_{k-1}}^+ \wedge e_i^+ \wedge e_{i_k}^+ \wedge \dots \wedge e_{i_l}^+ \wedge e_i^- \wedge e_{i_1}^- \wedge \dots \wedge e_{i_l}^- . \end{aligned}$$

Since  $j' \neq i_r$  and  $i' \neq i_r$  for  $r \neq k$  then we can move  $e_j^+$  and  $e_i^+$  to the left and  $e_{j'}^+$  and  $e_i^-$  to the right and we obtain

$$\begin{aligned} e_I^+ \wedge e_i^+ \wedge e_i^- \wedge e_I^- \\ = (-q)^{-(l-k)} \nu_q \sum_{1 \leq j_s < i} \bar{\gamma}_{ij} q^{-(j_s-i+1)+\alpha_{is}} e_{i_1}^+ \wedge \dots \wedge e_{j_s}^+ \wedge \dots \wedge e_{j'_s}^+ \wedge \dots \wedge e_{i_l}^+ \wedge e_{i_1}^- \wedge \dots \wedge e_i^- \wedge \dots \wedge e_{i_l}^- \\ + (-1)^l q^{l-1+2\beta_i} e_{i_1}^+ \wedge \dots \wedge e_i^+ \wedge \dots \wedge e_{i_k}^+ \wedge \dots \wedge e_{i_l}^+ \wedge e_{i_1}^- \wedge \dots \wedge e_i^- \wedge \dots \wedge e_{i_k}^- \wedge \dots \wedge e_{i_l}^- , \end{aligned}$$

where  $\alpha_{is} \in \mathbb{Z}$  and  $\beta_i \in \mathbb{N}_0$ . Now if  $i = i' = n$  and  $n \in I$  then by (35) and (39) we have

$$\begin{aligned} e_I^+ \wedge e_n^+ \wedge e_n^- \wedge e_I^- &= (-1)^{l-1} q^{l-1-2\gamma_n} e_{i_1}^+ \wedge \dots \wedge e_n^+ \wedge e_n^+ \wedge \dots \wedge e_{i_l}^+ \wedge e_{i_1}^- \wedge \dots \wedge e_n^- \wedge e_n^- \wedge \dots \wedge e_{i_l}^- \\ &= (-1)^l q^{l-1-2\gamma_i} \nu_{q^{1/2}}^2 \sum_{1 \leq j, r < n} \bar{\gamma}_{nj} \gamma_{nr} q^{r-j} e_{i_1}^+ \wedge \dots \wedge e_j^+ \wedge e_{j'}^+ \wedge \dots \wedge e_{i_l}^+ \wedge e_{i_1}^- \wedge \dots \wedge e_r^- \wedge e_{r'}^- \wedge \dots \wedge e_{i_l}^- \\ &= (-1)^l q^{l-1-2\gamma_i} \nu_{q^{1/2}}^2 \sum_{1 \leq j < n} |\gamma_{nj}|^2 e_{i_1}^+ \wedge \dots \wedge e_j^+ \wedge e_{j'}^+ \wedge \dots \wedge e_{i_l}^+ \wedge e_{i_1}^- \wedge \dots \wedge e_j^- \wedge e_{j'}^- \wedge \dots \wedge e_{i_l}^- \\ &\quad + (-1)^l q^{l-1-2\gamma_i} \nu_{q^{1/2}}^2 \sum_{\substack{1 \leq j, r < n \\ j \neq r}} \bar{\gamma}_{nj} \gamma_{nr} q^{r-j} e_{i_1}^+ \wedge \dots \wedge e_j^+ \wedge e_{j'}^+ \wedge \dots \wedge e_{i_l}^+ \wedge e_{i_1}^- \wedge \dots \wedge e_r^- \wedge e_{r'}^- \wedge \dots \wedge e_{i_l}^- , \end{aligned}$$

where  $\gamma_n \in \mathbb{Z}$ . Analogously, we can move  $e_j^+, e_j^-$  to the left and  $e_{j'}^+$  and  $e_{j'}^-$  to the right by using (34) and (38), respectively.

**Case  $i > i_l$  and  $i = i'_k$  for some  $k$ :** then we can move  $e_i^+$  to the right by using (34) and we get

$$\begin{aligned} e_{i_1}^+ \wedge \dots \wedge e_{i_l}^+ \wedge e_i^+ \wedge e_i^- \wedge e_{i_1}^- \wedge \dots \wedge e_{i_l}^- \\ = (-1)^{k-1} q^{k-1} e_{i_1}^+ \wedge \dots \wedge e_{i_l}^+ \wedge e_i^+ \wedge e_{i_1}^- \wedge \dots \wedge e_i^- \wedge e_{i_k}^- \wedge \dots \wedge e_{i_l}^- \\ = (-1)^{k-1} q^{k-1} \nu_q \sum_{j < i_k} \gamma_{i_k j} q^{j-i_k+1} e_{i_1}^+ \wedge \dots \wedge e_{i_l}^+ \wedge e_i^+ \wedge e_{i_1}^- \wedge \dots \wedge e_j^- \wedge e_{j'}^- \wedge \dots \wedge e_{i_l}^- \\ \quad - (-1)^{k-1} q^{k-1} e_{i_1}^+ \wedge \dots \wedge e_{i_l}^+ \wedge e_i^+ \wedge e_{i_1}^- \wedge \dots \wedge e_{i_k}^- \wedge e_i^- \wedge \dots \wedge e_{i_l}^- \\ = (-1)^{k-1} q^{k-1} \nu_q \sum_{j_s < i_k} \gamma_{i_k j_s} q^{j_s-i_k+1+\theta_{is}} e_{i_1}^+ \wedge \dots \wedge e_{i_l}^+ \wedge e_i^+ \wedge e_{i_1}^- \wedge \dots \wedge e_{j_s}^- \wedge \dots \wedge e_{j'_s}^- \wedge \dots \wedge e_{i_l}^- \\ \quad + (-1)^l q^{l-1} e_{i_1}^+ \wedge \dots \wedge e_{i_l}^+ \wedge e_i^+ \wedge e_{i_1}^- \wedge \dots \wedge e_{i_k}^- \wedge \dots \wedge e_{i_l}^- \wedge e_i^- . \end{aligned}$$

where  $\theta_{is} \in \mathbb{Z}$ .

**Case  $I \neq J$ :** in the same way it can be show that  $e_I^+ \wedge e_i^+ \wedge e_i^- \wedge e_J^-$  is a linear combination of elements of the form  $e_P^+ \wedge e_P^-$  and  $e_S^+ \wedge e_T^-$  for  $S \neq T$  where  $P, S, T \in \Theta(l+1)$ .

Finally, if  $I' \in \Theta(l+1)$  then from the previous cases we see that the coefficient  $f_{l+1, I', I'}(q)$  of  $e_{I'}^+ \wedge e_{I'}^-$  is of the form

$$(-1)^l \sum_t f_{l, I_t, I_t}(q) q^t + (q - q^{-1}) \xi(q) + (q^{1/2} - q^{-1/2})^2 \zeta(q) + \sum_r \chi_r(q) f_{l, I_r, J_r}(q),$$

where  $I_t, I_r, J_r \in \Theta(l)$  and  $\xi(q), \zeta(q), \chi_r(q) \in \mathbb{C}[q, q^{-1}]$ . Since all the  $f_{l, I_t, I_t}(1)$  have the same sign and  $f_{l, I_r, J_r}(1) = 0$ , then  $f_{l+1, I', I'}(1) = (-1)^l \sum_t f_{l, I_t, I_t}(1) \neq 0$  and

$$\text{sgn}(f_{l+1, I', I'}(1)) = (-1)^l \text{sgn}\left(\sum_t f_{l, I_t, I_t}(1)\right) = (-1)^l (-1)^{\frac{l(l-1)}{2}} = (-1)^{\frac{l(l+1)}{2}}.$$

□

## 4 Spectrum of the $\bar{\partial}$ -Laplace operator on zero forms

In this last section we present the proof of the main result of the paper by using previous results and the representation theory of the quantum group  $U_q(\mathfrak{g})$ .

**Lemma 4.1.** *There exist constants  $\theta, \theta', \theta''$  and  $\theta'''$  such that*

- (i)  $-*_{\kappa}(\partial y \wedge *_{\kappa}(\bar{\partial} y)) = \theta y^2,$
- (ii)  $-*_{\kappa}(*_{\kappa}(\bar{\partial} y) \wedge \partial z) = \theta' y z,$
- (iii)  $-*_{\kappa}(\partial y \wedge *_{\kappa}(\bar{\partial} z)) = \theta'' y z,$
- (iv)  $-*_{\kappa}(*_{\kappa}(\bar{\partial} z) \wedge \partial z) = \theta''' z^2,$

where the constants  $\theta, \theta'$  and  $\theta'''$  are non-zero.

*Proof.* We first note that both sides of the equations (i)-(iv) are highest weight vectors with the same weight. Since the decomposition

$$\Omega^{(0,0)} = \mathcal{O}_q(\mathbf{Q}_N) = \bigoplus_{k, l \in \mathbb{N}_0} U_q(\mathfrak{g}) \blacktriangleright z^k y^l$$

is multiplicity-free we have that there are constants  $\theta, \theta', \theta'', \theta'''$  such that equations (i)-(iv) hold.

To show that  $\theta$  is non-zero, we note first that  $\partial y \wedge *_{\kappa}(\bar{\partial} y) \neq 0$  if and only if  $\partial y \wedge \bar{\partial} y$  is not primitive. This follows from the fact that

$$*_{\kappa}(\bar{\partial} y) \wedge \partial y = -i^{-1} \frac{1}{(n-1)!} L^{n-1}(\bar{\partial} y) \wedge \partial y = -i^{-1} \frac{1}{(n-1)!} L^{n-2+1}(\bar{\partial} y \wedge \partial y).$$

A similar argument shows that  $*_{\kappa}(\bar{\partial} y) \wedge \partial z$ ,  $\partial y \wedge *_{\kappa}(\bar{\partial} z)$  and  $*_{\kappa}(\bar{\partial} z) \wedge \partial z$  are non-zero if and only if  $\bar{\partial} y \wedge \partial z$ ,  $\partial y \wedge \bar{\partial} z$  and  $\bar{\partial} z \wedge \partial z$  are not primitive, respectively.

Since the Lefschetz map is a right  $U_q(\mathfrak{g})$ -module map, if  $\bar{\partial}z \wedge \partial z$  is primitive then  $X \blacktriangleright (\bar{\partial}z \wedge \partial z)$  is primitive for any  $X \in U_q(\mathfrak{g})$ . We claim that there exists  $X \in U_q(\mathfrak{g})$  such that

$$[X \blacktriangleright (\bar{\partial}y \wedge \partial y)] = \gamma [\bar{\partial}z_{Ni}] \wedge [\partial z_{jN}] \quad \text{for some } i, j \in \{2, \dots, N-1\} \text{ and } \gamma \neq 0.$$

or equivalently there exists  $X' \in U_q(\mathfrak{g})$  and  $\gamma' \neq 0$  such that

$$[(\bar{\partial}y \wedge \partial) \triangleleft X'] = \gamma' [\bar{\partial}z_{Ni}] \wedge [\partial z_{iN}].$$

We show this for the case  $\mathfrak{so}_{2n+1}$  since for  $\mathfrak{so}_{2n}$  the proof is similar. We define the sequence

$$\begin{aligned} X_1 &:= F_2, X_2 := F_3, \dots, X_{n-1} := F_n, X_n := F_n, X_{n+1} := F_{n-1}, \dots, X_{2n-2} := F_2, \\ X_{2n-1} &:= F_2, \dots, X_{3n-3} := F_n, X_{3n-2} := F_n, X_{3n-1} := F_{n-1}, \dots, X_{4n-4} := F_2, \\ X_{4n-3} &:= F_1, X_{4n-2} := F_1. \end{aligned}$$

If  $1 \leq i_1 < \dots < i_r \leq 4n-2$  then by using the formulas (13)-(15) it can be shown that if  $y \triangleleft X_{i_1} \cdots X_{i_r}$  is non-zero then it is a multiple of one element of the following list

- (i')  $z_{i,N} - q^{\gamma_i} z_{1,i'}$ ,  $2 \leq i \leq N-1$ ,  $\gamma_i \in \mathbb{Z}$ ,
- (ii')  $z_{N,N} - q^{\eta_1} z_{N-1,N-1} + q^{\eta_2} z_{2,2} - q^{\eta_3} z_{1,1}$ ,  $\eta_i \in \mathbb{Z}$ ,
- (iii')  $z_{i,N-1} - q^{a_i} z_{2,i'}$ ,  $2 \leq i \leq N-2$ ,  $a_i \in \mathbb{Z}$ ,
- (iv')  $z_{N,N-1} - \mu z_{2,1}$ ,  $\mu < 0$ ,

and the only way to obtain  $z_{N,N-1} - \mu z_{2,1}$  is by acting by  $F_2 \cdots F_n^2 \cdots F_2 F_1^2$  on  $y = q(z_{2,N} - q^{-2} z_{1,N-1})$  from the right. If we set  $X := X_1 \cdots X_{4n-2}$  then by using the previous list (i')-(iv') and Lemma 3.3 (ii), we obtain

$$\begin{aligned} [(\partial(y) \wedge \bar{\partial}(y)) \triangleleft X] &= \sum_{\substack{r=0 \\ p \in \mathcal{P}_{4n-2,r}^{r=0}}}^{4n-2} q^{\alpha_r} [\partial(y \triangleleft X_{p(1)} \cdots X_{p(r)}) \wedge [\bar{\partial}(y \triangleleft X_{p(r+1)} \cdots X_{p(4n-2)})]] \\ &= q^\alpha \partial(y \triangleleft F_2 \cdots F_n^2 \cdots F_2) \wedge [\bar{\partial}(y \triangleleft F_2 \cdots F_n^2 \cdots F_2 F_1^2)] \\ &= \gamma [\partial(z_{N-1,N} - q^{\gamma} z_{1,2})] \wedge [\bar{\partial}(z_{N,N-1} - \mu z_{2,1})] \\ &= \gamma e_{N-2}^+ \wedge e_{N-2}^-, \end{aligned} \tag{41}$$

where  $\gamma < 0$  and  $\mathcal{P}_{4n-2,r}$  is the set of  $(r, 4n-2-r)$ -shuffles, that is, the set of permutations  $p \in \mathcal{P}_{4n-2}$  such that  $p(1) < \dots < p(r)$  and  $p(r+1) < \dots < p(4n-2)$ . On the other hand, the form  $e_{N-2}^+ \wedge e_{N-2}^-$  is not primitive since by Lemma 3.5 we have

$$\begin{aligned} [\kappa^{N-3}] \wedge e_{N-2}^+ \wedge e_{N-2}^- &= e_{N-2}^+ \wedge [\kappa]^{N-3} \wedge e_{N-2}^- \\ &= \sum_{I, J \in \Theta(N-3)} f_{N-3, I, J}(q) e_{N-2}^+ \wedge e_I^+ \wedge e_J^- \wedge e_{N-2}^- \\ &= \sum_{I \in \Theta(N-3)} f_{N-3, I, J}(q) e_{N-2}^+ \wedge e_I^+ \wedge e_I^- \wedge e_{N-2}^- \\ &\quad + \sum_{I \neq J \in \Theta(N-3)} f_{N-3, I, J}(q) e_{N-2}^+ \wedge e_I^+ \wedge e_J^- \wedge e_{N-2}^- \\ &= f_{N-3, J, J}(q) e_{N-2}^+ \wedge e_1^+ \wedge \cdots \wedge e_{N-3}^+ \wedge e_1^- \wedge \cdots \wedge e_{N-3}^- \wedge e_{N-2}^- \\ &= (-1)^{N-3} q^{-(N-3)} f_{N-3, J, J}(q) e_1^+ \wedge \cdots \wedge e_{N-2}^+ \wedge e_1^- \wedge \cdots \wedge e_{N-2}^-, \end{aligned}$$

where  $J = \{1, \dots, N-3\}$  and  $f_{N-3,J,J}(q) \neq 0$  if  $q$  belongs to a suitable interval around 1. Since  $e_{N-2}^+ \wedge e_{N-2}^-$  is not primitive then  $(\partial(y) \wedge \bar{\partial}(y)) \triangleleft X_1 \cdots X_{4n-2}$  is not primitive and therefore  $\partial(y) \wedge \bar{\partial}(y)$  is not primitive. The calculation for  $\bar{\partial}y \wedge \partial z$  is similar by using the fact that

$$(\bar{\partial}y \wedge \partial z) \triangleleft F_1 = q\bar{\partial}(z_{2,N} - q^{-2}z_{1,N-1}) \wedge \partial(z_{2,N} + qz_{1,N-1}),$$

and we can proceed by acting with  $X$  and obtain

$$\begin{aligned} [(\bar{\partial}y \wedge \partial z) \triangleleft F_1 X] &= [(\bar{\partial}y \wedge \partial z) \triangleleft F_1 (F_2 \cdots F_n^2 \cdots F_2)^2 F_1^2] \\ &= q^\beta [\bar{\partial}(y \triangleleft F_2 \cdots F_n^2 \cdots F_2 F_1^2)] \wedge [\partial((z_{2,N} + qz_{1,N-1}) \triangleleft F_2 \cdots F_n^2 \cdots F_2)] \\ &= \gamma' [\bar{\partial}(z_{N,N-1} - \mu z_{2,1})] \wedge [\partial(z_{N-1,N} + q^{\gamma'_{N-1}} z_{1,2})] \\ &= \gamma' e_{N-2}^- \wedge e_{N-2}^+, \end{aligned}$$

where  $\gamma' \neq 0$ . Similarly, it can be shown that  $e_{N-2}^- \wedge e_{N-2}^+$  is not primitive by using the fact that the Kähler form  $\kappa$  is non-zero multiple of

$$i \sum_{i,j=1}^N q^{2(N-i)} \bar{\partial}z_{ij} \wedge \partial z_{ji}$$

satisfying analogous results as those of Lemma 3.5, i.e., there exist  $g_{l,I,J}(q) \in \mathbb{C}[q, q^{-1}]$  such that

$$[\kappa^l] = i^l \sum_{I,J \in \Theta(l)} g_{l,I,J}(q) e_I^- \wedge e_J^+,$$

where  $\text{sgn}(g_{l,I,I}(1)) = (-1)^{\frac{l(l-1)}{2}}$  for all  $I \in \Theta(l)$  and  $g_{l,I,J}(1) = 0$ ,  $I \neq J$ .

Much in the same way is the calculation for  $\bar{\partial}(z) \wedge \partial z$ : if we act with  $F_1$  from the right we obtain

$$\begin{aligned} (\bar{\partial}z_{1,N} \wedge \partial z_{1,N}) \triangleleft F_1^2 &= \alpha \bar{\partial}(z_{2,N} + qz_{1,N-1}) \wedge \partial(z_{2,N} + qz_{1,N-1}) + \beta \bar{\partial}(z_{1,N}) \wedge \partial(z_{2,N-1}) \\ &\quad + \zeta \bar{\partial}(z_{2,N-1}) \wedge \partial(z_{1,N}), \end{aligned} \quad (42)$$

where  $\alpha, \beta, \zeta > 0$ . Now a routine calculation shows that

$$[(\bar{\partial}(z_{1,N}) \wedge \partial(z_{2,N-1})) \triangleleft (F_2 \cdots F_n^2 \cdots F_2)^2 F_1^2] = 0, \quad (43)$$

$$\begin{aligned} (\bar{\partial}(z_{2,N-1}) \wedge \partial(z_{1,N})) \triangleleft (F_2 \cdots F_n^2 \cdots F_2)^2 F_1^2 &= \chi (-\bar{\partial}(z_{N,2} + q^\alpha z_{N-1,1}) \wedge \partial(z_{2,N} + q^\beta z_{1,N-1}) \\ &\quad + \vartheta_1 \bar{\partial}(z_{N-1,2}) \wedge \partial(z_{2,N-1}) + \vartheta_2 \bar{\partial}(z_{N,1}) \wedge \partial(z_{1,N})), \end{aligned} \quad (44)$$

for a certain real  $\chi > 0$  and real  $\vartheta_1, \vartheta_2$ . On the other hand, similar calculations as given in (41) show that

$$[(\bar{\partial}(z_{2,N} + qz_{1,N-1}) \wedge \partial(z_{2,N} + qz_{1,N-1})) \triangleleft (F_2 \cdots F_n^2 \cdots F_2)^2 F_1^2] = -\gamma' e_{N-2}^- \wedge e_{N-2}^+, \quad (45)$$

for a real  $\gamma' > 0$ . Now combining equations (42)-(45) we have

$$[(\bar{\partial}z_{1,N} \wedge \partial z_{1,N}) \triangleleft F_1^2 (F_2 \cdots F_n^2 \cdots F_2)^2 F_1^2] = -\gamma' e_{N-2}^- \wedge e_{N-2}^+ - \chi e_1^- \wedge e_1^+$$

which is not primitive since  $-\gamma'$  and  $-\chi$  are both negative and we are assuming that  $q$  belongs to a suitable open interval around 1. Finally, similar calculations work for the case  $\mathfrak{so}_{2n}$  where the element  $X$  is given by  $X := (F_2 \cdots F_n F_{n-2} \cdots F_2)^2 F_1^2$ .  $\square$

With this lemma and results of previous sections we are now ready to calculate the eigenvalues of the Laplace operator  $\Delta_{\bar{\partial}}$  on zero forms.

**Proposition 4.2.** *The eigenvalue of the Dolbeault Laplace operator  $\Delta_{\bar{\partial}}$  on zero forms goes to infinity with finite multiplicity and therefore the operator has compact resolvent.*

*Proof.* Combining Lemma 3.4, Lemma 4.1 and the identities (29) we have

$$\begin{aligned}
\Delta_{\bar{\partial}}(y^k z^l) &= \bar{\partial}^\dagger \bar{\partial}(y^k z^l) \\
&= \bar{\partial}^\dagger ((k)_{q^2} y^{k-1} \bar{\partial} y z^l + (l)_{q-2} y^k \bar{\partial} z z^{l-1}) \\
&= - *_{\kappa} \circ \partial ((k)_{q^2} y^{k-1} *_{\kappa} (\bar{\partial} y) z^l + (l)_{q-2} y^k *_{\kappa} (\bar{\partial} z) z^{l-1}) \\
&= - *_{\kappa} ((k)_{q^2} y^{k-2} \partial y \wedge *_{\kappa} (\bar{\partial} y) z^l + (k)_{q^2} y^{k-1} (\partial \circ *_{\kappa} \circ \bar{\partial} y) z^l \\
&\quad + (k)_{q^2} (l)_{q^2} y^{k-1} (*_{\kappa} \circ \bar{\partial} y) \wedge \partial z z^l + (l)_{q-2} (k)_{q-2} y^{k-1} \partial y \wedge *_{\kappa} (\bar{\partial} z) z^{l-1} \\
&\quad + (l)_{q-2} y^k \partial \circ *_{\kappa} (\bar{\partial} z) z^{l-1} + (l)_{q-2} (l-1)_{q^2} y^k *_{\kappa} (\bar{\partial} z) \wedge \partial z z^{l-2}) \\
&= -(k)_{q^2} y^{k-2} *_{\kappa} (\partial y \wedge *_{\kappa} (\bar{\partial} y)) z^l - (k)_{q^2} y^{k-1} *_{\kappa} (\partial \circ *_{\kappa} \circ \bar{\partial} y) z^l \\
&\quad - (k)_{q^2} (l)_{q^2} y^{k-1} *_{\kappa} ((*_{\kappa} \circ \bar{\partial} y) \wedge \partial z) z^l - (l)_{q-2} (k)_{q-2} y^{k-1} *_{\kappa} (\partial y \wedge *_{\kappa} (\bar{\partial} z)) z^{l-1} \\
&\quad - (l)_{q-2} y^k *_{\kappa} (\partial \circ *_{\kappa} (\bar{\partial} z)) z^{l-1} - (l)_{q-2} (l-1)_{q^2} y^k *_{\kappa} (*_{\kappa} (\bar{\partial} z) \wedge \partial z) z^{l-2} \\
&= [\theta (k)_{q^2} (k-1)_{q-2} + (k)_{q^2} \mu_y + (l)_{q^2} (k)_{q^2} \theta' + (l)_{q-2} (k)_{q-2} \theta'' + (l)_{q-2} \mu_z \\
&\quad + (l)_{q-2} (l-1)_{q^2} \theta'''] y^k z^l,
\end{aligned}$$

where  $\mu_z$  and  $\mu_y$  are the eigenvalues of  $z$  and  $y$  respectively. A differential calculus  $(\Omega^{(\bullet, \bullet)}, \partial, \bar{\partial})$  is said to be *connected* if

$$\text{Ker}(\partial : \Omega^{(0,0)} \rightarrow \Omega^{(1,0)}) = \text{Ker}(\bar{\partial} : \Omega^{(0,0)} \rightarrow \Omega^{(0,1)}) = \mathbb{C}1.$$

Since the Heckengerger–Kolb calculus is connected [1] and  $\ker(\Delta_{\bar{\partial}}) = \ker(\partial) \cap \ker(\bar{\partial})$  [24], then  $\mu_z$  and  $\mu_y$  are nonzero. Now we note that  $(l)_{q-2} (k)_{q-2} \theta''$  and  $(l)_{q-2} \mu_z$  have finite limit as  $k, l \rightarrow \infty$  and therefore they do not contribute in the asymptotic behaviour of the eigenvalue. By Lemma 4.1 we have that  $\theta' \neq 0$ . It is not difficult to see that if  $\theta' < 0$  then the eigenvalue would tend to  $-\infty$  if  $k$  is fixed and  $l \rightarrow \infty$ . Therefore  $\theta' > 0$ .

Now we can consider the sum

$$(\theta \mu_y^{-1} (k-1)_{q-2} + 1) (k)_{q^2} \mu_y + (l)_{q-2} (l-1)_{q^2} \theta'''. \quad (46)$$

First we note that

$$\theta < -(1 - q^{-2}) \mu_y \Rightarrow (1 + (k-1)_{q-2} \theta \mu_y^{-1}) < 0, \quad \text{as } k \rightarrow \infty,$$

and this would imply that the eigenvalues of  $\Delta_{\bar{\partial}}$  on  $y^k$  are negative. Therefore  $\theta \geq -(1 - q^{-2}) \mu_y$ . If  $\theta = -(1 - q^{-2}) \mu_y$  then  $(\theta \mu_y^{-1} (k-1)_{q-2} + 1) (k)_{q^2} \mu_y$  converges when  $k \rightarrow \infty$ . On the other hand, if  $\theta > -(1 - q^{-2}) \mu_y$  then  $(\theta \mu_y^{-1} (k-1)_{q-2} + 1) (k)_{q^2} \mu_y$  tends to infinity.

Now if  $\theta''' < 0$  then

$$\Delta_{\bar{\partial}}(z^l) = \theta''' (l)_{q-2} (l-1)_{q^2} + (l)_{q-2} \mu_z = (\mu_z^{-1} \theta''' (l-1)_{q^2} + 1) \mu_z (l)_{q-2} < 0,$$

as  $l \rightarrow \infty$  which contradicts the fact the eigenvalues of  $z^l$  are non-negatives. Then  $\theta''' \geq 0$ . By Lemma 4.1 we have that  $\theta''' > 0$  and this implies that the second term of (46) tends to infinity as  $l \rightarrow \infty$ . Combining the above results we can conclude that the eigenvalues of  $\Delta_{\overline{\mathcal{F}}}$  tend to infinity, as  $k, l \rightarrow \infty$ .  $\square$

## Acknowledgments

FDG was supported by the Charles University PRIMUS grant *Spectral Noncommutative Geometry of Quantum Flag Manifolds* PRIMUS/21/SCI/026. The author would like to thank Réamonn Ó Buachalla for his support and useful discussions during the preparation of this paper.

## References

- [1] A. Carotenuto, F. Díaz García, R. Ó Buachalla: *A Borel–Weil theorem for the irreducible quantum flag manifolds*. To appear in International Mathematics Research Notices.
- [2] F. D’Andrea, L. Dąbrowski: *Dirac operators on quantum projective spaces*. Commun. Math. Phys. **295**, 731-790, 2010.
- [3] F. D’Andrea, L. Dąbrowski, G. Landi: *The noncommutative geometry of the quantum projective plane*. Rev. Math. Phys. **20**, 979-1006, 2008.
- [4] F. D’Andrea, L. Dąbrowski, G. Landi, E. Wagner: *Dirac operators on all Podleś quantum spheres*. J. Noncommut. Geom. **1**, 213-239, 2007.
- [5] L. Dąbrowski, G. Landi, A. Sitarz, W. van Suijlekom, J.C. Várilly: *The Dirac operator on  $SU_q(2)$* . Commun. Math. Phys. **259**, 729-759, 2005.
- [6] L. Dąbrowski, A. Sitarz: *Dirac operator on the standard Podleś quantum sphere*. Banach Cent. Publ. **61**, 49-58, 2003.
- [7] B. Das, R. Ó Buachalla, P. Somberg: *A Dolbeault–Dirac spectral triple for quantum projective space*. Doc. Math. **25**, 1079-1157, 2020.
- [8] B. Das, R. Ó Buachalla, P. Somberg: *Compact quantum homogeneous Kähler spaces*. arXiv:1910.14007.
- [9] B. Das, R. Ó Buachalla, P. Somberg: *Spectral gaps for twisted Dolbeault–Dirac operators over the irreducible quantum flag manifolds*. arXiv:2206.10719.
- [10] F. Díaz García, R. Ó Buachalla, E. Wagner: *A Dolbeault–Dirac spectral triple for the  $B_2$ -irreducible quantum flag manifold*. arXiv:2109.09885.
- [11] F. Díaz García, A. Krutov, R. Ó Buachalla, P. Somberg, K. R. Strung: *Positive line bundles over the irreducible quantum flag manifolds*. arXiv:1912.08802.
- [12] I. Heckenberger, S. Kolb: *The locally finite part of the dual coalgebra of irreducible quantum flag manifolds*. Proc. London Math. Soc. **89**, 457-484, 2004.
- [13] I. Heckenberger, S. Kolb: *De Rham complex for the quantized irreducible flag manifolds*. J. Algebra, **305**, 704-741, 2006.
- [14] I. Heckenberger, S. Kolb: *Differential forms via the Bernstein–Gelfand–Gelfand resolution for quantized irreducible flag manifolds*. J. Geom. Phys. **57**, 2316-2344, 2007.

- [15] J. Humphreys: *An introduction to Lie algebras and representation theory*. Springer-Verlag, New York, 1972.
- [16] D. Huybrechts: *Complex geometry: an introduction*. Universitext, Springer-Verlag, Heidelberg, 2005.
- [17] A. Klimyk, K. Schmüdgen: *Quantum groups and their representations*. Texts and Monograph in Physics, Springer-Verlag, Berlin, 1998.
- [18] U. Krähmer, M. Tucker-Simmons: *On the Dolbeault–Dirac operator on quantized symmetric spaces*. Trans. London Math. Soc. **2**, 33-56, 2015.
- [19] M. Krämer: *Sphärische Untergruppen in kompakten zusammenhängenden Liegruppen*. Compositio Math. **38**, 129-153, 1979.
- [20] S. Majid: *Noncommutative Riemannian and spin geometry of the standard  $q$ -sphere*. Comm. Math. Phys. **256**, 255-285, 2005.
- [21] M. Matassa: *Kähler structures on quantum irreducible flag manifolds*. J. Geom. Phys. **145** (2019), 103477.
- [22] M. Matassa: *The Parthasarathy formula and a spectral triple for the quantum Lagrangian Grassmannians of rank two*. Lett. Math. Phys. **109**, 1703-1734, 2019.
- [23] S. Neshveyev, L. Tuset: *The Dirac operator on compact quantum groups*. J. Reine. Angew. Math. **641**, 1-20, 2010.
- [24] R. Ó Buachalla: *Noncommutative Kähler structures of quantum homogeneous spaces*. Adv. Math. **322**, 892-939, 2017.
- [25] R. Ó Buachalla: *Noncommutative complex structures on quantum homogeneous spaces*. J. Geom. Phys. **99**, 154-173, 2016.
- [26] M. Takeuchi: *Relative Hopf modules - equivalence and freeness criteria*. J. Algebra, **60**, 452-471, 1979.
- [27] A.L. Onishchick, E. B. Vinberg: *Lie groups and algebraic groups*. Springer-Verlag, Heidelberg, 1990.