

ON POLYNOMIAL CARLESON OPERATORS ALONG QUADRATIC HYPERSURFACES

THERESA C. ANDERSON, DOMINIQUE MALDAGUE AND LILLIAN B. PIERCE

ABSTRACT. We prove that a maximally modulated singular oscillatory integral operator along a hypersurface defined by $(y, Q(y)) \subseteq \mathbb{R}^{n+1}$, for an arbitrary non-degenerate quadratic form Q , admits an *a priori* bound on L^p for all $1 < p < \infty$, for each $n \geq 2$. This operator takes the form of a polynomial Carleson operator of Radon-type, in which the maximally modulated phases lie in the real span of $\{p_2, \dots, p_d\}$ for any set of fixed real-valued polynomials p_j such that p_j is homogeneous of degree j , and p_2 is not a multiple of $Q(y)$. The general method developed in this work applies to quadratic forms of arbitrary signature, while previous work considered only the special positive definite case $Q(y) = |y|^2$.

1. INTRODUCTION

In this work, we study maximally modulated operators of Radon-type. Precisely, let integers $n \geq 1$ and $d \geq 2$ be fixed. For each fixed real-valued polynomial P on \mathbb{R}^n , define an operator, initially acting on Schwartz functions, by

$$(1.1) \quad R_P f(x) = \int_{\mathbb{R}^n} f(x - \gamma(y)) e^{iP(y)} K(y) dy,$$

where $\gamma(y) = (y, Q(y)) \subset \mathbb{R}^{n+1}$ is a hypersurface defined by a fixed non-degenerate quadratic form $Q : \mathbb{R}^n \rightarrow \mathbb{R}$ in n variables, and K is a Calderón-Zygmund kernel (see (2.1)). (Recall that a quadratic form Q on \mathbb{R}^n is said to be non-degenerate if the associated $n \times n$ symmetric matrix, also denoted Q , such that $Q(y) = y^t Q y$, has the property that $\det Q \neq 0$.) We study the corresponding maximally modulated operator, defined by

$$(1.2) \quad f(x) \mapsto \sup_{P \in \mathcal{P}} |R_P f(x)|,$$

in which \mathcal{P} is a chosen set of polynomials. It is reasonable to expect that the operator (1.2) satisfies an *a priori* bound on $L^p(\mathbb{R}^n)$ for all $1 < p < \infty$, for any class \mathcal{P} of polynomials of bounded degree. We prove this for arbitrary quadratic hypersurfaces $\gamma(y) = (y, Q(y))$ for all $n \geq 2$, when \mathcal{P} is a class of polynomials defined as the real span of a set of homogeneous polynomials, whose quadratic component is not Q .

Theorem 1.1. *Fix integers $n \geq 2$ and $d \geq 2$. Let $\gamma : \mathbb{R}^n \mapsto \mathbb{R}^{n+1}$ be defined by*

$$\{(y, Q(y)) : y \in \mathbb{R}^n\}$$

for a non-degenerate quadratic form $Q \in \mathbb{R}[X_1, \dots, X_n]$. Let $p_2(y), \dots, p_d(y)$ be a fixed set of real-valued polynomials on \mathbb{R}^n such that each $p_j(y)$ is homogeneous of degree j and $p_2(y) \neq CQ(y)$ for any $C \neq 0$. Let \mathcal{Q}_d denote the class of real-valued polynomials

$$(1.3) \quad \mathcal{Q}_d = \text{Span}_{\mathbb{R}} \{p_2, p_3, \dots, p_d\}.$$

For each $P \in \mathcal{Q}_d$, define the Carleson operator R_P as in (1.6). For each $1 < p < \infty$ the following a priori inequality holds for all Schwartz class functions $f \in \mathcal{S}(\mathbb{R}^{n+1})$:

$$(1.4) \quad \left\| \sup_{P \in \mathcal{Q}_d} |R_P f| \right\|_{L^p(\mathbb{R}^{n+1})} \leq A \|f\|_{L^p(\mathbb{R}^{n+1})},$$

with a constant A that may depend on n, d, p, Q and the fixed polynomials p_2, \dots, p_d .

A simple change of variables (see §2) shows that it suffices to prove Theorem 1.1 in the case that

$$Q(y) = \theta_1 y_1^2 + \cdots + \theta_n y_n^2, \quad \theta_i \in \{+1, -1\}.$$

If r of the signs θ_i are positive, the form is said to have signature $(r, n - r)$. The special case of Theorem 1.1 for signature $(n, 0)$ with $Q(y) = |y|^2$, so that $\gamma(y) = (y, |y|^2)$ defines a paraboloid, recovers the main theorem of Pierce and Yung [PY19], which introduced the study of Carleson operators of Radon-type. The new work of this paper enables us to treat Radon-type behavior defined by quadratic forms of *arbitrary* signature. This is significantly more difficult than the special positive definite case considered by Pierce and Yung. To treat hyperbolic cases of arbitrary signature, we develop a new perspective that provides a strong, versatile framework for extracting decay from an oscillatory integral operator at the heart of the argument. This perspective replaces ad hoc arguments in the earlier work of Pierce and Yung, and will be useful to study even broader classes of Carleson operators of Radon-type. We now describe the context for these operators, and then describe the key innovations of the present work.

1.1. Relation to previous work. Carleson operators have their genesis in Carleson's 1966 work [Car66] on the convergence of Fourier series. Carleson's theorem can now be summarized as the statement that the operator $f \mapsto \sup_{\lambda \in \mathbb{R}} |T_\lambda f(x)|$ satisfies an L^2 bound, where for each $\lambda \in \mathbb{R}$ we define

$$T_\lambda f(x) = \text{p.v.} \int_{\mathbb{T}} f(x - y) e^{i\lambda y} \frac{dy}{y}.$$

Carleson's work was followed closely by the proof of L^p bounds for $1 < p < \infty$ by Hunt [Hun68], an \mathbb{R}^n version by Sjölin [Sjö71], and importantly, a new method of proof by C. Fefferman [Fef73]. These approaches then led to other celebrated works such as that of Lacey and Thiele [LT00] on the bilinear Hilbert transform.

Elias M. Stein expanded the field of vision by asking whether L^p bounds hold for all $1 < p < \infty$ for polynomial Carleson operators. For each fixed real-valued polynomial P on \mathbb{R}^n , define

$$T_P f(x) = \int_{\mathbb{R}^n} f(x - y) e^{iP(y)} K(y) dy,$$

for K a Calderón-Zygmund kernel (see (2.1)). The corresponding Carleson operator, over a class \mathcal{P} of polynomials, takes the form

$$(1.5) \quad f \mapsto \sup_{P \in \mathcal{P}} |T_P f(x)|.$$

When \mathcal{P} is the class \mathcal{P}_d of all real-valued polynomials on \mathbb{R}^n of degree at most d (for a fixed d), this is called the polynomial Carleson operator. Stein initially proved (1.5) is bounded on $L^2(\mathbb{R})$ for $\mathcal{P} = \text{Span}\{y^2\}$ [Ste95]; this was based on special properties of Gaussians. Next, Stein and Wainger developed a TT^* argument to prove (1.5) is bounded on $L^p(\mathbb{R}^n)$ for $1 < p < \infty$ when $\mathcal{P} = \text{Span}\{y^\alpha : \alpha \in \mathbb{Z}_{\geq 0}^n, 2 \leq |\alpha| \leq d\}$; that is, \mathcal{P} is comprised of all polynomials of degree at most d , with no linear terms [SW01]. In dimension $n = 1$, Lie's breakthrough work finally proved L^p bounds for all $1 < p < \infty$ for the full class \mathcal{P}_d with no restrictions [Lie09, Lie20]. Finally, independent work of Lie and Zorin-Kranich resolved Stein's question in arbitrary dimensions [Lie17, ZK21]; see also the comprehensive survey of Lie [Lie19].

The present manuscript fits in the broader context of a further question raised by Stein: whether an L^p bound holds for a Carleson operator of Radon-type. Let K again be a Calderón-Zygmund kernel and let $\gamma : \mathbb{R}^n \rightarrow \mathbb{R}^m$ define an n -dimensional submanifold in \mathbb{R}^m of finite type (that is, such that it has at most a finite order of contact with any affine hyperplane). Then define for each fixed real-valued polynomial P on \mathbb{R}^n , an operator initially acting on Schwartz functions by

$$(1.6) \quad R_P^\gamma f(x) = \int_{\mathbb{R}^n} f(x - \gamma(y)) e^{iP(y)} K(y) dy.$$

The corresponding maximally modulated Carleson operator is then of the form

$$(1.7) \quad f \mapsto \sup_{P \in \mathcal{P}} |R_P^\gamma f(x)|,$$

in which \mathcal{P} is a chosen class of polynomials, such as \mathcal{P}_d . This setting was first studied by Pierce and Yung [PY19], who in particular considered the paraboloid $\gamma(y) = (y, |y|^2) \subset \mathbb{R}^{n+1}$ for $n \geq 2$. They proved that (1.7) satisfies an *a priori* bound on $L^p(\mathbb{R}^{n+1})$ for all $1 < p < \infty$ for the class $\mathcal{P} = \text{Span}\{p_2, p_3, \dots, p_d\}$ where p_2, p_3, \dots, p_d are $d - 1$ fixed polynomials with the property that p_j is homogeneous of degree j , and $p_2 \notin \text{Span}_{\mathbb{R}}\{|y|^2\}$. This work generalized the TT^* methods of Stein and Wainger [SW01], while also introducing techniques from Littlewood-Paley theory, square functions, and delicate arguments with oscillatory integral estimates. The restriction that $p_2 \notin \text{Span}_{\mathbb{R}}\{|y|^2\}$ was a natural consequence of applying TT^* methods to a maximally modulated operator: such methods only succeed when the phase has sufficient independence from the defining function of the submanifold $\gamma(y)$.

1.2. New strategies. In this paper, we prove L^p bounds for hypersurfaces $\gamma(y) = (y, Q(y))$ for quadratic forms Q of arbitrary signature. Our method also has a TT^* argument at its core, and a crucial part of the argument is to extract decay from an oscillatory integral operator. Carrying this out in the hyperbolic setting of Theorem 1.1 requires two significant innovations. First, we present a new perspective that frames an argument for bounding an oscillatory integral from above in terms of bounding the rank of a linear transformation from below (see §7.1). This new perspective greatly simplifies the present work, and we expect it will be essential in other settings not amenable to certain ad hoc arguments that were sufficient in [PY19]. Second, working with arbitrary signature introduces essential new difficulties, which did not arise in the special positive definite case $Q(y) = |y|^2$, because now we must control interactions between multiple terms in the phase of an oscillatory integral. To tackle this, we work coordinate by coordinate to show that either the oscillatory integral does exhibit decay (which completes the proof), or the phase “looks like” the forbidden case $p_2(y) = Q(y)$ in that coordinate. This strategy succeeds since, by hypothesis in the setting of Theorem 1.1, the phase cannot “look like” this forbidden case $p_2(y) = Q(y)$ in all coordinates simultaneously. (Remark 7.22 outlines precisely why hyperbolic cases are more difficult than the positive definite case.) We provide a roadmap of the entire proof in §1.3.

The study of Carleson operators of Radon-type, of the form (1.7) for a given submanifold γ , has only recently begun, after they were introduced in [PY19]. Several intrinsic difficulties of bounding such operators have been described in [PY19, §2]; here, we remark specifically on the restrictions present in Theorem 1.1. First, the case of dimension $n = 1$ is out of reach of the methods of this paper, effectively due to dimension counting; see Remark 5.1. For $n = 1$, it is natural to ask whether for a monomial curve $\gamma(y) = (y, y^m)$, the operator (1.7) is bounded on L^p for $1 < p < \infty$ for $\mathcal{P} = \mathcal{P}_d$ the class of polynomials of degree at most d . For $m = 1$ this can be reduced to an instance of the original Carleson theorem; see [GPRY17, p. 2980]. This question is open for degree $m \geq 2$; the first results in this direction, weaker than Theorem 1.1, were established by Guo, Pierce, Roos and Yung [GPRY17]. Recently, other authors have obtained interesting results on a cluster of closely related questions (Carleson operators with anisotropic scalings or with fewnomial phases), e.g. [Guo17, Roo19, Ram21a, Ram21b, Bec22]. Nevertheless, fundamental questions about operators of the form (1.7) remain open for $n = 1$. When $\gamma(y) = (y, Q(y))$ is defined by a non-degenerate quadratic form and $n \geq 2$, our methods impose that $\text{Span}\{p_2, \dots, p_d\}$ contains no linear polynomials, and that the (homogeneous) quadratic contribution is not in the span of $Q(y)$. These restrictions are a natural consequence of applying TT^* methods; one might expect that time-frequency methods must be applied in order to avoid such restrictions. (We describe exactly how these restrictions arise in our method in Remark 7.29 and Remark 7.30, respectively.) More generally, it is interesting to ask whether the class \mathcal{Q}_d in Theorem 1.1 could be allowed to

have more degrees of freedom, and it would be very interesting to obtain results for (1.7) for more general submanifolds γ .

1.3. Overview of the argument. We divide the proof of Theorem 1.1 into five steps. Steps 1, 2, 3 and 4 are natural generalizations of the method developed in [PY19], and although we provide a self-contained treatment of these generalizations, we are efficient in our exposition, which comprises Part I of the paper. In contrast, the remaining work, Step 5, departs significantly from the approach of [PY19] and is the novel technical heart of the paper; we present this in sections §7 and §8, which we denote as Part II of the paper. For the convenience of the reader, we give a concise roadmap of these steps. We focus on the case of γ with signature $(r, n - r)$ with $r \geq 1$, not previously covered by the work of [PY19]; for simplicity we will just refer to γ as a hyperboloid.

Step 1 (§3) reduces the proof of Theorem 1.1 to proving an L^2 estimate for an auxiliary operator. Fix a set of homogeneous polynomials p_1, \dots, p_M , and let $P_\lambda(y) = \sum_j \lambda_j p_j(y)$, $\|\lambda\| = \sum_j |\lambda_j|$. Define for η in an appropriate class of C^1 functions and real scaling parameter $a > 0$, the operator

$$(1.8) \quad {}^{(\eta)}I_a^\lambda f(x, t) = \int_{\mathbb{R}^n} f(x - y, t - Q(y)) e^{iP_\lambda(y/a)} \frac{1}{a^n} \eta\left(\frac{y}{a}\right) dy, \quad (x, t) \in \mathbb{R}^{n+1}.$$

This integrates over a portion of the hyperboloid. Roughly speaking, we show that Theorem 1.1 holds if there exists some $\delta > 0$ such that for every $r \geq 1$,

$$(1.9) \quad \left\| \sup_{r \leq \|\lambda\| \leq 2r} \sup_{k \in \mathbb{Z}} |{}^{(\eta)}I_{2^k}^\lambda f| \right\|_{L^2(\mathbb{R}^{n+1})} \ll r^{-\delta} \|f\|_{L^2(\mathbb{R}^{n+1})}.$$

This reduction is encapsulated in Theorem 3.1, and its proof uses results for maximally truncated singular Radon transforms over γ , which we recall in §2. Since it is an interesting open question to study a maximally modulated operator of the form (1.7), over an appropriate class \mathcal{P} of phases, for other types of submanifolds γ , we demonstrate this initial step in a quite general setting. (As the steps progress, we indicate precisely when we need to specialize further, both in terms of the submanifold γ , and the specific polynomials p_1, \dots, p_M used in the polynomial phase.)

Step 2 (§4) introduces a maximal version of a “flatter” auxiliary operator. Rather than integrating over a hyperboloid, this auxiliary operator integrates over \mathbb{R}^{n+1} but with anisotropic scaling compatible with the hyperboloid. Define for each C^1 bump function η and real scaling parameter $a > 0$, the operator

$${}^{(\eta)}J_a^\lambda f(x, t) = \int_{\mathbb{R}^{n+1}} f(x - y, t - s) e^{iP_\lambda(y/a)} \frac{1}{a^n} \eta\left(\frac{y}{a}\right) \frac{1}{a^2} \zeta\left(\frac{s}{a^2}\right) dy ds, \quad (x, t) \in \mathbb{R}^{n+1}.$$

This operator satisfies an L^2 bound analogous to (1.9), by the work of [SW01]. The heart of the second step is to introduce a square function that encapsulates the comparison of the auxiliary operator (1.8) to this flatter analogue. Roughly speaking, the square function, say $S_r(f)$, takes the following form:

$$(1.10) \quad \left(\sum_{k \in \mathbb{Z}} \left(\sup_{r \leq \|\lambda\| < 2r} |({}^{(\eta_k)}I_{2^k}^\lambda - {}^{(\eta_k)}J_{2^k}^\lambda) f| \right)^2 \right)^{1/2}.$$

(More precisely, it is defined using a Littlewood-Paley decomposition that localizes the Fourier support of f in the last variable.) Bounding this square function is then the key to proving (1.9). The central goal is to prove that there exists $\delta > 0$ such that for any Schwartz function f on \mathbb{R}^{n+1} , and any $r \geq 1$,

$$(1.11) \quad \|S_r(f)\|_{L^2(\mathbb{R}^{n+1})} \ll r^{-\delta} \|f\|_{L^2(\mathbb{R}^{n+1})}.$$

In order to verify this, one can see from the crude definition of $S_r(f)$ suggested by (1.10) that it is natural to aim to prove that uniformly in $k \in \mathbb{Z}$,

$$(1.12) \quad \left\| \sup_{r \leq \|\lambda\| < 2r} |({}^{(\eta_k)}I_{2^k}^\lambda - {}^{(\eta_k)}J_{2^k}^\lambda) f| \right\|_{L^2(\mathbb{R}^{n+1})} \ll r^{-\delta} 2^{-\varepsilon|k|} \|f\|_{L^2(\mathbb{R}^{n+1})},$$

for some $\delta > 0, \varepsilon > 0$. (More precisely, one must also prove an estimate that takes into account the Littlewood-Paley decomposition.) Theorem 4.3 formally records that if an appropriate bound roughly of the form (1.12) holds, then the desired square function estimate (1.11) holds on L^2 . Thus in order to complete the proof of our main theorem, we must verify that the hypothesis (1.12) of Theorem 4.3 is true. (Like Step 1, Step 2 also works for more general manifolds γ .)

Step 3 (§5) defines a crucial change of variables. Roughly speaking, fix $k \in \mathbb{Z}$ and let T denote the operator

$$f \mapsto \sup_{r \leq \|\lambda\| < 2r} |(\eta_k) I_{2^k}^\lambda - (\eta_k) J_{2^k}^\lambda f|.$$

(More precisely, to define T we linearize this operator, using stopping-times.) Our general strategy for proving (1.12) is to bound TT^* on L^2 , with a norm that exhibits decay like $r^{-\delta} 2^{-\varepsilon|k|}$ for some $\delta, \varepsilon > 0$. In Step 3, we define a change of variables that will allow us to show that $TT^*f(x, t)$ can be written as a sum of n convolutions $(f * K_{\sharp, l}^{\nu, \mu})(x, t)$ for an appropriate kernel $K_{\sharp, l}^{\nu, \mu}$, for each $1 \leq l \leq n$. Roughly speaking, if TT^*f is initially an integral over $(u, z) \in \mathbb{R}^n \times \mathbb{R}^n$, for each $1 \leq l \leq n$ we construct a change of variables to $(u, \tau, \sigma) \in \mathbb{R}^n \times \mathbb{R} \times \mathbb{R}^{n-1}$ so that

$$K_{\sharp, l}^{\nu, \mu}(u, \tau) = \int_{\mathbb{R}^{n-1}} e^{iP_\nu(u+z) - iP_\mu(z)} \Psi(u, z) d\sigma, \quad (u, \tau) \in \mathbb{R}^{n+1},$$

where $\sigma \in \mathbb{R}^{n-1}$ is defined implicitly in terms of u, z for every $z \in \mathbb{R}^n$, $\Psi(u, z)$ is a C^1 bump function, and ν, μ are stopping-times. We do this for each $1 \leq l \leq n$, depending on whether u lies in the l -th sector of a partition of \mathbb{R}^n ; this guarantees the compact support of σ in the region of integration. To define the change of variables in Step 3, we crucially use the fact that in our main theorem we assume that $\gamma(y) = (y, Q(y))$ for a non-degenerate *quadratic* form Q .

Step 4 (§6) shows that if an appropriate upper bound holds pointwise for the kernel $K_{\sharp, l}^{\nu, \mu}$ defined in Step 3, then the hypothesis (1.12) of Theorem 4.3 is true. This deduction is encapsulated in Theorem 6.1 This is the step in which TT^* methods are clearly applied. The conclusion of Step 4 ends Part I. After Step 4, what is left in order to complete the proof of Theorem 1.1 is to prove that the desired pointwise upper bound for $K_{\sharp, l}^{\nu, \mu}$ does hold.

Step 5 (§7 and §8) comprises Part II of the paper. Step 5 proves that the desired bound for $K_{\sharp, l}^{\nu, \mu}$ holds, as stated in Theorem 7.1. Step 5 crucially uses the assumption (1.3) on the class of polynomials \mathcal{Q}_d over which the maximal modulation occurs in Theorem 1.1. As mentioned before, Step 5 is the most novel and technically difficult work of the paper, and we split it into two sections. In §7, we use the explicit change of variables chosen in Step 3 in order to express $K_{\sharp, l}^{\nu, \mu}$ as an oscillatory integral with a phase that depends on stopping-time functions. We show that this phase is sufficiently large (almost all the time), so that the oscillatory integral has a satisfactory upper bound (via a van der Corput estimate), as long as certain polynomials, defined in terms of Q and the fixed polynomials p_2, \dots, p_d defining the class \mathcal{Q}_d , do not vanish identically. To accomplish this, we develop a new framework which differs conceptually from [PY19], and applies more generally. We verify these delicate polynomial properties in §8, and here we must overcome substantial difficulties not present in the setting of [PY19]; see Remark 7.22 for a precise description of why the analysis in [PY19] was simpler at this step.

Part I

2. PRELIMINARIES

In this section, we briefly record well-known properties of singular and maximal Radon transforms, which we will call upon in §3.

2.1. Calderón-Zygmund kernels. We work with Calderón-Zygmund kernels K defined as follows: K is a tempered distribution that agrees with a C^1 function $K(x)$ for $x \neq 0$, such that

$$(2.1) \quad |\partial_x^\alpha K(x)| \ll |x|^{-n-|\alpha|}, \quad 0 \leq |\alpha| \leq 1,$$

and \hat{K} is an L^∞ function. Such a kernel admits a decomposition

$$K(x) = \sum_{j=-\infty}^{\infty} 2^{-nj} \phi_j(2^{-j}x) =: \sum_{j=-\infty}^{\infty} K_j(x)$$

where each ϕ_j has the following properties (see [SW01, §5]):

- (i) ϕ_j is a C^1 function with support in $1/4 < |x| \leq 1$,
- (ii) $|\partial_x^\alpha \phi_j(x)| \leq C$ for $0 \leq |\alpha| \leq 1$ for some constant C that is uniform in j ,
- (iii) $\int_{\mathbb{R}^n} \phi_j(x) dx = 0$ for every j .

2.2. Singular and maximal Radon transforms. Let $\gamma \subset \mathbb{R}^m$ be a submanifold described by a polynomial mapping $\gamma = (\gamma_1, \dots, \gamma_m) : \mathbb{R}^n \rightarrow \mathbb{R}^m$ where for each i , $\gamma_i : \mathbb{R}^n \rightarrow \mathbb{R}$ is a polynomial with real coefficients. The singular Radon transform

$$(2.2) \quad \mathcal{T}f(x) = \int_{\mathbb{R}^n} f(x - \gamma(y))K(y)dy, \quad x \in \mathbb{R}^m,$$

initially defined for f of Schwartz class, extends to a bounded operator on $L^p(\mathbb{R}^m)$ for $1 < p < \infty$. This follows from [Ste93, Ch. XI §4.4-4.5]. (Precisely, the argument presented in §4.5 of that text assumes that K is homogeneous, but substituting the decomposition K_j described above for the functions K_j in that text, and using the properties (i), (ii), (iii) of the functions ϕ_j confirms that the resulting measures $\{dm^j\}$ satisfy the formalism in §4.4, and the argument can proceed verbatim.)

Next, define the maximally truncated singular Radon transform

$$(2.3) \quad \sup_{t>0} |\mathcal{T}_t f(x)| = \sup_{t>0} \left| \int_{|y| \geq t} f(x - \gamma(y))K(y)dy \right|.$$

When $\gamma(y) = (y, Q(y))$ with $Q(y) = |y|^2$, [PY19, Appendix] proves that this operator extends to a bounded operator on $L^p(\mathbb{R}^n)$ for all $1 < p < \infty$; the same argument applies (with only superficial changes) for any diagonal form $Q(y) = \sum_{1 \leq i \leq n} \theta_i y_i^2$ for any signs $\theta_i \in \{\pm 1\}$; this is the only case we require to prove Theorem 1.1. But in §3 of our work, it is no additional trouble to consider more general submanifolds defined by any polynomial mapping $\gamma : \mathbb{R}^n \rightarrow \mathbb{R}^m$. Thus for such γ , we note that the operator (2.3) extends to a bounded operator on $L^p(\mathbb{R}^n)$ for all $1 < p < \infty$ by [MSZK20a, Thm. 1.30] (although with slightly different constraints on K , see Remark 3.2); see also [DdF86, JSW08] for related results.

Finally, define for a Schwartz function f ,

$$(2.4) \quad \mathcal{M}_\gamma f(x) = \sup_{a>0} \int_{\mathbb{R}^n} |f(x - \gamma(y))| \frac{1}{a^n} \chi_{B_1}\left(\frac{y}{a}\right) dy.$$

For γ an n -dimensional submanifold in \mathbb{R}^m of finite type (this is satisfied if γ is defined by a polynomial mapping), \mathcal{M}_γ satisfies an *a priori* bound on L^p for all $1 < p \leq \infty$ by [Ste93, Ch. XI Thm. 1].

2.3. Reduction to the simplest type of quadratic form. In our main theorem we assume $\gamma(y) = (y, Q(y))$ for a non-degenerate quadratic form Q . We can restrict to proving Theorem 1.1 for

$$(2.5) \quad Q(y) = \sum_{i=1}^n \theta_i y_i^2, \quad \theta_i \in \{\pm 1\}.$$

Indeed, suppose first of all that $Q(y)$ is a non-degenerate quadratic form in $\mathbb{R}[X_1, \dots, X_n]$, with corresponding real symmetric $n \times n$ matrix Q such that $Q(y) = y^T Q y$. Then by the spectral theorem, there exists an orthogonal matrix B such that $BQB^{-1} = A = \text{diag}(a_1, \dots, a_n)$ with all $a_i \neq 0$. Then for $P_\lambda(y) = \sum_{1 \leq j \leq M} \lambda_j p_j(y)$, our operator of interest is

$$\begin{aligned} R_{P_\lambda, K}^Q f(x, t) &:= \int_{\mathbb{R}^n} f(x - y, t - Q(y)) e^{iP_\lambda(y)} K(y) dy \\ &= \int_{\mathbb{R}^n} f(x - y, t - (By)^T A (By)) e^{iP_\lambda(y)} K(y) dy. \end{aligned}$$

Define $\delta_B f(x, t) = f(Bx, t)$; $P_\lambda^B(y) = \sum_{1 \leq j \leq M} \lambda_j p_j^B(y)$ where $p_j^B(y) = p_j(B^{-1}y)$; and $K^B(y) = K(B^{-1}y)$, which is also a Calderón-Zygmund kernel as in (2.1). Note that $p_2 \notin \text{Span}(Q)$ if and only if $p_2^B \notin \text{Span}(A)$. After a change of variables $u = By$, we see that as an operator

$$R_{P_\lambda, K}^Q f = \delta_B \circ R_{P_\lambda^B, K^B}^A \circ \delta_{B^{-1}} f.$$

Thus, to prove that for any fixed Calderón-Zygmund kernel K , and any fixed polynomials p_1, \dots, p_M , the operator $f \mapsto \sup_{P \in \text{Span}\{p_1, \dots, p_M\}} |R_{P, K}^Q f|$ is bounded on L^p , it suffices to prove that for any fixed Calderón-Zygmund kernel K , and any fixed polynomials p_1, \dots, p_M the operator $f \mapsto \sup_{P \in \text{Span}\{p_1, \dots, p_M\}} |R_{P, K}^A f|$ is bounded on L^p . Similarly, another change of variables shows that we may rescale to the case where each $a_i = \pm 1$, so that from now on we may assume that $Q(y)$ takes the form (2.5). This specific form of Q only becomes useful in Step 3 (§5), when it motivates an explicit change of variables.

3. STEP 1: REDUCTION TO AN L^2 ESTIMATE

We begin by introducing an auxiliary operator, and show that an L^2 bound for this operator implies L^p bounds for the Carleson operator (1.7) for $1 < p < \infty$. For our main theorem, we only apply this in the case that $\gamma(y) = (y, Q(y)) \subset \mathbb{R}^{n+1}$, but it is no trouble to work more generally in this section, for readers interested in the case where $\gamma : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is an n -dimensional submanifold in \mathbb{R}^m defined by polynomials.

We now fix the class of polynomial phases we will consider. Let p_1, \dots, p_M be fixed homogeneous polynomials in $\mathbb{R}[X_1, \dots, X_n]$. For each m , we will set d_m to be the degree of p_m . We assume that the polynomials p_m are linearly independent over \mathbb{R} , which causes no limitations in our ultimate results, which involve a supremum over polynomials in the span of p_1, \dots, p_M . Precisely, we set

$$(3.1) \quad \mathcal{P} = \text{Span}_{\mathbb{R}}\{p_1, \dots, p_M\}$$

so that any $P \in \mathcal{P}$ can be expressed as

$$P_\lambda(y) = \sum_{m=1}^M \lambda_m p_m(y),$$

for a certain $\lambda = (\lambda_1, \dots, \lambda_M) \in \mathbb{R}^M$. We do not yet need to make any further assumptions about the polynomials p_m .

Next we define an auxiliary operator. Fix a C^1 bump function η supported in the unit ball B_1 in \mathbb{R}^n and a real number $a > 0$. Then define the operator

$$(3.2) \quad {}^{(\eta)}I_a^\lambda f(x) = \int_{\mathbb{R}^n} f(x - \gamma(y)) e^{iP_\lambda(y/a)} \frac{1}{a^n} \eta\left(\frac{y}{a}\right) dy.$$

This is dominated pointwise by the maximal Radon transform \mathcal{M}_γ defined in (2.4), so it admits an *a priori* estimate for $1 < p \leq \infty$. Indeed, if the bump function η varies over a family $\{\eta_k\}_k$ of

C^1 bump functions supported in B_1 , with uniformly bounded C^1 norm, then

$$(3.3) \quad \left\| \sup_{\lambda \in \mathbb{R}^M} \sup_{k \in \mathbb{Z}} |^{(\eta_k)} I_{2^k}^\lambda f| \right\|_{L^p(\mathbb{R}^m)} \ll \|f\|_{L^p(\mathbb{R}^m)}, \quad 1 < p \leq \infty.$$

This will serve as a trivial bound. However, it is possible to use oscillation within $^{(\eta_k)} I_{2^k}^\lambda$ to prove that the L^2 norm exhibits decay, measured in terms of the “size” of λ . For this purpose, we define the isotropic norm for $\lambda \in \mathbb{R}^M$:

$$(3.4) \quad \|\lambda\| = \sum_{m=1}^M |\lambda_m|.$$

The main result of this section is:

Theorem 3.1. *Let $p_1(y), \dots, p_M(y)$ be a fixed set of linearly independent, homogeneous polynomials in $\mathbb{R}[X_1, \dots, X_n]$, and for any $\lambda \in \mathbb{R}^M$ set $P_\lambda = \sum_{m=1}^M \lambda_m p_m$. Let $\gamma = (y, Q(y)) \subset \mathbb{R}^m$, $m = n + 1$. Suppose that for any family $\{\eta_k\}_{k \in \mathbb{Z}}$ of C^1 bump functions supported in the unit ball $B_1(\mathbb{R}^n)$ with C^1 norm uniformly bounded by 1, there exists a fixed $\delta > 0$ such that for any Schwartz function f on \mathbb{R}^{n+1} and any $r \geq 1$,*

$$(3.5) \quad \left\| \sup_{\substack{\lambda \in \mathbb{R}^M \\ r \leq \|\lambda\| < 2r}} \sup_{k \in \mathbb{Z}} |^{(\eta_k)} I_{2^k}^\lambda f(x, t)| \right\|_{L^2(\mathbb{R}^m)} \ll r^{-\delta} \|f\|_{L^2(\mathbb{R}^m)},$$

in which the implicit constant may depend only on n, m and the fixed polynomials p_1, \dots, p_M . Then for each $1 < p < \infty$, we have the a priori estimate

$$(3.6) \quad \left\| \sup_{\lambda \in \mathbb{R}^M} |R_{P_\lambda}^\gamma f| \right\|_{L^p(\mathbb{R}^m)} \ll \|f\|_{L^p(\mathbb{R}^m)},$$

in which the implicit constant may depend only on p, n, m , and the fixed polynomials p_1, \dots, p_M .

Note that in the notation of Theorem 1.1, the left-hand side of (3.6) is $\|\sup_{P \in \mathcal{P}} |R_P^\gamma f|\|_{L^p(\mathbb{R}^{n+1})}$ for the class \mathcal{P} in (3.1). (For simplicity we have stated this theorem for values $r \geq 1$, but of course the same method of proof applies for $r \geq c$ for any fixed universal constant c , as we apply in (3.9).) The theorem also holds for any submanifold defined by a polynomial mapping $\gamma : \mathbb{R}^n \rightarrow \mathbb{R}^m$ under appropriate assumptions on K (summarized in Remark 3.2), by the same argument we now describe.

The proof closely follows the original ideas of Stein and Wainger [SW01] as adapted by Pierce and Yung for the paraboloid [PY19, §5]; we generalize those steps briefly here. We recall that the operator $R_{P_\lambda}^\gamma$ is defined in (1.6), and we apply the dyadic decomposition $K = \sum_j K_j$ from (2.1). Given a fixed $\lambda \in \mathbb{R}^M$, we divide the indices j according to whether the phase polynomial $P_\lambda(y)$ is small or large on the support of $K_j(y)$. For this purpose, it is useful to let d_m denote $\deg p_m$ for each $1 \leq m \leq M$, and to define the nonisotropic norm

$$N(\lambda) = \sum_{m=1}^M |\lambda_m|^{1/d_m}.$$

Note that there exists a constant $c_0 = c_0 > 0$, depending only on M and the multi-set of degrees $\{d_1, \dots, d_M\}$, such that

$$(3.7) \quad N(\lambda) \leq c_0 \|\lambda\| \quad \text{for } N(\lambda) \geq 1.$$

Additionally, for any $1 \leq m \leq M$, $|\lambda_m| \leq N(\lambda)^{d_m}$.

For each fixed $\lambda \in \mathbb{R}^M$, we then set

$$K_\lambda^+ = \sum_{2^j \geq 1/N(\lambda)} K_j, \quad K_\lambda^- = \sum_{2^j < 1/N(\lambda)} K_j.$$

For each $\lambda \in \mathbb{R}^M$, we split $R_{P_\lambda}^\gamma f = R_{P_\lambda}^{\gamma,+} f + R_{P_\lambda}^{\gamma,-} f$, where

$$R_{P_\lambda}^{\gamma,\pm} f(x, t) = \int_{\mathbb{R}^n} f(x - \gamma(y)) e^{iP_\lambda(y)} K_\lambda^\pm(y) dy.$$

3.1. The case of large oscillation. We first bound the operator $\sup_\lambda |R_{P_\lambda}^{\gamma,+} f(x, t)|$, which corresponds to the case in which the phase $P_\lambda(y)$ is large, and oscillation plays a role. Define the nonisotropic scaling

$$2^j \circ \lambda = (2^{jd_m} \lambda_m)_m.$$

Here we use the assumption that each polynomial p_m is homogeneous of degree d_m , so that $P_{2^j \circ \lambda}(y/2^j) = P_\lambda(y)$. Moreover, the nonisotropic norm is homogeneous with respect to this scaling: $N(2^j \circ \lambda) = 2^j N(\lambda)$. Consequently, for each fixed λ , the operator decomposes as

$$R_{P_\lambda}^{\gamma,+} = \sum_{N(2^j \circ \lambda) \geq 1} (\phi_j) I_{2^j}^{2^j \circ \lambda}.$$

Thus, for a Schwartz function f , we have the pointwise inequality

$$\begin{aligned} \sup_\lambda |R_{P_\lambda}^{\gamma,+} f(x)| &\leq \sup_\lambda \sum_{N(2^j \circ \lambda) \geq 1} \sup_{k \in \mathbb{Z}} |(\phi_k) I_{2^k}^{2^j \circ \lambda} f(x, t)| \\ (3.8) \quad &\leq \sum_{l=0}^{\infty} \sup_{2^l \leq N(\lambda') < 2^{l+1}} \sup_{k \in \mathbb{Z}} |(\phi_k) I_{2^k}^{\lambda'} f(x, t)|. \end{aligned}$$

The hypothesis of Theorem 3.1 provides an L^2 estimate for each operator on the right-hand-side. Precisely, by (3.7), for $r \geq 1$, $\{\lambda : N(\lambda) \geq r\} \subseteq \{\lambda : c_0 \|\lambda\| \geq r\}$. Thus for any $r \geq 1$,

$$\sup_{N(\lambda') \geq r} \sup_{k \in \mathbb{Z}} |(\phi_k) I_{2^k}^{\lambda'} f(x, t)| \leq \sup_{\|\lambda'\| \geq r/c_0} \sup_{k \in \mathbb{Z}} |(\phi_k) I_{2^k}^{\lambda'} f(x, t)| \leq \sum_{2^v \geq r/c_0} \sup_{2^v \leq \|\lambda'\| < 2^{v+1}} \sup_{k \in \mathbb{Z}} |(\phi_k) I_{2^k}^{\lambda'} f(x, t)|.$$

By the key hypothesis (3.5), for any $r \geq 1$, the L^2 norm thus satisfies

$$(3.9) \quad \left\| \sup_{N(\lambda') \geq r} \sup_{k \in \mathbb{Z}} |(\phi_k) I_{2^k}^{\lambda'} f| \right\|_{L^2} \leq \sum_{2^v \geq r/c_0} (2^v)^{-\delta} \|f\|_{L^2} \ll r^{-\delta} \|f\|_{L^2}.$$

On the other hand, recall the trivial bound (3.3), valid for all $1 < p < \infty$. Interpolation of (3.9) and (3.3) shows that for each $1 < p < \infty$, there exists $\delta(p)$ such that for any $r \geq 1$,

$$(3.10) \quad \left\| \sup_{N(\lambda') \geq r} \sup_{k \in \mathbb{Z}} |(\phi_k) I_{2^k}^{\lambda'} f| \right\|_{L^p} \ll r^{-\delta(p)} \|f\|_{L^p}.$$

We remark that this interpolation argument leading to (3.10) requires several standard intermediate steps in order to pass from the *a priori* inequalities to appropriate versions for all simple functions, which can then be interpolated. We refer to the detailed treatments of these intermediate steps for the exemplary case of maximal operators over curves, in [SW78, Part II §5]. Applying (3.10) in (3.8) after taking norms, shows that for each $1 < p < \infty$,

$$\left\| \sup_\lambda |R_{P_\lambda}^{\gamma,+} f(x)| \right\|_{L^p} \leq \sum_{l=0}^{\infty} 2^{-l\delta(p)} \|f\|_{L^p} \ll \|f\|_{L^p}.$$

3.2. The case of small oscillation. Next we consider the portion $R_{P_\lambda}^{\gamma; -}$ of the operator $R_{P_\lambda}^\gamma$, for which $P_\lambda(y)$ is small, and oscillation does not play an important role. The support of the kernel $K_\lambda^-(y)$ is contained in $|y| \leq 1/N(\lambda)$. On this set,

$$|e^{iP(y)} - 1| \ll |P(y)| \leq \sum_{m=1}^M |\lambda_m| |p_m(y)| \ll \sum_{m=1}^M (N(\lambda)|y|)^{d_m} \leq \sum_{m=1}^M N(\lambda)|y|,$$

where in the last inequality we applied the fact that $N(\lambda)|y| \leq 1$. The implicit constants depend only the set of fixed polynomials $\{p_1, \dots, p_M\}$. For fixed λ , we can then write

(3.11)

$$R_{P_\lambda}^{\gamma; -} f(x, t) = \int_{|y| \leq 1/N(\lambda)} f(x - \gamma(y)) K_\lambda^-(y) dy + O\left(N(\lambda) \int_{|y| \leq 1/N(\lambda)} |y|^{-n+1} |f(x - \gamma(y))| dy\right).$$

The first term is controlled by a maximally truncated singular Radon transform, which is bounded on $L^p(\mathbb{R}^n)$ for all $1 < p < \infty$ by the observations recorded for (2.3). The second term is controlled by the maximal operator \mathcal{M}_γ defined in (2.4), since

$$\begin{aligned} N(\lambda) \sum_{2^l \leq N(\lambda)^{-1}} 2^{-l(n-1)} \int_{2^{l-1} \leq |y| < 2^l} |f(x - \gamma(y))| dy \\ \ll N(\lambda) \sum_{2^l \leq N(\lambda)^{-1}} 2^l \frac{1}{|B_{2^l}|} \int |f(x - \gamma(y))| \chi_{B_{2^l}}(y) dy \ll N(\lambda) N(\lambda)^{-1} \mathcal{M}_\gamma f(x). \end{aligned}$$

In conclusion, for each $1 < p < \infty$, $\|\sup_\lambda |R_{P_\lambda}^{\gamma; -} f|\|_{L^p} \ll \|f\|_{L^p}$. This completes the proof of Theorem 3.1.

Remark 3.2. For readers interested in the more general case when γ is a polynomial mapping, we summarize how the fact that the operator (2.3) extends to a bounded operator on $L^p(\mathbb{R}^n)$ for $1 < p < \infty$ is deduced from [MSZK20a]. Their work applies to kernels K with the properties: (a) (boundedness) $|K(x)| \leq C|x|^{-n}$ for all nonzero $x \in \mathbb{R}^n \setminus \{0\}$; (b) (cancellation) $\int_{r \leq |y| \leq R} K(y) dy = 0$ for $0 < r < R < \infty$; (c) (smoothness) $|K(x) - K(x+y)| \leq \omega_K(|y|/|x|)|x|^{-n}$ for all x, y such that $|y| \leq |x|/2$; here ω_K is a modulus of continuity. For such a kernel, [MSZK20a, Thm. 1.30] states that if $\|\omega_K^\theta\|_{\log \text{Dini}} + \|\omega_K^{\theta/2}\|_{\text{Dini}} < \infty$ for some $\theta \in (0, 1]$ then for every $p \in \{1 + \theta, (1 + \theta)'\}$ and $f \in L^p(\mathbb{R}^n)$, $J_2^p((\mathcal{T}_t f)_{t>0} : \mathbb{R}^n \rightarrow \mathbb{C}) \ll_{n,p} \|f\|_{L^p}$. Here the jump norm is

$$\begin{aligned} J_2^p((\mathcal{T}_t f)_{t>0} : \mathbb{R}^n \rightarrow \mathbb{C}) &= \sup_{\lambda>0} \|\lambda(N_\lambda(\mathcal{T}_t f) : t > 0)\|_{L^p}^{1/2} \\ &= \sup_{\lambda>0} \|\lambda(\sup\{J : \min_{0 < j \leq J} |\mathcal{T}_{t_j} f(\cdot) - \mathcal{T}_{t_{j-1}} f(\cdot)| \geq \lambda\})\|_{L^p}^{1/2} \end{aligned}$$

in which the minimum is over all $0 < j \leq J$ and all $t_0 < \dots < t_J$ with $t_j > 0$. In particular, by [MSZK20b, Lemma 2.3], for a given $1 < p < \infty$ for which Theorem 1.30 in [MSZK20a] holds, then for $r \in (2, \infty]$ the r -variational semi-norm $V^r(\mathcal{T}_t f : t > 0)$ satisfies

$$(3.12) \quad \|V^r(\mathcal{T}_t f : t > 0)\|_{L^{p,\infty}} \ll_{p,r} J_2^p(\mathcal{T}_t f : t > 0) \ll_{n,p} \|f\|_{L^p}.$$

If for example K is homogeneous of degree $-n$, smooth away from the origin, and satisfies the cancellation condition $\int_{|y|=1} K(y) d\sigma(y) = 0$, then it certainly satisfies the above properties (a), (b), (c) with $\omega_K(t) = ct$ for an appropriate constant $c \in (0, \infty)$. Similarly if K is C^1 away from the origin, satisfies (2.1) and (b) (which is somewhat stronger than the assumption $\hat{K} \in L^\infty$) then again all three properties are satisfied with $\omega_K(t) = ct$ for an appropriate constant $c \in (0, \infty)$. For $w_K(t) = ct$, the Dini and log-Dini norms (see [MSZK20a, Eqn. (1.24)]) of $(\omega_K(t))^\theta = c^\theta t^\theta$ are finite for every $\theta \in (0, 1]$. Therefore, by (3.12) we conclude that for each $p \in (1, \infty)$ and $r \in (2, \infty]$

$$(3.13) \quad \|V^r(\mathcal{T}_t f : t > 0)\|_{L^{p,\infty}} \ll_{p,r} \|f\|_{L^p}.$$

For $r = \infty$ we have the pointwise inequality

$$\sup_{t>0} |\mathcal{T}_t f(x)| \leq |\mathcal{T}f(x)| + V^r(\mathcal{T}_t f(x) : t > 0).$$

The operator $\mathcal{T}f(x)$ extends to a bounded operator on $L^p(\mathbb{R}^n)$ for $1 < p < \infty$ (recall (2.2)), and we apply (3.13) to the second term on the right-hand side. Finally, the strong L^p boundedness of $f \mapsto \sup_{t>0} |\mathcal{T}_t f|$ follows after invoking the Marcinkiewicz interpolation theorem with any exponents p_1, p_2 satisfying $1 < p_1 < p < p_2 < \infty$.

4. STEP 2: PASSAGE TO A SQUARE FUNCTION

In this section, we show that the key hypothesis (3.5) of Theorem 3.1 can be verified if a related square function has an L^2 bound with decay in the norm $\|\lambda\|$ defined in (3.4). We only require the case that $\gamma(y) = (y, Q(y))$, but the reader will note that the argument in this section may be adapted to the case where $\gamma(y)$ is a hypersurface in \mathbb{R}^{n+1} , of the form

$$(4.1) \quad \gamma(y) = (y, q(y)) \quad \text{for a homogeneous polynomial } q \text{ of degree } d_q \geq 1.$$

We now denote f as a function of $(x, t) \in \mathbb{R}^{n+1}$, so that along the hypersurface γ , we study the function $f(x - y, t - Q(y))$. We continue to assume homogeneous polynomials p_1, \dots, p_M have been fixed as in (3.1), but do not yet need further assumptions about them.

4.1. A flat analogue: the J -operator. The square function compares the I -operator defined in (3.2) to a smoother “flat” analogue (with a compatible nonisotropic scaling), defined as follows. Fix a C^1 function ζ with $\|\zeta\|_{C^1} \leq 1$, and with ζ supported on $B_1(\mathbb{R})$. Then for any $P_\lambda \in \mathcal{P} = \text{Span}\{p_1, \dots, p_M\}$, any C^1 bump function η supported in the unit ball, and any $a > 0$, define the operator, acting on a Schwartz function f , by

$$(4.2) \quad {}^{(n)}J_a^\lambda f(x, t) = \int_{\mathbb{R}^{n+1}} f(x - y, t - s) e^{iP_\lambda(y/a)} \frac{1}{a^n} \eta\left(\frac{y}{a}\right) \frac{1}{a^2} \zeta\left(\frac{s}{a^2}\right) dy ds.$$

The comparison to the operator ${}^{(n)}I_a^\lambda$ is clear if we write

$${}^{(n)}I_a^\lambda f(x, t) = \int_{\mathbb{R}^{n+1}} f(x - y, t - s) e^{iP_\lambda(y/a)} \frac{1}{a^n} \eta\left(\frac{y}{a}\right) \delta_{s=Q(y)} dy ds.$$

From now on, we will study the J -operator defined with a choice of bump function ζ such that

$$(4.3) \quad \int_{\mathbb{R}} \zeta(s) ds = 1.$$

This is useful, because if we temporarily let $I_a^\lambda(y, s)$ and $J_a^\lambda(y, s)$ represent the kernels of the operators, respectively, then for each fixed $\lambda \in \mathbb{R}^M$, $a > 0$ and $y \in \mathbb{R}^n$, we have

$$\int_{\mathbb{R}} (I_a^\lambda(y, s) - J_a^\lambda(y, s)) ds = 0.$$

(This will be used in (6.10) below, to introduce a derivative to a Littlewood-Paley projection, enabling the application of Lemma 6.3 (iii); this ultimately leads to (6.3).)

4.2. Littlewood-Paley decomposition. In order to introduce the square function built from differences of the I -operator (3.2) and the J -operator (4.2), we will employ a Littlewood-Paley decomposition constructed in [PY19]; we summarize its properties, all of which are proved explicitly in [PY19, §4]. There exists a family of Schwartz functions on \mathbb{R} , denoted $\Delta_j(t)$ for $j \in \mathbb{Z}$, with the following properties: $(\Delta_j)^\wedge(t)$ is supported in an annulus where $|t| \approx 2^{-2j}$,

$$(4.4) \quad \int_{\mathbb{R}} \Delta_j(t) dt = 0, \quad \text{for all } j,$$

and $\sum_j (\Delta_j)^\wedge(t) = 1$ for all $t \neq 0$. There is a second family of Schwartz functions on \mathbb{R} , denoted $\tilde{\Delta}_j$, with $(\tilde{\Delta}_j)^\wedge(t)$ supported on a slightly wider annulus $|t| \approx 2^{-2j}$ and with $(\tilde{\Delta}_j)^\wedge(t) \equiv 1$ on the support of $(\Delta_j)^\wedge(t)$ for each j . This family of Schwartz functions satisfies the analogue of (4.4) and $\sum_j (\tilde{\Delta}_j)^\wedge(t) = 1$ for all $t \neq 0$. Define the operator, also denoted Δ_j , acting on any Schwartz function f on \mathbb{R}^{n+1} by

$$\Delta_j f(x, t) = \int_{\mathbb{R}^n} f(x - y, t - s) \Delta_j(s) ds,$$

and define the operator $\tilde{\Delta}_j$ analogously.

For $P_j = \Delta_j$ or $P_j = \tilde{\Delta}_j$, for every $f \in L^2$,

$$f = \sum_{j=-\infty}^{\infty} P_j f,$$

with convergence of the partial sums holding in the L^2 sense, and

$$(4.5) \quad \left\| \left(\sum_{j=-\infty}^{\infty} |P_j f|^2 \right)^{1/2} \right\|_{L^2} \ll \|f\|_{L^2}.$$

Consequently, if we define the operator

$$L_N f = \sum_{|j| \leq N} \Delta_j \tilde{\Delta}_j f,$$

acting on f of Schwartz class, then $L_N f$ is a Schwartz function, and $L_N f$ converges to f in L^2 norm as $N \rightarrow \infty$. This concludes the summary of the properties cited from [PY19, §4].

We now fix N , so that all sums over j below are finite. Since the estimates below are independent of N , we can take $N \rightarrow \infty$ at the end of the argument. In particular,

$$\left\| \sup_{\substack{r \leq \|\lambda\| < 2r \\ k \in \mathbb{Z}}} |^{(\eta_k)} I_{2^k}^\lambda f \right\|_{L^2} \leq \left\| \sup_{\substack{r \leq \|\lambda\| < 2r \\ k \in \mathbb{Z}}} |^{(\eta_k)} I_{2^k}^\lambda L_N f \right\|_{L^2} + \left\| \sup_{\substack{r \leq \|\lambda\| < 2r \\ k \in \mathbb{Z}}} |^{(\eta_k)} I_{2^k}^\lambda (f - L_N f) \right\|_{L^2}.$$

By applying the trivial L^2 bound of (3.3) to the last term, and the fact that $L_N f$ converges to f in L^2 norm, we see that if we can prove

$$(4.6) \quad \left\| \sup_{\substack{r \leq \|\lambda\| < 2r \\ k \in \mathbb{Z}}} |^{(\eta_k)} I_{2^k}^\lambda L_N f \right\|_{L^2} \ll r^{-\delta} \|f\|_{L^2},$$

uniformly in N , then the key hypothesis (3.5) of Theorem 3.1 immediately follows. Thus our main goal is to prove (4.6), for all $r \geq 1$, uniformly in N .

4.3. The square function. Fix a family $\{\eta_k\}_{k \in \mathbb{Z}}$ of C^1 bump functions supported in the unit ball $B_1(\mathbb{R}^n)$ with C^1 norm uniformly bounded by 1. The key to proving (4.6) is a square function, defined each $r \geq 1$, for f of Schwartz class, by

$$S_r(f) = \left(\sum_{k \in \mathbb{Z}} \left(\sup_{r \leq \|\lambda\| < 2r} |^{(\eta_k)} I_{2^k}^\lambda - ^{(\eta_k)} J_{2^k}^\lambda \right) L_N f \right)^2)^{1/2}.$$

We aim to prove that there exists $\delta > 0$ such that for any Schwartz function f on \mathbb{R}^{n+1} , and any $r \geq 1$,

$$(4.7) \quad \|S_r(f)\|_{L^2(\mathbb{R}^{n+1})} \ll r^{-\delta} \|f\|_{L^2(\mathbb{R}^{n+1})},$$

uniformly in N . Let us see why this implies (4.6), and hence (3.5). First note that for any $r \geq 1$,

$$(4.8) \quad \sup_{\substack{r \leq \|\lambda\| < 2r \\ k \in \mathbb{Z}}} |^{(\eta_k)} I_{2^k}^\lambda L_N f| \leq \sup_{\substack{r \leq \|\lambda\| < 2r \\ k \in \mathbb{Z}}} |^{(\eta_k)} J_{2^k}^\lambda L_N f| + S_r(f).$$

Thus if we have proved (4.7), to deduce (4.6) it suffices to bound the first term on L^2 with decay in r . For this we cite the following, an immediate consequence of the work of Stein and Wainger in [SW01, Thm. 1]:

Theorem 4.1. *Let $p_1(y), \dots, p_M(y)$ be a fixed set of linearly independent, homogeneous polynomials in $\mathbb{R}[X_1, \dots, X_n]$, each of degree at least 2. For any family $\{\eta_k\}_{k \in \mathbb{Z}}$ of C^1 bump functions supported in the unit ball $B_1(\mathbb{R}^n)$ with C^1 norm uniformly bounded by 1, there exists a fixed $\delta > 0$ such that for any Schwartz function f on \mathbb{R}^{n+1} and any $r \geq 1$,*

$$(4.9) \quad \left\| \sup_{\substack{\lambda \in \mathbb{R}^M \\ r \leq \|\lambda\| < 2r}} \sup_{k \in \mathbb{Z}} |^{(\eta_k)} J_{2k}^\lambda f(x, t)| \right\|_{L^2(\mathbb{R}^{n+1})} \leq Ar^{-\delta} \|f\|_{L^2(\mathbb{R}^{n+1})},$$

in which the norm A may depend on p, n , and the fixed polynomials p_1, \dots, p_M .

Applying this theorem to the first term in (4.8), and (4.7) to the second term, then implies (4.6) and hence (3.5).

Theorem 4.1 follows from Stein and Wainger's original work with only a few minimal observations. The case we apply to prove Theorem 1.1 is explicitly proved in [PY19, Thm. 6.1], so we do not repeat the proof. (The same proof works verbatim for any $d_q \geq 1$ in the notation of (4.1).) We will, however, explicitly state an intermediate step from Stein and Wainger's proof, because we will apply it again in §6 (see (6.17)). We recall from [SW01] that their main strategy to prove Theorem 4.1 is to linearize and consider an operator $Tf(x, t) = ^{(\eta_k)} J_{2k}^\lambda f(x, t)$ for measurable stopping-time functions $k(x, t) : \mathbb{R}^{n+1} \rightarrow \mathbb{Z}$ and $\lambda(x, t) : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^M$. Then the key is to prove that TT^* is bounded on L^2 with a norm that decays in r , as long as $\lambda(x, t)$ takes values with $\|\lambda\| \approx r$. To prove this, Stein and Wainger compute the kernel of TT^* , and prove that it is compactly supported and has an upper bound that exhibits decay in r (in an appropriate sense).

More precisely, the kernel of TT^* takes the following form. Fix homogeneous polynomials p_1, \dots, p_M in $\mathbb{R}[X_1, \dots, X_n]$. Let $\Psi(u, z)$ be a C^1 bump function supported in $B_2 \times B_1 \subset \mathbb{R}^n \times \mathbb{R}^n$, and define

$$\|\Psi\|_{C^1} := \sup_{(u, z) \in B_2(\mathbb{R}^n) \times B_1(\mathbb{R}^n)} (|\Psi(u, z)| + |\nabla_z \Psi(u, z)|).$$

Define for stopping-times ν, μ taking values in \mathbb{R}^M the function

$$K_b^{\nu, \mu}(u) = \int_{\mathbb{R}^n} e^{iP_\nu(u+z) - iP_\mu(z)} \Psi(u, z) dz,$$

where $P_\nu(y) = \sum_{j=1}^M \nu_j p_j(y)$ and $P_\mu(y) = \sum_{j=1}^M \mu_j p_j(y)$. Trivially, $|K_b^{\nu, \mu}(u)| \ll \chi_{B_2}(u) \|\Psi\|_{C^1}$. The following nontrivial bound holds:

Proposition 4.2. *Let $p_1(y), \dots, p_M(y)$ be a fixed set of linearly independent, homogeneous polynomials in $\mathbb{R}[X_1, \dots, X_n]$, each of degree at least 2. Let $\Psi(u, z)$ be a C^1 bump function supported in $B_2 \times B_1 \subset \mathbb{R}^n \times \mathbb{R}^n$ with $\|\Psi\|_{C^1} \leq 1$. There exists a fixed $\delta > 0$ such that for any $r \geq 1$, if ν, μ satisfy*

$$r \leq \|\nu\|, \|\mu\| \leq 2r$$

then there exists a measurable set $G^\nu \subset B_2(\mathbb{R}^n)$ (which depends only on ν , and not on μ or Ψ), such that $|G^\nu| \ll r^{-\delta}$ and

$$|K_b^{\nu, \mu}(u)| \ll r^{-\delta} \chi_{B_2}(u) + \chi_{G^\nu}(u).$$

All implied constants are dependent only on n and the fixed polynomials p_1, \dots, p_M , and are independent of ν, μ, Ψ .

This is simply a reformulation of [SW01, Lemma 4.1] as stated in [PY19, Prop. 3.4]. It is crucial in Stein and Wainger's proof by the TT^* method that none of the polynomials p_j contains a linear term, and this restriction thus applies in Proposition 4.2 as well as in our main theorem.

4.4. Strategy to prove the square function estimate. We have established that to prove (4.6), the key is to prove (4.7). As a final result of this section, we record the strategy to prove the square function estimate (4.7):

Theorem 4.3. *Suppose that for any family $\{\eta_k\}_{k \in \mathbb{Z}}$ of C^1 bump functions supported in the unit ball $B_1(\mathbb{R}^n)$ with $\|\eta_k\|_{C^1} \leq 1$, there exists a fixed $\delta_0 > 0$ and $\varepsilon_0 > 0$ such that for any Schwartz function F on \mathbb{R}^{n+1} and any $r \geq 1$,*

$$(4.10) \quad \left\| \sup_{r \leq \|\lambda\| < 2r} |({}^{(\eta_k)} I_{2^k}^\lambda - {}^{(\eta_k)} J_{2^k}^\lambda) \Delta_j F| \right\|_{L^2} \ll r^{-\delta_0} 2^{-\varepsilon_0 |j-k|} \|F\|, \quad \text{for all } j, k \in \mathbb{Z}.$$

Then for any such family $\{\eta_k\}$, there exists $\delta > 0$ such that for any Schwartz function f on \mathbb{R}^{n+1} , and any $r \geq 1$,

$$(4.11) \quad \|S_r(f)\|_{L^2(\mathbb{R}^{n+1})} \ll r^{-\delta} \|f\|_{L^2(\mathbb{R}^{n+1})},$$

uniformly in N , and hence (3.5) holds.

Indeed, we recall that

$$|S_r(f)|^2 \leq \sum_{k \in \mathbb{Z}} \left(\sum_{|j| \leq N} \sup_{\|\lambda\| \approx r} |({}^{(\eta_k)} I_{2^k}^\lambda - {}^{(\eta_k)} J_{2^k}^\lambda) \Delta_j \tilde{\Delta}_j f| \right)^2.$$

By inserting $2^{-\varepsilon_1 |j-k|} 2^{+\varepsilon_1 |j-k|}$ for some $\varepsilon_1 < \varepsilon_0$, and applying Cauchy-Schwarz to the sum over j , we see that

$$|S_r(f)|^2 \leq \sum_{k \in \mathbb{Z}} \left(\sum_{|j| \leq N} 2^{-2\varepsilon_1 |j-k|} \right) \left(\sum_{|j| \leq N} 2^{2\varepsilon_1 |j-k|} \sup_{\|\lambda\| \approx r} |({}^{(\eta_k)} I_{2^k}^\lambda - {}^{(\eta_k)} J_{2^k}^\lambda) \Delta_j \tilde{\Delta}_j f|^2 \right).$$

The first sum over j is $\ll 1$ uniformly in k . Taking L^2 norms, we apply the hypothesis (4.10) to each $F = \tilde{\Delta}_j f$ to see that

$$\|S_r(f)\|_{L^2}^2 \ll r^{-2\delta_0} \sum_{k \in \mathbb{Z}} \sum_{|j| \leq N} 2^{2\varepsilon_1 |j-k|} 2^{-2\varepsilon_0 |j-k|} \|\tilde{\Delta}_j f\|_{L^2}^2.$$

We use the fact that $\varepsilon_1 < \varepsilon_0$ to sum over k , and apply the property (4.5) of the projections $\tilde{\Delta}_j$ to conclude that $\|S_r(f)\|_{L^2}^2 \ll r^{-2\delta_0} \|f\|_{L^2}^2$, and the theorem is proved.

The rest of the paper will focus on proving the hypothesis (4.10). Note that we now need to prove (4.10) for each fixed k rather than for an operator involving a supremum over k ; fixing k is a key advantage we have gained by working with the square function. (The later utility of having k be fixed is visible in (6.13), and subsequent similar places.)

5. STEP 3: CHANGE OF VARIABLES TO ISOLATE THE KERNEL $K_{\sharp, l}^{\nu, \mu}$

Our strategy to prove the main hypothesis (4.10) in Theorem 4.3 is to apply the TT^* method to an operator T that is (roughly) of the form

$$f \mapsto \sup_{r \leq \|\lambda\| < 2r} |({}^{(\eta_k)} I_{2^k}^\lambda - {}^{(\eta_k)} J_{2^k}^\lambda) \Delta_j f|.$$

To do so precisely, we will linearize this operator by replacing the supremum over λ by a stopping-time function $\lambda(x, t)$ taking values with $\|\lambda(x, t)\| \in [r, 2r)$, and then we will compute $TT^* f(x, t)$ for a Schwartz function f . Our goal is to write $TT^* f(x, t)$ as a convolution of f with a kernel, say K , that is itself defined by an oscillatory integral.

Remark 5.1. Since $TT^* f(x, t)$ is an integral over \mathbb{R}^{2n} , and a convolution of $f(x, t)$ with a kernel is an integral over $n+1$ variables, at least formally we would expect K to be an integral over $n-1$ variables. This is ultimately why the present paper is limited to the consideration of $n \geq 2$.

From now on, the fact that we work with $\gamma(y) = (y, Q(y))$ with $Q(y)$ quadratic becomes crucial to our method, and in particular we assume from now on that

$$(5.1) \quad Q(y) = \sum_{i=1}^n \theta_i y_i^2, \quad \theta_i \in \{\pm 1\}.$$

Recall from §2 that this suffices to prove Theorem 1.1.

At least intuitively, if Tf integrates $f(x - y, t - Q(y))$ over $y \in \mathbb{R}^n$, then we expect $TT^*f(x, t)$ to integrate $f(x - (y - z), t - (Q(y) - Q(z)))$ over $y \in \mathbb{R}^n, z \in \mathbb{R}^n$. It is then natural to set $u = y - z$, so we face an integral of $f(x - u, t - (Q(u + z) - Q(z)))$ over $u \in \mathbb{R}^n, z \in \mathbb{R}^n$. To isolate the kernel K , we must “free” $n - 1$ variables so that only $n + 1$ coordinates appear in the argument of f , that is, so that $Q(u + z) - Q(z)$ only involves $n + 1$ coordinates. This motivates the change of variables we now describe, which crucially uses the quadratic behavior of $\gamma(y)$. It is an interesting open problem to develop an argument for more general hypersurfaces.

For Q as in (5.1), define for any $u \in \mathbb{R}^n$, that

$$\tilde{u} = (\theta_1 u_1, \dots, \theta_n u_n),$$

where the signs θ_i are determined once and for all by (5.1). This notation will be used extensively for the remainder of the paper. Then define

$$\langle u, z \rangle = \sum_{i=1}^n \theta_i u_i z_i = \tilde{u} \cdot z.$$

Consequently,

$$Q(u + z) - Q(z) = Q(u) + 2|u| \frac{\langle u, z \rangle}{|u|}.$$

This motivates us to replace z by defining new coordinates τ, σ , where $\tau \in \mathbb{R}$, with $\tau = \langle u, z \rangle / |u|$, and $\sigma \in \mathbb{R}^{n-1}$. To define σ appropriately, we require a notational convention: for $z \in \mathbb{R}^n$ and $l \in \{1, \dots, n\}$, define

$$z^{(l)} = (z_1, \dots, \hat{z}_l, \dots, z_n) \in \mathbb{R}^{n-1}, \quad \text{that is, omitting the } l\text{-th coordinate.}$$

Now for a fixed $l \in \{1, \dots, n\}$ we can define the change of variables $z \mapsto (\tau, \sigma) \in \mathbb{R} \times \mathbb{R}^{n-1}$ by

$$(5.2) \quad \tau = \frac{\langle u, z \rangle}{|u|} \quad \text{and} \quad \sigma = \frac{\tau \tilde{u}^{(l)} - |u| z^{(l)}}{\theta_l u_l}.$$

The formula for z in terms of (τ, σ) is

$$(5.3) \quad z^{(l)} = \frac{\tau \tilde{u}^{(l)} - \theta_l u_l \sigma}{|u|} \quad \text{and} \quad z_l = \frac{\theta_l u_l \tau + \tilde{u}^{(l)} \cdot \sigma}{|u|},$$

or equivalently,

$$\begin{bmatrix} z^{(l)} \\ z_l \end{bmatrix} = \frac{1}{|u|} \begin{bmatrix} -\theta_l u_l \text{Id}_{n-1} & \tilde{u}^{(l)} \\ (\tilde{u}^{(l)})^T & \theta_l u_l \end{bmatrix} \begin{bmatrix} \sigma \\ \tau \end{bmatrix}.$$

It will be important in what follows that if (u, z) lies in $B_2 \times B_1 \subset \mathbb{R}^n \times \mathbb{R}^n$, then $(\tau, \sigma) \in \mathbb{R}^n$ also lies in a compact region. To achieve this, we will use the fact that for any $u \in \mathbb{R}^n$, we can choose for which l we perform the above change of variables. That is, we fix a small $c_0 > 0$ once and for all, and choose a partition of unity

$$(5.4) \quad 1 = \sum_{1 \leq l \leq n} W_l(s), \quad s \in S^{n-1},$$

where each $W_l \in C_c^\infty(S^{n-1})$ is supported where $|s_l|/|s| \geq c_0$. For u such that $u/|u|$ lies in the support of W_l , it follows that $|u_l|/|u| \geq c_0$, and we can apply the change of variables (5.2) for this choice l .

5.1. Definition of $K_{\sharp,l}^{\nu,\mu}$ kernel. Now, with this key change of variables in mind, we can define the oscillatory integral that is the heart of the matter. Let $\Psi(u, z)$ be a C^1 bump function supported in $B_2 \times B_1 \subset \mathbb{R}^n \times \mathbb{R}^n$. Define, for $u \in \mathbb{R}^n$ and for $1 \leq l \leq n$ such that $u/|u|$ lies in the support of W_l , the oscillatory integral

$$(5.5) \quad K_{\sharp,l}^{\nu,\mu}(u, \tau) = \int_{\mathbb{R}^{n-1}} e^{iP_\nu(u+z) - iP_\mu(z)} \Psi(u, z) d\sigma,$$

where z is defined implicitly in terms of u, τ, σ as in (5.2). (For $u \in B_2$ such that $u/|u|$ lies outside the support of W_l , by convention we set $K_{\sharp,l}^{\nu,\mu}(u, \tau) \equiv 0$ for $\tau \in B_1$, so the upper bounds we prove below for $K_{\sharp,l}^{\nu,\mu}$ will hold for all $(u, \tau) \in B_2 \times B_1$.) Here ν, μ are stopping-times taking values in \mathbb{R}^M , which will be described momentarily. Given a function $\Psi(u, z)$ as above, we use the notation

$$(5.6) \quad \|\Psi\|_{C^1(\sigma)} := \sup_{(u,z) \in B_2(\mathbb{R}^n) \times B_1(\mathbb{R}^n)} (|\Psi(u, z)| + |\nabla_\sigma \Psi(u, z)|),$$

which is well-defined, for z defined in terms of u, τ, σ by (5.2).

We now check that if u, z are each compactly supported in $B_2 \times B_1 \subset \mathbb{R}^n \times \mathbb{R}^n$ and $u/|u|$ lies in the support of W_l , then $K_{\sharp,l}^{\nu,\mu}(u, \tau)$ is compactly supported with respect to $(u, \tau) \in \mathbb{R}^n \times \mathbb{R}$, and the region of integration over $\sigma \in \mathbb{R}^{n-1}$ in (5.5) is compactly supported. To see this, note that

$$\begin{aligned} |z|^2 &= |z^{(l)}|^2 + z_l^2 = |u|^{-2} [|\tau \tilde{u}^{(l)} - \theta_l u_l \sigma|^2 + |\theta_l u_l \tau + \tilde{u}^{(l)} \cdot \sigma|^2] \\ &= |u|^{-2} \left[\sum_{i \neq l} (\tau \theta_i u_i - \theta_l u_l \sigma_i)^2 + (\tau \theta_l u_l + \tilde{u}^{(l)} \cdot \sigma)^2 \right] \\ &= |u|^{-2} \left[\sum_{i \neq l} (\tau^2 u_i^2 - 2\tau \theta_i u_i \theta_l u_l \sigma_i + u_l^2 \sigma_i^2) + \tau^2 u_l^2 + 2\tau \theta_l u_l (\tilde{u}^{(l)} \cdot \sigma) + (\tilde{u}^{(l)} \cdot \sigma)^2 \right]. \end{aligned}$$

We conclude that

$$|z|^2 = \tau^2 + (u_l/|u|)^2 |\sigma|^2 + |u|^{-2} (\tilde{u}^{(l)} \cdot \sigma)^2.$$

Since all terms in this relation are non-negative, it follows that $\tau^2 \leq |z|^2 \leq 1$. We also see that

$$(5.7) \quad |\sigma|^2 \leq (u_l/|u|)^{-2} \leq c_0^{-2},$$

since $u/|u|$ is in the support of W_l . In particular, since $\Psi(u, z)$ has support in $B_2 \times B_1 \subset \mathbb{R}^n \times \mathbb{R}^n$, this proves the trivial upper bound

$$(5.8) \quad |K_{\sharp,l}^{\nu,\mu}(u, \tau)| \ll_{c_0} \chi_{B_2}(u) \chi_{B_1}(\tau).$$

Once we prove a far stronger bound for $K_{\sharp,l}^{\nu,\mu}$, one that exhibits decay in r when $\|\mu\|, \|\nu\| \approx r$, then we can complete the proof of the key hypothesis of Theorem 4.3, and hence complete the proof of the main theorem of this paper. This deduction is the content of the next section.

6. STEP 4: REDUCTION OF THE SQUARE FUNCTION ESTIMATE TO BOUNDING $K_{\sharp,l}^{\nu,\mu}$

The main result of this section is the following theorem:

Theorem 6.1. *Let $\Psi(u, z)$ be a C^1 function supported on $B_2 \times B_1 \subset \mathbb{R}^n \times \mathbb{R}^n$, with $\|\Psi\|_{C^1(\sigma)} \leq 1$ as in (5.6). For each $1 \leq l \leq n$, for $u \in \mathbb{R}^n$ such that $u/|u|$ lies in the support of W_l , consider the kernel $K_{\sharp,l}^{\nu,\mu}$ as defined in (5.5). Suppose that there exists a small $\delta > 0$ such that for all $r \geq 1$, if the stopping times ν, μ satisfy*

$$r \leq \|\nu\|, \|\mu\| \leq 2r,$$

then there exists a measurable set $G^\nu \subset B_2(\mathbb{R}^n)$ (depending on ν but independent of μ, r, Ψ), and for each $u \in B_2(\mathbb{R}^n)$ there exists a measurable set $F_u^\nu \subset B_1(\mathbb{R})$ (depending on u and ν but independent of μ, r, Ψ), such that

$$(6.1) \quad |G^\nu| \ll r^{-\delta}, \quad |F_u^\nu| \ll r^{-\delta},$$

and

$$(6.2) \quad |K_{\sharp,l}^{\nu,\mu}(u, \tau)| \ll r^{-\delta} \chi_{B_2}(u) \chi_{B_1}(\tau) + \chi_{G^\nu}(u) \chi_{B_1}(\tau) + \chi_{B_2}(u) \chi_{F_u^\nu}(\tau),$$

in which all implicit constants depend only on n and the fixed polynomials p_1, \dots, p_M . Then the hypothesis (4.10) of Theorem 4.3 holds.

Remark 6.2. In the proof of this theorem, it is crucial that in the hypothesis, the sets in (6.1) and (6.2) are completely independent of μ , although they are allowed to depend on ν . This is because when we apply these bounds within the proof of Theorem 6.1 in the steps (6.11) and (6.22) below, $\nu = \lambda(x, t)$ does not depend on variables of integration, while $\mu = \lambda(x - u, t - \theta)$ does. Consequently, a dependence on ν can be controlled by an averaging operator (Lemma 6.4), while a dependence on μ cannot. Proving that such sets G^ν and F_u^ν do exist, independent of μ , was a key difficulty in the previous work [PY19] and is also the main difficulty of this paper; it is handled in Part II when we prove Theorem 7.1.

After we prove Theorem 6.1 in this section, all that remains to complete the proof of our main theorem is to show that the estimate (6.2) for $K_{\sharp,l}^{\nu,\mu}$ holds; this we accomplish in Part II of the paper.

To prove that (4.10) holds, as claimed in Theorem 6.1, it suffices to prove three bounds, under the hypotheses of the theorem:

$$(6.3) \quad \left\| \sup_{\|\lambda\| \approx r} |({}^{(\eta_k)} I_{2^k}^\lambda - {}^{(\eta_k)} J_{2^k}^\lambda) \Delta_j F| \right\|_{L^2} \ll r^{-\delta_0} 2^{-\varepsilon_0(j-k)} \|F\|_{L^2} \quad \text{for } j \geq k$$

$$(6.4) \quad \left\| \sup_{\|\lambda\| \approx r} |({}^{(\eta_k)} I_{2^k}^\lambda \Delta_j F| \right\|_{L^2} \ll r^{-\delta_0} 2^{\varepsilon_0(j-k)} \|F\|_{L^2} \quad \text{for } j < k$$

$$(6.5) \quad \left\| \sup_{\|\lambda\| \approx r} |({}^{(\eta_k)} J_{2^k}^\lambda \Delta_j F| \right\|_{L^2} \ll r^{-\delta_0} 2^{\varepsilon_0(j-k)} \|F\|_{L^2} \quad \text{for } j < k.$$

The bound (6.5) for the smoother ‘‘flat’’ operator $J_{2^k}^\lambda$ has already been proved in full by [PY19, Prop. 7.2]. We will prove (6.3) and (6.4) by the method of TT^* . While we give a complete proof, we do not elaborate on the motivation for each step, for which we refer the reader to the more extensive exposition in [PY19].

6.1. Computing the kernel of TT^* to prove (6.3). Fix $r \geq 1$. Fix $k \in \mathbb{Z}$ and set $a = 2^k$. We also fix a bump function η_k , and for simplicity suppress it in the notation, as follows. Let $\lambda(x, t)$ be a measurable stopping-time function that takes values in $r \leq \lambda(x, t) \leq 2r$. For each $j \in \mathbb{Z}$, define the operator

$$T = ({}^{(\eta_k)} I_{2^k}^\lambda - {}^{(\eta_k)} J_{2^k}^\lambda) \Delta_j = (I_{2^k}^\lambda - J_{2^k}^\lambda) \Delta_j,$$

so that

$$Tf(x, t) = \int_{\mathbb{R}^{n+2}} f(x - y, t - \theta) e^{iP_{\lambda(x,t)}(\frac{y}{a})} \frac{1}{a^n} \eta\left(\frac{y}{a}\right) \left[\delta_{s=Q(y)} - \frac{1}{a^2} \zeta\left(\frac{s}{a^2}\right) \right] \Delta_j(\theta - s) dy ds d\theta.$$

The dual T^* is

$$T^*g(x, t) = \int_{\mathbb{R}^{n+2}} g(x + z, t + \omega) e^{-iP_{\lambda(x+z,t+\omega)}(\frac{z}{a})} \frac{1}{a^n} \eta\left(\frac{z}{a}\right) \left[\delta_{\xi=Q(z)} - \frac{1}{a^2} \zeta\left(\frac{\xi}{a^2}\right) \right] \Delta_j(\omega - \xi) dz d\xi d\omega.$$

It then follows that

$$\begin{aligned} TT^*f(x, t) &= \int_{\mathbb{R}^{2n+4}} f(x - y + z, t - \theta + \omega) e^{i[P_{\lambda(x,t)}(\frac{y}{a}) - P_{\lambda(x+z-y,t-\theta+\omega)}(\frac{z}{a})]} \frac{1}{a^{2n}} \eta\left(\frac{y}{a}\right) \eta\left(\frac{z}{a}\right) \\ &\quad \cdot \left[\delta_{s=Q(y)} - \frac{1}{a^2} \zeta\left(\frac{s}{a^2}\right) \right] \left[\delta_{\xi=Q(z)} - \frac{1}{a^2} \zeta\left(\frac{\xi}{a^2}\right) \right] \Delta_j(\theta - s) \Delta_j(\omega - \xi) dz d\xi d\omega dy ds d\theta. \end{aligned}$$

As a first step, to verify (6.3) under the hypotheses of Theorem 6.1, we need to show that

$$(6.6) \quad \|TT^*f\|_{L^2} \ll r^{-2\delta_0} 2^{-2\varepsilon_0|j-k|} \|f\|_{L^2}.$$

To study TT^* , we require further Littlewood-Paley projections closely related to Δ_j by taking convolutions, antiderivatives, or derivatives. We quote the key properties from [PY19, Lemmas 4.3, 4.4, 4.5], which we summarize as a lemma:

Lemma 6.3. (i) *Convolutions: Define*

$$\underline{\Delta}_j(t) = \int_{\mathbb{R}} \Delta_j(w+t)\Delta_j(w)dw.$$

For each $j \in \mathbb{Z}$,

$$(6.7) \quad \int_{\mathbb{R}} \underline{\Delta}_j(t)dt = 0.$$

For each $a = 2^k$, $\underline{\Delta}_j(t) = \frac{1}{a^2}\underline{\Delta}_{j-k}(\frac{t}{a^2})$. Uniformly in j , $\|\underline{\Delta}_j\|_{L^1} \ll 1$.

(ii) *Antiderivatives: There exist Schwartz functions $\tilde{\Delta}$ and $\tilde{\underline{\Delta}}$ on \mathbb{R} such that upon defining $\tilde{\Delta}_j(t) = 2^{-2j}\tilde{\Delta}(2^{-2j}t)$ and $\tilde{\underline{\Delta}}_j(t) = 2^{-2j}\tilde{\underline{\Delta}}(2^{-2j}t)$, then*

$$(6.8) \quad \Delta_j(t) = 2^{2j}\left(\frac{d}{dt}\tilde{\Delta}_j\right)(t), \quad \underline{\Delta}_j(t) = 2^{2j}\left(\frac{d}{dt}\tilde{\underline{\Delta}}_j\right)(t), \quad j \in \mathbb{Z}.$$

Finally, $\|\tilde{\Delta}_j\|_{L^1} \ll 1$ and $\|\tilde{\underline{\Delta}}_j\|_{L^1} \ll 1$ uniformly in j .

(iii) *Mean value: Define $\psi(t) = (1+t^2)^{-1}$. For each $j \in \mathbb{Z}$ define $\psi_j(t) = 2^{-2j}\psi(2^{-2j}t)$, so that ψ_j is a non-negative function on \mathbb{R} with $\|\psi_j\|_{L^1} \ll 1$, uniformly in j . Then $|\underline{\Delta}'(t)| \ll 2^{-2j}\psi_j(t)$, and for any $|\xi| \leq 2$, $|\underline{\Delta}_j(t+\xi) - \underline{\Delta}_j(t)| \ll 2^{-2j}|\xi|\psi_j(t)$.*

The dilations employed in the lemma are intended to be compatible with parabolic rescaling. We will apply (iii) in §6.2 and §6.3 to prove (6.3), and (i) and (ii) in §6.4 to prove (6.4).

A short computation using the definition of $\underline{\Delta}_{j-k}$ (entirely analogous to [PY19, (7.10)]), shows that the operator TT^* given above can be written as

$$(6.9) \quad TT^*f(x,t) = \int_{\mathbb{R}^{n+1}} f(x-u, t-\theta) \frac{1}{a^{n+2}} K^{\lambda(x,t), \lambda(x-u, t-\theta)} \left(\frac{u}{a}, \frac{\theta}{a^2}\right) dud\theta$$

where for each $\nu, \mu \in \mathbb{R}^{d-1}$, we define the kernel

$$\begin{aligned} K^{\nu, \mu}(u, \theta) &= \int_{\mathbb{R}^{n+2}} e^{iP_\nu(u+z) - iP_\mu(z)} \eta(u+z)\eta(z) \\ &\quad \cdot [\delta_{s=Q(u+z)} - \zeta(s)] [\delta_{\xi=Q(z)} - \zeta(\xi)] \underline{\Delta}_{j-k}(\theta - s + \xi) dz d\xi ds \\ &= \int_{\mathbb{R}^{n+1}} e^{iP_\nu(u+z) - iP_\mu(z)} \eta(u+z)\eta(z) [\delta_{\xi=Q(z)} - \zeta(\xi)] \underline{\Delta}_{j-k}(\theta - Q(u+z) + \xi) dz d\xi \\ &\quad + \int_{\mathbb{R}^{n+2}} e^{iP_\nu(u+z) - iP_\mu(z)} \eta(u+z)\eta(z) \zeta(s) [\delta_{\xi=Q(z)} - \zeta(\xi)] \underline{\Delta}_{j-k}(\theta - s + \xi) dz d\xi ds \\ &=: \mathbf{I} + \mathbf{II}. \end{aligned}$$

To verify (6.6), it suffices to prove the bound separately, for the contribution of the term \mathbf{I} to the kernel, and the contribution of \mathbf{II} .

6.2. The contribution of the term \mathbf{I} . To analyze the contribution of \mathbf{I} , we write

$$\mathbf{I} = \sum_{l=1}^n \mathbf{I}_l$$

where for each $1 \leq l \leq n$,

$$\begin{aligned} \mathbf{I}_l &= W_l \left(\frac{u}{|u|} \right) \int_{\mathbb{R}^n} e^{iP_\nu(u+z) - iP_\mu(z)} \eta(u+z) \eta(z) \underline{\Delta}_{j-k}(\theta - Q(u+z) + Q(z)) dz \\ &\quad - W_l \left(\frac{u}{|u|} \right) \int_{\mathbb{R}^{n+1}} e^{iP_\nu(u+z) - iP_\mu(z)} \eta(u+z) \eta(z) \zeta(\xi) \underline{\Delta}_{j-k}(\theta - Q(u+z) + \xi) dz d\xi. \end{aligned}$$

Using the fact that $\int \zeta(\xi + Q(z)) d\xi = 1$, this can be re-written as

$$(6.10) \quad \mathbf{I}_l = W_l \left(\frac{u}{|u|} \right) \int_{\mathbb{R}^{n+1}} e^{iP_\nu(u+z) - iP_\mu(z)} \eta(u+z) \eta(z) \zeta(\xi + Q(z)) \cdot (\underline{\Delta}_{j-k}(\theta - Q(u+z) + Q(z)) - \underline{\Delta}_{j-k}(\theta - Q(u+z) + Q(z) + \xi)) dz d\xi.$$

For each fixed $1 \leq l \leq n$, we make the l -th change of variables (5.2), so that

$$\mathbf{I}_l = \left(\frac{|u_l|}{|u|} \right)^{n-2} W_l \left(\frac{u}{|u|} \right) \int_{\mathbb{R}^2} K_{\sharp, l}^{\nu, \mu}(u, \tau; \xi) (\underline{\Delta}_{j-k}(\theta - Q(u) - 2|u|\tau) - \underline{\Delta}_{j-k}(\theta - Q(u) - 2|u|\tau + \xi)) d\tau d\xi,$$

in which

$$(6.11) \quad K_{\sharp, l}^{\nu, \mu}(u, \tau; \xi) = \int_{\mathbb{R}^{n-1}} e^{iP_\nu(u+z) - iP_\mu(z)} \eta(u+z) \eta(z) \zeta(\xi + Q(z)) d\sigma,$$

where z is defined implicitly in terms of u, τ, σ , and we recall from (6.9) that

$$(6.12) \quad \nu = \lambda(x, t), \quad \mu = \lambda(x - u, t - \theta)$$

are stopping-times with values such that $r \leq \|\nu\|, \|\mu\| < 2r$. Note that the ξ -support is in $|\xi| \leq 2$. By the mean-value property of Lemma 6.3 (iii),

$$|\mathbf{I}_l| \ll 2^{-2(j-k)} \chi_{B_2}(u) \int_{\mathbb{R}^2} |K_{\sharp, l}^{\nu, \mu}(u, \tau; \xi)| \chi_{B_1}(\tau) \chi_{B_2}(\xi) \psi_{j-k}(\theta - Q(u) - 2|u|\tau) d\tau d\xi.$$

For each fixed $|\xi| \leq 2$, $K_{\sharp, l}^{\nu, \mu}(u, \tau; \xi)$ is a kernel of the form (5.5) for

$$\Psi(u, z) = \Psi_\xi(u, z) = \eta(u+z) \eta(z) \zeta(\xi + Q(z)).$$

We now apply the hypothesis of Theorem 6.1 to bound $K_{\sharp, l}^{\nu, \mu}(u, \tau; \xi)$ as in (6.2). Note that the resulting bound is uniform in ξ , and so we can then integrate trivially over $\xi \in B_2(\mathbb{R})$. We conclude that the contribution of \mathbf{I}_l to TT^*f in (6.9) is bounded by

$$(6.13) \quad \ll 2^{-2(j-k)} \iint_{\mathbb{R}^{n+1}} |f(x - u, t - \theta)| \chi((x, t), (u, \theta)) du d\theta$$

in which we define the function $\chi((x, t), (u, \theta))$ to be

$$\int_{\mathbb{R}} \frac{1}{a^n} \left\{ r^{-\delta} \chi_{B_2}\left(\frac{u}{a}\right) \chi_{B_1}(\tau) + \chi_{G^{\lambda(x, t)}}\left(\frac{u}{a}\right) \chi_{B_1}(\tau) + \chi_{B_2}\left(\frac{u}{a}\right) \chi_{F_{\left(\frac{u}{a}\right)}^{\lambda(x, t)}}(\tau) \right\} \frac{1}{a^2} \psi_{j-k} \left(\frac{\theta - Q(u) - 2a|u|\tau}{a^2} \right) d\tau.$$

We recall that $a = 2^k$ is fixed, so that (6.13) is an averaging operator, not a maximal operator. Next we will apply [PY19, Lemma 7.5], which we quote:

Lemma 6.4. *Let $\chi(w, y)$ be an integrable function on $\mathbb{R}^m \times \mathbb{R}^m$ with*

$$(6.14) \quad \left\| \sup_{w \in \mathbb{R}^m} |\chi(w, y)| \right\|_{L^1(\mathbb{R}^m(dy))} \leq C, \quad \sup_{w \in \mathbb{R}^m} \|\chi(w, y)\|_{L^1(\mathbb{R}^m(dy))} \leq \lambda.$$

Then

$$\left\| \int_{\mathbb{R}^m} f(w - y) \chi(w, y) dy \right\|_{L^p(\mathbb{R}^m(dw))} \ll_{C, p} \lambda^{1-1/p} \|f\|_{L^p(\mathbb{R}^m)}, \quad 1 \leq p \leq \infty.$$

To verify the first condition in (6.14) for our choice of $\chi((x, t), (u, \theta))$ above, we simply note that by the hypothesis (6.1), the small sets satisfy $G^{\lambda(x, t)} \subset B_2(\mathbb{R}^n)$ and $F_{(\frac{u}{a})}^{\lambda(x, t)} \subset B_1(\mathbb{R})$ for all (x, t) , so that

$$(6.15) \quad \sup_{(x, t)} |\chi((x, t), (u, \theta))| \ll \int_{\mathbb{R}} \frac{1}{a^n} \chi_{B_2}\left(\frac{u}{a}\right) \chi_{B_1}(\tau) \frac{1}{a^2} \psi_{j-k} \left(\frac{\theta - Q(u) - 2a|u|\tau}{a^2} \right) d\tau.$$

We compute the $L^1(dud\theta)$ norm of the above function of (u, θ) by integrating first in θ , and we see that the first condition in (6.14) holds, uniformly in j, k . For the second condition in (6.14), we will use the small measures of the sets $G^{\lambda(x, t)}$ and $F_{(\frac{u}{a})}^{\lambda(x, t)}$. For each fixed $(x, t) \in \mathbb{R}^{n+1}$, we compute the $L^1(dud\theta)$ norm of $\chi((x, t), (u, \theta))$ by integrating in θ, τ, u (in that order), and the second condition holds with $\lambda = r^{-\delta}$, by the hypothesis (6.1). Hence applying Lemma 6.4 with $\lambda = r^{-\delta}$ and $p = 2$, shows that the $L^2(dxdt)$ norm of the operator (6.13) is $\ll r^{-\delta/2} 2^{-2(j-k)}$. This proves the conclusion of (6.3) for the contribution of \mathbf{I}_l , with $\delta_0 = \delta/4$ and $\varepsilon_0 = 1$. Adding the contributions for $1 \leq l \leq n$ thus proves the conclusion of (6.3) for the contribution of \mathbf{I} .

6.3. The contribution of the term \mathbf{II} . To analyze the contribution of \mathbf{II} to the kernel $K^{\nu, \mu}(u, \theta)$ of TT^* in (6.9), the argument is similar, but simpler, since we do not require the change of variables (5.2) to study this term. A brief argument using $\int_{\mathbb{R}} \zeta(\xi + Q(z)) d\xi = 1$ and linear changes of variables (entirely analogous to [PY19, §7.3.2]) shows that we may write the contribution of \mathbf{II} to the kernel $K^{\nu, \mu}(u, \theta)$ as

$$(6.16) \quad \mathbf{II} = \iint_{\mathbb{R}^{n+2}} K_b^{\lambda(x, t), \lambda(x-u, t-\theta)}(u; \xi, s) (\underline{\Delta}_{j-k}(\theta - s) - \underline{\Delta}_{j-k}(\theta - s + \xi)) ds d\xi,$$

where we have defined for any $\nu, \mu \in \mathbb{R}^{d-1}$ the function

$$K_b^{\nu, \mu}(u; \xi, s) = \int_{\mathbb{R}^n} e^{iP_\nu(u+z) - iP_\mu(z)} \eta(u+z) \eta(z) \zeta(\xi + Q(z)) \zeta(s + Q(z)) dz.$$

Since η and ζ are supported in $B_1(\mathbb{R}^n)$, it follows that the integral is over $z \in B_1(\mathbb{R}^n)$, and $K_b^{\nu, \mu}(u; \xi, s)$ is supported where $u \in B_2(\mathbb{R}^n)$, $\xi \in B_2(\mathbb{R})$, $s \in B_2(\mathbb{R})$. We apply the mean value of Lemma 6.3 (iii) to bound the difference of $\underline{\Delta}_{j-k}$ terms in (6.16) by $\ll 2^{-2(j-k)} |\xi| \psi_{j-k}(\theta - s)$. We then apply Proposition 4.2 to bound $K_b^{\nu, \mu}(u; \xi, s)$, uniformly in $\xi, s \in B_2(\mathbb{R})$, and integrate trivially over $\xi \in B_2(\mathbb{R})$. Consequently, we see that the contribution of \mathbf{II} to $TT^*f(x, t)$ in (6.9) is bounded by

$$(6.17) \quad \ll 2^{-2(j-k)} \iint_{\mathbb{R}^{n+1}} |f(x-u, t-\theta)| \chi((x, t), (u, \theta)) dud\theta$$

in which we define the function $\chi((x, t), (u, \theta))$ to be

$$\chi((x, t), (u, \theta)) = \frac{1}{a^n} (r^{-\delta} \chi_{B_2}\left(\frac{u}{a}\right) + \chi_{G^{\lambda(x, t)}}\left(\frac{u}{a}\right)) \frac{1}{a^2} \int_{\mathbb{R}} \psi_{j-k}\left(\frac{\theta}{a^2} - s\right) \chi_{B_2}(s) ds.$$

We will again apply Lemma 6.4 to bound the $L^2(\mathbb{R}^{n+1})$ norm of the averaging operator (6.17). The first condition in (6.14) holds because

$$\sup_{(x, t)} |\chi((x, t), (u, \theta))| \leq \frac{1}{a^n} \chi_{B_2}\left(\frac{u}{a}\right) \frac{1}{a^2} \int_{\mathbb{R}} \psi_{j-k}\left(\frac{\theta}{a^2} - s\right) \chi_{B_2}(s) ds.$$

This has bounded $L^1(dud\theta)$ norm, uniformly in j, k , by integrating first in θ (using Lemma 6.3 (iii)), then in s and u . The second condition in (6.14) is verified by using the fact that for each fixed (x, t) , $|G^{\lambda(x, t)}| \leq r^{-\delta}$ by Proposition 4.2, so that uniformly in j, k

$$\sup_{(x, t)} \|\chi((x, t), (u, \theta))\|_{L^1(dud\theta)} \ll r^{-\delta}.$$

Then applying Lemma 6.4 with $\lambda = r^{-\delta}$ and $p = 2$ shows that the $L^2(dx, dt)$ norm of the operator (6.17) is $\ll r^{-\delta/2}2^{-2(j-k)}$, so that the L^2 norm of the contribution of \mathbf{II} to (6.9) is $\ll r^{-\delta/2}2^{-2(j-k)}$. This completes the proof of (6.3), with $\delta_0 = \delta/4$ and $\varepsilon_0 = 1$.

6.4. Proof of (6.4). The last step to complete the proof of Theorem 6.1 is to verify the L^2 bound in (6.4) in the case $j < k$. It suffices to prove two bounds: first, a bound with decay in r ,

$$(6.18) \quad \left\| \sup_{\|\lambda\| \approx r} |I_{2^k}^\lambda \Delta_j F| \right\|_{L^2(\mathbb{R}^{n+1})} \ll r^{-\delta_0} \|F\|_{L^2(\mathbb{R}^{n+1})}, \quad j < k, \quad \text{some } \delta_0 > 0,$$

and second, a bound with acceptable growth in r and a decay factor 2^{j-k} ,

$$(6.19) \quad \left\| \sup_{\|\lambda\| \approx r} |I_{2^k}^\lambda \Delta_j F| \right\|_{L^2(\mathbb{R}^{n+1})} \ll r^{1/2} 2^{j-k} \|F\|_{L^2(\mathbb{R}^{n+1})}, \quad j < k.$$

Taking the geometric mean of these bounds then shows that for any real $0 \leq \theta \leq 1$,

$$\left\| \sup_{\|\lambda\| \approx r} |I_{2^k}^\lambda \Delta_j F| \right\|_{L^2(\mathbb{R}^{n+1})} \ll r^{(1-\theta)/2 - \theta\delta_0} 2^{(j-k)(1-\theta)} \|F\|_{L^2(\mathbb{R}^{n+1})}.$$

This suffices to show (6.4), as long as $0 < \theta < 1$ is chosen sufficiently close to 1 that $(1-\theta)/2 - \theta\delta_0 < 0$; such a choice of course exists, since $\delta_0 > 0$.

We will prove (6.18) and (6.19) by a TT^* argument for the operator

$$T_I f(x, t) = I_{2^k}^\lambda \Delta_j f(x, t),$$

where $\lambda(x, t)$ is a measurable stopping-time taking values in $[r, 2r)$. We compute that

$$(6.20) \quad T_I T_I^* f(x, t) = \int_{\mathbb{R}^{n+1}} f(x-u, t-\theta) \frac{1}{a^{n+2}} K_I^{\lambda(x,t), \lambda(x-u, t-\theta)} \left(\frac{u}{a}, \frac{\theta}{a^2} \right) dud\theta,$$

with kernel

$$K_I^{\nu, \mu}(u, \theta) = \int_{\mathbb{R}^n} e^{iP_\nu(u+z) - iP_\mu(z)} \eta(u+z) \eta(z) \underline{\Delta}_{j-k}(\theta - Q(u+z) + Q(z)) dz.$$

We again employ the partition of unity (5.4), writing

$$K_I^{\nu, \mu}(u, \theta) = \sum_{l=1}^n W_l \left(\frac{u}{|u|} \right) K_{I,l}^{\nu, \mu}(u, \theta) =: \sum_{l=1}^n K_{I,l}^{\nu, \mu}(u, \theta).$$

For each l we perform the l -th change of coordinates (5.2), so that we can then write

$$(6.21) \quad K_{I,l}^{\nu, \mu}(u, \theta) = \left(\frac{|u|}{|u|} \right)^{n-2} W_l \left(\frac{u}{|u|} \right) \int_{\mathbb{R}} K_{\sharp, l}^{\nu, \mu}(u, \tau) \underline{\Delta}_{j-k}(\theta - Q(u) - 2|u|\tau) d\tau,$$

where for $W_l(u/|u|) \neq 0$,

$$(6.22) \quad K_{\sharp, l}^{\nu, \mu}(u, \tau) = \int_{\mathbb{R}^{n-1}} e^{iP_\nu(u+z) - iP_\mu(z)} \eta(u+z) \eta(z) d\sigma,$$

in which z is implicitly defined in terms of u, τ, σ as in (5.3). Here, $\nu = \lambda(x, t)$ and $\mu = \lambda(x-u, t-\theta)$. For $W_l(u/|u|) = 0$, we set $K_{\sharp, l}^{\nu, \mu}(u, \tau) = 0$. We note for later reference that $K_{\sharp, l}^{\nu, \mu}(u, \tau)$ is supported on $u \in B_2(\mathbb{R}^n), \tau \in B_1(\mathbb{R})$ (and the integral restricts to $|\sigma| \leq c_0^{-1}$ as explained in (5.7)), so that a trivial bound is

$$(6.23) \quad |K_{\sharp, l}^{\nu, \mu}(u, \tau)| \leq c_0^{-(n-1)} \chi_{B_2}(u) \chi_{B_1}(\tau).$$

Moreover, by the hypothesis of Theorem 6.1, a far stronger bound of the form (6.2) holds for $K_{\sharp, l}^{\nu, \mu}$. We apply this strong bound to see that the contribution of the l -th term to $T_I T_I^* f$ in (6.20) is bounded by

$$(6.24) \quad \ll \iint_{\mathbb{R}^{n+1}} |f(x-u, t-\theta)| \chi((x, t), (u, \theta)) dud\theta,$$

where $\chi((x, t), (u, \theta))$ is the function

$$\int_{\mathbb{R}} \frac{1}{a^n} \left\{ r^{-\delta} \chi_{B_2}\left(\frac{u}{a}\right) \chi_{B_1}(\tau) + \chi_{G^{\lambda(x,t)}}\left(\frac{u}{a}\right) \chi_{B_1}(\tau) + \chi_{B_2}\left(\frac{u}{a}\right) \chi_{F_{\left(\frac{u}{a}\right)}^{\lambda(x,t)}}(\tau) \right\} \frac{1}{a^2} \underline{\Delta}_{j-k} \left(\frac{\theta - Q(u) - 2a|u|\tau}{a^2} \right) d\tau.$$

By Lemma 6.3 (i), $\underline{\Delta}_{j-k}(\theta)$ has bounded $L^1(d\theta)$ norm, uniformly in j, k . We may thus apply Lemma 6.4, using the small measures of the sets $G^{\lambda(x,t)}$ and $F_{\left(\frac{u}{a}\right)}^{\lambda(x,t)}$ and arguing as we did in (6.15). We conclude that the L^2 norm of the operator (6.24) is $\ll r^{-\delta/2}$, which suffices to prove (6.18) with $\delta_0 = \delta/4$.

To prove the last piece of the puzzle, namely (6.19), we again work with $T_I T_I^*$, since this allows us to isolate an integral in τ , and then use properties of antiderivatives of $\underline{\Delta}_j$ from Lemma 6.3 (ii). (See [PY19, §7.4.1 and §10.2] for an in-depth motivation for this approach.) Precisely, we return to examine the l -th part of the kernel of $T_I T_I^*$ defined in (6.21). By the antiderivative property (ii) of Lemma 6.3,

$$\underline{\Delta}_{j-k}(\theta - 2|u|\tau - Q(u)) = -\frac{2^{2(j-k)}}{2|u|} \frac{d}{d\tau} \tilde{\Delta}_{j-k}(\theta - 2|u|\tau - Q(u)).$$

Thus after integration by parts in τ ,

$$(6.25) \quad K_{I,l}^{\nu,\mu}(u, \theta) = \frac{2^{2(j-k)}}{2|u|} \left(\frac{|u_l|}{|u|} \right)^{n-2} W_l \left(\frac{u}{|u|} \right) \int_{\mathbb{R}} \partial_{\tau} K_{\sharp,l}^{\nu,\mu}(u, \tau) \tilde{\Delta}_{j-k}(\theta - 2|u|\tau - Q(u)) d\tau.$$

Recall the trivial bound (6.23) for $K_{\sharp,l}^{\nu,\mu}(u, \tau)$. In fact, from the definition (6.22), $K_{\sharp,l}^{\nu,\mu}(u, \tau)$ is a smooth function of τ , upon recalling the definition of z in terms of u, τ, σ in (5.3). After taking a derivative with respect to τ , we get a trivial upper bound for $\partial_{\tau} K_{\sharp,l}^{\nu,\mu}(u, \tau)$ similar to (6.23) but now with a factor of r , which appears because the coefficients of the phase $P_{\nu}(u+z) - P_{\mu}(z)$ (as a function of τ) are of size $\|\nu\|, \|\mu\| \approx r$. Precisely, we see that

$$|\partial_{\tau} K_{\sharp,l}^{\nu,\mu}(u, \tau)| \ll r \cdot c_0^{-(n-1)} \chi_{B_2}(u) \chi_{B_1}(\tau),$$

in which the implied constant depends on the fixed polynomials $\{p_1, \dots, p_M\}$. Applying this in (6.25), it follows that uniformly in l ,

$$|K_{I,l}^{\nu,\mu}(u, \theta)| \ll r \frac{2^{2(j-k)}}{2|u|} \chi_{B_2}(u) \int_{\mathbb{R}} \chi_{B_1}(\tau) \tilde{\Delta}_{j-k}(\theta - Q(u) - 2|u|\tau) d\tau.$$

Now we compute the contribution of this term to $T_I T_I^* f$ in (6.20): this portion of the operator is bounded by

$$\ll r 2^{2(j-k)} \iint_{\mathbb{R}^{n+1}} |f(x-u, t-\theta)| \chi((x, t), (u, \theta)) du d\theta,$$

now with the function

$$\chi((x, t), (u, \theta)) = \int_{\mathbb{R}} \left(\frac{2|u|}{a} \right)^{-1} \frac{1}{a^n} \chi_{B_2} \left(\frac{u}{a} \right) \chi_{B_1}(\tau) \frac{1}{a^2} \tilde{\Delta}_{j-k} \left(\frac{\theta}{a^2} - 2a|u| \frac{\tau}{a^2} - \frac{Q(u)}{a^2} \right) d\tau.$$

(Note in this case that the function χ defined above is independent of (x, t) since no stopping-times are present.) To apply Lemma 6.4 we must check what values of C and λ are valid in the hypothesis (6.14) of the lemma. We recall that $a = 2^k$ is fixed, and compute the integral in θ first, to see that

$$\| \sup_{(x,t)} |\chi((x, t), (u, \theta))| \|_{L^1(du d\theta)} \leq \int_{\mathbb{R}^n} \left(\frac{2|u|}{a} \right)^{-1} \frac{1}{a^n} \chi_{B_2} \left(\frac{u}{a} \right) du \cdot \int_{\mathbb{R}} \chi_{B_1}(\tau) d\tau \cdot \| \tilde{\Delta}_{j-k} \|_{L^1}.$$

We recall from Lemma 6.3 (ii) that the L^1 norm of $\tilde{\Delta}_{j-k}$ is bounded uniformly in j, k . Then, since $|u|^{-1}$ is locally integrable in \mathbb{R}^n for $n \geq 2$, we see that the integral over u is bounded, uniformly in $a = 2^k$. Hence we may apply Lemma 6.4 for $p = 2$ with finite (but not ‘‘small’’) constants C and $\lambda = C'$. (Here we do not expect to benefit from any decay in r , since no ‘‘small sets’’ are present.)

We conclude that the contribution of $K_{I,l}^{\nu,\mu}$ to the operator $T_I T_I^* f$ in (6.20) has L^2 operator norm bounded by $\ll r^{2^{j-k}}$. Summing over $1 \leq l \leq n$, we see that in total $T_I T_I^*$ has L^2 operator norm $\ll r^{2^{j-k}}$, and consequently (6.19) holds. This completes the proof Theorem 6.1.

To finish the proof of our main result, Theorem 1.1, we next verify that the key hypothesis of Theorem 6.1 holds; that is, that $K_{\sharp,l}^{\nu,\mu}$ admits the appropriate upper bound (6.2). This is our focus in Part II of the paper.

Part II

7. STEP 5: BOUNDING THE OSCILLATORY INTEGRAL $K_{\sharp,l}^{\nu,\mu}$: OVERVIEW

In this section, we outline our strategy for proving the oscillatory integral bound (6.2) required in Theorem 6.1. We specify that $\gamma(y) = (y, Q(y))$ for $Q(y) = \sum_{1 \leq j \leq n} \theta_j y_j^2$, $\theta_j \in \{\pm 1\}$. At this point, we also restrict our attention to an appropriate class of polynomials for the $\text{Span}\{p_1, \dots, p_M\}$, as specified in Theorem 1.1. Our main result for the oscillatory integral kernel is as follows:

Theorem 7.1. *Let $\Psi(u, z)$ be a C^1 function supported on $B_2 \times B_1 \subset \mathbb{R}^n \times \mathbb{R}^n$, with $\|\Psi\|_{C^1(\sigma)} \leq 1$ as in (5.6). Fix $d \geq 2$ and fix polynomials p_2, \dots, p_d such that each p_j is homogeneous of degree j , and $p_2(y) \neq CQ(y)$ for any $C \neq 0$. Let*

$$\Lambda = \{2 \leq j \leq d : p_j \neq 0\}.$$

For each $\lambda \in \mathbb{R}^{|\Lambda|}$ set

$$P_\lambda(y) = \sum_{j \in \Lambda} \lambda_j p_j(y).$$

For each $1 \leq l \leq n$, for $u \in \mathbb{R}^n$ such that $u/|u|$ lies in the support of the partition W_l defined in (5.4), consider the kernel $K_{\sharp,l}^{\nu,\mu}(u, \tau)$ as defined in (5.5). Then there exists a small $\delta > 0$ such that for all $r \geq 1$, if $\nu, \mu \in \mathbb{R}^{|\Lambda|}$ satisfy

$$r \leq \|\nu\|, \|\mu\| \leq 2r,$$

then there exists a measurable set $G^\nu \subset B_2(\mathbb{R}^n)$ (depending on ν but independent of μ, r, Ψ), and for each $u \in B_2(\mathbb{R}^n)$ there exists a measurable set $F_u^\nu \subset B_1(\mathbb{R})$ (depending on u and ν but independent of μ, r, Ψ), such that

$$(7.1) \quad |G^\nu| \ll r^{-\delta}, \quad |F_u^\nu| \ll r^{-\delta},$$

and

$$(7.2) \quad |K_{\sharp,l}^{\nu,\mu}(u, \tau)| \ll r^{-\delta} \chi_{B_2}(u) \chi_{B_1}(\tau) + \chi_{G^\nu}(u) \chi_{B_1}(\tau) + \chi_{B_2}(u) \chi_{F_u^\nu}(\tau),$$

in which all implicit constants depend only on n and the fixed polynomials p_2, \dots, p_d .

Remark 7.2. For polynomials p_2, \dots, p_d fixed as in Theorem 7.1, then Theorem 7.1 proves that the hypothesis of Theorem 6.1 holds, so the hypothesis of Theorem 4.3 holds, so the hypothesis of Theorem 3.1 holds, and finally Theorem 1.1 holds. In this section and §8 we prove Theorem 7.1, and then the proof of Theorem 1.1 is complete.

Remark 7.3. It is significant that the sets in (7.1) and (7.2) are completely independent of μ , although they are allowed to depend on ν ; recall Remark 6.2. Thus a key point of the strategy in the proof of Theorem 7.1 is to avoid any dependence on μ .

The proof of Theorem 7.1 contains the main novel work of this paper. To begin the proof of the theorem, we now fix polynomials p_j homogeneous of degree j for $j \in \Lambda$. Fix $1 \leq l \leq n$, and perform the l -th change of variables from §5, so that z is defined implicitly in terms of u, τ, σ as in (5.3), for $u/|u|$ in the support of W_l . (In all that follows, we only consider such u , since $K_{\sharp,l}^{\nu,\mu}(u, \tau) \equiv 0$ for $u/|u|$ outside the support of W_l .) We recall the definition of $K_{\sharp,l}^{\nu,\mu}(u, \tau)$ from (5.5), as an oscillatory

integral in σ , with phase $P_\nu(u+z) - P_\mu(z)$, which is a polynomial in σ . We set a notation for the coefficient of each monomial σ^γ in this polynomial, by defining

$$P_\nu(u+z) - P_\mu(z) = \sum_{0 \leq |\gamma| \leq d} C[\sigma^\gamma](u, \tau) \sigma^\gamma,$$

where γ varies over multi-indices $\gamma \in \mathbb{Z}_{\geq 0}^{n-1}$. We define a norm for the coefficients of non-constant terms in this polynomial in σ :

$$\|P_\nu(u+z) - P_\mu(z)\|_\sigma := \sum_{1 \leq |\gamma| \leq d} |C[\sigma^\gamma](u, \tau)|.$$

By a standard van der Corput estimate, which we recall below in Lemma 7.4, for each (u, τ) such that

$$(7.3) \quad \|P_\nu(u+z) - P_\mu(z)\|_\sigma \gg r^\varepsilon,$$

then $|K_{\sharp, l}^{\nu, \mu}(u, \tau)| \ll r^{-\delta}$ for $\delta = \varepsilon/d$. We recall the estimate:

Lemma 7.4. [*Prop. 2.1 of [SW01]*] Let $Q_\lambda(x) = \sum_{0 \leq |\alpha| \leq d} \lambda_\alpha x^\alpha$ be a real-valued polynomial for $x \in \mathbb{R}^m$. Define $\|\lambda\| = \sum_{1 \leq |\alpha| \leq d} |\lambda_\alpha|$ (omitting the constant term in Q_λ). For any C^1 function ψ defined on $B_1(\mathbb{R}^m)$ such that $\|\psi\|_{C^1} \leq 1$, and for any convex subset $\Omega \subseteq B_1(\mathbb{R}^m)$,

$$\left| \int_\Omega e^{iQ_\lambda(x)} \psi(x) dx \right| \leq_{m,d} \|\lambda\|^{-1/d},$$

where the implicit constant is independent of ψ, Ω .

Our main goal is to prove that when $\|\nu\|, \|\mu\| \approx r$, for “most” (u, τ) , $\|P_\nu(u+z) - P_\mu(z)\|_\sigma \gg r^\varepsilon$ so that $|K_{\sharp, l}^{\nu, \mu}(u, \tau)| \ll r^{-\delta}$. However, for certain $(u, \tau) \in \mathbb{R}^{n+1}$ we cannot prove that $\|P_\nu(u+z) - P_\mu(z)\|_\sigma \gg r^\varepsilon$; it could be that for each γ the coefficient $C[\sigma^\gamma](u, \tau)$ is small. But these coefficients contain expressions that are polynomial in $u/|u|$ and τ , so instead we show that the set of such (u, τ) has small measure. This uses the following modification of a Van der Corput estimate, which we will apply to coefficients of u and τ inside $C[\sigma^\gamma](u, \tau)$:

Lemma 7.5. [*Lemma 3.3 of [PY19]*] Let $Q_\lambda(x) = \sum_{0 \leq |\alpha| \leq d} \lambda_\alpha x^\alpha$ be a real-valued polynomial for $x \in \mathbb{R}^m$. Define $\llbracket \lambda \rrbracket = \sum_{1 \leq |\alpha| \leq d} |\lambda_\alpha| + |Q_\lambda(0)|$ (including the constant term in Q_λ). For every $\rho > 0$,

$$|\{x \in B_1(\mathbb{R}^m) : |Q_\lambda(x)| \leq \rho\}| \ll_{m,d} \rho^{1/d} \llbracket \lambda \rrbracket^{-1/d}.$$

Finally, we need for these small sets of (u, τ) to be independent of μ , and this dictates the strategy we now describe.

We begin our study of $P_\nu(u+z) - P_\mu(z)$, aiming to prove (7.3) whenever possible. Since our final results must be independent of μ , our first step is to write each coefficient $C[\sigma^\gamma](u, \tau)$ as a sum of three terms: the first term isolates all differences $\nu_j - \mu_j$; the second term depends only on ν_j and is constant in τ , and the third term depends only on ν_j and is linear or higher-order in τ . We summarize this as Lemma 7.7 below, proved (in §8) by a simple Taylor expansion. By convention, $p_1 \equiv 0$ and $\nu_1 = \mu_1 = 0$, since throughout this paper we assume the class of phases (1.3) in our main theorem has no linear terms. In particular, if $|\gamma| = 1$ in the lemma below, the first sum is only over $j \geq 2$. Finally, we recall the notation first encountered in §5,

$$\tilde{u} = (\theta_1 u_1, \dots, \theta_n u_n),$$

where $\theta_i \in \{\pm 1\}$ are the eigenvalues of the fixed quadratic form Q . Since after the l -th change of variables σ has $(n-1)$ coordinates, but the polynomials p_j act on n coordinates, we need a notational convention for book-keeping, which we now define.

Definition 7.6. Fix a dimension $n \geq 2$ and distinguished coordinate $1 \leq l \leq n$. For $w \in \mathbb{R}^{n-1}$, $\lambda \in \mathbb{R}$, write

$$(w; \lambda) = (w_1, \dots, w_{l-1}, \lambda, w_l, \dots, w_{n-1}) \in \mathbb{R}^n,$$

that is, with λ in the l -th coordinate. We apply this notation to multi-indices; for example given a multi-index $\alpha = (\alpha_1, \dots, \alpha_{n-1})$, for $u \in \mathbb{R}^n$, $u^{(\alpha; 3)} = u_1^{\alpha_1} \dots u_l^3 \dots u_n^{\alpha_{n-1}}$. We also apply this notation to indicate derivatives; for example, given a multi-index $\alpha = (\alpha_1, \dots, \alpha_{n-1})$, $\partial_{(\alpha; 3)}$ indicates taking α_1 derivatives in the first variable, α_2 derivatives in the second variable, 3 derivatives in the l -th variable, and so on.

Lemma 7.7. For each multi-index $\gamma \in \mathbb{Z}_{\geq 0}^{n-1}$ with $1 \leq |\gamma| \leq d$,

$$C[\sigma^\gamma](u, \tau) = \sum_{j=|\gamma|}^d (\nu_j - \mu_j) \tau^{j-|\gamma|} B_{j,\gamma} \left(\frac{\tilde{u}}{|u|} \right) + \sum_{j=|\gamma|+1}^d \nu_j D_{j,\gamma} \left(\frac{\tilde{u}}{|u|} \right) |u|^{j-|\gamma|} + \sum_{j=|\gamma|+1}^d \nu_j E_{j,\gamma}(u, \tau)$$

in which we define

$$(7.4) \quad B_{j,\gamma}(w) = \sum_{\alpha \leq \gamma} \frac{(-1)^{|\alpha|}}{\alpha!(\gamma - \alpha)!} (\partial_{(\alpha; |\gamma - \alpha|)} p_j)(w) w^{(\gamma - \alpha; |\alpha|)}, \quad j \geq |\gamma|,$$

$$B_{j,\gamma}(w) \equiv 0, \quad j < |\gamma|.$$

For $j \geq |\gamma| + 1$ we define

$$D_{j,\gamma}(\tilde{u}) = \sum_{\alpha \leq \gamma} \sum_{|\beta|=j-|\gamma|} \frac{(-1)^{|\alpha|}}{\alpha! \beta!(\gamma - \alpha)!} (\partial_{(\alpha; |\gamma - \alpha|) + \beta} p_j)(\tilde{u}) u^\beta (\tilde{u})^{(\gamma - \alpha; |\alpha|)}$$

$$E_{j,\gamma}(u, \tau) = \sum_{\alpha \leq \gamma} \sum_{0 < |\beta| < j - |\gamma|} \frac{(-1)^{|\alpha|}}{\alpha! \beta!(\gamma - \alpha)!} (\partial_{(\alpha; |\gamma - \alpha|) + \beta} p_j)(\tilde{u}) u^\beta \left(\frac{\tau}{|u|} \right)^{j - |\gamma| - |\beta|} \left(\frac{\tilde{u}}{|u|} \right)^{(\gamma - \alpha; |\alpha|)}.$$

For $j \leq |\gamma|$ we define

$$D_{j,\gamma}(\tilde{u}) \equiv 0, \quad E_{j,\gamma}(u, \tau) \equiv 0.$$

For later reference, we summarize a few properties of these functions, which follow immediately from the definitions:

Lemma 7.8.

- (1) $B_{j,\gamma}$ only depends on p_j and γ (and not any other p_m , $m \neq j$). If $p_j = 0$, then $B_{j,\gamma} \equiv 0$.
- (2) $D_{j,\gamma}$ only depends on p_j and γ (and not any other p_m , $m \neq j$). If $p_j = 0$, then $D_{j,\gamma} \equiv 0$.
- (3) For all γ , for all $j \geq |\gamma| + 1$, every term in $E_{j,\gamma}$ has at least order 1 in τ .
- (4) There is a constant C depending only on n, d and the fixed polynomials p_2, \dots, p_d such that for all $1 \leq j \leq d$ and all $j \geq |\gamma|$, $|B_{j,\gamma}(w)|, |D_{j,\gamma}(\tilde{w})|, |E_{j,\gamma}(w, \tau)| \leq C$ for all $w \in B_2(\mathbb{R}^n), \tau \in B_1(\mathbb{R}^n)$.

7.1. Sketch of the strategy. We now give an overview of how we will show that for “most” (u, τ) , there is at least one multi-index γ such that the coefficient $C[\sigma^\gamma](u, \tau)$ is “large”, namely $|C[\sigma^\gamma](u, \tau)| \gg r^\varepsilon$, so that (7.3) holds. Let

$$\Lambda = \{2 \leq j \leq d : p_j \not\equiv 0\}, \quad L = |\Lambda|.$$

Let ν, μ be considered as column vectors of length L , indexed by $j \in \Lambda$ in increasing order. Let D denote the number of possible nonzero multi-indices for σ , that is

$$\mathcal{D} = \{\beta \in \mathbb{Z}_{\geq 0}^{n-1} : 1 \leq |\beta| \leq d\}, \quad D = |\mathcal{D}|.$$

Fix lexicographic order for \mathcal{D} once and for all (starting with all $|\beta| = 1$, all $|\beta| = 2$, and so on). Note that for each $n \geq 2$,

$$D > L.$$

Let $\mathbf{C}(u, \tau)$ denote a column vector of length D with entries $C[\sigma^\gamma](u, \tau)$ for $1 \leq |\gamma| \leq d$. We will define $D \times L$ matrices $\mathbf{B}(u, \tau)$, $\mathbf{D}(u)$ and $\mathbf{E}(u, \tau)$ so that the identity in Lemma 7.7 can be written simultaneously for all $1 \leq |\gamma| \leq d$ by

$$(7.5) \quad \mathbf{C}(u, \tau) = \mathbf{B}(u, \tau)(\nu - \mu) + \mathbf{D}(u)\nu + \mathbf{E}(u, \tau)\nu.$$

Rewrite this as

$$(7.6) \quad \mathbf{B}(u, \tau)(\nu - \mu) = -\mathbf{D}(u)\nu - \mathbf{E}(u, \tau)\nu + \mathbf{C}(u, \tau).$$

We will use this formulation to carry out three ideas simultaneously: on the one hand, we show that we can “invert” the matrix $\mathbf{B}(u, \tau)$ in order to solve for the column vector $\nu - \mu$ in terms of the right-hand side, using Cramer’s rule. For each $j \in \Lambda$, this is a first step toward replacing each occurrence of $\nu_j - \mu_j$ in the phase $P_\nu(u + z) - P_\mu(z)$ by a “harmless” expression only involving ν . Since $\nu - \mu$ is only an L -long vector, we only need to invert an $L \times L$ submatrix of $\mathbf{B}(u, \tau)$. This allows us to focus initially on a version of the relation (7.6) that only involves $L \times L$ submatrices.

On the other hand, suppose that some entry $C[\sigma^\gamma](u, \tau)$ in the column vector $\mathbf{C}(u, \tau)$ is large, that is $\gg r^\varepsilon$. Then (7.3) already holds, and we are finished. Thus in our $L \times L$ restricted version of (7.6), we can suppose that each of the L relevant entries in $\mathbf{C}(u, \tau)$ is small, $\ll r^\varepsilon$. We use this to show that these entries in (7.6) play the role of an $O(r^\varepsilon)$ error term on the right-hand side, uniformly in μ . Then, after inverting a chosen $L \times L$ submatrix of $\mathbf{B}(u, \tau)$, we see that up to an acceptable small error, each appearance of $\nu_j - \mu_j$ in $P_\nu(u + z) - P_\mu(z)$ can be replaced by expressions only involving $(\mathbf{D}(u) + \mathbf{E}(u, \tau))\nu$ (and a harmless determinant coming from Cramer’s rule).

Finally, the last idea is that we take this information, and plug it into a *different* entry in the vector $\mathbf{C}(u, \tau)$; such an entry exists because by construction, $\mathbf{C}(u, \tau)$ is D -long, and $D > L$. In a carefully chosen entry of $\mathbf{C}(u, \tau)$, we input all the previous information to express $C[\sigma^\gamma](u, \tau)$ in a way that is independent of μ , and then show that it is large for almost all (u, τ) . Ultimately this will show that $P_\nu(u + z) - P_\mu(z)$ has at least one large coefficient, aside from u, τ lying in a sufficiently small set, that is defined *independently* of μ .

To accomplish all this, the crux is to prove a lower bound for the rank of $\mathbf{B}(u, \tau)$. This perspective on the proof of Theorem 7.1 is new, and we anticipate it will also be useful for future work on more general maximally modulated Radon transforms.

7.2. Definition of the matrices. Define a $D \times L$ matrix $\mathbf{B}(u, \tau)$ in which the rows are indexed by $\gamma \in \mathbb{Z}_{\geq 0}^{n-1}$ with $1 \leq |\gamma| \leq d$ (the possible multi-indices of monomials σ^γ) and the columns are indexed by $j \in \Lambda \subset \{2, 3, \dots, d\}$:

$$(7.7) \quad \mathbf{B}(u, \tau) = \left(B_{j, \gamma} \left(\frac{\tilde{u}}{|u|} \right) \tau^{j-|\gamma|} \right)_{\substack{\gamma \in \mathcal{D} \\ j \in \Lambda}}.$$

Remark 7.9. For later reference, we notice immediately that in any row of $\mathbf{B}(u, \tau)$ indexed by multi-indices γ with $|\gamma| = 1$, every nonzero entry is at least order 1 in τ (since $1 \notin \Lambda$ so that $j - |\gamma| \geq 1$ for all $j \in \Lambda$). Similarly, if $2 \notin \Lambda$, then in any row of $\mathbf{B}(u, \tau)$ indexed by γ with $|\gamma| = 2$, every nonzero entry is at least order 1 in τ .

Next, define a $D \times L$ matrix $\mathbf{D}(u)$ by

$$(7.8) \quad \mathbf{D}(u) = \left(D_{j, \gamma} \left(\frac{\tilde{u}}{|u|} \right) |u|^{j-|\gamma|} \right)_{\substack{\gamma \in \mathcal{D} \\ j \in \Lambda}}.$$

Finally, define a $D \times L$ matrix $\mathbf{E}(u, \tau)$ by

$$\mathbf{E}(u, \tau) = \left(E_{j, \gamma}(u, \tau) \right)_{\substack{\gamma \in \mathcal{D} \\ j \in \Lambda}}.$$

We regard each such matrix over the ring $\mathbb{R}[u_1/|u|, \dots, u_n/|u|, \tau]$, and we say a determinant of a matrix over this ring is nonzero if it is not identically zero as a polynomial in $u/|u|, \tau$; similarly we say a minor of a matrix is nonzero if it is not identically zero as a polynomial in $u/|u|, \tau$.

Fix $(u, \tau) \in B_2(\mathbb{R}^n) \times B_1(\mathbb{R})$. We want to invert a submatrix of $\mathbf{B}(u, \tau)$ in order to solve for the column vector $\nu - \mu$ in terms of the right-hand side of (7.6). We will show that for each $j \in \Lambda$ with $j \geq 3$, there exists some multi-index $\gamma(j)$ with $|\gamma(j)| = j$, so that $B_{j, \gamma(j)} \neq 0$ (see property (2) in Lemma 7.16), and we will use this to solve for $\nu_j - \mu_j$. For $j = 2$, if $j \in \Lambda$, depending on a division into cases, we will either use some multi-index $\gamma(1)$ with $|\gamma(1)| = 1$ so that $B_{2, \gamma(1)} \neq 0$, or some multi-index $\gamma(2)$ with $|\gamma(2)| = 2$ so that $B_{2, \gamma(2)} \neq 0$, to solve for $\nu_2 - \mu_2$ (see properties (4) and (5) in Lemma 7.16).

Depending on the cases A, B1, B2 defined in Definition 7.15 below, we will fix a set of distinguished exponents, named

$$(7.9) \quad \mathcal{D}^* = \begin{cases} \{\gamma(j) : j \in \Lambda, j \geq 2\}, & \text{case A} \\ \{\gamma(j) : j \in \Lambda, j \geq 3\} & \text{case B1} \\ \{\gamma(j) : j \in \Lambda, j \geq 3\} \cup \{\gamma(1)\} & \text{case B2.} \end{cases}$$

We show precisely how to choose \mathcal{D}^* when we treat each case. In each case, $|\mathcal{D}^*| = |\Lambda| = L$. (For transparency, one sub-case of B2 is slightly different: we take $\mathcal{D}^* = \{\gamma(j) : j \in \Lambda, j \geq 3\}$ which has cardinality $L - 1$, and adapt the argument that follows, accordingly; see the explanation of case (1) in Lemma 7.20.)

Define $\mathbf{B}^*(u, \tau)$ to be the $L \times L$ submatrix obtained from $\mathbf{B}(u, \tau)$ by choosing the L rows corresponding to $\gamma \in \mathcal{D}^*$. By this construction, $\mathbf{B}^*(u, \tau)$ is an upper triangular square matrix with determinant that is a polynomial in $\tau, \tilde{u}/|u|$ that is $\neq 0$. Let $\mathbf{D}^*, \mathbf{E}^*$ correspondingly denote the $L \times L$ submatrices of \mathbf{D}, \mathbf{E} corresponding to these rows with $\gamma \in \mathcal{D}^*$. Finally, let $\mathbf{C}^*(u, \tau)$ denote the L -long column vector plucked from $\mathbf{C}(u, \tau)$ by taking the rows indexed by $\gamma \in \mathcal{D}^*$. We now continue to specify the rest of the strategy, whose conclusion we summarize in Proposition 7.11.

7.3. Application of Cramer's rule. Fix $\varepsilon_1 > 0$. Suppose that for some $\gamma \in \mathcal{D}^*$, $|C[\sigma^\gamma](u, \tau)| \geq r^{\varepsilon_1}$. Then $\|P_\nu(u + z) - P_\mu(z)\|_\sigma \geq r^{\varepsilon_1}$ so that by Lemma 7.4, $|K_{\sharp, l}^{\nu, \mu}(u, \tau)| \ll r^{-\varepsilon_1/d}$, which suffices for Theorem 7.1. Thus from now on we assume on the contrary that $|C[\sigma^\gamma](u, \tau)| \leq r^{\varepsilon_1}$ for all $\gamma \in \mathcal{D}^*$. Taking the $L \times L$ restriction of (7.6) we see that

$$\mathbf{B}^*(u, \tau)(\nu - \mu) = -\mathbf{D}^*(u)\nu - \mathbf{E}^*(u, \tau)\nu + \mathbf{C}^*(u, \tau),$$

where by our assumption, each entry in the L -long column vector $\mathbf{C}^*(u, \tau)$ is $O(r^{\varepsilon_1})$. Now we apply Cramer's rule to solve for the entries in $\nu - \mu$. Temporarily, for each $j \in \Lambda$ let

$$M^j(u, \tau) = \text{the matrix } \mathbf{B}^*(u, \tau) \text{ with } j\text{-th column replaced by } -(\mathbf{D}^*(u) + \mathbf{E}^*(u, \tau))\nu + \mathbf{C}^*(u, \tau).$$

More importantly, for each $j \in \Lambda$, let

$$(7.10) \quad \mathbf{B}^{*,j}(u, \tau) = \text{the matrix } \mathbf{B}^*(u, \tau) \text{ with } j\text{-th column replaced by } -(\mathbf{D}^*(u) + \mathbf{E}^*(u, \tau))\nu.$$

Then by Cramer's rule, for each $j \in \Lambda$,

$$(7.11) \quad \det \mathbf{B}^*(u, \tau)(\nu_j - \mu_j) = \det M^j(u, \tau) = \det \mathbf{B}^{*,j}(u, \tau) + O(r^{\varepsilon_1}).$$

The last equality follows because we use the uniform upper bound in Lemma 7.8 property (4) for all the polynomials $B_{j, \gamma}$ (when $u \in B_2, \tau \in B_1$) as well as the uniform upper bound $O(r^{\varepsilon_1})$ for each element in the column vector $\mathbf{C}^*(u, \tau)$. The impact of (7.11) is that up to a uniform $O(r^{\varepsilon_1})$ error, we can now replace any occurrence of $(\nu_j - \mu_j)$ by an expression involving $\det \mathbf{B}^{*,j}(u, \tau)$, which depends only on ν , not on μ .

Finally, we use the fact that $|\mathcal{D}^*| \leq d - 1$, while $|\mathcal{D}| \geq d$. So there exists at least one index $\gamma \in \mathcal{D} \setminus \mathcal{D}^*$. For each such γ , applying (7.11) to the γ -th row of the vector identity (7.5), we see

that

$$(7.12) \quad \det \mathbf{B}^*(u, \tau) C[\sigma^\gamma](u, \tau) = \underline{\mathbf{B}}_\gamma(u, \tau) (\det \mathbf{B}^{*,j}(u, \tau))_{j \in \Lambda} \\ + \det \mathbf{B}^*(u, \tau) \underline{\mathbf{D}}_\gamma(u) \nu + \det \mathbf{B}^*(u, \tau) \underline{\mathbf{E}}_\gamma(u, \tau) \nu + O(r^{\varepsilon_1}),$$

in which we let $\underline{\mathbf{B}}_\gamma$ denote the row indexed by γ in the matrix $\mathbf{B}(u, \tau)$, and similarly for the other matrices. Here we have again applied the uniform upper bound in Lemma 7.8 property (4) for all the $B_{j,\gamma}$ functions, when $(u, \tau) \in B_2 \times B_1$, so that the error term is still $O(r^{\varepsilon_1})$ after multiplying (7.5) through by $\det \mathbf{B}^*(u, \tau)$. Importantly, for any $\gamma \in \mathcal{D} \setminus \mathcal{D}^*$, all the main terms on the right-hand side of (7.12) are now *independent* of μ , and the error term is uniform in μ .

For any $\gamma \in \mathcal{D} \setminus \mathcal{D}^*$, define the function

$$(7.13) \quad R_{\nu,u}^\gamma(\tau) := \underline{\mathbf{B}}_\gamma(u, \tau) (\det \mathbf{B}^{*,j}(u, \tau))_{j \in \Lambda} + \det \mathbf{B}^*(u, \tau) \underline{\mathbf{D}}_\gamma(u) \nu + \det \mathbf{B}^*(u, \tau) \underline{\mathbf{E}}_\gamma(u, \tau) \nu.$$

This is a polynomial in τ whose coefficients depend on u and ν . Again by applying Lemma 7.8 property (4), there exists some uniform constant C such that $|\det \mathbf{B}^*(u, \tau)| \leq C$ for all $(u, \tau) \in B_2 \times B_1$, so that

$$(7.14) \quad |C[\sigma^\gamma](u, \tau)| \geq C^{-1} |\det \mathbf{B}^*(u, \tau) C[\sigma^\gamma](u, \tau)|.$$

Let C_0 be an absolute constant that is chosen sufficiently large relative to the $O(r^{\varepsilon_1})$ error term in (7.12) so that if $|R_{\nu,u}^\gamma(\tau)| \geq C_0 r^{\varepsilon_1}$ then

$$(7.15) \quad |R_{\nu,u}^\gamma(\tau) + O(r^{\varepsilon_1})| \geq (C_0/2) r^{\varepsilon_1}.$$

Then if we can prove $|R_{\nu,u}^\gamma(\tau)| \geq C_0 r^{\varepsilon_1}$ for a given (u, τ) , we can conclude by Lemma 7.4 that $|K_{\sharp,l}^{\nu,\mu}(u, \tau)| \ll r^{-\varepsilon_1/d}$.

Remark 7.10. The degree of this polynomial $R_{\nu,u}^\gamma(\tau)$ in τ is bounded above in d ; very crudely we can see the first term is at most degree d^{d+1} in τ while the second term is at most degree 1 in τ , and the last term is at most of degree $d+1$ in τ . Thus $\deg_\tau R_{\nu,u}^\gamma(\tau) \leq d^{d+1}$; this is relevant for applications of Lemma 7.5. We will ultimately focus on showing $R_{\nu,u}^\gamma(\tau)$ has a large coefficient for a term that is either constant or linear in τ .

7.4. Choice of a distinguished index. Our goal now is to show that we can choose some $\gamma \in \mathcal{D} \setminus \mathcal{D}^*$, such that $|R_{\nu,u}^\gamma(\tau)| \geq C_0 r^{\varepsilon_1}$. However, no matter the choice of γ , since $R_{\nu,u}^\gamma(\tau)$ is a polynomial in τ it may take small values for some τ ; Lemma 7.5 shows that the set of such τ will be of small measure (say $\ll r^{-\varepsilon'_1}$) if at least one coefficient of $R_{\nu,u}^\gamma(\tau)$ is large (say $\gg r^{\varepsilon_1}$). On the other hand, since the coefficients of $R_{\nu,u}^\gamma(\tau)$ are functions of u , they may take small values for some u . We will show they are controlled by polynomials in u , and we will verify that at least one such polynomial is not identically zero; since these polynomials in u have coefficients involving the stopping-time ν and by assumption $\|\nu\| \approx r$, we will be able to deduce from a second application of Lemma 7.5, now in the variable u , that the set of u on which the coefficients of $R_{\nu,u}^\gamma(\tau)$ are inconveniently small (say $\ll r^{\varepsilon'_1}$) itself has small measure (say $\ll r^{-\varepsilon_1}$).

We now make this precise. Fix any $\gamma \in \mathcal{D} \setminus \mathcal{D}^*$. Fix a large constant C_0 such that (7.15) holds if $|R_{\nu,u}^\gamma(\tau)| \geq C_0 r^{\varepsilon_1}$. Consider for any fixed $u \in B_2(\mathbb{R}^n)$ the set

$$(7.16) \quad F_u^\nu(\gamma) := \{\tau \in B_1(\mathbb{R}) : |R_{\nu,u}^\gamma(\tau)| \leq C_0 r^{\varepsilon_1}\}.$$

Let $\llbracket R_{\nu,u}^\gamma \rrbracket_\tau$ denote the norm (including constant term) of the coefficients of $R_{\nu,u}^\gamma$ as a polynomial in τ (as defined in Lemma 7.5). We will show that for a certain low degree d_0 , the coefficient of τ^{d_0} in $R_{\nu,u}^\gamma(\tau)$ is a function of u , denoted $\widetilde{W}_\nu^\gamma(u) = |u|^{-s_1} W_\nu^\gamma(u)$ for some $s_1 \geq 1$, where $W_\nu^\gamma(u)$ is a polynomial in u of degree $s_0 \geq 1$ (that in particular is not the zero polynomial); note that $s_0, s_1 \ll_d 1$. Let $\llbracket W_\nu^\gamma \rrbracket_u$ denote the norm (including constant term) of the coefficients of W_ν^γ as a polynomial in u . Consequently, for all $u \in B_2(\mathbb{R}^n)$,

$$\llbracket R_{\nu,u}^\gamma \rrbracket_\tau = |\widetilde{W}_\nu^\gamma(u)| \geq |u|^{-s_1} |W_\nu^\gamma(u)| \gg |W_\nu^\gamma(u)|.$$

Fix some ε_2 with $\varepsilon_1 < \varepsilon_2 < 1$ and define the set

$$G^\nu(\gamma) := \{u \in B_2(\mathbb{R}^n) : |W_\nu^\gamma(u)| \leq r^{\varepsilon_2}\}.$$

For each $u \notin G^\nu(\gamma)$ define $F_u^\nu(\gamma)$ as in (7.16); for each $u \in G^\nu(\gamma)$ set $F_u^\nu(\gamma) = \emptyset$.

Our conclusion is that for all $(u, \tau) \in B_2(\mathbb{R}^n) \times B_1(\mathbb{R})$ such that $u \notin G^\nu(\gamma)$ and $\tau \notin F_u^\nu(\gamma)$, we have $|R_{\nu, u}^\gamma(\tau)| \geq C_0 r^{\varepsilon_1}$, so that $|K_{\sharp, \ell}^{\nu, \mu}(u, \tau)| \ll r^{-\varepsilon_1/d}$ by (7.14). On the other hand, if u or τ is in one of these ‘‘bad sets,’’ we cannot come to this conclusion, but the bad sets are themselves small. Precisely, for each $u \notin G^\nu(\gamma)$, $\llbracket R_{\nu, u} \rrbracket_\tau \geq |u|^{-s_1} |W_\nu^\gamma(u)| \gg r^{\varepsilon_2}$ so that by Lemma 7.5 and Remark 7.10,

$$|F_u^\nu(\gamma)| \ll r^{-(\varepsilon_2 - \varepsilon_1)/d^{d+1}}.$$

Finally, if $\llbracket W_\nu^\gamma(u) \rrbracket_u \geq r$, then by Lemma 7.5,

$$|G^\nu(\gamma)| \ll r^{-(1-\varepsilon_2)/s_0}.$$

We document this discussion in the following proposition, which records both the hypotheses and the conclusions of the discussion:

Proposition 7.11. *Fix a set \mathcal{D}^* as in (7.9), chosen so that $\det \mathbf{B}^*(u, \tau) \neq 0$. Fix $\gamma \in \mathcal{D} \setminus \mathcal{D}^*$ and a degree $d_0 \geq 0$, and let the function $\widehat{W}_\nu^\gamma(u) = |u|^{-s_1} W_\nu^\gamma(u)$ (where $W_\nu^\gamma(u)$ is a polynomial in u of degree s_0) denote the coefficient of τ^{d_0} in*

$$(7.17) \quad \underline{\mathbf{B}}_\gamma(u, \tau) (\det \mathbf{B}^{*,j}(u, \tau))_{j \in \Lambda} + \det \mathbf{B}^*(u, \tau) \underline{\mathbf{D}}_\gamma(u) \nu + \det \mathbf{B}^*(u, \tau) \underline{\mathbf{E}}_\gamma(u, \tau) \nu.$$

Suppose that $\gamma \in \mathcal{D} \setminus \mathcal{D}^*$ and $d_0 \geq 0$ have been chosen to have the property that $\llbracket W_\nu^\gamma(u) \rrbracket_u \geq r$.

Then there exists $\delta > 0$ such that the following holds:

- (i) there exists a set $G^\nu(\gamma) \subset B_2(\mathbb{R}^n)$ with measure $|G^\nu(\gamma)| \ll r^{-\delta}$;
- (ii) for each $u \in B_2(\mathbb{R}^n) \setminus G^\nu(\gamma)$ there exists a set $F_u^\nu(\gamma) \subset B_1(\mathbb{R})$ depending only on u, ν and with measure $|F_u^\nu(\gamma)| \ll r^{-\delta}$;
- (iii) for all $(u, \tau) \in B_2(\mathbb{R}^n) \times B_1(\mathbb{R})$ such that $u \notin G^\nu(\gamma)$ and $\tau \notin F_u^\nu(\gamma)$,

$$|K_{\sharp, \ell}^{\nu, \mu}(u, \tau)| \ll r^{-\delta}.$$

All these sets and all implied constants may depend on ν but are independent of μ . (We may take δ to be the minimum of ε_1/d , $(1-\varepsilon_2)/s_0$, and $(\varepsilon_2 - \varepsilon_1)/d^{d+1}$.) The discussion above has proved the following corollary:

Corollary 7.12. *If the hypotheses of Proposition 7.11 hold for a choice of \mathcal{D}^* , multi-index $\gamma \in \mathcal{D} \setminus \mathcal{D}^*$ and a degree d_0 , then Theorem 7.1 is true.*

Remark 7.13. We will apply Proposition 7.11 for d_0 chosen to be the lowest degree (in τ) of any term in $\det \mathbf{B}^*(u, \tau)$, because then only the first two terms in (7.17) can potentially contribute terms of order d_0 in τ . This is because by property (3) of Lemma 7.8, every entry in the row $\underline{\mathbf{E}}_\gamma(u, \tau)$ has at least order 1 in τ , so every term in the last expression in (7.17) is at least order $d_0 + 1$ in τ .

7.5. Defining the three cases. In the remainder of this paper, we prove that the hypotheses of this proposition hold. We will need to consider three cases, which are based on the two following notions.

Definition 7.14. *Given a quadratic form $Q(y) = \sum_j \theta_j y_j^2$, we say that $p_j \neq 0$ is Q -type if $p_j(y) = CQ(y)^{j/2}$ for a nonzero constant C ; note that j is necessarily even. We say that $p_j \neq 0$ is parabolic if $p_j(y) = C(|y|^2)^{j/2}$ for a nonzero constant C ; note that j is necessarily even.*

Given a stopping-time value $\nu \in \mathbb{R}^L$ with $r \leq |\nu| \leq 2r$, we will say that a Q -type term dominates in $P_\nu = \sum_{j \in \Lambda} \nu_j p_j$ if there exists j such that p_j is Q -type and $|\nu_j| \gg r$. Similarly, a non- Q -type term dominates if there exists j such that p_j is non- Q -type and $|\nu_j| \gg r$.

For a fixed ν , it is possible for a Q -type term and a non- Q -type term both to be dominant, and this causes no contradiction in our method.

Definition 7.15. *The three cases for $P_\nu(y) = \sum_{j \in \Lambda} \nu_j p_j(y)$ we are led to consider are as follows: for a given stopping-time value $\nu \in \mathbb{R}^L$ with $r \leq \|\nu\| \leq 2r$,*

- (1) *Case A: a non- Q -type term dominates*
- (2) *Case B1: a Q -type term dominates and $p_2 \equiv 0$*
- (3) *Case B2: a Q -type term dominates and $p_2 \not\equiv 0$ is not Q -type.*

The definition of these cases are analogous to cases in [PY19]; however, our strategy for treating case B2 must be substantially more sophisticated; see Remark 7.22. (Note that these cases do not allow the Q -type term that dominates to be p_2 . If p_2 is Q -type and the only dominant term, our argument could not work; see Remark 7.30. This is why we must rule out $p_2(y) = CQ(y)$ in our main theorem.)

We conclude this section by stating the fundamental properties of the polynomials $B_{j,\gamma}$ and $D_{j,\gamma}$, which will play a key role in verifying the hypotheses of Proposition 7.11 are satisfied in the three cases. Note that in the functions $D_{j,\gamma}$ and $E_{j,\gamma}$ defined in Lemma 7.7, both \tilde{u} and u appear, while the definition of the $B_{j,\gamma}$ polynomial is insensitive to the “twisting” coefficients θ_i of the fixed quadratic form Q . For this reason, the properties of the $B_{j,\gamma}$ polynomials depend on whether p_j is parabolic, not whether it is Q -type.

Lemma 7.16 (Properties of $B_{j,\gamma}$).

- (1) $B_{j,\gamma}(w)$ is a homogeneous polynomial in w of degree j .
- (2) If $j \in \Lambda$, then there exists $|\gamma| = j$ such that $B_{j,\gamma} \not\equiv 0$.
- (3) If $j \in \Lambda$ and p_j is parabolic, then $B_{j,\gamma} \equiv 0$ whenever $|\gamma|$ is odd.
- (4) If $j \in \Lambda$ and p_j is not parabolic, then there exists some γ with $|\gamma| = 1$ such that $B_{j,\gamma} \not\equiv 0$.
- (5) If $j \in \Lambda$ and p_j is parabolic, then there exists some γ with $|\gamma| = 2$ such that $B_{j,\gamma} \not\equiv 0$.

Properties (3) and (4) together show that p_j is parabolic if and only if $B_{j,\gamma} \equiv 0$ for all $|\gamma| = 1$.

Lemma 7.17 (Properties of $D_{j,\gamma}$).

- (1) $D_{j,\gamma}(w)$ is a homogeneous polynomial in w of degree j .
- (2) If $j \in \Lambda$, then p_j is Q -type if and only if $D_{j,\gamma} \equiv 0$ for all $|\gamma| = 1$.
- (3) If $j \in \Lambda$ and p_j is Q -type, then there exists some γ with $|\gamma| = 2$ such that $D_{j,\gamma} \not\equiv 0$.

In order to focus now on the main argument, we defer the proof of these lemmas to Section 8.

7.6. Construction of the set \mathcal{D}^* and computation of $\det \mathbf{B}^*(u, \tau)$. In order to verify the hypothesis of Proposition 7.11 in each of the three cases, we first specify and justify the choice of the set \mathcal{D}^* in (7.9). Regardless of the case, for each $j \in \Lambda$ with $j \geq 3$, we always use property (2) of Lemma 7.16 to see there exists $\gamma(j)$ with $|\gamma(j)| = j$ such that $B_{j,\gamma(j)} \not\equiv 0$, and we take this as the j -th entry in \mathcal{D}^* . Now we discuss how to choose the $j = 2$ entry of \mathcal{D}^* , working case by case.

In case A, we simply again use property (2) of Lemma 7.16 to see there exists $\gamma(2)$ with $|\gamma(2)| = 2$ such that $B_{2,\gamma(2)} \not\equiv 0$, and we take this as the $j = 2$ entry in \mathcal{D}^* . In case B1, $p_2 \equiv 0$, so that $\nu_2 \equiv \mu_2 \equiv 0$, and the set \mathcal{D}^* does not need any entry that corresponds to solving for $\nu_2 - \mu_2$. In case B2, we will later need to consider four sub-cases (see Lemma 7.20 below). We will consider all γ with $|\gamma| = 1$, and denote such γ by $\gamma^m(1)$ if it has a 1 in the m th place. If p_2 has the property that $B_{2,\gamma^m(1)} \not\equiv 0$, we will take $\gamma^m(1)$ as the entry in \mathcal{D}^* that will help us solve for $\nu_2 - \mu_2$. If on the other hand $B_{2,\gamma^m(1)} \equiv 0$, we will not need to solve for $\nu_2 - \mu_2$, and there will be no corresponding entry included in \mathcal{D}^* .

Now that we have fixed the set \mathcal{D}^* , we define the square $L \times L$ matrix $\mathbf{B}^*(u, \tau)$ as in the discussion above, with columns indexed by $j \in \Lambda$ and rows indexed by $\gamma \in \mathcal{D}^*$. We mentioned earlier that

$\mathbf{B}^*(u, \tau)$ is an upper triangular matrix with determinant that is a polynomial in τ , $\tilde{u}/|u|$ that is $\neq 0$. We now show this. In case A, with the choice $\mathcal{D}^* = \{\gamma(j) : j \in \Lambda, j \geq 2\}$,

$$(7.18) \quad \mathbf{B}^*(u, \tau) = \begin{bmatrix} B_{2,\gamma(2)}\left(\frac{\tilde{u}}{|u|}\right)\tau^{2-2} & B_{3,\gamma(2)}\left(\frac{\tilde{u}}{|u|}\right)\tau^{3-2} & \cdots & B_{d,\gamma(2)}\left(\frac{\tilde{u}}{|u|}\right)\tau^{d-2} \\ 0 & B_{3,\gamma(3)}\left(\frac{\tilde{u}}{|u|}\right)\tau^{3-3} & \cdots & B_{d,\gamma(3)}\left(\frac{\tilde{u}}{|u|}\right)\tau^{d-3} \\ \vdots & & \ddots & \vdots \\ 0 & 0 & & B_{d,\gamma(d)}\left(\frac{\tilde{u}}{|u|}\right)\tau^{d-d} \end{bmatrix}.$$

(If $\Lambda \subsetneq \{2, \dots, d\}$ we simply omit the j -th column and $\gamma(j)$ -th row for $j \notin \Lambda$.) For this choice of \mathcal{D}^* , $\mathbf{B}^*(u, \tau)$ is an upper triangular matrix and $\det \mathbf{B}^*(u, \tau)$ is *constant* in τ , namely

$$(7.19) \quad \det \mathbf{B}^*(u, \tau) = \prod_{j \in \Lambda, j \geq 2} B_{j,\gamma(j)}\left(\frac{\tilde{u}}{|u|}\right).$$

Similarly, in case B1,

$$(7.20) \quad \det \mathbf{B}^*(u, \tau) = \prod_{j \in \Lambda, j \geq 3} B_{j,\gamma(j)}\left(\frac{\tilde{u}}{|u|}\right).$$

In case B2, when we choose $\mathcal{D}^* = \{\gamma(j) : j \in \Lambda, j \geq 3\} \cup \{\gamma^m(1)\}$ for some $|\gamma^m(1)| = 1$ with a 1 in the m -th place, an analogous observation shows that $\mathbf{B}^*(u, \tau)$ is an upper triangular matrix and $\det \mathbf{B}^*(u, \tau)$ is *linear* (homogeneous) in τ , namely

$$(7.21) \quad \det \mathbf{B}^*(u, \tau) = \tau \cdot B_{2,\gamma^m(1)}\left(\frac{\tilde{u}}{|u|}\right) \prod_{j \in \Lambda, j \geq 3} B_{j,\gamma(j)}\left(\frac{\tilde{u}}{|u|}\right).$$

We will next verify the key hypotheses of Proposition 7.11 by studying the coefficient of τ^{d_0} in the expression (7.17) with either the choice $d_0 = 0$ or $d_0 = 1$ (recall Remark 7.13).

7.7. Case A: A non- Q -type term dominates. In case A there exists $m_0 \in \Lambda$ such that p_{m_0} is not Q -type and $|\nu_{m_0}| \sim r$. We will verify the hypotheses of Proposition 7.11 for $\mathcal{D}^* = \{\gamma(j) : j \in \Lambda, j \geq 2\}$, $d_0 = 0$ and a choice of $\gamma \in \mathcal{D} \setminus \mathcal{D}^*$ with $|\gamma| = 1$. Under the hypothesis that p_{m_0} is not Q -type, there is a choice of γ with $|\gamma| = 1$ such that $D_{m_0,\gamma}(\tilde{u}) \neq 0$, by property (2) of Lemma 7.17. We choose such a γ and denote it $\gamma(1)$.

Now that $\gamma = \gamma(1)$ of order 1 has been fixed, we study the τ^{d_0} contribution to (7.17) with $d_0 = 0$. Note that every term in the first expression in (7.17) has order at least 1 in τ ; this follows immediately from the observation below (7.7). Second, every term in the third expression in (7.17) has order at least 1 in τ by property (3) of Lemma 7.8. Thus the constant in τ portion of (7.17) is precisely

$$\widetilde{W}_\nu^{\gamma(1)}(u) := \det \mathbf{B}^*(u, \tau) \underline{\mathbf{D}}_{\gamma(1)}(u) \nu = \det \mathbf{B}^*(u, \tau) \sum_{\substack{|\gamma(1)|+1 \leq j \leq d \\ j \in \Lambda}} \nu_j D_{j,\gamma(1)}\left(\frac{\tilde{u}}{|u|}\right) |u|^{j-|\gamma(1)|}.$$

Here we follow the nomenclature of Proposition 7.11. Using the homogeneity described in properties (1) of Lemma 7.16 and (1) of Lemma 7.17, we see that $\widetilde{W}_\nu^{\gamma(1)}(u) = |u|^{-s_1} W_\nu^{\gamma(1)}(u)$ where $s_1 = 1 + \sum_{j \in \Lambda} j$, and the polynomial is defined by

$$W_\nu^{\gamma(1)}(u) = \left(\prod_{j \in \Lambda} B_{j,\gamma(j)}(\tilde{u}) \right) \cdot \sum_{j \in \Lambda} \nu_j D_{j,\gamma(1)}(\tilde{u}).$$

By construction (using property (2) of Lemma 7.16), the leading product is not the zero polynomial.

Finally, we observe that $\llbracket W_\nu^{\gamma(1)} \rrbracket_u \gg r$. This is because in case A, with the index m_0 such that $|\nu_{m_0}| \sim r$ and p_{m_0} is not Q -type, we chose $\gamma(1)$ so that $D_{m_0,\gamma(1)} \neq 0$. Moreover, for $j \neq m_0$, the

polynomials $D_{j,\gamma(1)}$ are homogeneous of degree $j \neq m_0$, so they cannot cause complete cancellation with $D_{m_0,\gamma(1)}$ which is degree m_0 . Thus as a polynomial, $\llbracket W_\nu^{\gamma(1)} \rrbracket_u \gg r$, so that the hypotheses of Proposition 7.11 are verified, and hence Theorem 7.1 holds in case A.

7.8. Case B1: A Q -type term dominates and $p_2 = 0$. In case B1, there exists $m_0 \in \Lambda$ such that p_{m_0} is Q -type and $|\nu_{m_0}| \sim r$, and $p_2 = 0$, so that $2 \notin \Lambda$. (Indeed, by the B1 hypothesis, m_0 is an even integer, $m_0 \geq 4$.) We will verify the hypotheses of Proposition 7.11 for $\mathcal{D}^* = \{\gamma(j) : j \in \Lambda, j \geq 3\}$, $d_0 = 0$ and a choice of $\gamma \in \mathcal{D} \setminus \mathcal{D}^*$ with $|\gamma| = 2$. There is a choice of γ with $|\gamma| = 2$ such that $D_{m_0,\gamma}(\tilde{u}) \neq 0$, by property (3) of Lemma 7.17. We choose such a γ and denote it $\gamma(2)$.

We study the τ^{d_0} contribution in (7.17) with $d_0 = 0$. Now with $\gamma = \gamma(2)$ of order 2, note that every term in the first expression in (7.17) has order at least 1 in τ since $2 \notin \Lambda$; this follows immediately from the observation below (7.7). Additionally, every term in the third expression in (7.17) has order at least 1 in τ by property (3) of Lemma 7.8. Thus the constant in τ portion of (7.17) is precisely

$$\widetilde{W}_\nu^{\gamma(2)}(u) := \det \mathbf{B}^*(u, \tau) \underline{\mathbf{D}}_{\gamma(2)}(u) \nu = \det \mathbf{B}^*(u, \tau) \sum_{\substack{|\gamma(2)|+1 \leq j \leq d \\ j \in \Lambda}} \nu_j D_{j,\gamma(2)}\left(\frac{\tilde{u}}{|u|}\right) |u|^{j-|\gamma(2)|}.$$

We again follow the nomenclature of Proposition 7.11. Using the homogeneity described in properties (1) of Lemma 7.16 and (1) of Lemma 7.17, we see that $\widetilde{W}_\nu^{\gamma(2)}(u) = |u|^{-s_1} W_\nu^{\gamma(2)}(u)$ where $s_1 = 2 + \sum_{j \in \Lambda} j$, and the polynomial is defined by

$$W_\nu^{\gamma(2)}(u) = \left(\prod_{j \in \Lambda, j \geq 3} B_{j,\gamma(j)}(\tilde{u}) \right) \cdot \sum_{j \in \Lambda} \nu_j D_{j,\gamma(2)}(\tilde{u}).$$

By construction (using property (2) of Lemma 7.16), the leading product is not the zero polynomial.

Finally, we observe that $\llbracket W_\nu^{\gamma(2)} \rrbracket_u \gg r$. This is because in case B1, we chose $\gamma(2)$ so that $D_{m_0,\gamma(2)} \neq 0$. Moreover, for $j \neq m_0$, the polynomials $D_{j,\gamma(2)}$ are homogeneous of degree $j \neq m_0$. Thus as a polynomial in u , $\llbracket W_\nu^{\gamma(2)} \rrbracket_u \gg r$, so that the hypotheses of Proposition 7.11 are verified, and hence Theorem 7.1 holds in case B1.

7.9. Case B2: a Q -type term dominates and $p_2 \neq 0$ is not Q -type. In this case, there exists $m_0 \in \Lambda$ such that p_{m_0} is Q -type and $|\nu_{m_0}| \sim r$, and $p_2 \neq 0$ is not Q -type. Note that by the B2 hypothesis, m_0 is an even integer, $m_0 \geq 4$, and $2 \in \Lambda$.

To fix notation, we now write $\gamma^m(1) \in \mathbb{Z}_{\geq 0}^{n-1}$ with $|\gamma^m(1)| = 1$ for the index that has a 1 in the m -th place. (Note that by construction, $m \neq n$, since $\gamma^m(1) \in \mathbb{Z}_{\geq 0}^{n-1}$ is a multi-index with $(n-1)$ places.) We will frequently use such $(n-1)$ -coordinate multi-indices in this section and §8, and we need to adopt a notational convention that depends on whether m falls before or after the distinguished coordinate l .

Definition 7.18. Recall that the l -th coordinate in \mathbb{R}^n is distinguished, according to which change of variables (5.2) we have used. We let $\mathbf{e}_1, \dots, \mathbf{e}_n$ denote the standard coordinate vectors in \mathbb{R}^n .

$$(7.22) \quad \text{Let } m(l) = m \text{ if } m < l \text{ and let } m(l) = m + 1 \text{ if } m \geq l.$$

Then for example $(\gamma^m(1); 0) = \mathbf{e}_{m(l)}$ and $(\gamma^m(1); 1) = \mathbf{e}_{m(l)} + \mathbf{e}_l$.

Case B2 is the most difficult case in which to verify that the hypotheses of Proposition 7.11 hold for appropriate choices of \mathcal{D}^* , $\gamma \in \mathcal{D} \setminus \mathcal{D}^*$ and d_0 . We will check each $\gamma^m(1)$ for $1 \leq m \leq n-1$ and show that either the hypotheses of the proposition are verified for the choice $\gamma = \gamma^m(1)$, or p_2 “looks like Q in that coordinate”. As a result we’ll show that the hypotheses must be verified for *some* m , because otherwise p_2 would be Q -type, which is not allowed in case B2. To make this precise, we make the following definition:

Definition 7.19. Recall that the l -th coordinate in \mathbb{R}^n is distinguished, according to which change of variables (5.2) we have used. Throughout this section, we fix a notation for the coefficients of p_2 by writing $p_2(u) = \sum_{|\omega|=2} d_\omega u^\omega$. For each coordinate $1 \leq i \leq n$ we say p_2 is Q -type in the coordinate i if $\theta_l d_{2l} = \theta_i d_{2e_i}$ and $d_{e_i+e_j} = 0$ for all $j \neq i$.

Notice that p_2 is Q -type if and only if p_2 is Q -type for $i = l$ and for $i = m(l)$ for each $m \in \{1, \dots, n-1\}$. Indeed, if the latter holds then

$$(7.23) \quad p_2(u) = \theta_l d_{2e_l} (\theta_{1(l)} u_{1(l)}^2 + \dots + \theta_{(n-1)(l)} u_{(n-1)(l)}^2 + \theta_l u_l^2) = C \sum_{i=1}^n \theta_i u_i^2$$

with $C = \theta_l d_{2e_l}$, so that p_2 is Q -type. If p_2 is Q -type, then the conclusion follows via the same type of identity.

Our task is to prove that in case B2, the hypotheses of Proposition 7.11 hold for some appropriate choices of the set \mathcal{D}^* , an index $\gamma \in \mathcal{D} \setminus \mathcal{D}^*$ and a degree $d_0 \geq 0$. The key is to prove the following lemma:

Lemma 7.20. Suppose we are in case B2. Let any $m \in \{1, \dots, n-1\}$ be given.

- (1) If $B_{2, \gamma^m(1)} \equiv 0$ and $\theta_l \neq \theta_{m(l)}$, then the hypotheses of Proposition 7.11 hold with $\mathcal{D}^* = \{\gamma(j) : j \in \Lambda, j \geq 3\}$, $\gamma^m(1) \in \mathcal{D} \setminus \mathcal{D}^*$, and $d_0 = 1$.
- (2) If $B_{2, \gamma^m(1)} \not\equiv 0$ and $\theta_l = \theta_{m(l)}$, then the hypotheses of Proposition 7.11 hold with $\mathcal{D}^* = \{\gamma(j) : j \in \Lambda, j \geq 3\} \cup \{\gamma^m(1)\}$, a choice of $\gamma \in \mathcal{D} \setminus \mathcal{D}^*$ with $|\gamma| = 2$ provided by Proposition 7.26, and $d_0 = 1$.
- (3) If $B_{2, \gamma^m(1)} \equiv 0$ and $\theta_l = \theta_{m(l)}$ then p_2 is Q -type in coordinate l and in coordinate $m(l)$.
- (4) If $B_{2, \gamma^m(1)} \not\equiv 0$ and $\theta_l \neq \theta_{m(l)}$: either the hypotheses of Proposition 7.11 hold with $\mathcal{D}^* = \{\gamma(j) : j \in \Lambda, j \geq 3\} \cup \{\gamma^m(1)\}$, a choice of $\gamma \in \mathcal{D} \setminus \mathcal{D}^*$ with $|\gamma| = 2$ provided by Proposition 7.28, and $d_0 = 1$, or p_2 is Q -type in coordinate l and in coordinate $m(l)$.

In combination, the four cases of this lemma imply:

Proposition 7.21. Recall that in case B2, there exists $m_0 \in \Lambda$ such that p_{m_0} is Q -type and $|\nu_{m_0}| \sim r$, and $p_2 \not\equiv 0$ is not Q -type. In case B2, the hypotheses of Proposition 7.11 hold, for some choice of $\mathcal{D}^*, \gamma \in \mathcal{D} \setminus \mathcal{D}^*, d_0$. Hence in case B2, Theorem 7.1 holds.

Proof. Suppose we are in case B2. By Lemma 7.20, either there is some choice of $m \in \{1, \dots, n-1\}$ for which the hypotheses of Proposition 7.11 hold (with the appropriate choices for $\mathcal{D}^*, \gamma \in \mathcal{D} \setminus \mathcal{D}^*, d_0$ specified in the lemma), or for all $m \in \{1, \dots, n-1\}$, p_2 is Q -type in coordinate l and in coordinate $m(l)$. But then p_2 has the form (7.23), so that p_2 is either identically zero or Q -type, which contradicts the definition of Case B2. Thus there is some choice of $m \in \{1, \dots, n-1\}$ for which the hypotheses of Proposition 7.11 are satisfied. We apply Proposition 7.11 with this choice, and by Corollary 7.12, Theorem 7.1 holds. \square

Remark 7.22. It is worth specifying how the case B2 is significantly simpler when $Q(y) = |y|^2$ is positive definite, as considered by Pierce and Yung in [PY19]. By (3) and (4) of Lemma 7.16, p_2 is parabolic if and only if $B_{2, \gamma} \equiv 0$ for all $|\gamma| = 1$. If $Q(y) = |y|^2$ then in case B2, p_2 is not parabolic, so that there exists some $\gamma^m(1)$ with $|\gamma^m(1)| = 1$ such that $B_{2, \gamma^m(1)} \not\equiv 0$. Meanwhile, because $Q(y) = |y|^2$, it is true for all m that $\theta_l = \theta_{m(l)}$, so the only case we need prove of Lemma 7.20 is case (2). Moreover, it is much easier to prove case (2) when $Q(y) = |y|^2$. The key to proving case (2) is Proposition 7.26 (stated below). If $Q(y) = |y|^2$, it follows from (3) of Lemma 7.16 that $B_{m_0, \gamma^m(1)} \equiv 0$ for the even index $m_0 \geq 4$ for which p_{m_0} is Q -type. Consequently in Proposition 7.26 the whole second term vanishes. The first term can clearly be made nonzero by applying (3) of Lemma 7.17, which ensures there exists some $\gamma(2)$ with $|\gamma(2)| = 2$ such that $D_{m_0, \gamma(2)} \not\equiv 0$. Thus

the case B2 can be resolved immediately when $Q(y) = |y|^2$, simply by considering properties of the $B_{j,\gamma}$ and $D_{j,\gamma}$ polynomials, in isolation. (Analogous remarks apply when $Q(y) = -|y|^2$.) When $Q(y) \neq \pm|y|^2$, the situation is much more tricky, and we must consider possible interactions of these polynomials; and this leads to the algorithm carried out by applying Lemma 7.20, as described in Proposition 7.21.

We now focus on proving Lemma 7.20.

7.9.1. *Case (1) of Lemma 7.20:* $B_{2,\gamma^{m(1)}} \equiv 0$ and $\theta_l \neq \theta_{m(l)}$. Recall that in case B2, there exists $m_0 \in \Lambda$ such that p_{m_0} is Q -type and $|\nu_{m_0}| \sim r$ with $m_0 \geq 4$ (and even). We take $\mathcal{D}^* = \{\gamma(j) : j \in \Lambda, j \geq 3\}$. This is the only case in which \mathcal{D}^* has $L - 1$ elements instead of L elements; in the present case $B_{2,\gamma^{m(1)}} \equiv 0$ so $\nu_2 - \mu_2$ will not appear in the coefficient $C[\sigma^{\gamma^{m(1)}}](u, \tau)$ we consider, so we do not need to solve for $\nu_2 - \mu_2$ in terms of other expressions. Here is the precise argument.

We define the square $(L - 1) \times (L - 1)$ matrix we denote by $\mathbf{B}_{(2)}^*(u, \tau)$, with rows indexed by $\gamma \in \mathcal{D}^*$ (as usual) and columns indexed only by $j \geq 3, j \in \Lambda$. Then

$$(7.24) \quad \det \mathbf{B}_{(2)}^*(u, \tau) = \prod_{j \in \Lambda, j \geq 3} B_{j,\gamma(j)} \left(\frac{\tilde{u}}{|u|} \right),$$

which is constant in τ . We now use Cramer's rule as in §7.3, but only for the restricted matrix $\mathbf{B}_{(2)}^*(u, \tau)$ and restricted vector $\nu^{(2)} - \mu^{(2)} \in \mathbb{R}^{L-1}$, which omits the $\nu_2 - \mu_2$ entry. This yields the expression (7.11) for $\nu_j - \mu_j$ for each $j \geq 3$, now in terms of the matrix $\mathbf{B}_{(2)}^{*,j}(u, \tau)$ defined analogously to (7.10). We use this expression for each $\nu_j - \mu_j$ with $j \geq 3$, to conclude in (7.12) that

$$\begin{aligned} \det \mathbf{B}_{(2)}^*(u, \tau) C[\sigma^{\gamma^{m(1)}}](u, \tau) &= \sum_{j \in \Lambda, j \geq 3} B_{j,\gamma^{m(1)}} \left(\frac{\tilde{u}}{|u|} \right) \tau^{j-1} \det \mathbf{B}_{(2)}^{*,j}(u, \tau) \\ &\quad + \det \mathbf{B}_{(2)}^*(u, \tau) B_{2,\gamma^{m(1)}} \left(\frac{\tilde{u}}{|u|} \right) \tau (\nu_2 - \mu_2) \\ &\quad + \det \mathbf{B}_{(2)}^*(u, \tau) \underline{\mathbf{D}}_{\gamma^{m(1)}}(u) \nu + \det \mathbf{B}_{(2)}^*(u, \tau) \underline{\mathbf{E}}_{\gamma^{m(1)}}(u, \tau) \nu + O(r^{\varepsilon_1}). \end{aligned}$$

By assumption in case (1), $B_{2,\gamma^{m(1)}} \equiv 0$, so the coefficient of $(\nu_2 - \mu_2)$ vanishes identically. The rest of the argument leading to Proposition 7.11 proceeds verbatim, continuing onward from this identity in place of (7.12), and replacing (7.17) by its direct analogue,

$$(7.25) \quad \sum_{j \in \Lambda, j \geq 3} B_{j,\gamma^{m(1)}} \left(\frac{\tilde{u}}{|u|} \right) \tau^{j-1} \det \mathbf{B}_{(2)}^{*,j}(u, \tau) + \det \mathbf{B}_{(2)}^*(u, \tau) \underline{\mathbf{D}}_{\gamma^{m(1)}}(u) \nu + \det \mathbf{B}_{(2)}^*(u, \tau) \underline{\mathbf{E}}_{\gamma^{m(1)}}(u, \tau) \nu.$$

Thus we will say we can verify the hypotheses of Proposition 7.11 also in this case.

We will look at the coefficient of τ^1 and verify the case $d_0 = 1$ in Proposition 7.11. (We cannot pick $d_0 = 0$ and work with the constant term in τ in (7.25), or equivalently, (7.17). In fact the only constant in τ in (7.25) is $\det \mathbf{B}_{(2)}^*(u, \tau) \underline{\mathbf{D}}_{\gamma^{m(1)}}(u) \nu$. Since p_{m_0} is Q -type, $D_{m_0,\gamma^{m(1)}} \equiv 0$ by property (2) in Lemma 7.17. Consequently, the coefficient of ν_{m_0} in the constant term in τ is identically zero, so we cannot take advantage of the fact that $|\nu_{m_0}| \sim r$ in that term.)

We now compute the coefficient of τ in (7.25). Note that every term in the first sum in (7.25) visibly has order at least 2 in τ . The second term in (7.25) is constant in τ by (7.24). Thus the terms in (7.17) that are linear in τ can only come from the last expression, namely

$$\det \mathbf{B}_{(2)}^*(u, \tau) \underline{\mathbf{E}}_{\gamma^{m(1)}}(u, \tau) \nu = \det \mathbf{B}_{(2)}^*(u, \tau) \sum_{j \in \Lambda} \nu_j E_{j,\gamma^{m(1)}}(u, \tau).$$

By the definition of $E_{j,\gamma^m(1)}$ we see that linear-in- τ contribution of the sum is precisely

$$\sum_{j \in \Lambda} \nu_j \sum_{\alpha \leq \gamma^m(1)} \sum_{|\beta|=j-|\gamma^m(1)|-1} \frac{(-1)^{|\alpha|}}{\alpha! \beta! (\gamma^m(1) - \alpha)!} (\partial_{\alpha; |\gamma^m(1) - \alpha| + \beta} p_j)(\tilde{u}) u^\beta \left(\frac{\tau}{|u|} \right)^1 \left(\frac{\tilde{u}}{|u|} \right)^{(\gamma^m(1) - \alpha; |\alpha|)}.$$

The middle sum is only over the two terms $\alpha = 0$ and $\alpha = \gamma^m(1)$, and this verifies that the coefficient of τ in (7.25) is the function $\widetilde{W}_\nu^{\gamma^m(1)}(u)$ defined to be

$$\frac{1}{|u|^2} \left(\prod_{j \in \Lambda, j \geq 3} B_{j,\gamma(j)} \left(\frac{\tilde{u}}{|u|} \right) \right) \cdot \sum_{j \in \Lambda} \nu_j \sum_{|\beta|=j-|\gamma^m(1)|-1} \left(\frac{\theta_{m(l)} u^{\beta + \mathbf{e}_{m(l)}}}{\beta!} (\partial_{\beta + \mathbf{e}_l} p_j)(\tilde{u}) - \frac{\theta_l u^{\beta + \mathbf{e}_l}}{\beta!} (\partial_{\beta + \mathbf{e}_{m(l)}} p_j)(\tilde{u}) \right).$$

Now we record that $\widetilde{W}_\nu^{\gamma^m(1)}(u)$ fulfills the hypotheses of Proposition 7.11, using the notation $(\gamma; 0) = \mathbf{e}_{m(l)}$.

Proposition 7.23. *In case (1) of Lemma 7.20 when $\theta_l \neq \theta_{m(l)}$, we confirm that when p_{m_0} is Q -type,*

$$(7.26) \quad \sum_{|\beta|=m_0-|\gamma^m(1)|-1} \left(\frac{\theta_{m(l)} u^{\beta + \mathbf{e}_{m(l)}}}{\beta!} (\partial_{\beta + \mathbf{e}_l} p_{m_0})(\tilde{u}) - \frac{\theta_l u^{\beta + \mathbf{e}_l}}{\beta!} (\partial_{\beta + \mathbf{e}_{m(l)}} p_{m_0})(\tilde{u}) \right) \neq 0.$$

We defer the proof until §8, and finish case (1) of Lemma 7.20. We define $\widetilde{W}_\nu^{\gamma^m(1)}(u) = |u|^{-s_1} W_\nu^{\gamma^m(1)}(u)$, where $s_1 = 2 + \sum_{j \in \Lambda} j$ and $W_\nu^{\gamma^m(1)}(u)$ is a polynomial in u of degree $s_0 = d + \sum_{j \in \Lambda} j$. By (7.26) and the assumption in case B2 that $|\nu_{m_0}| \gg r$ for some m_0 such that p_{m_0} is Q -type, $\llbracket W_\nu^{\gamma^m(1)} \rrbracket_u \gg |\nu_{m_0}| \gg r$. This completes the proof of case (1) of Lemma 7.20, up to the verification of Proposition 7.23, which we complete in §8.

7.9.2. *Preliminary work for cases (2) and (4) of Lemma 7.20.* All the work of this section is equally applicable to cases (2) and (4) of Lemma 7.20, which share the property that $B_{2,\gamma^m(1)} \neq 0$. In both case (2) and case (4), we choose

$$(7.27) \quad \mathcal{D}^* = \{\gamma(j) : j \in \Lambda, j \geq 3\} \cup \{\gamma^m(1)\}.$$

We recall from (7.21) that $\det \mathbf{B}^*(u, \tau)$ is now linear in τ . Thus in cases (2) and (4) we will work with $d_0 = 1$ and consider terms in (7.17) that are linear in τ . By property (3) of Lemma 7.8, every term in the third expression in (7.17) has at least order 2 in τ , so we can ignore all these terms. The following lemma shows that only one term (corresponding to $j = 2$) in the first expression in (7.17) is linear in τ . It is convenient here to set $\min \deg_\tau(f)$ to denote the minimal degree of any monomial in a function f as a polynomial in τ .

Lemma 7.24. *Suppose $m \in \{1, \dots, n-1\}$ is fixed and $B_{2,\gamma^m(1)} \neq 0$; define \mathcal{D}^* as in (7.27). Let $\gamma \in \mathcal{D} \setminus \mathcal{D}^*$ be such that $|\gamma| = 2$. For each $j \in \Lambda$ with $j \geq 3$, either $\det \mathbf{B}^{*,j}(u, \tau) \equiv 0$, in which case this term does not contribute to (7.17), or $\min \deg_\tau \det \mathbf{B}^{*,j}(u, \tau) \geq 1$, in which case the corresponding term in $\underline{\mathbf{B}}_\gamma(u, \tau) (\det \mathbf{B}^{*,j}(u, \tau))_{j \in \Lambda}$ has at least degree 2 in τ .*

Proof. It is helpful to write $\mathbf{B}^*(u, \tau)$ explicitly, with columns indexed by $j = 2, 3, \dots, d$:

$$(7.28) \quad \mathbf{B}^*(u, \tau) = \begin{bmatrix} B_{2,\gamma^m(1)} \left(\frac{\tilde{u}}{|u|} \right) \tau^{2-1} & B_{3,\gamma^m(1)} \left(\frac{\tilde{u}}{|u|} \right) \tau^{3-1} & \cdots & B_{d,\gamma^m(1)} \left(\frac{\tilde{u}}{|u|} \right) \tau^{d-1} \\ 0 & B_{3,\gamma(3)} \left(\frac{\tilde{u}}{|u|} \right) \tau^{3-3} & \cdots & B_{d,\gamma(3)} \left(\frac{\tilde{u}}{|u|} \right) \tau^{d-3} \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & B_{d,\gamma(d)} \left(\frac{\tilde{u}}{|u|} \right) \tau^{d-d} \end{bmatrix}$$

(In general, for any $\Lambda \subset \{2, \dots, d\}$ we would see that $\mathbf{B}^*(u, \tau)$ is a submatrix of the above, omitting the column and row containing $B_{j,\gamma(j)}$ if $j \notin \Lambda$.)

We recall that for each $j \in \Lambda$, $\mathbf{B}^{*,j}(u, \tau)$ is an $L \times L$ matrix formed by replacing the j -th column of $\mathbf{B}^*(u, \tau)$ by the column vector $-(\mathbf{D}^*(u) + \mathbf{E}^*(u, \tau))\nu$. Note that the $j = 2$ column in $\mathbf{B}^*(u, \tau)$ (visually, the left-most column) has a single entry, namely $B_{2, \gamma^m(1)}(\tilde{u}/|u|)\tau^{2-1}$, where by the hypothesis of the lemma, $B_{2, \gamma^m(1)} \neq 0$. Thus for any $j \geq 3$, to compute $\det \mathbf{B}^{*,j}(u, \tau)$ we can expand along the left-most column, and thus every term in the determinant includes a factor of τ . Thus either it is identically zero, or it has minimum degree at least 1 in τ . In the latter case, since every entry in the row vector $\underline{\mathbf{B}}_\gamma(u, \tau)$ has degree at least 1 in τ , we see the contribution of this term to $\underline{\mathbf{B}}_\gamma(u, \tau)(\det \mathbf{B}^{*,j}(u, \tau))_{j \in \Lambda}$ has $\min \deg_\tau \geq 2$. \square

Consequently, for any $\gamma \in \mathcal{D} \setminus \mathcal{D}^*$ with $|\gamma| = 2$, the terms that are linear in τ in (7.17) are precisely the terms that are linear in τ in

$$(7.29) \quad B_{2, \gamma} \left(\frac{\tilde{u}}{|u|} \right) \det \mathbf{B}^{*,2}(u, \tau) + \det \mathbf{B}^*(u, \tau) \underline{\mathbf{B}}_\gamma(u) \nu.$$

We claim that for any $\gamma \in \mathcal{D} \setminus \mathcal{D}^*$ with $|\gamma| = 2$, the coefficient of τ in (7.29) (and hence in (7.17)) is the function

$$(7.30) \quad \widetilde{W}_\nu^\gamma(u) := \left(\prod_{\substack{j \in \Lambda \\ j \geq 3}} B_{j, \gamma(j)} \left(\frac{\tilde{u}}{|u|} \right) \right) \cdot \sum_{j \in \Lambda} \left[B_{2, \gamma^m(1)} \left(\frac{\tilde{u}}{|u|} \right) D_{j, \gamma} \left(\frac{\tilde{u}}{|u|} \right) |u|^{j-|\gamma|} - B_{2, \gamma} \left(\frac{\tilde{u}}{|u|} \right) \Xi_{j, \gamma^m(1)}(\tilde{u}) \right] \nu_j,$$

where for each $j \in \Lambda$ we define

$$(7.31) \quad \Xi_{j, \gamma^m(1)}(\tilde{u}) := \sum_{\alpha=0, \gamma^m(1)} \sum_{|\beta|=j-2} \frac{(-1)^{|\alpha|}}{\alpha! \beta! (\gamma^m(1) - \alpha)!} (\partial_{(\alpha; |\gamma^m(1) - \alpha| + \beta p_j)})(\tilde{u}) u^\beta \left(\frac{1}{|u|} \right)^1 \left(\frac{\tilde{u}}{|u|} \right)^{(\gamma^m(1) - \alpha; |\alpha|)}.$$

(See (8.17) for a somewhat more intuitive expression of this function.) For the record, note that for each j , $|u|^2 \Xi_{j, \gamma^m(1)}(\tilde{u})$ is a polynomial in u . In particular, $\widetilde{W}_\nu^\gamma(u) = |u|^{-s_1} W_\nu^\gamma(u)$ where $W_\nu^\gamma(u)$ is a polynomial in u , and $s_1 = 2 + \sum_{j \in \Lambda} j$. We will explicitly use the expression (7.30) in the proof of both case (2) and case (4) of Lemma 7.20. Furthermore, we will prove:

Lemma 7.25. *If j is even and p_j is of Q -type, then for any fixed index $\gamma^m(1) \in \mathbb{Z}_{\geq 0}^{n-1}$ with $|\gamma^m(1)| = 1$ and a 1 in the m th place,*

$$(7.32) \quad |u|^2 \Xi_{j, \gamma^m(1)}(\tilde{u}) = -\theta_l \theta_{m(l)} B_{j, \gamma^m(1)}(\tilde{u}).$$

This allows us to write the expression for $\widetilde{W}_\nu^\gamma(u)$ in (7.30) in a considerably simpler form, when p_j is Q -type. In particular, if p_j is Q -type, then for any $\gamma \in \mathcal{D} \setminus \mathcal{D}^*$ with $|\gamma| = 2$, (7.30) becomes:

$$(7.33) \quad \left(\prod_{\substack{j \in \Lambda \\ j \geq 3}} B_{j, \gamma(j)} \left(\frac{\tilde{u}}{|u|} \right) \right) \cdot \sum_{j \in \Lambda} \left[B_{2, \gamma^m(1)} \left(\frac{\tilde{u}}{|u|} \right) D_{j, \gamma} \left(\frac{\tilde{u}}{|u|} \right) |u|^{j-|\gamma|} + \theta_l \theta_{m(l)} B_{2, \gamma} \left(\frac{\tilde{u}}{|u|} \right) B_{j, \gamma^m(1)} \left(\frac{\tilde{u}}{|u|} \right) |u|^{j-|\gamma|} \right] \nu_j.$$

This is the expression we will use for $\widetilde{W}_\nu^\gamma(u)$, in order to complete the proofs of cases (2) and (4) of Lemma 7.20.

We defer the proof of Lemma 7.25 to §8. In this section we verify the above expression (7.30) for $\widetilde{W}_\nu^\gamma(u)$ is correct; Recall that the matrix $\mathbf{D}(u)$ defined in (7.8) is independent of τ . Let $\widetilde{W}_\nu^\gamma(u)$ denote the coefficient of τ^1 in (7.17). The formula (7.20) for $\det \mathbf{B}^*(u, \tau)$ shows it is linear in τ ,

and thus the contribution of $\det \mathbf{B}^*(u, \tau) \underline{\mathbf{D}}_\gamma(u) \nu$ to $\widetilde{W}_\nu^\gamma(u)$ is precisely

$$\left(\prod_{\substack{j \in \Lambda \\ j \geq 3}} B_{j, \gamma(j)} \left(\frac{\tilde{u}}{|u|} \right) \right) B_{2, \gamma^m(1)} \left(\frac{\tilde{u}}{|u|} \right) \underline{\mathbf{D}}_\gamma \left(\frac{\tilde{u}}{|u|} \right) \nu.$$

This yields the first part of (7.30).

We must only isolate the part of $B_{2, \gamma}(\tilde{u}/|u|) \det \mathbf{B}^{*,2}(u, \tau)$ that is linear in τ . We focus on $\det \mathbf{B}^{*,2}(u, \tau)$. Consider the linear functional \mathcal{L} taking $\mathbf{w} = (w_j)_{j \in \Lambda} \in \mathbb{R}^L$ to

$$\mathcal{L}(\mathbf{w}) = \det \begin{bmatrix} w_2 & B_{3, \gamma^m(1)} \left(\frac{\tilde{u}}{|u|} \right) \tau^2 & \cdots & B_{d, \gamma^m(1)} \left(\frac{\tilde{u}}{|u|} \right) \tau^{d-1} \\ w_3 & B_{3, \gamma(3)} \left(\frac{\tilde{u}}{|u|} \right) & \cdots & B_{d, \gamma(3)} \left(\frac{\tilde{u}}{|u|} \right) \tau^{d-3} \\ & & \ddots & \\ w_d & 0 & \cdots & B_{d, \gamma(d)} \left(\frac{\tilde{u}}{|u|} \right) \end{bmatrix}.$$

(Here we have written it out for $\Lambda = \{2, \dots, d\}$ but in general it would have L rows.) Then note that

$$(7.34) \quad \det \mathbf{B}^{*,2}(u, \tau) = -\mathcal{L}(\mathbf{D}^*(u) \nu) - \mathcal{L}(\mathbf{E}^*(u, \tau) \nu).$$

Let $q_j(u, \tau)$ for $j \in \Lambda$ denote the functions such that

$$\mathcal{L}(\mathbf{w}) = \sum_{j \in \Lambda} w_j q_j(u, \tau).$$

Visibly, q_2 is constant in τ , namely

$$(7.35) \quad q_2(u, \tau) = q_2(u) = \prod_{j \in \Lambda, j \geq 3} B_{j, \gamma(j)}(\tilde{u}/|u|).$$

Next, we claim that for each $j \in \Lambda$ with $j \geq 3$, $\min \deg_\tau q_j \geq 2$. This is also true by inspection, since in this case we compute q_j as a minor that omits the first column and a row that is *not the top row*. But that minor can be computed by expanding along its top row, and each entry in its top row is visibly of degree at least 2 in τ .

Consequently, $\mathcal{L}(\mathbf{D}^*(u) \nu)$ only has terms that are constant in τ or degree at least 2 in τ , so it contributes no linear terms in τ to (7.34). Second, in $\mathcal{L}(\mathbf{E}^*(u, \tau) \nu)$ the only terms that can be linear in τ must come from terms in the top entry of the column vector $\mathbf{E}^*(u, \tau) \nu$ that are themselves linear in τ . The top entry of the column vector $\mathbf{E}^*(u, \tau) \nu$ is $\underline{\mathbf{E}}_{\gamma^m(1)}(u, \tau) \nu$, where $\gamma^m(1)$ is the distinguished index of order 1 in \mathcal{D}^* . To summarize, the terms in (7.34) that are linear in τ are precisely the terms that are linear in τ in the polynomial

$$-q_2(u) \underline{\mathbf{E}}_{\gamma^m(1)}(u, \tau) \nu = -q_2(u) \sum_{j \in \Lambda} E_{j, \gamma^m(1)}(u, \tau) \nu_j.$$

In the definition of $E_{j, \gamma^m(1)}$ in Lemma 7.7, since $|\gamma^m(1)| = 1$, we only sum over $\alpha = 0$ and $\alpha = \gamma^m(1)$, and the terms that are linear in τ come from β with $|\beta| = j - 2$, so that the coefficient of τ in $E_{j, \gamma^m(1)}(u, \tau)$ is precisely the function $\Xi_{j, \gamma^m(1)}$ defined in (7.31). With this in hand, we see from (7.34) that the coefficient of τ in $\det \mathbf{B}^{*,2}(u, \tau)$ is $-q_2(u) \sum_{j \in \Lambda} \Xi_{j, \gamma^m(1)}(\tilde{u}) \nu_j$, and the formula (7.30) for $\widetilde{W}_\nu^\gamma(u)$ is proved. This completes our preliminary work for cases (2) and (4) of Lemma 7.20.

7.9.3. Case (2) of Lemma 7.20: $B_{2, \gamma^m(1)} \neq 0$ and $\theta_l = \theta_{m(l)}$. Now we specifically focus on case (2) of Lemma 7.20. We record that there is a choice of γ with $|\gamma| = 2$ such that $\widetilde{W}_\nu^\gamma(u)$ fulfills the hypotheses of Proposition 7.11.

Proposition 7.26. *In case (2) of Lemma 7.20, when p_{m_0} is Q -type, there is a choice of $\gamma(2)$ with $|\gamma(2)| = 2$ such that*

$$(7.36) \quad B_{2,\gamma^m(1)}\left(\frac{\tilde{u}}{|u|}\right)D_{m_0,\gamma(2)}\left(\frac{\tilde{u}}{|u|}\right)|u|^{m_0-|\gamma(2)|} + \theta_l\theta_{m(l)}B_{2,\gamma(2)}\left(\frac{\tilde{u}}{|u|}\right)B_{m_0,\gamma^m(1)}\left(\frac{\tilde{u}}{|u|}\right)|u|^{m_0-|\gamma(2)|} \neq 0.$$

(We note, by using the homogeneity of the $B_{j,\gamma}$ functions, that $|u|^4$ times the function in (7.36) is a polynomial in u , so that verifying (7.36) only involves a property of polynomials.) We delay the proof of Proposition 7.26 to §8, and complete the proof of case (2) in Lemma 7.20. We apply Proposition 7.26 to choose $\gamma(2)$ and define $\widetilde{W}_\nu^\gamma(u)$ as in (7.30) for this choice of $\gamma = \gamma(2)$. Then we see $\widetilde{W}_\nu^{\gamma(2)}(u) = |u|^{-s_1}W_\nu^{\gamma(2)}(u)$, where $s_1 = 2 + \sum_{j \in \Lambda} j$. By Proposition 7.26, and the assumption in case B2 that $|\nu_{m_0}| \gg r$, we have $\llbracket W_\nu^{\gamma(2)} \rrbracket_u \gg |\nu_{m_0}| \gg r$. Thus the hypotheses of Proposition 7.11 are satisfied.

7.9.4. *Case (3) of Lemma 7.20: $B_{2,\gamma^m(1)} \equiv 0$ and $\theta_l = \theta_{m(l)}$.* In this case we want to conclude that p_2 is Q -type in coordinate l and in coordinate $m(l)$; this is an immediate consequence of the following lemma.

Lemma 7.27. *Suppose $\theta_l = \theta_{m(l)}$. Then $B_{2,\gamma^m(1)} \equiv 0$ if and only if p_2 is Q -type in the coordinate $m(l)$ and in the coordinate l .*

Proof. We denote the coefficients of p_2 by writing $p_2 = \sum_{|\omega|=2} d_\omega u^\omega$. By definition,

$$\begin{aligned} B_{2,\gamma^m(1)}(w) &= \sum_{\alpha \leq \gamma^m(1)} \frac{(-1)^{|\alpha|}}{\alpha!(\gamma^m(1) - \alpha)!} (\partial_{(\alpha; |\gamma^m(1) - \alpha|)} p_2)(w) w^{(\gamma^m(1) - \alpha; |\alpha|)} \\ &= (\partial_{\mathbf{e}_l} p_2)(w) w^{(\gamma^m(1); 0)} + (-1)(\partial_{\mathbf{e}_{m(l)}} p_2)(w) w^{(0; 1)} \\ &= \sum_{\substack{1 \leq i \leq n \\ i \neq l}} d_{\mathbf{e}_l + \mathbf{e}_i} w^{\mathbf{e}_i + \mathbf{e}_{m(l)}} + 2d_{2\mathbf{e}_l} w^{\mathbf{e}_l + \mathbf{e}_{m(l)}} - \sum_{\substack{1 \leq i \leq n \\ i \neq m(l)}} d_{\mathbf{e}_{m(l)} + \mathbf{e}_i} w^{\mathbf{e}_i + \mathbf{e}_l} - 2d_{2\mathbf{e}_{m(l)}} w^{\mathbf{e}_{m(l)} + \mathbf{e}_l}. \end{aligned}$$

By matching unique powers of w , we see that $B_{2,\gamma^m(1)} \equiv 0$ is equivalent to

$$d_{2\mathbf{e}_l} = d_{2\mathbf{e}_{m(l)}}, \quad d_{\mathbf{e}_l + \mathbf{e}_i} = 0 \quad \forall i \neq l, \quad \text{and} \quad d_{\mathbf{e}_{m(l)} + \mathbf{e}_i} = 0 \quad \forall i \neq m(l).$$

Thus p_2 is Q -type in coordinates $m(l)$ and l according to Definition 7.19, since $\theta_l = \theta_{m(l)}$. \square

7.9.5. *Case (4) of Lemma 7.20: $B_{2,\gamma^m(1)} \neq 0$ and $\theta_l \neq \theta_{m(l)}$.* In this case, precisely as in case (2), we choose

$$\mathcal{D}^* = \{\gamma(j) : j \in \Lambda, j \geq 3\} \cup \{\gamma^m(1)\}$$

and we will work with $d_0 = 1$ and choose some $\gamma \in \mathcal{D} \setminus \mathcal{D}^*$ with $|\gamma| = 2$. We recall from the preliminary work in §7.9.2 that the coefficient of τ in (7.17) was computed to be the expression (7.30), which simplified to (7.33). To complete the proof of case (4) of the lemma, it suffices to verify the following:

Proposition 7.28. *In case (4) of Lemma 7.20, when p_{m_0} is Q -type, either there is a choice of $\gamma(2)$ with $|\gamma(2)| = 2$ such that*

$$(7.37) \quad B_{2,\gamma^m(1)}\left(\frac{\tilde{u}}{|u|}\right)D_{m_0,\gamma(2)}\left(\frac{\tilde{u}}{|u|}\right)|u|^{m_0-|\gamma(2)|} + \theta_l\theta_{m(l)}B_{2,\gamma(2)}\left(\frac{\tilde{u}}{|u|}\right)B_{m_0,\gamma^m(1)}\left(\frac{\tilde{u}}{|u|}\right)|u|^{m_0-|\gamma(2)|} \neq 0,$$

or p_2 is Q -type in coordinate l and in coordinate $m(l)$.

(We note that $|u|^4$ times the function in (7.37) is a polynomial in u , so that verifying (7.37) only involves a property of polynomials.) We delay the proof of Proposition 7.28 to §8, and complete the proof of case (4) in Lemma 7.20. Apply Proposition 7.28 to choose $\gamma(2)$; then define $\widetilde{W}_\nu^\gamma(u)$ as in (7.33) for this choice of $\gamma = \gamma(2)$. Then we see $\widetilde{W}_\nu^{\gamma(2)}(u) = |u|^{-s_1} W_\nu^{\gamma(2)}(u)$, where $s_1 = 2 + \sum_{j \in \Lambda} j$. By (7.37), and the assumption in case B2 that $|\nu_{m_0}| \gg r$, we have $\|W_\nu^{\gamma(2)}\|_u \gg |\nu_{m_0}| \gg r$. Thus the hypotheses of Proposition 7.11 are satisfied. In the alternative case, p_2 is Q -type in coordinate l and in coordinate $m(l)$, and case (4) of the lemma is proved.

To summarize, up until this point, we have completed the proof of Theorem 7.1 (and hence of our main result Theorem 1.1) except for the verification of Lemma 7.7, Lemma 7.16, Lemma 7.17, and the results we require to complete the proof of Lemma 7.20: Proposition 7.23, Lemma 7.25, Proposition 7.26, and Proposition 7.28. All of these we complete in §8.

7.10. Concluding remarks: forbidden cases. It is useful to see why the methods of this section cannot accommodate a linear polynomial within the class $\mathcal{P} = \text{Span}\{p_2, \dots, p_d\}$, or the case when $p_2(y) = Q(y)$. To illustrate this, we consider these two special cases in isolation.

Remark 7.29. Suppose that $\mathcal{P} = \text{Span}\{p_1\}$ where p_1 is a linear polynomial, so that $\Lambda = \{1\}$. (This is a valid reduction of the problem, since in general if $\mathcal{P} = \text{Span}\{p_1, p_2, \dots, p_d\}$, it could be the case that λ with $\|\lambda\| = r$ has $(\lambda_1, \lambda_2, \dots, \lambda_d) = (r, 0, \dots, 0)$; in particular if the method works when \mathcal{P} includes p_1 , it certainly must work for $\mathcal{P} = \text{Span}\{p_1\}$.) Roughly speaking, our method does not work in this case because our general strategy (e.g. in case A when a non- Q -type term of order j dominates) is to solve for $(\nu_j - \mu_j)$ within a coefficient of σ^γ for some $|\gamma| = j$, and then plug it into a coefficient of $\sigma^{\gamma'}$ of lower order where ν_j is present; then we use the large size of ν_j to conclude. If \mathcal{P} contains a linear polynomial, we can only solve for $(\nu_1 - \mu_1)$ inside a coefficient of σ^γ for some $|\gamma| = 1$, and there is no lower order (nonconstant) term in σ that contains ν_1 in which we can apply this information to eliminate $\nu_1 - \mu_1$ and then use the large size of ν_1 in isolation. We briefly present a precise demonstration of this difficulty, for the convenience of the reader.

Suppose that $|\nu_1|, |\mu_1| \approx r$. Fix a distinguished index $1 \leq l \leq n$. For $v \in \mathbb{R}^n$, let $v^{(l)} \in \mathbb{R}^{n-1}$ denote v with the l -th coordinate removed. We calculate directly using (5.3) that

$$\begin{aligned} P_\nu(u+z) - P_\mu(z) &= \nu_1 p_1(u+z) - \mu_1 p_1(z) = (\nu_1 - \mu_1) p_1(z) + \nu_1 p_1(u) \\ &= (\nu_1 - \mu_1) p_1\left(\frac{\tau \tilde{u}^{(l)} - \theta_l u_l \sigma}{|u|}; \frac{\theta_l u_l \tau + \tilde{u}^{(l)} \cdot \sigma}{|u|}\right) + \nu_1 p_1(u). \end{aligned}$$

For concreteness, suppose that $l = n$. Write $p_1(y) = c \cdot y = \sum_{i=1}^n c_i y_i$ so that

$$P_\nu(u+z) - P_\mu(z) = (\nu_1 - \mu_1) \frac{1}{|u|} \left[(-\theta_n u_n c^{(n)} + c_n \tilde{u}^{(n)}) \cdot \sigma \right] + (\nu_1 - \mu_1) \frac{1}{|u|} \left[c \cdot \tilde{u} \right] \tau + \nu_1 c \cdot u.$$

In this subsection, we temporarily let \mathbf{e}_i denote the i -th unit vector in \mathbb{R}^{n-1} . In the language of Lemma 7.7, for each $\gamma = \mathbf{e}_i$ with a 1 in the i -th entry,

$$C[\sigma^\gamma](u, \tau) = (\nu_1 - \mu_1) B_{1, \gamma} \left(\frac{\tilde{u}}{|u|} \right),$$

with $B_{1, \gamma} \left(\frac{\tilde{u}}{|u|} \right) = \frac{1}{|u|} (-\theta_n u_n c_i + c_n \theta_i u_i) \neq 0$. Note that since $j = d = 1$, no $D_{j, \gamma}, E_{j, \gamma}$ functions are present in this coefficient of σ^γ . (This is a problem, since is typically through the nonvanishing of the $D_{j, \gamma}$ or $E_{j, \gamma}$ functions that we are able to apply the large size of ν_j to our advantage.)

Now if $|C[\sigma_1]| \geq r^\varepsilon$, then by Lemma 7.4, we have the desired estimate $|K_{\sharp, l}^{\nu, \mu}(u, \tau)| \ll r^{-\varepsilon}$. Therefore, we may assume that $|C[\sigma_1]| \leq r^\varepsilon$, and consequently $(\nu_1 - \mu_1) B_{1, \mathbf{e}_1} \left(\frac{\tilde{u}}{|u|} \right) = O(r^\varepsilon)$. For any $j \neq 1$, let $\gamma = \mathbf{e}_j$, and multiply the expression for $C[\sigma^\gamma](u, \tau)$ through by $B_{1, \mathbf{e}_1} \left(\frac{\tilde{u}}{|u|} \right)$ to eliminate

the presence of $(\nu_1 - \mu_1)$; observe this shows that

$$B_{1,\gamma}\left(\frac{\tilde{u}}{|u|}\right)C[\sigma_j](u, \tau) = -(\nu_1 - \mu_1)B_{1,\mathbf{e}_1}\left(\frac{\tilde{u}}{|u|}\right)B_{1,\gamma}\left(\frac{\tilde{u}}{|u|}\right) = O(r^\varepsilon)B_{1,\gamma}\left(\frac{\tilde{u}}{|u|}\right) = O(r^\varepsilon).$$

That is, if the coefficient of one σ_i term is small then the coefficients of all the σ_j terms are small, and there is no term left containing ν_1 on its own, for us to exploit the fact that $|\nu_1| \approx r$. Thus we cannot conclude that $|K_{\sharp,l}^{\nu,\mu}(u, \tau)| \ll r^{-\varepsilon}$ when $\mathcal{P} = \text{Span}\{p_1\}$. This problem would remain even if \mathcal{P} also contained higher degree polynomials, since ν_1 will never appear in the expansion of any term σ^γ for any $|\gamma| \geq 2$. When ν_j is large, we can only exploit this by working in powers of σ that have order $< j$.

Remark 7.30. Suppose that $\mathcal{P} = \text{Span}\{p_2\}$ where $p_2 = Q$, so that $\Lambda = \{2\}$. Roughly speaking, our general strategy (e.g. in the umbrella case B2 when a Q -type term dominates of order j) is to solve for $(\nu_j - \mu_j)$ within a coefficient of σ^γ for some γ of order j , and then plug it into a coefficient of $\sigma^{\gamma'}$ for some γ' of lower order, where ν_j is present; then we use the large size of ν_j to conclude. By the expansion for $C[\sigma^\gamma](u, \tau)$ in Lemma 7.7, $(\nu_2 - \mu_2)$ only appears inside the coefficient of σ^γ for some $|\gamma| = 1$ or $|\gamma| = 2$, so potentially one could isolate it in either type of coefficient, and then use this information to isolate a term depending only on ν_2 . However, by Lemma 7.7 again, ν_2 only appears in isolation in a coefficient of $\sigma^{\gamma'}$ for $|\gamma'| = 1$; we'd aim to show that $C[\sigma^{\gamma'}]$ is large using that $|\nu_2| \approx r$. This method fails when p_2 is a multiple of Q because the coefficients of ν_2 in the expression for $C[\sigma^{\gamma'}]$ vanish for all $|\gamma'| = 1$, which follows from (2) of Lemma 7.17 and the vanishing of $E_{1,\gamma'}$ recorded in Lemma 7.7. For the convenience of the reader, we demonstrate this explicitly.

Suppose that $|\nu_2|, |\mu_2| \approx r$ and $p_2(y) = Q(y)$. Calculate directly using (5.3) that

$$\begin{aligned} P_\nu(u+z) - P_\mu(z) &= (\nu_2 - \mu_2)Q(z) + \nu_2(Q(u+z) - Q(z)) \\ &= (\nu_2 - \mu_2) \left(\sum_{\substack{i=1 \\ i \neq l}}^n \theta_i \left(\frac{\tau\theta_i u_i - \theta_l u_l \sigma_i}{|u|} \right)^2 + \theta_l \left(\frac{\theta_l u_l \tau + \tilde{u}^{(l)} \cdot \sigma}{|u|} \right)^2 \right) + \nu_2(Q(u) + 2|u|\tau) \\ &= (\nu_2 - \mu_2) \frac{1}{|u|^2} \left(Q(u)\tau^2 - 2u_l \tau (\theta_l u^{(l)} - \tilde{u}^{(l)} \cdot \sigma) + Q(u_l \sigma; \tilde{u}^{(l)} \cdot \sigma) \right) + \nu_2(Q(u) + 2|u|\tau). \end{aligned}$$

For concreteness, we will again assume that $l = n$. In this subsection, we temporarily let \mathbf{e}_i denote the i -th unit vector in \mathbb{R}^{n-1} . The expression for $P_\nu(u+z) - P_\mu(z)$ above reveals the coefficients $C[\sigma^\gamma]$ for each index γ with $|\gamma| = 2$ and $|\gamma| = 1$. There are three types of γ that we may use to solve for $(\nu_2 - \mu_2)$. The first is $\gamma = 2\mathbf{e}_i$, for which we note that

$$C[\sigma^{2\mathbf{e}_i}](u, \tau) = (\nu_2 - \mu_2)B_{2,2\mathbf{e}_i}\left(\frac{\tilde{u}}{|u|}\right), \quad B_{2,2\mathbf{e}_i}\left(\frac{\tilde{u}}{|u|}\right) = \theta_i \frac{\tilde{u}_n^2}{|u|^2} + \theta_n \frac{\tilde{u}_i^2}{|u|^2}.$$

Next, if $\gamma = \mathbf{e}_i + \mathbf{e}_j$ and $i \neq j$,

$$C[\sigma^{\mathbf{e}_i + \mathbf{e}_j}](u, \tau) = (\nu_2 - \mu_2)B_{2,\mathbf{e}_i + \mathbf{e}_j}\left(\frac{\tilde{u}}{|u|}\right), \quad B_{2,\mathbf{e}_i + \mathbf{e}_j}\left(\frac{\tilde{u}}{|u|}\right) = \frac{1}{|u|^2}(2\theta_n \tilde{u}_i \tilde{u}_j).$$

Finally, if $\gamma = \mathbf{e}_i$, then

$$C[\sigma^{\mathbf{e}_i}](u, \tau) = (\nu_2 - \mu_2)B_{1,\mathbf{e}_i}\left(\frac{\tilde{u}}{|u|}\right), \quad B_{1,\mathbf{e}_i}\left(\frac{\tilde{u}}{|u|}\right) = \frac{2\theta_i \theta_n \tilde{u}_i \tilde{u}_n \tau}{|u|^2} (\theta_i - \theta_n).$$

It suffices by Lemma 7.4 to assume that one of the three coefficients above is small, meaning $|C[\sigma^\gamma]| \leq r^\varepsilon$, otherwise we already have $|K_{\sharp,l}^{\nu,\mu}(u, \tau)| \ll r^{-\varepsilon/2}$. But while we can for example multiply the expression for any coefficient of a σ -term by $B_{2,\gamma}\left(\frac{\tilde{u}}{|u|}\right) \neq 0$ and use the assumption that $|(\nu_2 - \mu_2)B_2\left(\frac{\tilde{u}}{|u|}\right)| \leq r^\varepsilon$ to eliminate $\nu_2 - \mu_2$, the fact that no coefficient of a σ -term in

$P_\nu(u+z) - P_\mu(z)$ includes ν_2 on its own means that this argument cannot conclude that any coefficient of a σ -term is large, although $|\nu_2| \approx r$.

8. PROOF OF THE STATEMENTS APPLIED IN SECTION 7

In this section we gather the proofs of six results we called upon in Section 7 in order to prove Theorem 7.1: Lemma 7.7, Lemma 7.16, Lemma 7.17, and the statements of Proposition 7.23, Lemma 7.25, Proposition 7.26, and Proposition 7.28 that we applied to prove Lemma 7.20.

8.1. Proof of Lemma 7.7. Recall that Lemma 7.7 provides an expansion of the coefficient $C[\sigma^\gamma](u, \tau)$ of σ^γ in the polynomial $P_\nu(u+z) - P_\mu(z)$, for each multi-index $\gamma \in \mathbb{Z}_{>0}^{n-1}$. This follows from a direct computation: we recall that an index $1 \leq l \leq n$ is fixed, and apply the change of variables (5.2) in order to express z in terms of u, τ, σ inside $P_\nu(u+z) - P_\mu(z)$, and then we group terms according to the power of σ . Throughout, for a multi-index $\alpha = (\alpha_1, \dots, \alpha_m)$, we write $|\alpha| = \alpha_1 + \dots + \alpha_m$, $\partial_\alpha = \partial_{x_1}^{\alpha_1} \dots \partial_{x_m}^{\alpha_m}$, $\alpha! = \alpha_1! \dots \alpha_m!$, and $w^\alpha = w_1^{\alpha_1} \dots w_m^{\alpha_m}$. For multi-indices α, β in the same number of coordinates, we will say $\alpha \leq \beta$ if $\alpha_j \leq \beta_j$ for every coordinate j . For $v \in \mathbb{R}^n$, let $v^{(l)} \in \mathbb{R}^{n-1}$ denote v with the l -th coordinate removed.

By (5.3), the formula for z in terms of u, τ, σ is

$$z^{(l)} = \frac{\tau \tilde{u}^{(l)} - \theta_l u_l \sigma}{|u|} \quad \text{and} \quad z_l = \frac{\theta_l u_l \tau + \tilde{u}^{(l)} \cdot \sigma}{|u|}.$$

Then

$$\begin{aligned} u_i + z_i &= u_i + \frac{1}{|u|} (\tau \theta_i u_i - \theta_l u_l \sigma_i) = \frac{u_i}{|u|} (|u| + \theta_i \tau) - \frac{\theta_l u_l}{|u|} \sigma_i \quad \text{for } i \neq l \\ u_l + z_l &= u_l + \frac{1}{|u|} (\theta_l u_l \tau + \tilde{u}^{(l)} \cdot \sigma) = \frac{u_l}{|u|} (|u| + \theta_l \tau) + \frac{\tilde{u}^{(l)}}{|u|} \cdot \sigma \quad \text{for } i = l. \end{aligned}$$

Recall $P_\lambda(y) := \sum \lambda_j p_j(y)$, where each p_j is homogeneous of degree $j \leq d$. By Taylor expansion (that is, a finite multinomial expansion), for each j ,

$$(8.1) \quad p_j(v+w) = \sum_{0 \leq |\alpha| \leq d} \frac{(\partial_\alpha p_j)(v)}{\alpha!} w^\alpha.$$

(Of course, $\partial_\alpha p_j \equiv 0$ if $|\alpha| > j$.) Thus for example

$$\begin{aligned} p_j(u+z) &= p_j\left(u^{(l)} + \frac{\tilde{u}^{(l)}}{|u|} \tau - \frac{\theta_l u_l}{|u|} \sigma; u_l + \frac{\theta_l u_l}{|u|} \tau + \frac{\tilde{u}^{(l)}}{|u|} \cdot \sigma\right) \\ &= \sum_{0 \leq |\alpha| \leq d} \frac{1}{\alpha!} \left[(\partial_\alpha p_j)\left(u + \frac{\tilde{u}}{|u|} \tau\right) \right] \left(-\frac{\theta_l u_l}{|u|} \sigma; \frac{\tilde{u}^{(l)}}{|u|} \cdot \sigma \right)^\alpha. \end{aligned}$$

We can then apply (8.1) a second time to expand each term of the form $(\partial_\alpha p_j)(u + (\tilde{u}/|u|)\tau)$. We now isolate a part in $P_\nu(u+z) - P_\mu(z)$ that contains factors $\nu_j - \mu_j$, and a part that depends only

on ν_j , by applying expansions as above:

$$\begin{aligned}
P_\nu(u+z) - P_\mu(z) &= \sum_{j=2}^d \nu_j p_j(u+z) - \sum_{j=2}^d \mu_j p_j(z) \\
&= \sum_{j=2}^d (\nu_j - \mu_j) p_j(z) + \sum_{j=2}^d \nu_j (p_j(u+z) - p_j(z)) \\
&= \sum_{j=2}^d (\nu_j - \mu_j) \left[\sum_{0 \leq |\alpha| \leq d} \frac{1}{\alpha!} \left[(\partial_\alpha p_j) \left(\frac{\tilde{u}}{|u|} \tau \right) \right] \left(-\frac{\theta_l u_l}{|u|} \sigma; \frac{\tilde{u}^{(l)}}{|u|} \cdot \sigma \right)^\alpha \right] \\
&\quad + \sum_{j=2}^d \nu_j \left[\sum_{0 \leq |\alpha| \leq d} \sum_{0 < |\beta| \leq d} \frac{1}{\alpha! \beta!} (\partial_{\alpha+\beta} p_j) \left(\frac{\tilde{u}}{|u|} \tau \right) u^\beta \left(-\frac{\theta_l u_l}{|u|} \sigma; \frac{\tilde{u}^{(l)}}{|u|} \cdot \sigma \right)^\alpha \right] \\
&= \Sigma_1 + \Sigma_2,
\end{aligned}$$

say. Now we need to group terms according to the power of σ . By the standard multinomial expansion, for the index $\alpha_l \geq 0$,

$$(\tilde{u}^{(l)} \cdot \sigma)^{\alpha_l} = \sum_{|\gamma|=\alpha_l} \frac{\alpha_l!}{\gamma!} (\tilde{u}^{(l)})^\gamma \sigma^\gamma, \quad \text{in which } \gamma \in \mathbb{Z}_{\geq 0}^{n-1}.$$

Thus (recalling $\alpha \in \mathbb{Z}_{\geq 0}^n$ while $\alpha^{(l)}, \gamma \in \mathbb{Z}_{\geq 0}^{n-1}$):

$$\begin{aligned}
\Sigma_1 &= \sum_{j=2}^d (\nu_j - \mu_j) \sum_{0 \leq |\alpha| \leq d} \frac{1}{\alpha!} (\partial_\alpha p_j) \left(\frac{\tilde{u}}{|u|} \tau \right) |u|^{-|\alpha|} (-\theta_l u_l)^{|\alpha^{(l)}|} \sigma^{\alpha^{(l)}} \sum_{|\gamma|=\alpha_l} \frac{\alpha_l!}{\gamma!} (\tilde{u}^{(l)})^\gamma \sigma^\gamma \\
&= \sum_{j=2}^d (\nu_j - \mu_j) \sum_{0 \leq |\alpha^{(l)}| \leq d} \sum_{0 \leq |\gamma| \leq d - |\alpha^{(l)}|} \frac{(-1)^{|\alpha^{(l)}|}}{\alpha^{(l)}! \gamma!} (\partial_{(\alpha^{(l)}; |\gamma|)} p_j) \left(\frac{\tilde{u}}{|u|} \tau \right) |u|^{-|\gamma + \alpha^{(l)}|} (\theta_l u_l)^{|\alpha^{(l)}|} (\tilde{u}^{(l)})^\gamma \sigma^{\gamma + \alpha^{(l)}} \\
&= \sum_{j=2}^d (\nu_j - \mu_j) \sum_{0 \leq |\gamma'| \leq d} \sum_{\alpha^{(l)} \leq \gamma'} \frac{(-1)^{|\alpha^{(l)}|}}{\alpha^{(l)}! (\gamma' - \alpha^{(l)})!} (\partial_{(\alpha^{(l)}; |\gamma' - \alpha^{(l)}|)} p_j) \left(\frac{\tilde{u}}{|u|} \tau \right) \left(\frac{\tilde{u}}{|u|} \right)^{(\gamma' - \alpha^{(l)}; |\alpha^{(l)}|)} \sigma^{\gamma'},
\end{aligned}$$

where in the last line we set $\gamma' = \gamma + \alpha^{(l)} \in \mathbb{Z}_{\geq 0}^{n-1}$. We regroup terms in Σ_2 by a similar procedure, and conclude that

$$\Sigma_2 = \sum_{j=2}^d \nu_j \sum_{0 \leq |\gamma'| \leq d} \sum_{\alpha^{(l)} \leq \gamma'} \sum_{0 < |\beta| \leq d} \frac{(-1)^{|\alpha^{(l)}|}}{\alpha^{(l)}! \beta! (\gamma' - \alpha^{(l)})!} (\partial_{(\alpha^{(l)}; |\gamma' - \alpha^{(l)}| + \beta)} p_j) \left(\frac{\tilde{u}}{|u|} \tau \right) u^\beta \left(\frac{\tilde{u}}{|u|} \right)^{(\gamma' - \alpha^{(l)}; |\alpha^{(l)}|)} \sigma^{\gamma'}.$$

Here we note that γ' is a multi-index in $n-1$ coordinates; now for simplicity we relabel it as γ , and we relabel $\alpha^{(l)} \in \mathbb{Z}_{\geq 0}^{n-1}$ as α . Also, note that the sum over β is naturally restricted to $|\beta| \leq j - |\gamma|$, since higher order derivatives applied to p_j will vanish identically. Finally, since each p_j is homogeneous of degree j , we note that in Σ_1 ,

$$(\partial_{(\alpha; |\gamma - \alpha|)} p_j) \left(\frac{\tilde{u}}{|u|} \tau \right) = (\partial_{(\alpha; |\gamma - \alpha|)} p_j) \left(\frac{\tilde{u}}{|u|} \right) \tau^{j - |\gamma|}.$$

Similarly in Σ_2 we use

$$(\partial_{(\alpha; |\gamma - \alpha| + \beta)} p_j) \left(\frac{\tilde{u}}{|u|} \tau \right) = (\partial_{(\alpha; |\gamma - \alpha| + \beta)} p_j) \left(\frac{\tilde{u}}{|u|} \right) \tau^{j - |\gamma| - |\beta|}.$$

For each multi-index γ we now recognize that the coefficient $C[\sigma^\gamma](u, \tau)$ of σ^γ in $P_\nu(u+z) - P_\mu(z)$ has a contribution from Σ_1 and a contribution from Σ_2 . In the nomenclature of Lemma 7.7, the contribution to $C[\sigma^\gamma](u, \tau)$ from Σ_1 is

$$\sum_{j=|\gamma|}^d (\nu_j - \mu_j) \tau^{j-|\gamma|} B_{j,\gamma}(\tilde{u}/|u|),$$

with the polynomial $B_{j,\gamma}$ as defined in the lemma. The contribution to $C[\sigma^\gamma](u, \tau)$ from Σ_2 we now break into two parts: a part that is constant in τ (contributed by $|\beta| = j - |\gamma|$), and a part that has every term at least order 1 in τ (contributed by $|\beta| < j - |\gamma|$). In total these contributions to $C[\sigma^\gamma](u, \tau)$ take the form

$$\sum_{j=|\gamma|+1}^d \nu_j D_{j,\gamma}(\tilde{u}/|u|) |u|^{j-|\gamma|} + \sum_{j=|\gamma|+1}^d \nu_j E_{j,\gamma}(u, \tau),$$

as defined in the lemma. This completes the verification of Lemma 7.7.

Remark 8.1. Recall that for $Q(y) = \sum_i \theta_i y_i^2$, we have defined $\tilde{u} = (\theta_1 u_1, \theta_2 u_2, \dots, \theta_n u_n)$. Inside the functions $D_{j,\gamma}(\tilde{u})$ and $E_{j,\gamma}(u, \tau)$ as they are defined in Lemma 7.7, we explicitly see u and \tilde{u} both appearing as variables—that is, the “twist” by the signs θ_i plays a significant role. However, inside the function $B_{j,\gamma}(\tilde{u})$, only the variable \tilde{u} appears. This is convenient; in particular, as a function of w , the function $B_{j,\gamma}(w)$ given in (7.4) is identical to the corresponding function encountered in [PY19, Eqn. (9.17)], as we exploit (merely for efficiency) in the next section.

8.2. Proof of Lemma 7.16 for $B_{j,\gamma}$. Recall the definition (7.4) of the polynomial $B_{j,\gamma}(w)$ for each $1 \leq j \leq d$ and multi-index $\gamma \in \mathbb{Z}_{\geq 0}^{n-1}$. In this section, we will simply observe that $B_{j,\gamma}$ is identical to the function playing an analogous role in [PY19, Eqn. (9.17)], where it is called $B_{m,\beta}$ for any integer $1 \leq m \leq d$ and multi-index $\beta \in \mathbb{Z}_{\geq 0}^{n-1}$. Then all the properties of Lemma 7.16 hold immediately, since they have been proved in [PY19, Lemma 9.4] for the function $B_{m,\beta}$.

Let us verify that our function $B_{j,\gamma}(w)$ agrees with the function defined in [PY19, Eqn. (9.17)], which for the moment we notate as $\mathbb{B}_{j,\gamma}(w)$, until we confirm it is identical to our function $B_{j,\gamma}(w)$. Recall that in the setting of that paper (with $Q(y) = |y|^2$), it is shown that

$$(8.2) \quad C[\sigma^\gamma](u, \tau) = \sum_{j=|\beta|}^d (\nu_j - \mu_j) \tau^{j-|\gamma|} \mathbb{B}_{j,\gamma}(u/|u|) + \sum_{j=|\beta|}^d \nu_j ((|u| + \tau)^{j-|\gamma|} - \tau^{j-|\gamma|}) \mathbb{B}_{j,\gamma}(u/|u|).$$

In the notation of that paper, $p_j(y) = \sum_{|\alpha|=j} c_\alpha y^\alpha$ is a polynomial with fixed coefficients denoted c_α , and

$$\mathbb{B}_{j,\gamma}(w) = \sum_{|\alpha|=j} c_\alpha A_{\alpha,\gamma}(w) = \sum_{|\alpha|=j} \frac{\partial_\alpha p_j(w)}{\alpha!} A_{\alpha,\gamma}(w),$$

in which $A_{\alpha,\gamma}$ is defined for each $\gamma \in \mathbb{Z}_{\geq 0}^{n-1}$ with $|\gamma| \leq |\alpha|$ by

$$\sum_{|\gamma| \leq |\alpha|} A_{\alpha,\gamma}(w) \sigma^\gamma = (w_l + w^{(l)} \cdot \sigma)^{\alpha_l} (w^{(l)} - w_l \sigma)^{\alpha^{(l)}} = (w + (-w_l \sigma; w^{(l)} \cdot \sigma))^\alpha.$$

By comparison, we have shown in Lemma 7.7 that in the case where $Q(y) = \sum \theta_i y_i^2$ and $\tilde{u} = (\theta_1 u_1, \dots, \theta_n u_n)$,

$$C[\sigma^\gamma](u, \tau) = \sum_{j=|\gamma|}^d (\nu_j - \mu_j) \tau^{j-|\gamma|} B_{j,\gamma}\left(\frac{\tilde{u}}{|u|}\right) + \sum_{j=|\gamma|+1}^d \nu_j D_{j,\gamma}\left(\frac{\tilde{u}}{|u|}\right) |u|^{j-|\gamma|} + \sum_{j=|\gamma|+1}^d \nu_j E_{j,\gamma}(u, \tau).$$

In particular, in the case that $\theta_i = +1$ for all i , so that $\tilde{u} = u$, comparing this to (8.2) shows that $B_{j,\gamma}(\tilde{u}/|u|) = B_{j,\gamma}(u/|u|) = \mathbb{B}_{j,\gamma}(u/|u|)$, for each j and γ . Thus in particular as a function of w , $B_{j,\gamma}(w) = \mathbb{B}_{j,\gamma}(w)$, for each j and γ . Consequently, each statement of Lemma 7.16 for the function $B_{j,\gamma}(w)$ holds because the corresponding statement for $\mathbb{B}_{j,\gamma}(w)$ has been proved in [PY19, Lemma 9.4]. (Alternatively, one can compute the functions $A_{\alpha,\gamma}(w)$ from the definition above, and show formally, after suitable rearrangements, that the two formulas agree; finally, a direct proof of Lemma 7.16 also works, modelled on the proof of [PY19, Lemma 9.4].)

8.3. Proof of Lemma 7.17 for $D_{j,\gamma}$. Recall the definition:

$$D_{j,\gamma}(\tilde{u}) = \sum_{\alpha \leq \gamma} \sum_{|\beta|=j-|\gamma|} \frac{(-1)^{|\alpha|}}{\alpha! \beta! (\gamma - \alpha)!} (\partial_{(\alpha; |\gamma - \alpha|) + \beta} p_j)(\tilde{u}) u^\beta(\tilde{u})^{(\gamma - \alpha; |\alpha|)}.$$

8.3.1. Proof of property (1). The fact that $D_{j,\gamma}(w)$ is homogeneous in w of degree j follows directly from the definition. Indeed, note that p_j is homogeneous of degree j , and in each term we differentiate to a total order of $|\alpha| + |\gamma - \alpha| + |\beta|$ and then multiply by factors of total degree $|\beta| + |\gamma - \alpha| + |\alpha|$, thus preserving the total degree j .

8.3.2. Proof of property (2): Fix $j \geq 2$. We wish to show that p_j is Q -type if and only if $D_{j,\gamma} \equiv 0$ for all $|\gamma| = 1$. Recall that $D_{j,\gamma}$ was defined after fixing a distinguished index $1 \leq l \leq n$ and carrying out the change of variables (5.3). Fix some γ with $|\gamma| = 1$ and observe that within $D_{j,\gamma}$ the sum over $\alpha \leq \gamma$ has only two terms, corresponding to $\alpha = 0$ and $\alpha = \gamma$, so that

$$D_{j,\gamma}(\tilde{u}) = \sum_{|\beta|=j-1} \frac{1}{\beta!} (\partial_{(0;1) + \beta} p_j)(\tilde{u}) u^\beta(\tilde{u})^{(\gamma;0)} - \sum_{|\beta|=j-1} \frac{1}{\beta!} (\partial_{(\gamma;0) + \beta} p_j)(\tilde{u}) u^\beta(\tilde{u})^{(0;1)}.$$

For each $1 \leq i \leq n$, let \mathbf{e}_i be the i -th standard unit basis vector in \mathbb{R}^n . The property that $D_{j,\gamma} \equiv 0$ for all $|\gamma| = 1$ is thus equivalent to the property

$$(8.3) \quad \sum_{|\beta|=j-1} \frac{1}{\beta!} (\partial_{\mathbf{e}_i + \beta} p_j)(\tilde{u}) u^\beta \theta_i u_i = \sum_{|\beta|=j-1} \frac{1}{\beta!} (\partial_{\mathbf{e}_i + \beta} p_j)(\tilde{u}) u^\beta \theta_l u_l \quad \text{for each } i \neq l.$$

Write $p_j(u) = \sum_{|\alpha|=j} c_\alpha u^\alpha$. We may assume $p_j \not\equiv 0$ since otherwise $D_{j,\gamma} \equiv 0$ for all γ . We will show that (8.3) is equivalent to the property that $p_j(u)$ is of the form

$$p_j(u) = C(\theta_1 u_1^2 + \cdots + \theta_n u_n^2)^k = C \sum_{|\beta|=k} \frac{k!}{\beta!} (\theta_1 u_1^2)^{\beta_1} \cdots (\theta_n u_n^2)^{\beta_n},$$

where $2k = j$ and $C \neq 0$ is some constant. In fact, we claim (8.3) is equivalent to:

(2a): j is even, say $j = 2k$, and for all α with an odd entry, $c_\alpha = 0$;

(2b): for all multi-indices with every entry being even, say, written 2α for some $|\alpha| = k$, we have $c_{2\alpha} = c_{j\mathbf{e}_l} \theta_l^k \theta_\alpha^k / \alpha!$, that is, $C = c_{j\mathbf{e}_l} \theta_l^k$.

To prove this, note that

$$\partial_\beta u^\alpha = \begin{cases} \frac{\alpha!}{(\alpha - \beta)!} u^{\alpha - \beta} & \text{if } \beta \leq \alpha \\ 0 & \text{otherwise.} \end{cases}$$

Using this in (8.3) for each term in each derivative gives that for each $i \neq l$,

$$(8.4) \quad \sum_{|\alpha|=j} c_\alpha \sum_{\substack{|\beta|=j-1 \\ \mathbf{e}_i + \beta \leq \alpha}} \frac{\alpha!}{\beta! (\alpha - \beta - \mathbf{e}_i)!} \tilde{u}^{\alpha - \beta - \mathbf{e}_i} u^\beta \theta_i u_i = \sum_{|\alpha|=j} c_\alpha \sum_{\substack{|\beta|=j-1 \\ \mathbf{e}_i + \beta \leq \alpha}} \frac{\alpha!}{\beta! (\alpha - \beta - \mathbf{e}_i)!} \tilde{u}^{\alpha - \beta - \mathbf{e}_i} u^\beta \theta_l u_l.$$

Note that since $|\alpha| = j$ and $|\beta| = j-1$, each sum over β includes (at most) one term, with $\beta = \alpha - \mathbf{e}_l$ on the left-hand side and $\beta = \alpha - \mathbf{e}_i$ on the right-hand side. We learn that for each $i \neq l$,

$$(8.5) \quad \sum_{\substack{|\alpha|=j \\ \mathbf{e}_l \leq \alpha}} c_\alpha \frac{\theta_i \alpha!}{(\alpha - \mathbf{e}_l)!} u^{\alpha - \mathbf{e}_l + \mathbf{e}_i} = \sum_{\substack{|\alpha|=j \\ \mathbf{e}_i \leq \alpha}} c_\alpha \frac{\theta_l \alpha!}{(\alpha - \mathbf{e}_i)!} u^{\alpha - \mathbf{e}_i + \mathbf{e}_l}.$$

Upon relabeling $\tilde{\alpha} = \alpha - \mathbf{e}_l + \mathbf{e}_i$ on the left-hand side and $\tilde{\alpha} = \alpha - \mathbf{e}_i + \mathbf{e}_l$ on the right-hand side,

$$(8.6) \quad \sum_{\substack{|\tilde{\alpha}|=j \\ \mathbf{e}_i \leq \tilde{\alpha}}} c_{\tilde{\alpha} + \mathbf{e}_l - \mathbf{e}_i} \frac{\theta_i (\tilde{\alpha} + \mathbf{e}_l - \mathbf{e}_i)!}{(\tilde{\alpha} - \mathbf{e}_i)!} u^{\tilde{\alpha}} = \sum_{\substack{|\tilde{\alpha}|=j \\ \mathbf{e}_l \leq \tilde{\alpha}}} c_{\tilde{\alpha} + \mathbf{e}_i - \mathbf{e}_l} \frac{\theta_l (\tilde{\alpha} + \mathbf{e}_i - \mathbf{e}_l)!}{(\tilde{\alpha} - \mathbf{e}_l)!} u^{\tilde{\alpha}}, \quad \text{for each } i \neq l.$$

Inspecting this identity, we can conclude that the hypothesis $D_{j,\gamma} \equiv 0$ for all $|\gamma| = 1$ is equivalent to: for each multi-index $\alpha \in \mathbb{Z}_{\geq 0}^n$ with $|\alpha| = j$, and each $i \neq l$,

$$(8.7) \quad c_{\alpha + \mathbf{e}_l - \mathbf{e}_i} (\alpha_l + 1) \theta_i = c_{\alpha + \mathbf{e}_i - \mathbf{e}_l} (\alpha_i + 1) \theta_l \quad \text{if } \mathbf{e}_i + \mathbf{e}_l \leq \alpha$$

$$(8.8) \quad c_{\alpha + \mathbf{e}_l - \mathbf{e}_i} = 0 \quad \text{if } \mathbf{e}_i \leq \alpha, \mathbf{e}_l \not\leq \alpha,$$

$$(8.9) \quad c_{\alpha + \mathbf{e}_i - \mathbf{e}_l} = 0 \quad \text{if } \mathbf{e}_l \leq \alpha, \mathbf{e}_i \not\leq \alpha.$$

It is convenient to abbreviate (8.7) in the following way: it shows that

$$(8.10) \quad c_\beta \neq 0 \iff c_{\beta - 2\mathbf{e}_i + 2\mathbf{e}_l} \neq 0, \quad \text{for any } i \neq l \text{ such that } \beta \geq 2\mathbf{e}_i.$$

Thus we can increase the l -th coordinate of the multi-index by 2 and simultaneously decrease the i -th coordinate by 2, as long as the i -th coordinate has value at least 2 to begin with.

We use this to prove the property (2a) by ‘‘chaining’’ from one nonzero coefficient to another. Fix a multi-index α . Suppose that the coefficient $c_\alpha \neq 0$; recall that for all α under consideration, $|\alpha| = j \geq 2$. By successive applications of (8.10) to increase the l -th coordinate of the multi-index as much as possible, while decreasing the i -th coordinate for each $i \neq l$, we see that $c_\beta \neq 0$, for the multi-index

$$\beta := \sum_{\substack{\alpha_k \text{ odd} \\ k \neq l}} \mathbf{e}_k + \left(\sum_{\substack{\alpha_k \text{ odd} \\ k \neq l}} (\alpha_k - 1) + \sum_{\substack{\alpha_k \text{ even} \\ k \neq l}} \alpha_k + \alpha_l \right) \mathbf{e}_l.$$

If α_k is odd for some $k \neq l$, then $\beta - \mathbf{e}_k + \mathbf{e}_l$ satisfies the condition of (8.9), implying that

$$c_{(\beta - \mathbf{e}_k + \mathbf{e}_l) + \mathbf{e}_k - \mathbf{e}_l} = 0.$$

But this states that $c_\beta = 0$, which is a contradiction to the fact that $c_\beta \neq 0$. Thus for all $k \neq l$, α_k is even, so that β is actually equal to

$$\beta = \left(\sum_{\substack{\alpha_k \text{ even} \\ k \neq l}} \alpha_k + \alpha_l \right) \mathbf{e}_l = j \mathbf{e}_l$$

(since $|\alpha| = j$) and in particular we learn that $c_{j\mathbf{e}_l} \neq 0$.

We have shown that α_k must be even for each $k \neq l$. In particular, since $|\alpha| = |\alpha^{(l)}| + |\alpha_l| = j$ this means that α_l and j have the same parity. If we next suppose that α_l is odd, then j is odd, so $j \geq 3$. Note that (8.8) tells us that $c_{j\mathbf{e}_i - \mathbf{e}_i + \mathbf{e}_l} = 0$ for all $i \neq l$. But then successive applications of (8.10) show that for each $i \neq l$,

$$0 = c_{(j-1)\mathbf{e}_i + \mathbf{e}_l} = c_{(j-3)\mathbf{e}_i + 3\mathbf{e}_l} = \cdots = c_{j\mathbf{e}_l},$$

which is a contradiction. Therefore, α_l must be even, so that j must be even. This concludes the proof of property (2a).

Next we prove property (2b), for $j = 2k$. Fix any multi-index α with $|\alpha| = k$ and any $i \neq l$ such that $\alpha_i \neq 0$, and consider the coefficient $c_{2\alpha}$ in p_j . Successively apply (8.7) to obtain

$$\begin{aligned} c_{2\alpha} &= c_{2\alpha+2\mathbf{e}_l-2\mathbf{e}_i} \frac{\theta_i(\alpha_l+1)}{\theta_l\alpha_i} \\ &= c_{2\alpha+4\mathbf{e}_l-4\mathbf{e}_i} \frac{\theta_i(\alpha_l+2)}{\theta_l(\alpha_i-1)} \frac{\theta_i(\alpha_l+1)}{\theta_l\alpha_i} \\ &= \cdots = c_{2\alpha+2\alpha_i\mathbf{e}_l-2\alpha_i\mathbf{e}_i} \frac{\theta_i(\alpha_l+\alpha_i)}{\theta_l(1)} \cdots \frac{\theta_i(\alpha_l+2)}{\theta_l(\alpha_i-1)} \frac{\theta_i(\alpha_l+1)}{\theta_l\alpha_i} \\ &= c_{2\alpha+2\alpha_i\mathbf{e}_l-2\alpha_i\mathbf{e}_i} \left(\frac{\theta_i}{\theta_l}\right)^{\alpha_i} \frac{(\alpha_l+\alpha_i)!}{\alpha_i!\alpha_l!}. \end{aligned}$$

Iterate the above process for successive i , $i \neq l$ to obtain

$$\begin{aligned} c_{2\alpha} &= c_{j\mathbf{e}_l} \left(\prod_{i \neq l} \left(\frac{\theta_i}{\theta_l}\right)^{\alpha_i} \right) \frac{(\alpha_l+\alpha_{i_1})!}{\alpha_{i_1}!\alpha_l!} \frac{(\alpha_l+\alpha_{i_1}+\alpha_{i_2})!}{\alpha_{i_2}!(\alpha_l+\alpha_{i_1})!} \cdots \frac{(\alpha_l+\alpha_{i_1}+\cdots+\alpha_{i_{n-1}})!}{\alpha_{i_{n-1}}!(\alpha_l+\alpha_{i_1}+\cdots+\alpha_{i_{n-2}})!} \\ &= c_{j\mathbf{e}_l} \theta_l^k \theta^\alpha \frac{k!}{\alpha!}. \end{aligned}$$

Here we used that each $\theta_k = \pm 1$ and $\alpha_l = k - \alpha_{i_1} - \cdots - \alpha_{i_{n-1}}$. This proves (2b).

On the other hand, if j is even and each $c_{2\alpha}$ takes this form, while $c_\alpha = 0$ for any α with an odd entry, then the above identities show immediately that the key relationships (8.7), (8.8), (8.9) hold, thus completing the proof of property (2) in Lemma 7.17.

8.3.3. Proof of property (3): Fix $j \geq 2$. Here we suppose that p_j is Q -type, that is $j = 2k$ and $p_j(y) = CQ(y)^k$ for some $C \neq 0$. We will show that there is some γ with $|\gamma| = 2$ such that $D_{j,\gamma} \neq 0$. By normalizing, we may assume $C = 1$ and check property (3) for

$$(8.11) \quad p_j(w) = (\theta_1 w_1^2 + \cdots + \theta_n w_n^2)^k = \sum_{|\alpha|=k} c_{2\alpha} w^{2\alpha},$$

where $c_{2\alpha} \neq 0$ for each $|\alpha| = k$. Without loss of generality, for simplicity of notation we can suppose $l \neq 1$, and demonstrate that $\gamma = (2, 0, \dots, 0) \in \mathbb{R}^{n-1}$ yields $D_{j,\gamma} \neq 0$. We again let \mathbf{e}_i denote the standard i -th coordinate vector in \mathbb{R}^n . The formula for $D_{2k,\gamma}$ has three terms:

$$(8.12) \quad \begin{aligned} D_{2k,\gamma}(\tilde{u}) &= \sum_{|\beta|=2k-2} \frac{1}{2!|\beta|!} (\partial_{2\mathbf{e}_1+\beta} p_j)(\tilde{u}) u^\beta(\tilde{u})^{2\mathbf{e}_1} - \sum_{|\beta|=2k-2} \frac{1}{\beta!} (\partial_{\mathbf{e}_1+\mathbf{e}_l+\beta} p_j)(\tilde{u}) u^\beta(\tilde{u})^{\mathbf{e}_1+\mathbf{e}_l} \\ &\quad + \sum_{|\beta|=2k-2} \frac{1}{2!|\beta|!} (\partial_{2\mathbf{e}_l+\beta} p_j)(\tilde{u}) u^\beta(\tilde{u})^{2\mathbf{e}_l}. \end{aligned}$$

We claim that the monomial $u^{2k\mathbf{e}_l}$ only occurs once (in particular, in the first of these three sums), and it has a nonzero coefficient, which suffices to show that $D_{2k,\gamma} \neq 0$. To see this, examine the effect of the derivative $\partial_{2\mathbf{e}_1+\beta}$ on p_j of the above form (8.11), when $|\beta| = 2k - 2$. We see that for such β , the only term in $p_j(w)$ for which $\partial_{2\mathbf{e}_1+\beta}(w^{2\alpha}) \neq 0$ corresponds to the index α satisfying the relation $2\alpha = 2\mathbf{e}_1 + \beta$ (and $c_{2\alpha} \neq 0$ if β has all even entries). Thus the first sum in (8.12) is over indices β with $|\beta| = 2k - 2$ and all even entries, and each such β contributes precisely one term, which is a monomial of the form $u^{\beta+2\mathbf{e}_1}$ with nonzero coefficient. In particular, the index $\beta = (2k - 2)\mathbf{e}_l$ contributes precisely one monomial $u^{2k\mathbf{e}_l}$ in this way, and this has a nonzero coefficient (since $c_{2\mathbf{e}_1+\beta} \neq 0$, since p_j is Q -type).

On the other hand, the monomial $u^{2k\mathbf{e}_l}$ cannot arise in either the second or the third sum in (8.12). Indeed, within the second sum, we can reason similarly to examine the effect of the derivative $\partial_{\mathbf{e}_1+\mathbf{e}_l+\beta}$ on p_j when $|\beta| = 2k - 2$; the only term in $p_j(w)$ for which $\partial_{\mathbf{e}_1+\mathbf{e}_l+\beta}(w^{2\alpha}) \neq 0$

must have $2\alpha = \mathbf{e}_1 + \mathbf{e}_l + \beta$, and ultimately all monomials appearing in the second sum are of the form $u^{\beta+\mathbf{e}_1+\mathbf{e}_l}$ for some $|\beta| = 2k - 2$, which cannot ever have full degree $2k$ in the l -th coordinate. Similarly, all monomials appearing in the third sum are of the form $u^{\beta+2\mathbf{e}_1}$ for some $|\beta| = 2k - 2$, which cannot ever have full degree $2k$ in the l -th coordinate. This verifies the claim, and hence property (3). This completes the proof of Lemma 7.17.

8.4. Proof of Proposition 7.23 for case (1) in Lemma 7.20. In this proposition, we are in case B2 (for some $m_0 \geq 4$, p_{m_0} is Q -type) and under the assumptions of (1) in Lemma 7.20, so that $B_{2,\gamma^m(1)} \equiv 0$ and $\theta_l \neq \theta_{m(l)}$. Recall $\gamma^m(1)$ has a 1 in the m -th place for some $1 \leq m \leq n - 1$, and that $(\gamma(1); 0) = \mathbf{e}_{m(l)}$ for the $m(l)$ -th standard coordinate vector in \mathbb{R}^n . It is equivalent to verify (7.26) is not identically zero when we plug in u everywhere in place of \tilde{u} , so we now will verify that

$$(8.13) \quad \sum_{|\beta|=m_0-2} \left(\frac{\theta^\beta u^{\beta+\mathbf{e}_{m(l)}}}{\beta!} (\partial_{\beta+\mathbf{e}_l} p_{m_0})(u) - \frac{\theta^\beta u^{\beta+\mathbf{e}_l}}{\beta!} (\partial_{\beta+\mathbf{e}_{m(l)}} p_{m_0})(u) \right) \neq 0.$$

Since p_{m_0} is Q -type, say $m_0 = 2k$, then $p_{m_0}(u) = CQ(u)^k$; by normalizing (8.13) we may assume $C = 1$. We define a notation for the coefficients of p_{m_0} by

$$p_{m_0}(u) = \left(\sum_{j=1}^n \theta_j u_j^2 \right)^k = \sum_{|\omega|=k} c_{2\omega} u^{2\omega}$$

and note that each $c_{2\omega} \neq 0$. (In fact, we will also use the precise formula $c_{2\omega} = \frac{k!}{\omega!} \theta^\omega$.) We claim that the monomial $u^{m_0\mathbf{e}_l - \mathbf{e}_l + \mathbf{e}_{m(l)}}$ appears in (8.13) with the coefficient

$$(8.14) \quad \theta_l^k \left[m_0 - \theta_l \theta_{m(l)} m_0 \right].$$

In case (1) we assume $\theta_l \theta_{m(l)} = -1$, so that the coefficient (8.14) is nonzero, and hence the expression (8.13) is not identically zero, and the proposition is proved.

All that remains is to compute the claimed coefficient (8.14) of the monomial $u^{m_0\mathbf{e}_l - \mathbf{e}_l + \mathbf{e}_{m(l)}}$ in (8.13). First, for each fixed β with $|\beta| = 2k - 2$ we study the effect of the derivative $\partial_{\beta+\mathbf{e}_l}$ on p_{m_0} . We see that for such β , when $|\omega| = k$ the only indices ω such that u^ω survives in $\partial_{\beta+\mathbf{e}_l} p_{m_0}$ satisfy the relation $2\omega = \beta + \mathbf{e}_l + \mathbf{e}_g$ for each $1 \leq g \leq n$, and these lead to monomials in the first sum in (8.13) of the form $u^{\beta+\mathbf{e}_{m(l)}+\mathbf{e}_g}$, for each $1 \leq g \leq n$. For this exponent to be precisely the exponent $m_0\mathbf{e}_l - \mathbf{e}_l + \mathbf{e}_{m(l)}$ that we seek, we must have $\beta = m_0\mathbf{e}_l - \mathbf{e}_l - \mathbf{e}_g$ for some $1 \leq g \leq n$ and simultaneously $\beta \geq 0$. This can only occur if $g = l$, and we see that the only term of interest from this first sum comes from $\beta = (m_0 - 2)\mathbf{e}_l$ (and a corresponding term $2\omega = m_0\mathbf{e}_l$ in p_{m_0}).

We can analyze the contribution from the second sum in (8.13) similarly. For each fixed β in the sum, when $|\omega| = k$ the only indices ω such that u^ω survives in $\partial_{\beta+\mathbf{e}_{m(l)}} p_{m_0}$ satisfy the relation $2\omega = \beta + \mathbf{e}_{m(l)} + \mathbf{e}_g$ for each $1 \leq g \leq n$, and these lead to monomials in the second sum in (8.13) of the form $u^{\beta+\mathbf{e}_l+\mathbf{e}_g}$, for each $1 \leq g \leq n$. For this exponent to be precisely the exponent $m_0\mathbf{e}_l - \mathbf{e}_l + \mathbf{e}_{m(l)}$ that we seek, we must have $\beta = m_0\mathbf{e}_l - 2\mathbf{e}_l + \mathbf{e}_{m(l)} - \mathbf{e}_g$ for some $1 \leq g \leq n$ and simultaneously $\beta \geq 0$. This can only occur if $g = m(l)$ or if $g = l$ (in which case we use $m_0 \geq 4$). We see that the two terms of interest from this second sum come from $\beta = (m_0 - 2)\mathbf{e}_l$ (for a term $2\omega = (m_0 - 2)\mathbf{e}_l + 2\mathbf{e}_{m(l)}$ in p_{m_0}) and from $\beta = (m_0 - 3)\mathbf{e}_l + \mathbf{e}_{m(l)}$ (for a term $2\omega = (m_0 - 2)\mathbf{e}_l + 2\mathbf{e}_{m(l)}$ in p_{m_0}), respectively.

Now we compute the coefficients arising in these three terms and apply the precise formula for each coefficient $c_{2\omega}$. We see the total coefficient of $u^{m_0\mathbf{e}_l - \mathbf{e}_l + \mathbf{e}_{m(l)}}$ in (8.13) is precisely

$$\begin{aligned} &= c_{m_0\mathbf{e}_l}m_0(m_0 - 1) - c_{(m_0-2)\mathbf{e}_l+2\mathbf{e}_{m(l)}}\theta_{m(l)}\theta_l(m_0 - 2)(2) - c_{(m_0-2)\mathbf{e}_l+2\mathbf{e}_{m(l)}}(2) \\ &= \theta_l^k m_0(m_0 - 1) - \theta_l^{k-1}\theta_{m(l)}\frac{m_0}{2}\theta_{m(l)}\theta_l(m_0 - 2)(2) - \theta_l^{k-1}\theta_{m(l)}\frac{m_0}{2}(2) \\ &= \theta_l^k \left[m_0(m_0 - 1) - m_0(m_0 - 2) - \theta_l\theta_{m(l)}m_0 \right] = \theta_l^k \left[m_0 - \theta_l\theta_{m(l)}m_0 \right]. \end{aligned}$$

This proves (8.14) and hence Proposition 7.23 is proved, so that case (1) of Lemma 7.20 is proved in full.

8.5. Proof of Lemma 7.25: preliminary work for cases (2) and (4) in Lemma 7.20. In Lemma 7.20, recall that we suppose for some even $m_0 \geq 4$ that p_{m_0} is Q -type. For both case (2) and case (4) we have to apply Lemma 7.25, which simplifies the function $\Xi_{j,\gamma^{m(1)}}(\tilde{u})$, initially defined in (7.31). This lemma is an immediate consequence of the following lemma: for a certain type of polynomial p_j (and in particular for Q -type polynomials), the $\Xi_{j,\gamma^{m(1)}}$ function is closely related to a $B_{j,\gamma^{m(1)}}$ function, up to an ‘‘error term’’ that is a sum of derivatives of $D_{j,\gamma^{m(1)}}$ functions.

Lemma 8.2. *Suppose $p_j = \sum_{|\omega|=j} c_\omega u^\omega$ has the property that for any ω with $c_\omega \neq 0$, all entries in ω are even. Let $\gamma(1)$ be any fixed multi-index in $\mathbb{Z}_{\geq 0}^{n-1}$ such that $|\gamma(1)| = 1$. Then*

$$(8.15) \quad |u|^2 \Xi_{j,\gamma(1)}(\tilde{u}) - \left(\sum_{i=1}^n \theta_i u_i \partial_{u_i} \right) [D_{j,\gamma(1)}(\tilde{u})] = -\theta^{(\gamma(1);1)} B_{j,\gamma(1)}(\tilde{u}).$$

Thus in particular if j is even and p_j is of Q -type, then

$$(8.16) \quad |u|^2 \Xi_{j,\gamma(1)}(\tilde{u}) = -\theta^{(\gamma(1);1)} B_{j,\gamma(1)}(\tilde{u}).$$

The second identity in the lemma follows immediately from property (2) in Lemma 7.17 since $D_{j,\gamma(1)} \equiv 0$ for all $|\gamma(1)| = 1$ if p_j is Q -type. Thus it only remains to prove the first identity (8.15), and then Lemma 8.2, and hence Lemma 7.25 is proved. We first write down expressions for the relevant functions, and then prove the lemma.

8.5.1. Expressions for the main players. We now gather certain definitions in preparation for proving Lemma 8.2. It is again convenient to suppose that $\mathbf{e}_1, \dots, \mathbf{e}_n$ are the standard basis vectors in \mathbb{R}^n . To fix notation, we will assume throughout this section that the $(n-1)$ -coordinate multi-index $\gamma(1)$ has a 1 in the m -th place. As before in our convention (7.22), let $m(l) = m$ if $m < l$ and let $m(l) = m + 1$ if $m \geq l$. Then for example $(\gamma(1); 1) = \mathbf{e}_{m(l)} + \mathbf{e}_l$. With this notation, we can write

$$(8.17) \quad |u|^2 \Xi_{j,\gamma(1)}(\tilde{u}) = \sum_{|\beta|=j-2} \frac{1}{\beta!} \left[\theta^{\mathbf{e}_{m(l)}} (\partial_{\beta+\mathbf{e}_l} p_j)(\tilde{u}) u^{\beta+\mathbf{e}_{m(l)}} - \theta^{\mathbf{e}_l} (\partial_{\beta+\mathbf{e}_{m(l)}} p_j)(\tilde{u}) u^{\beta+\mathbf{e}_l} \right].$$

For concreteness, we denote the coefficients of p_j by $p_j(u) = \sum_{|\omega|=j} c_\omega u^\omega$, but we make no assumptions about the coefficients in this subsection. For any $|\beta| = j - 2$, since $|\omega| = j$, the only terms in p_j that survives $\partial_{\beta+\mathbf{e}_l}$ will correspond to ω that satisfy the relation $\omega = \beta + \mathbf{e}_l + \mathbf{e}_i$ for each $1 \leq i \leq n$. Similarly, in the second sum above, the only terms in p_j that survive $\partial_{\beta+\mathbf{e}_{m(l)}}$ will

correspond to $\omega = \beta + \mathbf{e}_{m(l)} + \mathbf{e}_i$ for each $1 \leq i \leq n$. Consequently, we can write

$$(8.18) \quad |u|^2 \Xi_{j,\gamma(1)}(\tilde{u}) = \sum_{|\omega|=j} \left[\sum_{\substack{i=1 \\ \omega - \mathbf{e}_l - \mathbf{e}_i \geq 0}}^n \theta^{\mathbf{e}_{m(l)} - \mathbf{e}_i} \frac{\omega!}{(\omega - \mathbf{e}_l - \mathbf{e}_i)!} u^{\omega + \mathbf{e}_{m(l)} - \mathbf{e}_i} \right. \\ \left. - \sum_{\substack{i=1 \\ \omega - \mathbf{e}_{m(l)} - \mathbf{e}_i \geq 0}}^n \theta^{\mathbf{e}_l - \mathbf{e}_i} \frac{\omega!}{(\omega - \mathbf{e}_{m(l)} - \mathbf{e}_i)!} u^{\omega - \mathbf{e}_{m(l)} + \mathbf{e}_i} \right] c_\omega.$$

Next, we recall that by definition,

$$D_{j,\gamma(1)}(\tilde{u}) = \sum_{|\beta|=j-1} \frac{1}{\beta!} \left[\theta^{\mathbf{e}_{m(l)}} (\partial_{\beta + \mathbf{e}_l} p_j)(\tilde{u}) u^{\beta + \mathbf{e}_{m(l)}} - \theta^{\mathbf{e}_l} (\partial_{\beta + \mathbf{e}_{m(l)}} p_j)(\tilde{u}) u^{\beta + \mathbf{e}_l} \right].$$

Since $|\beta| = j - 1$ while $|\omega| = j$ in each term of p_j , for each fixed β the only terms in p_j that can survive $\partial_{\beta + \mathbf{e}_l}$ in the first term must satisfy $\omega = \beta + \mathbf{e}_l$ (which only occurs if $\omega - \mathbf{e}_l \geq 0$); and the only terms in p_j that can survive $\partial_{\beta + \mathbf{e}_{m(l)}}$ in the second term must satisfy $\omega = \beta + \mathbf{e}_{m(l)}$ (which only occurs if $\omega - \mathbf{e}_{m(l)} \geq 0$). We get

$$D_{j,\gamma(1)}(\tilde{u}) = \sum_{|\omega|=j} \left[\theta^{\mathbf{e}_{m(l)}} \frac{\omega!}{(\omega - \mathbf{e}_l)!} u^{\omega + \mathbf{e}_{m(l)} - \mathbf{e}_l} - \theta^{\mathbf{e}_l} \frac{\omega!}{(\omega - \mathbf{e}_{m(l)})!} u^{\omega - \mathbf{e}_{m(l)} + \mathbf{e}_l} \right] c_\omega.$$

Here we use the convention that if $\omega - \mathbf{e}_l < 0$, the first term is omitted, and similarly for if $\omega - \mathbf{e}_{m(l)} < 0$, the second term is omitted. In any case we can see that

$$\omega! / (\omega - \mathbf{e}_l)! = \omega^{\mathbf{e}_l} = \omega_l,$$

since this coordinate is zero if $\omega - \mathbf{e}_l < 0$, and is $\omega_l \geq 1$ if $\omega - \mathbf{e}_l \geq 0$. This allows us to write in all cases that

$$(8.19) \quad D_{j,\gamma(1)}(\tilde{u}) = \sum_{|\omega|=j} \left[\omega^{\mathbf{e}_l} \theta^{\mathbf{e}_{m(l)}} u^{\omega + \mathbf{e}_{m(l)} - \mathbf{e}_l} - \omega^{\mathbf{e}_{m(l)}} \theta^{\mathbf{e}_l} u^{\omega - \mathbf{e}_{m(l)} + \mathbf{e}_l} \right] c_\omega.$$

Consequently, applying $\sum_i \theta_i u_i \partial_i$ to both sides,

$$(8.20) \quad \left(\sum_{i=1}^n \theta_i u_i \partial_{u_i} \right) [D_{j,\gamma(1)}(\tilde{u})] = \sum_{i=1}^n \sum_{|\omega|=j} \left[\omega^{\mathbf{e}_l} (\omega + \mathbf{e}_{m(l)} - \mathbf{e}_l)^{\mathbf{e}_i} \theta^{\mathbf{e}_{m(l)} + \mathbf{e}_i} u^{\omega + \mathbf{e}_{m(l)} - \mathbf{e}_l} \right. \\ \left. - \omega^{\mathbf{e}_{m(l)}} (\omega - \mathbf{e}_{m(l)} + \mathbf{e}_l)^{\mathbf{e}_i} \theta^{\mathbf{e}_l + \mathbf{e}_i} u^{\omega - \mathbf{e}_{m(l)} + \mathbf{e}_l} \right] c_\omega.$$

Here we note, for example, that for any i such that $\partial_{u_i} u^{\omega + \mathbf{e}_m - \mathbf{e}_n} \equiv 0$, we have $(\omega + \mathbf{e}_{m(l)} - \mathbf{e}_l)^{\mathbf{e}_i} = 0$, so that this expression is correct.

Finally, for reference, we merely recall that by definition,

$$(8.21) \quad B_{j,\gamma(1)}(\tilde{u}) = (\partial_{\mathbf{e}_l} p_j)(\tilde{u}) \tilde{u}^{\mathbf{e}_{m(l)}} - (\partial_{\mathbf{e}_{m(l)}} p_j)(\tilde{u}) \tilde{u}^{\mathbf{e}_l} \\ = \sum_{|\omega|=j} \left[\theta^{\omega + \mathbf{e}_{m(l)} - \mathbf{e}_l} \omega^{\mathbf{e}_l} u^{\omega + \mathbf{e}_{m(l)} - \mathbf{e}_l} - \theta^{\omega - \mathbf{e}_{m(l)} + \mathbf{e}_l} \omega^{\mathbf{e}_{m(l)}} u^{\omega - \mathbf{e}_{m(l)} + \mathbf{e}_l} \right] c_\omega.$$

8.5.2. *Proof of Lemma 8.2.* We suppose that $p_j(y) = \sum_{|\omega|=j} c_\omega y^\omega$ is a homogeneous degree j polynomial as stated, but make no further assumptions yet. Throughout the argument that follows,

we will use the fact that θ is a vector of entries ± 1 , so that θ^β only depends on the parity of each entry in β , and not on the sign of the entries in β . Now we consider the function

$$|u|^{2\Xi_{j,\gamma(1)}(\tilde{u})} - \left(\sum_{i=1}^n \theta_i u_i \partial_{u_i} \right) [D_{j,\gamma(1)}(\tilde{u})].$$

By comparing the expansions of (8.18) and (8.20), observe that the contributions from $i \neq m(l), l$ completely cancel. Indeed, first observe that the contribution to (8.18) from all $i \neq m(l), l$ is

$$(8.22) \quad \sum_{|\omega|=j} \sum_{i \neq m(l), l} [\theta^{\mathbf{e}_{m(l)} - \mathbf{e}_i} \omega^{\mathbf{e}_i + \mathbf{e}_l} u^{\omega + \mathbf{e}_{m(l)} - \mathbf{e}_l} - \theta^{\mathbf{e}_l - \mathbf{e}_i} \omega^{\mathbf{e}_i + \mathbf{e}_{m(l)}} u^{\omega - \mathbf{e}_{m(l)} + \mathbf{e}_l}] c_\omega.$$

(Initially in (8.18) we restricted to $\omega - \mathbf{e}_l - \mathbf{e}_i \geq 0$ when we evaluate $\omega! / (\omega - \mathbf{e}_l - \mathbf{e}_i)!$, for example. In that case, this ratio (for $i \neq l$) is $\omega_l \omega_i = \omega^{\mathbf{e}_l + \mathbf{e}_i}$. But indeed we can still see this is a valid expression for the ratio if $\omega - \mathbf{e}_l - \mathbf{e}_i < 0$, in which case by convention the ratio is 0, and also $\omega_l \omega_i = \omega^{\mathbf{e}_l + \mathbf{e}_i} = 0$. Thus the expression we wrote above is exact, without any need to state further restrictions.)

On the other hand, the contribution from all $i \neq m(l), l$ to (8.20) is

$$\sum_{i \neq m(l), l} \sum_{|\omega|=j} [\theta^{\mathbf{e}_{m(l)} + \mathbf{e}_i} \omega^{\mathbf{e}_l + \mathbf{e}_i} u^{\omega + \mathbf{e}_{m(l)} - \mathbf{e}_l} - \theta^{\mathbf{e}_l + \mathbf{e}_i} \omega^{\mathbf{e}_{m(l)} + \mathbf{e}_i} u^{\omega - \mathbf{e}_{m(l)} + \mathbf{e}_l}] c_\omega.$$

Note that since θ is a vector with entries ± 1 , $\theta^{\mathbf{e}_l + \mathbf{e}_i} = \theta^{\mathbf{e}_l - \mathbf{e}_i}$ and similarly for other expressions involving θ . Thus we see the contribution from $i \neq m(l), l$ in $(\sum_{i=1}^n \theta_i u_i \partial_{u_i}) [D_{j,\gamma(1)}(\tilde{u})]$ is identical to the $i \neq m(l), l$ terms in $|u|^{2\Xi_{j,\gamma(1)}}$, as expressed in (8.22).

What remains is to show the contributions from $i = l, m(l)$ yield the “error term” on the right-hand side of (8.15). The contribution from $i = l$ to (8.18) (after simplifying the powers of θ) is

$$\sum_{|\omega|=j} \left[\theta^{\mathbf{e}_{m(l)} + \mathbf{e}_l} \frac{\omega!}{(\omega - 2\mathbf{e}_l)!} u^{\omega + \mathbf{e}_{m(l)} - \mathbf{e}_l} - \frac{\omega!}{(\omega - \mathbf{e}_{m(l)} - \mathbf{e}_l)!} u^{\omega - \mathbf{e}_{m(l)} + \mathbf{e}_l} \right] c_\omega,$$

again understanding that a term appears only if the factorial in the denominator is evaluated at a non-negative number. We write this as

$$(8.23) \quad \sum_{|\omega|=j} [\theta^{\mathbf{e}_{m(l)} + \mathbf{e}_l} \omega^{\mathbf{e}_l} (\omega - \mathbf{e}_l)^{\mathbf{e}_l} u^{\omega + \mathbf{e}_{m(l)} - \mathbf{e}_l} - \omega^{\mathbf{e}_{m(l)} + \mathbf{e}_l} u^{\omega - \mathbf{e}_{m(l)} + \mathbf{e}_l}] c_\omega.$$

The contribution from $i = m(l)$ to (8.18) (after simplifying the powers of θ) is

$$\sum_{|\omega|=j} \left[\frac{\omega!}{(\omega - \mathbf{e}_{m(l)} - \mathbf{e}_l)!} u^{\omega + \mathbf{e}_{m(l)} - \mathbf{e}_l} - \theta^{\mathbf{e}_{m(l)} + \mathbf{e}_l} \frac{\omega!}{(\omega - 2\mathbf{e}_{m(l)})!} u^{\omega - \mathbf{e}_{m(l)} + \mathbf{e}_l} \right] c_\omega.$$

We write this as

$$(8.24) \quad \sum_{|\omega|=j} [\omega^{\mathbf{e}_{m(l)} + \mathbf{e}_l} u^{\omega + \mathbf{e}_{m(l)} - \mathbf{e}_l} - \theta^{\mathbf{e}_{m(l)} + \mathbf{e}_l} \omega^{\mathbf{e}_{m(l)}} (\omega - \mathbf{e}_{m(l)})^{\mathbf{e}_{m(l)}} u^{\omega - \mathbf{e}_{m(l)} + \mathbf{e}_l}] c_\omega.$$

On the other hand, let us consider the $i = l$ contribution to (8.20), which is

$$(8.25) \quad \sum_{|\omega|=j} [\theta^{\mathbf{e}_{m(l)} + \mathbf{e}_l} \omega^{\mathbf{e}_l} (\omega - \mathbf{e}_l)^{\mathbf{e}_l} u^{\omega + \mathbf{e}_{m(l)} - \mathbf{e}_l} - \omega^{\mathbf{e}_{m(l)}} (\omega + \mathbf{e}_l)^{\mathbf{e}_l} u^{\omega - \mathbf{e}_{m(l)} + \mathbf{e}_l}] c_\omega,$$

where we used that $\theta^{2\mathbf{e}_l} = 1$. The $i = m(l)$ contribution to (8.20) is

$$(8.26) \quad \sum_{|\omega|=j} [\omega^{\mathbf{e}_l} (\omega + \mathbf{e}_{m(l)})^{\mathbf{e}_{m(l)}} u^{\omega + \mathbf{e}_{m(l)} - \mathbf{e}_l} - \theta^{\mathbf{e}_{m(l)} + \mathbf{e}_l} \omega^{\mathbf{e}_{m(l)}} (\omega - \mathbf{e}_{m(l)})^{\mathbf{e}_{m(l)}} u^{\omega - \mathbf{e}_{m(l)} + \mathbf{e}_l}] c_\omega,$$

where we used that $\theta^{2\mathbf{e}_{m(l)}} = 1$.

Taking the $i = m(l), l$ contributions to Ξ in (8.23), (8.24) and subtracting the $i = m(l), l$ contributions to the D function in (8.25), (8.26), we see that there is some further cancellation. What is left is the statement:

$$|u|^2 \Xi_{j, \gamma(1)}(\tilde{u}) - \left(\sum_{i=1}^n \theta_i u_i \partial_{u_i} \right) [D_{j, \gamma(1)}(\tilde{u})] = \sum_{|\omega|=j} \left[(\omega^{\mathbf{e}_{m(l)}} (\omega + \mathbf{e}_l)^{\mathbf{e}_l} - \omega^{\mathbf{e}_{m(l)+\mathbf{e}_l}}) u^{\omega - \mathbf{e}_{m(l)+\mathbf{e}_l}} + (\omega^{\mathbf{e}_{m(l)+\mathbf{e}_l}} - \omega^{\mathbf{e}_l} (\omega + \mathbf{e}_{m(l)})^{\mathbf{e}_{m(l)}}) u^{\omega + \mathbf{e}_{m(l)-\mathbf{e}_l}} \right] c_\omega.$$

Now note that

$$\omega^{\mathbf{e}_{m(l)}} (\omega + \mathbf{e}_l)^{\mathbf{e}_l} - \omega^{\mathbf{e}_{m(l)+\mathbf{e}_l}} = \omega_{m(l)} (\omega_l + 1) - \omega_{m(l)} \omega_l = \omega_{m(l)}.$$

Similarly

$$\omega^{\mathbf{e}_{m(l)+\mathbf{e}_l}} - \omega^{\mathbf{e}_l} (\omega + \mathbf{e}_{m(l)})^{\mathbf{e}_{m(l)}} = \omega_{m(l)} \omega_l - \omega_l (\omega_{m(l)} + 1) = -\omega_l.$$

Thus we have

$$(8.27) \quad |u|^2 \Xi_{j, \gamma(1)}(\tilde{u}) - \left(\sum_{i=1}^n \theta_i u_i \partial_{u_i} \right) [D_{j, \gamma(1)}(\tilde{u})] = \sum_{|\omega|=j} \left[\omega_{m(l)} u^{\omega - \mathbf{e}_{m(l)+\mathbf{e}_l}} - \omega_l u^{\omega + \mathbf{e}_{m(l)-\mathbf{e}_l}} \right] c_\omega.$$

Now recall that by (8.21),

$$(8.28) \quad -\theta^{\mathbf{e}_{m(l)+\mathbf{e}_l}} B_{j, \gamma(1)}(\tilde{u}) = \sum_{|\omega|=j} \theta^\omega \left[\omega_{m(l)} u^{\omega - \mathbf{e}_{m(l)+\mathbf{e}_l}} - \omega_l u^{\omega + \mathbf{e}_{m(l)-\mathbf{e}_l}} \right] c_\omega.$$

Finally, we apply the hypothesis we assumed for p_j : for all ω such that $c_\omega \neq 0$, all entries in ω are even, so that $\theta^\omega = 1$. Under this condition, (8.28) is equal to (8.27) and Lemma 8.2 is proved. Thus Lemma 7.25 is proved.

8.6. Proof of Proposition 7.26 for case (2) in Lemma 7.20. Suppose that $j \geq 4$ is even, p_j is Q -type and $p_2 \neq 0$. Let $\gamma^m(1)$ of order 1 be fixed, with the property that $B_{2, \gamma^m(1)} \neq 0$ and $\theta_l = \theta_{m(l)}$. We now prove that there is a choice of $\gamma(2)$ with $|\gamma(2)| = 2$ such that

$$(8.29) \quad B_{2, \gamma^m(1)}(\tilde{u}) D_{j, \gamma(2)}(\tilde{u}) + \theta^{(\gamma^m(1); 1)} B_{2, \gamma(2)}(\tilde{u}) B_{j, \gamma^m(1)}(\tilde{u}) \neq 0.$$

By hypothesis $p_j(y) = CQ(y)^k$; by normalizing (8.29) we may assume $C = 1$. The second term in (8.29) contains $B_{j, \gamma^m(1)}(\tilde{u})$, which we notice has the following formula, when $p_j(u)$ is Q -type:

$$(8.30) \quad \begin{aligned} B_{j, \gamma^m(1)}(\tilde{u}) &= (\partial_{\mathbf{e}_1} p_j)(\tilde{u}) \tilde{u}^{\mathbf{e}_{m(l)}} - (\partial_{\mathbf{e}_{m(l)}} p_j)(\tilde{u}) \tilde{u}^{\mathbf{e}_l} \\ &= \tilde{u}^{\mathbf{e}_{m(l)}} \frac{j}{2} \left(\sum_{i=1}^n \theta_i u_i^2 \right)^{\frac{j}{2}-1} 2\theta_l (\theta_l u_l) - \tilde{u}^{\mathbf{e}_l} \frac{j}{2} \left(\sum_{i=1}^n \theta_i u_i^2 \right)^{\frac{j}{2}-1} 2\theta_{m(l)} (\theta_{m(l)} u_{m(l)}) \\ &= (\theta_{m(l)} - \theta_l) j \left(\sum_{i=1}^n \theta_i u_i^2 \right)^{\frac{j}{2}-1} u_{m(l)} u_l. \end{aligned}$$

Under the assumption that $\theta_l = \theta_{m(l)}$ we thus see that $B_{j, \gamma^m(1)} \equiv 0$ so that the expression (8.29) simplifies to $B_{2, \gamma^m(1)}(\tilde{u}) D_{j, \gamma(2)}(\tilde{u})$. By hypothesis, the first factor is $\neq 0$, and by property (3) of Lemma 7.17, since p_j is Q -type, there is some choice of $\gamma(2)$ with $|\gamma(2)| = 2$ such that $D_{j, \gamma(2)} \neq 0$. This completes the proof of Proposition 7.26, and hence case (2) of Lemma 7.20 is complete.

8.7. Proof of Proposition 7.28 for case (4) in Lemma 7.20. Suppose that $j \geq 4$ is even, p_j is Q -type and $p_2 \neq 0$. Let $\gamma^m(1)$ of order 1 be fixed, with the property that $B_{2,\gamma^m(1)} \neq 0$ and $\theta_l \neq \theta_{m(l)}$. We now prove that there is a choice of $\gamma(2)$ with $|\gamma(2)| = 2$ such that

$$(8.31) \quad B_{2,\gamma^m(1)}(\tilde{u})D_{j,\gamma(2)}(\tilde{u}) + \theta^{(\gamma^m(1);1)}B_{2,\gamma(2)}(\tilde{u})B_{j,\gamma^m(1)}(\tilde{u}) \neq 0,$$

or p_2 is Q -type in coordinate l and Q -type in coordinate $m(l)$.

We now set notation that we will use throughout the proof. Since p_j is Q -type, $p_j(y) = CQ(y)$, and by normalizing (8.31) we may assume $C = 1$. We will specify the coefficients of p_2 and p_j by

$$(8.32) \quad p_2(u) = \sum_{|\omega|=2} d_\omega u^\omega, \quad p_j(u) = \sum_{|\omega|=j} c_\omega u^\omega = \left(\sum_{i=1}^n \theta_i u_i^2 \right)^k = \sum_{|\beta|=k} \frac{k!}{\beta!} \theta^\beta u^{2\beta}.$$

We have the hypothesis that $\theta_{m(l)} \neq \theta_l$ or equivalently

$$(8.33) \quad \theta^{(\gamma^m(1);1)} = \theta^{\mathbf{e}_{m(l)}} \theta^{\mathbf{e}_l} = -1.$$

Lemma 8.3. *Given p_2 and p_j as in (8.32), and with the hypothesis (8.33), choose $\gamma(2)$ to have a 2 in the m th-place for the same m , so that*

$$(8.34) \quad (\gamma(2); 0) = 2\mathbf{e}_{m(l)}.$$

Within the homogeneous polynomial (8.31) of degree $j+2$, for this choice of $\gamma(2)$, we isolate three monomials, which temporarily we denote by

$$Au_{m(l)}^{j+2}, \quad Bu_{m(l)}^{j+1}u_l, \quad Cu_{m(l)}u_l^{j+1}.$$

For each index i in the range $1 \leq i \leq n$ with $i \neq l$ and $i \neq m(l)$, we also isolate the monomials

$$D_i u_i u_{m(l)}^{j+1} \quad \text{and} \quad E_i u_i u_l^{j+1}.$$

We claim these coefficients have the expressions

$$\begin{aligned} A &= d_{\mathbf{e}_{m(l)} + \mathbf{e}_l} c_{2\mathbf{e}_l + (j-2)\mathbf{e}_{m(l)}}, & c_{2\mathbf{e}_l + (j-2)\mathbf{e}_{m(l)}} &\neq 0 \\ B &= (d_{2\mathbf{e}_l} - d_{2\mathbf{e}_{m(l)}}) \theta_{m(l)}^k j - (\theta_{m(l)} - \theta_l) j \theta_{m(l)}^{k-1} d_{2\mathbf{e}_l} \\ C &= (d_{2\mathbf{e}_l} - d_{2\mathbf{e}_{m(l)}}) \theta_l^k j - (\theta_{m(l)} - \theta_l) j \theta_l^{k-1} d_{2\mathbf{e}_{m(l)}} \\ D_i &= d_{\mathbf{e}_i + \mathbf{e}_l} c_{2\mathbf{e}_l + (j-2)\mathbf{e}_{m(l)}} \theta_{m(l)} \theta_i, & c_{2\mathbf{e}_l + (j-2)\mathbf{e}_{m(l)}} &\neq 0, \quad i \neq l \\ E_i &= -d_{\mathbf{e}_i + \mathbf{e}_{m(l)}} c_{2\mathbf{e}_{m(l)} + (j-2)\mathbf{e}_l} \theta_l \theta_i, & c_{2\mathbf{e}_{m(l)} + (j-2)\mathbf{e}_l} &\neq 0, \quad i \neq m(l). \end{aligned}$$

Corollary 8.4. *In case (4) of Lemma 7.20 where $\theta_l \neq \theta_{m(l)}$, if (8.31) vanishes identically for this choice of $\gamma(2)$, then $d_{\mathbf{e}_{m(l)} + \mathbf{e}_l} = 0$ and $d_{2\mathbf{e}_l} = -d_{2\mathbf{e}_{m(l)}}$, and $d_{\mathbf{e}_i + \mathbf{e}_l} = 0$ for all $i \neq l$ and $d_{\mathbf{e}_i + \mathbf{e}_{m(l)}} = 0$ for all $i \neq m(l)$. In particular, then p_2 is Q -type in coordinates l and $m(l)$.*

Proof of the corollary. Of course if we assume that (8.31) vanishes identically, then $A = B = C = D_i = E_i = 0$, for all i . Observe that $A = 0$ forces $d_{\mathbf{e}_{m(l)} + \mathbf{e}_l} = 0$. Second, if $D_i = 0$ for all $i \neq l$ and $E_i = 0$ for all $i \neq m(l)$, then $d_{\mathbf{e}_i + \mathbf{e}_l} = 0$ for all $i \neq l$ and $d_{\mathbf{e}_i + \mathbf{e}_{m(l)}} = 0$ for all $i \neq m(l)$.

Finally, by studying differences of B and C we claim that $d_{2\mathbf{e}_l} = -d_{2\mathbf{e}_{m(l)}}$. Precisely, if k is even then $B - C = 0$ and the fact that $\theta_l \theta_{m(l)} = -1$ leads to this conclusion, since

$$B - C = -(\theta_{m(l)} - \theta_l) j [\theta_{m(l)} d_{2\mathbf{e}_l} - \theta_l d_{2\mathbf{e}_{m(l)}}].$$

If k is odd, then $B + C = 0$ and $\theta_l + \theta_{m(l)} = 0$ again leads to this conclusion, since

$$B + C = (d_{2\mathbf{e}_l} - d_{2\mathbf{e}_{m(l)}}) (\theta_{m(l)} + \theta_l) j - (\theta_{m(l)} - \theta_l) j [d_{2\mathbf{e}_l} + d_{2\mathbf{e}_{m(l)}}] = -(\theta_{m(l)} - \theta_l) j [d_{2\mathbf{e}_l} + d_{2\mathbf{e}_{m(l)}}].$$

Since we are in case (4) of Lemma 7.20 where $\theta_l \neq \theta_{m(l)}$, $d_{2\mathbf{e}_l} = -d_{2\mathbf{e}_{m(l)}}$ is equivalent to $\theta_l d_{2\mathbf{e}_l} = \theta_{m(l)} d_{2\mathbf{e}_{m(l)}}$, and p_2 must be Q -type in the coordinate l and the coordinate $m(l)$. \square

By the corollary, in case (4) of Lemma 7.20 either (8.31) is not identically zero for this choice of $\gamma(2)$, or p_2 is Q -type in coordinates l and $m(l)$. Thus the corollary implies Proposition 7.28, which was our goal in this section. All that remains is to prove Lemma 8.3.

8.7.1. *Proof of Lemma 8.3.* To compute the coefficients A, B, C and D_i, E_i we examine (8.31), which we write as $I + II$, with $I = B_{2,\gamma^m(1)}(\tilde{u})D_{j,\gamma(2)}(\tilde{u})$ and $II = -B_{2,\gamma(2)}(\tilde{u})B_{j,\gamma^m(1)}(\tilde{u})$.

We assemble expressions for the four main players. In I, by (8.21),

$$\begin{aligned} B_{2,\gamma^m(1)}(\tilde{u}) &= (\partial_{\mathbf{e}_l} p_2)(\tilde{u})\tilde{u}^{\mathbf{e}_{m(l)}} - (\partial_{\mathbf{e}_{m(l)}} p_2)(\tilde{u})\tilde{u}^{\mathbf{e}_l} \\ &= \sum_{i=1}^n d_{\mathbf{e}_i + \mathbf{e}_l}(\mathbf{e}_i + \mathbf{e}_l)_l \tilde{u}^{\mathbf{e}_{m(l)} + \mathbf{e}_i} - \sum_{i=1}^n d_{\mathbf{e}_i + \mathbf{e}_{m(l)}}(\mathbf{e}_i + \mathbf{e}_{m(l)})_{m(l)} \tilde{u}^{\mathbf{e}_l + \mathbf{e}_i}. \end{aligned}$$

(Here for example $(\mathbf{e}_i + \mathbf{e}_l)_l = 2$ if $i = l$ and 1 if $i \neq l$.) Next, by definition, using our choice of $\gamma(2)$ as in (8.34),

$$\begin{aligned} D_{j,\gamma(2)}(\tilde{u}) &= \sum_{\alpha \leq \gamma(2)} \sum_{|\beta|=j-2} \frac{(-1)^{|\alpha|}}{\alpha! \beta! (\gamma(2) - \alpha)!} \theta^{(\gamma(2) - \alpha; |\alpha|)} (\partial_{(\alpha; |\gamma(2) - \alpha|) + \beta} p_j)(\tilde{u}) u^{\beta + (\gamma(2) - \alpha; |\alpha|)} \\ &= \sum_{|\beta|=j-2} \frac{1}{\beta!} u^\beta \left(\frac{1}{2} u^{2\mathbf{e}_{m(l)}} (\partial_{2\mathbf{e}_l + \beta} p_j)(\tilde{u}) + u^{\mathbf{e}_{m(l)} + \mathbf{e}_l} (\partial_{\mathbf{e}_{m(l)} + \mathbf{e}_l + \beta} p_j)(\tilde{u}) + \frac{1}{2} u^{2\mathbf{e}_l} (\partial_{2\mathbf{e}_{m(l)} + \beta} p_j)(\tilde{u}) \right) \\ &= \sum_{|\beta|=j-2} \frac{1}{\beta!} \left(\frac{1}{2} (2\mathbf{e}_l + \beta)! c_{2\mathbf{e}_l + \beta} u^{2\mathbf{e}_{m(l)} + \beta} + (\mathbf{e}_{m(l)} + \mathbf{e}_l + \beta)! c_{\mathbf{e}_{m(l)} + \mathbf{e}_l + \beta} u^{\mathbf{e}_{m(l)} + \mathbf{e}_l + \beta} \right. \\ (8.35) \quad &\left. + \frac{1}{2} (2\mathbf{e}_{m(l)} + \beta)! c_{2\mathbf{e}_{m(l)} + \beta} u^{2\mathbf{e}_l + \beta} \right). \end{aligned}$$

In the second line we used that $\theta^{\mathbf{e}_{m(l)} + \mathbf{e}_l} = -1$ and $\theta^{2\mathbf{e}_{m(l)}} = \theta^{2\mathbf{e}_l} = 1$. In the third line we used that p_j is homogeneous of order j , with coefficients $p_j(w) = \sum_{|\omega|=j} c_\omega w^\omega$, so the only term that survives a derivative ∂_μ when $|\mu| = j$ corresponds precisely to $\omega = \mu$. (Moreover, since p_j is Q -type, we could reduce the above formula further since we know c_ω vanishes unless all entries in c_ω are even, in which case we have an explicit formula for c_ω .)

In II, by definition, and recalling the d_ω notation for coefficients of p_2 in (8.32),

$$\begin{aligned} B_{2,\gamma(2)}(\tilde{u}) &= \sum_{\alpha \leq \gamma(2)} \frac{(-1)^{|\alpha|}}{\alpha! (\gamma(2) - \alpha)!} \theta^{(\gamma(2) - \alpha; |\alpha|)} (\partial_{(\alpha; |\gamma(2) - \alpha|)} p_j)(\tilde{u}) u^{(\gamma(2) - \alpha; |\alpha|)} \\ &= \frac{1}{2} \theta^{2\mathbf{e}_{m(l)}} u^{2\mathbf{e}_{m(l)}} (\partial_{2\mathbf{e}_l} p_2)(\tilde{u}) - \theta^{\mathbf{e}_l + \mathbf{e}_{m(l)}} u^{\mathbf{e}_{m(l)} + \mathbf{e}_l} (\partial_{\mathbf{e}_{m(l)} + \mathbf{e}_l} p_2)(\tilde{u}) + \frac{1}{2} u^{2\mathbf{e}_l} (\partial_{2\mathbf{e}_{m(l)}} p_2)(\tilde{u}) \\ (8.36) \quad &= d_{2\mathbf{e}_l} u_{m(l)}^2 + d_{\mathbf{e}_{m(l)} + \mathbf{e}_l} u_{m(l)} u_l + d_{2\mathbf{e}_{m(l)}} u_l^2, \end{aligned}$$

where we used $\theta^{\mathbf{e}_{m(l)} + \mathbf{e}_l} = -1$. Finally, we already computed $B_{j,\gamma^m(1)}(\tilde{u})$ in (8.30). In what follows, recall that j is even; we will write $j = 2k$.

Contributions to A : Since every term in $B_{j,\gamma^m(1)}$ contains a factor u_l , term II does not contribute to the coefficient A . Within the product that defines I, the first sum in $B_{2,\gamma^m(1)}$ will contribute to A via the term $i = m(l)$ while the second sum does not contribute to A . Within the product that defines I, the only contribution to A from $D_{j,\gamma(2)}$ comes from the first summand when $\beta = (j - 2)\mathbf{e}_{m(l)}$. In total the contribution from the term I to A yields (upon recalling that II does not contribute to A):

$$A = d_{\mathbf{e}_{m(l)} + \mathbf{e}_l} \cdot \frac{1}{((j - 2)\mathbf{e}_{m(l)})!} \frac{1}{2} (2\mathbf{e}_l + (j - 2)\mathbf{e}_{m(l)})! \cdot c_{2\mathbf{e}_l + (j - 2)\mathbf{e}_{m(l)}} = d_{\mathbf{e}_{m(l)} + \mathbf{e}_l} c_{2\mathbf{e}_l + (j - 2)\mathbf{e}_{m(l)}}.$$

Now we note that under the assumption that p_j is Q -type, $c_{2\mathbf{e}_l+(j-2)\mathbf{e}_{m(l)}} \neq 0$, and the expression for A has been verified.

Contributions to B : The term II contributes to B via the product of the $i = m(l)$ term in (8.30) and the first term in (8.36). Precisely, the contribution of II to B is $-(\theta_{m(l)} - \theta_l)j\theta_{m(l)}^{k-1}d_{2\mathbf{e}_l}$. The contribution from the term I to B is

$$-2(d_{2\mathbf{e}_l} - d_{2\mathbf{e}_{m(l)}})c_{2\mathbf{e}_l+(j-2)\mathbf{e}_{m(l)}} = (d_{2\mathbf{e}_l} - d_{2\mathbf{e}_{m(l)}})\theta^{k\mathbf{e}_{m(l)}}j,$$

where we used $\theta^{\mathbf{e}_l+\mathbf{e}_{m(l)}} = -1$ and the fact that p_j is Q -type to show the coefficients c_ω vanish any time ω has an odd entry, and the rest of the coefficients have explicit formulae.

Contributions to C : The term II contributes to C via the product of the term $i = l$ in (8.30) and the last term in (8.36). Precisely, the contribution of II to C is $-(\theta_{m(l)} - \theta_l)j\theta_l^{k-1}d_{2\mathbf{e}_{m(l)}}$. The contribution from the term I to C is

$$-2(d_{2\mathbf{e}_l} - d_{2\mathbf{e}_{m(l)}})c_{2\mathbf{e}_{m(l)}+(j-2)\mathbf{e}_l} = (d_{2\mathbf{e}_l} - d_{2\mathbf{e}_{m(l)}})\theta^{k\mathbf{e}_l}j,$$

where we used $\theta^{\mathbf{e}_l+\mathbf{e}_{m(l)}} = -1$ and the fact that p_j is Q -type to show the coefficients c_ω vanish any time ω has an odd entry, and the rest of the coefficients have explicit formulae.

Contributions to D_i : In this case we use that $i \neq l, m(l)$. Since every term in $B_{j,\gamma^m(1)}$ contains a factor u_l , term II does not contribute to the coefficient D_i . Within term I, $B_{2,\gamma^m(1)}(\tilde{u})$ contributes a factor $d_{\mathbf{e}_i+\mathbf{e}_l}\tilde{u}^{e_{m(l)}+e_i}$ and $D_{j,\gamma(2)}(\tilde{u})$ contributes a factor $c_{2\mathbf{e}_l+(j-2)\mathbf{e}_{m(l)}}u_{m(l)}^j$ to the relevant monomial; there can be no other contributions since the coefficients c_ω in p_j vanish unless each entry of ω is even. This leads to

$$D_i = d_{\mathbf{e}_i+\mathbf{e}_l}\theta_{m(l)}\theta_i c_{2\mathbf{e}_l+(j-2)\mathbf{e}_{m(l)}}.$$

Contributions to E_i : We again use that $i \neq l, m(l)$. Since every term in $B_{j,\gamma(1)}$ contains a factor of $u_{m(l)}$, term II does not contribute to the coefficient E_i . Within the term I, $B_{2,\gamma^m(1)}(\tilde{u})$ contributes a factor $-d_{\mathbf{e}_i+\mathbf{e}_{m(l)}}\tilde{u}_l\tilde{u}_i$ and $D_{j,\gamma(2)}(\tilde{u})$ contributes a factor $c_{2\mathbf{e}_{m(l)}+(j-2)\mathbf{e}_l}u_l^j$ to the relevant monomial; there can be no other contributions since the coefficients c_ω in p_j vanish unless each entry of ω is even. This leads to

$$E_i = -d_{\mathbf{e}_i+\mathbf{e}_{m(l)}}\theta_l\theta_i c_{2\mathbf{e}_{m(l)}+(j-2)\mathbf{e}_l}.$$

This verifies Lemma 8.3.

We have completed the proofs of all the statements we deferred from Section 7. In particular, the proof of Theorem 7.1 is now complete, and consequently Theorem 1.1 is proved.

ACKNOWLEDGEMENTS

Anderson has been partially supported by NSF DMS-2231990, DMS-1502464, and an NSF Graduate Research Fellowship. She thanks Andreas Seeger for helpful conversations related to this project. Maldague is supported by the National Science Foundation under Award No. 2103249. Pierce has been partially supported by NSF CAREER grant DMS-1652173, DMS-2200470, a Sloan Research Fellowship, and a Joan and Joseph Birman Fellowship during portions of this work, and thanks the Hausdorff Center for Mathematics for a productive visit as a Bonn Research Chair in 2022.

REFERENCES

- [Bec22] L. Becker. Maximal polynomial modulations of singular Radon transforms, arxiv.2210.17263, 2022.
- [Car66] L. Carleson. On convergence and growth of partial sums of Fourier series. *Acta Math.*, 116:135–157, 1966.
- [DdF86] J. Duoandikoetxea and J. L. Rubio de Francia. Maximal and singular integral operators via Fourier transform estimates. *Invent. Math.*, 84:541–561, 1986.
- [Fef73] C. Fefferman. Pointwise convergence of Fourier series. *Ann. Math.*, 98(3):551–571, 1973.
- [GPRY17] S. Guo, L. B. Pierce, J. Roos, and P.-L. Yung. Polynomial Carleson operators along monomial curves in the plane. *J. Geom. Anal.*, 27(4):2977–3012, 2017.

- [Guo17] S. Guo. A remark on oscillatory integrals associated with fewnomials. *New York J. Math.*, 23:1733–1738, 2017.
- [Hun68] R. A. Hunt. On the convergence of Fourier series. Orthogonal expansions and their continuous analogues. In *Proceedings of the Conference held at Southern Illinois University, Edwardsville, 1967*. SIU Press, Carbondale, Illinois,, 1968.
- [JSW08] R. L. Jones, A. Seeger, and J. Wright. Strong variational and jump inequalities in harmonic analysis. *Trans. Amer. Math. Soc.*, 360(12):6711–6742, 2008.
- [Lie09] V. Lie. The weak- L^2 boundedness of the quadratic Carleson operator. *Geom. Funct. Anal.*, 19:457–497, 2009.
- [Lie17] V. Lie. A note on the polynomial Carleson operator in higher dimensions, arxiv:1712.03092, 2017.
- [Lie19] V. Lie. A unified approach to three themes in harmonic analysis (I & II), arxiv:1902.03807v2. 2019.
- [Lie20] V. Lie. The polynomial Carleson operator. *Ann. of Math. (2)*, 192(1):47–163, 2020.
- [LT00] M. Lacey and C. Thiele. A proof of boundedness of the Carleson operator. *Math. Res. Lett.*, 7(4):361–370, 2000.
- [MSZK20a] M. Mirek, E. M. Stein, and P. Zorin-Kranich. A bootstrapping approach to jump inequalities and their applications. *Anal. PDE*, 13(2):527–558, 2020.
- [MSZK20b] M. Mirek, E. M. Stein, and P. Zorin-Kranich. Jump inequalities via real interpolation. *Math. Ann.*, 376(1-2):797–819, 2020.
- [PY19] L. B. Pierce and P.-L. Yung. A polynomial Carleson operator along the paraboloid. *Rev. Mat. Iberoam.*, 35(2):339–422, 2019.
- [Ram21a] J. P. G. Ramos. The Hilbert transform along the parabola, the polynomial Carleson theorem and oscillatory singular integrals. *Math. Ann.*, 379(1-2):159–185, 2021.
- [Ram21b] J. P. G. Ramos. Uniform bounds for oscillatory and polynomial Carleson operators. *J. Fourier Anal. Appl.*, 27(1):Paper No. 5, 14, 2021.
- [Roo19] J. Roos. Bounds for anisotropic Carleson operators. *J. Fourier Anal. Appl.*, 25(5):2324–2355, 2019.
- [Sjö71] P. Sjölin. Convergence almost everywhere of certain singular integrals and multiple Fourier series. *Ark. Mat.*, 9:65–90, 1971.
- [Ste93] E. M. Stein. *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals*. Princeton University Press, Princeton NJ, 1993.
- [Ste95] E. M. Stein. Oscillatory integrals related to Radon-like transforms. In *Proceedings of the Conference in Honor of Jean-Pierre Kahane, Orsay*, volume special issue of *J. Fourier Anal. Appl.* 1995, pages 535–551, 1995.
- [SW78] E. M. Stein and S. Wainger. Problems in harmonic analysis related to curvature. *Bull. Amer. Math. Soc.*, 84:1239–1295, 1978.
- [SW01] E. M. Stein and S. Wainger. Oscillatory integrals related to Carleson’s theorem. *Math. Res. Lett.*, 8:789–800, 2001.
- [ZK21] P. Zorin-Kranich. Maximal polynomial modulations of singular integrals. *Adv. Math.*, 386:Paper No. 107832, 40, 2021.

CARNEGIE MELLON UNIVERSITY WEAN HALL, HAMMERSCHLAG DR., PITTSBURGH, PA 15213 USA

Email address: `tanders2@andrew.cmu.edu`

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MA 02142-4307 USA

Email address: `dmal@mit.edu`

DEPARTMENT OF MATHEMATICS, DUKE UNIVERSITY, 120 SCIENCE DRIVE, DURHAM NC 27708 USA

Email address: `pierce@math.duke.edu`