

# SPECTRAL ANALYSIS FOR A CLASS OF BOUNDED LINEAR OPERATORS

TEYLAMA HERVE MIABEY

ABSTRACT. We study the Spectral Analysis for a class of bounded linear operators  $T = D + u_1 \otimes v_1 + u_2 \otimes v_2 + \cdots + u_m \otimes v_m$  in a non Archimedean Hilbert space  $E_\omega$ , where  $D$  is a diagonal linear operator and  $F = u_1 \otimes v_1 + u_2 \otimes v_2 + \cdots + u_m \otimes v_m$  is a finite rank linear operator. In fact, our study is a generalization of the work of Diagana and McNeal. In their joint paper, Diagana and McNeal study the Spectral Analysis for rank 1 perturbation of a diagonal operator in  $E_\omega$  defined by  $A = D_\lambda + u \otimes v$ . Under some suitable assumption, they prove that

$$\sigma(A) = \{\theta_j\}_{j \geq 0} \cup \sigma_p(A)$$

where  $\sigma_p(A)$  is the set of all the eigenvalues of  $A$  and  $\theta = (\theta_j)_{j \in \mathbb{N}}$ ,  $\theta_j = \lambda_j + \omega_j \alpha_j \beta_j$ , with all  $j \in \mathbb{N}$ .

In this study of the Spectral Analysis, we use extensively the Theory of Fredholm Operators to deduce some of our main results. In a forthcoming paper, we will apply these methods to study a similar class of unbounded linear operators.

## 1. INTRODUCTION AND BACKGROUND

**Definition 1.1.** Let  $\mathbb{K}$  be an arbitrary field. We define an *absolute value* on  $\mathbb{K}$  to be the function

$$|\cdot| : \mathbb{K} \rightarrow [0, +\infty)$$

subject to the following conditions:

- (1)  $|x| = 0$  iff  $x = 0 \quad \forall x \in \mathbb{K}$
- (2)  $|xy| = |x||y| \quad \forall x, y \in \mathbb{K}$
- (3)  $|x + y| \leq |x| + |y| \quad \forall x, y \in \mathbb{K}$ .

$\mathbb{K}$  endowed with the aforementioned absolute value  $(\mathbb{K}, |\cdot|)$  is called a *non Archimedean value field* if condition (3) is replaced by the non Archimedean triangle inequality

$$(3') \quad |x + y| \leq \max\{|x|, |y|\}.$$

If  $|x| \neq |y|$ , then  $|x + y| = \max\{|x|, |y|\}$ .

*Remark 1.2.* Note that  $|x + y| \leq \max\{|x|, |y|\} \leq |x| + |y|$ , which implies that condition (3') is stronger than condition (3).

**Definition 1.3.** Let  $\mathbb{K}$  be a field. A non Archimedean metric  $d$  on  $\mathbb{K}$  is a function

$$d : \mathbb{K} \times \mathbb{K} \rightarrow \mathbb{R}$$

such that

- (1)  $d(x, y) \geq 0$  for all  $x, y \in \mathbb{K}$
- (2)  $d(x, y) = 0$  iff  $x = y$
- (3)  $d(x, y) = d(y, x)$  for all  $x, y \in \mathbb{K}$

$$(4) \quad d(x, y) \leq \max \{d(x, z), d(z, y)\} \text{ for all } x, y \text{ and } z \in \mathbb{K}.$$

The pair  $(\mathbb{K}, d)$  is called a non Archimedean metric space.

**Definition 1.4.** Let  $\mathbb{K}$  be a field and  $|\cdot|$  be a non Archimedean absolute value on  $\mathbb{K}$ . The non Archimedean distance between two elements  $x, y$  of  $\mathbb{K}$  is defined by  $d(x, y) = |x - y|$ .

*Remark 1.5.* (1) If  $|\cdot|$  is a non Archimedean absolute value on  $\mathbb{K}$ , then  $d(x, y) = |x - y|$  is a non Archimedean metric on  $\mathbb{K}$ .

(2) Clearly, the non Archimedean absolute value induces a metric  $d$  on  $\mathbb{K}$  defined above. Therefore, we can consider the notion of convergence in  $\mathbb{K}$ . A sequence  $(x_n)_{n \in \mathbb{N}} \subset \mathbb{K}$  is said to converge to  $x \in \mathbb{K}$  if  $d(x_n, x) \rightarrow 0$  as  $n \rightarrow \infty$ . The non Archimedean value field  $(\mathbb{K}, |\cdot|)$  is complete if the metric space  $(\mathbb{K}, d)$  is complete.  $(\mathbb{K}, d)$  is complete iff all the Cauchy sequences in  $\mathbb{K}$  converge in  $\mathbb{K}$ .

**Proposition 1.6.** Let  $\mathbb{K}$  be a field and  $|\cdot|$  be a normed value field. Then the following conditions are equivalent:

- (1)  $|\cdot|$  is a non Archimedean absolute value
- (2)  $|\cdot|$  is bounded in  $\mathbb{Z}$ .
- (3)  $|x| \leq 1$  for all  $x \in \mathbb{Z}$ .

*Proof.* (1)  $\Rightarrow$  (3):

Note that since  $|\cdot|$  is non Archimedean, it implies that  $|x| \leq 1$  since for any  $x \in \mathbb{Z}$

$$\begin{aligned} |x| &= |x \cdot 1| = \underbrace{|1 + 1 + \dots + 1|}_{x \text{ times}} \\ &\leq \max\{|1|, |1|, \dots, |1|\} \\ &\Rightarrow |x| \leq 1, \text{ since } \max\{|1|, |1|, \dots, |1|\} = 1 \end{aligned}$$

(3)  $\Rightarrow$  (2):

Since  $|x| \leq 1$  for any  $x \in \mathbb{Z}$ , it implies that  $|\cdot|$  is bounded in  $\mathbb{Z}$ . Therefore to complete the proof, it is sufficient to show that if  $|\cdot|$  is bounded in  $\mathbb{Z}$ , then  $|\cdot|$  is a non Archimedean absolute value, which is (2)  $\Rightarrow$  (1).

Suppose that  $|m| \leq M$  for all  $m \in Z$ . If  $x, y \in \mathbb{K}$  with  $n \in \mathbb{Z}$ , then

$$\begin{aligned} |x + y|^n &= |(x + y)^n| = \left| \sum_{i=0}^n \binom{n}{i} x^i y^{n-i} \right| \\ &\leq \sum_{i=0}^n \left| \binom{n}{i} \right| |x|^i |y|^{n-i} \quad \left( \text{Note that } \binom{n}{i} \leq M. \right) \\ &\leq M \sum_{i=0}^n (\max\{|x|, |y|\})^n = (n + 1)M \max\{|x|, |y|\}^n \end{aligned}$$

Therefore,  $|x + y|^n \leq (n + 1)M \max\{|x|, |y|\}^n$

Hence,  $\sqrt[n]{|(x + y)^n|} \leq (n + 1)^{1/n} M^{1/n} \max\{|x|, |y|\}$

$$|x + y| \leq \sqrt[n]{(n + 1)M} \max\{|x|, |y|\}$$

$$\text{but } \lim_{n \rightarrow \infty} \sqrt[n]{(n + 1)M} = 1$$

Therefore,  $|x + y| \leq \max\{|x|, |y|\}$ .

□

*Remark 1.7.* The difference between the Archimedean and non Archimedean Absolute values can be established in the following way: An Archimedean absolute value implies that for every  $x, y \in \mathbb{K}$  such that  $x \neq 0$ , there exists  $n \in \mathbb{Z}^+$  such that  $|nx| > |y|$ , i.e.  $\sup\{|n| : n \in \mathbb{Z}\} = +\infty$ . However the non Archimedean absolute value implies that  $\sup\{|n| : n \in \mathbb{Z}\} = 1$ .

**Proposition 1.8.** *Let  $(\mathbb{K}, |\cdot|)$  be a non Archimedean value field. Let  $|x| \neq |y|$ , then  $|x + y| = \max\{|x|, |y|\}$ .*

*Proof.* Let  $x, y \in \mathbb{K}$  and suppose that  $|x| > |y|$ . Then

$$\begin{aligned} |x + y| &\leq \max\{|x|, |y|\} \\ &\leq |x| = |(x + y) - y| \leq \max\{|x + y|, |y|\} \end{aligned}$$

And since  $|y| < |x|$

$$\max\{|x + y|, |y|\} = |x + y|$$

$$\text{but } |x + y| \leq \max\{|x|, |y|\} = |x|$$

We have thus established that  $|x + y| \leq |x|$  and  $|x| \leq |x + y|$ . Then it is clear that  $|x + y| = |x|$ ,  $\Rightarrow$  if  $|x| \neq |y|$  then  $|x + y| = \max\{|x|, |y|\}$ . □

**Proposition 1.9.** *Let  $(\mathbb{K}, |\cdot|)$  be a metric space on a non Archimedean value field and let  $x, y$  and  $z \in \mathbb{K}$  such that  $|x - z| \neq |z - y|$ . Then  $|x - y| = \max\{|x - z|, |z - y|\}$*

*Proof.* Suppose that  $|z - y| < |x - z|$ , then we have  $|x - y| \leq \max\{|x - z|, |z - y|\} = |x - z|$  and suppose now that  $|x - y| < |x - z|$ . Consequently,

$$|x - z| \leq \max\{|x - y|, |y - z|\} < |x - z|$$

which gives

$$|x - z| < |x - z|, \quad \text{which is a contradiction}$$

$$\therefore |x - z| = \max\{|x - z|, |z - y|\}$$

□

**Corollary 1.10.** *In the non Archimedean space, all triangles are isosceles.*

*Proof.* Consider  $x, y$  and  $z$  to be the vertices of a given triangle. Note that  $|x-y| = |(x-z)+(z-y)|$ . Suppose now  $|x-z| = |z-y|$ . Then two of the sides of the triangle have equal lengths, which implies that the triangle is isosceles. Suppose now  $|x-z| \neq |z-y|$  and that  $|x-z| > |z-y|$ .

Since we are in a non Archimedean field, we have  $|x-y| \leq \max\{|x-z|, |z-y|\}$  with equality when  $|x-z| \neq |z-y|$ . Therefore  $|x-y| = \max\{|x-z|, |z-y|\} = |x-z|$ . Therefore  $|x-y| = |x-z|$  and again two of the sides of the triangle have equal lengths. Therefore all triangles in the non Archimedean field are isosceles. □

**Definition 1.11.** Let  $(\mathbb{K}, |\cdot|)$  be a non Archimedean value field.

- (1) A sequence  $(x_n)_{n \in \mathbb{N}} \subset \mathbb{K}$  converges to  $x \in \mathbb{K}$  if for each  $\varepsilon > 0$ ,  $\exists \eta_0 \in \mathbb{N}$  such that  $|x_n - x| < \varepsilon$  whenever  $n \geq \eta_0$ .
- (2) A sequence  $(x_n)_{n \in \mathbb{N}} \subset \mathbb{K}$  is called a Cauchy sequence if for every  $\varepsilon > 0$ ,  $\exists \eta_0 \in \mathbb{N}$  such that  $|x_n - x_m| < \varepsilon$  whenever  $m \geq n \geq \eta_0$ .

Due to the non Archimedean triangle inequality, we conclude the following proposition:

**Proposition 1.12.** *If  $(\mathbb{K}, |\cdot|)$  is a non Archimedean value field,*

*a sequence  $(x_n)_{n \in \mathbb{N}} \subset \mathbb{K}$  is a Cauchy sequence iff*

$$|x_{n+1} - x_n| \rightarrow 0 \text{ as } n \rightarrow \infty.$$

*Proof.* ( $\Rightarrow$ ):

Suppose that  $(x_n)_{n \in \mathbb{N}}$  is a Cauchy sequence, then by the definition of the Cauchy sequence, for every  $\varepsilon > 0$ , there exists a  $\eta_0 \in \mathbb{N}$  such that  $|x_n - x_m| < \varepsilon$  whenever  $m \geq n \geq \eta_0$  which implies that  $|x_n - x_m| \rightarrow 0$  as  $n \rightarrow \infty$ .

( $\Leftarrow$ ): Suppose that  $|x_{n+1} - x_n| \rightarrow 0$  as  $n \rightarrow \infty$  and suppose that  $m \geq n$ . We have

$x_m - x_n = x_m - x_{m-1} + x_{m-1} - x_{m-2} + x_{m-2} - \cdots + x_{n+1} - x_n$ . Combining each pair of terms of the form  $x_t - x_{t-1}$ ,  $t = m, \dots, n$  and based on the earlier definition, we conclude that  $|x_m - x_n| \rightarrow 0$  as  $n, m \rightarrow \infty$ . Therefore  $(x_n)_{n \in \mathbb{N}}$  is a Cauchy sequence in  $\mathbb{K}$ . □

**Proposition 1.13.** *Let  $(\mathbb{K}, |\cdot|, d)$  be a complete non Archimedean field. The series  $\sum_{n=0}^{\infty} x_n$  converges to  $x \in \mathbb{K}$  iff  $x_n \rightarrow 0$  as  $n \rightarrow \infty$ .*

*Proof.* Set  $S_n = \sum_{r=0}^n x_r$ . It follows that  $x_n = S_n - S_{n-1}$ . Then if  $S_n \rightarrow S$ , then  $x_n \rightarrow 0$  in  $\mathbb{K}$  as  $n \rightarrow \infty$ . Conversely, if  $x_n \rightarrow 0$ , then  $(S_n)_{n \in \mathbb{N}}$  is a Cauchy sequence by the previous proposition. We know that  $\mathbb{K}$  is complete, hence there exists an element  $s \in \mathbb{K}$  such that  $S_n \rightarrow S$  as  $n \rightarrow \infty$ , which completes the proof. □

**Definition 1.14.** Let  $(\mathbb{K}, |\cdot|)$  be a non Archimedean value field.

- (1) An open ball of radius  $r$  and center  $x$  is defined by

$$B(x, r) = \{y \in \mathbb{K} : |x - y| < r\}.$$

- (2) A closed ball of radius  $r$  and center  $x$  is defined by

$$\overline{B(x, r)} = \{y \in \mathbb{K} : |x - y| \leq r\}.$$

- (3) A sphere in  $\mathbb{K}$  of radius  $r$  and center  $x$  is defined by  $S(x, r) = \{y \in \mathbb{K} : |x - y| = r\}$ .

**Proposition 1.15.** *Let  $(\mathbb{K}, |\cdot|)$  be a non Archimedean value field. Then*

- (1) *Every point that is contained in an open or closed ball in  $(\mathbb{K}, |\cdot|)$  is a center of the ball.*  
(2) *Every ball is both open and closed. These balls are called clopen.*  
(3) *If  $a, b$  are elements of  $\mathbb{K}$  and  $r_1, r_2 \in \mathbb{R}_+^\times$ , then  $B(a, r_1) \cap B(b, r_2) \neq \emptyset$  iff  $B(a, r_1) \subset B(b, r_2)$  or  $B(a, r_1) \supset B(b, r_2)$ , which implies that any two balls are either disjoint or one ball contains another.*

*Proof.* For a proof of (3), see Gouvea [1] pg 34.

- (1) Let  $B(a, r)$  be a ball in  $\mathbb{K}$  and let  $y \in B(a, r)$  and  $z \in B(y, r)$ . Therefore

$$\begin{aligned} |z - a| &= |z - y + y - a| \leq \max\{|z - y|, |y - a|\} \\ |z - a| &\leq \max\{|z - y|, |y - a|\} < r \end{aligned}$$

Therefore  $z \in B(a, r) \Rightarrow B(y, r) \subset B(a, r)$ .

$$\begin{aligned} w \in B(a, r) &\Rightarrow |w - y| = |w - a + a - y| \\ &\leq \max\{|w - a|, |a - y|\} < r \\ &\Rightarrow B(a, r) \subset B(y, r) \Rightarrow B(y, r) = B(a, r) \end{aligned}$$

Therefore every point in the closed or open ball is a center of the ball.

- (2) Let  $B(a, r)$  be a ball in  $(\mathbb{K}, |\cdot|)$ .  $z \in B(a, r), \exists \varepsilon > 0, \exists w \in B(a, r) \cap B(z, r)$ .

$$\text{Let } \varepsilon \leq r \Rightarrow |w - a| < r, |w - z| < \varepsilon$$

$$\Rightarrow |z - a| = |z - w + w - a| \leq \max\{|z - w|, |w - a|\} < r.$$

Since  $|z - w| < \varepsilon$  and  $|w - a| < r$ ,  $\Rightarrow z \in B(a, r)$ . Let  $\overline{B(a, r)}$  be a closed ball,  $w \in \overline{B(a, r)}$  and let  $\varepsilon \leq r$ . Then

$$\begin{aligned} z \in B(w, \varepsilon) &\Rightarrow |z - a| = |z - w + w - a| \\ |z - a| &\leq \max\{|z - w|, |w - a|\} < r. \end{aligned}$$

Since  $|z - w| < \varepsilon$  and  $|w - a| < r$ , then  $|z - a| \leq r \Rightarrow z \in \overline{B(a, r)}$ .

□

**Proposition 1.16.** *Let  $(\mathbb{K}, |\cdot|)$  be a non Archimedean field. Then*

- (1)  $\overline{B(0, 1)} = \{x \in \mathbb{K} : |x| \leq 1\}$  is a subring of  $\mathbb{K}$ .  
(2)  $B(0, 1) = \{x \in \mathbb{K} : |x| < 1\}$  is an ideal of  $\overline{B(0, 1)}$ .  
(3)  $B(0, 1)$  is a unique maximal ideal of  $\overline{B(0, 1)}$ .

(4) The Quotient  $\overline{B(0,1)}/B(0,1)$  is a field. It is the residue class field of  $|\cdot|$ .

(5)  $\mathbb{K}$  is a field of fractions of  $\overline{B(0,1)}$ .

*Proof.* See Gouvea [1]. □

## 2. THE VECTOR SPACE $\mathbb{K}^t$

**Definition 2.1.** Let  $(\mathbb{K}, |\cdot|)$  be a non Archimedean field. Just as in the classical case of  $\mathbb{R}$  and  $\mathbb{C}$ , there is a vector space  $\mathbb{K}^t$  which is associated to  $\mathbb{K}$  and defined as follows:

$$\mathbb{K}^t = \{ x = (x_1, x_2, \dots, x_t) : x_r \in \mathbb{K}, r = 1, \dots, t \}.$$

Note that the norm defined by  $\|x\|_t = \max_{1 \leq r \leq t} |x_r|$  for all  $x = (x_1, x_2, \dots, x_t) \in \mathbb{K}^t$  is a non Archimedean norm over  $\mathbb{K}^t$ . The inner product on  $\mathbb{K}^t$  is defined by

$$\langle x, y \rangle_t = \sum_{r=1}^t x_r y_r = x_1 y_1 + x_2 y_2 + \dots + x_t y_t$$

where  $x = (x_1, x_2, \dots, x_t)$  and  $y = (y_1, y_2, \dots, y_t) \in \mathbb{K}^t$ .

The inner product satisfies the Cauchy-Schwartz inequality i.e.  $|\langle x, y \rangle_t| \leq \|x\|_t \|y\|_t, x, y \in \mathbb{K}^t$ , and any element  $x = (x_1, x_2, \dots, x_t)$  can be written uniquely as  $x = \sum_{r=1}^t x_r e_r$ , where  $e_r = (0, 0, \dots, 1, 0, \dots, 0)$  and  $(e_r)_{r=1, \dots, t}$  is the orthonormal base of  $\mathbb{K}^t$ .

## 3. A VALUATION

**Definition 3.1.** A valuation on an arbitrary field  $\mathbb{K}$  is a function  $V : \mathbb{K} \rightarrow \mathbb{R} \cup \{\infty\}$  such that for every  $x, y \in \mathbb{K}$ , we have the following properties:

- (1)  $V(x) = \infty$  iff  $x = 0$
- (2)  $V(xy) = V(x) + V(y)$
- (3)  $V(x + y) \geq \min\{V(x), V(y)\}$

*Remark 3.2.* For any element  $c \in \mathbb{K}$  such that  $c > 1$ ,  $c^{-V(\cdot)} = |x|$  defines a non Archimedean absolute value on  $\mathbb{K}$ . The valuation is called non trivial if there exists an element  $a \in \mathbb{K}^*$  such that  $V(a) \neq 0$ .

**Definition 3.3.** A valuation  $V$  on  $\mathbb{K}$  is said to be a discrete valuation if the totally ordered group  $V(\mathbb{K}^*)$  is isomorphic to  $\mathbb{Z}$ .

**Lemma 3.4.** Let  $V$  be a discrete valuation,  $x, y \in \mathbb{K}$ ,  $V(x) \neq V(y)$ . Then we have

$$V(x + y) = \min\{V(x), V(y)\}$$

*Proof.* Since  $V(x) \neq V(y)$ , suppose without loss of generality that  $V(x) < V(y)$ . Suppose now that  $V(x + y) \neq \min\{V(x), V(y)\}$ . We have  $V(x + y) > V(x)$ .

Note that  $V(x) = V[(x + y) - y] \geq \min\{V(x + y), V(y)\}$ , since  $V(-y) = V(y)$ . Therefore,  $V(x) = V[(x + y) + y] \geq \min\{V(x + y), V(y)\} > V(x)$ . Hence

$V(x) > V(x)$ , a contradiction. Therefore  $V(x + y) = \min\{V(x), V(y)\}$ . □

**Example 3.5.** Let  $\mathcal{F} = \mathbb{K}(x)$ . Define  $V_{p(x)} : \mathcal{F} \rightarrow \mathbb{Z} \cup \{\infty\}$ , satisfying the conditions of Definition 3.1, where  $p(x) \in \mathbb{K}[x]$  is an irreducible monic polynomial.  $V_{p(x)}$  is a discrete valuation. We define the discrete valuation ring as the set

$$\mathcal{O}_{p(x)} = \left\{ \frac{f(x)}{g(x)} : f(x), g(x) \in \mathbb{K}[x], p(x) \nmid g(x) \right\}$$

and its unique maximal ideal

$$P_{p(x)} = \left\{ \frac{f(x)}{g(x)} : f(x), g(x) \in \mathbb{K}[x], p(x) \mid f(x) \text{ and } p(x) \nmid g(x) \right\}.$$

There is another valuation ring on  $\mathbb{K}(x)/\mathbb{K}$ :

$$\mathcal{O}_\infty = \left\{ \frac{f(x)}{g(x)} : f(x), g(x) \in \mathbb{K}[x], \deg f(x) \leq \deg g(x) \right\}$$

and its maximal ideal

$$P_\infty = \left\{ \frac{f(x)}{g(x)} : f(x), g(x) \in \mathbb{K}[x], \deg f(x) < \deg g(x) \right\}.$$

#### 4. THE FIELD $\mathbb{Q}_p$

**Definition 4.1.** A p-adic valuation on  $\mathbb{Z}$  is a function  $V_p : \mathbb{Z} - \{0\} \rightarrow \mathbb{R}$  where  $p \in \mathbb{Z}$  is a prime number and  $V_p(t)$  is a unique positive number such that  $t = p^{V_p(t)} \cdot q$  and  $(p, q) = 1$ .

*Remark 4.2.* The valuation  $V_p$  on  $\mathbb{Z}$  can be extended to the field of rational numbers  $\mathbb{Q}$ , i.e. if  $x \in \mathbb{Q} - \{0\}$ , then  $x = t/s$ , where  $(t, s) \in \mathbb{Z}^2 - \{0\}$ .

$$V_p(x) = V_p\left(\frac{t}{s}\right) = V_p\left(t \cdot \frac{1}{s}\right) = V_p(t) - V_p(s). \text{ If } x = 0, \text{ we have } V_p(0) = \infty.$$

**Corollary 4.3.** Consider  $(x, y) \in \mathbb{Z}^2$ . We have the following properties of the valuation:

- (1)  $V_p(x) = \infty$  iff  $x = 0$ .
- (2)  $V_p(x \cdot y) = V_p(x) + V_p(y)$
- (3)  $V_p(x + y) \geq \min\{V_p(x), V_p(y)\}$ , with equality if  $V_p(x) \neq V_p(y)$ .

*Proof.* (1) is deduced from the definition.

- (2) Let  $x, y \in \mathbb{Z}$ . Then  $x = p^r x'$ ,  $y = p^s y'$ , where  $x', y'$  are integers and  $p$  is relatively prime with respect to  $x'$  and  $y'$  and  $r, s \in \mathbb{Z}$ . Suppose that  $r > s$ .

$$\begin{aligned} x \cdot y &= p^r \cdot x' \cdot p^s \cdot y' \\ &= p^r \cdot p^s \cdot x' y' \\ &= p^{r+s} \cdot x' y'. \\ V_p(x, y) &= V_p(p^{r+s} \cdot x' y') \\ &= V_p(p^{r+s}) + V_p(x' y') \\ &= r + s, \quad \text{since } V_p(x) = r \text{ and } V_p(y) = s \\ \therefore V_p(x \cdot y) &= V_p(x) + V_p(y) \end{aligned}$$

- (3) In this case, suppose that  $x$  and  $y$  are rational numbers. Then we have  $x = p^r \frac{a}{b}$  and  $y = p^s \frac{c}{d}$ , where  $a, b, c$  and  $d \in \mathbb{Z}$ ,  $p$  is relatively prime with respect to  $a, b, c$  and  $d$  and  $r, s \in \mathbb{Z}$ . Now suppose that  $r = s$ . Then

$$\begin{aligned} x + y &= p^r \left( \frac{a}{b} + \frac{c}{d} \right) \\ &= p^r \cdot \left( \frac{ad + bc}{bd} \right) \\ \Rightarrow V_p(x + y) &= V_p \left[ p^r \left( \frac{ad + bc}{bd} \right) \right] \\ \Rightarrow V_p(x + y) &\geq r \quad \text{since } (p, bd) = 1. \end{aligned}$$

Assume that  $r \neq s$  and set  $s > r$ . Then we have

$$\begin{aligned} x + y &= p^r \frac{a}{b} + p^s \frac{c}{d} \\ &= p^r \left( \frac{a}{b} + p^{s-r} \cdot \frac{c}{d} \right) \\ &= p^r \left( \frac{ad + bc \cdot p^{s-r}}{bd} \right). \end{aligned}$$

Note that since  $s > r \Rightarrow s - r > 0$  and since  $(p, bd) = 1 \Rightarrow V_p(x + y) = r = \min\{V_p(x), V_p(y)\}$ .

□

**Definition 4.4.** For  $x \in \mathbb{Q}$ , the p-adic absolute value of  $x$  is given by

$$|x|_p = \begin{cases} p^{-V_p(x)} & \text{if } x \neq 0 \\ p^{-\infty} = 0 & \text{if } x = 0 \end{cases}$$

where  $V_p(s)$  is a valuation of  $x \in \mathbb{Q}$ .

*Remark 4.5.* The p-adic absolute value is a non Archimedean valuation.

**Proposition 4.6.** Consider the function  $|\cdot|_p : \mathbb{Q} \rightarrow \mathbb{R}^+ \cup \{0\}$ . Then we have the following properties:

- (1)  $|x|_p = 0$  iff  $x = 0$ .
- (2)  $|xy|_p = |x|_p |y|_p$ .
- (3)  $|x + y|_p \leq \max\{|x|_p, |y|_p\}$ .
- (4)  $|x|_p \neq |y|_p \Rightarrow |x + y|_p = \max\{|x|_p, |y|_p\}$ .

Hence  $(\mathbb{Q}, |\cdot|_p)$  is a non Archimedean value field.

*Remark 4.7.* The p-adic absolute value  $|\cdot|_p$  induces a metric  $d$  on  $\mathbb{Q} \times \mathbb{Q}$  defined by  $d(x, y) = |x - y|_p$  for all  $x, y \in \mathbb{Q}$ .

PROOF OF PROPOSITION (4.6)

- (1)  $|x|_p = 0 = p^{-V_p(x)} \Rightarrow V_p(x) = +\infty \Rightarrow x = 0$ .



(2)

$$\begin{aligned}
|xy|_p &= |p^{V_p(x)} \cdot q \cdot p^{V_p(y)} \cdot q'|_p \\
&= |p^{V_p(x)+V_p(y)}|_p = p^{-(V_p(x)+V_p(y))} \\
&= p^{-V_p(x)} \cdot p^{-V_p(y)} \\
&= |x|_p \cdot |y|_p
\end{aligned}$$

(3)

$$\begin{aligned}
|x+y|_p &= |p^{V_p(x)} \cdot q + p^{V_p(y)} \cdot q'|_p \\
&\quad \text{Note that by definition,} \\
|x+y|_p &= p^{-V_p(x+y)} \\
\text{but } p^{V_p(x+y)} &\leq p^{-\min\{V_p(x), V_p(y)\}} \\
&= \max \left\{ p^{-V_p(x)}, p^{-V_p(y)} \right\} \\
&= \max \{|x|_p, |y|_p\}
\end{aligned}$$

Hence  $|x+y| \leq \max\{|x|_p, |y|_p\}$ .

## 5. CONSTRUCTION OF $\mathbb{Q}_p$

**Proposition 5.1.** *For all primes  $p \in \mathbb{Z}$ , the field  $\mathbb{Q}$  is not complete with respect to the non Archimedean absolute value  $|\cdot|_p$ .*

*Proof.* See Gouvea[1], page 50. □

**Definition 5.2.** The field of p-adic numbers  $\mathbb{Q}_p$  is the completion of the non Archimedean value field  $(\mathbb{Q}, |\cdot|_p)$ .

**Lemma 5.3.** *Let  $|\cdot|_p$  be a non Archimedean absolute value on  $\mathbb{Q}$ . Then the extension of  $|\cdot|_p$  to  $\mathbb{Q}_p$  is also a non Archimedean absolute value.*

*Proof.* Let  $a, b \in \mathbb{Q}_p$  and let  $\{a_t\}_{t \in \mathbb{N}}$  and  $\{b_t\}_{t \in \mathbb{N}}$  be sequences that converge to  $a$  and  $b$  respectively, i.e.  $\lim_{t \rightarrow \infty} \{a_t\} = a$  and  $\lim_{t \rightarrow \infty} \{b_t\} = b$ . We then have

$$\begin{aligned}
\lim_{t \rightarrow \infty} |a_t + b_t|_p &\leq \max(|a_t|, |b_t|) \\
&\Rightarrow |a + b| \leq \max\{|a|, |b|\}
\end{aligned}$$

which concludes the proof. □

*Remark 5.4.* There are several methods that are used to complete  $(\mathbb{Q}, |\cdot|_p)$ . We will cover three methods of completion: the analytic method, the semi algebraic method and the algebraic method.

(1) Analytic Method:

Let  $\mathcal{C}$  be the set of all Cauchy sequences in  $\mathbb{Q}$  with respect to the Absolute value  $|\cdot|_p$ . We define the following relation: Two sequences  $\{x_n\}$  and  $\{y_n\}$  of  $\mathcal{C}$  are related, i.e.  $\{x_n\} \sim \{y_n\}$  iff  $\forall \varepsilon > 0, \exists N > 0$  such that  $n \geq N \Rightarrow |x_n - y_n| < \varepsilon$ .

We can easily verify that  $\sim$  is an equivalence relation. We define  $\mathbb{Q}_p = \mathcal{C} / \sim$ , where addition is defined by  $\overline{\{x_n\}} + \overline{\{y_n\}} = \overline{\{x_n\} + \{y_n\}}$  and multiplication by  $\overline{\{x_n\}} \cdot \overline{\{y_n\}} = \overline{\{x_n\} \cdot \{y_n\}}$ .

Therefore the elements of  $\mathbb{Q}_p$  are the equivalence classes of the Cauchy sequences.

## 6. COMPUTATION OF THE ESSENTIAL SPECTRUM OF $D$ , $\sigma_e D$ WHERE $D$ IS A BOUNDED LINEAR OPERATOR

**Definition 6.1.** Let  $\mathbb{K}$  be a non Archimedean field and let  $E$  and  $F$  be vector spaces over  $\mathbb{K}$  such that  $f : E \rightarrow F$  is a linear map. Let  $\lambda \in \mathbb{K}$ . We define the *weak multiplicity*  $\lambda$  relative to  $f$  to be the dimension of the kernel  $N(\lambda I - D)$ . The multiplicity  $\lambda$  is said to be finite if the dimension of the kernel  $N(\lambda I - D)$  is finite. The multiplicity  $\lambda$  is said to be infinite if the dimension of  $N(\lambda I - D)$  is infinite. We denote  $\eta(\lambda I - D)$  to be the dimension of the kernel of  $\lambda I - D$ .

**Theorem 6.2.** Let  $D : E_\omega \rightarrow E_\omega$  be a bounded diagonal operator such that  $D(u) = \sum \lambda_i u_i e_i$  with  $u = \sum u_i e_i$  for  $\lambda_i \in \mathbb{K}$ ,  $\forall i \in \mathbb{N}$ . Let  $\lambda$  be an eigenvalue of  $D$  with finite multiplicity with respect to  $D$ . Then  $\lambda I - D$  is a Fredholm operator of index 0.

*Proof.* 1. We want to show that  $\eta(\lambda I - D)$  is equal to the number of  $\lambda_i \in \Lambda = \{\lambda_k\}_{k \in \mathbb{N}}$  such that  $\lambda_k = \lambda$ .

$$\begin{aligned} \text{Let } u \in N(\lambda I - D) &\Rightarrow (\lambda I - D)(u) = 0 \\ &\Rightarrow \sum (\lambda u_i - \lambda_i u_i) e_i = 0 \\ &\Rightarrow \sum (\lambda - \lambda_i) u_i e_i = 0 \\ &\Leftrightarrow (\lambda - \lambda_i) u_i = 0 \quad \forall i \in \mathbb{N} \end{aligned}$$

For  $k \in \mathbb{K}$ , if  $\lambda_k \neq \lambda$ ,  $u_k = 0$ . This implies that  $N(\lambda I - D)$  is generated by  $e_i$  such that  $\lambda_i = \lambda \Rightarrow N(\lambda I - D) = \text{Cardinal}\{e_i | \lambda_i = \lambda\}$

2. We want to compute the dimension of the cokernel of  $(\lambda I - D)$ , denoted by  $\delta(\lambda I - D)$ . Let  $v \in \text{Im}(\lambda I - D) \Rightarrow \exists u \in E_\omega$  such that

$$\begin{aligned} (\lambda I - D)(u) = v &\Leftrightarrow \sum_{i \in \mathbb{N}} (\lambda - \lambda_i) u_i e_i = \sum_{i \in \mathbb{N}} v_i e_i \\ &\Leftrightarrow (\lambda - \lambda_i) u_i = v_i \text{ if } \lambda = \lambda_i \Rightarrow v_i = 0 \forall i \in \mathbb{N} \end{aligned}$$

This implies  $\text{Im}(\lambda I - D)$  is generated by the  $e_k$  such that  $\lambda_k \neq \lambda \Rightarrow E_\omega / \text{Im}(\lambda I - D)$  is of the dimension equal to the number of  $\{e_i | \lambda_i = \lambda\}$  or  $\dim(E_\omega / \text{Im}(\lambda I - D)) = \delta(\lambda I - D)$ . Therefore  $\delta(\lambda I - D) = \#\{e_i | \lambda_i = \lambda\}$

3. We want to show that if  $\lambda$  is an eigenvalue with finite multiplicity with respect to  $D$ , then the range, denoted by  $R(\lambda I - D)$  is closed in  $E_\omega$ . Let  $F$  be a vector subspace of  $E_\omega$ . We define the orthogonal set of  $F$ , denoted  $F^\perp$  to be the set of all  $v \in E_\omega$  such that  $(u, v) = 0 \forall u \in F$ , i.e.  $F^\perp = \{v \in E_\omega | (u, v) = 0 \forall u \in F\}$ .  $\square$

**Lemma 6.3.**  $F^\perp$  is a closed vector subspace of  $E_\omega$  for all vector subspaces  $F$  of  $E_\omega$ .

*Proof.* Let  $v_n$  be a sequence of  $F^\perp$  that converges in  $E_\omega$  to some  $v \in E_\omega$ , i.e.  $v = \lim_{n \rightarrow \infty} (v_n)$ . Then we have that  $\forall u \in F$ ,  $(u, v) = \lim_{n \rightarrow \infty} (u, v_n)$ .  $(u, v_n) = 0 \forall n \in \mathbb{N}$ , since  $v_n \in F^\perp$ . This implies  $(u, v) = 0, \Rightarrow v \in F^\perp$ . Therefore  $F^\perp$  is closed in  $E_\omega$ .  $\square$

**Lemma 6.4.** Let  $\lambda$  be an eigenvalue of  $D$  with finite multiplicity with respect to  $D$ . Then  $R(\lambda I - D) = N(\lambda I - D)^\perp$ , therefore  $R(\lambda I - D)$  is closed in  $E_\omega$ .

*Proof.* We have shown in Theorem 1 (2) that  $R(\lambda I - D)$  is generated by  $E_k$  such that  $\lambda_k \neq \lambda$  and in the first part of Theorem 1 (1) that  $N(\lambda I - D)$  is generated by the  $e_i$  such that  $\lambda_i = \lambda$ .

(1°) Let  $v \in R(\lambda I - D)$  with  $v = \sum v_i e_i$  and  $v_i = 0$  if  $\lambda_i = \lambda$ . Let  $u \in N(\lambda I - D)$  with  $u = \sum u_j e_j$  with  $u_j = 0$  if  $\lambda_j \neq \lambda$ .  $(u, v) = \sum u_k v_k$  if  $\lambda_k = \lambda \Rightarrow v_k = 0$ , and if  $\lambda_k \neq \lambda$ ,  $u_k = 0 \Rightarrow (u, v) = 0 \Rightarrow v \in N(\lambda I - D)^\perp$ . Therefore  $R(\lambda I - D) \subseteq N(\lambda I - D)^\perp$ .

(2°) Let  $v \in N(\lambda I - D)^{perp}$ ,  $v = \sum_{i \in \mathbb{N}} v_i e_i$ . Let  $e_j$  be such that  $\lambda_j = \lambda \Rightarrow e_j \in N(\lambda I - D)$ . Then we have

$$\begin{aligned} (e_j, v) &= 0. \text{ Since} \\ (e_j, v) &= \sum v_i (e_j, e_i) \text{ and} \\ (e_j, e_i) &= \begin{cases} 0 & \text{if } i \neq j \\ 1 & \text{if } i = j \end{cases} \\ \Rightarrow (e_j, v) &= v_j \Rightarrow v_j = 0 \end{aligned}$$

Therefore  $v$  belongs to the subspace generated by the  $e_k$  such that  $\lambda_k \neq \lambda$ . Therefore we have  $N(\lambda I - D)^\perp \subseteq R(\lambda I - D)$ . Therefore  $N(\lambda I - D)^\perp = R(\lambda I - D) \Rightarrow R(\lambda I - D)$  is closed in  $E_\omega$ .

Since  $\eta(\lambda I - D)$  and  $\delta(\lambda I - D)$  are finite such that  $\eta(\lambda I - D) = \delta(\lambda I - D) = \#\{e_i \mid \lambda_i = \lambda\} \Rightarrow X(\lambda I - D) = 0$ . Since  $R(\lambda I - D)$  is closed,  $(\lambda I - D)$  is a Fredholm operator of index 0.

□

**Definition 6.5.** We define the *proper spectrum* of  $D$  to be the set of all  $\lambda_k \in \sigma(D)$  with multiplicity 0 with respect to  $D$  denoted by

$$\begin{aligned} \sigma'_e(D) &= \{\lambda \in \mathbb{K}, (\lambda I - D) \text{ is injective but not surjective}\} \\ N(\lambda I - D) &= \{0\} \end{aligned}$$

Note that if  $\lambda \in \sigma(D)$  and  $\lambda \notin \sigma_p(D) = \Lambda$ ,  $\lambda$  is not an eigenvalue  $\Rightarrow N(\lambda I - D) = \{0\} \Rightarrow \lambda$  is of multiplicity 0 with respect to  $D$ .

**Definition 6.6.** We define the *improper essential spectrum* of  $D$  to be the set of all  $\lambda_k (k \in \mathbb{N})$  with infinite multiplicity with respect to  $D$ , denoted by

$$\sigma''_e(D) = \{\lambda \in \mathbb{K}, \eta(\lambda I - D) \text{ is infinite}\}$$

where  $\eta(\lambda I - D) = \dim N(\lambda I - D)$ .

**Theorem 6.7.** Let  $D : E_\omega \rightarrow E_\omega$  be a bounded diagonal operator such that  $D(u) = \sum \lambda_i u_i e_i$  with  $u = \sum u_i e_i$ . Then  $\sigma_e(D) = \sigma'_e(D) \cup \sigma''_e(D)$ .

*Proof.* (1) By the definition of  $\sigma''_e(D)$ , we conclude that  $\sigma''_e(D) \subseteq \sigma_e(D)$ , since  $\sigma_e(D)$  contains all the  $\lambda$  with infinite multiplicity with respect to  $D$  and all the  $\lambda$  with multiplicity 0 that are not surjective with respect to  $D$ .

(2)  $\lambda \in \sigma'_e(D) \Rightarrow \lambda \in \sigma_e(D)$  is of multiplicity 0 with respect to  $D$ . This implies that  $\lambda \notin \sigma_p(D) \Rightarrow \lambda$  is not an eigenvalue. But we know that  $\sigma(D) = \sigma_p(D) \cup \sigma_e(D) \Rightarrow \lambda \in \sigma_e(D) \Rightarrow \sigma'_e(D) \subseteq \sigma_e(D)$ . Therefore  $\sigma'_e(D) \cup \sigma''_e(D) \subseteq \sigma_e(D)$ .

(3) Let  $\lambda \in \sigma_e(D)$ . There are two cases:

Case 1: Let  $\lambda \in \sigma_e(D)$ . Then  $\lambda I - D$  is not a Fredholm operator of index 0  $\Rightarrow \lambda$  is of infinite multiplicity  $\Leftrightarrow \lambda \in \sigma_e''(D)$ .

Case 2:  $\lambda \notin \sigma_p(D) = \Lambda \Rightarrow N(\lambda I - D) = \{0\}$  and  $\lambda I - D$  is not surjective. Then  $\lambda$  is an element of  $\sigma(D)$  with multiplicity 0. Then  $\lambda \in \sigma_e'(D)$ .

$$\begin{aligned} \therefore \sigma(D) &\subseteq \sigma_e'(D) \cup \sigma_e''(D) \\ \text{Hence } \sigma_e(D) &\subseteq \sigma_e'(D) \cup \sigma_e''(D) \end{aligned}$$

□

**Theorem 6.8.** Let  $D : E_\omega \rightarrow E_\omega$  be a bounded diagonal operator, i.e.  $D(u) = \sum \lambda_i u_i e_i$  with  $u = \sum u_i e_i$  for  $\lambda_i \in \mathbb{K} \forall i \in \mathbb{N}$ . Let  $\Lambda = \{\lambda_k\}_{k \in \mathbb{N}} - \lambda_k$  are the eigenvalues. Then  $\sigma_e'(D) = \partial\Lambda = \bar{\Lambda} - \Lambda$ .

*Proof.* (1) Let  $\lambda \in \sigma_e'(D)$ . Then  $\lambda \in \sigma(D)$  and  $\lambda$  is of multiplicity 0 with respect to  $D$ . This implies that  $\lambda \notin \sigma_p(D) = \Lambda$  or  $\sigma(D) = \bar{\Lambda}$ . Thus  $\lambda \in \bar{\Lambda} - \Lambda = \partial\Lambda$ . Therefore  $\sigma_e'(D) = \partial\Lambda$ .

(2) Now let  $\lambda \in \partial\Lambda \Rightarrow \lambda \notin \Lambda = \sigma_p(D)$ . This implies  $\lambda$  is of multiplicity 0 with respect to  $D$  and  $\lambda \in \sigma(D) = \bar{\Lambda} \supseteq \partial\Lambda \Rightarrow \lambda \in \sigma_e'$ . Therefore  $\partial\Lambda \subseteq \sigma_e'(D)$ .

Based on 1 and 2, we conclude that

$$\sigma_e'(D) = \partial\Lambda = \bar{\Lambda} - \Lambda.$$

□

**Theorem 6.9.** Let  $D : E_\omega \rightarrow E_\omega$  be a bounded diagonal operator. Then  $\sigma_e''(D) \subseteq \sigma_p(D)$ .

*Proof.* Every element of  $\sigma_e''(D)$  is of infinite multiplicity with respect to  $D \Leftrightarrow N(\lambda I - D) \neq 0 \Rightarrow \lambda \in \sigma_p(D)$ . Therefore  $\sigma_e''(D) \subseteq \sigma_p(D)$ . □

**Corollary 6.10.** If every eigenvalue of  $D$  is of finite multiplicity, then  $\sigma_e(D) = \partial\Lambda$ .

*Proof.* Since  $\sigma_e''(D) = \phi$  and  $\sigma_e(D) = \sigma_e'(D) \cup \sigma_e''(D) \Rightarrow \sigma_e'(D) = \sigma_e(D)$  □

Note that  $\sigma_e''(D) = \phi$  since every eigenvalue of  $D$  is of finite multiplicity and  $\sigma_e''(D)$  is the set of all  $\lambda$  with infinite multiplicity.

**Theorem 6.11.** Let  $D : E_\omega \rightarrow E_\omega$  be a linear and diagonal operator of finite rank. Then

- (1) The spectrum of  $D$  is finite, and the spectrum of  $D$  equals  $\sigma_p(D)$ .
- (2) 0 is an element of the essential spectrum of  $D$ .

*Proof.* (1)  $D$  is of finite rank, which implies that  $\sigma_p(D)$  is finite. Hence  $\sigma_e(D) = \sigma_p(D)$ , based on the example (3.2) of DMR

- (2) Since  $E_\omega$  is infinite dimensional, it implies that  $\ker(D)$  is infinite dimensional since  $E_\omega$  is supposed to be of finite rank. This implies that  $D$  is not a Fredholm operator (FO). Therefore  $D$  is not a FO of index 0. Thus  $0I - D = -D$  is not a Fredholm operator of index 0. Therefore 0 belongs to the essential spectrum of  $D$ . □

**Corollary 6.12.** *Let  $D$  be a linear diagonal operator of finite rank and let  $\lambda \in K$  be a nonzero element belonging to  $\sigma_p(D)$ . Then  $\lambda I - D$  is a Fredholm operator of index 0. Therefore  $\lambda$  does not belong to the essential spectrum of  $D$ .*

*Proof.* The multiplicity of lambda is finite since the spectrum o. Therefore  $\lambda I - D$  is a Fredholm operator of index 0, by Theorem 1. We conclude that  $\lambda$  does not belong to the essential spectrum of  $D$ . □

**Corollary 6.13.** *If  $D$  is a linear diagonal operator of finite rank then the essential spectrum of  $D$  is equal to 0.*

*Proof.* Since the rank of  $D$  is finite,  $\ker(D)$  is infinite dimensional. This implies that  $D$  is not a Fredholm operator, hence  $0I - D = -D$  is not a Fredholm operator of index 0. Therefore 0 belongs to the essential spectra of  $D$ . By Theorem 1 and Corollary 1, we conclude that the essential spectrum of  $D$  is equal to 0. □

**Example 6.14.** Let  $\alpha_n, n \in \mathbb{N}^*$  be distinct non zero elements of  $\mathbb{K}$  with  $||\alpha_i|| < 1$ . For each  $n \in \mathbb{N}$  let  $\beta(n, k) = \alpha_n$  with  $k \in \mathbb{N}$ . Now set  $\lambda_i = \beta(\varphi_2(i))$  where  $\varphi_2 : \mathbb{N} \rightarrow \mathbb{N} \times \mathbb{N}$  is a bijection. Let  $D : E_\omega \rightarrow E_\omega$  be defined by  $D(u) = \sum_{i \in \mathbb{N}} \lambda_i u_i e_i$  for  $u = \sum_{i \in \mathbb{N}} u_i e_i \in E_\omega$ .

For  $i_0 \in \mathbb{N}$ , if  $\varphi_2(i_0) = (n_0, k_0) \in \mathbb{N} \times \mathbb{N}$ , we have  $\lambda_{i_0} = \beta(\varphi_2(i_0)) = \beta(n_0, k_0) = \alpha_{n_0}$ . We also have that  $\forall k \in \mathbb{N}, \beta(n_0, k) = \alpha_{n_0}$ . For each  $k \in \mathbb{N}, \{\varphi_2^{-1}(n_0, k)\} = B$  is a subset of  $\mathbb{N}$  but  $\lambda_{j_k} = \lambda_{i_0} = \alpha_{n_0}$  where  $j_k = \varphi_2^{-1}(n_0, k)$ . This implies that  $N(\lambda_{i_0} I - D)$  for all  $k \in \mathbb{N}$ . This implies  $D(e_{j_k}) = \lambda_{j_k} e_k \Rightarrow D(e_{j_k}) = \lambda_{i_0} e_k$ . Then  $(\lambda_{i_0} I - D)(e_{j_k}) = 0$ .

Since the kernel of a Fredholm operator is finite dimensional,  $\lambda_{i_0} I - D$  is not a Fredholm operator of index 0. This means that  $\lambda_{i_0} \in \sigma_e(D) \forall i_0 \in \mathbb{N}$ .  $\therefore \Lambda \subseteq \sigma_e(D)$  since  $\partial\Lambda \subseteq \sigma_e(D)$  by Theorem 2, which implies that  $\sigma_e(D) \supseteq \Lambda \cup \partial\Lambda = \overline{\Lambda} = \sigma(D)$ . But  $\sigma(D) = \sigma_p(D) \cup \sigma_e(D) \Rightarrow \sigma_e(D) \subseteq \sigma(D)$ .  $\therefore \sigma_e(D) = \sigma(D)$ .

## REFERENCES

- [1] F. Gouvêa, *p-adic numbers: An introduction*. Springer-Verlag, Berlin, 1993.
- [2] J. Araujo, C. Perez-Garcia, and S. Vega, Preservation of the index of p-adic linear operators under compact perturbations. *Compositio Mathematica* **118** (1999), pp. 291–303.
- [3] S. Basu, T. Diagana, and F. Ramarosan, A p-adic version of Hilbert-Schmidt operators and applications. *J. Anal. Appl.* **2** (2004), no. 3, pp. 173–188.
- [4] T. Diagana, Towards a theory of some unbounded linear operators on p-adic Hilbert spaces and applications, *Ann. Math. Blaise Pascal* **12** (2005), no. 1, pp. 205–222.
- [5] T. Diagana, *Non-archimedean linear operators and applications*. Nova Science Publishers, Inc., Huntington, NY, 2007.
- [6] T. Diagana and G. D. McNeal, Spectral analysis for rank one perturbations of diagonal operators in non-archimedean Hilbert space. *Comment. Math. Univ. Carolin.* **50** (2009), No. 3, pp. 385–400.
- [7] T. Diagana and G. D. McNeal, Corrigendum to Spectral analysis for rank one perturbations of diagonal operators in non-archimedean Hilbert space. *Comment. Math. Univ. Carolin.* **50** (2009), No. 4, pp. 637–638.

- [8] T. Diagana, *An Introduction to classical and  $p$ -adic theory of linear operators and applications*, Nova Science Publishers, New York, 2006.
- [9] B. Diarra, An operator on some ultrametric Hilbert Spaces, *J. Analysis* **6** (1998), pp. 55–74.
- [10] B. Diarra, *Geometry of the  $p$ -adic Hilbert spaces*, Preprint, 1999.
- [11] B. Diarra, Bounded linear operators on ultrametric Hilbert spaces. *Afr Diaspora J. Math.* **8** (2009), no. 2, pp. 173–181.
- [12] Q. Fang and J. Xia, Invariant subspaces for certain finite-rank perturbations of diagonal operators. *J. Func. Anal.* **263** (2012), Issue 5, pp. 1356–1377.
- [13] C. Foias, I. B. Jung, E. Ko, C. Pearcy, On rank one perturbations of normal operators. *J. Func. Anal.* **253** (2007), Issue 2, pp. 628–646.
- [14] I. Gohberg, S. Goldberg, and M. A. Kaashoek, *Basic classes of linear operators*. Birkhäuser, 2003.
- [15] E. Ionascu, Rank-one perturbations of diagonal operators. *Integral Equations Operator Theory* **39** (2001), pp. 421–440.
- [16] H. A. Keller and H. Ochsenius, Bounded operators on non-archimedean orthomodular spaces,. *Math. Slovaca* **45** (1995), no. 4, pp. 413–434.
- [17] J. Martínez-Maurica, T. Pellon, C. Perez-Garcia, Some characterizations of  $p$ -adic semi-Fredholm operators *Ann. Mat. Pura Appl.* **156** (1990), pp. 243–251.
- [18] G. D. McNeal, Spectral analysis for rank one perturbations of diagonal operators in non-archimedean Hilbert space. PhD Thesis, Howard University, 2009.
- [19] H. Ochsenius, W. H. Schikhof, *Banach spaces over fields with an infinite rank valuation,  $p$ -adic Functional Analysis* (Poznan,1998), Dekker, New York, 1999, pp. 233–293.
- [20] C. Perez-Garcia and S. Vega, Perturbation theory of  $p$ -adic Fredholm and semi-Fredholm operators. *Indag. Math. (N.S.)* **15** (2004), Issue 1, pp. 115–127.
- [21] C. Perez-Garcia, Semi-Fredholm operators and the Calkin algebra in  $p$ -adic analysis. I, II. *Bull. Soc. Math. Belg. Sér. B* **42** (1990), no. 1, pp. 69–101.
- [22] J. P. Serre, Completely continuous endomorphisms of  $p$ -adic Banach spaces. *Publ. Math. I.H.E.S.* No. **12** (1962), pp. 69–85.
- [23] S. Śliwa, On Fredholm operators between non-archimedean Fréchet spaces. *Compositio Mathematica* **139** (2003), pp. 113–118.
- [24] A.C.M. van Rooij, *Non-archimedean functional analysis*, Marcel Dekker Inc, New York (1978).
- [25] H. Stichtenoth, *Algebraic function fields and codes*, Springer-Verlag, New York (1993).