

# Simplified Veltman frames for sublogics of **IL**

Yuya Okawa

## Abstract

We study finite model property for several sublogics of **IL** with respect to simplified Veltman frames. Among other things, we prove that the logic **CL** of conservativity has finite model property with respect to that frames. This is an affirmative solution to Ignatiev’s problem.

## 1 Introduction

Throughout the introduction,  $T$  and  $U$  stand for consistent recursively enumerable theories extending Peano Arithmetic **PA**. The notion of interpretability has been investigated in modal logic. The language of interpretability logics is obtained by adding the binary modal operator  $\triangleright$  to the usual language of propositional modal logic. The modal formula  $A \triangleright B$  is intended to mean “ $T + B$  is interpretable in  $T + A$ ”. The logic **IL** of interpretability is a base logic to investigate several interpretability logics. Berarducci [1] and Shavrukov [8] independently proved that the extension **ILM** of **IL** is arithmetically complete with respect to arithmetical interpretation for any  $\Sigma_1$ -sound theory  $T$ . It is known that the logic **ILM** is also the logic of  $\Pi_1$ -conservativity (cf. Hájek and Montagna [3]). Ignatiev [4] further investigated the logic of  $\Gamma$ -conservativity where  $\Gamma \in \{\Sigma_n, \Pi_n : n \geq 1\}$ . He introduced the logic **CL** of conservativity, that is a sublogic of **IL**, as a base logic to investigate several conservativity logics.

The logics **IL** and **CL** are complete with respect to Veltman semantics. We say that a triple  $(W, R, \{S_x\}_{x \in W})$  is an **IL-frame** if  $W$  is a non-empty set,  $R$  is a transitive and conversely well-founded binary relation on  $W$ , and for each  $x \in W$ ,  $S_x$  is a transitive and reflexive binary relation on  $R[x] := \{y \in W : xRy\}$  satisfying the condition ( $\dagger$ ):  $(\forall x, y, z \in W)(xRy \ \& \ yRz \Rightarrow yS_xz)$ . Also, a **CL-frame** is obtained by removing the condition ( $\dagger$ ) from the definition of an **IL-frame**. Then, de Jongh and Veltman [2] proved that **IL** is sound and complete with respect to the class of all finite **IL-frames**. Moreover, Ignatiev [4] proved that **CL** is sound and complete with respect to the class of all finite **CL-frames**.

Visser [10] introduced and investigated the notion of simplified Veltman frames. A *simplified **IL-frame*** is a triple  $(W, R, S)$  where  $(W, R)$  satisfies the same condition as in the definition of Veltman frames and  $S$  is a transitive and reflexive binary relation on  $W$  satisfying the condition ( $\dagger'$ ):  $R \subseteq S$ . A *simplified **CL-frame*** is obtained by removing the condition ( $\dagger'$ ) from the definition of a simplified **IL-frame**. Then, Visser [10] proved that **IL** is sound and complete

with respect to the class of all simplified **IL**-frames. On the other hand, unlike the case of Veltman frames, Visser [11] proved that **IL** does not have finite model property with respect to simplified **IL**-frames. Ignatiev [4] proved that **CL** is sound and complete with respect to the class of all simplified **CL**-frames<sup>1</sup>. Then, he proposed the following problem.

**Problem 1.1** (Ignatiev [4]). *Does the logic **CL** have finite model property with respect to simplified **CL**-frames?*

By using the completeness for **CL** with respect to simplified Veltman frames, Ignatiev [4] also proved the arithmetical completeness for **CL**<sup>2</sup>. Namely, he proved that **CL** is exactly the intersection of all logics of  $\Gamma$ -conservativity for suitable sets  $\Gamma$ . However, the author at least think that there is one fault in his proof. His proof is proceeded by embedding a counter simplified **CL**-model  $(W, R, S, \Vdash)$  obtained by his completeness theorem into arithmetic as in the proof of Solovay [9], where  $W$  is an infinite set. Then, Ignatiev stated for each  $x \in W$ , finite set  $X \subset W$ , and modal formula  $C$ , the following relation is represented by a  $\Delta_0$  formula:

$$(\exists z \in W)(xRz \wedge (\exists y \in X)(ySz) \wedge z \Vdash C).$$

However, since  $W$  is an infinite set, the formula “ $\exists z \in W$ ” is not a  $\Delta_0$  formula in general. In addition, the author at least think that this formula is not represented by  $\Delta_0$  formula as long as we use his model. On the other hand, if **CL** has finite model property with respect to simplified Veltman frames, then the above formula is obviously represented by a  $\Delta_0$  formula because  $W$  is a finite set. Therefore, if Problem 1.1 is affirmatively solved, then Ignatiev’s proof is completely correct.

Veltman semantics was also introduced for sublogics of **IL** other than **CL**. Visser [10] introduced **IL**<sup>-</sup>-frames (or *Veltman prestructures*) which are general notions of **IL**-frames and **CL**-frames. Kurahashi and Okawa [7] introduced the sublogic **IL**<sup>-</sup> of **IL**, and proved that **IL**<sup>-</sup> is sound and complete with respect to the class of all finite **IL**<sup>-</sup>-frames. In [7], several sublogics of **IL** were introduced by adding several axioms to **IL**<sup>-</sup>, and their completeness with respect to **IL**<sup>-</sup>-frames was investigated.

In this context, we investigate finite model property with respect to simplified Veltman frames for the several sublogics of **IL**. Table 1 summarizes soundness, completeness, and finite model property for these sublogics with respect to simplified Veltman frames obtained in this paper. Among other things, we affirmatively solve Problem 1.1. To consider the completeness with respect to simplified Veltman frames for several sublogics of **IL**, we introduce several new frames. As a basis, we consider simplified Veltman frames  $(W, R, S)$  where  $(W, R)$  satisfies the same condition as in the definition of Veltman frames and  $S$

<sup>1</sup>Ignatiev actually proved a stronger result that **CL** is sound and complete with respect to the class of all simplified **CL**-frames where  $S$  is symmetric.

<sup>2</sup>In his proof of the arithmetical completeness for **CL**, it is sufficient to consider the completeness of simplified **CL**-frames.

Table 1: Soundness, completeness, and finite model property with respect to simplified Veltman frames for sublogics of  $\mathbf{IL}$

	Soundness and completeness	Finite model property
$\mathbf{IL}^-(\mathbf{J4}_+)$	✓ (Theorem 3.1)	✓ (Theorem 3.1)
$\mathbf{IL}^-(\mathbf{J1}, \mathbf{J4}_+)$	✓ (Theorem 3.1)	✓ (Theorem 3.1)
$\mathbf{IL}^-(\mathbf{J4}_+, \mathbf{J5})$	✓ (Theorem 3.1)	✓ (Theorem 3.1)
$\mathbf{IL}^-(\mathbf{J1}, \mathbf{J4}_+, \mathbf{J5})$	✓ (Theorem 3.1)	✓ (Theorem 3.1)
$\mathbf{IL}^-(\mathbf{J2}_+)$	✓ (Theorem 3.3)	✓ (Theorem 3.3)
$\mathbf{CL}$	✓ (Theorem 3.3)	✓ (Theorem 3.3)
$\mathbf{IL}^-(\mathbf{J2}_+, \mathbf{J5})$	✓ (Theorem 3.4)	× (Corollary 3.5)
$\mathbf{IL}$	✓ [10]	× [11]

is merely a binary relation on  $W$ . Thus, our basic simplified frames are general notions of simplified  $\mathbf{IL}$ -frames and  $\mathbf{CL}$ -frames. However, our basic simplified Veltman frames do not characterize the logic  $\mathbf{IL}^-$ . On the other hand, we show that the extension  $\mathbf{IL}^-(\mathbf{J4}_+)$  of  $\mathbf{IL}^-$  is sound and complete with respect to the class of all finite our basic simplified Veltman frames. Hence, in this paper, we call our basic simplified Veltman frames *simplified  $\mathbf{IL}^-(\mathbf{J4}_+)$ -frames*.

## 2 Preliminaries

In this section, we introduce the logic  $\mathbf{IL}$ , sublogics of it, and two relational semantics: Veltman frames and simplified Veltman frames. Then, we introduce and prove several basic properties for these logics and frames.

The language of interpretability logics consists of propositional variables  $p, q, r, \dots$ , the logical constants  $\top, \perp$ , the connectives  $\rightarrow, \vee, \wedge$ , and the modal operators  $\Box$  and  $\triangleright$ . The modal formulas are given by

$$A ::= \perp \mid \top \mid p \mid A \circ A \mid \Box A \mid A \triangleright A$$

where  $\circ \in \{\rightarrow, \vee, \wedge\}$ . We define  $\Diamond A := \neg \Box \neg A$ . The binary modal operator  $\triangleright$  binds stronger than  $\rightarrow$ , and weaker than  $\neg, \wedge, \vee, \Box$ , and  $\Diamond$ .

**Definition 2.1.** The logic  $\mathbf{IL}$  of interpretability has the following axioms and rules:

- L0:** Tautologies;
- L1:**  $\Box(A \rightarrow B) \rightarrow (\Box A \rightarrow \Box B)$ ;
- L2:**  $\Box(\Box A \rightarrow A) \rightarrow \Box A$ ;
- J1:**  $\Box(A \rightarrow B) \rightarrow A \triangleright B$ ;
- J2:**  $(A \triangleright B) \wedge (B \triangleright C) \rightarrow A \triangleright C$ ;

**J3:**  $(A \triangleright C) \wedge (B \triangleright C) \rightarrow (A \vee B) \triangleright C$ ;

**J4:**  $A \triangleright B \rightarrow (\diamond A \rightarrow \diamond B)$ ;

**J5:**  $\diamond A \triangleright A$ ;

(Modus Ponens)  $\frac{A \quad A \rightarrow B}{B}$ ;

(Necessitation)  $\frac{A}{\Box A}$ .

**Definition 2.2.** The logic **CL** of conservativity is obtained by removing **J5** from **IL**.

**Definition 2.3.** The logic **IL**<sup>-</sup> has the axioms **L0**, **L1**, **L2**, **J3**, and

**J6:**  $\Box \neg A \leftrightarrow A \triangleright \perp$ ,

and has the rules Modus Ponens, Necessitation,

**R1:**  $\frac{A \rightarrow B}{C \triangleright A \rightarrow C \triangleright B}$ , and **R2:**  $\frac{A \rightarrow B}{B \triangleright C \rightarrow A \triangleright C}$ .

The logic **IL**<sup>-</sup> is sound and complete with respect to the class of the following all (finite) **IL**<sup>-</sup>-frames which were originally introduced by Visser [10].

**Definition 2.4.** A triple  $\mathcal{F} = (W, R, \{S_x\}_{x \in W})$  is said to be an **IL**<sup>-</sup>-frame if the following conditions hold:

- $W$  is a non-empty set;
- $R$  is a transitive and conversely well-founded binary relation on  $W$ ;
- For each  $x \in W$ ,  $S_x \subseteq R[x] \times W$  where  $R[x] := \{y \in W : xRy\}$ .

We say that  $\mathcal{F}$  is finite if  $W$  is a finite set.

**Definition 2.5.** A quadruple  $(W, R, \{S_x\}_{x \in W}, \Vdash)$  is said to be an **IL**<sup>-</sup>-model if the following conditions hold.

- $(W, R, \{S_x\}_{x \in W})$  is an **IL**<sup>-</sup>-frame.
- $\Vdash$  is a usual forcing relation between  $W$  and the set of all modal formulas with the following clauses:

- $x \Vdash \Box A \iff (\forall y \in W)(xRy \Rightarrow y \Vdash A)$ ;
- $x \Vdash A \triangleright B \iff (\forall y \in W)(xRy \ \& \ y \Vdash A \Rightarrow \exists z(yS_x z \ \& \ z \Vdash B))$ .

Let  $A$  be any modal formula and let  $\mathcal{F} = (W, R, \{S_x\}_{x \in W})$  be any **IL**<sup>-</sup>-frame. We say that  $A$  is *valid* in  $\mathcal{F}$  ( $\mathcal{F} \models A$ ) if for any **IL**<sup>-</sup>-model  $(W, R, \{S_x\}_{x \in W}, \Vdash)$  and  $x \in W$ ,  $x \Vdash A$ .

Here, we introduce the following axioms  $\mathbf{J2}_+$  and  $\mathbf{J4}_+$  which were introduced by Kurahashi and Okawa [7] and Visser [10], respectively.

$$\mathbf{J2}_+: A \triangleright (B \vee C) \wedge B \triangleright C \rightarrow A \triangleright C.$$

$$\mathbf{J4}_+: \Box(A \rightarrow B) \rightarrow (C \triangleright A \rightarrow C \triangleright B).$$

Let  $L(\Sigma_1, \dots, \Sigma_n)$  denote the logic obtained by adding the axioms  $\Sigma_1, \dots, \Sigma_n$  to the logic  $L$ . Then, the following facts hold (for a proof, see [7]).

**Fact 2.6.**

1.  $\mathbf{IL}^-(\mathbf{J2}_+) \vdash \mathbf{J2}$ .
2.  $\mathbf{IL}^-(\mathbf{J1}, \mathbf{J2}) \vdash \mathbf{J2}_+$ .
3.  $\mathbf{IL}^-(\mathbf{J2}_+) \vdash \mathbf{J4}_+$ .
4.  $\mathbf{IL}^-(\mathbf{J4}_+) \vdash \mathbf{J4}$ .
5.  $\mathbf{IL}^-(\mathbf{J1}, \mathbf{J4}) \vdash \mathbf{J4}_+$ .
6. *The logics  $\mathbf{IL}$  and  $\mathbf{CL}$  are deductively equivalent to  $\mathbf{IL}^-(\mathbf{J1}, \mathbf{J2}, \mathbf{J5})$  and  $\mathbf{IL}^-(\mathbf{J1}, \mathbf{J2})$ , respectively*

**Fact 2.7.** *Let  $\mathcal{F} = (W, R, \{S_x\}_{x \in W})$  be any  $\mathbf{IL}^-$ -frame.*

1.  $\mathcal{F} \models \mathbf{J1} \iff (\forall x, y \in W)(xRy \Rightarrow yS_xy)$ .
2.  $\mathcal{F} \models \mathbf{J2}_+ \iff \mathcal{F} \models \mathbf{J4}_+ \ \& \ (\forall x, y, z, v \in W)(yS_xz \ \& \ zS_xv \Rightarrow yS_xv)$ .
3.  $\mathcal{F} \models \mathbf{J4}_+ \iff (\forall x, y, z \in W)(yS_xz \Rightarrow xRz)$ .
4.  $\mathcal{F} \models \mathbf{J5} \iff (\forall x, y, z \in W)(xRy \ \& \ yRz \Rightarrow yS_xz)$ .

**Fact 2.8.** *Let  $L$  be any logic appearing in Figure 1. Then the following are equivalent:*

1.  $L \vdash A$ .
2.  *$A$  is valid in any (finite)  $\mathbf{IL}^-$ -frame  $\mathcal{F}$  in which all axioms of  $L$  are valid.*

*Here, the logics in Figure 1 extend as it goes to the right along each of lines.*

**Remark 2.9.** From Fact 2.6, for any extension  $L$  of  $\mathbf{IL}^-(\mathbf{J1})$  closed under the rules of  $\mathbf{IL}^-$ , we always identify the axioms  $\mathbf{J2}_+$  and  $\mathbf{J4}_+$  with  $\mathbf{J2}$  and  $\mathbf{J4}$ , respectively. Also, we always identify  $\mathbf{IL}$  and  $\mathbf{CL}$  with  $\mathbf{IL}^-(\mathbf{J1}, \mathbf{J2}, \mathbf{J5})$  and  $\mathbf{IL}^-(\mathbf{J1}, \mathbf{J2})$ , respectively.

In the next section, we investigate the completeness for several sublogics of  $\mathbf{IL}$  in Figure 1 with respect to simplified Veltman frames which were originally introduced by Visser [10]. In this paper, we consider the following *simplified  $\mathbf{IL}^-(\mathbf{J4}_+)$ -frames* as the basis, which are general notion of Visser's simplified Veltman frames.




Figure 1: Sublogics of  $\mathbf{IL}$  complete with respect to a corresponding class of  $\mathbf{IL}^-$ -frames

**Definition 2.10.** A triple  $(W, R, S)$  is said to be a *simplified  $\mathbf{IL}^-(\mathbf{J4}_+)$ -frame* if the following conditions hold:

- $W$  is a non-empty set;
- $R$  is a transitive and conversely well-founded binary relation on  $W$ ;
- $S$  is a relation on  $W$ .

**Definition 2.11.** A quadruple  $(W, R, S, \Vdash)$  is said to be a *simplified  $\mathbf{IL}^-(\mathbf{J4}_+)$ -model* if the following conditions hold:

- $(W, R, S)$  is a simplified  $\mathbf{IL}^-(\mathbf{J4}_+)$ -frame;
- As in Definition 2.5, we similarly define a forcing relation  $\Vdash$  with the following clause:

$$x \Vdash A \triangleright B \iff (\forall y \in W)(xRy \ \& \ y \Vdash A \Rightarrow (\exists z \in W)(xRz \ \& \ ySz \ \& \ z \Vdash B)).$$

For any modal formula  $A$  and simplified  $\mathbf{IL}^-(\mathbf{J4}_+)$ -frame  $\mathcal{F}$ , we define  $\mathcal{F} \models A$  as in Definition 2.5.

**Remark 2.12.** For a simplified  $\mathbf{IL}^-(\mathbf{J4}_+)$ -model  $(W, R, S, \Vdash)$ , it may be natural to consider a forcing relation with the following clause:

$$x \Vdash A \triangleright B \iff (\forall y \in W)(xRy \ \& \ y \Vdash A \Rightarrow (\exists z \in W)(ySz \ \& \ z \Vdash B)).$$

However, if we adopt this definition, then the axiom

$$\mathbf{P}: A \triangleright B \rightarrow \Box(A \triangleright B)$$

is valid in all simplified Veltman frames (for details, see [6]). Therefore, in this paper, we adopt the definition in Definition 2.11 which was originally adopted in [10]. On the other hand, our definition causes that  $\mathbf{J4}_+$  is valid in all simplified  $\mathbf{IL}^-(\mathbf{J4}_+)$ -frames (see Proposition 2.13.2). Therefore, we focus on and investigate the sublogics of  $\mathbf{IL}$  proving  $\mathbf{J4}_+$  in the next section.

**Proposition 2.13.** *Let  $\mathcal{F} = (W, R, S)$  be any simplified  $\mathbf{IL}^-(\mathbf{J4}_+)$ -frame.*

1.  $\mathcal{F} \models \mathbf{J3}$ .
2.  $\mathcal{F} \models \mathbf{J4}_+$ .
3.  $\mathcal{F} \models \mathbf{J6}$ .
4.  $\mathcal{F} \models \mathbf{J1} \iff (\forall x, y \in W)(xRy \Rightarrow ySy)$ .
5.  $\mathcal{F} \models \mathbf{J2}_+ \iff (\forall x \in W)(\forall y, z, v \in R[x])(ySz \ \& \ zSv \Rightarrow ySv)$ .
6.  $\mathcal{F} \models \mathbf{J5} \iff (\forall x, y, z \in W)(xRy \ \& \ yRz \Rightarrow ySz)$ .

*Proof.* 1: Let  $x \in W$ . Suppose  $x \Vdash (A \triangleright C) \wedge (B \triangleright C)$ . We show  $x \Vdash (A \vee B) \triangleright C$ . Let  $y \in W$  with  $xRy$  and  $y \Vdash A \vee B$ . If  $y \Vdash A$ , then there exists  $z \in W$  such that  $xRz$ ,  $ySz$ , and  $z \Vdash C$  by  $x \Vdash A \triangleright C$ . If  $y \Vdash B$ , then there exists  $z \in W$  such that  $xRz$ ,  $ySz$ , and  $z \Vdash C$  by  $x \Vdash B \triangleright C$ . In any case, we obtain  $x \Vdash (A \vee B) \triangleright C$ .

2: Let  $x \in W$ . Suppose  $x \Vdash \Box(A \rightarrow B) \wedge C \triangleright A$ . We show  $x \Vdash C \triangleright B$ . Let  $y \in W$  with  $xRy$  and  $y \Vdash C$ . Since  $x \Vdash C \triangleright A$ , there exists  $z \in W$  such that  $xRz$ ,  $ySz$ , and  $z \Vdash A$ . Since  $x \Vdash \Box(A \rightarrow B)$ , we have  $z \Vdash B$ . Therefore,  $x \Vdash C \triangleright B$ .

3: Let  $x \in W$ . Suppose  $x \Vdash \Box \neg A$ . Then, there is no  $y \in W$  such that  $xRy$  and  $y \Vdash A$ . Thus,  $x \Vdash A \triangleright \perp$ . Conversely, if  $x \Vdash A \triangleright \perp$  and there exists  $y$  such that  $xRy$  and  $y \Vdash A$ , then there exists  $z \in W$  such that  $z \Vdash \perp$ . Thus,  $x \Vdash A \triangleright \perp$  implies  $x \Vdash \Box \neg A$ .

4: ( $\implies$ ): Let  $x, y \in W$  with  $xRy$ . Let  $(W, R, S, \Vdash)$  be a simplified  $\mathbf{IL}^-(\mathbf{J4}_+)$ -model such that  $a \Vdash p \iff a = y$  and  $a \Vdash q \iff a = y$ . Then, we have  $x \Vdash \Box(p \rightarrow q)$ . Thus,  $x \Vdash p \triangleright q$ . Since  $xRy$  and  $y \Vdash p$ , there exists  $y' \in W$  such that  $xRy'$ ,  $ySy'$ , and  $y' \Vdash q$ . Then,  $y' = y$ , and hence  $ySy$ .

4: ( $\impliedby$ ): Let  $x \in W$  with  $x \Vdash \Box(A \rightarrow B)$ . We show  $x \Vdash A \triangleright B$ . Let  $y \in W$  with  $xRy$  and  $y \Vdash A$ . Since  $xRy$ ,  $y \Vdash B$  and  $ySy$ , and hence  $x \Vdash A \triangleright B$ .

5: ( $\implies$ ): Let  $x \in W$  and  $y, z, v \in R[x]$  with  $ySz$  and  $zSv$ . Let  $(W, R, S, \Vdash)$  be a simplified  $\mathbf{IL}^-(\mathbf{J4}_+)$ -model such that  $a \Vdash p \iff a = y$ ,  $a \Vdash q \iff a = z$ , and  $a \Vdash r \iff a = v$ . We show  $x \Vdash p \triangleright (q \vee r)$  and  $x \Vdash q \triangleright r$ . Let  $y' \in W$  with  $xRy'$  and  $y' \Vdash p$ . Then,  $y' = y$ . Since  $xRz$ ,  $ySz$ , and  $z \Vdash q$ , we have  $x \Vdash p \triangleright (q \vee r)$ . Let  $z' \in W$  with  $xRz'$  and  $z' \Vdash q$ . Then,  $z' = z$ . Since  $xRv$ ,  $zSv$ , and  $v \Vdash r$ , we have  $x \Vdash q \triangleright r$ . Therefore,  $x \Vdash p \triangleright r$ . Since  $xRy$  and  $y \Vdash p$ , there exists  $v' \in W$  such that  $xRv'$ ,  $ySv'$ , and  $v' \Vdash r$ . Thus,  $v' = v$ , and hence  $ySv$ .

5: ( $\impliedby$ ): Let  $x \in W$  with  $x \Vdash A \triangleright (B \vee C)$  and  $x \Vdash B \triangleright C$ . We show  $x \Vdash A \triangleright C$ . Let  $y \in W$  with  $xRy$  and  $y \Vdash A$ . Since  $x \Vdash A \triangleright (B \vee C)$ , there exists  $z \in W$  such that  $xRz$ ,  $ySz$ , and  $z \Vdash B \vee C$ . If  $z \Vdash C$ , then we are done.

Assume  $z \Vdash B$ . Since  $x \Vdash B \triangleright C$ , there exists  $v \in W$  such that  $xRv$ ,  $zSv$ , and  $v \Vdash C$ . Since  $y, z, v \in R[x]$ ,  $ySz$ , and  $zSv$ , we obtain  $ySv$ . Therefore, we have  $x \Vdash A \triangleright C$ .

6: ( $\implies$ ): Let  $x, y, z \in W$  with  $xRy$  and  $yRz$ . Let  $(W, R, S, \Vdash)$  be a simplified  $\mathbf{IL}^-(\mathbf{J4}_+)$ -model such that  $a \Vdash p \iff a = z$ . Then,  $y \Vdash \Diamond p$ . Since  $x \Vdash \Diamond p \triangleright p$  and  $xRy$ , there exists  $z' \in W$  such that  $xRz'$ ,  $ySz'$ , and  $z' \Vdash p$ . Then,  $z' = z$ , and hence  $ySz$ .

6: ( $\impliedby$ ): Let  $x \in W$ . We show  $x \Vdash \Diamond A \triangleright A$ . Let  $y \in W$  with  $xRy$  and  $y \Vdash \Diamond A$ . Then, there exists  $z \in W$  such that  $yRz$  and  $z \Vdash A$ . Since  $xRy$  and  $yRz$ , we have  $xRz$  and  $ySz$ . Therefore,  $x \Vdash \Diamond A \triangleright A$ .  $\square$

**Definition 2.14.** Let  $L$  be any logic in Figure 1 between  $\mathbf{IL}^-(\mathbf{J4}_+)$  and  $\mathbf{IL}$ , and let  $\mathcal{F} = (W, R, S)$  be any simplified  $\mathbf{IL}^-(\mathbf{J4}_+)$ -frame. We say that  $\mathcal{F}$  is a *simplified  $L$ -frame* if the following conditions hold:

- If  $\mathbf{J1}$  is an axiom of  $L$ , then  $S$  is reflexive;
- If  $\mathbf{J2}_+$  is an axiom of  $L$ , then  $S$  is transitive;
- If  $\mathbf{J5}$  is an axiom of  $L$ , then  $R \subseteq S$ .

For the logics  $\mathbf{IL}$  and  $\mathbf{CL}$ , Visser [10] and Ignatiev [4] proved their soundness and completeness with respect to simplified Veltman frames, respectively. Moreover, Visser [11] proved that  $\mathbf{IL}$  does not have finite model property with respect to simplified Veltman frames.

**Fact 2.15.** *Let  $A$  be any modal formula.*

1. (Visser [10])  $\mathbf{IL} \vdash A \iff A$  is valid in all simplified  $\mathbf{IL}$ -frames.
2. (Visser [11])  $\mathbf{IL}$  does not have finite model property with respect to simplified  $\mathbf{IL}$ -frames.
3. (Ignatiev [4])  $\mathbf{CL} \vdash A \iff A$  is valid in all simplified  $\mathbf{CL}$ -frames.

Our main purpose of this paper is to prove the finite model property for  $\mathbf{CL}$  with respect to simplified Veltman frames (Theorem 3.3).

### 3 Completeness with respect to Simplified $\mathbf{IL}^-(\mathbf{J4}_+)$ -frames

In this section, we study whether each sublogic of  $\mathbf{IL}$  has the finite model property with respect to simplified Veltman frames or not. Among other things, we prove that the logic  $\mathbf{CL}$  has finite model property. As we stated in Remark 2.12, we investigate the logics in Figure 1 between  $\mathbf{IL}^-(\mathbf{J4}_+)$  and  $\mathbf{IL}$ .

Firstly, we investigate the logics  $\mathbf{IL}^-(\mathbf{J4}_+)$ ,  $\mathbf{IL}^-(\mathbf{J1}, \mathbf{J4}_+)$ ,  $\mathbf{IL}^-(\mathbf{J4}_+, \mathbf{J5})$ , and  $\mathbf{IL}^-(\mathbf{J1}, \mathbf{J4}_+, \mathbf{J5})$ .



**Theorem 3.1.** *Let  $L$  be either  $\mathbf{IL}^-(\mathbf{J4}_+)$ ,  $\mathbf{IL}^-(\mathbf{J1}, \mathbf{J4}_+)$ ,  $\mathbf{IL}^-(\mathbf{J4}_+, \mathbf{J5})$ , or  $\mathbf{IL}^-(\mathbf{J1}, \mathbf{J4}_+, \mathbf{J5})$ . For any modal formula  $A$ , the following are equivalent.*

1.  $L \vdash A$ .
2.  $A$  is valid in all (finite) simplified  $L$ -frames.

*Proof.* 1  $\implies$  2: Let  $\mathcal{F}$  be any simplified  $L$ -frame. By the definition of  $\mathcal{F}$  and Proposition 2.13, all axioms of  $L$  are valid in  $\mathcal{F}$ . Also, the validity of formulas in  $\mathcal{F}$  is preserved by every rule of  $\mathbf{IL}^-$ .

2  $\implies$  1: Suppose  $L \not\vdash A$ . Then, there exist an  $\mathbf{IL}^-$ -model  $(W, R, \{S_x\}_{x \in W}, \Vdash)$  and  $w \in W$  such that  $w \not\Vdash A$  and all axioms of  $L$  are valid in  $(W, R, \{S_x\}_{x \in W})$  by Fact 2.8. We define the following set  $W'$  of sequences of elements of  $W$ .

$$W' := \{\langle x_1, \dots, x_n \rangle : n \geq 1, \text{ for any } i \leq n \text{ and } j < n, x_i \in W \text{ and } x_j R x_{j+1}\}.$$

Since  $R$  is transitive and irreflexive, the length of each sequence of  $W'$  is less than or equal to the number of elements of  $W$ . Therefore,  $W'$  is a finite set.

Let  $\epsilon$  denote the empty sequence, and let  $\mathbf{x}, \mathbf{y}, \mathbf{z}, \dots$  denote elements of  $W' \cup \{\epsilon\}$ . Then, we define several notations. In the following, suppose  $\mathbf{x} = \langle x_1, \dots, x_n \rangle$  and  $\mathbf{y} = \langle y_1, \dots, y_m \rangle$  where we regard  $\mathbf{x} = \epsilon$  if  $n = 0$ .

- $\mathbf{x} - 1 := \begin{cases} \langle x_1, \dots, x_{n-1} \rangle & (n \geq 1), \\ \epsilon & (n = 0). \end{cases}$
- If  $n \geq 1$ , then  $\mathbf{x}^e := x_n$ .
- For each  $x \in W$ ,  $\mathbf{x} * \langle x \rangle := \langle x_1, \dots, x_n, x \rangle$ .
- $\mathbf{x} \subset \mathbf{y} : \iff \mathbf{x}$  is a proper initial segment of  $\mathbf{y}$ . That is,  $n < m$  and for any  $i$  with  $1 \leq i \leq n$ ,  $x_i = y_i$ .
- $\mathbf{x} \subseteq \mathbf{y} : \iff \mathbf{x} \subset \mathbf{y}$  or  $\mathbf{x} = \mathbf{y}$ .
- $\mathbf{y} \cap \mathbf{z}$  is the maximum element of  $\{\mathbf{x} \in W' \cup \{\epsilon\} : \mathbf{x} \subseteq \mathbf{y} \ \& \ \mathbf{x} \subseteq \mathbf{z}\}$  with respect to  $\subseteq$ .
- $\text{Setm}(\mathbf{y}, \mathbf{x}) := \{y_i : 1 \leq i \leq m\} \setminus \{x_i : 1 \leq i \leq n\}$ .

**Remark 3.2.** Note that if  $\mathbf{x} \neq \epsilon$  and  $\mathbf{x} \subset \mathbf{y}$ , then  $\mathbf{x}^e \in \text{Setm}(\mathbf{y} - 1, \mathbf{x} - 1)$  because  $\mathbf{x} \subseteq (\mathbf{y} - 1)$  and  $R$  is transitive and irreflexive.

Then, we define the binary relations  $R'$  and  $S'$  on  $W'$  and the forcing relation  $\Vdash'$  as follows:

- $\mathbf{x} R' \mathbf{y} : \iff \mathbf{x} \subset \mathbf{y}$ ;
- $\mathbf{y} S' \mathbf{z} : \iff$  For  $\mathbf{x} := (\mathbf{y} - 1) \cap (\mathbf{z} - 1)$ , either  $(\mathbf{x} = \epsilon \text{ and } \mathbf{y} \subseteq \mathbf{z})$  or the following conditions hold:
  - $\mathbf{x} \neq \epsilon$ ;

– there exists  $v \in \text{Setm}(\mathbf{y} - 1, \mathbf{x} - 1)$  such that  $\mathbf{y}^e S_v \mathbf{z}^e$ .

- $\mathbf{x} \Vdash' p \iff \mathbf{x}^e \Vdash p$ .

Since  $R'$  is obviously transitive and irreflexive,  $\mathcal{F} = (W', R', S')$  is a finite simplified  $\mathbf{IL}^-(\mathbf{J4}_+)$ -frame. We show that  $\mathcal{F}$  is a finite simplified  $L$ -frame. We distinguish the following three cases.

Case 1:  $L = \mathbf{IL}^-(\mathbf{J1}, \mathbf{J4}_+)$ . We show that  $S'$  is reflexive. Let  $\mathbf{y} \in W'$  and let  $\mathbf{x} = (\mathbf{y} - 1) \cap (\mathbf{y} - 1)$ , that is,  $\mathbf{x} = (\mathbf{y} - 1)$ . If  $\mathbf{x} = \epsilon$ , we obtain  $\mathbf{y} S' \mathbf{y}$  by the definition of  $S'$ . Assume  $\mathbf{x} \neq \epsilon$ . Then, we have  $\mathbf{x}^e \in \text{Setm}(\mathbf{y} - 1, \mathbf{x} - 1)$ . Since  $\mathbf{x}^e R \mathbf{y}^e$ , we have  $\mathbf{y}^e S_{\mathbf{x}^e} \mathbf{y}^e$  by Fact 2.7.1. Therefore, we obtain  $\mathbf{y} S' \mathbf{y}$ .

Case 2:  $L = \mathbf{IL}^-(\mathbf{J4}_+, \mathbf{J5})$ . We show that  $R' \subseteq S'$ . Let  $\mathbf{y}, \mathbf{z} \in W'$  with  $\mathbf{y} R' \mathbf{z}$ . Let  $\mathbf{x} = (\mathbf{y} - 1) \cap (\mathbf{z} - 1)$ . Since  $(\mathbf{y} - 1) \subseteq (\mathbf{z} - 1)$ , we have  $\mathbf{x} = (\mathbf{y} - 1)$ . If  $\mathbf{x} = \epsilon$ , we obtain  $\mathbf{y} S' \mathbf{z}$  by the definition of  $S'$ . Assume  $\mathbf{x} \neq \epsilon$ . Then,  $\mathbf{x}^e \in \text{Setm}(\mathbf{y} - 1, \mathbf{x} - 1)$ . Since  $\mathbf{x} \subseteq \mathbf{y} \subseteq \mathbf{z}$ , we have  $\mathbf{x}^e R \mathbf{y}^e$  and  $\mathbf{y}^e R \mathbf{z}^e$ . By Fact 2.7.4,  $\mathbf{y}^e S_{\mathbf{x}^e} \mathbf{z}^e$ . Therefore, we obtain  $\mathbf{y} S' \mathbf{z}$ .

Case 3: Otherwise. This is proved in a similar way as above.

Next, we prove the following claim.

**Claim 1.** For any modal formula  $A'$  and  $\mathbf{x} \in W'$ ,

$$\mathbf{x} \Vdash' A' \iff \mathbf{x}^e \Vdash A'.$$

*Proof.* We show the claim by induction on the construction of  $A'$ . We only prove the case  $A' \equiv B \triangleright C$ .

$\implies$ : Assume  $\mathbf{x} \Vdash' B \triangleright C$ . We show  $\mathbf{x}^e \Vdash B \triangleright C$ . Let  $y \in W$  with  $\mathbf{x}^e R y$  and  $y \Vdash B$ . Let  $\mathbf{y} := \mathbf{x} * \langle y \rangle$ . Then, obviously,  $\mathbf{y} \in W'$  and  $\mathbf{x} R' \mathbf{y}$ . By the induction hypothesis,  $\mathbf{y} \Vdash' B$ . Thus, by our assumption, there exists  $\mathbf{z} \in W'$  such that  $\mathbf{x} R' \mathbf{z}$ ,  $\mathbf{y} S' \mathbf{z}$ , and  $\mathbf{z} \Vdash' C$ . Then,  $(\mathbf{y} - 1) \cap (\mathbf{z} - 1) = \mathbf{x}$  because  $\mathbf{x} = (\mathbf{y} - 1)$  and  $\mathbf{x} \subseteq (\mathbf{z} - 1)$ . Since  $\text{Setm}(\mathbf{y} - 1, \mathbf{x} - 1) = \{\mathbf{x}^e\}$ , we obtain  $\mathbf{y}^e S_{\mathbf{x}^e} \mathbf{z}^e$  from  $\mathbf{y} S' \mathbf{z}$ . By the induction hypothesis, we have  $\mathbf{z}^e \Vdash C$ , and hence we obtain  $\mathbf{x}^e \Vdash B \triangleright C$ .

$\impliedby$ : Assume  $\mathbf{x}^e \Vdash B \triangleright C$ . We show  $\mathbf{x} \Vdash' B \triangleright C$ . Let  $\mathbf{y} \in W'$  with  $\mathbf{x} R' \mathbf{y}$  and  $\mathbf{y} \Vdash' B$ . Then, we have  $\mathbf{x}^e R \mathbf{y}^e$  and  $\mathbf{y}^e \Vdash B$  by the induction hypothesis. By our assumption, there exists  $z \in W$  such that  $\mathbf{y}^e S_{\mathbf{x}^e} z$  and  $z \Vdash C$ . Let  $\mathbf{z} := \mathbf{x} * \langle z \rangle$ . By Fact 2.7.3,  $\mathbf{x}^e R z$ , and hence we have  $\mathbf{z} \in W'$  and  $\mathbf{x} R' \mathbf{z}$ . Then,  $(\mathbf{y} - 1) \cap (\mathbf{z} - 1) = \mathbf{x}$  because  $\mathbf{x} = (\mathbf{z} - 1)$  and  $\mathbf{x} \subseteq (\mathbf{y} - 1)$ . Since  $\mathbf{x}^e \in \text{Setm}(\mathbf{y} - 1, \mathbf{x} - 1)$  and  $\mathbf{y}^e S_{\mathbf{x}^e} \mathbf{z}^e$ , we obtain  $\mathbf{y} S' \mathbf{z}$ . By the induction hypothesis,  $\mathbf{z} \Vdash' C$ , and hence  $\mathbf{x} \Vdash' B \triangleright C$ .  $\square$

From  $w \not\Vdash' A$ , we obtain  $\langle w \rangle \not\Vdash' A$  by Claim 1. Thus,  $A$  is not valid in  $\mathcal{F}$ .  $\square$

Secondly, we show that the logics  $\mathbf{IL}^-(\mathbf{J2}_+)$  and  $\mathbf{CL}$  have finite model property with respect to simplified Veltman frames. This is an affirmative solution of Problem 1.1.

**Theorem 3.3.** Let  $L$  be either  $\mathbf{IL}^-(\mathbf{J2}_+)$  or  $\mathbf{CL}$ . For any modal formula  $A$ , the following are equivalent.

1.  $L \vdash A$ .

2.  $A$  is valid in all (finite) simplified  $L$ -frames.

*Proof.* We only show the implication  $2 \implies 1$ . Suppose  $L \not\vdash A$ . By Fact 2.8, there exist a finite  $\mathbf{IL}^-$ -model  $(W, R, \{S_x\}_{x \in W}, \Vdash)$  and  $w \in W$  such that  $w \not\Vdash A$  and all axioms of  $L$  are valid in this frame. Then, we define  $W'$ ,  $R'$ , and  $\Vdash'$  as in the proof of Theorem 3.1. We define the binary relation  $S'$  on  $W'$  as follows:

- $\mathbf{y}S'\mathbf{z} :\iff$  For  $\mathbf{x} := (\mathbf{y} - 1) \cap (\mathbf{z} - 1)$ , either  $(\mathbf{x} = \epsilon$  and  $\mathbf{y} = \mathbf{z})$  or the following conditions hold:
  - $\mathbf{x} \neq \epsilon$ ;
  - $\text{Setm}(\mathbf{y} - 1, \mathbf{x} - 1) \supseteq \text{Setm}(\mathbf{z} - 1, \mathbf{x} - 1)$ ;
  - there exist  $l \geq 1, v_1, \dots, v_l \in \text{Setm}(\mathbf{y} - 1, \mathbf{x} - 1)$ , and  $a_1, \dots, a_{l+1} \in W$  such that  $a_1 = \mathbf{y}^e$ ,  $a_{l+1} = \mathbf{z}^e$ , and  $a_i S_{v_i} a_{i+1}$  for any  $i \leq l$ .

Then,  $\mathcal{F} = (W', R', S')$  is a finite simplified  $\mathbf{IL}^-$ -( $\mathbf{J4}_+$ )-frame. We show that  $L$  is a finite simplified  $L$ -frame. We only consider the case  $L = \mathbf{CL}$ .

- We show that  $S'$  is reflexive. Let  $\mathbf{y} \in W'$  and let  $\mathbf{x} = (\mathbf{y} - 1) \cap (\mathbf{y} - 1) = (\mathbf{y} - 1)$ . If  $\mathbf{x} = \epsilon$ , we obtain  $\mathbf{y}S'\mathbf{y}$  by the definition of  $S'$ . Assume  $\mathbf{x} \neq \epsilon$ . The condition  $\text{Setm}(\mathbf{y} - 1, \mathbf{x} - 1) \supseteq \text{Setm}(\mathbf{y} - 1, \mathbf{x} - 1)$  obviously holds. Since  $\mathbf{x}^e \in \text{Setm}(\mathbf{y} - 1, \mathbf{x} - 1)$  and  $\mathbf{y}^e S_{\mathbf{x}^e} \mathbf{y}^e$  by Fact 2.7.1, we obtain  $\mathbf{y}S'\mathbf{y}$ .
- We show that  $S'$  is transitive. Let  $\mathbf{y}, \mathbf{z}, \mathbf{v} \in W'$  with  $\mathbf{y}S'\mathbf{z}$  and  $\mathbf{z}S'\mathbf{v}$ , and let  $\mathbf{x}_1 = (\mathbf{y} - 1) \cap (\mathbf{z} - 1)$ ,  $\mathbf{x}_2 = (\mathbf{z} - 1) \cap (\mathbf{v} - 1)$ , and  $\mathbf{x} = (\mathbf{y} - 1) \cap (\mathbf{v} - 1)$ . If  $\mathbf{x}_1 = \epsilon$  or  $\mathbf{x}_2 = \epsilon$ , then  $\mathbf{y} = \mathbf{z}$  or  $\mathbf{z} = \mathbf{v}$  because  $\mathbf{y}S'\mathbf{z}$  and  $\mathbf{z}S'\mathbf{v}$ , and hence we obtain  $\mathbf{y}S'\mathbf{v}$ . Assume  $\mathbf{x}_1 \neq \epsilon$  and  $\mathbf{x}_2 \neq \epsilon$ . Since  $\mathbf{x}_1$  and  $\mathbf{x}_2$  are initial segments of  $(\mathbf{z} - 1)$ , there are three possibilities  $\mathbf{x}_1 \subset \mathbf{x}_2$ ,  $\mathbf{x}_2 \subset \mathbf{x}_1$ , and  $\mathbf{x}_1 = \mathbf{x}_2$ . Therefore,  $\mathbf{x}' := \mathbf{x}_1 \cap \mathbf{x}_2$  equals to  $\mathbf{x}_1$  or  $\mathbf{x}_2$ , and hence  $\mathbf{x}' \neq \epsilon$ . Since  $\mathbf{x}'$  is an initial segment of  $(\mathbf{y} - 1)$  and  $(\mathbf{v} - 1)$ , we have  $\mathbf{x}' \subseteq \mathbf{x}$  by the maximality of  $\mathbf{x}$ . Thus,  $\mathbf{x} \neq \epsilon$ . Then, we distinguish the following three cases.

Case 1:  $\mathbf{x}_1 \subset \mathbf{x}_2$ . Let

$$\mathbf{x}_1 = \langle x_1, \dots, x_m \rangle,$$

$$\mathbf{x}_2 = \langle x_1, \dots, x_m, x_{m+1}, \dots, x_k \rangle$$

where  $0 < m < k$ , and let

$$\mathbf{y} - 1 = \langle x_1, \dots, x_m, y_{m+1}, \dots, y_{n_1} \rangle,$$

$$\mathbf{z} - 1 = \langle x_1, \dots, x_m, x_{m+1}, \dots, x_k, z_{k+1}, \dots, z_{n_2} \rangle,$$

$$\mathbf{v} - 1 = \langle x_1, \dots, x_m, x_{m+1}, \dots, x_k, v_{k+1}, \dots, v_{n_3} \rangle$$

where  $n_1 \geq m$ ,  $n_2 \geq k$ , and  $n_3 \geq k$ . Even if there exists  $y_{m+1}$ , then  $y_{m+1} \neq x_{m+1}$  by the maximality of  $\mathbf{x}_1$ . Thus,  $\mathbf{x} = (\mathbf{y} - 1) \cap (\mathbf{v} - 1) = \mathbf{x}_1$ . From  $\mathbf{y}S'\mathbf{z}$  and  $\mathbf{z}S'\mathbf{v}$ , we have

$$\text{Setm}(\mathbf{y} - 1, \mathbf{x}_1 - 1) \supseteq \text{Setm}(\mathbf{z} - 1, \mathbf{x}_1 - 1),$$

$$\text{Setm}(\mathbf{z} - 1, \mathbf{x}_2 - 1) \supseteq \text{Setm}(\mathbf{v} - 1, \mathbf{x}_2 - 1).$$

That is,

$$\{x_m, y_{m+1}, \dots, y_{n_1}\} \supseteq \{x_m, x_{m+1}, \dots, x_k, z_{k+1}, \dots, z_{n_2}\},$$

$$\{x_k, z_{k+1}, \dots, z_{n_2}\} \supseteq \{x_k, v_{k+1}, \dots, v_{n_3}\}.$$

Since  $\text{Setm}(\mathbf{v} - 1, \mathbf{x}_1 - 1) = \{x_m, x_{m+1}, \dots, x_k, v_{k+1}, \dots, v_{n_3}\}$ , we obtain

$$\text{Setm}(\mathbf{y} - 1, \mathbf{x} - 1) \supseteq \text{Setm}(\mathbf{v} - 1, \mathbf{x} - 1).$$

On the other hand, from  $\mathbf{y}S'\mathbf{z}$  and  $\mathbf{z}S'\mathbf{v}$ ,

– there exist  $w_1, \dots, w_l \in \text{Setm}(\mathbf{y} - 1, \mathbf{x}_1 - 1)$  and  $a_1, \dots, a_{l+1} \in W$  such that  $a_1 = \mathbf{y}^e, a_{l+1} = \mathbf{z}^e$  and  $a_i S_{w_i} a_{i+1}$  for any  $i \leq l$ ,

– there exist  $w'_1, \dots, w'_{l'} \in \text{Setm}(\mathbf{z} - 1, \mathbf{x}_2 - 1)$  and  $b_1, \dots, b_{l'+1} \in W$  such that  $b_1 = \mathbf{z}^e, b_{l'+1} = \mathbf{v}^e$  and  $b_i S_{w'_i} b_{i+1}$  for any  $i \leq l'$ .

Since  $\text{Setm}(\mathbf{z} - 1, \mathbf{x}_2 - 1) \subseteq \text{Setm}(\mathbf{z} - 1, \mathbf{x}_1 - 1)$ , we have

$$\text{Setm}(\mathbf{z} - 1, \mathbf{x}_2 - 1) \subseteq \text{Setm}(\mathbf{y} - 1, \mathbf{x}_1 - 1).$$

Hence, there exist  $w''_1, \dots, w''_{l''} \in \text{Setm}(\mathbf{y} - 1, \mathbf{x} - 1)$  and  $c_1, \dots, c_{l''+1} \in W$  such that

$$c_1 = \mathbf{y}^e, c_{l''+1} = \mathbf{v}^e \text{ and } c_i S_{w''_i} c_{i+1} \text{ for any } i \leq l''.$$

Then, we conclude  $\mathbf{y}S'\mathbf{v}$ .

Case 2:  $\mathbf{x}_2 \subset \mathbf{x}_1$ . Let

$$\mathbf{x}_1 = \langle x_1, \dots, x_m, x_{m+1}, \dots, x_k \rangle,$$

$$\mathbf{x}_2 = \langle x_1, \dots, x_m \rangle$$

where  $0 < m < k$ , and let

$$\mathbf{y} - 1 = \langle x_1, \dots, x_m, x_{m+1}, \dots, x_k, y_{k+1}, \dots, y_{n_1} \rangle,$$

$$\mathbf{z} - 1 = \langle x_1, \dots, x_m, x_{m+1}, \dots, x_k, z_{k+1}, \dots, z_{n_2} \rangle,$$

$$\mathbf{v} - 1 = \langle x_1, \dots, x_m, v_{m+1}, \dots, v_{n_3} \rangle.$$

where  $n_1 \geq k$ ,  $n_2 \geq k$ , and  $n_3 \geq m$ . Then,  $\mathbf{x} = \mathbf{x}_2$ . From

$$\text{Setm}(\mathbf{y} - 1, \mathbf{x}_1 - 1) \supseteq \text{Setm}(\mathbf{z} - 1, \mathbf{x}_1 - 1),$$

$$\text{Setm}(\mathbf{z} - 1, \mathbf{x}_2 - 1) \supseteq \text{Setm}(\mathbf{v} - 1, \mathbf{x}_2 - 1),$$

we have

$$\{x_k, y_{k+1}, \dots, y_{n_1}\} \supseteq \{x_k, z_{k+1}, \dots, z_{n_2}\},$$

$$\{x_m, x_{m+1}, \dots, x_k, z_{k+1}, \dots, z_{n_2}\} \supseteq \{x_m, v_{m+1}, \dots, v_{n_3}\}.$$

Since  $\text{Setm}(\mathbf{y} - 1, \mathbf{x}_2 - 1) = \{x_m, x_{m+1}, \dots, x_k, y_{k+1}, \dots, y_{n_1}\}$ , we obtain

$$\text{Setm}(\mathbf{y} - 1, \mathbf{x} - 1) \supseteq \text{Setm}(\mathbf{v} - 1, \mathbf{x} - 1).$$

On the other hand, we have  $\text{Setm}(\mathbf{y} - 1, \mathbf{x}_1 - 1) \subseteq \text{Setm}(\mathbf{y} - 1, \mathbf{x} - 1)$  and  $\text{Setm}(\mathbf{z} - 1, \mathbf{x}_2 - 1) \subseteq \text{Setm}(\mathbf{y} - 1, \mathbf{x} - 1)$ . Therefore, from  $\mathbf{y}S'\mathbf{z}$  and  $\mathbf{z}S'\mathbf{v}$ , there exist  $w_1, \dots, w_l \in \text{Setm}(\mathbf{y} - 1, \mathbf{x} - 1)$  and  $a_1, \dots, a_{l+1} \in W$  such that

$$a_1 = \mathbf{y}^e, a_{l+1} = \mathbf{v}^e \text{ and } a_i S_{w_i} a_{i+1} \text{ for any } i \leq l.$$

Thus, we conclude  $\mathbf{y}S'\mathbf{v}$ .

Case 3:  $\mathbf{x}_1 = \mathbf{x}_2$ . Let

$$\mathbf{x}_1 = \mathbf{x}_2 = \langle x_1, \dots, x_m \rangle$$

where  $m > 0$ . We already showed that  $\mathbf{x}_1 \cap \mathbf{x}_2 \subseteq \mathbf{x}$ . Thus, let

$$\mathbf{x} = \langle x_1, \dots, x_k \rangle$$

where  $k \geq m$ , and let

$$\mathbf{y} - 1 = \langle x_1, \dots, x_m, x_{m+1}, \dots, x_k, y_{k+1}, \dots, y_{n_1} \rangle,$$

$$\mathbf{z} - 1 = \langle x_1, \dots, x_m, z_{m+1}, \dots, z_{n_2} \rangle,$$

$$\mathbf{v} - 1 = \langle x_1, \dots, x_m, x_{m+1}, \dots, x_k, v_{k+1}, \dots, v_{n_3} \rangle$$

where  $n_1 \geq k$ ,  $n_2 \geq m$ , and  $n_3 \geq k$ . We show  $k = m$ . Towards a contradiction, assume  $k > m$ . Then, there exists  $x_{m+1}$ . From

$$\text{Setm}(\mathbf{y} - 1, \mathbf{x}_1 - 1) \supseteq \text{Setm}(\mathbf{z} - 1, \mathbf{x}_1 - 1),$$

$$\text{Setm}(\mathbf{z} - 1, \mathbf{x}_2 - 1) \supseteq \text{Setm}(\mathbf{v} - 1, \mathbf{x}_2 - 1),$$

we have

$$\{x_m, x_{m+1}, \dots, x_k, y_{k+1}, \dots, y_{n_1}\} \supseteq \{x_m, z_{m+1}, \dots, z_{n_2}\},$$

$$\{x_m, z_{m+1}, \dots, z_{n_2}\} \supseteq \{x_m, x_{m+1}, \dots, x_k, v_{k+1}, \dots, v_{n_3}\}.$$

Then,

$$x_{m+1} \in \{x_m, z_{m+1}, \dots, z_{n_2}\}.$$

Since  $x_m R x_{m+1}$ , we have  $x_m \neq x_{m+1}$ . Thus,  $x_{m+1} \in \{z_{m+1}, \dots, z_{n_2}\}$ , and hence there exists  $z_{m+1}$ . Also, by the maximality of  $\mathbf{x}_2$ ,  $x_{m+1} \neq z_{m+1}$ . Therefore, we have  $z_{m+1} R x_{m+1}$ . On the other hand,

$$z_{m+1} \in \{x_m, x_{m+1}, \dots, x_k, y_{k+1}, \dots, y_{n_1}\}.$$

Since  $x_m R z_{m+1}$ , we have  $z_{m+1} \neq x_m$ . Also, since  $z_{m+1} \neq x_{m+1}$ , we obtain  $x_{m+1} R z_{m+1}$ . Then,  $x_{m+1} R x_{m+1}$ . This is a contradiction. Therefore, we have  $k = m$  and  $\mathbf{x}_1 = \mathbf{x}_2 = \mathbf{x}$ . Then, we obtain

$$\text{Setm}(\mathbf{y} - 1, \mathbf{x} - 1) \supseteq \text{Setm}(\mathbf{v} - 1, \mathbf{x} - 1).$$

Also, since  $\text{Setm}(\mathbf{z} - 1, \mathbf{x}_2 - 1) \subseteq \text{Setm}(\mathbf{y} - 1, \mathbf{x}_1 - 1)$ , there exist  $w_1, \dots, w_l \in \text{Setm}(\mathbf{y} - 1, \mathbf{x} - 1)$  and  $a_1, \dots, a_{l+1} \in W$  such that

$$a_1 = \mathbf{y}^e, a_{l+1} = \mathbf{v}^e \text{ and } a_i S_{w_i} a_{i+1} \text{ for any } i \leq l.$$

Therefore, we conclude  $\mathbf{y} S' \mathbf{v}$ .

Next, we show the following claim.

**Claim 2.** For any modal formula  $A'$  and  $\mathbf{x} \in W'$ ,

$$\mathbf{x} \Vdash' A' \iff \mathbf{x}^e \Vdash A'.$$

*Proof.* We show the claim by induction on the construction of  $A'$ . We only prove the case  $A' \equiv B \triangleright C$ .

( $\implies$ ): Assume  $\mathbf{x} \Vdash' B \triangleright C$ . We show  $\mathbf{x}^e \Vdash B \triangleright C$ . Let  $y \in W$  with  $\mathbf{x}^e R y$  and  $y \Vdash B$ . Then, let  $\mathbf{y} := \mathbf{x} * \langle y \rangle$ . Obviously,  $\mathbf{y} \in W'$  and  $\mathbf{x} R' \mathbf{y}$ . By the induction hypothesis,  $\mathbf{y} \Vdash' B$ . By our assumption, there exists  $\mathbf{z} \in W'$  such that  $\mathbf{x} R' \mathbf{z}$ ,  $\mathbf{y} S' \mathbf{z}$ , and  $\mathbf{z} \Vdash' C$ . Since  $(\mathbf{y} - 1) = \mathbf{x}$  and  $\mathbf{x} \subseteq (\mathbf{z} - 1)$ , we have  $(\mathbf{y} - 1) \cap (\mathbf{z} - 1) = \mathbf{x}$ . Also, since  $\text{Setm}(\mathbf{y} - 1, \mathbf{x} - 1) = \{\mathbf{x}^e\}$ , we obtain

$$(\exists a_1, \dots, a_{l+1} \in W)(a_1 = \mathbf{y}^e \ \& \ a_{l+1} = \mathbf{z}^e \ \& \ (\forall i \leq l)(a_i S_{\mathbf{x}^e} a_{i+1}))$$

from  $\mathbf{y} S' \mathbf{z}$ . By Proposition 2.7.2,  $S_{\mathbf{x}^e}$  is transitive, and hence  $\mathbf{y}^e S_{\mathbf{x}^e} \mathbf{z}^e$ . By the induction hypothesis,  $\mathbf{z}^e \Vdash C$ , and therefore  $\mathbf{x}^e \Vdash B \triangleright C$ .

( $\impliedby$ ): Assume  $\mathbf{x}^e \Vdash B \triangleright C$ . We show  $\mathbf{x} \Vdash' B \triangleright C$ . Let  $\mathbf{y} \in W'$  with  $\mathbf{x} R' \mathbf{y}$  and  $\mathbf{y} \Vdash' B$ . Then,  $\mathbf{x}^e R \mathbf{y}^e$  and  $\mathbf{y}^e \Vdash B$  by the induction hypothesis. By our assumption, there exists  $z \in W$  such that  $\mathbf{y}^e S_{\mathbf{x}^e} z$  and  $z \Vdash C$ . Let  $\mathbf{z} := \mathbf{x} * \langle z \rangle$ . By Propositions 2.7.2 and 3,  $\mathbf{x}^e R z$ . Thus,  $\mathbf{z} \in W'$  and  $\mathbf{x} R' \mathbf{z}$ . Since  $(\mathbf{z} - 1) = \mathbf{x}$  and  $\mathbf{x} \subseteq (\mathbf{y} - 1)$ , we have  $(\mathbf{y} - 1) \cap (\mathbf{z} - 1) = \mathbf{x}$ . Then,  $\{\mathbf{x}^e\} = \text{Setm}(\mathbf{z} - 1, \mathbf{x} - 1)$ , and hence  $\text{Setm}(\mathbf{y} - 1, \mathbf{x} - 1) \supseteq \text{Setm}(\mathbf{z} - 1, \mathbf{x} - 1)$ . Also, since  $\mathbf{x}^e \in \text{Setm}(\mathbf{y} - 1, \mathbf{x} - 1)$  and  $\mathbf{y}^e S_{\mathbf{x}^e} \mathbf{z}^e$ , we obtain  $\mathbf{y} S' \mathbf{z}$ . By the induction hypothesis,  $\mathbf{z} \Vdash' C$ , and therefore we conclude  $\mathbf{x} \Vdash' B \triangleright C$ .  $\square$

By Claim 2, we conclude  $\langle w \rangle \not\Vdash A$ .  $\square$

Thirdly, we investigate the logic  $\mathbf{IL}^-(\mathbf{J2}_+, \mathbf{J5})$ . The logic  $\mathbf{IL}^-(\mathbf{J2}_+, \mathbf{J5})$  is complete with respect to the class of all simplified  $\mathbf{IL}^-(\mathbf{J2}_+, \mathbf{J5})$ -frames, however, it does not have finite model property as  $\mathbf{IL}$ . Our proof of the completeness of  $\mathbf{IL}^-(\mathbf{J2}_+, \mathbf{J5})$  is based on the idea in Ignatiev's proofs of the completeness of several logics with respect to simplified Veltman frames [4].

**Theorem 3.4.** *For any modal formula  $A$ , the following are equivalent:*

1.  $\mathbf{IL}^-(\mathbf{J2}_+, \mathbf{J5}) \vdash A$ .
2.  $A$  is valid in all simplified  $\mathbf{IL}^-(\mathbf{J2}_+, \mathbf{J5})$ -frames.

*Proof.* We only show the implication  $2 \implies 1$ . Suppose  $\mathbf{IL}^-(\mathbf{J2}_+, \mathbf{J5}) \not\vdash A$ . By Fact 2.8, there exist a finite  $\mathbf{IL}^-$ -model  $(W, R, \{S_x\}_{x \in W}, \Vdash)$  and  $w \in W$  such that  $w \not\Vdash A$  and  $\mathbf{J2}_+$  and  $\mathbf{J5}$  are valid in this frame. Assume  $0 \notin W$ . For each sequence  $\Gamma$  of elements of  $W \cup \{0\}$ , let  $|\Gamma|$  denote the length of  $\Gamma$ . Let  $W'$  be the following set.

- $$W' := \{(\Gamma, \Delta) : \begin{array}{l} 1) \ \Gamma \text{ is a finite sequence of elements of } W \text{ and } \Gamma \neq \epsilon, \\ 2) \ \Delta \text{ is a finite sequence of elements of } W \cup \{0\} \text{ and} \\ \quad \Delta = \epsilon \text{ is allowed,} \\ 3) \ |\Delta| + 1 = |\Gamma|, \\ 4) \ \text{for } \Gamma = \langle x_1, \dots, x_n \rangle \text{ and } \Delta = \langle v_1, \dots, v_{n-1} \rangle, \\ \quad (\forall i < n)[(v_i = 0 \Rightarrow x_i R x_{i+1}) \wedge (v_i \neq 0 \Rightarrow x_i S_{v_i} x_{i+1})] \}. \end{array}$$

Let  $\mathbf{x}, \mathbf{y}, \mathbf{z}, \dots$  denote elements of  $W'$ . Then, we define several notations. Let  $\mathbf{x} = (\langle x_1, \dots, x_n \rangle, \langle v_1, \dots, v_{n-1} \rangle)$ ,  $\mathbf{y} = (\langle y_1, \dots, y_m \rangle, \langle w_1, \dots, w_{m-1} \rangle) \in W'$ .

- $\mathbf{x} \subset \mathbf{y} := \Leftrightarrow n < m$ ,  $(\forall i \leq n)(x_i = y_i)$ , and  $(\forall j < n)(v_j = w_j)$ .
- $R(\mathbf{x}) := (\langle x_1, \dots, x_k \rangle, \langle v_1, \dots, v_{k-1} \rangle)$  where  $k := \min\{j > 0 : \forall i(j \leq i < n \Rightarrow v_i \neq 0)\}$ .
- $|\mathbf{x}| := n$ .
- $\mathbf{x}^e := x_n$ .

Then, we define the binary relations  $R'$  and  $S'$  on  $W'$  and the forcing relation  $\Vdash'$  as follows:

- $\mathbf{x} R' \mathbf{y} := \Leftrightarrow \mathbf{x} \subset \mathbf{y}$  and for  $\mathbf{x} = (\langle x_1, \dots, x_n \rangle, \langle v_0, \dots, v_{n-1} \rangle)$  and  $\mathbf{y} = (\langle x_1, \dots, x_m \rangle, \langle v_0, \dots, v_{m-1} \rangle)$ ,
$$\forall i(n \leq i \leq m-1 \ \& \ v_i \neq 0 \implies \exists k(n \leq k < i \ \& \ v_i = x_k)). \quad (1)$$
- $\mathbf{y} S' \mathbf{z} := \Leftrightarrow R(\mathbf{y}) \subseteq R(\mathbf{z})$  and  $|\mathbf{y}| < |\mathbf{z}|$ .
- $\mathbf{x} \Vdash' p := \Leftrightarrow \mathbf{x}^e \Vdash p$ .

We show that  $\mathcal{F} := (W', R', S')$  is a simplified  $\mathbf{IL}^-(\mathbf{J2}_+, \mathbf{J5})$ -frame. Since  $S'$  is obviously transitive and satisfies  $R' \subseteq S'$ , it suffices to prove that  $R'$  is transitive and conversely well-founded.

Firstly, we show that  $R'$  is transitive. Let  $\mathbf{x}, \mathbf{y}, \mathbf{z} \in W'$  with  $\mathbf{x}R'\mathbf{y}$  and  $\mathbf{y}R'\mathbf{z}$  and let  $\mathbf{x} = (\langle x_0, \dots, x_n \rangle, \langle v_0, \dots, v_{n-1} \rangle)$ ,  $\mathbf{y} = (\langle x_0, \dots, x_m \rangle, \langle v_0, \dots, v_{m-1} \rangle)$ ,  $\mathbf{z} = (\langle x_0, \dots, x_l \rangle, \langle v_0, \dots, v_{l-1} \rangle)$  with  $n < m < l$ . Obviously,  $\mathbf{x} \subset \mathbf{z}$ . From  $\mathbf{x}R'\mathbf{y}$  and  $\mathbf{y}R'\mathbf{z}$ ,

$$\forall i(n \leq i \leq m-1 \ \& \ v_i \neq 0 \Rightarrow \exists k(n \leq k < i \ \& \ v_i = x_k)),$$

$$\forall i(m \leq i \leq l-1 \ \& \ v_i \neq 0 \Rightarrow \exists k(m \leq k < i \ \& \ v_i = x_k)).$$

Since  $n < m < l$ , we have

$$\forall i(n \leq i \leq l-1 \ \& \ v_i \neq 0 \Rightarrow \exists k(n \leq k < i \ \& \ v_i = x_k)).$$

Therefore,  $\mathbf{x}R'\mathbf{z}$ .

Next, we show that  $R'$  is conversely well-founded. We show the following claim.

**Claim 3.** *Let  $\mathbf{x}, \mathbf{y} \in W'$  with  $\mathbf{x}R'\mathbf{y}$ ,  $\mathbf{x} = (\langle x_1, \dots, x_n \rangle, \langle v_1, \dots, v_{n-1} \rangle)$ , and  $\mathbf{y} = (\langle x_1, \dots, x_m \rangle, \langle v_1, \dots, v_{m-1} \rangle)$ . Then, for any  $i$  with  $n+1 \leq i \leq m$ ,  $x_n R x_i$ .*

*Proof.* Assume for any  $j$  with  $n+1 \leq j < i$ ,  $x_n R x_j$ . Then, we show  $x_n R x_i$ . We distinguish the following two cases.

Case 1:  $v_{i-1} = 0$ . Then,  $x_{i-1} R x_i$ . If  $i-1 = n$ , then we are done. If  $i-1 \geq n+1$ , then we have  $x_n R x_i$  by our assumption and the transitivity of  $R$ .

Case 2:  $v_{i-1} \neq 0$ . Then,  $x_{i-1} S_{v_{i-1}} x_i$ . By the definition of  $R'$ , there exists  $k$  such that  $n \leq k < i-1$  and  $v_{i-1} = x_k$ . Thus,  $x_{i-1} S_{x_k} x_i$ , and hence we have  $x_k R x_i$  by Facts 2.7.2 and 3. Since  $k = n$  or  $k \geq n+1$ , we have  $x_n R x_i$  by our assumption and the transitivity of  $R$ .  $\square$

By Claim 3, if  $\mathbf{x}R'\mathbf{y}$ , then  $\mathbf{x}^e R \mathbf{y}^e$ . Since  $R$  is conversely well-founded,  $R'$  is also conversely well-founded. Moreover, the following claim holds.

**Claim 4.** *Let  $\mathbf{x}, \mathbf{y} \in W'$  with  $\mathbf{x}R'\mathbf{y}$ ,  $\mathbf{x} = (\langle x_1, \dots, x_n \rangle, \langle v_1, \dots, v_{n-1} \rangle)$ , and  $\mathbf{y} = (\langle x_1, \dots, x_m \rangle, \langle v_1, \dots, v_{m-1} \rangle)$ . Then, for any  $i$  with  $n+2 \leq i \leq m$ ,  $x_{n+1} S_{x_n} x_i$ .*

*Proof.* Assume for any  $j$  with  $n+2 \leq j < i$ ,  $x_{n+1} S_{x_n} x_j$ . Then, we show  $x_{n+1} S_{x_n} x_i$ . We distinguish the following two cases.

Case 1:  $v_{i-1} = 0$ . Then,  $x_{i-1} R x_i$ . On the other hand,  $x_n R x_{i-1}$  by Claim 3. By Fact 2.7.4,  $x_{i-1} S_{x_n} x_i$ . If  $i-1 = n+1$ , then we are done. Let  $i-1 \geq n+2$ . By our assumption,  $x_{n+1} S_{x_n} x_{i-1}$ . Then, we obtain  $x_{n+1} S_{x_n} x_i$  by Fact 2.7.2.



Case 2:  $v_{i-1} \neq 0$ . Then,  $x_{i-1}S_{v_{i-1}}x_i$ . By the definition of  $R'$ , there exists a natural number  $k$  such that  $n \leq k < i-1$  and  $v_{i-1} = x_k$ . If  $k = n$ , then we obtain  $x_{n+1}S_{x_n}x_i$  as in the case of  $v_{i-1} = 0$ . Let  $k \geq n+1$ . Then,  $x_nRx_k$  by Claim 3. On the other hand,  $x_kRx_i$  by Facts 2.7.2 and 3. Thus, we have  $x_kS_{x_n}x_i$  by Fact 2.7.4. If  $k = n+1$ , then we are done. Even if  $k \geq n+2$ , then we obtain  $x_{n+1}S_{x_n}x_i$  by our assumption and Fact 2.7.2.  $\square$

Then, we show the following claim.

**Claim 5.** *For any modal formula  $A'$  and  $\mathbf{x} \in W'$ ,*

$$\mathbf{x} \Vdash' A' \iff \mathbf{x}^e \Vdash A'.$$

*Proof.* Let  $\mathbf{x} = (\langle x_1, \dots, x_n \rangle, \langle v_1, \dots, v_{n-1} \rangle)$ . We show the claim by induction on the construction of  $A'$ . We only prove the case  $A' \equiv B \triangleright C$ .

( $\implies$ ): Assume  $\mathbf{x} \Vdash' B \triangleright C$ . We show  $\mathbf{x}^e \Vdash B \triangleright C$ . Let  $y \in W$  with  $\mathbf{x}^e Ry$  and  $y \Vdash B$ , and let  $\mathbf{y} := (\langle x_1, \dots, x_n, y \rangle, \langle v_1, \dots, v_{n-1}, 0 \rangle)$ . Then,  $\mathbf{y} \in W'$  and  $\mathbf{x} \subset \mathbf{y}$ . Also, the condition (1) in the definition of  $R'$  obviously holds, and hence  $\mathbf{x}R'\mathbf{y}$ . Since  $\mathbf{y} \Vdash' B$  by the induction hypothesis, there exists  $\mathbf{z} \in W'$  such that  $\mathbf{x}R'\mathbf{z}$ ,  $\mathbf{y}S'\mathbf{z}$ , and  $\mathbf{z} \Vdash' C$  by our assumption. Since  $\mathbf{x} \subset \mathbf{z}$ , we have  $\mathbf{z} = (\langle x_1, \dots, x_n, x_{n+1}, \dots, x_m \rangle, \langle v_1, \dots, v_{n-1}, v_n, \dots, v_{m-1} \rangle)$  with  $n < m$ . Also, since  $\mathbf{y} = R(\mathbf{y})$  and  $R(\mathbf{y}) \subseteq R(\mathbf{z})$  by the definition of  $S'$ , we have  $x_{n+1} = y$ . Moreover, since  $|\mathbf{y}| < |\mathbf{z}|$ ,  $m \geq n+2$ . By Claim 4, we obtain  $yS_{\mathbf{x}^e}\mathbf{z}^e$ . By the induction hypothesis,  $\mathbf{z}^e \Vdash C$ , and hence  $\mathbf{x}^e \Vdash B \triangleright C$ .

( $\impliedby$ ): Assume  $\mathbf{x}^e \Vdash B \triangleright C$ . We show  $\mathbf{x} \Vdash' B \triangleright C$ . Let  $\mathbf{y} \in W'$  with  $\mathbf{x}R'\mathbf{y}$  and  $\mathbf{y} \Vdash' B$ , and let  $\mathbf{y} = (\langle x_1, \dots, x_n, x_{n+1}, \dots, x_m \rangle, \langle v_1, \dots, v_{n-1}, v_n, \dots, v_{m-1} \rangle)$ . By Claim 3 and the induction hypothesis, we have  $\mathbf{x}^e R\mathbf{y}^e$  and  $\mathbf{y}^e \Vdash B$ . By our assumption, there exists  $z \in W$  such that  $\mathbf{y}^e S_{\mathbf{x}^e}z$  and  $z \Vdash C$ . Let  $\mathbf{z} := (\langle x_1, \dots, x_m, z \rangle, \langle v_1, \dots, v_{m-1}, \mathbf{x}^e \rangle)$ . Then, obviously,  $\mathbf{z} \in W'$  and  $\mathbf{x}R'\mathbf{z}$ . Since  $R(\mathbf{y}) = R(\mathbf{z})$  and  $|\mathbf{y}| < |\mathbf{z}|$ , we obtain  $\mathbf{y}S'\mathbf{z}$ . By the induction hypothesis,  $\mathbf{z} \Vdash' C$ , and hence  $\mathbf{x} \Vdash' B \triangleright C$ .  $\square$

By Claim 5, we conclude  $(\langle w \rangle, \epsilon) \not\Vdash A'$ .  $\square$

Finally, we show that  $\mathbf{IL}^-(\mathbf{J2}_+, \mathbf{J5})$  loses the finite model property.

**Corollary 3.5.** *The logic  $\mathbf{IL}^-(\mathbf{J2}_+, \mathbf{J5})$  does not have finite model property with respect to simplified  $\mathbf{IL}^-(\mathbf{J2}_+, \mathbf{J5})$ -frames.*

*Proof.* Suppose that  $\mathbf{IL}^-(\mathbf{J2}_+, \mathbf{J5})$  has finite model property with respect to simplified  $\mathbf{IL}^-(\mathbf{J2}_+, \mathbf{J5})$ -frames. Then, we show that  $\mathbf{IL}$  also has finite model property with respect to simplified  $\mathbf{IL}$ -frames. This contradicts Fact 2.15.2.

Let  $A$  be any modal formula, and suppose  $\mathbf{IL} \not\Vdash A$ . It suffices to show that there exists a finite simplified  $\mathbf{IL}$ -frame in which  $A$  is not valid. We have

$$\mathbf{IL}^-(\mathbf{J2}_+, \mathbf{J5}) \not\Vdash \bigwedge \{(C \triangleright C) \wedge \Box(C \triangleright C) : C \in \text{Sub}(A)\} \rightarrow A$$

where  $\text{Sub}(A)$  is the set of all subformulas of  $A$ . By our supposition, there exist a finite simplified  $\mathbf{IL}^-(\mathbf{J2}_+, \mathbf{J5})$ -frame  $\mathcal{F} = (W, R, S)$  and  $w \in W$  such that

$w \Vdash \bigwedge \{(C \triangleright C) \wedge \Box(C \triangleright C) : C \in \text{Sub}(A)\}$ , and  $w \not\Vdash A$ . Here, we may assume that  $w$  is the root of  $W$ , that is, for any  $x \in W$ ,  $wRx$  or  $w = x$ . Hence, for any  $x \in W$  and any  $C \in \text{Sub}(A)$ ,  $x \Vdash C \triangleright C$ . We define the binary relation  $S'$  and the forcing relation  $\Vdash'$  as follows:

- $yS'z : \iff ySz$  or  $y = z$ .
- $x \Vdash' p : \iff x \Vdash p$ .

Obviously,  $\mathcal{F}' = (W, R, S')$  is a finite simplified **IL**-frame. Then, we prove the following claim.

**Claim 6.** *For any modal formula  $A' \in \text{Sub}(A)$  and  $x \in W$ ,*

$$x \Vdash' A' \iff x \Vdash A'.$$

*Proof.* We show the claim by induction on the construction of  $A'$ . We only prove the case  $A' \equiv B \triangleright C$ .

( $\implies$ ): Assume  $x \Vdash' B \triangleright C$  and we show  $x \Vdash B \triangleright C$ . Let  $y \in W$  with  $xRy$  and  $y \Vdash B$ . By the induction hypothesis,  $y \Vdash' B$ , and hence there exists  $z \in W$  such that  $xRz$ ,  $yS'z$ , and  $z \Vdash' C$ . By the induction hypothesis,  $z \Vdash C$ . If  $ySz$ , then we are done. If  $y = z$ , then since  $x \Vdash C \triangleright C$ , there exists  $z' \in W$  such that  $xRz'$ ,  $ySz'$ , and  $z' \Vdash C$ . Therefore, we obtain  $x \Vdash B \triangleright C$ .

( $\impliedby$ ): This is trivial. □

Since  $w \not\Vdash A$ ,  $w \not\Vdash' A$  by Claim 6. Therefore, we conclude  $\mathcal{F}' \not\equiv A$ . □

## 4 Concluding remarks

In this paper, we investigated finite model property for several sublogics of **IL** with respect to simplified Veltman frames. Among other things, we proved that **CL** has finite model property with respect to simplified **CL**-frames (Theorem 3.3). Then, Ignatiev's proof of the arithmetical completeness for **CL** is verified.

Moreover, our result is applicable to completeness with respect to topological semantics. In [5], Iwata and Kurahashi proved that **CL** is complete with respect to their topological semantics by translating a simplified **CL**-frame to a bitopological space. Since Ignatiev's counter simplified **CL**-model is infinite, Iwata and Kurahashi asked whether **CL** has finite model property with respect to topological semantics or not. By Theorem 3.3, we affirmatively solve this problem.

**Corollary 4.1.** *The logic **CL** has finite model property with respect to Iwata and Kurahashi's topological semantics.*

## Acknowledgement

The research was supported by Foundation of Research Fellows, The Mathematical Society of Japan.

## Declaration

**Conflict of interest:** The author declares that he has no conflict of interest.

## References

- [1] Alessandro Berarducci. The interpretability logic of Peano arithmetic. *The Journal of Symbolic Logic*, 55(3):1059–1089, 1990.
- [2] Dick de Jongh and Frank Veltman. Provability logics for relative interpretability. In *Mathematical logic*, pages 31–42, New York, 1990. Plenum Press.
- [3] Petr Hájek and Franco Montagna. The logic of  $\Pi_1$ -conservativity. *Archive for Mathematical Logic*, 30(2):113–123, 1990.
- [4] Konstantin N. Ignatiev. Partial conservativity and modal logics. *ITLI Publication Series X-91-04*, 1991.
- [5] Sohei Iwata and Taishi Kurahashi. Topological semantics of conservativity and interpretability logics. *Journal of Logic and Computation*, 31(7):1716–1739, 2021.
- [6] Sohei Iwata, Taishi Kurahashi, and Yuya Okawa. The persistence principle over the logic  $\mathbf{IL}^-$ . *arXiv preprint arXiv:2203.02183*, 2022.
- [7] Taishi Kurahashi and Yuya Okawa. Modal completeness of sublogics of the interpretability logic  $\mathbf{IL}$ . *Mathematical Logic Quarterly*, 67(2):164–185, 2021.
- [8] V. Yu. Shavrukov. The logic of relative interpretability over Peano arithmetic (in Russian). Technical Report 5, Steklov Mathematical Institute, Moscow, 1988.
- [9] Robert M. Solovay. Provability interpretations of modal logic. *Israel Journal of Mathematics*, 25(3-4):287–304, 1976.
- [10] Albert Visser. Preliminary notes on interpretability logic. Technical Report 29, Department of Philosophy, Utrecht University, 1988.
- [11] Albert Visser. An overview of interpretability logic. In *Advances in modal logic*, volume 1, pages 307–359, 1997.