

A VARIATION OF CONTINUITY IN  $n$ -NORMED SPACES

SIBEL ERSAN  
MALTEPE UNIVERSITY  
TURKEY  
SIBELERSAN@MALTEPE.EDU.TR

**ABSTRACT.** The  $s$ -th forward difference sequence that tends to zero, inspired by the consecutive terms of a sequence approaching zero, is examined in this study. Functions that take sequences satisfying this condition to sequences satisfying the same condition are called  $s$ -ward continuous. Inclusion theorems that are related to this kind of uniform continuity and continuity are also considered. Additionally, the concept of  $s$ -ward compactness of a subset of  $X$  via  $s$ -quasi-Cauchy sequences are investigated. One finds out that the uniform limit of any sequence of  $s$ -ward continuous function is  $s$ -ward continuous and the set of  $s$ -ward continuous functions is a closed subset of the set of continuous functions.

## 1. INTRODUCTION AND PRELIMINARIES

Although some evaluations was first made about the axioms of an abstract  $n$ -dimensional metric, the main developments regarding the definition of the 2-metric, 2-normed spaces and their topological properties were described by Gähler [22] then the results of these concepts were extended to the most generalized case  $n$ -metric and  $n$ -normed spaces where  $n$  is a natural number by Gähler[23]. Shortly after the concept of  $n$ -normed space is introduced, the concept of 2-inner product space is also defined in [3]. Afterwards many authors have done lots of impressive improvements in  $n$ -normed spaces or in 2-inner product spaces ([2, 20, 19, 13, 15, 11, 16]).

Firstly we recall the notion of  $n$ -normed space:

**Definition 1.1.** An  $n$ -norm on a real vector space  $X$  of dimension  $d$ , where  $2 \leq n \leq d$  is a real valued function  $\|\cdot, \dots, \cdot\|$  on  $X^n$  which satisfies the properties:

- (1)  $\|\zeta_1, \zeta_2, \dots, \zeta_n\| = 0$  if and only if  $\zeta_1, \zeta_2, \dots, \zeta_n$  are linearly dependent,
- (2)  $\|\zeta_1, \zeta_2, \dots, \zeta_n\| = \|\zeta_{k_1}, \dots, \zeta_{k_n}\|$  for every permutation  $(k_1, \dots, k_n)$  of  $(1, \dots, n)$ ,
- (3)  $\|\zeta_1, \zeta_2, \dots, \delta\zeta_n\| = |\delta| \|\zeta_1, \zeta_2, \dots, \zeta_n\|$  for any real number  $\delta$ ,
- (4)  $\|m + n, \zeta_1, \dots, \zeta_{n-1}\| \leq \|m, \zeta_1, \dots, \zeta_{n-1}\| + \|n, \zeta_1, \dots, \zeta_{n-1}\|$ .

A set  $X$  is an  $n$ -normed space with an  $n$ -norm  $\|\cdot, \dots, \cdot\|$ .

In [12],

$$(1) \quad \|\zeta_1, \dots, \zeta_n\|_p = \left[ \frac{1}{n!} \sum_{t_1} \dots \sum_{t_n} \det(\zeta_{it_k})^p \right]^{1/p}.$$

---

2020 *Mathematics Subject Classification.* 40A05, 40A25, 40A30, 54C35.  
*Key words and phrases.* compactness, continuity,  $n$ -normed space.

is given as an example of an  $n$ -norm on  $l^p \times \dots \times l^p$  space for  $1 \leq p < \infty$ . Also if  $p = \infty$ , the  $n$ -norm on  $l^\infty \times \dots \times l^\infty$  is given as [1]

$$(2) \quad \|\zeta_1, \dots, \zeta_n\|_\infty = \sup_{t_1} \dots \sup_{t_n} \det(x_{it_k}).$$

**Definition 1.2.** A sequence  $(x_k)$  converges to an  $\zeta \in X$  in an  $n$ -normed space  $X$  if for each  $\epsilon > 0$ , there exists a positive integer  $\tilde{k}$  such that for every  $k \geq \tilde{k}$

$$(3) \quad \|x_k - \zeta, \mu_1, \dots, \mu_{n-1}\| < \epsilon, \quad \forall \mu_1, \dots, \mu_{n-1} \in X.$$

**Definition 1.3.** A sequence  $(x_k)$  is a Cauchy sequence if for each  $\epsilon > 0$ , there exists a positive integer  $t_0$  such that for every  $k, m \geq t_0$

$$(4) \quad \|x_k - x_m, \mu_1, \dots, \mu_{n-1}\| < \epsilon, \quad \forall \mu_1, \dots, \mu_{n-1} \in X.$$

If each Cauchy sequence in  $X$  converges to an element of  $X$ , we call  $X$  is complete and if  $X$  is complete, then it is called an  $n$ -Banach space.

In recent times, the notion of quasi-Cauchy sequences is given in [4]. The distance between consecutive terms of a sequence tending to zero is expressed by Burton and Coleman with the idea of quasi-Cauchy sequence. Then using this idea, different types of continuities were defined for real functions in [6, 7] as ward continuity, statistically ward continuity, lacunary ward continuity and etc. They were also studied in 2-normed space in [24, 9, 10].

The aim of this research is to give a generalization of the notions of a quasi-Cauchy sequence and ward continuity of a function to the notions of an  $s$ -quasi-Cauchy sequence and  $s$ -ward continuity of a function in an  $n$ -normed space for any fixed positive integer  $s$ . Also interesting theorems related to ordinary continuity, uniform continuity, compactness and  $s$ -ward continuity are proved. This paper contains not only an extension of results of [24] to an  $n$ -normed space, but also includes new results in 2-normed spaces as a special case for  $n = 2$ .

## 2. MAIN RESULTS

In this paper  $X$ ,  $\mathbb{R}$  and  $s$  will denote a first countable  $n$ -normed space with an  $n$ -norm  $\|\cdot, \dots, \cdot\|$ , the set of all real numbers and a fixed positive integer, respectively. Now we give the notion of  $s$ -quasi Cauchyness of a sequence in  $X$ :

**Definition 2.1.** A sequence  $(x_k)$  of points in  $X$  is said to be  $s$ -quasi-Cauchy if for all  $\mu_1, \mu_2, \dots, \mu_{n-1} \in X$  it satisfies

$$(5) \quad \lim_{k \rightarrow \infty} \|\Delta_s x_k, \mu_1, \mu_2, \dots, \mu_{n-1}\| = 0$$

where  $\Delta_s x_k = x_{k+s} - x_k$  for each positive integer  $k$ .

If one chooses  $s = 1$ , the sequences returns to the ordinary quasi-Cauchy sequences and also using the equality

$$x_{k+s} - x_k = x_{k+s} - x_{k+s-1} + x_{k+s-1} - x_{k+s-2} \dots - x_{k+2} + x_{k+2} - x_{k+1} + x_{k+1} - x_k,$$

we see that any quasi-Cauchy sequence is  $s$ -quasi-Cauchy, however the converse is not true.

Any Cauchy sequence is  $s$ -quasi-Cauchy, so is any convergent sequence. A sequence of partial sums of a convergent series is  $s$ -quasi-Cauchy. One notes that the

set  $\Delta_s(X)$ , the set of  $s$ -quasi-Cauchy sequences in  $X$ , is a vector space. If  $(x_k), (y_k)$  are  $s$ -quasi-Cauchy sequences in  $X$  so

$$(6) \quad \lim_{k \rightarrow \infty} \|\Delta_s x_k, \mu_1, \mu_2, \dots, \mu_{n-1}\| = 0 \quad \text{and}$$

$$(7) \quad \lim_{k \rightarrow \infty} \|\Delta_s y_k, \mu_1, \mu_2, \dots, \mu_{n-1}\| = 0.$$

for all  $\mu_1, \mu_2, \dots, \mu_{n-1} \in X$ . Therefore

$$(8) \quad \begin{aligned} \lim_{k \rightarrow \infty} \|\Delta_s(x_k + y_k), \mu_1, \mu_2, \dots, \mu_{n-1}\| &\leq \lim_{k \rightarrow \infty} \|\Delta_s x_k, \mu_1, \mu_2, \dots, \mu_{n-1}\| \\ &+ \lim_{k \rightarrow \infty} \|\Delta_s y_k, \mu_1, \mu_2, \dots, \mu_{n-1}\| = 0. \end{aligned}$$

So the sum of two  $s$ -quasi-Cauchy sequence is again  $s$ -quasi-Cauchy, it is clear that  $(ax_k)$  is an  $s$ -quasi-Cauchy sequence in  $X$  for any constant  $a \in \mathbb{R}$ .

**Definition 2.2.** *A subset  $A$  of  $X$  is called  $s$ -ward compact if any sequence in the set  $A$  has an  $s$ -quasi-Cauchy subsequence.*

If a set  $A$  is an  $s$ -ward compact subset of  $X$ , then any subset of  $A$  is  $s$ -ward compact. Moreover any ward compact subset of  $X$  is  $s$ -ward compact. Union of finite number of  $s$ -ward compact subset of  $X$  is  $s$  ward compact. Any sequentially compact subset of  $X$  is  $s$ -ward compact.

For each real number  $\alpha > 0$ , an  $\alpha$ -ball with center  $a$  in  $X$  is defined as

$$(9) \quad B_\alpha(a, x_1, \dots, x_{n-1}) = \{x \in X : \|a - x, x_1 - x, \dots, x_{n-1} - x\| < \alpha\}$$

for  $x_1, \dots, x_{n-1} \in X$ . The family of all sets  $W_i(a) = B_{\alpha_i}(a, x_{i_1}, \dots, x_{i_{(n-1)}})$  where  $i = 1, 2, \dots$  is an open basis in  $a$ . Let  $\beta_{n-1}$  be the collection of linearly independent sets  $B$  with  $n - 1$  elements. For  $B \in \beta_{n-1}$ , the mapping

$$p_B(x) = \|x, x_1, \dots, x_{n-1}\|, \text{ for } x \in X, \quad x_1, \dots, x_{n-1} \in B$$

defines a seminorm on  $X$  and the collection  $\{p_B : B \in \beta_{n-1}\}$  of seminorms makes  $X$  a locally convex topological vector space. For each  $x \in X$ , different from zero, there exists  $x_1, \dots, x_{n-1} \in B$  such that  $x, x_1, x_2, \dots, x_{n-1}$  are linearly independent so  $\|x, x_1, \dots, x_{n-1}\| \neq 0$ , which makes  $X$  a Hausdorff space. A neighborhood of origin for this topology is in a form of a finite intersection

$$\bigcap_{i=1}^n \{x \in X : \|x, x_{i_1} - x, \dots, x_{i_{(n-1)}} - x\| < \epsilon\}$$

where  $\epsilon > 0$ .

Now the following theorem characterizes totally boundedness not only valid for  $n$ -normed spaces but also valid for the 2-normed spaces. It extends the results for quasi-Cauchy sequences given in [24] for 2-normed valued sequences to  $n$ -normed valued  $s$ -quasi-Cauchy sequences in which  $s = 1$  gives earlier results given for 2-normed spaces. It should be noted that Theorem 3 in [8] can not be obtained just by putting  $n = 1$  in the  $n$ -normed space to get in a normed space, which is awkward, whereas the following theorem is interesting as a point of studying a new space.

**Lemma 2.1.** *If a subset of  $X$  is totally bounded then every sequence in  $A$  contains an  $s$ -quasi-Cauchy subsequence.*

*Proof.* Let  $A$  be totally bounded. Let  $(x_n)$  be any sequence in  $A$ . Since  $A$  is covered by a finite number of balls of  $X$  of diameter less than 1. One of these sets, which we denote by  $A_1$ , must contain  $x_n$  for infinitely many values of  $n$ . Choose a positive integer  $n_1$  such that  $x_{n_1} \in A_1$ . Since  $A_1$  is totally bounded, it is covered by a finite number of balls of diameter less than  $1/2$ . One of these balls, which we denote by  $A_2$ , must contain  $x_n$  for infinitely many  $n$ . Let  $n_2$  be a positive integer such that  $n_2 > n_1$  and  $x_{n_2} \in A_2$ . Since  $A_2 \subset A_1$ , it follows that  $x_{n_2} \in A_1$ . Continuing in this way, a ball  $A_k$  of  $A_{k-1}$  of diameter less than  $1/k$  and a term  $x_{n_k} \in A_k$  of the sequence  $(x_n)$ , where  $n_k > n_{k-1}$  for any positive integer  $k$ . Since  $x_{n_k}, x_{n_{k+1}}, \dots, x_{n_{k+s}}, \dots$  lie in  $A_k$  and diameter ( $A_k$ ) less than  $1/k$ , then  $x_{n_k}$  is an  $s$ -quasi-Cauchy subsequence of  $(x_n)$ .  $\square$

**Theorem 2.2.** *A subset of  $X$  is totally bounded if and only if it is  $s$ -ward compact for any positive integer  $s$ .*

*Proof.* If  $A$  is totally bounded, then every sequence of  $A$  has an  $s$ -quasi-Cauchy subsequence by Lemma 2.1. So the set  $A$  is  $s$ -ward compact for any fixed positive integer  $s$ . For the converse, think  $A$  is not a totally bounded set. Choose any  $x_1 \in A$  and  $\alpha > 0$ . Since  $A$  is not totally bounded, the neighborhood of a point  $x_1$  in  $A$  which is defined by  $B_\alpha(x_1, \mu_1^1, \dots, \mu_{n-1}^1) = \{y \in A; \|x_1 - y, \mu_1^1 - y, \dots, \mu_{n-1}^1 - y\| < \alpha\}$  is not equal to  $A$ . There is an  $x_2 \in A$  such that  $x_2 \notin B_\alpha(x_1, \mu_1^1, \dots, \mu_{n-1}^1)$ , that is,  $\|x_1 - x_2, \mu_1^1 - x_2, \dots, \mu_{n-1}^1 - x_2\| \geq \alpha$ . Since  $A$  is not totally bounded  $B_\alpha(x_1, \mu_1^1, \dots, \mu_{n-1}^1) \cup B_\alpha(x_2, \mu_1^2, \dots, \mu_{n-1}^2) \neq A$  where  $B_\alpha(x_2, \mu_1^2, \dots, \mu_{n-1}^2)$  is the neighborhood of a point  $x_2$  in  $A$ . Continuing the procedure, a sequence  $(x_k)$  of points in  $A$  can be obtained as  $x_{k+s} \notin \bigcup_{i=1}^{k+s-1} B_\alpha(x_i, \mu_1^i, \dots, \mu_{n-1}^i)$ . So  $\|x_{k+s} - x_k, \mu_1^i - x_k, \dots, \mu_{n-1}^i - x_k\| \geq \alpha$  and all nonzero  $\mu_1^i, \dots, \mu_{n-1}^i \in A$  where  $i = 1, \dots, k+s-1$ . So the sequence  $(x_k)$  has not any  $s$ -quasi-Cauchy subsequence. Therefore  $A$  is not  $s$ -ward compact.  $\square$

**Definition 2.3.** *A function  $f$  is called  $s$ -ward continuous on a subset  $A$  of  $X$  if*

$$(10) \quad \lim_{k \rightarrow \infty} \|\Delta_s x_k, \mu_1, \mu_2, \dots, \mu_{n-1}\| = 0$$

*is satisfied for all  $\mu_1, \mu_2, \dots, \mu_{n-1} \in X$ , then*

$$(11) \quad \lim_{k \rightarrow \infty} \|\Delta_s f(x_k), f(\mu_1), f(\mu_2), \dots, f(\mu_{n-1})\| = 0.$$

In the following we give that any  $s$ -ward continuous function is continuous.

**Theorem 2.3.** *Any  $s$ -ward continuous function on a subset  $A$  of  $X$  is continuous on  $A$ .*

*Proof.* Let the function  $f$  be  $s$ -ward continuous on  $A \subset X$  and any sequence  $(x_k)$  in  $A$  converge to  $\zeta$ , that is

$$(12) \quad \lim_{k \rightarrow \infty} \|x_k - \zeta, \mu_1, \mu_2, \dots, \mu_{n-1}\| = 0$$

for all  $\mu_1, \mu_2, \dots, \mu_{n-1} \in X$ . Let us write a new sequence using some terms of the sequence  $(x_k)$  as

$$(t_m) = (x_1, \dots, x_1, \zeta, \dots, \zeta, x_2, \dots, x_2, \zeta, \dots, \zeta, \dots, x_n, \dots, x_n, \zeta, \dots, \zeta, \dots).$$

where same terms repeated  $s$ -times. Every convergent sequences is Cauchy and moreover any Cauchy sequence is  $s$ -quasi-Cauchy, then it follows that

$$\lim_{m \rightarrow \infty} \|\Delta_s t_m, \mu_1, \mu_2, \dots, \mu_{n-1}\| = \lim_{m \rightarrow \infty} \|t_{m+s} - t_m, \mu_1, \mu_2, \dots, \mu_{n-1}\| = 0$$

in which either

$$\lim_{m \rightarrow \infty} \|t_{m+s} - \zeta, \mu_1, \mu_2, \dots, \mu_{n-1}\| = 0$$

or

$$\lim_{m \rightarrow \infty} \|\zeta - t_m, \mu_1, \mu_2, \dots, \mu_{n-1}\| = 0$$

for every  $\mu_1, \mu_2, \dots, \mu_{n-1}$ . This result implies  $(t_m)$  is an  $s$ -quasi Cauchy sequence. Since the function  $f$  is assumed to be  $s$ -ward continuous, using this assumption we get

$$(13) \quad \begin{aligned} & \lim_{m \rightarrow \infty} \|\Delta_s f(t_m), f(\mu_1), f(\mu_2), \dots, f(\mu_{n-1})\| \\ &= \lim_{m \rightarrow \infty} \|f(t_{m+s}) - f(t_m), f(\mu_1), f(\mu_2), \dots, f(\mu_{n-1})\| = 0 \end{aligned}$$

in which either

$$\lim_{m \rightarrow \infty} \|f(t_{m+s}) - f(\zeta), f(\mu_1), f(\mu_2), \dots, f(\mu_{n-1})\| = 0$$

or

$$\lim_{m \rightarrow \infty} \|f(\zeta) - f(t_m), f(\mu_1), f(\mu_2), \dots, f(\mu_{n-1})\| = 0.$$

So  $(f(x_k))$  converges to  $f(\zeta)$ .  $\square$

As the sum of two  $s$ -ward continuous function on  $A$  is  $s$ -ward continuous and  $cf$  is  $s$ -ward continuous for any constant real number  $c$ , the set of  $s$ -ward continuous functions on  $A$  is a vector subspace of vector space of all continuous function on  $A$ .

**Theorem 2.4.** *Every  $s$ -ward continuous function on  $A \subset X$  is ward continuous on  $A$ .*

*Proof.* Assume that  $(x_k)$  is a quasi-Cauchy sequence in  $A$  and  $f$  is any  $s$ -ward continuous function on  $A$ . If  $s = 1$ , the result is obvious. Let  $s > 1$  and a sequence

$$(t_m) = \underbrace{(x_1, x_1, \dots, x_1)}_{s\text{-times}}, \underbrace{(x_2, x_2, \dots, x_2)}_{s\text{-times}}, \dots, \underbrace{(x_n, x_n, \dots, x_n)}_{s\text{-times}}, \dots$$

be  $s$ -quasi-Cauchy, i.e.

$$(14) \quad \lim_{m \rightarrow \infty} \|\Delta_s t_m, \mu_1, \mu_2, \dots, \mu_{n-1}\| = 0.$$

We have

$$(15) \quad \lim_{m \rightarrow \infty} \|\Delta_s f(t_m), f(\mu_1), f(\mu_2), \dots, f(\mu_{n-1})\| = 0$$

by using the  $s$ -ward continuity of the function  $f$ . Therefore

$$(16) \quad \lim_{m \rightarrow \infty} \|\Delta f(t_m), f(\mu_1), f(\mu_2), \dots, f(\mu_{n-1})\| = 0$$

So  $s$ -ward continuity of the function  $f$  implies that the ward continuity of  $f$  on  $A \subset X$ .  $\square$

**Theorem 2.5.** *The image of an  $s$ -ward compact subset of  $X$  by an  $s$ -ward continuous function is  $s$ -ward compact.*

*Proof.* Assume that  $f$  is an  $s$ -ward continuous function and  $A$  is an  $s$ -ward compact subset of  $X$ . Choose a sequence  $t$  as  $t = (t_k) \in f(A)$  and say  $(t_k) = f(x_k)$  where  $x_k \in A$ .  $A$  is  $s$ -ward compact so there is a subsequence  $(x_m)$  of  $(x_k)$  with

$$(17) \quad \lim_{m \rightarrow \infty} \|\Delta_s x_m, \mu_1, \mu_2, \dots, \mu_{n-1}\| = 0$$

for all  $\mu_1, \mu_2, \dots, \mu_{n-1} \in X$ . Using the  $s$ -ward continuity of  $f$  we have

$$(18) \quad \lim_{m \rightarrow \infty} \|\Delta_s f(x_m), f(\mu_1), f(\mu_2), \dots, f(\mu_{n-1})\| = 0,$$

so there is an  $s$ -quasi-Cauchy subsequence  $(f(x_m))$  of  $t$ . The result implies that the subset  $f(A) \subset X$  is  $s$ -ward compact.  $\square$

$s$ -ward continuous image of any compact subset of  $X$  is compact. It is easily evaluated from Theorem 2.3.

**Theorem 2.6.** *If  $f$  is uniformly continuous on  $A \subset X$ , then it is  $s$ -ward continuous on  $A$ .*

*Proof.* Let  $f$  be a uniformly continuous function on  $A$ , and the sequence  $(x_k)$  be an  $s$ -quasi-Cauchy sequence in  $A$ . Our aim is to prove the sequence  $(f(x_k))$  is also an  $s$ -quasi-Cauchy sequence in  $A$ . Take any  $\varepsilon > 0$ . There exists a  $\delta > 0$  such that if

$$(19) \quad \|x - y, \mu_1, \mu_2, \dots, \mu_{n-1}\| < \delta \quad \text{then} \quad \|f(x) - f(y), f(\mu_1), f(\mu_2), \dots, f(\mu_{n-1})\| < \varepsilon.$$

There exists an  $\tilde{k} = \tilde{k}(\delta)$  for this  $\delta > 0$  such that

$$(20) \quad \|\Delta_s x_k, \mu_1, \mu_2, \dots, \mu_{n-1}\| < \delta$$

for every  $\mu_1, \mu_2, \dots, \mu_{n-1} \in X$  whenever  $k > \tilde{k}$ . Uniform continuity of  $f$  on  $A$  for every  $k > \tilde{k}$  implies

$$(21) \quad \|\Delta_s f(x_k), f(\mu_1), f(\mu_2), \dots, f(\mu_{n-1})\| < \varepsilon$$

for every  $f(\mu_1), f(\mu_2), \dots, f(\mu_{n-1}) \in X$ . The sequence  $(f(x_k))$  is  $s$ -quasi-Cauchy so the function  $f$  is  $s$ -ward continuous.  $\square$

**Theorem 2.7.** *Uniform limit of a sequence of  $s$ -ward continuous function is  $s$ -ward continuous.*

*Proof.* Let  $(f_t)$  be a sequence of  $s$ -ward continuous functions and it be uniformly convergent sequence to a function  $f$ . Pick an  $s$ -quasi-Cauchy sequence  $(x_k)$  in  $A$  and choose any  $\varepsilon > 0$ . There is an integer  $N \in \mathbb{Z}^+$  such that

$$(22) \quad \|f_t(x) - f(x), f(\mu_1), f(\mu_2), \dots, f(\mu_{n-1})\| < \frac{\varepsilon}{3}$$

for every  $x \in A$ , for all  $f(\mu_1), f(\mu_2), \dots, f(\mu_{n-1}) \in X$  whenever  $t \geq N$ . Using the  $s$ -ward continuity of  $f_N$ , there is a positive integer  $N_1(\varepsilon) > N$  such that

$$(23) \quad \|\Delta_s f_t(x_k), f(\mu_1), f(\mu_2), \dots, f(\mu_{n-1})\| < \frac{\varepsilon}{3}$$

for every  $t \geq N_1$ . Now for  $t \geq N_1$  we have

$$(24) \quad \begin{aligned} & \|\Delta_s f(x_k), f(\mu_1), f(\mu_2), \dots, f(\mu_{n-1})\| \\ &= \|f(x_{k+s}) - f(x_k), f(\mu_1), f(\mu_2), \dots, f(\mu_{n-1})\| \\ &\leq \|f(x_{k+s}) - f_N(x_{k+s}), f(\mu_1), f(\mu_2), \dots, f(\mu_{n-1})\| \\ &\quad + \|\Delta_s f_N(x_k), f(\mu_1), f(\mu_2), \dots, f(\mu_{n-1})\| \\ &+ \|f_N(x_k) - f(x_k), f(\mu_1), f(\mu_2), \dots, f(\mu_{n-1})\| < \frac{\varepsilon}{3} + \frac{\varepsilon}{3} + \frac{\varepsilon}{3} = \varepsilon. \end{aligned}$$

So the function  $f$  is  $s$ -ward continuous on  $A$ .  $\square$

**Theorem 2.8.** *The collection of the  $s$ -ward continuous functions on  $A \subset X$  is a closed subset of the collection of every continuous functions on  $A$ .*

*Proof.* Let  $E$  be a collection of all  $s$ -ward continuous functions on  $A \subset X$  and  $\overline{E}$  is the closure of  $E$ .  $\overline{E}$  is defined as for every  $x \in X$  there exists  $x_k \in E$  with  $\lim_{k \rightarrow \infty} x_k = x$  and  $E$  is closed if  $E = \overline{E}$ . It is obvious that  $E \subseteq \overline{E}$ . Let  $f$  be any element of the set of all closure points of  $E$  which means there exists a sequence of points  $f_t$  in  $E$  as

$$(25) \quad \lim_{t \rightarrow \infty} \|f_t - f, f(\mu_1), f(\mu_2), \dots, f(\mu_{n-1})\| = 0$$

for all  $f(\mu_1), f(\mu_2), \dots, f(\mu_{n-1}) \in X$  and also  $f_t$  is a  $s$ -ward continuous function. Choose the sequence  $(x_k)$  as any  $s$ -quasi-Cauchy sequence. Since  $(f_t)$  converges to  $f$ , for every  $\varepsilon > 0$  and  $x \in E$ , there is any  $N_0$  such that for every  $t \geq N_0$ ,

$$(26) \quad \|f(x) - f_t(x), f(\mu_1), f(\mu_2), \dots, f(\mu_{n-1})\| < \frac{\varepsilon}{3}.$$

As  $f_N$  is  $p$ -ward continuous,  $N_1 > N_0$  exists such that for all  $t \geq N_1$ ,

$$(27) \quad \|\Delta_s f_N(x_k), f(\mu_1), f(\mu_2), \dots, f(\mu_{n-1})\| < \frac{\varepsilon}{3}.$$

Hence for all  $t \geq N_1$ ,

$$(28) \quad \begin{aligned} & \|\Delta_s f(x_k), f(\mu_1), f(\mu_2), \dots, f(\mu_{n-1})\| \\ &= \|f(x_{k+s}) - f(x_k), f(\mu_1), f(\mu_2), \dots, f(\mu_{n-1})\| \\ &\leq \|f(x_{k+s}) - f_N(x_{k+s}), f(\mu_1), f(\mu_2), \dots, f(\mu_{n-1})\| \\ &\quad + \|f(x_k) - f_N(x_k), f(\mu_1), f(\mu_2), \dots, f(\mu_{n-1})\| \\ &+ \|\Delta_s f_N(x_k), f(\mu_1), f(\mu_2), \dots, f(\mu_{n-1})\| < \frac{\varepsilon}{3} + \frac{\varepsilon}{3} + \frac{\varepsilon}{3} = \varepsilon. \end{aligned}$$

Since the function  $f$  is  $s$ -ward continuous in  $E$  then  $E = \overline{E}$ , using Theorem 2.3, it ends the proof.  $\square$

### 3. CONCLUSION

The notion of an  $n$ -normed space was given by thinking if there is a problem where  $n$ -norm topology works however norm topology doesn't. As an application of the notion of  $n$ -norm, we can examine that if a term in the definition of  $n$ -norm shows the change of shape then the  $n$ -norm stands for the associated volume of this surface. Suppose that for any particular output one needs  $n$ -inputs but with one main input and other  $(n-1)$ -inputs as dummy inputs required which accomplish the operation, so this concept may be used as an application in many areas of science. The generalization of the notions of quasi-Cauchy sequences and ward continuous functions to the notions of  $s$ -quasi-Cauchy sequences and  $s$ -ward continuous functions in  $n$ -normed spaces are investigated in this paper. Also we find out some interesting inclusion theorems related to the concepts of ordinary continuity, uniform continuity,  $s$ -ward continuity, and  $s$ -ward compactness. We prove that the uniform limit of a sequence of  $s$ -ward continuous function is  $s$ -ward continuous and the set of  $s$ -ward continuous functions is a closed subset of the set of continuous functions. We recommend research  $s$ -quasi-Cauchy sequences of points and fuzzy functions in an  $n$ -normed fuzzy space as a further study. However, due to the different structure, the methods of proof will not be similar to the one in this study. (see [14], [21]). Also we recommend investigate  $s$ -quasi-Cauchy sequences of double sequences in  $n$ -normed spaces as another further study (see [17], [18]).

## DECLARATIONS

**Ethical Approval** Not applicable

**Competing interests** Not applicable

**Authors' contributions** Not applicable

**Funding** Not applicable

**Availability of data and materials** Not applicable

## REFERENCES

- [1] A. Malceski,  $l^\infty$  as  $n$ -normed space, Mat. Bilten., 21, (1997), 103-110.
- [2] A. Misiak,  $n$ -inner product spaces, Mathematische Nachrichten, 140, (1989), 299-319.
- [3] C. Diminnie, S. Gähler and A. White, 2-inner product spaces, Demonstratio Math., 6, (1973), 525-536.
- [4] D. Burton, J. Coleman, Quasi-Cauchy sequences, Amer. Math. Monthly, 117 4 (2010), 328-333.
- [5] F. Lael, K. Nourouzi, Compact Operators Defined on 2-Normed and 2-Probabilistic Normed Spaces, Mathematical Problems in Engineering, 2009, (2009), Article Number: 950234.
- [6] H. Cakalli, Forward continuity, J. Comput. Anal. Appl., 13, 2, (2011), 225-230. MR 2012c:26004
- [7] H. Cakalli, Variations on Quasi-Cauchy Sequences, Filomat, 29:1, (2015), 13-19.
- [8] H. Cakalli, Statistical quasi-Cauchy Sequences, Mathematical and Computer Modelling, 54, (2011), 1620-1624.
- [9] H. Cakalli, S. Ersan, Strongly lacunary ward continuity in 2-normed spaces, The Scientific World Journal, 2014 (2014), 5 pages, Article ID 479679.
- [10] H. Cakalli, S. Ersan, Lacunary ward continuity in 2-normed spaces, Filomat, 29 (10) (2015), 2257-2263.
- [11] H. Dutta, On some  $n$ -normed linear space valued difference sequences, Journal of the Franklin Institute, 348, (2011), 2876-2883.
- [12] H. Gunawan, The space of  $p$ -summable sequences and its natural  $n$ -norm, Bull. Austral. Math. Soc., 64, (2001), 137-147.
- [13] H. Gunawan, M. Mashadi, On  $n$ -normed spaces, International Journal of Mathematics and Mathematical Sciences, 27(10), (2001), 631-639.
- [14] Lj D. R., Kocinac, V. A., Khan, K. M. A. S., Alshloul and H., Altaf, On some topological properties of intuitionistic 2-fuzzy  $n$ -normed linear spaces Hacettepe Journal of Mathematics and Statistics, 49 1, (2020), 208-220.
- [15] M. Gurdal, A. Sahnier, Ideal convergence in  $n$ -normal spaces and some new sequence spaces via  $n$ -norm, Malaysian Journal of Fundamental and Applied Sciences, 4(1), (2014).
- [16] M. Gurdal, N. Sari and E. Savas,  $A$ -statistically localized sequences in  $n$ -normed spaces, Commun. Fac. Sci. Univ. Ank. Ser. A1 Math. Stat., 69, 2, (2020), 1484-1497.
- [17] M. Mursaleen and SK, Sharma, Riesz Lacunary Almost Convergent Double Sequence Spaces Defined By Sequence Of Orlicz Functions Over  $N$ -Normed Spaces, TWMS Journal Of Pure And Applied Mathematics, 8 1, (2017),43-63.
- [18] N. Khan, Classes of  $I$ -Convergent Double Sequences over  $n$ -Normed Spaces, Journal of Function Spaces, (2016), Article Number7594031.
- [19] R. Malceski, Strong  $n$ -convex  $n$ -normed spaces, Mat. Bilten, 21, (1997), 81-102.
- [20] S.S. Kim and Y. J. Cho, Strict convexity in linear  $n$ -normed spaces, Demonstratio Math., 29 4, (1996),739-744.
- [21] S. Altundag and E. Kamber, Lacunary Delta-statistical convergence in intuitionistic fuzzy  $n$ -normed space, Journal Of Inequalities And Applications, 40, (2014).
- [22] S. Gähler, 2-metrische Räume und ihre topologische Struktur, Math. Nachr., 26, (1963), 115-148.
- [23] S. Gähler, Untersuchungen über verallgemeinerte  $m$ -metrische Räume I, Math. Nachr., 40, (1969), 165-189.
- [24] S. Ersan and H. Cakalli, Ward continuity in 2-normed Spaces, Filomat, 29:7, (2015), 1507-1513.



SIBEL ERSAN, FACULTY OF ENGINEERING AND NATURAL SCIENCES, MALTEPE UNIVERSITY, ISTANBUL, TURKEY

*Email address:* `sibelersan@maltepe.edu.tr`