

Cartier smoothness in prismatic cohomology

Vincent Bouis

Abstract

We introduce the notion of a p -Cartier smooth algebra. It generalises that of a smooth algebra and includes valuation rings over a perfectoid base. We give several characterisations of p -Cartier smoothness in terms of prismatic cohomology, and deduce a comparison theorem between syntomic and étale cohomologies under this hypothesis.

CONTENTS

1	Introduction	1
2	Prismatic cohomology of Cartier smooth algebras	4
3	Valuation rings are Cartier smooth	17
4	The syntomic-étale comparison theorem	28

1 INTRODUCTION

Let p be a prime number. The Cartier isomorphism is a fundamental cohomological property of smooth \mathbb{F}_p -algebras. It states that for any smooth \mathbb{F}_p -algebra A , the inverse Cartier map

$$C^{-1} : \begin{cases} \Omega_{A/\mathbb{F}_p}^n \longrightarrow \mathrm{H}^n(\Omega_{A/\mathbb{F}_p}^\bullet) \\ fdg_1 \wedge \cdots \wedge dg_n \longmapsto f^p g_1^{p-1} \cdots g_n^{p-1} dg_1 \wedge \cdots \wedge dg_n \end{cases}$$

is an isomorphism of \mathbb{F}_p -vector spaces for each $n \geq 0$. Assuming resolution of singularities in characteristic p , any \mathbb{F}_p -algebra would be locally smooth in the cdh-topology. The local rings in the cdh-topology are valuation rings, so valuation rings of characteristic p should be filtered colimits of smooth \mathbb{F}_p -algebras, and in particular should satisfy the Cartier isomorphism. Without assuming resolution of singularities, Gabber proved [KST21, Appendix] that valuation rings of characteristic p actually satisfy the Cartier isomorphism. Motivated by this result and the general lack of understanding of valuation rings, Kelly and Morrow [KM21] introduced *Cartier smooth algebras* as \mathbb{F}_p -algebras satisfying the Cartier isomorphism, and study their algebraic K -theory.

We develop here an analogue of this story in mixed-characteristic. The inverse Cartier map, as an extension of the Frobenius morphism, is specific to characteristic p , and we define p -Cartier smoothness for general ring homomorphisms essentially in terms of their reductions modulo p . This more general notion coincides with that introduced by Kelly–Morrow [KM21] in the special case that the base ring is \mathbb{F}_p . For an \mathbb{F}_p -algebra R , we denote by ϕ_R its Frobenius endomorphism.

Definition 1.1 (Cartier smoothness). *(1) A morphism $R \rightarrow S$ of \mathbb{F}_p -algebras is Cartier smooth, or S is a Cartier smooth R -algebra, if it is quasismooth, i.e., it is flat and its cotangent complex $\mathbb{L}_{S/R}$ is a flat S -module in degree 0 given by $\Omega_{S/R}^1$, and if the inverse Cartier map*

$$C^{-1} : \Omega_{S/R}^n \otimes_{R, \phi_R} R \longrightarrow \mathrm{H}^n(\Omega_{S/R}^\bullet)$$

is an isomorphism of R -modules for each $n \geq 0$.

- (2) A morphism $R \rightarrow S$ of commutative rings is p -Cartier smooth, or S is a p -Cartier smooth R -algebra, if R has bounded p^∞ -torsion (i.e., the p -primary torsion is killed by a fixed power of p), the natural map $S \otimes_R^{\mathbb{L}} R/p \rightarrow S/p$ is an equivalence in the derived category $\mathcal{D}(R/p)$ and the reduction $R/p \rightarrow S/p$ modulo p is Cartier smooth.

Said another way, a morphism $R \rightarrow S$ of commutative rings is p -Cartier smooth if it is p -quasismooth (Definition 2.1) and if its reduction $R/p \rightarrow S/p$ modulo p satisfies the Cartier isomorphism. The p -quasismoothness hypothesis is necessary not to lose control, when specialising to characteristic p , on the invariants we will study. Any smooth morphism of commutative rings is p -Cartier smooth (for any prime p). Regarding valuation rings, an extension of valuation rings (i.e., an injective morphism of valuation rings) in mixed-characteristic is in general not p -Cartier smooth. For instance, the morphism $\mathbb{Z}_p \rightarrow \overline{\mathbb{Z}}_p$, where $\overline{\mathbb{Z}}_p$ is the ring of integers of an algebraic closure $\overline{\mathbb{Q}}_p$ of \mathbb{Q}_p , is not p -quasismooth and does not satisfy the Cartier isomorphism: its cotangent complex is in degree 0, given by the torsion $\overline{\mathbb{Z}}_p$ -module $\overline{\mathbb{Q}}_p/\overline{\mathbb{Z}}_p$, and $\Omega_{(\overline{\mathbb{Z}}_p/p)/\mathbb{F}_p}^0 = \overline{\mathbb{Z}}_p/p$ contains nilpotent elements; in fact the de Rham complex of $\mathbb{F}_p \rightarrow \overline{\mathbb{Z}}_p/p$ is zero in positive degrees, and thus does not convey a lot of information. However, if the base valuation ring is sufficiently ramified, these obstructions vanish and we can prove the following general result. This generalises a theorem of Gabber in characteristic p , which is valid over perfect valuation rings (Theorem 3.4 below). We refer the reader to [BMS18, Definition 3.5] for the definition of perfectoid rings.

Theorem A (see Theorem 3.1). *Let C^+ be a valuation ring whose p -adic completion is a perfectoid ring. Let E^+ be a valuation ring extension of C^+ . Then the morphism $C^+ \rightarrow E^+$ is p -Cartier smooth.*

The proof proceeds by reduction to the case of valuation rings of characteristic p , and uses almost mathematics and deformation theory.

With the examples of valuation rings and smooth algebras in mind, we study the algebraic K -theory of general p -Cartier smooth algebras over a perfectoid ring. Recall that in certain cases algebraic K -theory is equipped with a motivic filtration with graded pieces $\mathbb{Z}(i)(-) \in \mathcal{D}(\mathbb{Z})$, for $i \geq 0$, called motivic cohomology. The motivic filtration, with its expected properties, was defined by Voevodsky for smooth varieties over a field. In characteristic p the Beilinson–Lichtenbaum conjecture, proved by Suslin and Voevodsky as a consequence of the Bloch–Kato conjecture [SV00], computes the ℓ -adic part of motivic cohomology in terms of the étale cohomology. More precisely, given a field k of characteristic p , a smooth k -variety X and a prime ℓ different from p , it states that there is a natural quasi-isomorphism

$$\mathbb{Z}/\ell^n(i)(X) \simeq R\Gamma_{Zar}(X, \tau^{\leq i} R\alpha_* \mu_{\ell^n}^{\otimes i}),$$

where μ_{ℓ^n} is the étale sheaf of ℓ^n -roots of unity and $\alpha : (Sm/k)_{\acute{e}t} \rightarrow (Sm/k)_{Zar}$ is the projection of sites.

The Beilinson–Lichtenbaum conjecture can be interpreted as a comparison between Zariski motivic cohomology and étale motivic cohomology. The étale sheaf μ_{ℓ^n} is indeed naturally identified with the étale sheafification $\mathbb{Z}/\ell^n(i)^{\acute{e}t}$ of the (Zariski) motivic sheaf $\mathbb{Z}/\ell^n(i)$, and in fact with the étale sheafification of $K_{2i}(-; \mathbb{Z}/\ell^n)$ by rigidity results from [Sus83, Gab92].

In characteristic p , to describe the p -adic part of motivic cohomology, and thus the p -adic part of algebraic K -theory, one needs to replace μ_{ℓ^n} (which is zero on smooth varieties when $\ell = p$) by the logarithmic de Rham–Witt sheaf $W_n \Omega_{-, \log}^i$, see [GL00]. This description of p -adic algebraic K -theory in terms of the logarithmic de Rham–Witt sheaf is generalised in [KM21] to all Cartier smooth \mathbb{F}_p -algebras.

In mixed-characteristic, the ℓ -adic part of algebraic K -theory can often be computed by Gabber’s rigidity theorem [Gab92], but much less is known about the p -adic part. The syntomic complexes $\mathbb{Z}/p^n(i)^{\text{syn}}$, introduced in [BMS19, Section 7.4] and generalised in [BL22, Section 8], provide a well-behaved theory of p -adic étale motivic cohomology for arbitrary schemes. On p -quasisyntomic rings S (a nonnoetherian generalisation of local complete intersection rings) the complex $\mathbb{Z}/p^n(i)^{\text{syn}}(S)$ is concentrated in degrees $[0; i + 1]$ by [AMMN22, Theorem 5.1]. The top degree $i + 1$ of $\mathbb{Z}/p^n(i)^{\text{syn}}$ can be computed in terms of the logarithmic de Rham–Witt forms by [AMMN22, Corollary 5.43], and vanishes étale locally. We provide a description in smaller degrees of $\mathbb{Z}_p(i)^{\text{syn}}(S)$ in terms of the étale

cohomology of the generic fibre of S , when S is p -Cartier smooth over a perfectoid ring. Note that Bhatt and Mathew introduce in [BM22] the notion of F -smoothness in terms of absolute prismatic cohomology, and prove the following result for p -torsionfree F -smooth rings. Over a perfectoid base ring, F -smoothness and p -Cartier smoothness coincide by Theorem 2.18.

Theorem B (Syntomic-étale comparison theorem, see Theorems 4.12 and 4.14). *Let R be a perfectoid ring, and S a p -complete p -Cartier smooth R -algebra. Then for any integers $i \geq 0$ and $n \geq 1$, the map*

$$\mathbb{Z}/p^n(i)^{\text{syn}}(S) \longrightarrow R\Gamma_{\text{ét}}(\text{Spec}(S[\frac{1}{p}]), \mu_{p^n}^{\otimes i})$$

is an isomorphism on cohomology in degrees $< i - 1$, and is injective on H^{i-1} . If the perfectoid ring R is p -torsionfree, this map is an isomorphism on cohomology in degrees $< i$, and is injective on H^i .

The proof relies on the comparison between étale cohomology and prismatic cohomology [BS22, Theorem 9.1], and a property for p -Cartier smooth algebras of the Nygaard filtration on relative prismatic cohomology (Theorem C (\mathcal{N}^{\geq}) below). To prove this property, we extend various results on the relative prismatic cohomology of smooth algebras to general p -Cartier smooth algebras. It is a priori surprising that these properties are true assuming only a property on the special fibre (namely, the Cartier isomorphism); we prove that some of them even characterise p -Cartier smoothness.

Theorem C (see Theorem 2.16). *Let (A, I) be a bounded prism, and S a p -quasismooth A/I -algebra. The following are equivalent:*

(CSm) S is p -Cartier smooth over A/I .

($\mathbb{L}\Omega = \widehat{\mathbb{L}\Omega}$) *The canonical map $(\mathbb{L}\Omega_{S/(A/I)})_p^\wedge \rightarrow (\widehat{\mathbb{L}\Omega}_{S/(A/I)})_p^\wedge$ is an equivalence in the derived category $\mathcal{D}(A/I)$, where $(\widehat{\mathbb{L}\Omega}_{S/(A/I)})_p^\wedge$ is the Hodge-completion of the p -completed derived de Rham complex.*

($\mathbb{L}\Omega = \Omega$) *The counit map $(\mathbb{L}\Omega_{S/(A/I)})_p^\wedge \rightarrow (\Omega_{S/(A/I)})_p^\wedge$ is an equivalence in the derived category $\mathcal{D}(A/I)$, where $(\mathbb{L}\Omega_{S/(A/I)})_p^\wedge$ is the p -completed derived de Rham complex.*

(dR) *The de Rham comparison map $\Delta_{S/A}^{(1)} \otimes_A^{\mathbb{L}} A/I \rightarrow (\Omega_{S/(A/I)})_p^\wedge$ is an equivalence in the derived category $\mathcal{D}(A/I)$.*

($\Delta = \widehat{\Delta}$) *The canonical map $\text{can} : \Delta_{S/A}^{(1)} \rightarrow \widehat{\Delta}_{S/A}^{(1)}$ is an equivalence in the derived category $\mathcal{D}(A)$, where $\widehat{\Delta}_{S/A}^{(1)}$ is the Nygaard-completion of the prismatic complex.*

($L\eta$) *The Frobenius map $\tilde{\phi} : \Delta_{S/A}^{(1)} \rightarrow L\eta_I \Delta_{S/A}$ is an equivalence in the derived category $\mathcal{D}(A)$.*

(\mathcal{N}^{\geq}) *The Frobenius map $\tau^{\leq i} \phi : \tau^{\leq i} \mathcal{N}^{\geq i} \Delta_{S/A}^{(1)} \rightarrow \tau^{\leq i} I^i \Delta_{S/A}$ is an equivalence in the derived category $\mathcal{D}(A)$ for all $i \geq 0$.*

Outline.

In Section 2 we study the relative prismatic cohomology of p -Cartier smooth algebras and prove Theorem C (Theorem 2.16). We also provide a comparison between p -Cartier smoothness and F -smoothness over a perfectoid base (subsection 2.4). In Section 3 we prove Theorem A on the p -Cartier smoothness of certain valuation ring extensions (Theorem 3.1). In Section 4 we prove the syntomic-étale comparison Theorem B (Theorems 4.12 and 4.14), applying the results of Section 2 to the case of a perfectoid base.

Notation.

Given a commutative ring R , denote by $\mathcal{D}(R)$ the derived ∞ -category of R -modules, in which we use the cohomological convention for the indices. Also denote by $\mathcal{DF}(R) := \text{Fun}((\mathbb{Z}, \geq)^{\text{op}}, \mathcal{D}(R))$ the filtered derived ∞ -category of R -modules, where the filtrations are decreasing. Given a filtered complex $\text{Fil}^* C \in \mathcal{DF}(R)$ and for each integer $n \in \mathbb{Z}$, let $\text{gr}^n C \in \mathcal{D}(R)$ be the homotopy cofibre of the transition map $\text{Fil}^{n+1} C \rightarrow \text{Fil}^n C$. A filtered complex $\text{Fil}^* C \in \mathcal{DF}(R)$ is said to be complete if the limit $\lim_n \text{Fil}^n C$ vanishes in the derived ∞ -category $\mathcal{D}(R)$. For a nonzerodivisor d in R , the (derived) reduction C/d modulo d of a complex $C \in \mathcal{D}(R)$ is the homotopy cofibre of the map $d : C \rightarrow C$ induced by multiplication by d , *i.e.*, $C/d := C \otimes_R^{\mathbb{L}} R/d$. If C is given by an R -module M in degree 0, we also call derived reduction of M modulo d the complex $M \otimes_R^{\mathbb{L}} R/d$; it is given by M/d (the classical reduction of M modulo d) in degree 0 and by $M[d]$ (the d -torsion of M) in degree -1 . The (derived) d -adic completion of a complex $C \in \mathcal{D}(R)$ is the derived limit of the complexes C/d^n over $n \geq 1$. An R -module M is said to have bounded d^∞ -torsion if there exists an integer $n \geq 1$ such that $M[d^m] = M[d^n]$ for all $m \geq n$; this assumption guarantees that the derived d -adic completion of M is in degree 0, given by the classical d -adic completion of M . Given a complex $C \in \mathcal{D}(R)$ and some integers $a, b \in \mathbb{Z}$, there is a naturally associated complex $\tau^{[a;b]} C \in \mathcal{D}(R)$ which is concentrated in degrees $[a; b]$ and with the same cohomology groups as C in degrees $[a; b]$. A complex $C \in \mathcal{D}(R)$ is said to have Tor-amplitude in degrees $[a; b]$ if for any R -module N , the complex $C \otimes_R^{\mathbb{L}} N$ is in degrees $[a; b]$. Similarly, C is said to have d -complete Tor-amplitude in degrees $[a; b]$ if $C \otimes_R^{\mathbb{L}} R/d \in \mathcal{D}(R/d)$ has Tor-amplitude in degrees $[a; b]$. Given an ideal $I \subset R$, a complex $C \in \mathcal{D}(R)$ is I -adically complete if it is d -adically complete for every element $d \in I$.

Given a commutative ring R , an ∞ -category \mathcal{D} which admits sifted colimits (*e.g.*, $\mathcal{D}(R)$ or $\mathcal{DF}(R)$), and a functor

$$\mathcal{F} : \text{Sm}_R := \{\text{smooth } R\text{-algebras}\} \longrightarrow \mathcal{D},$$

define

$$\begin{aligned} \mathbb{L}\mathcal{F} : R\text{-Alg} &\longrightarrow \mathcal{D} \\ S &\longmapsto \text{colim}_{P \rightarrow S} \mathcal{F}(P), \end{aligned}$$

where the colimit is taken over all free R -algebras P mapping to S . The functor $\mathbb{L}\mathcal{F}$ is called the left Kan extension (from free R -algebras) of \mathcal{F} . For instance, the cotangent complex $\mathbb{L}_{-/R} := \mathbb{L}\Omega_{-/R}^1$ is the left Kan extension of the module of Kähler differentials $\Omega_{-/R}^1$, and the derived de Rham complex $\mathbb{L}\Omega_{-/R}$ is the left Kan extension of the de Rham complex $\Omega_{-/R}$. The Hodge filtration on the de Rham complex is given for each $n \in \mathbb{Z}$ by $\text{Fil}_{\text{Hod}}^n \Omega_{-/R} := \Omega_{-/R}^{\geq n}$; the Hodge filtration on the derived de Rham complex is defined as the left Kan extension of this filtration, and is not always complete.

Acknowledgments.

I am very grateful to Matthew Morrow for suggesting this project to me, sharing many insights and for careful readings of this manuscript. I would also like to thank Elden Elmanto and Arnab Kundu for helpful discussions, and Javier Fresán, Akhil Mathew and Mohamed Moakher for comments on a draft of this paper. This project has received funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (grant agreement No. 101001474).

2 PRISMATIC COHOMOLOGY OF CARTIER SMOOTH ALGEBRAS

Let p be a prime number. In this section we introduce the notion of p -Cartier smoothness for morphisms of commutative rings (Definition 2.5), and give several characterisations in terms of prismatic cohomology (Theorem 2.16).

The p -Cartier smooth morphisms generalise smooth morphisms, and behave as if they were smooth from the point of view of p -adic étale motivic cohomology. More precisely, the main interest of this article is in syntomic cohomology (see Section 4), which provides a general theory of p -adic étale motivic cohomology. Syntomic cohomology can be defined as Frobenius eigenspaces of prismatic cohomology, and this section is devoted to the prismatic cohomology of p -Cartier smooth algebras. Our main result

is Theorem 2.16. We also give a comparison with the notion of F -smoothness introduced in [BM22] (see subsection 2.4).

2.1 Definitions

For a commutative ring R , the cotangent complex

$$\mathbb{L}_{-/R} : R\text{-Alg} \longrightarrow \mathcal{D}(R)$$

is the natural derived version of the module of Kähler differentials

$$\Omega_{-/R}^1 : R\text{-Alg} \longrightarrow R\text{-Mod.}$$

It controls information on the relative prismatic complex via the Hodge–Tate comparison theorem [BS22, Theorem 4.11]. The following definition axiomatises some properties satisfied by the cotangent complex in the smooth case.

Definition 2.1 (Quasismoothness, [BMS19]). *(1) A morphism $R \rightarrow S$ of commutative rings is quasismooth¹, or S is a quasismooth R -algebra, if it is flat and the cotangent complex $\mathbb{L}_{S/R}$ has Tor-amplitude in $[0; 0]$, i.e., $\Omega_{S/R}^1$ is a flat S -module and $H_n(\mathbb{L}_{S/R}) = 0$ for all $n > 0$.*

(2) A morphism $R \rightarrow S$ of commutative rings is p -quasismooth, or S is a p -quasismooth R -algebra, if R has bounded p^∞ -torsion (i.e., the p -primary torsion is killed by a fixed power of p), the natural map $S \otimes_{\mathbb{L}_R} R/p \rightarrow S/p$ is an equivalence and the reduction $R/p \rightarrow S/p$ modulo p is quasismooth.

Remark 2.2. A morphism $R \rightarrow S$ of commutative rings is p -quasismooth if and only if its p -adic completion $R_p^\wedge \rightarrow S_p^\wedge$ is p -quasismooth. Moreover, if $R \rightarrow S$ is p -quasismooth, then S has bounded p^∞ -torsion by [BMS19, Corollary 4.8 (1)].

The condition that $S \otimes_{\mathbb{L}_R} R/p \in \mathcal{D}(R/p)$ is in degree 0 and base change for the cotangent complex imply that the morphism $\mathbb{L}_{S/R} \otimes_{\mathbb{L}_R} R/p \rightarrow \mathbb{L}_{(S/p)/(R/p)}$ is an equivalence. This allows one to lift properties of the cotangent complex of $R/p \rightarrow S/p$ to similar properties on the p -completed cotangent complex of $R \rightarrow S$.

Definition 2.3. *A morphism $R \rightarrow S$ of commutative rings is p -discrete if the derived tensor product $S \otimes_{\mathbb{L}_R} R/p \in \mathcal{D}(R/p)$ is concentrated in degree 0, where it is given by S/p .*

For instance, a morphism $R \rightarrow S$ of commutative rings is p -discrete if either R and S are p -torsionfree rings or S is a flat R -algebra. A morphism $R \rightarrow S$ of commutative rings is called p -flat (resp. p -faithfully flat) if it is p -discrete and its reduction $R/p \rightarrow S/p$ is flat (resp. faithfully flat)².

Examples 2.4. (1) A smooth morphism of commutative rings is quasismooth, because its cotangent complex is in degree 0, where it is given by the locally free module of differential forms.

(2) A p -flat morphism of perfectoid rings is p -quasismooth ([BMS19, Proposition 4.19 (3)] and [BMS18, Lemma 3.14]).

(3) A filtered colimit of (p -)quasismooth algebras is (p -)quasismooth, because the cotangent complex commutes with filtered colimits.

(4) Let C^+ be a valuation ring with perfect fraction field, and E^+ a valuation ring extension of C^+ (see section 3). The morphism $C^+ \rightarrow E^+$ has cotangent complex concentrated in degree 0 ([GR03, Theorem 6.5.8 (ii)]). If C^+ contains a perfect field, then the morphism $C^+ \rightarrow E^+$ is quasismooth ([GR03, Corollary 6.5.21]). If the p -adic completion of C^+ is a perfectoid ring, then the morphism $C^+ \rightarrow E^+$ is p -quasismooth (Proposition 3.8 below).

¹As in [BMS19], this should not be confused with other existing notions of quasismooth.

²More generally, a morphism $R \rightarrow S$ of commutative rings, possibly with bounded p^∞ -torsion, will be called p -something if it is p -discrete and if its reduction $R/p \rightarrow S/p$ modulo p is something. We distinguish this from similar notions on p -complete rings, often called p -completely something.

Several properties of prismatic cohomology in the smooth case –such as the comparison between prismatic cohomology and de Rham cohomology [BS22, Corollary 15.4]– do not hold for general p -quasismooth morphisms $R \rightarrow S$. We will prove (Theorem 2.16) that the obstruction to satisfy these properties vanishes exactly when the morphism $R/p \rightarrow S/p$ satisfies the Cartier isomorphism. For an \mathbb{F}_p -algebra R , we denote by ϕ_R its Frobenius endomorphism.

Definition 2.5 (Cartier smoothness). *(1) A morphism $R \rightarrow S$ of \mathbb{F}_p -algebras is Cartier smooth, or S is a Cartier smooth R -algebra, if it is quasismooth and if the inverse Cartier map*

$$C^{-1} : \Omega_{S/R}^n \otimes_{R, \phi_R} R \longrightarrow H^n(\Omega_{S/R}^\bullet)$$

is an isomorphism of R -modules for each $n \geq 0$.

(2) A morphism $R \rightarrow S$ of commutative rings is p -Cartier smooth, or S is a p -Cartier smooth R -algebra, if R has bounded p^∞ -torsion, it is p -discrete and its reduction $R/p \rightarrow S/p$ modulo p is Cartier smooth.

Equivalently, a morphism $R \rightarrow S$ of commutative rings is p -Cartier smooth if it is p -quasismooth and if the morphism $R/p \rightarrow S/p$ satisfies the Cartier isomorphism. For a morphism $R \rightarrow S$ of \mathbb{F}_p -algebras, recall that the inverse Cartier map

$$C^{-1} : \Omega_{S/R}^n \otimes_{R, \phi_R} R \longrightarrow H^n(\Omega_{S/R}^\bullet),$$

for any integer $n \geq 0$, is defined as the unique R -linear map satisfying

$$C^{-1}(fdg_1 \wedge \cdots \wedge dg_n \otimes 1) = f^p g_1^{p-1} \cdots g_n^{p-1} dg_1 \wedge \cdots \wedge dg_n.$$

Cartier smoothness when $R = \mathbb{F}_p$ was already studied in [KM21].

Examples 2.6. A morphism $R \rightarrow S$ of commutative rings is p -Cartier smooth in the following cases.

- (1) The morphism $R \rightarrow S$ is smooth, and R has bounded p^∞ -torsion. Then the cotangent complex $\mathbb{L}_{S/R}$ is a flat S -module in degree 0 and S is a flat R -algebra, so the morphism $R \rightarrow S$ is p -discrete and the morphism $R/p \rightarrow S/p$ is quasismooth. The Cartier isomorphism is a standard property of smooth morphisms in characteristic p (see for instance [DI87, Theorem 1.2]).
- (2) S is a filtered colimit of smooth R -algebras, and R has bounded p^∞ -torsion. Filtered colimits of p -quasismooth algebras are p -quasismooth, and the inverse Cartier map commutes with filtered colimits.
- (3) $R = \mathbb{F}_p$ and S is a valuation ring. The quasismoothness is a result of Gabber–Ramero (Example 2.4(4)). The Cartier isomorphism is a result of Gabber ([KST21, Corollary A.4]).
- (4) $R \rightarrow S$ is a p -flat morphism of perfectoid rings. Any perfectoid ring has bounded p^∞ -torsion ([BMS19, Proposition 4.19(3)]). The cotangent complex of the morphism $R/p \rightarrow S/p$ is zero ([BMS18, Lemma 3.14]), so the Cartier isomorphism is trivial for $n \geq 1$. If R and S are perfect \mathbb{F}_p -algebras and $n = 0$ the Cartier isomorphism holds; in general the inverse Cartier map for $n = 0$ is the reduction modulo p^b of the inverse Cartier map associated to the morphism $R^b \rightarrow S^b$ of perfect \mathbb{F}_p -algebras, and is thus an isomorphism.
- (5) R is a valuation ring whose p -adic completion is a perfectoid ring, and S is a valuation ring extension of R (Theorem 3.1 below).

Example 2.7. An imperfect \mathbb{F}_p -algebra S whose cotangent complex $\mathbb{L}_{S/\mathbb{F}_p}$ is trivial, is quasismooth but not Cartier smooth. Indeed, it is quasismooth because any algebra over the field \mathbb{F}_p is flat and because the zero complex of S -modules has Tor-amplitude in $[0; 0]$. In degree 0, the inverse Cartier map for the map $\mathbb{F}_p \rightarrow S$ is the Frobenius map $S \rightarrow S$, and is thus not an isomorphism; so the \mathbb{F}_p -algebra S is not Cartier smooth. Remark that imperfect \mathbb{F}_p -algebras with trivial cotangent complex exist, by [Bha13].

Lemma 2.8 (Base change for Cartier smoothness). *Let $R \rightarrow S$ be a p -Cartier smooth morphism, and R' an R -algebra with bounded p^∞ -torsion. Then the morphism $R' \rightarrow S \otimes_R R'$ is p -Cartier smooth.*

Proof. The p -flatness is preserved by base change. The Cartier isomorphism depends only on the reduction modulo p of the morphism $R' \rightarrow S \otimes_R R'$, so we can assume that R, R' and S are \mathbb{F}_p -algebras. Let $n \geq 0$ be an integer. There is a canonical isomorphism of R' -modules

$$\Omega_{(S \otimes_R R')/R'}^n \cong \Omega_{S/R}^n \otimes_R R'.$$

By Cartier smoothness of the morphism $R \rightarrow S$, the R -module $H^n(\Omega_{S/R}^\bullet)$ is flat, so there is a canonical isomorphism of R' -modules

$$H^n(\Omega_{S/R}^\bullet) \otimes_R R' \cong H^n(\Omega_{S/R}^\bullet \otimes_R^{\mathbb{L}} R').$$

Moreover, $\Omega_{S/R}^\bullet$ is a complex of flat R -modules by quasismoothness of the morphism $R \rightarrow S$, so there is a canonical isomorphism of R' -modules

$$H^n(\Omega_{S/R}^\bullet \otimes_R^{\mathbb{L}} R') \cong H^n(\Omega_{(S \otimes_R R')/R'}^\bullet)$$

and the morphism $R' \rightarrow S \otimes_R R'$ satisfies the Cartier isomorphism. \square

Lemma 2.9 (Transitivity of Cartier smoothness). *Let $R \rightarrow S$ and $S \rightarrow T$ be p -Cartier smooth morphisms. Then the composite $R \rightarrow T$ is also p -Cartier smooth.*

Proof. The p -quasismoothness and the Cartier isomorphism for $n = 1$ follow from the transitivity sequence for the cotangent complex. The Cartier isomorphism follows in general from the transitivity property of the derived de Rham complex ([Bha12, Proposition 3.22]). \square

Corollary 2.10. *Let $R \rightarrow S$ and $R \rightarrow S'$ be p -Cartier smooth morphisms. Then the morphism $R \rightarrow S \otimes_R S'$ is also p -Cartier smooth.*

Proof. The morphism $S \rightarrow S \otimes_R S'$ is p -Cartier smooth by Lemma 2.8. The morphism $R \rightarrow S \otimes_R S'$ is then p -Cartier smooth, as the composite of two p -Cartier smooth morphisms (Lemma 2.9). \square

2.2 Review of relative prismatic cohomology

Prisms are defined in [BS22] as pairs (A, I) , where A is a δ -ring (roughly³, a $\mathbb{Z}_{(p)}$ -algebra with a lift of Frobenius $\phi_A : A \rightarrow A$) and $I \subset A$ is an ideal defining a Cartier divisor in $\text{Spec}(A)$, such that A is derived (p, I) -complete and $p \in I + \phi_A(I)A$. In all the cases of interest the ideal $I \subset A$ will be a principal ideal generated by a nonzerodivisor. A prism (A, I) is called bounded if the ring A/I has bounded p^∞ -torsion. Restricting to bounded prisms avoids complications involving derived completions: for instance, the underlying ring A of a bounded prism (A, I) is (p, I) -complete in the classical sense.

Following [BMS19, Section 4] or [BL22, Appendix C], a morphism $R \rightarrow S$ of commutative rings is p -quasisyntomic if R has bounded p^∞ -torsion, $R \rightarrow S$ is p -flat and $\mathbb{L}_{(S/p)/(R/p)}$ has Tor-amplitude in $[-1; 0]$. For instance, any noetherian local complete intersection ring S is p -quasisyntomic over \mathbb{Z} by [Avr99, Theorem 1.2]. Any p -quasismooth, and in particular any p -Cartier smooth morphism, is p -quasisyntomic.

Given a bounded prism (A, I) and a p -quasisyntomic⁴ A/I -algebra S , [BS22, Section 4.1] defines the prismatic site $(S/A)_\Delta$ as the site having as objects the prisms (B, IB) over (A, I) with a map $S \rightarrow B/IB$, and covers given by flat covers. The relative prismatic complex $\Delta_{S/A} \in \mathcal{D}(A)$ is defined as the cohomology of the sheaf

$$\mathcal{O}_\Delta : (S/A)_\Delta \longrightarrow A\text{-Alg}, (B, IB) \longmapsto B.$$

³More precisely, a δ -ring is a $\mathbb{Z}_{(p)}$ -algebra A equipped with a map $\delta_A : A \rightarrow A$ such that $\phi_A : A \rightarrow A, x \mapsto x^p + p\delta_A(x)$ is a ring homomorphism. If A is a p -torsionfree $\mathbb{Z}_{(p)}$ -algebra, a δ -ring structure on A is the same as a ring endomorphism lifting the Frobenius on A/p . If A is a general $\mathbb{Z}_{(p)}$ -algebra, a δ -ring structure on A is the same as a lift of Frobenius in the derived sense ([BS22, Remark 2.5]).

⁴For a general A/I -algebra S , prismatic cohomology is defined by left Kan extension from the smooth case. The compatibility between the two definitions in the p -quasisyntomic case is proved in [BL22, Section 4].

Similarly, the Hodge–Tate complex $\overline{\Delta}_{S/A} \in \mathcal{D}(A/I)$ is defined as the cohomology of the sheaf

$$\overline{\mathcal{O}}_{\Delta} : (S/A)_{\Delta} \longrightarrow A/I\text{-Alg}, (B, IB) \longmapsto B/IB.$$

For instance, if k is a perfect field of characteristic p and S is a (p) -quasisyntomic k -algebra, then $(W(k), (p))$ is an example of bounded prism and $\Delta_{S/W(k)}$ recovers the crystalline cohomology of S ([BL22, Section 4.6]).

The prismatic complex $\Delta_{S/A}$ naturally bears a ϕ_A -semilinear Frobenius endomorphism

$$\phi : \Delta_{S/A}^{(1)} \longrightarrow \Delta_{S/A},$$

where $\Delta_{S/A}^{(1)} := \Delta_{S/A} \widehat{\otimes}_{A, \phi_A}^{\mathbb{L}} A$ is the Frobenius-twisted prismatic complex. Our main result on the prismatic cohomology of p -Cartier smooth algebras describes the image of this Frobenius endomorphism ϕ , and is formulated in terms of the functor $L\eta_I$. Following [BMS18, Section 6] and for A a commutative ring, $I \subset A$ an ideal defining a Cartier divisor in $\text{Spec}(A)$ and $C \in \mathcal{D}(A)$ a complex, the object $L\eta_I C \in \mathcal{D}(A)$ is defined as follows. The complex C in the derived category $\mathcal{D}(A)$ is represented by a complex (C^{\bullet}, d) of A -modules such that for all $i \in \mathbb{Z}$, C^i is I -torsionfree (i.e., $C^i[I] = 0$). Define the complex $\eta_I C^{\bullet}$ with terms

$$\eta_I C^i := \{x \in I^i C^i \mid dx \in I^{i+1} C^{i+1}\}$$

and differential induced by the differential of C^{\bullet} . As an object of the derived category $\mathcal{D}(A)$ this construction does not depend on the choice of (C^{\bullet}, d) ([BMS18, Corollary 6.5]), and we call this object $L\eta_I C \in \mathcal{D}(A)$. Following [BMS19, Section 5], an object $\text{Fil}^* C$ of the filtered derived category $\mathcal{DF}(A)$ is called *connective for the Beilinson t -structure* if the graded pieces $\text{gr}^i C \in \mathcal{D}(A)$ are in degree $\leq i$ for all $i \in \mathbb{Z}$. The *connective cover for the Beilinson t -structure* of a filtered complex $\text{Fil}^* C$ is the universal connective filtered complex with a map to $\text{Fil}^* C$.

Proposition 2.11. *Let A be a commutative ring, $I \subset A$ an ideal defining a Cartier divisor in $\text{Spec}(A)$ and C an object in the derived category $\mathcal{D}(A)$.*

(1) ([BMS18, Lemma 6.4]) *For each integer $i \in \mathbb{Z}$, there is a canonical isomorphism of A -modules*

$$H^i(L\eta_I C) \cong (H^i(C)/H^i(C)[I]) \otimes_A I^i.$$

(2) ([BMS19, Proposition 5.8]) *$L\eta_I C \in \mathcal{D}(A)$ is the complex underlying the connective cover for the Beilinson t -structure of the I -adically filtered object $I^* C \in \mathcal{DF}(A)$.*

The upshot is that the image of the Frobenius $\phi : \Delta_{S/A}^{(1)} \rightarrow \Delta_{S/A}$ will be identified with $L\eta_I \Delta_{S/A}$, when S is a p -Cartier smooth A/I -algebra. By the previous proposition, the complex $L\eta_I \Delta_{S/A}$ is characterised by a universal property in terms of the I -adic filtration on the prismatic complex $\Delta_{S/A}$. Following [BS22, Section 15], define the Nygaard filtration $\mathcal{N}^* \Delta_{S/A}^{(1)}$ on the Frobenius-twisted prismatic complex $\Delta_{S/A}^{(1)}$ as the preimage⁵ of the I -adic filtration $I^* \Delta_{S/A}$ via the morphism ϕ . The Frobenius

$$\phi : \Delta_{S/A}^{(1)} \longrightarrow \Delta_{S/A}$$

naturally upgrades to a map of filtered complexes

$$\phi : \mathcal{N}^{\geq *} \Delta_{S/A}^{(1)} \longrightarrow I^* \Delta_{S/A} \tag{1}$$

in the filtered derived category $\mathcal{DF}(A)$. Beware that the filtration $\mathcal{N}^{\geq *} \Delta_{S/A}^{(1)}$ is in general not complete, and we denote by $\widehat{\Delta}_{S/A}^{(1)} \in \mathcal{D}(A)$ (resp. $\mathcal{N}^i \Delta_{S/A}^{(1)} \in \mathcal{D}(A/I)$, for $i \geq 0$) the Nygaard-completed prismatic

⁵To make sense of this preimage, one needs to restrict to large p -quasisyntomic A/I -algebras –for which the prismatic complex $\Delta_{S/A}$ is concentrated in degree 0–, and then generalise the definition using descent on the p -quasisyntomic site.

complex (resp. the Nygaard graded pieces). On the Nygaard graded pieces, the Frobenius (1) can be rewritten as a map

$$\phi : \mathcal{N}^i \Delta_{S/A}^{(1)} \longrightarrow \overline{\Delta}_{S/A}\{i\}$$

for each $i \geq 0$, where

$$\overline{\Delta}_{S/A}\{i\} := \overline{\Delta}_{S/A} \otimes_{A/I} (I/I^2)^{\otimes i}.$$

To describe the image of this Frobenius map, define the conjugate filtration $\mathrm{Fil}_*^{\mathrm{conj}} \overline{\Delta}_{S/A}\{i\}$ on the Hodge–Tate complex $\overline{\Delta}_{S/A}\{i\}$ as the left Kan extension from smooth A/I -algebras of the increasing Postnikov filtration $\tau^{\leq *}\overline{\Delta}_{-/A}\{i\}$.

Theorem 2.12 ([BS22, BL22]). *Let (A, I) be a bounded prism, and S a p -quasisyntomic A/I -algebra.*

(1) *For each $i \geq 0$, the Frobenius map $\phi : \mathcal{N}^i \Delta_{S/A}^{(1)} \rightarrow \overline{\Delta}_{S/A}\{i\}$ factors as*

$$\mathcal{N}^i \Delta_{S/A}^{(1)} \xrightarrow{\tilde{\phi}} \mathrm{Fil}_i^{\mathrm{conj}} \overline{\Delta}_{S/A}\{i\} \xrightarrow{\mathrm{can}} \overline{\Delta}_{S/A}\{i\},$$

and the map $\tilde{\phi} : \mathcal{N}^i \Delta_{S/A}^{(1)} \rightarrow \mathrm{Fil}_i^{\mathrm{conj}} \overline{\Delta}_{S/A}\{i\}$ is an equivalence. In particular, the complex $\mathcal{N}^i \Delta_{S/A}^{(1)} \in \mathcal{D}(A/I)$ is concentrated in degrees $\leq i$.

(2) *The Frobenius map $\phi : \Delta_{S/A}^{(1)} \rightarrow \Delta_{S/A}$ factors as*

$$\Delta_{S/A}^{(1)} \xrightarrow{\tilde{\phi}} L\eta_I \Delta_{S/A} \xrightarrow{\mathrm{can}} \Delta_{S/A}.$$

Proof. The first part of (1) is a special case of [BL22, Proposition 5.1.1 and Remark 5.1.3], where the result is formulated for a general animated commutative A/I -algebra S . All the objects are left Kan extended from the smooth case, where the result is also given by [BS22, Theorem 15.3]. The functor $\tau^{\leq i}\overline{\Delta}_{-/A}\{i\}$ is concentrated in degrees $\leq i$, so its left Kan extension (defined as a sifted colimit) is also concentrated in degrees $\leq i$.

(2) The filtered complex $\mathcal{N}^{\geq *}\Delta_{S/A}^{(1)} \in \mathcal{DF}(A)$ is connective for the Beilinson t -structure by (1). So the map $\phi : \mathcal{N}^{\geq *}\Delta_{S/A}^{(1)} \rightarrow I^*\Delta_{S/A}$ factors through the connective cover for the Beilinson t -structure of the target. The result follows from Proposition 2.11 (2), by looking at the underlying complexes. \square

We recall some important features of the relative prismatic complex in the smooth case, following [BS22].

Theorem 2.13 (Prismatic cohomology in the smooth case, [BS22]). *Let (A, I) be a bounded prism, and S a smooth A/I -algebra.*

(1) *($L\eta_I$ comparison) The Frobenius map*

$$\tilde{\phi} : \Delta_{S/A}^{(1)} \xrightarrow{\sim} L\eta_I \Delta_{S/A}$$

of Theorem 2.12 (2) is an equivalence in the derived category $\mathcal{D}(A)$.

(2) *(Hodge–Tate comparison) There is a canonical isomorphism*

$$(\Omega_{S/(A/I)}^*)^{\wedge}_p \xrightarrow{\cong} \mathrm{H}^*(\overline{\Delta}_{S/A}\{*\}) := \bigoplus_{i \geq 0} \mathrm{H}^i(\overline{\Delta}_{S/A}\{i\})$$

of differential graded A/I -algebras, where the differential on $\mathrm{H}^(\overline{\Delta}_{S/A}\{*\})$ is the Bockstein operator induced by the I -adic filtration on $\Delta_{S/A}$.*

(3) (*de Rham comparison*) *There is a canonical equivalence*

$$\mathbb{A}_{S/A}^{(1)} \otimes_A^{\mathbb{L}} A/I \xrightarrow{\sim} (\Omega_{S/(A/I)})_p^\wedge$$

in the derived category $\mathcal{D}(A/I)$.

Proof. By [BL22, Corollary 4.1.14], the prismatic cohomology of S is the same as the prismatic cohomology of the p -adic completion of S . (1) and (2) then respectively follow from [BS22, Theorem 15.3] and [BS22, Theorem 4.11]. Using [BMS18, Proposition 6.12], (3) is a consequence of (1) and (2). \square

2.3 Prismatic cohomology of Cartier smooth algebras

In this subsection we prove Theorem C (Theorem 2.16), characterising p -Cartier smooth algebras in terms of their prismatic cohomology.

Any p -Cartier smooth algebra is p -quasismooth by definition. We first extend some properties of smooth algebras to general p -quasismooth algebras. Given a morphism $R \rightarrow S$ of commutative rings, denote by $\widehat{\mathbb{L}}\Omega_{S/R}$ the Hodge-completed derived de Rham complex, and by $(\widehat{\mathbb{L}}\Omega_{S/R})_p^\wedge$ its p -adic completion.

Proposition 2.14. *Let (A, I) be a bounded prism, and S a p -quasismooth A/I -algebra.*

- (1) *The canonical map $(\widehat{\mathbb{L}}\Omega_{S/(A/I)})_p^\wedge \rightarrow (\Omega_{S/(A/I)})_p^\wedge$ is an equivalence in the derived category $\mathcal{D}(A/I)$.*
- (2) *The conjugate filtration $\mathrm{Fil}_*^{\mathrm{conj}} \overline{\Delta}_{S/A}$ on the Hodge–Tate complex $\overline{\Delta}_{S/A} \in \mathcal{D}(A/I)$ coincides with the Postnikov filtration $\tau^{\leq *}_* \overline{\Delta}_{S/A}$. In particular for each $i \geq 0$, the Frobenius map*

$$\phi : \mathcal{N}^i \mathbb{A}_{S/A}^{(1)} \longrightarrow \overline{\Delta}_{S/A}\{i\}$$

induces an equivalence

$$\tilde{\phi} : \mathcal{N}^i \mathbb{A}_{S/A}^{(1)} \xrightarrow{\sim} \tau^{\leq i} \overline{\Delta}_{S/A}\{i\}$$

in the derived category $\mathcal{D}(A/I)$.

- (3) *There is a canonical isomorphism*

$$(\Omega_{S/(A/I)}^*)_p^\wedge \xrightarrow{\cong} \mathrm{H}^*(\overline{\Delta}_{S/A}\{*\})$$

of differential graded A/I -algebras, where the differential on $\mathrm{H}^(\overline{\Delta}_{S/A}\{*\})$ is the Bockstein operator induced by the I -adic filtration on $\Delta_{S/A}$.*

- (4) *The Frobenius map $\tilde{\phi} : \mathbb{A}_{S/A}^{(1)} \rightarrow L\eta_I \mathbb{A}_{S/A}$ of Theorem 2.12(2) factors as*

$$\mathbb{A}_{S/A}^{(1)} \xrightarrow{\mathrm{can}} \widehat{\mathbb{A}}_{S/A}^{(1)} \xrightarrow{\tilde{\phi}} L\eta_I \mathbb{A}_{S/A},$$

and the map $\tilde{\phi} : \widehat{\mathbb{A}}_{S/A}^{(1)} \rightarrow L\eta_I \mathbb{A}_{S/A}$ is an equivalence in the derived category $\mathcal{D}(A)$.

Proof. (1) By the derived Nakayama lemma ([Sta19, 091N]), it suffices to prove the result after derived reduction modulo p . By base change for the Hodge-completed derived de Rham complex and definition of p -quasismoothness, this is equivalent to proving that the canonical map

$$\widehat{\mathbb{L}}\Omega_{(S/p)/(A/(p,I))} \longrightarrow \Omega_{(S/p)/(A/(p,I))}$$

is an equivalence in the derived category $\mathcal{D}(A/(p, I))$. Both sides are complete for the Hodge filtration, so it suffices to prove that this canonical map is an equivalence on the Hodge graded pieces. The corresponding map

$$\mathbb{L}_{(S/p)/(A/(p,I))}^n[-n] \longrightarrow \Omega_{(S/p)/(A/(p,I))}^n[-n]$$

is the shift of a wedge power of the counit map

$$\mathbb{L}_{(S/p)/(A/(p,I))} \longrightarrow \Omega_{(S/p)/(A/(p,I))}^1[0]$$

for all $n \geq 0$, which is an equivalence by p -quasismoothness of the morphism $A/I \rightarrow S$.

(2) By left Kan extension of the Hodge–Tate comparison Theorem 2.13 (2) (or [BL22, Remark 4.1.7]), there is a canonical equivalence

$$(\mathbb{L}_{S/(A/I)}^n)_p^\wedge[-n]\{-n\} \xrightarrow{\sim} \mathrm{gr}_n^{\mathrm{conj}} \overline{\Delta}_{S/A}$$

in the derived category $\mathcal{D}(A/I)$, for each $n \geq 0$. By the derived Nakayama lemma ([Sta19, 091N]) and p -quasismoothness of the morphism $A/I \rightarrow S$, the complex $(\mathbb{L}_{S/(A/I)}^n)_p^\wedge \in \mathcal{D}(A/I)$ is in degree 0 and the conjugate graded piece $\mathrm{gr}_n^{\mathrm{conj}} \overline{\Delta}_{S/A}$ is thus in degree n . By induction on $n \geq 0$ and using the long exact sequence in cohomology groups associated to the homotopy cofibre sequence

$$\mathrm{Fil}_{n-1}^{\mathrm{conj}} \overline{\Delta}_{S/A} \longrightarrow \mathrm{Fil}_n^{\mathrm{conj}} \overline{\Delta}_{S/A} \longrightarrow \mathrm{gr}_n^{\mathrm{conj}} \overline{\Delta}_{S/A},$$

we deduce that $\mathrm{Fil}_n^{\mathrm{conj}} \overline{\Delta}_{S/A}$ belongs to $\mathcal{D}^{[0;n]}(A/I)$, that the induced morphism

$$\mathrm{Fil}_{n-1}^{\mathrm{conj}} \overline{\Delta}_{S/A} \longrightarrow \tau^{\leq n-1} \mathrm{Fil}_n^{\mathrm{conj}} \overline{\Delta}_{S/A}$$

is an equivalence and that the morphism

$$\mathrm{H}^n(\mathrm{Fil}_n^{\mathrm{conj}} \overline{\Delta}_{S/A}) \longrightarrow \mathrm{H}^n(\mathrm{gr}_n^{\mathrm{conj}} \overline{\Delta}_{S/A})$$

is an isomorphism of A/I -modules. The conjugate filtration $\mathrm{Fil}_*^{\mathrm{conj}} \overline{\Delta}_{S/A}$ is moreover exhaustive, as a left Kan extension of the exhaustive Postnikov filtration $\tau^{\leq *}\overline{\Delta}_{S/A}$. So the canonical map

$$\mathrm{Fil}_n^{\mathrm{conj}} \overline{\Delta}_{S/A} \longrightarrow \tau^{\leq n} \lim_k (\mathrm{Fil}_{n+k}^{\mathrm{conj}} \overline{\Delta}_{S/A}) \simeq \tau^{\leq n} \overline{\Delta}_{S/A}$$

is an equivalence for each $n \geq 0$, and the conjugate filtration $\mathrm{Fil}_*^{\mathrm{conj}} \overline{\Delta}_{S/A}$ coincides with the Postnikov filtration $\tau^{\leq *}\overline{\Delta}_{S/A}$. The last statement is a consequence of Theorem 2.12 (1).

(3) By left Kan extension of the Hodge–Tate comparison Theorem 2.13 (2) (or [BL22, Remark 4.1.7]) and (2), there is for each $n \geq 0$ a canonical isomorphism

$$(\Omega_{S/(A/I)}^n)_p^\wedge \xrightarrow{\cong} \mathrm{H}^n(\overline{\Delta}_{S/A}\{n\})$$

of A/I -modules. To prove that the de Rham differential d coincides via these isomorphisms with the Bockstein operator β on $\mathrm{H}^*(\overline{\Delta}_{S/A}\{*\})$, it suffices to prove it when S is a polynomial A/I -algebra, where this is Theorem 2.13 (2).

(4) For each $k \geq 0$, the map of filtered complexes $\phi : \mathcal{N}^{\geq *}\Delta_{S/A}^{(1)} \rightarrow I^*\Delta_{S/A}$ induces a map of filtered complexes

$$\phi : \mathcal{N}^{\geq *}\Delta_{S/A}^{(1)} / \mathcal{N}^{\geq *+k}\Delta_{S/A}^{(1)} \longrightarrow I^*\Delta_{S/A} / I^{*+k}\Delta_{S/A} \simeq \Delta_{S/A} \otimes_A^{\mathbb{L}} I^* / I^{*+k}.$$

Taking the inverse limit over $k \geq 0$ defines a map of filtered complexes

$$\phi : \mathcal{N}^{\geq *}\widehat{\Delta}_{S/A}^{(1)} \longrightarrow I^*\Delta_{S/A}.$$

The canonical map $\mathcal{N}^{\geq *}\Delta_{S/A}^{(1)} \rightarrow \mathcal{N}^{\geq *}\widehat{\Delta}_{S/A}^{(1)}$ is an equivalence on graded pieces, so the same argument as in Theorem 2.12 (2) proves that

$$\Delta_{S/A}^{(1)} \xrightarrow{\mathrm{can}} \widehat{\Delta}_{S/A}^{(1)} \xrightarrow{\phi} \Delta_{S/A}$$

factors as

$$\Delta_{S/A}^{(1)} \xrightarrow{\mathrm{can}} \widehat{\Delta}_{S/A}^{(1)} \xrightarrow{\tilde{\phi}} L\eta_I \Delta_{S/A} \xrightarrow{\mathrm{can}} \Delta_{S/A},$$

where the composite $\tilde{\phi} : \Delta_{S/A}^{(1)} \rightarrow L\eta_I \Delta_{S/A}$ of the first two maps is the map of Theorem 2.12 (2). Remark that the I -adic filtration on $L\eta_I \Delta_{S/A}$ is complete by [BL22, Remark D.10]. To prove that $\tilde{\phi} : \widehat{\Delta}_{S/A}^{(1)} \rightarrow L\eta_I \Delta_{S/A}$ is an equivalence, it suffices by completeness to prove that it is an equivalence on graded pieces. Proposition 2.11 (2) (more precisely, [BMS19, Theorem 5.4 (2)]) identifies the i^{th} graded piece of $L\eta_I \Delta_{S/A}$ with the truncation $\tau^{\leq i}$ of the I -adic graded piece $\overline{\Delta}_{S/A}\{i\}$. Together with the identification $\mathcal{N}^i \Delta_{S/A}^{(1)} \xrightarrow{\sim} \mathcal{N}^i \widehat{\Delta}_{S/A}^{(1)}$, it then suffices to prove that

$$\tilde{\phi} : \mathcal{N}^i \Delta_{S/A}^{(1)} \longrightarrow \tau^{\leq i} \overline{\Delta}_{S/A}\{i\}$$

is an equivalence for all $i \geq 0$, which is (2). \square

Remark 2.15. The factorisation part of Proposition 2.14 (4) and its proof hold more generally for (A, I) a bounded prism and S a p -quasisyntomic A/I -algebra.

Theorem 2.16. *Let (A, I) be a bounded prism, and S a p -quasismooth A/I -algebra. The following are equivalent:*

(CSm) *The inverse Cartier map*

$$\Omega_{(S/p)/(A/(p,I))}^n \otimes_{A/(p,I), \phi_{A/(p,I)}} A/(p, I) \xrightarrow{C^{-1}} \mathbf{H}^n(\Omega_{(S/p)/(A/(p,I))}^\bullet)$$

is an isomorphism of $A/(p, I)$ -modules for all $n \geq 0$, i.e., S is p -Cartier smooth over A/I .

($\mathbb{L}\Omega = \widehat{\mathbb{L}\Omega}$) *The Hodge-completion map $(\mathbb{L}\Omega_{S/(A/I)})_p^\wedge \rightarrow (\widehat{\mathbb{L}\Omega}_{S/(A/I)})_p^\wedge$ is an equivalence in the derived category $\mathcal{D}(A/I)$.*

($\mathbb{L}\Omega = \Omega$) *The counit map $(\mathbb{L}\Omega_{S/(A/I)})_p^\wedge \rightarrow (\Omega_{S/(A/I)})_p^\wedge$ is an equivalence in the derived category $\mathcal{D}(A/I)$.*

(dR) *The de Rham comparison map $\Delta_{S/A}^{(1)} \otimes_A^{\mathbb{L}} A/I \rightarrow (\Omega_{S/(A/I)})_p^\wedge$ is an equivalence in the derived category $\mathcal{D}(A/I)$.*

($\overline{\Delta}$) *The canonical map $\Delta_{S/A}^{(1)} \otimes_A^{\mathbb{L}} A/I \rightarrow \mathbf{H}^*(\overline{\Delta}_{S/A}\{*\})$ is an equivalence in the derived category $\mathcal{D}(A/I)$.*

($\Delta = \widehat{\Delta}$) *The Nygaard-completion map $\Delta_{S/A}^{(1)} \rightarrow \widehat{\Delta}_{S/A}^{(1)}$ is an equivalence in the derived category $\mathcal{D}(A)$.*

($L\eta$) *The Frobenius map $\tilde{\phi} : \Delta_{S/A}^{(1)} \rightarrow L\eta_I \Delta_{S/A}$ is an equivalence in the derived category $\mathcal{D}(A)$.*

(\mathcal{N}) *The canonical map $\mathbf{H}_B^0 : \Delta_{S/A}^{(1)} \otimes_A^{\mathbb{L}} A/I \rightarrow \mathbf{H}^*(\mathcal{N}^* \Delta_{S/A}^{(1)})$ is an equivalence in the derived category $\mathcal{D}(A/I)$.*

(\mathcal{N}^{\geq}) *The Frobenius map $\tau^{\leq i} \phi : \tau^{\leq i} \mathcal{N}^{\geq i} \Delta_{S/A}^{(1)} \rightarrow \tau^{\leq i} I^i \Delta_{S/A}$ is an equivalence in the derived category $\mathcal{D}(A)$ for all $i \geq 0$.*

Proof. (CSm) \Leftrightarrow ($\mathbb{L}\Omega = \Omega$) By derived Nakayama ([Sta19, 091N]), the counit map

$$(\mathbb{L}\Omega_{S/(A/I)})_p^\wedge \longrightarrow (\Omega_{S/(A/I)})_p^\wedge$$

is an equivalence in the derived category $\mathcal{D}(A/I)$ if and only if the counit map

$$\mathbb{L}\Omega_{(S/p)/(A/(p,I))} \longrightarrow \Omega_{(S/p)/(A/(p,I))}$$

is an equivalence in the derived category $\mathcal{D}(A/(p, I))$, i.e., if the induced map

$$\mathbf{H}^n(\mathbb{L}\Omega_{(S/p)/(A/(p,I))}) \longrightarrow \mathbf{H}^n(\Omega_{(S/p)/(A/(p,I))})$$

is an isomorphism of $A/(p, I)$ -modules for each $n \geq 0$. By [Bha12, Proposition 3.5], the derived de Rham complex $\mathbb{L}\Omega_{(S/p)/(A/(p, I))}$ is equipped with an exhaustive \mathbb{N} -indexed increasing filtration $\mathrm{Fil}_\star^{\mathrm{conj}} \mathbb{L}\Omega_{(S/p)/(A/(p, I))}$, whose graded pieces are given by

$$C^{-1} : \wedge^n (\mathbb{L}_{(S/p)/(A/(p, I))} \otimes_{A/(p, I), \phi_{A/(p, I)}}^{\mathbb{L}} A/(p, I))[-n] \xrightarrow{\sim} \mathrm{gr}_n^{\mathrm{conj}} \mathbb{L}\Omega_{(S/p)/(A/(p, I))},$$

where C^{-1} is the left Kan extension of the inverse Cartier map. By quasismoothness of the morphism $A/(p, I) \rightarrow S/p$, the graded piece $\mathrm{gr}_n^{\mathrm{conj}} \mathbb{L}\Omega_{(S/p)/(A/(p, I))}$ is thus concentrated in degree n for each $n \geq 0$. Arguing by induction on $n \geq 0$ and by exhaustiveness of the filtration $\mathrm{Fil}_\star^{\mathrm{conj}} \mathbb{L}\Omega_{(S/p)/(A/(p, I))}$, there is a canonical isomorphism

$$\mathrm{H}^n(\mathrm{gr}_n^{\mathrm{conj}} \mathbb{L}\Omega_{(S/p)/(A/(p, I))}) \xrightarrow{\cong} \mathrm{H}^n(\mathbb{L}\Omega_{(S/p)/(A/(p, I))}).$$

In particular the counit map $\mathbb{L}\Omega_{(S/p)/(A/(p, I))} \rightarrow \Omega_{(S/p)/(A/(p, I))}$ is an equivalence in the derived category $\mathcal{D}(A/(p, I))$ if and only if the inverse Cartier map

$$C^{-1} : \Omega_{(S/p)/(A/(p, I))}^n \otimes_{A/(p, I), \phi_{A/(p, I)}} A/(p, I) \longrightarrow \mathrm{H}^n(\Omega_{(S/p)/(A/(p, I))}^\bullet)$$

is an isomorphism of $A/(p, I)$ -modules for all $n \geq 0$.

$(\mathbb{L}\Omega = \widehat{\mathbb{L}}\Omega) \Leftrightarrow (\mathbb{L}\Omega = \Omega) \Leftrightarrow (dR) \Leftrightarrow (\overline{\Delta}) \Leftrightarrow (\Delta = \widehat{\Delta}) \Leftrightarrow (L\eta)$ By [BL22, Proposition 5.2.3] there is a commutative diagram

$$\begin{array}{ccccc}
 & & \widehat{\Delta}_{S/A}^{(1)} & & \\
 & \text{can} \nearrow & \downarrow & \searrow \tilde{\phi} & \\
 \Delta_{S/A}^{(1)} & & & & L\eta_I \Delta_{S/A} \\
 & \searrow \tilde{\phi} & & & \downarrow \\
 & & (\widehat{\mathbb{L}}\Omega_{S/(A/I)})_p^\wedge & & \\
 & \text{can} \nearrow & \downarrow & \searrow \text{can} & \\
 (\mathbb{L}\Omega_{S/(A/I)})_p^\wedge & & & & (\Omega_{S/(A/I)})_p^\wedge \\
 & \searrow & & \searrow \text{HT} & \\
 & & & & \mathrm{H}^*(\overline{\Delta}_{S/A}\{*\})
 \end{array}$$

in the derived category $\mathcal{D}(A)$, where the dashed arrows are derived reduction modulo I ([BL22, Proposition 5.2.5 and Corollary 5.2.8] and [BMS18, Proposition 6.12]). The map $(\mathbb{L}\Omega_{S/(A/I)})_p^\wedge \rightarrow \mathrm{H}^*(\overline{\Delta}_{S/A}\{*\})$ is defined as the composite

$$(\mathbb{L}\Omega_{S/(A/I)})_p^\wedge \longrightarrow (\Omega_{S/(A/I)})_p^\wedge \longrightarrow \mathrm{H}^*(\overline{\Delta}_{S/A}\{*\}).$$

The three equivalences in this diagram hold for any p -quasismooth A/I -algebra S by Proposition 2.14. By derived Nakayama ([Sta19, 091N]) and commutativity of this diagram, the conditions $(\mathbb{L}\Omega = \widehat{\mathbb{L}}\Omega)$, $(\mathbb{L}\Omega = \Omega)$, (dR) , $(\overline{\Delta})$, $(\Delta = \widehat{\Delta})$ and $(L\eta)$ are then equivalent.

$(L\eta) \Leftrightarrow (\mathcal{N})$ There is a commutative diagram

$$\begin{array}{ccc}
 \Delta_{S/A}^{(1)} \otimes_A^{\mathbb{L}} A/I & \xrightarrow{\mathrm{H}_B^0} & \mathrm{H}^*(\mathcal{N}^* \Delta_{S/A}^{(1)}) \\
 \downarrow \tilde{\phi} & & \sim \downarrow \tilde{\phi} \\
 (L\eta_I \Delta_{S/A}) \otimes_A^{\mathbb{L}} A/I & \xrightarrow[\sim]{\mathrm{H}_B^0} & \mathrm{H}^*(\overline{\Delta}_{S/A}\{*\})
 \end{array}$$

in the derived category $\mathcal{D}(A/I)$, where the maps $\tilde{\phi}$ are defined in Theorem 2.12, and the functor H_B^0 (where B refers to the Beilinson t -structure) is defined in [BMS19, Theorem 5.4]. More precisely, the functor H_B^0 sends the filtered complex

$$\mathcal{N}^{\geq *}\Delta_{S/A}^{(1)} \in \mathcal{DF}(A)$$

to

$$H^*(\mathcal{N}^*\Delta_{S/A}^{(1)}) \in \mathcal{D}(A) \simeq \mathcal{DF}(A)^\heartsuit,$$

where $\mathcal{DF}(A)^\heartsuit$ is the heart of the filtered derived category $\mathcal{DF}(A)$ for its Beilinson t -structure. Because $\mathcal{N}^*\Delta_{S/A}^{(1)}$ is naturally an object of the derived category $\mathcal{D}(A/I)$, this induces a map

$$\Delta_{S/A}^{(1)} \otimes_A^{\mathbb{L}} A/I \xrightarrow{H_B^0} H^*(\mathcal{N}^*\Delta_{S/A}^{(1)})$$

on the underlying complexes. For any p -quasismooth A/I -algebra S , the right and bottom maps of the previous diagram are equivalences (Proposition 2.14 (2) and [BMS18, Proposition 6.12]). By derived Nakayama and commutativity of the diagram, the Frobenius map

$$\tilde{\phi} : \Delta_{S/A}^{(1)} \longrightarrow L\eta_I \Delta_{S/A}$$

is an equivalence in the derived category $\mathcal{D}(A)$ if and only if the canonical map

$$H_B^0 : \Delta_{S/A}^{(1)} \otimes_A^{\mathbb{L}} A/I \longrightarrow H^*(\mathcal{N}^*\Delta_{S/A}^{(1)})$$

is an equivalence in the derived category $\mathcal{D}(A/I)$.

(\mathcal{N}^{\geq}) \Rightarrow ($L\eta$) Assume that the Frobenius map

$$\tau^{\leq i} \phi : \tau^{\leq i} \mathcal{N}^{\geq i} \Delta_{S/A}^{(1)} \longrightarrow \tau^{\leq i} I^i \Delta_{S/A}$$

is an equivalence for each $i \geq 0$, and in particular that the Frobenius map

$$\phi : H^i(\mathcal{N}^{\geq i} \Delta_{S/A}^{(1)}) \longrightarrow H^i(I^i \Delta_{S/A})$$

is an isomorphism of A -modules for each $i \geq 0$. The homotopy cofibre sequence

$$\mathcal{N}^{\geq i} \Delta_{S/A}^{(1)} \longrightarrow \mathcal{N}^{\geq i-1} \Delta_{S/A}^{(1)} \longrightarrow \mathcal{N}^{i-1} \Delta_{S/A}^{(1)}$$

induces an exact sequence

$$H^{n-1}(\mathcal{N}^{i-1} \Delta_{S/A}^{(1)}) \longrightarrow H^n(\mathcal{N}^{\geq i} \Delta_{S/A}^{(1)}) \longrightarrow H^n(\mathcal{N}^{\geq i-1} \Delta_{S/A}^{(1)}) \longrightarrow H^n(\mathcal{N}^{i-1} \Delta_{S/A}^{(1)})$$

for each integer n . By Proposition 2.14 (2), the graded piece $\mathcal{N}^{i-1} \Delta_{S/A}^{(1)}$ is in degrees $[0; i-1]$ and the Frobenius map $\phi : \mathcal{N}^{i-1} \Delta_{S/A}^{(1)} \rightarrow \overline{\Delta}_{S/A}\{i-1\}$ is an isomorphism in degrees $\leq i-1$. In particular the morphism $H^n(\mathcal{N}^{\geq i} \Delta_{S/A}^{(1)}) \rightarrow H^n(\mathcal{N}^{\geq i-1} \Delta_{S/A}^{(1)})$ is an isomorphism for all $i, n \geq 0$ satisfying $i < n$, and the canonical morphism

$$H^i(\mathcal{N}^{\geq i-1} \Delta_{S/A}^{(1)}) \longrightarrow H^i(\mathcal{N}^{\geq 0} \Delta_{S/A}^{(1)}) = H^i(\Delta_{S/A}^{(1)})$$

is an isomorphism for all $i \geq 0$. The A -module $H^i(L\eta_I \Delta_{S/A})$ is canonically identified with the image of the morphism $H^i(I^i \Delta_{S/A}) \rightarrow H^i(I^{i-1} \Delta_{S/A})$ by Proposition 2.11 (1). The Frobenius map

$$\phi : \mathcal{N}^{\geq *} \Delta_{S/A}^{(1)} \longrightarrow I^* \Delta_{S/A}$$

thus induces a map of short exact sequences

$$\begin{array}{ccccccc}
 \mathrm{H}^{i-1}(\mathcal{N}^{\geq i-1}\Delta_{S/A}^{(1)}) & \longrightarrow & \mathrm{H}^{i-1}(\mathcal{N}^{i-1}\Delta_{S/A}^{(1)}) & \longrightarrow & \mathrm{H}^i(\mathcal{N}^{\geq i}\Delta_{S/A}^{(1)}) & \longrightarrow & \mathrm{H}^i(\Delta_{S/A}^{(1)}) \longrightarrow 0 \\
 \downarrow \phi & & \downarrow \phi & & \downarrow \phi & & \downarrow \tilde{\phi} \\
 \mathrm{H}^{i-1}(I^{i-1}\Delta_{S/A}) & \longrightarrow & \mathrm{H}^{i-1}(\overline{\Delta}_{S/A}\{i-1\}) & \longrightarrow & \mathrm{H}^i(I^i\Delta_{S/A}) & \longrightarrow & \mathrm{H}^i(L\eta_I\Delta_{S/A}) \rightarrow 0.
 \end{array}$$

The first three vertical maps are isomorphisms by assumption, thus so is the right vertical map, for each $i \geq 0$. So the Frobenius map $\tilde{\phi} : \Delta_{S/A}^{(1)} \rightarrow L\eta_I\Delta_{S/A}$ is an equivalence.

$(L\eta) \Rightarrow (\mathcal{N}^{\geq})$ Assume the Frobenius map $\tilde{\phi} : \Delta_{S/A}^{(1)} \rightarrow L\eta_I\Delta_{S/A}$ is an equivalence. We prove by induction on $i \geq 0$ that for every integer $n \leq i$, the Frobenius morphism

$$\phi : \mathrm{H}^n(\mathcal{N}^{\geq i}\Delta_{S/A}^{(1)}) \longrightarrow \mathrm{H}^n(I^i\Delta_{S/A})$$

is an isomorphism of A -modules. For $i = 0$, it suffices to prove that the Frobenius morphism

$$\phi : \mathrm{H}^0(\Delta_{S/A}^{(1)}) \longrightarrow \mathrm{H}^0(\Delta_{S/A})$$

is an isomorphism, or equivalently that the canonical map

$$\mathrm{H}^0(L\eta_I\Delta_{S/A}) \cong \mathrm{H}^0(\Delta_{S/A})/\mathrm{H}^0(\Delta_{S/A})[I] \longrightarrow \mathrm{H}^0(\Delta_{S/A})$$

is an isomorphism. For any prism (B, IB) over (A, I) , the A -algebra B is I -torsionfree [BS22, Lemma 3.5], so

$$\mathrm{H}^0(\Delta_{S/A}) = \lim_{(B, IB) \in (S/A)_{\Delta}} B$$

is also I -torsionfree and the morphism $\mathrm{H}^0(L\eta_I\Delta_{S/A}) \rightarrow \mathrm{H}^0(\Delta_{S/A})$ is the identity. Assume i is now a nonnegative integer for which the result holds. Using the equivalence

$$\tilde{\phi} : \Delta_{S/A}^{(1)} \xrightarrow{\sim} L\eta_I\Delta_{S/A},$$

the first, second and fourth vertical maps of the previous diagram are isomorphisms, so the morphism

$$\phi : \mathrm{H}^i(\mathcal{N}^{\geq i}\Delta_{S/A}^{(1)}) \longrightarrow \mathrm{H}^i(I^i\Delta_{S/A})$$

is also an isomorphism. For $n < i$, the Frobenius $\phi : \mathcal{N}^{\geq *}\Delta_{S/A}^{(1)} \rightarrow I^*\Delta_{S/A}$ induces a map of exact sequences

$$\begin{array}{ccccccccc}
 \mathrm{H}^{n-1}(\mathcal{N}^{\geq i}\Delta_{S/A}^{(1)}) & \rightarrow & \mathrm{H}^{n-1}(\mathcal{N}^i\Delta_{S/A}^{(1)}) & \rightarrow & \mathrm{H}^n(\mathcal{N}^{\geq i+1}\Delta_{S/A}^{(1)}) & \rightarrow & \mathrm{H}^n(\mathcal{N}^{\geq i}\Delta_{S/A}^{(1)}) & \rightarrow & \mathrm{H}^n(\mathcal{N}^i\Delta_{S/A}^{(1)}) \\
 \cong \downarrow \phi & & \cong \downarrow \phi & & \downarrow \phi & & \cong \downarrow \phi & & \cong \downarrow \phi \\
 \mathrm{H}^{n-1}(I^{i-1}\Delta_{S/A}) & \rightarrow & \mathrm{H}^{n-1}(\overline{\Delta}_{S/A}\{i\}) & \rightarrow & \mathrm{H}^n(I^{i+1}\Delta_{S/A}) & \rightarrow & \mathrm{H}^n(I^i\Delta_{S/A}) & \rightarrow & \mathrm{H}^n(\overline{\Delta}_{S/A}\{i\})
 \end{array}$$

where the isomorphisms are given by Proposition 2.14 (2) and the induction hypothesis. It follows that the morphism $\phi : \mathrm{H}^n(\mathcal{N}^{\geq i}\Delta_{S/A}^{(1)}) \rightarrow \mathrm{H}^n(I^i\Delta_{S/A})$ is also an isomorphism, which concludes the proof. \square

2.4 Comparison with F -smoothness

In this subsection we compare the absolute notion of F -smoothness, introduced by Bhatt–Mathew [BM22], with our relative notion of p -Cartier smoothness, in the case when the base is perfectoid (Theorem 2.18). F -smoothness is a variant of (p -adic) smoothness designed to capture smoothness in an absolute sense. For instance, regular rings are F -smooth ([BM22, Theorem 4.15]).

Absolute prismatic cohomology can be defined using an absolute version of the relative prismatic site introduced in subsection 2.2. We recall only the necessary notation, and refer the reader to [BL22] for the general theory. Following [BMS19, Section 4] or [BL22, Appendix C], a commutative ring S is p -quasisyntomic if S has bounded p^∞ -torsion and $\mathbb{L}_{(S/p)/\mathbb{F}_p} \in \mathcal{D}(S/p)$ has Tor-amplitude in $[-1; 0]$. Following [BL22, Sections 4 and 5] and for any p -quasisyntomic ring S , the absolute prismatic site $(S)_\Delta$ is the site having as objects the prisms (B, J) with a map $S \rightarrow B/J$ and covers given by flat covers. The absolute prismatic complex $\Delta_S \in \mathcal{D}(\mathbb{Z}_p)$ is the cohomology of the sheaf

$$\mathcal{O}_\Delta : (S)_\Delta \longrightarrow A\text{-Alg}, (B, J) \longmapsto B.$$

It is equipped with a Nygaard filtration $\mathcal{N}^{\geq *}\Delta_S$ and a map of filtered complexes

$$\phi : \mathcal{N}^{\geq *}\Delta_S \longrightarrow \Delta_S^{[*]},$$

where the filtered complex $\Delta_S^{[*]}$ is an absolute version of the I -adic filtration on relative prismatic cohomology. This Frobenius map is compatible with the Frobenius map on relative prismatic cohomology when S is defined over a base prism.

Definition 2.17 (F -smoothness, [BM22]). *A p -quasisyntomic ring S is F -smooth if for each integer $i \geq 0$, the Nygaard filtration on $\Delta_S\{i\}$ is complete and the homotopy cofibre $\text{hocofib}(\phi)$ of the Frobenius map*

$$\phi : \mathcal{N}^i\Delta_S \longrightarrow \overline{\Delta}_S\{i\},$$

where $\overline{\Delta}_S\{i\} := \text{gr}^i\Delta_S^{[*]}$, has p -complete Tor-amplitude in degrees $\geq i + 1$.

Following [BS22, Section 3], a bounded prism (A, I) is *perfect* if its Frobenius ϕ_A is an isomorphism. The functor $(A, I) \mapsto A/I$ induces an equivalence between the category of perfect prisms and the category of perfectoid rings ([BS22, Theorem 3.10]). Given a perfect prism (A, I) and a p -quasisyntomic A/I -algebra S , the canonical maps

$$\Delta_S \longrightarrow \Delta_{S/A} \longrightarrow \Delta_{S/A}^{(1)}$$

are equivalences and their composite can be refined into an equivalence of filtered complexes ([BL22, Theorem 5.6.2])

$$\mathcal{N}^{\geq *}\Delta_S \xrightarrow{\sim} \mathcal{N}^{\geq *}\Delta_{S/A}^{(1)}.$$

Theorem 2.18. *Let (A, I) be a perfect prism, and S a p -quasisyntomic A/I -algebra. Then S is F -smooth if and only if S is p -Cartier smooth over A/I .*

Proof. Assume first that S is p -Cartier smooth over A/I . The ring S is p -quasisyntomic ([BMS19, Lemma 4.15 and Proposition 4.19]). Let $i \geq 0$ be an integer. The Frobenius map

$$\phi : \mathcal{N}^i\Delta_S \longrightarrow \overline{\Delta}_S\{i\}$$

is naturally identified with the Frobenius map

$$\phi : \mathcal{N}^i\Delta_{S/A}^{(1)} \longrightarrow \overline{\Delta}_{S/A}\{i\},$$

whose homotopy cofibre is $\tau^{\geq i+1}\overline{\Delta}_{S/A}\{i\}$ by Theorem 2.14 (2). For all $n \geq i + 1$, the cohomology groups $H^n(\tau^{\geq i+1}\overline{\Delta}_{S/A}\{i\})$ are canonically isomorphic to $(\Omega_{S/(A/I)}^n)_p^\wedge$ for all $n \geq i + 1$ by the Hodge–Tate comparison (Proposition 2.14 (3)). By p -quasismoothness of the A/I -algebra S the S -modules $(\Omega_{S/(A/I)}^n)_p^\wedge$ are p -completely flat for all n , hence the complex $\tau^{\geq i+1}\overline{\Delta}_{S/A}$ has p -complete Tor-amplitude in degrees $\geq i + 1$. Moreover, the Nygaard filtration on $\overline{\Delta}_S\{i\}$ is canonically identified with the filtration $(\mathcal{N}^{\geq *}\Delta_{S/A}^{(1)}) \otimes_A^L I^i/I^{i+1}$, which is complete by Theorem 2.16 ($\Delta = \widehat{\Delta}$).

Assume now that S is F -smooth. In particular the Frobenius map

$$\phi : \mathcal{N}^1\Delta_{S/A}^{(1)} \longrightarrow \overline{\Delta}_{S/A}\{1\}$$

has homotopy cofibre in degrees ≥ 2 , and $\mathcal{N}^1 \Delta_{S/A}^{(1)}$ is in degrees ≤ 1 (Theorem 2.12 (1)), so there is an equivalence

$$\tilde{\phi} : \mathcal{N}^1 \Delta_{S/A}^{(1)} \longrightarrow \tau^{\leq 1} \overline{\Delta}_{S/A} \{1\}.$$

The Frobenius factors through the equivalence

$$\tilde{\phi} : \mathcal{N}^1 \Delta_{S/A}^{(1)} \xrightarrow{\sim} \mathrm{Fil}_1^{\mathrm{conj}} \overline{\Delta}_{S/A} \{1\}$$

of Theorem 2.12 (1), so the canonical map

$$\mathrm{Fil}_1^{\mathrm{conj}} \overline{\Delta}_{S/A} \{1\} \xrightarrow{\sim} \tau^{\leq 1} \overline{\Delta}_{S/A} \{1\}$$

is an equivalence. Moreover, $\mathrm{gr}_0^{\mathrm{conj}} \overline{\Delta}_{S/A} \{1\}$ is naturally identified with $H^0(\overline{\Delta}_{S/A} \{1\})$ by the Hodge–Tate comparison. So, using the Hodge–Tate comparison $\mathrm{gr}_1^{\mathrm{conj}} \overline{\Delta}_{S/A} \{1\} \simeq (\mathbb{L}_{S/(A/I)})_p^\wedge[-1]$, the cotangent complex $(\mathbb{L}_{S/(A/I)})_p^\wedge$ is in degree 0 and is naturally identified with $H^1(\overline{\Delta}_{S/A} \{1\})$. Using the F -smoothness condition for $i = 0, 1$, the complexes $\tau^{\geq 1} \overline{\Delta}_{S/A} \{1\} := \tau^{\geq 1} \overline{\Delta}_{S/A} \otimes_{A/I} I/I^2$ and $\tau^{\geq 2} \overline{\Delta}_{S/A} \{1\}$ have p -complete Tor-amplitude in degrees ≥ 1 and ≥ 2 respectively. So $H^1(\overline{\Delta}_{S/A} \{1\})[-1]$, as the homotopy cofibre of the morphism

$$\tau^{\geq 1} \overline{\Delta}_{S/A} \{1\} \longrightarrow \tau^{\geq 2} \overline{\Delta}_{S/A} \{1\},$$

has p -complete Tor-amplitude in degrees ≥ 1 . In particular the cotangent complex $(\mathbb{L}_{S/(A/I)})_p^\wedge$ has p -complete Tor-amplitude in degrees ≥ 0 , or equivalently it is a p -completely flat S -module in degree 0. So the morphism $A/I \rightarrow S$ is p -quasismooth. The Nygaard filtration on $\overline{\Delta}_S$ (which is complete by assumption) is naturally identified with the filtration $(\mathcal{N}^{\geq *}\Delta_{S/A}^{(1)}) \otimes_A^{\mathbb{L}} A/I$. In particular the Nygaard filtration on $\Delta_{S/A}^{(1)}$ is complete by derived Nakayama. So S is p -Cartier smooth over A/I by Theorem 2.16 ($\Delta = \widehat{\Delta} \Rightarrow (CSm)$). \square

Remark 2.19. In general, given a p -quasisyntomic morphism $R \rightarrow S$ of commutative rings where R has bounded p^∞ -torsion, the notions of F -smoothness for S and p -Cartier smoothness for $R \rightarrow S$ do not agree. For instance, the ring $\mathbb{Z}_p[p^{1/p}]$ is regular noetherian and is thus F -smooth ([BM22, Theorem 4.15]), but the morphism $\mathbb{Z}_p \rightarrow \mathbb{Z}_p[p^{1/p}]$ is not p -Cartier smooth (it is not p -quasismooth). On the other hand, the identity endomorphism of R is always p -Cartier smooth, but not all rings R with bounded p^∞ -torsion are F -smooth: any F -smooth p -complete noetherian ring is regular ([BM22, Theorem 4.15]). If S is a p -torsionfree p -Cartier smooth \mathbb{Z}_p -algebra, then S is F -smooth ([BM22, Corollary 4.17]).

Remark 2.20. Let (A, I) be a bounded prism, and S a p -quasisyntomic A/I -algebra. Then S is p -Cartier smooth over A/I if and only if it satisfies the following relative version of F -smoothness: for each integer $i \geq 0$ (or equivalently $i \in \{0, 1\}$), the homotopy cofibre of the Frobenius map $\phi : \mathcal{N}^i \Delta_{S/A}^{(1)} \rightarrow \overline{\Delta}_{S/A} \{i\}$ has p -complete Tor-amplitude in degrees $\geq i + 1$ and the Nygaard filtration on $\Delta_{S/A}^{(1)}$ is complete.

3 VALUATION RINGS ARE CARTIER SMOOTH

Recall that a valuation ring is an integral domain C^+ such that for any elements f and g in C^+ , either $f \in gC^+$ or $g \in fC^+$. The valuation on the fraction field C of C^+ is defined as the canonical group homomorphism $v : C^\times \rightarrow C^\times / (C^+)^\times =: \Gamma_C$, where Γ_C is naturally equipped with a structure of totally ordered abelian group, and is called the value group of C . The value group Γ_C (resp. the valuation ring C^+) is said to be discrete if it is isomorphic to the ring of integers \mathbb{Z} . The rank of a valuation ring is defined as its number of nonzero prime ideals. In particular, a valuation ring C^+ has rank at most 1 if and only if its value group Γ_C can be embedded in the ordered group of real numbers \mathbb{R} . A valuation ring extension E^+ of C^+ is a valuation ring E^+ equipped with a flat

ring morphism $C^+ \rightarrow E^+$. The flat modules over a valuation ring C^+ are exactly the torsionfree C^+ -modules, so a morphism $C^+ \rightarrow E^+$ of valuation rings is flat if and only if it is injective.

Valuation rings arise in various contexts in p -adic geometry, *e.g.*, in [BS17, Hub96, Sch12, BM21]. The goal of this section is to prove the following result.

Theorem 3.1. *Let C^+ be a valuation ring whose p -adic completion is a perfectoid ring. Let E^+ be a valuation ring extension of C^+ . Then the morphism $C^+ \rightarrow E^+$ is p -Cartier smooth.*

Corollary 3.2. *Let C^+ be a p -torsionfree valuation ring whose p -adic completion is a perfectoid ring, and E^+ be a valuation ring extension of C^+ . Then for any integers $i \geq 0$ and $n \geq 1$, the map*

$$\mathbb{Z}/p^n(i)^{\text{syn}}(E^+) \longrightarrow R\Gamma_{\text{ét}}(\text{Spec}(E^+[\frac{1}{p}]), \mu_{p^n}^{\otimes i})$$

is an isomorphism on cohomology in degrees $< i$, and is injective on H^i .

Proof. This is a direct consequence of Theorems 3.1 and 4.14. □

Examples of valuation rings C^+ satisfying the hypothesis of Theorem 3.1 include the ring of integers $\overline{\mathbb{Z}}_p$ of an algebraic closure $\overline{\mathbb{Q}}_p$ of \mathbb{Q}_p , the ring of integers $\mathcal{O}_{\mathbb{C}_p}$ of the p -adic complex numbers \mathbb{C}_p and the ring $\mathbb{Z}_p[p^\infty]$ (Example 3.6 below).

Remark 3.3. An extension of valuation rings $C^+ \rightarrow E^+$ is p -Cartier smooth if and only if the morphism $C^+/p \rightarrow E^+/p$ is Cartier smooth, *i.e.*, if $C^+/p \rightarrow E^+/p$ satisfies the Cartier isomorphism and the cotangent complex $\mathbb{L}_{(E^+/p)/(C^+/p)}$ is a flat (E^+/p) -module in degree 0. Indeed, any valuation ring C^+ has bounded p^∞ -torsion: either p is zero in C^+ , in which case $C^+[p^\infty] = C^+[p] = C^+$; or p is nonzero in C^+ , in which case $C^+[p^\infty] = C^+[p] = 0$. Any extension $C^+ \rightarrow E^+$ of valuation rings is flat by definition, and thus p -discrete.

We will distinguish three cases. If p is invertible in the valuation ring C^+ , then the rings C^+/p and E^+/p are zero⁶ and the result is trivial. If p is zero in the valuation ring C^+ , the result is essentially due to Gabber–Ramero and Gabber, as we recall now. We will then focus on the remaining case, *i.e.*, that of mixed-characteristic.

Assume that $p = 0$ in the valuation ring C^+ . By [BMS18, Example 3.15] an \mathbb{F}_p -algebra, or equivalently its p -adic completion, is a perfectoid ring if and only if it is perfect (*i.e.*, its Frobenius endomorphism is an isomorphism). Theorem 3.1 is then due to Gabber–Ramero and Gabber. Remark that Cartier smoothness in characteristic p , including its relation to valuation rings and algebraic K -theory, was previously studied in [KM21].

Theorem 3.4 (Cartier smoothness of characteristic p valuation rings, [GR03, KST21]). *Let C^+ be a perfect valuation ring of characteristic p , and E^+ a valuation ring extension of C^+ . Then the morphism $C^+ \rightarrow E^+$ is Cartier smooth. Equivalently:*

- (1) *The cotangent complex \mathbb{L}_{E^+/C^+} is concentrated in degree 0, and Ω_{E^+/C^+}^1 is a flat E^+ -module.*
- (2) *The inverse Cartier map*

$$C^{-1} : \Omega_{E^+/C^+}^n \otimes_{C^+, \phi_{C^+}} C^+ \longrightarrow H^n(\Omega_{E^+/C^+}^\bullet)$$

is an isomorphism of C^+ -modules for each $n \geq 0$.

Proof. (1) The cotangent complexes \mathbb{L}_{E^+/C^+} and $\mathbb{L}_{E^+/\mathbb{F}_p}$ are concentrated in degree 0 ([GR03, Theorem 6.5.8 (ii)]) where they are given by Ω_{E^+/C^+}^1 and $\Omega_{E^+/\mathbb{F}_p}^1$, and $\Omega_{E^+/\mathbb{F}_p}^1$ is a torsionfree E^+ -module ([GR03, Corollary 6.5.21]). There is a homotopy transitivity fibre sequence

$$\mathbb{L}_{C^+/\mathbb{F}_p} \otimes_{C^+}^{\mathbb{L}} E^+ \longrightarrow \mathbb{L}_{E^+/\mathbb{F}_p} \longrightarrow \mathbb{L}_{E^+/C^+},$$

⁶There is a slight ambiguity whether the zero ring is perfectoid or not. We include this case to avoid any ambiguity.

and $\mathbb{L}_{C^+/\mathbb{F}_p} \simeq 0$ because C^+ is a perfect \mathbb{F}_p -algebra. So the natural morphism

$$\Omega_{E^+/\mathbb{F}_p}^1 \longrightarrow \Omega_{E^+/C^+}^1$$

is an isomorphism of E^+ -modules. Torsionfree modules over a valuation ring are flat, so Ω_{E^+/C^+}^1 is a flat E^+ -module.

(2) The morphism $\mathbb{F}_p \rightarrow E^+$ satisfies the Cartier isomorphism ([KST21, Corollary A.4]). By the previous paragraph, there is a canonical isomorphism

$$\Omega_{E^+/\mathbb{F}_p}^1 \xrightarrow{\cong} \Omega_{E^+/C^+}^1$$

of E^+ -modules, so the morphism $C^+ \rightarrow E^+$ also satisfies the Cartier isomorphism. \square

Assume now, for the rest of this section, that p is neither invertible nor zero in the valuation ring C^+ ; we say in this case that C^+ is a *mixed-characteristic* valuation ring. The hypothesis on C^+ in Theorem 3.1 can be reformulated as follows.

Lemma 3.5. *Let C^+ be a mixed-characteristic valuation ring. The following are equivalent:*

- (1) *The p -adic completion of the valuation ring C^+ is a perfectoid ring.*
- (2) *The ring C^+/p has a nonzero nilpotent element, and its Frobenius endomorphism is surjective.*

Proof. The second condition depends only on the ring C^+/p , so we can assume the valuation ring C^+ is p -adically complete. Because p is nonzero in the valuation ring C^+ , the ring C^+ is in particular p -torsionfree.

By [BMS18, Lemmas 3.9 and 3.10], a p -torsionfree ring R is perfectoid if and only if R is p -adically complete, the Frobenius on R/p is surjective and $up \in R$ admits a compatible system of p -power roots for some unit $u \in R^\times$. From now on, let C^+ be a nonzero p -adically complete and p -torsionfree valuation ring, such that the Frobenius on C^+/p is surjective.

(1) \Rightarrow (2) Let $u \in (C^+)^\times$ be a unit such that up admits a compatible system $((up)^{1/p^n})_{n \in \mathbb{N}}$ of p -power roots in the valuation ring C^+ . Because C^+ is p -torsionfree and not the zero ring, p is nonzero in C^+ and neither is $(up)^{1/p}$. The element $(up)^{1/p} \in C^+$ thus defines a nonzero nilpotent element in the ring C^+/p .

(2) \Rightarrow (1) Assume the valuation ring C^+ has an element $\pi \in C^+$ defining a nonzero nilpotent element in C^+/p . Because the Frobenius on C^+/p is surjective, we can assume that π^p is nonzero in C^+/p . Because C^+ is a valuation ring, this implies that π^p divides p and that p divides some power of π in C^+ . In particular the valuation ring C^+ is π -adically complete for this element $\pi \in C^+$. By [BMS18, Lemma 3.9], there exists a unit $u \in (C^+)^\times$ such that $up \in C^+$ admits a compatible system of p -power roots in C^+ . \square

Example 3.6. The p -adic completion of the ring $\mathbb{Z}_p[p^{1/p^\infty}]$ is a perfectoid valuation ring, because it satisfies the hypotheses of Lemma 3.5 (2). In particular, any valuation ring extension E^+ of \mathbb{Z}_p containing a compatible system of p -power roots of p is p -Cartier smooth over $\mathbb{Z}_p[p^{1/p^\infty}]$ by Theorem 3.1. Beware that p -adic completion (or even taking the p -adically separated quotient) does not preserve the value group of a valuation ring. For instance, the localisation of the ring $\mathbb{Z}_p[X/p^n, n \geq 0]$ at the ideal $(X/p^n, n \geq 0) \subset \mathbb{Z}_p[X/p^n, n \geq 0]$ is a valuation ring, with fraction field $\mathbb{Q}_p(X)$ and with p -adic completion \mathbb{Z}_p ; an alternative description of this valuation ring is the fiber product $\mathbb{Z}_p \times_{\mathbb{Q}_p} \mathbb{Q}_p[X]_{(X)}$. Similarly, the localisation of the ring $\mathbb{Z}_p[p^{1/p^\infty}, X/p^n, n \geq 0]$ at the ideal $(X/p^n, n \geq 0) \subset \mathbb{Z}_p[p^{1/p^\infty}, X/p^n, n \geq 0]$ is a valuation ring, with fraction field $\mathbb{Q}_p(p^{1/p^\infty}, X)$ and p -adically separated quotient $\mathbb{Z}_p[p^{1/p^\infty}]$.

To prove Theorem 3.1 for the morphism $C^+ \rightarrow E^+$, it suffices to prove that the morphism $C^+/p \rightarrow E^+/p$ is quasismooth (*i.e.*, its cotangent complex $\mathbb{L}_{(E^+/p)/(C^+/p)}$ is given by a flat E^+/p -module in degree 0) and satisfies the Cartier isomorphism (Remark 3.3). We first prove the quasismoothness.

Lemma 3.7. *Let E^+ be a valuation ring in which p is nonzero, and M be an E^+ -module. If the E^+ -module M is p -torsionfree, then its derived reduction $M \otimes_{E^+}^{\mathbb{L}} E^+/p$ modulo p is a flat E^+/p -module in degree 0.*

Proof. Because p is a nonzerodivisor in the valuation ring E^+ , the derived reduction $M \otimes_{E^+}^{\mathbb{L}} E^+/p$ modulo p of any E^+ -module M is concentrated in degree 0 if and only if M is p -torsionfree.

Assuming M is a p -torsionfree E^+ -module, we prove now that M/p is a flat E^+/p -module. Consider the p -adically separated quotient $E^{+'}$ of E^+ , *i.e.*, the quotient of E^+ by its ideal $\bigcap_{n \geq 0} p^n E^+$. The ring $E^{+'}$ is a valuation ring, because it is an integral domain and satisfies the divisibility condition of valuation rings. Define M' as the $E^{+'}$ -module $M \otimes_{E^+} E^{+'}$. The morphism $M \rightarrow M'$ of E^+ -modules is surjective, with kernel given by a quotient of the E^+ -module $M \otimes_{E^+} \bigcap_{n \geq 0} p^n E^+$. The E^+ -module $M \otimes_{E^+} \bigcap_{n \geq 0} p^n E^+$ is p -divisible, so the morphism $M \rightarrow M'$ becomes an isomorphism after reduction modulo p . Because $E^{+'}$ is a p -adically separated valuation ring, any p -torsionfree $E^{+'}$ -module is torsionfree and thus flat. In particular the $E^{+'}$ -module M' is flat, and the $E^{+'}/p$ -module M'/p is also flat. So the E^+/p -module M/p is flat. \square

Proposition 3.8 (Quasismoothness of the morphism $C^+/p \rightarrow E^+/p$). *Let C^+ be a mixed-characteristic valuation ring whose p -adic completion is a perfectoid ring. Let E^+ be a valuation ring extension of C^+ . Then the E^+ -module Ω_{E^+/C^+}^1 is p -torsionfree and the cotangent complex $\mathbb{L}_{(E^+/p)/(C^+/p)}$ is a flat E^+/p -module in degree 0.*

Proof. Note the equivalence $\mathbb{L}_{(E^+/p)/(C^+/p)} \simeq \mathbb{L}_{E^+/C^+} \otimes_{E^+}^{\mathbb{L}} E^+/p$. The cotangent complex of any extension of valuation rings with characteristic 0 fraction fields is concentrated in degree 0 ([GR03, Theorem 6.5.8 (ii)]⁷). In particular the cotangent complex \mathbb{L}_{E^+/C^+} is concentrated in degree 0, given by the E^+ -module Ω_{E^+/C^+}^1 . If the E^+ -module Ω_{E^+/C^+}^1 is p -torsionfree, then the cotangent complex $\mathbb{L}_{(E^+/p)/(C^+/p)}$ is a flat E^+/p -module in degree 0 (Lemma 3.7).

We prove now that the E^+ -module Ω_{E^+/C^+}^1 is p -torsionfree. Let E be the fraction field of the valuation ring E^+ , and \overline{E} an algebraic closure of E . We fix a valuation on \overline{E} extending the valuation of the valued field E , and denote by \overline{E}^+ the corresponding ring of integers. Applying again [GR03, Theorem 6.5.8 (ii)] to C^+ , E^+ and \overline{E}^+ and because the morphism $E^+ \rightarrow \overline{E}^+$ is flat, the transitivity sequence for the cotangent complex can be rewritten as a short exact sequence of \overline{E}^+ -modules:

$$0 \longrightarrow \Omega_{E^+/C^+}^1 \otimes_{E^+} \overline{E}^+ \longrightarrow \Omega_{\overline{E}^+/C^+}^1 \longrightarrow \Omega_{\overline{E}^+/E^+}^1 \longrightarrow 0.$$

An E^+ -module (resp. \overline{E}^+ -module) M is p -torsionfree if and only if the multiplication by p morphism $p : M \rightarrow M$ is injective. The morphism $E^+ \rightarrow \overline{E}^+$ being faithfully flat, the E^+ -module Ω_{E^+/C^+}^1 is thus p -torsionfree if and only if the \overline{E}^+ -module $\Omega_{\overline{E}^+/C^+}^1 \otimes_{E^+} \overline{E}^+$ is p -torsionfree. By the previous short exact sequence, it then suffices to prove the \overline{E}^+ -module $\Omega_{\overline{E}^+/C^+}^1$ is p -torsionfree. The p -adic completions $(C^+)_p^\wedge$ and $(\overline{E}^+)_p^\wedge$ of C^+ and \overline{E}^+ are perfectoid rings, respectively by assumption and because the field \overline{E} is algebraically closed. In particular the cotangent complex $\mathbb{L}_{(\overline{E}^+)_p^\wedge/p/((C^+)_p^\wedge)/p}$ is zero ([BMS18, Lemma 3.14]). Equivalently, the cotangent complex $\mathbb{L}_{(\overline{E}^+/p)/(C^+/p)}$ is zero. In particular, the derived reduction $\Omega_{\overline{E}^+/C^+}^1 \otimes_{\overline{E}^+}^{\mathbb{L}} \overline{E}^+/p$ modulo p of the \overline{E}^+ -module $\Omega_{\overline{E}^+/C^+}^1$ is in degree 0, or equivalently the \overline{E}^+ -module $\Omega_{\overline{E}^+/C^+}^1$ is p -torsionfree. \square

It remains to prove the Cartier isomorphism for the reduction modulo p of valuation ring extensions $C^+ \rightarrow E^+$, where C^+ is a mixed-characteristic valuation ring whose p -adic completion is a perfectoid ring. The proof of the Cartier isomorphism for positive characteristic valuation rings [KST21, Corollary A.4] relies on subtle approximation results of Gabber, which do not immediately pass to mixed-characteristic. Our strategy of proof in mixed-characteristic is to reduce to this result in positive characteristic. We are immediately faced with the following issue: if C^+ is a mixed-characteristic

⁷More precisely, [GR03, Theorem 6.5.8 (ii)] is formulated for extensions of valued fields. An extension of valuation rings $C^+ \hookrightarrow E^+$ is the composition of the localisation $C^+ \hookrightarrow C_{\mathfrak{p}}^+$ at the prime ideal $\mathfrak{p} := C^+ \cap \mathfrak{m}_{E^+}$ and the morphism $C_{\mathfrak{p}}^+ \hookrightarrow E^+$. The cotangent complex of the first morphism is trivial and the second morphism is induced by an extension of valued fields.

valuation ring, the ring C^+/p is in general not an integral domain, and in particular not a valuation ring.

Here we use the perfectoid assumption on the base C^+ to remark that there is a perfect valuation ring C^{+b} of characteristic p (the tilt of C^+) whose reduction modulo some element $d \in C^{+b}$ is naturally isomorphic to C^+/p . The Cartier isomorphism is preserved by base change (Lemma 2.8). So it would suffice to find a valuation ring extension E^{+b} of C^{+b} whose reduction modulo d is E^+/p to prove the Cartier isomorphism for $C^+/p \rightarrow E^+/p$. To try to construct such a valuation ring E^{+b} over the d -adically complete lift C^{+b} of C^+/p , we use the deformation theory of Illusie [Ill71, III.2.1.2.3] (see also [Sch12, Theorem 5.11]). Namely, we will need the following result, where R is a ring, $I \subset R$ is an ideal such that $I^2 = 0$, and S_0 is a flat $R_0 := R/I$ -algebra.

Theorem 3.9 ([Ill71]). *There is an obstruction class $\omega(S_0)$ in the group $\mathrm{Ext}_{S_0}^2(\mathbb{L}_{S_0/R_0}, S_0 \otimes_{R_0} I)$ which vanishes precisely when there exists a flat R -algebra S such that $S \otimes_R R_0 = S_0$. If there exists such a deformation, then the set of all isomorphism classes of such deformations forms a torsor under $\mathrm{Ext}_{S_0}^1(\mathbb{L}_{S_0/R_0}, S_0 \otimes_{R_0} I)$, and every deformation has automorphism group $\mathrm{Hom}_{S_0}(\mathbb{L}_{S_0/R_0}, S_0 \otimes_{R_0} I)$.*

To apply this deformation result recursively up to a d -adic deformation E^{+b} of the flat C^+/p -algebra E^+/p , we need to have control on the cotangent complex $\mathbb{L}_{(E^+/p)/(C^+/p)}$. More precisely, if the cotangent complex $\mathbb{L}_{(E^+/p)/(C^+/p)}$ was a projective E^+/p -module in degree 0 then the higher Ext-groups in this deformation result would vanish and we could construct the lift E^b . By Proposition 3.8, the cotangent complex $\mathbb{L}_{(E^+/p)/(C^+/p)}$ is concentrated in degree 0, given by the flat E^+/p -module $\Omega_{(E^+/p)/(C^+/p)}^1$. The E^+/p -module $\Omega_{(E^+/p)/(C^+/p)}^1$ will not be projective in general, but only a filtered colimit of almost projective modules, as we explain now.

By Proposition 3.8, the E^+ -module Ω_{E^+/C^+}^1 is p -torsionfree; for simplicity, assume for a moment that E^+ is p -adically separated, so that Ω_{E^+/C^+}^1 is a torsionfree E^+ -module. In particular the morphism $\Omega_{E^+/C^+}^1 \rightarrow \Omega_{E^+/C}^1$ of E^+ -module is injective. The inverse Cartier map commutes with filtered colimits, so we can assume the field extension $C \rightarrow E$ induced by the valuation ring extension $C^+ \rightarrow E^+$ is of finite type to prove the Cartier isomorphism. In particular $\Omega_{E^+/C}^1$ is a finite dimensional E -vector space, and $\Omega_{(E^+/p)/(C^+/p)}^1$ is a finite type E^+/p -module. The latter is thus projective if and only if it is free. The upshot is that there exist E^+/p -modules constructed like this which are not free E^+/p -modules. For instance, when $E \cong C(X)$, it follows from the proof of [GR03, Proposition 6.5.6] that the E^+ -module Ω_{E^+/C^+}^1 can be isomorphic to $\mathfrak{m}_{C^+} E^+$, where \mathfrak{m}_{C^+} is the maximal ideal of C^+ .

Although the E^+/p -module $\Omega_{(E^+/p)/(C^+/p)}^1$ is not projective in general, we will prove it is in fact almost projective in the sense of [GR03]. More precisely, almost mathematics are defined over a pair (V, \mathfrak{m}) , where V is a ring and $\mathfrak{m} \subset V$ is an ideal satisfying $\mathfrak{m} = \mathfrak{m}^2$. A morphism of V -modules is said to be an almost isomorphism if its kernel and cokernel are killed by multiplication by every element of the ideal $\mathfrak{m} \subset V$. The next lemma proves almost mathematics make sense over the pair $(C^+, \sqrt{(p)})$, where $\sqrt{(p)}$ is the radical of the ideal $(p) \subset C^+$. We will prove the E^+/p -module $\Omega_{(E^+/p)/(C^+/p)}^1$ is almost projective with respect to this pair $(C^+, \sqrt{(p)})$. For instance, in the previous paragraph, the morphism $\mathfrak{m}_{C^+} E^+ \rightarrow E^+$ is an almost isomorphism, so $\mathfrak{m}_{C^+} E^+$ is almost free of rank 1. We refer the reader to [GR03, AGT16] for the foundations of almost homological algebra.

Lemma 3.10. *Let C^+ be a mixed-characteristic valuation ring whose p -adic completion is a perfectoid ring. Then the prime ideal $\sqrt{(p)} \subset C^+$ satisfies $\sqrt{(p)} = \sqrt{(p)^2}$.*

Proof. We want to prove any element x of the ideal $\sqrt{(p)}$ can be written as the product of two elements of the ideal $\sqrt{(p)}$. It suffices to prove that x is divisible by such a product. By Lemma 3.5 there exists $\pi \in \sqrt{(p)}$ such that p does not divide π in C^+ . By surjectivity of the Frobenius on C^+/p , we can even assume that π^p divides p . If p divides x , then $\pi \cdot \pi^{p-1}$ also divides x , and π and π^{p-1} are elements of the ideal $\sqrt{(p)}$. If p does not divide x , then by surjectivity of the Frobenius on C^+/p , there exist $y \in \sqrt{(p)}$ and $t \in C^+$ such that $y^p + pt = x$. Because p does not divide x , p can not divide y^p , so y^p divides p . In particular y^p divides x and we conclude similarly. \square

Notation 3.11 (Almost mathematics). *Let C^+ be a mixed-characteristic valuation ring whose p -adic completion is a perfectoid ring. Denote by*

$$C^{+\flat} := \lim_{x \rightarrow x^p} C^+ / p$$

the tilt of this perfectoid ring. The ring $C^{+\flat}$ is a perfect valuation ring of characteristic p , which is d -adically complete for some element $d \in C^{+\flat}$ such that there is a natural ring isomorphism

$$C^+ / p \cong C^{+\flat} / d.$$

We will use almost mathematics either with respect to the pair $(C^+, \sqrt{(p)})$ or with respect to the pair $(C^{+\flat}, \sqrt{(d)})$ (which satisfies the analogue of Lemma 3.10), depending on the context. The two are equivalent over $C^+ / p \cong C^{+\flat} / d$.

Lemma 3.12. *Let C^+ be a mixed-characteristic valuation ring whose p -adic completion is a perfectoid ring. Let E^+ be a valuation ring extension of C^+ , with fraction field E . If M is a torsionfree E^+ -module such that $M \otimes_{E^+} E$ is a finite dimensional E -vector space, then the E^+ / p -module M / p is almost free of rank at most $\dim_E(M \otimes_{E^+} E)$.*

Proof. The morphism of E^+ -modules $M \rightarrow M \otimes_{E^+} E$ is injective because the E^+ -module M is torsionfree. We identify the E^+ -module M with a submodule of the E^+ -module $M \otimes_{E^+} E$ via this morphism. We prove the result by induction on the dimension n of the E -vector space $M \otimes_{E^+} E$.

For $n = 0$, the E^+ / p -module M / p is free of rank 0. For $n = 1$, the E^+ -module M is isomorphic to a submodule of E , and we consider two cases. If multiplication by p is an automorphism of the E^+ -module M , then the E^+ / p -module M / p is free of rank 0. If multiplication by p is not an automorphism of the E^+ -module M , then it is injective by torsionfreeness of M , but not surjective. In this case, the E^+ -module M is not equal to E and in particular, up to shifting M multiplicatively by an element of E^+ , we can assume that M is contained in E^+ , i.e., that M is an ideal of E^+ . Every ideal of a ring is the filtered union of its finitely generated subideals, and every such ideal is principal in a valuation ring. So the E^+ -module M can be written as a filtered union of free rank 1 submodules $b_i E^+ \subset E^+$, $i \in I$. Multiplication by p is not surjective on M if and only if for all $\varepsilon \in \sqrt{(p)}$, multiplication by ε is not surjective on M . In terms of the identification $M = \cup_i b_i E^+ \subset E^+$, this means that for all $\varepsilon \in \sqrt{(p)}$ there is some index $i_0 \in I_0$ such that for all higher indices $i \geq i_0$, the element εb_i does not divide b_{i_0} in the valuation ring E^+ . In particular for any $\varepsilon \in \sqrt{(p)}$, the cokernel of the inclusion $b_{i_0} E^+ \hookrightarrow M$ is killed by multiplication by ε . So the E^+ -module M is almost free of rank 1, and the E^+ / p -module M / p is almost free of rank 1.

Fix an integer $N \geq 1$, assume the result is proved for all integers $n \leq N$, and let M be an E^+ -submodule of the $(N + 1)$ -dimensional E -vector space $\oplus_{i=1}^{N+1} E e_i$. There is an exact sequence of E^+ -modules

$$0 \longrightarrow M \cap \oplus_{i=1}^N E e_i \longrightarrow M \longrightarrow M_{e_{N+1}} \longrightarrow 0,$$

where $M_{e_{N+1}}$ is the image of M under the projection $\oplus_{i=1}^{N+1} E e_i \rightarrow E e_{N+1}$. All the E^+ -modules in this exact sequence are torsionfree and thus flat over the valuation ring E^+ . In particular the exactness is preserved modulo p :

$$0 \longrightarrow (M \cap \oplus_{i=1}^N E e_i) / p \longrightarrow M / p \longrightarrow M_{e_{N+1}} / p \longrightarrow 0.$$

Let ε be an element of the prime ideal $\sqrt{(p)}$. The induction hypothesis implies that the E^+ / p -modules $(M \cap \oplus_{i=1}^N E e_i) / p$ and $(M_{e_{N+1}}) / p$ are almost free. Let M_1 and M_2 be free E^+ / p -modules equipped with morphisms $M_1 \rightarrow (M \cap \oplus_{i=1}^N E e_i) / p$ and $M_2 \rightarrow (M_{e_{N+1}}) / p$ of E^+ / p -modules whose kernels and cokernels are killed by multiplication by ε . There is a commutative diagram of E^+ / p -modules

$$\begin{array}{ccccccc} 0 & \longrightarrow & M_1 & \longrightarrow & M_1 \oplus M_2 & \longrightarrow & M_2 \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & (M \cap \oplus_{i=1}^N E e_i) / p & \longrightarrow & M / p & \longrightarrow & (M_{e_{N+1}}) / p \longrightarrow 0 \end{array}$$

where the morphism $M_2 \rightarrow M/p$ is defined by choosing arbitrarily a basis of the free E^+/p -module M_2 and, using this basis, a lift of the morphism $M_2 \rightarrow (M_{e_{N+1}})/p$ via the surjection $M/p \rightarrow (M_{e_{N+1}})/p$. By the snake lemma, the kernel and cokernel of the vertical map $M_1 \oplus M_2 \rightarrow M/p$ are killed by multiplication by ε^2 . More precisely, for any element

$$x \in \text{Ker}(M_1 \oplus M_2 \rightarrow M/p) \quad (\text{resp. } x \in \text{Coker}(M_1 \oplus M_2 \rightarrow M/p)),$$

its multiplication εx by ε maps to 0 in

$$\text{Ker}(M_2 \rightarrow (M_{e_{N+1}})/p) \quad (\text{resp. } \text{Coker}(M_2 \rightarrow (M_{e_{N+1}})/p)).$$

So

$$\varepsilon x \in \text{Ker}(M_1 \oplus M_2 \rightarrow M/p) \quad (\text{resp. } \varepsilon x \in \text{Coker}(M_1 \oplus M_2 \rightarrow M/p))$$

is the image of an element of

$$\text{Ker}(M_1 \rightarrow (M \cap \bigoplus_{i=1}^N Ee_i)/p) \quad (\text{resp. } \text{Coker}(M_1 \rightarrow (M \cap \bigoplus_{i=1}^N Ee_i)/p)),$$

and is thus killed by multiplication by ε . By Lemma 3.10, this implies that the E^+/p -module M is almost free of rank $\leq N + 1$. \square

Corollary 3.13. *Let C^+ be a p -adically separated mixed-characteristic valuation ring whose p -adic completion is a perfectoid ring. Let E^+ be a p -adically separated valuation ring extension of C^+ , whose induced field extension $C \rightarrow E$ is of finite type. Then the E^+/p -module $\Omega_{(E^+/p)/(C^+/p)}^1$ is almost free of finite rank.*

Proof. We apply Lemma 3.12 to the E^+ -module Ω_{E^+/C^+}^1 . The E^+ -module Ω_{E^+/C^+}^1 is p -torsionfree (Proposition 3.8), and thus torsionfree since E^+ is p -adically separated. The localisation at a prime ideal has trivial cotangent complex, so the canonical map $\Omega_{E^+/C^+}^1 \otimes_{E^+} E \rightarrow \Omega_{E/C}^1$ is an isomorphism. Moreover, the E -vector space $\Omega_{E/C}^1$ is finite dimensional since $C \rightarrow E$ is a field extension of finite type. The reduction modulo p of the E^+ -module Ω_{E^+/C^+}^1 is the E^+/p -module $\Omega_{(E^+/p)/(C^+/p)}^1$, hence the conclusion. \square

We now use this almost projectivity result to almost lift the C^+/p -algebra E^+/p to a valuation ring extension E^{+b} of C^{+b} (see Notation 3.11).

We will need to use the functor $(-)_*$ to pass from the almost setting to the honest setting. More precisely, the localisation functor $(-)^a$ from C^+ -modules to almost C^+ -modules has a right adjoint given by:

$$(-)_* : \begin{cases} (C^+)^a\text{-Mod} & \longrightarrow C^+\text{-Mod} \\ M & \longmapsto \text{Hom}_{(C^+)^a}((\sqrt{(p)})^a, M) \end{cases}$$

By [GR03, Proposition 2.2.13 (ii)], this functor also restricts to a functor

$$(-)_* : (C^+)^a\text{-Alg} \longrightarrow C^+\text{-Alg}$$

which is right adjoint to the localisation functor from C^+ -algebras to almost C^+ -algebras. Given an almost C^+ -algebra A , the unit in $A_* := \text{Hom}_{(C^+)^a}(\mathfrak{m}_{C^+}^a, A)$ is given by the restriction of the structural map $(C^+)^a \rightarrow A$ to $\mathfrak{m}_{C^+}^a$, and the product of elements in A_* is given by multiplication of maps.

The composite functor

$$C^+\text{-Alg} \xrightarrow{(-)^a} (C^+)^a\text{-Alg} \xrightarrow{(-)_*} C^+\text{-Alg}$$

is idempotent, because almost isomorphisms are sent to isomorphisms by the functor $(-)_*$. In general, given a C^+ -algebra A , the unit morphism $A \rightarrow A_* := (A^a)_*$ of C^+ -algebras is an almost isomorphism, and not an isomorphism. For instance, if $A = C^+$, the unit morphism $C^+ \rightarrow (C^+)_*$ is an isomorphism if and only if the maximal ideal of C^+ is equal to $\sqrt{(p)}$. The functor $(-)_*$ has a natural analogue over the pair $(C^{+b}, \sqrt{(d)})$.

Lemma 3.14. *Let C^+ be a mixed-characteristic valuation ring whose p -adic completion is a perfectoid ring. If E^+ is a valuation ring extension of C^+ , then the C^+ -algebra $(E^+)_* := (E^{+a})_*$ is also a valuation ring extension of C^+ . More precisely: if p is not invertible in E^+ , the ideal*

$$\sqrt{(p)}E^+ := \{\varepsilon x \in E^+ \mid \varepsilon \in \sqrt{(p)} \subset C^+, x \in E^+\}$$

is a prime ideal of the valuation ring E^+ , and the unit morphism $E^+ \rightarrow (E^+)_$ is naturally identified with the localisation at this prime ideal; if p is invertible in E^+ , then the morphism $E^+ \rightarrow (E^+)_*$ is the identity. The same holds true for valuation ring extensions of C^{+b} .*

Proof. First assume that p is not invertible in E^+ ; in this case, no element of the ideal $\sqrt{(p)}E^+ \subset E^+$ is invertible. Each element ε of the prime ideal $\sqrt{(p)} \subset C^+$ is the product of two elements of $\sqrt{(p)} \subset C^+$ by Lemma 3.10, and the same holds true for the ideal $\sqrt{(p)}E^+ \subset E^+$ via the embedding $C^+ \hookrightarrow E^+$. An element of the valuation ring E^+ is not in the ideal $\sqrt{(p)}E^+ \subset E^+$ if and only if it is not divisible by any element of $\sqrt{(p)}E^+$, or equivalently if it divides every element of $\sqrt{(p)}E^+$. In particular the product of two elements of E^+ which are not in $\sqrt{(p)}E^+$ is not in $\sqrt{(p)}E^+$, and the ideal $\sqrt{(p)}E^+$ is a prime ideal of E^+ .

Every morphism $\alpha : \sqrt{(p)} \rightarrow E^+$ of C^+ -modules extends uniquely, by torsionfreeness of the C^+ -module E^+ , to a morphism $\tilde{\alpha} : C^+ \rightarrow E^+[1/p]$ of C^+ -modules. A morphism $\beta : C^+ \rightarrow E^+[1/p]$ of C^+ -modules then comes from a morphism $\alpha : \sqrt{(p)} \rightarrow E^+$ of C^+ -modules if and only if $\varepsilon\beta(1)$ belongs to $E^+ \subset E^+[1/p]$ for every $\varepsilon \in \sqrt{(p)}$, or equivalently if $\beta(1)$ is in $E^+_{\sqrt{(p)}E^+}$. In particular the C^+ -module $(E^+)_* := \text{Hom}_{C^+}(\sqrt{(p)}, E^+)$ is naturally identified with the localisation $E^+_{\sqrt{(p)}E^+}$ of E^+ at the prime ideal $\sqrt{(p)}E^+$. As a localisation of the valuation ring E^+ , the ring $(E^+)_*$ is a valuation ring extension of E^+ , and thus of C^+ .

Similarly, if p is invertible in E^+ then any morphism $\alpha : \sqrt{(p)} \rightarrow E^+$ of C^+ -modules extends uniquely to a morphism $\tilde{\alpha} : C^+ \rightarrow E^+[1/p] = E^+$ of C^+ -modules. In particular the morphism $E^+ \rightarrow (E^+)_*$ is the identity. \square

Remark 3.15. A mixed-characteristic valuation ring has rank 1 if and only if it is p -adically separated and if its maximal ideal is the prime ideal $\sqrt{(p)}$. In particular if C^+ is a p -adically separated mixed-characteristic valuation ring, then C^+ has rank 1 if and only if the morphism $C^+ \rightarrow (C^+)_*$ is an isomorphism (Lemma 3.14). If C^+ is a rank 1 mixed-characteristic valuation ring, note that the morphism $C^+/p \rightarrow (C^+/p)_*$ is not an isomorphism of C^+ -algebras in general; it is if C^+ is spherically complete.

We will need the following to ensure that the flat lift E^{+b} produced by deformation theory is actually a valuation ring.

Lemma 3.16. *Let C^+ be a mixed-characteristic valuation ring whose p -adic completion is a perfectoid ring. Let E^+ be a valuation ring extension of C^+ and E^{+b} a d -complete flat C^{+b} -algebra such that the morphism $E^{+b} \rightarrow (E^{+b})_*$ is an isomorphism. Assume there is an almost isomorphism of $(C^{+b}/d)^a = (C^+/p)^a$ -algebras $(E^{+b}/d)^a \cong (E^+/p)^a$. Then E^{+b} is a valuation ring extension of C^{+b} .*

Proof. We want to prove that E^{+b} is an integral domain such that for any elements f and g in E^{+b} , either f divides g or g divides f . The morphism $E^+ \rightarrow (E^+)_*$ of C^+ -algebras is an almost isomorphism, so its reduction modulo p is an almost isomorphism of C^+/p -algebras. The C^+ -algebra $(E^+)_*$ is a valuation ring extension of C^+ (Lemma 3.14). In particular one can replace E^+ by $(E^+)_*$, and we assume now that the morphism $E^+ \rightarrow (E^+)_*$ is an isomorphism of C^+ -algebras.

The almost isomorphism $(E^{+b}/d)^a \cong (E^+/p)^a$, corresponding to an isomorphism between $(E^{+b}/d)_*$ and $(E^+/p)_*$, will be used in the following way. The functor $(-)_*$ is left exact and the morphism $E^+ \rightarrow (E^+)_*$ is an isomorphism, so $(-)_*$ applied to the short exact sequence of C^+ -modules

$$0 \longrightarrow E^+ \xrightarrow{p} E^+ \longrightarrow E^+/p \longrightarrow 0$$

implies that the morphism $E^+/p \rightarrow (E^+/p)_*$ of C^+/p -algebras is injective. Similarly, the morphism $E^{+b}/d \rightarrow (E^{+b}/d)_*$ of C^{+b}/d -algebras is injective. These two injective morphisms are almost isomorphisms by definition, meaning here that their cokernels are killed by any $\varepsilon \in \sqrt{(p)}$ (resp. $\varepsilon \in \sqrt{(d)}$). In particular we will identify, via these morphisms, the multiplication by ε of any element of E^+/p (resp. E^{+b}/d) with an element of E^+/p (resp. E^{+b}/d).

The valuation ring C^{+b} is perfect by definition. Denote by $(d^{1/p^n})_{n \in \mathbb{N}} \in (C^{+b})^{\mathbb{N}}$ a compatible system of p -power roots of $d \in C^{+b}$. A module over the valuation ring C^{+b} is flat if and only if it is torsionfree, so the C^{+b} -module E^{+b} is d^{1/p^n} -torsionfree for each $n \geq 0$. In the following we identify the rings C^+/p and C^{+b}/d ; in particular any (rational) power d^α of $d \in C^{+b}$ defines an element of C^+/p , which we also denote by d^α .

Let f and g be elements of the ring E^{+b} . Let us prove that either f divides g or g divides f . The ring E^{+b} is d -torsionfree, so we can assume that d does not divide one of f and g . We first assume that d does not divide f in E^{+b} , and prove that f divides d . Because the morphism $E^{+b} \rightarrow (E^{+b})_*$ is an isomorphism, there exists $\varepsilon \in \sqrt{(d)}$ such that εf is not divisible by d in E^{+b} : otherwise the element $f/d \in E^{+b}[1/d]$ would be in $(E^{+b})_*$ and not in E^{+b} . Then εf is naturally a nonzero element of the ring E^+/p , and because the morphism $E^+ \rightarrow (E^+)_*$ is an isomorphism there exists $\alpha \in \mathbb{Z}[1/p]$, $\alpha < 1$, such that d^α does not divide εf in E^+/p , and in particular εf divides d^α in E^+/p . This implies that there exist $h, t \in E^{+b}$ such that $\varepsilon fh = d^\alpha + dt$. Because E^{+b} is d -adically complete it is also $d^{(1-\alpha)}$ -adically complete. In particular $1 + d^{(1-\alpha)}t$ is a unit in E^{+b} , so f divides d^α . By construction d^α divides d , so f divides d .

Going back to the elements f and g of E^{+b} , we deduce the following: if f is not divisible by d and g is divisible by d , then f divides g . We now assume that d does not divide f or g . Because the morphism $E^{+b} \rightarrow (E^{+b})_*$ is an isomorphism of C^{+b} -algebras, there exists $\varepsilon \in \sqrt{(d)}$ such that $\varepsilon^2 f$ and $\varepsilon^2 g$ are not divisible by d . We identify εf and εg with nonzero elements of the ring E^+/p . Without loss of generality and because E^+ is a valuation ring, there is an element $h \in E^+/p$ such that $\varepsilon f = \varepsilon gh$. Multiply this equality by ε and identify εh with an element of E^{+b}/d . We can then write $\varepsilon^2 f = \varepsilon g \tilde{h} + dt$ for some elements $\tilde{h}, t \in E^{+b}$ (where \tilde{h} is mapped to εh by the morphism $E^{+b} \rightarrow E^{+b}/d$). Because d does not divide $\varepsilon^2 f$, in particular d does not divide ε and ε must divide d in C^{+b} . The element $\varepsilon h \in E^{+b}/d$ is zero modulo ε , so the element \tilde{h} is also zero modulo ε , i.e., \tilde{h} is divisible by ε . By the previous paragraph and because d does not divide $\varepsilon^2 g$ by assumption, we know that $\varepsilon^2 g$ divides d . So $\varepsilon^2 g$ divides $\varepsilon^2 f$, and g divides f by ε -torsionfreeness.

We now prove that the ring E^{+b} is an integral domain. Given two elements f and g of the ring E^{+b} , then without loss of generality we can assume that g divides f using the divisibility property we just proved. Assume now $fg = 0$. The valuation ring E^{+b} is d -adically separated, so if f is nonzero then f divides d^n for some integer $n \geq 1$. But then $d^{2n} = 0$, which is impossible since E^{+b} is d -torsionfree. So $f = 0$, and the ring E^{+b} is an integral domain. By assumption E^{+b} is a flat C^{+b} -algebra, so E^{+b} is a valuation ring extension of C^{+b} . \square

The next result will be used to deal with compatibility between d -adic completion and the functor $(-)_*$ in the proof of Theorem 3.18.

Lemma 3.17. *Let C^+ be a mixed-characteristic valuation ring whose p -adic completion is a perfectoid ring, and M be an almost flat $(C^+)^a$ -module. For any $\varepsilon_1, \varepsilon_2, \varepsilon_3 \in \mathfrak{m}_{C^+}$, the reduction modulo ε_1 of the morphism $(M/\varepsilon_1 \varepsilon_2 \varepsilon_3)_* \rightarrow (M/\varepsilon_1 \varepsilon_2)_*$ is an isomorphism of C^+ -modules. The same holds true over C^{+b} .*

Proof. This is a direct consequence of [Sch12, Lemma 5.3 (iii)]. More precisely, the proof for almost mathematics with respect to the pair (K°, \mathfrak{m}) , where K° is a rank 1 perfectoid valuation ring with maximal ideal \mathfrak{m} , adapts directly to almost mathematics with respect to the pair $(C^+, \sqrt{(p)})$. \square

Theorem 3.18. *Let C^+ be a mixed-characteristic valuation ring whose p -adic completion is a perfectoid ring, and E^+ be a valuation ring extension of C^+ . If the E^+/p -module $\Omega^1_{(E^+/p)/(C^+/p)}$ is almost projective, then there exists a unique d -complete rank 1 valuation ring extension E^{+b} of C^{+b} such that there is an almost isomorphism $(E^{+b}/d)^a \cong (E^+/p)^a$.*

The almost projectivity hypothesis is satisfied in particular if the induced field extension $C \rightarrow E$ is of finite type, (Corollary 3.13). It is also satisfied for the p -adic completion of such an extension, as it depends only on its reduction modulo p .

Proof. This result depends only on the morphism $C^+/p \rightarrow E^+/p$, so we can replace C^+ and E^+ by their p -adically separated quotients $C^+/\bigcap_{n \geq 0} p^n C^+$ and $E^+/\bigcap_{n \geq 0} p^n E^+$. We first prove by induction on $n \geq 1$ that there are unique, up to isomorphism, almost flat $(C^{+b}/d^m)^a$ -algebras E_m^{+a} with isomorphisms $E_m^{+a}/d \xrightarrow{\cong} (E^+/p)^a$ for all $1 \leq m \leq n$. By the almost variant [GR03, Corollary 3.2.11] of Theorem 3.9, the vanishing of the obstructions and ambiguities in this process can be expressed in terms of the almost cotangent complexes $\mathbb{L}_{E_m^{+a}/(C^{+b}/d^m)^a}^a$. For $n = 1$, define E_1^{+a} as $(E^+/p)^a$: E^+ is a flat C^+ -algebra, so E^+/p is a flat C^+/p -algebra, and E_1^{+a} is thus an almost flat $(C^{+b}/d)^a = (C^+/p)^a$ -algebra.

Now let $n \geq 1$, and assume we are given for each $1 \leq m \leq n$ an almost flat $(C^{+b}/d^m)^a$ -algebras E_m^{+a} , with isomorphisms $E_m^{+a}/d^{m-1} \cong E_{m-1}^{+a}$ for all $2 \leq m \leq n$. We claim the almost cotangent complex $\mathbb{L}_{E_n^{+a}/(C^{+b}/d^n)^a}^a$ is in degree 0, given by an almost projective E_n^{+a} -module. Tensoring the short exact sequence

$$0 \longrightarrow E_1^{+a} \xrightarrow{d^{n-1}} E_n^{+a} \longrightarrow E_{n-1}^{+a} \longrightarrow 0 \quad (2)$$

of $(C^{+b}/d^n)^a$ -modules by the almost cotangent complex $\mathbb{L}_{E_n^{+a}/(C^{+b}/d^n)^a}^a$ induces a distinguished triangle

$$\mathbb{L}_{(E^+/p)/(C^+/p)^a}^a \longrightarrow \mathbb{L}_{(E_n^{+a})/(C^{+b}/d^n)^a}^a \longrightarrow \mathbb{L}_{(E_{n-1}^{+a})/(C^{+b}/d^{n-1})^a}^a \longrightarrow$$

by [GR03, Theorem 2.5.32]. The discreteness and the almost projectivity of $\mathbb{L}_{E_n^{+a}/(C^{+b}/d^n)^a}^a$ thus reduces inductively to the case of $n = 1$. For $n = 1$, $\mathbb{L}_{(E^+/p)/(C^+/p)}^a$ is in degree 0 by Proposition 3.8, and the E^+/p -module $\Omega_{(E^+/p)/(C^+/p)}^1$ is almost projective by assumption.

In particular the Ext-groups in the almost deformation theory [GR03, Corollary 3.2.11] vanish, and so do the deformation classes. This implies that there is a unique, up to isomorphism, almost flat $(C^{+b}/d^{n+1})^a$ -algebra E_{n+1}^{+a} with an isomorphism $E_{n+1}^{+a}/d^n \xrightarrow{\cong} E_n^{+a}$. This concludes the induction.

We then define for each $n \geq 1$ the C^{+b}/d^n -algebras $E_n^+ := (E_n^{+a})_*$. Beware that the functor $(-)_*$ is not right exact in general, and in particular the morphisms $E_n^+ \rightarrow E_{n-1}^+$ are only almost surjective, even if E^+ is a rank 1 valuation ring. Let E^{+b} be the inverse limit of the system $(E_n^+)_{n \geq 1}$ along these maps. The functor $(-)_*$ being right-adjoint to the localisation functor it commutes with limits, and the morphism $E^{+b} \rightarrow (E^{+b})_*$ is an isomorphism. In particular the C^{+b} -algebra E^{+b} is the unique C^{+b} -algebra which is almost flat, almost d -complete, whose reduction modulo d is almost isomorphic to E^+/p and such that the morphism $E^{+b} \rightarrow (E^{+b})_*$ is an isomorphism. Let us prove that this unique C^{+b} -algebra is necessarily flat, d -complete and a valuation ring extension of C^{+b} .

By Lemma 3.17 and for any $n \geq 1$, the image of E_{n+k}^+ in E_n^+ is the same for all $k \geq 1$. In particular the C^{+b} -algebra E^{+b} is d -adically complete. The module M_* associated to an almost flat C^{+b} -module M is not necessarily flat. However proving that the algebra E^{+b} over the valuation ring C^{+b} is flat is equivalent to proving that it is a torsionfree C^{+b} -module, which is again equivalent to proving that it is a d -torsionfree C^{+b} -module. To prove this, consider the short exact sequence of almost C^{+b} -module:

$$0 \longrightarrow E_{n-1}^{+a} \xrightarrow{d} E_n^{+a} \longrightarrow E_1^{+a} \longrightarrow 0.$$

Applying the functor $(-)_*$ induces an exact sequence of C^{+b} -modules:

$$0 \longrightarrow E_{n-1}^+ \xrightarrow{d} E_n^+ \longrightarrow E_1^+,$$

and then passing to the inverse limit over $n \geq 1$ implies that the C^{+b} -module E^{+b} has no d -torsion. The C^{+b} -algebra E^{+b} is thus d -adically complete and flat, and the morphism $E^{+b} \rightarrow (E^{+b})_*$ is an isomorphism. By Lemma 3.16, E^{+b} is thus a valuation ring extension of C^{+b} . \square

Before proving the Cartier isomorphism for general valuation rings E^+ , let us reformulate the deformation result Theorem 3.18.

Corollary 3.19. *Let C^+ be a perfectoid valuation ring, and C^{+b} the tilt of C^+ . There is a natural bijection between:*

- (1) *isomorphism classes of p -complete rank 1 valuation ring extensions E^+ of C^+ such that the E^+/p -module $\Omega_{(E^+/p)/(C^+/p)}^1$ is almost projective;*
- (2) *isomorphism classes of d -complete rank 1 valuation ring extensions E^{+b} of C^{+b} such that the E^{+b}/d -module $\Omega_{(E^{+b}/d)/(C^{+b}/d)}^1$ is almost projective.*

This bijection is characterised by the following: two valuation rings E^+ and E^{+b} correspond via this bijection if and only if there is an almost isomorphism between E^+/p and E^{+b}/d .

Beware that this bijection is not functorial, because the automorphisms on both sides are not identified. As a consequence, filtered systems of C^+/p -algebras do not lift naturally to filtered systems of C^{+b} -algebras, and Theorem 3.18 and Corollary 3.19 do not hold directly for arbitrary valuation rings.

Proof. This is a direct consequence of the proof of Theorem 3.18, where the d -adic and p -adic objects play symmetric roles. \square

If the almost isomorphism in Theorem 3.18 is an honest isomorphism (*e.g.*, if the p -adic completion of E^+ is a perfectoid ring or if E^+ is spherically complete), then $C^+/p \rightarrow E^+/p$ satisfies the Cartier isomorphism. More precisely, the Cartier isomorphism for $C^{+b} \rightarrow E^{+b}$ (Theorem 3.4) implies the Cartier isomorphism for $C^+/p \rightarrow E^+/p$ by base change (Lemma 2.8) along the morphism $C^{+b} \rightarrow C^+/d$.

In general the almost isomorphism of Theorem 3.18 will be sufficient to deduce the Cartier isomorphism for the morphism $C^+/p \rightarrow E^+/p$.

Proof of Theorem 3.1. As a flat morphism of integral domains, the valuation ring extension $C^+ \rightarrow E^+$ is p -Cartier smooth if and only if the morphism $C^+/p \rightarrow E^+/p$ is Cartier smooth (Remark 3.3). The result in characteristic p is Theorem 3.4. In mixed-characteristic, the quasismoothness of the morphism $C^+/p \rightarrow E^+/p$ is Proposition 3.8. The inverse Cartier map depends only on the morphism $C^+/p \rightarrow E^+/p$, so we can replace C^+ and E^+ by their p -adically separated quotients to prove the Cartier isomorphism. We assume now that C^+ is a mixed-characteristic p -adically separated valuation ring whose p -adic completion is a perfectoid ring, that E^+ is a p -adically separated valuation ring extension of C^+ , and prove the Cartier isomorphism for the morphism $C^+/p \rightarrow E^+/p$. The fraction field E of the valuation ring E^+ is the filtered union of the finite type field extensions E_i of C contained in E . The valuation ring E^+ is the filtered union of the associated valuation rings E_i^+ and the inverse Cartier map commutes with filtered colimits, so we can assume that the field extension $C \rightarrow E$ is of finite type.

Proving the Cartier isomorphism for the morphism $C^+/p \rightarrow E^+/p$ is equivalent to proving that the map

$$\mathbb{L}\Omega_{(E^+/p)/(C^+/p)} \longrightarrow \Omega_{(E^+/p)/(C^+/p)}$$

is an equivalence in the derived category $\mathcal{D}(C^+/p)$ (Theorem 2.16). The map

$$\mathbb{L}\Omega_{E^{+b}/C^{+b}} \longrightarrow \Omega_{E^{+b}/C^{+b}}$$

and thus its reduction modulo d

$$\mathbb{L}\Omega_{(E^{+b}/d)/(C^{+b}/d)} \longrightarrow \Omega_{(E^{+b}/d)/(C^{+b}/d)}$$

are equivalences for any valuation ring extension E^{+b} of C^{+b} (Theorems 2.16 and 3.4). We apply this to the valuation ring E^{+b} corresponding to E^+ by Theorem 3.18, for which there is an almost isomorphism between E^+/p and E^{+b}/d . By [GR03, Theorem 2.5.36] and using transivity for the cotangent complex, the first and last maps in

$$\mathbb{L}_{(E^+/p)/(C^+/p)} \longrightarrow \mathbb{L}_{(E^+/p)_*/(C^+/p)} \simeq \mathbb{L}_{(E^{+b}/d)_*/(C^{+b}/d)} \longleftarrow \mathbb{L}_{(E^{+b}/d)/(C^{+b}/d)}$$

are almost equivalences. Taking wedge powers, the same is true for \mathbb{L}^n , for each $n \geq 0$. The first and last maps in

$$\mathbb{L}\Omega_{(E^+/p)/(C^+/p)} \longrightarrow \mathbb{L}\Omega_{(E^+/p)_*/(C^+/p)} \simeq \mathbb{L}\Omega_{(E^{+\flat}/d)_*/(C^{+\flat}/d)} \longleftarrow \mathbb{L}\Omega_{(E^{+\flat}/d)/(C^{+\flat}/d)}$$

are thus almost equivalences, because they are almost equivalences on the graded pieces of the exhaustive conjugate filtration. This defines an almost quasi-isomorphism between $\mathbb{L}\Omega_{(E^+/p)/(C^+/p)}$ and $\mathbb{L}\Omega_{(E^{+\flat}/d)/(C^{+\flat}/d)}$. In particular the map

$$\mathbb{L}\Omega_{(E^+/p)/(C^+/p)} \longrightarrow \Omega_{(E^+/p)/(C^+/p)}$$

is an almost equivalence in the derived category $\mathcal{D}(C^+/p)$, *i.e.*, its homotopy fibre F has almost zero cohomology groups. Let us prove that this homotopy fibre F is actually zero.

There there is an integer $n \geq 0$ such that the E^+/p -module $\Omega_{(E^+/p)/(C^+/p)}^1$ is flat and almost free of rank at most n (Proposition 3.8). The wedge power of a flat module is flat, so the E^+/p -module $\Omega_{(E^+/p)/(C^+/p)}^k$ is flat for any $k \geq 0$, and almost zero if $k \geq n+1$. A flat and almost zero E^+/p -module M is necessarily zero: for any $\varepsilon \in \mathfrak{m}_{C^+}$ such that ε^2 divides p in C^+ , any element of the module M is almost zero and in particular killed by multiplication by ε ; by flatness, any such element is also divisible by ε , and thus zero. In particular the E^+/p -module $\Omega_{(E^+/p)/(C^+/p)}^k$ is zero for any $k \geq n+1$. By quasismoothness of the morphism $C^+/p \rightarrow E^+/p$ and using the conjugate filtration on the derived de Rham complex, this implies that the complex $\mathbb{L}\Omega_{(E^+/p)/(C^+/p)}$ is also concentrated in degrees $[0; n]$. The homotopy fibre F is thus almost zero, and possibly nonzero only in degrees $[0; n+1]$.

We now use that for an almost zero C^+/p -module M , the canonical map $M \rightarrow M/\sqrt{(p)}$ is an isomorphism of C^+/p -modules. Assume F is nonzero, and let $i \geq 0$ be the maximal integer such that $H^i(F)$ is nonzero. Tensoring the fibre sequence

$$F \longrightarrow \mathbb{L}\Omega_{(E^+/p)/(C^+/p)} \longrightarrow \Omega_{(E^+/p)/(C^+/p)}$$

with $C^+/\sqrt{(p)}$ induces a fibre sequence

$$F \otimes_{C^+/p}^{\mathbb{L}} C^+/\sqrt{(p)} \longrightarrow \mathbb{L}\Omega_{(E^+/\sqrt{(p)})/(C^+/\sqrt{(p)})} \longrightarrow \Omega_{(E^+/\sqrt{(p)})/(C^+/\sqrt{(p)})}.$$

The ring $C^+/\sqrt{(p)}$ is a perfect valuation ring of characteristic p , and $E^+/\sqrt{(p)}$ is a valuation ring extension of $C^+/\sqrt{(p)}$. By Theorems 3.4 and 2.16, the homotopy fibre $F \otimes_{C^+/p}^{\mathbb{L}} C^+/\sqrt{(p)}$ is thus zero. Because F is concentrated in degrees $[0; i]$ and almost zero, there are canonical identifications

$$H^i(F \otimes_{C^+/p}^{\mathbb{L}} C^+/\sqrt{(p)}) \cong H^i(F)/\sqrt{(p)} \cong H^i(F),$$

which contradicts the fact that $H^i(F)$ is nonzero. So the homotopy fibre F is zero, and the morphism $C^+/p \rightarrow E^+/p$ satisfies the Cartier isomorphism. \square

4 THE SYNTOMIC-ÉTALE COMPARISON THEOREM

In this section we prove Theorem B (Theorems 4.12 and 4.14), comparing the syntomic cohomology of a p -Cartier smooth algebra over a perfectoid ring to the étale cohomology of its generic fibre. This comparison theorem was also proved in [KM21, LM21] in characteristic p , and in [BM22] for p -torsionfree F -smooth schemes, using different methods.

4.1 Syntomic cohomology

The syntomic complexes $\mathbb{Z}_p(i)^{\text{syn}}$ were first defined in [BMS19, Section 7.4] for p -complete p -quasisyntomic rings in terms of p -completed topological cyclic homology. Another equivalent definition was given in [BS22] in terms of absolute prismatic cohomology, and a generalisation for general schemes was developed in [BL22, Section 8.4]. We will be interested only in p -complete p -quasisyntomic algebras over

a perfectoid ring, whose syntomic complexes can be defined in terms of relative prismatic cohomology (Definition 4.1).

The ideal $I \subset A$ of a perfect prism (subsection 2.4 or [BS22, Definition 3.2 (2)]) is necessarily principal, generated by a nonzerodivisor $d \in A$. When defining a perfect prism $(A, (d))$ in this section, we implicitly fix a choice of generator $d \in A$. When the base prism (A, I) is perfect, one can define a Nygaard filtration $\mathcal{N}^{\geq *}\Delta_{S/A}$ on the prismatic complex $\Delta_{S/A}$ (without Frobenius twist), and we denote by

$$\phi : \mathcal{N}^{\geq *}\Delta_{S/A} \xrightarrow{\sim} \mathcal{N}^{\geq *}\Delta_{S/A}^{(1)} \xrightarrow{\phi} I^*\Delta_{S/A}$$

the Frobenius on the relative prismatic complex (see subsection 2.2 for the definition of the second map). Following [BS22, Section 12] and for each $i \geq 0$, a divided Frobenius map

$$\phi_i = \left\langle \frac{\phi}{d^i} \right\rangle : \mathcal{N}^{\geq i}\Delta_{S/A} \longrightarrow \Delta_{S/A}$$

is defined and sits in the commutative diagram

$$\begin{array}{ccc} \mathcal{N}^{\geq i}\Delta_{S/A} & \xrightarrow{\phi_i} & \Delta_{S/A} \\ & \searrow \phi & \downarrow d^i \\ & & d^i\Delta_{S/A}. \end{array}$$

Definition 4.1 (Syntomic complexes). *Let $(A, (d))$ be a perfect prism, and S a p -complete p -quasi-syntomic A/d -algebra. For each integer $i \geq 0$, the syntomic complex $\mathbb{Z}_p(i)^{\text{syn}}(S) \in \mathcal{D}^{\geq 0}(\mathbb{Z}_p)$ is*

$$\mathbb{Z}_p(i)^{\text{syn}}(S) := \text{hofib}(\phi_i - 1 : \mathcal{N}^{\geq i}\Delta_{S/A} \longrightarrow \Delta_{S/A}).$$

For each integer $n \geq 1$, also define $\mathbb{Z}/p^n(i)^{\text{syn}}(S) := \mathbb{Z}_p(i)^{\text{syn}}(S) \otimes_{\mathbb{Z}_p}^{\mathbb{L}} \mathbb{Z}/p^n$.

Over a scheme in which p is invertible, the object $\mathbb{Z}_p(i)$, called *p -adic étale Tate twist*, will denote the (pro-)étale sheaf defined as the inverse limit over $n \geq 1$ of the étale sheaves $\mu_{p^n}^{\otimes i}$. Following [BL22, Section 8.3], there is map comparing syntomic complexes and p -adic étale Tate twists.

Construction 4.2 (The syntomic-étale comparison map, [BL22]). *Let S be a p -complete p -quasi-syntomic ring (e.g., a p -complete p -quasisyntomic algebra over a perfectoid ring). For every integer $i \geq 0$, there is a canonical map*

$$\gamma_{\text{syn}}^{\text{ét}}\{i\} : \mathbb{Z}_p(i)^{\text{syn}}(S) \longrightarrow R\Gamma_{\text{proét}}(\text{Spec}(S[\frac{1}{p}]), \mathbb{Z}_p(i)).$$

For every integer $n \geq 1$, one can also consider the derived reduction modulo p^n of this canonical map

$$\gamma_{\text{syn}}^{\text{ét}}\{i\}/p^n : \mathbb{Z}/p^n(i)^{\text{syn}}(S) \longrightarrow R\Gamma_{\text{ét}}(\text{Spec}(S[\frac{1}{p}]), \mu_{p^n}^{\otimes i}).$$

Lemma 4.3. *Let S be a perfectoid ring. Then for any integer $n \geq 1$, the canonical map*

$$\mu_{p^n}(S) \longrightarrow \mu_{p^n}(S[\frac{1}{p}])$$

is a group isomorphism.

Proof. The map $S \rightarrow S[\frac{1}{p}]$ factors as

$$S \rightarrow S/S[p] \hookrightarrow (S/S[p])[\frac{1}{p}] \cong S[\frac{1}{p}],$$

where $S/S[p]$ is a p -torsionfree perfectoid ring ([BMS19, Proposition 4.19 (3)]). The map

$$\mu_{p^n}(S/S[p]) \longrightarrow \mu_{p^n}((S/S[p])[\frac{1}{p}])$$

is injective because the ring $S/S[p]$ is p -torsionfree, and surjective because $S/S[p]$ is integrally closed in $(S/S[p])[\frac{1}{p}]$ ([BMS18, Lemma 3.21]). The map

$$\mu_{p^n}(S) \longrightarrow \mu_{p^n}(S/S[p])$$

is injective: suppose some element $x \in \mu_{p^n}(S)$ is sent to 1 in $\mu_{p^n}(S/S[p])$; then it is also sent to 1 in $\mu_{p^n}(S/\sqrt{(p)})$ since $S/\sqrt{(p)}$ is a perfect \mathbb{F}_p -algebra; the cartesian square ([ČS21, 2.1.3])

$$\begin{array}{ccc} S & \longrightarrow & S/\sqrt{(p)} \\ \downarrow & & \downarrow \\ S/S[p] & \longrightarrow & (S/S[p])/\sqrt{p(S/S[p])} \end{array} \quad (3)$$

then implies that x is equal to 1. It is also surjective: using the previous cartesian square, any element of $\mu_{p^n}(S/S[p])$ can be lifted to an element x in S whose image in $S/\sqrt{(p)}$ is equal to 1; the p^n -power of $x \in S$ is equal to 1 by construction, *i.e.*, $x \in \mu_{p^n}(S)$. \square

Remark 4.4. Let $(A, (d))$ be a perfect prism, and S a p -quasisyntomic A/d -algebra. For each $i \geq 0$, the canonical map

$$(\mathcal{N}^{\geq i} \Delta_{S/A}[\frac{1}{d}]_p^\wedge \longrightarrow \Delta_{S/A}[\frac{1}{d}]_p^\wedge$$

is an equivalence in the derived category $\mathcal{D}(A)$. It suffices to prove it locally on the p -quasisyntomic site, where there are natural inclusions of (p, d) -completely flat A -modules $\phi_A^{-1}(d)^i \Delta_{S/A} \subseteq \mathcal{N}^{\geq i} \Delta_{S/A} \subseteq \Delta_{S/A}$. The result follows by derived Nakayama and using the equality $(\phi_A^{-1}(d))^p = d$ in A/p .

Theorem 4.5 (Prismatic-étale comparison). *Let $(A, (d))$ be a perfect prism, and S a p -complete p -quasisyntomic A/d -algebra. Then for any integers $i \geq 0$ and $n \geq 1$, there are canonical identifications*

$$R\Gamma_{\text{ét}}(\text{Spec}(S[\frac{1}{p}]), \mu_{p^n}^{\otimes i}) \xrightarrow{\sim} \text{hofib}(\phi_i - 1 : \Delta_{S/A}[\frac{1}{d}]/p^n \longrightarrow \Delta_{S/A}[\frac{1}{d}]/p^n)$$

and

$$R\Gamma_{\text{proét}}(\text{Spec}(S[\frac{1}{p}]), \mathbb{Z}_p(i)) \xrightarrow{\sim} \text{hofib}(\phi_i - 1 : \Delta_{S/A}[\frac{1}{d}]_p^\wedge \longrightarrow \Delta_{S/A}[\frac{1}{d}]_p^\wedge).$$

Proof. For $i = 0$ this is [BS22, Theorem 9.1], and we adapt the proof to the general case $i \geq 0$. We prove the result for torsion coefficients; taking limits over $n \geq 1$ implies the result for integral coefficients. For any object $C \in \mathcal{D}(\mathbb{Z}/p^n)$ equipped with a map $F : C \rightarrow C$, denote by $C^{F=1}$ the homotopy fibre of the map $F - 1 : C \rightarrow C$.

We first prove that the constructions

$$S \longmapsto R\Gamma_{\text{ét}}(\text{Spec}(S[\frac{1}{p}]), \mu_{p^n}^{\otimes i}), (\Delta_{S/A}[\frac{1}{d}]/p^n)^{\phi_i=1}$$

from p -complete p -quasisyntomic A/d -algebras S to $\mathcal{D}(\mathbb{Z}/p^n)$, are arc-sheaves. For the first one this is [BM21, Corollary 6.17]. For the second one, remark that the construction is zero whenever A/d has characteristic p ; base change for prismatic cohomology and the cartesian square (3) then allow us to reduce to the case of a p -torsionfree perfectoid base A/d . Assume first that this p -torsionfree perfectoid ring A/d admits a compatible system of p -power roots of unity, defining an element $[\varepsilon] \in A = W(R^\flat)$. Up to modifying $d \in A$ by a unit, the element $\mu := [\varepsilon] - 1$ is related to $d \in A$ by the formula $d = \mu/\phi_A^{-1}(\mu)$. Multiplication by μ^{-i} induces an equivalence

$$(\Delta_{S/A}[\frac{1}{d}]/p^n)^{\phi_i=1} \simeq (\mu^{-i} \Delta_{S/A}[\frac{1}{d}]/p^n)^{\phi=1},$$

and the map

$$(\mu^{-i} \Delta_{S/A}[\frac{1}{d}]/p^n)^{\phi=1} \longrightarrow (\mu^{-i} \Delta_{S/A, \text{perf}}[\frac{1}{d}]/p^n)^{\phi=1}$$

is a quasi-isomorphism: it suffices to prove it when $n = 1$, where this is a consequence of the equality $d = \mu^{\frac{p-1}{p}}$ and [BS22, Lemma 9.2]. The construction $S \mapsto \Delta_{S/A, \text{perf}}$ is an arc-sheaf because $S \mapsto S_{\text{perfd}}$

is by [BS22, Corollary 8.11]. So the construction $S \mapsto (\Delta_{S/A}[\frac{1}{d}]/p^n)^{\phi_i=1}$ is an arc-sheaf. In general, *i.e.*, without assuming that the p -torsionfree perfectoid ring A/d admits a compatible system of p -power roots of unity, we use the p -quasisyntomic descent property for prismatic cohomology ([BL22, Lemma 4.3.12]) to reduce to the previous case. More precisely, consider a perfect prism $(A', (d))$ over $(A, (d))$ such that A'/d admits a compatible system of p -power roots and such that $A/d \rightarrow A'/d$ is a p -quasisyntomic cover ([BS22, Theorem 7.14]). There are canonical equivalences

$$\Delta_{-/A} \xleftarrow{\sim} \Delta_- \xrightarrow{\sim} \Delta_{-/A'}$$

on A'/d -algebras ([BL22, Theorem 5.6.2]), so the construction

$$S \mapsto (\Delta_{S/A}[\frac{1}{d}]/p^n)^{\phi_i=1}$$

from p -complete p -quasisyntomic A'/d -algebras to $\mathcal{D}(\mathbb{Z}/p^n)$ is an arc-sheaf by the previous argument. The same construction from p -complete p -quasisyntomic A/d -algebras S to $\mathcal{D}(\mathbb{Z}/p^n)$ is a p -quasisyntomic sheaf. There exists a p -quasisyntomic cover $S \rightarrow S'$, where S' has a structure of A'/d -algebra – consider for instance the map

$$S \longrightarrow S' := S \widehat{\otimes}_{A/d} A'/d,$$

which is a p -quasisyntomic cover by [BMS19, Lemma 4.16 (2)] – so it is also an arc-sheaf.

We now construct a natural map

$$R\Gamma_{\text{ét}}(\text{Spec}(S[\frac{1}{p}]), \mu_{p^n}^{\otimes i}) \longrightarrow (\Delta_{S/A}[\frac{1}{d}]/p^n)^{\phi_i=1}$$

for all p -complete p -quasisyntomic A/d -algebras S . By the previous paragraph, it suffices to do so locally in the arc-topology, *e.g.*, for all perfectoid A/d -algebras S ([BS22, Lemma 8.8]). The construction

$$S \mapsto R\Gamma_{\text{ét}}(\text{Spec}(S[\frac{1}{p}]), \mu_{p^n}^{\otimes i})$$

is discrete locally in the arc-topology: any perfectoid ring S has an arc-cover by an absolutely integrally closed perfectoid ring S' ([BS22, Theorem 7.14]), whose generic fibre has étale cohomology concentrated in degree 0 ([Sta19, Proposition 59.81.5]). So it suffices to construct a natural map

$$\mu_{p^n}^{\otimes i}(S[\frac{1}{p}]) \mapsto H^0(\Delta_{S/A}[\frac{1}{d}]/p^n)^{\phi_i=1}$$

for perfectoid A/d -algebras S ; then one can compose it with

$$H^0(\Delta_{S/A}[\frac{1}{d}]/p^n)^{\phi_i=1} \longrightarrow (\Delta_{S/A}[\frac{1}{d}]/p^n)^{\phi_i=1},$$

and arc-sheafify the construction. For $i = 0$, this map is constructed in the proof of [BS22, Theorem 9.1]. Using the multiplicative structure on both sides, it then suffices to construct this map for $i = 1$. There is a canonical identification

$$\mu_{p^n}(S) \cong H^0(\Delta_{S/A}/p^n)^{\phi_1=1}$$

given by the prismatic first Chern class for any p -complete p -quasisyntomic A/d -algebra S ([BL22, Theorem 7.5.6]), and thus a natural map

$$\mu_{p^n}(S) \cong H^0(\Delta_{S/A}/p^n)^{\phi_1=1} \longrightarrow H^0(\Delta_{S/A}[\frac{1}{d}]/p^n)^{\phi_1=1}$$

for all p -complete p -quasisyntomic A/d -algebras S . The canonical map

$$\mu_{p^n}(S) \longrightarrow \mu_{p^n}(S[\frac{1}{p}])$$

is a group isomorphism for any perfectoid ring S (Lemma 4.3), hence we get a natural map

$$\mu_{p^n}(S[\frac{1}{p}]) \longrightarrow H^0(\Delta_{S/A}[\frac{1}{d}]/p^n)^{\phi_1=1}$$

for all perfectoid A/d -algebras S .

We finally prove that the natural map

$$R\Gamma_{\text{ét}}(\text{Spec}(S[\frac{1}{p}]), \mu_{p^n}^{\otimes i}) \longrightarrow (\Delta_{S/A}[\frac{1}{d}]/p^n)^{\phi_i=1}$$

is an equivalence in the derived category $\mathcal{D}(\mathbb{Z}/p^n)$. Using arc-localisation, one can assume that S is an absolutely integrally closed valuation ring of rank ≤ 1 , in which case $R\Gamma_{\text{ét}}(\text{Spec}(S[\frac{1}{p}]), \mu_{p^n}^{\otimes i})$ is concentrated in degree 0, and so is $(\Delta_{S/A}[\frac{1}{d}]/p^n)^{\phi_i=1}$ by Artin–Scheier–Witt theory. If the valuation ring S has characteristic p , then both sides of the morphism are zero. Otherwise, the valuation ring S is p -torsionfree and admits a compatible system of p -power roots of unity. Multiplying the morphism by μ^{-i} then reduces to the case of $i = 0$, which is [BS22, Theorem 9.1]. \square

4.2 The syntomic-étale comparison theorem in characteristic p

In this subsection we review the description of the syntomic complexes in characteristic p in terms of logarithmic de Rham–Witt forms (following [KM21, LM21]), and its consequence on the syntomic-étale comparison map (Construction 4.2).

Proposition 4.6 (Syntomic-étale comparison theorem in characteristic p). *Let R be perfect \mathbb{F}_p -algebra, and S a Cartier smooth R -algebra. Then for any integers $i \geq 0$ and $n \geq 1$, there are canonical identifications*

$$\mathbb{Z}/p^n(i)^{\text{syn}}(S) \simeq R\Gamma_{\text{ét}}(\text{Spec}(S), W_n\Omega_{\log}^i)[-i]$$

and

$$\mathbb{Z}_p(i)^{\text{syn}}(S) \simeq R\Gamma_{\text{proét}}(\text{Spec}(S), W\Omega_{\log}^i)[-i].$$

Proof. The first equivalence follows from [LM21, Proposition 5.1 (ii)], the second by taking limits over $n \geq 1$. \square

Corollary 4.7. *Let R be a perfect \mathbb{F}_p -algebra, and S a Cartier smooth R -algebra. Then for any integers $i \geq 0$ and $n \geq 1$, the comparison maps*

$$\gamma_{\text{syn}}^{\text{ét}}\{i\}/p^n : \mathbb{Z}/p^n(i)^{\text{syn}}(S) \longrightarrow R\Gamma_{\text{ét}}(\text{Spec}(S[\frac{1}{p}]), \mu_{p^n}^{\otimes i})$$

and

$$\gamma_{\text{syn}}^{\text{ét}}\{i\} : \mathbb{Z}_p(i)^{\text{syn}}(S) \longrightarrow R\Gamma_{\text{proét}}(\text{Spec}(S[\frac{1}{p}]), \mathbb{Z}_p(i))$$

have homotopy cofibres in degrees $\geq i - 1$.

Proof. The syntomic complexes $\mathbb{Z}/p^n(i)^{\text{syn}}(S)$ and $\mathbb{Z}_p(i)^{\text{syn}}(S)$ are in degrees $\geq i$ by Proposition 4.6, and the generic fibre $S[\frac{1}{p}]$ of S is zero. \square

4.3 The syntomic-étale comparison theorem

In this subsection we prove the syntomic-étale comparison theorem over a general perfectoid base ring (Theorem 4.12), generalising Corollary 4.7.

The following result is a direct consequence of Theorem 2.16 (\mathcal{N}^{\geq}) when the base prism is perfect.

Lemma 4.8. *Let $(A, (d))$ be a perfect prism, and S a p -Cartier smooth A/d -algebra. Then for every integer $i \geq 0$, the map*

$$\tau^{\leq i}\mathcal{N}^{\geq i}\Delta_{S/A} \xrightarrow{\phi_i} \tau^{\leq i}\Delta_{S/A}$$

is an equivalence.

Proof. The diagram

$$\begin{array}{ccc} \mathcal{N}^{\geq i}\Delta_{S/A} & \xrightarrow{\phi_i} & \Delta_{S/A} \\ & \searrow \phi & \downarrow d^i \\ & & d^i\Delta_{S/A} \end{array}$$

is commutative for any p -quasisyntomic A/d -algebra S and the map $d^i : \Delta_{S/A} \rightarrow d^i \Delta_{S/A}$ is an equivalence. Locally on the p -quasisyntomic site $\Delta_{S/A}$ is a d -torsionfree A -module in degree 0 and this is by definition of the divided Frobenius map ϕ_i , and in general this is true by descent on the p -quasisyntomic site. When S is p -Cartier smooth over A/d , the Frobenius map $\phi : \mathcal{N}^{\geq i} \Delta_{S/A} \rightarrow d^i \Delta_{S/A}$ is an isomorphism in degrees $\leq i$, hence the result. \square

Lemma 4.9. *Let A be a commutative ring, $i \in \mathbb{Z}$ an integer, $f : C \rightarrow C'$ and $g : D \rightarrow D'$ two maps of p -complete objects in the derived category $\mathcal{D}(A)$, and*

$$\begin{array}{ccc} \tau^{\leq i} C & \longrightarrow & \tau^{\leq i} D \\ \downarrow \tau^{\leq i} f & & \downarrow \tau^{\leq i} g \\ \tau^{\leq i} C' & \longrightarrow & \tau^{\leq i} D' \end{array}$$

a commutative diagram. If the map

$$\mathrm{hocolim}(\tau^{\leq i-1}(f/p)) \longrightarrow \mathrm{hocolim}(\tau^{\leq i-1}(g/p)),$$

induced by reduction modulo p of the canonical map $\mathrm{hocolim}(\tau^{\leq i} f) \rightarrow \mathrm{hocolim}(\tau^{\leq i} g)$, is an isomorphism in degrees $\leq i-2$, then the canonical map $\mathrm{hocolim}(\tau^{\leq i} f) \rightarrow \mathrm{hocolim}(\tau^{\leq i} g)$ is an isomorphism in degrees $\leq i-3$ and is injective in degree $i-2$.

Proof. Let $h : E \rightarrow E'$ be either $f : C \rightarrow C'$ or $g : D \rightarrow D'$. For any $j \leq i-3$, the cohomology of $\mathrm{hocolim}(\tau^{\leq i-1} h/p)$ in degree j is naturally identified with that of $\mathrm{hocolim}(\tau^{\leq i-1}(h/p))$:

$$\begin{aligned} \mathrm{H}^j(\mathrm{hocolim}(\tau^{\leq i-1} h/p)) &\cong \mathrm{H}^j(\mathrm{hocolim}(\tau^{\leq i-2}(\tau^{\leq i-1} h/p))) \\ &\cong \mathrm{H}^j(\mathrm{hocolim}(\tau^{\leq i-2}(h/p))) \\ &\cong \mathrm{H}^j(\mathrm{hocolim}(\tau^{\leq i-1}(h/p))). \end{aligned}$$

If the map

$$\mathrm{hocolim}(\tau^{\leq i-1}(f/p)) \longrightarrow \mathrm{hocolim}(\tau^{\leq i-1}(g/p))$$

is an isomorphism in degrees $\leq i-3$, then so is the canonical map

$$\mathrm{hocolim}(\tau^{\leq i} f) \longrightarrow \mathrm{hocolim}(\tau^{\leq i} g)$$

by derived Nakayama.

In degree $j = i-2$, the cohomology group of $\mathrm{hocolim}(\tau^{\leq i-1} h/p)$ naturally embeds into that of $\mathrm{hocolim}(\tau^{\leq i-1}(h/p))$ because of the commutative diagram

$$\begin{array}{ccccccc} \mathrm{H}^{i-2}(\tau^{\leq i-1} E/p) & \rightarrow & \mathrm{H}^{i-2}(\tau^{\leq i-1} E'/p) & \rightarrow & \mathrm{H}^{i-2}(\mathrm{hocolim}(\tau^{\leq i-1} h)/p) & \rightarrow & \mathrm{H}^{i-1}(\tau^{\leq i-1} E/p) \\ \downarrow \cong & & \downarrow \cong & & \downarrow & & \downarrow \\ \mathrm{H}^{i-2}(\tau^{\leq i-1}(E/p)) & \rightarrow & \mathrm{H}^{i-2}(\tau^{\leq i-1}(E'/p)) & \rightarrow & \mathrm{H}^{i-2}(\mathrm{hocolim}(\tau^{\leq i-1}(h/p))) & \rightarrow & \mathrm{H}^{i-1}(\tau^{\leq i-1}(E/p)) \end{array}$$

where the two horizontal lines are exact sequences. If the map

$$\mathrm{hocolim}(\tau^{\leq i-1}(f/p)) \longrightarrow \mathrm{hocolim}(\tau^{\leq i-1}(g/p))$$

is an isomorphism in degree $i-2$, then the canonical map

$$\mathrm{hocolim}(\tau^{\leq i} f) \longrightarrow \mathrm{hocolim}(\tau^{\leq i} g)$$

is injective in degree $i-2$ by derived Nakayama. \square

Lemma 4.10. *Let $(A, (d))$ be a perfect prism, and S a p -Cartier smooth A/d -algebra. For all integers $i \geq 0$ and $k \leq i - 1$, define the map*

$$\phi_i^{-1} : \mathbf{H}^k(\Delta_{S/A}/p) \longrightarrow \mathbf{H}^k(\Delta_{S/A}/p)$$

as the composite

$$\mathbf{H}^k(\Delta_{S/A}/p) \xrightarrow{\phi_i^{-1}} \mathbf{H}^k((\mathcal{N}^{\geq i} \Delta_{S/A})/p) \xrightarrow{\text{can}} \mathbf{H}^k(\Delta_{S/A}/p)$$

where the first map is given by the derived reduction modulo p of Lemma 4.8. Then for every $k \leq i - 2$, the map ϕ_i^{-1} is zero on the d -torsion subgroup $\mathbf{H}^k(\Delta_{S/A}/p)[d]$. For $k = i - 1$, the map $\phi_i^{-1} \circ \phi_i^{-1}$ is zero on the d -torsion subgroup $\mathbf{H}^{i-1}(\Delta_{S/A}/p)[d]$.

Proof. First assume that $k \leq i - 2$. In this case, the map

$$\phi_i^{-1} : \mathbf{H}^k(\Delta_{S/A}/p) \longrightarrow \mathbf{H}^k(\Delta_{S/A}/p)$$

can be rewritten as the map $d^{1/p} \phi_{i-1}^{-1}$, where ϕ_{i-1}^{-1} is defined as ϕ_i^{-1} . Let x be a d -torsion element of the A/p -module $\mathbf{H}^k(\Delta_{S/A}/p)$. Then

$$\phi_i^{-1}(x) = d^{1/p} \phi_{i-1}^{-1}(x) = \phi_{i-1}^{-1}(dx) = 0,$$

hence the result.

Now assume that $k = i - 1$, and let us prove that $(\phi_i^{-1})^2 := \phi_i^{-1} \circ \phi_i^{-1}$ is the zero map on the A/p -module $\mathbf{H}^{i-1}(\Delta_{S/A}/p)[d]$. There is a natural short exact sequence of A/p -modules

$$0 \longrightarrow \mathbf{H}^{i-1}(\Delta_{S/A})/p \longrightarrow \mathbf{H}^{i-1}(\Delta_{S/A}/p) \longrightarrow \mathbf{H}^i(\Delta_{S/A})[p] \longrightarrow 0,$$

and compatible maps ϕ_i^{-1} on each of its terms. This short exact sequence induces, by the snake lemma, an exact sequence of A/p -modules

$$0 \longrightarrow (\mathbf{H}^{i-1}(\Delta_{S/A})/p)[d] \longrightarrow \mathbf{H}^{i-1}(\Delta_{S/A}/p)[d] \longrightarrow (\mathbf{H}^i(\Delta_{S/A})[p])[d],$$

and the maps ϕ_i^{-1} restrict to these A/p -modules. It suffices to prove that the map ϕ_i^{-1} is zero on both $(\mathbf{H}^{i-1}(\Delta_{S/A})/p)[d]$ and $(\mathbf{H}^i(\Delta_{S/A})[p])[d]$. Indeed, if x is an element of $\mathbf{H}^{i-1}(\Delta_{S/A}/p)[d]$, then $\phi_i^{-1}(x)$ is naturally in the kernel of the canonical map

$$\mathbf{H}^{i-1}(\Delta_{S/A}/p)[d] \longrightarrow (\mathbf{H}^i(\Delta_{S/A})[p])[d].$$

So $\phi_i^{-1}(x)$ is in the subgroup $(\mathbf{H}^{i-1}(\Delta_{S/A})/p)[d] \subseteq \mathbf{H}^{i-1}(\Delta_{S/A}/p)[d]$, and thus $\phi_i^{-1}(\phi_i^{-1}(x)) = 0$.

On the A/p -module $\mathbf{H}^{i-1}(\Delta_{S/A})/p$, the map ϕ_i^{-1} can be rewritten as the map $d^{1/p} \phi_{i-1}^{-1}$, and is thus zero on $(\mathbf{H}^{i-1}(\Delta_{S/A})/p)[d]$, arguing as in the first paragraph of this proof. We now prove that the map ϕ_i^{-1} is zero on the A/p -module $(\mathbf{H}^i(\Delta_{S/A})[p])[d]$. By definition of the map ϕ_i^{-1} , it is equivalent to prove that the canonical map

$$\text{can} : \mathbf{H}^i(\mathcal{N}^{\geq i} \Delta_{S/A})[p] \longrightarrow \mathbf{H}^i(\Delta_{S/A})[p]$$

is zero on the $d^{1/p}$ -torsion subgroup $(\mathbf{H}^i(\mathcal{N}^{\geq i} \Delta_{S/A})[p])[d^{1/p}]$. By Theorem 2.16, there is a commutative diagram

$$\begin{array}{ccc} \mathbf{H}^i(\mathcal{N}^{\geq i} \Delta_{S/A})[p] & \xrightarrow[\cong]{\phi} & \mathbf{H}^i(d^i \Delta_{S/A})[p] \\ \downarrow \text{can} & & \downarrow \text{can} \\ \mathbf{H}^i(\Delta_{S/A})[p] & \xrightarrow[\cong]{\phi} & \mathbf{H}^i(L\eta_d \Delta_{S/A})[p]. \end{array}$$

By Proposition 2.11 (1), the right vertical map of this diagram is surjective, with kernel given by the d -torsion subgroup of $\mathbf{H}^i(d^i \Delta_{S/A})[p]$. So the left vertical map is also surjective, with kernel given by the $d^{1/p}$ -torsion subgroup $(\mathbf{H}^i(\mathcal{N}^{\geq i} \Delta_{S/A})[p])[d^{1/p}]$. Hence the result. \square

Remark 4.11. In the previous lemma, the result can be slightly improved if the perfectoid base ring A/d is p -torsionfree. In this case, the map ϕ_i^{-1} is zero on $H^k(\Delta_{S/A}/p)[d]$, for any integer $k \leq i-1$; see the proof of Theorem 4.14 below.

Theorem 4.12 (Syntomic-étale comparison theorem). *Let R be a perfectoid ring, and S a p -Cartier smooth R -algebra. Then for any integers $i \geq 0$ and $n \geq 1$, the homotopy cofibres of the maps*

$$\gamma_{\text{syn}}^{\text{ét}}\{i\}/p^n : \mathbb{Z}/p^n(i)^{\text{syn}}(S) \longrightarrow R\Gamma_{\text{ét}}(\text{Spec}(S[\frac{1}{p}]), \mu_{p^n}^{\otimes i})$$

and

$$\gamma_{\text{syn}}^{\text{ét}}\{i\} : \mathbb{Z}_p(i)^{\text{syn}}(S) \longrightarrow R\Gamma_{\text{proét}}(\text{Spec}(S[\frac{1}{p}]), \mathbb{Z}_p(i))$$

are in degrees $\geq i-1$.

Proof. We prove the result for integral coefficients; reducing modulo p^n implies the result for torsion coefficients. Let $i \geq 0$ be an integer, $(A, (d))$ the perfect prism corresponding to the perfectoid ring R and $R\Phi(S, \mathbb{Z}_p(i)) \in \mathcal{D}(\mathbb{Z}_p)$ the homotopy cofibre of the comparison map

$$\gamma_{\text{syn}}^{\text{ét}}\{i\} : \mathbb{Z}_p(i)^{\text{syn}}(S) \longrightarrow R\Gamma_{\text{proét}}(\text{Spec}(S[\frac{1}{p}]), \mathbb{Z}_p(i)).$$

By Theorem 4.5 and Remark 4.4, there is a commutative diagram

$$\begin{array}{ccccc} \mathbb{Z}_p(i)^{\text{syn}}(S) & \longrightarrow & \mathcal{N}^{\geq i} \Delta_{S/A} & \xrightarrow{\phi_i^{-1}} & \Delta_{S/A} \\ \downarrow & & \downarrow \lambda_i & & \downarrow \lambda \\ R\Gamma_{\text{proét}}(\text{Spec}(S[\frac{1}{p}]), \mathbb{Z}_p(i)) & \longrightarrow & \Delta_{S/A}[\frac{1}{d}]_p^\wedge & \xrightarrow{\phi_i^{-1}} & \Delta_{S/A}[\frac{1}{d}]_p^\wedge \\ \downarrow & & \downarrow & & \downarrow \\ R\Phi(S, \mathbb{Z}_p(i)) & \longrightarrow & \text{hocofib}(\lambda_i) & \xrightarrow{\phi_i^{-1}} & \text{hocofib}(\lambda) \end{array}$$

where all the horizontal maps are homotopy fibre sequences. The maps λ_i and λ correspond to inverting d and then p -adically completing.

We want to prove that $R\Phi(S, \mathbb{Z}_p(i)) \in \mathcal{D}^{\geq i-1}(\mathbb{Z}_p)$, i.e., that $R\Phi(S, \mathbb{Z}_p(i))$ is zero in degrees $\leq i-2$. This statement depends only on

$$\tau^{\leq i-2} \text{hocofib}(\lambda_i) \xrightarrow{\phi_i^{-1}} \tau^{\leq i-2} \text{hocofib}(\lambda),$$

and thus only on the commutative diagram

$$\begin{array}{ccc} \tau^{\leq i} \mathcal{N}^{\geq i} \Delta_{S/A} & \xrightarrow{\phi_i^{-1}} & \tau^{\leq i} \Delta_{S/A} \\ \downarrow \tau^{\leq i} \lambda_i & & \downarrow \tau^{\leq i} \lambda \\ \tau^{\leq i} (\Delta_{S/A}[\frac{1}{d}]_p^\wedge) & \xrightarrow{\phi_i^{-1}} & \tau^{\leq i} (\Delta_{S/A}[\frac{1}{d}]_p^\wedge). \end{array}$$

In terms of this diagram, we want to prove that the map

$$\text{hocofib}(\tau^{\leq i} \lambda_i) \xrightarrow{\phi_i^{-1}} \text{hocofib}(\tau^{\leq i} \lambda)$$

is an isomorphism in degrees $\leq i-3$ and is injective in degree $i-2$.

By Lemma 4.8, the divided Frobenius map $\phi_i : \mathcal{N}^{\geq i} \Delta_{S/A} \rightarrow \Delta_{S/A}$ is an isomorphism in degrees $\leq i$. Define

$$1 - \phi_i^{-1} : \tau^{\leq i} \Delta_{S/A} \longrightarrow \tau^{\leq i} \Delta_{S/A}$$

as the map $\phi_i - 1 : \tau^{\leq i} \mathcal{N}^{\geq i} \Delta_{S/A} \rightarrow \tau^{\leq i} \Delta_{S/A}$ precomposed with the inverse of the divided Frobenius map $\phi_i : \tau^{\leq i} \mathcal{N}^{\geq i} \Delta_{S/A} \xrightarrow{\sim} \tau^{\leq i} \Delta_{S/A}$. Similarly the divided Frobenius map $\phi_i : \Delta_{S/A}[\frac{1}{d}]_p^\wedge \rightarrow \Delta_{S/A}[\frac{1}{d}]_p^\wedge$ is an equivalence by Theorem 2.16 ($L\eta$), and we define

$$1 - \phi_i^{-1} : \tau^{\leq i} (\Delta_{S/A}[\frac{1}{d}]_p^\wedge) \longrightarrow \tau^{\leq i} (\Delta_{S/A}[\frac{1}{d}]_p^\wedge)$$

as the map $\phi_i - 1 : \tau^{\leq i}(\Delta_{S/A}[\frac{1}{d}]_p^\wedge) \rightarrow \tau^{\leq i}(\Delta_{S/A}[\frac{1}{d}]_p^\wedge)$ precomposed with the inverse of the divided Frobenius map $\phi_i : \tau^{\leq i}(\Delta_{S/A}[\frac{1}{d}]_p^\wedge) \xrightarrow{\sim} \tau^{\leq i}(\Delta_{S/A}[\frac{1}{d}]_p^\wedge)$. We thus want to prove that the induced map

$$\text{hocofib}(\tau^{\leq i} \lambda) \xrightarrow{1 - \phi_i^{-1}} \text{hocofib}(\tau^{\leq i} \lambda)$$

is an isomorphism in degrees $\leq i - 3$ and is injective in degree $i - 2$. By Lemma 4.9, it suffices to prove that

$$\text{hocofib}(\tau^{\leq i-1}(\lambda/p)) \xrightarrow{1 - \phi_i^{-1}} \text{hocofib}(\tau^{\leq i-1}(\lambda/p))$$

is an isomorphism in degrees $\leq i - 2$.

Let k be an integer $\leq i - 2$. There is a short exact sequence of A/p -modules:

$$0 \longrightarrow \text{Coker}(\mathbf{H}^k(\lambda/p)) \longrightarrow \mathbf{H}^k(\text{hocofib}(\lambda/p)) \longrightarrow \text{Ker}(\mathbf{H}^{k+1}(\lambda/p)) \longrightarrow 0.$$

To prove that $1 - \phi_i^{-1}$ is an isomorphism on $\mathbf{H}^k(\text{hocofib}(\lambda/p))$, let us prove that it is an isomorphism on both $\text{Ker}(\mathbf{H}^{k+1}(\lambda/p))$ and $\text{Coker}(\mathbf{H}^k(\lambda/p))$.

First consider the map

$$\mathbf{H}^{k+1}(\lambda/p) : \mathbf{H}^{k+1}(\Delta_{S/A}/p) \longrightarrow \mathbf{H}^{k+1}(\Delta_{S/A}/p)[\frac{1}{d}].$$

Its kernel is given by the d -power torsion subgroup of $\mathbf{H}^{k+1}(\Delta_{S/A}/p)$, so we want to prove that the map $1 - \phi_i^{-1}$ is an isomorphism on the A/p -module $\mathbf{H}^{k+1}(\Delta_{S/A}/p)[d^\infty]$. The relation $\phi_A(d) = d^p$ holds in the ring A/p . So for any integer $j \geq 1$ and any element $x \in \mathbf{H}^{k+1}(\Delta_{S/A}/p)[d^{p^j}]$:

$$d^{p^{j-1}} \phi_i^{-1}(x) = \phi_i^{-1}(d^{p^j} x) = 0,$$

and thus $\phi_i^{-1}(x) \in \mathbf{H}^{k+1}(\Delta_{S/A}/p)[d^{p^{j-1}}]$. By Lemma 4.10, the map $(\phi_i^{-1})^2$ is zero on the A/p -module $\mathbf{H}^{k+1}(\Delta_{S/A}/p)[d]$. So the map $1 - \phi_i^{-1}$ is an isomorphism on the A/p -module $\mathbf{H}^{k+1}(\Delta_{S/A}/p)[d^{p^j}]$, for each integer $j \geq 1$, with inverse given by the map

$$1 + \phi_i^{-1} + \dots + (\phi_i^{-1})^{j+1}.$$

So the map $1 - \phi_i^{-1}$ is an isomorphism on the A/p -module $\mathbf{H}^{k+1}(\Delta_{S/A}/p)[d^\infty]$.

Now consider the map

$$\mathbf{H}^k(\lambda/p) : \mathbf{H}^k(\Delta_{S/A}/p) \longrightarrow \mathbf{H}^k(\Delta_{S/A}/p)[\frac{1}{d}].$$

It is the filtered colimit over $m \geq 0$ of the maps

$$\lambda^{(m)}/p : \Delta_{S/A}/p \longrightarrow \Delta_{S/A}/p$$

where $\lambda^{(m)} : \Delta_{S/A} \rightarrow \Delta_{S/A}$ is the multiplication by d^m . The A/p -module $\text{Coker}(\mathbf{H}^k(\lambda/p))$ and the map $1 - \phi_i^{-1}$ acting on it can be rewritten as the colimit over $m \geq 0$ of the A/p -modules $\mathbf{H}^k(\Delta_{S/A}/p)/d^m$ with maps $1 - d^{\frac{(p-1)m}{p}} \phi_i^{-1}$. Let $m \geq 0$ be an integer. We claim that the map

$$d^{\frac{(p-1)m}{p}} \phi_i^{-1} : \mathbf{H}^k(\Delta_{S/A}/p)/d^m \longrightarrow \mathbf{H}^k(\Delta_{S/A}/p)/d^m$$

is nilpotent. Because $k \leq i - 2$, this map $d^{\frac{(p-1)m}{p}} \phi_i^{-1}$ is naturally identified with the map

$$d^{\frac{(p-1)m}{p} + \frac{1}{p}} \phi_{i-1}^{-1} : \mathbf{H}^k(\Delta_{S/A}/p)/d^m \longrightarrow \mathbf{H}^k(\Delta_{S/A}/p)/d^m,$$

where

$$\phi_{i-1} : \tau^{\leq i-1} \mathcal{N}^{\geq i-1} \Delta_{S/A} \longrightarrow \tau^{\leq i-1} \Delta_{S/A}$$

is the equivalence of Lemma 4.8. Composing with itself k times for some integer $k \geq 1$ and using the ϕ_A^{-1} -linearity of ϕ_{i-1}^{-1} gives the map $d^{\frac{(p^k-1)m}{p^k} + \frac{p^k-1}{p^k(p-1)}} (\phi_{i-1}^{-1})^k$. This map is zero for any integer $k \geq 1$ satisfying $\frac{(p^k-1)m}{p^k} + \frac{p^k-1}{p^k(p-1)} \geq m$, that is $p^k \geq m(p-1) + 1$. The map $1 - d^{\frac{(p-1)m}{p}} \phi_i^{-1}$ is thus a sum of an isomorphism and a nilpotent map, so it is an equivalence. Taking colimit over $m \geq 0$, the map $1 - \phi_i^{-1}$ is an isomorphism on $\text{Coker}(\mathbf{H}^k(\lambda/p))$. Hence the result. \square

4.4 The syntomic-étale comparison theorem over a p -torsionfree base

In this subsection we prove a refined version of the syntomic-étale comparison theorem (Theorem 4.14), assuming the perfectoid base ring is p -torsionfree.

Lemma 4.13. *Let $(A, (d))$ be a perfect prism such that A/d is p -torsionfree, and S a p -Cartier smooth A/d -algebra. Then for every integer $i \geq 0$, the Frobenius maps and divided Frobenius map*

$$\begin{aligned} \phi &: \mathbb{A}_{S/A}/p \longrightarrow L\eta_d(\mathbb{A}_{S/A}/p) \\ \phi &: \tau^{\leq i}(\mathcal{N}^{\geq i}\mathbb{A}_{S/A}/p) \longrightarrow \tau^{\leq i}(d^i\mathbb{A}_{S/A}/p) \\ \phi_i &: \tau^{\leq i}(\mathcal{N}^{\geq i}\mathbb{A}_{S/A}/p) \longrightarrow \tau^{\leq i}(\mathbb{A}_{S/A}/p) \end{aligned}$$

are equivalences.

Proof. By Theorem 2.16 ($L\eta$), the Frobenius map

$$\phi : \mathbb{A}_{S/A} \longrightarrow L\eta_d\mathbb{A}_{S/A}$$

is an equivalence, thus so is its derived reduction modulo p . Note that p is a nonzerodivisor in the ring A ([BS22, Lemma 2.28 (1)]). Moreover, S is p -quasismooth over A/d , so the groups $(\Omega_{S/(A/d)}^n)_p^\wedge$ are p -flat modules over the p -torsionfree ring A/d and are in particular p -torsionfree. The groups $\mathrm{H}^n(\mathbb{A}_{S/A})$ are thus also p -torsionfree (Proposition 2.14 (3)), and the natural map

$$(L\eta_d\mathbb{A}_{S/A})/p \longrightarrow L\eta_d(\mathbb{A}_{S/A}/p)$$

is an equivalence in the derived category $\mathcal{D}(A/p)$ ([Bha18, Lemma 5.16]), which proves the first statement.

The proof of Theorem 2.16 ($L\eta$) \Rightarrow (\mathcal{N}^{\geq}), where we use that

$$\mathrm{H}^0(\mathbb{A}_{S/A}/p) = \lim_{(B, (d)B) \in (S/A)_\Delta} B/p$$

is d -torsionfree, then adapts readily to prove that the Frobenius map

$$\phi : \tau^{\leq i}(\mathcal{N}^{\geq i}\mathbb{A}_{S/A}/p) \longrightarrow \tau^{\leq i}(d^i\mathbb{A}_{S/A}/p)$$

is an equivalence in the derived category $\mathcal{D}(A/p)$. The proof of the third statement is the same as in Lemma 4.8. \square

Theorem 4.14 (Syntomic-étale comparison theorem over a p -torsionfree base). *Let R be a p -torsionfree perfectoid ring, and S a p -Cartier smooth R -algebra. Then for any integers $i \geq 0$ and $n \geq 1$, the homotopy cofibres of the maps*

$$\gamma_{\mathrm{syn}}^{\acute{\mathrm{e}}\mathrm{t}}\{i\}/p^n : \mathbb{Z}/p^n(i)^{\mathrm{syn}}(S) \longrightarrow R\Gamma_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(S[\frac{1}{p}]), \mu_{p^n}^{\otimes i})$$

and

$$\gamma_{\mathrm{syn}}^{\acute{\mathrm{e}}\mathrm{t}}\{i\} : \mathbb{Z}_p(i)^{\mathrm{syn}}(S) \longrightarrow R\Gamma_{\mathrm{pro}\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(S[\frac{1}{p}]), \mathbb{Z}_p(i))$$

are in degrees $\geq i$.

Proof. We keep the same notation as in the proof of Theorem 4.12. We want to prove that

$$R\Phi(S, \mathbb{Z}_p(i)) \in \mathcal{D}^{\geq i}(\mathbb{Z}_p),$$

i.e., that $R\Phi(S, \mathbb{Z}_p(i))$ is zero in degrees $\leq i - 1$. By derived Nakayama this is equivalent to proving that $R\Phi(S, \mathbb{Z}_p(i))/p$ is zero in degrees $\leq i - 1$. This statement depends only on

$$\tau^{\leq i-1}\mathrm{hocofib}(\lambda_i/p) \xrightarrow{\phi_{i-1}} \tau^{\leq i-1}\mathrm{hocofib}(\lambda/p),$$

and thus only on the commutative diagram

$$\begin{array}{ccc} \tau^{\leq i}(\mathcal{N}^{\geq i}\Delta_{S/A}/p) & \xrightarrow{\phi_i^{-1}} & \tau^{\leq i}(\Delta_{S/A}/p) \\ \downarrow \tau^{\leq i}(\lambda_i/p) & & \downarrow \tau^{\leq i}(\lambda/p) \\ \tau^{\leq i}(\Delta_{S/A}[\frac{1}{d}]/p) & \xrightarrow{\phi_i^{-1}} & \tau^{\leq i}(\Delta_{S/A}[\frac{1}{d}]/p). \end{array}$$

In terms of this diagram, we want to prove that the map

$$\mathrm{hocofib}(\tau^{\leq i}(\lambda_i/p)) \xrightarrow{\phi_i^{-1}} \mathrm{hocofib}(\tau^{\leq i}(\lambda/p))$$

is an isomorphism in degrees $\leq i-2$ and is injective in degree $i-1$.

The divided Frobenius map $\phi_i : \mathcal{N}^{\geq i}\Delta_{S/A}/p \rightarrow \Delta_{S/A}/p$ is an isomorphism in degrees $\leq i$ by Lemma 4.13. Define

$$1 - \phi_i^{-1} : \tau^{\leq i}(\Delta_{S/A}/p) \rightarrow \tau^{\leq i}(\Delta_{S/A}/p)$$

as the map $\phi_i - 1 : \tau^{\leq i}(\mathcal{N}^{\geq i}\Delta_{S/A}/p) \rightarrow \tau^{\leq i}(\Delta_{S/A}/p)$ precomposed with the inverse of the divided Frobenius map $\phi_i : \tau^{\leq i}(\mathcal{N}^{\geq i}\Delta_{S/A}/p) \xrightarrow{\sim} \tau^{\leq i}(\Delta_{S/A}/p)$. Similarly the divided Frobenius map $\phi_i : \Delta_{S/A}[\frac{1}{d}]/p \rightarrow \Delta_{S/A}[\frac{1}{d}]/p$ is an equivalence (Theorem 2.16 ($L\eta$)), and we define

$$1 - \phi_i^{-1} : \tau^{\leq i}(\Delta_{S/A}[\frac{1}{d}]/p) \rightarrow \tau^{\leq i}(\Delta_{S/A}[\frac{1}{d}]/p)$$

as the map $\phi_i - 1 : \tau^{\leq i}(\Delta_{S/A}[\frac{1}{d}]/p) \rightarrow \tau^{\leq i}(\Delta_{S/A}[\frac{1}{d}]/p)$ precomposed with the inverse of the divided Frobenius map $\phi_i : \tau^{\leq i}(\Delta_{S/A}[\frac{1}{d}]/p) \xrightarrow{\sim} \tau^{\leq i}(\Delta_{S/A}[\frac{1}{d}]/p)$. We thus want to prove that the induced map

$$\mathrm{hocofib}(\tau^{\leq i}(\lambda/p)) \xrightarrow{1 - \phi_i^{-1}} \mathrm{hocofib}(\tau^{\leq i}(\lambda/p))$$

is an isomorphism in degrees $\leq i-2$ and is injective in degree $i-1$. We prove that it is an isomorphism in degrees $\leq i-1$.

Let k be an integer $\leq i-1$. There is a short exact sequence of A/p -modules:

$$0 \rightarrow \mathrm{Coker}(\mathrm{H}^k(\lambda/p)) \rightarrow \mathrm{H}^k(\mathrm{hocofib}(\lambda/p)) \rightarrow \mathrm{Ker}(\mathrm{H}^{k+1}(\lambda/p)) \rightarrow 0.$$

To prove that $1 - \phi_i^{-1}$ is an isomorphism on $\mathrm{H}^k(\mathrm{hocofib}(\lambda/p))$, let us prove that it is an isomorphism on both $\mathrm{Coker}(\mathrm{H}^k(\lambda/p))$ and $\mathrm{Ker}(\mathrm{H}^{k+1}(\lambda/p))$.

For $\mathrm{Coker}(\mathrm{H}^k(\lambda/p))$, the argument is the same as in the proof of Theorem 4.12, where we need Lemma 4.13 for the case of $k = i-1$.

For $\mathrm{Ker}(\mathrm{H}^{k+1}(\lambda/p))$, we follow the lines of the proof of Theorem 4.12. It suffices to prove the case of $k = i-1$, *i.e.*, that $1 - \phi_i^{-1}$ is an isomorphism on the A/p -module $\mathrm{H}^i(\Delta_{S/A}/p)[d^\infty]$. It then suffices to prove that ϕ_i^{-1} is nilpotent on the A/p -module $\mathrm{H}^i(\Delta_{S/A}/p)[d]$; we prove that it is zero. By definition of the map ϕ_i^{-1} , it is equivalent to proving that the canonical map

$$\mathrm{can} : \mathrm{H}^i((\mathcal{N}^{\geq i}\Delta_{S/A})/p) \rightarrow \mathrm{H}^i(\Delta_{S/A}/p)$$

is zero on the $d^{1/p}$ -torsion subgroup $\mathrm{H}^i((\mathcal{N}^{\geq i}\Delta_{S/A})/p)[d^{1/p}]$. By Lemma 4.13, there is a commutative diagram

$$\begin{array}{ccc} \mathrm{H}^i((\mathcal{N}^{\geq i}\Delta_{S/A})/p) & \xrightarrow[\cong]{\phi} & \mathrm{H}^i(d^i\Delta_{S/A}/p) \\ \downarrow \mathrm{can} & & \downarrow \mathrm{can} \\ \mathrm{H}^i(\Delta_{S/A}/p) & \xrightarrow[\cong]{\phi} & \mathrm{H}^i(L\eta_d(\Delta_{S/A}/p)). \end{array}$$

By Proposition 2.11 (1), the right vertical map of this diagram is surjective, with kernel given by the d -torsion subgroup of $\mathrm{H}^i(d^i\Delta_{S/A}/p)$. So the left vertical map is also surjective, with kernel given by the $d^{1/p}$ -torsion subgroup $\mathrm{H}^i((\mathcal{N}^{\geq i}\Delta_{S/A})/p)[d^{1/p}]$. Hence the result. \square

Remark 4.15. Without assuming that the perfectoid base is p -torsionfree, the previous result would be false: for instance, in characteristic p , the homotopy cofibre of the syntomic-étale comparison map is typically nonzero in degree $i-1$ (Proposition 4.6).

REFERENCES

- [AGT16] Ahmed Abbes, Michel Gros, and Takeshi Tsuji. *The p -adic Simpson correspondence*, volume 193 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2016.
- [AMMN22] Benjamin Antieau, Akhil Mathew, Matthew Morrow, and Thomas Nikolaus. On the Beilinson fiber square. *Duke Math. J.*, 2022. to appear.
- [Avr99] Luchezar L. Avramov. Locally complete intersection homomorphisms and a conjecture of Quillen on the vanishing of cotangent homology. *Ann. of Math.*, 150(2):455–487, 1999.
- [Bha12] Bhargav Bhatt. p -adic derived de Rham cohomology. <https://arxiv.org/abs/1204.6560>, 2012.
- [Bha13] Bhargav Bhatt. An imperfect ring with trivial cotangent complex. <https://www.math.ias.edu/~bhatt/math/trivial-cc.pdf>, 2013.
- [Bha18] Bhargav Bhatt. Specializing varieties and their cohomology from characteristic 0 to characteristic p . *Algebraic Geometry: Salt Lake City 2015*, 2018.
- [BL22] Bhargav Bhatt and Jacob Lurie. Absolute prismatic cohomology. <https://arxiv.org/abs/2201.06120>, 2022.
- [BLM21] Bhargav Bhatt, Jacob Lurie, and Akhil Mathew. Revisiting the de Rham–Witt complex. *Astérisque*, (424), 2021.
- [BM21] Bhargav Bhatt and Akhil Mathew. The arc topology. *Duke Math. J.*, 170(9):1899–1988, 2021. .
- [BM22] Bhargav Bhatt and Akhil Mathew. Syntomic complexes and p -adic étale Tate twists. <https://arxiv.org/abs/2202.04818>, 2022.
- [BMS18] Bhargav Bhatt, Matthew Morrow, and Peter Scholze. Integral p -adic Hodge theory. *Publ. Math. Inst. Hautes Études Sci.*, 128:219–397, 2018.
- [BMS19] Bhargav Bhatt, Matthew Morrow, and Peter Scholze. Topological Hochschild homology and integral p -adic Hodge theory. *Publ. Math. Inst. Hautes Études Sci.*, 129:199–310, 2019.
- [BS17] Bhargav Bhatt and Peter Scholze. Projectivity of the Witt vector affine Grassmannian. *Invent. Math.*, 209(2):329–423, 2017.
- [BS21] Bhargav Bhatt and Peter Scholze. Prismatic F -crystals and crystalline Galois representations. <https://arxiv.org/abs/2106.14735>, 2021.
- [BS22] Bhargav Bhatt and Peter Scholze. Prisms and prismatic cohomology. *Ann. of Math.*, 2022. to appear.
- [CMM21] Dustin Clausen, Akhil Mathew, and Matthew Morrow. K -theory and topological cyclic homology of henselian pairs. *J. Amer. Math. Soc.*, 34(2):411–473, 2021.
- [ČS21] Kęstutis Česnavičius and Peter Scholze. Purity for flat cohomology. <https://arxiv.org/abs/1912.10932>, 2021.
- [DI87] Pierre Deligne and Luc Illusie. Relèvements modulo p^2 et décomposition du complexe de de Rham. *Invent. Math.*, 89(2):247–270, 1987.
- [EHIK21] Elden Elmanto, Marc Hoyois, Ryomei Iwasa, and Shane Kelly. Cdh descent, cdarc descent, and Milnor excision. *Math. Ann.*, 379(3-4):1011–1045, 2021.

- [Gab92] Ofer Gabber. K -theory of Henselian local rings and Henselian pairs. In *Algebraic K-theory, commutative algebra, and algebraic geometry (Santa Margherita Ligure, 1989)*, volume 126 of *Contemp. Math.*, pages 59–70. Amer. Math. Soc., Providence, RI, 1992.
- [GL00] Thomas Geisser and Marc Levine. The K -theory of fields in characteristic p . *Invent. Math.*, 139(3):459–493, 2000.
- [GL01] Thomas Geisser and Marc Levine. The Bloch–Kato conjecture and a theorem of Suslin–Voevodsky. *J. Reine Angew. Math.*, 530:55–103, 2001.
- [GR03] Ofer Gabber and Lorenzo Ramero. *Almost ring theory*, volume 1800 of *Lecture Notes in Math.* Springer-Verlag, Berlin, 2003.
- [HK18] Annette Huber and Shane Kelly. Differential forms in positive characteristic, II: cdh-descent via functorial Riemann–Zariski spaces. *Algebra Number Theory*, 12(3):649–692, 2018.
- [Hub96] Roland Huber. Adic spaces. In *Étale cohomology of rigid analytic varieties and adic spaces*, volume E30 of *Aspects of Mathematics*, pages 36–107. Friedr. Vieweg & Sohn, Braunschweig, 1996.
- [Ill71] Luc Illusie. *Complexe cotangent et déformations. I*, volume 239 of *Lecture Notes in Math.* Springer-Verlag, Berlin-New York, 1971.
- [KM21] Shane Kelly and Matthew Morrow. K -theory of valuation rings. *Compos. Math.*, 157(6):1121–1142, 2021.
- [KST21] Moritz Kerz, Florian Strunk, and Georg Tamme. Towards Vorst’s conjecture in positive characteristic. *Compos. Math.*, 157(6):1143–1171, 2021.
- [LM21] Morten Lüdgers and Matthew Morrow. Milnor K -theory of p -adic rings. <https://arxiv.org/abs/2101.01092>, 2021.
- [Nyg81] Niels O. Nygaard. Slopes of powers of Frobenius on crystalline cohomology. *Ann. Sci. École Norm. Sup.*, 14(4):369–401, 1981.
- [Sch12] Peter Scholze. Perfectoid spaces. *Publ. Math. Inst. Hautes Études Sci.*, 116:245–313, 2012.
- [Sta19] The Stacks Project Authors. *Stacks Project*. <https://stacks.math.columbia.edu>, 2019.
- [Sus83] Andrei Suslin. On the K -theory of algebraically closed fields. *Invent. Math.*, 73(2):241–245, 1983.
- [SV00] Andrei Suslin and Vladimir Voevodsky. Bloch–Kato conjecture and motivic cohomology with finite coefficients. In *The arithmetic and geometry of algebraic cycles (Banff, AB, 1998)*, volume 548 of *NATO Sci. Ser. C Math. Phys. Sci.*, pages 117–189. Kluwer Acad. Publ., Dordrecht, 2000.