

STABILITY CONDITIONS ON CYCLIC CATEGORIES I: BASIC DEFINITIONS AND EXAMPLES

YUCHENG LIU

ABSTRACT. A triangulated category \mathcal{C} with a canonical Bott's isomorphism $[2] \xrightarrow{\sim} id$ is called a cyclic category in this paper. We give a new notion of stability conditions on a k -linear Krull-Schmidt cyclic category. Given such a stability condition σ , we can assign a Maslov index to each basic loop in such a category. If all Maslov indexes vanish, we get \mathcal{C}', σ' as the \mathbb{Z} -lifts of \mathcal{C}, σ respectively such that \mathcal{C}' is a \mathbb{Z} -graded triangulated category and σ' is a Bridgeland stability condition on \mathcal{C}' . Moreover, we showed that there is an isomorphism

$$Stab^{0,e}(\mathcal{C}) \xrightarrow{\cong} BStab(\mathcal{C}')$$

where $Stab^{0,e}(\mathcal{C})$ denotes the equivalence classes of stability conditions which are deformation equivalent to σ , and $BStab(\mathcal{C}')$ denotes the space of Bridgeland stability conditions on \mathcal{C}' .

We provide examples of stability conditions on a simple cyclic category. We also discuss some interesting phenomena in these examples, such as the chirality symmetry breaking phenomenon and nontrivial monodromy. The chirality symmetry breaking phenomenon involves stability conditions which can not be lifted to Bridgeland stability conditions.

1. INTRODUCTION

The phenomenon $[2] = [0]$ has a relatively long history in mathematics, and can be found in many different branches of mathematics and physics. For examples: the celebrated Tate cohomology of cyclic groups and Bott's periodicity; in early 1980s, Eisenbud discovered that every finitely generated S module admits a free resolution which will eventually become 2-periodic, where S is the algebra of functions of a hypersurface with an isolated singularity (see [Eis80]); at almost the same time, people were developing the theory of cyclic homology (e.g. [Con85],[Goo85],[CQ97],[NS18]), the periodic cyclic homology played a role in such a theory; Kapustin and Li used the category of matrix factorizations to describe the D-branes in Landau-Ginzburg models by following a proposal of Kontsevich (see e.g. [KL03b], [KL03a]), which was further studied by mathematicians (e.g. [Dyc11], [DM12], [PV12], [Mur13]); We could also find the phenomenon of $[2]=[0]$ in Treumann's paper [Tre19].

On the other hand, the story of stability conditions on triangulated categories is relatively new. Motivated by Douglas's work on D-branes and II stability (see e.g. [Dou02]),

2010 *Mathematics Subject Classification.* 14F08, 14J40.

Key words and phrases. Bridgeland stability conditions, Matrix factorizations, Maslov index, Chirality symmetry breaking.

Bridgeland introduced a general theory of stability conditions on triangulated categories in [Bri07]; the theory was further studied by Kontsevich and Soibelman in [KS08].

Let \mathcal{C} be a k -linear triangulated category, if there exists a canonical isomorphism $\beta : [2] \simeq id$ between two functors, it is easy to see that there is no t -structures on \mathcal{C} . Hence no Bridgeland stability conditions exists on \mathcal{C} either. However, as Bridgeland stability condition can be viewed as an \mathbb{R} -grading refinement of a \mathbb{Z} -grading (t -structure) on a triangulated category, we expect that there exists a notion of S^1 -grading which refines a $\mathbb{Z}/2\mathbb{Z}$ -grading of a cyclic category.

Unlike on the real line, there are many homotopy non-equivalent paths to connects two points on S^1 , we need to introduce a new data to distinguish these paths. This new data is the degree function of a real decomposition on \mathcal{C} (see Definition 3.5 for the precise definition).

Roughly speaking, given a real decomposition on \mathcal{C} and any two indecomposable objects $E, F \in \mathcal{C}$, we have following decomposition

$$Hom_{\mathcal{C}}(E, F) = \bigoplus_{a \in \mathbb{R}} Hom_a(E, F),$$

which satisfy some natural conditions. A morphism $f \in Hom_a(E, F)$ is called a homogeneous morphism of degree a . We get a degree function

$$q : \{Nontrivial\ homogeneous\ morphisms\} \rightarrow \mathbb{R}.$$

This enables us to define the notion of connecting path (see Definition 3.10) and liftable commutative diagram (see Definition 3.12), which are the basic notion in our definitions and results. We use these notion to define what is a stability condition on a cyclic category in Section 3.

In Section 4, we define what is a basic loop and its Maslov index. This notion of Maslov index plays a similar role as its namesake in Fukaya categories. Indeed, we have the following lifting theorem.

Theorem 1.1. *Given a stability condition $\sigma = (\mathcal{Q}, Z, \phi, q)$ on a k -linear Krull-Schmidt cyclic category \mathcal{C} , we assume that the Maslov indexes of all basic loops are zero. There are \mathbb{Z} -lifts of σ and \mathcal{C} , which we denote by σ' and \mathcal{C}' respectively, such that σ' is a Bridgeland stability condition on \mathcal{C}' .*

Here the data (Z, ϕ, q) in a stability condition σ is called a charge triple, and in fact, the \mathbb{Z}' -lift \mathcal{C}' only depends on the charge triple. We say that \mathcal{C} is liftable with respect to a charge triple $R_1 := (Z, \phi, q)$ if the condition in Theorem 1.1 is satisfied. Furthermore, one can define when two charge triples are deformation equivalent (see Definition 4.10). And we proved the following result in Section 4.

Proposition 1.2. *Let \mathcal{C} be a k -linear Krull-Schmidt connective cyclic category, and $R_1 = (Z_1, \phi_1, q_1)$ and $R_2 = (Z_2, \phi_2, q_2)$ be two charge triples on \mathcal{C} . Suppose that R_1, R_2 are deformation equivalent and \mathcal{C} is liftable with respect to R_1 . Then \mathcal{C} is also liftable with respect to R_2 .*

Moreover, if we denote the \mathbb{Z} -lifts by \mathcal{C}_1 and \mathcal{C}_2 respectively, there exists an equivalence (not canonical) $H : \mathcal{C}_1 \simeq \mathcal{C}_2$ such that the following diagram is commutative.

$$\begin{array}{ccc} \mathcal{C}_1 & \xrightarrow{H} & \mathcal{C}_2 \\ & \searrow \pi_1 & \swarrow \pi_2 \\ & \mathcal{C} & \end{array}$$

A consistent choice of these equivalences forms a connection of \mathbb{Z} -lifts of \mathcal{C} fibered over the set of charge triples which are deformation equivalent to R_1 . Hence Theorem 1.1 and Proposition 1.2 provide us a map

$$Stab^0(\mathcal{C}) \rightarrow BStab(\mathcal{C}'),$$

where $Stab^0(\mathcal{C})$ consists of stability conditions whose charge triples are deformation equivalent to R_1 , and \mathcal{C}' is the associated \mathbb{Z} -lift of \mathcal{C} with respect to R_1 . We use $BStab(\mathcal{C}')$ to denote the space of Bridgeland stability conditions on \mathcal{C}' .

And a close look at this map provides us the following comparison theorem, which is proved in Section 5.

Theorem 1.3. *There is an isomorphism*

$$Stab^0(\mathcal{C}) / \simeq \rightarrow BStab(\mathcal{C}') / \mathbb{Z},$$

where the equivalence relation is defined in Definition 5.6 and the \mathbb{Z} action is given by $k \mapsto [2k]$ for any $k \in \mathbb{Z}$.

However, there are many stability conditions which can not be lifted to Bridgeland stability conditions. In Section 6, we provide some examples of stability conditions on the category of $\mathbb{Z}/3\mathbb{Z}$ -equivariant matrix factorizations of $w = x^3$. In this section, we also discuss some interesting phenomenon in these examples, for instance, the chirality symmetry breaking phenomena and the non-trivial monodromy of the map from stability condition to its central charge. The chirality symmetry breaking phenomenon involves stability conditions which can not be lifted to Bridgeland stability conditions.

1.1. Outline of this paper. In Section 2, we briefly review some basic definitions and results of matrix factorizations. In Section 3, we firstly give some preliminary definitions, such as real decompositions, connecting paths, liftable commutative diagrams and so on. Then we give our definition of stability conditions on cyclic categories in the end of this section. In Section 4, we introduced our notion of basic loop and its Maslov index, and prove the lifting theorem in this section. In Section 5, we prove the uniqueness of Harder-Narasimhan filtration under the assumption that all Maslov indexes are non-negative. Furthermore, we prove the comparison theorem. In Section 6, we give the examples of stability conditions on the category of $\mathbb{Z}/3\mathbb{Z}$ -equivariant matrix factorizations of $w = x^3$. We also discuss the chirality symmetry breaking phenomenon and nontrivial monodromy in these examples.

1.2. Acknowledgement. I would like to thank Emanuel Scheidegger for teaching me the background in physics, many helpful discussions and his guidance to the literature. I also thank Alex Martsinkovsky and Amnon Neeman for answering my questions via email correspondences. I also thank Qingyuan Bai, Hanfei Guo, Gang Han, Chunyi Li, Zhiyu Liu, Yongbin Ruan, Zhiyu Tian, Shizhuo Zhang, Xiaolei Zhao for helpful discussions. Theorem 5.9 is motivated by Zhiyu Tian's question. The final write-up of this paper was done while the author was visiting Zhejiang University, whose hospitality is gratefully acknowledged.

1.3. Notation and convention. A cyclic category, usually denoted by \mathcal{C} in this paper, is always assumed to be k -linear, Krull-Schmidt and essentially small.

2. MATRIX FACTORIZATIONS

2.1. Matrix Factorizations. Matrix factorizations was firstly introduced by Eisenbud in [Eis80]. It got the attention from physicists because of a proposal of Kontsevich, which suggests using them to describe the D-branes on Landau-Ginzburg model (see e.g. [KL03b] and [KL03a]). Mathematicians further studied them (see e.g. [Dyc11], [PV12], [Mur13], [Orl04], and [Orl09]). All the materials in this section are taken from these references.

Let us briefly recall the story of matrix factorizations. Suppose that (R, m) is a regular local ring and M is a finitely generated R -module. The famous Auslander-Buchsbaum formula is

$$pd(M) = dim(R) - depth(M).$$

In particular, if the depth of M is equal to the Krull dimension of R , then M is free.

However, if we consider a ring $S = R/w$ of hypersurface of singularity, where w is singular at m . The Auslander-Buchsbaum formula fails, and the condition

$$depth(M) = dim(S)$$

no longer implies that M is free. A module satisfying such condition is called a maximal Cohen-Macaulay module.

Given a maximal Cohen-Macaulay S -module M , we can consider M as an R -module. Then by Auslander-Buchsbaum formula we know that there is a length 1 R -free resolution of M . Suppose the following short exact sequence

$$0 \rightarrow F^1 \xrightarrow{f} F^0 \rightarrow M \rightarrow 0$$

is the R -free resolution of M . Since multiplication by w on M is trivial, there exists a homotopy g such that the diagram

$$\begin{array}{ccc} F^1 & \xrightarrow{f} & F^0 \\ \downarrow w & \swarrow g & \downarrow w \\ F^1 & \xrightarrow{f} & F^0 \end{array}$$

commutes. We use the pair (f, g) to define a matrix factorization of w . Note that the original maximal Cohen-Macaulay M is isomorphic to $coker(f)$.

Therefore, it is natural to bring up the following definition, where we assume that R is a commutative ring over a field k and $w \in R$ is an element.

Definition 2.1. A matrix factorization of the potential w over R is a pair

$$(E, \delta_E) = (E^0 \xrightarrow{\delta_0} E^1 \xrightarrow{\delta_1} E_0),$$

- $E = E^0 \oplus E^1$ is a $\mathbb{Z}/2$ -graded finitely generated projective R -module, and
- $\delta_E = \begin{pmatrix} 0 & \delta_1 \\ \delta_0 & 0 \end{pmatrix} \in \text{End}_R^1(E)$ is an odd (i.e. of degree $1 \in \mathbb{Z}/2$) endomorphism of E , such that $\delta_E^2 = w \cdot \text{id}_E$.

Remark 2.2. The map δ_E is usually called a twisted differential in the literature. We will adopt this terminology in this paper. Note that $\delta_E^2 = w \cdot \text{id}_E$ is also equivalent to $\delta_0 \delta_1 = \delta_1 \delta_0 = w \cdot I$.

In the case when $R = k[[x_1, x_2, \dots, x_n]]$ is the ring of formal power series in n variables over a field k , E_0 and E_1 are free R -modules. Moreover, E_0, E_1 are of the same rank over R by the requirement $\delta_0 \delta_1 = \delta_1 \delta_0 = w \cdot I$, hence all the maps δ_0, δ_1 and δ_E can be represented as matrices of elements in R . For all explicit examples in this paper, we will take R to be a ring of formal power series.

To a potential $w \in R$ we can associated a $\mathbb{Z}/2$ dg-category $MF(w) := MF(R, w)$ whose objects are matrix factorizations of w over R . The morphisms from $\bar{E} = (E, \delta_E)$ to $\bar{F} = (F, \delta_F)$ are elements of the $\mathbb{Z}/2$ -graded module of R -linear homomorphisms

$$\mathcal{H}om_w(\bar{E}, \bar{F}) := \text{Hom}_{\text{Mod}_R}(E, F) = \text{Hom}_{\mathbb{Z}/2\text{-Mod}_R}(E, F) \oplus \text{Hom}_{\mathbb{Z}/2\text{-Mod}_R}(E, F[1]).$$

The $\mathbb{Z}/2$ -graded dg-structure is given by the following differential on $f \in \mathcal{H}om_w(\bar{E}, \bar{F})$

$$df = \delta_F \circ f - (-1)^{|f|} f \circ \delta_E.$$

We use

$$HMF(R, w) = H^0 MF(R, w)$$

to denote the associated homotopy category of $MF(R, w)$, i.e, the space of morphisms in this category are chain maps up to homotopy. The homotopy category $HMF(R, w)$ is naturally triangulated (see e.g. [Orl04]) with the shift functor

$$T : (E^0 \xrightarrow{\delta_0} E^1 \xrightarrow{\delta_1} E_0) \mapsto (E^1 \xrightarrow{-\delta_1} E^0 \xrightarrow{-\delta_0} E_1).$$

Remark 2.3. Eisenbud proved that (see [Eis80]) in the case when (R, m) is a regular local ring and $S = R/w$, where w is singular at the closed point m , the functor coker induces an equivalence

$$\text{coker} : HMF(R, w) \xrightarrow{\sim} \underline{MCM}(S).$$

Here $\underline{MCM}(S)$ is the stable category of maximal Cohen-Macaulay S -modules. The objects are maximal Cohen-Macaulay modules, the morphisms are defined by

$$\underline{\text{Hom}}_S(M, M') = \text{Hom}_S(M, M')/P,$$

where P denotes the set of S -linear homomorphisms factoring through some free S -module.

In such a case, Buchweitz proved that there is another equivalence

$$\underline{MCM}(S) \rightarrow D_{sing}^b(S),$$

where $D_{sing}^b(S)$ is the Verdier quotient

$$D_{sing}^b(S) := D^b(S)/D_{perf}^b(S).$$

Here $D^b(S)$ is the derived category of all complexes of S -modules with finitely generated total cohomology. Such a complex is called perfect if it is isomorphic in $D^b(S)$ to a bounded complex of free S -modules. The full triangulated subcategory formed by the perfect complexes is denoted by $D_{perf}^b(S)$. See also the paper [Orl04] for the proofs of such equivalences.

Moreover, in such a case, there is a non-degenerate pairing on the morphism spaces in $HMF(R, w)$. This beautiful formula was firstly introduced by Kapustin and Li using the path integral method when $k = \mathbb{C}$ (see e.g. [KL03b]). This formula was mathematically proved in [Mur13] and [DM12].

For simplicity, we can assume $R = k[[x_1, \dots, x_n]]$. For the explicit meaning of the notation in the formula, please consult [DM12, Sections 1-3].

Theorem 2.4. [DM12, Theorem 3.4] *The pairing*

$$Hom(X, Y) \otimes_R Hom(Y, X[n]) \rightarrow k,$$

$$(F, G) \mapsto (-1)^{\binom{n+1}{2}} \frac{1}{n!} Res_{[\partial_1 w, \partial_2 w, \dots, \partial_n w]} [tr(FG(dQ)^n)] = (-1)^{\binom{n+1}{2}} \frac{1}{n!} Res_{[\partial_1 w, \partial_2 w, \dots, \partial_n w]} [tr(GF(dP)^n)]$$

provides a homologically non-degenerate pairing on the morphism complexes of the category $MF(R, w)$ associated to the germ of an isolated hypersurface singularity. Here P, Q are the twisted differentials of X and Y respectively.

Proof. The formula involving the twisted differential Q of Y is proved in [DM12], and the formula involving the twisted differential P of X can be proved by the same method. \square

We conclude this section by briefly recalling the G -equivariant version of Matrix factorizations (see e.g. [PV12, Section 2.1]). Indeed, if G is a finite group of automorphisms of R which fixes the potential w , one defines the G -equivariant $\mathbb{Z}/2$ -graded ad-category of matrix factorizations $MF_G(w)$ (and the corresponding homotopy category $HMF_G(w)$), by requiring that all modules and morphisms should be G -equivariant. In other words, the module E in Definition 2.1 should be a $\mathbb{Z}/2$ -graded finitely generated projective R -module equipped with a compatible G -action, and δ_E has to be G -equivariant. Morphisms between G -equivariant factorizations \bar{E} and \bar{F} should also be compatible with the action of G , i.e.

$$Hom_{MF_G(w)}(\bar{E}, \bar{F}) = Hom_{MF(w)}(\bar{E}, \bar{F})^G.$$

3. STABILITY CONDITIONS ON CYCLIC CATEGORIES

3.1. Bridgeland stability conditions. The theory of Bridgeland stability conditions was introduced by Bridgeland in [Bri07], motivated by Douglas's work on D-branes and Π -stability [Dou02]. This theory was further studied by Kontsevich and Soibelman in [KS08]. In this section, we will review some basic notions in the theory of stability conditions (see [BBD82], [Bri07], [KS08] and [BLMS17]).

The first notion is t -structures on triangulated categories, which was firstly introduced in [BBD82].

Definition 3.1. Let \mathcal{D} be an triangulated category. A t -structure on \mathcal{D} is a pair of full subcategories $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ satisfying the condition (i), (ii) and (iii) below. We denote $\mathcal{D}^{\leq n} = \mathcal{D}^{\leq 0}[-n]$, $\mathcal{D}^{\geq n} = \mathcal{D}^{\geq 0}[-n]$ for every $n \in \mathbb{Z}$. Then the conditions are:

- (i) $\text{Hom}(E, F) = 0$ for every $E \in \mathcal{D}^{\leq 0}$ and $F \in \mathcal{D}^{\geq 1}$;
- (ii) $\mathcal{D}^{\leq -1} \subset \mathcal{D}^{\leq 0}$ and $\mathcal{D}^{\geq 1} \subset \mathcal{D}^{\geq 0}$.
- (iii) every object $E \in \mathcal{D}$ fits into an exact triangle

$$\tau^{\leq 0} E \rightarrow E \rightarrow \tau^{\geq 1} E \rightarrow \dots$$

with $\tau^{\leq 0} E \in \mathcal{D}^{\leq 0}$, $\tau^{\geq 1} E \in \mathcal{D}^{\geq 1}$.

The heart of the t -structure is $\mathcal{A} = \mathcal{D}^{\leq 0} \cap \mathcal{D}^{\geq 0}$. It is an abelian category (see [HT07, Theorem 8.1.9]). The associated cohomology functors are defined by $H^0(E) = \tau^{\leq 0} \tau^{\geq 0} E$, $H^i(E) = H^0(E[i])$. We will also need the notation $\mathcal{D}^{[a,b]} = \mathcal{D}^{\leq b} \cap \mathcal{D}^{\geq a}$.

Combining this definition with Harder-Narasimhan filtrations, Bridgeland defined the notion of stability conditions on a triangulated category in [Bri07].

Definition 3.2. A Bridgeland stability condition (\mathcal{P}, Z) on a triangulated category \mathcal{D} consists of a group homomorphism $Z : K_0(\mathcal{D}) \rightarrow \mathbb{C}$, which factors through a fixed group homomorphism $K_0(\mathcal{D}) \rightarrow \Lambda$, where Λ a lattice of finite rank. This group homomorphism is called a central charge. And full subcategories $\mathcal{P}(\phi) \in \mathcal{D}$ for each $\phi \in \mathbb{R}$, satisfying the following axioms:

- (a) if $E \in \mathcal{P}(\phi)$ is a nonzero object, then $Z(E) = m(E) \exp(i\pi\phi)$ for some $m(E) \in \mathbb{R}_{>0}$,
- (b) for all $\phi \in \mathbb{R}$, $\mathcal{P}(\phi + 1) = \mathcal{P}(\phi)[1]$,
- (c) if $\phi_1 > \phi_2$ and $A_j \in \mathcal{P}(\phi_j)$ then $\text{Hom}_{\mathcal{D}}(A_1, A_2) = 0$,
- (d) for every $0 \neq E \in \mathcal{D}$ there exist a finite sequence of real numbers

$$\phi_1 > \phi_2 > \dots > \phi_m$$

and a sequence of morphisms

$$0 = E_0 \xrightarrow{f_1} E_1 \xrightarrow{f_2} \dots \xrightarrow{f_m} E_m = E$$

such that the cone of f_j is in $\mathcal{P}(\phi_j)$ for all j .

Remark 3.3. (i) If we allow $m(E)$ to be 0 for $\phi \in \mathbb{Z}$ in (a), then the pair (\mathcal{P}, Z) is called a weak stability condition. In [KS08], the authors require the pair (\mathcal{P}, Z) to satisfy one extra

condition (support property) to be a stability condition. We do not include this condition because it is not needed in this paper.

(ii) This notion of Bridgeland stability conditions is categorical. Indeed, suppose that $H : \mathcal{D}_1 \xrightarrow{\sim} \mathcal{D}_2$ is an exact equivalence between two triangulated categories, and $\sigma = (\mathcal{P}, Z)$ is a Bridgeland stability condition on \mathcal{D}_2 . Then we have a Bridgeland stability condition $H^*\sigma = (H^*\mathcal{P}, H^*Z)$ on \mathcal{D}_1 , which is defined in the following way:

- $H^*\mathcal{P}(\phi) = \{E \mid H(E) \in \mathcal{P}(\phi)\}$, for any $\phi \in \mathbb{R}$.
- $H^*Z(E) = Z(H(E))$ for any E in \mathcal{D}_1 .

It is easy to check this is a Bridgeland stability condition on \mathcal{D}_1 .

The data \mathcal{P} of full subcategories $\mathcal{P}(\phi)$ is called a slicing on \mathcal{D} , a slicing can be viewed as a refinement of a t -structure on a triangulated category. Indeed, one can easily check that a slicing on \mathcal{D} gives us a lot of t -structures on \mathcal{D} : for any $\phi \in \mathbb{R}$, we have a t -structure $(\mathcal{P}(> \phi - 1), \mathcal{P}(\leq \phi))$ on \mathcal{D} .

In particular, a slicing \mathcal{P} of \mathcal{D} provides us a heart $\mathcal{P}(0, 1] = \mathcal{P}(> 0) \cap \mathcal{P}(\leq 1)$. Hence, a stability condition (\mathcal{P}, Z) gives us a pair (\mathcal{A}, Z) , where \mathcal{A} is an abelian category. This construction results in an equivalent definition of stability conditions.

In this paper, we are interested in the triangulated categories with the special property $[2] \simeq [0]$. Hence we have the following terminology.

Definition 3.4. A triangulated category \mathcal{C} is called a cyclic category if there is canonical isomorphism between two functors

$$\beta : [2] \simeq id.$$

We usually call β a Bott's isomorphism.

It is easy to see that there exist no t -structures on any nontrivial cyclic categories. Indeed, assume that there is a t -structure on \mathcal{C} . Then $[1] = [-1]$ implies that $id_{A[1]} \in Hom(A[1], A[1]) \simeq Hom(A[1], A[-1]) = 0$ for any object A in the heart, which implies that the heart is trivial. As a slicing is a refinement of a t -structure, any nontrivial cyclic categories do not admit any Bridgeland stability conditions.


However, if one intuitively think a slicing as a helix parametrization of the structure of a triangulated category, one would expect to have the notion of S^1 -gradings on cyclic categories, which is a circle parametrization of the structure of a cyclic category (see Figure 1 for this intuition).

3.2. Real decompositions and liftable commutative diagrams. To make this intuition precise, we need to do some preliminary work.

From here on, we will assume that our cyclic category \mathcal{C} is Krull-Schmidt, i.e. every object decomposes into direct sum of indecomposable objects in an unique way up to isomorphism, and the endomorphism ring of an indecomposable object is local.

Definition 3.5. Let \mathcal{C} be a k -linear Krull-Schmidt cyclic category. We say that \mathcal{C} admits a real decomposition if every space $Hom_{\mathcal{C}}(E, F)$ admits a grading by \mathbb{R} , i.e.

$$Hom_{\mathcal{C}}(E, F) = \bigoplus_{a \in \mathbb{R}} Hom_a(E, F)$$



A helix grading for a triangulated category.

An S^1 grading for a cyclic category.

FIGURE 1

for any two indecomposable objects $E, F \in \mathcal{C}$. And these gradings should satisfy the following conditions.

- (1) For any morphism $f \in \text{Hom}_a(E, F)$, we also have $f[1] \in \text{Hom}_a(E[1], F[1])$.
- (2) For any two morphisms f, g with

$$f \in \text{Hom}_a(E, F), \quad g \in \text{Hom}_b(F, G),$$

we have

$$g \circ f \in \text{Hom}_{(a+b)}(E, G).$$

- (3) For any indecomposable object $E \in \mathcal{C}$, the morphisms $id_E \in \text{Hom}_0(E, E)$ and $\beta_E \in \text{Hom}_0(E[2], E)$.
- (4) A morphism $f \in \text{Hom}_a(E, F)$ is called a homogeneous morphism of degree a . For any isomorphism f , if f is homogeneous, its degree is 0.

This decomposition also defines a function

$$q : \{\text{Nontrivial homogeneous morphisms}\} \rightarrow \mathbb{R},$$

which sends a nontrivial homogeneous morphism to its degree. The function q is called the degree function associated with the real decomposition of \mathcal{C} .

Remarks 3.6. (i) The degree function q is also called R -charge of homogeneous morphism in physics literature (see e.g. [Wal05]). The real decomposition endows a graded local algebra structure on $\text{Hom}_{\mathcal{C}}(E, E)$ for any indecomposable object $E \in \mathcal{C}$, and a graded $\text{Hom}_{\mathcal{C}}(E, E) - \text{Hom}_{\mathcal{C}}(F, F)$ bi-module structure on $\text{Hom}_{\mathcal{C}}(E, F)$ for any two indecomposable objects E, F .

(ii) By the definition, we know that a degree function q satisfy the following basic properties.

(1) For any nontrivial homogeneous morphism f , we have $q(f) = q(f[1])$.

(2) If f, g, gf are three nontrivial homogeneous morphisms, we have $q(gf) = q(g) + q(f)$.

(iii) A real decomposition on \mathcal{C} also induces a natural \mathbb{R} -grading on $\text{Hom}_{\mathcal{C}}(E, F)$, where E, F are not necessarily indecomposable. Indeed, let $E = \bigoplus_{i=1}^n E_i, F = \bigoplus_{j=1}^m F_j$, where E_i, F_j are indecomposable objects for any $1 \leq i \leq n, 1 \leq j \leq m$. One can decompose $\text{Hom}_{\mathcal{C}}(E, F)$ in the following way:

$$\text{Hom}_{\mathcal{C}}(E, F) = \bigoplus_{a \in \mathbb{R}} \text{Hom}_a(E, F),$$

where

$$\text{Hom}_a(E, F) = \bigoplus_{i,j} \text{Hom}_a(E_i, F_j).$$

And a morphism $f \in \text{Hom}_a(E, F)$ is also called a homogeneous morphism of degree a .

We have the following lemma from the definition of real decomposition.

Lemma 3.7. If \mathcal{C} admits a real decomposition and E, F are two isomorphic indecomposable objects, then there exists an isomorphism $f \in \text{Hom}_0(E, F)$. And f induces an isomorphism between \mathbb{R} -graded vector spaces

$$\text{Hom}_{\mathcal{C}}(F, G) \xrightarrow{\circ f} \text{Hom}_{\mathcal{C}}(E, G).$$

Proof. Since \mathcal{C} is Krull-Schmidt. The sum of two non-invertible morphisms in $\text{Hom}(E, F)$ is still not invertible. Hence any isomorphism must have a homogeneous summand f which is also an isomorphism. By condition (4) in Definition 3.5, f is of degree 0. Condition (2) in Definition 3.5 implies the lemma. \square

Remark 3.8. Take G to be E in the lemma, we get that f^{-1} is also a homogeneous morphism of degree 0.

Given a real decomposition of \mathcal{C} , we can define quasi-homogeneous morphisms and liftable quasi-homogeneous morphisms.

Definition 3.9. Suppose a morphism $f : A \rightarrow B$ can be written in the following way

$$f : \bigoplus_{i=1}^l A_i \rightarrow \bigoplus_{j=1}^m B_j,$$

where A_i, B_j are nontrivial indecomposable objects for all i, j . Then f is called quasi-homogeneous if all its direct summand $f_{ji} : A_i \rightarrow B_j$ are homogeneous morphisms for arbitrary i, j .

We also need the following terminology to define liftable quasi-homogeneous morphisms.

Definition 3.10. (1) Given a real decomposition on \mathcal{C} , we call the following diagram

$$E_0 \xleftrightarrow{f_0} E_1 \xleftrightarrow{f_1} \dots \xleftrightarrow{f_{n-1}} E_n$$

a connecting path from E_0 to E_n if E_i is an indecomposable object for any $0 \leq i \leq n$ and f_j is a nontrivial homogeneous morphism in either $\text{Hom}_{\mathcal{C}}(E_j, E_{j+1})$ or $\text{Hom}_{\mathcal{C}}(E_{j+1}, E_j)$ for any $0 \leq j \leq n-1$. We usually use l to denote a connecting path, and let

$$q(l) := \sum_{i=0}^{n-1} \text{Sign}(f_i) q(f_i),$$

where

$$\text{Sign}(f_i) = \begin{cases} 1, & \text{if } f_i \in \text{Hom}_{\mathcal{C}}(E_i, E_{i+1}); \\ -1 & \text{if } f_i \in \text{Hom}_{\mathcal{C}}(E_{i+1}, E_i). \end{cases}$$

In the special case when $E_0 = E_n$, we say that l is a connecting loop. A connecting path l is called simple if it contains no connecting loops.

(2) The cyclic category \mathcal{C} is called connective if for any two indecomposable objects E, F , there exists a connecting path between them.

(3) Let \mathbf{D} be a commutative digram in \mathcal{C} , if all the morphisms in \mathbf{D} are quasi-homogeneous. One could say that a connecting path

$$l : E_0 \xleftrightarrow{f_0} E_1 \xleftrightarrow{f_1} \dots \xleftrightarrow{f_{n-1}} E_n$$

is in \mathbf{D} if any homogeneous morphism f_i is a homogeneous direct summand of a quasi-homogeneous morphism in that commutative diagram.

Remarks 3.11. (1) *The degree of a connecting path depends on its direction. Indeed, a connecting path l from E to F can also be viewed as a connecting path from F to E , which we denote it by \bar{l} . We have*

$$q(l) = -q(\bar{l}).$$

(2) *The property of being connective is independent of the real decomposition, it is a property of the category \mathcal{C} itself.*

(3) *For any commutative digram \mathbf{D} in \mathcal{C} , there is an associated directed graph $G(\mathbf{D})$ whose vertices are objects in \mathbf{D} , and edges are the morphisms in \mathbf{D} . We are only interested in commutative diagrams whose associated graph is simple, i.e. with neither self loops nor multiple edges. So every commutative diagram in this paper is assumed to be of such kind.*

All diagrams below are assumed to be in a k -linear Krull-Schmidt cyclic category \mathcal{C} which admits a real decomposition. We introduce two special kinds of such commutative diagrams .

Definition 3.12. Let \mathbf{D} be a commutative diagram in \mathcal{C} . We say that \mathbf{D} is liftable if the following conditions are satisfied:

is also liftable.

Proof. For any $1 \leq i \leq n$, we can write

$$A_0 = \bigoplus_{j=1}^m A_{0,j}, \quad A_i = \bigoplus_{k=1}^l A_{i,k}.$$

Let $f_{kj} : A_{0,j} \rightarrow A_{i,k}$ to denote the corresponding summand in $f_{i-1} \circ \cdots \circ f_0$. The morphism f_{kj} can be written as the sum of homogeneous morphisms from $A_{0,j}$ to $A_{i,k}$ in \mathbf{D} . As \mathbf{D} is liftable, these homogeneous morphisms are of the same degree. Hence any nontrivial morphism f_{kj} is a homogeneous morphism of degree equal to $q(l)$, where $l : A_{0,j} \dashrightarrow A_{i,k}$ is a connecting path in \mathbf{D} .

One can easily show that the commutative diagram is liftable by this observation. \square

Remark 3.16. In fact, the same argument can show that the associated $(n+1)$ -complete commutative digram is also liftable.

Lemma 3.17. *Consider the following diagram*

$$\begin{array}{ccc} \bigoplus_{i=1}^l A_i & \xrightarrow{f} & \bigoplus_{j=1}^m B_j \\ \downarrow g & & \\ \bigoplus_{k=1}^n C_k & & \end{array}$$

where

$$f = \begin{pmatrix} f_{11} & \cdots & f_{1l} \\ \cdots & \cdots & \cdots \\ f_{m1} & \cdots & f_{ml} \end{pmatrix} \quad g = \begin{pmatrix} g_{11} & \cdots & g_{1l} \\ \cdots & \cdots & \cdots \\ g_{n1} & \cdots & g_{nl} \end{pmatrix}$$

are quasi-homogeneous morphisms. Suppose that this diagram is a sub-digram of a liftable commutative diagram \mathbf{D} , and can be completed into a commutative diagram by a morphism h .

$$\begin{array}{ccc} \bigoplus_{i=1}^l A_i & \xrightarrow{f} & \bigoplus_{j=1}^m B_j \\ \downarrow g & \swarrow h & \\ \bigoplus_{k=1}^n C_k & & \end{array}$$

Then there is a quasi-homogeneous morphism \bar{h} such that if we add \bar{h} to the digram \mathbf{D} , we get a liftable digram $\bar{\mathbf{D}}$ and the following diagram

$$\begin{array}{ccc} \bigoplus_{i=1}^l A_i & \xrightarrow{f} & \bigoplus_{j=1}^m B_j \\ \downarrow g & \swarrow \bar{h} & \\ \bigoplus_{k=1}^n C_k & & \end{array}$$

is a liftable commutative diagram.

Proof. If we write

$$h = \begin{pmatrix} h_{11} & \cdots & h_{1m} \\ \cdots & \cdots & \cdots \\ h_{n1} & \cdots & h_{nm} \end{pmatrix}$$

we claim that if we take

$$\bar{h} = \begin{pmatrix} \bar{h}_{11} & \cdots & \bar{h}_{1m} \\ \cdots & \cdots & \cdots \\ \bar{h}_{n1} & \cdots & \bar{h}_{nm} \end{pmatrix}$$

where

$$\bar{h}_{kj} = \begin{cases} \text{Summand of } h_{kj} \text{ in degree } q(l), & \text{if there is a connecting path } l : B_j \dashrightarrow C_k \in \mathbf{D}; \\ 0, & \text{otherwise.} \end{cases}$$

Then \bar{h} satisfy the statement in the lemma.

First of all, the new diagram is liftable by our choice. Indeed, for any connecting loop l in \mathbf{D} , we can replace the segments \bar{h}_{kj} in l by a connecting path $l : B_j \dashrightarrow C_k \in \mathbf{D}$. This does not change the degree of the loop by the definition of \bar{h}_{kj} . Hence we get a new loop $l' \in \mathbf{D}$ with $q(l) = q(l') = 0$.

The next thing is to check that $g = \bar{h} \circ f$. It suffices to prove that

$$(1) \quad (\bar{h}_{k1} \quad \cdots \quad \bar{h}_{km}) \begin{pmatrix} f_{11} & \cdots & f_{1l} \\ \cdots & \cdots & \cdots \\ f_{m1} & \cdots & f_{ml} \end{pmatrix} = (g_{k1} \quad \cdots \quad g_{kl})$$

for any $1 \leq k \leq n$.

For any given $1 \leq i_1 \leq l$, we let

$$J := \{j_1, j_2, \dots, j_s\} = \{1 \leq j \leq m \mid f_{ji_1} \neq 0\}.$$

The case $J = \emptyset$ is trivial. We assume that $J \neq \emptyset$. If there is no connecting paths from B_{j_r} to C_k in \mathbf{D} for any $1 \leq r \leq s$, the morphism g_{ki_1} must be 0. Hence

$$\sum_{j=1}^m \bar{h}_{kj} f_{ji_1} = 0 = g_{ki_1}.$$

On the other hand, if there is an integer $1 \leq r_1 \leq s$ such that there is a connecting path from $B_{j_{r_1}}$ to C_k in \mathbf{D} . Then there is a connecting path

$$B_{j_r} \xleftarrow{f_{j_r i_1}} A_{i_1} \xrightarrow{f_{j_{r_1} i_1}} B_{j_{r_1}} \dashrightarrow C_k$$

for any $1 \leq r \leq s$. Hence we have that $\bar{h}_{kj_r} f_{j_r i_1}$ are homogeneous morphisms of same degree for any $1 \leq r \leq s$, and this degree equals to $q(g_{ki_1})$ if $g_{ki_1} \neq 0$. This implies that $\sum_{j=1}^m \bar{h}_{kj} f_{ji_1} = g_{ki_1}$ if $g_{ki_1} \neq 0$. If $g_{ki_1} = 0$, we know that $\sum_{j=1}^m \bar{h}_{kj} f_{ji_1}$ is a homogeneous summand of $\sum_{j=1}^m h_{kj} f_{ji_1} = g_{ki_1} = 0$, which is also 0. This completes the proof. \square

Remark 3.18. One thing worth noticing is that, as we can see in the proof, the morphism \bar{h} may depend on the ambient commutative liftable diagram \mathbf{D} .

Corollary 3.20. *Let \mathbf{D} be any liftable commutative diagram, suppose that we can add a morphism h into \mathbf{D} such that the new diagram \mathbf{D}' is still commutative. Then there exists a quasi-homogeneous morphism \bar{h} such that, if we add \bar{h} into \mathbf{D} in the same position, the new diagram $\bar{\mathbf{D}}$ is a liftable commutative diagram.*

Proof. We can construct \bar{h} as in the proof of Lemma 3.17, the new diagram is obviously liftable. And the commutativity follows from Lemma 3.15 and Proposition 3.19. \square

Sometimes, we need to glue two liftable commutative diagrams along a common sub-diagram. In general the glued diagram will not be liftable. However, we have following easy lemma.

Lemma 3.21. *Let \mathbf{D}_1 and \mathbf{D}_2 be two liftable diagrams, \mathbf{D}_3 be a common sub-diagram of \mathbf{D}_1 and \mathbf{D}_2 , and \mathbf{D} be the glued diagram of \mathbf{D}_1 and \mathbf{D}_2 along \mathbf{D}_3 . If \mathbf{D}_3 is connective, then \mathbf{D} is liftable.*

Proof. We can focus on the loops which are not in \mathbf{D}_1 or \mathbf{D}_2 . Any loop of such kind $l : E \dashrightarrow E$ in \mathbf{D} can be written as $l_1 \circ l_2 \circ \cdots \circ l_n$ where $l_i : E_i \dashrightarrow E_{i+1}$ is a connecting path in \mathbf{D}_1 or \mathbf{D}_2 depending on the parity of i , and the indecomposable objects E_i are in the diagram \mathbf{D}_3 with $E_0 = E_{n+1} = E$.

As \mathbf{D}_3 is connective, we have connecting paths $l'_i : E_i \dashrightarrow E_{i+1}$ in \mathbf{D}_3 . Therefore, one can get

$$q(l) = \sum_{i=1}^n q(l_i) = \sum_{i=1}^n q(l'_i) = q(l'_1 \circ l'_2 \circ \cdots \circ l'_n) = 0.$$

The second equation follows from the assumption that \mathbf{D}_1 and \mathbf{D}_2 are liftable. Hence, the lemma is proved. \square

3.3. Definition of stability conditions. In the following, we will discuss the compatibility between the triangulated structure of \mathcal{C} and the real decomposition on \mathcal{C} .

Definition 3.22. Let \mathcal{C} be a k -linear Krull-Schmidt cyclic category which admits a real decomposition. We say that \mathcal{C} is pre-liftable with respect to the real decomposition if the following conditions holds:

- (1) for any liftable quasi-homogeneous morphism $f : A \rightarrow B$, we can complete it to a distinguished triangle

$$A \xrightarrow{f} B \xrightarrow{g} C \xrightarrow{h} A[1]$$

such that the following diagram

$$\cdots \xrightarrow{g[-1]} A[-1] \xrightarrow{h[-1]} A \xrightarrow{f} B \xrightarrow{g} C \xrightarrow{h} A[1] \xrightarrow{f[1]} B[1] \xrightarrow{g[1]} \cdots$$

is liftable,

(2) if we have the following liftable commutative diagram

$$\begin{array}{ccccccc}
 X & \xrightarrow{id} & X & & & & \\
 \downarrow u & & \downarrow v \circ u & & & & \\
 Y & \xrightarrow{v} & Z & \xrightarrow{l} & X' & \xrightarrow{i} & Y[1] \\
 \downarrow j & & \downarrow m & & & & \\
 Z' & & Y' & & & & \\
 \downarrow k & & \downarrow n & & & & \\
 X[1] & \xrightarrow{id} & X[1] & & & &
 \end{array}$$

where all row and columns and are distinguished triangle, then it can be completed into the following liftable commutative diagram

$$\begin{array}{ccccccc}
 X & \xrightarrow{id} & X & & & & \\
 \downarrow u & & \downarrow v \circ u & & & & \\
 Y & \xrightarrow{v} & Z & \xrightarrow{l} & X' & \xrightarrow{i} & Y[1] \\
 \downarrow j & & \downarrow m & & \downarrow id & & \downarrow j[1] \\
 Z' & \xrightarrow{f} & Y' & \xrightarrow{g} & X' & \xrightarrow{h} & Z[1] \\
 \downarrow k & & \downarrow n & & & & \\
 X[1] & \xrightarrow{id} & X[1] & & & &
 \end{array}$$

where all the columns and rows are distinguished triangles.

Now, we are ready to define stability conditions on a k -linear Krull-Schmidt cyclic category.

Definition 3.23. A stability condition on a k -linear Krull-Schmidt cyclic category \mathcal{C} consists of four parts $(\mathcal{Q}, Z, \phi, q)$, where \mathcal{Q} is a circle slicing, i.e. full subcategories $\mathcal{Q}(\psi)$ for any $\psi \in (0, 2]$, a central charge $Z : K_0(\mathcal{C}) \xrightarrow{v} \Lambda \rightarrow \mathbb{C}$, a map ϕ sending every indecomposable object E to its phase $\phi(E) \in (0, 2]$, and q a degree function of a real decomposition of \mathcal{C} . Here \mathcal{C} is pre-liftable with respect to the real decomposition and these data satisfy the following compatibility conditions.

- (1) We have that $\phi(E[1]) \equiv \phi(E) + 1 \pmod{2\mathbb{Z}}$.
- (2) For any indecomposable object E , the central charge can be written as

$$Z(E) = m(E)e^{i\pi\phi(E)},$$

where $m(E) \in \mathbb{R}_{\geq 0}$ and $\phi(E) \in (0, 2]$.

- (3) For any homogeneous morphism $f : E_1 \rightarrow E_2$ between two indecomposable objects, we have

$$q(f) \equiv \phi(E_2) - \phi(E_1) \pmod{2\mathbb{Z}}.$$

- (4) $\mathcal{Q}(\psi)[1] = \mathcal{Q}(\psi')$, where $\psi' \equiv \psi + 1 \pmod{2\mathbb{Z}}$ and $\psi, \psi' \in (0, 2]$.
 (5) For any object $E \in \mathcal{Q}(\phi)$, we have $m(E) > 0$ and $\phi(E) = \phi$.
 (6) For any nontrivial homogeneous morphism $f : E_1 \rightarrow E_2$, if $E_k \in \mathcal{Q}(\phi_k)$ for $k = 1, 2$, we have $q(f) > 0$.
 (7) For any indecomposable object $E \in \mathcal{C}$, we have the following filtration:

$$\begin{array}{ccccccc}
 & & E_0 & & E_1 & & \cdots & & E_{n-2} & & E_{n-1} \\
 & \nearrow id & \downarrow f_1 & \nearrow id & \downarrow f_2 & \nearrow id & \cdots & \nearrow id & \downarrow f_{n-1} & \nearrow id & \downarrow f_n \\
 0 = E_0 & & E_1 & & E_2 & & \cdots & & E_{n-1} & & E_n = E \\
 & & \downarrow p_1 & & \downarrow p_2 & & \cdots & & \downarrow p_{n-1} & & \downarrow p_n \\
 & & Q_1 & & Q_2 & & \cdots & & Q_{n-1} & & Q_n
 \end{array}$$

such that it satisfies the following conditions:

- the whole diagram is connective and liftable,
- for any $1 \leq i \leq n$, we have $Q_i \in \mathcal{Q}(\phi_i)$ being nontrivial semi-stable objects,
- the diagram can be completed into the following liftable diagram,

$$\begin{array}{ccccccc}
 & & \cdots & & \cdots & & \cdots & & \cdots & & \cdots \\
 & & \downarrow t_1[-1] & & \downarrow t_2[-1] & & \cdots & & \downarrow t_{n-1}[-1] & & \downarrow t_n[-1] \\
 & \nearrow id & E_0 & \nearrow id & E_1 & \nearrow id & \cdots & \nearrow id & E_{n-2} & \nearrow id & E_{n-1} \\
 & & \downarrow f_1 & & \downarrow f_2 & & \cdots & & \downarrow f_{n-1} & & \downarrow f_n \\
 0 = E_0 & & E_1 & & E_2 & & \cdots & & E_{n-1} & & E_n \\
 & & \downarrow p_1 & & \downarrow p_2 & & \cdots & & \downarrow p_{n-1} & & \downarrow p_n \\
 & & Q_1 & & Q_2 & & \cdots & & Q_{n-1} & & Q_n \\
 & & \downarrow t_1 & & \downarrow t_2 & & \cdots & & \downarrow t_{n-1} & & \downarrow t_n \\
 & & E_0[1] & & E_1[1] & & \cdots & & E_{n-2}[1] & & E_{n-1}[1] \\
 & \nearrow id & \downarrow f_1[1] & \nearrow id & \downarrow f_2[1] & \nearrow id & \cdots & \nearrow id & \downarrow f_{n-1}[1] & \nearrow id & \downarrow f_n[1] \\
 & & E_1[1] & & E_2[1] & & \cdots & & E_{n-1}[1] & & E_n[1] \\
 & & \downarrow p_1[1] & & \downarrow p_2[1] & & \cdots & & \downarrow p_{n-1}[1] & & \downarrow p_n[1] \\
 & & \cdots & & \cdots & & \cdots & & \cdots & & \cdots
 \end{array}$$

where the sequences

$$E_i \xrightarrow{f_{i+1}} E_{i+1} \xrightarrow{p_{i+1}} Q_{i+1} \xrightarrow{t_{i+1}} E_i[1]$$

are distinguished triangles for all $0 \leq i \leq n - 1$. We call this diagram the Harder-Narasimhan diagram of E ,

- for any $2 \leq i \leq n$, there exists a real number $c_i < 0$ such that for any two indecomposable summands $Q_{i-1,1}$ and $Q_{i,1}$ of Q_{i-1} and Q_i respectively, there exists a simple connecting path from $Q_{i-1,1}$ to $Q_{i,1}$ in the diagram and for any such simple connecting path l , we have

$$q(l) = c_i < 0.$$

Remarks 3.24. (i) Note that in condition (2), instead of simply requiring that $m(E) > 0$ for all indecomposable objects, we define $\phi(E)$ even when $Z(E) = 0$, this is because we want condition (3) to hold even when one of the central charges of E_1, E_2 is zero.

(ii) A filtration of E as in condition (7) is also called a Harder-Narasimhan filtration of E , Q_i is called the i -th Harder-Narasimhan factor of such a filtration. Its uniqueness will be discussed in Section 5. Also note that we do not require E_i, Q_i to be indecomposable.

(iii) Sometimes, such a stability condition may contain more information than we need. Though these extra information could be encoded in a geometric picture, we still want to have an equivalence relation among such stability conditions. This equivalence relation will also be introduced in Section 5.

(iv) From the definition, one can easily show that for any integer i and any connecting path l between two indecomposable summands of Q_i in condition (7), we have $q(l) = 0$.

(v) To distinguish two different terms of stability conditions, we will call the notion in Definition 3.23 stability conditions, while call the notion in Definition 3.2 Bridgeland stability conditions. Usually, the distinction will be clear in context, depending whether we assume the triangulated category to be cyclic or not. In the next section, we will see that these two notions of stability conditions though different in general, are closely related to each other.

4. THE \mathbb{Z} -LIFTS OF CYCLIC CATEGORIES AND STABILITY CONDITIONS

Let $Stab(\mathcal{C})$ to denote the set of stability conditions on a k -linear Krull-Schmidt cyclic category. In this section, we will investigate some basic properties of $Stab(\mathcal{C})$ and other closely related topological spaces.

We start with the definition of charge triples and charge pairs.

Definition 4.1. Assume that \mathcal{C} is a k -linear Krull-Schmidt cyclic category, then a charge triple $R = (Z, \phi, q)$ consists of a central charge $Z : K_0(\mathcal{C}) \rightarrow \Lambda \rightarrow \mathbb{C}$, a map ϕ sending every indecomposable object E to its phase $\phi(E) \in (0, 2]$, and q a degree function of a real decomposition on \mathcal{C} , which satisfy conditions (1), (2) and (3) in Definition 3.23.

A pair (Z, q) is called a charge pair if for any two indecomposable objects E_1, E_2 with $Z(E_i) = m(E_i)e^{i\theta_i} \neq 0$ for $i = 1, 2$, we have that

$$q(l) = \theta_2 - \theta_1 \pmod{2\mathbb{Z}}$$

for any connecting path $l : E_1 \dashrightarrow E_2$.

Remark 4.2. As we will see Lemma 4.3, in most cases a charge triple (Z, ϕ, q) is determined by its charge pair (Z, q) .

Let us assume that the set of homogeneous morphisms is fixed. And denote the set of charge triples and charge pairs with such a fixed set of homogeneous morphisms on \mathcal{C} by $\mathbf{T}(\mathcal{C})$ and $\mathbf{P}(\mathcal{C})$ respectively. There are natural topologies on these two sets, which are the coarsest topologies such that the forgetful maps

$$\mathbf{T}(\mathcal{C}) \rightarrow \text{Hom}(\Lambda, \mathbb{C}) \simeq \mathbb{C}^{\text{rank}(\Lambda)}, \quad \mathbf{P}(\mathcal{C}) \rightarrow \text{Hom}(\Lambda, \mathbb{C}) \simeq \mathbb{C}^{\text{rank}(\Lambda)},$$

$$\text{where } (Z, \phi, q) \mapsto Z, \quad (Z, q) \mapsto Z,$$

$$\mathbf{T}(\mathcal{C}) \rightarrow \mathbb{R}, \quad \mathbf{P}(\mathcal{C}) \rightarrow \mathbb{R},$$

$$\text{where } (Z, \phi, q) \mapsto q(f), \quad (Z, q) \mapsto q(f),$$

and

$$\mathbf{T}(\mathcal{C}) \rightarrow (0, 2] \xrightarrow{e^{i\pi x}} S^1,$$

$$\text{where } (Z, \phi, q) \mapsto \phi(E)$$

are continuous for any indecomposable object E and any homogeneous morphism f . Here \mathbb{R} , S^1 and $\mathbb{C}^{\text{rank}(\Lambda)}$ are endowed with the standard Euclidean topology. There is also a continuous forgetful map

$$\mathbf{T}(\mathcal{C}) \rightarrow \mathbf{P}(\mathcal{C}).$$

Lemma 4.3. *Let \mathcal{C} be a k -linear Krull-Schmidt connective cyclic category, the continuous map*

$$h : \mathbf{T}(\mathcal{C}) \rightarrow \mathbf{P}(\mathcal{C})$$

is an isomorphism except on the locus where the central charge Z is trivial.

Proof. Suppose that there exists an indecomposable object E with $Z(E) \neq 0$, then $\phi(E)$ is determined by $Z(E)$. For any other indecomposable object F , by the assumption that \mathcal{C} is connective, there exists a connecting path from E to F . Hence $\phi(E)$ and $q(l)$ determines $\phi(F)$. \square

Remark 4.4. One can easily show that on the locus where the central charge is trivial, the map h is an S^1 bundle map. By condition (5) in Definition 3.23, there is no stability conditions above this locus.

In the next subsection, we will show that there are distinguished components in $\mathbf{T}(\mathcal{C})$ with vanishing Maslov indexes, above where the stability conditions can be lifted to be Bridgeland stability conditions.

4.1. Maslov index and \mathbb{Z} liftings. Given a real decomposition on \mathcal{C} , and assume that \mathcal{C} is pre-liftable with respect to the real decomposition. We can define what is a basic loop in such a real decomposition.

Definition 4.5. Assume that \mathcal{C} admits a real decomposition and q is the associated degree function, a fundamental distinguished hexagon in \mathcal{C} is the following locally liftable diagram

$$\begin{array}{ccccc}
 & & \bigoplus_{j=1}^m B_j & \xrightarrow{g} & \bigoplus_{k=1}^n C_k & & \\
 & \nearrow f & & & & \searrow h & \\
 \bigoplus_{i=1}^l A_i & & & & & & \bigoplus_{i=1}^l A_i[1] \\
 & \nwarrow \beta_A \circ h[1] & & & & \swarrow f[1] & \\
 & & \bigoplus_{k=1}^n C_k[1] & \xleftarrow{g[1]} & \bigoplus_{j=1}^m B_j[1] & &
 \end{array}$$

where

$$\bigoplus_{i=1}^l A_i \xrightarrow{f} \bigoplus_{j=1}^m B_j \xrightarrow{g} \bigoplus_{k=1}^n C_k \xrightarrow{h} \bigoplus_{i=1}^l A_i[1]$$

is a distinguished triangle in \mathcal{C} .

A basic loop l in \mathcal{C} is $\beta_{A_i} \circ f[1] \circ f$, where $f : A_i \dashrightarrow A_i[1]$ is a connecting path in the diagram of the distinguished triangle.

Remark 4.6. The same loop can also be defined as basing at B_j or C_k as β is a natural transformation.

We can define what is the Maslov index of a basic loop.

Definition 4.7. Given a basic loop in \mathcal{C} as in Definition 4.5, its Maslov index is

$$M(l) := \frac{q(f) - 1}{2}.$$

As in the definition of Fukaya categories, we will show that if all the Maslov indexes vanish, there is a suitable \mathbb{Z} -lift of \mathcal{C} .

Theorem 4.8. *Given a stability condition $\sigma = (\mathcal{Q}, \mathcal{Z}, \phi, q)$ on a k -linear Krull-Schmidt cyclic category \mathcal{C} , we assume that \mathcal{C} is pre-liftable and the Maslov indexes of all basic loops are zero. There are \mathbb{Z} -lifts of σ and \mathcal{C} , which we denote by σ' and \mathcal{C}' respectively, such that σ' is a Bridgeland stability condition on \mathcal{C}' .*

Proof. The natural \mathbb{Z} -lift of \mathcal{C} is defined in the following way: the indecomposable objects are pairs (E, ϕ) , where E is an indecomposable object in \mathcal{C} and $\phi \in \mathbb{R}$ such that

$$\phi(E) \equiv \phi \pmod{2\mathbb{Z}}.$$

The morphism space between two indecomposable objects is defined by the following formula

$$\text{Hom}_{\mathcal{C}'}((E_1, \phi_1), (E_2, \phi_2)) = \{f \in \text{Hom}_{\mathcal{C}}(E_1, E_2) \mid f \text{ is homogeneous and } q(f) = \phi_2 - \phi_1\}.$$

Moreover, one can easily show that the composition is well defined because of the condition (2) of real decomposition. Note that we can define a functor $\pi : \mathcal{C}' \rightarrow \mathcal{C}$ in a natural way. It sends objects (E, ϕ) to E and be the natural inclusion on the spaces of morphisms.

The triangulated structure on \mathcal{C}' is defined in the following way. The shift functor $[1]$ sends (E, ϕ) to $(E[1], \phi + 1)$, and is the natural isomorphism on maps due to condition (1) of real decomposition.

$$\text{Hom}_{\mathcal{C}'}((E_1, \phi_1), (E_2, \phi_2)) \xrightarrow{\sim} \text{Hom}_{\mathcal{C}'}((E_1[1], \phi_1 + 1), (E_2[1], \phi_2 + 1)).$$

Obviously, one has $\pi \circ [1] = [1] \circ \pi$. We define a triangle in \mathcal{C}' to be a distinguished triangle if and only if its image under π is a distinguished triangle.

We need to check that this indeed define a triangulated category \mathcal{C}' (for the definition of triangulated category, we refer to [Nee01, Chapter 1]).

First of all, assume that we have a morphism

$$f : \bigoplus_{i=1}^l (A_i, \phi_i) \rightarrow \bigoplus_{j=1}^m (B_j, \psi_j)$$

in \mathcal{C}' . The image of f under π is a liftable quasi-homogeneous morphism, hence by the assumption that \mathcal{C} is pre-liftable, there exists a distinguished triangle

$$\bigoplus_{i=1}^l A_i \xrightarrow{\pi(f)} \bigoplus_{j=1}^m B_j \xrightarrow{g} \bigoplus_{k=1}^n C_k \xrightarrow{h} \bigoplus_{i=1}^l A_i[1]$$

in \mathcal{C} , and the diagram is liftable. Therefore, there exists a pre-image of such distinguished triangle in \mathcal{C}'

$$\bigoplus_{i=1}^l (A_i, \phi_i) \xrightarrow{f} \bigoplus_{i=1}^m (B_j, \psi_j) \xrightarrow{g} \bigoplus_{1 \leq k \leq n, d \in \mathbb{Z}} (C_k, \phi(C_k) + 2d) \xrightarrow{h} \bigoplus_{i=1}^l (A[1], \phi_i + 1).$$

The last term can be written as $\bigoplus_{i=1}^l (A[1], \phi_i + 1)$ since all Maslov indexes of basic loops vanish.

Hence axioms **TR1** is satisfied by our assumption that \mathcal{C} is pre-liftable and all Maslov indexes vanish. By Corollary 3.20, we know that **TR3** is also satisfied. The axiom **TR2** hold in \mathcal{C}' since they already hold in \mathcal{C} . The octahedral axiom **TR4** follows from condition (2) in Definition 3.22.

For $\sigma' = (\mathcal{P}, Z)$ on \mathcal{C}' , we define the central charge to be the same one as in σ . And $\mathcal{P}(\phi)$ to be the full subcategory consists of objects (E, ϕ) such that $E \in \mathcal{Q}(\phi')$ where $\phi' \equiv \phi \pmod{2\mathbb{Z}}$. One can easily check that σ' is a Bridgeland stability condition on \mathcal{C}' by unwinding the definitions. \square

Remarks 4.9. (1) *The Octahedral Axiom (TR 4) of \mathcal{C}' holds by the second condition in Definition 3.22. However, in practice, this condition could be difficult to check. The reader could ignore that condition, and take \mathcal{C}' to be just a pre-triangulated category. All the results in this paper holds in that situation, as we do not use the Octahedral Axiom in any of our proofs.*

(2) *We call the functor $\pi : \mathcal{C}' \rightarrow \mathcal{C}$ a \mathbb{Z} -covering since it is similar to the \mathbb{Z} -covering of topological spaces. We will use $\pi^*(\sigma)$ to denote σ' . Also note that the \mathbb{Z} -lift \mathcal{C}' only depends on the charge triple. It is easy to see that the functor π is exact and faithful.*

(3) The functor π induce a surjective homomorphism on Grothendieck groups $\pi_0 : K_0(\mathcal{C})' \twoheadrightarrow K_0(\mathcal{C})$. Hence the central charge $Z : K_0(\mathcal{C})' \xrightarrow{\pi_0} K_0(\mathcal{C}) \xrightarrow{v} \Lambda$ also factors through the given lattice Λ . In the following, we will suppress the dependence of stability conditions on the given lattice Λ . But the reader should be aware of that there is always a implicitly given lattice Λ .

According to Theorem 4.8, the following definitions is natural.

Definition 4.10. (1) We say that \mathcal{C} is liftable with respect to $R = (Z, \phi, q)$ if \mathcal{C} is pre-liftable with respect to the real decomposition in R and all the Maslov indexes vanish.

(2) Let $R_1 = (Z_1, \phi_1, q_1)$ and $R_2 = (Z_2, \phi_2, q_2)$ be two charge triples on \mathcal{C} , if for any two connecting paths $l_1, l_2 : E \dashrightarrow F$, we have that l_1, l_2 are homogeneous with respect to R_1 if and only if l_1, l_2 are homogeneous with respect to R_2 , and

$$q_1(l_1) - q_1(l_2) = q_2(l_1) - q_2(l_2).$$

Then we say that R_1, R_2 are deformation equivalent.

Proposition 4.11. *Let \mathcal{C} be a k -linear Krull-Schmidt connective cyclic category, and $R_1 = (Z_1, \phi_1, q_1)$ and $R_2 = (Z_2, \phi_2, q_2)$ be two charge triples on \mathcal{C} . Suppose that R_1, R_2 are deformation equivalent and \mathcal{C} is liftable with respect to R_1 . Then \mathcal{C} is also liftable with respect to R_2 .*

Moreover, if we denote the \mathbb{Z} -lifts by \mathcal{C}_1 and \mathcal{C}_2 respectively, there exists an equivalence (not canonical) $H : \mathcal{C}_1 \simeq \mathcal{C}_2$.

Proof. By definition, it is easy to see that \mathcal{C} is pre-liftable with respect to R_2 . Moreover, for any basic loop $l_1 : A \dashrightarrow A$ in \mathcal{C} , take $l_2 = id_A$, we get $0 = q_1(l_1) = q_2(l_2)$. Hence \mathcal{C} is also liftable with respect to R_2 .

To construct an equivalence H , let E be a nontrivial indecomposable object in \mathcal{C} , we can define $H : \mathcal{C}_1 \rightarrow \mathcal{C}_2$ in the following way. First of all

$$H((E, \phi_1(E) + 2k)) = (E, \phi_2(E) + 2k),$$

for any $k \in \mathbb{Z}$.

Take another indecomposable object F , there exists a connecting path l starting from E ending at F in \mathcal{C}_1 . We set

$$H((F, \phi_1(E) + q_1(l) + 2k)) := (F, \phi_2(E) + q_2(l) + 2k)$$

for any $k \in \mathbb{Z}$. Our assumption

$$q_1(l_1) - q_1(l_2) = q_2(l_1) - q_2(l_2)$$

ensures that it is independent of the choice of connecting paths. As \mathcal{C} is connective, there exists a connecting path $l' : F \dashrightarrow F[1]$. As $l'' := \bar{\beta} \circ l'[1] : F \dashrightarrow F[1]$ is another connecting from F to $F[1]$. By assumption we know that

$$2q_1(l') = q_1(l') - q_1(l'') = q_2(l') - q_2(l'') = 2q_2(l'),$$

which implies that $q_1(l') = q_2(l')$. Hence we get

$$\begin{aligned} H((F, \phi_1(E) + q_1(l) + q_1(l') + 2k - 1)[1]) &= H((F[1], \phi_1(E) + q_1(l' \circ l) + 2k)) \\ &= (F[1], \phi_2(E) + q_2(l') + q_2(l) + 2k) \\ &= (F, \phi_2(E) + q_1(l') + q_2(l) + 2k - 1)[1] \end{aligned}$$

This implies that $H \circ [1] = [1] \circ H$ on objects.

For the morphisms, assume that

$$f \in \text{Hom}_{\mathcal{C}}(G, G')$$

is a homogeneous morphism between two indecomposable objects $G, G' \in \mathcal{C}$, and

$$l : E \dashrightarrow G$$

is a connecting path from E to G . The morphism f can be lifted in

$$f' \in \text{Hom}_{\mathcal{C}_1}((G, \phi_1(E) + q_1(l) + 2k), (G', \phi_1(E) + q_1(l) + q_1(f) + 2k)),$$

and we define

$$H(f') \in \text{Hom}_{\mathcal{C}_2}((G, \phi_2(E) + q_1(f) + 2k), (G', \phi_2(E) + q_2(l) + q_2(f) + 2k)).$$

From the definition of distinguished triangles in \mathcal{C}_1 and \mathcal{C}_2 , it is easy to see that $H : \mathcal{C}_1 \rightarrow \mathcal{C}_2$ sends distinguished triangles to distinguished triangles. Therefore, H is an exact functor between triangulated categories.

By the symmetry of R_1, R_2 , we can define a functor $G : \mathcal{C}_2 \rightarrow \mathcal{C}_1$ starting from the same indecomposable object $E \in \mathcal{C}$. It is easy to check that H, G are inverse to each other as exact functors between triangulated categories. \square

Remarks 4.12. (i) As we can see in the proof, different choices of E may result in different equivalences, hence the equivalence is not canonical. By the assumption that \mathcal{C} is connective, one can show that there are \mathbb{Z} -copies of equivalences in total.

(ii) This equivalence induces a commutative diagram between coverings.

$$\begin{array}{ccc} \mathcal{C}_1 & \xrightarrow{H} & \mathcal{C}_2 \\ \pi_1 \searrow & & \swarrow \pi_2 \\ & \mathcal{C} & \end{array}$$

(iii) Consistent choices of such equivalences can be viewed as a connection on the \mathbb{Z} -lifts fibered over a connected component of $\mathbf{T}(\mathcal{C})$ where \mathcal{C} is liftable with respect to the charge triples in that component.

5. UNIQUENESS OF HARDER-NARASIMHAN FILTRATION

In this section, we will discuss the uniqueness of Harder-Narasimhan filtration. We start with the following lemma.

Let \mathcal{C} be a k -linear Krull-Schmidt cyclic category, $\sigma = (\mathcal{Q}, Z, \phi, q)$ be a stability condition on \mathcal{C} .

Lemma 5.1. *Let \mathcal{C} be a k -linear Krull-Schmidt connective cyclic category, $\sigma = (\mathcal{Q}, Z, \phi, q)$ be a stability condition on \mathcal{C} . Assume that all the Maslov indexes of basic loops are non-negative and let*

$$\begin{array}{ccccccccc}
 & & \cdots & & \cdots & & \cdots & & \cdots & & \cdots & & \cdots & & \cdots \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 & & E_0 & & E_1 & & \cdots & & E_{n-2} & & E_{n-1} & & & & \\
 \nearrow id & & \downarrow f_1 & \nearrow id & \downarrow f_2 & \nearrow id & & \nearrow id & \downarrow f_{n-1} & \nearrow id & \downarrow f_n & & & & \\
 0 = E_0 & & E_1 & & E_2 & & \cdots & & E_{n-1} & & E_n = E & & & & \\
 & & \downarrow p_1 & & \downarrow p_2 & & \cdots & & \downarrow p_{n-1} & & \downarrow p_n & & & & \\
 & & Q_1 & & Q_2 & & \cdots & & Q_{n-1} & & Q_n & & & & \\
 & & \downarrow & & \downarrow & & \cdots & & \downarrow & & \downarrow & & & & \\
 & & \cdots & & \cdots & & \cdots & & \cdots & & \cdots & & & &
 \end{array}$$

be a Harder-Narasimhan filtration of an indecomposable object E as in condition (7) of Definition 3.23 with $n \geq 2$. Further assume that we have the following liftable commutative diagram between distinguished triangles,

$$\begin{array}{ccccccccccc}
 \cdots & \longrightarrow & E_{i-1} & \xrightarrow{f_i} & E_i & \xrightarrow{p_i} & Q_i & \xrightarrow{t_i} & E_{i-1}[1] & \longrightarrow & \cdots \\
 & & \alpha \downarrow & & \downarrow id & & \downarrow \delta & & \downarrow \alpha[1] & & \\
 \cdots & \longrightarrow & E_{i-1} & \xrightarrow{f_i} & E_i & \xrightarrow{p_i} & Q_i & \xrightarrow{t_i} & E_{i-1}[1] & \longrightarrow & \cdots
 \end{array}$$

and if we glue two copies of Harder-Narasimhan diagrams along each rows in this diagram respectively, we get a liftable commutative diagram. Then $\delta = id$ and $\alpha = id$.

Proof. Firstly, we claim that: there is a connecting path $l : E_{n-1,1} \dashrightarrow E_{n-1,1}[1]$ in the following distinguished triangle

$$E_{n-1} \xrightarrow{f_n} E_n \xrightarrow{p_n} Q_n \xrightarrow{t_n} E_{n-1}[1],$$

where $E_{n-1,1}$ is an indecomposable summand of E_{n-1} .

Let $E_{n-1,1}$ be an arbitrary indecomposable summand of E_{n-1} , if we assume that $pot_n = 0$ where $p : E_{n-1}[1] \rightarrow E_{n-1,1}[1]$ is the natural projection map. We have the following diagram

$$\begin{array}{ccccccc}
 E_{n-1} & \xrightarrow{f_n} & E_n & \xrightarrow{p_n} & Q_n & \xrightarrow{t_n} & E_{n-1}[1] \\
 \downarrow p & & \downarrow f & & \downarrow id & & \downarrow p[1] \\
 E_{n-1,1} & \longrightarrow & Q_n \oplus E_{n-1,1} & \longrightarrow & Q_n & \xrightarrow{0} & E_{n-1,1}[1]
 \end{array}$$

If we can precompose the natural inclusion $i : E_{n-1,1} \rightarrow E_{n-1}$ and postcompose the natural projection $p : Q_n \oplus E_{n-1,1} \rightarrow E_{n-1,1}$ with the diagram, we can see that $E_{n-1,1}$ is a direct summand of E_n . As E_n is indecomposable, we get $E_{n-1,1} \simeq E_n$. Then one can easily show

that there is a morphism $i' : E_n \rightarrow E_{n-1}$ such that $f_n \circ i' = id$, this implies that $p_n = 0$, which contradicts the first condition in Definition 3.23.(7). Hence we show that $p \circ t_n \neq 0$ for any indecomposable direct summand $E_{n-1,1}$ of E_{n-1} .

This easily implies the claim. Indeed, by the connectivity condition in Definition 3.23.(7), there is an indecomposable summand $E_{n-1,1}$ with $f_n|_{E_{n-1,1}} \neq 0$, and any summand p' of p_n is nonzero. Hence we get a connecting loop $l : E_{n-1,1} \dashrightarrow E_{n-1,1}[1]$. By the assumption that all Maslov index are nonnegative, we have $q(l) \geq 1$.

We use \mathbf{D} to denote the glued diagram between Harder-Narasimhan diagrams. As in Remark 3.24.(iv), it is easy to see that every summand of δ is a homogeneous morphism of degree 0. Hence, if we replace δ by $\delta - id$ in \mathbf{D} , the new diagram is still liftable.

By assumption, we get that $(id - \delta) \circ p_i = 0$. Hence $id - \delta$ is in the image of the map

$$Hom_{\mathcal{C}}(E_{i-1}[1], Q_i) \xrightarrow{\circ t_i} Hom_{\mathcal{C}}(Q_i, Q_i),$$

i.e. we can write $id - \delta = \delta_i \circ t_i$, where δ_i can be chosen to be a quasi-homogeneous morphism such that the following diagram is commutative and liftable by Corollary 3.20.

$$\begin{array}{ccccccc} \cdots & \longrightarrow & Q_i & \xrightarrow{t_i} & E_{i-1}[1] & \longrightarrow & \cdots \\ & & \downarrow id-\delta & \swarrow \delta_i & & & \\ & & Q_i & & & & \end{array}$$

where the first row is the Harder-Narasimhan diagram of E .

We assume that $id - \delta \neq 0$, hence $t_i \neq 0, \delta_i \neq 0$. We have the following liftable commutative diagram

$$\begin{array}{ccccc} E_{i-2}[1] & \xrightarrow{f_{i-1}[1]} & E_{i-1}[1] & \xrightarrow{p_{i-1}[1]} & Q_{i-1}[1] \\ & & \downarrow \delta_i & & \\ & & Q_i & & \end{array}$$

Our second claim is that δ_i is not in the image of

$$Hom_{\mathcal{C}}(Q_{i-1}[1], Q_i) \xrightarrow{\circ p_{i-1}[1]} Hom_{\mathcal{C}}(E_{i-1}[1], Q_i).$$

Suppose the contrary, i.e. there exists a nonzero morphism $\epsilon \in Hom_{\mathcal{C}}(Q_{i-1}[1], Q_i)$ with $\delta_i = \epsilon \circ p_{i-1}[1]$. By Corollary 3.20, we can choose ϵ such that the following diagram is commutative and liftable.

$$\begin{array}{ccccc} E_{i-2}[1] & \xrightarrow{f_{i-1}[1]} & E_{i-1}[1] & \xrightarrow{p_{i-1}[1]} & Q_{i-1}[1] \\ & & \downarrow \delta_i & \swarrow \epsilon & \\ & & Q_i & & \end{array}$$

Let $Q_{i-1,1}[1], Q_{i,1}$ be indecomposable summands of $Q_{i-1}[1], Q_i$ respectively such that the homogeneous summand $\epsilon_{11} : Q_{i-1,1}[1] \rightarrow Q_{i,1}$ is nonzero. By the proof of Lemma 3.17,

this implies that there is an indecomposable summand $E_{i-1,1}[1]$ with a connecting path

$$l_1 : Q_{i-1,1}[1] \dashrightarrow E_{i-1,1}[1]$$

in the liftable morphism $E_{i-1}[1] \xrightarrow{p_{i-1}[1]} Q_{i-1}[1]$, and a connecting path

$$l_2 : E_{i-1,1}[1] \dashrightarrow Q_{i,1}$$

in the Harder-Narasimhan diagram of E .

By the connectivity condition in Definition 3.23.(7), one can find a connecting path

$$l_3 : E_{i-1,1} \dashrightarrow Q_{i,1}$$

in the Harder-Narasimhan diagram of E .

By the last condition of Definition 3.23.(7), we know that

$$(2) \quad 0 > c_i = q(l_3 \circ l_1[-1]) = q(l_3) + q(l_1).$$

Once again by the connectivity assumption, one can find a connecting path

$$l_4 : E_{i-1,1} \dashrightarrow E_{n,1}$$

in the Harder-Narasimhan diagram of E . Then we have the following connecting loop.

$$\begin{array}{ccc} E_{i-1,1} & \xrightarrow{\bar{l}_2 \circ l_3} & E_{i-1,1}[1] \\ \downarrow l_4 & & \downarrow l_4[1] \\ E_{n,1} & \xrightarrow{l} & E_{n,1}[1] \end{array}$$

As the Harder-Narasimhan diagram of E is liftable, we know that

$$(3) \quad -q(l_2) + q(l_3) = q(\bar{l}_2 \circ l_3) = q(l) \geq 1 > 0.$$

The inequalities (2) and (3) implies that

$$q(\epsilon_{11}) = q(l_1) + q(l_2) < 0,$$

this implies that $\epsilon_{11} = 0$ by Definition 3.23, which contradicts the assumption $\epsilon_{11} \neq 0$. Hence our second claim is proved. Therefore, we get that

$$\delta_i \circ f_{i-1}[1] \neq 0.$$


We can the same argument as in the proof of our second claim to do induction, at the end, we will get that

$$\delta_i \circ f_{i-1}[1] \circ \cdots \circ f_2[1] \circ f_1[1] \neq 0,$$

which is impossible since $f_1 = 0$. Hence we proved that $\delta - id = 0$.

The proof for $\alpha = id$ is similar. □

Proposition 5.2. *Let \mathcal{C} be a k -linear Krull-Schmidt connective cyclic category, $\sigma = (\mathcal{Q}, Z, \phi, q)$ be a stability condition on \mathcal{C} . Assume that all the Maslov indexes of basic loops are nonnegative and let*




be a Harder-Narasimhan filtration of an indecomposable object E as in condition (7) of Definition 3.23.

Then the compositions $f_j \circ f_{j-1} \circ \cdots \circ f_i \neq 0$ for any $2 \leq i < j \leq n$.

Proof. Without loss of generality, we can let $j = n$. By induction, we can assume that $f_n \circ f_{n-1} \circ \cdots \circ f_{i+1} \neq 0$, and it suffices to prove that $f_n \circ f_{n-1} \circ \cdots \circ f_i \neq 0$. Assume the contrary, i.e. the composition $f_n \circ f_{n-1} \circ \cdots \circ f_i$ is trivial.

Since $f_n \circ f_{n-1} \circ \cdots \circ f_i = 0$, we have the following liftable commutative diagram.



One can easily see that $p_n \circ h_n = 0$ by Definition 3.23. Hence h_n factors through E_{n-1} , we have following liftable diagram

$$\begin{array}{ccccccc}
 & & \cdots & & \cdots & & \cdots & & \cdots & & \cdots \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 & & E_{i-1} & & \cdots & & E_{n-2} & & E_{n-1} & & \cdots \\
 E_{i-1} & \nearrow^{id} & \downarrow f_i & \nearrow^{id} & \cdots & \nearrow^{id} & \downarrow f_{n-1} & \nearrow^{id} & \downarrow f_n & & \cdots \\
 & & E_i & & \cdots & & E_{n-1} & & E_n = E & & \cdots \\
 & & \downarrow p_i & & \nearrow^{h_{n-1}} & & \nearrow^{h_n} & & \downarrow p_n & & \cdots \\
 & & Q_i & & \cdots & & \cdots & & Q_n & & \cdots \\
 & & \downarrow & & \cdots & & \downarrow & & \downarrow & & \cdots \\
 & & \cdots & & \cdots & & \cdots & & \cdots & & \cdots
 \end{array}$$

We claim that this diagram is commutative. It suffices to prove that

$$h_{n-1} \circ p_i = f_{n-1} \circ \cdots \circ f_{i+1}.$$

Denote the difference $h_{n-1} \circ p_i - f_{n-1} \circ \cdots \circ f_{i+1}$ by δ , we have $f_n \circ \delta = 0$. Hence we have the following liftable commutative diagram


$$\begin{array}{ccccccc}
 \cdots & \longrightarrow & E_{i-1} & \xrightarrow{f_i} & E_i & \xrightarrow{p_i} & Q_i & \longrightarrow & \cdots \\
 & & & \swarrow \epsilon & \downarrow \delta & & & & \\
 \cdots & \longrightarrow & Q_n[-1] & \xrightarrow{t} & E_{n-1} & \xrightarrow{f_n} & E_n & \longrightarrow & \cdots
 \end{array}$$

where both rows are in the Harder-Narasimhan diagram of E . By the same argument in the proof of Lemma 5.1, we can show that $\delta = 0$. Hence, the claim is proved.

Continuing the same argument, we get two liftable commutative diagrams

$$\begin{array}{ccccccc}
 & & \cdots & & \cdots & & \cdots & & \cdots & & \cdots \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 & & E_{i-1} & & E_i & & \cdots & & E_{n-1} & & \cdots \\
 E_{i-1} & \nearrow^{id} & \downarrow f_i & \nearrow^{id} & \downarrow f_{i+1} & & \cdots & \nearrow^{id} & \downarrow f_n & & \cdots \\
 & & E_i & & E_{i+1} & & \cdots & & E_n = E & & \cdots \\
 & & \downarrow p_i & & \nearrow^{h_{i+1}} & & \nearrow^{h_n} & & \downarrow p_n & & \cdots \\
 & & Q_i & & \cdots & & \cdots & & Q_n & & \cdots \\
 & & \downarrow & & \cdots & & \downarrow & & \downarrow & & \cdots \\
 & & \cdots & & \cdots & & \cdots & & \cdots & & \cdots
 \end{array}$$

and




Hence we get $f_{i+1} \circ (h_i \circ p_i - id) = 0$. By the same argument in the proof of Lemma 5.1, one can show that $h_i \circ p_i = id$. Hence E_i is a direct summand of Q_i , which implies that $f_i = 0$. This contradicts the connectivity condition. Hence, the proposition is proved. \square

Theorem 5.3. *Let \mathcal{C} be a k -linear Krull-Schmidt connective cyclic category, $\sigma = (\mathcal{Q}, Z, \phi, q)$ be a stability condition on \mathcal{C} . Assume that all the Maslov indexes of basic loops are non-negative.*

Then the Harder-Narasimhan filtration of any indecomposable object E is unique up an isomorphism.

Proof. Let the following liftable diagram



be a Harder-Narasimhan diagram of an indecomposable object E . By Definition 3.23, we know that for any indecomposable summand $Q_{i,1}$ in Q_i , there is a connecting path $l : Q_{i,1} \dashrightarrow E$ in the diagram. By Definition 3.23.(7), $q(l)$ is independent of the path, only depends on the integer i . Hence, we can denote this degree by ϕ_i . Moreover, we also have

$$\phi_1 < \phi_2 < \cdots < \phi_n$$

by definition.

Suppose we have another liftable diagram

$$\begin{array}{cccccc}
 \dots & & \dots & & \dots & & \dots & & \dots \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 & \nearrow id & E_0 & \nearrow id & \dots & \nearrow id & E'_{m-2} & \nearrow id & E'_{m-1} \\
 & & \downarrow f'_1 & & \downarrow f'_2 & & \downarrow f'_{m-1} & & \downarrow f'_m \\
 0 = E_0 & & E'_1 & & E'_2 & & \dots & & E_m = E \\
 & & \downarrow p'_1 & & \downarrow p'_2 & & \downarrow p'_{m-1} & & \downarrow p'_m \\
 & & Q'_1 & & Q'_2 & & \dots & & Q'_m \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \dots & & \dots & & \dots & & \dots & & \dots
 \end{array}$$

which is another Harder-Narasimhan diagram of an indecomposable object E . Similarly we have a increasing sequence of real numbers

$$\phi'_1 > \phi'_2 > \dots > \phi'_m,$$

where ϕ'_i denotes the degree of a connecting path from an indecomposable summand $Q'_{i,1}$ in Q'_i to E in this diagram.

As E is an indecomposable object, we have following liftable diagram.

$$\begin{array}{ccccccc}
 \dots & \longrightarrow & E_{n-1} & \xrightarrow{f_n} & E & \xrightarrow{p_n} & Q_n & \longrightarrow & \dots \\
 & & & & \downarrow id & & & & \\
 \dots & \longrightarrow & E'_{m-1} & \xrightarrow{f_m} & E & \xrightarrow{p'_m} & Q'_m & \longrightarrow & \dots
 \end{array}$$

where the first row is in the first Harder-Narasimhan diagram, and the second row is in the second Harder-Narasimhan diagram.

We claim that $\phi_n = \phi'_m$. To prove the claim, let us assume that $\phi_n < \phi'_m$ first. Under such assumption, one can show that $p'_m \circ f_n = 0$. Indeed, if $p'_m \circ f_n \neq 0$, by the proof in Proposition 5.2 and the assumption that $\phi_n < \phi'_m$, we conclude that

$$p'_m \circ f_n \circ f_{n-1} \circ \dots \circ f_1 \neq 0,$$

which is a contradiction. Hence we get $p'_m \circ f_n = 0$. Therefore, we have a liftable commutative diagram

$$\begin{array}{ccccccc}
 \dots & \longrightarrow & E_{n-1} & \xrightarrow{f_n} & E & \xrightarrow{p_n} & Q_n & \longrightarrow & \dots \\
 & & & & \downarrow id & & \downarrow t & & \\
 \dots & \longrightarrow & E'_{m-1} & \xrightarrow{f_m} & E & \xrightarrow{p'_m} & Q'_m & \longrightarrow & \dots
 \end{array}$$

However, by the assumption $\phi_n < \phi'_m$, we know that any homogeneous summand of t is of negative degree, hence $t = 0$, which implies $p'_m = 0$. This contradicts the connectivity

condition. Therefore, we proved $\phi_n \geq \phi'_m$. And by a symmetric argument, we get $\phi_n = \phi'_m$. This proves the claim.

In the case $\phi_n = \phi'_m$, the proof of Lemma 5.1 could show that $p_n \circ f_m = 0$ and $p'_m \circ f_n = 0$. Hence we have the following liftable and commutative diagram.

$$\begin{array}{ccccccc}
 \cdots & \longrightarrow & E_{n-1} & \xrightarrow{f_n} & E & \xrightarrow{p_n} & Q_n & \longrightarrow & \cdots \\
 & & \downarrow \alpha & & \downarrow id & & \downarrow t & & \\
 \cdots & \longrightarrow & E'_{m-1} & \xrightarrow{f_m} & E & \xrightarrow{p'_m} & Q'_m & \longrightarrow & \cdots \\
 & & \downarrow \alpha' & & \downarrow id & & \downarrow t' & & \\
 \cdots & \longrightarrow & E_{n-1} & \xrightarrow{f_n} & E & \xrightarrow{p_n} & Q_n & \longrightarrow & \cdots \\
 & & \downarrow \alpha & & \downarrow id & & \downarrow t & & \\
 \cdots & \longrightarrow & E'_{m-1} & \xrightarrow{f_m} & E & \xrightarrow{p'_m} & Q'_m & \longrightarrow & \cdots
 \end{array}$$

By Lemma 5.1, we get that $Q_n \simeq Q'_m$ and $E_{n-1} \simeq E'_{m-1}$.

Inductively using the similar argument, one can conclude that $n = m$, $\phi_i = \phi'_i$, $E_i \simeq E'_i$, $Q_i \simeq Q'_i$, and the last two class of isomorphisms are compatible. Hence these two Harder-Narasimhan filtrations are isomorphic. \square

This theorem leads to the following natural definition of stability conditions.

Definition 5.4. A stability condition σ is called a good stability condition if the Harder-Narasimhan filtration of any indecomposable object is unique. We denote such set by $Stab^u(\mathcal{C})$.

Remark 5.5. By Theorem 5.3, we know that $Stab^u(\mathcal{C})$ differs from $Stab(\mathcal{C})$ only on the locus where there are some basic loops with negative Maslov indexes.

Also note that, although the non-negativity pf Maslov indexes is a sufficient condition for the uniqueness of Harder-Narasimhan filtration, it is not a necessary condition.

As mentioned in Remark 3.24.(iii), the space of good stability condition as in Definition 3.23 may contain more information than we need. Therefore, we have following equivalence relation between good stability conditions.

Definition 5.6. Let $\sigma_1 = (\mathcal{Q}_1, Z_1, \phi_1, q_1)$ and $\sigma_2 = (\mathcal{Q}_2, Z_2, \phi_2, q_2)$ be two good stability conditions on \mathcal{C} , we say that σ_1 is equivalent to σ_2 if the following conditions are satisfied:

- (1) The circle slicings and central charges are the same, i.e. $\mathcal{Q}_1 = \mathcal{Q}_2$ and $Z_1 = Z_2$.
- (2) A morphism f is homogeneous with respect to σ_1 if and only if it is homogeneous with respect to σ_2 .
- (3) For any indecomposable objects E , the HN filtration of E with respect to σ_1 is isomorphic to the HN filtration of E with respect to σ_2 .
- (4) For any connecting path l from a semi-stable object to another semi-stable object, we have $q_1(l) = q_2(l)$.

Remark 5.7. It is obviously an equivalence relation. We use $\sigma_1 \simeq \sigma_2$ to denote such a relation.

We denote the equivalent classes of good stability conditions by $Stab^{u,e}(\mathcal{C})$. We have the following diagram.

$$Stab^{u,e}(\mathcal{C}) \leftarrow Stab^u(\mathcal{C}) \hookrightarrow Stab(\mathcal{C}).$$

The following two results are the main reason of introducing this equivalence relation. The notations are the same as in Proposition 4.11 and Remarks 4.12.

Proposition 5.8. *Let \mathcal{C} be a k -linear Krull-Schmidt connective cyclic category, and σ_1, σ_2 be two equivalent good stability conditions on \mathcal{C} . Then the charge triples R_1, R_2 of σ_1 and σ_2 are deformation equivalent. In particular, the Maslov indexes of a basic loop with respect to σ_1, σ_2 are the same.*

Moreover, if \mathcal{C} is liftable with respect to σ_1 (hence also liftable with respect to σ_2). Then there exists a canonical equivalence $H : \mathcal{C}_1 \rightarrow \mathcal{C}_2$ between two \mathbb{Z} -coverings of \mathcal{C} , such that

$$H^* \pi_2^* \sigma_2 = \pi_1^* \sigma_1,$$

where π_1, π_2 are projections in the following commutative diagram.

$$\begin{array}{ccc} \mathcal{C}_1 & \xrightarrow{H} & \mathcal{C}_2 \\ & \searrow \pi_1 & \swarrow \pi_2 \\ & \mathcal{C} & \end{array}$$

Proof. Let l be a connecting path from E to F , and consider the Harder-Narasimhan filtrations of E and F respectively. There are connecting paths $l_1 : Q_{1,1} \dashrightarrow E$ and $l_2 : F \dashrightarrow Q'_{m,1}$, where $Q_{1,1}$ is an indecomposable summand in the first Harder-Narasimhan factor of E and $Q'_{m,1}$ is an indecomposable summand in the last Harder-Narasimhan factor of F . Therefore, we get a connecting path

$$l_2 \circ l \circ l_1 : Q_{1,1} \dashrightarrow Q'_{m,1}.$$

By Definition 5.6.(4), we get

$$q_1(l_2 \circ l \circ l_1) = q_2(l_2 \circ l \circ l_1),$$

which implies that $q_1(l) - q_2(l)$ is a constant real number for any connecting path $l : E \dashrightarrow F$. This proves that R_1, R_2 are deformation equivalent.

For the second half of the proposition, we assume that \mathcal{C} is liftable with respect to σ_1 . Our equivalence H is constructed in the following way.

We start with a semi-stable object $E \in \mathcal{Q}(\phi)$, then proceed as in the proof of Proposition 4.11. One can easily show that

$$H^* \pi_2^* \sigma_2 = \pi_1^* \sigma_1$$

just by unwinding the definitions. We leave the details to the reader. \square

We have the following theorem relating the stability conditions on \mathcal{C} to Bridgeland stability conditions on its \mathbb{Z} -lift.

Theorem 5.9. *Let σ be a stability condition on a connective cyclic category \mathcal{C} such that \mathcal{C} is liftable with respect to σ . By Theorem 4.8, we have a \mathbb{Z} -lift \mathcal{C}' of the cyclic category \mathcal{C} . We use $\text{Stab}^0(\mathcal{C})$ to denote the set of stability conditions whose charge triples are deformation equivalent to the charge triple of σ , and $\text{Stab}^{0,e}(\mathcal{C})$ to denote the equivalent classes $\text{Stab}^0(\mathcal{C})/\simeq$.*

Then there is an isomorphism

$$B\text{Stab}(\mathcal{C}')/\mathbb{Z} \xrightarrow{\sim} \text{Stab}^{0,e}(\mathcal{C}),$$

where $B\text{Stab}(\mathcal{C}')$ denotes the space of Bridgeland stability conditions on \mathcal{C}' , and the \mathbb{Z} action is given by $k \mapsto [2k]$ for any $k \in \mathbb{Z}$.

Proof. Let us first give a map

$$\mathbb{L} : \text{Stab}^{0,e}(\mathcal{C}) \rightarrow B\text{Stab}(\mathcal{C}')/\mathbb{Z}.$$

Fix an indecomposable object $G \in \mathcal{C}$, for any stability condition $\sigma_1 \in \text{Stab}^0(\mathcal{C})$, we have an equivalence

$$H : \mathcal{C}' \rightarrow \mathcal{C}_1$$

by Proposition 4.11, where \mathcal{C}_1 is the \mathbb{Z} -lift of \mathcal{C} with respect to σ_1 . We define

$$\mathbb{L}(\sigma_1) = H^* \pi_1^* \sigma_1,$$

where $\pi_1 : \mathcal{C}_1 \rightarrow \mathcal{C}$ is the natural covering functor. This map \mathbb{L} is well defined on equivalence classes by Proposition 5.8 and the fact that these equivalent functors form a connection (this fact is because we fix an indecomposable object G and \mathcal{C} is connective, see Remark 4.12).

On the other hand, we need to construct a map

$$\mathbb{P} : B\text{Stab}(\mathcal{C}')/\mathbb{Z} \rightarrow \text{Stab}^{0,e}(\mathcal{C}).$$

Given a Bridgeland stability condition $\sigma'_1 = (\mathcal{P}, Z')$ on \mathcal{C}' , we can define a stability condition $\sigma_1 = (\mathcal{Q}, Z, \phi, q)$ on \mathcal{C} in the following way.

For the circle slicing \mathcal{Q} , let $\mathcal{Q}(\phi)$ be the full subcategory with objects $\pi(\mathcal{P}(\phi))$ for any $\phi \in (0, 2]$.

For the central charge, we let $Z(\pi(E')) := Z'(E')$ for any object $E' \in \mathcal{C}'$. As π is surjective on objects, this is defined on all objects in \mathcal{C} .

For the phase function ϕ , let E be an indecomposable object in \mathcal{C} , if

$$Z(E) = m(E)e^{i\pi\phi} \neq 0$$

and $\phi \in (0, 2]$, we let $\phi(E) = \phi$. In the other case, if $Z(E) = 0$, we let $\phi(E)$ to be any real number in $(0, 2]$ and $\phi(E[1]) \equiv \phi(E) + 1 \pmod{2\mathbb{Z}}$. In the end, we will show that this is well defined, i.e. all the possible stability conditions defined in this way are equivalent to each other.

For the real decomposition, one can show that for any two indecomposable objects $E', F' \in \mathcal{C}'$, we define

$$\text{Hom}_{\mathcal{C}}(\pi(E'), \pi(F')) = \bigoplus_{k \in \mathbb{Z}} \text{Hom}_{\mathcal{C}'}(E', F'[2k]).$$

This defines the homogeneous morphisms in \mathcal{C} , we need to assign appropriate degree of each homogeneous morphism. Let $f : E' \rightarrow F'$ be a nonzero morphism between two indecomposable objects in \mathcal{C}' . In the case when $E' \in \mathcal{P}(\phi_1), F' \in \mathcal{P}(\phi_2)$, we define $q(f) = \phi_2 - \phi_1$, which is a positive real number by definition. In other cases, let us take S to be the set of unstable indecomposable objects in \mathcal{C}' such that, any unstable indecomposable object in \mathcal{C}' is isomorphic to $A'[k]$, where $A' \in S$ and $k \in \mathbb{Z}$, and for any two different objects $A', B' \in S$, we have that A' is not isomorphic to $B'[k]$ for any $k \in \mathbb{Z}$.

In the case when $E' \in \mathcal{P}(\phi_1)$, and $F' \simeq A'[k]$ where $A' \in S$. Let us take $A' \xrightarrow{p_n} Q'_n$ to be the natural morphism from A' to its last Harder-Narasimhan factor Q'_n . Then we have the following diagram

$$E' \in \mathcal{P}(\phi_1) \xrightarrow{f} F' \xrightarrow{p_n[k]} Q'_n[k] \in \mathcal{P}(k + \phi_2)$$

We define

$$q(\pi(p_{n,i})) = \phi_2 - \phi(A'), \quad q(\pi(f)) = k + \phi(A') - \phi_1,$$

where $p_{n,i}$ is a summand of the morphism of p_n .

The case when $F' \in \mathcal{P}(\phi_2)$ and $E' \simeq A'[k]$ where $A' \in S$ is similar. Indeed, we have the following diagram in \mathcal{C}'

$$\begin{array}{ccc} E' & \xrightarrow{f} & F' \in \mathcal{P}(\phi_2) \\ \downarrow p_n[k] & & \\ Q'_n[k] & \in & \mathcal{P}(\phi_1 + k) \end{array}$$

where $p_n : A' \rightarrow Q'_n$ is the natural morphism from A' to its last Harder-Narasimhan factor $Q'_n \in \mathcal{P}(\phi_1)$. We define

$$q(\pi(p_{n,i})) = \phi_1 - \phi(A'), \quad q(\pi(f)) = \phi_2 - \phi(A') - k.$$

The last case is when $E' \simeq A'[k_1]$ and $F' \simeq B'[k_2]$, we have the following diagram

$$\begin{array}{ccc} E' & \xrightarrow{f} & F' \\ \downarrow p_n[k_1] & & \downarrow p_m[k_2] \\ Q'_n[k] \in \mathcal{P}(\phi_1 + k_1) & & Q'_m \in \mathcal{P}(\phi_2 + k_2) \end{array}$$

where $p_n : A' \rightarrow Q'_n$, $p_m : B' \rightarrow Q'_m$ are the natural morphisms from A', B' to its last Harder-Narasimhan factor $Q'_n \in \mathcal{P}(\phi_1), Q'_m \in \mathcal{P}(\phi_2)$ respectively. We define

$$q(\pi(f)) = k_2 - k_1 + \phi(B') - \phi(A').$$

This gives the degree function of the real decomposition on \mathcal{C} . Thus, we got our data $\sigma_1 = (\mathcal{Q}, \mathbb{Z}, \phi, q)$ on \mathcal{C} . Before proving that σ is indeed a stability condition on \mathcal{C} and our map \mathbb{P} is well defined, let us state an easy consequence of the construction of degrees.

For any connecting path

$$l : E' \in \mathcal{P}(\phi_1) \dashrightarrow F' \in \mathcal{P}(\phi_2)$$

in \mathcal{C}' , where E', F' are semi-stable indecomposable objects. We have

$$q(\pi(l)) = \phi_2 - \phi_1.$$

This follows directly from the construction of degree function q , we leave the direct check to the reader. We call such property as the invariance of semi-stable path.

There are several things to check for the construction of σ_1 , we list them below.

- (1) The data $\sigma_1 = (\mathcal{Q}, Z, \phi, q)$ is a stability condition on \mathcal{C} ,
- (2) the charge triple (Z, ϕ, q) is deformation equivalent to the charge triple of σ ,
- (3) and the equivalent class of σ_1 is independent of the choices of $\phi(E)$ for $Z(E) = 0$ and the set S .

For (1), most conditions in Definition 3.23 are direct consequences of the construction. We only need to check that the Harder-Narasimhan diagram is connective and liftable. We prove the connectivity by the induction on the number of Harder-Narasimhan factors.

Firstly, if E' is semi-stable in \mathcal{C}' , the connectivity is obvious. Assume that the connectivity condition is true for any indecomposable object E' with $n - 1$ Harder-Narasimhan factors. We consider an indecomposable object E' with n Harder-Narasimhan factors. Let the following distinguished triangle

$$E'_{n-1} \xrightarrow{f_n} E'_n = E' \xrightarrow{p_n} Q'_n \rightarrow E'_{n-1}[1]$$

be the last triangle in the Harder-Narasimhan filtration of E' with respect to σ'_1 . We claim that every summand of f_n or p_n is non-trivial. The proof is essentially included in the proof of Lemma 5.1. We prove the case of f_n for readers' convenience. If there is an indecomposable summand $E'_{n-1,1}$ of E'_{n-1} with $f_n|_{E'_{n-1,1}} = 0$. Then we have the following commutative diagram

$$\begin{array}{ccccccc} E'_{n-1,1} & \xrightarrow{0} & E'_n & \longrightarrow & E'_{n-1,1}[1] \oplus E'_n & \longrightarrow & E'_{n-1,1}[1] \\ \downarrow i & & \downarrow id & & \downarrow & & \downarrow i[1] \\ E'_{n-1} & \xrightarrow{f_n} & E'_n & \xrightarrow{p_n} & Q'_n & \longrightarrow & E'_{n-1}[1] \end{array}$$

This implies that $E'_{n-1,1}[1]$ is a direct summand of Q'_n , but this is impossible unless $E'_{n-1,1} = 0$, as every Harder-Narasimhan factor of E'_{n-1} had bigger phase the phase of Q'_n . Thus the claim is proved. The connectivity is also proved, as the Harder-Narasimhan diagram of any indecomposable summand of E'_{n-1} is connective by induction, and every such summand is connected to E' and the summands of Q'_n .

The liftability of Harder-Narasimhan diagram follows from the connectivity and the invariance of semi-stable path. Indeed, let $l_1, l_2 : A' \dashrightarrow B'$ be two connecting path in the Harder-Narasimhan diagram of E' . There are connecting paths $l_3 : Q_1 \dashrightarrow A'$ and $l_4 : B' \dashrightarrow Q_n$ by connectivity condition. By the invariance of semi-stable path, we get

$$q(l_4 \circ l_1 \circ l_3) = q(l_4 \circ l_2 \circ l_3),$$

hence $q(l_1) = q(l_2)$ and (1) is proved.

For (2), let $l_1, l_2 : E \dashrightarrow F$ be two connecting paths in \mathcal{C} . They can be lifted to connecting paths $l'_1 : E' \dashrightarrow F'$ and $l'_2 : E' \dashrightarrow F'[2k]$ in \mathcal{C}' where $2k = q_0(l_2) - q_0(l_1)$ (q_0 is the degree function in σ). From the definition of q , one can easily show that $q(l_2) - q(l_1) = 2k$. Hence (2) is proved.

For (3), it follows directly from the invariance of semi-stable path. Hence \mathbb{P} is well defined.

By unwinding the definitions, it is easy to see that $\mathbb{L} \circ \mathbb{P} = id$ and $\mathbb{P} \circ \mathbb{L} = id$. Hence, the theorem is proved. \square

Remark 5.10. This equivalent relation is the reason why the phases of unstable objects in Bridgeland stability conditions are not well defined in general. We will explain it more concretely in Section 6.

We end this section by gently touching another aspect of the space $Stab(\mathcal{C})$: chirality of objects and morphisms.

5.1. Chirality of morphisms and objects. There is a natural chirality of morphisms and objects from the definition of charge triples.

Definition 5.11. Given a real decomposition on a cyclic category \mathcal{C} , let f be a homogeneous morphism, we say that f is left chiral if $q(f) > 0$, and f is right chiral if $q(f) < 0$.

Let E be an indecomposable object in \mathcal{C} , we say that E is left chiral if the Maslov indexes of all basic loops involving E are non-negative. Similarly, E is right chiral if the Maslov indexes of all basic loops involving E is negative. In such cases, we say that E has a well-defined chirality.

Lemma 5.12. *Let \mathcal{C} be a k -linear Krull-Schmidt connective cyclic category, and q be a real decomposition on \mathcal{C} such that \mathcal{C} is liftable with respect to q . If moreover, every indecomposable object has a well-defined chirality. Then all indecomposable objects has the same chirality.*

Proof. Suppose that E is a left chiral indecomposable object. For any other indecomposable object F , there is a connecting path between E and F .

Since any nontrivial homogeneous morphism $f : M \rightarrow N$ between two indecomposable objects can be completed to a liftable diagram

$$\cdots \rightarrow M \xrightarrow{f} N \xrightarrow{g} Cone(f) \xrightarrow{h} M[1] \rightarrow \cdots .$$

Then every summand of f and g are nontrivial. As the proof has already appeared twice in the proofs of Lemma 5.1 and Theorem 5.9, we leave it to the reader. This non-triviality implies that the chirality of M is the same as the chirality of N .

Hence, by chasing the connecting path, we can conclude that F is also left chiral. \square

There is a natural involution map τ on the space $\mathbf{T}(\mathcal{C})$, which is defined in the following way:

$$\tau(Z, \phi, q) = (-\bar{Z}, \phi', -q),$$

where \bar{Z} is the conjugate of the central charge Z and $\phi'(E) + \phi(E) = 1 \pmod{2\mathbb{Z}}$ for any indecomposable object E . Obviously, this involution changes the chirality of every homogeneous morphism to its opposite.

Notice that the stability condition breaks the chirality symmetry of real decompositions. In fact, for the forgetful map

$$\text{Stab}(\mathcal{C}) \xrightarrow{\alpha} \mathbf{T}(\mathcal{C}),$$

as we will see in the next section, there are charge triples R with nontrivial preimage $\alpha^{-1}(R)$ but $\alpha^{-1}(\tau(R)) = \emptyset$.

6. STABILITY CONDITIONS ON EQUIVARIANT MATRIX FACTORIZATIONS OF A_2 TYPE

In this section, we will present some examples of stability conditions on cyclic categories. We start with a Schur type lemma in any k -linear category.

Lemma 6.1. *Let \mathcal{T} be any k -linear category, and E, F are two simple objects in \mathcal{T} , i.e. $\text{Hom}_{\mathcal{T}}(E, E) = \text{Hom}_{\mathcal{T}}(F, F) = k$. Then the composition*

$$\text{Hom}_{\mathcal{T}}(E, F) \times \text{Hom}_{\mathcal{T}}(F, E) \rightarrow \text{Hom}_{\mathcal{T}}(E, E) \xrightarrow{\sim} k$$

is trivial unless E is isomorphic to F .

Proof. Suppose that the composition is not trivial, i.e. there exist morphisms $a \in \text{Hom}_{\mathcal{T}}(E, F)$ and $b \in \text{Hom}_{\mathcal{T}}(F, E)$ such that their composition $b \circ a = x \cdot \text{id}_E$, where $x \in k$ is nonzero.

We claim that $a \circ b = x \cdot \text{id}_F$. Indeed, denote $a \circ b := y \cdot \text{id}_F$. Then

$$x^2 \cdot \text{id}_E = (x \cdot \text{id}_E) \circ (x \cdot \text{id}_E) = b \circ a \circ b \circ a = b \circ (y \cdot \text{id}_F) \circ a = xy \cdot \text{id}_E.$$

Hence, we get $x^2 = xy$, which implies $x = y$ since x is nonzero. Therefore, a and $x^{-1}b$ are inverse to each other. \square

Corollary 6.2. *Let \mathcal{C} be a k -linear Krull-Schmidt category, and E be a simple object in \mathcal{C} . Then either the composition*

$$\text{Hom}(E, E[1]) \times \text{Hom}(E, E[1]) \rightarrow \text{Hom}(E, E[2]) \xrightarrow{\sim} k$$

is trivial or $E \xrightarrow{\sim} E[1]$.

Remark 6.3. As one can easily see from the definition of stability conditions, the existence of non-trivial indecomposable objects $E \simeq E[1]$ forms an obstruction of the existence of stability conditions on \mathcal{C} . Unfortunately, this obstruction is common.

However, some finite group action may resolve such obstructions. Let us begin with weighted homogeneous polynomial $w \in \mathbb{C}[[x_1, \dots, x_N]]$ and their Abelian automorphisms (the following definitions are taken from [FJR13, Section 2]).

Definition 6.4. A weighted homogeneous polynomial $w \in \mathbb{C}[[x_1, \dots, x_N]]$ is a polynomial for which there exist positive rational numbers $q_1, \dots, q_n \in \mathbb{Q}_{>0}$, such that for any $\lambda \in \mathbb{C}^*$

$$w(\lambda^{q_1} x_1, \dots, \lambda^{q_N} x_N) = \lambda w(x_1, \dots, x_N).$$

We will call q_j the weight of x_j . We define d and n_i to be the unique positive integers such that $(q_1, \dots, q_N) = (n_1/d, \dots, n_N/d)$ with $\text{gcd}(d, n_1, \dots, n_N) = 1$.

Definition 6.5. We call w nondegenerate if

- (1) the polynomial w contains non monomial of the form $x_i x_j$, for $i \neq j$ and
- (2) the hypersurface defined by $w = 0$ in weighted projective space is non-singular, or equivalently, the affine hypersurface defined by $w = 0$ has isolated singularity at the origin.

Definition 6.6. There are special groups associated with the polynomial w . The first one G_w is defined in the following way

$$G_w := \{(\alpha_1, \dots, \alpha_N) \in (\mathbb{C}^*)^N \mid w(\alpha_1 x_1, \dots, \alpha_N x_N) = w(x_1, \dots, x_N)\}.$$

There is special element $J \in G_w$ which is defined to be

$$J := (\exp(2\pi i q_1), \dots, \exp(2\pi i q_N)),$$

where the q_i are the weights defined in Definition 6.4. For any group G with $\langle J \rangle \leq G \leq G_w$, we call G an admissible subgroup of G_w .

Recall the following famous *ADE* examples.

Example 6.7. *The simple singularities can be classified in the following way.*

- $A_n : w = x^{n+1}, n \geq 1;$
- $D_n : w = x^{n-1} + xy^2, n \geq 4;$
- $E_6 : w = x^3 + y^4;$
- $E_7 : w = x^3 + xy^3;$
- $E_8 : w = x^3 + y^5.$

By [Kn87] and [BGS87], we know that the category $HMF(R, w)$ has only finitely many indecomposable objects if and only if w is of simple singularity.

In the following example, we present some examples of stability conditions on the homotopy category of $\mathbb{Z}/3\mathbb{Z}$ -equivariant matrix factorizations of x^3 . And we use results from the paper [Wal05, Section 4.6, Section 7.1] as our starting point, readers should consult these two sections for the details.


Example 6.8. *We will draw pictures of deforming stability conditions on the homotopy category of $\mathbb{Z}/3\mathbb{Z}$ -equivariant matrix factorizations of A_2 case. In the top left corner of Figure 2, we start with Walcher's point as described in [Wal05]. In the deformation procedure, we keep the central charges $Z(M_1^1)$ and $Z(M_2^1)$ unchanged, move $Z(M_1^2)$ along the dotted path to $Z(M_1^3)$ (central charge of other indecomposable objects change correspondingly), and imagine that there is a pillar standing on the origin such that the homogeneous morphisms could go around the pillar but could not pass through the pillar along deformation.*

In Figure 2, we observe the phenomenon of chirality symmetry breaking. In fact, the charge triples on the left hand side and right hand side are mirror symmetric to each other. But the stability conditions are not symmetric, as we can see in Figure 1, our definition of stability conditions has an implicate orientation, which breaks the symmetry of charge triples.


Chirality symmetry breaking

On the left hand side, at the starting point, every indecomposable object is semi-stable.

On the right hand side, at the starting point, no indecomposable objects are semi-stable. Hence no stability conditions over this charge triple.



On the left hand side, at the end point, M_1^1 and M_2^1 become unstable.



On the right hand side, at the end point, there are two stability conditions over this charge triple. Indeed, we can choose either M_2^3, M_1^3 or M_1^2, M_2^2 to be unstable.

FIGURE 2

Also note that, the charge triples on the left hand side are liftable, while the charge triples on the right hand side are not.

The Figure 3 illustrates the nontrivial monodromy of the map

$$\text{Stab}(\mathcal{C}) \rightarrow \text{Hom}(\Lambda, \mathbb{C}) \simeq \mathbb{C}^{\text{rank}(\Lambda)}.$$

Indeed, if we compare Figure 3 with the left hand side of Figure 2, we see that different paths result in different stability conditions. If we denote the nontrivial monodromy by H , it is easy to see that $H^3 = \text{id}$ in $\text{Stab}(\mathcal{C})$. However, if we go to the lifted space, we get $H^3 = [2]$.




FIGURE 3

REFERENCES

[BBD82] AA Beilinson, J Bernstein, and P Deligne. Faisceaux pervers, analyse et topologie sur les espaces singuliers (i) cirm, 6–10 juillet 1981. *Astérisque*, 100, 1982.

[BGS87] R.-O. Buchweitz, G.-M. Greuel, and F.-O. Schreyer. Cohen-Macaulay modules on hypersurface singularities. II. *Invent. Math.*, 88(1):165–182, 1987.

[BLMS17] Arend Bayer, Martí Lahoz, Emanuele Macrì, and Paolo Stellari. Stability conditions on Kuznetsov components. *arXiv preprint arXiv:1703.10839*, 2017.

- [Bri07] Tom Bridgeland. Stability conditions on triangulated categories. *Ann. of Math. (2)*, 166(2):317–345, 2007.
- [Con85] Alain Connes. Noncommutative differential geometry. *Inst. Hautes Études Sci. Publ. Math.*, (62):257–360, 1985.
- [CQ97] Joachim Cuntz and Daniel Quillen. Excision in bivariant periodic cyclic cohomology. *Invent. Math.*, 127(1):67–98, 1997.
- [DM12] Tobias Dyckerhoff and Daniel Murfet. The Kapustin-Li formula revisited. *Adv. Math.*, 231(3-4):1858–1885, 2012.
- [Dou02] Michael R. Douglas. Dirichlet branes, homological mirror symmetry, and stability. In *Proceedings of the International Congress of Mathematicians, Vol. III (Beijing, 2002)*, pages 395–408. Higher Ed. Press, Beijing, 2002.
- [Dyc11] Tobias Dyckerhoff. Compact generators in categories of matrix factorizations. *Duke Math. J.*, 159(2):223–274, 2011.
- [Eis80] David Eisenbud. Homological algebra on a complete intersection, with an application to group representations. *Trans. Amer. Math. Soc.*, 260(1):35–64, 1980.
- [FJR13] Huijun Fan, Tyler Jarvis, and Yongbin Ruan. The Witten equation, mirror symmetry, and quantum singularity theory. *Ann. of Math. (2)*, 178(1):1–106, 2013.
- [Goo85] Thomas G. Goodwillie. Cyclic homology, derivations, and the free loop space. *Topology*, 24(2):187–215, 1985.
- [HT07] Ryoshi Hotta and Toshiyuki Tanisaki. *D-modules, perverse sheaves, and representation theory*, volume 236. Springer Science & Business Media, 2007.
- [KL03a] Anton Kapustin and Yi Li. D-branes in Landau-Ginzburg models and algebraic geometry. *J. High Energy Phys.*, (12):005, 44, 2003.
- [KL03b] Anton Kapustin and Yi Li. Topological correlators in Landau-Ginzburg models with boundaries. *Adv. Theor. Math. Phys.*, 7(4):727–749, 2003.
- [Kn87] Horst Knörrer. Cohen-Macaulay modules on hypersurface singularities. I. *Invent. Math.*, 88(1):153–164, 1987.
- [KS08] Maxim Kontsevich and Yan Soibelman. Stability structures, motivic Donaldson-Thomas invariants and cluster transformations. *arXiv preprint arXiv:0811.2435*, 2008.
- [Mur13] Daniel Murfet. Residues and duality for singularity categories of isolated Gorenstein singularities. *Compos. Math.*, 149(12):2071–2100, 2013.
- [Nee01] Amnon Neeman. *Triangulated categories*, volume 148 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2001.
- [NS18] Thomas Nikolaus and Peter Scholze. On topological cyclic homology. *Acta Math.*, 221(2):203–409, 2018.
- [Orl04] D. O. Orlov. Triangulated categories of singularities and D-branes in Landau-Ginzburg models. *Tr. Mat. Inst. Steklova*, 246(Algebr. Geom. Metody, Svyazi i Prilozh.):240–262, 2004.
- [Orl09] Dmitri Orlov. Derived categories of coherent sheaves and triangulated categories of singularities. In *Algebra, arithmetic, and geometry: in honor of Yu. I. Manin. Vol. II*, volume 270 of *Progr. Math.*, pages 503–531. Birkhäuser Boston, Boston, MA, 2009.
- [PV12] Alexander Polishchuk and Arkady Vaintrob. Chern characters and Hirzebruch-Riemann-Roch formula for matrix factorizations. *Duke Math. J.*, 161(10):1863–1926, 2012.
- [Tre19] David Treumann. Smith theory and geometric Hecke algebras. *Math. Ann.*, 375(1-2):595–628, 2019.
- [Wal05] Johannes Walcher. Stability of Landau-Ginzburg branes. *J. Math. Phys.*, 46(8):082305, 29, 2005.

BEIJING INTERNATIONAL CENTER FOR MATHEMATICAL RESEARCH, PEKING UNIVERSITY, NO.5 YI-HEYUAN ROAD HAIDIAN DISTRICT, BEIJING, 100871, P.R.CHINA

Email address: liuyucheng@bicmr.pku.edu.cn