

# THE QUERMASSTEGAL PRESERVING MEAN CURVATURE FLOW IN THE SPHERE

ESTHER CABEZAS-RIVAS AND JULIAN SCHEUER

**ABSTRACT.** We introduce a mean curvature flow with global term of convex hypersurfaces in the sphere, for which the global term can be chosen to keep any quermassintegral fixed. Then, starting from a strictly convex initial hypersurface, we prove that the flow exists for all times and converges smoothly to a geodesic sphere. This provides a workaround to an issue present in the volume preserving mean curvature flow in the sphere introduced by Huisken in 1987. We also classify solutions for some constant curvature type equations in space forms, as well as solitons in the sphere and in the upper branch of the De Sitter space.

## 1. INTRODUCTION AND STATEMENT OF MAIN RESULTS

Let  $n \geq 2$  and  $M^n \subset \mathbb{M}_K^{n+1}$  be a smooth, closed, embedded hypersurface in a simply connected space form  $\mathbb{M}_K^{n+1}$  of constant curvature  $K \in \mathbb{R}$ , given by the embedding  $x_0$ . We consider a family of embeddings  $x = x(t, \cdot)$  satisfying the following mean curvature type flow with a global forcing term

$$(1.1) \quad \partial_t x = (\mu(t)c_K(r) - H)\nu,$$

which has initial condition  $x(0, \cdot) = x_0$ . Here  $H$  is the mean curvature and  $\nu$  the outward unit normal of the evolving hypersurfaces  $M_t$ . For convex hypersurfaces (i.e. with  $\kappa_1 \geq 0$ , where  $\kappa_1 \leq \dots \leq \kappa_n$  denote the principal curvatures), the sign conventions are taken so that  $-H\nu$  points inwards. Moreover, let  $r$  denote the radial distance to a given point  $\mathcal{O} \in \mathbb{M}_K^{n+1}$ , which we call *origin* in the sequel. This means that the flow (1.1) depends on the choice of the origin and, in fact, along the flow we will change the origin in a controlled way. We use the notation

$$(1.2) \quad c_K(r) = s'_K(r), \quad \text{where} \quad s_K(r) = \begin{cases} K^{-\frac{1}{2}} \sin(\sqrt{K}r), & \text{if } K > 0 \\ r, & \text{if } K = 0 \\ |K|^{-\frac{1}{2}} \sinh(\sqrt{|K|}r), & \text{if } K < 0. \end{cases}$$

---

*Date:* December 1, 2022. The work of the first author is partially supported by the AEI (Spain) and FEDER project PID2019-105019GB-C21, and by the GVA project AICO 2021 21/378.01/1.

2020 *Mathematics Subject Classification.* 53E10, 53C21.

If  $\sigma_\ell$  represents the  $\ell$ -th elementary symmetric function, we define the time-dependent term by

$$(1.3) \quad \mu(t) = \frac{\int_M H \sigma_\ell dV_t}{\int_M c_K \sigma_\ell dV_t},$$

for each  $\ell = 0, 1, \dots, n$ , where  $dV_t$  denotes the volume element of  $M_t$ . This choice guarantees that (1.1) gives a family of globally constrained mean curvature flows, where  $\mu$  can be chosen to preserve any of the  $n + 1$  quermassintegrals  $W_\ell(\Omega_t)$  of the evolving hypersurfaces  $M_t$  (see section 2 for a review of the quermassintegrals). Here  $\Omega_t$  denotes the convex region enclosed by  $M_t = x(t, \mathbb{S}^n)$  (note that we may assume that the common domain of the embeddings is  $\mathbb{S}^n$ , due to convexity).

Let us stress that for  $\ell = 0$  and  $K = 0$  the flow (1.1) with non-local term as in (1.3) coincides with the *volume-preserving mean curvature flow* (VPMCF) introduced by Huisken in 1987. He proved that strictly convex hypersurfaces in  $\mathbb{R}^{n+1}$  remain convex and embedded under the flow, the solution exists for all times and converges to a round sphere smoothly as  $t \rightarrow \infty$ . Since then it was still an open question of extending the result to an  $(n + 1)$ -dimensional sphere  $\mathbb{S}_K^{n+1}$ ,  $K > 0$ , where convexity can be lost under VPMCF, as pointed out by Huisken [26, page 38].

Our main result settles this question by proposing the flow (1.1) as the most natural generalization of the VPMCF to a space form with positive curvature. Indeed, such a definition preserves convexity under the flow, and allows us to prove the following version of Huisken's original result within the half sphere, where for a point  $p \in \mathbb{S}_K^{n+1}$ ,  $\mathcal{H}(p)$  denotes the open hemisphere around  $p$ .

**1.1. Theorem.** *Let  $n \geq 2$  and  $M_0 \subset \mathbb{S}_K^{n+1}$  be a strictly convex hypersurface enclosing a domain  $\Omega_0$ . Then there exists a finite system of origins  $(\mathcal{O}_i)_{0 \leq i \leq m}$  and numbers  $0 = t_0 < t_1 < \dots < t_m < t_{m+1} = \infty$ , such that the problem*

$$\begin{aligned} \partial_t x &= (\mu_i(t)c_K(r_i) - H)\nu, \quad t \in [t_i, t_{i+1}), \quad 0 \leq i \leq m, \\ x(0, M) &= M_0 \\ x(t_i, M) &= \lim_{t \nearrow t_i} M_t, \quad 1 \leq i \leq m, \end{aligned}$$

where  $r_i$  is the distance to  $\mathcal{O}_i$  and  $\mu_i$  is given as in (1.3) to keep the quermassintegral  $W_\ell(\Omega_t)$  for any  $\ell = 0, 1, \dots, n$  fixed, has a solution

$$x: [0, \infty) \times \mathbb{S}^n \rightarrow \mathbb{S}_K^{n+1}.$$

For every  $t \geq 0$ , the embeddings  $x(t, \cdot)$  smoothly map  $\mathbb{S}^n$  to strictly convex hypersurfaces with

$$\mathcal{O}_i \in \Omega_t \quad \text{and} \quad M_t \subset \mathcal{H}(\mathcal{O}_i) \quad \forall t \in [t_i, t_{i+1})$$

and satisfy spatial  $C^\infty$ -estimates which are uniform in time. The restriction

$$x: [t_m, \infty) \times \mathbb{S}^n \rightarrow \mathbb{S}_K^{n+1}$$

is smooth and converges for  $t \rightarrow \infty$  in  $C^\infty$  to a geodesic sphere around  $\mathcal{O}_m$  with radius determined by  $W_\ell(B_r) = W_\ell(\Omega_0)$ .

At this stage we should mention that Guan and Li [24] invented a *purely local* mean curvature type flow in the sphere, which is volume-preserving and drives starshaped hypersurfaces to geodesic spheres. There is also a flow of Guan-Li type that preserves  $W_{\ell-1}(\Omega_t)$  and decreases  $W_\ell(\Omega_t)$ , which has so far refused to allow curvature estimates (see [12] for an overview over known results).

However, notice that non-local flows are more challenging than their corresponding local counterparts, as the evolution depends heavily on the global shape of the hypersurface  $M_t$  and the presence of the term  $\mu(t)$  in all the relevant evolution equations causes a plethora of extra complications; e.g. comparison principles and preservation of key properties fail (cf. [11]), as well as embedded hypersurfaces may develop self-intersections (cf. [33]).

In this framework, our flow (1.1) is, to the best of our knowledge, the first known curvature flow in the sphere, which preserves any desired quermass-integral by a suitable choice of the global term  $\mu$  and which enjoys smooth convergence to a geodesic sphere.

**Elliptic counterpart: rigidity results.** As a byproduct of the pinching estimates (see Proposition 4.3) that we require to prove Theorem 1.1, we can also classify hypersurfaces in a space form  $\mathbb{M}_K^{n+1}$  which have a rotationally symmetric curvature function, under suitable assumptions on the sectional curvature.

With this goal, we work with curvature functions more general than  $H$ . Let  $\Gamma \subset \mathbb{R}^n$  be a symmetric, open cone containing the positive cone

$$\Gamma_+ = \{\kappa \in \mathbb{R}^n : \kappa_i > 0, \text{ for all } i = 1, \dots, n\},$$

and consider a symmetric function  $f \in C^2(\Gamma)$ . Let

$$F(A) = f(\kappa_1, \dots, \kappa_n)$$

be the corresponding operator dependent function, where  $A$  denotes the Weingarten or shape operator. We assume the following:

**1.2. Conditions.** Let  $f(\kappa) = F(A)$  be a  $C^2$  symmetric function, defined on an open, symmetric cone  $\Gamma \supset \Gamma_+$ . We ask further that

- (a)  $f$  is strictly increasing in each argument.
- (b)  $f$  is homogeneous of degree 1.
- (c)  $f$  is normalized so that  $f(1, \dots, 1) = n$ .

Notice that (a) implies that  $F$  defines a strictly elliptic operator on  $M$ , as proved in [28]. We say that  $f$  is *inverse concave/convex* if the dual function

$$\tilde{f}(\kappa_1, \dots, \kappa_n) = f(\kappa_1^{-1}, \dots, \kappa_n^{-1})$$

is concave/convex (see section 8 for a more detailed introduction).

A classical result by Alexandrov [1] says that if a compact hypersurface embedded in  $\mathbb{R}^{n+1}$  has  $H = \text{constant}$ , it must be a round sphere. Later on,

Ros [38] extended this result to the constancy of higher order symmetric functions  $\sigma_\ell$ . Hypersurfaces in a model space  $\mathbb{M}_K^{n+1}$  which have a constant curvature function  $F$  were often called *Weingarten hypersurfaces* in the previous literature. It was shown in [16, Theorem 28] that round spheres are the only examples of compact Weingarten hypersurfaces in the hyperbolic space  $\mathbb{H}^{n+1}$ . In this spirit, we obtain similar rigidity results for  $F$  radially symmetric instead of constant:

**1.3. Theorem.** *Let  $n \geq 2$ ,  $K \in \mathbb{R}$ ,  $|\alpha| \geq 1$  and  $M^n \subset \mathbb{M}_K^{n+1}$  be a convex hypersurface, which is located in the northern hemisphere for  $K > 0$ , such that*

$$\text{Sec}_M \geq -\alpha K.$$

*Suppose that  $F$  is a convex function satisfying [Conditions 1.2](#), which is a solution to*

$$F = \gamma c_K^\alpha$$

*for some constant  $\gamma$ . Then  $M$  is a geodesic sphere, which is centred at the origin provided  $K \neq 0$ . If  $\alpha = 1$ , the convexity assumption on  $M$  can be dropped, while for  $\alpha = -1$  the convexity of  $F$  may be replaced by inverse concavity.*

Such results have been obtained for  $\sigma_\ell$  by integral methods: for instance, [46] studies constant linear combinations of higher order mean curvatures, [45] analyses constancy of  $c_K \sigma_\ell$  in  $\mathbb{H}^{n+1}$ , and [29] deals with hypersurfaces having radially symmetric higher order mean curvatures in general  $\mathbb{M}_K^{n+1}$  under mild convexity assumptions. But those integral techniques are restricted to the  $\sigma_\ell$  because they are divergence free in spaceforms. Our maximum principle approach enables us to relax the assumptions on the curvature functions, at the cost of having to impose a condition on the sectional curvature of the hypersurface. However, notice that if  $(1 + \alpha)K > 0$  this assumption is weaker than convexity, while if this product is  $\leq 0$  the condition already implies convexity.

**Classification of solitons.** In the study of singularity formation along curvature flows, especially the mean curvature flow, the class of *self-shrinking solutions*, simply called *solitons* subsequently, plays an important role. For the mean curvature flow in Euclidean space they arise as blow-up limits of type-I singularities (see [27]), and they satisfy the elliptic equation

$$H = \langle x, \nu \rangle.$$

Huisken [27] showed that the only compact mean-convex solitons are spheres.

A similar recent result when  $H$  is replaced by the Gauss curvature  $K$ , see [9], settled the long standing open problem of whether the flow by certain powers of the Gauss curvature of  $n$ -dimensional hypersurfaces,  $n \geq 3$ , converges to a round sphere; the convergence to a soliton had already been proved in [7].

The study of solitons for more general curvature functions has received plenty of attention, also in spaceforms; see e.g. [17, 18, 19, 35]. Here one considers the general equation

$$(1.4) \quad F^\beta = u,$$

where  $\beta \in \mathbb{R}$ ,  $F$  is a function of the principal curvatures with suitable assumptions and

$$(1.5) \quad u = s_K(r) \langle \partial_r, \nu \rangle$$

is the generalised support function. From a well-known duality relation by means of the Gauss map for hypersurfaces of the sphere and itself, respectively a duality relation between hypersurfaces of the hyperbolic and De Sitter space, from [Theorem 1.3](#) we can deduce a new classification result for convex solitons in the sphere  $\mathbb{S}_1^{n+1}$ , as well as in the upper branch of the  $(n+1)$ -dimensional De Sitter space  $\mathbb{S}^{n,1}$  with sectional curvature  $K = 1$ , i.e.

$$\mathbb{S}^{n,1} = \{y \in \mathbb{R}^{n+2}: -(y^0)^2 + \sum_{i=1}^{n+1} (y^i)^2 = 1, y^0 > 0\}.$$

More precisely, with the notation

$$(1.6) \quad \text{sgn}(\mathbb{M}) = \begin{cases} 1, & \mathbb{M} = \mathbb{S}_1^{n+1}, \\ -1, & \mathbb{M} = \mathbb{S}^{n,1}, \end{cases}$$

we prove the following result

**1.4. Corollary.** *Let  $n \geq 2$ ,  $|\beta| \leq 1$ ,  $\beta \neq 0$  and  $\mathbb{M}$  be either  $\mathbb{S}_1^{n+1}$  or  $\mathbb{S}^{n,1}$ . Consider  $M^n \subset \mathbb{M}$  a closed strictly convex hypersurface, and in case  $\frac{1-\beta}{\beta} \text{sgn}(\mathbb{M}) > 0$ , we assume further that*

$$\text{Sec}_M \leq \frac{\text{sgn}(\mathbb{M})}{1-\beta}.$$

*If  $F$  is an inverse convex function satisfying [Conditions 1.2](#), which is a solution to the soliton equation (1.4), then  $M$  is a geodesic sphere centred at the origin. In case  $\beta = 1$ , the inverse convexity may be replaced by concavity.*

**1.5. Remark.** [Corollary 1.4](#) is remarkable in several ways. Firstly, to our knowledge this is the first such result, where the  $\beta$ -regime ranges down to zero. This is surprising, as in the Euclidean space, for  $F = K^{1/n}$  and  $\beta \leq n/(n+2)$  the result is false, see [4, 9]. Note however that  $K^{1/n}$  is not inverse convex. Secondly, in all of the previous results of this type, the inverse concavity of  $F$  was exploited crucially. The duality approach allows us to deal with a further class of curvature functions, which could not be treated by earlier methods.

Notice that, while Weingarten hypersurfaces are known to be geodesic spheres in  $\mathbb{S}^{n,1}$  (see [37]), we are not aware of rigidity results for solitons in this setting. On the other hand, to have some model examples in mind,

$F = |A|$  satisfies the assumptions of [Theorem 1.3](#) and the harmonic mean curvature is suitable for [Corollary 1.4](#).

**The problem of extending a non-local flow to curved spaces.** As said before, Huisken [\[26\]](#) introduced the VPMCF of convex hypersurfaces in the Euclidean space,

$$(1.7) \quad \partial_t x = (\mu(t) - H) \nu,$$

where the global term is the average mean curvature  $\mu = \overline{H} = \int_{M_t} H$ . Taking  $\mu$  as in [\(1.3\)](#) for  $K = 0$ , McCoy [\[34\]](#) obtained convergence of convex hypersurfaces in  $\mathbb{R}^{n+1}$  to round spheres under a flow that preserves any quermassintegral (which in the Euclidean case coincide with the mixed volumes, see [section 2](#)).

Huisken already pointed out that an interesting problem is to extend his result to non-Euclidean ambient spaces, with the warning that the generalization will not be straightforward because [\(1.7\)](#) does not preserve convexity in general Riemannian manifolds, due to terms with an unfavourable sign in the evolution equation of the second fundamental form. In particular, for hypersurfaces in  $\mathbb{M}_K^{n+1}$  the Weingarten matrix  $h_j^i$  evolves according to

$$(\partial_t - \Delta)h_j^i = (|A|^2 - nK)h_j^i + 2KH\delta_j^i - \mu(h_\ell^i h_j^\ell + K\delta_j^i).$$

Notice that for  $K < 0$  the bad term is  $2KH\delta_j^i$ , which comes from the background geometry and causes that convexity is not preserved in general. This failure is independent of the non-local nature of the flow; indeed, if we replace convexity by  $h$ -convexity ( $\kappa_1 > |K|$ ), Miquel and the first author [\[10\]](#) proved that  $h$ -convex hypersurfaces can be deformed under [\(1.7\)](#) to a geodesic sphere; this was extended by Andrews and Wei [\[8\]](#) for a class of quermassintegral preserving flows. The curvature condition was relaxed to positive sectional curvature ( $\kappa_1\kappa_2 > |K|$ ) by Andrews, Chen and Wei in [\[6\]](#) in the volume-preserving case.

Notice that the complication for  $K > 0$  is of completely different nature, since the fatal term is now  $-\mu K\delta_j^i$ , and thus comes directly from the global term. Indeed, Huisken illustrated this [\[26\]](#) with an intuitive example: if the flow starts with a convex hypersurface of  $\mathbb{S}^{n+1}$  with a portion  $M^*$   $C^2$ -close to the equator, then in this region  $\overline{H} \gg H$  and hence  $M^*$  moves in the outward direction crossing the equator, and thus the evolving hypersurface becomes non-convex.

This obstruction to the preservation of convexity in an ambient sphere supports the claim that the flow [\(1.7\)](#) is, geometrically, not the most natural generalisation of the same flow in Euclidean case to the spherical ambient space. Indeed, our alternative flow [\(1.1\)](#) does preserve pinching of the principal curvatures and hence it succeeds in driving any convex initial hypersurface to a geodesic sphere. Notice that Huisken's example is actually the motivation for the definition of [\(1.1\)](#), as the effect of multiplying the global

term by  $c_K(r)$  is to slow down the motion as the hypersurfaces approach the equator.

In short, to extend Huisken's results to the hyperbolic space one needs to strengthen the notion of convexity, whereas for the ambient sphere we propose a different generalization of the flow (notice that (1.1) and (1.7) coincide for the Euclidean space), which works for convex hypersurfaces.

**The isoperimetric nature of the flow.** In addition, under (1.7) the surface area is non-increasing, and hence Huisken's theorem provides an alternative proof of the isoperimetric inequality for convex hypersurfaces of  $\mathbb{R}^{n+1}$ . An interesting side-effect of the extra term in (1.1) is that this flow is no longer of isoperimetric nature in the classical sense, because if we choose  $\mu$  to preserve enclosed volume, the surface area is no longer decreasing necessarily.

However, the flow (1.1) with global term chosen to preserve the weighted volume  $\int_{\Omega_t} c_K$ , has decreasing surface area, which suggests that in principle it is the right flow to prove the isoperimetric type inequality

$$\int_{\Omega_0} c_K \leq \phi(|M_0|)$$

with equality if and only if  $\Omega_0$  is a ball centred at the origin. Here  $\phi$  is a function that gives equality on the slices. This was originally shown in [23, Proposition 4] by other means and hence we do not pursue any further investigation in this matter here.

This reinforces the idea that our new flow has a geometric meaning beyond the generalization of Huisken's result, and we hope that in the future some interesting new applications will follow.

**Structure of the paper.** The contents of this paper are organized as follows. We first introduce in section 2 the basic notation and evolution equations that ensure that our flow preserves the quermassintegrals, while section 3 gathers new estimates for strictly convex hypersurfaces in the sphere, which may be of independent interest, like a refined outradius bound (Theorem 3.3) or inradius control in terms of pinching (Corollary 3.2). Then in section 4 we prove that the *pinching deficit* decreases exponentially under the flow as time evolves, which is the key to get convergence of the evolving hypersurfaces. To achieve upper curvature bounds, we perform a technically intricate process in section 5, which includes a delicate iterative changing of origin to ensure an optimal configuration that enables us to gain some uniform bound on the global term for some controlled time interval. This is a novel method, providing an alternative to proving initial value independent curvature bounds after a waiting time. To finish the proof of Theorem 1.1, in section 6 we establish long time existence and convergence to a geodesic sphere is done in section 7. Finally, the elliptic results are proved in section 8.

## 2. NOTATION, CONVENTIONS AND PRELIMINARY RESULTS

**Hypersurfaces in space forms.** Let  $x: M \hookrightarrow \mathbb{M}_K^{n+1}$  be an embedded smooth hypersurface in a simply connected space form  $\mathbb{M}_K^{n+1}$  enclosing a bounded domain  $\Omega$ . Then the metric in polar coordinates is given by

$$\bar{g} = dr^2 + s_K^2(r)\sigma,$$

where  $r$  is the radial distance to a fixed point  $\mathcal{O} \in \mathbb{M}_K^{n+1}$  and  $\sigma$  is the round metric on  $\mathbb{S}^n$ .

The trigonometric functions in (1.2) satisfy the following computational rules:

$$c'_K = -K s_K, \quad c_K^2 + K s_K^2 = 1.$$

We will also use the related notation  $\text{co}_K(r) = c_K(r)/s_K(r)$ .

For the outward pointing unit normal  $\nu$ , we define the second fundamental form  $h = (h_{ij})$  by

$$\bar{\nabla}_X Y = \nabla_X Y - h(X, Y)\nu,$$

where  $\bar{\nabla}$  is the Levi-Civita connection of the metric  $\bar{g} = \langle \cdot, \cdot \rangle$  on  $\mathbb{M}_K^{n+1}$  and  $X, Y$  are vector fields on  $M$ . We adopt the summation convention throughout and latin indices indicate components with respect to a coordinate frame  $(\partial_i)_{1 \leq i \leq n}$  on the domain of the embedding  $x$ .

If the induced metric on  $M$  is denoted by  $g$ , then we write  $\Delta$  for its Laplace-Beltrami operator and define the Weingarten operator  $A = (h_j^i)$  via

$$h_{ij} = g(A(\partial_i), \partial_j) = g_{ik}h_j^k.$$

Recall that the symmetry of  $h$  and the *Codazzi equations*

$$\nabla_i h_{jk} = \nabla_j h_{ik}$$

imply that the tensor  $\nabla A$  is totally symmetric. Moreover, one can relate the geometry of a hypersurface  $M$  with the ambient manifold  $\mathbb{M}_K^{n+1}$  by means of the *Gauss equation*

$$(2.1) \quad R_{ijkl} = h_{ik}h_{jl} - h_{il}h_{jk} + K(g_{ik}g_{jl} - g_{il}g_{jk}).$$

On the other hand, if

$$\kappa_1 \leq \dots \leq \kappa_n$$

denote the eigenvalues of the operator  $A$ , that is, the principal curvatures of  $M$ , we consider the normalised mean curvatures  $H_\ell$  defined as

$$H_\ell = \binom{n}{\ell}^{-1} \sigma_\ell, \quad \text{with} \quad \sigma_\ell = \sum_{1 \leq i_1 < \dots < i_\ell \leq n} \kappa_{i_1} \dots \kappa_{i_\ell}.$$

In particular,  $H_1 = H/n$  and  $H_n$  equals the Gauss curvature. We use the convention that  $H_0 = 1$ . For convex hypersurfaces, these symmetric functions satisfy the *Newton-MacLaurin* inequalities [44]:

$$(2.2) \quad H_{\ell-1}H_k \geq H_\ell H_{k-1} \quad \text{for} \quad 1 \leq k < \ell \leq n.$$



We will also use the following *Hsiung-Minkowski* identities [24]:

$$(2.3) \quad (\ell + 1) \int_M u \sigma_{\ell+1} = (n - \ell) \int_M c_K \sigma_\ell$$

for  $\ell = 0, \dots, n - 1$ .

Later on, we need control of the support function from below, given that there is a uniform ball enclosed by the evolving domain. Fortunately, for strictly convex domains such control is easy to obtain. We furnish quantities like  $r$  and  $u$  with a hat, if their reference point is not the origin. The right reference point will then be displayed as a subscript, and in cases where the reference point is clear by context, it is suppressed

**2.1. Lemma.** *Let  $\Omega \subset \mathbb{M}_K^{n+1}$  be a strictly convex domain with  $p \in \Omega$  and  $M = \partial\Omega$ . Then the support function*

$$\hat{u}_p = s_K(\hat{r}_p) \langle \partial_{\hat{r}_p}, \nu \rangle,$$

where  $\hat{r}_p$  is the distance to the point  $p$ ,

$$\hat{u}_p \geq \min_M \hat{u}_p = \min_M s_K(\hat{r}_p) = s_K(\text{dist}(p, M)).$$

*Proof.* At a global minimum of the support function we have  $\nabla \hat{u}_p = 0$ . It is well known (cf. [24]) that

$$(2.4) \quad \nabla_i \hat{u}_p = h_i^j \nabla_j \left( \frac{1 - c_K}{K} \right) = -h_i^j \frac{c'_K}{K} \nabla_j \hat{r}_p = s_K(\hat{r}_p) h_i^j \nabla_j \hat{r}_p.$$

Accordingly, due to the invertability of  $A$ , we also have  $\nabla \hat{r}_p = 0$ . Hence at such a point, and for  $K > 0$ ,

$$\hat{u}_p = s_K(\hat{r}_p) \geq \min(s_K(\min_M \hat{r}_p), s_K(\max_M \hat{r}_p)),$$

due to the concavity of  $s_K$  within the interval  $[0, \pi/\sqrt{K}]$ . In case  $K \leq 0$ ,  $s_K$  is increasing, so in this case we are done. Now suppose that

$$(2.5) \quad s_K(\max_M \hat{r}_p) < s_K(\min_M \hat{r}_p).$$

Due to the symmetries of the sine function we must then have

$$\min_M \hat{r}_p > \frac{\pi}{\sqrt{K}} - \max_M \hat{r}_p.$$

At a point  $\xi \in M$ , where  $\hat{r}_p$  is maximized, the geodesic which connects  $p$  and  $\xi$ , intersects  $M$  in another point, say  $\zeta \in M$ . Due to the convexity of  $M$  there holds

$$\frac{\pi}{\sqrt{K}} > \text{dist}(\zeta, \xi) = \hat{r}_p(\zeta) + \hat{r}_p(\xi) \geq \hat{r}_p(\zeta) + \frac{\pi}{\sqrt{K}} - \min_M \hat{r}_p \geq \frac{\pi}{\sqrt{K}},$$

a contradiction. Hence (2.5) can not be valid and the proof is complete.  $\square$

**Mixed volumes and quermassintegrals.** We define the *curvature integrals* or *mixed volumes* as follows:

$$V_{n-\ell}(\Omega) = \int_M H_\ell dV, \quad \text{for } \ell = 0, \dots, n.$$

On the other hand, for any connected domain  $\Omega \subset \mathbb{M}_K^{n+1}$  bounded by a compact hypersurface of class  $C^3$ , the *quermassintegrals* are given by (see [42] or [41, chapter 17]):

$$(2.6) \quad W_\ell(\Omega) = \frac{(n+1-\ell)\omega_{\ell-1} \cdots \omega_0}{(n+1)\omega_{n-1} \cdots \omega_{n-\ell}} \int_{\mathcal{L}_\ell} \chi(L_\ell \cap \Omega) dL_\ell, \quad \ell = 1, \dots, n.$$

Here  $\mathcal{L}_\ell$  represents the space of  $\ell$ -dimensional totally geodesic subspaces  $L_\ell$  in  $\mathbb{M}_K^{n+1}$ , where one can define a natural invariant measure  $dL_\ell$ , and  $\omega_n = |\mathbb{S}^n|$  is the area of  $n$ -dimensional unit sphere in  $\mathbb{R}^{n+1}$ . If  $\Omega$  is a convex set, then the function  $\chi$  is equal to 1 if  $L_\ell \cap \Omega \neq \emptyset$ , and 0 otherwise.

One typically sets

$$W_0(\Omega) = |\Omega| \quad \text{and} \quad W_{n+1}(\Omega) = \frac{\omega_n}{n+1}.$$

Moreover, using the Cauchy-Crofton formula (cf. [41]) we recover the area of the hypersurface as

$$|\partial\Omega| = (n+1)W_1(\Omega).$$

Accordingly, volume and area-preserving flows can be regarded as particular cases of quermassintegral preserving flows.

Mixed volumes and quermassintegrals are related (see [42, Proposition 7]) in a space of constant curvature  $\mathbb{M}_K^{n+1}$  by means of

$$(2.7) \quad \begin{aligned} \frac{1}{n+1}V_{n-\ell}(\Omega) &= W_{\ell+1}(\Omega) - K \frac{\ell}{n+2-\ell} W_{\ell-1}(\Omega), \quad \ell = 1, \dots, n, \\ V_n(\Omega) &= (n+1)W_1(\Omega) = |\partial\Omega|. \end{aligned}$$

Notice that in  $\mathbb{R}^{n+1}$  the mixed volumes coincide with the quermassintegrals, up to a constant factor.

The following result gathers the evolution equations of the above defined quantities under a normal variation.

**2.2. Lemma.** *If  $M_t$  is a hypersurface of  $\mathbb{M}_K^{n+1}$  evolving along a flow given by  $\partial_t x = \varphi \nu$ , then*

- (a)  $\partial_t \text{Vol}(\Omega_t) = \int_M \varphi dV_t \quad \text{and} \quad \partial_t |M_t| = \int_M \varphi H dV_t.$
- (b)  $\partial_t \int_M H_\ell dV_t = \int_M \varphi ((n-\ell)H_{\ell+1} - K\ell H_{\ell-1}) dV_t, \quad \ell = 0, \dots, n.$
- (c)  $\partial_t W_\ell(\Omega_t) = \frac{n+1-\ell}{n+1} \int_M \varphi H_\ell dV_t, \quad \ell = 0, \dots, n.$
- (d)  $\partial_t g_{ij} = 2\varphi h_{ij}.$
- (e)  $\partial_t h_j^i = -g^{ik} \nabla_{kj}^2 \varphi - \varphi h_k^i h_j^k - K\varphi \delta_j^i.$

*Proof.* Formulas in (a) and (b) were deduced in [36]. The evolution in (c) follows arguing by induction on  $\ell$  and using the relation (2.7); this was done in [44, Proposition 3.1] for  $K = -1$ . The evolution for the metric and the Weingarten operator are standard, e.g. [20, chapter 2].  $\square$

**2.3. Corollary.** *If the global term in (1.1) is chosen as in (1.3), then the quermassintegral  $W_\ell(\Omega_t)$  is constant along the flow (1.1).*

The fact that  $\mu(t) > 0$  for strictly convex hypersurfaces is heavily used within the proof of **Theorem 1.1**.

**2.4. Remark.** Notice that a global term given by

$$\mu(t) = \frac{\int_M H((n-\ell)H_{\ell+1} - K\ell H_{\ell-1}) dV_t}{\int_M c_K((n-\ell)H_{\ell+1} - K\ell H_{\ell-1}) dV_t}$$

leads to a flow that preserves the mixed volume  $V_{n-\ell}(\Omega_t)$ . Unlike the quermassintegral-preserving case, this term does not have a sign for convex hypersurfaces if  $K > 0$ . For  $K < 0$  this difficulty disappears, but another type of mixed volume preserving curvature flows for  $h$ -convex hypersurfaces in the hyperbolic space was already studied in [31].

We use the following conventions for the use of constants. Indexed letters  $C$ , i.e.  $C_0, C_1$ , etc. will retain a specific meaning throughout the whole paper, while the letter  $C$  denotes a generic constant, which is always allowed to change from line to line and depends on the quantities listed in the formulation of the lemma or theorem. Capital letters also stand for “large” constants. A similar convention holds for lower case letters, which stand for “small” constants. The only exception from this convention concerns the use of various versions of the letter  $t$ , like  $T, \tau, \hat{\tau}$  etc. Those always refer somehow to time and  $t$  denotes the time variable, while  $T, \tau, \hat{\tau}$  etc. will, once defined, not change value.

### 3. GEOMETRY AND LOCATION OF STRICTLY CONVEX HYPERSURFACES IN THE SPHERE

This section presents some geometric results for strictly convex hypersurfaces of the sphere, which are required to prove **Theorem 1.1**. In particular, we obtain inradius estimates in terms of pinching, as well as a suitable out-ball configuration in terms of pinching and the value of any given  $W_\ell(\Omega)$ . **Throughout section 3, section 4 and section 5, we make the standing assumption that  $M \subset \mathbb{S}_K^{n+1}$  is a strictly convex hypersurface enclosing a domain  $\Omega$ .**

Let  $B_r$  denote a geodesic ball of radius  $r$  in  $\mathbb{S}_K^{n+1}$ . The outer radius of  $\Omega$  is given by

$$\rho_+(\Omega) = \inf \{R > 0 \mid \Omega \subset B_R(q) \text{ for some } q \in \mathbb{S}_K^{n+1}\},$$

and the inner radius is given by

$$\rho_-(\Omega) = \sup \{\rho > 0 \mid B_\rho(p) \subset \Omega \text{ for some } p \in \mathbb{S}_K^{n+1}\}.$$

In [2], it was shown that pinched hypersurfaces of the Euclidean space satisfy a uniform control of outer by inner radius, and a version for a positive ambient space can be found in [22, Sec. 6]. The proof of this version relied on uniform positivity of the smallest principle curvature, which is insufficient for our purposes. Hence we provide a more general version in the following proposition.

**3.1. Proposition.** *If for some number  $C_0 > 0$  there holds the pinching estimate  $\kappa_n \leq C_0 \kappa_1$  in  $M$ , then the outer radius is estimated from above according to*

$$\rho_+(\Omega) \leq C_1 \rho_-(\Omega),$$

for some positive constant  $C_1 = C_1(n, K, C_0)$ .

*Proof.* For simplicity but without loss of generality, we assume  $K = 1$ . Due to a classical result [15],

$$0 < \rho_+(\Omega) < \frac{\pi}{2},$$

because  $M$  lies in some open hemisphere. Hence there is a centre  $q$  such that

$$\Omega \subset B_{\rho_+}(q),$$

and it is true that  $q \in \bar{\Omega}$  (cf. [40, p. 455]). By moving  $q$  slightly inwards, we can achieve  $M \subset B_{\frac{\pi}{2}}(q)$  and it is starshaped around  $q$ .

Now consider the stereographic projection from the antipodal point  $-q$ , where  $q$  is mapped to the origin  $0 \in \mathbb{R}^{n+1}$ . It follows (see [22, (6.15)]) that the metric  $\bar{g}$  of  $\mathbb{S}^{n+1}$  is conformal to the Euclidean metric; more precisely,

$$\bar{g} = e^{2\psi}(dr^2 + r^2\sigma), \quad \text{with} \quad \psi(r) = -\ln(1 + r^2/4).$$

Hereafter, we denote by tilde the Euclidean geometric quantities. On  $\bar{B}_{\frac{\pi}{2}}(q)$  the metric  $\bar{g}$  is uniformly equivalent to the Euclidean metric.

Next, from [20, (1.1.51)] we get

$$e^\psi \kappa_i = \tilde{\kappa}_i + d\psi(\tilde{\nu})$$

and hence from our pinching assumption,

$$0 < C_0^{-1} \leq \frac{\kappa_1}{\kappa_n} = \frac{\tilde{\kappa}_1 + d\psi(\tilde{\nu})}{\tilde{\kappa}_n + d\psi(\tilde{\nu})} \leq 1.$$

Thus

$$\tilde{\kappa}_1 + d\psi(\tilde{\nu}) \geq C_0^{-1}(\tilde{\kappa}_n + d\psi(\tilde{\nu}))$$

and

$$\tilde{\kappa}_1 \geq C_0^{-1}\tilde{\kappa}_n + (C_0^{-1} - 1)\psi'(r) \langle \partial_r, \tilde{\nu} \rangle \geq C_0^{-1}\tilde{\kappa}_n,$$

because  $\psi' < 0$  and  $\tilde{M}$  is starshaped, i.e.  $\langle \partial_r, \tilde{\nu} \rangle > 0$ . Therefore the Euclidean hypersurface  $\tilde{M} \subset \tilde{B}_2(0)$  is pinched, which from [2, Lemma 5.4] leads to

$$\rho_+(\Omega) \leq C\tilde{\rho}_+(\Omega) \leq C\tilde{\rho}_-(\Omega) \leq C_1\rho_-(\Omega),$$

where we have used the uniform equivalence of the ambient metrics.  $\square$

**3.2. Corollary.** *If for some number  $C_0 > 0$  there holds the pinching estimate  $\kappa_n \leq C_0\kappa_1$  in  $M$ , then one can find positive constants  $d_1$  and  $C_2$ , depending on  $n, K, C_0$  and  $W_\ell(\Omega)$ , such that*

$$d_1 \leq \rho_-(\Omega) \leq C_2 < \frac{\pi}{2\sqrt{K}}.$$

*Proof.* By the definition of inner and outer radius, we can find points  $p, q \in \mathbb{S}_K^{n+1}$  such that

$$B_{\rho_-(\Omega)}(p) \subset \Omega \subset B_{\rho_+(\Omega)}(q).$$

From (2.6), the quermassintegrals  $W_\ell$  are clearly monotone under the inclusion of convex domains, and hence

$$W_\ell(B_{\rho_-(\Omega)}(p)) \leq W_\ell(\Omega) \leq W_\ell(B_{\rho_+(\Omega)}(q)).$$

We obtain with Proposition 3.1,

$$C_2 := f_\ell^{-1}(W_\ell(\Omega)) \geq \rho_-(\Omega) \geq C_1^{-1}\rho_+(\Omega) \geq C_1^{-1}f_\ell^{-1}(W_\ell(\Omega)) =: d_1,$$

where  $f_\ell$  denotes the increasing function given by  $f_\ell(r) = W_\ell(B_r)$ .  $\square$

The trivial outer radius estimate

$$\rho_+(\Omega) < \frac{\pi}{2\sqrt{K}}$$

is not good enough for our purposes. Now we present a refined estimate which should be of independent interest in the future.

**3.3. Theorem.** *The outer radius satisfies*

$$\rho_+(\Omega) \leq \frac{\pi}{2\sqrt{K}} - \frac{\log W_\ell(\mathcal{H}) - \log W_\ell(\Omega)}{(n+1-\ell)\max_M H} =: \frac{\pi}{2\sqrt{K}} - \frac{d_2}{\max_M H},$$

where  $\mathcal{H}$  is an open hemisphere.

*Proof.* From the initial hypersurface  $M_0 = M$ , we start the curvature flow

$$(3.1) \quad \begin{aligned} x: [0, T^*) \times \mathbb{S}^n &\rightarrow \mathbb{S}_K^{n+1} \\ \partial_t x &= \frac{H_{\ell-1}}{H_\ell} \nu. \end{aligned}$$

Then Lemma 2.2 (c) ensures that  $W_\ell$  evolves in time according to

$$\partial_t W_\ell(\Omega_t) = \frac{n+1-\ell}{n+1} V_{n-\ell+1}(\Omega_t) \leq (n+1-\ell)W_\ell(\Omega_t),$$

where the inequality follows from (2.7).

From [22, 32], we know that (3.1) preserves the strict convexity and the solution converges smoothly to an equator, while we also have the estimate

$$W_\ell(\Omega_t) \leq W_\ell(\Omega)e^{(n+1-\ell)t}.$$

Hence the maximal existence time of (3.1) is at least

$$T^* \geq \frac{1}{n+1-\ell} (\log W_\ell(\mathcal{H}) - \log W_\ell(\Omega)),$$

because we know that  $W_\ell(\Omega)$  must converge to  $W_\ell(\mathcal{H})$  at  $T^*$ .

Now [32, Lemma 4.7] leads to the curvature bound

$$\max_{M_t} H \leq \max_{M_0} H \quad \text{for all } 0 \leq t < T^*.$$

Due to the convexity this implies a full second fundamental form bound, as well as a bound

$$(3.2) \quad \frac{H_\ell}{H_{\ell-1}} \leq H_1 \leq \max_{M_0} H,$$

which follows by application of (2.2) for  $k = 1$ .

Next let  $E = \partial\mathcal{H}$  be the limiting equator of the flow and let  $r$  be the radial distance from the centre of  $\mathcal{H}$ , which contains all  $M_t$ . Define

$$\tilde{r}(t) = \max_{\mathbb{S}^n} r(t, \cdot) = r(t, \xi_t),$$

where  $\xi_t$  is any choice of point where the maximum is realised. The function  $\tilde{r}$  is Lipschitz and hence differentiable a.e. At times, where  $\tilde{r}$  is differentiable, we have

$$\frac{d}{dt} \tilde{r}(t) = \frac{H_{\ell-1}}{H_\ell}(t, \xi_t),$$

where we used that  $\nu(t, \xi_t) = \partial_r$ . Integration and (3.2) yield

$$\frac{\pi}{2\sqrt{K}} - \tilde{r}(0) = \tilde{r}(T^*) - \tilde{r}(0) = \int_0^{T^*} \frac{d}{dt} \tilde{r} \geq \frac{\log W_\ell(\mathcal{H}) - \log W_\ell(\Omega)}{(n+1-\ell) \max_{M_0} H}.$$

Hence

$$\max_{M_0} r = \tilde{r}(0) \leq \frac{\pi}{2\sqrt{K}} - \frac{\log W_\ell(\mathcal{H}) - \log W_\ell(\Omega)}{(n+1-\ell) \max_{M_0} H}.$$

Accordingly,  $M_0$  fits into a neighbourhood of the origin of size given by the right hand side of the latter inequality, and therefore the outer radius is controlled by the very same quantity.  $\square$

**3.4. Remark.** A lune, i.e. the intersection of two hemispheres, shows that an outer radius bound in terms of  $W_\ell(\Omega)$  can not be independent of  $\max H$ .

**Theorem 3.3** enables us to find, for a given strictly convex hypersurface of the sphere, a suitable origin, which allows a ball of controlled size within  $\Omega$  and at the same time ensures a controlled positive distance of  $M$  to the equator.

**3.5. Lemma.** *There exists an origin  $\mathcal{O} \in \Omega$ , such that with the constant  $d_2$  from [Theorem 3.3](#) there holds*

$$(3.3) \quad B_{4\epsilon}(\mathcal{O}) \subset \Omega \quad \text{and} \quad \max_M r \leq \frac{\pi}{2\sqrt{K}} - 4\epsilon$$

$$\text{for all } \epsilon \leq \frac{1}{4} \min \left( \frac{d_2}{2 \max_M H}, \frac{\frac{\pi}{2} - \tan^{-1}(\max_M H/\sqrt{K})}{2\sqrt{K}} \right).$$

*Proof.* If  $B_{\rho_+(\Omega)}$  denotes an outball for  $\Omega$  with center  $\mathcal{O}$ , then we know that  $\mathcal{O} \in \bar{\Omega}$ . The distance of focal points from  $M$  can be calculated from the evolution of the Weingarten operator in [Lemma 2.2](#) along the normal variation with speed  $\varphi = -1$ . Then the largest principal curvature is controlled by the solution to the ODE

$$\begin{aligned} y' &= y^2 + K \\ y(0) &= \max_M H, \end{aligned}$$

which exists for all  $t < t_0$ , with

$$t_0 := \frac{\pi}{2\sqrt{K}} - \frac{\tan^{-1}(\max_M H/\sqrt{K})}{\sqrt{K}}.$$

Hence, around all points belonging to the set  $\{x \in \Omega : \text{dist}(x, M) = t_0/2\}$ , there exists an interior ball of radius  $t_0/2$ . In addition, if we shift  $\mathcal{O}$  by a distance of  $d_2/(2 \max_M H)$  in any direction, there still is the same amount of space between  $M$  and the new equator. Therefore, if we shift  $\mathcal{O}$  into  $\Omega$  along a perpendicular geodesic only by the amount

$$\epsilon \leq \epsilon_0 = \frac{1}{4} \min \left( \frac{d_2}{2 \max_M H}, \frac{t_0}{2} \right),$$

then [\(3.3\)](#) holds.  $\square$

#### 4. MONOTONICITY OF THE PINCHING DEFICIT

The geometric results from [section 3](#) depend on the quality of the pinching and the size of the quermassintegral. In the following we investigate how these quantities behave under the flow [\(1.1\)](#). As this flow is defined to be quermassintegral preserving, the key ingredient for proving [Theorem 1.1](#) is the pinching estimate to be proven in this section.

Again, we assume  $K > 0$ , unless stated otherwise. As a prior step, we need the following evolution equations.

**4.1. Lemma.** *For every choice of origin  $\mathcal{O}$ , along [\(1.1\)](#) the induced metric  $g$  and second fundamental form  $h$  satisfy the following evolution equations:*

$$\begin{aligned} \partial_t g_{ij} &= 2(\mu c_K - H)h_{ij} \\ \partial_t h_j^i &= \Delta h_j^i + (|A|^2 - nK)h_j^i + 2KH\delta_j^i - \mu(c_K h_k^i h_j^k + K u h_j^i), \end{aligned}$$

where  $u$  is the generalised support function in [\(1.5\)](#) of  $M_t$  with respect to the origin  $\mathcal{O}$ .

*Proof.* The evolution of the metric comes from [Lemma 2.2](#) (d). For the evolution of  $A$ , we depart from the standard evolution equation

$$\partial_t h_{ij} = \nabla_{ij}^2(H - \mu c_K) + (\mu c_K - H)(h_{ik}h_j^k - Kg_{ij}),$$

see [\[3, Theorem 3-15\]](#) and use the following Simon's type identity:

$$\nabla_{ij}^2 H = \Delta h_{ij} + (|A|^2 - nK)h_{ij} + H(Kg_{ij} - h_i^k h_{jk}).$$

Now we expand the second derivatives of  $c_K$ :

$$(4.1) \quad -\nabla_{ij}^2 c_K = dc_K(\nu)h_{ij} - \bar{\nabla}^2 c_K(x_i, x_j) = c'_K \langle \partial_r, \nu \rangle h_{ij} + Kc_K g_{ij},$$

where we have used

$$\bar{\nabla}^2 c_K = -Kc_K \bar{g}.$$

The proof is complete, using the evolution of the metric to revert to  $h_j^i$ .  $\square$

**4.2. Remark.** Notice that the strong maximum principle for tensors applied to the evolution of  $h_j^i$  already implies that the property of strict convexity is preserved for all times. Accordingly,  $H > 0$  and the quotient  $\frac{\kappa_1}{H}$  is well-defined as long as the flow exists.

Next we deduce an evolution equation that is the key to convergence of the flow.

**4.3. Proposition.** *Let  $\kappa_1$  be the smallest eigenvalue of  $A$ . Then for every choice of origin  $\mathcal{O}$ , under the flow [\(1.1\)](#) with initial data  $M$ , the function*

$$p = \frac{\kappa_1}{H}$$

*is a supersolution to the following evolution equation in viscosity sense, as long as the flow exists:*

$$\begin{aligned} \partial_t p - \Delta p &= \frac{2}{H} \sum_{k=1}^n \sum_{j>D} \frac{(\nabla_k h_j^1)^2}{\kappa_j - \kappa_1} + 2dp(\nabla \log H) + \frac{\mu}{H} c_K |A|^2 p \\ &\quad + \mu c_K \kappa_1 \left(\frac{1}{n} - p\right) + 2nK \left(\frac{1}{n} - p\right), \end{aligned}$$

where  $D$  is the multiplicity of  $\kappa_1$ .

*Proof.* Assume that the flow is defined on a maximal time interval  $[0, T)$ . Let  $(t_0, \xi_0) \in (0, T) \times M$  and let  $\eta$  be a smooth lower support of  $p$  at  $(t_0, \xi_0)$ , i.e.  $\eta$  is defined on a spacetime neighbourhood  $\mathcal{U}$  of  $(t_0, \xi_0)$  and there holds

$$\eta(t_0, \xi_0) = p(t_0, \xi_0), \quad \eta \leq p|_{\mathcal{U}}.$$

Hence  $\varphi = H\eta$  is a smooth lower support for  $\kappa_1$ .

Now we take coordinates with the properties

$$g_{ij} = \delta_{ij}, \quad h_j^i = \kappa_j \delta_j^i \quad \text{at } (t_0, \xi_0).$$

If we denote by  $D$  the multiplicity of  $\kappa_1(t_0, \xi_0)$ , then at the point  $(t_0, \xi_0)$  and for all  $1 \leq i, j \leq D$  it is satisfied (see [\[9, Lemma 5\]](#)) that

$$\partial_t h_j^i = \delta_j^i \partial_t \varphi, \quad \nabla_k h_j^i = \delta_j^i \nabla_k \varphi$$



and

$$(4.2) \quad \nabla_{kk}^2 \varphi \leq \nabla_{kk}^2 h_1^1 - 2 \sum_{j>D} \frac{(\nabla_k h_j^1)^2}{\kappa_j - \kappa_1}.$$

Next, write

$$G_j^i = \partial_t h_j^i - \Delta h_j^i,$$

and compute at  $(t_0, \xi_0)$

$$\begin{aligned} \partial_t \eta &= \frac{\partial_t h_1^1}{H} - \frac{\varphi}{H^2} \partial_t H \\ &= \frac{G_1^1 + g^{kl} \nabla_{kl}^2 h_1^1}{H} - \frac{\varphi}{H^2} \partial_t H \\ &\geq \frac{1}{H} \sum_{k=1}^n \left( \nabla_{kk}^2 \varphi + 2 \sum_{j>D} \frac{(\nabla_k h_j^1)^2}{\kappa_j - \kappa_1} \right) + \frac{G_1^1}{H} - \frac{\varphi}{H^2} \partial_t H \\ &= \frac{2}{H} \sum_{k=1}^n \sum_{j>D} \frac{(\nabla_k h_j^1)^2}{\kappa_j - \kappa_1} + \frac{G_1^1}{H} + \Delta \eta + 2d\eta(\nabla \log H) - \frac{\eta}{H} G_k^k. \end{aligned}$$

On the other hand, we know by [Lemma 4.1](#) that

$$G_j^i = (|A|^2 - nK) h_j^i + 2KH \delta_j^i - \mu(c_K h_k^i h_j^k + K u h_j^i).$$

Accordingly, we get

$$G_1^1 - \eta G_k^k = -\mu c_K \kappa_1^2 + \eta \mu c_K |A|^2 + 2KH(1 - n\eta).$$

Finally, by means of

$$|A|^2 = |\dot{A}|^2 + \frac{1}{n} H^2,$$

we obtain

$$G_1^1 - \eta G_k^k = \eta \mu c_K |\dot{A}|^2 + \mu c_K \kappa_1 H \left( \frac{1}{n} - \eta \right) + 2KH(1 - n\eta),$$

which completes the proof.  $\square$

**4.4. Corollary.** *For every choice of origin  $\mathcal{O}$ , for which  $M \subset \mathcal{H}(\mathcal{O})$ , the flow (1.1) with initial data  $M$  stays in  $\mathcal{H}(\mathcal{O})$  and it improves every pinching, i.e. the pinching deficit*

$$\omega(t) = \frac{1}{n} - \min_{M_t} \frac{\kappa_1}{H}$$

is exponentially decreasing,

$$\omega(t) \leq \omega(s) e^{-2nK(t-s)} \quad \text{for all } 0 \leq s \leq t < T,$$

where  $T$  is the maximal time of existence of the flow with initial data  $M$ .

*Proof.* We need the evolution equation of  $c_K$ . From (1.1) and (4.1) we obtain

$$\partial_t c_K = c'_K \partial_t r = -K s_K dr(\partial_t x) = Ku(H - \mu c_K) = \Delta c_K + Kc_K(n - \mu u).$$

The preservation of  $c_K > 0$  follows immediately by strong maximum principle, as long as the flow exists.

The strict convexity is also preserved from Remark 4.2. The statement about the pinching deficit follows from the strong maximum principle for viscosity solutions, e.g. see [13], and from the fact that

$$\partial_t \omega \leq -2nK\omega,$$

where we can discard the terms including  $\mu$  because  $c_K > 0$  and  $\mu(t) > 0$  by convexity.  $\square$

In particular, for  $s = 0$  we reach a pinching relation between the biggest and smallest principal curvatures of  $M_t$ :

$$(4.3) \quad \kappa_1 \geq \left( \min_{M_0} \frac{\kappa_1}{H} \right) H \geq C_0^{-1} \kappa_n,$$

provided that an origin is chosen such that the strictly convex initial hypersurface is contained in the open hemisphere centred at that origin.

## 5. UPPER CURVATURE BOUNDS

Notice that, unlike in previous treatments of quermassintegral preserving curvature flows, an upper bound for the global term does not come automatically from an upper bound for  $H$ , since the  $c_K$  in the denominator of  $\mu$  is not uniformly bounded away from zero, at least not without further work. Moreover, we need some uniform control of  $\mu$  to get bounds for  $H$ .

To overcome these difficulties, the idea is to choose the origin such that a configuration as in Lemma 3.5 is achieved, which will allow us to deduce uniform bounds on the curvature and the global term in a short but controlled interval  $[0, \tau(\epsilon)]$  (Lemma 5.4). Then, since the pinching is at least as good as at the beginning, we can repeat this process as often as needed, in order to keep the flow going as long as we like (Lemma 6.1). During this evolution, the pinching improves exponentially and at some point will be so strong, that  $M_t$  is very close to a sphere. From here the flow is very easy to estimate and no further shifting of the origin is necessary. Now we implement all required steps to make this argument rigorous.

A key idea to obtain a curvature bound is to adapt a well-known trick from [43], which consists in a suitable combination of the generalised support function with the mean curvature. For this we need control on the size of inballs during the flow.

**A lower bound for the support function of an arbitrary inball.** For a domain  $\Omega$  and a point  $p \in \Omega$ , we say that  $B$  is an *inball at  $p$* , if  $p$  is the center of  $B$  and  $B$  has maximal radius with the property that  $B \subset \Omega$ . In the sequel we are going to prove, that along the flow, the radii of inballs at  $p$  don't decrease too quickly. We give a quantitative estimate.

We need to be careful because now we are dealing with two different support functions: we denote by  $u$  the support function with respect to the origin  $\mathcal{O}$  that is implicit in the flow equation, while for a given point  $p \in \Omega$ ,

$$\hat{u} \equiv \hat{u}_p = s_K(\hat{r}_p) \langle \partial_{\hat{r}_p}, \nu \rangle$$

takes another interior point  $p$  as origin of distances. Accordingly,  $r$  and  $\hat{r}$  mean distance from  $\mathcal{O}$  and  $p$ , respectively. Similar notations will apply to the corresponding trigonometric functions, i.e.

$$\hat{s}_K := s_K(\hat{r}_p) \quad \text{and} \quad \hat{c}_K := c_K(\hat{r}_p).$$

Note that for brevity we suppressed the dependence on the point  $p$  within the notations  $\hat{u}$  and  $\hat{r}$ .

**5.1. Lemma.** *For every choice of origin, along (1.1) the evolution equations of the mean curvature  $H$  and the support function  $\hat{u} = \hat{u}_p$  are given by*

$$\begin{aligned} \partial_t H &= \Delta H + H(|A|^2 + Kn) - \mu(c_K |A|^2 + uKH), \\ \partial_t \hat{u} &= \Delta \hat{u} + \hat{u}|A|^2 + (\mu c_K - 2H)\hat{c}_K + \mu K s_K \hat{s}_K \langle \nabla r, \nabla \hat{r} \rangle. \end{aligned}$$

*Proof.* The formula for  $H$  follows directly by taking the trace in the evolution equation for  $h_j^i$  from Lemma 4.1. On the other hand, a standard calculation leads to

$$\begin{aligned} \partial_t \hat{u} &= \langle \bar{\nabla}_t(\hat{s}_K \partial_{\hat{r}}), \nu \rangle + \langle \hat{s}_K \partial_{\hat{r}}, \bar{\nabla}_t \nu \rangle \\ &= (\mu c_K - H)\hat{c}_K + \hat{s}_K \langle \partial_{\hat{r}}, \nabla(H - \mu c_K) \rangle \\ &= \Delta \hat{u} + (\mu c_K - 2H)\hat{c}_K + \hat{u}|A|^2 - \mu \hat{s}_K \langle \partial_{\hat{r}}, \nabla c_K \rangle, \end{aligned}$$

where we applied well-known formulas for  $\Delta \hat{u}$  and  $\bar{\nabla}_t(\hat{s}_K \partial_{\hat{r}})$  on  $\mathbb{M}_K^{n+1}$  (cf. [10, (4.6) and (4.11)]). The stated formula follows by realizing that

$$\langle \partial_{\hat{r}}, \nabla c_K \rangle = -K s_K \langle \partial_{\hat{r}}, \nabla r \rangle.$$

□

**5.2. Proposition.** *For every choice of origin  $\mathcal{O}$ , for which  $M \subset \mathcal{H}(\mathcal{O})$ , and for every  $p \in \Omega$  and radius  $\rho$  with the property  $B_\rho(p) \subset \Omega$ , the solution  $M_t = \partial\Omega_t$  of (1.1), with initial data  $M$  and maximal existence time  $T > 0$ , satisfies the following:*

(i) *There is a positive constant  $\tilde{\tau} = \tilde{\tau}(n, K, \rho)$  with the property*

$$B_{\rho/4}(p) \subset \Omega_t \quad \text{for all} \quad t \in [0, \min(\tilde{\tau}, T)).$$

(ii) *One can find positive constants  $d_3 \leq \frac{1}{2\sqrt{K}}$  and  $\tau$ , depending on  $n, K, C_0$  and  $W_\ell(\Omega)$ , with the property*

$$\hat{u}_{p_\Omega} - 2d_3 \geq 2d_3 > 0 \quad \text{for all} \quad t \in [0, \min(\tau, T)),$$

*where  $p_\Omega$  is the center of an inball corresponding to the inradius  $\rho_-(\Omega)$ .*

*Proof.* (i) Let us first obtain the evolution of the distance  $\hat{r} = \hat{r}_p$  from the fixed point  $p$  to the points on  $M_t$  under the flow (1.1):

$$(5.1) \quad \partial_t \hat{r} = d\hat{r}(\partial_t x) = (\mu c_K - H) \langle \nu, \partial_{\hat{r}} \rangle.$$

On the other hand,  $\mathbf{r}(t)$  denotes the radius of a geodesic sphere centred at  $p$  that moves under the ordinary mean curvature flow starting at  $\mathbf{r}(0) = \rho/2$ , that is,

$$\mathbf{r}'(t) = -n \operatorname{co}_K(\mathbf{r}(t)),$$

whose solution is given by

$$c_K(\mathbf{r}(t)) = e^{Knt} c_K(\rho/2) \quad \text{for } t \geq 0.$$

As  $c_K$  is a decreasing function,

$$\mathbf{r}(t) \geq \rho/4 \quad \text{if and only if} \quad e^{Knt} c_K(\rho/2) \leq c_K(\rho/4),$$

meaning that

$$\mathbf{r}(t) \geq \rho/4 \quad \text{if and only if} \quad t \leq \frac{1}{Kn} \log \frac{c_K(\rho/4)}{c_K(\rho/2)} =: \tilde{\tau}.$$

Set  $f(t, \cdot) = \hat{r}(t, \cdot) - \mathbf{r}(t)$  for  $t \in [0, \min(\tau, T))$ . Then  $f(0, \cdot) > 0$  and  $f$  evolves according to

$$\partial_t f = (\mu c_K - H) \langle \nu, \partial_{\hat{r}} \rangle + n \operatorname{co}_K(\mathbf{r}(t)).$$

If there exists a first time  $t_1$  such that the geodesic sphere  $B_{\mathbf{r}(t_1)}$  touches the hypersurface  $M_{t_1}$  at some point  $x_1$ , then at this first minimum for  $f$  it holds  $H(x_1, t_1) \leq n \operatorname{co}_K(\mathbf{r}(t_1))$ ,  $\langle \partial_{\hat{r}}, \nu \rangle = 1$  and  $\partial_t f(x_1, t_1) \leq 0$ . Consequently, taking into account that  $c_K > 0$  and strict convexity is preserved, we have

$$\partial_t f(x_1, t_1) \geq \mu(t_1) c_K(\mathbf{r}(t_1)) > 0,$$

which is a contradiction, and hence the statement follows.

(ii) Apply (i) with  $p = p_\Omega$  and  $\rho = \rho_-(\Omega)$  and obtain the desired  $\tau$  due to [Corollary 3.2](#). Using [Lemma 2.1](#) we get

$$(5.2) \quad \hat{u}_{p_\Omega} \geq s_K(\rho_-(\Omega)/4) \geq s_K(d_1/4) =: 4d_3,$$

where we applied the lower bound in [Corollary 3.2](#). □

**An upper bound for the mean curvature.** To estimate the mean curvature along a solution with suitably located initial data, we use the well known auxiliary function

$$\Phi_p = \frac{H}{\hat{u}_p - d_3},$$

which, after choosing the origin  $\mathcal{O}$  as in [Proposition 5.2](#), is well-defined for a while for some suitable  $p \in \Omega$ . Routine computations lead to the evolution of  $\Phi$  where we suppress the dependence on  $p$  within the notation.

**5.3. Lemma.** *Under the assumptions of Proposition 5.2, along the flow (1.1) the function  $\Phi$  evolves according to*

$$\begin{aligned} \partial_t \Phi &= \Delta \Phi + \frac{2}{\hat{u} - d_3} \langle \nabla \Phi, \nabla \hat{u} \rangle + \Phi \left( nK - \frac{d_3}{\hat{u} - d_3} |A|^2 \right) + 2\Phi^2 \hat{c}_K \\ &\quad - \mu K \Phi u - \frac{\mu}{\hat{u} - d_3} \left( c_K |A|^2 + \Phi (K s_K \hat{s}_K \langle \nabla r, \nabla \hat{r} \rangle + c_K \hat{c}_K) \right). \end{aligned}$$

Unlike in previous literature, we cannot neglect all the terms including  $\mu$ , as some of them do not have a sign. Nor is it known at this point that  $\mu$  is bounded. The novelty about our approach is to make use of Lemma 3.5, a configuration which enables us to gain some control on  $\mu$  and then get an estimate on  $\Phi$  and  $H$ .

Another complication arises from the necessity of using an iterative change of origin. The configuration of Lemma 3.5 depends on curvature. Hence we need a very precise estimate of curvature as the flow progresses and it is insufficient to estimate the curvature by a multiple of its initial value, as then our time interval, along which  $B_\epsilon(\mathcal{O}) \subset \Omega_t$  is valid, would decrease and we would not be able to prove long-time existence.

For this reason, we introduce a novel method, which also gives an interesting alternative to proving initial value independent curvature bounds after a waiting time as for example in [34, Equation (17)]. It provides a bound on  $\Phi$ , which is uniform in  $p$  lying within a certain region.

With this purpose, we define a modified auxiliary function

$$\begin{aligned} \Psi: [0, T) \times \mathbb{S}^n \times \mathbb{S}_K^{n+1} &\rightarrow \mathbb{R} \\ (t, \xi, p) &\mapsto \begin{cases} (\text{dist}(p, M_t) - 2d_3)\Phi_p, & p \in V_t \\ 0, & \text{else,} \end{cases} \end{aligned}$$

with

$$V_t := \{p \in \Omega_t: \text{dist}(p, M_t) > 2d_3\},$$

and where for  $p \notin \bar{\Omega}$  the distance to  $M_t$  is defined to be negative. In short,

$$\Psi(t, \xi, p) = \max(0, \min_{M_t} \hat{r}_p - 2d_3)\Phi_p.$$

Note that  $\Psi$  is Lipschitz, because when  $\min_{M_t} \hat{r}_p = \text{dist}(p, M_t) \geq 2d_3$ , then by Lemma 2.1

$$\hat{u}_p \geq s_K(\min_{M_t} \hat{r}_p) \geq s_K(2d_3),$$

due to  $\min_{M_t} \hat{r}_p < \pi/(2\sqrt{K})$ . Furthermore,

$$\hat{u}_p \geq s_K(2d_3) = \frac{1}{\sqrt{K}} \sin(2d_3\sqrt{K}) \geq \frac{6}{5}d_3,$$

where we used  $\sin x \geq \frac{3}{5}x$  for  $x \in [0, \pi/2]$ . Hence  $\Phi_p$  is well-defined for  $p \in \bar{V}_t$ . In the sequel we write, for brevity,

$$\max_{M_t} \Psi = \max_{\mathbb{S}^n \times \mathbb{S}_K^{n+1}} \Psi(t, \cdot).$$

**5.4. Lemma.** *There exists an origin  $\mathcal{O} \in \Omega$ , a constant  $C_3(n, K, C_0, W_\ell(\Omega))$ , and constants  $\tau$  and  $\epsilon_1$ , depending on  $n, K, C_0, W_\ell(\Omega), \max(C_3, \max_{M_0} \Psi)$ , such that for the solution  $M_t = \partial\Omega_t$  of (1.1) with initial data  $M_0 = M$  and maximal existence time  $T > 0$  there hold:*

- (i)  $M_t \subset \mathcal{H}(\mathcal{O})$  for all  $t \in [0, \min(\tau, T)]$ .
- (ii)  $\hat{u}_{p_\Omega} \geq 4d_3$  and  $B_{\epsilon_1}(\mathcal{O}) \subset \Omega_t$  for all  $t \in [0, \min(\tau, T)]$ .
- (iii) The function  $\Psi$  satisfies

$$\max(C_3, \max_{M_t} \Psi) \leq \max(C_3, \max_M \Psi) \quad \text{for all } t \in [0, \min(\tau, T)].$$

*Proof.* Define the functions

$$\epsilon_0(y) = \frac{1}{4} \min \left( \frac{d_2 d_3 \sqrt{K}}{2y}, \frac{\frac{\pi}{2} - \tan^{-1}(\frac{y}{d_3 K})}{2\sqrt{K}} \right), \quad \text{for } y > 0,$$

and

$$q(y) = \frac{n\pi C_0^{\ell+1}}{\sqrt{K}} \left( 1 + \frac{5\pi}{\sqrt{K}d_3} \right) \frac{y}{\epsilon_0(y)} + \frac{n\pi^2}{4}y + \frac{\pi}{\sqrt{K}}y^2 - \frac{d_3^2}{n}y^3,$$

where  $d_3$  and  $C_0$  are the constants from [Proposition 5.2](#) and [\(4.3\)](#) respectively. It is clear that  $q(y)$  converges to  $-\infty$  as  $y \rightarrow \infty$  and hence it has a largest zero  $\bar{y}$ , which only depends on  $n, K, C_0$  and  $W_\ell(\Omega)$ . Let  $\psi_0 = \max_{t=0} \Psi$  and define

$$\epsilon_1 = \epsilon_0(\max(\bar{y}, \psi_0)).$$

(i) & (ii): With  $p_\Omega$  from [Proposition 5.2](#) (ii) and by definition of  $d_3$  in [\(5.2\)](#), along  $M_0 = M$  there holds

$$H = (\hat{u}_{p_\Omega} - d_3)\Phi_{p_\Omega} \leq \frac{\Phi_{p_\Omega}}{\sqrt{K}} \leq \frac{s_K(\min_M \hat{r}_{p_\Omega}) - 2d_3}{2d_3} \frac{\Phi_{p_\Omega}}{\sqrt{K}} \leq \frac{1}{d_3\sqrt{K}}\psi_0$$

and hence

$$\epsilon_1 \leq \epsilon_0(d_3\sqrt{K} \max_M H) < \frac{\pi}{2\sqrt{K}},$$

where the latter estimate is due to the definition of  $\epsilon_0$ . This exactly the threshold required to apply [Lemma 3.5](#).

Then we can apply [Lemma 3.5](#) with  $\epsilon = \epsilon_1$  in order to obtain a suitable origin  $\mathcal{O} \in \Omega$  with the property [\(3.3\)](#). From the first part of [Proposition 5.2](#) applied to  $p = \mathcal{O}$  and  $\rho = 4\epsilon_1$ , as well as from the second part of [Proposition 5.2](#), we obtain  $\tau = \tau(n, K, C_0, W_\ell(\Omega), \epsilon_1)$ , up to which the claimed properties of (ii) are satisfied. Property (i) is then clear from the fact that at the equator  $c_K = 0$  and hence  $\frac{d}{dt} \max r < 0$ .

(iii) Next, we bound the function  $\Psi$ . Suppose  $\Psi$  attains a positive maximum over the set  $[0, \bar{t}] \times \mathbb{S}^n \times \mathbb{S}_K^{n+1}$  at some  $(\bar{t}, \bar{\xi}, \bar{p})$ . Define the Lipschitz function

$$\bar{\Psi}_{\bar{p}}(t) = \max_{\mathbb{S}^n} \Psi(t, \cdot, \bar{p}),$$

which is positive in some small interval  $J := [\bar{t} - \delta, \bar{t}]$ . Thus we have

$$\text{dist}(\bar{p}, M_t) > 2d_3 \quad \forall t \in J.$$

Hence in  $J$  the function  $\Phi_{\bar{p}}$  is smooth and

$$\bar{\Psi}_{\bar{p}}(t) = \left( \min_{M_t} \hat{r}_{\bar{p}} - 2d_3 \right) \max_{M_t} \Phi_{\bar{p}}$$

is differentiable almost everywhere in  $J$ .

There holds for almost every  $t \in J$ , using (5.1) and  $\langle \partial_{\hat{r}_{\bar{p}}}, \nu \rangle > 0$  due to  $\bar{p} \in \Omega_t$  for all  $t \in J$ ,

$$\begin{aligned} \frac{d}{dt} \bar{\Psi}_{\bar{p}} &= \frac{d}{dt} \min_{M_t} \hat{r}_{\bar{p}} \max_{M_t} \Phi_{\bar{p}} + \left( \min_{M_t} \hat{r}_{\bar{p}} - 2d_3 \right) \frac{d}{dt} \max_{M_t} \Phi_{\bar{p}} \\ &= (c_K(r)\mu - H(\bar{t}, x(\bar{t}, \text{argmin}_{M_{\bar{t}}} \hat{r}_{\bar{p}}))) \langle \partial_{\hat{r}_{\bar{p}}}, \nu \rangle \max_{M_t} \Phi_{\bar{p}} \\ &\quad + \left( \min_{M_{\bar{t}}} \hat{r}_{\bar{p}} - 2d_3 \right) \frac{d}{dt} \max_{M_t} \Phi_{\bar{p}} \\ &\leq \mu \max_{M_t} \Phi_{\bar{p}} + \left( \min_{M_{\bar{t}}} \hat{r}_{\bar{p}} - 2d_3 \right) \frac{d}{dt} \max_{M_t} \Phi_{\bar{p}}. \end{aligned}$$

Inserting the evolution equation from Lemma 5.3, discarding two good terms, and using  $\nabla \Phi_{\bar{p}}(t, \xi_t) = 0$  at all maximisers  $\xi_t$ , we obtain

$$\begin{aligned} \frac{d}{dt} \bar{\Psi}_{\bar{p}} &\leq \mu \max_{M_t} \Phi_{\bar{p}} + \left( \min_{M_{\bar{t}}} \hat{r}_{\bar{p}} - 2d_3 \right) \left( \max_{M_t} \Phi_{\bar{p}} \left( nK - \frac{d_3}{\hat{u}_{\bar{p}} - d_3} |A|^2 \right) + 2 \max_{M_t} \Phi_{\bar{p}}^2 \right. \\ &\quad \left. - \frac{\mu}{\hat{u}_{\bar{p}} - d_3} \left( \max_{M_t} \Phi_{\bar{p}} (K s_K \hat{s}_K \langle \nabla r, \nabla \hat{r}_{\bar{p}} \rangle + c_K \hat{c}_K) \right) \right), \end{aligned}$$

where we used that  $u$  and  $c_K$  are positive due to  $M_t \subset \mathcal{H}(\mathcal{O})$  and  $\mathcal{O} \in \Omega_t$ . Using  $s_K \leq K^{-1/2}$  and

$$\frac{\min_{M_{\bar{t}}} \hat{r}_{\bar{p}} - 2d_3}{\hat{u}_{\bar{p}} - d_3} (K s_K \hat{s}_K \langle \nabla r, \nabla \hat{r}_{\bar{p}} \rangle + c_K \hat{c}_K) \leq 2 \cdot \frac{\pi / (2\sqrt{K})}{\frac{1}{5}d_3} = \frac{5\pi}{d_3\sqrt{K}},$$

we get

$$\frac{d}{dt} \bar{\Psi}_{\bar{p}} \leq \left( 1 + \frac{5\pi}{\sqrt{K}d_3} \right) \mu \max_{M_t} \Phi_{\bar{p}} + nK \bar{\Psi}_{\bar{p}} + 2\bar{\Psi}_{\bar{p}} \max_{M_t} \Phi_{\bar{p}} - \frac{d_3^2}{n} \bar{\Psi}_{\bar{p}} \max_{M_t} \Phi_{\bar{p}}^2.$$

By means of the Hsiung-Minkowski identity (2.3) and Lemma 2.1, we estimate for all  $0 \leq t \leq \tau$ ,

$$(5.3) \quad \mu(t) \leq n \binom{n}{\ell+1} \frac{\int_{M_t} \kappa_n^{\ell+1}}{\int_{M_t} u \sigma_{\ell+1}} \leq \frac{n}{s_K(\epsilon_1)} \frac{\int_{M_t} \kappa_n^{\ell+1}}{\int_{M_t} \kappa_1^{\ell+1}} \leq \frac{2nC_0^{\ell+1}}{\epsilon_1},$$

where the constant  $C_0$  comes from the pinching (4.3). Hence

$$\begin{aligned} \frac{d}{dt} \bar{\Psi}_{\bar{p}} &\leq 2nC_0^{\ell+1} \left( 1 + \frac{5\pi}{\sqrt{K}d_3} \right) \frac{\max_{M_t} \Phi_{\bar{p}}}{\epsilon_0(\max(\bar{y}, \psi_0))} + nK\bar{\Psi}_{\bar{p}} \\ &\quad + 2\bar{\Psi}_{\bar{p}} \max_{M_t} \Phi_{\bar{p}} - \frac{d_3^2}{n} \bar{\Psi}_{\bar{p}} \max_{M_t} \Phi_{\bar{p}}^2 \\ &= \frac{2nC_0^{\ell+1}}{\min_{M_{\bar{t}}} \hat{r}_{\bar{p}} - 2d_3} \left( 1 + \frac{5\pi}{\sqrt{K}d_3} \right) \frac{\bar{\Psi}_{\bar{p}}}{\epsilon_0(\max(\bar{y}, \psi_0))} + nK\bar{\Psi}_{\bar{p}} \\ &\quad + \frac{2}{\min_{M_{\bar{t}}} \hat{r}_{\bar{p}} - 2d_3} \bar{\Psi}_{\bar{p}}^2 - \frac{d_3^2}{n(\min_{M_{\bar{t}}} \hat{r}_{\bar{p}} - 2d_3)^2} \bar{\Psi}_{\bar{p}}^3. \end{aligned}$$

Multiplication with  $(\min_{M_{\bar{t}}} \hat{r}_{\bar{p}} - 2d_3)^2$  gives for almost every  $t \in J$ ,

$$\begin{aligned} (\min_{M_{\bar{t}}} \hat{r}_{\bar{p}} - 2d_3)^2 \frac{d}{dt} \bar{\Psi}_{\bar{p}} &\leq \frac{n\pi C_0^{\ell+1}}{\sqrt{K}} \left( 1 + \frac{5\pi}{\sqrt{K}d_3} \right) \frac{\bar{\Psi}_{\bar{p}}}{\epsilon_0(\max(\bar{y}, \psi_0))} \\ &\quad + \frac{n\pi^2}{4} \bar{\Psi}_{\bar{p}} + \frac{\pi}{\sqrt{K}} \bar{\Psi}_{\bar{p}}^2 - \frac{d_3^2}{n} \bar{\Psi}_{\bar{p}}^3. \end{aligned}$$

which is strictly negative whenever  $\max(\bar{y}, \psi_0) < \bar{\Psi}_{\bar{p}}(t)$ . To conclude the argument, suppose that

$$\Psi(\bar{t}, \bar{\xi}, \bar{p}) = \max_{[0, \bar{t}] \times \mathbb{S}^n \times \mathbb{S}_K^{n+1}} \Psi > \max(\bar{y}, \psi_0).$$

We know that

$$\bar{\Psi}_{\bar{p}}(t) \leq \Psi(\bar{t}, \bar{\xi}, \bar{p})$$

with equality at  $t = \bar{t}$ . But before we have shown that, for  $t$  close to  $\bar{t}$ , we have  $\frac{d}{dt} \bar{\Psi}_{\bar{p}} < 0$  almost everywhere, which is impossible. Hence we obtain the desired estimate with  $C_3 = \bar{y}$ .  $\square$

**Higher order curvature bounds.** We use the estimate from Lemma 5.4 to control the global term and estimate the derivatives of curvature.

**5.5. Lemma.** *For the origin  $\mathcal{O} \in \Omega$  from Lemma 5.4 and the solution  $M_t = \partial\Omega_t$  of (1.1) with initial data  $M$  and maximal existence time  $T > 0$  and for all  $m \in \mathbb{N}$ , there exists  $C_4 = C_4(n, m, K, C_0, W_\ell(\Omega), \max(C_3, \max_M \Psi))$  with the property*

$$\mu(t) + |\nabla^m A| \leq C_4 \quad \forall t \in [0, \min(\tau, T)].$$

where  $\tau$  is the number from Lemma 5.4. In particular there holds  $T \geq \tau$  and the flow exists smoothly on  $[0, \tau]$ .

*Proof.* Up to the time  $\min(\tau, T)$ , (5.3) holds, i.e. with the notation from the proof of Lemma 5.4 we have

$$\mu(t) \leq \frac{2nC_0^{\ell+1}}{\epsilon_1} = \frac{2nC_0^{\ell+1}}{\epsilon_0(\max(C_3, \psi_0))}.$$



The curvature derivative bound can be proved by a well-known induction argument, as for example in [25]. First, due to convexity and Lemma 5.4 (ii) and (iii),

$$|A|^2 \leq H^2 \leq \frac{\Phi_{p\Omega}^2}{K} \leq \frac{1}{d_3^2 K} (\max(C_3, \max_M \Psi))^2.$$

Assuming that all derivatives up to order  $m-1$  are bounded by a constant of the form  $C_4$ , we obtain the evolution equation of  $|\nabla^m A|$ ,

$$\begin{aligned} \partial_t |\nabla^m A|^2 &\leq \Delta |\nabla^m A|^2 - 2|\nabla^{m+1} A|^2 + C(\mu u + K)|\nabla^m A|^2 \\ &\quad + C \sum_{i+j+k=m} \left( \mu |\nabla^i c_K| + |\nabla^i A| \right) |\nabla^j A| |\nabla^k A| |\nabla^m A|, \end{aligned}$$

where we used that  $\nabla u = A * \nabla c_K$  (see (2.4)). Here  $S * T$  denotes any linear combination of tensors formed by contracting  $S$  and  $T$  by means of  $g$ .

Then we claim

$$(5.4) \quad |\nabla^m A|^2 \leq C \quad \forall t \in [0, \min(\tau, T)].$$

As  $\mu$  and  $u$  are bounded, we get

$$(\partial_t - \Delta) |\nabla^m A|^2 \leq -2|\nabla^{m+1} A|^2 + C|\nabla^m A|^2 + C \left( \sum_{i=0}^m |\nabla^i c_K| + 1 \right) |\nabla^m A|.$$

Notice that  $c_K(r)$  and  $|\nabla c_K|$  are bounded as well. Moreover, from (4.1), one has for  $\ell \geq 0$  the following covariant derivatives

$$\nabla^{\ell+2} c_K = \nabla^\ell c_K * K + u * \nabla^\ell A + \sum_{i+j+k=\ell} \nabla^i c_K * \nabla^j A * \nabla^k A,$$

which are controlled by uniform constants arguing by induction. In short, we reach

$$(\partial_t - \Delta) |\nabla^m A|^2 \leq -2|\nabla^{m+1} A|^2 + C|\nabla^m A| (1 + |\nabla^m A|),$$

which leads to (5.4) by standard maximum principle arguments, as for example in the proof of [26, Theorem 4.1].

As the right hand side of the flow equation and all higher derivatives of the curvature remain uniformly bounded, we conclude (as in [25, pages 257 ff.]) that, if  $T < \tau$ , then  $M_t$  converges (as  $t \rightarrow T$ , in the  $C^\infty$ -topology) to a unique, smooth and strictly convex hypersurface<sup>1</sup>. Now we can apply short time existence to continue the solution after  $T$ , contradicting the maximality of  $T$ . Hence the solution of (1.1) starting at a strictly convex hypersurface exists on  $[0, \tau)$ . On this interval we have uniform smooth estimates, and hence the flow also exists on  $[0, \tau]$ .  $\square$

<sup>1</sup>Note that due to the bound on  $\mu$ , it can be seen from Lemma 5.1 that  $H$  is bounded from below on every finite time interval.

## 6. CONSTRUCTION OF A GLOBAL SOLUTION

In the previous section we achieved existence and uniform estimates of any solution to (1.1) with strictly convex initial data  $M$  on a time interval  $[0, \tau]$  from Lemma 5.4, the length of which only depends on preserved data of the problem. Those are in particular the hemisphere  $\mathcal{H}(\mathcal{O})$ , the pinching constant  $C_0$ , the quermassintegral  $W_\ell(\Omega)$  and the number  $\max(C_3, \max_M \Psi)$ . The full curvature derivative bounds also only depend on those quantities.

Hence we can start an iteration process and shift, at time  $i\tau$  with  $i \in \mathbb{N}$ , the origin according to Lemma 5.4 applied to the new strictly convex initial hypersurface  $M_{i\tau}$ . The constant  $C_4$  from Lemma 5.5 is then uniform among the integers  $i$ , because it only depends on quantities which are always preserved. The following lemma makes this precise.

**6.1. Lemma.** *Let  $M_0 \subset \mathbb{S}_K^{n+1}$  be a strictly convex hypersurface enclosing a domain  $\Omega$ . Then there exists a sequence of origins  $(\mathcal{O}_i)_{i \in \mathbb{N} \cup \{0\}}$  and positive numbers  $\tau_0, \epsilon_1(0)$  depending only on  $n, K, C_0, W_\ell(\Omega_0), \max(C_3, \max_{M_0} \Psi)$ , such that the problem*

$$\begin{aligned} \partial_t x &= (\mu_i(t)c_K(r_i) - H)\nu, \quad \forall t \in [i\tau_0, (i+1)\tau_0) \\ x(0, \mathbb{S}^n) &= M_0 \\ x((i+1)\tau_0, \mathbb{S}^n) &= \lim_{t \nearrow (i+1)\tau_0} M_t, \end{aligned}$$

where  $r_i$  is the distance to  $\mathcal{O}_i$  and  $\mu_i$  is given as in (1.3) to keep the quermassintegral  $W_\ell(\Omega_t)$  for any  $\ell = 0, 1, \dots, n$  fixed, has a solution

$$x: [0, \infty) \times \mathbb{S}^n \rightarrow \mathbb{S}_K^{n+1}.$$

For every  $t \geq 0$ ,  $M_t$  is strictly convex and there holds

$$(6.1) \quad B_{\epsilon_1(0)}(\mathcal{O}_i) \subset \Omega_t \quad \text{and} \quad M_t \subset \mathcal{H}(\mathcal{O}_i) \quad \forall t \in [i\tau_0, (i+1)\tau_0).$$

The mappings  $x(t, \cdot)$  satisfy spatial  $C^\infty$ -estimates which are uniform in time.

*Proof.* For  $M_0$ , pick  $\mathcal{O}_0$  according to Lemma 5.4. From Lemma 5.5 we conclude that the solution  $M_t$  of (1.1) with initial data  $M_0$  exists on  $[0, \tau_0]$ , where  $\tau_0$  and  $\epsilon_1(0)$  depend on  $n, K, C_0, W_\ell(\Omega_0), \max(C_3, \max_{M_0} \Psi)$ . The derivatives of  $A$  are bounded by

$$C_4 = C_4 \left( n, m, K, C_0, W_\ell(\Omega_0), \max(C_3, \max_{M_0} \Psi) \right).$$

Now suppose that for  $i \geq 0$  the hypersurface  $M_{i\tau_0}$ , the origin  $\mathcal{O}_i$  and the solution  $(M_t)_{t \in [i\tau_0, (i+1)\tau_0)}$  were already constructed, such that (6.1),

$$\max(C_3, \max_{M_t} \Psi) \leq \max(C_3, \max_{M_0} \Psi),$$

as well as

$$\mu_i(t) + |\nabla^m A| \leq C_4 \left( n, m, K, C_0, W_\ell(\Omega_0), \max(C_3, \max_{M_0} \Psi) \right)$$

all hold for all  $t \in [i\tau_0, (i+1)\tau_0]$ . Then apply [Lemma 5.4](#) to the initial hypersurface  $M_{(i+1)\tau_0}$  and obtain an origin  $\mathcal{O}_{i+1}$ , such that the solution  $M_t$  of [\(1.1\)](#) with initial data  $M_{(i+1)\tau_0}$  satisfies [\(6.1\)](#),

$$\max(C_3, \max_{M_t} \Psi) \leq \max(C_3, \max_{M_{(i+1)\tau_0}} \Psi) \leq \max(C_3, \max_{M_0} \Psi)$$

and

$$\mu_{i+1}(t) + |\nabla^m A| \leq C_4 \left( n, m, K, C_0, W_\ell(\Omega_0), \max(C_3, \max_{M_0} \Psi) \right)$$

during the interval  $[(i+1)\tau_0, (i+2)\tau_0]$  and with the same  $\epsilon_1(0)$ . Here we also used that  $C_0$  and  $W_\ell$  are preserved. This means that the construction can be carried out infinitely often to obtain the desired long-time solution.  $\square$

## 7. ASYMPTOTIC ESTIMATES AND CONVERGENCE TO A SPHERICAL CAP

In the previous sections we have put ourselves into a position where we have a strictly convex flow  $(M_t)_{0 \leq t < \infty}$  in the sphere. This flow is not necessarily smooth in time, but it satisfies spatial  $C^k$ -estimates which are uniform with respect to time and it has a uniformly bounded global term, due to the proof of [Lemma 6.1](#).

Additionally, by means of [Corollary 4.4](#) and the curvature bounds, we get

$$\sum_{i=1}^n (\kappa_i - \kappa_1) = H - n\kappa_1 \leq nH\omega(t) \leq nCe^{-2nKt},$$

which implies exponential decay of the traceless second fundamental form:

$$(7.1) \quad |\mathring{A}|^2 = \frac{1}{n} \sum_{i < j} (\kappa_j - \kappa_i)^2 \leq Ce^{-4nKt}.$$

Using this property, in the following we are going to apply some recent estimates of almost-umbilical type due to De Rosa and Gioffré [\[14\]](#), to show that the process of picking new origins actually terminates after finitely many steps and that the flow will then converge to a geodesic sphere of a uniquely determined radius. The crucial ingredient is the following result.

**7.1. Theorem.** [\[14, Thm. 1.3\]](#) *Let  $n \geq 2$ , let  $\Sigma$  be a closed hypersurface in  $\mathbb{R}^{n+1}$  and let  $p > n$  be given. We assume that there exists  $c_0 > 0$  such that  $\Sigma$  satisfies the conditions*

$$|\Sigma| = |\mathbb{S}^n|, \quad \|A\|_{L^p(\Sigma)} \leq c_0.$$

*There exist positive numbers  $\delta, C > 0$ , depending only on  $n, p, c_0$ , with the following property: if*

$$\|\mathring{A}\|_{L^p(\Sigma)} \leq \delta,$$

then there exists a vector  $c = c(\Sigma)$ , such that  $\Sigma - c$  is a graph over the sphere, namely there exists a parametrization

$$\psi: \mathbb{S}^n \rightarrow \Sigma, \quad \psi(x) = e^{f(x)}x + c,$$

and  $f$  satisfies the estimate

$$\|f\|_{W^{2,p}(\mathbb{S}^n)} \leq C\|\mathring{A}\|_{L^p(\Sigma)}.$$

In the following we will use this result to prove that the surfaces become exponentially  $C^2$ -close to geodesic spheres and that the necessity to pick new origins vanishes.

**7.2. Lemma.** *In the situation of Lemma 6.1, there exists an integer  $m > 0$  depending on  $n, K$  and  $M$ , such that in Lemma 6.1 the origins  $\mathcal{O}_i$ ,  $i > m$ , may be chosen constantly equal to  $\mathcal{O}_m$ .*

*Proof.* Let  $m$  be a positive integer to be specified during the proof. Let  $\mathcal{O}_m$  be the flow origin associated to the interval  $I_m := [m\tau_0, (m+1)\tau_0)$ . By stereographic projection from the antipodal point of  $\mathcal{O}_m$ , the family  $(M_t)_{t \in I_m}$  can be viewed as a flow in the Euclidean space, then denoted by  $(\tilde{M}_t)_{t \in I_m}$ . Geometric quantities of this flow, denoted by a tilde, are related to the original ones as follows, see [20, Prop. 1.1.11], where  $e^{2\varphi}$  is the conformal factor:

$$g = e^{2\varphi}\tilde{g}, \quad \nu = e^{-\varphi}\tilde{\nu}$$

$$e^\varphi A = \tilde{A} + d\varphi(\tilde{\nu}) \text{ id}, \quad e^\varphi H = \tilde{H} + n d\varphi(\tilde{\nu}).$$

In particular, we obtain

$$\mathring{\tilde{A}} = e^\varphi \mathring{A}.$$

The surface areas of  $M_t$  and  $\tilde{M}_t$  are related by

$$|M_t| = \int_{M_t} 1 = \int_{\tilde{M}_t} e^{n\varphi}$$

and hence

$$C^{-1}|M_t| \leq |\tilde{M}_t| \leq |M_t|,$$

where  $C$  depends on  $|\varphi|_{C^0(M_t)}$ . Now define the scaled hypersurface

$$\hat{M}_t = \lambda \tilde{M}_t, \quad \text{with} \quad \lambda^n = \frac{|\mathbb{S}^n|}{|\tilde{M}_t|},$$

so that  $|\hat{M}_t| = |\mathbb{S}^n|$ . Now the associated Weingarten operator is

$$\hat{A} = \lambda^{-1} \tilde{A} = \frac{e^\varphi}{\lambda} A - \lambda^{-1} d\varphi(\tilde{\nu}) \text{ id}$$

and similarly for the traceless Weingarten operator. Hence

$$\|\hat{A}\|_{L^\infty(\hat{M}_t)} \leq C(\|A\|_{L^\infty(M_t)} + 1)$$

and

$$\|\mathring{A}\|_{L^\infty(\hat{M}_t)} \leq C \|\mathring{A}\|_{L^\infty(M_t)},$$

where  $C$  depends on  $n$ ,  $|\varphi|_{C^1(M_t)}$  and  $|M_t|$ .

Then  $\|\mathring{A}\|_{L^\infty(\hat{M}_t)}$  is bounded by [Lemma 5.5](#) and [\(7.1\)](#) ensures that  $\|\mathring{A}\|_{L^\infty}$  is as small as needed for  $m$  big enough. Therefore, we can apply [Theorem 7.1](#) for sufficiently large  $m$ , to get a function  $\hat{f}$  which, from the embedding theorems of Sobolev spaces into Hölder spaces, satisfies

$$\|\hat{f}\|_{C^1(\mathbb{S}^n)} \leq C \|\mathring{A}\|_{L^\infty(\hat{M}_t)} \leq C \|\mathring{A}\|_{L^\infty(M_t)} \leq C e^{-2nKt} \quad \text{for all } t \in I_m.$$

Then, due to our curvature bounds [\(5.4\)](#), we have full  $C^k$ -bounds on  $\hat{f}$  for all  $k$ . By iteration of interpolation arguments for  $C^k$  bounds (see [\[21, Corollary 6.2\]](#)), this implies that

$$\|\hat{f}\|_{C^k(\mathbb{S}^n)} \leq C e^{-2nKt} \quad \text{for all } t \in I_m.$$

In other words,  $\hat{M}_t$  is exponentially  $C^k$ -close to a sphere  $\hat{\mathcal{S}}_t$  for all  $k \in \mathbb{N}$  and for all  $t \in I_m$ . As the area along the  $M_t$  is uniformly bounded above and below by [Corollary 3.2](#), we get  $C^k$  bounds for the conformal factor as well, and this property of closeness to a sphere  $\mathcal{S}_t$  carries over to the original flow in  $\mathbb{S}_K^{n+1}$ . Note that the radii of the spheres  $\mathcal{S}_t$  converge to a well defined limit, which is strictly less than  $\pi/(2\sqrt{K})$ , determined by the initial value of  $W_\ell(\Omega_0)$ . Hence the curvature of  $M_t$  is uniformly bounded from below.

On the other hand, the radial distance to the origin  $\mathcal{O}_m$  satisfies

$$(7.2) \quad \partial_t r = (\mu c_K - H)v^{-1}, \quad \text{with} \quad v^2 = 1 + s_K^{-2}|dr|_\sigma^2.$$

Hence, for an error  $\delta_m$  that converges to zero when  $m \rightarrow \infty$ ,

$$(7.3) \quad \partial_t r = \left( \frac{\int_{M_t} \sigma_\ell H}{\int_{M_t} \sigma_\ell c_K} c_K - H \right) v^{-1} \leq \left( \frac{c_K}{f_{\mathcal{S}_t} c_K} - 1 \right) \frac{H_{\mathcal{S}_t}}{v} + \delta_m,$$

where  $H_{\mathcal{S}_t}$  is the mean curvature of the sphere  $\mathcal{S}_t$ . At points which maximize  $r$ ,  $c_K$  is minimized. At such points the first term on the RHS of [\(7.3\)](#) is strictly negative if  $c_K$  is not constant. Hence, for large  $m$ , if  $\mathcal{S}_t$  is uniformly off-center,  $\max_{M_t} r$  is decreasing and a similar estimate shows that  $\min_{M_t} r$  is increasing. Hence, from then on, there is no need to adjust the origin anymore.  $\square$

**Convergence to a spherical cap.** To complete the proof of [Theorem 1.1](#), it only remains to show that the immortal solution coming from [Lemma 6.1](#) actually converges to a limit geodesic sphere. After  $m$ -fold picking of a new origin, we now may without loss of generality assume that origins have not been changed at all. We will exploit the  $C^\infty$ -estimates for the flow hypersurfaces  $M_t$  coming from [\(5.4\)](#). We already know from [Lemma 7.2](#), that every limit point of the flow must be round sphere.

Now we prove that only the sphere centred at the origin can arise as a limit. Notice that the radius of  $R$  of any limit sphere is determined by the initial hypersurfaces by means of the equality  $W_\ell(B_R) = W_\ell(\Omega_0)$ . Denote by  $H_R$  the mean curvature of such a sphere  $S_R$ . Hence, from the evolution (7.2) of the radial distance and for an error  $\delta$  that converges to zero when  $M_{t_k} \rightarrow S_R$ , we get

$$(7.4) \quad \partial_t r = \left( \frac{c_K}{\int_M c_K} - 1 \right) \frac{H_R}{v} + \delta.$$

As above, at points which maximize  $r$ ,  $c_K$  is minimized, thus at such points the first term on the RHS of (7.4) is strictly negative if  $c_K$  is not constant. Therefore the function  $\max_{M_t} r$  is strictly decreasing in sufficiently small  $C^2$ -neighbourhoods of any non-centred sphere, which excludes those as limits. Thus subsequential limits are unique and the whole flow must converge.

## 8. THE ELLIPTIC CASE: RIGIDITY RESULTS

In order to prove [Theorem 1.3](#), we first have to get the elliptic viscosity equation of the pinching deficit for general curvature function  $F$ . Let us first gather some prerequisites about these functions.

**Symmetric curvature functions.** If  $M$  is a hypersurface of  $\mathbb{M}_K^{n+1}$ , then we set

$$F(x) = f(\kappa_1(x), \dots, \kappa_n(x)),$$

which can be alternatively seen as a function defined on the diagonalisable endomorphisms,  $F = F(A)$ , or as a function of a symmetric and a positive definite bilinear form,  $F = F(g, h)$ . In the latter case, we write

$$F^{ij} = \frac{\partial F}{\partial h_{ij}}, \quad F^{ij,kl} = \frac{\partial^2 F}{\partial h_{ij} \partial h_{kl}}.$$

With these conventions, the covariant derivatives are given by

$$(8.1) \quad \nabla_i F = F^{jk} \nabla_i h_{jk}.$$

On strictly convex hypersurfaces  $M$ , we can define the so-called *inverse curvature function* by

$$\tilde{F}(A) = \frac{1}{F(A^{-1})}.$$

A curvature function  $F$  is called *inverse concave*, if  $\tilde{F}$  is concave. Notice that concavity/convexity with respect to the matrix variables is equivalent to the same property with respect to the eigenvalues (see [5, 20] for deeper information). Next we gather several useful properties:

**8.1. Lemma.** (a) *If  $F$  is inverse concave and  $M$  is strictly convex, then*

$$(8.2) \quad F^{ij,kl} \nabla_1 h_{ij} \nabla_1 h_{kl} + 2 \sum_j \frac{F^{ii}}{\kappa_j} (\nabla_1 h_{ij})^2 \geq \frac{2}{F} (\nabla_1 F)^2.$$

(b) Under *Conditions 1.2*, it holds:

- (i) If  $F$  is convex, then it is inverse concave (cf. [20, Lemma 2.2.12, Lemma 2.2.14])
- (ii) Euler's formula  $F^{ii}\kappa_i = F$ , implies that  $F$  is strictly positive.

**Rigidity for radial curvature functions.** We start with a result that contains an elliptic version of *Proposition 4.3*.

**8.2. Lemma.** *Suppose  $F$  satisfies *Conditions 1.2*, then:*

(i) *The Weingarten operator satisfies the following elliptic equation:*

$$\begin{aligned} -F^{rs}\nabla_{rs}^2 h_{ij} &= F^{pq,rs}\nabla_i h_{pq}\nabla_j h_{rs} - \nabla_{ij}^2 F \\ &\quad + F^{rs}h_{ms}h_r^m h_{ij} + KFg_{ij} - Fh_{mj}h_i^m - KF^{rs}g_{rs}h_{ij}. \end{aligned}$$

(ii) *For the function  $p = \frac{\kappa_1}{F}$ , it holds*

$$\begin{aligned} -FF^{kl}\nabla_{kl}^2 p &\geq 2\sum_{j>D} \frac{F^{kk}(\nabla_k h_j^1)^2}{\kappa_j - \kappa_1} + F^{pq,rs}\nabla_1 h_{pq}\nabla_1 h_{rs} + 2F^{kl}\nabla_k p\nabla_l F \\ &\quad + pF^{kl}\nabla_{kl} F - \nabla_{11} F + F^{ii}\text{Sec}_{i1}(\kappa_i - \kappa_1) \end{aligned}$$

*in viscosity sense, where  $D$  is the multiplicity of  $\kappa_1$ .*

*Proof.* (i) We differentiate (8.1), to get

$$\nabla_{ij}^2 F = F^{pq,rs}\nabla_i h_{pq}\nabla_j h_{rs} + F^{rs}\nabla_{ij}^2 h_{rs}.$$

Now combine the commutator formula for second covariant derivatives with Gauss equation (2.1) to deduce:

$$\begin{aligned} F^{rs}\nabla_{ij}^2 h_{rs} &= F^{rs}(\nabla_{rs}^2 h_{ij} + R_{isj}^m h_{mr} + R_{rsj}^m h_{mi}) \\ &= F^{rs}\nabla_{rs}^2 h_{ij} + F^{rs}(h_{ms}h_{ij} - h_{mj}h_{is} + Kg_{ms}g_{ij} - Kg_{mj}g_{is})h_r^m \\ &\quad + F^{rs}(h_{ms}h_{rj} - h_{mj}h_{rs} + Kg_{ms}g_{rj} - Kg_{mj}g_{rs})h_i^m \\ &= F^{rs}(\nabla_{rs}^2 h_{ij} + h_{ms}h_r^m h_{ij}) + KFg_{ij} - Fh_{mj}h_i^m - KF^{rs}g_{rs}h_{ij}. \end{aligned}$$

(ii) As in the proof of *Proposition 4.3*, let  $\eta$  be a smooth lower support of  $p$  at  $\xi_0 \in M$  and define  $\varphi = \eta F$ . Then, in coordinates with

$$(8.3) \quad g_{ij} = \delta_{ij}, \quad h_{ij} = \kappa_i \delta_{ij}$$

we have inequality (4.2), which combined with (i) leads to

$$\begin{aligned} -FF^{kl}\nabla_{kl}^2 \eta &= -F^{kl}\nabla_{kl}^2 \varphi + \frac{\varphi}{F}F^{kl}\nabla_{kl}^2 F + 2F^{kl}\nabla_k \eta \nabla_l F \\ &\geq 2F^{kl}\nabla_k \eta \nabla_l F + 2\sum_{j>D} \frac{F^{kk}(\nabla_k h_j^1)^2}{\kappa_j - \kappa_1} + F^{pq,rs}\nabla_1 h_{pq}\nabla_1 h_{rs} \\ &\quad + \eta F^{kl}\nabla_{kl}^2 F - \nabla_{11}^2 F + F^{ii}(\kappa_i \kappa_1 + K)(\kappa_i - \kappa_1). \end{aligned}$$

□

In order to prove [Theorem 1.3](#), let  $F = \gamma c_K^\alpha$ , then taking derivatives and using [\(4.1\)](#) we get

$$\begin{aligned}\nabla_{kl}^2 F &= \alpha \gamma c_K^{\alpha-1} \nabla_{kl}^2 c_K + \alpha(\alpha-1) \gamma c_K^{\alpha-2} \nabla_k c_K \nabla_l c_K \\ &= K \left( \frac{u}{c_K} h_{kl} - g_{kl} \right) \alpha F + \frac{2\epsilon}{F} \nabla_k F \nabla_l F,\end{aligned}$$

where

$$(8.4) \quad \epsilon = \frac{\alpha-1}{2\alpha} = \frac{1}{2} - \frac{1}{2\alpha}.$$

This implies

$$\begin{aligned}(8.5) \quad pF^{kl} \nabla_{kl}^2 F - \nabla_{11}^2 F &= p\alpha KF \left( \frac{u}{c_K} F - F^{kl} g_{kl} \right) - \alpha KF \left( \frac{u}{c_K} \kappa_1 - 1 \right) \\ &\quad + \frac{2\epsilon p}{F} F^{kl} \nabla_k F \nabla_l F - \frac{2\epsilon}{F} (\nabla_1 F)^2 \\ &= K\alpha F^{ii} (\kappa_i - \kappa_1) + \frac{2\epsilon}{F} (pF^{kl} \nabla_k F \nabla_l F - (\nabla_1 F)^2).\end{aligned}$$

Here we have used Euler's relation and computations are done in the coordinates [\(8.3\)](#).

*Proof of [Theorem 1.3](#).* As  $|\alpha| \geq 1$ , we have that  $\epsilon \in [0, 1]$  for  $\epsilon$  defined as in [\(8.4\)](#). Then using that  $M$  is convex, we can estimate

$$2 \sum_{j>D} \frac{F^{kk} (\nabla_k h_j^1)^2}{\kappa_j - \kappa_1} \geq 2\epsilon \sum_{j=1}^n \frac{F^{kk} (\nabla_k h_j^1)^2}{\kappa_j} - 2\epsilon \frac{p}{F} F^{kl} \nabla_k F \nabla_l F + T * \nabla p,$$

where we also used the following (see [\[9, Lemma 5\]](#)):

$$\nabla_k \varphi \delta_{ij} = \nabla_k h_{ij} \quad \text{for all } 1 \leq i, j \leq D.$$

Plugging this in [Lemma 8.2](#) (ii) and using the convexity of  $F$ , we get

$$\begin{aligned}-FF^{kl} \nabla_{kl}^2 p &\geq 2\epsilon \sum_{j=1}^n \frac{F^{kk} (\nabla_k h_j^1)^2}{\kappa_j} - 2\epsilon \frac{p}{F} F^{kl} \nabla_k F \nabla_l F + \epsilon F^{pq,rs} \nabla_1 h_{pq} \nabla_1 h_{rs} \\ &\quad + pF^{kl} \nabla_{kl} F - \nabla_{11} F + F^{ii} \text{Sec}_{i1} (\kappa_i - \kappa_1) + T * \nabla p \\ (8.5) \quad &\geq 2\epsilon \sum_{j=1}^n \frac{F^{kk} (\nabla_k h_j^1)^2}{\kappa_j} + \epsilon F^{pq,rs} \nabla_1 h_{pq} \nabla_1 h_{rs} - \frac{2\epsilon}{F} (\nabla_1 F)^2 \\ &\quad + F^{ii} (\text{Sec}_{i1} + \alpha K) (\kappa_i - \kappa_1) + T * \nabla p.\end{aligned}$$

Notice that in the case  $\alpha = -1$ , then  $\epsilon = 1$  and we do not need the convexity of  $F$  in the first inequality above.

By [Lemma 8.1](#) (b) our  $F$  is inverse concave, and hence [\(8.2\)](#) leads to

$$-FF^{kl} \nabla_{kl}^2 p + T * \nabla p \geq F^{ii} (\text{Sec}_{i1} + \alpha K) (\kappa_i - \kappa_1).$$



This completes the proof using the strong maximum principle for viscosity solutions and  $M$  has to be centred at the origin provided  $K \neq 0$ , since only in this case can  $c_K$  be constant.  $\square$

8.3. *Remark.* Note that this approach also provides a direct maximum principle proof of Liebmann’s soap bubble theorem (the convex case of Alexandrov’s theorem), [30].

**Solitons.** We complete this paper by proving [Corollary 1.4](#).

*Proof of [Corollary 1.4](#).* For the given hypersurface  $M$ , there is a dual hypersurface  $\tilde{M} \subset \mathbb{M}_K^{n+1}$ , where  $K = \text{sgn}(\mathbb{M})$  as in (1.6), and with the properties

$$\tilde{\kappa}_i = \frac{1}{\kappa_i}, \quad \tilde{c}_K = u,$$

see [20, Thm. 10.4.4, Thm. 10.4.9] and [39]. Hence  $\tilde{M}$  satisfies the equation

$$\tilde{c}_K^{\frac{1}{\beta}} = F(\kappa_i) = F(\tilde{\kappa}_i^{-1}) = \frac{1}{\tilde{F}(\tilde{\kappa}_i)},$$

i.e. with  $\alpha = -1/\beta$  we have

$$\tilde{F}|_{\tilde{M}} = \tilde{c}_K^\alpha,$$

where  $\tilde{F}$  is the inverse curvature function of  $F$ . Therefore to complete the proof it only remains to check the conditions of [Theorem 1.3](#) for  $\tilde{M}$ , which holds provided that for any  $\tilde{g}$ -orthonormal frame we have

$$\widetilde{\text{Sec}}_{ij} \geq -\alpha K.$$

In coordinates that diagonalize  $\tilde{A}$ , the Gauss equation (2.1) for  $\tilde{M}$  gives

$$\tilde{R}_{ijij} + \alpha K = \tilde{h}_{ii}\tilde{h}_{jj} - \tilde{h}_{ij}\tilde{h}_{ij} + (1 + \alpha)K = \frac{1}{\kappa_i\kappa_j} + (1 + \alpha)K \geq 0,$$

provided

$$\frac{1}{\kappa_i\kappa_j} \geq \frac{1 - \beta}{\beta} K.$$

Notice that if  $\frac{1 - \beta}{\beta} K \leq 0$  this is guaranteed by convexity of  $M$ ; otherwise, the inequality follows by the assumption on  $\text{Sec}_M$ . Hence the statement follows by direct application of [Theorem 1.3](#).  $\square$

## REFERENCES

- [1] Alexandr Alexandroff, *A characteristic property of spheres*, Ann. Mat. Pura Appl. **58** (1962), no. 4, 303–315.
- [2] Ben Andrews, *Contraction of convex hypersurfaces in Euclidean space*, Calc. Var. Partial Differ. Equ. **2** (1994), no. 2, 151–171.
- [3] Ben Andrews, *Contraction of convex hypersurfaces in Riemannian spaces*, J. Differential Geom. **39** (1994), no. 2, 407–431.
- [4] Ben Andrews, *Motion of hypersurfaces by Gauss curvature*, Pac. J. Math. **195** (2000), no. 1, 1–34.

- [5] Ben Andrews, *Pinching estimates and motion of hypersurfaces by curvature functions*, J. Reine Angew. Math. **608** (2007), 17–33.
- [6] Ben Andrews, Xuzhong Chen, and Yong Wei, *Volume preserving flow and Alexandrov-Fenchel type inequalities in hyperbolic space*, J. Eur. Math. Soc. **23** (2021), no. 7, 2467–2509.
- [7] Ben Andrews, Pengfei Guan, and Lei Ni, *Flow by powers of the Gauss curvature*, Adv. Math. **299** (2016), 174–201.
- [8] Ben Andrews and Yong Wei, *Quermassintegral preserving curvature flow in hyperbolic space*, Geom. Funct. Anal. **28** (2018), no. 5, 1183–1208.
- [9] Simon Brendle, Kyeongsu Choi, and Panagiota Daskalopoulos, *Asymptotic behavior of flows by powers of the Gaussian curvature*, Acta Math. **219** (2017), no. 1, 1–16.
- [10] Esther Cabezas-Rivas and Vicente Miquel, *Volume preserving mean curvature flow in the hyperbolic space*, Indiana Univ. Math. J. **56** (2007), no. 5, 2061–2086.
- [11] Esther Cabezas-Rivas and Vicente Miquel, *Non-preserved curvature conditions under constrained mean curvature flows*, Differential Geom. Appl. **49** (2016), 287–300.
- [12] Chuanqiang Chen, Pengfei Guan, Junfang Li, and Julian Scheuer, *A fully-nonlinear flow and quermassintegral inequalities in the sphere*, Pure Appl. Math. Quarterly **18** (2022), no. 2, 437–461.
- [13] Francesca Da Lio, *Remarks on the strong maximum principle for viscosity solutions to fully nonlinear parabolic equations*, Commun. Pure Appl. Math. **3** (2004), no. 3, 395–415.
- [14] Antonio De Rosa and Stefano Gioffré, *Absence of bubbling phenomena for non-convex anisotropic nearly umbilical and quasi-Einstein hypersurfaces*, J. Reine Angew. Math. **780** (2021), 1–40.
- [15] Manfredo Do Carmo and Frank Warner, *Rigidity and convexity of hypersurfaces in spheres*, J. Differ. Geom. **4** (1970), no. 2, 133–144.
- [16] Jose Espinar, Jose Galvez, and Pablo Mira, *Hypersurfaces in  $\mathbb{H}^{n+1}$  and conformally invariant equations; the generalized Christoffel and Nirenberg problems*, J. Eur. Math. Soc. **11** (2009), no. 4, 903–939.
- [17] Shanze Gao, Haizhong Li, and Hui Ma, *Uniqueness of closed self-similar solutions to  $\sigma_k^\alpha$ -curvature flow*, Nonlinear Differ. Equ. Appl. **25** (2018), art. 45.
- [18] Shanze Gao, Haizhong Li, and Xianfeng Wang, *Self-similar solutions to fully nonlinear curvature flows by high powers of curvature*, J. Reine Angew. Math. **783** (2022), 135–157.
- [19] Shanze Gao and Hui Ma, *Self-similar solutions of curvature flows in warped products*, Differ. Geom. Appl. **62** (2019), 234–252.
- [20] Claus Gerhardt, *Curvature problems*, Series in Geometry and Topology, vol. 39, International Press of Boston Inc., Somerville, 2006.
- [21] Claus Gerhardt, *Inverse curvature flows in Hyperbolic space*, J. Differ. Geom. **89** (2011), 487–527.
- [22] Claus Gerhardt, *Curvature flows in the sphere*, J. Differ. Geom. **100** (2015), no. 2, 301–347.
- [23] Frederico Girao and Neilha Pinheiro, *An Alexandrov-Fenchel-type inequality for hypersurfaces in the sphere*, Ann. Glob. Anal. Geom. **52** (2017), no. 4, 413–424.
- [24] Pengfei Guan and Junfang Li, *A mean curvature type flow in space forms*, Int. Math. Res. Not. **2015** (2015), no. 13, 4716–4740.
- [25] Gerhard Huisken, *Flow by mean curvature of convex surfaces into spheres*, J. Differ. Geom. **20** (1984), no. 1, 237–266.
- [26] Gerhard Huisken, *The volume preserving mean curvature flow*, J. Reine Angew. Math. **382** (1987), 35–48.
- [27] Gerhard Huisken, *Asymptotic behavior for singularities of the mean curvature flow*, J. Differ. Geom. **31** (1990), no. 1, 285–299.

- [28] Gerhard Huisken and Alexander Polden, *Geometric evolution equations for hypersurfaces*, Calculus of variations and geometric evolution problems (Stefan Hildebrandt and Michael Struwe, eds.), Lecture notes in mathematics, vol. 1713, Springer, Cetraro, 1996, pp. 45–84.
- [29] Kwok-Kun Kwong, Hoojoo Lee, and Juncheol Pyo, *Weighted Hsiung-Minkowski formulas and rigidity of umbilical hypersurfaces*, Math. Res. Lett. **25** (2018), no. 2, 297–616.
- [30] Heinrich Liebmann, *Über die Verbiegung der geschlossenen Flächen positiver Krümmung*, Math. Ann. **53** (1900), no. 1–2, 81–112.
- [31] Matthias Makowski, *Mixed volume preserving curvature flows in hyperbolic space*, unpublished, 2012.
- [32] Matthias Makowski and Julian Scheuer, *Rigidity results, inverse curvature flows and Alexandrov-Fenchel type inequalities in the sphere*, Asian J. Math. **20** (2016), no. 5, 869–892.
- [33] Uwe Mayer and Gieri Simonett, *Self-intersections for the surface diffusion and the volume-preserving mean curvature flow*, Differ. Integr. Equ. **13** (2000), no. 7–9, 1189–1199.
- [34] James McCoy, *The mixed volume preserving mean curvature flow*, Math. Z. **246** (2004), no. 1, 155–166.
- [35] James McCoy, *Self-similar solutions of fully nonlinear curvature flows*, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) **X** (2011), no. 2, 317–333.
- [36] Robert Reilly, *Variational properties of functions of the mean curvatures for hypersurfaces in space forms*, J. Differ. Geom. **8** (1973), no. 3, 465–477.
- [37] Ángela Roldán, *Hypersurfaces with prescribed curvatures in de Sitter space*, Rev. Mat. Iberoam. **38** (2022), no. 1, 269–294.
- [38] Antonio Ros, *Compact hypersurfaces with constant higher order mean curvatures*, Rev. Mat. Iberoam. **3** (1987), no. 3–4, 447–453.
- [39] Julian Scheuer, *The Minkowski inequality in de Sitter space*, Pac. J. Math. **314** (2021), no. 2, 425–449.
- [40] Luis A. Santaló, *Convex regions on the  $n$ -dimensional spherical surface*. Ann. of Math. (2) **47** (1946), 448–459.
- [41] Luis A. Santaló, *Integral geometry and geometric probability*. Second edition. Cambridge Mathematical Library. Cambridge University Press, Cambridge, 2004.
- [42] Gil Solanes, *Integral geometry and the Gauss-Bonnet theorem in constant curvature spaces*, Trans. Am. Math. Soc. **358** (2006), no. 3, 1105–1115.
- [43] Kaising Tso, *Deforming a hypersurface by its Gauss-Kronecker curvature*, Commun. Pure Appl. Math. **38** (1985), no. 6, 867–882.
- [44] Guofang Wang and Chao Xia, *Isoperimetric type problems and Alexandrov-Fenchel type inequalities in the hyperbolic space*, Adv. Math. **259** (2014), 532–556.
- [45] Jie Wu, *A new characterization of geodesic spheres in the hyperbolic space*, Proc. Am. Math. Soc. **144** (2016), no. 7, 3077–3084.
- [46] Jie Wu and Chao Xia, *On rigidity of hypersurfaces with constant curvature functions in warped product manifolds*, Ann. Glob. Anal. Geom. **46** (2014), no. 1, 1–22.

UNIVERSITAT DE VALÈNCIA, DEPARTAMENT DE MATEMÀTIQUES, DR MOLINER. 50,  
46100 BURJASSOT, VALÈNCIA (SPAIN)

*Email address:* Esther.Cabezas-Rivas@uv.es

GOETHE-UNIVERSITÄT, INSTITUT FÜR MATHEMATIK, ROBERT-MAYER-STR. 10, 60325  
FRANKFURT, GERMANY

*Email address:* scheuer@math.uni-frankfurt.de