

A family of diameter perfect constant-weight codes from Steiner systems*

Minjia Shi[†], Yuhong Xia[‡], Denis S. Krotov[§]

Abstract

If S is a transitive metric space, then $|C| \cdot |A| \leq |S|$ for any distance- d code C and a set A , “anticode”, of diameter less than d . For every Steiner $S(t, w, n)$ system S , we show the existence of a q -ary constant-weight code C of length n , weight w (or $n - w$), and distance $d = 2w - t + 1$ (respectively, $d = n - t + 1$) and an anticode A of diameter $d - 1$ such that the pair (C, A) attains the code–anticode bound and the supports of the codewords of C are the blocks of S (respectively, the complements of the blocks of S). We study the problem of estimating the minimum value of q for which such a code exists, and find that minimum for small values of t .

Keywords: diameter perfect codes, anticodes, constant-weight codes, code–anticode bound, Steiner systems.

1. Introduction

Diameter perfect codes are natural generalizations of perfect codes, fascinating structures in coding theory. They can be defined for different metrics,

*This research is supported in part by National Natural Science Foundation of China (12071001); the work of D. S. Krotov is supported with the state contract of the Sobolev Institute of Mathematics (Project FWNF-2022-0017).

[†]M. J. Shi is with the Key Laboratory of Intelligent Computing Signal Processing, Ministry of Education, School of Mathematical Sciences, Anhui University, Hefei 230601, China.

[‡]Y. H. Xia is with the School of Mathematical Sciences, Anhui University, Hefei 230601, China.

[§]D. S. Krotov is with the Sobolev Institute of Mathematics, Novosibirsk 630090, Russia.

especially, for the Hamming metric. In this paper, we discuss diameter perfect codes in the subspaces of the Hamming metric space restricted by the words whose weight is constant.

The concept of a diameter perfect codes [1] is based on the the code–anticode bound, which is a generalization of the sphere-packing bound; the most famous referenced result with that bound is Delsarte’s [7, Theorem 3.9]. However, the main importance of [7, Theorem 3.9] is that the code–anticode bound is proved there for any distance-regular graph (more generally for an association scheme), without any assumption on its symmetries. For symmetric metric spaces, such as Hamming and Johnson schemes, the bound has a much simpler proof, and probably was known earlier (in particular, the key lemma in the proof of the Elias–Bassalygo bound [4] is based on the same generalized pigeonhole principle).

Lemma 1 (code–anticode bound). *Let \mathcal{V} be a finite metric space whose automorphism group $\text{Iso}(\mathcal{V})$ acts transitively on its points. Let \mathcal{C} be a subset of \mathcal{V} (a code) with minimum distance d between elements, and let \mathcal{A} be a subset of \mathcal{V} of diameter D (an anticode), where $D < d$. Then*

$$|\mathcal{C}| \cdot |\mathcal{A}| \leq |\mathcal{V}|. \tag{1}$$

Proof. Denote $\overline{\mathcal{A}} = \{\pi(\mathcal{A}) : \pi \in \text{Iso}(\mathcal{V})\}$. Because of the transitivity, each point of \mathcal{V} belongs to exactly s sets in $\overline{\mathcal{A}}$, where $s = |\overline{\mathcal{A}}| \cdot |\mathcal{A}| / |\mathcal{V}|$. From the hypothesis on the diameter of \mathcal{A} and minimum distance of \mathcal{C} , we see that each set from $\overline{\mathcal{A}}$ contains at most one element of \mathcal{C} . On the other hand, each element of \mathcal{C} is contained in s elements of $\overline{\mathcal{A}}$. Therefore, $|\overline{\mathcal{A}}| \geq s|\mathcal{C}|$. Substituting $s = |\overline{\mathcal{A}}| \cdot |\mathcal{A}| / |\mathcal{V}|$, we get (1). \square

If (1) holds with equality, then \mathcal{C} is called a *diameter perfect distance- d code* or *D -diameter perfect code*. It is easy to see that in this case \mathcal{A} is a maximum diameter- D anticode. The space $\mathcal{J}_q(n, w)$ of weight- w words of length n over the q -ary alphabet, as was shown in [10], belongs to the class of spaces for which the code–anticode bound holds. In [10], Etzion classified known families of non-binary diameter perfect constant-weight codes as follows (below, q is the alphabet size, n is the length, w is the weight and d is the minimum distance):

- F1 Non-binary diameter-perfect constant-weight codes for which $w = n$. Essentially, they are diameter-perfect codes in $H(q - 1, n)$ (perfect codes, extended perfect codes, MDS codes).

- F2 Non-binary diameter perfect constant-weight codes for which $w = n - 1$. Known constructions [11], [13], [14], [15], [18], [20] are related to perfect codes in $H(q - 1, n - 1)$, $q = 2^m + 1$, and their extensions.
- F3 Generalized Steiner systems [9], for which every codeword of weight t , $t < w$, is at distance $w - t$ from exactly one codeword and $d = 2(w - t) + 1$.
- F4,F6 $d = w$ or $d < w$, respectively. Codes of size $\binom{n}{w}(q - 1)^{w-d+1}$, where all codewords of one of $\binom{n}{w}$ possible supports form an unrestricted distance- d MDS code of length w over the $(q - 1)$ -ary alphabet $\{1, \dots, q - 1\}$.
- F5 Codes of size $\binom{n}{w}$ with $d = w + 1$, where the supports of the codewords form the complete design (all w -subsets of $\{1, \dots, n\}$).

Actually, family F5 also falls in the definition of F4 and F6 with $d = w + 1$. However, we keep it separate in the list because our goal is to extend it. In this paper, we further explore diameter perfect constant-weight codes over non-binary alphabets and maximum anticodes corresponding to those codes. We consider two families of codes, F5' and F5'', generalizing F5 (F5 corresponds to F5' with $t = w$ and to F5'' with $n - t = w$):

- F5' Codes of size $\binom{n}{t}/\binom{w}{t}$ with $d = 2w - t + 1$, where the supports of the codewords form a Steiner system $S(t, w, n)$, $w \leq n/2$.
- F5'' Codes of size $\binom{n}{t}/\binom{n-w}{t}$ with $d = n - t + 1$, where the complements of the supports of the codewords form a Steiner system $S(t, n - w, n)$, $w \geq n/2$.

It is notable that F5' and F5'' are related to the two known families of binary diameter perfect constant-weight codes, constructed from Steiner systems [10, Sect. III]. While in the binary case, the codes from the two families are connected by a trivial operation (swapping the role of 0 and 1), the codes from F5' and F5'' are essentially different, except for the special case $w = n/2$.

The structure of the paper is as follows. In Section 2, we give main definitions and basic results. In Section 3, we construct a class of anticodes corresponding to families F5', F5'' and show that they are maximum for large q (Theorems 1 and 1), also showing the same for a more general class

of anticodes (Proposition 1). In Section 4, we observe that the existence of a Steiner system (with the corresponding parameters) implies the existence, for sufficiently large q , of diameter perfect codes that attain the code–anticode bound with the new anticodes (Theorem 3), and define q'_0 and q''_0 as the smallest possible q (corresponding to families F5' and F5'', respectively) for which such codes exist.

In Section 4.1, we find q'_0 in the case of Steiner systems of strength 2 (Theorems 4) and 3 (Theorems 5, Corollaries 2 and 3). Further, in Sections 4.2 and 4.3, we derive an upper bound on q'_0 in the general case (Theorems 6) and consider special cases related to the Steiner systems $S(4, 5, 11)$ (Proposition 2), $S(3, 4, 14)$ and $S(3, 4, 20)$ (Proposition 3). In Section 4.4, we consider q''_0 ; we derive a lower bound (Proposition 4, Corollary 5) and solve the special cases related to affine and projective planes (Corollaries 6 and 7). Section 5 contains concluding remarks and conjectures.

2. Preliminaries

In this section, we present definitions and notations, as well as several results to facilitate understanding throughout the paper.

By \mathbb{Z}_q , we denote the set $\{0, \dots, q-1\}$ (we do not need to endow it by any algebraic operation, such as addition or multiplication); we will also use the notation $[i, j]$ for $\{i, i+1, \dots, j\}$, $i \leq j$, and $[n]$ for $[1, n]$. Given a positive integer n , let \mathbb{Z}_q^n be the set of all words of length n over the alphabet \mathbb{Z}_q ; for each word $\bar{x} \in \mathbb{Z}_q^n$, we have $\bar{x} = (x_i)_{i \in [n]}$ and $x_i \in \mathbb{Z}_q$ for any $i \in [n]$. The *support* of a word $\bar{x} \in \mathbb{Z}_q^n$ is the set $\text{supp}(\bar{x}) = \{i \in [n] : x_i \neq 0\}$. The *weight* of a word $\bar{x} \in \mathbb{Z}_q^n$, denoted by $\text{wt}(\bar{x})$, is equal to $|\text{supp}(\bar{x})|$. The *distance* (Hamming distance) $d(\bar{x}, \bar{y})$ between two words \bar{x}, \bar{y} in \mathbb{Z}_q^n is defined to be the number of positions in which \bar{x} and \bar{y} differ. By $\mathcal{J}_q(n, w)$, we denote the subset of \mathbb{Z}_q^n that is restricted by words of weight w , and the corresponding metric subspace of the Hamming space.

Definition 1 (codes). Given a metric space, a *code* as a subset with at least two vertices, referred to as *codewords*. A code C is said to have *distance* d if the distance between any two distinct codewords is not less than d . A code C is a *q -ary code of length n* if $C \subseteq \mathbb{Z}_q^n$. If, additionally, $C \subseteq \mathcal{J}_q(n, w)$, then it is a *constant-weight*, or *constant-weight- w* , code. For brevity, we call a q -ary constant-weight- w code of distance d a $(n, d, w; |C|)_q$ code, or a $(n, d, w; \cdot)_q$ code if the cardinality is not specified or important.

Definition 2 (anticoes and diameter perfect codes). An *anticode* of diameter D in a metric space \mathcal{V} is a nonempty subset \mathcal{A} of \mathcal{V} such that the distance between any \bar{x} and \bar{y} in \mathcal{A} is not greater than D . A distance- d code \mathcal{C} in \mathcal{V} is called *diameter perfect* if it attains the code–anticode bound (1) for some anticode \mathcal{A} of diameter smaller than d .

Definition 3 (code matrix). For a code \mathcal{C} in \mathbb{Z}_q^n , we define the *code matrix* whose rows are the codewords of \mathcal{C} . To be explicit, we can require that the rows are ordered lexicographically; however, we will never use this order in our considerations.

Definition 4 (Steiner systems). A (Steiner) system $S(t, k, n)$, $0 < t \leq k < n$, is a pair $S = (N, B)$, where N is an n -set (whose elements are often called *points*) and B is a set of k -subsets (called *blocks*) of N , where each t -subset of N is contained in exactly one block of B . Systems $S(2, 3, n)$ are also denoted $\text{STS}(n)$ and called *Steiner triple systems*, *STS*. Systems $S(3, 4, n)$ are also denoted $\text{SQS}(n)$ and called *Steiner quadruple systems*, *SQS*.

Definition 5 (derived systems). For a Steiner $S(t, k, n)$ system $S = (N, B)$ and a set of points $\alpha \subset N$, $0 < |\alpha| < t$, the system $(N \setminus \alpha, \{\beta \setminus \alpha : \alpha \subset \beta \in B\})$ is a Steiner $S(t - |\alpha|, k - |\alpha|, n - |\alpha|)$ system, called *derived* from S .

Definition 6 (resolvable systems). A Steiner $S(t, k, n)$ system $S = (N, B)$ is called *l -resolvable*, $1 \leq l < t$, (if $l = 1$, just *resolvable*) if the set of blocks B can be partitioned into subsets each of which forms an $S(l, k, n)$ on the same point set N . If $l = 1$, those subsets are referred to as *parallel classes*.

Definition 7 (minimum-distance graph and chromatic number). The *minimum-distance graph* of an $S(t, k, n)$ is a graph whose vertices are the blocks of the system, with two vertices joined by an edge if and only if the corresponding blocks intersect in $t - 1$ points. For a Steiner system $S = (N, B)$, we denote by $\chi(S)$ the chromatic number of its minimum-distance graph (i.e. the minimum number of cells in a partition of B such that two distinct blocks from the same cell intersect in at most $t - 2$ points).

In the rest of this section, we list some known facts in the form of lemmas, to be utilized in the proofs of our results.

Lemma 2 (see, e.g., [6]). *The number of blocks in a Steiner system $S(t, k, n)$ is $\binom{n}{t} / \binom{k}{t}$ and every point in N is contained in $\binom{n-1}{t-1} / \binom{k-1}{t-1}$ blocks.*

Lemma 3 ([6, Sect. 5.4]). *For a Steiner system $S(t, k, n)$, the number $\Lambda_i(t, k, n)$ of blocks that intersect a fixed block B in exactly i elements does not depend on the choice of B and equals $\binom{k}{i} \lambda_{i, k-i}$ where $\lambda_{i, j}$, $0 \leq i+j \leq k$ are recursively defined as follows:*

$$\begin{aligned} \lambda_{i,0} &= \binom{n-i}{t-i} / \binom{k-i}{t-i}, & 0 \leq i \leq t, \\ \lambda_{i,0} &= 1, & t < i \leq k, \\ \lambda_{i,j} &= \lambda_{i,j-1} - \lambda_{i+1,j-1}, & 0 \leq i < i+j \leq k. \end{aligned}$$

Lemma 4. *Assume positive integers m and s are given, and we consider representations of m as the sum of s nonnegative integers: $m = m_1 + \dots + m_s$. The value $\sum_i \binom{m_i}{2}$ reaches its minimum if and only if $m_1, \dots, m_s \in \{a, a+1\}$, where $a = \lfloor m/s \rfloor$.*

Proof. Otherwise, $m_l \leq m_j + 2$ for some l, j . In this case, obviously, $\binom{m_l}{2} + \binom{m_j}{2} > \binom{m_l+1}{2} + \binom{m_j-1}{2}$, and $\sum \binom{m_i}{2}$ is not as small as possible. \square

3. Anticodes

Generalizing the anticode $\mathcal{A}^m(n, w, w)$ from [10, IV.DE], we define these two sets:

$$\begin{aligned} \mathcal{A}'_q(n, w, t) &= \{(a_1, \dots, a_t, a_{t+1}, \dots, a_n) \in \mathbb{Z}_q^n : \\ &\quad \text{wt}((a_1, \dots, a_t)) = t, \text{wt}((a_{t+1}, \dots, a_n)) = w - t\}, \end{aligned}$$

where $t \leq w$ and $2w - t \leq n$, and

$$\mathcal{A}''_q(n, w, t) = \{(0, \dots, 0, a_{t+1}, \dots, a_n) \in \mathbb{Z}_q^n : \text{wt}((a_{t+1}, \dots, a_n)) = w\},$$

where $t \leq n - w$ and $2w + t \geq n$. The main theorem in this section establishes that $\mathcal{A}'_q(n, w, t)$ and $\mathcal{A}''_q(n, w, t)$ are maximum anticodes in $\mathcal{J}_q(n, w)$, assuming some conditions on the parameters hold.

Theorem 1. *The set $\mathcal{A}'_q(n, w, t)$, $t \leq 2w - t \leq n$, satisfies the following:*

- (1') $\mathcal{A}'_q(n, w, t)$ has $\binom{n-t}{w-t} (q-1)^w$ words;
- (2') $\mathcal{A}'_q(n, w, t)$ is a diameter- $(2w - t)$ anticode;

(3') if $n \geq (w - t + 1)(t + 1)$ and q is large enough, then $\mathcal{A}'_q(n, w, t)$ is a maximum diameter- $(2w - t)$ anticode in $\mathcal{J}_q(n, w)$.

Theorem 2. The set $\mathcal{A}''_q(n, w, t)$, $n + w \leq n \leq 2w + t$, satisfies the following:

(1'') $\mathcal{A}''_q(n, w, t)$ is a diameter- $(n - t)$ anticode;

(2'') $\mathcal{A}''_q(n, w, t)$ has $\binom{n-t}{w}(q - 1)^w$ words;

(3'') if $n \geq (n - w - t + 1)(t + 1)$ (i.e., $n \leq (w + t - 1)(t + 1)/t$) and q is large enough, then $\mathcal{A}''_q(n, w, t)$ is a maximum diameter- $(n - t)$ anticode in $\mathcal{J}_q(n, w)$.

Claims (1'), (1''), (2'), and (2'') are obvious. To show (3') and (3''), we firstly prove a more general result, of independent interest.

Proposition 1. Let T be a maximum family of different w -subsets of $[n]$ such that every two subsets from T intersect in at least s elements. Let \mathcal{A} be the set of all $|T|(q - 1)^w$ words over \mathbb{Z}_q with supports in T . If q is large enough, then \mathcal{A} is a maximum diameter- $(2w - s)$ anticode.

Proof. Let \mathcal{B} be a diameter- $(2w - s)$ anticode.

If for every two words \bar{a}, \bar{b} in \mathcal{B} it holds $|\text{supp}(\bar{a}) \cap \text{supp}(\bar{b})| \geq s$, then the words from \mathcal{B} have at most $|T|$ different supports, and $|\mathcal{B}| \leq |\mathcal{A}|$.

Assume \mathcal{B} contains \bar{a} and \bar{b} such that $|\text{supp}(\bar{a}) \cap \text{supp}(\bar{b})| < s$. Denote by B the set of all supports of words in \mathcal{B} ; we divide it into two disjoint subsets B_1, B_2 :

$$B_1 = \{\beta \in B : |\beta \cap \alpha| < s \text{ for some } \alpha \in B\},$$

$$B_2 = \{\beta \in B : |\beta \cap \alpha| \geq s \text{ for all } \alpha \in B\}.$$

We state two claims.

- For every β in B_1 , the number of words in \mathcal{B} with support β is smaller than $(q - 1)^w - (q - 2)^w$. Indeed, by the definition of B_1 , there is a word \bar{x} in \mathcal{B} such that $|\beta \cap \text{supp}(\bar{x})| < s$. Any word with support β that differs with \bar{x} in all positions of $\beta \cap \text{supp}(\bar{x})$ cannot belong to \mathcal{B} because of the diameter of \mathcal{B} . The number of such words is larger than $(q - 2)^w$, and the claim follows.

- It holds $|B_2| \leq |T| - 1$. Indeed, by the hypothesis, B contains at least one element α not in $|B_2|$. By the definition of B_2 , each every two distinct sets in $B_2 \cup \{\alpha\}$ intersect in at least s points. Since T is a maximum family with this property, we get $|B_2 \cup \{\alpha\}| = |B_2| + 1 \leq |T|$.

Therefore, we have

$$\begin{aligned} |\mathcal{B}| &< |B_2|(q-1)^w + |B_1|((q-1)^w - (q-2)^w), \\ &< (|T| - 1)(q-1)^w + \binom{n}{w}((q-1)^w - (q-2)^w), \end{aligned}$$

Since the degree of the polynomial $(q-1)^w - (q-2)^w$ is less than w , we have $|\mathcal{B}| < |T|(q-1)^w$ for sufficiently large q , and the result follows. \square

The next lemma implies that the set of supports of the words from $\mathcal{A}'_q(n, w, t)$ or $\mathcal{A}''_q(n, w, t)$ satisfies the hypothesis of Proposition 1.

Lemma 5. [Erdős–Ko–Rado theorem [8, 21] for s -intersecting families] Let $0 \leq s \leq w \leq n$, and let T be a family of w -subsets of $[n]$ such that every two subsets from T intersect in at least s elements.

- (i) If $n \geq (w-s+1)(s+1)$, then $|T| \leq \binom{n-s}{w-s}$.
- (ii) If $2w-s \leq n \leq \frac{(w-s+1)(2w-s-1)}{w-s}$, then $|T| \leq \binom{2w-s}{w}$.

Actually, the classic Erdős–Ko–Rado theorem, proved in [8] for $n \geq s + (k-s)\binom{k}{s}^3$ and in [21] for $n \geq (w-s+1)(s+1)$, does not state (ii), which is a direct corollary of (i), obtained by replacing the sets from T by their complements with respect to $[n]$. All possible maximum s -intersecting families of w -subsets of $[n]$ are described in [2]. Note that each of them, by Proposition 1, produces a sequence of maximum constant-weight anticodes over nonbinary alphabet. We focus on only two types of such anticodes, $\mathcal{A}'_q(n, w, t)$ and $\mathcal{A}''_q(n, w, t)$, because corresponding diameter perfect codes can be constructed.

Proof of (3') and (3''). From Lemma 5(i) we see that the set of supports from $\mathcal{A}'_q(n, w, t)$, under the conditions of Theorem 1, forms a maximum t -intersecting family; hence, (3') follows from Proposition 1.

If we denote $s = 2w - n + t$, then we see that the set of supports from $\mathcal{A}''_q(n, w, t)$ forms an s -intersecting family. The hypothesis $n \leq \frac{(w-s+1)(2w-s-1)}{w-s}$ of Lemma 5(ii) is equivalent to the condition $n \geq (n-w-t+1)(t+1)$ of Theorem 2, while $2w-s \leq n$ is equivalent to the trivial $0 \leq t$. Therefore, we can apply Lemma 5(ii) and conclude that we have a maximum s -intersecting family. Proposition 1 completes the proof of (3''). \square

4. Codes and bounds for the smallest q

Theorem 3. *Let $t, k,$ and n be integers such that $0 < t \leq k < n$. If there exists a Steiner system $S(t, k, n)$, then the following assertions hold.*

- (i) *For $w = k$ and $q \geq \frac{\binom{n-1}{t-1}}{\binom{w-1}{t-1}} + 1$, there is an $(n, 2w - t + 1, w; \binom{n}{t} / \binom{w}{t})_q$ code, which attains the code-anticode bound with the anticode $\mathcal{A}'_q(n, w, t)$.*
- (ii) *For $w = n - k$ and $q \geq \frac{\binom{n}{k}}{\binom{n-1}{k}} - \frac{\binom{n-1}{t-1}}{\binom{k-1}{t-1}} + 1$, there is an $(n, n - t + 1, w; \binom{n}{t} / \binom{n-w}{t})_q$ code, which attains the code-anticode bound with the anticode $\mathcal{A}'_q(n, w, t)$.*

If there is no $S(t, k, n)$, then $(n, 2k - t + 1, k; \binom{n}{t} / \binom{k}{t})_q$ codes and $(n, n - t + 1, n - k; \binom{n}{t} / \binom{k}{t})_q$ codes do not exist for any q .

Proof. (i) Assume there exists a Steiner system $S(t, w, n)$ with point set $[n]$ and blocks $\Delta_1, \Delta_2, \dots, \Delta_M$, $M = \binom{n}{t} / \binom{w}{t}$. Let \bar{x}^i , $i = 1, \dots, M$, be the length- n word containing 0 in the j th position if $j \notin \Delta_i$ and containing a positive value a in the j th position, $j = 1, \dots, n$, if Δ_i is the a th block with j (that is, if $j \in \Delta_i$ and $a = |\{l \in \{1, \dots, i\} : j \in \Delta_l\}|$). According to that definition, $\text{wt}(\bar{x}^i) = w$, and two different words \bar{x}^i and \bar{x}^l coincide only in positions where both have zeros. By the definition of Steiner system, the supports of such \bar{x}^i and \bar{x}^l have no more than $t - 1$ common elements; hence their union has size at least $2w - (t - 1)$, and $d(\bar{x}^i, \bar{x}^l) \geq 2w - t + 1$. It remains to note that by Lemma 2 all codewords involve $\binom{n-1}{t-1} / \binom{w-1}{t-1}$ nonzero symbols. We conclude that $\{\bar{x}^1, \dots, \bar{x}^M\}$ can be treated as an $(n, 2w - t + 1, w; \binom{n}{t} / \binom{w}{t})_q$ code for every $q \geq \frac{\binom{n-1}{t-1}}{\binom{w-1}{t-1}} + 1$.

(ii) Similarly, assume there exists a Steiner system $S(t, n - w, n)$ with point set $[n]$ and blocks $\Delta_1, \Delta_2, \dots, \Delta_M$, $M = \binom{n}{t} / \binom{n-w}{t}$. Similarly to (i), we can construct a code such that the codeword supports are the complements of the Steiner system and for each coordinate all the nonzero elements in the codewords have distinct symbols. All codewords have weight w , and any two different codewords \bar{x} and \bar{y} coincide only in positions where both have zeros. Since such \bar{x} and \bar{y} have no more than $t - 1$ common zero positions, their union has size at least $n - (t - 1)$, and $d(\bar{x}, \bar{y}) \geq n - t + 1$. It remains to note that by Lemma 2 all codewords involve $\binom{n}{t} / \binom{w}{t} - \frac{\binom{n-1}{t-1}}{\binom{w-1}{t-1}}$ nonzero symbols. We conclude that $\{\bar{x}^1, \dots, \bar{x}^M\}$ can be treated as an $(n, n - t + 1, w; \binom{n}{t} / \binom{n-w}{t})_q$ code for every $q \geq \frac{\binom{n}{t}}{\binom{w}{t}} - \frac{\binom{n-1}{t-1}}{\binom{w-1}{t-1}} + 1$.

It remains to prove the necessity of a Steiner system. Assume there exists an $(n, 2w - t + 1, w; \binom{n}{t}/\binom{w}{t})_q$ code. For any two different codewords, the union of their supports cannot have less elements than the code distance, $2w - (t - 1)$; hence, the intersection of the supports have at most $t - 1$ elements in common. Each codeword support contains w elements and includes $\binom{w}{t}$ subsets of size t . Hence, all $\binom{n}{t}/\binom{w}{t}$ codeword supports cover all $\binom{n}{t}$ different t -subsets of $[n]$. We have an $S(t, w, n)$.

Similarly, assume an $(n, n - t + 1, w; \binom{n}{t}/\binom{n-w}{t})_q$ code exists. From the code distance $n - (t - 1)$, we see that two different codewords have at most $t - 1$ common positions with zeros. That is, the complements of their supports intersect in less than t elements. Again, from the code cardinality we see that we have an $S(t, n - w, n)$. \square

Remark 1. The special cases $w = t$ and $n - w = t$ of Theorem 3 (cases (i) and (ii), respectively) correspond to [10, Theorem 16], where the code distance is $w + 1$. If $w > t$ and $n - w > t$, the code distances in Theorem 3 are larger than $w + 1$.

The bounds for q given in Theorem 3 are not best possible. For each t , k , and n such that there is a Steiner system $S(t, k, n)$, we denote

- by $q'_0(t, k, n)$ the smallest q such that an $(n, 2k - t + 1, k; \binom{n}{t}/\binom{k}{t})_q$ code exists
- by $q''_0(t, k, n)$ the smallest q such that an $(n, n - t + 1, n - k; \binom{n}{t}/\binom{k}{t})_q$ code exists

We emphasize that in the notation $q'_0(t, k, n)$ the parameter k coincides with the weight of the corresponding constant-weight codes; that is why we will often see w at the place of that parameter. The situation is different with $q''_0(t, k, n)$, where the corresponding weight is $n - k$. From Theorem 3, we get the following bounds for q'_0 and q''_0 , which will be called *trivial* and can be further improved.

Corollary 1. $q'_0(t, w, n) \leq \frac{\binom{n-1}{t-1}}{\binom{w-1}{t-1}} + 1$; $q''_0(t, k, n) \leq \frac{\binom{n}{t}}{\binom{k}{t}} - \frac{\binom{n-1}{t-1}}{\binom{k-1}{t-1}} + 1$.

Below, we find some values of q'_0 and q''_0 for small t , improve the trivial bound for arbitrary t , and consider in details some concrete parameter collections (t, w, n) .

4.1. Bounds for q'_0 : $t = 2$ and $t = 3$

Theorem 4. *If there exists an $S(2, w, n)$, then $q'_0(2, w, n) = \frac{n-1}{w-1} + 1$.*

Proof. According to Corollary 1, $q'_0(2, w, n) \leq \frac{n-1}{w-1} + 1$.

On the other hand, assume we have an $(n, 2w - 1, w; \cdot)_q$ code C constructed from $S(n, w, 2)$. Take any two codewords \bar{x}, \bar{y} in C . By the code parameters, $\text{wt}(x) = \text{wt}(y) = w$ and $d(\bar{x}, \bar{y}) \geq 2w - 1$. Let

$$k = |\{i \in [n] : x_i = y_i \neq 0\}|.$$

We have $2w - 1 \leq d(\bar{x}, \bar{y}) \leq 2(w - k)$; hence $k = 0$. By Lemma 2, at any position, there are $\frac{n-1}{w-1}$ codewords in C that have a non-zero symbol at this position. From $k = 0$, we see that all those non-zero symbols are distinct. Therefore, $q'_0(2, w, n) \geq \frac{n-1}{w-1} + 1$. \square

Remark 2. For $t = w = 2$, we have $q'_0(2, 2, n) = n$; this case was considered in [10, Corollary 11].

Example 1. From the Steiner system $S(2, 3, 7)$, with the blocks

$$\{1, 2, 3\}, \{1, 4, 5\}, \{1, 6, 7\}, \{2, 4, 6\}, \{2, 5, 7\}, \{3, 4, 7\}, \{3, 5, 6\},$$

we can construct the following $(7, 5, 3; 7)_q$ code, $q \geq 4$:

$$\{1110000, 2001100, 3000011, 0202020, 0300202, 0023003, 0030330\}.$$

The anticode $\mathcal{A}'_q(7, 3, 2)$ has diameter 4 and size $5(q - 1)^3$. It follows from the existence of the code that $\mathcal{A}'_q(7, 3, 2)$ is a maximum diameter-4 anticode of length 7 and weight 3 over \mathbb{Z}_q , $q \geq 4$.

Theorem 5. *If there exists an $S(3, w, n)$, then*

$$q'_0(3, w, n) = \min_{S: S(3, w, n)} \max_{D \in S'} \chi(D) + 1,$$

where S' is the set of all $S(2, w - 1, n - 1)$ derived from S .

Proof. Assume there exists a Steiner system $S(3, w, n)$ with point set $[n]$ and block set $B = \{\Delta_1, \dots, \Delta_M\}$, $M = \binom{n}{3} / \binom{w}{3}$. Denote by S_i the $S(2, w - 1, n - 1)$ system on $[n] \setminus \{i\}$ derived system from the $S(3, w, n)$, and by λ_i its chromatic number $\chi(S_i)$, $i = 1, \dots, n$. What is more, we denote by B_i the

set of all blocks from B containing i . Next, according to the definition of $\chi(S_i)$, we can partition B_i into λ_i disjoint subsets:

$$B_i = B_i^1 \cup B_i^2 \cup \dots \cup B_i^{\lambda_i}$$

where B_i^r , $r = 1, \dots, \lambda_i$, can be viewed as a cell, and the blocks of B_i from the same cell mutually intersect in exactly one element i .

Let \bar{x}^i , $i = 1, \dots, M$, be the length- n word containing 0 in the j th position if $j \notin \Delta_i$ and containing a positive value s in the j th position, if $j \in \Delta_i$ and $\Delta_i \in B_j^s$ for some $s \in [\lambda_j]$ (in fact, if $j \in \Delta_i$, then $\Delta_i \in B_j$ and there must exist an integer s such that $\Delta_i \in B_j^s$). Therefore, $\text{wt}(\bar{x}^i) = w$, and take any two different words \bar{x}^i and \bar{x}^l , they coincide either only in positions where both have zeros, or in positions where have nonzero. The first case was analyzed in the proof of Theorem 3; at this time, $d(\bar{x}^i, \bar{x}^l) \geq 2w - 2$. For the second case, assume \bar{x}^i and \bar{x}^l both have a positive value s , $s \in [\lambda_j]$, in the j th position. By the definition of \bar{x}^i , we have $j \in \Delta_i \in B_j^s$ and $j \in \Delta_l \in B_j^s$; that means Δ_i, Δ_l are in the same cell B_j^s and $\Delta_i \cap \Delta_l = \{j\}$, at this time, $d(\bar{x}^i, \bar{x}^l) = 2w - 2$. Thus all codewords involve $\max_{D \in S'} \chi(D)$ nonzero symbols, and $\{\bar{x}^1, \dots, \bar{x}^M\}$ can be treated as an $(n, 2w - 2, w; \binom{n}{3} / \binom{w}{3})_q$ code for $q = \max_{D \in S'} \chi(D) + 1$, and it is easy to verify that q reaches the minimum for given $S(3, w, n)$, thus for any Steiner system $S(3, w, n)$, the result follows. \square

Corollary 2. *If there is an $S(3, w, n)$ such that each its derived $S(2, w - 1, n - 1)$ is resolvable, then*

$$q'_0(3, w, n) = \frac{n - 2}{w - 2} + 1. \quad (2)$$

Proof. For an $S(3, w, n)$, each its derived $S(2, w - 1, n - 1)$ is resolvable, that is to say, the set of blocks B of each derived $S(2, w - 1, n - 1)$ can be partitioned into subsets each of which forms an $S(1, w - 1, n - 1)$ on the same $s \subset N$. In other words, assume S' is the set of all $S(2, w - 1, n - 1)$ derived from S , then for any $D \in S'$, we have $\chi(D) = \frac{\binom{n-1}{2}}{\binom{w-1}{2}} \cdot \frac{w-1}{n-1} = \frac{n-2}{w-2}$, and according to Theorem 5, $q'_0(3, w, n) = \frac{n-2}{w-2} + 1$. \square

Remark 3. The case $t = w = 3$ was solved in [10, Theorem 17]: if n is odd then $q'_0(3, 3, n) = n - 1$, and $q'_0(3, 3, n) = n$ if even.

Trivially, all $S(t-1, w-1, n-1)$ systems derived from a 2-resolvable $S(t, w, n)$ are resolvable. It is proved in [3] that there are 2-resolvable $S(3, 4, 4^m)$ for each $m \geq 1$. In [19], 2-resolvable Steiner systems $S(3, 4, n)$ are constructed for $n = 2 \cdot p^m + 2$, $p \in \{7, 31, 127\}$, $m \geq 1$. Moreover, from Keevash's theory [12] we know that for given w and n large enough, 2-resolvable $S(3, w, n)$ systems exist if and only if the obviously necessary divisibility conditions

$$w(w-1) \mid n(n-1), \quad (w-1) \mid (n-1), \quad (w-2) \mid (n-2)$$

are satisfied. In particular, for $w = 4$, those conditions are equivalent to $n \equiv 4 \pmod{12}$. However, for Corollary 2, the condition $n \equiv 4 \pmod{12}$ (as well as the 2-divisibility of $S(3, 4, n)$) is not necessary; for example all $S(2, 3, 9)$ derived from $S(3, 4, 10)$ are resolvable, while $10 \not\equiv 4 \pmod{12}$ and $S(3, 4, 10)$ is not 2-resolvable.

Corollary 3. *If $n = 4^m$ or $n = 2 \cdot p^m + 2$, $p \in \{7, 31, 127\}$, $m \in \{1, 2, \dots\}$, or n large enough satisfying $n \equiv 4 \pmod{12}$, or $n = 10$, then $q'_0(3, 4, n) = n/2$.*

Remark 4. In the case of $S(3, 4, n)$, (2) turns to $q_0(3, 4, n) = n/2$. In [23, Lemma 2.12], 2-resolvable $S(3, 4, n)$ are used to construct optimal $(n, 6, 4; (q-1)n(n-1)/12)_q$ codes for every q in $\{3, \dots, n/2\}$. For $q = n/2$, this gives the same result as in Corollary 3, except the case $n = 10$.

4.2. Bounds for q'_0 : arbitrary t

In this section, we generalize Theorem 5 to arbitrary t . It is hard to expect that the generalization gives a good evaluation of the real value of q'_0 for $t \geq 3$; however, it is important to conclude that the trivial bound from Corollary 1 is never tight. It is also a little step towards the research direction pointed in [10] as Problem 4.

Theorem 6. *If there exists an $S(t, w, n)$,*

$$q'_0(t, w, n) \leq \frac{\binom{n-1}{t-1}}{\binom{w-1}{t-1}} - \frac{\binom{n-t+2}{2}}{\binom{w-t+2}{2}} + \min_{S: S(n, w, t)} \max_{D \in S'} \chi(D) + 1,$$

where S' is the set of all $S(2, w-t+2, n-t+2)$ derived from S .

Proof. Assume there exist a Steiner system $S(t, w, n)$ with point set $[n]$ and block set $B = \{\Delta_1, \dots, \Delta_M\}$, $M = \binom{n}{t} / \binom{w}{t}$. For convenience, we consider all

operations over $[n]$ modulo n . Denote by S_i the $S(2, w-t+2, n-t+2)$ system on $[n] \setminus \{i-t+3, \dots, i\}$ derived from the $S(3, w, n)$, and by λ_i its chromatic number $\chi(S_i)$, $i = 1, \dots, n$. We denote B_i as a subset of B which is the set of all blocks that include the subset $\{i-t+3, \dots, i\}$. By the definition of $\chi(S_i)$, we can partition B_i into λ_i disjoint subsets:

$$B_i = B_i^1 \cup B_i^2 \cup \dots \cup B_i^{\lambda_i}$$

where B_i^r , $r = 1, \dots, \lambda_i$, can be viewed as a cell, and the blocks of B_i from the same cell mutually intersect exactly in $\{i-t+3, \dots, i\}$. Let \bar{x}^i , $i = 1, \dots, M$, be the length- n word containing 0 in the j th position if $j \notin \Delta_i$, containing a positive value s , in the j th position, if $j \in \Delta_i$ and $\Delta_i \in B_j^s$ for some $s \in [\lambda_j]$, and containing a positive value a , in the j th position, $j \in [n]$, if $j \in \Delta_i$ and $\Delta_i \notin B_j$, where $a = |\{l \in \{1, \dots, i\} : j \in \Delta_l\}| + \lambda_j$. Therefore, $\text{wt}(\bar{x}^i) = w$, and take any two different words \bar{x}^i and \bar{x}^l , they coincide either only in positions where both have zeros, or in positions where have nonzero. According to the proof of Theorem 3, we just consider the case that two distinct words \bar{x}^i and \bar{x}^l both have a positive value s , $s \in [\lambda_j]$, in the j th position. By the definition of \bar{x}^i , we have that Δ_i, Δ_l are in the same cell B_j^s and $\Delta_i \cap \Delta_l = \{j-t+3, \dots, j\}$, and it is easy to verify that \bar{x}^i, \bar{x}^l only coincide in j ; at this time, $d(\bar{x}^i, \bar{x}^l) = 2w - t + 1$. And all codewords involve $\binom{n-1}{t-1} / \binom{w-1}{t-1} - \binom{n-t+2}{2} / \binom{w-t+2}{2} + \min_{S: S(n, w, t)} \max_{D \in S'} \chi(D)$ nonzero symbols, and $\{\bar{x}^1, \dots, \bar{x}^M\}$ can be treated as an $(n, 2w - t + 1, w; \binom{n}{t} / \binom{w}{t})_q$ code for every $q \geq \binom{n-1}{t-1} / \binom{w-1}{t-1} - \binom{n-t+2}{2} / \binom{w-t+2}{2} + \min_{S: S(n, w, t)} \max_{D \in S'} \chi(D) + 1$. \square

4.3. Bounds for q'_0 : S(4,5,11), SQS(14), and SQS(20)

In this section, we evaluate $q'_0(4, 5, 11)$, related to a unique $S(4, 5, 11)$, derived from the small Witt design $S(5, 6, 12)$ [22], and find $q'_0(3, 4, 14)$ and $q'_0(3, 4, 20)$.

Proposition 2. *It holds $9 \leq q'_0(4, 5, 11) \leq 11$.*

Proof. Computing $\chi(S')$ for the unique (up to equivalence) $S(3, 4, 10)$, which is derived from $S(4, 5, 11)$, gives $q'_0(4, 5, 11) \geq \chi(S') + 1 = 9$.

An example of a $(11, 7, 5; 66)_{11}$ code, which shows that $q'_0(4, 5, 11) \leq 11$, is the cyclic code with representatives 0000a100615, 00080901087, 00103009406, 00022013005, 00070040726, 0000035480a (each representative generates 11 codewords by cyclic shifting). \square

Proposition 3. *The values $q'_0(3, 4, 14)$ and $q'_0(3, 4, 20)$ equal 9 and 11, respectively.*

Proof. There are two STS(13), up to permutation of points, and each of them has chromatic number 8 (a computational result). It follows that $q'_0(3, 4, 14) = 8 + 1$.

The number of blocks in an STS(19) S' is 57, and the maximum number of mutually disjoint blocks is $\lfloor 19/3 \rfloor = 6$. So, $\chi(S') \geq \lceil 57/6 \rceil = 10$ and hence $q'_0(3, 4, 20) \geq 11$.

Consider the cyclic SQS(20) (II.A.1) from [16]. It consists of the blocks $\{1, 6, 11, 16\}$, $\{1, 3, 11, 13\}$, $\{1, 5, 11, 15\}$, $\{1, 4, 9, 17\}$, $\{1, 4, 7, 14\}$, $\{1, 2, 3, 12\}$, $\{1, 5, 7, 10\}$, $\{1, 3, 7, 8\}$, $\{1, 3, 4, 10\}$, $\{1, 6, 7, 9\}$, $\{1, 3, 9, 14\}$, $\{1, 2, 6, 10\}$, $\{1, 2, 9, 18\}$, $\{1, 2, 7, 13\}$, $\{1, 2, 5, 14\}$, $\{1, 4, 6, 8\}$ and all their cyclic shifts $\{a_1, a_2, a_3, a_4\} \rightarrow \{a_1 + i, a_2 + i, a_3 + i, a_4 + i\}$, where $+$ is modulo 20. Because of the cyclicity, all derived STS are isomorphic, and we consider the derived STS S' on the point set [19]. The block set of the derived STS is partitioned into 10 cells with mutually disjoint blocks in each cell:

$\{\{1, 2, 11\}, \{3, 5, 7\}, \{4, 10, 14\}, \{6, 17, 19\}, \{8, 12, 15\}, \{13, 16, 18\}\}$,
 $\{\{1, 3, 15\}, \{2, 10, 12\}, \{5, 13, 17\}, \{6, 11, 18\}, \{7, 16, 19\}, \{8, 9, 14\}\}$,
 $\{\{1, 4, 13\}, \{2, 15, 18\}, \{3, 12, 19\}, \{5, 6, 8\}, \{7, 14, 17\}, \{9, 10, 11\}\}$,
 $\{\{1, 5, 9\}, \{2, 8, 13\}, \{3, 14, 18\}, \{4, 7, 12\}, \{6, 10, 16\}, \{11, 15, 17\}\}$,
 $\{\{1, 6, 12\}, \{2, 4, 17\}, \{3, 8, 16\}, \{5, 10, 15\}, \{9, 18, 19\}, \{11, 13, 14\}\}$,
 $\{\{1, 7, 18\}, \{2, 5, 16\}, \{3, 10, 17\}, \{4, 8, 19\}, \{6, 14, 15\}, \{9, 12, 13\}\}$,
 $\{\{1, 8, 17\}, \{2, 14, 19\}, \{3, 6, 13\}, \{4, 5, 18\}, \{7, 9, 15\}, \{11, 12, 16\}\}$,
 $\{\{2, 6, 7\}, \{3, 4, 11\}, \{5, 12, 14\}, \{8, 10, 18\}, \{9, 16, 17\}, \{13, 15, 19\}\}$,
 $\{\{1, 10, 19\}, \{2, 3, 9\}, \{4, 15, 16\}, \{7, 8, 11\}, \{12, 17, 18\}\}$,
 $\{\{1, 14, 16\}, \{4, 6, 9\}, \{5, 11, 19\}, \{7, 10, 13\}\}$.

So, $\chi(S') = 10$ and $q'_0(20, 4, 3) = \chi(S') + 1 = 11$. □

4.4. Bounds for q''_0

The trivial upper bound on $q''_0(t, k, n)$ in Corollary 1 comes from a construction where in each column of the code matrix all nonzero symbols are distinct. Merging some symbols results in decreasing the distance between some codewords. However, this does not always lead to decreasing the code distance, and in general a code with the same parameters and a smaller number of symbols used can be found. This is not the case if $t = 1$ because the distance

between any two codewords in the trivial construction already coincides with the minimum distance of the code.

Corollary 4. $q_0''(1, k, n) = n/k$, i.e., an $(n, n, n - k; n/k)_q$ code exists if and only if k divides n (necessary and sufficient condition for the existence of $S(1, k, n)$) and $q \geq n/k$.

In the following proposition, we estimate the number of symbol pairs that can be merged if $t > 1$. The formula there looks a bit complicate, but as we see from the corollaries below, for special cases it becomes much simpler.

Proposition 4. Assume there exists an $(n, n - t + 1, w; M)_q$ code C , where $w = n - k$ and $M = \binom{n}{t} / \binom{k}{t}$.

- (i) C is a diameter perfect code in $\mathcal{J}_q(n, w)$;
- (ii) the supports of the codeword of C are the complements of the blocks of an $S(t, k, n)$ system S ;
- (iii) it holds

$$\begin{aligned}
 (q - 1) \binom{a}{2} + ba &\leq \lfloor \tilde{P} \rfloor, & \text{where} \\
 a &= \lfloor R / (q - 1) \rfloor, & b = R - a(q - 1), \\
 R &= \binom{n}{t} / \binom{k}{t} - \binom{n - 1}{t - 1} / \binom{k - 1}{t - 1}, \\
 \tilde{P} &= \frac{M}{2n} \sum_{i=0}^{t-2} (t - 1 - i) \Lambda_i(t, k, n),
 \end{aligned}$$

and $\Lambda_i(t, k, n) = \binom{k}{i} \lambda_{i, k-i}$ is from Lemma 3;

- (iii') in particular, $q - 1 \geq R - \lfloor \tilde{P} \rfloor$.

Proof. (i) Since C meets the code–anticode bound with the anticode $\mathcal{A}_q''(n, n - k, t)$, it is diameter perfect.

(ii) The cardinality of the code coincides the number of blocks in $S(t, k, n)$. Each codeword has exactly k zeros. Each two codewords have at most $t - 1$ common position with a zero. The claim is now straightforward.

(iii) The code matrix is an $M \times n$ matrix. Each row contains k zeros and $n - k$ nonzeros. Each column contains $\binom{n-1}{t-1} / \binom{k-1}{t-1}$ zeros (Lemma 2) and

$$R = \binom{n}{t} / \binom{k}{t} - \binom{n-1}{t-1} / \binom{k-1}{t-1} = \frac{(n-1)!(k-t)!(n-k)}{(n-t)!k!}$$

nonzeros. The set of zero positions of each row corresponds to some block of S .

We will say that we have a *collision* (a, b) if two different cells a, b in the same column contain the same nonzero value. Let us evaluate the maximum number of collisions. If two rows correspond to blocks with i common positions, then they have at most $t - 1 - i$ collisions because the code distance is $n - t + 1$. We denote the number of pairs of such blocks by P_i . Utilizing Lemma 3 we find that $P_i = \frac{M}{2} \cdot \Lambda_i(t, k, n)$. So, the number of collisions does not exceed $P = \sum_{i=1}^{t-1} (t - 1 - i)P_i$. The average number of collisions in one column does not exceed P/n , and there is a column with at most $\tilde{P} = \lfloor P/n \rfloor$ collisions.

On the other hand, the number of nonzeros in each column is R , while the number of different nonzero symbols is $q - 1$. With those constants, the minimum number of collisions is possible if each nonzero symbol occurs a or $a + 1$ times (see Lemma 4), where $a = \lfloor R/(q - 1) \rfloor$. This gives the left part of the inequality in (iii). (iii') coincides with (iii) if $a = 1$ (i.e., when \tilde{P} is not more than the half of the number of ones in a column of the code matrix); otherwise (iii') is weaker. \square

Corollary 5. *In the case $t = 2$, Proposition 4 holds with*

$$R = \frac{(n-1)(n-k)}{k(k-1)} \quad \text{and} \quad \tilde{P} = \frac{(n-1)((n-k^2)(n-1) + k(k-1)^2)}{2k^2(k-1)^2}.$$

Proof. Substituting $t = 2$, we get

$$M = \frac{n(n-1)}{k(k-1)}, \quad R = \frac{(n-1)(n-k)}{k(k-1)}, \quad \text{and} \quad \tilde{P} = \frac{M}{2n} \cdot \lambda_{0,k}.$$

From the definition of $\lambda_{i,j}$ in Lemma 3, we find that for $j \geq 1$

$$\begin{aligned} \lambda_{i,j} &= \lambda_{i,j-1} - \lambda_{i+1,j-1} = 0 \quad \text{if } i \geq t; \\ \lambda_{t-1,j} &= \lambda_{t-1,j-1} - \lambda_{t,j-1} = \frac{n-t+1}{k-t+1} - 1; \end{aligned}$$

$$\begin{aligned}\lambda_{t-2,j} &= \lambda_{t-2,j-1} - \lambda_{t-1,j-1} \\ &= \frac{(n-t+2)(n-t+1)}{(k-t+2)(k-t+1)} - j \cdot \frac{n-t+1}{k-t+1} - (j-1).\end{aligned}$$

Substituting $t = 2$ and $j = k$, we get

$$\lambda_{0,k} = \frac{n(n-1)}{k(k-1)} - k \cdot \frac{n-1}{k-1} - (k-1) = \frac{n(n-1) - k^2(n-1) - k(k-1)^2}{k(k-1)},$$

It is easy to check now that $\frac{M}{2n}\lambda_{0,k}$ is exactly the expression for \tilde{P} from the claim of the corollary. \square

The next claim is about Steiner systems related to affine planes.

Corollary 6. *If k is a prime power, then $q_0''(2, k, k^2) = k^2 - \lfloor \frac{k-1}{2} \rfloor$.*

Proof. For $n = k^2$, \tilde{P} turns to $\frac{k^2-1}{2k}$, and $\lfloor \tilde{P} \rfloor = \lfloor \frac{k}{2} - \frac{1}{2k} \rfloor = \lfloor \frac{k-1}{2} \rfloor$. At the same time, $\frac{(n-1)(n-k)}{k(k-1)} = k^2 - 1$. We see that (iii') turns to $q \geq k^2 - \lfloor \frac{k-1}{2} \rfloor$.

Next, we are going to construct a $(k^2, k^2 - 1, k^2 - k; k^2 + k)_q$ code such that $q = k^2 - \lfloor \frac{k-1}{2} \rfloor$ and the supports of codewords form an $S(2, k, k^2)$ system. For the role of the system, we take the affine plane over the finite field \mathbb{F}_k of order k . The points are the pairs from \mathbb{F}_k^2 (to treat the points as indices, we identify them with the integers from $[k^2]$, in an arbitrary fixed order); the lines (blocks of the $S(2, k, k^2)$) are sets of points (x, y) satisfying one of the equations $x + ay = b$, where $a, b \in \mathbb{F}_w$, or $y = b$, where $a \in \mathbb{F}_w$. We want the code matrix to have $k^2 \cdot \lfloor \frac{k-1}{2} \rfloor$ collisions in total, each column to have $\lfloor \frac{k-1}{2} \rfloor$ pairwise disjoint collisions, and each two rows no more than one collision. We consider two cases, depending on the parity of k .

Odd k , $\lfloor \frac{k-1}{2} \rfloor = \frac{k-1}{2}$. Let S be a subset of \mathbb{F}_k of size $\frac{k-1}{2}$ such that $0, 1 \notin S$ and for every s from \mathbb{F}_k^* it holds $|S \cap \{s, -s\}| = 1$.

For each x_0, y_0 from \mathbb{F}_k and s from S , we make a collision in the column corresponding to the point (x_0, y_0) and the lines

$$\{(x, y) : x + (y_0 + s)y = x_0 + (y_0 + s)y_0 - 1\}, \quad (3)$$

$$\{(x, y) : x + (y_0 + s)y = x_0 + (y_0 + s)y_0 - 1 + s\}. \quad (4)$$

We now check the required properties.

- First of all, we see that the point (x_0, y_0) does not lie on any of the two lines (3), (4), hence the code matrix has nonzeros in the corresponding two cells and we can make a collision there.

- Next, each point corresponds to $|S| = \lfloor \frac{k-1}{2} \rfloor$ collisions and the collisions are pairwise disjoint (for different s , the coefficients at y are different, and hence the lines are different and the corresponding rows are different too). So, after making all these collisions, the number of symbols used in each column becomes $k^2 - \lfloor \frac{k-1}{2} \rfloor$, as required.
- Finally, for any two parallel lines $x+ay = b$ and $x+ay = c$, we uniquely find s from $s \in \{b-c, c-b\}$, then y_0 from $y_0 + s = a$ and x_0 from c or b . This means that the rows corresponding to such two lines have exactly one collision.

Additionally, any two intersecting lines have no collisions, and we conclude that the code distance is $k^2 - 1$, as required.

Even k , $\lfloor \frac{k-1}{2} \rfloor = \frac{k}{2} - 1$. Let $S_{x_0}, S_{x_0} \subset \mathbb{F}_k$, be defined for each x_0 from \mathbb{F}_k in such a way that:

- (i) $0, 1 \notin S_{x_0}$,
- (ii) for every s from $\mathbb{F}_k \setminus \{0, 1\}$, it holds $|S_{x_0} \cap \{s, s+1\}| = 1$,
- (iii) if $s \in S_{x_0}$, then $s \notin S_{x_0+s}$.

Let us show that such $S_{x_0}, x_0 \in \mathbb{F}_k$, exist. For every pair $\{s, s+1\}$, $s \in \mathbb{F}_k \setminus \{0, 1\}$, we divide \mathbb{F}_k into $k/4$ pairs quadruples of form $\{z, z+s, z+1, z+s+1\}$. Then, for each such quadruple, we include s in S_z and S_{z+1} , but not in S_{z+s} or S_{z+s+1} , and include $s+1$ in S_{z+s} and S_{z+s+1} , but not in S_z or S_{z+1} . This rule depends on the choice of the representative z in each quadruple, but once the representatives are chosen, it uniquely specifies for each x_0 , which element of $\mathbb{F}_k \setminus \{0, 1\}$ belongs to S_{x_0} and which element does not belong. Moreover, we see that (ii) and (iii) are satisfied automatically and we can require (i).

Now we proceed similarly to the case of odd k , but with S_{x_0} instead of S . For each x_0, y_0 from \mathbb{F}_k and s from S_{x_0} , we make a collision in the column corresponding to the point (x_0, y_0) and lines (3), (4). The properties are also checked similarly, except the last one, verifying of which differs a bit:

- For any two parallel lines $x+ay = b$ and $x+ay = c$, we uniquely find s from $s = b-c$, then y_0 from $y_0 + s = a$ and finally x_0 from $x_0 + (y_0 + s)y_0 - 1 \in \{c, b\}$ and (iii).

□

Remark 5. We did not make any collisions with lines of form $y = b$. Potentially, they can also be used to decrease the number of used symbols in some columns, but it is not enough to decrease the alphabet in all columns.

Another special case is $S(2, k, k^2 - k + 1)$ known as projective plane of order $k - 1$. It is known that in a projective plane every two blocks (lines) have a common point. It follows that in a column of the code matrix all nonzero symbols are distinct (alternatively, one can substitute $n = k^2 - k + 1$ and see that $P = 0$ in that case).

Corollary 7. *If there exists a projective plane of order s , then*

$$q_0''(2, s + 1, s^2 + s + 1) = s^2 + 1.$$

5. Conclusion

In this paper, we explore two new classes of diameter perfect constant-weight nonbinary codes and estimated the smallest values q_0' and q_0'' of the alphabet size for which such codes exist. Further studying of these values, for special parameters, is related to some problems in the design theory, related to some weak form of resolvability. Below, we formulate those problems in the form of conjectures. Another interesting connection of weakly resolvable designs with optimal non-binary constant-weight codes is shown in [5]

It is well known that resolvable STS(v) exist if and only if $v \equiv 3 \pmod{6}$ [17]. To show that $q_0'(3, 4, n) = n/2$, (Corollary 2) we need the following:

Conjecture 1. *If $n \equiv 4 \pmod{6}$, then there is an SQS(n) such that all its derived STS are resolvable.*

Up to now, we only know that the conjecture is true for $n = 10$ and in the cases when 2-resolvable SQS(n) are known to exist: for $n = 4^m$ [3]; for $n = 2 \cdot p^m + 2$, $p \in \{7, 31, 127\}$ [19]; for sufficiently large n , $n \equiv 4 \pmod{12}$ [12].

If $v \equiv 1 \pmod{6}$, then an STS(v) S' cannot be resolvable, and the maximum number of mutually disjoint blocks in such a system is $(v - 1)/3$. So, $\chi(S') \geq \lceil \frac{v(v-1)/6}{(v-1)/3} \rceil = \frac{v+1}{2}$.

Conjecture 2. *If $20 \leq n \equiv 2 \pmod{6}$, then there is an SQS(n) such that all its derived STS have chromatic number $n/2$, i.e., can be partitioned into $n/2$ partial parallel classes (equivalently, $q_0'(3, 4, n) = n/2 + 1$).*

References

- [1] R. Ahlswede, H. K. Aydinian, and L. H. Khachatrian. On perfect codes and related concepts. *Des. Codes Cryptography*, 22(3):221–237, 2001. DOI: 10.1023/A:1008394205999.
- [2] R. Ahlswede and L. H. Khachatrian. The complete intersection theorem for systems of finite sets. *Eur. J. Comb.*, 18(2):125–136, 1997. DOI: 10.1006/eujc.1995.0092.
- [3] R. D. Baker. Partitioning the planes of $AG_{2^m}(2)$ into 2-designs. *Discrete Math.*, 15(3):205–211, 1976. DOI: 10.1016/0012-365X(76)90025-X.
- [4] L. A. Bassalygo. New upper bounds for error correcting codes. *Probl. Inf. Transm.*, 1(4):32–35, 1965. Translated from *Probl. Peredachi Inf.*, 1(4): 41–44, 1965.
- [5] L. A. Bassalygo, V. A. Zinoviev, and V. S. Lebedev. Weakly resolvable block designs and nonbinary codes meeting the johnson bound. *Probl. Inf. Transm.*, 58(1):3–15, 2022. DOI: 10.1134/S003294602201001X, translated from *Probl. Peredachi Inf.*, 58(1):84–90, 2022.
- [6] C. J. Colbourn and R. Matheron. Steiner systems. In C. J. Colbourn and J. H. Dinitz, editors, *Handbook of Combinatorial Designs*, Discrete Mathematics and Its Applications, pages 102–108. Chapman & Hall/CRC, Boca Raton, London, New York, second edition, 2006.
- [7] P. Delsarte. *An Algebraic Approach to Association Schemes of Coding Theory*, volume 10 of *Philips Res. Rep., Supplement*. N.V. Philips’ Gloeilampenfabrieken, Eindhoven, 1973.
- [8] P. Erdős, C. Ko, and R. Rado. Intersection theorems for systems of finite sets. *Q. J. Math., Oxf. II. Ser.*, 12:313–320, 1961. DOI: 10.1093/qmath/12.1.313.
- [9] T. Etzion. Optimal constant weight codes over Z_k and generalized designs. *Discrete Math.*, 169(1-3):55–82, 1997. DOI: 10.1016/0012-365X(95)00333-R.
- [10] T. Etzion. Non-binary diameter perfect constant-weight codes. *IEEE Trans. Inf. Theory*, 68(2):891–904, 2022. DOI: 10.1109/TIT.2021.3125277.

- [11] T. Etzion and J. H. van Lint. On perfect constant weight codes. In *Codes and Association Schemes*, volume 56 of *DIMACS, Ser. Discrete Math. Theor. Comput. Sci.*, pages 131–134. DIMACS/AMS, 2001. DOI: 10.1090/dimacs/056.
- [12] P. Keevash. The existence of designs. E-print 1401.3665, arXiv.org, 2014. Available at <http://arxiv.org/abs/1401.3665>.
- [13] D. S. Krotov. Inductive constructions of perfect ternary constant-weight codes with distance 3. *Probl. Inf. Transm.*, 37(1):1–9, 2001. DOI: 10.1023/A:1010424208992.
- [14] D. S. Krotov. On diameter perfect constant-weight ternary codes. *Discrete Math.*, 308(14):3104–3114, 2008. DOI: 10.1016/j.disc.2007.08.037.
- [15] D. S. Krotov, P. R. J. Östergård, and O. Pottönen. Non-existence of a ternary constant weight $(16, 5, 15; 2048)$ diameter perfect code. *Adv. Math. Commun.*, 10(2):393–399, May 2016. DOI: 10.3934/amc.2016013.
- [16] K. T. Phelps. On cyclic Steiner systems $S(3,4,20)$. In *Topics on Steiner Systems*, volume 7 of *Ann. Discrete Math.*, pages 277–300. 1980.
- [17] D. K. Ray-Chaudhuri and R. M. Wilson. Solution of Kirkman’s school-girl problem. In *Combinatorics, Proc. Sympos. Pure Math., Univ. California, Los Angeles, Calif., 1968*, volume 19, pages 187–203, 1971.
- [18] M. Svanström. A class of perfect ternary constant-weight codes. *Des. Codes Cryptography*, 18(1-3):223–229, 1999. DOI: 10.1023/A:1008361925021.
- [19] L. Teirlinck. Some new 2-resolvable Steiner quadruple systems. *Des. Codes Cryptography*, 4(1):5–10, 1994. DOI: 10.1007/BF01388556.
- [20] J. van Lint and L. Tolhuizen. On perfect ternary constant-weight codes. *Des. Codes Cryptography*, 18(1-3):231–234, 1999. DOI: 10.1023/A:1008314009092.
- [21] R. M. Wilson. The exact bound in the Erdős–Ko–Rado theorem. *Combinatorica*, 4(2-3):247–257, June 1984. DOI: 10.1007/BF02579226.
- [22] V. E. Witt. Über Steinersche systeme. *Abh. Math. Semin. Univ. Hamb.*, 12(1):265–275, 1937. DOI: 10.1007/BF02948948.

- [23] H. Zhang and G. Ge. Optimal ternary constant-weight codes of weight four and distance six. *IEEE Trans. Inf. Theory*, 56(5):2188–2203, May 2010. DOI: 10.1109/TIT.2010.2043870.