

# AN INEQUALITY OF MULTIPLE INTEGRAL OF $s$ NORM OF VECTORS IN $\mathbb{R}^n$

T. AGAMA

ABSTRACT. In this paper, we prove some new inequalities. To facilitate this proof, we introduce the notion of the local product on a sheet and associated space. We apply this notion to prove the main result in this paper.

## 1. Introduction

There is hardly a formal introduction to the concept of an inner product and associated space in the literature. The inner product space is usually a good place to go for a wide range of mathematical results, from identities to inequalities. In this situation, the best potential result is frequently obtained. The Cauchy-Schwartz inequality obtained in the case of the Hilbert space [1] is a good example. In [2], the notion of the **outer product** and associated **outer product space** was studied and used to prove some spectral inequalities. In this study, however, the notion of the local product and the induced local product space are introduced and developed. This space reveals itself to be a unique form of complicated inner product space. The following inequality is obtained by utilizing this space.

**Theorem 1.1.** *Let  $\vec{a} = (a_1, a_2, \dots, a_n), \vec{b} = (b_1, b_2, \dots, b_n) \in \mathbb{R}^n$  such that  $0 < \langle \vec{a}, \vec{b} \rangle$  and  $\langle \vec{a}, \vec{b} \rangle \neq 1$ , then the lower bound holds*

$$\begin{aligned} & \left| \int_{|a_n|}^{|b_n|} \int_{|a_{n-1}|}^{|b_{n-1}|} \cdots \int_{|a_1|}^{|b_1|} \sqrt[4s]{\sum_{i=1}^n x_i^{4s}} dx_1 dx_2 \cdots dx_n \right| \\ & \leq \frac{|\langle \vec{a}, \vec{b} \rangle|}{2\pi |\log(\langle \vec{a}, \vec{b} \rangle)|} \times (||\vec{a}||^{4s+1} + ||\vec{b}||^{4s+1}) \times \left| \prod_{i=1}^n |b_i| - |a_i| \right|. \end{aligned}$$

for all  $s \in \mathbb{N}$ , where  $\langle, \rangle$  denotes the inner product.

**Theorem 1.2.** *Let  $\vec{a} = (a_1, a_2, \dots, a_n), \vec{b} = (b_1, b_2, \dots, b_n) \in \mathbb{R}^n$  with  $a_j, b_j > 0$  and  $\langle, \rangle$  denotes the inner product such that  $0 < \langle a, b \rangle \leq e$  with  $\langle a, b \rangle \neq 1$ , then we have*

$$\int_{|a_n|}^{|b_n|} \int_{|a_{n-1}|}^{|b_{n-1}|} \cdots \int_{|a_1|}^{|b_1|} \frac{1}{\sqrt[4s+3]{\sum_{j=1}^n x_j^{4s+3}}} dx_1 dx_2 \cdots dx_n \geq \frac{2\pi \times |\log(\langle a, b \rangle)| \left| \prod_{j=1}^n |b_j| - |a_j| \right|}{||\vec{a}||^{4s+4} + ||\vec{b}||^{4s+4}}.$$

*Date:* December 2, 2022.

*2010 Mathematics Subject Classification.* Primary 11Pxx, 11Bxx; Secondary 11Axx, 11Gxx.

*Key words and phrases.* Local product; local product space; sheet.

The concept of the local product and associated space is often thought of as a black box for quickly establishing a huge class of mathematical inequalities that are difficult to prove using traditional mathematical methods. It operates by traveling into the space and selecting appropriate sheets as functions that are present in the anticipated inequality, as well as satisfying some local requirements with the appropriate support. The local product and associated space could be useful for more than just demonstrating complex mathematical inequalities. As a bi-variate map that assigns any two vectors in a complex inner product space to a complex number, they could be fascinating in and of themselves. It's a unique subspace in many ways. The  $k^{\text{th}}$  local product space over a sheet  $f : \mathbb{C} \rightarrow \mathbb{C}$  is an inner product space equipped with the local product  $\mathcal{G}_f^k(\cdot)$  over a fixed sheet.

## 2. The local product and associated space

In this note we introduce and study the notion of the **local product** and associated space.

**Definition 2.1.** Let  $\vec{a}, \vec{b} \in \mathbb{C}^n$  and  $f : \mathbb{C} \rightarrow \mathbb{C}$  be continuous on  $\cup_{j=1}^n [|a_j|, |b_j|]$ . Let  $(\mathbb{C}^n, \langle, \rangle)$  be a complex inner product space. Then by the  $k^{\text{th}}$  local product of  $\vec{a}$  with  $\vec{b}$  on the sheet  $f$ , we mean the bi-variate map  $\mathcal{G}_f^k : (\mathbb{C}^n, \langle, \rangle) \times (\mathbb{C}^n, \langle, \rangle) \rightarrow \mathbb{C}$  such that

$$\mathcal{G}_f^k(\vec{a}; \vec{b}) = f(\langle \vec{a}, \vec{b} \rangle) \int_{|a_n|}^{|b_n|} \int_{|a_{n-1}|}^{|b_{n-1}|} \cdots \int_{|a_1|}^{|b_1|} f \circ \mathbf{e} \left( (i)^k \frac{\sqrt[k]{\sum_{j=1}^n x_j^k}}{\|\vec{a}\|^{k+1} + \|\vec{b}\|^{k+1}} \right) dx_1 dx_2 \cdots dx_n$$

where  $\langle, \rangle$  denotes the inner product and where  $\mathbf{e}(q) = e^{2\pi i q}$ . We denote an inner product space with a  $k^{\text{th}}$  **local product** defined over a sheet  $f$  as the  $k^{\text{th}}$  local product space over a sheet  $f$ . We denote this space with the triple  $(\mathbb{C}^n, \langle, \rangle, \mathcal{G}_f^k(\cdot))$ .

The  $k^{\text{th}}$  local product is in some sense a universal map induced by a sheet. In other words a local product can be constructed by carefully choosing the sheet. By taking our sheet to be the constant function  $f := 1$  we obtain the local product

$$\begin{aligned} \mathcal{G}_1^k(\vec{a}; \vec{b}) &= \int_{|a_n|}^{|b_n|} \int_{|a_{n-1}|}^{|b_{n-1}|} \cdots \int_{|a_1|}^{|b_1|} dx_1 dx_2 \cdots dx_n \\ &= \prod_{i=1}^n |b_i| - |a_i|. \end{aligned}$$

Similarly, if we take our sheet to be  $f = \log$ , then under the condition that  $\langle \vec{a}, \vec{b} \rangle \neq 0$ , we obtain the induced local product

$$\mathcal{G}_{\log}^k(\vec{a}; \vec{b}) = 2\pi \times (i)^{k+1} \frac{\log(\langle \vec{a}, \vec{b} \rangle)}{\|\vec{a}\|^{k+1} + \|\vec{b}\|^{k+1}} \int_{|a_n|}^{|b_n|} \int_{|a_{n-1}|}^{|b_{n-1}|} \cdots \int_{|a_1|}^{|b_1|} \sqrt[k]{\sum_{j=1}^n x_j^k} dx_1 dx_2 \cdots dx_n.$$

By taking the sheet  $f = \text{Id}$  to be the identity function, then we obtain in this setting the associated local product

$$\mathcal{G}_{\text{Id}}^k(\vec{a}; \vec{b}) = \langle \vec{a}, \vec{b} \rangle \int_{|a_n|}^{|b_n|} \int_{|a_{n-1}|}^{|b_{n-1}|} \cdots \int_{|a_1|}^{|b_1|} \mathbf{e} \left( \frac{(i)^k \sqrt[k]{\sum_{j=1}^n x_j^k}}{\|\vec{a}\|^{k+1} + \|\vec{b}\|^{k+1}} \right) dx_1 dx_2 \cdots dx_n.$$

Again, by taking the sheet  $f = \text{Id}^{-1}$  with  $\langle a, b \rangle \neq 0$ , then we obtain the corresponding induced  $k^{\text{th}}$  local product

$$\mathcal{G}_{\text{Id}^{-1}}^k(\vec{a}; \vec{b}) = \frac{1}{\langle \vec{a}, \vec{b} \rangle} \int_{|a_n|}^{|b_n|} \int_{|a_{n-1}|}^{|b_{n-1}|} \cdots \int_{|a_1|}^{|b_1|} \mathbf{e} \left( - \frac{(i)^k \sqrt[k]{\sum_{j=1}^n x_j^k}}{\|\vec{a}\|^{k+1} + \|\vec{b}\|^{k+1}} \right) dx_1 dx_2 \cdots dx_n.$$

### 3. Properties of the local product

In this section we study some properties of the local product on a fixed sheet.

**Proposition 3.1.** *The following holds*

(i) *If  $f$  is linear such that  $\langle a, b \rangle = -\langle b, a \rangle$  then*

$$\mathcal{G}_f^k(\vec{a}; \vec{b}) = (-1)^{n+1} \mathcal{G}_f^k(\vec{b}; \vec{a}).$$

(ii) *Let  $f, g : \mathbb{R} \rightarrow \mathbb{R}^+$  such that  $f(t) \leq g(t)$  for any  $t \in [1, \infty)$ . Then  $|\mathcal{G}_f^k(\vec{a}; \vec{b})| \leq |\mathcal{G}_g^k(\vec{a}; \vec{b})|$ .*

*Proof.* (i) By the linearity of  $f$ , we can write

$$\begin{aligned} \mathcal{G}_f^k(\vec{a}; \vec{b}) &= f(\langle \vec{a}, \vec{b} \rangle) \int_{|a_n|}^{|b_n|} \int_{|a_{n-1}|}^{|b_{n-1}|} \cdots \int_{|a_1|}^{|b_1|} f \circ \mathbf{e} \left( (i)^k \frac{\sqrt[k]{\sum_{j=1}^n x_j^k}}{\|\vec{a}\|^{k+1} + \|\vec{b}\|^{k+1}} \right) dx_1 dx_2 \cdots dx_n \\ &= f(\langle \vec{a}, \vec{b} \rangle) \int_{|a_n|}^{|b_n|} \int_{|a_{n-1}|}^{|b_{n-1}|} \cdots \int_{|a_1|}^{|b_1|} f \circ \mathbf{e} \left( (i)^k \frac{\sqrt[k]{\sum_{j=1}^n x_j^k}}{\|\vec{a}\|^{k+1} + \|\vec{b}\|^{k+1}} \right) dx_1 dx_2 \cdots dx_n \\ &= f(-\langle b, a \rangle) (-1)^n \int_{|b_n|}^{|a_n|} \int_{|b_{n-1}|}^{|a_{n-1}|} \cdots \int_{|b_1|}^{|a_1|} f \circ \mathbf{e} \left( (i)^k \frac{\sqrt[k]{\sum_{j=1}^n x_j^k}}{\|\vec{a}\|^{k+1} + \|\vec{b}\|^{k+1}} \right) dx_1 dx_2 \cdots dx_n \\ &= (-1)^{n+1} f(\langle b, a \rangle) \int_{|b_n|}^{|a_n|} \int_{|b_{n-1}|}^{|a_{n-1}|} \cdots \int_{|b_1|}^{|a_1|} f \circ \mathbf{e} \left( (i)^k \frac{\sqrt[k]{\sum_{j=1}^n x_j^k}}{\|\vec{a}\|^{k+1} + \|\vec{b}\|^{k+1}} \right) dx_1 dx_2 \cdots dx_n \\ &= (-1)^{n+1} \mathcal{G}_f^k(\vec{b}; \vec{a}). \end{aligned}$$

(ii) Property (ii) follows very easily from the inequality  $f(t) \leq g(t)$ .  $\square$

#### 4. Applications of the local product

In this section we explore some applications of the local product.

**Theorem 4.1.** *Let  $\vec{a} = (a_1, a_2, \dots, a_n), \vec{b} = (b_1, b_2, \dots, b_n) \in \mathbb{R}^n$  such that  $0 < \langle \vec{a}, \vec{b} \rangle$  and  $\langle \vec{a}, \vec{b} \rangle \neq 1$ , then the lower bound holds*

$$\begin{aligned} & \left| \int_{|a_n|}^{|b_n|} \int_{|a_{n-1}|}^{|b_{n-1}|} \cdots \int_{|a_1|}^{|b_1|} \sqrt[4s]{\sum_{i=1}^n x_i^{4s}} dx_1 dx_2 \cdots dx_n \right| \\ & \leq \frac{|\langle \vec{a}, \vec{b} \rangle|}{2\pi |\log(\langle \vec{a}, \vec{b} \rangle)|} \times (||\vec{a}||^{4s+1} + ||\vec{b}||^{4s+1}) \times \left| \prod_{i=1}^n |b_i| - |a_i| \right|. \end{aligned}$$

for all  $s \in \mathbb{N}$ , where  $\langle \cdot, \cdot \rangle$  denotes the inner product.

*Proof.* Let  $f : \mathbb{R} \rightarrow \mathbb{R}^+$  and  $\vec{a}, \vec{b} \in \mathbb{R}^n$  such that  $0 < \langle \vec{a}, \vec{b} \rangle$  and  $\langle \vec{a}, \vec{b} \rangle \neq 1$ . We note that

$$\mathcal{G}_{\log}^{4s}(\vec{a}; \vec{b}) = 2\pi \times (i)^{4s+1} \frac{\log(\langle \vec{a}, \vec{b} \rangle)}{||\vec{a}||^{4s+1} + ||\vec{b}||^{4s+1}} \int_{|a_n|}^{|b_n|} \int_{|a_{n-1}|}^{|b_{n-1}|} \cdots \int_{|a_1|}^{|b_1|} \sqrt[4s]{\sum_{j=1}^n x_j^{4s}} dx_1 dx_2 \cdots dx_n$$

by taking  $k = 4s$  for any  $s \in \mathbb{N}$ . Also by taking the sheet  $f := |\cdot|$  to be the distance function, then we obtain in this setting the associated local product

$$\begin{aligned} \mathcal{G}_{|\cdot|}^{4s}(\vec{a}; \vec{b}) &= |\langle \vec{a}, \vec{b} \rangle| \int_{|a_n|}^{|b_n|} \int_{|a_{n-1}|}^{|b_{n-1}|} \cdots \int_{|a_1|}^{|b_1|} dx_1 dx_2 \cdots dx_n \\ &= |\langle \vec{a}, \vec{b} \rangle| \times \left| \prod_{i=1}^n |b_i| - |a_i| \right|. \end{aligned}$$

Since  $\log < |\cdot|$  on  $(1, \infty)$  the claim inequality is a consequence by appealing to Proposition 3.1.  $\square$

**Theorem 4.2.** *Let  $\vec{a} = (a_1, a_2, \dots, a_n), \vec{b} = (b_1, b_2, \dots, b_n) \in \mathbb{R}^n$  with  $a_j, b_j > 0$  and  $\langle \cdot, \cdot \rangle$  denotes the inner product such that  $0 < \langle a, b \rangle \leq e$  with  $\langle a, b \rangle \neq 1$ , then we have*

$$\left| \int_{|a_n|}^{|b_n|} \int_{|a_{n-1}|}^{|b_{n-1}|} \cdots \int_{|a_1|}^{|b_1|} \frac{1}{\sqrt[4s+3]{\sum_{j=1}^n x_j^{4s+3}}} dx_1 dx_2 \cdots dx_n \right| \geq \frac{2\pi \times |\log(\langle a, b \rangle)| \left| \prod_{j=1}^n |b_j| - |a_j| \right|}{||\vec{a}||^{4s+4} + ||\vec{b}||^{4s+4}}.$$

*Proof.* Let  $f : \mathbb{R} \rightarrow \mathbb{R}^+$  and  $\vec{a}, \vec{b} \in \mathbb{R}^n$ . We note that

$$\mathcal{G}_{\frac{1}{\log}}^k(\vec{a}; \vec{b}) = \frac{1}{\log(\langle a, b \rangle)} \times (||\vec{a}||^{k+1} + ||\vec{b}||^{k+1}) \times \frac{1}{2j^{k+1}\pi} \times \int_{|a_n|}^{|b_n|} \int_{|a_{n-1}|}^{|b_{n-1}|} \cdots \int_{|a_1|}^{|b_1|} \frac{1}{\sqrt[k]{\sum_{j=1}^n x_j^k}} dx_1 dx_2 \cdots dx_n$$

by taking  $k = 4s + 3$  for any  $s \in \mathbb{N}$ . Also by taking the sheet  $f := 1$  to be the constant function, then we obtain in this setting the associated local product

$$\begin{aligned} \mathcal{G}_1^{4s}(\vec{a}; \vec{b}) &= \int_{|a_n|}^{|b_n|} \int_{|a_{n-1}|}^{|b_{n-1}|} \cdots \int_{|a_1|}^{|b_1|} dx_1 dx_2 \cdots dx_n \\ &= \prod_{i=1}^n |b_i| - |a_i|. \end{aligned}$$

Since  $\frac{1}{\log} \geq 1$  on  $(1, e)$  the claim inequality is a consequence by appealing to Proposition 3.1 and the requirement  $0 < \langle a, b \rangle \leq e$  with  $\langle a, b \rangle \neq 1$ .  $\square$

---

#### REFERENCES

1. Rudin, W. *Real and complex analysis*, Tata McGraw-hill education, 2006.
2. Agama, T. *The Outer product and Outer product space*, Operators and Matrices, vol. 15:4, 2021, pp 1575-1580.

DEPARTMENT OF MATHEMATICS, AFRICAN INSTITUTE FOR MATHEMATICAL SCIENCES, GHANA.  
*Email address:* Theophilus@aims.edu.gh/emperordagama@yahoo.com