

# Existence and Uniqueness of Constraint Minimizers for the Planar Schrödinger-Poisson System with Logarithmic Potentials

Yujin Guo\*, Wenning Liang† and Yan Li‡

## Abstract

In this paper, we study constraint minimizers  $u$  of the planar Schrödinger-Poisson system with a logarithmic convolution potential  $\ln|x|*u^2$  and a logarithmic external potential  $V(x) = \ln(1+|x|^2)$ , which can be described by the  $L^2$ -critical constraint minimization problem with a subcritical perturbation. We prove that there is a threshold  $\rho^* \in (0, \infty)$  such that constraint minimizers exist if and only if  $0 < \rho < \rho^*$ . In particular, the local uniqueness of positive constraint minimizers as  $\rho \nearrow \rho^*$  is analyzed by overcoming the sign-changing property of the logarithmic convolution potential and the non-invariance under translations of the logarithmic external potential.

*Keywords:* Schrödinger-Poisson system; constraint minimizers; logarithmic convolution potential; logarithmic external potential; local uniqueness

## 1 Introduction

In this paper, we consider the following Schrödinger-Poisson system

$$\begin{cases} i\psi_t - \Delta\psi + V(x)\psi + \gamma w\psi = |\psi|^{p-2}\psi & \text{in } \mathbb{R}^N \times \mathbb{R}, \\ \Delta w = |\psi|^2 & \text{in } \mathbb{R}^N \times \mathbb{R}, \end{cases} \quad (1.1)$$

where  $\psi : \mathbb{R}^N \times \mathbb{R} \rightarrow \mathbb{C}$  ( $N \geq 2$ ) is the time-dependent wave function,  $x \mapsto V(x)$  is a real external potential,  $p > 2$  and  $\gamma \in \mathbb{R}$  is a parameter. The function  $w$  represents an internal potential for a nonlocal self-interaction of the wave function  $\psi$ , and the nonlinear term  $|\psi|^{p-2}\psi$  is frequently used to model the interaction among particles [8, 19, 51]. Over the past few decades, the system (1.1) has attracted considerable attentions due to its physical relevance: for instance, it arises in quantum mechanics (see [3, 7–10, 13, 25, 26, 42, 44, 48–50]) and particularly semiconductor physics (see [12, 45, 54]). We refer the reader to [2, 22, 24, 36] for more details on its physical aspects.

Under the standing wave ansatz  $\psi(x, t) = e^{i\mu t}u(x)$ ,  $\mu \in \mathbb{R}$ , the Schrödinger-Poisson system (1.1) is reduced to the following problem

$$-\Delta u + (V(x) - \mu)u + \gamma[\Phi_N * u^2]u = |u|^{p-2}u \quad \text{in } \mathbb{R}^N, \quad (1.2)$$

where

$$\Phi_N(x) := \begin{cases} \frac{1}{N(2-N)\omega_N}|x|^{2-N}, & N \geq 3, \\ \frac{1}{2\pi} \ln|x|, & N = 2, \end{cases} \quad (1.3)$$

\*School of Mathematics and Statistics, and Hubei Key Laboratory of Mathematical Sciences, Central China Normal University, P.O. Box 71010, Wuhan 430079, P. R. China. Email: yguo@ccnu.edu.cn. Y. J. Guo is partially supported by NSFC under Grants 12225106 and 11931012.

†School of Mathematics and Statistics, Central China Normal University, P.O. Box 71010, Wuhan 430079, P. R. China. Email: wnliliang@ccnu.edu.cn.

‡School of Mathematics and Statistics, Central China Normal University, P.O. Box 71010, Wuhan 430079, P. R. China. Email: yanlimath@mails.ccnucn.edu.cn.

and  $\omega_N > 0$  denotes the volume of the unit ball in  $\mathbb{R}^N$ . The case  $N \geq 3$  of (1.2) was extensively studied in the past few years. In particular, variational methods were employed (cf. [1, 5, 18, 46, 52, 53]) to analyze the existence and multiplicity results of solutions for (1.2). We also refer the reader to [6, 14, 37, 38, 41, 44] for analyzing normalized solutions  $u$  of (1.2), in the sense that  $u$  satisfies  $\|u\|_2^2 = \rho$  for some given  $\rho > 0$ .

However, the approaches mentioned above for the case  $N \geq 3$  cannot be directly adapted to the case  $N = 2$ , due to the fact that the Newtonian potential  $\Phi_2(x)$  defined in (1.3) is sign-changing and presents singularities as  $|x|$  approaches to zero and infinity. Therefore, the existing works [4, 16, 17, 23, 40, 55] show that it is more delicate to study (1.2) for the case  $N = 2$ . More precisely, Cingolani and Weth [17] derived a strong compactness condition for Cerami sequences, based on which they established various existence results of solutions for (1.2) under the assumptions  $V(x) \in L^\infty(\mathbb{R}^2)$  and  $p \geq 4$ . Further, Du and Weth [23] generalized the above results to the rest case  $p \in (2, 4)$  by exploring a more complicated underlying functional geometry with a different variational method. On the other hand, by analyzing normalized solutions, Stubbe [55] studied the existence and uniqueness of ground states for (1.2) with  $N = 2$ , where the external potential  $V(x)$  and the nonlinear term were both ignored. After Stubbe's work, Cingolani and Jeanjean [16] obtained several existence and multiplicity results of normalized solutions for (1.2) with  $N = 2$ , where  $V(x) \equiv 0$  and  $p \in (2, 4)$  were considered, see also [40] for related works. Moreover, when  $N = 2$ , the problem (1.2) with a different nonlinear term was recently analyzed in [21] and the references therein, where the authors considered the following logarithmic external potential

$$V(x) = \ln(1 + |x|^2) \quad \text{in } \mathbb{R}^2. \quad (1.4)$$

Motivated by above facts, in this paper we focus on normalized solutions  $u$  of (1.2) with the external potential  $V(x)$  satisfying (1.4), together with  $N = 2$ ,  $p = 4$  and  $\gamma = 2\pi$ , which can be obtained as (constraint) minimizers of the following variational problem:

$$e(\rho) := \inf_{u \in S_\rho} E(u), \quad \rho > 0, \quad (1.5)$$

where the energy functional  $E(u)$  satisfies

$$\begin{aligned} E(u) := & \frac{1}{2} \int_{\mathbb{R}^2} \left[ |\nabla u(x)|^2 + \ln(1 + |x|^2) u^2(x) \right] dx \\ & + \frac{1}{4} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x - y| u^2(x) u^2(y) dx dy - \frac{1}{4} \int_{\mathbb{R}^2} u^4(x) dx. \end{aligned} \quad (1.6)$$

We notice that  $E(u)$  involves both a logarithmic potential term and a logarithmic convolution term, which unfortunately are not well defined in  $H^1(\mathbb{R}^2)$ . Stimulated by [17, 55], we consider the space  $S_\rho$  of  $e(\rho)$  satisfying

$$S_\rho := \left\{ u \in X : \|u\|_2^2 = \rho \right\}, \quad X = \left\{ u \in H^1(\mathbb{R}^2) : \int_{\mathbb{R}^2} \ln(1 + |x|^2) u^2(x) dx < \infty \right\},$$

which is endowed with the norm

$$\|u\|_X = \left\{ \int_{\mathbb{R}^2} \left[ |\nabla u|^2 + \left(1 + \ln(1 + |x|^2)\right) u^2(x) \right] dx \right\}^{\frac{1}{2}}, \quad u \in X.$$

One can note that the norm of  $X$  and the functional  $E(\cdot)$  are not invariant under translations in  $\mathbb{R}^2$ , and the quadratic part of the energy functional (1.6) is not coercive on  $X$ . All these lead to some extra difficulties in the analysis of (1.5), compared with the existing works [32–34] and the references therein. By analyzing carefully two different types of logarithmic terms involved in  $E(\cdot)$ , the main purpose of this paper is to discuss the existence and local uniqueness of minimizers for  $e(\rho)$ .

If the logarithmic external potential  $V(x) = \ln(1 + |x|^2)$  of (1.2) is ignored, it was proved in [16, 40] that  $e(\rho)$  admits minimizers if and only if  $0 < \rho < \rho^*$ , where  $\rho^* := \|Q\|_2^2$ , and  $Q = Q(|x|) > 0$ , up to translations, is the unique (cf. [27, 39]) positive solution of the following elliptic equation:

$$-\Delta u + u - u^3 = 0 \quad \text{in } \mathbb{R}^2, \quad u \in H^1(\mathbb{R}^2). \quad (1.7)$$

In addition, the function  $Q > 0$  decays exponentially as  $|x| \rightarrow \infty$ , in the sense that

$$Q(x), \quad |\nabla Q(x)| = O\left(|x|^{-\frac{1}{2}} e^{-|x|}\right) \quad \text{as } |x| \rightarrow \infty. \quad (1.8)$$

Moreover, we have the following classical Gagliardo-Nirenberg inequality:

$$\int_{\mathbb{R}^2} u^4 dx \leq \frac{2}{\|Q\|_2^2} \int_{\mathbb{R}^2} |\nabla u|^2 dx \int_{\mathbb{R}^2} u^2 dx, \quad u \in H^1(\mathbb{R}^2), \quad (1.9)$$

where the identity is achieved at  $u = Q$ , and  $Q = Q(x) > 0$  is the unique positive solution of (1.7). The following theorem shows that the existence and non-existence of minimizers for  $e(\rho)$  are the same as those of the case addressed in [16, 40], where the external potential  $V(x) = \ln(1 + |x|^2)$  is ignored.

**Theorem 1.1.** *Let  $Q = Q(|x|)$  be the unique positive solution of (1.7). We have*

1. *If  $0 < \rho < \rho^* := \|Q\|_2^2$ , then there exists at least one minimizer of  $e(\rho)$ ;*
2. *If  $\rho \geq \rho^*$ , then there is no minimizer of  $e(\rho)$ .*

Moreover, we have  $\lim_{\rho \nearrow \rho^*} e(\rho) = -\infty$ .

The proof of Theorem 1.1 is similar to [16, 32, 40], which is given in Section 2 for completion. Our analysis shows that the proof of Theorem 1.1 needs the Gagliardo-Nirenberg inequality (1.9) and the following Hardy-Littlewood-Sobolev inequality (cf. [43]):

$$\int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{1}{|x-y|} |u(x)||v(y)| dx dy \leq C \|u\|_{\frac{4}{3}} \|v\|_{\frac{4}{3}}, \quad \text{where } u, v \in L^{\frac{4}{3}}(\mathbb{R}^2). \quad (1.10)$$

Actually, the inequality (1.10) is often used throughout this paper to handle with the logarithmic convolution term of  $E(u)$ .

Suppose  $u_\rho$  is a minimizer of  $e(\rho)$  for any  $\rho \in (0, \rho^*)$ , then the variational theory gives that  $u_\rho$  satisfies the following Euler-Lagrange equation:

$$-\Delta u_\rho + \ln(1 + |x|^2)u_\rho + \left( \int_{\mathbb{R}^2} \ln|x-y| u_\rho^2(y) dy \right) u_\rho = \mu_\rho u_\rho + |u_\rho|^2 u_\rho \quad \text{in } \mathbb{R}^2, \quad (1.11)$$

where  $\mu_\rho \in \mathbb{R}$  is the associated Lagrange multiplier. We obtain from [43, Theorem 6.17] that  $E(u_\rho) \geq E(|u_\rho|)$ , which implies that  $|u_\rho|$  is also a minimizer of  $e(\rho)$ . By the strong maximum principle, it further follows from (1.11) that  $|u_\rho| > 0$  in  $\mathbb{R}^2$ . Therefore,  $u_\rho$  must be either strictly positive or strictly negative. Without loss of generality, we next focus on positive minimizers  $u_\rho > 0$  of  $e(\rho)$ .

The uniqueness of positive minimizers for  $e(\rho)$  is an interesting problem in physics. Stimulated by Refs. [29, 30] and the references therein, the main result of this paper is concerned with the following local uniqueness of positive minimizers for  $e(\rho)$ .

**Theorem 1.2.** *There exists a unique positive minimizer of  $e(\rho)$  as  $\rho$  is sufficiently close to  $\rho^*$  from below.*

It is our conjecture that  $e(\rho)$  admits a unique positive minimizer for all  $\rho \in (0, \rho^*)$ , which however seems a more challenging problem. We now sketch the main idea of proving Theorem 1.2 as follows: To prove Theorem 1.2, by contradiction we assume that there exist two different minimizers  $u_\rho \not\equiv v_\rho$ . We then define the following difference function:

$$\eta_\rho(x) := \frac{u_\rho - v_\rho}{\|u_\rho - v_\rho\|_{L^\infty(\mathbb{R}^2)}} \quad \text{in } \mathbb{R}^2.$$

On one hand, since  $\|\eta_\rho(x)\|_{L^\infty(\mathbb{R}^2)} \equiv 1$ , we shall show that  $\eta_\rho(x)$  cannot vanish as  $\rho \nearrow \rho^*$ . On the other hand, we shall prove that under suitable scaling,  $\eta_\rho(x)$  converges to zero in  $L^\infty(\mathbb{R}^2)$  as  $\rho \nearrow \rho^*$ , a contradiction, by constructing various types of Pohozaev identities (cf. [11, 20, 29, 30]). We comment that our argument is different from that of [55], where the uniqueness of minimizers was essentially obtained by analyzing the corresponding elliptic system.

However, the above procedure depends on the refined estimates of  $\eta_\rho(x)$  as  $\rho \nearrow \rho^*$ , for which it is necessary to establish the following limiting behavior of positive minimizers for  $e(\rho)$  as  $\rho \nearrow \rho^*$ .

**Theorem 1.3.** *Assume  $u_\rho$  is a positive minimizer of  $e(\rho)$  for  $\rho \in (0, \rho^*)$ , then we have*

$$\lim_{\rho \nearrow \rho^*} \frac{2(\rho^* - \rho)^{\frac{1}{2}}}{\rho^*} u_\rho \left( \frac{2(\rho^* - \rho)^{\frac{1}{2}}}{\rho^*} x + x_\rho \right) = Q(x) \quad \text{in } X \cap L^\infty(\mathbb{R}^2),$$

where  $x_\rho$  is the unique maximal point of  $u_\rho$  as  $\rho \nearrow \rho^*$  and satisfies

$$\lim_{\rho \nearrow \rho^*} \frac{x_\rho}{(\rho^* - \rho)^{\frac{1}{2}}} = 0.$$

We remark that even though the limiting behavior of ground states for Bose-Einstein condensates was widely studied recently in [32–34] and the references therein, as far as we know, there exist almost no similar results for the planar Schrödinger-Poisson system. Moreover, comparing with the existing works mentioned above, there exist some extra difficulties in the proof of Theorem 1.3. Firstly, the energy functional of  $e(\rho)$  is involved with the challenging logarithmic convolution term

$$\int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x - y| u^2(x) u^2(y) dx dy. \quad (1.12)$$

To overcome the difficulties arising from (1.12), we shall derive in (3.12) the following important estimate: for any given  $x \in \mathbb{R}^2$ , there exists a constant  $C > 0$  such that for any  $\rho \in (0, \rho^*)$ ,

$$\int_{\mathbb{R}^2} \ln \left( 1 + \frac{1}{|x - y|^2} \right) w_\rho^2(y) dy \leq C,$$

where  $w_\rho > 0$  is a suitably scaled function of the minimizer  $u_\rho$ . Secondly, since the norm of  $X$  is not invariant under translations, different from [32–34], we cannot directly obtain the uniform boundedness of the scaled minimizers  $w_\rho$  as  $\rho \nearrow \rho^*$ . To overcome this difficulty, we shall borrow the idea from [31]. Finally, because  $\lim_{\rho \nearrow \rho^*} e(\rho) = -\infty$  and the external potential term  $\int_{\mathbb{R}^2} \ln(1 + |x|^2) u_\rho^2 dx$  is not invariant under translations, in Section 3 we need to investigate new techniques to derive the refined information of the maximal point  $x_\rho$  as  $\rho \nearrow \rho^*$ .

This paper is organized as follows. In Section 2, we first give some preliminary results, after which we then prove Theorem 1.1 on the existence and non-existence of minimizers for  $e(\rho)$ . In Section 3, we shall prove Theorem 1.3 on the refined limiting profiles of positive minimizers for  $e(\rho)$  as  $\rho \nearrow \rho^*$ . Theorem 1.2 is then proved in Section 4 for the local uniqueness of positive minimizers. We finally give in Appendix A the detailed proofs of some results used in the proof of Theorems 1.2 and 1.3.

## 2 Preliminaries

In this section, we first establish various preliminary results which are often used throughout the whole paper. We then follow them to finish the proof of Theorem 1.1 in Subsection 2.1.

As performed in [17], we first define the following symmetric bilinear forms

$$B_1(u, v) = \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln(1 + |x - y|^2) u(x) v(y) dx dy,$$

$$B_2(u, v) = \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln \left( 1 + \frac{1}{|x - y|^2} \right) u(x) v(y) dx dy - \frac{1}{2} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln \left( 1 + \frac{1}{|x - y|^2} \right) u(x) v(y) dx dy,$$

and

$$B_0(u, v) = \frac{1}{2} [B_1(u, v) - B_2(u, v)] = \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x - y| u(x) v(y) dx dy, \quad u, v \in X.$$

Since

$$\ln(1 + |x - y|^2) \leq \ln(1 + 2|x|^2 + 2|y|^2) \leq 2\ln(1 + |x|^2) + 2\ln(1 + |y|^2), \quad x, y \in \mathbb{R}^2,$$

we have the estimate

$$\begin{aligned} |B_1(uv, wz)| &\leq 2 \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} [\ln(1 + |x|^2) + \ln(1 + |y|^2)] |u(x)v(x)w(y)z(y)| dx dy \\ &\leq 2\|u\|_* \|v\|_* \|w\|_2 \|z\|_2 + 2\|u\|_2 \|v\|_2 \|w\|_* \|z\|_*, \quad u, v, w, z \in X, \end{aligned} \quad (2.1)$$

where

$$\|u\|_* := \left\{ \int_{\mathbb{R}^2} \ln(1 + |x|^2) u^2(x) dx \right\}^{\frac{1}{2}}, \quad u \in X.$$

Since  $0 < \ln(1 + r) < r$  holds for all  $r > 0$ , it follows from the Hardy-Littlewood-Sobolev inequality (1.10) that there exists a constant  $C > 0$  such that

$$\begin{aligned} \frac{1}{2} |B_2(u, v)| &\leq \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln\left(1 + \frac{1}{|x-y|}\right) |u(x)v(y)| dx dy \\ &\leq \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{1}{|x-y|} |u(x)v(y)| dx dy \\ &\leq C \|u\|_{\frac{4}{3}} \|v\|_{\frac{4}{3}}, \quad u, v \in L^{\frac{4}{3}}(\mathbb{R}^2). \end{aligned} \quad (2.2)$$

By the continuous embedding from  $X$  into  $L^q(\mathbb{R}^2)$  for  $q \in [2, \infty)$ , we derive from (2.1) and (2.2) that the functionals  $B_i(u^2, v^2)$  are well defined on  $X \times X$  for  $i = 0, 1, 2$ . Moreover, denote functionals on  $X$  as follows:

$$\begin{aligned} V_1(u) &= B_1(u^2, u^2) = \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln(1 + |x-y|^2) u^2(x) u^2(y) dx dy, \\ V_2(u) &= B_2(u^2, u^2) = \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln\left(1 + \frac{1}{|x-y|^2}\right) u^2(x) u^2(y) dx dy, \end{aligned}$$

and

$$V_0(u) = B_0(u^2, u^2) = \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| u^2(x) u^2(y) dx dy.$$

Using above notations, we have the following crucial properties of  $V_i(\cdot)$  for  $i = 0, 1, 2$ .

**Lemma 2.1.** *The space  $X$  and the functionals  $V_i(\cdot)$  ( $i = 0, 1, 2$ ) have the following properties:*

1. *The space  $X$  is compactly embedded into  $L^s(\mathbb{R}^2)$  for all  $s \in [2, \infty)$ ;*
2. *The functionals  $V_0(\cdot), V_1(\cdot), V_2(\cdot)$  and  $E(\cdot)$  defined in (1.6) are of class  $C^1$  on  $X$ , and*

$$\langle V'_i(u), v \rangle = 4B_i(u^2, uv) \quad \text{for } u, v \in X, \quad i = 0, 1, 2;$$

3.  *$V_2(\cdot)$  is continuously differentiable on  $L^{\frac{8}{3}}(\mathbb{R}^2)$ ;*
4.  *$V_1(\cdot)$  is weakly lower semicontinuous on  $H^1(\mathbb{R}^2)$ ;*
5.  *$V_0(\cdot)$  is weakly lower semicontinuous on  $X$ .*

Since the proof of Lemma 2.1 is similar to that of [17, Lemma 2.2], we omit the details for simplicity.

## 2.1 The existence and non-existence of minimizers for $e(\rho)$

Applying above preliminary results, the main purpose of this subsection is to address the proof of Theorem 1.1 on the existence and non-existence of minimizers for  $e(\rho)$ . For the unique positive solution  $Q = Q(|x|) > 0$  of (1.7), we note from [15, Lemma 8.12] that  $Q$  satisfies the following identities

$$\int_{\mathbb{R}^2} |\nabla Q|^2 dx = \int_{\mathbb{R}^2} Q^2 dx = \frac{1}{2} \int_{\mathbb{R}^2} Q^4 dx. \quad (2.3)$$

**Proof of Theorem 1.1.** 1. We first prove the existence of minimizers for  $e(\rho)$  with  $0 < \rho < \rho^*$ . If  $u \in X$  and  $\|u\|_2^2 = \rho$  with  $\rho \in (0, \rho^*)$ , then we deduce from the Gagliardo-Nirenberg inequality ([56]) that for any  $p \geq 2$ ,

$$\|u\|_p \leq \left( \frac{p\rho}{2\|Q\|_2^{p-2}} \right)^{\frac{1}{p}} \left( \int_{\mathbb{R}^2} |\nabla u|^2 dx \right)^{\frac{p-2}{2p}}.$$

By Hardy-Littlewood-Sobolev inequality (1.10), we derive from above that there exists a constant  $C > 0$  such that

$$\int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{u^2(x)u^2(y)}{|x-y|} dx dy \leq C \left( \int_{\mathbb{R}^2} |u|^{\frac{8}{3}} dx \right)^{\frac{3}{2}} \leq C \rho^{\frac{3}{2}} \left( \int_{\mathbb{R}^2} |\nabla u|^2 dx \right)^{\frac{1}{2}}. \quad (2.4)$$

It then follows from (2.2) and (2.4) that

$$\begin{aligned} & \frac{1}{2} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln \left( 1 + \frac{1}{|x-y|^2} \right) u^2(x)u^2(y) dx dy \\ & \leq \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{1}{|x-y|} u^2(x)u^2(y) dx dy \\ & \leq C \rho^{\frac{3}{2}} \left( \int_{\mathbb{R}^2} |\nabla u|^2 dx \right)^{\frac{1}{2}}. \end{aligned} \quad (2.5)$$

Together with (1.9), we get from (2.5) that

$$E(u) \geq \frac{\rho^* - \rho}{2\rho^*} \int_{\mathbb{R}^2} |\nabla u|^2 dx + \frac{1}{2} \int_{\mathbb{R}^2} \ln(1 + |x|^2) u^2 dx - \frac{1}{4} C \rho^{\frac{3}{2}} \left( \int_{\mathbb{R}^2} |\nabla u|^2 dx \right)^{\frac{1}{2}}, \quad (2.6)$$

which implies that  $E(u)$  is bounded from below.

Letting  $\{u_n\} \subset S_\rho$  be a minimizing sequence of  $e(\rho)$  for  $\rho \in (0, \rho^*)$ , we see from (2.6) that  $\int_{\mathbb{R}^2} |\nabla u_n|^2 dx$  and  $\int_{\mathbb{R}^2} \ln(1 + |x|^2) u_n^2 dx$  are bounded uniformly in  $n$ . Since  $\|u_n\|_2^2 = \rho$ , we know that  $\{u_n\}$  is bounded uniformly in  $X$ . According to Lemma 2.1 (1), there exists a function  $u \in X$  such that

$$u_n \rightharpoonup u \text{ in } X \text{ and } u_n \rightarrow u \text{ strongly in } L^s(\mathbb{R}^2), \quad s \in [2, \infty),$$

which implies that  $\|u\|_2^2 = \rho$ . We hence obtain that  $u \in S_\rho$ . Moreover, we note from (1) and (5) of Lemma 2.1 that

$$\int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| u^2(x)u^2(y) dx dy \leq \liminf_{n \rightarrow \infty} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| u_n^2(x)u_n^2(y) dx dy,$$

and

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^2} u_n^4 dx = \int_{\mathbb{R}^2} u^4 dx.$$

By the weakly lower semicontinuity, we then obtain from above that

$$e(\rho) \leq E(u) \leq \liminf_{n \rightarrow \infty} E(u_n) = e(\rho).$$

Thus,  $E(u) = e(\rho)$ , that is,  $u$  is a minimizer of  $e(\rho)$  for  $0 < \rho < \rho^*$ .

2. We next prove the non-existence of minimizers of  $e(\rho)$  for  $\rho \geq \rho^*$ . In this case, consider the test function

$$u_\tau(x) = \frac{\tau \rho^{\frac{1}{2}}}{\|Q\|_2} Q(\tau x), \quad \tau > 0,$$

so that  $u_\tau \in S_\rho$  holds for all  $\tau > 0$ . We then deduce from (2.3) that

$$\begin{aligned} e(\rho) \leq E(u_\tau) &= \frac{\rho(\rho^* - \rho)}{2\rho^*} \tau^2 + \frac{\rho}{2\rho^*} \int_{\mathbb{R}^2} \ln \left( 1 + \left| \frac{x}{\tau} \right|^2 \right) Q^2(x) dx \\ &+ \frac{\rho^2}{4(\rho^*)^2} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| Q^2(x)Q^2(y) dx dy - \frac{1}{4} \rho^2 \ln \tau, \quad \tau > 0. \end{aligned} \quad (2.7)$$

Together with (1.8), (2.1) and (2.5), we derive from (2.7) that  $e(\rho) \leq \lim_{\tau \rightarrow \infty} E(u_\tau) = -\infty$  holds for  $\rho \geq \rho^*$ .

Thus, there is no minimizer of  $e(\rho)$  for  $\rho \geq \rho^*$ .

Moreover, for  $\rho \in (0, \rho^*)$ , taking  $\tau = (\rho^* - \rho)^{-\frac{1}{2}}$  into (2.7) yields that

$$\lim_{\rho \nearrow \rho^*} e(\rho) \leq \lim_{\rho \nearrow \rho^*} E(u_\tau) = -\infty,$$

and hence  $\lim_{\rho \nearrow \rho^*} e(\rho) = -\infty$ . This completes the proof of Theorem 1.1.  $\square$

### 3 Limiting Behavior of Minimizers for $e(\rho)$ as $\rho \nearrow \rho^*$

In this section, we shall establish Theorem 1.3 on the limiting behavior of minimizers for  $e(\rho)$  as  $\rho \nearrow \rho^*$ . We first derive some estimates of the minimizers for  $e(\rho)$  as  $\rho \nearrow \rho^*$ .

**Lemma 3.1.** *Let  $u_\rho$  be a positive minimizer of  $e(\rho)$  for  $\rho \in (0, \rho^*)$ , and define*

$$\varepsilon_\rho = \left( \int_{\mathbb{R}^2} |\nabla u_\rho|^2 dx \right)^{-\frac{1}{2}} > 0. \quad (3.1)$$

Then we have

1.  $\varepsilon_\rho > 0$  satisfies

$$\varepsilon_\rho \rightarrow 0 \quad \text{and} \quad \mu_\rho \varepsilon_\rho^2 \rightarrow -\frac{1}{\rho^*} \quad \text{as} \quad \rho \nearrow \rho^*.$$

2. Denote

$$w_\rho(x) := \varepsilon_\rho u_\rho(\varepsilon_\rho x + x_\rho) > 0 \quad \text{in} \quad \mathbb{R}^2, \quad (3.2)$$

where  $x_\rho$  is a global maximal point of  $u_\rho$ . Then there exists a constant  $\eta > 0$ , independent of  $0 < \rho < \rho^*$ , such that

$$\int_{B_2(0)} w_\rho^2 dx \geq \eta > 0 \quad \text{as} \quad \rho \nearrow \rho^*. \quad (3.3)$$

3.  $w_\rho$  satisfies

$$w_\rho(x) \rightarrow \frac{1}{\sqrt{\rho^*}} Q\left(\frac{1}{\sqrt{\rho^*}}|x|\right) \quad \text{in} \quad H^1(\mathbb{R}^2) \quad \text{as} \quad \rho \nearrow \rho^*. \quad (3.4)$$

**Proof.** 1. We first prove that  $\varepsilon_\rho \rightarrow 0$  as  $\rho \nearrow \rho^*$ . Following (1.9) and (2.5), we derive that for  $\rho \in (0, \rho^*)$ ,

$$\begin{aligned} e(\rho) &= E(u_\rho) \\ &\geq \frac{\rho^* - \rho}{2\rho^*} \int_{\mathbb{R}^2} |\nabla u_\rho|^2 dx + \frac{1}{2} \int_{\mathbb{R}^2} \ln(1 + |x|^2) u_\rho^2(x) dx \\ &\quad + \frac{1}{8} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln(1 + |x - y|^2) u_\rho^2(x) u_\rho^2(y) dx dy - \frac{1}{4} C \rho^{\frac{3}{2}} \left( \int_{\mathbb{R}^2} |\nabla u_\rho|^2 dx \right)^{\frac{1}{2}} \\ &\geq \frac{\rho^* - \rho}{2\rho^*} \varepsilon_\rho^{-2} - \frac{1}{4} C \rho^{\frac{3}{2}} \varepsilon_\rho^{-1} \geq -\frac{1}{4} C \rho^{\frac{3}{2}} \varepsilon_\rho^{-1}. \end{aligned} \quad (3.5)$$

Moreover, we have  $\lim_{\rho \nearrow \rho^*} e(\rho) = -\infty$  in view of Theorem 1.1, together with (3.5), which yields that  $\varepsilon_\rho \rightarrow 0$  as  $\rho \nearrow \rho^*$ .

We next prove that  $\mu_\rho \varepsilon_\rho^2 \rightarrow -\frac{1}{\rho^*}$  as  $\rho \nearrow \rho^*$ . By the definition of  $\varepsilon_\rho$ , we have

$$\begin{aligned} \varepsilon_\rho^2 e(\rho) &= \frac{1}{2} + \frac{\varepsilon_\rho^2}{2} \int_{\mathbb{R}^2} \ln(1 + |x|^2) u_\rho^2(x) dx + \frac{\varepsilon_\rho^2}{4} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x - y| u_\rho^2(x) u_\rho^2(y) dx dy - \frac{\varepsilon_\rho^2}{4} \int_{\mathbb{R}^2} u_\rho^4(x) dx \\ &= \frac{1}{2} + \frac{\varepsilon_\rho^2}{2} \int_{\mathbb{R}^2} \ln(1 + |x|^2) u_\rho^2(x) dx + \frac{\varepsilon_\rho^2}{8} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln(1 + |x - y|^2) u_\rho^2(x) u_\rho^2(y) dx dy \\ &\quad - \frac{\varepsilon_\rho^2}{8} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln\left(1 + \frac{1}{|x - y|^2}\right) u_\rho^2(x) u_\rho^2(y) dx dy - \frac{\varepsilon_\rho^2}{4} \int_{\mathbb{R}^2} u_\rho^4(x) dx. \end{aligned} \quad (3.6)$$

Since  $\varepsilon_\rho \rightarrow 0$  as  $\rho \nearrow \rho^*$ , we also derive from (2.5) and that

$$\frac{1}{2} \varepsilon_\rho^2 \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln\left(1 + \frac{1}{|x - y|^2}\right) u_\rho^2(x) u_\rho^2(y) dx dy \leq \varepsilon_\rho^2 \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{u_\rho^2(x) u_\rho^2(y)}{|x - y|} dx dy \leq C \rho^{\frac{3}{2}} \varepsilon_\rho \rightarrow 0 \quad \text{as} \quad \rho \nearrow \rho^*. \quad (3.7)$$

By the fact that  $\lim_{\rho \nearrow \rho^*} e(\rho) = -\infty$ , we deduce from (1.9), (3.6) and (3.7) that

$$\begin{aligned} 0 \geq \limsup_{\rho \nearrow \rho^*} \varepsilon_\rho^2 e(\rho) &= \limsup_{\rho \nearrow \rho^*} \frac{1}{2} \left( 1 - \frac{\varepsilon_\rho^2}{2} \int_{\mathbb{R}^2} u_\rho^4(x) dx \right) + \limsup_{\rho \nearrow \rho^*} \frac{\varepsilon_\rho^2}{2} \int_{\mathbb{R}^2} \ln(1 + |x|^2) u_\rho^2(x) dx \\ &\quad + \limsup_{\rho \nearrow \rho^*} \frac{\varepsilon_\rho^2}{8} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln(1 + |x - y|^2) u_\rho^2(x) u_\rho^2(y) dx dy \geq 0, \end{aligned}$$

which implies that

$$\begin{aligned} \lim_{\rho \nearrow \rho^*} \varepsilon_\rho^2 \int_{\mathbb{R}^2} u_\rho^4 dx &= 2, \quad \lim_{\rho \nearrow \rho^*} \varepsilon_\rho^2 \int_{\mathbb{R}^2} \ln(1 + |x|^2) u_\rho^2 dx = 0, \\ \lim_{\rho \nearrow \rho^*} \varepsilon_\rho^2 \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln(1 + |x - y|^2) u_\rho^2(x) u_\rho^2(y) dx dy &= 0, \quad \lim_{\rho \nearrow \rho^*} \varepsilon_\rho^2 e(\rho) = 0. \end{aligned} \tag{3.8}$$

Because  $u_\rho$  is a positive minimizer of  $e(\rho)$ , it satisfies the Euler-Lagrange equation (1.11), where the Lagrange multiplier  $\mu_\rho \in \mathbb{R}$  satisfies

$$\mu_\rho \rho = 2e(\rho) + \frac{1}{2} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x - y| u_\rho^2(x) u_\rho^2(y) dx dy - \frac{1}{2} \int_{\mathbb{R}^2} u_\rho^4(x) dx. \tag{3.9}$$

We thus obtain from (3.7)–(3.9) that

$$\mu_\rho \varepsilon_\rho^2 \rightarrow -\frac{1}{\rho^*} \quad \text{as } \rho \nearrow \rho^*. \tag{3.10}$$

2. By the definition of  $w_\rho$ , we reduce from (1.11) that  $w_\rho$  satisfies

$$\begin{aligned} -\Delta w_\rho + \varepsilon_\rho^2 \ln[1 + |\varepsilon_\rho x + x_\rho|^2] w_\rho + \varepsilon_\rho^2 \left( \int_{\mathbb{R}^2} \ln|x - y| w_\rho^2(y) dy \right) w_\rho + \rho \varepsilon_\rho^2 \ln \varepsilon_\rho w_\rho \\ = \mu_\rho \varepsilon_\rho^2 w_\rho + w_\rho^3 \quad \text{in } \mathbb{R}^2. \end{aligned} \tag{3.11}$$

Since  $w_\rho$  is bounded uniformly in  $H^1(\mathbb{R}^2)$ , a straightforward calculation shows that there exists a constant  $C > 0$  such that for any  $x \in \mathbb{R}^2$  and  $\rho \in (0, \rho^*)$ ,

$$\begin{aligned} &\frac{1}{2} \int_{\mathbb{R}^2} \ln\left(1 + \frac{1}{|x - y|^2}\right) w_\rho^2(y) dy \\ &\leq \int_{\mathbb{R}^2} \frac{1}{|x - y|} w_\rho^2(y) dy \\ &= \int_{|x - y| < 1} \frac{1}{|x - y|} w_\rho^2(y) dy + \int_{|x - y| \geq 1} \frac{1}{|x - y|} w_\rho^2(y) dy \\ &\leq \left( \int_{|x - y| < 1} \frac{1}{|x - y|^{\frac{3}{2}}} dy \right)^{\frac{2}{3}} \left( \int_{|x - y| < 1} w_\rho^6 dy \right)^{\frac{1}{3}} + \int_{|x - y| \geq 1} w_\rho^2(y) dy \\ &\leq C \|w_\rho\|^2 \leq C, \end{aligned} \tag{3.12}$$

where and below we denote

$$\|u\| := \|u\|_{H^1(\mathbb{R}^2)} = \left\{ \int_{\mathbb{R}^2} (|\nabla u|^2 + u^2) dx \right\}^{\frac{1}{2}}, \quad u \in H^1(\mathbb{R}^2). \tag{3.13}$$

We derive from (3.10)–(3.12) that

$$-\Delta w_\rho + \frac{1}{2\rho^*} w_\rho - w_\rho^3 \leq 0 \quad \text{in } \mathbb{R}^2 \quad \text{as } \rho \nearrow \rho^*. \tag{3.14}$$

Since  $x_\rho$  is a global maximal point of  $u_\rho$ , the origin is a maximal point of  $w_\rho$  for all  $\rho \in (0, \rho^*)$ . This implies that  $-\Delta w_\rho(0) \geq 0$  for  $\rho \in (0, \rho^*)$ . We then infer from (3.14) that there exists some constant  $\alpha > 0$ ,



independent of  $\rho$ , such that  $w_\rho(0) \geq \alpha > 0$  as  $\rho \nearrow \rho^*$ . Applying the De Giorgi-Nash-Moser theory [35, Theorem 4.1], we deduce from (3.14) that there exists a constant  $C > 0$  such that

$$\int_{B_2(0)} w_\rho^2 dx \geq C \max_{x \in B_1(0)} w_\rho(x) \geq C(\alpha) := \eta > 0 \text{ as } \rho \nearrow \rho^*,$$

which thus completes the proof of (3.3).

3. Since the sequence  $\{w_\rho\}$  is bounded uniformly in  $H^1(\mathbb{R}^2)$ , up to a subsequence if necessary, there exists a function  $w_0 \in H^1(\mathbb{R}^2)$  such that  $w_\rho \rightharpoonup w_0$  in  $H^1(\mathbb{R}^2)$ ,  $w_\rho \rightarrow w_0$  in  $L^q_{loc}(\mathbb{R}^2)$  for  $q \in [2, \infty)$ , and  $w_\rho \rightarrow w_0$  almost everywhere in  $\mathbb{R}^2$  as  $\rho \nearrow \rho^*$ . Moreover, we have  $w_0 \not\equiv 0$  in view of (3.3). Note from the Brézis-Lieb lemma ([57]) that

$$\|w_\rho\|_2^2 = \|w_0\|_2^2 + \|w_\rho - w_0\|_2^2 + o(1) \text{ as } \rho \nearrow \rho^*,$$

$$\|w_\rho\|_4^4 = \|w_0\|_4^4 + \|w_\rho - w_0\|_4^4 + o(1) \text{ as } \rho \nearrow \rho^*,$$

and

$$\|\nabla w_\rho\|_2^2 = \|\nabla w_0\|_2^2 + \|\nabla w_\rho - \nabla w_0\|_2^2 + o(1) \text{ as } \rho \nearrow \rho^*.$$

Combining (3.8), it yields from (1.9) that

$$\begin{aligned} 0 &= \lim_{\rho \nearrow \rho^*} \left( \int_{\mathbb{R}^2} |\nabla w_\rho|^2 dx - \frac{1}{2} \int_{\mathbb{R}^2} w_\rho^4 dx \right) \\ &= \int_{\mathbb{R}^2} |\nabla w_0|^2 dx - \frac{1}{2} \int_{\mathbb{R}^2} w_0^4 dx + \lim_{\rho \nearrow \rho^*} \left( \int_{\mathbb{R}^2} |\nabla w_\rho - \nabla w_0|^2 dx - \frac{1}{2} \int_{\mathbb{R}^2} |w_\rho - w_0|^4 dx \right) \\ &\geq \frac{1}{2} \left( \rho^* \|w_0\|_2^{-2} - 1 \right) \int_{\mathbb{R}^2} |w_0|^4 dx + \lim_{\rho \nearrow \rho^*} \left( \int_{\mathbb{R}^2} |\nabla w_\rho - \nabla w_0|^2 dx - \frac{1}{2} \int_{\mathbb{R}^2} |w_\rho - w_0|^4 dx \right) \\ &\geq \lim_{\rho \nearrow \rho^*} \left( 1 - \frac{1}{\rho^*} \int_{\mathbb{R}^2} |w_\rho - w_0|^2 dx \right) \int_{\mathbb{R}^2} |\nabla w_\rho - \nabla w_0|^2 dx \geq 0, \end{aligned} \tag{3.15}$$

from which we obtain that  $\|w_0\|_2^2 = \rho^*$  and

$$\int_{\mathbb{R}^2} |\nabla w_\rho - \nabla w_0|^2 dx \rightarrow 0 \text{ as } \rho \nearrow \rho^*,$$

and hence

$$w_\rho \rightarrow w_0 \text{ strongly in } H^1(\mathbb{R}^2) \text{ as } \rho \nearrow \rho^*.$$

On the other hand, the second equality of (3.15) yields that

$$\int_{\mathbb{R}^2} |\nabla w_0|^2 dx = \frac{1}{2} \int_{\mathbb{R}^2} w_0^4 dx,$$

together with the fact  $\|w_0\|_2^2 = \rho^*$ , which implies that the identity of the inequality (1.9) is achieved by  $w_0$ . Thus we derive that  $w_0$  satisfies

$$-\Delta w_0 + \frac{1}{\rho^*} w_0 - w_0^3 = 0 \text{ in } \mathbb{R}^2,$$

and furthermore there exists  $y_0 \in \mathbb{R}^2$  such that, up to a subsequence if necessary,

$$w_\rho(x) \rightarrow w_0(x) = \frac{1}{\sqrt{\rho^*}} Q\left(\frac{1}{\sqrt{\rho^*}} |x + y_0|\right) \text{ in } H^1(\mathbb{R}^2) \text{ as } \rho \nearrow \rho^*.$$

Moreover, since the origin is a global maximal point of  $w_\rho$  for  $\rho \in (0, \rho^*)$ , it is also a global maximal point of  $w_0$ . This implies that  $y_0 = 0$ , and hence

$$w_\rho(x) \rightarrow w_0(x) = \frac{1}{\sqrt{\rho^*}} Q\left(\frac{1}{\sqrt{\rho^*}} |x|\right) \text{ in } H^1(\mathbb{R}^2) \text{ as } \rho \nearrow \rho^*. \tag{3.16}$$

Since the convergence (3.16) is independent of the choice of the subsequence, we deduce that (3.16) holds essentially true for the whole sequence. This therefore completes the proof of Lemma 3.1.  $\square$

The following lemma is concerned with the exponential decay and  $L^\infty$ -convergence of  $w_\rho$  as  $\rho \nearrow \rho^*$ .

**Lemma 3.2.** *Assume that  $u_\rho$  is a positive minimizer of  $e(\rho)$  for  $\rho \in (0, \rho^*)$ , and  $w_\rho > 0$  is defined by (3.2). Then we have*

1. *There exists a large constant  $R > 0$  such that*

$$|w_\rho(x)| \leq Ce^{-\frac{2|x|}{3\sqrt{\rho^*}}} \quad \text{and} \quad |\nabla w_\rho(x)| \leq Ce^{-\frac{|x|}{2\sqrt{\rho^*}}} \quad \text{for } |x| \geq R \text{ as } \rho \nearrow \rho^*, \quad (3.17)$$

where the constant  $C > 0$  is independent of  $\rho \in (0, \rho^*)$ .

2. *The function  $w_\rho$  satisfies*

$$w_\rho(x) \rightarrow \frac{1}{\sqrt{\rho^*}} Q\left(\frac{1}{\sqrt{\rho^*}}|x|\right) \quad \text{in } L^\infty(\mathbb{R}^2) \text{ as } \rho \nearrow \rho^*. \quad (3.18)$$

Moreover, the global maximal point  $x_\rho$  of  $u_\rho$  is unique and satisfies  $x_\rho \rightarrow 0$  as  $\rho \nearrow \rho^*$ .

**Proof.** 1. Since  $\varepsilon_\rho \ln \varepsilon_\rho \rightarrow 0$  as  $\rho \nearrow \rho^*$ , inserting (3.10) and (3.12) into (3.11) yields that

$$-\Delta w_\rho + \frac{5}{9\rho^*} w_\rho - w_\rho^3 \leq 0 \quad \text{in } \mathbb{R}^2 \text{ as } \rho \nearrow \rho^*. \quad (3.19)$$

Applying De Giorgi-Nash-Moser theory [35, Theorem 4.1], we then derive from (3.19) that there exists a large constant  $R > 0$  such that

$$w_\rho(x) \leq \frac{1}{3\sqrt{\rho^*}} \quad \text{uniformly for } |x| \geq R \text{ as } \rho \nearrow \rho^*, \quad (3.20)$$

due to the fact that  $w_\rho \rightarrow w_0$  in  $L^2(\mathbb{R}^2)$  as  $\rho \nearrow \rho^*$ . We now derive from (3.19) and (3.20) that

$$-\Delta w_\rho + \frac{4}{9\rho^*} w_\rho \leq 0 \quad \text{uniformly for } |x| \geq R \text{ as } \rho \nearrow \rho^*. \quad (3.21)$$

Applying the comparison principle to (3.21) gives that there exists a constant  $C > 0$  such that

$$|w_\rho(x)| \leq Ce^{-\frac{2|x|}{3\sqrt{\rho^*}}} \quad \text{uniformly for } |x| \geq R \text{ as } \rho \nearrow \rho^*. \quad (3.22)$$

Since the proof of exponential decay for  $|\nabla w_\rho|$  is more involved, we leave it to Lemma A.1 in the Appendix. This completes the proof of (3.17).

2. To obtain the estimate of  $x_\rho$  as  $\rho \nearrow \rho^*$ , we first claim that

$$\int_{\mathbb{R}^2} \ln(1 + |x|^2) u_\rho^2(x) dx \rightarrow 0 \quad \text{as } \rho \nearrow \rho^*. \quad (3.23)$$

In fact, by the the exponential decays (1.8) and (3.17), it follows from (3.4) that

$$\begin{aligned} \int_{\mathbb{R}^2} \ln(1 + |x|^2) u_\rho^2(x + x_\rho) dx &= \int_{\mathbb{R}^2} \ln(1 + \varepsilon_\rho^2 |x|^2) w_\rho^2(x) dx \\ &\leq \varepsilon_\rho^2 \int_{\mathbb{R}^2} |x|^2 w_\rho^2(x) dx \\ &= (1 + o(1)) \rho^* \varepsilon_\rho^2 \int_{\mathbb{R}^2} |x|^2 Q^2(x) dx \quad \text{as } \rho \nearrow \rho^*, \end{aligned}$$

which shows that  $u_\rho(\cdot + x_\rho) \in S_\rho$  as  $\rho \nearrow \rho^*$ . By the definition of  $e(\rho)$ , we have

$$e(\rho) = E(u_\rho) \leq E(u_\rho(\cdot + x_\rho)),$$

which implies that

$$\begin{aligned} 0 &\leq \int_{\mathbb{R}^2} \ln(1 + |x|^2) u_\rho^2(x) dx \leq \int_{\mathbb{R}^2} \ln(1 + |x|^2) u_\rho^2(x + x_\rho) dx \\ &\leq (1 + o(1)) \rho^* \varepsilon_\rho^2 \int_{\mathbb{R}^2} |x|^2 Q^2(x) dx \rightarrow 0 \quad \text{as } \rho \nearrow \rho^*, \end{aligned} \quad (3.24)$$

and hence the claim (3.23) is proved.

Based on (3.23), we next prove that any maximal point  $x_\rho$  of  $u_\rho$  satisfies  $x_\rho \rightarrow 0$  as  $\rho \nearrow \rho^*$ . On the contrary, we assume that there exist a constant  $\gamma > 0$  and a subsequence  $\{\rho_k\}$ , where  $\rho_k \nearrow \rho^*$  as  $k \rightarrow \infty$ , such that

$$|x_{\rho_k}| \geq \gamma \text{ as } k \rightarrow \infty. \quad (3.25)$$

By Fatou's Lemma, it then follows from (3.3) and (3.25) that

$$\begin{aligned} \liminf_{k \rightarrow \infty} \int_{\mathbb{R}^2} \ln(1 + |x|^2) u_{\rho_k}^2(x) dx &= \liminf_{k \rightarrow \infty} \int_{\mathbb{R}^2} \ln(1 + |\varepsilon_{\rho_k} x + x_{\rho_k}|^2) w_{\rho_k}^2(x) dx \\ &\geq \liminf_{k \rightarrow \infty} \int_{B_2(0)} \ln(1 + |\varepsilon_{\rho_k} x + x_{\rho_k}|^2) w_{\rho_k}^2(x) dx \\ &\geq \frac{1}{2} \ln(1 + \gamma^2) \eta > 0, \end{aligned}$$

which is a contradiction in view of (3.23). Hence, we have  $x_\rho \rightarrow 0$  as  $\rho \nearrow \rho^*$ .

In the following, we prove that

$$w_\rho \rightarrow w_0 = \frac{1}{\sqrt{\rho^*}} Q\left(\frac{1}{\sqrt{\rho^*}} |x|\right) \text{ in } L^\infty(\mathbb{R}^2) \text{ as } \rho \nearrow \rho^*.$$

In view of the exponential decays (1.8) and (3.17), it suffices to show the  $L^\infty$ -uniform convergence of  $w_\rho$  on any compact domain of  $\mathbb{R}^2$  as  $\rho \nearrow \rho^*$ . Denote

$$\begin{aligned} G_\rho(x) &:= -\varepsilon_\rho^2 \ln(1 + |\varepsilon_\rho x + x_\rho|^2) w_\rho - \varepsilon_\rho^2 \left( \int_{\mathbb{R}^2} \ln|x-y| w_\rho^2(y) dy \right) w_\rho \\ &\quad - \rho \varepsilon_\rho^2 \ln \varepsilon_\rho w_\rho + \mu_\rho \varepsilon_\rho^2 w_\rho + w_\rho^3 \text{ in } \mathbb{R}^2. \end{aligned}$$

We then get from (3.11) that

$$-\Delta w_\rho = G_\rho \text{ in } \mathbb{R}^2. \quad (3.26)$$

Because  $\{w_\rho\}$  is bounded uniformly in  $H^1(\mathbb{R}^2)$  and satisfies the exponential decay (3.17), we obtain that  $G_\rho$  is also bounded uniformly in  $L_{loc}^2(\mathbb{R}^2)$  as  $\rho \nearrow \rho^*$ , in view of the fact that  $x_\rho \rightarrow 0$  as  $\rho \nearrow \rho^*$ . For any  $R > 0$ , it follows from [28, Theorem 8.8] that

$$\|w_\rho\|_{H^2(B_R)} \leq C \left( \|w_\rho\|_{H^1(B_{R+1})} + \|G_\rho\|_{L^2(B_{R+1})} \right),$$

where  $C > 0$  is independent of  $\rho$  and  $R$ . Therefore,  $\{w_\rho\}$  is also bounded uniformly in  $H_{loc}^2(\mathbb{R}^2)$  as  $\rho \nearrow \rho^*$ . Since the embedding  $H^2(B_R) \hookrightarrow L^\infty(B_R)$  is compact (cf. [28, Theorem 7.26]), we deduce from (3.16) that there exists a subsequence of  $\{w_\rho\}$ , still denoted by  $\{w_\rho\}$ , such that

$$\lim_{\rho \nearrow \rho^*} w_\rho(x) = w_0(x) \text{ in } L^\infty(B_R).$$

Since  $R > 0$  is arbitrary, we get that

$$\lim_{\rho \nearrow \rho^*} w_\rho(x) = w_0(x) \text{ in } L_{loc}^\infty(\mathbb{R}^2). \quad (3.27)$$

Because the above convergence is independent of the subsequence that we choose, (3.27) holds essentially for the whole sequence. This shows that (3.18) holds true.

We now prove the uniqueness of  $x_\rho$  as  $\rho \nearrow \rho^*$ , where  $x_\rho$  is a global maximal point of  $u_\rho$ . Since  $\{w_\rho\}$  is bounded uniformly in  $H^1(\mathbb{R}^2)$ , we get that  $\{w_\rho\}$  is bounded uniformly in  $L_{loc}^q(\mathbb{R}^2)$  as  $\rho \nearrow \rho^*$  for  $q \in [2, \infty)$ . Applying the  $L^p$  estimate [28, Theorem 9.11] to (3.26), we then derive that  $\{w_\rho\}$  is bounded uniformly in  $W_{loc}^{2,q}(\mathbb{R}^2)$  as  $\rho \nearrow \rho^*$ . By a standard Sobolev embedding, it holds that

$$\{w_\rho\} \text{ is bounded uniformly in } C_{loc}^{1,\alpha}(\mathbb{R}^2) \text{ as } \rho \nearrow \rho^*. \quad (3.28)$$

Furthermore, since it gives from [17, Proposition 2.3] that

$$\varepsilon_\rho^2 \ln(1 + |\varepsilon_\rho x + x_\rho|^2) \in C_{loc}^\alpha(\mathbb{R}^2), \quad \varepsilon_\rho^2 \int_{\mathbb{R}^2} \ln|x-y| w_\rho^2(y) dy \in C_{loc}^{2,\alpha}(\mathbb{R}^2),$$

we have  $G_\rho$  is bounded uniformly in  $C^\alpha(B_R(0))$ , where  $R > 0$  is large enough. Applying the Schauder estimate [28, Theorem 6.2], it yields from (3.26) that  $\{w_\rho\}$  is bounded uniformly in  $C^{2,\alpha}(B_R(0))$  as  $\rho \nearrow \rho^*$ . Therefore, up to a subsequence if necessary, there exists a function  $\tilde{w}_0 \in C^2(B_R(0))$  such that

$$w_\rho \rightarrow \tilde{w}_0 \quad \text{in } C^2(B_R(0)) \quad \text{as } \rho \nearrow \rho^*,$$

and we thus deduce from (3.18) that  $\tilde{w}_0 = w_0$ , so that

$$w_\rho \rightarrow w_0 \quad \text{in } C^2(B_R(0)) \quad \text{as } \rho \nearrow \rho^*, \quad (3.29)$$

where  $R > 0$  is a constant large enough. Since the convergence (3.29) is independent of the subsequence that we choose, it holds essentially for the whole sequence.

Because the origin is the unique global maximal point of  $Q(x)$ , the convergence of (3.29) shows that all local maximal points of  $w_\rho$  must approach the origin and hence stay in a small ball  $B_\delta(0)$  as  $\rho \nearrow \rho^*$  for some  $\delta > 0$ . Since  $Q''(0) < 0$ , we conclude that  $Q''(r) < 0$  for  $r \in [0, \delta)$ . It then follows from [47, Lemma 4.2] that as  $\rho \nearrow \rho^*$ , each  $w_\rho$  has a unique global maximum point which is exactly the origin. This further proves the uniqueness of global maximal points for  $u_\rho$  as  $\rho \nearrow \rho^*$ . This completes the proof of Lemma 3.2.  $\square$

Following the  $L^\infty$ -uniform convergence of positive minimizers  $u_\rho$  for  $e(\rho)$  as  $\rho \nearrow \rho^*$ , the rest part of this section is to complete the proof of Theorem 1.3. In order to reach this aim, more importantly, we shall set up the refined energy estimates, based on which we shall analyze the blow up point  $x_\rho$  and the blow up rate  $\varepsilon_\rho^{-1}$  of  $u_\rho$  as  $\rho \nearrow \rho^*$ . In addition, the convergence of  $u_\rho$  in  $X$  can be obtained in view of the exponential decay.

**Proof of Theorem 1.3.** To complete the proof of Theorem 1.3, we note from Lemmas 3.1 and 3.2 that the rest is to prove

$$\bar{\varepsilon}_\rho := \sqrt{\rho^*} \varepsilon_\rho = \frac{2}{\rho^*} (\rho^* - \rho)^{\frac{1}{2}} (1 + o(1)) \quad \text{as } \rho \nearrow \rho^*, \quad (3.30)$$

and

$$\frac{x_\rho}{\bar{\varepsilon}_\rho} \rightarrow 0 \quad \text{as } \rho \nearrow \rho^*. \quad (3.31)$$

We first obtain the upper estimate of  $e(\rho)$  as  $\rho \nearrow \rho^*$ . Similar to (2.7), one can get that for any  $\tau > 0$ ,

$$\begin{aligned} e(\rho) \leq E(u_\tau) &\leq \frac{\rho(\rho^* - \rho)}{2\rho^*} \tau^2 - \frac{1}{4} \rho^2 \ln \tau + \frac{\rho}{2\rho^* \tau^2} \int_{\mathbb{R}^2} |x|^2 Q^2(x) dx \\ &\quad + \frac{\rho^2}{4(\rho^*)^2} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| Q^2(x) Q^2(y) dx dy. \end{aligned} \quad (3.32)$$

Setting

$$\tau = \left[ \frac{\rho \rho^*}{4(\rho^* - \rho)} \right]^{\frac{1}{2}} > 0$$

into (3.32), we deduce that

$$\begin{aligned} e(\rho) &\leq \frac{(\rho^*)^2}{8} - \frac{(\rho^*)^2}{4} \ln \rho^* + \frac{(\rho^*)^2}{8} \ln [4(\rho^* - \rho)] \\ &\quad + \frac{1}{4} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| Q^2(x) Q^2(y) dx dy + o(1) \quad \text{as } \rho \nearrow \rho^*. \end{aligned} \quad (3.33)$$

To derive the lower estimate of  $e(\rho)$  as  $\rho \nearrow \rho^*$ , take  $u_\rho \in X$  to be a positive minimizer of  $e(\rho)$  satisfying  $\|u_\rho\|_2^2 = \rho < \rho^*$ . Denote

$$\bar{w}_\rho(x) = \bar{\varepsilon}_\rho u_\rho(\bar{\varepsilon}_\rho x + x_\rho) = \sqrt{\rho^*} \varepsilon_\rho u_\rho(\sqrt{\rho^*} \varepsilon_\rho x + x_\rho) > 0 \quad \text{in } \mathbb{R}^2,$$

where  $x_\rho$  is the maximal point of  $u_\rho$ . By the exponential decays (1.8) and (3.17), we obtain that for any  $\epsilon > 0$ , there exists a constant  $R > 0$  such that

$$2 \int_{B_R^c} \ln(1 + |x|^2) \bar{w}_\rho^2(x) dx + 2 \int_{B_R^c} \ln(1 + |x|^2) Q^2(x) dx < \frac{\epsilon}{2},$$

from which and (3.16) we deduce that

$$\begin{aligned} & \int_{\mathbb{R}^2} \ln(1 + |x|^2) \left( \bar{w}_\rho(x) - Q(x) \right)^2 dx \\ & \leq \ln(1 + R^2) \int_{B_R} \left( \bar{w}_\rho(x) - Q(x) \right)^2 dx + 2 \int_{B_R^c} \ln(1 + |x|^2) \bar{w}_\rho^2(x) dx + 2 \int_{B_R^c} \ln(1 + |x|^2) Q^2(x) dx \\ & < \epsilon \text{ as } \rho \nearrow \rho^*. \end{aligned}$$

This then implies that  $\bar{w}_\rho \rightarrow Q$  in  $X$  as  $\rho \nearrow \rho^*$ . Moreover, combining (3.12) with (3.16) yields that

$$\begin{aligned} & \left| \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| \bar{w}_\rho^2(x) \bar{w}_\rho^2(y) dx dy - \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| Q^2(x) Q^2(y) dx dy \right| \\ & \leq \left| \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| \left( \bar{w}_\rho^2(x) - Q^2(x) \right) \bar{w}_\rho^2(y) dx dy \right| \\ & \quad + \left| \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| Q^2(x) \left( \bar{w}_\rho^2(y) - Q^2(y) \right) dx dy \right| \\ & \leq \rho \int_{\mathbb{R}^2} \ln(1 + |x|^2) \left| \bar{w}_\rho^2(x) - Q^2(x) \right| dx + \int_{\mathbb{R}^2} \left| \bar{w}_\rho^2(x) - Q^2(x) \right| dx \int_{\mathbb{R}^2} \ln(1 + |y|^2) \bar{w}_\rho^2(y) dy \\ & \quad + \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{1}{|x-y|} \left| \bar{w}_\rho^2(x) - Q^2(x) \right| \bar{w}_\rho^2(y) dx dy \\ & \quad + \int_{\mathbb{R}^2} \ln(1 + |x|^2) Q^2(x) dx \int_{\mathbb{R}^2} \left| \bar{w}_\rho^2(y) - Q^2(y) \right| dy + \rho^* \int_{\mathbb{R}^2} \ln(1 + |y|^2) \left| \bar{w}_\rho^2(y) - Q^2(y) \right| dy \\ & \quad + \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{1}{|x-y|} Q^2(x) \left| \bar{w}_\rho^2(y) - Q^2(y) \right| dx dy \rightarrow 0 \text{ as } \rho \nearrow \rho^*. \end{aligned} \tag{3.34}$$

We then deduce from (1.9), (3.4) and (3.34) that

$$\begin{aligned} e(\rho) &= E(u_\rho) \\ &= \frac{1}{2} \bar{\epsilon}_\rho^{-2} \left( \int_{\mathbb{R}^2} |\nabla \bar{w}_\rho|^2 dx - \frac{\rho^*}{2\rho} \int_{\mathbb{R}^2} \bar{w}_\rho^4 dx \right) + \frac{1}{2} \int_{\mathbb{R}^2} \ln(1 + |\bar{\epsilon}_\rho x + x_\rho|^2) \bar{w}_\rho^2(x) dx \\ & \quad + \frac{1}{4} \rho^2 \ln \bar{\epsilon}_\rho + \frac{1}{4} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| \bar{w}_\rho^2(x) \bar{w}_\rho^2(y) dx dy + \frac{1}{4} \left( \frac{\rho^*}{\rho} - 1 \right) \bar{\epsilon}_\rho^{-2} \int_{\mathbb{R}^2} \bar{w}_\rho^4 dx \\ & \geq \frac{\rho^* - \rho}{4\rho} \bar{\epsilon}_\rho^{-2} \int_{\mathbb{R}^2} \bar{w}_\rho^4 dx + \frac{1}{4} \rho^2 \ln \bar{\epsilon}_\rho + \frac{1}{4} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| \bar{w}_\rho^2(x) \bar{w}_\rho^2(y) dx dy \\ & = \frac{\rho^* (\rho^* - \rho)}{2\rho} (1 + o(1)) \bar{\epsilon}_\rho^{-2} + \frac{1}{4} \rho^2 \ln \bar{\epsilon}_\rho + \frac{1}{4} (1 + o(1)) \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| Q^2(x) Q^2(y) dx dy \\ & \geq \frac{(\rho^*)^2}{8} - \frac{(\rho^*)^2}{4} \ln \rho^* + \frac{(\rho^*)^2}{8} \ln [4(\rho^* - \rho)] \\ & \quad + \frac{1}{4} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| Q^2(x) Q^2(y) dx dy + o(1) \text{ as } \rho \nearrow \rho^*, \end{aligned} \tag{3.35}$$

where the identity of the last inequality is achieved at  $\bar{\epsilon}_\rho > 0$  satisfying (3.30), i.e.,  $\bar{\epsilon}_\rho = \frac{2}{\rho^*} (\rho^* - \rho)^{\frac{1}{2}} (1 + o(1))$  as  $\rho \nearrow \rho^*$ . Together with (3.33) and (3.35), we now conclude that

$$\begin{aligned} e(\rho) &\approx \frac{(\rho^*)^2}{8} - \frac{(\rho^*)^2}{4} \ln \rho^* + \frac{(\rho^*)^2}{8} \ln [4(\rho^* - \rho)] \\ & \quad + \frac{1}{4} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| Q^2(x) Q^2(y) dx dy \text{ as } \rho \nearrow \rho^*, \end{aligned}$$

and  $\bar{\varepsilon}_\rho > 0$  satisfies (3.30). In addition, since  $\bar{w}_\rho \rightarrow Q$  in  $X$  as  $\rho \nearrow \rho^*$ , we derive from Lemmas 3.1 and 3.2 that

$$\lim_{\rho \nearrow \rho^*} \frac{2(\rho^* - \rho)^{\frac{1}{2}}}{\rho^*} u_\rho \left( \frac{2(\rho^* - \rho)^{\frac{1}{2}}}{\rho^*} x + x_\rho \right) = Q(x) \quad \text{in } X \cap L^\infty(\mathbb{R}^2).$$

We finally prove that  $\bar{\varepsilon}_\rho^{-1} x_\rho \rightarrow 0$  as  $\rho \nearrow \rho^*$ . It follows from (3.24) that

$$\int_{\mathbb{R}^2} \ln(1 + |x|^2) u_\rho^2(x + x_\rho) dx \leq (1 + o(1)) \bar{\varepsilon}_\rho^2 \int_{\mathbb{R}^2} |x|^2 Q^2(x) dx \quad \text{as } \rho \nearrow \rho^*. \quad (3.36)$$

On the other hand, we note that  $x_\rho \rightarrow 0$  and  $\bar{w}_\rho \rightarrow Q$  in  $H^1(\mathbb{R}^2)$  as  $\rho \nearrow \rho^*$ . Since  $u_\rho$  is a positive minimizer of  $e(\rho)$ , we now have

$$\begin{aligned} & \int_{\mathbb{R}^2} \ln(1 + |x|^2) u_\rho^2(x) dx \\ & \geq \int_{B_{\frac{1}{\sqrt{\bar{\varepsilon}_\rho}}}(0)} \ln(1 + |\bar{\varepsilon}_\rho x + x_\rho|^2) \bar{w}_\rho^2(x) dx \\ & = (1 + o(1)) \bar{\varepsilon}_\rho^2 \int_{B_{\frac{1}{\sqrt{\bar{\varepsilon}_\rho}}}(0)} \left| x + \frac{x_\rho}{\bar{\varepsilon}_\rho} \right|^2 Q^2(x) dx \quad \text{as } \rho \nearrow \rho^*. \end{aligned} \quad (3.37)$$

Similar to (3.24), we deduce from (3.36) and (3.37) that

$$\bar{\varepsilon}_\rho^2 \int_{B_{\frac{1}{\sqrt{\bar{\varepsilon}_\rho}}}(0)} \left| x + \frac{x_\rho}{\bar{\varepsilon}_\rho} \right|^2 Q^2(x) dx \leq (1 + o(1)) \bar{\varepsilon}_\rho^2 \int_{\mathbb{R}^2} |x|^2 Q^2(x) dx \quad \text{as } \rho \nearrow \rho^*. \quad (3.38)$$

On the contrary, we now assume that (3.31) is false, i.e.,  $\frac{x_\rho}{\bar{\varepsilon}_\rho} \not\rightarrow 0$  as  $\rho \nearrow \rho^*$ . Then there exist a constant  $\tau > 0$  and a subsequence  $\{\rho_k\}$ , where  $\rho_k \nearrow \rho^*$  as  $k \rightarrow \infty$ , such that

$$\frac{|x_{\rho_k}|}{\bar{\varepsilon}_{\rho_k}} \geq \tau > 0 \quad \text{as } k \rightarrow \infty. \quad (3.39)$$

By Fatou's Lemma, it then follows from (3.38) and (3.39) that

$$\int_{\mathbb{R}^2} |x|^2 Q^2 dx < \int_{\mathbb{R}^2} \liminf_{k \rightarrow \infty} \left| x + \frac{x_{\rho_k}}{\bar{\varepsilon}_{\rho_k}} \right|^2 Q^2(x) dx \leq \int_{\mathbb{R}^2} |x|^2 Q^2 dx,$$

which is a contradiction. This implies that (3.31) holds true, and the proof of Theorem 1.3 is therefore complete.  $\square$

## 4 Local Uniqueness of Minimizers for $e(\rho)$

Based on the limiting behavior and asymptotic properties of positive minimizers established in Theorem 1.3, this section is devoted to the proof of Theorem 1.2 on the local uniqueness of positive minimizers  $u_\rho$  for  $e(\rho)$  as  $\rho \nearrow \rho^*$ . Throughout the whole section, we denote  $u_k$  to be a positive minimizer of  $e(\rho_k)$  for  $0 < \rho_k < \rho^*$ , and define

$$\varepsilon_k = \frac{2(\rho^* - \rho_k)^{\frac{1}{2}}}{\rho^*} > 0, \quad (4.1)$$

where  $\rho_k \nearrow \rho^*$  as  $k \rightarrow \infty$ . Note that the minimizer  $u_k$  of  $e(\rho_k)$  satisfies

$$-\Delta u_k + \ln(1 + |x|^2) u_k + \left( \int_{\mathbb{R}^2} \ln|x - y| u_k^2(y) dy \right) u_k = \mu_k u_k + u_k^3 \quad \text{in } \mathbb{R}^2, \quad (4.2)$$

where the associated Lagrange multiplier  $\mu_k \in \mathbb{R}$  satisfies

$$\mu_k \rho_k = 2e(\rho_k) + \frac{1}{2} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x - y| u_k^2(x) u_k^2(y) dx dy - \frac{1}{2} \int_{\mathbb{R}^2} u_k^4(x) dx. \quad (4.3)$$

Similar to (3.10), it holds that

$$\mu_k \varepsilon_k^2 \rightarrow -1 \quad \text{as } k \rightarrow \infty, \quad (4.4)$$

where  $\varepsilon_k > 0$  is as in (4.1).

Set

$$w_k(x) := \varepsilon_k u_k(\varepsilon_k x + x_k) > 0 \quad \text{in } \mathbb{R}^2, \quad (4.5)$$

where  $x_k$  is the unique maximal point of  $u_k$  as  $k \rightarrow \infty$ . We then get from (4.2) and (4.5) that  $w_k > 0$  satisfies

$$\begin{aligned} & -\Delta w_k + \varepsilon_k^2 \ln(1 + |\varepsilon_k x + x_k|^2) w_k + \varepsilon_k^2 \left( \int_{\mathbb{R}^2} \ln|x-y| w_k^2(y) dy \right) w_k + \rho_k \varepsilon_k^2 \ln \varepsilon_k w_k \\ & = \mu_k \varepsilon_k^2 w_k + w_k^3 \quad \text{in } \mathbb{R}^2. \end{aligned}$$

Note from Lemma 3.2 (1) that there exist constants  $C > 0$  and  $R > 0$  large enough such that  $w_k$  satisfies

$$|w_k(x)| \leq C e^{-\frac{2|x|}{3}}, \quad |\nabla w_k(x)| \leq C e^{-\frac{|x|}{2}} \quad \text{uniformly for } |x| \geq R \quad (4.6)$$

as  $k \rightarrow \infty$ . We also define the linearized operator  $\mathcal{L}$  of (1.7):

$$\mathcal{L} := -\Delta + (1 - 3Q^2) \quad \text{in } \mathbb{R}^2,$$

where  $Q(x) = Q(|x|) > 0$  is the unique positive solution of (1.7) and satisfies the exponential decay (1.8). Recall from [39, 47] that

$$\ker(\mathcal{L}) = \text{span} \left\{ \frac{\partial Q}{\partial x_1}, \frac{\partial Q}{\partial x_2} \right\}. \quad (4.7)$$

We are now ready to address the local uniqueness of positive minimizers  $u_\rho$  for  $e(\rho)$  as  $\rho \nearrow \rho^*$ .

**Proof of Theorem 1.2.** Argue by contradiction, suppose that there exist two different positive minimizers  $u_{1,k}$  and  $u_{2,k}$  for  $e(\rho_k)$ , where  $\rho_k \nearrow \rho^*$  as  $k \rightarrow \infty$ . Let  $x_{1,k}$  and  $x_{2,k}$  be the unique maximal point of  $u_{1,k}$  and  $u_{2,k}$  as  $k \rightarrow \infty$ , respectively.

Denote

$$V(x) := \ln(1 + |x|^2).$$

Following (4.2), we have

$$-\Delta u_{i,k} + V(x) u_{i,k} + \left( \int_{\mathbb{R}^2} \ln|x-y| u_{i,k}^2(y) dy \right) u_{i,k} = \mu_{i,k} u_{i,k} + u_{i,k}^3 \quad \text{in } \mathbb{R}^2, \quad i = 1, 2. \quad (4.8)$$

Define

$$\bar{u}_{i,k}(x) := \varepsilon_k u_{i,k}(x) > 0 \quad \text{in } \mathbb{R}^2, \quad i = 1, 2. \quad (4.9)$$

One can check that  $\bar{u}_{i,k} > 0$  satisfies

$$-\varepsilon_k^2 \Delta \bar{u}_{i,k} + \varepsilon_k^2 V(x) \bar{u}_{i,k} + \left( \int_{\mathbb{R}^2} \ln|x-y| \bar{u}_{i,k}^2(y) dy \right) \bar{u}_{i,k} = \mu_{i,k} \varepsilon_k^2 \bar{u}_{i,k} + \bar{u}_{i,k}^3 \quad \text{in } \mathbb{R}^2, \quad i = 1, 2. \quad (4.10)$$

Because  $u_{1,k} \not\equiv u_{2,k}$  in  $\mathbb{R}^2$ , we define

$$\bar{\eta}_k(x) := \frac{u_{2,k}(x) - u_{1,k}(x)}{\|u_{2,k} - u_{1,k}\|_{L^\infty(\mathbb{R}^2)}} = \frac{\bar{u}_{2,k}(x) - \bar{u}_{1,k}(x)}{\|\bar{u}_{2,k} - \bar{u}_{1,k}\|_{L^\infty(\mathbb{R}^2)}} \quad \text{in } \mathbb{R}^2. \quad (4.11)$$

It then follows from (4.10) and (4.11) that  $\bar{\eta}_k$  satisfies the following equation:

$$-\varepsilon_k^2 \Delta \bar{\eta}_k + \varepsilon_k^2 V(x) \bar{\eta}_k = \mu_{2,k} \varepsilon_k^2 \bar{\eta}_k + \bar{g}_k(x) + \bar{f}_k(x) \quad \text{in } \mathbb{R}^2, \quad (4.12)$$

where

$$\bar{g}_k(x) = - \left( \int_{\mathbb{R}^2} \ln|x-y| \bar{u}_{2,k}^2(y) dy \right) \bar{\eta}_k + \left( \bar{u}_{2,k}^2 + \bar{u}_{2,k} \bar{u}_{1,k} + \bar{u}_{1,k}^2 \right) \bar{\eta}_k,$$

and

$$\begin{aligned}
\bar{f}_k(x) &= \frac{\mu_{2,k} - \mu_{1,k}}{\|\bar{u}_{2,k} - \bar{u}_{1,k}\|_{L^\infty(\mathbb{R}^2)}} \varepsilon_k^2 \bar{u}_{1,k} - \left[ \int_{\mathbb{R}^2} \ln|x-y| (\bar{u}_{2,k}(y) + \bar{u}_{1,k}(y)) \bar{\eta}_k(y) dy \right] \bar{u}_{1,k} \\
&= \frac{1}{2\rho_k \varepsilon_k^2} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| (\bar{u}_{2,k}(x) + \bar{u}_{1,k}(x)) \bar{\eta}_k(x) \bar{u}_{2,k}^2(y) dx dy \bar{u}_{1,k} \\
&\quad + \frac{1}{2\rho_k \varepsilon_k^2} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| \bar{u}_{1,k}^2(x) (\bar{u}_{2,k}(y) + \bar{u}_{1,k}(y)) \bar{\eta}_k(y) dx dy \bar{u}_{1,k} \\
&\quad - \frac{1}{2\rho_k \varepsilon_k^2} \int_{\mathbb{R}^2} (\bar{u}_{2,k}^2(x) + \bar{u}_{1,k}^2(x)) (\bar{u}_{2,k}(x) + \bar{u}_{1,k}(x)) \bar{\eta}_k(x) dx \bar{u}_{1,k} \\
&\quad - \left[ \int_{\mathbb{R}^2} \ln|x-y| (\bar{u}_{2,k}(y) + \bar{u}_{1,k}(y)) \bar{\eta}_k(y) dy \right] \bar{u}_{1,k}.
\end{aligned}$$

We first claim that for any  $x_0 \in \mathbb{R}^2$ , there exist a small constant  $\delta > 0$  and a constant  $C > 0$  such that

$$\int_{\partial B_\delta(x_0)} \left( \varepsilon_k^2 |\nabla \bar{\eta}_k|^2 + \varepsilon_k^2 V(x) \bar{\eta}_k^2 + \bar{\eta}_k^2 \right) dS \leq C \varepsilon_k^2 \quad \text{as } k \rightarrow \infty. \quad (4.13)$$

To prove the above claim, multiplying (4.12) by  $\bar{\eta}_k$  and integrating over  $\mathbb{R}^2$ , we obtain that

$$\begin{aligned}
&\varepsilon_k^2 \int_{\mathbb{R}^2} |\nabla \bar{\eta}_k|^2 dx + \varepsilon_k^2 \int_{\mathbb{R}^2} V(x) \bar{\eta}_k^2 dx - \mu_{2,k} \varepsilon_k^2 \int_{\mathbb{R}^2} \bar{\eta}_k^2 dx + \frac{1}{2} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln(1+|x-y|^2) \bar{\eta}_k^2(x) \bar{u}_{2,k}^2(y) dx dy \\
&= \frac{1}{2} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln\left(1 + \frac{1}{|x-y|^2}\right) \bar{\eta}_k^2(x) \bar{u}_{2,k}^2(y) dx dy + \int_{\mathbb{R}^2} (\bar{u}_{2,k}^2(x) + \bar{u}_{2,k}(x) \bar{u}_{1,k}(x) + \bar{u}_{1,k}^2(x)) \bar{\eta}_k^2(x) dx \\
&\quad + \frac{1}{2\rho_k \varepsilon_k^2} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| (\bar{u}_{2,k}(x) + \bar{u}_{1,k}(x)) \bar{\eta}_k(x) \bar{u}_{2,k}^2(y) dx dy \int_{\mathbb{R}^2} \bar{u}_{1,k}(x) \bar{\eta}_k(x) dx \\
&\quad + \frac{1}{2\rho_k \varepsilon_k^2} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| \bar{u}_{1,k}^2(x) (\bar{u}_{2,k}(y) + \bar{u}_{1,k}(y)) \bar{\eta}_k(y) dx dy \int_{\mathbb{R}^2} \bar{u}_{1,k}(x) \bar{\eta}_k(x) dx \\
&\quad - \frac{1}{2\rho_k \varepsilon_k^2} \int_{\mathbb{R}^2} (\bar{u}_{2,k}^2(x) + \bar{u}_{1,k}^2(x)) (\bar{u}_{2,k}(x) + \bar{u}_{1,k}(x)) \bar{\eta}_k(x) dx \int_{\mathbb{R}^2} \bar{u}_{1,k}(x) \bar{\eta}_k(x) dx \\
&\quad - \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| \bar{u}_{1,k}(x) \bar{\eta}_k(x) (\bar{u}_{2,k}(y) + \bar{u}_{1,k}(y)) \bar{\eta}_k(y) dx dy.
\end{aligned} \quad (4.14)$$

Lemma A.2 in Appendix shows that there exists a constant  $C > 0$  such that

$$I := \varepsilon_k^2 \int_{\mathbb{R}^2} |\nabla \bar{\eta}_k|^2 dx + \varepsilon_k^2 \int_{\mathbb{R}^2} V(x) \bar{\eta}_k^2 dx + \int_{\mathbb{R}^2} \bar{\eta}_k^2 dx \leq C \varepsilon_k^2 \quad \text{as } k \rightarrow \infty. \quad (4.15)$$

We thus derive from [11, Lemma 4.5] that for any  $x_0 \in \mathbb{R}^2$ , there exist a small constant  $\delta > 0$  and a constant  $M > 0$  such that

$$\int_{\partial B_\delta(x_0)} \left( \varepsilon_k^2 |\nabla \bar{\eta}_k|^2 + \varepsilon_k^2 V(x) \bar{\eta}_k^2 + \bar{\eta}_k^2 \right) dS \leq MI \leq MC \varepsilon_k^2 \quad \text{as } k \rightarrow \infty.$$

We next define

$$\eta_k(x) := \bar{\eta}_k(\varepsilon_k x + x_{2,k}), \quad k = 1, 2, \dots,$$

and

$$\tilde{u}_{i,k}(x) := \varepsilon_k u_{i,k}(\varepsilon_k x + x_{2,k}) > 0, \quad i = 1, 2. \quad (4.16)$$

According to Theorem 1.3, we get that  $\tilde{u}_{i,k} \rightarrow Q$  in  $L^\infty(\mathbb{R}^2)$  as  $k \rightarrow \infty$ . Based on the previous conclusions, we shall carry out the proof of Theorem 1.2 by deriving a contradiction through the following three steps.

Step 1. There exist some constants  $b_0, b_1$  and  $b_2$  such that up to a subsequence if necessary,  $\eta_k \rightarrow \eta_0$  in  $C_{loc}(\mathbb{R}^2)$  as  $k \rightarrow \infty$ , where  $\eta_0(x)$  satisfies

$$\eta_0(x) = b_0(Q + x \cdot \nabla Q) + \sum_{i=1}^2 b_i \frac{\partial Q}{\partial x_i} \quad \text{in } \mathbb{R}^2. \quad (4.17)$$



Note first from (4.8) that  $\eta_k$  satisfies

$$-\Delta\eta_k + C_k(x)\eta_k = D_k(x) \quad \text{in } \mathbb{R}^2, \quad (4.18)$$

where

$$C_k(x) = \varepsilon_k^2 V(\varepsilon_k x + x_{2,k}) + \frac{1}{2}\varepsilon_k^2 \int_{\mathbb{R}^2} \ln(1 + |x - y|^2) \tilde{u}_{2,k}^2(y) dy + \rho_k \varepsilon_k^2 \ln \varepsilon_k - \mu_{2,k} \varepsilon_k^2 + A_k(x), \quad (4.19)$$

$$A_k(x) = -\frac{1}{2}\varepsilon_k^2 \int_{\mathbb{R}^2} \ln\left(1 + \frac{1}{|x - y|^2}\right) \tilde{u}_{2,k}^2(y) dy - (\tilde{u}_{2,k}^2 + \tilde{u}_{2,k} \tilde{u}_{1,k} + \tilde{u}_{1,k}^2), \quad (4.20)$$

and

$$\begin{aligned} D_k(x) &= \frac{\mu_{2,k} - \mu_{1,k}}{\|\tilde{u}_{2,k} - \tilde{u}_{1,k}\|_{L^\infty(\mathbb{R}^2)}} \varepsilon_k^2 \tilde{u}_{1,k} - \varepsilon_k^2 \left[ \int_{\mathbb{R}^2} \ln|x - y| (\tilde{u}_{2,k}(y) + \tilde{u}_{1,k}(y)) \eta_k(y) dy \right] \tilde{u}_{1,k} \\ &= \frac{1}{2\rho_k} \varepsilon_k^2 \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x - y| (\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)) \eta_k(x) \tilde{u}_{2,k}^2(y) dx dy \tilde{u}_{1,k} \\ &\quad + \frac{1}{2\rho_k} \varepsilon_k^2 \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x - y| \tilde{u}_{1,k}^2(x) (\tilde{u}_{2,k}(y) + \tilde{u}_{1,k}(y)) \eta_k(y) dx dy \tilde{u}_{1,k} \\ &\quad - \frac{1}{2\rho_k} \left[ \int_{\mathbb{R}^2} (\tilde{u}_{2,k}^2 + \tilde{u}_{1,k}^2) (\tilde{u}_{2,k} + \tilde{u}_{1,k}) \eta_k dx \right] \tilde{u}_{1,k} \\ &\quad - \varepsilon_k^2 \left[ \int_{\mathbb{R}^2} \ln|x - y| (\tilde{u}_{2,k}(y) + \tilde{u}_{1,k}(y)) \eta_k(y) dy \right] \tilde{u}_{1,k}. \end{aligned} \quad (4.21)$$

Since  $|\eta_k| \leq 1$ , the standard elliptic regularity then implies that  $\|\eta_k\|_{C_{loc}^{1,\alpha}(\mathbb{R}^2)} \leq C$  for some  $\alpha \in (0, 1)$ , where the constant  $C > 0$  is independent of  $k$ , see Lemma A.3 in the Appendix for details. Therefore, there exists a subsequence of  $\{\rho_k\}$  (still denoted by  $\{\rho_k\}$ ) and a function  $\eta_0$  such that  $\eta_k \rightarrow \eta_0$  in  $C_{loc}(\mathbb{R}^2)$  as  $k \rightarrow \infty$ . We further deduce from Lemma A.4 that  $\eta_0$  satisfies

$$\mathcal{L}\eta_0 = -\Delta\eta_0 + (1 - 3Q^2)\eta_0 = -\frac{2}{\rho^*} \left( \int_{\mathbb{R}^2} Q^3 \eta_0 dx \right) Q \quad \text{in } \mathbb{R}^2. \quad (4.22)$$

Since  $\mathcal{L}(Q + x \cdot \nabla Q) = -2Q$ , we conclude from (4.7) and (4.22) that (4.17) holds for some constants  $b_0, b_1$  and  $b_2$ .

Step 2. The constants  $b_0 = b_1 = b_2 = 0$  in (4.17).

We first derive  $b_j = 0$  for  $j = 1, 2$ . Multiplying (4.10) by  $\frac{\partial \bar{u}_{i,k}}{\partial x_j}$ , where  $i, j = 1, 2$ , and integrating over  $B_\delta(x_{2,k})$ , where  $\delta > 0$  is small and given by (4.13), direct computations yield that

$$\begin{aligned} & -\varepsilon_k^2 \int_{B_\delta(x_{2,k})} \frac{\partial \bar{u}_{i,k}}{\partial x_j} \Delta \bar{u}_{i,k} dx + \varepsilon_k^2 \int_{B_\delta(x_{2,k})} V(x) \bar{u}_{i,k} \frac{\partial \bar{u}_{i,k}}{\partial x_j} dx \\ & + \int_{B_\delta(x_{2,k})} \int_{\mathbb{R}^2} \ln|x - y| \bar{u}_{i,k}(x) \frac{\partial \bar{u}_{i,k}(x)}{\partial x_j} \bar{u}_{i,k}^2(y) dy dx \\ & = \mu_{i,k} \varepsilon_k^2 \int_{B_\delta(x_{2,k})} \bar{u}_{i,k} \frac{\partial \bar{u}_{i,k}}{\partial x_j} dx + \int_{B_\delta(x_{2,k})} \bar{u}_{i,k}^3 \frac{\partial \bar{u}_{i,k}}{\partial x_j} dx \\ & = \frac{1}{2} \mu_{i,k} \varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} \bar{u}_{i,k}^2 \nu_j dS + \frac{1}{4} \int_{\partial B_\delta(x_{2,k})} \bar{u}_{i,k}^4 \nu_j dS, \end{aligned} \quad (4.23)$$

where  $\nu := (\nu_1, \nu_2)$  denotes the outward unit normal vector of  $\partial B_\delta(x_{2,k})$ . Note that

$$\begin{aligned} -\varepsilon_k^2 \int_{B_\delta(x_{2,k})} \frac{\partial \bar{u}_{i,k}}{\partial x_j} \Delta \bar{u}_{i,k} dx &= -\varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} \frac{\partial \bar{u}_{i,k}}{\partial x_j} \frac{\partial \bar{u}_{i,k}}{\partial \nu} dS + \varepsilon_k^2 \int_{B_\delta(x_{2,k})} \nabla \bar{u}_{i,k} \cdot \nabla \left( \frac{\partial \bar{u}_{i,k}}{\partial x_j} \right) dx \\ &= -\varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} \frac{\partial \bar{u}_{i,k}}{\partial x_j} \frac{\partial \bar{u}_{i,k}}{\partial \nu} dS + \frac{1}{2} \varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} |\nabla \bar{u}_{i,k}|^2 \nu_j dS, \end{aligned} \quad (4.24)$$

$$\varepsilon_k^2 \int_{B_\delta(x_{2,k})} V(x) \bar{u}_{i,k} \frac{\partial \bar{u}_{i,k}}{\partial x_j} dx = \frac{1}{2} \varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} V(x) \bar{u}_{i,k}^2 \nu_j dS - \frac{1}{2} \varepsilon_k^2 \int_{B_\delta(x_{2,k})} \frac{\partial V(x)}{\partial x_j} \bar{u}_{i,k}^2 dx, \quad (4.25)$$

and

$$\begin{aligned}
& \int_{B_\delta(x_{2,k})} \int_{\mathbb{R}^2} \ln|x-y| \bar{u}_{i,k}(x) \frac{\partial \bar{u}_{i,k}(x)}{\partial x_j} \bar{u}_{i,k}^2(y) dy dx \\
&= \frac{1}{2} \int_{\partial B_\delta(x_{2,k})} \int_{\mathbb{R}^2} \ln|x-y| \bar{u}_{i,k}^2(y) \bar{u}_{i,k}^2(x) \nu_j dy dS - \frac{1}{2} \int_{B_\delta(x_{2,k})} \int_{\mathbb{R}^2} \frac{x_j - y_j}{|x-y|^2} \bar{u}_{i,k}^2(x) \bar{u}_{i,k}^2(y) dy dx \\
&= \frac{1}{2} \int_{\partial B_\delta(x_{2,k})} \int_{\mathbb{R}^2} \ln|x-y| \bar{u}_{i,k}^2(y) \bar{u}_{i,k}^2(x) \nu_j dy dS + \frac{1}{2} \int_{B_\delta^c(x_{2,k})} \int_{\mathbb{R}^2} \frac{x_j - y_j}{|x-y|^2} \bar{u}_{i,k}^2(x) \bar{u}_{i,k}^2(y) dy dx,
\end{aligned} \tag{4.26}$$

where we have used the following fact

$$\int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{x_j - y_j}{|x-y|^2} \bar{u}_{i,k}^2(x) \bar{u}_{i,k}^2(y) dx dy = 0.$$

Submitting (4.24)–(4.26) into (4.23), we then deduce that

$$\begin{aligned}
& \varepsilon_k^2 \int_{B_\delta(x_{2,k})} \frac{\partial V(x)}{\partial x_j} \bar{u}_{i,k}^2(x) dx \\
&= -2\varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} \frac{\partial \bar{u}_{i,k}}{\partial x_j} \frac{\partial \bar{u}_{i,k}}{\partial \nu} dS + \varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} |\nabla \bar{u}_{i,k}|^2 \nu_j dS + \varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} V(x) \bar{u}_{i,k}^2 \nu_j dS \\
&+ \int_{\partial B_\delta(x_{2,k})} \int_{\mathbb{R}^2} \ln|x-y| \bar{u}_{i,k}^2(y) \bar{u}_{i,k}^2(x) \nu_j dy dS + \int_{B_\delta^c(x_{2,k})} \int_{\mathbb{R}^2} \frac{x_j - y_j}{|x-y|^2} \bar{u}_{i,k}^2(x) \bar{u}_{i,k}^2(y) dy dx \\
&- \mu_{i,k} \varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} \bar{u}_{i,k}^2 \nu_j dS - \frac{1}{2} \int_{\partial B_\delta(x_{2,k})} \bar{u}_{i,k}^4 \nu_j dS.
\end{aligned} \tag{4.27}$$

Following (4.27), it yields that

$$\begin{aligned}
& \varepsilon_k^2 \int_{B_\delta(x_{2,k})} \frac{\partial V(x)}{\partial x_j} (\bar{u}_{2,k}(x) + \bar{u}_{1,k}(x)) \bar{\eta}_k dx \\
&= -2\varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} \left( \frac{\partial \bar{\eta}_k}{\partial x_j} \frac{\partial \bar{u}_{2,k}}{\partial \nu} + \frac{\partial \bar{u}_{1,k}}{\partial x_j} \frac{\partial \bar{\eta}_k}{\partial \nu} \right) dS \\
&+ \varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} (\nabla \bar{u}_{2,k} + \nabla \bar{u}_{1,k}) \cdot \nabla \bar{\eta}_k \nu_j dS \\
&+ \varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} V(x) (\bar{u}_{2,k} + \bar{u}_{1,k}) \bar{\eta}_k \nu_j dS \\
&+ \int_{\partial B_\delta(x_{2,k})} \int_{\mathbb{R}^2} \ln|x-y| (\bar{u}_{2,k}(y) + \bar{u}_{1,k}(y)) \bar{\eta}_k(y) \bar{u}_{2,k}^2(x) \nu_j dy dS \\
&+ \int_{\partial B_\delta(x_{2,k})} \int_{\mathbb{R}^2} \ln|x-y| \bar{u}_{1,k}^2(y) (\bar{u}_{2,k}(x) + \bar{u}_{1,k}(x)) \bar{\eta}_k(x) \nu_j dy dS \\
&+ \int_{B_\delta^c(x_{2,k})} \int_{\mathbb{R}^2} \frac{x_j - y_j}{|x-y|^2} (\bar{u}_{2,k}(x) + \bar{u}_{1,k}(x)) \bar{\eta}_k(x) \bar{u}_{2,k}^2(y) dy dx \\
&+ \int_{B_\delta^c(x_{2,k})} \int_{\mathbb{R}^2} \frac{x_j - y_j}{|x-y|^2} \bar{u}_{1,k}^2(x) (\bar{u}_{2,k}(y) + \bar{u}_{1,k}(y)) \bar{\eta}_k(y) dy dx \\
&- \varepsilon_k^2 \mu_{2,k} \int_{\partial B_\delta(x_{2,k})} (\bar{u}_{2,k}(x) + \bar{u}_{1,k}(x)) \bar{\eta}_k(x) \nu_j dS \\
&- \frac{1}{2} \int_{\partial B_\delta(x_{2,k})} (\bar{u}_{2,k}^2 + \bar{u}_{1,k}^2) (\bar{u}_{2,k} + \bar{u}_{1,k}) \bar{\eta}_k \nu_j dS \\
&- \frac{\mu_{2,k} - \mu_{1,k}}{\|\bar{u}_{2,k} - \bar{u}_{1,k}\|_{L^\infty(\mathbb{R}^2)}} \varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} \bar{u}_{1,k}^2 \nu_j dS := \alpha_{j,k}.
\end{aligned} \tag{4.28}$$

Since  $V(x) = \ln(1 + |x|^2)$ , according to Lemma A.5, it follows from (3.31), (4.6) and (4.28) that there exists a constant  $a > 0$  such that

$$\begin{aligned}
o(e^{-\frac{a\delta}{\varepsilon_k}}) &= \varepsilon_k^2 \int_{B_\delta(x_{2,k})} \frac{\partial V(x)}{\partial x_j} (\tilde{u}_{2,k} + \bar{u}_{1,k}) \bar{\eta}_k dx \\
&= \varepsilon_k^4 \int_{B_{\frac{\delta}{\varepsilon_k}}(0)} \frac{\partial V(\varepsilon_k x + x_{2,k})}{\partial x_j} (\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)) \eta_k(x) dx \\
&= 2\varepsilon_k^4 \int_{B_{\frac{\delta}{\varepsilon_k}}(0)} \frac{\varepsilon_k x_j + x_{2,k}^j}{1 + |\varepsilon_k x + x_{2,k}|^2} (\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)) \eta_k(x) dx \\
&= O(\varepsilon_k^5) \int_{B_{\frac{\delta}{\varepsilon_k}}(0)} \left(x_j + \frac{x_{2,k}^j}{\varepsilon_k}\right) (\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)) \eta_k(x) dx \\
&= O(\varepsilon_k^5) \int_{B_{\frac{\delta}{\varepsilon_k}}(0)} \left[ \left(x_j + \frac{x_{2,k}^j}{\varepsilon_k}\right) (\tilde{u}_{2,k}(x) + \bar{u}_{1,k}(\varepsilon_k x + x_{1,k})) \eta_k(x) \right. \\
&\quad \left. + \left(x_j + \frac{x_{2,k}^j}{\varepsilon_k}\right) (\bar{u}_{1,k}(\varepsilon_k x + x_{2,k}) - \bar{u}_{1,k}(\varepsilon_k x + x_{1,k})) \eta_k(x) \right] dx \\
&= O(\varepsilon_k^5) \int_{B_{\frac{\delta}{\varepsilon_k}}(0)} \left(x_j + \frac{x_{2,k}^j}{\varepsilon_k}\right) (\tilde{u}_{2,k}(x) + \bar{u}_{1,k}(\varepsilon_k x + x_{1,k})) \eta_k(x) dx \quad \text{as } k \rightarrow \infty,
\end{aligned} \tag{4.29}$$

where  $x_{2,k} := (x_{2,k}^1, x_{2,k}^2) \in \mathbb{R}^2$ .

Since  $\eta_k \rightarrow \eta_0$  in  $C_{loc}(\mathbb{R}^2)$  as  $k \rightarrow \infty$ , in view of the exponential decays (1.8) and (A.11), we derive from (3.31), (4.6) and (A.12) that

$$\begin{aligned}
&\left| \int_{B_{\frac{\delta}{\varepsilon_k}}(0)} \left(x_j + \frac{x_{2,k}^j}{\varepsilon_k}\right) (\tilde{u}_{2,k}(x) + \bar{u}_{1,k}(\varepsilon_k x + x_{1,k})) \eta_k(x) dx - 2 \int_{\mathbb{R}^2} x_j Q(x) \eta_0(x) dx \right| \\
&\leq \left| \frac{x_{2,k}^j}{\varepsilon_k} \right| \int_{\mathbb{R}^2} |(\tilde{u}_{2,k}(x) + \bar{u}_{1,k}(\varepsilon_k x + x_{1,k})) \eta_k(x)| dx + \int_{\mathbb{R}^2} |x_j| |\tilde{u}_{2,k}(x) + \bar{u}_{1,k}(\varepsilon_k x + x_{1,k}) - 2Q(x)| |\eta_k(x)| dx \\
&\quad + 2 \int_{\mathbb{R}^2} |x_j| |Q| |\eta_k - \eta_0| dx + \int_{B_{\delta/\varepsilon_k}^c(0)} \left| \left(x_j + \frac{x_{2,k}^j}{\varepsilon_k}\right) (\tilde{u}_{2,k}(x) + \bar{u}_{1,k}(\varepsilon_k x + x_{1,k})) \eta_k(x) \right| dx \rightarrow 0 \quad \text{as } k \rightarrow \infty,
\end{aligned}$$

which implies that

$$\int_{B_{\frac{\delta}{\varepsilon_k}}(0)} \left(x_j + \frac{x_{2,k}^j}{\varepsilon_k}\right) (\tilde{u}_{2,k}(x) + \bar{u}_{1,k}(\varepsilon_k x + x_{1,k})) \eta_k(x) dx \rightarrow 2 \int_{\mathbb{R}^2} x_j Q(x) \eta_0(x) dx \quad \text{as } k \rightarrow \infty. \tag{4.30}$$

We thus deduce from (4.17), (4.29) and (4.30) that

$$0 = 2 \int_{\mathbb{R}^2} x_j Q \eta_0 dx = 2 \int_{\mathbb{R}^2} x_j Q \left[ b_0(Q + x \cdot \nabla Q) + \sum_{i=1}^2 b_i \frac{\partial Q}{\partial x_i} \right] dx = -b_j \int_{\mathbb{R}^2} Q^2 dx = -b_j \rho^*, \quad j = 1, 2,$$

which means that  $b_j = 0$  for  $j = 1, 2$ .

We next prove  $b_0 = 0$ . Using integration by parts, we get that

$$\begin{aligned}
&-\varepsilon_k^2 \int_{B_\delta(x_{2,k})} \Delta \bar{u}_{i,k} \left[ (x - x_{2,k}) \cdot \nabla \bar{u}_{i,k} \right] dx \\
&= -\varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} \frac{\partial \bar{u}_{i,k}}{\partial \nu} \left[ (x - x_{2,k}) \cdot \nabla \bar{u}_{i,k} \right] dS + \varepsilon_k^2 \int_{B_\delta(x_{2,k})} \nabla \bar{u}_{i,k} \cdot \nabla \left[ (x - x_{2,k}) \cdot \nabla \bar{u}_{i,k} \right] dx \\
&= -\varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} \frac{\partial \bar{u}_{i,k}}{\partial \nu} \left[ (x - x_{2,k}) \cdot \nabla \bar{u}_{i,k} \right] dS + \frac{1}{2} \varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} (x - x_{2,k}) \cdot \nu |\nabla \bar{u}_{i,k}|^2 dS.
\end{aligned} \tag{4.31}$$

Multiplying (4.10) by  $(x - x_{2,k}) \cdot \nabla \bar{u}_{i,k}$ ,  $i = 1, 2$ , and integrating over  $B_\delta(x_{2,k})$ , where  $\delta > 0$  is as in (4.13), we derive that

$$\begin{aligned}
& -\varepsilon_k^2 \int_{B_\delta(x_{2,k})} \Delta \bar{u}_{i,k} \left[ (x - x_{2,k}) \cdot \nabla \bar{u}_{i,k} \right] dx \\
& = \varepsilon_k^2 \int_{B_\delta(x_{2,k})} \left( \mu_{i,k} - V(x) \right) \bar{u}_{i,k}(x) \left[ (x - x_{2,k}) \cdot \nabla \bar{u}_{i,k}(x) \right] dx \\
& \quad - \int_{B_\delta(x_{2,k})} \int_{\mathbb{R}^2} \ln|x-y| \bar{u}_{i,k}(x) \left[ (x - x_{2,k}) \cdot \nabla \bar{u}_{i,k}(x) \right] \bar{u}_{i,k}^2(y) dy dx \\
& \quad + \int_{B_\delta(x_{2,k})} \bar{u}_{i,k}^3(x) \left[ (x - x_{2,k}) \cdot \nabla \bar{u}_{i,k}(x) \right] dx \\
& = -\frac{1}{2} \varepsilon_k^2 \int_{B_\delta(x_{2,k})} \left[ 2(\mu_{i,k} - V(x)) - (x - x_{2,k}) \cdot \nabla V(x) \right] \bar{u}_{i,k}^2(x) dx \\
& \quad + \frac{1}{2} \varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} (\mu_{i,k} - V(x)) (x - x_{2,k}) \cdot \nu \bar{u}_{i,k}^2(x) dS \\
& \quad + \frac{1}{2} \int_{B_\delta(x_{2,k})} \int_{\mathbb{R}^2} \frac{(x - x_{2,k}) \cdot (x - y)}{|x - y|^2} \bar{u}_{i,k}^2(x) \bar{u}_{i,k}^2(y) dy dx \\
& \quad + \int_{B_\delta(x_{2,k})} \int_{\mathbb{R}^2} \ln|x-y| \bar{u}_{i,k}^2(x) \bar{u}_{i,k}^2(y) dy dx \\
& \quad - \frac{1}{2} \int_{\partial B_\delta(x_{2,k})} \int_{\mathbb{R}^2} \ln|x-y| \bar{u}_{i,k}^2(x) (x - x_{2,k}) \cdot \nu \bar{u}_{i,k}^2(y) dy dS \\
& \quad - \frac{1}{2} \int_{B_\delta(x_{2,k})} \bar{u}_{i,k}^4(x) dx + \frac{1}{4} \int_{\partial B_\delta(x_{2,k})} (x - x_{2,k}) \cdot \nu \bar{u}_{i,k}^4(x) dS,
\end{aligned} \tag{4.32}$$

where  $\nu$  denotes the outward unit normal vector of  $\partial B_\delta(x_{2,k})$ . Combining (4.31) with (4.32) gives that

$$\begin{aligned}
& -\varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} \frac{\partial \bar{u}_{i,k}}{\partial \nu} \left[ (x - x_{2,k}) \cdot \nabla \bar{u}_{i,k} \right] dS + \frac{1}{2} \varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} (x - x_{2,k}) \cdot \nu |\nabla \bar{u}_{i,k}|^2 dS \\
& = -\varepsilon_k^2 \mu_{i,k} \int_{B_\delta(x_{2,k})} \bar{u}_{i,k}^2 dx + \varepsilon_k^2 \int_{B_\delta(x_{2,k})} V(x) \bar{u}_{i,k}^2 dx + \frac{1}{2} \varepsilon_k^2 \int_{B_\delta(x_{2,k})} (x - x_{2,k}) \cdot \nabla V(x) \bar{u}_{i,k}^2 dx \\
& \quad + \frac{\varepsilon_k^2}{2} \int_{\partial B_\delta(x_{2,k})} (\mu_{i,k} - V(x)) (x - x_{2,k}) \cdot \nu \bar{u}_{i,k}^2(x) dS \\
& \quad + \frac{1}{2} \int_{B_\delta(x_{2,k})} \int_{\mathbb{R}^2} \frac{(x - x_{2,k}) \cdot (x - y)}{|x - y|^2} \bar{u}_{i,k}^2(x) \bar{u}_{i,k}^2(y) dy dx \\
& \quad + \int_{B_\delta(x_{2,k})} \int_{\mathbb{R}^2} \ln|x-y| \bar{u}_{i,k}^2(x) \bar{u}_{i,k}^2(y) dy dx - \frac{1}{2} \int_{\partial B_\delta(x_{2,k})} \int_{\mathbb{R}^2} \ln|x-y| \bar{u}_{i,k}^2(x) (x - x_{2,k}) \cdot \nu \bar{u}_{i,k}^2(y) dy dS \\
& \quad - \frac{1}{2} \int_{B_\delta(x_{2,k})} \bar{u}_{i,k}^4 dx + \frac{1}{4} \int_{\partial B_\delta(x_{2,k})} (x - x_{2,k}) \cdot \nu \bar{u}_{i,k}^4 dS \\
& = -\varepsilon_k^2 \mu_{i,k} \int_{\mathbb{R}^2} \bar{u}_{i,k}^2 dx + \varepsilon_k^2 \int_{B_\delta(x_{2,k})} V(x) \bar{u}_{i,k}^2 dx + \frac{1}{2} \varepsilon_k^2 \int_{B_\delta(x_{2,k})} x \cdot \nabla V(x) \bar{u}_{i,k}^2(x) dx \\
& \quad - \frac{1}{2} \varepsilon_k^2 \int_{B_\delta(x_{2,k})} x_{2,k} \cdot \nabla V(x) \bar{u}_{i,k}^2(x) dx + \frac{1}{2} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{(x - x_{2,k}) \cdot (x - y)}{|x - y|^2} \bar{u}_{i,k}^2(x) \bar{u}_{i,k}^2(y) dx dy \\
& \quad + \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| \bar{u}_{i,k}^2(x) \bar{u}_{i,k}^2(y) dx dy - \frac{1}{2} \int_{\mathbb{R}^2} \bar{u}_{i,k}^4 dx \\
& \quad + \varepsilon_k^2 \mu_{i,k} \int_{B_\delta^c(x_{2,k})} \bar{u}_{i,k}^2(x) dx + \frac{1}{2} \varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} (\mu_{i,k} - V(x)) \bar{u}_{i,k}^2(x) (x - x_{2,k}) \cdot \nu dS
\end{aligned} \tag{4.33}$$

$$\begin{aligned}
& -\frac{1}{2} \int_{B_\delta^c(x_{2,k})} \int_{\mathbb{R}^2} \frac{(x-x_{2,k}) \cdot (x-y)}{|x-y|^2} \bar{u}_{i,k}^2(x) \bar{u}_{i,k}^2(y) dy dx - \int_{B_\delta^c(x_{2,k})} \int_{\mathbb{R}^2} \ln|x-y| \bar{u}_{i,k}^2(x) \bar{u}_{i,k}^2(y) dy dx \\
& -\frac{1}{2} \int_{\partial B_\delta(x_{2,k})} \int_{\mathbb{R}^2} \ln|x-y| \bar{u}_{i,k}^2(y) \bar{u}_{i,k}^2(x) (x-x_{2,k}) \cdot \nu dy dS \\
& +\frac{1}{2} \int_{B_\delta^c(x_{2,k})} \bar{u}_{i,k}^4 dx + \frac{1}{4} \int_{\partial B_\delta(x_{2,k})} \bar{u}_{i,k}^4(x) (x-x_{2,k}) \cdot \nu dS.
\end{aligned}$$

Note that

$$\begin{aligned}
& \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{(x-x_{2,k}) \cdot (x-y)}{|x-y|^2} \bar{u}_{i,k}^2(x) \bar{u}_{i,k}^2(y) dx dy \\
& = \varepsilon_k^4 \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{(x-y) \cdot x}{|x-y|^2} \tilde{u}_{i,k}^2(x) \tilde{u}_{i,k}^2(y) dx dy \\
& = \frac{1}{2} \varepsilon_k^4 \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{(x-y) \cdot x + (y-x) \cdot y}{|x-y|^2} \tilde{u}_{i,k}^2(x) \tilde{u}_{i,k}^2(y) dx dy \\
& = \frac{1}{2} \varepsilon_k^4 \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \tilde{u}_{i,k}^2(x) \tilde{u}_{i,k}^2(y) dx dy = \frac{1}{2} \rho_k^2 \varepsilon_k^4.
\end{aligned}$$

We then infer from (4.33) that

$$\begin{aligned}
& -\varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} \frac{\partial \bar{u}_{i,k}}{\partial \nu} \left[ (x-x_{2,k}) \cdot \nabla \bar{u}_{i,k} \right] dS + \frac{1}{2} \varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} |\nabla \bar{u}_{i,k}|^2 (x-x_{2,k}) \cdot \nu dS \\
& = -\varepsilon_k^2 \mu_{i,k} \int_{\mathbb{R}^2} \bar{u}_{i,k}^2 dx + \varepsilon_k^2 \int_{B_\delta(x_{2,k})} V(x) \bar{u}_{i,k}^2 dx + \frac{1}{2} \varepsilon_k^2 \int_{B_\delta(x_{2,k})} x \cdot \nabla V(x) \bar{u}_{i,k}^2 dx \\
& -\frac{1}{2} \varepsilon_k^2 \int_{B_\delta(x_{2,k})} x_{2,k} \cdot \nabla V(x) \bar{u}_{i,k}^2(x) dx + \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| \bar{u}_{i,k}^2(x) \bar{u}_{i,k}^2(y) dx dy \\
& -\frac{1}{2} \int_{\mathbb{R}^2} \bar{u}_{i,k}^4 dx + \frac{1}{4} \rho_k^2 \varepsilon_k^4 + \beta_{i,k},
\end{aligned} \tag{4.34}$$

where  $\beta_{i,k}$  satisfies

$$\begin{aligned}
\beta_{i,k} & = \varepsilon_k^2 \mu_{i,k} \int_{B_\delta^c(x_{2,k})} \bar{u}_{i,k}^2(x) dx + \frac{1}{2} \varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} (\mu_{i,k} - V(x)) \bar{u}_{i,k}^2(x) (x-x_{2,k}) \cdot \nu dS \\
& -\frac{1}{2} \int_{B_\delta^c(x_{2,k})} \int_{\mathbb{R}^2} \frac{(x-x_{2,k}) \cdot (x-y)}{|x-y|^2} \bar{u}_{i,k}^2(x) \bar{u}_{i,k}^2(y) dy dx \\
& -\int_{B_\delta^c(x_{2,k})} \int_{\mathbb{R}^2} \ln|x-y| \bar{u}_{i,k}^2(x) \bar{u}_{i,k}^2(y) dy dx \\
& -\frac{1}{2} \int_{\partial B_\delta(x_{2,k})} \int_{\mathbb{R}^2} \ln|x-y| \bar{u}_{i,k}^2(x) \bar{u}_{i,k}^2(y) (x-x_{2,k}) \cdot \nu dy dS \\
& +\frac{1}{2} \int_{B_\delta^c(x_{2,k})} \bar{u}_{i,k}^4 dx + \frac{1}{4} \int_{\partial B_\delta(x_{2,k})} \bar{u}_{i,k}^4(x) (x-x_{2,k}) \cdot \nu dS.
\end{aligned} \tag{4.35}$$

Noting from (4.3) that

$$\frac{1}{2} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| \bar{u}_{i,k}^2(x) \bar{u}_{i,k}^2(y) dx dy - \frac{1}{2} \int_{\mathbb{R}^2} \bar{u}_{i,k}^4 dx - \varepsilon_k^4 \mu_{i,k} \rho_k = -2\varepsilon_k^4 e(\rho_k),$$

we obtain from (4.34) that

$$\begin{aligned}
& 2\varepsilon_k^4 e(\rho_k) - \varepsilon_k^2 \int_{B_\delta(x_{2,k})} V(x) \bar{u}_{i,k}^2 dx - \frac{1}{2} \varepsilon_k^2 \int_{B_\delta(x_{2,k})} x \cdot \nabla V(x) \bar{u}_{i,k}^2 dx - \frac{1}{4} \rho_k^2 \varepsilon_k^4 \\
& -\frac{1}{2} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| \bar{u}_{i,k}^2(x) \bar{u}_{i,k}^2(y) dx dy
\end{aligned} \tag{4.36}$$

$$\begin{aligned}
&= \varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} \frac{\partial \bar{u}_{i,k}}{\partial \nu} \left[ (x - x_{2,k}) \cdot \nabla \bar{u}_{i,k} \right] dS - \frac{1}{2} \varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} |\nabla \bar{u}_{i,k}|^2 (x - x_{2,k}) \cdot \nu dS \\
&\quad - \frac{1}{2} \varepsilon_k^2 \int_{B_\delta(x_{2,k})} x_{2,k} \cdot \nabla V(x) \bar{u}_{i,k}^2 dx + \beta_{i,k}.
\end{aligned}$$

Since

$$\int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| \bar{u}_{i,k}^2(x) \bar{u}_{i,k}^2(y) dx dy = \varepsilon_k^4 \ln \varepsilon_k \rho_k^2 + \varepsilon_k^4 \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| \tilde{u}_{i,k}^2(x) \tilde{u}_{i,k}^2(y) dx dy,$$

we thus deduce from (4.13) and (4.36) that

$$\begin{aligned}
\gamma_k &:= -\varepsilon_k^2 \int_{B_\delta(x_{2,k})} V(x) (\bar{u}_{2,k} + \bar{u}_{1,k}) \bar{\eta}_k dx - \frac{1}{2} \varepsilon_k^2 \int_{B_\delta(x_{2,k})} x \cdot \nabla V(x) (\bar{u}_{2,k} + \bar{u}_{1,k}) \bar{\eta}_k dx \\
&\quad - \frac{1}{2} \varepsilon_k^4 \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| (\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)) \eta_k(x) \tilde{u}_{2,k}^2(y) dx dy \\
&\quad - \frac{1}{2} \varepsilon_k^4 \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| \tilde{u}_{1,k}^2(x) (\tilde{u}_{2,k}(y) + \tilde{u}_{1,k}(y)) \eta_k(y) dx dy \\
&= \frac{\beta_{2,k} - \beta_{1,k}}{\|\bar{u}_{2,k} - \bar{u}_{1,k}\|_{L^\infty(\mathbb{R}^2)}} - \frac{1}{2} \varepsilon_k^2 \int_{B_\delta(x_{2,k})} x_{2,k} \cdot \nabla V(x) (\bar{u}_{2,k} + \bar{u}_{1,k}) \bar{\eta}_k dx \\
&\quad + \varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} \frac{\partial \bar{\eta}_k}{\partial \nu} \left[ (x - x_{2,k}) \cdot \nabla \bar{u}_{2,k} \right] dS + \varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} \frac{\partial \bar{u}_{1,k}}{\partial \nu} \left[ (x - x_{2,k}) \cdot \nabla \bar{\eta}_k \right] dS \\
&\quad - \frac{1}{2} \varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} (\nabla \bar{u}_{2,k} + \nabla \bar{u}_{1,k}) \cdot \nabla \bar{\eta}_k (x - x_{2,k}) \cdot \nu dS \\
&= \frac{\beta_{2,k} - \beta_{1,k}}{\|\bar{u}_{2,k} - \bar{u}_{1,k}\|_{L^\infty(\mathbb{R}^2)}} - \frac{1}{2} \varepsilon_k^2 \int_{B_\delta(x_{2,k})} x_{2,k} \cdot \nabla V(x) (\bar{u}_{2,k} + \bar{u}_{1,k}) \bar{\eta}_k dx + o(e^{-\frac{a\delta}{\varepsilon_k}}),
\end{aligned} \tag{4.37}$$

where the exponential decay of  $|\nabla \tilde{u}_{i,k}|$  as  $|x| \rightarrow \infty$  is used.

Furthermore, applying Lemma A.6, we conclude from (4.37) that there exists a constant  $b \in (0, a)$  such that

$$\begin{aligned}
&\frac{1}{2} \varepsilon_k^4 \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| (\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)) \eta_k(x) \tilde{u}_{2,k}^2(y) dx dy \\
&\quad + \frac{1}{2} \varepsilon_k^4 \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| \tilde{u}_{1,k}^2(x) (\tilde{u}_{2,k}(y) + \tilde{u}_{1,k}(y)) \eta_k(y) dx dy + o(e^{-\frac{b\delta}{\varepsilon_k}}) \\
&= -\varepsilon_k^2 \int_{B_\delta(x_{2,k})} V(x) (\bar{u}_{2,k} + \bar{u}_{1,k}) \bar{\eta}_k dx - \frac{1}{2} \varepsilon_k^2 \int_{B_\delta(x_{2,k})} x \cdot \nabla V(x) (\bar{u}_{2,k} + \bar{u}_{1,k}) \bar{\eta}_k dx \\
&= -\varepsilon_k^4 \int_{B_{\frac{\delta}{\varepsilon_k}}(0)} V(\varepsilon_k x + x_{2,k}) (\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)) \eta_k(x) dx \\
&\quad - \frac{1}{2} \varepsilon_k^4 \int_{B_{\frac{\delta}{\varepsilon_k}}(0)} (\varepsilon_k x + x_{2,k}) \cdot \nabla V(\varepsilon_k x + x_{2,k}) (\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)) \eta_k(x) dx \\
&= -\varepsilon_k^4 \int_{B_{\frac{\delta}{\varepsilon_k}}(0)} \ln\left(1 + |\varepsilon_k x + x_{2,k}|^2\right) (\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)) \eta_k(x) dx \\
&\quad - \varepsilon_k^4 \int_{B_{\frac{\delta}{\varepsilon_k}}(0)} \frac{|\varepsilon_k x + x_{2,k}|^2}{1 + |\varepsilon_k x + x_{2,k}|^2} (\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)) \eta_k(x) dx \\
&= O(\varepsilon_k^6) \int_{B_{\frac{\delta}{\varepsilon_k}}(0)} \left|x + \frac{x_{2,k}}{\varepsilon_k}\right|^2 (\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)) \eta_k(x) dx \quad \text{as } k \rightarrow \infty.
\end{aligned} \tag{4.38}$$

Since  $\eta_k \rightarrow \eta_0$  in  $C_{loc}(\mathbb{R}^2)$  as  $k \rightarrow \infty$ , in view of the exponential decay (1.8) and (A.11), we derive from

(3.31) and (A.12) that

$$\int_{B_{\frac{\delta}{\varepsilon_k}}(0)} \left| x + \frac{x_{2,k}}{\varepsilon_k} \right|^2 (\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)) \eta_k(x) dx \rightarrow 2 \int_{\mathbb{R}^2} |x|^2 Q(x) \eta_0(x) dx \quad \text{as } k \rightarrow \infty.$$

Together with (4.38), this yields that

$$\begin{aligned} & \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| \eta_k(x) (\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)) \tilde{u}_{2,k}^2(y) dx dy \\ & + \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| \tilde{u}_{1,k}^2(x) \eta_k(y) (\tilde{u}_{2,k}(y) + \tilde{u}_{1,k}(y)) dx dy = O(\varepsilon_k^2) \quad \text{as } k \rightarrow \infty. \end{aligned} \quad (4.39)$$

Since

$$\int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{(x-y) \cdot x}{|x-y|^2} Q^2(x) Q^2(y) dx dy = \frac{1}{2} (\rho^*)^2,$$

applying Lemma A.7, we derive from (4.17) and (4.39) that

$$\begin{aligned} 0 &= 2 \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| \eta_0(x) Q(x) Q^2(y) dx dy \\ &= 2b_0 \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| Q(x) [Q(x) + x \cdot \nabla Q(x)] Q^2(y) dx dy \\ &= -b_0 \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{(x-y) \cdot x}{|x-y|^2} Q^2(x) Q^2(y) dx dy = -\frac{1}{2} b_0 (\rho^*)^2, \end{aligned}$$

which implies that  $b_0 = 0$ . Step 2 is thus proved.

Step 3.  $\eta_0 \equiv 0$  cannot occur.

By the exponential decay of  $\tilde{u}_{i,k}$  for  $i = 1, 2$ , there exists a point  $z_k \in \mathbb{R}^2$  such that  $|\eta_k(z_k)| = \|\eta_k\|_\infty = 1$ . Since both  $\tilde{u}_{i,k}$  and  $Q$  decay exponentially as  $|x| \rightarrow \infty$ , applying the maximal principle to (4.18) yields that  $|z_k| \leq C$  holds uniformly in  $k$ . Therefore, we have  $\eta_k \rightarrow \eta_0 \not\equiv 0$  in  $C_{loc}(\mathbb{R}^2)$  as  $k \rightarrow \infty$ , which however contradicts with the fact that  $\eta_0 \equiv 0$  in  $\mathbb{R}^2$ .  $\square$

## A Appendix

For the reader's convenience, in this appendix we shall give the detailed proofs of some results used in Theorems 1.2 and 1.3.

**Lemma A.1.** *Let  $u_\rho > 0$  be a positive minimizer of  $e(\rho)$  for  $\rho \in (0, \rho^*)$ , and define*

$$w_\rho(x) := \varepsilon_\rho u_\rho(\varepsilon_\rho x + x_\rho) > 0 \quad \text{in } \mathbb{R}^2,$$

where  $\varepsilon_\rho > 0$  is as in (3.1) and  $x_\rho$  is a maximal point of  $u_\rho$ . Then there exists a large constant  $R > 0$  such that

$$|\nabla w_\rho(x)| \leq C e^{-\frac{|x|}{2\sqrt{\rho^*}}} \quad \text{uniformly for } |x| \geq R \quad \text{as } \rho \nearrow \rho^*.$$

**Proof.** Denoting  $\partial_j w_\rho = \partial w_\rho / \partial x_j$  for  $j = 1, 2$ , it follows from (3.11) that

$$\begin{aligned} & -\Delta \partial_j w_\rho + \varepsilon_\rho^2 \ln[1 + |\varepsilon_\rho x + x_\rho|^2] \partial_j w_\rho + \frac{2\varepsilon_\rho^3 (\varepsilon_\rho x_j + x_\rho^j)}{1 + |\varepsilon_\rho x + x_\rho|^2} w_\rho \\ & + \varepsilon_\rho^2 \left( \int_{\mathbb{R}^2} \ln|x-y| w_\rho^2(y) dy \right) \partial_j w_\rho + \varepsilon_\rho^2 \left( \int_{\mathbb{R}^2} \frac{x_j - y_j}{|x-y|^2} w_\rho^2(y) dy \right) w_\rho \\ & + \rho \varepsilon_\rho^2 \ln \varepsilon_\rho \partial_j w_\rho = \mu_\rho \varepsilon_\rho^2 \partial_j w_\rho + 3w_\rho^2 \partial_j w_\rho \quad \text{in } \mathbb{R}^2, \end{aligned} \quad (\text{A.1})$$

where  $x_\rho := (x_\rho^1, x_\rho^2) \in \mathbb{R}^2$ . Multiplying (A.1) by  $\partial_j w_\rho$  yields that

$$\begin{aligned}
& -\frac{1}{2}\Delta |\partial_j w_\rho|^2 + |\nabla \partial_j w_\rho|^2 + \varepsilon_\rho^2 \ln[1 + |\varepsilon_\rho x + x_\rho|^2] |\partial_j w_\rho|^2 \\
& + \frac{2\varepsilon_\rho^3 (\varepsilon_\rho x_j + x_\rho^j)}{1 + |\varepsilon_\rho x + x_\rho|^2} w_\rho \partial_j w_\rho + \varepsilon_\rho^2 \left( \int_{\mathbb{R}^2} \ln|x-y| w_\rho^2(y) dy \right) |\partial_j w_\rho|^2 \\
& + \varepsilon_\rho^2 \left( \int_{\mathbb{R}^2} \frac{x_j - y_j}{|x-y|^2} w_\rho^2(y) dy \right) w_\rho \partial_j w_\rho + \rho \varepsilon_\rho^2 \ln \varepsilon_\rho |\partial_j w_\rho|^2 \\
& = \mu_\rho \varepsilon_\rho^2 |\partial_j w_\rho|^2 + 3w_\rho^2 |\partial_j w_\rho|^2 \quad \text{in } \mathbb{R}^2.
\end{aligned} \tag{A.2}$$

By Cauchy's inequality, we obtain from (3.12) and (3.22) that there exists a large constant  $R > 0$  such that

$$\frac{2\varepsilon_\rho^3 |\varepsilon_\rho x_j + x_\rho^j|}{1 + |\varepsilon_\rho x + x_\rho|^2} |w_\rho \partial_j w_\rho| \leq |w_\rho|^2 + \varepsilon_\rho^6 |\partial_j w_\rho|^2 \leq C e^{-\frac{4|x|}{3\sqrt{\rho^*}}} + \varepsilon_\rho^6 |\partial_j w_\rho|^2$$

and

$$\varepsilon_\rho^2 \left( \int_{\mathbb{R}^2} \frac{|x_j - y_j|}{|x-y|^2} w_\rho^2(y) dy \right) |w_\rho \partial_j w_\rho| \leq \varepsilon_\rho^2 \left( \int_{\mathbb{R}^2} \frac{1}{|x-y|} w_\rho^2(y) dy \right) |w_\rho \partial_j w_\rho| \leq C e^{-\frac{4|x|}{3\sqrt{\rho^*}}} + \varepsilon_\rho^4 |\partial_j w_\rho|^2$$

uniformly for  $|x| \geq R$  as  $\rho \nearrow \rho^*$ . In addition, we infer from (3.12) that

$$\varepsilon_\rho^2 \left[ \int_{\mathbb{R}^2} \ln \left( 1 + \frac{1}{|x-y|^2} \right) w_\rho^2(y) dy \right] |\partial_j w_\rho|^2 \leq C \varepsilon_\rho^2 \|w_\rho\|^2 |\partial_j w_\rho|^2 \quad \text{in } \mathbb{R}^2,$$

where and below  $\|\cdot\|$  is as in (3.13). Since  $\mu_\rho \varepsilon_\rho^2 \rightarrow -\frac{1}{\rho^*}$  as  $\rho \nearrow \rho^*$ , and

$$\begin{aligned}
& |\nabla \partial_j w_\rho|^2 + \varepsilon_\rho^2 \ln[1 + |\varepsilon_\rho x + x_\rho|^2] |\partial_j w_\rho|^2 \\
& + \frac{1}{2} \varepsilon_\rho^2 \left[ \int_{\mathbb{R}^2} \ln(1 + |x-y|^2) w_\rho^2(y) dy \right] |\partial_j w_\rho|^2 \geq 0 \quad \text{in } \mathbb{R}^2,
\end{aligned}$$

we conclude from (A.2) that

$$\left( -\frac{1}{2}\Delta + \frac{3}{4\rho^*} - 3w_\rho^2 \right) |\nabla w_\rho|^2 \leq C e^{-\frac{4|x|}{3\sqrt{\rho^*}}} \quad \text{uniformly for } |x| \geq R \text{ as } \rho \nearrow \rho^*. \tag{A.3}$$

Following De Giorgi-Nash-Moser theory [35, Theorem 4.1], we deduce from (A.3) that for any  $\xi \in \mathbb{R}^2$ , there exists a constant  $C > 0$  such that

$$\max_{B_1(\xi)} |\nabla w_\rho(x)|^2 \leq C \left( \int_{B_2(\xi)} |\nabla w_\rho|^2 dx + \left\| e^{-\frac{4|x|}{3\sqrt{\rho^*}}} \right\|_{L^2(B_2(\xi))} \right). \tag{A.4}$$

Moreover, because  $w_\rho \rightarrow \frac{1}{\sqrt{\rho^*}} Q(\frac{1}{\sqrt{\rho^*}}|x|)$  in  $H^1(\mathbb{R}^2)$  as  $\rho \nearrow \rho^*$ , it follows from (A.4) that

$$\lim_{|x| \rightarrow \infty} |\nabla w_\rho(x)| = 0 \quad \text{as } \rho \nearrow \rho^*. \tag{A.5}$$

Substituting (A.5) into (A.3), we infer that for sufficiently large  $R > 0$ ,

$$\left( -\Delta + \frac{4}{3\rho^*} \right) |\nabla w_\rho|^2 \leq C e^{-\frac{4|x|}{3\sqrt{\rho^*}}} \leq C e^{-\frac{|x|}{\sqrt{\rho^*}}} \quad \text{uniformly for } |x| \geq R \text{ as } \rho \nearrow \rho^*. \tag{A.6}$$

Applying the comparison principle to (A.6) yields that

$$|\nabla w_\rho(x)| \leq C e^{-\frac{|x|}{2\sqrt{\rho^*}}} \quad \text{uniformly for } |x| \geq R \text{ as } \rho \nearrow \rho^*.$$

This completes the proof of Lemma A.1.  $\square$



**Lemma A.2.** Let  $\varepsilon_k > 0$  and  $\bar{\eta}_k$  be as in (4.1) and (4.11), respectively. Then there exists a constant  $C > 0$ , independent of  $k$ , such that

$$\varepsilon_k^2 \int_{\mathbb{R}^2} |\nabla \bar{\eta}_k|^2 dx + \varepsilon_k^2 \int_{\mathbb{R}^2} V(x) \bar{\eta}_k^2 dx + \int_{\mathbb{R}^2} \bar{\eta}_k^2 dx \leq C \varepsilon_k^2 \quad \text{as } k \rightarrow \infty.$$

**Proof.** Define

$$\eta_k(x) := \bar{\eta}_k(\varepsilon_k x + x_{2,k}) \quad \text{in } \mathbb{R}^2,$$

and

$$\tilde{u}_{i,k}(x) := \varepsilon_k u_{i,k}(\varepsilon_k x + x_{2,k}) = \bar{u}_{i,k}(\varepsilon_k x + x_{2,k}) > 0 \quad \text{in } \mathbb{R}^2, \quad i = 1, 2,$$

where  $\bar{u}_{i,k}$  is as in (4.9) and  $x_{2,k}$  is the unique maximal point of  $u_{2,k}$  as  $k \rightarrow \infty$ . We now estimate the right hand side of (4.14) as follows.

The exponential decay of  $\tilde{u}_{i,k}$  in (4.6) yields that

$$\begin{aligned} & \int_{\mathbb{R}^2} (\bar{u}_{2,k}^2 + \bar{u}_{2,k} \bar{u}_{1,k} + \bar{u}_{1,k}^2) \bar{\eta}_k^2 dx - \frac{1}{2\rho_k \varepsilon_k^2} \int_{\mathbb{R}^2} (\bar{u}_{2,k}^2 + \bar{u}_{1,k}^2) (\bar{u}_{2,k} + \bar{u}_{1,k}) \bar{\eta}_k dx \int_{\mathbb{R}^2} \bar{u}_{1,k} \bar{\eta}_k dx \\ & \leq C \varepsilon_k^2 \quad \text{as } k \rightarrow \infty. \end{aligned}$$

It follows from (3.12) that there exists a constant  $C > 0$ , independent of  $x \in \mathbb{R}^2$  and  $k$ , such that

$$\int_{\mathbb{R}^2} \frac{1}{|x-y|} \tilde{u}_{2,k}^2(y) dy \leq C \|\tilde{u}_{2,k}\|^2 \leq C,$$

from which we deduce that

$$\begin{aligned} & \frac{1}{2} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln\left(1 + \frac{1}{|x-y|^2}\right) \bar{\eta}_k^2(x) \bar{u}_{2,k}^2(y) dx dy \\ & \leq \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{1}{|x-y|} \bar{\eta}_k^2(x) \bar{u}_{2,k}^2(y) dx dy \\ & \leq \varepsilon_k^3 \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{1}{|x-y|} \eta_k^2(x) \tilde{u}_{2,k}^2(y) dx dy \\ & = \varepsilon_k^3 \int_{\mathbb{R}^2} \eta_k^2(x) \int_{\mathbb{R}^2} \frac{1}{|x-y|} \tilde{u}_{2,k}^2(y) dy dx \\ & \leq C \varepsilon_k \|\tilde{u}_{2,k}\|^2 \int_{\mathbb{R}^2} \bar{\eta}_k^2(x) dx \leq C \varepsilon_k \int_{\mathbb{R}^2} \bar{\eta}_k^2(x) dx \quad \text{as } k \rightarrow \infty. \end{aligned}$$

In addition, since  $|\eta_k| \leq 1$ , we get from (3.12) and (4.6) that

$$\begin{aligned} & \frac{1}{2\rho_k \varepsilon_k^2} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| (\bar{u}_{2,k}(x) + \bar{u}_{1,k}(x)) \bar{\eta}_k(x) \bar{u}_{2,k}^2(y) dx dy \int_{\mathbb{R}^2} \bar{u}_{1,k}(x) \bar{\eta}_k(x) dx \\ & = \frac{\varepsilon_k^4 \ln \varepsilon_k}{2\rho_k} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} (\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)) \eta_k(x) \tilde{u}_{2,k}^2(y) dx dy \int_{\mathbb{R}^2} \tilde{u}_{1,k}(x) \eta_k(x) dx \\ & + \frac{\varepsilon_k^4}{4\rho_k} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln(1 + |x-y|^2) (\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)) \eta_k(x) \tilde{u}_{2,k}^2(y) dx dy \int_{\mathbb{R}^2} \tilde{u}_{1,k}(x) \eta_k(x) dx \\ & - \frac{\varepsilon_k^4}{4\rho_k} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln\left(1 + \frac{1}{|x-y|^2}\right) (\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)) \eta_k(x) \tilde{u}_{2,k}^2(y) dx dy \int_{\mathbb{R}^2} \tilde{u}_{1,k}(x) \eta_k(x) dx \\ & \leq \left| \frac{1}{2} \varepsilon_k^4 \ln \varepsilon_k \int_{\mathbb{R}^2} (\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)) \frac{\tilde{u}_{2,k}(x) - \tilde{u}_{1,k}(x)}{\|\tilde{u}_{2,k} - \tilde{u}_{1,k}\|_{L^\infty(\mathbb{R}^2)}} dx \int_{\mathbb{R}^2} \tilde{u}_{1,k}(x) \eta_k(x) dx \right| \\ & + \frac{\varepsilon_k^4}{4\rho_k} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \left[ \ln(1 + 2|x|^2 + 2|y|^2) + \ln\left(1 + \frac{1}{|x-y|^2}\right) \right] |\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)| \tilde{u}_{2,k}^2(y) dx dy \\ & \quad \cdot \left| \int_{\mathbb{R}^2} \tilde{u}_{1,k}(x) \eta_k(x) dx \right| \\ & \leq \left| \frac{(\rho_k - \rho_k) \varepsilon_k^4 \ln \varepsilon_k}{2 \|\tilde{u}_{2,k} - \tilde{u}_{1,k}\|_{L^\infty(\mathbb{R}^2)}} \int_{\mathbb{R}^2} \tilde{u}_{1,k}(x) \eta_k(x) dx \right| \end{aligned}$$

$$\begin{aligned}
& + \frac{\varepsilon_k^4}{4\rho_k} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \left[ \ln(1 + |x|^2)^2 + \ln(1 + |y|^2)^2 + \ln\left(1 + \frac{1}{|x-y|}\right)^2 \right] |\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)| \tilde{u}_{2,k}^2(y) dx dy \\
& \quad \cdot \left| \int_{\mathbb{R}^2} \tilde{u}_{1,k}(x) \eta_k(x) dx \right| \\
& \leq \frac{\varepsilon_k^4}{2\rho_k} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \left[ \ln(1 + |x|^2) + \ln(1 + |y|^2) + \frac{1}{|x-y|} \right] |\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)| \tilde{u}_{2,k}^2(y) dx dy \left| \int_{\mathbb{R}^2} \tilde{u}_{1,k}(x) \eta_k(x) dx \right| \\
& \leq \frac{\varepsilon_k^4}{2\rho_k} \left[ \rho_k \int_{\mathbb{R}^2} \ln(1 + |x|^2) |\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)| dx + \int_{\mathbb{R}^2} |\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)| dx \int_{\mathbb{R}^2} \ln(1 + |y|^2) \tilde{u}_{2,k}^2(y) dy \right. \\
& \quad \left. + \int_{\mathbb{R}^2} |\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)| \int_{\mathbb{R}^2} \frac{1}{|x-y|} \tilde{u}_{2,k}^2(y) dy dx \right] \left| \int_{\mathbb{R}^2} \tilde{u}_{1,k}(x) \eta_k(x) dx \right| \leq C\varepsilon_k^4 \text{ as } k \rightarrow \infty,
\end{aligned} \tag{A.7}$$

and

$$\frac{1}{2\rho_k \varepsilon_k^2} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| \bar{u}_{1,k}^2(x) (\bar{u}_{2,k}(y) + \bar{u}_{1,k}(y)) \bar{\eta}_k(y) dx dy \int_{\mathbb{R}^2} \bar{u}_{1,k}(x) \bar{\eta}_k(x) dx \leq C\varepsilon_k^4 \text{ as } k \rightarrow \infty.$$

Similarly, we also have

$$\begin{aligned}
& \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| \bar{u}_{1,k}(x) \bar{\eta}_k(x) (\bar{u}_{2,k}(y) + \bar{u}_{1,k}(y)) \bar{\eta}_k(y) dx dy \\
& \leq \varepsilon_k^4 \int_{\mathbb{R}^2} \ln(1 + |x|^2) |\bar{u}_{1,k}(x)| dx \int_{\mathbb{R}^2} |\bar{u}_{2,k}(y) + \bar{u}_{1,k}(y)| dy \\
& \quad + \varepsilon_k^4 \int_{\mathbb{R}^2} |\bar{u}_{1,k}(x)| dx \int_{\mathbb{R}^2} \ln(1 + |y|^2) |\bar{u}_{2,k}(y) + \bar{u}_{1,k}(y)| dy \\
& \quad + \varepsilon_k^4 \int_{\mathbb{R}^2} |\bar{u}_{1,k}(x) \bar{\eta}_k(x)| \left( \int_{\mathbb{R}^2} \frac{1}{|x-y|} |\bar{u}_{2,k}(y) + \bar{u}_{1,k}(y)|^2 dy \right)^{\frac{1}{2}} \left( \int_{\mathbb{R}^2} \frac{1}{|x-y|} \bar{\eta}_k^2(y) dy \right)^{\frac{1}{2}} dx.
\end{aligned} \tag{A.8}$$

The same argument of (3.12) gives that

$$\left( \int_{\mathbb{R}^2} \frac{1}{|x-y|} |\bar{u}_{2,k}(y) + \bar{u}_{1,k}(y)|^2 dy \right)^{\frac{1}{2}} \left( \int_{\mathbb{R}^2} \frac{1}{|x-y|} \bar{\eta}_k^2(y) dy \right)^{\frac{1}{2}} \leq C \|\bar{u}_{1,k} + \bar{u}_{2,k}\| \|\bar{\eta}_k\|. \tag{A.9}$$

We then deduce from (A.8) and (A.9) that

$$\begin{aligned}
& \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| \bar{u}_{1,k}(x) \bar{\eta}_k(x) (\bar{u}_{2,k}(y) + \bar{u}_{1,k}(y)) \bar{\eta}_k(y) dx dy \\
& \leq C\varepsilon_k^4 + C\varepsilon_k^4 \rho_k^{\frac{1}{2}} \|\bar{u}_{2,k} + \bar{u}_{1,k}\| \|\bar{\eta}_k\|^2 \\
& \leq C\varepsilon_k^4 + C\varepsilon_k^2 \int_{\mathbb{R}^2} \bar{\eta}_k^2 dx + C\varepsilon_k^4 \int_{\mathbb{R}^2} |\nabla \bar{\eta}_k|^2 dx \text{ as } k \rightarrow \infty.
\end{aligned}$$

Applying (4.14), we infer from above that there exists a constant  $C > 0$ , independent of  $k$ , such that

$$\begin{aligned}
& \varepsilon_k^2 \int_{\mathbb{R}^2} |\nabla \bar{\eta}_k|^2 dx + \varepsilon_k^2 \int_{\mathbb{R}^2} V(x) \bar{\eta}_k^2 dx - \mu_{2,k} \varepsilon_k^2 \int_{\mathbb{R}^2} \bar{\eta}_k^2 dx \\
& \leq C \left( \varepsilon_k \int_{\mathbb{R}^2} \bar{\eta}_k^2 dx + \varepsilon_k^4 \int_{\mathbb{R}^2} |\nabla \bar{\eta}_k|^2 dx + \varepsilon_k^2 \right) \text{ as } k \rightarrow \infty.
\end{aligned}$$

Following (4.4), we obtain from above that

$$\varepsilon_k^2 \int_{\mathbb{R}^2} |\nabla \bar{\eta}_k|^2 dx + \varepsilon_k^2 \int_{\mathbb{R}^2} V(x) \bar{\eta}_k^2 dx + \int_{\mathbb{R}^2} \bar{\eta}_k^2 dx \leq C\varepsilon_k^2 \text{ as } k \rightarrow \infty.$$

This completes the proof of Lemma A.2.  $\square$

**Lemma A.3.** *Let  $\eta_k$  be a solution of (4.18), i.e.,  $\eta_k$  satisfies*

$$-\Delta\eta_k + C_k(x)\eta_k = D_k(x) \quad \text{in } \mathbb{R}^2,$$

where  $C_k(x)$  and  $D_k(x)$  are as in (4.19) and (4.21), respectively. Then the sequence  $\{\eta_k\}$  is bounded uniformly in  $C_{loc}^{1,\alpha}(\mathbb{R}^2)$  as  $k \rightarrow \infty$  for some  $\alpha \in (0, 1)$ .

**Proof.** Multiplying (4.18) by  $\eta_k$ , we deduce that

$$\begin{aligned} & -\frac{1}{2}\Delta\eta_k^2 + |\nabla\eta_k|^2 + \varepsilon_k^2 V(\varepsilon_k x + x_{2,k})\eta_k^2 + \rho_k \varepsilon_k^2 \ln \varepsilon_k \eta_k^2 - \mu_{2,k} \varepsilon_k^2 \eta_k^2 \\ & + \frac{1}{2} \varepsilon_k^2 \left( \int_{\mathbb{R}^2} \ln(1 + |x - y|^2) \tilde{u}_{2,k}^2(y) dy \right) \eta_k^2 + A_k(x) \eta_k^2 = D_k(x) \eta_k \quad \text{in } \mathbb{R}^2, \end{aligned} \quad (\text{A.10})$$

where  $A_k(x)$  and  $D_k(x)$  are as in (4.20) and (4.21), respectively.

We first claim that there exists a large constant  $R > 0$  such that

$$|\eta_k(x)| \leq C e^{-\frac{|x|}{3}} \quad \text{uniformly for } |x| \geq R \quad \text{as } k \rightarrow \infty. \quad (\text{A.11})$$

Indeed, one can obtain from (4.15) that there exists a constant  $C > 0$ , independent of  $k$ , such that

$$\|\eta_k\|^2 = \varepsilon_k^{-2} \int_{\mathbb{R}^2} \bar{\eta}_k^2 dx + \int_{\mathbb{R}^2} |\nabla \bar{\eta}_k|^2 dx \leq C \quad \text{as } k \rightarrow \infty. \quad (\text{A.12})$$

Similar to (A), we derive that

$$\begin{aligned} & \varepsilon_k^2 \tilde{u}_{1,k} \eta_k \left[ \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x - y| (\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)) \eta_k(x) \tilde{u}_{2,k}^2(y) dx dy \right] \\ & \leq C \varepsilon_k^2 |\tilde{u}_{1,k} \eta_k| \leq \varepsilon_k^4 \eta_k^2 + C e^{-\frac{4|x|}{3}} \quad \text{uniformly for } |x| \geq R \quad \text{as } k \rightarrow \infty, \end{aligned} \quad (\text{A.13})$$

and

$$\begin{aligned} & \varepsilon_k^2 \tilde{u}_{1,k} \eta_k \left[ \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x - y| \tilde{u}_{1,k}^2(x) (\tilde{u}_{2,k}(y) + \tilde{u}_{1,k}(y)) \eta_k(y) dx dy \right] \\ & \leq \varepsilon_k^4 \eta_k^2 + C e^{-\frac{4|x|}{3}} \quad \text{uniformly for } |x| \geq R \quad \text{as } k \rightarrow \infty. \end{aligned} \quad (\text{A.14})$$

The same argument of (A.8) yields that

$$\begin{aligned} & \varepsilon_k^2 \tilde{u}_{1,k} \eta_k \left[ \int_{\mathbb{R}^2} \ln|x - y| (\tilde{u}_{2,k}(y) + \tilde{u}_{1,k}(y)) \eta_k(y) dy \right] \\ & \leq C \varepsilon_k^2 [1 + \ln(1 + |x|^2)] |\tilde{u}_{1,k} \eta_k| \leq \varepsilon_k^4 \eta_k^2 + C e^{-\frac{2|x|}{3}} \quad \text{uniformly for } |x| \geq R \quad \text{as } k \rightarrow \infty. \end{aligned} \quad (\text{A.15})$$

Moreover, it follows from (4.6) that

$$\tilde{u}_{1,k} \eta_k \int_{\mathbb{R}^2} (\tilde{u}_{2,k}^2 + \tilde{u}_{1,k}^2) (\tilde{u}_{2,k} + \tilde{u}_{1,k}) \eta_k dx \leq \frac{1}{4} \eta_k^2 + C e^{-\frac{4|x|}{3}} \quad \text{uniformly for } |x| \geq R \quad \text{as } k \rightarrow \infty, \quad (\text{A.16})$$

and

$$(\tilde{u}_{2,k}^2 + \tilde{u}_{2,k} \tilde{u}_{1,k} + \tilde{u}_{1,k}^2) \eta_k^2 \leq C e^{-\frac{4|x|}{3}} \quad \text{uniformly for } |x| \geq R \quad \text{as } k \rightarrow \infty. \quad (\text{A.17})$$

Using (4.4), we thus derive from (A.10) and (A.13)–(A.17) that there exists a large constant  $R > 0$  such that

$$-\frac{1}{2}\Delta\eta_k^2 + \frac{2}{3}\eta_k^2 \leq C e^{-\frac{2|x|}{3}} \quad \text{uniformly for } |x| \geq R \quad \text{as } k \rightarrow \infty. \quad (\text{A.18})$$

Applying the comparison principle to (A.18), the claim (A.11) is therefore proved.

We next prove that  $\{\eta_k\}$  is bounded uniformly in  $C_{loc}^{1,\alpha}(\mathbb{R}^2)$  as  $k \rightarrow \infty$  for some  $\alpha \in (0, 1)$ . Denote

$$F_k(x) = -C_k(x)\eta_k + D_k(x) \quad \text{in } \mathbb{R}^2,$$

so that  $\eta_k$  satisfies

$$-\Delta\eta_k = F_k \text{ in } \mathbb{R}^2.$$

We now claim that for any  $p \geq 2$ ,  $\{F_k\}$  is bounded uniformly in  $L^p_{loc}(\mathbb{R}^2)$  as  $k \rightarrow \infty$ . By the exponential decay of  $\tilde{u}_{i,k}$ , it is sufficient to prove that

$$\int_{\mathbb{R}^2} \ln|x-y| \tilde{u}_{2,k}^2(y) dy \eta_k(x)$$

is bounded uniformly in  $L^p_{loc}(\mathbb{R}^2)$  as  $k \rightarrow \infty$ .

Direct calculations yield that for any  $p \geq 2$ ,

$$\begin{aligned} & \int_{\mathbb{R}^2} \left| \int_{\mathbb{R}^2} \ln|x-y| \tilde{u}_{2,k}^2(y) dy \right|^p |\eta_k(x)|^p dx \\ &= \int_{\mathbb{R}^2} \left| \frac{1}{2} \int_{\mathbb{R}^2} \left( \ln(1+|x-y|^2) - \ln\left(1 + \frac{1}{|x-y|^2}\right) \right) \tilde{u}_{2,k}^2(y) dy \right|^p |\eta_k(x)|^p dx \\ &\leq \int_{\mathbb{R}^2} \left[ \frac{1}{2} \int_{\mathbb{R}^2} \left( \ln(1+2|x|^2+2|y|^2) + \ln\left(1 + \frac{1}{|x-y|^2}\right) \right) \tilde{u}_{2,k}^2(y) dy \right]^p |\eta_k(x)|^p dx \\ &\leq \int_{\mathbb{R}^2} \left[ \frac{1}{2} \int_{\mathbb{R}^2} \left( \ln(1+|x|^2)^2 + \ln(1+|y|^2)^2 + \ln\left(1 + \frac{1}{|x-y|^2}\right)^2 \right) \tilde{u}_{2,k}^2(y) dy \right]^p |\eta_k(x)|^p dx \\ &\leq \int_{\mathbb{R}^2} \left[ \int_{\mathbb{R}^2} \left( \ln(1+|x|^2) + \ln(1+|y|^2) + \frac{1}{|x-y|^2} \right) \tilde{u}_{2,k}^2(y) dy \right]^p |\eta_k(x)|^p dx \\ &= \int_{\mathbb{R}^2} \left[ \ln(1+|x|^2) \rho_k + \int_{\mathbb{R}^2} \ln(1+|y|^2) \tilde{u}_{2,k}^2(y) dy + \int_{\mathbb{R}^2} \frac{1}{|x-y|^2} \tilde{u}_{2,k}^2(y) dy \right]^p |\eta_k(x)|^p dx. \end{aligned} \tag{A.19}$$

We obtain from (3.12) and (4.6) that

$$\int_{\mathbb{R}^2} \ln(1+|y|^2) \tilde{u}_{2,k}^2(y) dy + \int_{\mathbb{R}^2} \frac{1}{|x-y|^2} \tilde{u}_{2,k}^2(y) dy \leq C + C \|\tilde{u}_{2,k}\|^2 \leq C. \tag{A.20}$$

Applying (A.11), we then deduce from (A.19) and (A.20) that

$$\int_{\mathbb{R}^2} \left| \int_{\mathbb{R}^2} \ln|x-y| \tilde{u}_{2,k}^2(y) dy \right|^p |\eta_k|^p dx \leq C \int_{\mathbb{R}^2} \left[ 1 + \ln(1+|x|^2) \right]^p |\eta_k(x)|^p dx \leq C.$$

This implies that  $\{F_k\}$  is bounded uniformly in  $L^p_{loc}(\mathbb{R}^2)$  as  $k \rightarrow \infty$  for  $p \in [2, \infty)$ . The same argument of (3.28) yields that  $\{\eta_k\}$  is bounded uniformly in  $C^{1,\alpha}_{loc}(\mathbb{R}^2)$  as  $k \rightarrow \infty$  for some  $\alpha \in (0, 1)$ . This completes the proof of Lemma A.3.  $\square$

**Lemma A.4.** *Assume  $\eta_k$  is a solution of (4.18) and  $\eta_k \rightarrow \eta_0$  in  $H^1(\mathbb{R}^2)$  as  $k \rightarrow \infty$ . Then  $\eta_0$  is a weak solution of the following equation:*

$$-\Delta\eta_0 + (1-3Q^2)\eta_0 = -\frac{2}{\rho^*} \left( \int_{\mathbb{R}^2} Q^3 \eta_0 dx \right) Q \text{ in } \mathbb{R}^2.$$

**Proof.** Since  $x_{2,k} \rightarrow 0$  and  $\tilde{u}_{i,k} \rightarrow Q$  in  $L^\infty(\mathbb{R}^2)$  as  $k \rightarrow \infty$ , we first derive from (1.8), (A.11) and (A.12) that for any  $\varphi \in H^1(\mathbb{R}^2)$ ,

$$\begin{aligned} & \int_{\mathbb{R}^2} \nabla\eta_k \cdot \nabla\varphi dx + \varepsilon_k^2 \int_{\mathbb{R}^2} V(\varepsilon_k x + x_{2,k}) \eta_k \varphi dx + \rho_k \varepsilon_k^2 \ln \varepsilon_k \int_{\mathbb{R}^2} \eta_k \varphi dx - \mu_{2,k} \varepsilon_k^2 \int_{\mathbb{R}^2} \eta_k \varphi dx \\ & - \int_{\mathbb{R}^2} (\tilde{u}_{2,k}^2 + \tilde{u}_{2,k} \tilde{u}_{1,k} + \tilde{u}_{1,k}^2) \eta_k \varphi dx + \frac{1}{2\rho_k} \int_{\mathbb{R}^2} (\tilde{u}_{2,k}^2 + \tilde{u}_{1,k}^2) (\tilde{u}_{2,k} + \tilde{u}_{1,k}) \eta_k dx \int_{\mathbb{R}^2} \tilde{u}_{1,k} \varphi dx \\ & \rightarrow \int_{\mathbb{R}^2} \nabla\eta_0 \cdot \nabla\varphi dx + \int_{\mathbb{R}^2} \eta_0 \varphi dx - 3 \int_{\mathbb{R}^2} Q^2 \eta_0 \varphi dx + \frac{2}{\rho^*} \left( \int_{\mathbb{R}^2} Q^3 \eta_0 dx \right) \int_{\mathbb{R}^2} Q \varphi dx \text{ as } k \rightarrow \infty. \end{aligned} \tag{A.21}$$

Similar to (A), we obtain from (3.12), (4.6), (A.11) and (A.12) that

$$\begin{aligned}
& \varepsilon_k^2 \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} |\ln|x-y||\eta_k(x)\varphi(x)|\tilde{u}_{2,k}^2(y)dydx \\
& \leq \rho_k \int_{\mathbb{R}^2} \ln(1+|x|^2)|\eta_k(x)\varphi(x)|dx + \left(\int_{\mathbb{R}^2} \eta_k^2 dx\right)^{\frac{1}{2}} \left(\int_{\mathbb{R}^2} \varphi^2 dx\right)^{\frac{1}{2}} \int_{\mathbb{R}^2} \ln(1+|y|^2)\tilde{u}_{2,k}^2(y)dy \\
& \quad + \int_{\mathbb{R}^2} |\eta_k(x)\varphi(x)| \int_{\mathbb{R}^2} \frac{1}{|x-y|} \tilde{u}_{2,k}^2(y)dydx \leq C\varepsilon_k^2 \rightarrow 0 \text{ as } k \rightarrow \infty.
\end{aligned} \tag{A.22}$$

Finally, the same arguments of (A.13), (A.14) and (A.15) give that

$$\begin{aligned}
& \frac{1}{2\rho_k} \varepsilon_k^2 \left[ \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y|(\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x))\eta_k(x)\tilde{u}_{2,k}^2(y)dx dy \right] \int_{\mathbb{R}^2} \tilde{u}_{1,k}(x)\varphi(x)dx \\
& \quad + \frac{1}{2\rho_k} \varepsilon_k^2 \left[ \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y|\tilde{u}_{1,k}^2(x)(\tilde{u}_{2,k}(y) + \tilde{u}_{1,k}(y))\eta_k(y)dx dy \right] \int_{\mathbb{R}^2} \tilde{u}_{1,k}(x)\varphi(x)dx \\
& \quad - \varepsilon_k^2 \int_{\mathbb{R}^2} \left[ \int_{\mathbb{R}^2} \ln|x-y|(\tilde{u}_{2,k}(y) + \tilde{u}_{1,k}(y))\eta_k(y)dy \right] \tilde{u}_{1,k}(x)\varphi(x)dx = o(1) \text{ as } k \rightarrow \infty.
\end{aligned} \tag{A.23}$$

The proof of Lemma A.4 is therefore complete in view of (A.21)–(A.23).  $\square$

**Lemma A.5.** *Let  $\alpha_{j,k}$  be as in (4.28) for  $j = 1, 2$ . Then there exists a constant  $a > 0$  such that for  $j = 1, 2$ ,*

$$\alpha_{j,k} = o\left(e^{-\frac{a\delta}{\varepsilon_k}}\right) \text{ as } k \rightarrow \infty,$$

where  $\varepsilon_k$  and  $\delta > 0$  are as in (4.1) and (4.13), respectively.

**Proof.** We now estimate each term of  $\alpha_{j,k}$  as  $k \rightarrow \infty$  as follows. Since  $|\nabla\tilde{u}_{i,k}|$  decays exponentially as  $k \rightarrow \infty$ , it follows from (4.13) that there exists a constant  $a > 0$  such that

$$\begin{aligned}
& \varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} \left| \frac{\partial \bar{\eta}_k}{\partial x_j} \frac{\partial \bar{u}_{2,k}}{\partial \nu} \right| dS \\
& \leq \varepsilon_k \left( \int_{\partial B_\delta(x_{2,k})} |\nabla \bar{u}_{2,k}|^2 dS \right)^{\frac{1}{2}} \left( \varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} |\nabla \bar{\eta}_k|^2 dS \right)^{\frac{1}{2}} \\
& \leq C\varepsilon_k \left( \varepsilon_k^{-1} \int_{\partial B_{\frac{\delta}{\varepsilon_k}}(0)} |\nabla \tilde{u}_{2,k}|^2 dS \right)^{\frac{1}{2}} \left( \varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} |\nabla \bar{\eta}_k|^2 dS \right)^{\frac{1}{2}} \\
& \leq C\varepsilon_k^2 e^{-\frac{a\delta}{\varepsilon_k}} \text{ as } k \rightarrow \infty,
\end{aligned} \tag{A.24}$$

$$\varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} \left| \frac{\partial \bar{u}_{1,k}}{\partial x_j} \frac{\partial \bar{\eta}_k}{\partial \nu} \right| dS \leq C\varepsilon_k^2 e^{-\frac{a\delta}{\varepsilon_k}} \text{ as } k \rightarrow \infty, \tag{A.25}$$

and

$$\varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} \left| (\nabla \bar{u}_{2,k} + \nabla \bar{u}_{1,k}) \cdot \nabla \bar{\eta}_k \nu_j \right| dS \leq C\varepsilon_k^2 e^{-\frac{a\delta}{\varepsilon_k}} \text{ as } k \rightarrow \infty. \tag{A.26}$$

In addition, applying the exponential decay of  $\tilde{u}_{i,k}$  in (4.6), we infer from (4.4) and (4.13) that

$$\begin{aligned}
& \left| \varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} \ln(1+|x|^2)(\bar{u}_{2,k} + \bar{u}_{1,k})\bar{\eta}_k \nu_j dS - \varepsilon_k^2 \mu_{2,k} \int_{\partial B_\delta(x_{2,k})} (\bar{u}_{2,k} + \bar{u}_{1,k})\bar{\eta}_k(x)\nu_j dS \right. \\
& \quad \left. - \frac{1}{2} \int_{\partial B_\delta(x_{2,k})} (\bar{u}_{2,k}^2 + \bar{u}_{1,k}^2)(\bar{u}_{2,k} + \bar{u}_{1,k})\bar{\eta}_k \nu_j dS \right| = o\left(e^{-\frac{a\delta}{\varepsilon_k}}\right) \text{ as } k \rightarrow \infty.
\end{aligned} \tag{A.27}$$

By the same argument of (A), it yields from (4.6), (4.13), (A.9), (A.11) and (A.12) that

$$\begin{aligned}
& \left| \int_{\partial B_\delta(x_{2,k})} \int_{\mathbb{R}^2} \ln|x-y| (\bar{u}_{2,k}(y) + \bar{u}_{1,k}(y)) \bar{\eta}_k(y) \bar{u}_{2,k}^2(x) \nu_j dy dS \right| \\
& \leq \varepsilon_k^3 \int_{\partial B_{\frac{\delta}{\varepsilon_k}}(0)} \ln(1+|x|^2) \tilde{u}_{2,k}^2(x) dS \int_{\mathbb{R}^2} |(\tilde{u}_{2,k}(y) + \tilde{u}_{1,k}(y)) \eta_k(y)| dy \\
& \quad + \varepsilon_k^3 \int_{\partial B_{\frac{\delta}{\varepsilon_k}}(0)} \tilde{u}_{2,k}^2(x) dS \int_{\mathbb{R}^2} \ln(1+|y|^2) |(\tilde{u}_{2,k}(y) + \tilde{u}_{1,k}(y)) \eta_k(y)| dy \\
& \quad + \varepsilon_k^3 \int_{\partial B_{\frac{\delta}{\varepsilon_k}}(0)} \tilde{u}_{2,k}^2(x) \left[ \int_{\mathbb{R}^2} \frac{1}{|x-y|} (\tilde{u}_{2,k}(y) + \tilde{u}_{1,k}(y))^2 dy \right]^{\frac{1}{2}} \left( \int_{\mathbb{R}^2} \frac{1}{|x-y|} \eta_k^2(y) dy \right)^{\frac{1}{2}} dS \\
& \leq C \rho_k^{\frac{1}{2}} \varepsilon_k^3 e^{-\frac{\alpha\delta}{\varepsilon_k}} \|\eta_k\|_2 + C \rho_k^{\frac{1}{2}} \varepsilon_k^3 e^{-\frac{\alpha\delta}{\varepsilon_k}} \left( \int_{\mathbb{R}^2} [\ln(1+|y|^2)]^2 \eta_k^2(y) dy \right)^{\frac{1}{2}} \\
& \quad + C \varepsilon_k^3 e^{-\frac{\alpha\delta}{\varepsilon_k}} (\|\tilde{u}_{2,k}\| + \|\tilde{u}_{1,k}\|) \|\eta_k\| \leq C \varepsilon_k^3 e^{-\frac{\alpha\delta}{\varepsilon_k}} \quad \text{as } k \rightarrow \infty,
\end{aligned} \tag{A.28}$$

and

$$\left| \int_{\partial B_\delta(x_{2,k})} \int_{\mathbb{R}^2} \ln|x-y| \bar{u}_{1,k}^2(y) (\bar{u}_{2,k}(x) + \bar{u}_{1,k}(x)) \bar{\eta}_k(x) \nu_j dy dS \right| \leq C \varepsilon_k^3 e^{-\frac{\alpha\delta}{\varepsilon_k}} \quad \text{as } k \rightarrow \infty. \tag{A.29}$$

Applying (4.6), we deduce from (3.12) and (A.12) that

$$\begin{aligned}
& \int_{B_\delta^c(x_{2,k})} \int_{\mathbb{R}^2} \left| \frac{x_j - y_j}{|x-y|^2} (\bar{u}_{2,k}(x) + \bar{u}_{1,k}(x)) \bar{\eta}_k(x) \bar{u}_{2,k}^2(y) \right| dy dx \\
& \leq \varepsilon_k^3 \int_{B_{\delta/\varepsilon_k}^c(0)} |(\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)) \eta_k(x)| \int_{\mathbb{R}^2} \frac{1}{|x-y|} \tilde{u}_{2,k}^2(y) dy dx \\
& \leq C \varepsilon_k^3 \|\tilde{u}_{2,k}\|^2 \left( \int_{B_{\delta/\varepsilon_k}^c(0)} |\tilde{u}_{2,k} + \tilde{u}_{1,k}|^2 dx \right)^{\frac{1}{2}} \left( \int_{\mathbb{R}^2} \eta_k^2 dx \right)^{\frac{1}{2}} \leq C \varepsilon_k^3 e^{-\frac{\alpha\delta}{\varepsilon_k}} \quad \text{as } k \rightarrow \infty,
\end{aligned} \tag{A.30}$$

and

$$\int_{B_\delta^c(x_{2,k})} \int_{\mathbb{R}^2} \left| \frac{x_j - y_j}{|x-y|^2} \bar{u}_{1,k}^2(x) (\bar{u}_{2,k}(y) + \bar{u}_{1,k}(y)) \bar{\eta}_k(y) \right| dy dx \leq C \varepsilon_k^3 e^{-\frac{\alpha\delta}{\varepsilon_k}} \quad \text{as } k \rightarrow \infty. \tag{A.31}$$

To estimate the last term of  $\alpha_{j,k}$  as  $k \rightarrow \infty$ , we first claim that

$$\begin{aligned}
\frac{\mu_{2,k} - \mu_{1,k}}{\|\bar{u}_{2,k} - \bar{u}_{1,k}\|_{L^\infty(\mathbb{R}^2)}} \varepsilon_k^2 &= \frac{1}{2\rho_k \varepsilon_k^2} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| (\bar{u}_{2,k}(x) + \bar{u}_{1,k}(x)) \bar{\eta}_k(x) \bar{u}_{2,k}^2(y) dx dy \\
& \quad + \frac{1}{2\rho_k \varepsilon_k^2} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| \bar{u}_{1,k}^2(x) (\bar{u}_{2,k}(y) + \bar{u}_{1,k}(y)) \bar{\eta}_k(y) dx dy \\
& \quad - \frac{1}{2\rho_k \varepsilon_k^2} \int_{\mathbb{R}^2} (\bar{u}_{2,k}^2(x) + \bar{u}_{1,k}^2(x)) (\bar{u}_{2,k}(x) + \bar{u}_{1,k}(x)) \bar{\eta}_k(x) dx
\end{aligned} \tag{A.32}$$

is bounded uniformly in  $k$  as  $k \rightarrow \infty$ . In fact, the same argument of (A) yields that

$$\begin{aligned}
& \left| \frac{1}{\varepsilon_k} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| (\bar{u}_{2,k}(x) + \bar{u}_{1,k}(x)) \bar{\eta}_k(x) \bar{u}_{2,k}^2(y) dx dy \right. \\
& \quad \left. + \frac{1}{\varepsilon_k} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| \bar{u}_{1,k}^2(x) (\bar{u}_{2,k}(y) + \bar{u}_{1,k}(y)) \bar{\eta}_k(y) dx dy \right| \leq C \varepsilon_k^2 \quad \text{as } k \rightarrow \infty.
\end{aligned} \tag{A.33}$$

It follows from (4.6) that

$$\begin{aligned}
& \frac{1}{2\varepsilon_k^2} \int_{\mathbb{R}^2} |(\bar{u}_{2,k}^2 + \bar{u}_{1,k}^2) (\bar{u}_{2,k} + \bar{u}_{1,k}) \bar{\eta}_k| dx \\
& = \frac{1}{2} \int_{\mathbb{R}^2} |(\tilde{u}_{2,k}^2 + \tilde{u}_{1,k}^2) (\tilde{u}_{2,k} + \tilde{u}_{1,k}) \eta_k| dx \leq C \quad \text{as } k \rightarrow \infty.
\end{aligned} \tag{A.34}$$

We then conclude from (A.33) and (A.34) that (A.32) is bounded uniformly in  $k$  as  $k \rightarrow \infty$ .

Moreover, by the exponential decay of  $\tilde{u}_{1,k}$  as  $k \rightarrow \infty$ , it yields from (A.32) that

$$\frac{\mu_{2,k} - \mu_{1,k}}{\|\bar{u}_{2,k} - \bar{u}_{1,k}\|_{L^\infty(\mathbb{R}^2)}} \varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} \bar{u}_{1,k}^2 \nu_j dS = o(e^{-\frac{a\delta}{\varepsilon_k}}) \quad \text{as } k \rightarrow \infty. \quad (\text{A.35})$$

Applying the estimates of (A.24)–(A.31) and (A.35), the proof of Lemma A.5 is therefore complete.  $\square$

**Lemma A.6.** *Let  $\gamma_k$  and  $\delta > 0$  be as in (4.37) and (4.13), respectively. Then there exists a constant  $b \in (0, a)$  such that*

$$\gamma_k = o(e^{-\frac{b\delta}{\varepsilon_k}}) \quad \text{as } k \rightarrow \infty.$$

**Proof.** Since  $x_{2,k} \rightarrow 0$  as  $k \rightarrow \infty$ , we note from the first identity of (4.29) that

$$\begin{aligned} & \frac{\varepsilon_k^2}{2} \int_{B_\delta(x_{2,k})} x_{2,k} \cdot \nabla V(x) (\bar{u}_{2,k} + \bar{u}_{1,k}) \bar{\eta}_k dx \\ &= \frac{\varepsilon_k^2}{2} \sum_{j=1}^2 x_{2,k}^j \int_{B_\delta(x_{2,k})} \frac{\partial V(x)}{\partial x_j} (\bar{u}_{2,k} + \bar{u}_{1,k}) \bar{\eta}_k(x) dx = o(e^{-\frac{b\delta}{\varepsilon_k}}) \quad \text{as } k \rightarrow \infty, \end{aligned} \quad (\text{A.36})$$

where  $x_{2,k} := \{x_{2,k}^1, x_{2,k}^2\} \in \mathbb{R}^2$ . By the definition of  $\gamma_k$  in (4.37), it is sufficient to prove that

$$\frac{\beta_{2,k} - \beta_{1,k}}{\|\bar{u}_{2,k} - \bar{u}_{1,k}\|_{L^\infty(\mathbb{R}^2)}} = o(e^{-\frac{b\delta}{\varepsilon_k}}) \quad \text{as } k \rightarrow \infty. \quad (\text{A.37})$$

It follows from (4.35) that

$$\begin{aligned} & \frac{\beta_{2,k} - \beta_{1,k}}{\|\bar{u}_{2,k} - \bar{u}_{1,k}\|_{L^\infty(\mathbb{R}^2)}} \\ &= \varepsilon_k^2 \mu_{2,k} \int_{B_\delta^c(x_{2,k})} (\bar{u}_{2,k} + \bar{u}_{1,k}) \bar{\eta}_k dx + \frac{\mu_{2,k} - \mu_{1,k}}{\|\bar{u}_{2,k} - \bar{u}_{1,k}\|_{L^\infty(\mathbb{R}^2)}} \varepsilon_k^2 \int_{B_\delta^c(x_{2,k})} \bar{u}_{1,k}^2 dx \\ &+ \frac{1}{2} \varepsilon_k^2 \mu_{2,k} \int_{\partial B_\delta(x_{2,k})} (\bar{u}_{2,k} + \bar{u}_{1,k}) \bar{\eta}_k(x - x_{2,k}) \cdot \nu dS \\ &+ \frac{1}{2} \frac{\mu_{2,k} - \mu_{1,k}}{\|\bar{u}_{2,k} - \bar{u}_{1,k}\|_{L^\infty(\mathbb{R}^2)}} \varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} \bar{u}_{1,k}^2(x - x_{2,k}) \cdot \nu dS \\ &- \frac{1}{2} \varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} V(x) (\bar{u}_{2,k} + \bar{u}_{1,k}) \bar{\eta}_k(x - x_{2,k}) \cdot \nu dS \\ &- \frac{1}{2} \int_{B_\delta^c(x_{2,k})} \int_{\mathbb{R}^2} \frac{(x - x_{2,k}) \cdot (x - y)}{|x - y|^2} \bar{u}_{2,k}^2(x) (\bar{u}_{2,k}(y) + \bar{u}_{1,k}(y)) \bar{\eta}_k(y) dy dx \\ &- \frac{1}{2} \int_{B_\delta^c(x_{2,k})} \int_{\mathbb{R}^2} \frac{(x - x_{2,k}) \cdot (x - y)}{|x - y|^2} (\bar{u}_{2,k}(x) + \bar{u}_{1,k}(x)) \bar{\eta}_k(x) \bar{u}_{1,k}^2(y) dy dx \\ &- \int_{B_\delta^c(x_{2,k})} \int_{\mathbb{R}^2} \ln|x - y| \bar{u}_{2,k}^2(x) (\bar{u}_{2,k}(y) + \bar{u}_{1,k}(y)) \bar{\eta}_k(y) dy dx \\ &- \int_{B_\delta^c(x_{2,k})} \int_{\mathbb{R}^2} \ln|x - y| (\bar{u}_{2,k}(x) + \bar{u}_{1,k}(x)) \bar{\eta}_k(x) \bar{u}_{1,k}^2(y) dy dx \\ &- \frac{1}{2} \int_{\partial B_\delta(x_{2,k})} \int_{\mathbb{R}^2} \ln|x - y| \bar{u}_{2,k}^2(x) (\bar{u}_{2,k}(y) + \bar{u}_{1,k}(y)) \bar{\eta}_k(y) (x - x_{2,k}) \cdot \nu dy dS \\ &- \frac{1}{2} \int_{\partial B_\delta(x_{2,k})} \int_{\mathbb{R}^2} \ln|x - y| (\bar{u}_{2,k}(x) + \bar{u}_{1,k}(x)) \bar{\eta}_k(x) \bar{u}_{1,k}^2(y) (x - x_{2,k}) \cdot \nu dy dS \\ &+ \frac{1}{2} \int_{B_\delta^c(x_{2,k})} (\bar{u}_{2,k}^2 + \bar{u}_{1,k}^2) (\bar{u}_{2,k} + \bar{u}_{1,k}) \bar{\eta}_k dx \\ &+ \frac{1}{4} \int_{\partial B_\delta(x_{2,k})} (\bar{u}_{2,k}^2 + \bar{u}_{1,k}^2) (\bar{u}_{2,k} + \bar{u}_{1,k}) \bar{\eta}_k(x - x_{2,k}) \cdot \nu dS. \end{aligned}$$

We next estimate the right hand side of (A) as follows. By the exponential decay of  $\tilde{u}_{i,k}$ , we obtain from (A.32) that

$$\begin{aligned}
o(e^{-\frac{b\delta}{\varepsilon_k}}) &= \varepsilon_k^2 \mu_{2,k} \int_{B_\delta^c(x_{2,k})} (\bar{u}_{2,k} + \bar{u}_{1,k}) \bar{\eta}_k dx + \frac{\mu_{2,k} - \mu_{1,k}}{\|\bar{u}_{2,k} - \bar{u}_{1,k}\|_{L^\infty(\mathbb{R}^2)}} \varepsilon_k^2 \int_{B_\delta^c(x_{2,k})} \bar{u}_{1,k}^2 dx \\
&+ \frac{1}{2} \varepsilon_k^2 \mu_{2,k} \int_{\partial B_\delta(x_{2,k})} (\bar{u}_{2,k} + \bar{u}_{1,k}) \bar{\eta}_k(x - x_{2,k}) \cdot \nu dS \\
&+ \frac{1}{2} \frac{\mu_{2,k} - \mu_{1,k}}{\|\bar{u}_{2,k} - \bar{u}_{1,k}\|_{L^\infty(\mathbb{R}^2)}} \varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} \bar{u}_{1,k}^2(x - x_{2,k}) \cdot \nu dS \\
&- \frac{1}{2} \varepsilon_k^2 \int_{\partial B_\delta(x_{2,k})} V(x) (\bar{u}_{2,k} + \bar{u}_{1,k}) \bar{\eta}_k(x - x_{2,k}) \cdot \nu dS \\
&+ \frac{1}{2} \int_{B_\delta^c(x_{2,k})} (\bar{u}_{2,k}^2 + \bar{u}_{1,k}^2) (\bar{u}_{2,k} + \bar{u}_{1,k}) \bar{\eta}_k dx \\
&+ \frac{1}{4} \int_{\partial B_\delta(x_{2,k})} (\bar{u}_{2,k}^2 + \bar{u}_{1,k}^2) (\bar{u}_{2,k} + \bar{u}_{1,k}) \bar{\eta}_k(x - x_{2,k}) \cdot \nu dS \quad \text{as } k \rightarrow \infty.
\end{aligned} \tag{A.38}$$

Similar to (A.30), we obtain that there exists a constant  $b \in (0, a)$  such that

$$\begin{aligned}
&\int_{B_\delta^c(x_{2,k})} \int_{\mathbb{R}^2} \frac{(x - x_{2,k}) \cdot (x - y)}{|x - y|^2} \bar{u}_{2,k}^2(x) (\bar{u}_{2,k}(y) + \bar{u}_{1,k}(y)) \bar{\eta}_k(y) dy dx \\
&+ \int_{B_\delta^c(x_{2,k})} \int_{\mathbb{R}^2} \frac{(x - x_{2,k}) \cdot (x - y)}{|x - y|^2} (\bar{u}_{2,k}(x) + \bar{u}_{1,k}(x)) \bar{\eta}_k(x) \bar{u}_{1,k}^2(y) dy dx = o(e^{-\frac{b\delta}{\varepsilon_k}}) \quad \text{as } k \rightarrow \infty.
\end{aligned} \tag{A.39}$$

The same argument of (A.28) yields that

$$\begin{aligned}
&- \int_{B_\delta^c(x_{2,k})} \int_{\mathbb{R}^2} \ln|x - y| \bar{u}_{2,k}^2(x) (\bar{u}_{2,k}(y) + \bar{u}_{1,k}(y)) \bar{\eta}_k(y) dy dx \\
&- \int_{B_\delta^c(x_{2,k})} \int_{\mathbb{R}^2} \ln|x - y| (\bar{u}_{2,k}(x) + \bar{u}_{1,k}(x)) \bar{\eta}_k(x) \bar{u}_{1,k}^2(y) dy dx \\
&- \frac{1}{2} \int_{\partial B_\delta(x_{2,k})} \int_{\mathbb{R}^2} \ln|x - y| \bar{u}_{2,k}^2(x) (\bar{u}_{2,k}(y) + \bar{u}_{1,k}(y)) \bar{\eta}_k(y) (x - x_{2,k}) \cdot \nu dy dS \\
&- \frac{1}{2} \int_{\partial B_\delta(x_{2,k})} \int_{\mathbb{R}^2} \ln|x - y| (\bar{u}_{2,k}(x) + \bar{u}_{1,k}(x)) \bar{\eta}_k(x) \bar{u}_{1,k}^2(y) (x - x_{2,k}) \cdot \nu dy dS = o(e^{-\frac{b\delta}{\varepsilon_k}}) \quad \text{as } k \rightarrow \infty.
\end{aligned} \tag{A.40}$$

We conclude from (A.38)–(A.40) that (A.37) holds true, and the proof of Lemma A.6 is therefore complete.  $\square$

**Lemma A.7.** *Let  $\tilde{u}_{i,k}$  be as in (4.16) for  $i = 1, 2$ , and suppose  $\eta_k$  is a solution of (4.18). Then we have*

$$\int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x - y| \eta_k(x) (\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)) \tilde{u}_{2,k}^2(y) dx dy \rightarrow 2 \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x - y| \eta_0(x) Q(x) Q^2(y) dx dy \quad \text{as } k \rightarrow \infty,$$

where  $\eta_0(x)$  is defined by (4.17).

**Proof.** We note that

$$\begin{aligned}
&\int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x - y| \left[ \eta_k(x) (\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)) \tilde{u}_{2,k}^2(y) - 2\eta_0(x) Q(x) Q^2(y) \right] dx dy \\
&= \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x - y| \left( \eta_k(x) - \eta_0(x) \right) (\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)) \tilde{u}_{2,k}^2(y) dx dy \\
&+ \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x - y| \eta_0(x) \left( \tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x) - 2Q(x) \right) \tilde{u}_{2,k}^2(y) dx dy \\
&+ 2 \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x - y| \eta_0(x) Q(x) (\tilde{u}_{2,k}^2(y) - Q^2(y)) dx dy := I_{1,k} + I_{2,k} + I_{3,k}.
\end{aligned}$$



To prove Lemma A.7, it is sufficient to prove that for  $i = 1, 2, 3$ ,

$$I_{i,k} \rightarrow 0 \quad \text{as } k \rightarrow \infty. \quad (\text{A.41})$$

By the exponential decay of  $\tilde{u}_{i,k}$ , since  $\tilde{u}_{2,k} \rightarrow Q$  in  $L^\infty(\mathbb{R}^2) \cap X$  as  $k \rightarrow \infty$ , we have for any  $\epsilon > 0$ , there exists a large constant  $R > 0$  such that

$$\begin{aligned} & 2\rho_k \int_{B_R^c(0)} \ln(1 + |x|^2) |\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)| dx + 2 \int_{B_R^c(0)} |\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)| dx \\ & \cdot \left( \|\tilde{u}_{2,k}\|_*^2 + C\|\tilde{u}_{2,k}\|^2 \right) < \frac{\epsilon}{2} \quad \text{as } k \rightarrow \infty. \end{aligned} \quad (\text{A.42})$$

For the fixed constant  $R > 0$  in (A.42), since  $\tilde{u}_{2,k} \rightarrow Q$  in  $L^\infty(\mathbb{R}^2) \cap X$  and  $\eta_k \rightarrow \eta_0$  in  $C_{loc}(\mathbb{R}^2)$  as  $k \rightarrow \infty$ , we obtain from (4.6) that

$$\begin{aligned} & \|\eta_k - \eta_0\|_{C(B_R(0))} \left( \rho_k \int_{B_R(0)} \ln(1 + |x|^2) |\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)| dx \right. \\ & \left. + \int_{B_R(0)} |\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)| dx \cdot \left( \|\tilde{u}_{2,k}\|_*^2 + C\|\tilde{u}_{2,k}\|^2 \right) \right) < \frac{\epsilon}{2} \quad \text{as } k \rightarrow \infty. \end{aligned} \quad (\text{A.43})$$

Similar to (A), we obtain from (3.12), (A.42) and (A.43) that for any  $\epsilon > 0$ , there exists a large constant  $R > 0$  such that

$$\begin{aligned} |I_{1,k}| &= \left| \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| \left( \eta_k(x) - \eta_0(x) \right) (\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)) \tilde{u}_{2,k}^2(y) dx dy \right| \\ &\leq \rho_k \int_{\mathbb{R}^2} \ln(1 + |x|^2) |\eta_k(x) - \eta_0(x)| |\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)| dx \\ &\quad + \int_{\mathbb{R}^2} |\eta_k(x) - \eta_0(x)| |\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)| dx \int_{\mathbb{R}^2} \ln(1 + |y|^2) \tilde{u}_{2,k}^2(y) dy \\ &\quad + \int_{\mathbb{R}^2} |\eta_k(x) - \eta_0(x)| |\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)| \int_{\mathbb{R}^2} \frac{1}{|x-y|} \tilde{u}_{2,k}^2(y) dy dx \\ &\leq \rho_k \|\eta_k - \eta_0\|_{C(B_R(0))} \int_{B_R(0)} \ln(1 + |x|^2) |\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)| dx \\ &\quad + 2\rho_k \int_{B_R^c(0)} \ln(1 + |x|^2) |\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)| dx \\ &\quad + \left( \|\eta_k - \eta_0\|_{C(B_R(0))} \int_{B_R(0)} |\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)| dx + 2 \int_{B_R^c(0)} |\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x)| dx \right) \\ &\quad \cdot \left( \|\tilde{u}_{2,k}\|_*^2 + C\|\tilde{u}_{2,k}\|^2 \right) < \epsilon \quad \text{as } k \rightarrow \infty. \end{aligned} \quad (\text{A.44})$$

We thus have  $I_{1,k} \rightarrow 0$  as  $k \rightarrow \infty$ .

Similarly, one can obtain from (1.8), (3.12) and (4.6) that

$$\begin{aligned} |I_{2,k}| &= \left| \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| \eta_0(x) (\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x) - 2Q(x)) \tilde{u}_{2,k}^2(y) dx dy \right| \\ &\leq \rho_k \int_{\mathbb{R}^2} \ln(1 + |x|^2) |\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x) - 2Q(x)| dx \\ &\quad + \int_{\mathbb{R}^2} |\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x) - 2Q(x)| dx \int_{\mathbb{R}^2} \ln(1 + |y|^2) \tilde{u}_{2,k}^2(y) dy \\ &\quad + \int_{\mathbb{R}^2} |\tilde{u}_{2,k}(x) + \tilde{u}_{1,k}(x) - 2Q(x)| \int_{\mathbb{R}^2} \frac{1}{|x-y|} \tilde{u}_{2,k}^2(y) dy dx \rightarrow 0 \quad \text{as } k \rightarrow \infty, \end{aligned} \quad (\text{A.45})$$

and

$$\begin{aligned}
|I_{3,k}| &= \left| 2 \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| \eta_0(x) Q(x) (\tilde{u}_{2,k}^2(y) - Q^2(y)) dx dy \right| \\
&= \left| \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \left( \ln(1+|x-y|^2) - \ln\left(1 + \frac{1}{|x-y|^2}\right) \right) \eta_0(x) Q(x) (\tilde{u}_{2,k}^2(y) - Q^2(y)) dx dy \right| \\
&\leq 2 \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \left( \ln(1+|x|^2) + \ln(1+|y|^2) + \frac{1}{|x-y|} \right) |\eta_0(x) Q(x)| |\tilde{u}_{2,k}^2(y) - Q^2(y)| dx dy \quad (\text{A.46}) \\
&\leq 2 \|\tilde{u}_{2,k} - Q\|_2 \|\tilde{u}_{2,k} + Q\|_2 \int_{\mathbb{R}^2} \ln(1+|x|^2) Q(x) dx + 2 \|\tilde{u}_{2,k} - Q\|_* \|\tilde{u}_{2,k} + Q\|_* \int_{\mathbb{R}^2} Q dx \\
&\quad + C \|\tilde{u}_{2,k} + Q\| \|\tilde{u}_{2,k} - Q\| \int_{\mathbb{R}^2} Q dx \rightarrow 0 \text{ as } k \rightarrow \infty,
\end{aligned}$$

where we have used the fact that  $\tilde{u}_{i,k} \rightarrow Q$  in  $L^\infty(\mathbb{R}^2) \cap X$  as  $k \rightarrow \infty$ .

Following (A.44)–(A.46), we obtain that (A.41) holds true. This completes the proof of Lemma A.7.  $\square$

## References

- [1] A. Ambrosetti and D. Ruiz, *Multiple bound states for the Schrödinger-Poisson problem*, Comm. Contemp. Math. **10** (2008), 391–404.
- [2] T. Ando, A. B. Fowler and F. Stern, *Electronic properties of two-dimensional systems*, Rev. Mod. Phys. **54** (1982).
- [3] A. Arnold, *Self-consistent relaxation-time models in quantum mechanics*, Comm. Partial Differential Equations **21** (1996), 473–506.
- [4] A. Azzollini, *The planar Schrödinger-Poisson system with a positive potential*, Nonlinearity **34** (2021), 5799–5820.
- [5] A. Azzollini and A. Pomponio, *Ground state solutions for the nonlinear Schrödinger-Maxwell equations*, J. Math. Anal. Appl. **345** (2008), 90–108.
- [6] J. Bellazzini, L. Jeanjean and T. J. Luo, *Existence and instability of standing waves with prescribed norm for a class of Schrödinger-Poisson equations*, Proc. Lond. Math. Soc. **107** (2013), 303–339.
- [7] V. Benci and D. Fortunato, *Solitary waves of the nonlinear Klein-Gordon equation coupled with Maxwell equations*, Rev. Math. Phys. **14** (2002), 409–420.
- [8] V. Benci and D. Fortunato, *An eigenvalue problem for the Schrödinger-Maxwell equations*, Topol. Methods Nonlinear Anal. **11** (1998), 283–293.
- [9] R. Benguria, H. Brézis and E. H. Lieb, *The Thomas-Fermi-von Weizsäcker theory of atoms and molecules*, Comm. Math. Phys. **79** (1981), 167–180.
- [10] O. Bokanowski, J. L. López and J. Soler, *On an exchange interaction model for the quantum transport; the Schrödinger-Poisson-slater term*, Math. Model Methods Appl. Sci. **13** (2003), 1397–1412.
- [11] D. M. Cao, S. L. Li and P. Luo, *Uniqueness of positive bound states with multi-bump for nonlinear Schrödinger equations*, Calc. Var. Partial Differential Equations **54** (2015), 4037–4063.
- [12] I. Catto, J. Dolbeault, O. Sánchez and J. Soler, *Existence of steady states for the Maxwell-Schrödinger-Poisson system: exploring the applicability of the concentration-compactness principle*, Math. Models Methods Appl. Sci. **23** (2013), 1915–1938.
- [13] I. Catto and P.-L. Lions, *Binding of atoms and stability of molecules in Hartree and Thomas-Fermi type theories*, Comm. Partial Differential Equations **18** (1993), 1149–1159.

- [14] F. Castella,  *$L^2$  solutions to the Schrödinger-Poisson system: existence, uniqueness, time behaviour, and smoothing effects*, Math. Models Methods Appl. Sci. **7** (1997), 1051–1083.
- [15] T. Cazenave, *Semilinear Schrödinger equations*, Courant Lecture Notes in Mathematics Vol. 10, Courant Institute of Mathematical Science/AMS, New York, 2003.
- [16] S. Cingolani and L. Jeanjean, *Stationary waves with prescribed  $L^2$ -norm for the planar Schrödinger-Poisson system*, SIAM J. Math. Anal. **51** (2019), 3533–3568.
- [17] S. Cingolani and T. Weth, *On the planar Schrödinger-Poisson system*, Ann. Inst. H. Poincaré Anal. Non Linéaire **33** (2016), 169–197.
- [18] C. Y. Chen, Y.-C. Kuo and T.-F. Wu, *Existence and multiplicity of positive solutions for the nonlinear Schrödinger-Poisson equations*, Proc. Roy. Soc. Edinburgh Sect. A **143** (2013), 745–764.
- [19] T. D’Aprile and D. Mugnai, *Solitary waves for nonlinear Klein-Gordon-Maxwell and Schrödinger-Maxwell equations*, Proc. Roy. Soc. Edinburgh Sect. A **134** (2004), 893–906.
- [20] Y. B. Deng, C.-S. Lin and S. S. Yan, *On the prescribed scalar curvature problem in  $\mathbb{R}^N$ , local uniqueness and periodicity*, J. Math. Pures Appl. **104** (2015), 1013–1044.
- [21] J. Dolbeault, R. L. Frank and L. Jeanjean, *Logarithmic estimates for mean-field models in dimension two and the Schrödinger-Poisson system*, Comptes Rendus. Mathématique **10** (2021), 1279–1293.
- [22] S. Doniach and B. A. Huberman, *Topological excitations in two-dimensional superconductors*, Phys. Review Lett. **42** (1979), 1169.
- [23] M. Du and T. Weth, *Ground states and high energy solutions of the planar Schrödinger-Poisson system*, Nonlinearity **30** (2017), 3492–3515.
- [24] D. S. Fisher, *Flux-lattice melting in thin-film superconductors*, Phys. Review B **22** (1980), 1190.
- [25] H. Fröhlich, *Theory of electrical breakdown in ionic crystals*, Proc. Royal Soc. London Series A Math. Phys. Sciences **160** (1937), 230–241.
- [26] J. Garnier, K. Baudin, A. Fusaro and A. Picozzi, *Incoherent localized structures and hidden coherent solitons from the gravitational instability of the Schrödinger-Poisson equation*, Phys. Review E **104** (2021).
- [27] B. Gidas, W. M. Ni and L. Nirenberg, *Symmetry of positive solutions of nonlinear elliptic equations in  $\mathbb{R}^N$* , Adv. in Math. Suppl. Stud. **7** (1981), 369–402.
- [28] D. Gilbarg and N. S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Springer, 1997.
- [29] Y. J. Guo, C. S. Lin and J. C. Wei, *Local uniqueness and refined spike profiles of ground states for two-dimensional attractive Bose-Einstein condensates*, SIAM J. Math. Anal. **49** (2017), 3671–3715.
- [30] Y. J. Guo, Y. Luo and S. J. Peng, *Local uniqueness of ground states for rotating Bose-Einstein condensates with attractive interactions*, Calc. Var. Partial Differential Equations **60** (2021).
- [31] Y. J. Guo, Y. Luo and W. Yang, *The nonexistence of vortices for rotating Bose-Einstein condensates with attractive interactions*, Arch. Rational Mech. Anal. **238** (2020), 1231–1281.
- [32] Y. J. Guo and R. Seiringer, *On the mass concentration for Bose-Einstein condensates with attractive interactions*, Lett. Math. Phys. **104** (2014), 141–156.
- [33] Y. J. Guo, Z.-Q. Wang, X. Y. Zeng and H.-S. Zhou, *Properties of ground states of attractive Gross-pitaevskii equations with multi-well potentials*, Nonlinearity **31** (2018), 957–979.
- [34] Y. J. Guo, X. Y. Zeng and H.-S. Zhou, *Energy estimates and symmetry breaking in attractive Bose-Einstein condensates with ring-shaped potentials*, Ann. Inst. H. Poincaré Anal. Non Linéaire **33** (2016), 809–828.

- [35] Q. Han and F. H. Lin, *Elliptic Partial Differential Equations*, Courant Lecture Note in Math. 1, Courant Inst. Math. Science/AMS, New York, 2011.
- [36] A. F. Hebard and A. T. Fiory, *Recent experimental results on vortex processes in thin-film superconductors*, *Ordering in Two Dimensions* (1980), 181–187.
- [37] L. Jeanjean and T. T. Le, *Multiple normalized solutions for a Sobolev critical Schrödinger-Poisson-Slater equation*, *J. Differential Equations* **303** (2021), 277–325.
- [38] L. Jeanjean and T. J. Luo, *Sharp nonexistence results of prescribed  $L^2$ -norm solutions for some class of Schrödinger-Poisson and quasi-linear equations*, *Z. Angew. Math. Phys.* **64** (2013), 937–954.
- [39] M. K. Kwong, *Uniqueness of positive solutions of  $\Delta u - u + u^p = 0$  in  $\mathbb{R}^N$* , *Arch. Rational Mech. Anal.* **105** (1989), 243–266.
- [40] Y. K. Li, W. B. Wang and W. Zhang, *Minimizers of the planar Schrödinger-Newton equations*, *Complex Var. Elliptic Equ.* **67** (2022), 151–161.
- [41] E. H. Lieb, *Existence and uniqueness of the minimizing solution of Choquard’s nonlinear equation*, *Studies in Applied Math.* **57** (1977), 93–105.
- [42] E. H. Lieb, *Thomas-Fermi and related theories of atoms and molecules*, *Rev. Mod. Phys.* **53** (1981), 263–301.
- [43] E. H. Lieb and M. Loss, *Analysis*, Second edition, Graduate Studies in Mathematics, vol. 14, American Math. Society, Providence, RI, 2001.
- [44] P.-L. Lions, *Solutions of Hartree-Fock equations for Coulomb systems*, *Comm. Math. Phys.* **109** (1987), 33–97.
- [45] P. A. Markowich, C. Ringhofer and C. Schmeiser, *Semiconductor Equations*, Springer, New York, 1990.
- [46] C. Mercuri, V. Moroz and J. Van Schaftingen, *Groundstates and radial solutions to nonlinear Schrödinger-Poisson-Slater equations at the critical frequency*, *Calc. Var. Partial Differential Equations* **55** (2016), 1–58.
- [47] W.-M. Ni and I. Takagi, *On the shape of least-energy solutions to a semilinear Neumann problem*, *Comm. Pure Appl. Math.* **44** (1991), 819–851.
- [48] S. I. Pekar, *Untersuchungen über die Elektronentheorie der Kristalle*, Akademie-Verlag, Berlin, 1954.
- [49] R. Penrose, *On gravity’s role in quantum state reduction*, *Gen. Rel. Grav.* **28** (1996), 581–600.
- [50] R. Penrose, *Quantum computation, entanglement and state reduction*, *R. Soc. Lond. Philos. Trans. Ser. A Math. Phys. Eng. Sci.* **356** (1998), 1927–1939.
- [51] D. Ruiz, *Semiclassical states for coupled Schrödinger-Maxwell equations: Concentration around a sphere*, *Math. Model. Methods Appl. Sci.* **15** (2005), 141–164.
- [52] D. Ruiz, *The Schrödinger-Poisson equation under the effect of a nonlinear local term*, *J. Func. Anal.* **237** (2006), 655–674.
- [53] D. Ruiz, *On the Schrödinger-Poisson-Slater system: behavior of minimizers, radial and nonradial cases*, *Arch. Ration. Mech. Anal.* **198** (2010), 349–368.
- [54] O. Sánchez and J. Soler, *Long-time dynamics of the Schrödinger-Poisson-Slater system*, *J. Statist. Phys.* **114** (2004), 197–204.
- [55] J. Stubbe, *Bound states of two-dimensional Schrödinger-Newton equations*, preprint, arXiv:0807.4059.
- [56] M. I. Weinstein, *Nonlinear Schrödinger equations and sharp interpolation estimates*, *Comm. Math. Phys.* **87** (1982/83), 567–576.
- [57] M. Willem, *Minimax theorems*, *Progress in Nonlinear Differential Equations and their Applications*, Birkhäuser Boston, Inc., Boston, MA, 1996.