

SIMPLICITY OF CUNTZ-PIMSNER ALGEBRAS

MENEVŞE ERYÜZLÜ AND MARK TOMFORDE

ABSTRACT. We introduce a property of C^* -correspondences, which we call Condition (S), to serve as an analogue of Condition (L) of graphs. We use Condition (S) to prove a Cuntz-Krieger Uniqueness Theorem for Cuntz-Pimsner algebras and obtain sufficient conditions for simplicity of Cuntz-Pimsner algebras. We also prove that if \mathcal{Q} is a topological quiver with no sinks and $X(\mathcal{Q})$ is the associated C^* -correspondence, then $X(\mathcal{Q})$ satisfies Condition (S) if and only if \mathcal{Q} satisfies Condition (L). We also consider several examples to compare and contrast Condition (S) with Schweizer’s aperiodicity condition.

1. INTRODUCTION

In 1997 Pimsner introduced a method for constructing a C^* -algebra \mathcal{O}_X from a C^* -correspondence ${}_A X_A$. These C^* -algebras have been named *Cuntz-Pimsner algebras*, and a great deal of research has focused on understanding how the structure of the Cuntz-Pimsner algebra \mathcal{O}_X is determined by properties of the C^* -correspondence ${}_A X_A$. As with any class of C^* -algebras, one wishes to understand the ideal structure of Cuntz-Pimsner algebras and, in particular, to characterize when Cuntz-Pimsner algebras are simple. Partial results for simplicity of Cuntz-Pimsner algebras were obtained early in the subject’s history, with Kajiwara, Pinzari, and Watatani providing sufficient conditions for simplicity when A is unital and ${}_A X_A$ is full and finite projective with injective left action by compact operators [4, Corollary 4.4], and Muhly and Solel providing sufficient conditions for simplicity when A is unital and ${}_A X_A$ is projective and essential with injective left action [9, Theorem 1]. In 2001 Schweizer introduced a property for C^* -correspondences called *aperiodicity* and proved that if A is unital and ${}_A X_A$ is full with injective left action, then \mathcal{O}_X is simple if and only if ${}_A X_A$ is aperiodic and A has no nontrivial invariant ideals [14, Theorem 3.9]. Subsequently, Muhly and Solel wrote in [10] that Schweizer’s result was “the last word on simplicity” for Cuntz-Pimsner algebras. In this paper we hope to convince the reader that there remain many open questions concerning simplicity of Cuntz-Pimsner algebras, and Schweizer’s result (while significant and elegant) is far from the last word on the topic.

In the development of Cuntz-Pimsner algebras, graph C^* -algebras have provided useful intuition and guidance for the investigation of more general Cuntz-Pimsner algebras. There are two fundamental structure theorems in the theory of graph C^* -algebras: the Gauge-Invariant Uniqueness Theorem and the Cuntz-Krieger Uniqueness Theorem. Katsura has successfully extended the Gauge-Invariant Uniqueness Theorem to all Cuntz-Pimsner algebras [6, Theorem 6.4] and used this result to

Date: December 13, 2022.

2020 Mathematics Subject Classification. 46L55, 46L08.

Key words and phrases. C^* -algebras, Cuntz-Pimsner algebras, C^* -correspondences.

This work was supported by a grant from the Simons Foundation (#527708 to Mark Tomforde).

classify the gauge-invariant ideals in any Cuntz-Pimsner algebra [7, Theorem 8.6]. By contrast, up to this point there has been no extension of the Cuntz-Krieger Uniqueness Theorem to general Cuntz-Pimsner algebras.

A graph is said to satisfy Condition (L) if every cycle in the graph has an exit. The Cuntz-Krieger Uniqueness Theorem for graph C^* -algebras states that if a graph E satisfies Condition (L) and $\phi : C^*(E) \rightarrow A$ is a $*$ -homomorphism that is nonzero on all vertex projections, then ϕ is injective. The Cuntz-Krieger Uniqueness Theorem is used to obtain a characterization of simplicity for graph C^* -algebras, namely $C^*(E)$ is simple if and only if E satisfies Condition (L) and E contains no nontrivial saturated hereditary subsets.

It would be advantageous to generalize the Cuntz-Krieger Uniqueness Theorem to all Cuntz-Pimsner algebras. Among other applications, such a theorem would provide sufficient conditions for simplicity of general Cuntz-Pimsner algebras as an almost immediate consequence. The main obstacle to developing a generalized Cuntz-Krieger Uniqueness Theorem has been an inability to formulate an appropriate analogue of Condition (L) for C^* -correspondences. While versions of the Cuntz-Krieger Uniqueness Theorem have been obtained by Katusra for topological graph C^* -algebras [5, Theorem 5.12] and by Muhly and the second author for topological quiver C^* -algebras [12, Theorem 6.16], in these situations the C^* -correspondence is built from an object (e.g., graph, topological graph, quiver) providing notions of edges, paths, and cycles for which Condition (L) can be formulated. Up to this point, it has been unclear how to translate Condition (L) to an appropriate property entirely in terms of the C^* -correspondence.

Comparing Schweizer's simplicity result (i.e., a Cuntz-Pimsner algebra is simple if and only if the C^* -correspondence is aperiodic and A has no nontrivial invariant ideals) to the simplicity result for graph C^* -algebras (i.e., a graph C^* -algebra is simple if and only if the graph satisfies Condition (L) and the graph has no nontrivial saturated hereditary subsets), and using the fact that invariant ideals in A correspond to saturated hereditary subsets in the graph, it is natural to conjecture that aperiodicity is the appropriate analogue of Condition (L) for C^* -correspondences. However, we shall show in this paper that this is not the case. In particular, in Section 6 we show that even under the hypotheses of Schweizer's simplicity theorem there are graphs that do not satisfy Condition (L) but whose C^* -correspondences are aperiodic. Therefore aperiodicity is definitively not the correct condition for formulating a Cuntz-Krieger Uniqueness Theorem.

In this paper we make some of the first progress in establishing an analogue of Condition (L) for C^* -correspondences that do not necessarily come from underlying objects such as graphs, topological graphs, or quivers. In particular, in Section 3 we define a property of C^* -correspondences that we call Condition (S). We use Condition (S) to prove a generalization of the Cuntz-Krieger Uniqueness Theorem (see Theorem 3.8 and Corollary 3.9) for Cuntz-Pimsner algebras of C^* -correspondences with injective left action: If ${}_A X_A$ satisfies Condition (S) and $\rho : \mathcal{O}_X \rightarrow B$ is a $*$ -homomorphism with $\rho|_{\pi_A(A)}$ injective, then ρ is injective. In Section 4 we use this theorem to deduce sufficient conditions for simplicity of Cuntz-Pimsner algebras; in particular, we show in Theorem 4.3 that if ${}_A X_A$ satisfies Condition (S) and A has no nontrivial invariant ideals, then \mathcal{O}_X is simple. We mention that Condition (S) implicitly implies that the left action of A on ${}_A X_A$ is injective, but other than this

assumption of injective left action our theorems hold without imposing any additional hypotheses on ${}_A X_A$ or A . As a consequence, our simplicity theorem applies to substantially broader classes of C^* -correspondences than Schweizer's simplicity result or other existing simplicity results.

In Section 5 we prove that if \mathcal{Q} is a topological quiver with no sinks and $X(\mathcal{Q})$ is the associated C^* -correspondence, then $X(\mathcal{Q})$ satisfies Condition (S) if and only if \mathcal{Q} satisfies Condition (L). (The corresponding results for topological graphs and graphs follow as special cases.) This gives additional credibility to Condition (S) as an appropriate analogue of Condition (L).

In Section 6 we examine Schweizer's aperiodicity condition in more detail to understand its relationship with Condition (S) and also Condition (L). We produce examples showing that even for finite graphs with no sinks and no sources, aperiodicity is distinct from Condition (L). We also provide examples showing that if any of the hypotheses of Schweizer's theorem are removed, the conclusion of the theorem need not hold. This shows that aperiodicity is an inappropriate notion for characterizing simplicity of Cuntz-Pimsner algebras when any of the hypotheses of Schweizer's theorem are dropped. The conclusions we draw from our examples are summarized in Remark 6.20.

We end the paper with two open questions related to Condition (S), outlining what remains to be accomplished to develop precisely the correct analogue of Condition (L) for all C^* -correspondences.

2. PRELIMINARIES

Given a (right) Hilbert A -module X we let $\mathcal{L}(X)$ denote the C^* -algebra of adjointable operators on X , and we let $\mathcal{K}(X)$ denote the closed two-sided ideal of compact operators given by

$$\mathcal{K}(X) := \overline{\text{span}}\{\Theta_{\xi,\eta} : \xi, \eta \in X\}$$

where $\Theta_{\xi,\eta}$ is defined by $\Theta_{\xi,\eta}(\zeta) := \xi\langle\eta, \zeta\rangle_A$.

Definition 2.1. If A is a C^* -algebra, a C^* -correspondence over A is a (right) Hilbert A -module X together with a $*$ -homomorphism $\phi_X : A \rightarrow \mathcal{L}(X)$. We consider ϕ_X as providing a left action of A on X by $a \cdot x := \phi_X(a)x$. When no confusion arises, we shall often drop the subscript and write $\phi := \phi_X$ for ease of notation.

Definition 2.2. If X is a C^* -correspondence over A , a *representation* of X into a C^* -algebra B is a pair (ψ, π) consisting of a linear map $\psi : X \rightarrow B$ and a $*$ -homomorphism $\pi : A \rightarrow B$ satisfying

- (i) $\psi(\xi)^*\psi(\eta) = \pi(\langle\xi, \eta\rangle_A)$
- (ii) $\psi(\phi(a)\xi) = \pi(a)\psi(\xi)$
- (iii) $\psi(\xi \cdot a) = \psi(\xi)\pi(a)$

for all $\xi, \eta \in X$ and $a \in A$. We often write $(\psi, \pi) : (X, A) \rightarrow B$ to denote this representation, and we let $C^*(\psi, \pi)$ denote the C^* -subalgebra of B generated by $\psi(X) \cup \pi(A)$. For a representation $(\psi, \pi) : (X, A) \rightarrow B$ the relation (i)–(ii) above imply

$$C^*(\pi, t) = \overline{\text{span}}\{\psi^n(\xi)\psi^m(\eta)^* : \xi \in X^{\otimes n}, \eta \in X^{\otimes m}, n, m \geq 0\}.$$

A representation (ψ, π) is said to be *injective* if π is injective. Note that in this case ψ will be isometric (and hence also injective) since

$$\|\psi(\xi)\|^2 = \|\psi(\xi)^*\psi(\xi)\| = \|\pi(\langle \xi, \xi \rangle_A)\| = \|\langle \xi, \xi \rangle_A\| = \|\xi\|^2.$$

Definition 2.3. For a representation $(\psi, \pi) : (X, A) \rightarrow B$ there exists a $*$ -homomorphism $\pi^{(1)} : \mathcal{K}(X) \rightarrow B$ with the property that

$$\pi^{(1)}(\Theta_{\xi, \eta}) = \psi(\xi)\psi(\eta)^*.$$

See [13, p. 202], [4, Lemma 2.2], and [3, Remark 1.7] for details on the existence of this $*$ -homomorphism. Also note that if (ψ, π) is an injective representation, then $\pi^{(1)}$ is injective as well [3, Proposition 1.6(2)].

Definition 2.4. For an ideal I in a C^* -algebra A we define

$$I^\perp := \{a \in A : ab = 0 \text{ for all } b \in I\}$$

and we refer to I^\perp as the *annihilator of I in A* . If X is a C^* -correspondence over A , we define an ideal $J(X)$ of A by

$$J(X) := \phi^{-1}(\mathcal{K}(X)).$$

We also define an ideal J_X of A by

$$J_X := J(X) \cap (\ker \phi)^\perp.$$

Note that $J_X = J(X)$ when ϕ is injective, and that J_X is the maximal ideal of $J(X)$ on which the restriction of ϕ is an injection into $\mathcal{K}(X)$.

Definition 2.5. If X is a C^* -correspondence over A , we say that a representation $(\psi, \pi) : (X, A) \rightarrow B$ is *covariant* if

$$\pi^{(1)}(\phi(a)) = \pi(a) \quad \text{for all } a \in J_X.$$

We say that a covariant representation $(\psi_X, \pi_A) : (X, A) \rightarrow B$ is *universal* if whenever $(\psi, \pi) : (X, A) \rightarrow C$ is a covariant representation there exists a $*$ -homomorphism $\rho_{(\psi, \pi)} : C^*(\psi_X, \pi_A) \rightarrow C$ with the property that $\psi = \rho_{(\psi, \pi)} \circ \psi_X$ and $\pi = \rho_{(\psi, \pi)} \circ \pi_A$.

Definition 2.6. If X is a C^* -correspondence over A , the *Cuntz-Pimsner algebra*, which we shall denote \mathcal{O}_X , is the C^* -algebra $C^*(\psi_X, \pi_A)$ where (ψ_X, π_A) is a universal covariant representation of X . The existence of \mathcal{O}_X is established in [2, Proposition 1.3], and the universal property ensures it is unique up to isomorphism.

Remark 2.7. If \mathcal{O}_X is the Cuntz-Pimsner algebra of a C^* -correspondence ${}_A X_A$, and if $(\psi_X, \pi_A) : (X, A) \rightarrow \mathcal{O}_X$ is a universal covariant representation, then for any $z \in \mathbb{T}$ we have that $(z\psi_X, \pi_A)$ is also a universal covariant representation. Hence by the universal property, there exists a homomorphism $\gamma_z : \mathcal{O}_X \rightarrow \mathcal{O}_X$ such that $\gamma_z(\pi_A(a)) = \pi_A(a)$ for all $a \in A$ and $\gamma_z(\psi_X(\xi)) = z\psi_X(\xi)$ for all $\xi \in X$. Since $\gamma_{z^{-1}}$ is an inverse for this homomorphism, we see that γ_z is an automorphism. Thus we have an action $\gamma : \mathbb{T} \rightarrow \text{Aut } \mathcal{O}_X$ with the property that $\gamma_z(\pi_A(a)) = \pi_A(a)$ for all $a \in A$ and $\gamma_z(\psi_X(\xi)) = z\psi_X(\xi)$ for all $\xi \in X$. Furthermore, a routine $\epsilon/3$ -argument shows that γ is strongly continuous. We call γ the *gauge action on \mathcal{O}_X* .

Because \mathbb{T} is compact, averaging over γ with respect to normalized Haar measure gives an expectation E of \mathcal{O}_X onto the fixed point algebra \mathcal{O}_X^γ by

$$E(x) := \int_{\mathbb{T}} \gamma_z(x) dz \quad \text{for } x \in \mathcal{O}_X$$

where the integral denotes vector-valued integration in \mathcal{O}_X and dz denotes normalized Haar measure on \mathbb{T} . The map E is completely positive, has norm 1, and is faithful in the sense that $E(a^*a) = 0$ implies $a = 0$.

Remark 2.8. If X is a C^* -correspondence over A , we let $X^{\otimes n} := X \otimes_A \dots \otimes_A X$ denote the n -fold A -balanced tensor product of X with itself. We note that $X^{\otimes n}$ is a C^* -correspondence over A with right action $(x_1 \otimes \dots \otimes x_n) \cdot a = x_1 \otimes \dots \otimes x_n \cdot a$, left action $a \cdot (x_1 \otimes \dots \otimes x_n) = a \cdot x_1 \otimes \dots \otimes x_n$, and inner product defined inductively by

$$\langle x_1 \otimes \dots \otimes x_n, y_1 \otimes \dots \otimes y_n \rangle := \langle x_2 \otimes \dots \otimes x_n, \langle x_1, y_1 \rangle \cdot y_2 \otimes \dots \otimes y_n \rangle.$$

If $(\psi, \pi) : (X, A) \rightarrow B$ is a representation of X in a C^* -algebra B , then it is shown in [2, Lemma 3.6] that for each $n \in \mathbb{N}$ there is a linear map $\psi^{\otimes n} : X^{\otimes n} \rightarrow B$ satisfying

$$\psi^{\otimes n}(x_1 \otimes \dots \otimes x_n) = \psi(x_1) \dots \psi(x_n) \quad \text{for } x_1, \dots, x_n \in X.$$

Furthermore, $(\psi^{\otimes n}, \pi) : (X^{\otimes n}, A) \rightarrow B$ is a representation and there exists a $*$ -homomorphism $\pi^{(n)} : \mathcal{K}(X^{\otimes n}) \rightarrow B$ such that

$$\pi^{(n)}(\Theta_{x,y}) = \psi^{\otimes n}(x)\psi^{\otimes n}(y)^* \quad \text{for } x, y \in X^{\otimes n}.$$

Given a correspondence ${}_A X_A$ with injective left action $\phi_X : A \rightarrow \mathcal{L}(X)$ and a covariant representation $(\psi, \pi) : (X, A) \rightarrow B$ we let 1^k denote the identity operator on $X^{\otimes k}$ and define

$$\mathcal{C}_n := A \otimes 1^n + \mathcal{K}(X) \otimes 1^{n-1} + \mathcal{K}(X^{\otimes 2}) \otimes 1^{n-2} + \dots + \mathcal{K}(X^{\otimes n}) \subseteq \mathcal{L}(X^{\otimes n}).$$

Furthermore, the homomorphism $\kappa_n^{\psi, \pi} : \mathcal{C}_n \rightarrow B$ given by

$$\kappa_n^{\psi, \pi}(k_0 \otimes 1^n + k_1 \otimes 1^{n-1} + \dots + k_n) := \pi(k_0) + \pi^{(1)}(k_1) + \dots + \pi^{(n)}(k_n)$$

is injective whenever π is injective [2, Proposition 4.6]. It is also a fact that if $\mathcal{C} := \varinjlim \mathcal{C}_n$ under the isometric homomorphisms $c \in \mathcal{C}_n \mapsto c \otimes 1 \in \mathcal{C}_{n+1}$, then the homomorphisms $\kappa_n^{\psi, \pi} : \mathcal{C}_n \rightarrow B$ induce a map $\kappa^{\psi, \pi} : \mathcal{C} \rightarrow B$, and [2, Corollary 4.8] shows that $\kappa^{\psi, \pi}$ is injective when π is injective. Furthermore, [2, Corollary 4.9] shows that for a universal covariant representation $(\psi_X, \pi_A) : (X, A) \rightarrow \mathcal{O}_X$ we have that $\kappa^{\psi_X, \pi_A} : \mathcal{C} \rightarrow \mathcal{O}_X$ is an injective map onto the fixed point algebra \mathcal{O}_X^γ .

3. A CUNTZ-KRIEGER UNIQUENESS THEOREM FOR CUNTZ-PIMSNER ALGEBRAS

Definition 3.1. Let X be a C^* -correspondence over A , and let $(\psi_X, \pi_A) : (X, A) \rightarrow \mathcal{O}_X$ be a universal covariant representation of (X, A) into \mathcal{O}_X . We define a vector $\zeta \in X^{\otimes m}$ with $m \in \mathbb{N}$ to be *nonreturning* if whenever $n \in \{1, \dots, m-1\}$ and $\xi \in X^{\otimes n}$, then

$$\psi_X^{\otimes m}(\zeta)^* \psi_X^{\otimes n}(\xi) \psi_X^{\otimes m}(\zeta) = 0.$$

Lemma 3.2. *Let X be a C^* -correspondence over A and let $\zeta \in X^{\otimes m}$ be a nonreturning vector. If $(\psi, \pi) : (X, A) \rightarrow B$ is any covariant representation, then for any $n \in \{1, \dots, m-1\}$ and $\xi \in X^{\otimes n}$, we have*

$$\psi^{\otimes m}(\zeta)^* \psi^{\otimes n}(\xi) \psi^{\otimes m}(\zeta) = 0.$$

Proof. Let $(\psi_X, \pi_A) : (X, A) \rightarrow \mathcal{O}_X$ be a universal covariant representation of (X, A) into \mathcal{O}_X . If $\rho := \rho_{(\psi, \pi)} : \mathcal{O}_X \rightarrow B$ denotes the $*$ -homomorphism induced by (ψ, π) , then

$$\begin{aligned} \psi^{\otimes m}(\zeta)^* \psi^{\otimes n}(\xi) \psi^{\otimes m}(\zeta) &= (\rho \circ \psi_X^{\otimes m})(\zeta)^* (\rho \circ \psi_X^{\otimes n})(\xi) (\rho \circ \psi_X^{\otimes m})(\zeta) \\ &= \rho(\psi_X^{\otimes m}(\zeta)^* \psi_X^{\otimes n}(\xi) \psi_X^{\otimes m}(\zeta)) = \rho(0) = 0. \quad \square \end{aligned}$$

Condition (S): We say that a C^* -correspondence X over a C^* -algebra A satisfies Condition (S) if whenever $a \in A$ with $a \geq 0$, then for all $n \in \mathbb{N}$ and for all $\epsilon > 0$ there exists $m > n$ and a nonreturning vector $\zeta \in X^{\otimes m}$ with $\|\zeta\| = 1$ and

$$\|\langle \zeta, a\zeta \rangle\| > \|a\| - \epsilon.$$

Remark 3.3. The (S) in Condition (S) is meant to stand for ‘‘simplicity’’.

Proposition 3.4. *If X is a C^* -correspondence over A satisfying Condition (S), then the left action $\phi_X : A \rightarrow \mathcal{L}(X)$ is injective.*

Proof. For each $n \in \mathbb{N}$ let $\phi_n : A \rightarrow \mathcal{L}(X^{\otimes n})$ denote the $*$ -homomorphism defined by $\phi_n(a)(x_1 \otimes \dots \otimes x_n) := a \cdot x_1 \otimes \dots \otimes x_n$. Also define $h : \mathcal{L}(X^{\otimes n}) \rightarrow \mathcal{L}(X^{\otimes n+1})$ by $h(T) := T \otimes I$. Then h is a $*$ -homomorphism, and hence $\|h(T)\| \leq \|T\|$ for all $T \in \mathcal{L}(X^{\otimes n})$. Furthermore, for any $a \in A$ and $n \in \mathbb{N}$, we have $h(\phi_n(a)) = \phi_{n+1}(a)$. Thus $\|\phi_{n+1}(a)\| \leq \|\phi_n(a)\|$. By induction, we conclude that

$$\|\phi_n(a)\| \leq \|\phi_1(a)\| = \|\phi_X(a)\| \quad \text{for all } a \in A \text{ and } n \in \mathbb{N}.$$

For any $b \in A$, let $a := b^*b \geq 0$. For any $\epsilon > 0$, Condition (S) implies there exists $m \in \mathbb{N}$ and a nonreturning vector $\zeta \in X^{\otimes m}$ with $\|\zeta\| = 1$ and

$$\|\langle \zeta, \phi_m(a)\zeta \rangle\| > \|a\| - \epsilon.$$

Since $\|\zeta\| = 1$, we have

$$\begin{aligned} \|\phi_X(b)\|^2 &= \|\phi_1(b)\|^2 \geq \|\phi_m(b)\|^2 \geq \|\phi_m(b)\zeta\|^2 = \|\langle \phi_m(b)\zeta, \phi_m(b)\zeta \rangle\| \\ &= \|\langle \zeta, \phi_m(b^*b)\zeta \rangle\| = \|\langle \zeta, \phi_m(a)\zeta \rangle\| > \|a\| - \epsilon = \|b^*b\| - \epsilon = \|b\|^2 - \epsilon. \end{aligned}$$

Since $\epsilon > 0$ was arbitrary, we deduce $\|\phi_X(b)\|^2 \geq \|b\|^2$ and $\|\phi_X(b)\| \geq \|b\|$. Moreover, because ϕ is a $*$ -homomorphism, and hence norm decreasing, we conclude $\|\phi_X(b)\| = \|b\|$, so that ϕ_X is isometric and hence also injective. \square

Remark 3.5. Proposition 3.4 shows that any C^* -correspondence satisfying Condition (S) necessarily has injective left action. The reader should keep this in mind, and note that in all forthcoming results the hypothesis that a C^* -correspondence satisfies Condition (S) implicitly requires the C^* -correspondence to have injective left action.

Lemma 3.6. *Let X be a C^* -correspondence over A that satisfies Condition (S), and let $(\psi_X, \pi_A) : (X, A) \rightarrow \mathcal{O}_X$ be a universal covariant representation of ${}_A X_A$ in \mathcal{O}_X . Also let $(\psi, \pi) : (X, A) \rightarrow B$ be a covariant representation of ${}_A X_A$ into a C^* -algebra B with the property that $\pi : A \rightarrow B$ is injective. If*

$$x = \sum_{i=1}^N \psi_X^{\otimes n_i}(\xi_i) \psi_X^{\otimes m_i}(\eta_i)^*$$

is positive in \mathcal{O}_X , then

$$y_0 := \sum_{n_i=m_i} \psi^{\otimes n_i}(\xi_i) \psi^{\otimes m_i}(\eta_i)^*$$

is positive in B , and if $\rho := \rho_{(\psi, \pi)} : \mathcal{O}_X \rightarrow B$ denotes the $*$ -homomorphism induced by (ψ, π) , then for all $\epsilon > 0$ there exists $a \in A$ and $b \in B$ with $\|a\| = \|y_0\|$, $\|b\| \leq 1$, and

$$\|b^* \rho(x) b - \pi(a)\| < \epsilon.$$

Proof. Let $E : \mathcal{O}_X \rightarrow \mathcal{O}_X^\gamma$ denote the conditional expectation determined by the gauge action γ on \mathcal{O}_X and given by $E(x) := \int_{\mathbb{T}} \gamma_z(x) dz$. Then

$$\begin{aligned} y_0 &= \sum_{n_i=m_i} \psi^{\otimes n_i}(\xi_i) \psi^{\otimes m_i}(\eta_i)^* = \sum_{n_i=m_i} (\rho \circ \psi_X)^{\otimes n_i}(\xi_i) (\rho \circ \psi_X)^{\otimes m_i}(\eta_i)^* \\ &= \rho \left(\sum_{n_i=m_i} \psi_X^{\otimes n_i}(\xi_i) \psi_X^{\otimes m_i}(\eta_i)^* \right) = \rho(E(x)). \end{aligned}$$

Since E is a conditional expectation, E is completely positive (as well as positive). Likewise, the $*$ -homomorphism ρ is positive. Since x is positive in \mathcal{O}_X , it follows that $y_0 = \rho(E(x))$ is positive in B .

Given $\epsilon > 0$, set $n := \max\{n_1, \dots, n_N, m_1, \dots, m_N\}$. By [2, Proposition 4.6] the $*$ -homomorphism $\kappa_n^{\psi, \pi} : \mathcal{C}_n \rightarrow B$ is injective and hence isometric. Therefore, since $y_0 = \sum_{n_i=m_i} \psi^{\otimes n_i}(\xi_i) \psi^{\otimes m_i}(\eta_i)^* \in \text{im } \kappa_n^{\psi, \pi}$, there exists $T \in \mathcal{C}_n \subseteq \mathcal{L}(X^{\otimes n})$ with $\kappa_n^{\psi, \pi}(T) = y_0$ and $\|T\| = \|y_0\|$. Moreover, since y_0 is positive in B , we may choose T positive in \mathcal{C}_n : since $y_0 \in \text{im } \kappa_n^{\psi, \pi} \subseteq B$ is positive in B , and hence positive in $\text{im } \kappa_n^{\psi, \pi}$, we may write $y_0 = v^* v$ for $v \in \text{im } \kappa_n^{\psi, \pi}$, choose $S \in \mathcal{C}_n$ with $\kappa_n^{\psi, \pi}(S) = v$, and let $T := S^* S \geq 0$. Then $\kappa_n^{\psi, \pi}(T) = \kappa_n^{\psi, \pi}(S^* S) = \kappa_n^{\psi, \pi}(S)^* \kappa_n^{\psi, \pi}(S) = v^* v = y_0$.

Since $\|T\| = \|y_0\|$ and T is positive, there exists $\xi \in X^{\otimes n}$ with $\|\xi\| = 1$ such that $g := \langle \xi, T\xi \rangle$ is a positive element of A with $\|g\| > \|y_0\| - \epsilon/2$. When $n_i > m_i$ we have

$$\psi^{\otimes n}(\xi)^* \psi^{\otimes n_i}(\xi_i) \psi^{\otimes m_i}(\eta_i)^* \psi^{\otimes n}(\xi) = \psi^{\otimes n'_i}(\xi'_i)$$

for some $\xi'_i \in X^{\otimes n'_i}$, where $n'_i := n_i - m_i$. Similarly, when $n_i < m_i$ we have

$$\psi^{\otimes n}(\xi)^* \psi^{\otimes n_i}(\xi_i) \psi^{\otimes m_i}(\eta_i)^* \psi^{\otimes n}(\xi) = \psi^{\otimes m'_i}(\eta'_i)^*$$

for some $\eta'_i \in X^{\otimes m'_i}$, where $m'_i := m_i - n_i$. Also, when $n_i = m_i$ we have

$$\begin{aligned} \psi^{\otimes n}(\xi)^* \left(\sum_{n_i=m_i} \psi^{\otimes n_i}(\xi_i) \psi^{\otimes m_i}(\eta_i)^* \right) \psi^{\otimes n}(\xi) &= \psi^{\otimes n}(\xi)^* y_0 \psi^{\otimes n}(\xi) \\ &= \psi^{\otimes n}(\xi)^* \kappa_n^{\psi, \pi}(T) \psi^{\otimes n}(\xi) = \psi^{\otimes n}(\xi)^* \psi^{\otimes n}(T\xi) = \pi(\langle \xi, T\xi \rangle) = \pi(g). \end{aligned}$$

Combining these three equations yields

$$\begin{aligned} (3.1) \quad \psi^{\otimes n}(\xi)^* \rho(x) \psi^{\otimes n}(\xi) &= \psi^{\otimes n}(\xi)^* \left(\sum_{i=1}^N \psi^{\otimes n_i}(\xi_i) \psi^{\otimes m_i}(\eta_i)^* \right) \psi^{\otimes n}(\xi) \\ &= \pi(g) + \sum_{n_i > m_i} \psi^{\otimes n'_i}(\xi'_i) + \sum_{n_i < m_i} \psi^{\otimes m'_i}(\eta'_i)^*. \end{aligned}$$

Since $g \geq 0$, by Condition (S) there exists $m > n$ and a nonreturning path $\zeta \in X^{\otimes m}$ with $\|\zeta\| = 1$ and $\|\langle \zeta, g\zeta \rangle\| > \|g\| - \epsilon/2$.

Let $c := \langle \zeta, g\zeta \rangle \in A$. Then

$$\|c\| = \|\langle \zeta, g\zeta \rangle\| > \|g\| - \epsilon/2 > (\|y_0\| - \epsilon/2) - \epsilon/2 = \|y_0\| - \epsilon.$$

In addition,

$$\|c\| = \|\langle \zeta, g\zeta \rangle\| \leq \|g\| \|\langle \zeta, \zeta \rangle\| = \|g\| = \|\langle \xi, T\xi \rangle\|$$

$$\leq \|T\| \|\langle \zeta, \zeta \rangle\| = \|T\| = \|y_0\| < \|y_0\| + \epsilon.$$

These two inequalities show that

$$\left| \|c\| - \|y_0\| \right| < \epsilon.$$

Let $b := \psi^{\otimes n}(\xi)\psi^{\otimes m}(\zeta)$. Then $\|b\| \leq \|\psi^{\otimes n}(\xi)\| \|\psi^{\otimes m}(\zeta)\| = \|\xi\| \|\zeta\| = 1 \cdot 1 = 1$. In addition,

$$\begin{aligned} b^* \rho(x) b &= \psi^{\otimes m}(\zeta)^* \psi^{\otimes n}(\xi)^* \rho(x) \psi^{\otimes n}(\xi) \psi^{\otimes m}(\zeta) \\ &= \psi^{\otimes m}(\zeta)^* \left(\pi(g) + \sum_{n_i > m_i} \psi^{\otimes n'_i}(\xi'_i) + \sum_{n_i < m_i} \psi^{\otimes m'_i}(\eta'_i)^* \right) \psi^{\otimes m}(\zeta) \quad (\text{by (3.1)}) \\ &= \psi^{\otimes m}(\zeta)^* \pi(g) \psi^{\otimes m}(\zeta) \quad (\text{since } \zeta \text{ is a nonreturning vector}) \\ &= \psi^{\otimes m}(\zeta)^* \psi^{\otimes m}(g\zeta) \\ &= \pi(\langle \zeta, g\zeta \rangle) \\ &= \pi(c). \end{aligned}$$

Let $a := \frac{\|y_0\|}{\|c\|}c$. Then $\|a\| = \|y_0\|$, and

$$\begin{aligned} \|b^* \rho(x) b - \pi(a)\| &= \|\pi(c) - \pi(a)\| = \|\pi(c - a)\| \leq \|c - a\| = \left\| c - \frac{\|y_0\|}{\|c\|}c \right\| \\ &= \left| 1 - \frac{\|y_0\|}{\|c\|} \right| \|c\| = \left| \|c\| - \|y_0\| \right| < \epsilon. \quad \square \end{aligned}$$

Lemma 3.7. *Let A_0 be a dense $*$ -subalgebra of a C^* -algebra A . If $x \in A$ and $x \geq 0$, then for every $\epsilon > 0$ there exists $x_0 \in A_0$ with $x_0 \geq 0$ and $\|x - x_0\| < \epsilon$.*

Proof. If $x = 0$, the result is trivial, so assume $x \neq 0$. Since $x \geq 0$, we may write $x = v^*v$ for some nonzero $v \in A$. Using the fact that A_0 is dense in A , choose $v_0 \in A_0$ with

$$\|v - v_0\| \leq \min \left\{ \frac{\epsilon}{3\|v\|}, \sqrt{\frac{\epsilon}{3}} \right\}.$$

Let $x_0 := v_0^*v_0 \in A_0$. Then $x_0 \geq 0$, and

$$\begin{aligned} \|x - x_0\| &= \|v^*v - v_0^*v_0\| \\ &\leq \|v^*v - v^*v_0\| + \|v^*v_0 - v_0^*v_0\| \\ &\leq \|v^*\| \|v - v_0\| + \|v^* - v_0^*\| \|v_0\| \\ &= \|v\| \|v - v_0\| + \|v - v_0\| \|v_0\| \\ &\leq \|v\| \frac{\epsilon}{3\|v\|} + \|v - v_0\| \|v_0 - v + v\| \\ &\leq \epsilon/3 + \|v - v_0\| (\|v_0 - v\| + \|v\|) \\ &\leq \epsilon/3 + \|v - v_0\|^2 + \|v\| \|v - v_0\| \\ &\leq \epsilon/3 + \left(\sqrt{\frac{\epsilon}{3}} \right)^2 + \|v\| \frac{\epsilon}{3\|v\|} \\ &= \epsilon/3 + \epsilon/3 + \epsilon/3 \\ &= \epsilon. \quad \square \end{aligned}$$

Theorem 3.8 (Cuntz-Krieger Uniqueness for Cuntz-Pimsner Algebras). *Let X be a C^* -correspondence over A satisfying Condition (S), and let $(\psi_X, \pi_A) : (X, A) \rightarrow \mathcal{O}_X$ be a universal covariant representation. If $\rho : \mathcal{O}_X \rightarrow B$ is a $*$ -homomorphism from \mathcal{O}_X into a C^* -algebra B with the property that the restriction $\rho|_{\pi_A(A)}$ is injective, then ρ is injective.*

Proof. Let γ denote the gauge action on \mathcal{O}_X , and let $E : \mathcal{O}_X \rightarrow \mathcal{O}_X^\gamma$ denote the conditional expectation given by

$$E(x) := \int_{\mathbb{T}} \gamma_z(x) dz.$$

We shall show the following statements hold:

- (a) ρ is injective on the fixed point algebra \mathcal{O}_X^γ , and
- (b) $\|\rho(E(x))\| \leq \|\rho(x)\|$ whenever $x \in \mathcal{O}_X$ and $x \geq 0$.

To see (a), let $\psi := \rho \circ \psi_X$ and $\pi := \rho \circ \pi_A$. Then $\rho = \rho_{(\psi, \pi)}$; that is, ρ equal to the $*$ -homomorphism induced by the representation (ψ, π) . For each $n \in \mathbb{N}$, we have $\kappa_n^{\psi, \pi} = \rho \circ \kappa_n^{\psi_X, \pi_A}$, and hence $\kappa^{\psi, \pi} = \rho \circ \kappa^{\psi_X, \pi_A}$. Since $\kappa^{\psi, \pi}$ is injective by [2, Corollary 4.8], it follows that ρ is injective on $\text{im } \kappa^{\psi_X, \pi_A} = \mathcal{O}_X^\gamma$, so (a) holds.

To see (b), note that since

$$\mathcal{O}_X = \overline{\text{span}}\{\psi_X^{\otimes n}(\xi)\psi_X^{\otimes m}(\eta)^* : \xi \in X^{\otimes n}, \eta \in X^{\otimes m}, \text{ and } n, m \geq 0\},$$

Lemma 3.7 implies that any positive element in \mathcal{O}_X may be approximated by a positive element from $\text{span}\{\psi_X^{\otimes n}(\xi)\psi_X^{\otimes m}(\eta)^* : \xi \in X^{\otimes n}, \eta \in X^{\otimes m}, \text{ and } n, m \geq 0\}$. Hence it suffices to prove (b) when x is positive and has the form

$$x = \sum_{i=1}^N \psi_X^{\otimes n_i}(\xi_i)\psi_X^{\otimes m_i}(\eta_i)^*.$$

Keeping the notation $(\psi, \pi) = (\rho \circ \psi_X, \rho \circ \pi_A)$, if we let

$$y := \rho(x) = \sum_{i=1}^N \psi^{\otimes n_i}(\xi_i)\psi^{\otimes m_i}(\eta_i)^*$$

and

$$y_0 := \sum_{n_i=m_i} \psi^{\otimes n_i}(\xi_i)\psi^{\otimes m_i}(\eta_i)^*,$$

then Lemma 3.6 shows that for all $\epsilon > 0$ there exists $a \in A$ and $b \in B$ such that $\|a\| = \|y_0\|$, $\|b\| \leq 1$, and $\|b^*yb - \pi(a)\| < \epsilon$. Thus

$$\begin{aligned} \|\rho(E(x))\| &= \|\rho\left(\sum_{n_i=m_i} \psi_X^{\otimes n_i}(\xi_i)\psi_X^{\otimes m_i}(\eta_i)^*\right)\| = \left\|\sum_{n_i=m_i} \psi^{\otimes n_i}(\xi_i)\psi^{\otimes m_i}(\eta_i)^*\right\| \\ &= \|y_0\| = \|a\| = \|\rho(\pi_A(a))\| = \|\pi(a)\| \leq \|b^*yb\| + \|\pi(a) - b^*yb\| \\ &\leq \|b^*\| \|y\| \|b\| + \epsilon \leq \|y\| + \epsilon = \|\rho(x)\| + \epsilon. \end{aligned}$$

Since this inequality holds for all $\epsilon > 0$, we have that $\|\rho(E(x))\| \leq \|\rho(x)\|$. Hence (b) holds.

Finally, given (a) and (b) we see that whenever $\rho(x) = 0$, we have $\rho(x^*x) = 0$ and (b) implies that $\rho(E(x^*x)) = 0$. Thus $E(x^*x) = 0$ by (a), and since E is a faithful conditional expectation, this implies $x = 0$. Thus ρ is injective. \square

Corollary 3.9. *Let X be a C^* -correspondence over A satisfying Condition (S). If $(\psi, \pi) : (X, A) \rightarrow B$ is a covariant representation with the property that π is injective, then the induced $*$ -homomorphism $\rho_{(\psi, \pi)} : \mathcal{O}_X \rightarrow B$ is injective.*

4. SIMPLICITY OF CUNTZ-PIMSNER ALGEBRAS

All ideals in C^* -algebras that we consider will be closed and two-sided. Let X be a C^* -correspondence over A . For any ideal I of A , we define

$$IX := \overline{\text{span}}\{ax : a \in I \text{ and } x \in X\}$$

and

$$XI := \overline{\text{span}}\{xa : a \in I \text{ and } x \in X\}.$$

We also define

$$X_I := \{x \in X : \langle y, x \rangle \in I \text{ for all } y \in X\}.$$

Lemma 4.1. *Let X be a C^* -correspondence over A . For any ideal I of A we have $XI = X_I$.*

Proof. Let $\xi \in XI := \overline{\text{span}}\{xa : a \in I \text{ and } x \in X\}$. Then $\xi = \lim_{n \rightarrow \infty} \sum_{i=1}^{N_n} x_i^n a_i^n$ with each $x_i^n \in X$ and each $a_i^n \in I$. For any $y \in X$ we have

$$\langle y, \xi \rangle = \langle y, \lim_{n \rightarrow \infty} \sum_{i=1}^{N_n} x_i^n a_i^n \rangle = \lim_{n \rightarrow \infty} \langle y, \sum_{i=1}^{N_n} x_i^n a_i^n \rangle = \lim_{n \rightarrow \infty} \sum_{i=1}^{N_n} \langle y, x_i^n \rangle a_i^n \in I.$$

Thus $\xi \in X_I$, and $XI \subseteq X_I$. For the reverse containment, let $\xi \in X_I = \{x \in X : \langle y, x \rangle \in I \text{ for all } y \in X\}$. By [8, Lemma 4.4] there exists $\eta \in X$ such that $\xi = \eta \langle \xi, \xi \rangle^{1/3}$. Since $\xi \in X_I$, we conclude $\langle \xi, \xi \rangle \in I$ and $\langle \xi, \xi \rangle^{1/3} \in I$. Thus $\xi = \eta \langle \xi, \xi \rangle^{1/3} \in XI$, and $X_I \subseteq XI$. \square

Let X be a C^* -correspondence over A . If I is an ideal in A , we say I is *hereditary* if $\phi(I)X \subseteq XI$. We also say I is *saturated* if for all $a \in J_X$, $\phi(a)X \subseteq XI$ implies $a \in I$. We call an ideal *invariant* when it is both saturated and hereditary.

Lemma 4.2. *Let X be a C^* -correspondence over A and let $(\psi_X, \pi_A) : (X, A) \rightarrow \mathcal{O}_X$ be a universal covariant representation. If \mathcal{I} is an ideal in \mathcal{O}_X , then $\pi_A^{-1}(\mathcal{I})$ is an invariant ideal in A .*

Proof. Let $I := \pi_A^{-1}(\mathcal{I})$. The fact that I is an ideal in A follows from the fact π_A is a $*$ -homomorphism. If $a \in I$ and $x \in X$, then for any $y \in X$ we have

$$\pi_A(\langle y, \phi(a)x \rangle) = \psi_X(y)^* \psi_X(ax) = \psi_X(y)^* \pi_A(a) \psi_X(x) \in \mathcal{I}.$$

Thus $\phi(a)x \in X_I = XI$. Hence $\phi(I)X \subseteq XI$, and I is hereditary.

In addition, let $a \in J_X$ with $\phi(a)X \subseteq XI$. Since

$$\begin{aligned} \pi_A^{(1)}(\mathcal{K}(X)) &= \pi_A^{(1)}(\overline{\text{span}}\{\theta_{x,y} : x, y \in X\}) \\ &= \overline{\text{span}}\{\pi_A^{(1)}(\theta_{x,y}) : x, y \in X\} = \overline{\text{span}}\{\psi_X(x) \psi_X(y)^* : x, y \in X\} \end{aligned}$$

we may choose an approximate unit $\{e_\lambda\}_{\lambda \in \Lambda}$ for $\pi^{(1)}(\mathcal{K}(X))$ with

$$e_\lambda := \sum_{i=1}^{N_\lambda} \psi_X(x_i^\lambda) \psi_X(y_i^\lambda)^* \quad \text{for each } \lambda \in \Lambda.$$

Furthermore, since $a \in J_X$, we conclude $\pi_A(a) = \pi^{(1)}(\phi(a)) \in \pi_A(\mathcal{K}(X))$, and

$$\pi_A(a) = \lim_{\lambda} e_{\lambda} \pi_A(a) e_{\lambda}.$$

Because $\phi(a)X \subseteq XI = X_I$, for any $x, y \in X$ we have $\phi(a)x \in X_I$ and $\langle y, \phi(a)x \rangle \in I$. Thus for all $x, y \in X$,

$$\psi_X(y)^* \pi_A(a) \psi_X(x) = \pi_A(\langle y, \phi(a)x \rangle) \in \mathcal{I}$$

and since \mathcal{I} is an ideal, it follows that $e_{\lambda} \pi_A(a) e_{\lambda} \in \mathcal{I}$ for all $\lambda \in \Lambda$. Hence $\pi_A(a) = \lim_{\lambda} e_{\lambda} \pi_A(a) e_{\lambda} \in \mathcal{I}$, so that $a \in I$ and I is saturated. Therefore I is invariant. \square

Theorem 4.3 (Simplicity of Cuntz-Pimsner Algebras). *If a C^* -correspondence ${}_A X_A$ satisfies Condition (S) and A has no invariant ideals other than A and $\{0\}$, then \mathcal{O}_X is simple.*

Proof. Let $(\psi_X, \pi_A) : (X, A) \rightarrow \mathcal{O}_X$ be a universal covariant representation. Suppose \mathcal{I} is an ideal in \mathcal{O}_X with $\mathcal{I} \neq \mathcal{O}_X$. Let $I := \pi_A^{-1}(\mathcal{I})$. By Lemma 4.2, I is an invariant ideal of A . Hence our hypothesis implies $I = A$ or $I = \{0\}$. Since $\mathcal{I} \neq \mathcal{O}_X$, we conclude $I \neq A$ and therefore $I = \{0\}$. Define $Q_{\mathcal{I}} : \mathcal{O}_X \rightarrow \mathcal{O}_X/\mathcal{I}$ to be the quotient map $Q_{\mathcal{I}}(x) = x + \mathcal{I}$. Then for any $a \in A$ we have

$$Q_{\mathcal{I}}(\pi_A(a)) = 0 \iff \pi_A(a) + \mathcal{I} = 0 + \mathcal{I} \iff \pi_A(a) \in \mathcal{I} \iff a \in I \iff a = 0.$$

Hence $Q_{\mathcal{I}}|_{\pi_A(A)}$ is injective, and Theorem 3.8 implies $Q_{\mathcal{I}} : \mathcal{O}_X \rightarrow \mathcal{O}_X/\mathcal{I}$ is injective. But the only way this can occur is if $\mathcal{I} = \{0\}$. Hence \mathcal{O}_X is simple. \square

Remark 4.4. Theorem 4.3 gives sufficient conditions for simplicity of a Cuntz-Pimsner algebra. However, at this time we do not know if Condition (S) is necessary for simplicity. If it is, then Condition (S) would be an appropriate generalization of Condition (L) to C^* -correspondences with injective left action.

5. TOPOLOGICAL QUIVERS AND CONDITION (L)

Definition 5.1. A topological quiver is a quintuple $\mathcal{Q} = (E^0, E^1, r, s, \lambda)$ consisting of a second countable locally compact Hausdorff space E^0 (whose elements are called vertices), a second countable locally compact Hausdorff space E^1 (whose elements are called edges), a continuous open map $r : E^1 \rightarrow E^0$, a continuous map $s : E^1 \rightarrow E^0$, and a family of Radon measures $\lambda = \{\lambda_v\}_{v \in E^0}$ on E^1 satisfying the following two conditions:

- (1) $\text{supp } \lambda_v = r^{-1}(v)$ for all $v \in E^0$, and
- (2) $v \mapsto \int_{E^1} \xi(\alpha) d\lambda_v(\alpha)$ is an element of $C_c(E^0)$ for all $\xi \in C_c(E^1)$.

Definition 5.2 (The quiver C^* -correspondence). If $\mathcal{Q} = (E^0, E^1, r, s, \lambda)$ is a quiver, we let $A := C_0(E^0)$ and define an A -valued inner product on $C_c(E^1)$ by

$$\langle \xi, \eta \rangle(v) := \int_{r^{-1}(v)} \overline{\xi(\alpha)} \eta(\alpha) d\lambda_v(\alpha) \quad \text{for } v \in E^0 \text{ and } \xi, \eta \in C_c(E^1).$$

We let $X(\mathcal{Q})$ denote the closure of $C_c(E^1)$ with respect to the norm arising from this inner product. We define a right action of A on $X(\mathcal{Q})$ by

$$(\xi \cdot f)(\alpha) := \xi(\alpha) f(r(\alpha)) \quad \text{for } \alpha \in E^1, \xi \in C_c(E^1), \text{ and } f \in C_0(E^0)$$

and extending this action to all of $X(\mathcal{Q})$. We define a left action $\phi : A \rightarrow \mathcal{L}(X(\mathcal{Q}))$ by

$$(\phi(f)\xi)(\alpha) := f(s(\alpha))\xi(\alpha) \quad \text{for } \alpha \in E^1, \xi \in C_c(E^1), \text{ and } f \in C_0(E^0)$$

and extending to all of $X(\mathcal{Q})$. With this inner product and these actions $X(\mathcal{Q})$ is a C^* -correspondence that we call the *quiver C^* -correspondence* of \mathcal{Q} .

Remark 5.3. Although $X(\mathcal{Q}) := \overline{C_c(E^1)}$, as described in [12, Remark 2.2] the elements of $X(\mathcal{Q})$ may be viewed as the continuous sections of a continuous field of Hilbert spaces. In particular, if $\xi \in X(\mathcal{Q})$, then for any $v \in E^0$ the vector ξ determines a function in $L^2(r^{-1}(v), \lambda_v)$. Thus for any vectors in $X(\mathcal{Q})$, the evaluation of the inner product via the integral $\int_{r^{-1}(v)} \overline{\xi(\alpha)} \eta(\alpha) d\lambda_v(\alpha)$ can be performed for all vectors $\xi, \eta \in X(\mathcal{Q})$ and not only those in $C_c(E^1)$.

Definition 5.4 (The quiver C^* -algebra). If $\mathcal{Q} = (E^0, E^1, r, s, \lambda)$ is a quiver, let $X(\mathcal{Q})$ denote the quiver C^* -correspondence of \mathcal{Q} . The *quiver C^* -algebra* $C^*(\mathcal{Q})$ is defined to be the Cuntz-Pimsner algebra $\mathcal{O}_{X(\mathcal{Q})}$.

Both topological graph C^* -algebras and graph C^* -algebras are special cases of topological quiver C^* -algebras. Topological graphs and their C^* -algebras are the special case when r is a local homeomorphism and each λ_v is counting measure on $r^{-1}(v)$. Graphs and their C^* -algebras are the special case when E^0 and E^1 both have the discrete topology and each λ_v is counting measure on $r^{-1}(v)$.

If $\mathcal{Q} = (E^0, E^1, r, s, \lambda)$ is a topological quiver, we define $E_{\text{sinks}}^0 := E^0 \setminus \overline{s(E^1)}$. An element of E_{sinks}^0 is called a *sink*, and we say \mathcal{Q} has no sinks if E_{sinks}^0 is empty (or, equivalently, if the image of $s : E^1 \rightarrow E^0$ is dense in E^0). It is shown in [12, Proposition 3.15(1)] that $\ker \phi_{X(\mathcal{Q})} = C_0(E_{\text{sinks}}^0)$, and thus the left action $\phi_{X(\mathcal{Q})}$ is injective if and only if \mathcal{Q} has no sinks.

If $\mathcal{Q} = (E^0, E^1, r, s, \lambda)$ is a topological quiver, a *path* in \mathcal{Q} is a finite sequence of edges $\alpha := e_1 \dots e_n$ with $r(e_i) = s(e_{i+1})$ for all $1 \leq i \leq n-1$. We say that the path α has length $|\alpha| := n$, and we let E^n denote the set of paths in \mathcal{Q} of length n . We may extend the maps r and s to $r^n : E^n \rightarrow E^0$ and $s^n : E^n \rightarrow E^0$ by setting $r^n(\alpha) := r(e_n)$ and $s^n(\alpha) := s(e_1)$. When no confusion arises, we shall omit the superscript and simply write r (respectively, s) for r^n (respectively, s^n).

We endow E^n with the topology it inherits as a subspace of the cartesian product $E^1 \times \dots \times E^1$. Since s is continuous and r is continuous and open, we see that s^n is continuous and r^n is continuous and open. We also define a Radon measure λ_v^n on E^n with support in $(r^n)^{-1}(v)$ inductively as follows: For $\xi : E^n \rightarrow \mathbb{C}$ we let

$$\begin{aligned} & \int_{(r^n)^{-1}(v)} \xi(e_1 \dots e_n) d\lambda_v^n(e_1 \dots e_n) \\ & := \int_{r^{-1}(s(e_2))} \dots \int_{r^{-1}(s(e_n))} \int_{r^{-1}(v)} \xi(e_1 \dots e_n) d\lambda_{s(e_2)}(e_1) \dots d\lambda_{s(e_n)}(e_{n-1}) d\lambda_v(e_n). \end{aligned}$$

Since the support of λ_v is $r^{-1}(v)$ for all $v \in E^0$, it follows that the support of λ_v^n is $(r^n)^{-1}(v)$.

If $\mathcal{Q} = (E^0, E^1, r, s, \lambda)$ is a topological quiver, then for any $n \in \mathbb{N}$ we have that $\mathcal{Q}^n := (E^0, E^n, r, s, \lambda^n)$ is a topological quiver as well. Furthermore, if $X(\mathcal{Q})$ is the quiver C^* -correspondence for \mathcal{Q} over $A := C_0(E^0)$, then $X(\mathcal{Q})^{\otimes n}$ is isomorphic to $X(\mathcal{Q}^n)$ (cf. [12, Section 6]). Moreover, we see that for $\xi, \eta \in X(\mathcal{Q})^{\otimes n}$ we have that $\langle \xi, \eta \rangle \in C_0(E^0)$ is the function with

$$\langle \xi, \eta \rangle(v) := \int_{(r^n)^{-1}(v)} \overline{\xi(\alpha)} \eta(\alpha) d\lambda_v^n(\alpha) \quad \text{for } v \in E^0.$$

A path $\alpha = e_1 \dots e_n$ is called a *cycle* if $s(e_1) = r(e_n)$, and we call the vertex $s(e_1) = r(e_n)$ the *base point* of the cycle. An *exit* for a cycle $\alpha = e_1 \dots e_n$ is an edge $f \in E^1$ with the property that $s(f) = s(e_i)$ and $f \neq e_i$ for some $1 \leq i \leq n$. (Note: In earlier literature, cycles we often called “loops”. In modern terminology, the typical convention is to use the term “loop” only for cycles of length one.) A cycle $\alpha = e_1 \dots e_n$ is called *simple* when $s(e_1) \neq r(e_i)$ for $1 \leq i \leq n-1$.

Condition (L): If \mathcal{Q} is a topological quiver, we say that \mathcal{Q} satisfies Condition (L) if the set of base points of cycles with no exits has empty interior in E^0 .

Note that a graph satisfies Condition (L) if and only if every cycle in the graph has an exit.

Definition 5.5. A path $\alpha := e_1 \dots e_n \in E^n$ is *returning* if $e_k = e_n$ for some $k \in \{1, 2, \dots, n-1\}$. Otherwise α is said to be *nonreturning*.

Definition 5.6. A nonempty subset $U \subseteq E^n$ is *nonreturning* if whenever $\alpha := e_1 \dots e_n \in U$ and $\beta := f_1 \dots f_n \in U$ we have that $\alpha_n \neq \beta_k$ for all $k \in \{1, \dots, n-1\}$.

Proposition 5.7. *Let $\mathcal{Q} = (E^0, E^1, r, s, \lambda)$ be a topological quiver with no sinks, and let $X(\mathcal{Q})$ be the associated quiver C^* -correspondence over $A := C_0(E^0)$. If \mathcal{Q} satisfies Condition (L), then $X(\mathcal{Q})$ satisfies Condition (S).*

Proof. Let $a \in A := C_0(E^0)$ with $a \geq 0$, let $n \in \mathbb{N}$, and let $\epsilon > 0$. Set

$$V := \{v \in E^0 : a(v) > \|a\| - \epsilon\} \subseteq E^0.$$

Since V is a nonempty open subset of E^0 , [12, Lemma 6.11] implies that there exists $m > n$ and a nonreturning path $\alpha \in (s^m)^{-1}(V)$. By [12, Lemma 6.12] there exists an open set $U \subseteq (s^m)^{-1}(V)$ such that $\alpha \in U$ and U is nonreturning. Using the fact that U is an open subset of the locally compact space E^m , choose a continuous compactly support function $\zeta : E^m \rightarrow [0, 1]$ with $\|\zeta\| = 1$ and with the support of ζ contained in U . It follows from [12, Lemma 6.11] that ζ is a nonreturning vector in $X(\mathcal{Q})^{\otimes m}$.

Since $\langle \zeta, \zeta \rangle$ is a positive continuous function in $C_0(E^0)$ with $\|\langle \zeta, \zeta \rangle\| = \|\zeta\|^2 = 1$, by the extreme value theorem there exists $v_0 \in E^0$ with $\langle \zeta, \zeta \rangle(v_0) = 1$. In addition, we have

$$\langle \zeta, a\zeta \rangle(v_0) = \int_{(r^m)^{-1}(v_0)} \overline{\zeta(\alpha)} a(s(\alpha)) \zeta(\alpha) d\lambda_{v_0}^n(\alpha).$$

Since ζ is supported in U , whenever $\zeta(\alpha) \neq 0$ we have $\alpha \in U$ and $s(\alpha) \in V$ so that $a(s(\alpha)) > \|a\| - \epsilon$. Thus

$$\begin{aligned} \langle \zeta, a\zeta \rangle(v_0) &= \int_{(r^m)^{-1}(v_0)} \overline{\zeta(\alpha)} a(s(\alpha)) \zeta(\alpha) d\lambda_{v_0}^n(\alpha) \\ &= \int_{(r^m)^{-1}(v_0)} a(s(\alpha)) |\zeta(\alpha)|^2 d\lambda_{v_0}^n(\alpha) \\ &\geq \int_{(r^m)^{-1}(v_0)} (\|a\| - \epsilon) |\zeta(\alpha)|^2 d\lambda_{v_0}^n(\alpha) \\ &= (\|a\| - \epsilon) \int_{(r^m)^{-1}(v_0)} |\zeta(\alpha)|^2 d\lambda_{v_0}^n(\alpha) \end{aligned}$$

$$\begin{aligned}
&= (\|a\| - \epsilon) \int_{(r^m)^{-1}(v_0)} \overline{\zeta(\alpha)} \zeta(\alpha) d\lambda_{v_0}^m(\alpha) \\
&= (\|a\| - \epsilon) \langle \zeta, \zeta \rangle(v_0) \\
&= (\|a\| - \epsilon) \cdot 1 \\
&= (\|a\| - \epsilon).
\end{aligned}$$

Hence $\|\langle \zeta, a\zeta \rangle\| \geq \|a\| - \epsilon$, and $X(\mathcal{Q})$ satisfies Condition (S). \square

Proposition 5.8. *Let $\mathcal{Q} = (E^0, E^1, r, s, \lambda)$ be a topological quiver, and let $X(\mathcal{Q})$ be the associated quiver C^* -correspondence over $A := C_0(E^0)$. If $X(\mathcal{Q})$ satisfies Condition (S), then \mathcal{Q} satisfies Condition (L).*

Proof. We shall establish the contrapositive: If \mathcal{Q} does not satisfy Condition (L), then $X(\mathcal{Q})$ does not satisfy Condition (S).

Suppose that \mathcal{Q} does not satisfy Condition (L). Then there exists a nonempty open subset $U \subseteq E^0$ such that every vertex in U is the base point of a cycle with no exits. Since E^0 is locally compact, there exists a nonempty precompact subset $V \subseteq U$ with $\overline{V} \subseteq U$. Because every point in U is the base point of a cycle with no exits, we see that $\overline{V} \subseteq U \subseteq \bigcup_{n=1}^{\infty} r((s^n)^{-1}(U))$. In addition, since s^n is continuous and r is open, for each $n \in \mathbb{N}$ the set $r((s^n)^{-1}(U))$ is open. Therefore $\{r((s^n)^{-1}(U)) : n \in \mathbb{N}\}$ is an open cover of \overline{V} , and by the compactness of \overline{V} there exists $N \in \mathbb{N}$ such that $\overline{V} \subseteq \bigcup_{n=1}^N r((s^n)^{-1}(U))$. Thus $V \subseteq \bigcup_{n=1}^N r((s^n)^{-1}(U))$, and hence every vertex in V is the base point of a cycle with no exits having length at most N .

To prove that $X(\mathcal{Q})$ does not satisfy Condition (S), we shall show that there exists $a \in A := C_0(E^0)$ with $a \geq 0$ and there exists $\epsilon > 0$ such that for all $m > N$ when $\zeta \in X(\mathcal{Q})^{\otimes m}$ with $\|\zeta\| = 1$ and $\|\langle \zeta, a\zeta \rangle\| > \|a\| - \epsilon$, then ζ is not a nonreturning vector.

Using the fact that E^0 is locally compact, we choose a nonzero compactly supported $a \in C_0(E^0)$ with $a \geq 0$ and the support of a contained in V . Also let $\epsilon := \|a\|/2$. Suppose $m > N$ and $\zeta \in X(\mathcal{Q})^{\otimes m}$ with $\|\zeta\| = 1$ and $\|\langle \zeta, a\zeta \rangle\| > \|a\| - \epsilon = \|a\|/2$. We shall show that ζ is not a nonreturning vector.

As described in Remark 5.3, ζ may be viewed as a complex-valued function. Choose $v_0 \in E^0$ with $\langle \zeta, a\zeta \rangle(v_0) > \|a\|/2$. Since

$$(5.1) \quad \langle \zeta, a\zeta \rangle(v_0) = \int_{(r^m)^{-1}(v_0)} \overline{\zeta(\alpha)} a(s(\alpha)) \zeta(\alpha) d\lambda_{v_0}^m(\alpha),$$

and a has support contained in V , there exists a path $\alpha_0 \in E^m$ with $s(\alpha_0) \in V$ and $r(\alpha_0) = v_0$. Because every vertex in V is the base point of a cycle with no exits of length at most N , and since $N < m$, we may write

$$\alpha_0 = \underbrace{\beta\beta \dots \beta}_k \gamma$$

k times

where β is the unique simple cycle based at $s(\alpha_0) \in W$ and γ is a subpath of β . However, we see that α_0 is then the unique path of length m with source in V and range equal to v_0 . Hence

$$\int_{(r^m)^{-1}(v_0)} \overline{\zeta(\alpha)} a(s(\alpha)) \zeta(\alpha) d\lambda_{v_0}^m(\alpha) = a(s(\alpha_0)) |\zeta(\alpha_0)|^2 \lambda_{v_0}^m(\{\alpha_0\})$$

and (5.1) implies

$$(5.2) \quad a(s(\alpha_0)) |\zeta(\alpha_0)|^2 \lambda_{v_0}^m(\{\alpha_0\}) = \langle \zeta, a\zeta \rangle(v_0) > \|a\|/2 > 0.$$

It follows that $\lambda_{v_0}^m(\{\alpha_0\}) > 0$. Let $n := |\beta|$, and note that $n \leq N < m$. Since $s(\beta) = s(\alpha_0) \in V$, we see that $\beta \in (s^n)^{-1}(V)$. Because E^n is locally compact, there exists a continuous compactly supported function $\xi_\beta : E^n \rightarrow [0, \infty)$ with $\xi_\beta(\beta) = a(s(\alpha_0))$ and the support of ξ_β contained in the open set $(s^n)^{-1}(V)$.

Let $(\psi_X, \pi_A) : (X(\mathcal{Q}), A) \rightarrow \mathcal{O}_{X(\mathcal{Q})} = C^*(\mathcal{Q})$ denote a universal covariant representation of ${}_A X(\mathcal{Q})_A$ into $C^*(\mathcal{Q})$. It follows from [12, Lemma 6.4] that

$$(5.3) \quad \psi_X^{\otimes m}(\zeta)^* \psi_X^{\otimes n}(\xi_\beta) \psi_X^{\otimes m}(\zeta) = \psi_X^{\otimes n}(\eta)$$

where $\eta : E^n \rightarrow \mathbb{C}$ is defined by

$$\eta(\alpha) := \int_{(r^m)^{-1}(s(\alpha))} \overline{\zeta(e_1 \dots e_m)} \xi_\beta(e_1 \dots e_n) \zeta(e_{n+1} \dots e_m \alpha) d\lambda_{s(\alpha)}^m(e_1 \dots e_m).$$

Recalling that γ is a subpath of the cycle β , we may write $\beta = \gamma\mu$, where μ is a subpath of β with $s(\mu) = r(\gamma)$ and $r(\mu) = s(\beta) = s(\gamma)$. Note that $\mu\gamma$ is a cycle of length n , which is based at $s(\mu) = r(\gamma)$.

As noted before, $\alpha_0 = \beta\beta \dots \beta\gamma$ is the unique path of length m with source in V and range equal to v_0 . Hence if we take $\mu\gamma \in E^n$, then α_0 is also the unique path of length m with source in V and range equal to $r(\mu\gamma) = r(\gamma) = v_0$. Recalling that ξ_β has support contained in $(s^n)^{-1}(V)$ (and hence ξ_β vanishes on any path of length n with source outside V), we conclude

$$\begin{aligned} \eta(\mu\gamma) &= \overline{\zeta(\alpha_0)} \xi_\beta(\beta) \zeta(\underbrace{\beta\beta \dots \beta}_{k-1 \text{ times}} \gamma)(\mu\gamma) \lambda_{v_0}^m(\{\alpha_0\}) \\ &= \overline{\zeta(\alpha_0)} a(s(\alpha_0)) \zeta(\underbrace{\beta\beta \dots \beta}_{k-1 \text{ times}} (\gamma\mu)\gamma) \lambda_{v_0}^m(\{\alpha_0\}) \\ &= \overline{\zeta(\alpha_0)} a(s(\alpha_0)) \zeta(\underbrace{(\beta\beta \dots \beta)\gamma}_{k \text{ times}}) \lambda_{v_0}^m(\{\alpha_0\}) \\ &= \overline{\zeta(\alpha_0)} a(s(\alpha_0)) \zeta(\alpha_0) \lambda_{v_0}^m(\{\alpha_0\}) \\ &= a(s(\alpha_0)) |\zeta(\alpha_0)|^2 \lambda_{v_0}^m(\{\alpha_0\}) \\ &> 0 \end{aligned}$$

where the last line follows from (5.2).

Thus $\eta \neq 0$. Since (ψ_X, π_A) is a universal covariant representation of ${}_A X(\mathcal{Q})_A$, both $\psi_X^{\otimes n}$ and π_A are injective, and hence $\psi_X^{\otimes n}(\eta) \neq 0$. It follows from (5.3) that $\psi_X^{\otimes m}(\zeta)^* \psi_X^{\otimes n}(\xi_\beta) \psi_X^{\otimes m}(\zeta) \neq 0$. Thus ζ is not a nonreturning vector. Hence $X(\mathcal{Q})$ does not satisfy Condition (S). \square

Theorem 5.9. *Let $\mathcal{Q} = (E^0, E^1, r, s, \lambda)$ be a topological quiver with no sinks, and let $X(\mathcal{Q})$ be the associated quiver C^* -correspondence over $A := C_0(E^0)$. Then \mathcal{Q} satisfies Condition (L) if and only if $X(\mathcal{Q})$ satisfies Condition (S).*

Proof. The result follows from Proposition 5.7 and Proposition 5.8. \square

6. AN EXAMINATION OF SCHWEIZER'S SIMPLICITY RESULT

Definition 6.1. Let X be a C^* -correspondence over A . We say X is *full* if

$$\overline{\text{span}}\{\langle x, y \rangle : x, y \in X\} = A.$$

Definition 6.2. Let X be a C^* -correspondence over A . We say ${}_A X_A$ is *aperiodic* if for all $n \in \mathbb{N} \cup \{0\}$, ${}_A X_A^{\otimes n} \cong {}_A A_A$ (as C^* -correspondences) implies $n = 0$. We say that a C^* -correspondence is *periodic* if it is not aperiodic.

Remark 6.3. In his paper [14], Schweizer used the term “nonperiodic” rather than “aperiodic”. We have chosen to use “aperiodic” since it appears in most English dictionaries, while the word “nonperiodic” does not.

Theorem 6.4. (Schweizer’s Simplicity Theorem [14, Theorem 3.9].) *Let X be a C^* -correspondence over A , and suppose that X is full, A is unital, and the left action $\phi_X : A \rightarrow \mathcal{L}(X)$ is injective. Then \mathcal{O}_X is simple if and only if X is aperiodic and A has no hereditary ideals other than $\{0\}$ and A .*

Remark 6.5. Although the hypothesis of the injectivity of the left action is not mentioned in the statement of [14, Theorem 3.9], it is used to apply [14, Theorem 3.8] in its proof. Our statement above of Schweizer’s theorem makes this hypothesis explicit.

Definition 6.6. Let $E = (E^0, E^1, r, s)$ be a directed graph. The graph correspondence is the nondegenerate C^* -correspondence $X(E)$ over $C_0(E^0)$ defined by

$$X(E) = \{\xi : E^1 \rightarrow \mathbb{C} : \text{the map } v \mapsto \sum_{s(e)=v} |\xi(e)|^2 \text{ is in } C_0(E^0)\},$$

with module actions and $C_0(E^0)$ -valued inner product given by


$$(a \cdot \xi)(e) = a(r(e))\xi(e), \quad (\xi \cdot b)(e) = \xi(e)b(s(e)), \quad \text{and} \quad \langle \xi, \eta \rangle(v) = \sum_{s(e)=v} \overline{\xi(e)}\eta(e)$$

for $a, b \in C_0(E^0)$, $\xi, \eta \in X(E)$, $e \in E^1$, and $v \in E^0$.

Definition 6.7. If $E = (E^0, E^1, r, s)$ is a graph, then for any $n \in \mathbb{N}$ we let E^n denote the set of paths in E of length n . We also let $E^{\times n} := (E^0, E^n, r, s)$ denote the graph whose vertices are the vertices of E and whose edges are the paths of length n of E .

Remark 6.8. It follows from [1, Proposition 3.3] that if E is a graph, then $X(E)^{\otimes n} \cong X(E^{\times n})$ as C^* -correspondences for all $n \in \mathbb{N}$.

Example 6.9. Let E be the graph




consisting of a simple cycle with n vertices for some $n \in \mathbb{N}$. Note that $A := C(E^0) = \mathbb{C}^n$. Since each vertex of E is the base point of a cycle of length n , we see that $E^{\times n}$ is the graph




so that $X(E^{\times n}) = \mathbb{C}^n$ with the left and right multiplication occurring component-wise. Thus $X(E)^{\otimes n} \cong X(E^{\times n}) \cong \mathbb{C}^n \cong A$ as C^* -correspondences over A , and $X(E)$ is periodic.

Example 6.10. Let E be the graph



consisting of the disjoint union of three simple cycles of lengths 2, 3, and 4, respectively. If we take the least common multiple of these lengths, $\text{lcm}(2, 3, 4) = 12$, then we see that each vertex is the base point of a cycle of length 12, and hence $E^{\times 12}$ is the graph



In particular, since there are 9 vertices, we have $X(E)^{\otimes 12} = X(E^{\times 12}) \cong \mathbb{C}^9 \cong A$ as C^* -correspondences over A , and thus $X(E)$ is periodic.

Proposition 6.11. *Let E be a graph with a finite number of vertices, no sinks, and no sources. (Note that E is not required to have a finite number of edges.) Then the following are equivalent:*

- (1) $X(E)$ is aperiodic.
- (2) E is not a disjoint union of simple cycles.

Proof. We shall establish the result by proving the contrapositives: $X(E)$ is periodic if and only if E is a disjoint union of cycles.

If E is disjoint union of cycles, then since E has a finite number of vertices there are a finite number of cycles in this disjoint union. Let n be the least common multiple of the lengths of these cycles. Then $E^{\times n}$ is a graph with a single cycle of length one at each vertex. Hence we have the C^* -correspondence isomorphisms $X(E)^{\otimes n} \cong X(E^{\times n}) \cong A$ (cf. Example 6.10). Thus $X(E)$ is periodic.

Conversely, suppose that $X(E)$ is periodic. Then there exists $n \in \mathbb{N}$ such that $X(E)^{\otimes n} \cong A$ as C^* -correspondences. For convenience of notation, label the vertices as $E^0 = \{1, \dots, k\}$. Then $C(E^0) = \mathbb{C}^k$, and we have the C^* -correspondence isomorphisms $X(E^{\times n}) \cong X(E)^{\otimes n} \cong \mathbb{C}^k \cong A$. Recalling that $X(E^{\times n}) := C(E^n)$, we conclude there exists a \mathbb{C}^k -linear isomorphism $h : C(E^n) \rightarrow C(E^0) = \mathbb{C}^k$. For each $\alpha \in E^n$, let $\delta_\alpha : E^n \rightarrow \mathbb{C}$ be the function with $\delta_\alpha(\alpha) = 1$ and $\delta_\alpha(\beta) = 0$ when $\beta \neq \alpha$. Also, for each $1 \leq i \leq k$, let e_i denote the standard basis element of \mathbb{C}^k having 1 in entry i and 0 in all other entries. Suppose $h(\delta_\alpha) = (z_1, \dots, z_k)$. Then

$$\begin{aligned} e_{r(\alpha)} &= \langle \delta_\alpha, \delta_\alpha \rangle = \langle h(\delta_\alpha), h(\delta_\alpha) \rangle = (z_1, \dots, z_k)^*(z_1, \dots, z_k) \\ &= (\overline{z_1}, \dots, \overline{z_k})(z_1, \dots, z_k) = (|z_1|^2, \dots, |z_k|^2). \end{aligned}$$

Consequently $z_i = 0$ for $i \neq r(\alpha)$, and $z_{r(\alpha)} \neq 0$. Thus $h(\delta_\alpha) = z_{r(\alpha)}e_{r(\alpha)}$. Hence

$$z_{r(\alpha)}e_{r(\alpha)} = h(\delta_\alpha) = h(e_{s(\alpha)} \cdot \delta_\alpha) = e_{s(\alpha)}h(\delta_\alpha) = e_{s(\alpha)} \cdot (z_1, \dots, z_k) = z_{s(\alpha)}e_{s(\alpha)}.$$

Since $z_{r(\alpha)} \neq 0$, the above equation implies $s(\alpha) = r(\alpha)$. Because α is an arbitrary element of E^n , we conclude every element of E^n is a cycle. Moreover, since $C(E^n) \cong C(E^0) \cong \mathbb{C}^k$, we deduce E^n has k elements. Thus $E^n = \{\alpha_1, \dots, \alpha_k\}$, where α_i is a cycle of length n for each $1 \leq i \leq k$. Therefore E is a disjoint union of simple cycles. \square

Definition 6.12. Let $E = (E^0, E^1, r, s)$ be a graph. We say E is *weakly connected* if for all $v, w \in E^0$ there exists a sequence of vertices $v_1, \dots, v_n \in E^0$ with $v_1 = v$, $v_n = w$, and for all $1 \leq i \leq n$ there is either an edge in E from v_i to v_{i+1} or there is an edge in E from v_{i+1} to v_i . We say E is *strongly connected* if for all $v, w \in E^0$ there is a path in E from v to w .

Corollary 6.13. *Let E be a weakly connected graph with a finite number of vertices, no sinks, and no sources. (Note that E is not required to have a finite number of edges.) Then the following are equivalent:*

- (1) $X(E)$ is aperiodic.
- (2) E is not a simple cycle.

Corollary 6.14. *Let E be a strongly connected graph with a finite number of vertices. (Note that E is not required to have a finite number of edges.) Then the following are equivalent:*


- (1) $X(E)$ is aperiodic.
- (2) E is not a simple cycle.
- (3) E satisfies Condition (L).

Proof. The equivalence (1) \iff (2) follows from Corollary 6.13.

To see (2) \implies (3), suppose E is not a simple cycle. If α is any cycle in E , then since E is not a simple cycle, there exists a vertex $w \in E^0$ that is not on α . Since E is strongly connected, there exists a path from $s(\alpha)$ to w , and some edge of this path must be an exit for α . Hence E satisfies Condition (L), and (3) holds.


To see (3) \implies (2), simply note that if every cycle in E has an exit, then E cannot be a simple cycle. \square

Example 6.15. If E is the graph




then E is weakly connected and does not satisfy Condition (L). In addition, for any $n \in \mathbb{N}$, we see that there are paths of length n in E that are not cycles. Hence, the correspondence $X(E^{\times n})$ is not isomorphic to the correspondence $C_0(E^0)$, so that the correspondence $X(E)^{\otimes n}$ is not isomorphic to ${}_A A_A$. Consequently, $X(E)$ is aperiodic.

Example 6.16. If E is the graph



then E is weakly connected and does not satisfy Condition (L). In addition, for any $n \in \mathbb{N}$, we see that there are paths of length n in E that are not cycles. Hence, the correspondence $X(E^{\times n})$ is not isomorphic to the correspondence $C_0(E^0)$, so that the correspondence $X(E)^{\otimes n}$ is not isomorphic to ${}_A A_A$. Consequently, $X(E)$ is aperiodic. Furthermore, E has no nontrivial saturated hereditary subsets, and hence $A = \mathbb{C}^2$ has no invariant ideals other than $\{0\}$ and A . Since E does not satisfy Condition (L), $\mathcal{O}_{X(E)} \cong C^*(E)$ is not simple. (In fact, $C^*(E) \cong M_2(C(\mathbb{T}))$, so $C^*(E)$ is Morita equivalent to $C(\mathbb{T})$, and $C^*(E)$ has uncountably many ideals.)

Example 6.17. If E is the graph



then E is weakly connected, has no sinks and no sources, and does not satisfy Condition (L). In addition, for any $n \in \mathbb{N}$, we see that there are paths of length n in E that are not cycles. Hence, the correspondence $X(E^{\times n})$ is not isomorphic to the correspondence $C_0(E^0)$, so that the correspondence $X(E)^{\otimes n}$ is not isomorphic to ${}_A A_A$. Consequently, $X(E)$ is aperiodic. Furthermore, E has no nontrivial saturated hereditary subsets, and hence $A = C_0(E^0)$ has no invariant ideals other than $\{0\}$ and A . Since E does not satisfy Condition (L), $\mathcal{O}_{X(E)} \cong C^*(E)$ is not simple. (In fact, $C^*(E) \cong C(\mathbb{T}) \otimes \mathcal{K}$, so $C^*(E)$ is Morita equivalent to $C(\mathbb{T})$, and $C^*(E)$ has uncountably many ideals.)

Example 6.18. For each $k \in \mathbb{N}$ let C_k be the graph that is a simple cycle of length k . If $E := C_1 \sqcup C_2 \sqcup \dots$, then for any $n \in \mathbb{N}$, we see that E contains a path of length n that is not a cycle. Hence, the correspondence $X(E^{\times n})$ is not isomorphic to the correspondence $C_0(E^0)$, so that the correspondence $X(E)^{\otimes n}$ is not isomorphic to ${}_A A_A$. Consequently, $X(E)$ is aperiodic. Thus $X(E)$ is aperiodic even though E is a disjoint union of simple cycles.

Example 6.19. Let E be a strongly connected graph with an infinite number of vertices. If α is any cycle in E , then there exists a vertex $w \in E^0$ that is not on α . Since E is strongly connected, there exists a path from $s(\alpha)$ to w , and some edge of this path must be an exit for α . Hence E satisfies Condition (L).

In addition, since E is strongly connected, for any $n \in \mathbb{N}$ there exists a path of length n in E that is not a cycle. Hence, the correspondence $X(E^{\times n})$ is not isomorphic to the correspondence $C_0(E^0)$, so that the correspondence $X(E)^{\otimes n}$ is not isomorphic to ${}_A A_A$. Consequently, $X(E)$ is aperiodic.

Therefore if E is any strongly connected graph with infinitely many vertices, then E satisfies Condition (L) and $X(E)$ is aperiodic.

Remark 6.20. For a graph E , let $X(E)$ denote the graph C^* -correspondence over $A := C_0(E^0)$. Our results and examples from this section can be summarized by the following points:

- Corollary 6.14 states that if E is strongly connected with finitely many vertices, aperiodicity of $X(E)$ is equivalent to E satisfying Condition (L).
- Example 6.15 shows that there exists a weakly connected finite graph with no sinks and no sources such that $X(E)$ is aperiodic, but E does not satisfy Condition (L). This shows that, even for finite graphs with no sinks and

no sources, aperiodicity of $X(E)$ is not equivalent to E satisfying Condition (L). Thus, even in the special case of graph C^* -algebras, aperiodicity is not the correct property to obtain a Cuntz-Krieger Uniqueness Theorem.

- Example 6.16 shows that there exists a weakly connected finite graph E with no sinks (but having a source) for which $X(E)$ is aperiodic and A has no invariant ideals other than $\{0\}$ and A , but E does not satisfy Condition (L). Since E does not satisfy Condition (L), $C^*(E)$ is not simple. Hence Schweizer’s simplicity theorem does not hold if the fullness of the C^* -correspondence (which is equivalent to “no sources” in the graph case) is removed. Thus aperiodicity is not the correct property to use for a simplicity result when a C^* -correspondence is not full.
- Example 6.17 shows that there exists a weakly connected graph E with no sinks and no sources (but having a countably infinite number of vertices) for which $X(E)$ is aperiodic and A has no invariant ideals other than $\{0\}$ and A , but E does not satisfy Condition (L). Since E does not satisfy Condition (L), $C^*(E)$ is not simple. Hence Schweizer’s simplicity theorem does not hold if the condition that A is unital (which is equivalent to “finite vertices” in the graph case) is removed. Thus aperiodicity is not the correct property to use for a simplicity result when a C^* -correspondence is over a nonunital C^* -algebra A .
- Finally, Example 6.19 shows that for strongly connected graphs with an infinite number of vertices, aperiodicity and Condition (L) are both automatic.

These observations show that while aperiodicity is useful for characterizing simplicity under the hypotheses of Schweizer’s simplicity theorem, it is not the correct notion for characterizing simplicity more generally. Moreover, even when Schweizer’s hypotheses hold, aperiodicity is inadequate for obtaining a Cuntz-Krieger Uniqueness Theorem. Indeed, we see this even in the special case of graph C^* -algebras.

This gives additional credibility to the notion of Condition (S) that we introduced in this paper. First of all, under the hypothesis that the left action is injective, Condition (S) coincides with Condition (L) for graphs, topological graphs, and topological quivers, suggesting it is a reasonable generalization of Condition (L) for general C^* -correspondences. Second, Condition (S) allows us to prove a simplicity result for Cuntz-Krieger algebras that removes Schweizer’s hypotheses that X is full and A is unital. Since X full and A unital are strong hypotheses, Condition (S) allows us to obtain sufficient conditions for simplicity that apply to a much wider class of Cuntz-Pimsner algebras. Third, unlike aperiodicity, we have been able to use Condition (S) to obtain a Cuntz-Krieger Uniqueness Theorem and obtain our simplicity result via this uniqueness theorem.

7. CONCLUDING REMARKS

We conclude with two questions that are important for extending the work in this paper.

Question 1: Let X be a C^* -correspondence over A with injective left action. If \mathcal{O}_X is simple, is it necessary that X satisfies Condition (S)?

When $X = X(\mathcal{Q})$ is the C^* -correspondence of a topological quiver \mathcal{Q} , we know that $\mathcal{O}_{X(E)} = C^*(\mathcal{Q})$ simple implies \mathcal{Q} satisfies Condition (L), which implies $X(\mathcal{Q})$ satisfies Condition (S). Thus we have an affirmative answer to Question (1) for C^* -correspondences of topological quivers, topological graphs, and graphs. If an affirmative answer to Question (1) is obtained in general, then Condition (S) is precisely the correct generalization of Condition (L) for C^* -correspondences with injective left action.

Question 2: How can Condition (S) be appropriately modified to obtain a Cuntz-Krieger Uniqueness Theorem for general C^* -correspondences whose left action is not necessarily injective?

As noted in Proposition 3.4, a C^* -correspondence satisfying Condition (S) necessarily has injective left action. It would be desirable to remove the hypothesis of injective left action and obtain an appropriate condition for general C^* -correspondences. While there are techniques for extending results for C^* -correspondences with injective left actions to the general situation (e.g., via the method of “adding tails” described in [11]), translating Condition (S) to the non-injective setting has been complicated and elusive, and we have been unable to apply these techniques in any fruitful way. Currently, we are uncertain how to modify Condition (S) when the left action is not injective.

If one is able to find an appropriate formulation of Condition (S) for general C^* -correspondences and prove a Cuntz-Krieger Uniqueness Theorem under this hypothesis, then the result will also provide sufficient conditions for simplicity. In particular, once a general Cuntz-Krieger Uniqueness Theorem is obtained, the proof of Theorem 4.3 will go through verbatim and the modified Condition (S) together with the hypothesis that A has no invariant ideals other than $\{0\}$ and A will imply simplicity of the Cuntz-Pimsner algebra.

REFERENCES

- [1] T.M. Carlsen, A. Dor-On and S. Eilers, *Shift equivalences through the lens of Cuntz-Krieger algebras*, arXiv.2011.10329v5.
- [2] N. Fowler, P.S. Muhly, and I. Raeburn, *Representations of Cuntz-Pimsner algebras*, Indiana Univ. Math. J. **52** (2003), 569–605.
- [3] N. Fowler and I. Raeburn, *The Toeplitz algebra of a Hilbert bimodule*, Indiana Univ. Math. J. **48** (1999), 155–181.
- [4] T. Kajiwara, C. Pinzari, Y. Watatani, *Ideal structure and simplicity of the C^* -algebras generated by Hilbert bimodules*, J. Funct. Anal. **159** (1998), 295–322.
- [5] T. Katsura, *A class of C^* -algebras generalizing both graph algebras and homeomorphism C^* -algebras. I. Fundamental results*, Trans. Amer. Math. Soc. **356** (2004), 4287–4322.
- [6] T. Katsura, *On C^* -algebras associated with C^* -correspondences*, J. Funct. Anal. **217** (2004), 366–401.
- [7] T. Katsura, *Ideal structure of C^* -algebras associated with C^* -correspondences*, Pacific J. Math. **230** (2007), 107–145.
- [8] E. C. Lance, *Hilbert C^* -modules. A toolkit for operator algebraists*. London Mathematical Society Lecture Note Series, **210**. Cambridge University Press, Cambridge, 1995.
- [9] P. S. Muhly and B. Solel, *On the simplicity of some Cuntz-Pimsner algebras*, Math. Scand. **83** (1998), 53–73.
- [10] P. S. Muhly and B. Solel, *Corrigendum to: “On the simplicity of some Cuntz-Pimsner algebras”*, Math. Scand. **91** (2002), 244–246.
- [11] P. S. Muhly and M. Tomforde, *Adding tails to C^* -correspondences*, Documenta Math. **9** (2004), 79–106.

- [12] P.S. Muhly and M. Tomforde, *Topological quivers*, Internat. J. Math. **16** (2005), no. 7, 693–755.
- [13] M. V. Pimsner, *A class of C^* -algebras generalizing both Cuntz-Krieger algebras and crossed products by \mathbb{Z}* , Fields Institute Comm. **12** (1997), 189–212.
- [14] J. Schweizer, *Dilations of C^* -correspondences and the simplicity of Cuntz-Pimsner algebras*, J. Funct. Anal. **180** (2001), 404–425.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF COLORADO BOULDER, BOULDER, CO 80309-0395, USA

Email address: `menevse.eryuzlu@colorado.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF COLORADO, COLORADO SPRINGS, CO 80918-3733, USA

Email address: `mtomford@uccs.edu`