

# Twice epi-differentiability of a class of non-amenable composite functions

Yulan Liu\* and Shaohua Pan<sup>†</sup>

January 2, 2023

## Abstract

This paper focuses on the twice epi-differentiability of a class of non-amenable functions, which are the composition of a piecewise twice differentiable (PWTD) function and a parabolically semidifferentiable mapping. Such composite functions frequently appear in constrained and composite optimization problems, disjunctive programs, and low-rank or/and sparsity optimization problems. To achieve their twice epi-differentiability, we first justify the proper twice epi-differentiability, parabolic epi-differentiability and regularity of PWTD functions, and then derive an upper and lower estimate for the second subderivatives of this class of composite functions in terms of a chain rule of their parabolic subderivatives. We employ the obtained upper and lower estimates to characterize the parabolic regularity, the second subderivatives, and so the proper twice epi-differentiability for several classes of popular functions, which include the compositions of PWTD outer functions and twice differentiable inner mappings, the regularized functions inducing group sparsity, and the indicator functions of the  $q$  ( $q > 1$ )-order cone.

## 1 Introduction

Let  $\mathbb{X}$  represent a finite dimensional real vector space endowed with the inner product  $\langle \cdot, \cdot \rangle$  and its induced norm  $\| \cdot \|$ . We are interested in the following composite function

$$f(x) := \vartheta(F(x)) \quad \text{for } x \in \mathbb{X}, \quad (1)$$

where  $F: \mathbb{X} \rightarrow \mathbb{R}^m$  and  $\vartheta: \mathbb{R}^m \rightarrow \overline{\mathbb{R}} := (-\infty, \infty]$  satisfy the basic assumption:

**Assumption 1 (i)**  $\vartheta$  is PWTD and is strictly continuous relative to its domain  $\text{dom}\vartheta$ ;

**(ii)**  $F$  is a parabolically semidifferentiable mapping on an open set  $\mathcal{O} \supset F^{-1}(\text{dom}\vartheta)$ .

---

\*ylliu@gdut.edu.cn, School of Mathematics and Statistics, Gangdong University of Technology, Guangzhou

<sup>†</sup>Corresponding author (shhpan@scut.edu.cn), School of Mathematics, South China University of Technology, Guangzhou.

By Definition 2.1 later, we assume that  $\emptyset \neq \text{dom } \vartheta = \bigcup_{i=1}^s C_i$  for polyhedral sets  $C_1, \dots, C_s$ , and relative to each  $C_i$ ,  $\vartheta = \vartheta_i$ , a twice differentiable function.

The optimization model involving the above  $f$  and a lower semicontinuous (lsc)  $f_0: \mathbb{X} \rightarrow \overline{\mathbb{R}}$  that is twice differentiable on an open set containing  $F^{-1}(\text{dom } \vartheta)$ , that is

$$\min_{x \in \mathbb{X}} \Phi(x) := f_0(x) + f(x), \quad (2)$$

encompasses major classes of constrained and composite optimization problems such as classical nonlinear programming problems, second-order cone and semidefinite programming problems, eigenvalue optimization problems [1–3], and amenable composite optimization problems [4]. Since the outer  $\vartheta$  is not required to be convex, some new problems are absorbed in model (2) such as disjunctive programs [5, 6] and nonconvex optimization problems arising from low-rank or/and sparsity problems; for example, the loss function  $\psi(\mathcal{A}(UV^\top))$ , where  $\mathcal{A}: \mathbb{R}^{n_1 \times n_2} \rightarrow \mathbb{R}^m$  is a sampling operator and  $UV^\top$  is a factorized form of low-rank matrices, precisely has the form of (1) if  $\psi: \mathbb{R}^m \rightarrow \mathbb{R}$  is chosen to be a piecewise linear-quadratic (PWLQ) function such as the SCAD or MCP function (see [7, 8]). This class of loss functions is suitable for low-rank optimization problems with heavily-tailed noises. In addition, the group SCAD and MCP functions in group sparsity optimization [9, 10], which were shown to be the equivalent DC surrogates of group zero-norm in [11], also take the form of (1).

Composite functions of type (1) constitute a convenient framework in variational analysis and continuous optimization for developing theoretical and algorithmic issues of constrained and composite optimizations. Standard assumptions under which this class of functions was investigated and applied in constrained optimization require that the inner  $F$  is twice continuously differentiable, that the function  $\vartheta$  is lsc and convex, and that the epigraphical multifunction associated to  $\vartheta$  is metrically regular around the point in question (see [12]). Compared with metric regularity, metric subregularity of a multifunction or calmness equivalent for its inverse mapping has more important application scenarios where its robust counterpart is out of reach. We refer the reader to the reference [13–20] for various developments on metric subregularity and calmness, and their applications to optimality and stability conditions, error bounds, and convergence of algorithms. Inspired by this, Mohammadi et al. [21] carried out the first-order and second-order variational analysis of composition (1) for a class of fully subamenable functions, which are same as fully amenable functions [4] except that metric regularity of epigraphical multifunctions required by the latter is replaced by metric subregularity qualification condition (MSQC) for system  $F(x) \in \text{dom } \vartheta$  at the reference point.

As discussed in [22], twice epi-differentiability of extended real-valued functions, as a carrier of vital second-order information, plays a significant role in second-order variational analysis of nonconvex and nonsmooth optimization, but its justification has been well recognized to be extremely difficult. Since Rockafellar’s landmark paper [4] where this property was verified for the fully amenable functions, there are few works on this topic except [23–25]. The papers [23, 24] extended Rockafellar’s results to the composition (1) with a proper lsc convex  $\vartheta$  and a twice differentiable  $F$ , but required a restrictive assumption on the second subderivative that does not hold for constrained

optimization problems, while the work [25] obtained upper and lower estimates for the second subderivative but did not discuss the twice epi-differentiability. Until recently, Mohammadi et al. [21, 22, 26] conducted a systematic study under MSQCs for the twice epi-differentiability of this class of compositions by leveraging their parabolic epi-differentiability and regularity. Among others, they fully analyzed parabolic regularity for constraint system  $\Omega \cap \mathcal{O} = \{x \in \mathcal{O} \mid g(x) \in \Theta\}$  in [26] so as to achieve the twice epi-differentiability of the indicator function of  $\Omega$ , where  $\mathcal{O}$  is a neighborhood of the reference point,  $g: \mathbb{R}^n \rightarrow \mathbb{R}^m$  is a mapping twice differentiable at the reference point, and  $\Theta \subset \mathbb{R}^m$  is a closed convex set; and later they showed in [22] that the twice epi-differentiability of this class of compositions can be guaranteed under parabolic regularity if the outer  $\vartheta$  is locally Lipschitz continuous relative to its domain.

In this work, we continue to promote this direction by investigating the twice epi-differentiability for a class of non-amenable functions, i.e., the composition (1) with  $\vartheta$  and  $F$  satisfying Assumption 1. Although the domain of the outer  $\vartheta$  has a certain polyhedrality, the associated optimization model (2) still involves nonpolyhedral conic optimization problems because the inner  $F$  is not required to be differentiable. In section 3, we conduct a systematic second-order variational analysis of PWTD functions, and confirm their proper twice epi-differentiability, parabolic epi-differentiability and parabolic regularity. Based on these properties of PWTD functions, we derive in section 4 an upper and lower estimate for second subderivatives of this class of compositions. In section 5 these estimates are applied to achieving the parabolic regularity and then twice epi-differentiability for several classes of common  $f$ , including the compositions of PWTD outer functions and twice differentiable inner mappings, the regularized functions inducing group sparsity, and the indicator functions of the  $q$  ( $q > 1$ )-order cone. To the best of our knowledge, this work is the first to explore the twice epi-differentiability of non-amenable functions. Mohammadi [27] recently studied the first-order variational analysis of non-amenable functions, but did not touch their second-order properties.

**Notation.** For a function  $h: \mathbb{X} \rightarrow (-\infty, \infty]$ , write  $\text{dom } h := \{x \in \mathbb{X} \mid h(x) < \infty\}$ , and denote by  $h^*$  the conjugate of  $h$ , i.e.,  $h^*(z) = \sup_{x \in \mathbb{X}} \{\langle x, z \rangle - h(x)\}$ . If a mapping  $g: \mathbb{X} \rightarrow \mathbb{R}^m$  is differentiable at  $\bar{x} \in \mathbb{X}$ ,  $\nabla g(\bar{x})$  denotes the transpose of  $g'(\bar{x})$ , the Jacobian of  $g$  at  $\bar{x}$ , and if  $g$  is twice differentiable at  $\bar{x}$ ,  $D^2g(\bar{x})$  denotes its second-order differential mapping at  $\bar{x}$ . For a closed set  $S \subset \mathbb{X}$ ,  $\delta_S$  denotes the indicator function of  $S$ , i.e.,  $\delta_S(x) = 0$  if  $x \in S$ , otherwise  $\delta_S(x) = \infty$ ; and  $\text{dist}(x, S)$  denotes the distance of  $x$  from  $S$  on the norm  $\|\cdot\|$ . For a given  $x \in \mathbb{X}$ ,  $\mathbb{B}(x, \delta)$  denotes the closed ball of radius  $\delta$  centered at  $x$  on the norm  $\|\cdot\|$ . For an integer  $m \geq 1$ , write  $[m] = \{1, \dots, m\}$ . For every  $q \in (1, +\infty)$ ,  $\|\cdot\|_q$  represents the  $\ell_q$ -norm of vectors.

## 2 Preliminaries

This section includes some necessary knowledge on variational analysis (see the standard texts such as [12] and [28] for more details) and some preliminary results that will be used later. First of all, we recall the concepts of PWTD functions and subdifferentials.

**Definition 2.1** A function  $h: \mathbb{R}^m \rightarrow \overline{\mathbb{R}}$  is said to be *PWTD* if  $\text{dom } h$  is nonempty and can be represented as a union of finitely many polyhedral sets, relative to each of which  $h$  takes a twice differentiable function.

**Definition 2.2** (see [12, Definition 8.3]) Consider a function  $h: \mathbb{X} \rightarrow \overline{\mathbb{R}}$  and a point  $x \in \text{dom } h$ . The regular subdifferential of  $h$  at  $x$  is defined as

$$\widehat{\partial}h(x) := \left\{ v \in \mathbb{X} \mid \liminf_{x' \neq x \rightarrow x} \frac{h(x') - h(x) - \langle v, x' - x \rangle}{\|x' - x\|} \geq 0 \right\},$$

and the basic (also known as the limiting or Morduhovich) subdifferential of  $h$  at  $x$  is

$$\partial h(x) := \left\{ v \in \mathbb{X} \mid \exists x^k \rightarrow x \text{ with } h(x^k) \rightarrow h(x) \text{ and } v^k \in \widehat{\partial}h(x^k) \text{ s.t. } v^k \rightarrow v \right\}.$$

When  $h = \delta_S$  for a nonempty closed set  $S \subseteq \mathbb{X}$ ,  $\widehat{\partial}h(x)$  and  $\partial h(x)$  for  $x \in S$  respectively reduce to the regular normal cone  $\widehat{\mathcal{N}}_S(x)$  and normal cone  $\mathcal{N}_S(x)$ .

## 2.1 Metric subregularity of epigraphical multifunctions

Recall that a multifunction  $\mathcal{F}: \mathbb{X} \rightrightarrows \mathbb{R}^m$  is said to have the (metric) subregularity at a point  $(\bar{x}, \bar{y}) \in \text{gph } \mathcal{F}$  if there exist  $\delta > 0$  and  $\kappa > 0$  such that for all  $x \in \mathbb{B}(\bar{x}, \delta)$ ,

$$\text{dist}(x, \mathcal{F}^{-1}(\bar{y})) \leq \kappa \text{dist}(\bar{y}, \mathcal{F}(x)).$$

The following proposition states the relation between the subregularity of an epigraphical multifunction and that of the associated domain multifunction, which improves the result of [21, Proposition 3.1] because the latter only confirms part (i) of Proposition 2.1.

**Proposition 2.1** Let  $\varphi: \mathbb{R}^m \rightarrow \overline{\mathbb{R}}$  be a proper function, and let  $g: \mathbb{X} \rightarrow \mathbb{R}^m$  be a single-valued map. Define the multifunctions  $\mathcal{G}: \mathbb{X} \times \mathbb{R} \rightrightarrows \mathbb{R}^m \times \mathbb{R}$  and  $\mathcal{H}: \mathbb{X} \rightrightarrows \mathbb{R}^m$  by

$$\mathcal{G}(x, \alpha) := (g(x), \alpha) - \text{epi } \varphi \quad \text{and} \quad \mathcal{H}(x) := g(x) - \text{dom } \varphi. \quad (3)$$

Consider any  $\bar{x} \in \mathbb{X}$  with  $\bar{\omega} = \varphi(g(\bar{x}))$  finite. The following statements hold.

- (i) If  $g$  is continuous at  $\bar{x}$  and  $\varphi$  is continuous at  $g(\bar{x})$  relative to its domain, then the subregularity of  $\mathcal{G}$  at  $((\bar{x}, \bar{\omega}), (0, 0))$  implies that of  $\mathcal{H}$  at  $(\bar{x}, 0)$ .
- (ii) If  $g$  is strictly continuous at  $\bar{x}$  and  $\varphi$  is strictly continuous at  $g(\bar{x})$  relative to its domain, then the subregularity of  $\mathcal{H}$  at  $(\bar{x}, 0)$  implies that of  $\mathcal{G}$  at  $((\bar{x}, \bar{\omega}), (0, 0))$ .

**Proof:** The proof of part (i) is similar to that of [21, Proposition 3.1], so it suffices to establish part (ii). Because the mapping  $\mathcal{H}$  is subregular at  $(\bar{x}, 0)$ , there exist  $\delta > 0$  and  $c > 0$  such that for all  $x \in \mathbb{B}(\bar{x}, \delta)$ ,

$$\text{dist}(x, \mathcal{H}^{-1}(0)) \leq c \text{dist}(g(x), \text{dom } \varphi). \quad (4)$$

Since  $\varphi$  is strictly continuous at  $g(\bar{x})$  relative to its domain, there exists  $\varepsilon_1 > 0$  such that

$$|\varphi(z) - \varphi(z')| \leq L_\varphi \|z - z'\|_2 \quad \forall z, z' \in \mathbb{B}(g(\bar{x}), \varepsilon_1) \cap \text{dom } \varphi. \quad (5)$$

Since  $g$  is strictly continuous at  $\bar{x}$ , there exist  $\varepsilon_2 > 0$  and  $L_g > 0$  such that

$$\|g(x) - g(\bar{x})\|_2 \leq \varepsilon_1/3 \quad \text{and} \quad \|g(x) - g(x')\|_2 \leq L_g \|x - x'\| \quad \forall x, x' \in \mathbb{B}(\bar{x}, \varepsilon_2).$$

Take  $\varepsilon = \min(\delta, \varepsilon_1, \varepsilon_2)/3$ . Consider any  $(x, \alpha) \in \mathbb{B}((\bar{x}, \bar{\omega}), \varepsilon)$ . Let  $(y^*, \alpha^*) \in \text{epi } \varphi$  be such that  $\text{dist}((g(x), \alpha), \text{epi } \varphi) = \sqrt{\|g(x) - y^*\|_2^2 + (\alpha - \alpha^*)^2}$ . Clearly, it holds that

$$\|g(x) - y^*\|_2 \leq \sqrt{\|g(x) - g(\bar{x})\|_2^2 + (\alpha - \bar{\omega})^2} \leq 2\varepsilon_1/3,$$

so  $\|y^* - g(\bar{x})\|_2 \leq \|g(x) - y^*\|_2 + \|g(x) - g(\bar{x})\|_2 \leq \varepsilon_1$ . We proceed the proof by two cases.

**Case 1:**  $\alpha^* = \varphi(y^*)$ . Let  $\hat{x} \in \mathcal{H}^{-1}(0)$  be such that  $\text{dist}(x, \mathcal{H}^{-1}(0)) = \|x - \hat{x}\|$ . Clearly,  $g(\hat{x}) \in \text{dom } \varphi$  and  $\|\hat{x} - \bar{x}\| \leq 2\|x - \bar{x}\| \leq \varepsilon_2$ . Then,  $\|g(x) - g(\hat{x})\|_2 \leq L_g \|x - \hat{x}\|$  and  $\|g(\hat{x}) - g(\bar{x})\|_2 \leq \varepsilon_1/3$ . The latter, along with  $\|y^* - g(\bar{x})\|_2 \leq \varepsilon_1$  and (5), implies that

$$|\varphi(y^*) - \varphi(g(\hat{x}))| \leq L_\varphi \|y^* - g(\hat{x})\|_2. \quad (6)$$

From (4) we have  $\|x - \hat{x}\| \leq c \text{dist}(g(x), \text{dom } \varphi)$ . Since  $y^* \in \text{dom } \varphi$ , it holds that

$$\|x - \hat{x}\| \leq c \|g(x) - y^*\|_2. \quad (7)$$

By combining this with  $(\hat{x}, \varphi(g(\hat{x}))) \in \mathcal{G}^{-1}(0, 0)$ , it is immediate to obtain that

$$\begin{aligned} \text{dist}((x, \alpha), \mathcal{G}^{-1}(0, 0)) &\leq \sqrt{\|x - \hat{x}\|^2 + |\alpha - \varphi(g(\hat{x}))|^2} \\ &\leq \sqrt{\|x - \hat{x}\|^2 + 2|\alpha - \alpha^*|^2 + 2|\varphi(y^*) - \varphi(g(\hat{x}))|^2} \\ &\leq \sqrt{\|x - \hat{x}\|^2 + 2|\alpha - \alpha^*|^2 + 2L_\varphi^2 \|y^* - g(\hat{x})\|_2^2} \\ &\leq \sqrt{\|x - \hat{x}\|^2 + 2|\alpha - \alpha^*|^2 + 4L_\varphi^2 \|y^* - g(x)\|_2^2 + 4L_\varphi^2 \|g(x) - g(\hat{x})\|_2^2} \\ &\leq \sqrt{(1 + 4L_\varphi^2 L_g^2) \|x - \hat{x}\|^2 + 2|\alpha - \alpha^*|^2 + 4L_\varphi^2 \|y^* - g(x)\|_2^2} \\ &\leq \kappa \sqrt{|\alpha - \alpha^*|^2 + \|y^* - g(x)\|_2^2} = \kappa \text{dist}((0, 0), \mathcal{G}(x, \alpha)) \end{aligned}$$

with  $\kappa = \max(\sqrt{2}, \sqrt{(1 + 4L_\varphi^2 L_g^2)c^2 + 4L_\varphi^2})$ , where the third inequality is due to (6).

**Case 2:**  $\alpha^* > \varphi(y^*)$ . In this case, we have  $\mathcal{N}_{\mathbb{R}_+}(\alpha^* - \varphi(y^*)) = \{0\}$ . Recall that

$$(y^*, \alpha^*) \in \arg \min_{(z, \omega) \in \mathbb{R}^m \times \mathbb{R}} \left\{ \frac{1}{2} \|(z, \omega) - (g(x), \alpha)\|_2^2 + \delta_{\mathbb{R}_+}(h(z, \omega)) \right\}$$

where  $h(z, \omega) := \omega - \varphi(z)$  for  $(z, \omega) \in \mathbb{R}^m \times \mathbb{R}$ . By [12, Theorem 10.1&10.49], we have

$$(0, 0) \in (y^*, \alpha^*) - (g(x), \alpha) + D^*h(y^*, \alpha^*)[\mathcal{N}_{\mathbb{R}_+}(\alpha^* - \varphi(y^*))],$$

which by  $\mathcal{N}_{\mathbb{R}_+}(\alpha^* - \varphi(y^*)) = \{0\}$  means that  $(g(x), \alpha) = (y^*, \alpha^*) \in \text{epi } \varphi$ , and then  $(x, \alpha) \in \mathcal{G}^{-1}(0, 0)$ . Thus,  $\text{dist}((x, \alpha), \mathcal{G}^{-1}(0, 0)) \leq \kappa \text{dist}((0, 0), \mathcal{G}(x, \alpha))$  for any  $\kappa > 0$ .  $\square$

To characterize the tangent cones and second-order tangent sets to  $\text{dom } f$  later, we need the MSQC for constraint system  $F(x) \in \text{dom } \vartheta$ , a mild condition used in [15, 29–31].

**Definition 2.3** The MSQC is said to hold for constraint system  $F(x) \in \text{dom}\vartheta$  at a point  $\bar{x} \in \text{dom}f$  if the mapping  $\mathcal{F}(x) := F(x) - \text{dom}\vartheta$  is metrically subregular at  $(\bar{x}, 0)$ .

## 2.2 Subderivative and tangent cones

**Definition 2.4** (see [12, Definition 8.1]) Consider a function  $h: \mathbb{X} \rightarrow \overline{\mathbb{R}}$  and a point  $x \in \text{dom}h$ . Let  $\Delta_\tau h(x): \mathbb{X} \rightarrow \overline{\mathbb{R}}$  denote the first-order difference quotients of  $h$  at  $x$ :

$$\Delta_\tau h(x)(w') := \tau^{-1}[h(x + \tau w') - h(x)] \quad \text{for } \tau > 0.$$

The subderivative function  $dh(x): \mathbb{X} \rightarrow [-\infty, \infty]$  of  $h$  at  $x$  is defined as

$$dh(x)(w) := \liminf_{\substack{\tau \downarrow 0 \\ w' \rightarrow w}} \Delta_\tau h(x)(w') \quad \text{for } w \in \mathbb{X},$$

and  $h$  is called (properly) epi-differentiable at  $x$  if  $\Delta_\tau h(x)$  epi-converge to the (proper) function  $dh(x)$  as  $\tau \downarrow 0$ . The function  $h$  is said to be semidifferentiable at  $x$  for  $w$  if the (possibly infinite) limit  $\lim_{\tau \downarrow 0, w' \rightarrow w} \Delta_\tau h(x)(w')$  exists.

In the sequel, a mapping  $g: \mathbb{X} \rightarrow \mathbb{R}^m$  is said to be semidifferentiable at  $x$  for  $w \in \mathbb{X}$  if its every component function  $g_i: \mathbb{X} \rightarrow \mathbb{R}$  is semidifferentiable at  $x$  for  $w$ , and we denote by  $dg(x)(w)$  the semiderivative of  $g$  at  $x$  for  $w$ . When  $h = \delta_S$  for a closed set  $S \subset \mathbb{X}$ , the subderivative  $dh(x)$  is the indicator function of the tangent cone to  $S$  at  $x$ . The tangent and inner tangent cones to  $S$  at  $x \in S$  are respectively defined as

$$\begin{aligned} \mathcal{T}_S(x) &:= \{w \in \mathbb{X} \mid \exists \tau_k \downarrow 0 \text{ s.t. } \text{dist}(x + \tau_k w, S) = o(\tau_k)\}, \\ \mathcal{T}_S^i(x) &:= \{w \in \mathbb{X} \mid \text{dist}(x + \tau w, S) = o(\tau) \quad \forall \tau \geq 0\}. \end{aligned}$$

The following lemma characterizes the tangent cone and inner tangent cone to  $\text{dom}f$ .

**Lemma 2.1** Fix any  $\bar{x} \in \text{dom}f$ . If the MSQC holds for system  $F(x) \in \text{dom}\vartheta$  at  $\bar{x}$ , then

$$\begin{aligned} \mathcal{T}_{\text{dom}f}(\bar{x}) &= \{w \in \mathbb{X} \mid dF(\bar{x})(w) \in \mathcal{T}_{\text{dom}\vartheta}(F(\bar{x}))\} \\ &= \left\{w \in \mathbb{X} \mid dF(\bar{x})(w) \in \bigcup_{i=1}^s \mathcal{T}_{C_i}(F(\bar{x}))\right\} = \mathcal{T}_{\text{dom}f}^i(\bar{x}). \end{aligned} \quad (8)$$

**Proof:** Since the MSQC holds for system  $F(x) \in \text{dom}\vartheta$  at  $\bar{x}$ , the first equality in (8) is implied by [32, Proposition 1]. From  $\text{dom}\vartheta = \bigcup_{i=1}^s C_i$  and [33, Proposition 3.37],

$$\bigcup_{i=1}^s \mathcal{T}_{C_i}^i(F(\bar{x})) \subseteq \mathcal{T}_{\text{dom}\vartheta}^i(F(\bar{x})) \subseteq \mathcal{T}_{\text{dom}\vartheta}(F(\bar{x})) = \bigcup_{i=1}^s \mathcal{T}_{C_i}(F(\bar{x})) = \bigcup_{i=1}^s \mathcal{T}_{C_i}^i(F(\bar{x})),$$

which implies that the second equality in (8) holds. By combining these relations with the first equality in (8) and  $\mathcal{T}_{\text{dom}f}^i(\bar{x}) \subseteq \mathcal{T}_{\text{dom}f}(\bar{x})$ , the rest only needs to prove that

$$\{w \in \mathbb{X} \mid dF(\bar{x})(w) \in \mathcal{T}_{\text{dom}\vartheta}^i(F(\bar{x}))\} \subseteq \mathcal{T}_{\text{dom}f}^i(\bar{x}). \quad (9)$$

Pick any  $w \in \mathbb{X}$  such that  $dF(\bar{x})(w) \in \mathcal{T}_{\text{dom}\vartheta}^i(F(\bar{x}))$ . By the definition of the inner tangent cone, for any sufficiently small  $\tau > 0$ ,  $\text{dist}(F(\bar{x}) + \tau dF(\bar{x})(w), \text{dom}\vartheta) = o(\tau)$ . Since the MSQC holds for system  $F(x) \in \text{dom}\vartheta$  at  $\bar{x}$ , for each sufficiently small  $\tau > 0$ , there exists  $x_\tau \in \text{dom}f$  and  $\kappa > 0$  such that  $\|\bar{x} + \tau w - x_\tau\| \leq \kappa \text{dist}(F(\bar{x} + \tau w), \text{dom}\vartheta)$ . Since  $F$  is parabolic semidifferentiable at  $\bar{x}$  (see Definition 2.8), it is semidifferentiable at  $\bar{x}$  for  $w$ , and consequently,  $F(\bar{x} + \tau w) - F(\bar{x}) - \tau dF(\bar{x})(w) = o(\tau)$ . Thus,

$$\|\bar{x} + \tau w - x_\tau\| \leq \kappa \text{dist}(F(\bar{x}) + \tau dF(\bar{x})(w), \text{dom}\vartheta) + o(\tau) = o(\tau).$$

Hence,  $\text{dist}(\bar{x} + \tau w, \text{dom}f) = o(\tau)$  and  $w \in \mathcal{T}_{\text{dom}f}^i(\bar{x})$ . The inclusion in (9) holds.  $\square$

With the subderivative functions, we follow the same line as in [26] to introduce the critical cone to an extended real-valued  $h: \mathbb{X} \rightarrow \bar{\mathbb{R}}$  at a point  $(x, v) \in \text{gph}\partial h$ :

$$\mathcal{C}_h(x, v) := \{w \in \mathbb{X} \mid dh(x)(w) = \langle v, w \rangle\}, \quad (10)$$

When  $h = \delta_S$  for a nonempty closed set  $S$ , by [12, Theorem 8.2]  $\mathcal{C}_h(x, v) = \mathcal{T}_S(x) \cap \{v\}^\perp$ .

### 2.3 Second and parabolic subderivatives

To introduce two kinds of second subderivatives for a function  $h: \mathbb{X} \rightarrow \bar{\mathbb{R}}$  at a point  $x \in \text{dom}h$ , we define two kinds of second-order difference quotients for  $h$  at  $x$ :

$$\begin{aligned} \Delta_\tau^2 h(x)(w') &:= \frac{2[h(x + \tau w') - h(x) - \tau dh(x)(w')]}{\tau^2} \quad \text{for } \tau > 0, \\ \Delta_\tau^2 h(x|v)(w') &:= \frac{2[h(x + \tau w') - h(x) - \tau \langle v, w' \rangle]}{\tau^2} \quad \text{for } \tau > 0, \end{aligned}$$

where  $\Delta_\tau^2 h(x)(w') := \infty$  whenever  $h(x + \tau w') = dh(x)(w') = \infty$  or  $-\infty$ .

**Definition 2.5** (see [12, Section 13.B]) Consider a function  $h: \mathbb{X} \rightarrow \bar{\mathbb{R}}$  and a point  $x \in \text{dom}h$ . The second subderivative of  $h$  at  $x$  for  $w$  is defined as

$$d^2 h(x)(w) := \liminf_{\substack{\tau \downarrow 0 \\ w' \rightarrow w}} \Delta_\tau^2 h(x)(w'),$$

while the second subderivative of  $h$  at  $x$  for  $v$  and  $w$  is defined as

$$d^2 h(x|v)(w) := \liminf_{\substack{\tau \downarrow 0 \\ w' \rightarrow w}} \Delta_\tau^2 h(x|v)(w').$$

The function  $h$  is said to be (properly) twice epi-differentiable at  $x$  for  $v$  if  $\Delta_\tau^2 h(x|v)$  epi-converge to the (proper) function  $d^2 h(x|v)$  as  $\tau \downarrow 0$ .

By [12, Proposition 13.5], the second subderivative  $d^2 h(x|v)$  for  $x \in \text{dom}h$  and  $v \in \mathbb{X}$  is lsc and positive homogenous of degree 2, and if it is proper, i.e.,  $d^2 h(x|v)(w) > -\infty$  for all  $w \in \mathbb{X}$  and  $\text{dom}d^2 h(x|v) \neq \emptyset$ ,  $\text{dom}d^2 h(x|v) \subseteq \{w \in \mathbb{X} \mid dh(x)(w) = \langle v, w \rangle\}$ . Parabolic regularity, as demonstrated in [22, 26], plays a prominent role in characterizing the precise formula of second subderivatives of extended real-valued functions and then justifying their epi-differentiability. Its formal definition is stated as follows.

**Definition 2.6** A function  $h : \mathbb{X} \rightarrow \overline{\mathbb{R}}$  is parabolically regular at  $\bar{x} \in \text{dom } h$  for  $\bar{v}$  if for any  $w \in \text{dom } d^2h(\bar{x}|\bar{v})$ , there exist, among the sequences  $\tau_k \downarrow 0$  and  $w^k \rightarrow w$  with  $\Delta_{\tau_k}^2 h(\bar{x}|\bar{v})(w^k) \rightarrow d^2h(\bar{x}|\bar{v})(w)$ , those with the additional property  $\limsup_{k \rightarrow \infty} \frac{\|w^k - w\|}{\tau_k} < \infty$ .

Next we recall the parabolic epi-differentiability of extended real-valued functions, which is another crucial property to achieve their twice epi-differentiability.

**Definition 2.7** (see [12, Section 13.J]) Consider a function  $h : \mathbb{X} \rightarrow \overline{\mathbb{R}}$ ,  $x \in \text{dom } h$ , and  $w$  with  $dh(x)(w)$  finite. Define the parabolic difference quotients of  $h$  at  $x$  for  $w$  by

$$\Delta_{\tau}^2 h(x)(w|z') := \frac{2[h(x + \tau w + \frac{1}{2}\tau^2 z') - h(x) - \tau dh(x)(w)]}{\tau^2} \quad \text{for } \tau > 0.$$

The parabolic subderivative of  $h$  at  $x$  for  $w$  with respect to (w.r.t.)  $z$  is defined as

$$d^2h(x)(w|z) := \liminf_{\substack{\tau \downarrow 0 \\ z' \rightarrow z}} \Delta_{\tau}^2 h(x)(w|z'),$$

and  $h$  is said to be parabolically epi-differentiable at  $x$  for  $w$  if  $\text{dom } d^2h(x)(w|\cdot) \neq \emptyset$  and  $\Delta_{\tau}^2 h(x)(w|\cdot)$  epi-converge to  $d^2h(x)(w|\cdot)$  as  $\tau \downarrow 0$ .

If  $h = \delta_S$  for a nonempty closed set  $S \subset \mathbb{X}$ , for any  $x \in S$  and  $w \in \mathcal{T}_S(x)$ ,  $d^2h(x)(w|\cdot)$  is the indicator of the second-order tangent set to  $S$  at  $x$  for  $w$ . The second-order tangent set and inner second-order tangent set to  $S$  at  $x \in S$  for  $w$  are respectively defined as

$$\begin{aligned} \mathcal{T}_S^2(x, w) &:= \{z \in \mathbb{X} \mid \exists \tau_k \downarrow 0 \text{ s.t. } \text{dist}(x + \tau_k w + \frac{1}{2}\tau_k^2 z, S) = o(\tau_k^2)\}, \\ \mathcal{T}_S^{i,2}(x, w) &:= \{z \in \mathbb{X} \mid \text{dist}(x + \tau w + \frac{1}{2}\tau^2 z, S) = o(\tau^2) \quad \forall \tau \geq 0\}. \end{aligned}$$

One can check that  $\mathcal{T}_S^{i,2}(x, w) = \emptyset$  if  $w \notin \mathcal{T}_S^i(x)$ , and  $\mathcal{T}_S^2(x, w) = \emptyset$  if  $w \notin \mathcal{T}_S(x)$ . The set  $S$  is said to be parabolically derivable at  $x$  for  $w \in \mathcal{T}_S(x)$  if  $\mathcal{T}_S^{i,2}(x, w) = \mathcal{T}_S^2(x, w) \neq \emptyset$ .

**Definition 2.8** A mapping  $g : \mathbb{X} \rightarrow \mathbb{R}^m$  is parabolically semidifferentiable at  $x$  if for every  $w \in \mathbb{X}$ , it is semidifferentiable at  $x$  for  $w$ , and for all  $z \in \mathbb{X}$  the limit

$$\lim_{\substack{\tau \downarrow 0 \\ z' \rightarrow z}} \frac{2[g(x + \tau w + \frac{1}{2}\tau^2 z') - g(x) - \tau dg(x)(w)]}{\tau^2} \text{ exists.}$$

We call this limit, denoted by  $d^2g(x)(w|z)$ , the parabolic semiderivative of  $g$  at  $x$  for  $w$ .

The parabolic semidifferentiability of a mapping, introduced as above, is actually the second-order directional differentiability in the sense of Hadamard [33, Section 2.2.3]. The rest of this section focuses on the characterization of the second-order tangent sets to  $\text{dom } f$ . This requires the following lemma that weakens the twice differentiability of the inner mapping in [26, Theorem 4.3] to be the parabolic semidifferentiability.



**Lemma 2.2** *Let  $h = \varphi \circ g$  where  $\varphi: \mathbb{R}^m \rightarrow \overline{\mathbb{R}}$  is a proper lsc function and  $g: \mathbb{X} \rightarrow \mathbb{R}^m$  is a mapping. Fix any  $\bar{x} \in \text{dom } h$ . For each  $w \in \mathbb{X}$  with  $dg(\bar{x})(w) \in \mathcal{T}_{\text{dom}\varphi}(g(\bar{x}))$ , define*

$$\mathcal{S}_w(p) := \{u \in \mathbb{X} \mid d^2g(\bar{x})(w|u) + p \in \mathcal{T}_{\text{dom}\varphi}^2(g(\bar{x}), dg(\bar{x})(w))\}. \quad (11)$$

*If  $g$  is parabolically semidifferentiable at  $\bar{x}$  and the mapping  $\mathcal{H}$  defined in (3) is subregular at  $(\bar{x}, 0)$  with modulus  $\kappa$ , then  $\mathcal{S}_w(p) \subseteq \mathcal{S}_w(0) + \kappa\|p\|_2\mathbb{B}_{\mathbb{X}}$  for all  $p \in \mathbb{R}^m$  uniformly in  $w$ .*

**Proof:** Fix any  $p \in \mathbb{R}^m$ . Pick any  $u \in \mathcal{S}_w(p)$ . From the expression of  $\mathcal{S}_w$ , it follows that  $d^2g(\bar{x})(w|u) + p \in \mathcal{T}_{\text{dom}\varphi}^2(g(\bar{x})|dg(\bar{x})(w))$ , so there exist  $\tau_k \downarrow 0$  and  $u^k \rightarrow d^2g(\bar{x})(w|u) + p$  such that  $g(\bar{x}) + \tau_k dg(\bar{x})(w) + \frac{1}{2}\tau_k^2 u^k \in \text{dom}\varphi$  for all  $k \in \mathbb{N}$ . For each  $k \in \mathbb{N}$ , write  $x^k := \bar{x} + \tau_k w + \frac{1}{2}\tau_k^2 u$ . By the parabolic semidifferentiability of  $g$  at  $\bar{x}$ , we have

$$g(x^k) = g(\bar{x}) + \tau_k dg(\bar{x})(w) + \frac{1}{2}\tau_k^2 d^2g(\bar{x})(w|u) + o(\tau_k^2).$$

By the subregularity of  $\mathcal{H}$  at  $(\bar{x}, 0)$ , for each sufficiently large  $k$ , there exists  $z^k \in \text{dom } h$  such that

$$\|x^k - z^k\| \leq \kappa \text{dist}(g(x^k), \text{dom}\varphi) \leq \kappa \|g(x^k) - g(\bar{x}) - \tau_k dg(\bar{x})(w) - \frac{1}{2}\tau_k^2 u^k\|_2 \leq \frac{\tau_k^2}{2} \kappa \|p\|_2 + o(\tau_k^2),$$

which implies that the sequence  $d^k := \frac{2(x^k - z^k)}{\tau_k^2}$  is bounded. Therefore, there exists  $d \in \mathbb{X}$  such that  $d^k \rightarrow d$  with  $\|d\| \leq \kappa\|p\|_2$ . On the other hand, for all sufficiently large  $k$ ,

$$\text{dom } h \ni z^k = \bar{x} + \tau_k w + (\tau_k^2/2)(u - d) + o(\tau_k^2),$$

which the parabolic semidifferentiability of  $g$  at  $\bar{x}$  implies that

$$\text{dom}\varphi \ni g(z^k) = g(\bar{x}) + \tau_k dg(\bar{x})(w) + (\tau_k^2/2)d^2g(\bar{x})(w|u - d) + o(\tau_k^2).$$

This means that  $d^2g(\bar{x})(w|u - d) \in \mathcal{T}_{\text{dom}\varphi}^2(g(\bar{x}), dg(\bar{x})(w))$  or  $u - d \in \mathcal{S}_w(0)$ . Hence,  $\text{dist}(u, \mathcal{S}_w(0)) \leq \|d\| \leq \kappa\|p\|_2$ . The result then follows.  $\square$

For the composite function  $h$  in Lemma 2.2, the following proposition establishes the bridge between the second-order tangent set to  $\text{dom } h$  and the one to  $\text{dom } \varphi$ . This result extends the conclusions of [26, Theorem 4.5] and [22, Proposition 4.3 (ii)] to the constraint system  $g(x) \in \text{dom } \varphi$ , where  $g$  is parabolically semidifferentiable rather than twice differentiable, and the Clarke regularity of  $\text{dom}\varphi$  is also removed.

**Proposition 2.2** *Let  $h$  be the same as the one in Lemma 2.2. Consider any  $\bar{x} \in \text{dom } h$ . Suppose that  $g$  is parabolically semidifferentiable at  $\bar{x}$ , and that the mapping  $\mathcal{H}$  in (3) is subregular at  $(\bar{x}, 0)$  with modulus  $\kappa$ . Then, for any  $w \in \mathcal{T}_{\text{dom } h}(\bar{x})$ ,*

$$z \in \mathcal{T}_{\text{dom } h}^2(\bar{x}, w) \iff d^2g(\bar{x})(w|z) \in \mathcal{T}_{\text{dom}\varphi}^2(g(\bar{x}), dg(\bar{x})(w)), \quad (12)$$

*and if in addition  $\text{dom}\varphi$  is parabolically derivable at  $g(\bar{x})$  for  $dg(\bar{x})(w)$ , so is  $\text{dom } h$  at  $\bar{x}$  for  $w$ .*

**Proof:** Fix any  $w \in \mathcal{T}_{\text{dom}h}(\bar{x})$ . Pick any  $z \in \mathcal{T}_{\text{dom}h}^2(\bar{x}, w)$ . Then there exist  $\tau_k \downarrow 0$  and  $z^k \rightarrow z$  such that  $\bar{x} + \tau_k w + \frac{1}{2}\tau_k^2 z^k \in \text{dom}h$  for each  $k \in \mathbb{N}$ . By the parabolic semidifferentiability of  $g$ , for all sufficiently large  $k$ , it holds that

$$\text{dom}\varphi \ni g(\bar{x} + \tau_k w + \frac{1}{2}\tau_k^2 z^k) = g(\bar{x}) + \tau_k dg(\bar{x})(w) + \frac{1}{2}\tau_k^2 d^2g(\bar{x})(w|z) + o(\tau_k^2).$$

Hence,  $d^2g(\bar{x})(w|z) \in \mathcal{T}_{\text{dom}\varphi}^2(g(\bar{x}), dg(\bar{x})(w))$ , and the implication  $\implies$  follows. For the converse inclusion, pick any  $z \in \mathbb{X}$  such that  $d^2g(\bar{x})(w|z) \in \mathcal{T}_{\text{dom}\varphi}^2(g(\bar{x}), dg(\bar{x})(w))$ . There exist  $\tau_k \downarrow 0$  and  $\xi^k \rightarrow d^2g(\bar{x})(w|z)$  such that  $g(\bar{x}) + \tau_k dg(\bar{x})(w) + \frac{1}{2}\tau_k^2 \xi^k \in \text{dom}\varphi$  for each  $k \in \mathbb{N}$ . Write  $x^k := \bar{x} + \tau_k w + \frac{1}{2}\tau_k^2 z$ . By the parabolic semidifferentiability of  $g$ ,  $g(x^k) = g(\bar{x}) + \tau_k dg(\bar{x})(w) + \frac{1}{2}\tau_k^2 d^2g(\bar{x})(w|z) + o(\tau_k^2)$  for all sufficiently large  $k$ . From the subregularity of  $\mathcal{H}$  at  $(\bar{x}, 0)$ , for each sufficiently large  $k$ , there is  $z^k \in \text{dom}h$  such that

$$\|x^k - z^k\| \leq \kappa \text{dist}(g(x^k), \text{dom}\varphi) \leq \kappa \|g(x^k) - [g(\bar{x}) + \tau_k dg(\bar{x})(w) + \frac{1}{2}\tau_k^2 \xi^k]\|_2 = o(\tau_k^2).$$

Thus, there exists a sequence  $\{z^k\} \subseteq \text{dom}h$  with  $z^k = \bar{x} + \tau_k w + \frac{1}{2}\tau_k^2(z + o(\tau_k^2)/\tau_k^2)$  for all  $k$  large enough. Consequently,  $z \in \mathcal{T}_{\text{dom}h}^2(\bar{x}, w)$ , and the implication  $\impliedby$  also holds.

Fix any  $z \in \mathcal{T}_{\text{dom}h}^2(\bar{x}, w)$ . From the subregularity of  $\mathcal{H}$  at  $(\bar{x}, 0)$ , for any sufficiently small  $\tau > 0$ , we have  $\text{dist}(\bar{x} + \tau w + \frac{1}{2}\tau^2 z, \text{dom}h) \leq \kappa \text{dist}(g(\bar{x} + \tau w + \frac{1}{2}\tau^2 z), \text{dom}\varphi)$  or

$$\text{dist}\left(z, \frac{\text{dom}h - \bar{x} - \tau w}{\frac{1}{2}\tau^2}\right) \leq \kappa \text{dist}\left(\Delta_\tau^2 g(\bar{x})(w|z), \frac{\text{dom}\varphi - g(\bar{x}) - \tau dg(\bar{x})(w)}{\frac{1}{2}\tau^2}\right). \quad (13)$$

Clearly,  $\Delta_\tau^2 g(\bar{x})(w|z) \rightarrow d^2g(\bar{x})(w|z) := \zeta$  as  $\tau \downarrow 0$ . Moreover, since  $\text{dom}\varphi$  is parabolically derivable at  $g(\bar{x})$  for  $dg(\bar{x})(w)$ , by invoking [12, Corollary 4.7], when  $\tau \downarrow 0$ ,

$$\text{dist}\left(\zeta, \frac{\text{dom}\varphi - g(\bar{x}) - \tau dg(\bar{x})(w)}{\frac{1}{2}\tau^2}\right) \rightarrow \text{dist}\left(\zeta, \mathcal{T}_{\text{dom}\varphi}^{i,2}(g(\bar{x}), dg(\bar{x})(w))\right) = 0.$$

Thus, when the right hand side of (13) tends to zero as  $\tau \downarrow 0$ , the left hand side must go to 0, which implies that  $z \in \mathcal{T}_{\text{dom}h}^{i,2}(\bar{x}, w)$ . Therefore, the parabolic derivability of  $\text{dom}\varphi$  at  $g(\bar{x})$  for  $dg(\bar{x})(w)$  implies that of  $\text{dom}h$  at  $\bar{x}$  for  $w$ , provided that the nonemptiness of  $\mathcal{T}_{\text{dom}\varphi}^2(g(\bar{x}), dg(\bar{x})(w))$  implies that of  $\mathcal{T}_{\text{dom}h}^2(\bar{x}, w)$ . Pick any  $u \in \mathcal{T}_{\text{dom}\varphi}^2(g(\bar{x}), dg(\bar{x})(w))$ , whose nonemptiness is implied by the parabolic derivability of  $\text{dom}\varphi$ . Pick any  $z \in \mathbb{X}$  and take  $p = u - d^2g(\bar{x})(w|z)$ . Clearly,  $z \in \mathcal{S}_w(p)$  with  $p = u - d^2g(\bar{x})(w|z)$ . By Lemma 2.2, there exists  $\tilde{z} \in \mathcal{S}_w(0)$  such that  $\|z - \tilde{z}\| \leq \kappa \|p\|_2$ . From  $\tilde{z} \in \mathcal{S}_w(0)$  and the previous equivalence,  $\tilde{z} \in \mathcal{T}_{\text{dom}h}^2(\bar{x}, w)$  and  $\mathcal{T}_{\text{dom}h}^2(\bar{x}, w) \neq \emptyset$ . The proof is completed.  $\square$

**Corollary 2.1** *Let  $h$  be the same as the one in Lemma 2.2. Consider any  $\bar{x} \in \text{dom}h$ . Suppose that  $g$  is parabolically semidifferentiable at  $\bar{x}$ , and that the mapping  $\mathcal{G}$  defined in (3) is subregular at  $((\bar{x}, h(\bar{x})), (0, 0))$ . Then, for any  $w \in \text{dom}dh(\bar{x})$ ,*

$$d^2h(\bar{x})(w|z) = d^2\varphi(g(\bar{x}))(dg(\bar{x})(w) | d^2g(\bar{x})(w|z)) \quad \forall z \in \mathbb{X}.$$

*and furthermore,  $h$  is parabolically epi-differentiable at  $\bar{x}$  for  $w$  if in addition  $\varphi$  is parabolically epi-differentiable at  $g(\bar{x})$  for  $dg(\bar{x})(w)$ .*

**Proof:** Fix any  $w \in \text{dom } dh(\bar{x})$ . Note that  $\text{epi } h = G^{-1}(\text{epi } \varphi)$  with  $G(x, \omega) = (g(x), \omega)$  for  $(x, \omega) \in \mathbb{X} \times \mathbb{R}$ . Let  $\alpha = h(\bar{x})$  and  $\beta = dh(\bar{x})(w)$ . By Proposition 2.2,

$$\begin{aligned} (z, t) \in \mathcal{T}_{\text{epi } h}^2((\bar{x}, \alpha), (w, \beta)) &\iff (d^2g(\bar{x})(w|z), t) \in \mathcal{T}_{\text{epi } \varphi}^2((g(\bar{x}), \alpha), (dg(\bar{x})(w), \beta)) \\ &\iff (d^2g(\bar{x})(w|z), t) \in \text{epi } d^2\varphi(g(\bar{x}))(dg(\bar{x})(w) | \cdot), \end{aligned}$$

This implies that the desired conclusion holds.  $\square$

For any  $w \in \mathbb{X}$  with  $dF(\bar{x})(w) \in \mathcal{T}_{\text{dom } \vartheta}(F(\bar{x}))$ , by [33, Proposition 3.37] we have

$$\begin{aligned} \bigcup_{i=1}^s \mathcal{T}_{C_i}^{i,2}(F(\bar{x}), dF(\bar{x})(w)) &\subseteq \mathcal{T}_{\text{dom } \vartheta}^{i,2}(F(\bar{x}), dF(\bar{x})(w)) \subseteq \mathcal{T}_{\text{dom } \vartheta}^2(F(\bar{x}), dF(\bar{x})(w)) \\ &= \bigcup_{i=1}^s \mathcal{T}_{C_i}^2(F(\bar{x}), dF(\bar{x})(w)) = \bigcup_{i=1}^s \mathcal{T}_{C_i}^{i,2}(F(\bar{x}), dF(\bar{x})(w)). \end{aligned}$$

Also,  $0 \in \mathcal{T}_{\text{dom } \vartheta}^2(F(\bar{x}), dF(\bar{x})(w))$ . From Proposition 2.2, the following corollary holds.

**Corollary 2.2** *Fix any  $\bar{x} \in \text{dom } f$ . If the MSQC holds for system  $F(x) \in \text{dom } \vartheta$  at  $\bar{x}$ , then  $\text{dom } f$  is parabolically derivable at  $\bar{x}$  for any  $w \in \mathcal{T}_{\text{dom } f}(\bar{x})$ , i.e., for any  $w \in \mathcal{T}_{\text{dom } f}(\bar{x})$ ,*

$$\emptyset \neq \mathcal{T}_{\text{dom } f}^2(\bar{x}, w) = \{z \in \mathbb{X} \mid d^2F(\bar{x})(w|z) \in \mathcal{T}_{\text{dom } \vartheta}^2(F(\bar{x}), dF(\bar{x})(w))\} = \mathcal{T}_{\text{dom } f}^{i,2}(\bar{x}, w).$$

Note that  $\text{epi } f = \mathcal{L}^{-1}(\text{dom } f)$  with  $\mathcal{L}(x, \alpha) = x$  for  $(x, \alpha) \in \mathbb{X} \times \mathbb{R}$ . Combining Corollary 2.2 and Proposition 2.2 with  $h = \delta_{\text{epi } f}$ ,  $\varphi = \delta_{\text{dom } f}$ ,  $g = \mathcal{L}$  yields Corollary 2.3.

**Corollary 2.3** *Consider any  $\bar{x} \in \text{dom } f$ . If the MSQC holds for system  $F(x) \in \text{dom } \vartheta$  at  $\bar{x}$ , then for any  $w \in \mathcal{T}_{\text{dom } f}(\bar{x})$ ,  $\mathcal{T}_{\text{epi } f}^2((\bar{x}, f(\bar{x})), (w, df(\bar{x})(w))) \neq \emptyset$ .*

### 3 Second-order variational properties of PWTD functions

Before characterizing the second and parabolic subderivatives of PWTD functions, we take a closer look at their subderivatives. The following proposition provides the characterization for them, and extends the result of [12, Proposition 10.21] to PWTD functions. Since its proof is similar to that of [12, Proposition 10.21], we do not include it.

**Proposition 3.1** *Consider a PWTD function  $h: \mathbb{R}^m \rightarrow \overline{\mathbb{R}}$  with  $\emptyset \neq \text{dom } h = \bigcup_{i=1}^s \Omega_i$  and  $h = h_i$ , a twice differentiable function, relative to each  $\Omega_i$ . Then,*

(i) *dom  $h$  is closed and  $h$  is continuous relative to dom  $h$  (hence lsc on  $\mathbb{R}^m$ );*

(ii) *for each  $y \in \text{dom } h$ ,  $dh(y)$  is a proper piecewise linear function with*

$$\text{dom } dh(y) = \mathcal{T}_{\text{dom } h}(y) = \bigcup_{i \in J_y} \mathcal{T}_{\Omega_i}(y) \quad \text{for } J_y := \{i \in [s] \mid y \in \Omega_i\},$$

*it takes the form of  $\langle \nabla h_i(y), \cdot \rangle$  relative to each  $\mathcal{T}_{\Omega_i}(y)$  for  $i \in J_y$ , and*

$$dh(y)(w) = \lim_{\tau \downarrow 0} \Delta_\tau h(y)(w) = \begin{cases} \langle \nabla h_i(y), w \rangle & \text{if } w \in \mathcal{T}_{\Omega_i}(y), i \in J_y, \\ \infty & \text{if } w \notin \mathcal{T}_{\text{dom } h}(y). \end{cases} \quad (14)$$

*Consequently,  $dh(y)$  is a properly epi-differentiable function.*

Now we characterize the second subderivatives of PWTD functions, and show that a PWTD function is twice epi-differentiable at any regular points of its domain, which extends [12, Proposition 13.9] for PWLQ convex functions to PWTD functions.

**Proposition 3.2** *Let  $h: \mathbb{R}^m \rightarrow \overline{\mathbb{R}}$  be the same as the one in Proposition 3.1. Consider any  $y \in \text{dom } h$ . Then, the following assertions hold.*

(i)  $d^2h(y)$  is a PWLQ function and at any  $w \in \mathbb{R}^m$  it takes the following form:

$$d^2h(y)(w) = \lim_{\tau \downarrow 0} \Delta_\tau^2 h(y)(w) = \begin{cases} \langle w, \nabla^2 h_i(y) w \rangle & \text{if } w \in \mathcal{T}_{\Omega_i}(y), i \in J_y, \\ \infty & \text{if } w \notin \mathcal{T}_{\text{dom}h}(y). \end{cases} \quad (15)$$

(ii) If  $\partial h(y) = \widehat{\partial}h(y)$ , for every  $v \in \partial h(y)$ ,  $d^2h(y|v)$  is a proper PWLQ function with

$$\begin{aligned} d^2h(y|v)(w) &= \begin{cases} \langle w, \nabla^2 h_k(y) w \rangle & \text{if } w \in \mathcal{T}_{\Omega_i}(y) \cap \mathcal{C}_h(y, v), i \in J_y, \\ \infty & \text{if } w \notin \mathcal{C}_h(y, v) \end{cases} \\ &= d^2h(y)(w) + \delta_{\mathcal{C}_h(y, v)}(w) = \lim_{\tau \downarrow 0} \Delta_\tau^2 h(y|v)(w), \end{aligned} \quad (16)$$

and consequently  $d^2h(y|v)$  is properly twice epi-differentiable at  $y$  for  $v$ .

**Proof:** Fix any  $w \in \mathbb{R}^m$  and define the index set  $J_{y, w} := \{j \in J_y \mid w \in \mathcal{T}_{\Omega_j}(y)\}$ .

(i) We first consider the case that  $w \in \mathcal{T}_{\text{dom}h}(y) = \bigcup_{i \in J_y} \mathcal{T}_{\Omega_i}(y)$ . By Definition 2.5, there exist  $\tau_\nu \downarrow 0$  and  $w^\nu \rightarrow w$  with  $w^\nu \in \tau_\nu^{-1}[\bigcup_{i \in J_y} \Omega_i - y]$  for each  $\nu \in \mathbb{N}$  such that

$$d^2h(y)(w) = \lim_{\nu \rightarrow \infty} \frac{2[h(y + \tau_\nu w^\nu) - h(y) - dh(y)(w^\nu)]}{\tau_\nu^2}.$$

Obviously, for each  $\nu \in \mathbb{N}$ ,  $y + \tau_\nu w^\nu \in \text{dom } h$  and  $w^\nu \in \text{dom } dh(y)$ . Then, there exist  $\bar{k} \in J_y$  and  $N \subset \mathbb{N}$  such that for all  $\nu \in N$ ,  $y + \tau_\nu w^\nu \in \Omega_{\bar{k}}$  and  $w^\nu \in \mathcal{T}_{\Omega_{\bar{k}}}(y)$ , and hence

$$d^2h(y)(w) = \lim_{\nu \rightarrow \infty} \frac{2[h_{\bar{k}}(y + \tau_\nu w^\nu) - h_{\bar{k}}(y) - \langle \nabla h_{\bar{k}}(y), w^\nu \rangle]}{\tau_\nu^2} = \langle w, \nabla^2 h_{\bar{k}}(y) w \rangle. \quad (17)$$

On the other hand, by the polyhedrality of each  $\Omega_i$ , for all sufficiently small  $\tau > 0$ ,  $y + \tau w \in \bigcap_{j \in J_{y, w}} \Omega_j$ . Consequently, for each  $j \in J_{y, w}$ , it holds that

$$\begin{aligned} \langle w, \nabla^2 h_j(y) w \rangle &= \lim_{\tau \downarrow 0} \frac{2[h_j(y + \tau w) - h_j(y) - \langle \nabla h_j(y), w \rangle]}{\tau^2} \\ &= \lim_{\tau \downarrow 0} \frac{2[h(y + \tau w) - h(y) - dh(y)(w)]}{\tau^2}. \end{aligned} \quad (18)$$

Note that  $\bar{k} \in J_{y, w}$ . By combining (18) for  $j = \bar{k}$  with (17), when  $w \in \mathcal{T}_{\Omega_{\bar{k}}}(y)$  for  $k \in J_y$ ,

$$d^2h(y)(w) = \langle w, \nabla^2 h_k(y) w \rangle = \lim_{\tau \downarrow 0} \Delta_\tau^2 h(y)(w).$$

When  $w \notin \mathcal{T}_{\text{dom}h}(y)$ , for any sufficiently small  $\tau > 0$  and any  $w'$  sufficiently close to  $w$ ,  $y + \tau w' \notin \text{dom} h$ , which implies that  $d^2h(y)(w) = \infty = \lim_{\tau \downarrow 0} \Delta_\tau^2 h(y)(w)$ . Thus,  $d^2h(y)$  has the expression as in (15) and hence is a PWLQ function.

(ii) Pick any  $v \in \partial h(y) = \widehat{\partial}h(y)$ . When  $w \notin \mathcal{T}_{\text{dom}h}(y)$ , there exists  $\delta > 0$  such that for any  $\tau \in (0, \delta)$  and any  $w'$  with  $\|w' - w\| \leq \delta$ ,  $w' \notin \tau^{-1}[\text{dom} h - y]$ . By the definition of second subderivatives,  $d^2h(y|v)(w) = \infty = d^2h(y)(w)$ , where the second equality is by part (i), so it suffices to consider that  $w \in \mathcal{T}_{\text{dom}h}(y)$ . From the definition of  $d^2h(y|v)$ , there exist  $\tau_\nu \downarrow 0$  and  $w^\nu \rightarrow w$  with  $w^\nu \in \tau_\nu^{-1}[\bigcup_{i \in J_y} \Omega_i - y]$  for each  $\nu \in \mathbb{N}$  such that

$$d^2h(y|v)(w) = \lim_{\nu \rightarrow \infty} \frac{2[h(y + \tau_\nu w^\nu) - h(y) - \tau_\nu \langle v, w^\nu \rangle]}{\tau_\nu^2}.$$

Obviously, for each  $\nu \in \mathbb{N}$ ,  $y + \tau_\nu w^\nu \in \text{dom} h$ . Then, there exist  $\tilde{k} \in J_y$  and  $N \subset \mathbb{N}$  such that for all  $\nu \in N$ ,  $y + \tau_\nu w^\nu \in \Omega_{\tilde{k}}$  and  $w^\nu \in \mathcal{T}_{\Omega_{\tilde{k}}}(y)$ , and consequently,

$$\begin{aligned} d^2h(y|v)(w) &= \lim_{\nu \rightarrow \infty} \frac{2[h_{\tilde{k}}(y + \tau_\nu w^\nu) - h_{\tilde{k}}(y) - \tau_\nu \langle v, w^\nu \rangle]}{\tau_\nu^2} \\ &= \lim_{\nu \rightarrow \infty} \left[ \langle w^\nu, \nabla^2 h_{\tilde{k}}(y) w^\nu \rangle + \frac{2(dh(y)(w^\nu) - \langle v, w^\nu \rangle)}{\tau_\nu} + \frac{o(\tau_\nu^2)}{\tau_\nu^2} \right] \\ &\geq \begin{cases} \langle w, \nabla^2 h_{\tilde{k}}(y) w \rangle & \text{if } w \in \mathcal{C}_h(y, v), \\ \infty & \text{if } w \in \mathcal{T}_{\text{dom}h}(y) \setminus \mathcal{C}_h(y, v), \end{cases} \end{aligned} \quad (19)$$

where the last inequality is due to  $dh(y)(w') \geq \langle v, w' \rangle$  for any  $w' \in \mathbb{R}^n$  implied by  $v \in \widehat{\partial}h(y)$  and the continuity of  $dh(y)$  relative to its domain by Proposition 3.1 (i). Note that  $\tilde{k} \in J_{y,w}$ . Combining equality (18) for  $j = \tilde{k}$  with (19) yields that

$$d^2h(y|v)(w) = \begin{cases} \langle w, \nabla^2 h_k(y) w \rangle & \text{if } w \in \mathcal{T}_{\Omega_k}(y) \cap \mathcal{C}_h(y, v), k \in J_y, \\ \infty & \text{if } w \in \mathcal{T}_{\text{dom}h}(y) \setminus \mathcal{C}_h(y, v). \end{cases}$$

Thus, we establish the first and the third equalities in (16). The second equality in (16) is due to the first one and part (i). The proof is completed.  $\square$

Next we provide the characterization of parabolic subderivatives of PWTD functions, and show that under the same condition as in Proposition 3.2, a PWTD function  $h$  is parabolically epi-differentiable at any  $y \in \text{dom} h$  for all  $w \in \text{dom} dh(y)$ .

**Proposition 3.3** *Let  $h: \mathbb{R}^m \rightarrow \overline{\mathbb{R}}$  be the same as the one in Proposition 3.1. Fix any  $y \in \text{dom} h$  and  $w \in \text{dom} dh(y)$ . Then, the following statements hold.*

(i)  $\mathcal{T}_{\text{dom}h}^2(y, w) = \bigcup_{j \in J_{y,w}} \mathcal{T}_{\Omega_j}^2(y, w) = \mathcal{T}_{\mathcal{T}_{\text{dom}h}(y)}(w)$  with  $J_{y,w} := \{i \in J_y \mid w \in \mathcal{T}_{\Omega_i}(y)\}$ .

(ii)  $d^2h(y)(w|z) < \infty$  if and only if  $z \in \mathcal{T}_{\text{dom}h}^2(y, w)$ .

(iii)  $h$  is proper and parabolically epi-differentiable at  $y$  for  $w$ , and for any  $z \in \mathbb{R}^m$ ,

$$d^2h(y)(w|z) = \lim_{\tau \downarrow 0} \frac{2[h(y + \tau w + \frac{1}{2}\tau^2 z) - h(y) - \tau dh(y)(w)]}{\tau^2}. \quad (20)$$

(iv) If  $h$  is regular at  $y$ , with  $\mathcal{A}_h(y, w) := \{\xi \in \partial h(y) \mid dh(y)(w) = \langle \xi, w \rangle\}$  it holds that

$$d^2h(y)(w|z) = d^2h(y)(w) + \sup_{\xi \in \mathcal{A}_h(y, w)} \langle \xi, z \rangle \quad \text{for } z \in \mathbb{R}^m. \quad (21)$$

**Proof:** (i) By Proposition 3.1,  $w \in \mathcal{T}_{\text{dom } h}(y)$ . From [33, Proposition 3.37], it follows that  $\mathcal{T}_{\text{dom } h}^2(y, w) = \bigcup_{i=1}^s \mathcal{T}_{\Omega_i}^2(y, w)$ . Because  $\mathcal{T}_{\Omega_j}^2(y, w) = \mathcal{T}_{\mathcal{T}_{\Omega_j}(y)}(w)$  for each  $\Omega_j$  by [12, Proposition 13.12], we have  $w \in \mathcal{T}_{\Omega_j}(y)$  if  $z \in \mathcal{T}_{\Omega_j}^2(y, w)$ . Then,  $\bigcup_{i=1}^s \mathcal{T}_{\Omega_i}^2(y, w) = \bigcup_{j \in J_{y, w}} \mathcal{T}_{\Omega_j}^2(y, w)$ . By [33, Proposition 3.37],  $\bigcup_{j \in J_{y, w}} \mathcal{T}_{\Omega_j}^2(y, w) = \mathcal{T}_{\bigcup_{j \in J_{y, w}} \mathcal{T}_{\Omega_j}(y)}(w) = \mathcal{T}_{\text{dom } h}(y)(w)$ .

(ii) Pick any  $z \in \mathbb{R}^m$  with  $d^2h(y)(w|z) < \infty$ . By Definition 3.3, there exist sequences  $\tau_\nu \downarrow 0$  and  $z^\nu \rightarrow z$  such that

$$\infty > d^2h(y)(w|z) = \lim_{\nu \rightarrow \infty} \frac{2[h(y + \tau_\nu w + \frac{1}{2}\tau_\nu^2 z^\nu) - h(y) - \tau_\nu dh(y)(w)]}{\tau_\nu^2},$$

which implies that  $y + \tau_\nu w + \frac{1}{2}\tau_\nu^2 z^\nu \in \text{dom } h$  for all sufficiently large  $\nu$ . Then, there must exist  $\bar{k} \in [s]$  and an index set  $N \subset \mathbb{N}$  such that  $\{y + \tau_\nu w + \frac{1}{2}\tau_\nu^2 z^\nu\}_{\nu \in N} \subseteq \Omega_{\bar{k}}$ . Obviously,  $w \in \mathcal{T}_{\Omega_{\bar{k}}}(y)$  and  $z \in \mathcal{T}_{\Omega_{\bar{k}}}^2(y, w) \subseteq \mathcal{T}_{\text{dom } h}^2(y, w)$ . Moreover, from the last equation,

$$\begin{aligned} d^2h(y)(w|z) &= \lim_{\nu \rightarrow \infty} \frac{2[h_{\bar{k}}(y + \tau_\nu w + \frac{1}{2}\tau_\nu^2 z^\nu) - h_{\bar{k}}(y) - \tau_\nu dh_{\bar{k}}(y)(w)]}{\tau_\nu^2} \\ &= \langle \nabla h_{\bar{k}}(y), z \rangle + \langle w, \nabla^2 h_{\bar{k}}(y)w \rangle. \end{aligned} \quad (22)$$

Conversely, pick any  $z \in \mathcal{T}_{\text{dom } h}^2(y, w)$ . There exists  $\tilde{k} \in [s]$  such that  $z \in \mathcal{T}_{\Omega_{\tilde{k}}}^2(y, w)$ . Since  $\Omega_{\tilde{k}}$  is polyhedral, for any  $\tau > 0$  enough small,  $y + \tau w + \frac{1}{2}\tau^2 z \in \Omega_{\tilde{k}}$ , which implies that

$$\begin{aligned} \lim_{\tau \downarrow 0} \frac{h(y + \tau w + \frac{1}{2}\tau^2 z) - h(y) - \tau dh(y)(w)}{\frac{1}{2}\tau^2} &= \lim_{\tau \downarrow 0} \frac{h_{\tilde{k}}(y + \tau w + \frac{1}{2}\tau^2 z) - h_{\tilde{k}}(y) - \tau dh_{\tilde{k}}(y)(w)}{\frac{1}{2}\tau^2} \\ &= \langle w, \nabla^2 h_{\tilde{k}}(y)w \rangle + \langle \nabla h_{\tilde{k}}(y), z \rangle. \end{aligned}$$

This shows that  $d^2h(y)(w|z) < \infty$ , and the desired equivalence then follows.

(iii) Fix any  $z \in \mathbb{R}^m$ . If  $z \notin \mathcal{T}_{\text{dom } h}^2(y, w)$ , for any sufficiently small  $\tau > 0$ , we have  $y + \tau w + \frac{1}{2}\tau^2 z \notin \text{dom } h$ . Together with  $w \in \text{dom } dh(y)$  and part (i), it holds that

$$d^2h(y)(w|z) = \infty = \lim_{\tau \downarrow 0} \frac{2[h(y + \tau w + \frac{1}{2}\tau^2 z) - h(y) - \tau dh(y)(w)]}{\tau^2}.$$

Next we consider the case that  $z \in \mathcal{T}_{\text{dom } h}^2(y, w) = \bigcup_{j \in J_{y, w}} \mathcal{T}_{\Omega_j}^2(y, w)$ . For convenience, write  $J_{y, w, z} := \{k \in J_{y, w} \mid z \in \mathcal{T}_{\Omega_k}^2(y, w)\}$ . For each  $k \in J_{y, w, z}$ , from  $z \in \mathcal{T}_{\Omega_k}^2(y, w)$ , the polyhedrality of each  $\Omega_i$ , and [12, Proposition 13.12], for any sufficiently small  $\tau > 0$ ,

$y + \tau w + \frac{1}{2}\tau^2 z \in \Omega_k$ . Thus, for each  $k \in J_{y,w,z}$ ,

$$\begin{aligned} \langle w, \nabla^2 h_k(y)w \rangle + \langle \nabla h_k(y), z \rangle &= \lim_{\tau \downarrow 0} \frac{2[h_k(y + \tau w + \frac{1}{2}\tau^2 z) - h_k(y) - \tau dh_k(y)(w)]}{\tau^2} \\ &= \lim_{\tau \downarrow 0} \frac{2[h(y + \tau w + \frac{1}{2}\tau^2 z) - h(y) - \tau dh(y)(w)]}{\tau^2}. \end{aligned} \quad (23)$$

On the other hand, from the arguments for part (ii), we have  $\bar{k} \in J_{y,w,z}$ . Now by combining equality (23) for  $k = \bar{k}$  with (22), we obtain (20). Note that  $0 \in \mathcal{T}_{\Omega_j}^2(y, w)$  for each  $j \in J_{y,w}$ . From part (ii), it follows that  $d^2 h(y)(w|0) < \infty$ . This, together with (20), shows that  $h$  is parabolically epi-differentiable at  $y$  for  $w$ .

(iv) Fix any  $z \in \mathbb{R}^m$ . Pick any  $k \in J_{y,w,z}$ . From the polyhedrality of every  $\Omega_i$ , for any sufficiently small  $\tau > 0$ ,  $y + \tau w + \frac{1}{2}\tau^2 z \in \Omega_k$ . Together with part (iii), it holds that

$$\begin{aligned} d^2 h(y)(w|z) &= \lim_{\tau \downarrow 0} \frac{2[h_k(y + \tau w + \frac{1}{2}\tau^2 z) - h_k(y) - \tau dh_k(y)(w)]}{\tau^2} \\ &= \langle w, \nabla^2 h_k(y)w \rangle + \langle \nabla h_k(y), z \rangle = d^2 h(y)(w) + \langle \nabla h_k(y), z \rangle \end{aligned}$$

where the second equality is using  $dh(y)(w) = \langle \nabla h_k(y), w \rangle$ , and the last one is due to Proposition 3.2 (i). By the proof of part (i),  $\text{dom } d^2 h(y)(w|\cdot) = \bigcup_{i \in J_{y,w}} \mathcal{T}_{\Omega_i}(y)(w)$ . Then

$$d^2 h(y)(w|z) = d^2 h(y)(w) + \begin{cases} \langle \nabla h_i(y), z \rangle & \text{if } z \in \mathcal{T}_{\Omega_i}(y)(w), i \in J_{y,w}, \\ \infty & \text{if } z \notin \mathcal{T}_{\text{dom } h}(y)(w). \end{cases} \quad (24)$$

Let  $h_y(w') := dh(y)(w')$  for  $w' \in \mathbb{R}^m$ . By Proposition 3.1 (ii),  $h_y$  is a proper piecewise linear function with  $\text{dom } h_y = \bigcup_{i \in J_y} \mathcal{T}_{\Omega_i}(w)$ . By applying Proposition 3.1 (ii) to  $h_y$ ,

$$dh_y(w)(z) = \begin{cases} \langle \nabla h_i(y), z \rangle & \text{if } z \in \mathcal{T}_{\Omega_i}(y)(w), i \in J_{y,w}, \\ \infty & \text{if } z \notin \mathcal{T}_{\text{dom } h}(y)(w). \end{cases}$$

Together with (24), it follows that  $d^2 h(y)(w|z) = d^2 h(y)(w) + dh_y(w)(z)$ . Because  $h$  is assumed to be regular at  $y$ , by invoking [12, Theorem 8.30] it holds that

$$h_y(w') = \sup_{v \in \partial h(y)} \langle v, w' \rangle \quad \text{for } w' \in \mathbb{R}^n.$$

Note that  $h_y$  is a closed convex function. Hence,  $\partial h_y(w) = \arg \max_{v \in \partial h(y)} \langle v, w \rangle$  and

$$dh_y(w)(z) = \sup_{\xi \in \partial h_y(w)} \langle \xi, z \rangle = \sup_{\xi \in \mathbb{R}^m} \left\{ \langle \xi, z \rangle \text{ s.t. } \xi \in \partial h(y), \langle \xi, w \rangle = dh(y)(w) \right\}.$$

Substituting this into  $d^2 h(y)(w|z) = d^2 h(y)(w) + dh_y(w)(z)$  and using the definition of  $\mathcal{A}_h(y, w)$ , we obtain the desired result. The proof is then completed.  $\square$

Proposition 3.3 extends the result of [12, Exercise 13.61] for PWLQ convex functions to PWTQ functions. From Propositions 3.2 and 3.3, we have the parabolic regularity of PWTQ functions, which partly extends the result of [12, Theorem 13.67].

**Proposition 3.4** *Let  $h: \mathbb{R}^m \rightarrow \overline{\mathbb{R}}$  be the same as the one in Proposition 3.1. Consider any  $y \in \text{dom } h$ . Suppose that  $h$  is regular at  $y$ . The following assertions hold.*

(i) *Consider any  $w \in \text{dom } dh(y)$  and define  $\varphi(z) := d^2h(y)(w|z)$  for  $z \in \mathbb{R}^m$ . Then,  $\varphi$  is a proper lsc positively homogenous convex function with*

$$\varphi^*(v') = -d^2h(y)(w) + \delta_{\mathcal{A}_h(y,w)}(v') \quad \text{for } v' \in \mathbb{R}^m.$$

(ii) *The function  $h$  is parabolically regular at  $y$  for every  $v \in \partial h(y)$ .*

**Proof:** (i) Since  $h$  is regular at  $y$ , by Proposition 3.3 (iv),  $\varphi$  is a sum of the support function of the closed convex set  $\mathcal{A}_h(y, w)$  and a constant  $d^2h(y)(w)$ , so is a proper lsc positively homogenous convex function. By the definition of the conjugate function,

$$\varphi^*(v') = -d^2h(y)(w) + \delta_{\mathcal{A}_h(y,w)}(v') \quad \text{for } v' \in \mathbb{R}^m.$$

(ii) Fix any  $v \in \partial h(y)$ . Consider any  $w \in \mathbb{R}^m$  with  $dh(y)(w) = \langle v, w \rangle$ . From part (i),

$$-\varphi^*(v) = d^2h(y)(w) - \delta_{\mathcal{A}_h(y,w)}(v) \leq d^2h(y)(w) = d^2h(y|v)(w),$$

where the last equality is due to Proposition 3.2 (ii). On the other hand, by invoking [12, Proposition 13.64],  $-\varphi^*(v) = \inf_{z \in \mathbb{R}^m} \{d^2h(y)(w|z) - \langle v, z \rangle\} \geq d^2h(y|v)(w)$ . Along with the last inequality,  $\inf_{z \in \mathbb{R}^m} \{d^2h(y)(w|z) - \langle v, z \rangle\} = d^2h(y|v)(w)$ . This shows that  $h$  is parabolically regular at  $y$  for  $v$ . By the arbitrariness of  $v \in \partial h(y)$ , we get the result.  $\square$

## 4 Estimates for second subderivative of $f$

To derive tight lower and upper estimates for the second subderivative of  $f$ , we need to characterize the critical cone of  $f$  and justify its parabolic epi-differentiability. From Assumption 1, [27, Theorem 3.4] and Proposition 3.1, we have the following result.

**Lemma 4.1** *Consider any  $\bar{x} \in \text{dom } f$ . If the MSQC holds for constraint system  $F(x) \in \text{dom } \vartheta$  at  $\bar{x}$ , then  $df(\bar{x})(w) = d\vartheta(F(\bar{x}))(dF(\bar{x})(w))$  for any  $w \in \mathbb{X}$ , and hence  $df(\bar{x})$  is a proper and piecewise positively homogeneous continuous function.*

**Proposition 4.1** *Consider any  $\bar{x} \in \text{dom } f$  and any  $\bar{v} \in \partial f(\bar{x})$ . Suppose that the MSQC holds for constraint system  $F(x) \in \text{dom } \vartheta$  at  $\bar{x}$ . Then, the following assertions hold.*

(i)  *$w \in \mathcal{C}_f(\bar{x}, \bar{v})$  iff  $d\vartheta(F(\bar{x}))(dF(\bar{x})(w)) = \langle \bar{v}, w \rangle$ , which implies that  $w \in \mathcal{T}_{\text{dom } f}(\bar{x})$ .*

(ii)  *$\mathcal{C}_f(\bar{x}, \bar{v}) \subseteq \text{dom } d^2f(\bar{x}|\bar{v})$ , and the converse inclusion also holds if  $\bar{v} \in \widehat{\partial}f(\bar{x})$ .*

(iii)  *$\mathcal{C}_f(\bar{x}, \bar{v}) \neq \emptyset$  whenever  $dF(\bar{x})(0) = 0$ .*



**Proof:** (i) The equivalence is due to the definition of  $\mathcal{C}_f(\bar{x}, \bar{v})$  and Lemma 4.1. For any  $w$  such that  $d\vartheta(F(\bar{x}))(dF(\bar{x})(w)) = \langle \bar{v}, w \rangle$ ,  $dF(\bar{x})(w) \in \text{dom } d\vartheta(F(\bar{x})) = \mathcal{T}_{\text{dom}\vartheta}(F(\bar{x}))$ , where the equality is by Proposition 3.1, and hence  $w \in \mathcal{T}_{\text{dom}f}(\bar{x})$  by Lemma 2.1.

(ii) Pick any  $w \in \mathcal{C}_f(\bar{x}, \bar{v})$ . By part (i),  $w \in \mathcal{T}_{\text{dom}f}(\bar{x})$ , which by Corollary 2.3 means that  $\mathcal{T}_{\text{epi}f}^2((\bar{x}, f(\bar{x})), (w, df(\bar{x})(w))) \neq \emptyset$ . Pick any  $(u, \varpi) \in \mathcal{T}_{\text{epi}f}^2((\bar{x}, f(\bar{x})), (w, df(\bar{x})(w)))$ . Then, there exist  $\tau_k \downarrow 0$  and  $(u^k, \varpi_k) \rightarrow (u, \varpi)$  such that  $(\bar{x}, f(\bar{x})) + \tau_k(w, df(\bar{x})(w)) + \frac{1}{2}\tau_k^2(u^k, \varpi_k) \in \text{epi}f$  for each  $k \in \mathbb{N}$ . Together with  $df(\bar{x})(w) = \langle \bar{v}, w \rangle$ , for each  $k \in \mathbb{N}$ ,

$$\begin{aligned} \varpi_k &\geq \frac{2[f(\bar{x} + \tau_k w + \frac{1}{2}\tau_k^2 u^k) - f(\bar{x}) - \tau_k df(\bar{x})(w)]}{\tau_k^2} \\ &= \frac{2[f(\bar{x} + \tau_k(w + \frac{1}{2}\tau_k u^k)) - f(\bar{x}) - \tau_k \langle \bar{v}, w + \frac{1}{2}\tau_k u^k \rangle]}{\tau_k^2} + \langle \bar{v}, u^k \rangle. \end{aligned}$$

Passing the limit  $k \rightarrow \infty$  to the last inequality yields that  $\varpi \geq d^2f(\bar{x}|\bar{v})(w) + \langle \bar{v}, u \rangle$ , which means that  $w \in \text{dom } d^2f(\bar{x}|\bar{v})$  and the inclusion  $\mathcal{C}_f(\bar{x}, \bar{v}) \subseteq \text{dom } d^2f(\bar{x}|\bar{v})$ .

When  $\bar{v} \in \widehat{\partial}f(\bar{x})$ , by [12, Exercise 8.4],  $df(\bar{x})(w') \geq \langle \bar{v}, w' \rangle$  for all  $w' \in \mathbb{X}$ . On the other hand, from [12, Proposition 13.5],  $\text{dom } d^2f(\bar{x}|\bar{v}) \subseteq \{w \in \mathbb{X} \mid df(\bar{x})(w) \leq \langle \bar{v}, w \rangle\}$ . The two sides show that  $\text{dom } d^2f(\bar{x}|\bar{v}) \subseteq \{w \in \mathbb{X} \mid df(\bar{x})(w) = \langle \bar{v}, w \rangle\} = \mathcal{C}_f(\bar{x}, \bar{v})$ .

(iii) When  $dF(\bar{x})(0) = 0$ , we have  $d\vartheta(F(\bar{x}))(dF(\bar{x})(0)) = 0$  by Proposition 3.1 (ii), which along with part (i) implies that  $0 \in \mathcal{C}_f(\bar{x}, \bar{v})$  and then  $\mathcal{C}_f(\bar{x}, \bar{v}) \neq \emptyset$ .  $\square$

Proposition 4.1 (ii) extends the conclusion of [21, Theorem 4.4] for the fully subamenable function to a large class of nonconvex composite functions  $f$ .

#### 4.1 Parabolic subderivative of $f$

The following proposition characterizes the parabolic subderivative of  $f$ , which extends the conclusion of [22, Theorem 4.4] to the composition of a PWTD outer function and a parabolic semidifferentiable inner mapping.

**Proposition 4.2** *Consider any  $\bar{x} \in \text{dom}f$ . Suppose that the MSQC holds for system  $F(x) \in \text{dom}\vartheta$  at  $\bar{x}$ . Fix any  $w \in \mathcal{T}_{\text{dom}f}(\bar{x})$ . Then, the following assertions hold.*

(i) *For any  $z \in \mathbb{X}$ ,  $d^2f(\bar{x})(w|z) = d^2\vartheta(F(\bar{x}))(dF(\bar{x})(w) \mid d^2F(\bar{x})(w|z))$  with*

$$\text{dom } d^2f(\bar{x})(w|\cdot) = \mathcal{T}_{\text{dom}f}^2(\bar{x}, w); \quad (25)$$

(ii) *The function  $f$  is parabolically epi-differentiable at  $\bar{x}$  for  $w$ .*

**Proof:** (i) Since  $w \in \mathcal{T}_{\text{dom}f}(\bar{x})$ , from Lemma 2.1 and Proposition 3.1 (ii) for  $h = \vartheta$ ,  $d\vartheta(F(\bar{x}))(dF(\bar{x})(w)) < \infty$ , which along with the properness of  $d\vartheta(F(\bar{x}))$  and Lemma 4.1 implies that  $df(\bar{x})(w)$  is finite. Then, by Definition 2.7, it is easy to check that

$$\text{dom } d^2f(\bar{x})(w|\cdot) \subseteq \mathcal{T}_{\text{dom}f}^2(\bar{x}, w). \quad (26)$$

Fix any  $z \in \mathbb{X}$ . By Definition 2.7, there exist sequences  $\tau_k \downarrow 0$  and  $z^k \rightarrow z$  such that

$$d^2f(\bar{x})(w|z) = \lim_{k \rightarrow \infty} \frac{2[\vartheta(F(\bar{x} + \tau_k w + \frac{1}{2}\tau_k^2 z^k)) - \vartheta(F(\bar{x})) - \tau_k d\vartheta(F(\bar{x}))(dF(\bar{x})(w))]}{\tau_k^2},$$

which together with the parabolic semidifferentiability of  $F$  at  $\bar{x}$  implies that

$$d^2f(\bar{x})(w|z) \geq d^2\vartheta(F(\bar{x}))(dF(\bar{x})(w)|d^2F(\bar{x})(w|z)).$$

To get the converse inequality, let  $u := d^2F(\bar{x})(w|z)$ . We proceed the proof by two cases.

**Case 1:**  $z \notin \mathcal{T}_{\text{dom}f}^2(\bar{x}, w)$ . From (26),  $d^2f(\bar{x})(w|z) = \infty$ . On the other hand, by Corollary 2.2,  $u \notin \mathcal{T}_{\text{dom}\vartheta}^2(F(\bar{x}), dF(\bar{x})(w))$ , which by Proposition 3.3 (ii) for  $h = \vartheta$  implies that  $d^2\vartheta(F(\bar{x}))(dF(\bar{x})(w)|u) = \infty$ . Then,  $d^2f(\bar{x})(w|z) = d^2\vartheta(F(\bar{x}))(dF(\bar{x})(w)|d^2F(\bar{x})(w|z))$ .

**Case 2:**  $z \in \mathcal{T}_{\text{dom}f}^2(\bar{x}, w)$ . By Corollary 2.2,  $z \in \mathcal{T}_{\text{dom}f}^{i,2}(\bar{x}, w)$ . Fix any  $\tau_k \downarrow 0$ . From the definition of inner second-order tangent set, there exists  $z^k \rightarrow z$  such that for each  $k \in \mathbb{N}$ ,  $\text{dom}f \ni x^k := \bar{x} + \tau_k w + \frac{1}{2}\tau_k^2 z^k$ . For each  $k$ , let  $y^k := F(\bar{x}) + \tau_k dF(\bar{x})(w) + \frac{1}{2}\tau_k^2 u$ . Note that  $u \in \mathcal{T}_{\text{dom}\vartheta}^2(F(\bar{x}), dF(\bar{x})(w))$  by Corollary 2.2. From Proposition 3.3 (ii) for  $h = \vartheta$ ,

$$\infty > d^2\vartheta(F(\bar{x}))(dF(\bar{x})(w)|u) = \lim_{k \rightarrow \infty} \frac{2[\vartheta(y^k) - \vartheta(F(\bar{x})) - \tau_k d\vartheta(F(\bar{x}))(dF(\bar{x})(w))]}{\tau_k^2},$$

which implies that  $y^k \in \text{dom}\vartheta$  for all sufficiently large  $k$ . Consequently, we have

$$\begin{aligned} d^2f(\bar{x})(w|z) &\leq \limsup_{k \rightarrow \infty} \frac{2[\vartheta(F(x^k)) - \vartheta(F(\bar{x})) - \tau_k d\vartheta(F(\bar{x}))(dF(\bar{x})(w))]}{\tau_k^2} \\ &= \lim_{k \rightarrow \infty} \frac{\vartheta(y^k) - \vartheta(F(\bar{x})) - \tau_k d\vartheta(F(\bar{x}))(dF(\bar{x})(w))}{\frac{1}{2}\tau_k^2} + \limsup_{k \rightarrow \infty} \frac{\vartheta(F(x^k)) - \vartheta(y^k)}{\frac{1}{2}\tau_k^2} \\ &\leq d^2\vartheta(F(\bar{x}))(dF(\bar{x})(w)|u) + L_\vartheta \limsup_{k \rightarrow \infty} \frac{2\|F(x^k) - y^k\|_2}{\tau_k^2} \quad \text{for some } L_\vartheta > 0 \\ &= d^2\vartheta(F(\bar{x}))(dF(\bar{x})(w)|u), \end{aligned}$$

where the second inequality is using Assumption 1 (i), and the last equality is due to the parabolic semidifferentiability of  $F$ . The desired inequality also holds for this case.

The above arguments show that the first equality of part (i) holds. To establish equality (25), it suffices to prove that the converse inclusion in (26) holds. Pick any  $z \in \mathcal{T}_{\text{dom}f}^2(\bar{x}, w)$ . By Corollary 2.2,  $u \in \mathcal{T}_{\text{dom}\vartheta}^2(F(\bar{x}), dF(\bar{x})(w))$ . From Proposition 3.3 (ii) and the first equality,  $d^2f(\bar{x})(w|z) = d^2\vartheta(F(\bar{x}))(dF(\bar{x})(w)|u) < \infty$ , which implies that  $z \in \text{dom}d^2f(\bar{x})(w|\cdot)$ . The converse inclusion in (26) follows.

(ii) By Corollary 2.2,  $\emptyset \neq \mathcal{T}_{\text{dom}f}^2(\bar{x}, w)$ , which by (25) implies that  $d^2f(\bar{x})(w|\cdot) \not\equiv \infty$ . From Lemma 2.1,  $dF(\bar{x})(w) \in \text{dom}d\vartheta(F(\bar{x}))$ . By invoking Proposition 3.3 (iii) with  $h = \vartheta$ ,  $\vartheta$  is parabolically epi-differentiable at  $F(\bar{x})$  for  $dF(\bar{x})(w)$ . Together part (i), we conclude that  $f$  is parabolically epi-differentiable at  $\bar{x}$  for  $w$ .  $\square$

## 4.2 Lower and upper estimates for second subderivative of $f$

Now we establish the upper and lower estimates for the second subderivative of  $f$ .

**Theorem 4.1** *Consider any  $\bar{x} \in \text{dom} f$  and  $\bar{v} \in \partial f(\bar{x})$ . Suppose that the MSQC holds for system  $F(x) \in \text{dom} \vartheta$  at  $\bar{x}$ . Then, for any  $w \in \mathcal{C}_f(\bar{x}, \bar{v})$ , it holds that*

$$d^2f(\bar{x}|\bar{v})(w) \leq \inf_{z \in \mathbb{X}} \{d^2\vartheta(F(\bar{x}))(dF(\bar{x})(w)|d^2F(\bar{x})(w|z)) - \langle \bar{v}, z \rangle\} < \infty, \quad (27)$$

$$d^2f(\bar{x}|\bar{v})(w) \geq \sup_{\xi \in \Lambda_{\bar{x}, \bar{v}}} \{d^2\vartheta(F(\bar{x})|\xi)(dF(\bar{x})(w)) + \langle \xi, d^2F(\bar{x})(w) \rangle\}, \quad (28)$$

where  $\Lambda_{\bar{x}, \bar{v}} := \{\xi \in \partial\vartheta(F(\bar{x})) \mid \langle \xi, dF(\bar{x})(w') \rangle \geq \langle \bar{v}, w' \rangle \ \forall w' \in \mathbb{X}\}$ . If in addition  $\partial f(\bar{x}) \subseteq \bigcup_{\xi \in \partial\vartheta(F(\bar{x}))} \sum_{i=1}^m \xi_i \partial F_i(\bar{x})$ , then for any  $w \in \mathcal{C}_f(\bar{x}, \bar{v})$  it also holds that

$$d^2f(\bar{x}|\bar{v})(w) \geq \sup_{(\xi, u) \in \Gamma_{\bar{x}, \bar{v}}} \left\{ d^2\vartheta(F(\bar{x})|\xi)(dF(\bar{x})(w)) + \sum_{i=1}^m \xi_i d^2 F_i(\bar{x}|u_i)(w) \right\} \quad (29)$$

where  $\Gamma_{\bar{x}, \bar{v}} := \{(\xi, u_1, \dots, u_m) \in \partial\vartheta(F(\bar{x})) \times \partial F_1(\bar{x}) \times \dots \times \partial F_m(\bar{x}) \mid \bar{v} = \sum_{i=1}^m \xi_i u_i\}$ .

**Proof:** Pick any  $w \in \mathcal{C}_f(\bar{x}, \bar{v})$ . By Propositions 4.1 (i) and 4.2 (i), for any  $z \in \mathbb{X}$ ,  $d^2f(\bar{x})(w|z) = d^2\vartheta(F(\bar{x}))(dF(\bar{x})(w)|d^2F(\bar{x})(w|z))$ . Together with [12, Proposition 13.64], we obtain the first inequality in (27). From Corollary 2.2 and (25),  $\text{dom} d^2f(\bar{x})(w|\cdot) \neq \emptyset$ , so there exists  $z \in \mathbb{X}$  such that  $d^2\vartheta(F(\bar{x}))(dF(\bar{x})(w)|d^2F(\bar{x})(w|z)) < \infty$ .

In order to achieve (28), it suffices to consider that  $\Lambda_{\bar{x}, \bar{v}} \neq \emptyset$ . Pick any  $\xi \in \Lambda_{\bar{x}, \bar{v}}$ . For any sufficiently small  $\tau > 0$  and any  $w'$  sufficiently close to  $w$ , it holds that

$$\begin{aligned} \Delta_\tau^2 f(\bar{x}|\bar{v})(w') &= \frac{2[\vartheta(F(\bar{x} + \tau w')) - \vartheta(F(\bar{x})) - \tau \langle \bar{v}, w' \rangle]}{\tau^2} \\ &\geq \frac{2[\vartheta(F(\bar{x}) + \tau \Delta_\tau F(\bar{x})(w')) - \vartheta(F(\bar{x})) - \tau \langle \xi, dF(\bar{x})(w') \rangle]}{\tau^2} \\ &= \frac{2[\vartheta(F(\bar{x}) + \tau \Delta_\tau F(\bar{x})(w')) - \vartheta(F(\bar{x})) - \tau \langle \xi, \Delta_\tau F(\bar{x})(w') \rangle]}{\tau^2} \\ &\quad + \frac{2[\langle \xi, F(\bar{x} + \tau w') - F(\bar{x}) - \tau dF(\bar{x})(w') \rangle]}{\tau^2} \end{aligned}$$

where  $\Delta_\tau F(\bar{x})(\cdot)$  denotes the first-order difference quotients of  $F$  at  $\bar{x}$ . Recall that  $F$  is semidifferentiable. By [12, Theorem 7.21],  $\lim_{\tau \downarrow 0, w' \rightarrow w} \Delta_\tau F(\bar{x})(w') = dF(\bar{x})(w)$ . Together with the last equality and the definition of  $d^2F(\bar{x})(w)$ ,

$$d^2f(\bar{x}|\bar{v})(w) \geq d^2\vartheta(F(\bar{x})|\xi)(dF(\bar{x})(w)) + \langle \xi, d^2F(\bar{x})(w) \rangle.$$

This, by the arbitrariness of  $\xi$  in  $\Lambda_{\bar{x}, \bar{v}}$ , implies that inequality (28) holds.

To achieve (29), it suffices to consider that  $\Gamma_{\bar{x}, \bar{v}} \neq \emptyset$ . Pick any  $(\xi, u) \in \Gamma_{\bar{x}, \bar{v}}$ . There exist  $(\xi, u_1, \dots, u_m) \in \partial\vartheta(F(\bar{x})) \times \partial F_1(\bar{x}) \times \dots \times \partial F_m(\bar{x})$  such that  $\bar{v} = \sum_{i=1}^m \xi_i u_i$ . Then,

for any sufficiently small  $\tau > 0$  and any  $w'$  sufficiently close to  $w$ , we have

$$\begin{aligned}\Delta_\tau^2 f(\bar{x}|\bar{v})(w') &= \frac{2[\vartheta(F(\bar{x}) + \tau\Delta_\tau F(\bar{x})(w')) - \vartheta(F(\bar{x})) - \tau\sum_{i=1}^m \xi_i \langle u_i, w' \rangle]}{\tau^2} \\ &= \frac{2[\vartheta(F(\bar{x}) + \tau\Delta_\tau F(\bar{x})(w')) - \vartheta(F(\bar{x})) - \tau\langle \xi, \Delta_\tau F(\bar{x})(w') \rangle]}{\tau^2} \\ &\quad + \frac{2}{\tau^2} \left[ \sum_{i=1}^m \xi_i (F(\bar{x} + \tau w') - F(\bar{x}) - \tau \langle u_i, w' \rangle) \right].\end{aligned}$$

Recall that  $\lim_{\tau \downarrow 0, w' \rightarrow w} \Delta_\tau F(\bar{x})(w') = dF(\bar{x})(w)$ . By the definition of  $d^2 F_i(\bar{x}|u_i)(w)$ ,

$$d^2 g(\bar{x}|\bar{v})(w) \geq d^2 \vartheta(F(\bar{x})|\xi)(dF(\bar{x})(w)) + \sum_{i=1}^m \xi_i d^2 F_i(\bar{x}|u_i)(w).$$

By the arbitrariness of  $(\xi, u) \in \Gamma_{\bar{x}, \bar{v}}$ , we obtain inequality (29).  $\square$

Theorem 4.1 extends the conclusion of [22, Proposition 5.1] for the second subderivative of the composition (1) with a parabolically epi-differentiable convex outer function and a twice differentiable inner mapping. The upper estimate in (27) is essentially same as the one in [22, Equa (5.3)], but the lower estimate in (28) or (29) is different from the one in [22, Equa (5.2)] because the inner  $F$  here is allowed to be nondifferentiable.

## 5 Twice epi-differentiability for several classes of $f$

The following theorem states that  $f$  is twice epi-differentiable at  $\bar{x} \in \text{dom } f$  for  $\bar{v} \in \widehat{\partial}f(\bar{x})$  if  $f$  is parabolically regular at  $\bar{x}$  for  $\bar{v}$ . Its proof is similar to that of [22, Theorem 3.8], and we include it for completeness.

**Theorem 5.1** *Consider any  $\bar{x} \in \text{dom } f$  and  $\bar{v} \in \widehat{\partial}f(\bar{x})$ . Suppose that the MSQC holds for system  $F(x) \in \text{dom } \vartheta$  at  $\bar{x}$ , and that  $\partial f(\bar{x}) = \widehat{\partial}f(\bar{x})$ . If  $f$  is parabolically regular at  $\bar{x}$  for  $\bar{v}$ , then it is twice epi-differentiable at  $\bar{x}$  for  $\bar{v}$  with*

$$d^2 f(\bar{x}|\bar{v})(w) = \begin{cases} \min_{z \in \mathbb{X}} \{d^2 f(\bar{x})(w|z) - \langle \bar{v}, z \rangle\} & \text{if } w \in \mathcal{C}_f(\bar{x}, \bar{v}), \\ \infty & \text{otherwise.} \end{cases}$$

**Proof:** Pick any  $w \in \mathbb{X}$ . We proceed the arguments by the following two cases.

**Case 1:**  $w \in \text{dom } d^2 f(\bar{x}|\bar{v})$ . From  $\partial f(\bar{x}) = \widehat{\partial}f(\bar{x})$  and Proposition 4.1 (ii),  $w \in \mathcal{C}_f(\bar{x}, \bar{v})$ . Pick an arbitrary sequence  $\tau_k \downarrow 0$ . Since  $f$  is parabolically regular at  $\bar{x}$  for  $\bar{v}$ , from the proof of [22, Proposition 3.6], there exists  $\bar{z} \in \text{dom } d^2 f(\bar{x})(w|\cdot)$  such that  $d^2 f(\bar{x}|\bar{v})(w) = d^2 f(\bar{x})(w|\bar{z}) - \langle \bar{v}, \bar{z} \rangle$ . By Proposition 4.2 (ii),  $f$  is parabolically epi-differentiable at  $\bar{x}$  for  $w$ , and we find a sequence  $z^k \rightarrow \bar{z}$  such that

$$\begin{aligned}d^2 f(\bar{x})(w|\bar{z}) &= \lim_{k \rightarrow \infty} \frac{2[f(\bar{x} + \tau_k w + \frac{1}{2}\tau_k^2 z^k) - f(\bar{x}) - \tau_k df(\bar{x})(w)]}{\tau_k^2} \\ &= \lim_{k \rightarrow \infty} \Delta_{\tau_k}^2 f(\bar{x}|\bar{v})(w^k) + \langle \bar{v}, z^k \rangle \quad \text{with } w^k = w + \frac{1}{2}\tau_k z^k.\end{aligned}$$

Together with  $d^2f(\bar{x})(w|\bar{z}) = d^2f(\bar{x}|\bar{v})(w) + \langle \bar{v}, \bar{z} \rangle$  and  $z^k \rightarrow \bar{z}$ , we conclude that the sequence  $w^k \rightarrow w$  is such that  $d^2f(\bar{x}|\bar{v})(w) = \lim_{k \rightarrow \infty} \Delta_{\tau_k}^2 f(\bar{x}|\bar{v})(w^k)$ .

**Case 2:**  $w \notin \text{dom } d^2f(\bar{x}|\bar{v})$ . In this case,  $d^2f(\bar{x}|\bar{v})(w) = \infty$ . Pick an arbitrary sequence  $\tau_k \downarrow 0$  and set  $w^k = w$  for all  $k \in \mathbb{N}$ . By Definition 2.5, it is immediate to have

$$\infty = d^2f(\bar{x}|\bar{v})(w) \leq \liminf_{k \rightarrow \infty} \Delta_{\tau_k}^2 f(\bar{x}|\bar{v})(w^k) \leq \limsup_{k \rightarrow \infty} \Delta_{\tau_k}^2 f(\bar{x}|\bar{v})(w^k) \leq \infty = d^2f(\bar{x}|\bar{v})(w),$$

which shows that the sequence  $w^k$  is such that  $d^2f(\bar{x}|\bar{v})(w) = \lim_{k \rightarrow \infty} \Delta_{\tau_k}^2 f(\bar{x}|\bar{v})(w^k)$ .

The above arguments show that  $f$  is twice epi-differentiable at  $\bar{x}$  for  $\bar{v}$ .  $\square$

Inspired by Theorem 5.1, we employ the lower and upper estimates in Theorem 4.1 to demonstrate the parabolic regularity for the following several classes of  $f$ :

- (I)  $F$  is twice differentiable on  $\mathcal{O}$  and  $\vartheta$  satisfies Assumption 1 (i).
- (II)  $F(x) = (\|x_{J_1}\|_q, \dots, \|x_{J_m}\|_q)$  ( $q > 1$ ) for  $x \in \mathbb{R}^n$ , where  $\{J_1, \dots, J_m\}$  is a partition of  $[n]$ , and  $\vartheta(z) = \sum_{i=1}^m \rho_\lambda(z_i)$  ( $\lambda > 0$ ) where  $\rho_\lambda: \mathbb{R} \rightarrow \overline{\mathbb{R}}$  satisfies conditions
  - (II.1)  $\rho_\lambda$  is a PWT function satisfying  $\rho_\lambda(0) = 0$ ;
  - (II.2)  $\rho_\lambda$  is differentiable at all  $t \neq 0$  and  $\partial\rho_\lambda(0) = \gamma[-\lambda, \lambda]$  for some  $\gamma > 0$ ;
  - (II.3)  $\rho_\lambda$  is regular and strictly continuous relative to its domain.

The function  $\vartheta$  induced by  $\rho_\lambda$  satisfying conditions (II.1)-(II.3) often appears in sparsity optimization; for example, the SCAD function [7] and MCP function [8]:

$$\rho_\lambda(t) := \begin{cases} \lambda|t| & \text{if } |t| \leq \lambda, \\ \frac{-t^2 + 2a\lambda|t| - \lambda^2}{2(a-1)} & \text{if } \lambda < |t| \leq a\lambda, \quad \text{with } a > 2; \\ \frac{(a+1)}{2}\lambda^2 & \text{if } |t| > a\lambda \end{cases} \quad (30)$$

$$\rho_\lambda(t) := \lambda \text{sign}(t) \int_0^{|t|} \left(1 - \frac{\omega}{\lambda b}\right)_+ d\omega \quad \text{with } b > 0.$$

- (III)  $F(x) = \|x_2\|_q - x_1$  with  $q > 1$  for  $x = (x_1, x_2) \in \mathbb{R} \times \mathbb{R}^{n-1}$ , and  $\vartheta(t) = \delta_{\mathbb{R}_-}(t)$  for  $t \in \mathbb{R}$ . Now  $f = \delta_K$  with  $K := \{(x_1, x_2) \in \mathbb{R} \times \mathbb{R}^{n-1} \mid \|x_2\|_q \leq x_1\}$ .
- (IV)  $F(x) = \lambda_1(x)$  for  $x \in \mathbb{S}^n$ , the space consisting of all  $n \times n$  symmetric matrices, and  $\vartheta(t) = \delta_{\mathbb{R}_-}(t)$  for  $t \in \mathbb{R}$ . Now  $f$  is the indicator of negative semidefinite cone  $\mathbb{S}_-^n$ .

## 5.1 Parabolic regularity for $f$ of type I

Because  $F$  is twice differentiable on the set  $\mathcal{O}$ , for any  $\bar{x} \in \text{dom } f$ ,  $dF(\bar{x})(w) = F'(\bar{x})w$  for all  $w \in \mathbb{R}^n$ . In this case, the result of Theorem 4.1 can be sharpened as follows.

**Proposition 5.1** Consider any  $\bar{x} \in \text{dom}f$  and  $\bar{v} \in \partial f(\bar{x})$ . Suppose that the MSQC holds for system  $F(x) \in \text{dom}\vartheta$  at  $\bar{x}$ , and that  $\vartheta$  is regular at  $F(\bar{x})$ . Then, for every  $w \in \mathcal{C}_f(\bar{x}, \bar{v})$ ,

$$\begin{aligned} d^2f(\bar{x}|\bar{v})(w) &= \inf_{z \in \mathbb{X}} \left\{ d^2\vartheta(F(\bar{x}))(F'(\bar{x})w | D^2F(\bar{x})(w, w) + F'(\bar{x})z) - \langle \bar{v}, z \rangle \right\}, \\ &= d^2\vartheta(F(\bar{x}))(F'(\bar{x})(w)) + \sup_{\xi \in \Lambda_{\bar{x}, \bar{v}}} \langle \xi, D^2F(\bar{x})(w, w) \rangle \end{aligned}$$

where  $\Lambda_{\bar{x}, \bar{v}} = \{ \xi \in \partial\vartheta(F(\bar{x})) \mid \nabla F(\bar{x})\xi = \bar{v} \}$ , and hence  $f$  is parabolic regular at  $\bar{x}$  for  $\bar{v}$ .

**Proof:** Fix any  $w \in \mathcal{C}_f(\bar{x}, \bar{v})$ . Since  $dF(\bar{x})(w) = F'(\bar{x})w$ , the set  $\Lambda_{\bar{x}, \bar{v}}$  defined in Theorem 4.1 takes the form as in Proposition 5.1. For every  $\xi \in \Lambda_{\bar{x}, \bar{v}}$ , from  $w \in \mathcal{C}_f(\bar{x}, \bar{v})$  and Proposition 4.1 (i),  $F'(\bar{x})w \in \mathcal{C}_\vartheta(F(\bar{x}), \xi)$ . By inequality (28) and Proposition 3.2 (ii),

$$d^2f(\bar{x}|\bar{v})(w) \geq d^2\vartheta(F(\bar{x}))(F'(\bar{x})(w)) + \sup_{\xi \in \Lambda_{\bar{x}, \bar{v}}} \langle \xi, D^2F(\bar{x})(w, w) \rangle. \quad (31)$$

For the convenience of discussion, we introduce the following optimal value function

$$\Upsilon(p) := \inf_{z \in \mathbb{X}} \sup_{u \in \mathcal{A}(F(\bar{x}), F'(\bar{x})w)} \left\{ \langle u, D^2F(\bar{x})(w, w) + p \rangle + \langle \nabla F(\bar{x})u - \bar{v}, z \rangle \right\} \quad (32)$$

with  $\mathcal{A}(F(\bar{x}), F'(\bar{x})w) := \{ u \in \partial\vartheta(F(\bar{x})) \mid d\vartheta(F(\bar{x}))(F'(\bar{x})w) = \langle \nabla F(\bar{x})u, w \rangle \}$ . Note that  $d^2F(\bar{x})(w|z) = D^2F(\bar{x})(w, w) + F'(\bar{x})z$  for any  $z \in \mathbb{R}^n$ . By (27) and Proposition 3.3 (iv),

$$\begin{aligned} d^2f(\bar{x}|\bar{v})(w) &\leq \inf_{z \in \mathbb{X}} \left\{ d^2\vartheta(F(\bar{x}))(F'(\bar{x})w | D^2F(\bar{x})(w, w) + F'(\bar{x})z) - \langle \bar{v}, z \rangle \right\} \\ &= \inf_{z \in \mathbb{X}} \sup_{u \in \mathcal{A}(F(\bar{x}), F'(\bar{x})w)} \left\{ \langle u, D^2F(\bar{x})(w, w) + F'(\bar{x})z \rangle - \langle \bar{v}, z \rangle \right\} \\ &\quad + d^2\vartheta(F(\bar{x}))(F'(\bar{x})w) \\ &= \Upsilon(0) + d^2\vartheta(F(\bar{x}))(F'(\bar{x})w). \end{aligned}$$

Together with (31), to achieve the desired two equalities, it suffices to argue that

$$\inf_{z \in \mathbb{X}} \sup_{u \in \mathcal{A}(F(\bar{x}), F'(\bar{x})w)} \left\{ \langle u, D^2F(\bar{x})(w, w) \rangle + \langle \nabla F(\bar{x})u - \bar{v}, z \rangle \right\} = \sup_{\xi \in \Lambda_{\bar{x}, \bar{v}}} \langle \xi, D^2F(\bar{x})(w, w) \rangle. \quad (33)$$

Note that  $\mathcal{A}(F(\bar{x}), F'(\bar{x})w)$  is a closed convex set because  $\partial\vartheta(F(\bar{x})) = \widehat{\partial}\vartheta(F(\bar{x}))$ . Hence, the minimization problem on the left hand side of (33) is a convex program whose dual, after an elementary calculation, is precisely the supremum problem on the right hand side of (33). By [33, Theorem 2.142], we only need to show that  $\partial\Upsilon(0) \neq \emptyset$ . By the definition of  $\Upsilon$  and the second inequality in (27),  $\Upsilon(0) < \infty$ . Together with [26, Proposition 5.1], it suffices to argue that there exists a constant  $c_1 > 0$  such that for all  $p \in \mathbb{R}^m$ ,

$$\Upsilon(p) \geq \Upsilon(0) - c_1 \|p\|_2.$$

Pick any  $(z, p) \in \mathbb{X} \times \mathbb{R}^m$  with  $u_p := D^2F(\bar{x})(w, w) + p + F'(\bar{x})z \in \text{dom}d^2\vartheta(F(\bar{x}))(F'(\bar{x})w|\cdot)$ . From (25),  $u_p \in \mathcal{T}_{\text{dom}\vartheta}^2(F(\bar{x}), F'(\bar{x})w)$ . Let  $\mathcal{S}_w$  be the multifunction defined in (11) with

the functions  $\varphi$  and  $g$  replaced by  $\vartheta$  and  $F$ , respectively. Obviously,  $z \in \mathcal{S}_w(p)$ . By invoking Lemma 2.2, there exist  $z^0 \in \mathcal{S}_w(0)$  and a vector  $b \in \mathbb{B}$  such that

$$z = z^0 + \kappa \|p\|_2 b \quad \text{and} \quad D^2F(\bar{x})(w, w) + F'(\bar{x})z^0 \in \text{dom } d^2\vartheta(F(\bar{x}))(F'(\bar{x})w | \cdot)$$

Note that  $d^2\vartheta(F(\bar{x}))(F'(\bar{x})w | \cdot)$  is proper by Proposition 3.3 (iii). By using Assumption 1 (i) and following the same arguments as those for the last part of [22, Proposition 4.5],  $d^2\vartheta(F(\bar{x}))(F'(\bar{x})w | \cdot)$  is Lipschitz continuous relative to its domain, and we denote by  $L$  the corresponding Lipschitz constant. Then, it holds that

$$\begin{aligned} & d^2\vartheta(F(\bar{x}))(F'(\bar{x})w | u_p) - \langle \bar{v}, z \rangle - d^2\vartheta(F(\bar{x}))(F'(\bar{x})w) \\ & \geq d^2\vartheta(F(\bar{x}))(F'(\bar{x})w | D^2F(\bar{x})(w, w) + F'(\bar{x})z^0) - d^2\vartheta(F(\bar{x}))(F'(\bar{x})w) \\ & \quad - \langle \bar{v}, z^0 \rangle - L(\|p\|_2 + \|F'(\bar{x})\| \|z - z^0\|) - \|\bar{v}\| \|z - z^0\| \\ & \geq \Upsilon(0) - L(\|p\|_2 + \kappa \|F'(\bar{x})\| \|p\|_2) - \kappa \|\bar{v}\| \|p\|_2, \end{aligned}$$

which implies that  $\Upsilon(p) \geq \Upsilon(0) - [\kappa \|\bar{v}\| + L(1 + \kappa \|F'(\bar{x})\|)] \|p\|_2$ . Hence,  $\partial\Upsilon(0) \neq \emptyset$ . The parabolic regularity of  $f$  holds by combining the first equality and Proposition 4.2 (i).  $\square$

**Remark 5.1** *By combining Proposition 2.1 (ii) with [15, Page 219], under the MSQC for system  $F(x) \in \text{dom } \vartheta$  at  $\bar{x}$ ,  $\partial f(\bar{x}) \subseteq \nabla F(\bar{x}) \partial \vartheta(F(\bar{x}))$ . Along with  $\widehat{\partial} f(\bar{x}) \supseteq \nabla F(\bar{x}) \widehat{\partial} \vartheta(F(\bar{x}))$ , the regularity of  $\vartheta$  at  $F(\bar{x})$  implies  $\widehat{\partial} f(\bar{x}) = \partial f(\bar{x})$ . Note that  $d\vartheta(F(\bar{x}))(0) = 0$ . By Proposition 4.1 (iii),  $\mathcal{C}_f(\bar{x}, \bar{v}) \neq \emptyset$ . Along with Proposition 5.1 and Theorem 5.1,  $f$  is properly twice epi-differentiable at  $\bar{x}$  for  $\bar{v}$  under the conditions of Proposition 5.1.*

By Remark 5.1 and [12, Theorem 13.24], we obtain the following second-order necessary and sufficient optimality conditions for (2) with  $\vartheta$  and  $F$  from type I, which extends the result of [21, Theorem 6.1] for the outer PWLQ convex  $\vartheta$  to nonconvex PWTQ  $\vartheta$ .

**Corollary 5.1** *Consider problem (2) with  $\vartheta$  and  $F$  from type (I). Consider any  $\bar{x} \in \text{dom } f$ . Suppose that the MSQC holds for system  $F(x) \in \text{dom } \vartheta$  at  $\bar{x}$ , and that  $\vartheta$  is regular at  $F(\bar{x})$ . Let  $\bar{v} := -\nabla f_0(\bar{x})$ . Then the following statements holds.*

(i) *If  $\bar{x}$  is a local optimal solution of (2), then  $\bar{v} \in \partial f(\bar{x})$  and for any  $w \in \mathcal{C}_f(\bar{x}, \bar{v})$ ,*

$$\langle w, \nabla^2 f_0(\bar{x}) w \rangle + d^2\vartheta(F(\bar{x}))(F'(\bar{x})w) + \sup_{\xi \in \Lambda_{\bar{x}, \bar{v}}} \langle \xi, D^2F(\bar{x})(w, w) \rangle \geq 0, \quad (34)$$

*where  $\Lambda_{\bar{x}, \bar{v}}$  is the same as the one in Proposition 5.1.*

(ii) *If  $\bar{x} \in (\partial f)^{-1}(\bar{v})$  and (34) holds with “ $\geq$ ” for all  $w \in \mathcal{C}_f(\bar{x}, \bar{v}) \setminus \{0\}$ , then it is a strong local optimal solution of problem (2), i.e., there exist  $c > 0$  and  $\varepsilon > 0$  such that*

$$\Phi(x) \geq \Phi(\bar{x}) + (c/2) \|x - \bar{x}\|^2 \quad \text{for all } x \in \mathbb{B}(\bar{x}, \varepsilon).$$

## 5.2 Parabolic regularity for $f$ of type II

In this section, we let  $p = 1 - 1/q$ , for each  $i \in [m]$ , write  $F_i(z) := \|z\|_q$  for  $z \in \mathbb{R}^{J_i}$ , and for every  $x \in \mathbb{R}^n$ , define  $\text{gs}(x) := \{i \in [m] \mid x_{J_i} \neq 0\}$  and  $\overline{\text{gs}}(x) := [m] \setminus \text{gs}(x)$ . Obviously, at any  $x \in \mathbb{R}^n$ , every  $F_i$  for  $i \in \text{gs}(x)$  is at least twice continuously differentiable at  $x_{J_i}$  with

$$\nabla F_i(x_{J_i}) = \text{sign}(x_{J_i}) \circ |x_{J_i}|^{q-1} \|x_{J_i}\|_q^{1-q}.$$

The mapping  $F$  is globally Lipschitz continuous and directionally differentiable on  $\mathbb{R}^n$ , and for any  $x, w \in \mathbb{R}^n$ ,  $dF(x)(w) = (dF_1(x_{J_1})(w_{J_1}), \dots, dF_m(x_{J_m})(w_{J_m}))^\top$  with

$$dF_i(x_{J_i})(w_{J_i}) = \begin{cases} \langle \nabla F_i(x_{J_i}), w_{J_i} \rangle & \text{if } i \in \text{gs}(x), \\ \|w_{J_i}\|_q & \text{otherwise.} \end{cases} \quad (35)$$

It is easy to check that  $F$  is parabolically semidifferentiable, and at any  $x, w \in \mathbb{R}^n$ ,  $d^2F(x)(w|z) = (d^2F_1(x_{J_1})(w_{J_1}|z_{J_1}), \dots, d^2F_m(x_{J_m})(w_{J_m}|z_{J_m}))^\top$  for  $z \in \mathbb{R}^n$  with

$$d^2F_i(x_{J_i})(w_{J_i}|z_{J_i}) = \begin{cases} \|z_{J_i}\|_q & \text{if } i \in \overline{\text{gs}}(x) \cap \overline{\text{gs}}(w), \\ \langle \nabla F_i(w_{J_i}), z_{J_i} \rangle & \text{if } i \in \overline{\text{gs}}(x) \cap \text{gs}(w), \\ \langle w_{J_i}, \nabla^2 F_i(x_{J_i}) w_{J_i} \rangle + \langle \nabla F_i(x_{J_i}), z_{J_i} \rangle & \text{if } i \in \text{gs}(x). \end{cases} \quad (36)$$

By conditions (II.1)-(II.3), it is easy to obtain the following properties of  $\vartheta$ , where conditions (II.2)-(II.3) imply that  $d\rho_\lambda(0)(\omega) = \gamma\lambda|\omega|$  for any  $\omega \in \mathbb{R}$  by [12, Theorem 8.30].

**Lemma 5.1** *Consider any  $y \in \mathbb{R}^m$ , the following statements hold for  $\vartheta$  in type II.*

(i) *For any  $w \in \mathbb{R}^m$ ,  $d\vartheta(y)(w) = \sum_{i \in \text{supp}(y)} \rho'_\lambda(y_i) w_i + \gamma\lambda \sum_{i \notin \text{supp}(y)} |w_i|$ .*

(ii)  *$\vartheta$  is regular at  $y$  with  $\emptyset \neq \widehat{\partial}\vartheta(y) = \partial\vartheta(y) = \partial\rho_\lambda(y_1) \times \dots \times \partial\rho_\lambda(y_m)$ , where*

$$\partial\rho_\lambda(y_i) = \begin{cases} [-\gamma\lambda, \gamma\lambda] & \text{if } y_i = 0, \\ \{\rho'_\lambda(y_i)\} & \text{if } y_i \neq 0 \end{cases} \quad \text{for each } i \in [m].$$

By Lemma 5.1 (ii), we have the following conclusion on the subdifferential of  $f$ .

**Lemma 5.2** *Consider any  $\bar{x} \in \mathbb{R}^n$ . Then, for every  $v \in \partial f(\bar{x})$ ,  $v_i = \rho'_\lambda(F_i(\bar{x}_{J_i})) \nabla F_i(\bar{x}_{J_i})$  for  $i \in \text{gs}(\bar{x})$ ; and for  $i \in \overline{\text{gs}}(\bar{x})$ , there exist  $\eta_i \in [-\gamma\lambda, \gamma\lambda]$  and  $\zeta_i \in \mathbb{R}^{|J_i|}$  such that  $v_i = \eta_i \zeta_i$ , where  $\|\zeta_i\|_p \leq 1$  if  $\eta_i \geq 0$ , otherwise  $\|\zeta_i\|_p = 1$ .*

**Proof:** Since  $v \in \partial f(\bar{x})$ , from the expression of  $F$  and [12, Proposition 10.5 & Theorem 10.49], there exist  $\eta \in \partial\vartheta(F(\bar{x}))$  such that  $v_i \in \partial(\eta_i F_i)(\bar{x}_{J_i})$  for each  $i \in [m]$ . By Lemma 5.1 (ii), for  $i \in \text{gs}(\bar{x})$ ,  $\eta_i = \rho'_\lambda(F_i(\bar{x}_{J_i}))$ , so  $v_i = \rho'_\lambda(F_i(\bar{x}_{J_i})) \nabla F_i(\bar{x}_{J_i})$ ; and for  $i \in \overline{\text{gs}}(\bar{x})$ ,  $\eta_i \in [-\gamma\lambda, \gamma\lambda]$ . Fix any  $i \in \overline{\text{gs}}(\bar{x})$ . If  $\eta_i \geq 0$ , we have  $\partial(\eta_i F_i)(\bar{x}_{J_i}) = \eta_i \partial F_i(\bar{x}_{J_i}) = \eta_i \{d \in \mathbb{R}^{|J_i|} \mid \|d\|_p \leq 1\}$ ; and if  $\eta_i < 0$ , by [12, Corollary 9.21],  $\partial(\eta_i F_i)(\bar{x}_{J_i}) = \partial(-|\eta_i| F_i)(\bar{x}_{J_i}) = \eta_i \{d \in \mathbb{R}^{|J_i|} \mid \|d\|_p = 1\}$ . This shows that the desired result holds.  $\square$

Now we are ready to show that the function  $f$  of type II is parabolically regular.



**Proposition 5.2** Consider any  $\bar{x} \in \mathbb{R}^n$  and  $\bar{v} \in \partial f(\bar{x})$ . Then, for every  $w \in \mathcal{C}_f(\bar{x}, \bar{v})$ ,

$$\begin{aligned} d^2f(\bar{x}|\bar{v})(w) &= \min_{z \in \mathbb{R}^n} \{d^2\vartheta(F(\bar{x}))(dF(\bar{x})(w) | d^2F(\bar{x})(w|z)) - \langle \bar{v}, z \rangle\} \\ &= d^2f(F(\bar{x}))(dF(\bar{x})(w)) + \langle \bar{\xi}, d^2F(\bar{x})(w) \rangle, \end{aligned} \quad (37)$$

where  $\bar{\xi} \in \mathbb{R}^m$  with  $\bar{\xi}_i = \rho'_\lambda(F_i(\bar{x}_{J_i}))$  for  $i \in \text{gs}(\bar{x})$  and  $\bar{\xi}_i = \gamma\lambda$  for  $i \in \overline{\text{gs}}(\bar{x})$ , and hence  $f$  is both parabolically regular and properly twice epi-differentiable at  $\bar{x}$  for  $\bar{v}$ .

**Proof:** Note that  $\text{dom } \vartheta = \mathbb{R}^m$ . The MSQC holds for system  $F(x) \in \text{dom } \vartheta$  at  $\bar{x}$ . We first argue that  $\bar{\xi} \in \Lambda_{\bar{x}, \bar{v}}$ . Clearly,  $\bar{\xi} \in \partial\vartheta(F(\bar{x})) = \widehat{\partial}\vartheta(F(\bar{x}))$  by Lemma 5.1 (ii). From  $\bar{v} \in \partial f(\bar{x})$  and Lemma 5.2,  $\bar{v}_i = \rho'_\lambda(F_i(\bar{x}_{J_i}))\nabla F_i(\bar{x}_{J_i})$  for each  $i \in \text{gs}(\bar{x})$ , and for each  $i \in \overline{\text{gs}}(\bar{x})$ , there exist  $\bar{\eta}_i \in [-\gamma\lambda, \gamma\lambda]$  and  $\bar{\zeta}_i \in \mathbb{R}^{|J_i|}$  with  $\|\bar{\zeta}_i\|_p \leq 1$  when  $\bar{\eta}_i \geq 0$  and  $\|\bar{\zeta}_i\|_p = 1$  when  $\bar{\eta}_i < 0$  such that  $\bar{v}_i = \bar{\eta}_i\bar{\zeta}_i$ . Consequently,

$$\langle \bar{v}, w' \rangle = \sum_{i \in \text{gs}(\bar{x})} \rho'_\lambda(F_i(\bar{x}_{J_i}))\langle \nabla F_i(\bar{x}_{J_i}), w'_{J_i} \rangle + \sum_{i \in \overline{\text{gs}}(\bar{x})} \bar{\eta}_i \langle \bar{\zeta}_i, w'_{J_i} \rangle \quad \forall w' \in \mathbb{R}^n. \quad (38)$$

On the other hand, from equation (35) it follows that for any  $w' \in \mathbb{R}^n$ ,

$$\langle \bar{\xi}, dF(\bar{x})(w') \rangle = \sum_{i \in \text{gs}(\bar{x})} \rho'_\lambda(F_i(\bar{x}_{J_i}))\langle \nabla F_i(\bar{x}_{J_i}), w'_{J_i} \rangle + \gamma\lambda \sum_{i \in \overline{\text{gs}}(\bar{x})} \|w'_{J_i}\|_q.$$

The last two equalities imply that  $\langle \bar{\xi}, dF(\bar{x})(w') \rangle \geq \langle \bar{v}, w' \rangle$  for all  $w' \in \mathbb{R}^n$ , so  $\bar{\xi} \in \Lambda_{\bar{x}, \bar{v}}$ . In particular, from  $\bar{\xi} \in \Lambda_{\bar{x}, \bar{v}}$  and  $\partial\vartheta(F(\bar{x})) = \widehat{\partial}\vartheta(F(\bar{x}))$ , it follows that for any  $w' \in \mathbb{R}^m$ ,

$$\langle \bar{v}, w' \rangle \leq \langle \bar{\xi}, dF(\bar{x})(w') \rangle \leq d\vartheta(F(\bar{x}))(dF(\bar{x})(w')),$$

which by Proposition 4.1 (i) implies that if  $w' \in \mathcal{C}_f(\bar{x}, \bar{v})$ , then  $dF(\bar{x})(w') \in \mathcal{C}_\vartheta(F(\bar{x}), \bar{\xi})$ .

Fix any  $w \in \mathcal{C}_f(\bar{x}, \bar{v})$ . From Proposition 4.1 (i) and Lemma 5.1 (i), it follows that

$$\langle \bar{v}, w \rangle = d\vartheta(F(\bar{x}))(dF(\bar{x})(w)) = \sum_{i \in \text{gs}(\bar{x})} \rho'_\lambda(F_i(\bar{x}_{J_i}))\langle \nabla F_i(\bar{x}_{J_i}), w_{J_i} \rangle + \gamma\lambda \sum_{i \in \overline{\text{gs}}(\bar{x})} \|w_{J_i}\|_q \quad (39)$$

which together with the above (38) implies that the following equality holds

$$\sum_{i \in \overline{\text{gs}}(\bar{x})} \gamma\lambda \|w_{J_i}\|_q = \sum_{i \in \overline{\text{gs}}(\bar{x})} \bar{\eta}_i \langle \bar{\zeta}_i, w_{J_i} \rangle \quad \text{or} \quad \sum_{i \in \overline{\text{gs}}(\bar{x}) \cap \text{gs}(w)} \gamma\lambda \|w_{J_i}\|_q = \sum_{i \in \overline{\text{gs}}(\bar{x}) \cap \text{gs}(w)} \bar{\eta}_i \langle \bar{\zeta}_i, w_{J_i} \rangle.$$

Because  $\langle u, \xi \rangle \leq \|u\|_p \|\xi\|_q$  for any  $u, \xi$  of the same dimension and the equality holds iff  $u = \text{sign}(\xi) \circ |\xi|^{q-1} \|\xi\|_q^{1-q}$ , from the last equality we deduce that for each  $i \in \overline{\text{gs}}(\bar{x}) \cap \text{gs}(w)$ ,

$$\text{sign}(\bar{\eta}_i)\bar{\zeta}_i = \text{sign}(w_{J_i}) \circ |w_{J_i}|^{q-1} \|w_{J_i}\|_q^{1-q} = \nabla F_i(w_{J_i}) \quad \text{and} \quad |\eta_i| = \gamma\lambda. \quad (40)$$

From inequality (28),  $\bar{\xi} \in \Lambda_{\bar{x}, \bar{v}}$ , Proposition 3.2 (ii) and  $dF(\bar{x})(w) \in \mathcal{C}_\vartheta(F(\bar{x}), \bar{\xi})$ , we have

$$\begin{aligned} d^2f(\bar{x}|\bar{v})(w) &\geq d^2\vartheta(F(\bar{x}) | \bar{\xi})(dF(\bar{x})(w)) + \langle \bar{\xi}, d^2F(\bar{x})(w) \rangle \\ &= d^2\vartheta(F(\bar{x}))(dF(\bar{x})(w)) + \langle \bar{\xi}, d^2F(\bar{x})(w) \rangle. \end{aligned}$$

On the other hand, from inequality (27) and Proposition 3.3 (iv), it follows that

$$\begin{aligned} d^2f(\bar{x}|\bar{v})(w) &\leq \inf_{z \in \mathbb{R}^n} \{d^2\vartheta(F(\bar{x}))(dF(\bar{x})(w)|d^2F(\bar{x})(w|z)) - \langle \bar{v}, z \rangle\} \\ &= d^2\vartheta(F(\bar{x}))(dF(\bar{x})(w)) + \inf_{z \in \mathbb{R}^n} \left\{ \sup_{u \in \mathcal{A}(F(\bar{x}), dF(\bar{x})(w))} \langle u, d^2F(\bar{x})(w|z) \rangle - \langle \bar{v}, z \rangle \right\} \end{aligned}$$

where  $\mathcal{A}(F(\bar{x}), dF(\bar{x})(w)) = \{u \in \partial\vartheta(F(\bar{x})) \mid d\vartheta(F(\bar{x}))(dF(\bar{x})(w)) = \langle u, dF(\bar{x})(w) \rangle\}$ . The above two equations show that to get the equalities in (37), it suffices to argue that

$$\Gamma_0 := \inf_{z \in \mathbb{R}^n} \left\{ \sup_{u \in \mathcal{A}(F(\bar{x}), dF(\bar{x})(w))} \langle u, d^2F(\bar{x})(w|z) \rangle - \langle \bar{v}, z \rangle \right\} = \langle \bar{\xi}, d^2F(\bar{x})(w) \rangle. \quad (41)$$

To establish equality (41), we first claim that  $u \in \mathcal{A}(F(\bar{x}), dF(\bar{x})(w))$  if and only if

$$u_i \in \begin{cases} \{\rho'_\lambda(F_i(\bar{x}_{J_i}))\} & \text{if } i \in \text{gs}(\bar{x}); \\ \{\gamma\lambda\} & \text{if } i \in \overline{\text{gs}}(\bar{x}) \cap \text{gs}(w); \\ [-\gamma\lambda, \gamma\lambda] & \text{if } i \in \overline{\text{gs}}(\bar{x}) \cap \overline{\text{gs}}(w) \end{cases} \quad \text{for each } i \in [m]. \quad (42)$$

Indeed, from the expression of  $\mathcal{A}(F(\bar{x}), dF(\bar{x})(w))$ , Lemma 5.1 (i), equations (35) and (39),  $u \in \mathcal{A}(F(\bar{x}), dF(\bar{x})(w))$  if and only if  $u \in \partial\vartheta(F(\bar{x}))$  and

$$\begin{aligned} \sum_{i \in \text{gs}(\bar{x})} u_i \langle \nabla F_i(\bar{x}_{J_i}), w_{J_i} \rangle + \sum_{i \in \overline{\text{gs}}(\bar{x})} u_i \|w_{J_i}\|_q &= \langle u, dF(\bar{x})(w) \rangle = d\vartheta(F(\bar{x}))(dF(\bar{x})(w)) \\ &= \sum_{i \in \text{gs}(\bar{x})} \rho'_\lambda(F_i(\bar{x}_{J_i})) \langle \nabla F_i(\bar{x}_{J_i}), w_{J_i} \rangle + \gamma\lambda \sum_{i \in \overline{\text{gs}}(\bar{x})} \|w_{J_i}\|_q. \end{aligned}$$

Then, by invoking Lemma 5.1 (ii), it is easy to obtain the claimed relation in (42). For every  $z \in \mathbb{R}^n$ , from equations (38) with  $w' = z$  and (40), it follows that

$$\begin{aligned} \langle \bar{v}, z \rangle &= \sum_{i \in \text{gs}(\bar{x})} \rho'_\lambda(F_i(\bar{x}_{J_i})) [\langle \nabla F_i(\bar{x}_{J_i}), z_{J_i} \rangle + \langle w_{J_i}, \nabla^2 F_i(\bar{x}_{J_i}) w_{J_i} \rangle] + \sum_{i \in \overline{\text{gs}}(\bar{x}) \cap \overline{\text{gs}}(w)} \bar{\eta}_i \langle \bar{\zeta}_i, z_{J_i} \rangle \\ &\quad + \gamma\lambda \sum_{i \in \overline{\text{gs}}(\bar{x}) \cap \text{gs}(w)} \langle \nabla F_i(w_{J_i}), z_{J_i} \rangle - \sum_{i \in \text{gs}(\bar{x})} \rho'_\lambda(F_i(\bar{x}_{J_i})) \langle w_{J_i}, \nabla^2 F_i(\bar{x}_{J_i}) w_{J_i} \rangle. \end{aligned}$$

Let  $\Xi(z)$  denote the sum of the first three terms on the right hand side. For the last term, we have  $\sum_{i \in \text{gs}(\bar{x})} \rho'_\lambda(F_i(\bar{x}_{J_i})) \langle w_{J_i}, \nabla^2 F_i(\bar{x}_{J_i}) w_{J_i} \rangle = \langle \bar{\xi}, d^2F(\bar{x})(w) \rangle$  by noting that  $d^2F_i(\bar{x})(w) = 0$  for  $i \in \overline{\text{gs}}(\bar{x})$  and  $d^2F_i(\bar{x})(w) = \langle w_{J_i}, \nabla^2 F_i(\bar{x}_{J_i}) w_{J_i} \rangle$  for  $i \in \text{gs}(\bar{x})$ . Thus,

$$\Gamma_0 = \inf_{z \in \mathbb{R}^n} \sup_{u \in \mathcal{A}(F(\bar{x}), dF(\bar{x})(w))} \left\{ \langle u, d^2F(\bar{x})(w|z) \rangle - \Xi(z) \right\} + \langle \bar{\xi}, d^2F(\bar{x})(w) \rangle. \quad (43)$$

Fix any  $z \in \mathbb{R}^n$ . For every  $u \in \mathcal{A}(F(\bar{x}), dF(\bar{x})(w))$ , by using equations (42) and (36) and comparing the expression of  $\Xi(z)$  with that of  $\langle u, d^2F(\bar{x})(w|z) \rangle$ , it follows that  $\langle u, d^2F(\bar{x})(w|z) \rangle - \Xi(z) = \sum_{i \in \overline{\text{gs}}(\bar{x}) \cap \overline{\text{gs}}(w)} [u_i \|z_{J_i}\|_q - \bar{\eta}_i \langle \bar{\zeta}_i, z_{J_i} \rangle]$ , which implies that

$$\sup_{u \in \mathcal{A}(F(\bar{x}), dF(\bar{x})(w))} \left\{ \langle u, d^2F(\bar{x})(w|z) \rangle - \Xi(z) \right\} = \sum_{i \in \overline{\text{gs}}(\bar{x}) \cap \overline{\text{gs}}(w)} \sup_{t \in [-\gamma\lambda, \gamma\lambda]} \left\{ t \|z_{J_i}\|_q - \bar{\eta}_i \langle \bar{\zeta}_i, z_{J_i} \rangle \right\}.$$

Together with the above equality (43), it is immediate to obtain that

$$\Gamma_0 = \sum_{i \in \overline{\text{gs}}(\bar{x}) \cap \overline{\text{gs}}(w)} \inf_{z \in \mathbb{R}^{J_i}} \sup_{t \in [-\gamma\lambda, \gamma\lambda]} \left\{ t \|z\|_q - \bar{\eta}_i \langle \bar{\zeta}_i, z \rangle \right\} + \langle \bar{\xi}, d^2 F(\bar{x})(w) \rangle.$$

Note that  $\inf_{z \in \mathbb{R}^{J_i}} \sup_{t \in [-\gamma\lambda, \gamma\lambda]} \{t \|z\|_q - \bar{\eta}_i \langle \bar{\zeta}_i, z \rangle\} \geq 0$  and the equality holds when  $z = 0$ . This means that the desired equality (41) holds, so  $f$  is parabolically regular at  $\bar{x}$  for  $\bar{v}$ .

By Lemma 5.1 (i),  $d\vartheta(F(\bar{x}))(dF(\bar{x})(0)) = 0$ . Hence,  $\mathcal{C}_f(\bar{x}, \bar{v}) \neq \emptyset$  by Proposition 4.1 (iii). From Theorem 5.1,  $f$  is properly twice epi-differentiable at  $\bar{x}$  for  $\bar{v}$ .  $\square$

By combining Proposition 5.2 and [12, Theorem 13.24], the following corollary holds.

**Corollary 5.2** *Consider problem (2) with  $\vartheta$  and  $F$  from type (II). Consider any  $\bar{x} \in \mathbb{R}^n$ .*

(i) *If  $\bar{x}$  is a local optimal solution, then  $-\nabla f_0(\bar{x}) \in \partial f(\bar{x})$  and for any  $w \in \mathcal{C}_f(\bar{x}, \bar{v})$ ,*

$$\langle w, \nabla^2 f_0(\bar{x})w \rangle + d^2 \vartheta(F(\bar{x}))(dF(\bar{x})(w)) + \langle \bar{\xi}, d^2 F(\bar{x})(w) \rangle \geq 0. \quad (44)$$

where  $\bar{\xi} \in \mathbb{R}^m$  with  $\bar{\xi}_i = \rho'_\lambda(F_i(\bar{x}_{J_i}))$  for  $i \in \text{gs}(\bar{x})$  and  $\bar{\xi}_i = \gamma\lambda$  for  $i \in \overline{\text{gs}}(\bar{x})$ .

(ii) *If  $-\nabla f_0(\bar{x}) \in \partial f(\bar{x})$  and (44) holds with “ $>$ ” for all  $w \in \mathcal{C}_f(\bar{x}, -\nabla f_0(\bar{x})) \setminus \{0\}$ , then  $\bar{x}$  is a strong local optimal solution of problem (2).*

### 5.3 Parabolic regularity for $f$ of type III

We establish the parabolic regularity of  $f = \delta_K$ , where  $K$  is the  $q$ -order cone.

**Proposition 5.3** *Let  $f$  be the composite function of type III. Consider any  $\bar{x} \in \text{dom} f$  and  $\bar{v} \in \partial f(\bar{x})$ . Then, for every  $w \in \mathcal{C}_f(\bar{x}, \bar{v})$ ,*

$$\begin{aligned} d^2 f(\bar{x}|\bar{v})(w) &= \inf_{z \in \mathbb{R}^n} \left\{ d^2 \vartheta(F(\bar{x}))(dF(\bar{x})(w) | d^2 F(\bar{x})(w|z)) - \langle \bar{v}, z \rangle \right\} \\ &= \begin{cases} 0 & \text{if } \bar{x} = 0 \text{ or } \bar{x} \in \text{int} K, \\ \frac{\|\bar{v}\|_2}{2^{1/p}} D^2 F(\bar{x})(w, w) & \text{if } \bar{x} \in \text{bd} K, \end{cases} \end{aligned} \quad (45)$$

and hence  $f$  is both parabolically regular and properly twice epi-differentiable at  $\bar{x}$  for  $\bar{v}$ .

**Proof:** Fix any  $w \in \mathcal{C}_f(\bar{x}, \bar{v})$ . Write  $w = (w_1, w_2) \in \mathbb{R} \times \mathbb{R}^{n-1}$  and  $\bar{v} = (\bar{v}_1, \bar{v}_2) \in \mathbb{R} \times \mathbb{R}^{n-1}$ . By Proposition 4.1 (i),  $d\vartheta(F(\bar{x}))(dF(\bar{x})(w)) = \langle \bar{v}, w \rangle$  and  $w \in \mathcal{T}_{\text{dom} f}(\bar{x})$ . The former means that  $dF(\bar{x})(w) \in \text{dom} d\vartheta(F(\bar{x})) = \mathcal{T}_{\mathbb{R}^-}(F(\bar{x}))$ . By [12, Theorem 8.2], we have

$$0 = d\vartheta(F(\bar{x}))(dF(\bar{x})(w)) = \langle \bar{v}, w \rangle. \quad (46)$$

We proceed the proof by the following three cases:  $\bar{x} \in \text{int} K$ ,  $\bar{x} = 0$  and  $\bar{x} \in \text{bd} K \setminus \{0\}$ .

**Case 1:**  $\bar{x} \in \text{int} K$ . Now  $F(\bar{x}) < 0$  and  $\partial f(\bar{x}) = \mathcal{N}_K(\bar{x}) = \{0\}$ . So  $\bar{v} = 0$ ,  $\partial \vartheta(F(\bar{x})) = \{0\}$  and  $\Lambda_{\bar{x}, \bar{v}} = \{0\}$ . Along with (46),  $dF(\bar{x})(w) \in \mathcal{C}_\vartheta(F(\bar{x}), 0)$ . By invoking (28), it follows

that  $d^2f(\bar{x}|0)(w) \geq d^2\vartheta(F(\bar{x})|0)(dF(\bar{x})(w)) = 0$  where the equality is by the polyhedrality of  $\vartheta = \delta_{\mathbb{R}_-}$ , while from (27) and Proposition 3.3 (iv),

$$d^2f(\bar{x}|0)(w) \leq \inf_{z \in \mathbb{R}^n} d^2\vartheta(F(\bar{x}))(dF(\bar{x})(w)|d^2F(\bar{x})(w|z)) = 0.$$

Thus,  $d^2f(\bar{x}|0)(w) = \inf_{z \in \mathbb{R}^n} \{d^2\vartheta(F(\bar{x}))(dF(\bar{x})(w)|d^2F(\bar{x})(w|z)) - \langle \bar{v}, z \rangle\} = 0$ .

**Case 2:**  $\bar{x} = 0$ . In this case,  $\partial f(\bar{x}) = K^\circ = \{u = (u_1, u_2) \in \mathbb{R} \times \mathbb{R}^{n-1} \mid -u_1 \geq \|u_2\|_p\}$  for  $p = \frac{q}{q-1}$ ,  $F(\bar{x}) = 0$ ,  $\partial\vartheta(F(\bar{x})) = \mathbb{R}_+$  and  $dF(\bar{x})(w) \in \mathcal{T}_{\mathbb{R}_-}(F(\bar{x})) = \mathbb{R}_-$ . From  $\bar{v} \in K^\circ$ ,  $-\bar{v}_1 \in \partial\vartheta(F(\bar{x})) = \mathbb{R}_+$ . For any  $z = (z_1, z_2) \in \mathbb{R} \times \mathbb{R}^{n-1}$ , since  $dF(\bar{x})(z) = \|z_2\|_q - z_1$ ,

$$(-\bar{v}_1)dF(\bar{x})(z) = -\bar{v}_1(-z_1 + \|z_2\|_q) = \bar{v}_1 z_1 - \bar{v}_1 \|z_2\|_q \geq \bar{v}_1 z_1 + \|\bar{v}_2\|_p \|z_2\|_q \geq \langle \bar{v}, z \rangle. \quad (47)$$

This shows that  $-\bar{v}_1 \in \Lambda_{\bar{x}, \bar{v}}$ . In addition, from (46)-(47),  $-\bar{v}_1 \in \mathbb{R}_+$ , and  $dF(\bar{x})(w) \in \mathbb{R}_-$ , we have  $0 \geq (-\bar{v}_1)dF(\bar{x})(w) \geq \langle \bar{v}, w \rangle = 0$ , which implies that  $dF(\bar{x})(w) \in \mathcal{C}_\vartheta(F(\bar{x}), \bar{v})$ . By using (28) and noting that  $d^2F(\bar{x})(w) = 0$ , it follows that

$$d^2f(\bar{x}|\bar{v})(w) \geq d^2\vartheta(F(\bar{x})|-\bar{v}_1)(dF(\bar{x})(w)) - \bar{v}_1 d^2F(\bar{x})(w) = 0.$$

In addition, since  $\text{dom } \vartheta = \mathbb{R}_-$ , we have  $0 \in \mathcal{T}_{\text{dom } \vartheta}^2(F(\bar{x}), dF(\bar{x})(w))$ . By Corollary 2.2 and [12, Example 13.62],  $d^2\vartheta(F(\bar{x}))(dF(\bar{x})(w)|d^2F(\bar{x})(w|0)) = 0$ . Then, from (27),

$$\begin{aligned} d^2f(\bar{x}|\bar{v})(w) &\leq \inf_{z \in \mathbb{R}^n} \{d^2\vartheta(F(\bar{x}))(dF(\bar{x})(w)|d^2F(\bar{x})(w|z)) - \langle \bar{v}, z \rangle\} \\ &\leq d^2\vartheta(F(\bar{x}))(dF(\bar{x})(w)|d^2F(\bar{x})(w|0)) = 0. \end{aligned}$$

Hence,  $d^2f(\bar{x}|\bar{v})(w) = \inf_{z \in \mathbb{R}^n} \{d^2\vartheta(F(\bar{x}))(dF(\bar{x})(w)|d^2F(\bar{x})(w|z)) - \langle \bar{v}, z \rangle\} = 0$ .

**Case 3:**  $\bar{x} \in \text{bd } K \setminus \{0\}$ . Now  $\|\bar{x}_2\|_q = \bar{x}_1 \neq 0$  and  $\partial f(\bar{x}) = \mathcal{N}_K(\bar{x}) = \{tu_p(\bar{x}) \mid t \in \mathbb{R}_+\}$  with  $u_p(\bar{x}) = (-1; \text{sign}(\bar{x}_2) \circ |\bar{x}_2|^{q-1} \bar{x}_1^{1-q})$ . Clearly,  $F$  is twice continuously differentiable in a neighborhood of  $\bar{x}$ . By invoking Proposition 5.1, for any  $w \in \mathcal{C}_f(\bar{x}, \bar{v})$ ,

$$\begin{aligned} d^2f(\bar{x}|\bar{v})(w) &= \inf_{z \in \mathbb{R}^n} \{d^2\vartheta(F(\bar{x}))(dF(\bar{x})(w)|d^2F(\bar{x})(w|z)) - \langle \bar{v}, z \rangle\} \\ &= d^2\vartheta(F(\bar{x}))(dF(\bar{x})(w)) + \langle w, \nabla^2 F(\bar{x})w \rangle \sup_{\xi \in \Lambda_{\bar{x}, \bar{v}}} \xi \end{aligned}$$

where  $\Lambda_{\bar{x}, \bar{v}} = \{\xi \in \partial\vartheta(F(\bar{x})) \mid \nabla F(\bar{x})\xi = \bar{v}\}$ . Notice that  $\|\nabla F(\bar{x})\|_2 = \|u_p(\bar{x})\|_2 = 2^{1/p}$  and  $\partial\vartheta(F(\bar{x})) = \mathbb{R}_+$ . Then, we have  $\Lambda_{\bar{x}, \bar{v}} = \{\frac{\|\bar{v}\|_2}{2^{1/p}}\}$ . Together with the last equation and  $d^2\vartheta(F(\bar{x}))(dF(\bar{x})(w)) = 0$ , the desired result follows. The proof is completed.  $\square$

**Remark 5.2** When  $q = 2$ , the parabolic regularity of  $\delta_K$  is implied by the second-order regularity of  $K$  by [33, Propositions 3.103 & 3.136] and [34, Lemma 15]; and Mohammadi et al. recently provided a direct proof for this fact (see [26, Example 5.8]) by using the developed chain rule for parabolic regularity and reformulating the second order cone as the smooth constraint system  $(\|x_2\|^2 - x_1^2, -x_1) \in \mathbb{R}_-^2$  for  $(x_1, x_2) \in \mathbb{R} \times \mathbb{R}^{n-1}$ . Proposition 5.3 verifies the parabolic regularity of  $\delta_K$  for any  $q > 1$  by applying the obtained upper and lower estimates to its composition form of (1).

From Proposition 5.3, we immediately obtain the parabolic regularity of the Cartesian product of several  $q$ -order cones, which is stated as follows.

**Corollary 5.3** *Let  $K = K_1 \times \dots \times K_m$  with  $K_i = \{(x_{i1}, x_{i2}) \in \mathbb{R} \times \mathbb{R}^{n_i-1} \mid \|x_{i2}\|_q \leq x_{i1}\}$ . Consider any  $\bar{x} = (\bar{x}_1; \bar{x}_2; \dots; \bar{x}_m) \in K$  and  $\bar{v} = (\bar{v}_1; \bar{v}_2; \dots; \bar{v}_m) \in \mathcal{N}_K(\bar{x})$ . Then,  $\delta_K$  is parabolically regular at  $\bar{x}$  for  $\bar{v}$  with  $d^2\delta_K(\bar{x}|\bar{v})(w) = \sum_{i=1}^m d^2\delta_{K_i}(\bar{x}_i|\bar{v}_i)(w_i)$ .*

#### 5.4 Parabolic regularity for $f$ of type IV

Although the parabolic regularity of  $f = \delta_{\mathbb{S}^n_-}$  is implied by the second-order regularity of  $\mathbb{S}^n_-$  (see [33, Propositions 3.103 and 3.136 & Example 3.140]), we here justify it to illustrate the usefulness of the estimates in (27) and (29). Note that the parabolic regularity of  $f$  was also obtained from a composite rule in [22, Example 3.7].

**Proposition 5.4** *Let  $f$  be the composite function of type IV. Consider any  $\bar{x} \in \mathbb{S}^n_-$  and any  $\bar{v} \in \partial f(\bar{x})$ . Then, for any  $w \in \mathcal{C}_f(\bar{x}, \bar{v})$ ,*

$$\begin{aligned} d^2f(\bar{x}|\bar{v})(w) &= \inf_{z \in \mathbb{S}^n} \{d^2\vartheta(F(\bar{x}))(dF(\bar{x})(w)|d^2F(\bar{x})(w|z)) - \langle \bar{v}, z \rangle\} \\ &= \begin{cases} 0 & \text{if } \lambda_1(\bar{x}) < 0, \\ -2\langle \bar{v}, w\bar{x}^\dagger w \rangle & \text{if } \lambda_1(\bar{x}) = 0 \end{cases} \end{aligned} \quad (48)$$

where  $\bar{x}^\dagger$  denotes the pseudo-inverse of  $\bar{x}$ , and hence  $f$  is both parabolically regular and properly twice epi-differentiable at  $\bar{x}$  for  $\bar{v}$ .

**Proof:** Fix any  $w \in \mathcal{C}_f(\bar{x}, \bar{v})$ . By Proposition 4.1 (i),  $d\vartheta(F(\bar{x}))(dF(\bar{x})(w)) = \langle \bar{v}, w \rangle$  and  $w \in \mathcal{T}_{\text{dom } f}(\bar{x})$ . The former means that  $dF(\bar{x})(w) \in \text{dom } d\vartheta(F(\bar{x})) = \mathcal{T}_{\mathbb{R}_-}(F(\bar{x}))$ , which along with  $\vartheta(\cdot) = \delta_{\mathbb{R}_-}(\cdot)$  and [12, Theorem 8.2 (b)] yields that

$$0 = d\vartheta(F(\bar{x}))(dF(\bar{x})(w)) = \langle \bar{v}, w \rangle. \quad (49)$$

We proceed the arguments by the following two cases:  $F(\bar{x}) < 0$  and  $F(\bar{x}) = 0$ .

**Case 1:**  $F(\bar{x}) < 0$ . Now  $\partial f(\bar{x}) = \mathcal{N}_{\mathbb{S}^n_-}(\bar{x}) = \{0\}$ , and hence  $\bar{v} = 0$ ,  $\partial\vartheta(F(\bar{x})) = \{0\}$  and  $\Lambda_{\bar{x}, \bar{v}} = \{0\}$ . From (49),  $dF(\bar{x})(w) \in \mathcal{C}_\vartheta(F(\bar{x}), \bar{v})$ . By invoking (28), it holds that

$$d^2f(\bar{x}|0)(w) \geq d^2\vartheta(F(\bar{x})|0)(dF(\bar{x})(w)) = 0$$

where the equality is by the polyhedrality of  $\vartheta$ , while from (27) and Proposition 3.3 (iv),

$$d^2f(\bar{x}|0)(w) \leq \inf_{z \in \mathbb{S}^n} d^2\vartheta(F(\bar{x}))(dF(\bar{x})(w)|d^2F(\bar{x})(w|z)) = 0.$$

Thus,  $d^2f(\bar{x}|0)(w) = \inf_{z \in \mathbb{S}^n} \{d^2\vartheta(F(\bar{x}))(dF(\bar{x})(w)|d^2F(\bar{x})(w|z)) - \langle \bar{v}, z \rangle\} = 0$ .

**Case 2:**  $F(\bar{x}) = 0$ . By [12, Theorem 10.49],  $\partial f(\bar{x}) = \mathcal{N}_{\mathbb{S}^n_-}(\bar{x}) = \{v \in \xi\partial\lambda_1(\bar{x}) \mid \xi \in \mathbb{R}_+\}$ . Since  $\bar{v} \in \partial f(\bar{x})$ , there exist  $\xi \in \mathbb{R}_+$  and  $u \in \partial\lambda_1(\bar{x})$  such that  $\bar{v} = \xi u$ . Hence,  $(\xi, u) \in \Gamma_{\bar{x}, \bar{v}}$ . Together with (29), it follows that

$$d^2\delta_{\mathbb{S}^n_-}(\bar{x}|\bar{v})(w) \geq d^2\vartheta(F(\bar{x})|\xi)(dF(\bar{x})(w)) + \xi d^2\lambda_1(\bar{x}|u)(w) = -2\langle \bar{v}, w\bar{x}^\dagger w \rangle,$$

where the equality is due to  $d^2\vartheta(F(\bar{x})|\xi)(dF(\bar{x})(w))$  by the polyhedrality of  $\vartheta$ , and is using [3, Theorem 2.1]. On the other hand, by [12, Proposition 13.64], it holds that

$$\begin{aligned} d^2f(\bar{x}|\bar{v})(w) &\leq \inf_{z \in \mathbb{S}^n} \{d^2f(\bar{x})(w|z) - \langle \bar{v}, z \rangle\} = \inf_{z \in \mathbb{S}^n} \{\delta_{\mathcal{T}_{\mathbb{S}^n}(\bar{x}, w)}(z) - \langle \bar{v}, z \rangle\} \\ &= - \sup_{z \in \mathcal{T}_{\mathbb{S}^n}(\bar{x}, w)} \langle \bar{v}, z \rangle = -2\langle \bar{v}, w\bar{x}^\dagger w \rangle. \end{aligned}$$

where the first equality is by [12, Example 13.62] and the last one is by [33, Page 487].

The above discussions show that the desired equalities hold. The parabolic regularity of  $f$  then follows by Proposition 4.2 (i). The proof is completed.  $\square$

## 6 Conclusions

For the composition  $f$  of type (1) with  $\vartheta$  and  $F$  satisfying Assumption 1, we have derived an upper and lower estimate for its second subderivate by characterizing the second-order variational properties of PWTD functions and establishing the chain rule for its parabolic subderivate, and employed these estimates to achieve its parabolic regularity and then twice epi-differentiability for several classes of popular composite  $f$ . An interesting topic is to explore second-order variational properties of the composition (1) involving other types of nonconvex and nonsmooth  $\vartheta$ .

## References

- [1] A. S. Lewis, M. L. Overton, Eigenvalue optimization, *Acta numerica* 5 (1996) 149–190.
- [2] C. Kan, W. Song, Second-order conditions for existence of augmented lagrange multipliers for eigenvalue composite optimization problems, *Journal of Global Optimization* 63 (1) (2015) 77–97.
- [3] M. Torki, First-and second-order epi-differentiability in eigenvalue optimization, *Journal of Mathematical Analysis and Applications* 234 (2) (1999) 391–416.
- [4] R. T. Rockafellar, Second-order optimality conditions in nonlinear programming obtained by way of epi-derivatives, *Mathematics of Operations Research* 14 (3) (1989) 462–484.
- [5] E. Balas, *Disjunctive Programming*, Springer, 2018.
- [6] H. Gfrerer, Optimality conditions for disjunctive programs based on generalized differentiation with application to mathematical programs with equilibrium constraints, *SIAM Journal on Optimization* 24 (2) (2014) 898–931.

- [7] J. Fan, R. Li, Variable selection via nonconcave penalized likelihood and its oracle properties, *Journal of the American statistical Association* 96 (456) (2001) 1348–1360.
- [8] C.-H. Zhang, Nearly unbiased variable selection under minimax concave penalty, *The Annals of Statistics* 38 (2) (2010) 894–942.
- [9] X. Guo, H. Zhang, Y. Wang, J.-L. Wu, Model selection and estimation in high dimensional regression models with group scad, *Statistics & Probability Letters* 103 (2015) 86–92.
- [10] J. O. Ogutu, H.-P. Piepho, Regularized group regression methods for genomic prediction: Bridge, mcp, scad, group bridge, group lasso, sparse group lasso, group mcp and group scad, in: *BMC proceedings*, Vol. 8, BioMed Central, 2014, pp. 1–9.
- [11] Y. Liu, S. Bi, S. Pan, Equivalent lipschitz surrogates for zero-norm and rank optimization problems, *Journal of Global Optimization* 72 (4) (2018) 679–704.
- [12] R. T. Rockafellar, R. J.-B. Wets, *Variational Analysis*, Vol. 317, Springer Science & Business Media, 2009.
- [13] F. J. A. Artacho, M. H. Geoffroy, Metric subregularity of the convex subdifferential in banach spaces, *Journal of Nonlinear and Convex Analysis* 15 (2014) 35–47.
- [14] H. Gfrerer, First order and second order characterizations of metric subregularity and calmness of constraint set mappings, *SIAM Journal on Optimization* 21 (4) (2011) 1439–1474.
- [15] A. D. Ioffe, J. V. Outrata, On metric and calmness qualification conditions in subdifferential calculus, *Set-Valued Analysis* 16 (2) (2008) 199–227.
- [16] H. Gfrerer, J. V. Outrata, On computation of generalized derivatives of the normal-cone mapping and their applications, *Mathematics of Operations Research* 41 (4) (2016) 1535–1556.
- [17] A. Y. Kruger, Error bounds and holder metric subregularity, *Set-Valued Analysis* 23 (4) (2015) 705–736.
- [18] J. Ye, X. Ye, Necessary optimality conditions for optimization problems with variational inequality constraints, *Mathematics of Operations Research* 22 (4) (1997) 977–997.
- [19] G. Li, B. S. Mordukhovich, Holder metric subregularity with applications to proximal point method, *SIAM Journal on Optimization* 22 (4) (2012) 1655–1684.
- [20] X. Y. Zheng, K. F. Ng, Metric subregularity and calmness for nonconvex generalized equations in banach spaces, *SIAM Journal on Optimization* 20 (5) (2010) 2119–2136.

- [21] A. Mohammadi, B. S. Mordukhovich, M. E. Sarabi, Variational analysis of composite models with applications to continuous optimization, *Mathematics of Operations Research* 47 (1) (2022) 397–426.
- [22] A. Mohammadi, M. E. Sarabi, Twice epi-differentiability of extended-real-valued functions with applications in composite optimization, *SIAM Journal on Optimization* 30 (3) (2020) 2379–2409.
- [23] A. Ioffe, Variational analysis of a composite function: A formula for the lower second order epi-derivative, *Journal of Mathematical Analysis and Applications* 160 (2) (1991) 379–405.
- [24] R. Cominetti, On pseudo-differentiability, *Transactions of the American Mathematical Society* 324 (2) (1991) 843–865.
- [25] A. B. Levy, Second-order epi-derivatives of composite functionals, *Annals of Operations Research* 101 (1) (2001) 267–281.
- [26] A. Mohammadi, B. S. Mordukhovich, M. Sarabi, Parabolic regularity in geometric variational analysis, *Transactions of the American Mathematical Society* 374 (3) (2021) 1711–1763.
- [27] A. Mohammadi, First-order variational analysis of non-amenable composite functions, *arXiv preprint arXiv:2204.01191* (2022).
- [28] B. S. Mordukhovich, *Variational Analysis and Applications*, Vol. 30, Springer, 2018.
- [29] H. Gfrerer, On directional metric regularity, subregularity and optimality conditions for nonsmooth mathematical programs, *Set-Valued and Variational Analysis* 21 (2) (2013) 151–176.
- [30] H. Gfrerer, J. V. Outrata, On lipschitzian properties of implicit multifunctions, *SIAM Journal on Optimization* 26 (4) (2016) 2160–2189.
- [31] M. Benko, H. Gfrerer, J. Ye, J. Zhang, J. Zhou, Second-order optimality conditions for general nonconvex optimization problems and variational analysis of disjunctive systems, *arXiv preprint arXiv:2203.10015* (2022).
- [32] R. Henrion, J. V. Outrata, Calmness of constraint systems with applications, *Mathematical Programming* 104 (2) (2005) 437–464.
- [33] J. F. Bonnans, A. Shapiro, *Perturbation Analysis of Optimization Problems*, Springer Science & Business Media, 2000.
- [34] J. F. Bonnans, H. Ramírez C, et al., Perturbation analysis of second-order cone programming problems, *Mathematical Programming* 104 (2) (2005) 205–227.