

# DECAY OF OPERATOR SEMIGROUPS, INFINITE-TIME ADMISSIBILITY, AND RELATED RESOLVENT ESTIMATES

MASASHI WAKAIKI

ABSTRACT. We study decay rates for bounded  $C_0$ -semigroups from the viewpoint of  $L^p$ -infinite-time admissibility and related resolvent estimates. In the Hilbert space setting, polynomial decay of semigroup orbits is characterized by the resolvent behavior in the open right half-plane. A similar characterization based on  $L^p$ -infinite-time admissibility is provided for multiplication semigroups on  $L^q$ -spaces with  $1 \leq q \leq p < \infty$ . We also give a sufficient condition for  $L^2$ -infinite-time admissibility for polynomially stable  $C_0$ -semigroups on Hilbert spaces.

## 1. INTRODUCTION

Consider the abstract Cauchy problem

$$\begin{cases} \dot{u}(t) = Au(t), & t \geq 0 \\ u(0) = x, & x \in X, \end{cases}$$

where  $A$  is the generator of a bounded  $C_0$ -semigroup  $(T(t))_{t \geq 0}$  on a Banach space  $X$ . Throughout this paper, we consider the domain  $D(A)$  of  $A$  to be equipped with the graph norm of  $A$ . Let us assume that  $A$  is invertible. Then the domain of  $A$  coincides with the range of  $A^{-1}$ . To obtain uniform decay rates of classical solutions, we study the quantitative behavior of the operator norm  $\|T(t)A^{-1}\|$  as  $t \rightarrow \infty$ . We concentrate mainly on the situation where  $\|T(t)A^{-1}\|$  decays polynomially, i.e.,  $\|T(t)A^{-1}\| = O(t^{-1/\alpha})$  as  $t \rightarrow \infty$  for some  $\alpha > 0$ .

The relation between polynomial rates of decay of  $\|T(t)A^{-1}\|$  and growth of  $(i\eta I - A)^{-1}$  on the imaginary axis  $i\mathbb{R}$  has been studied in, e.g., [3, 5, 6, 29]. In particular, it was shown in [6] that  $\|T(t)A^{-1}\| = O(t^{-1/\alpha})$  as  $t \rightarrow \infty$  is equivalent to  $\|(i\eta I - A)^{-1}\| = O(|\eta|^{-\alpha})$  as  $|\eta| \rightarrow \infty$  in the Hilbert space setting. We refer to [4, 5, 10, 33] for more general transference between semigroup decay and resolvent growth on  $i\mathbb{R}$ . The long-time asymptotic behavior of semigroup orbits is related to the resolvent of  $A$  on the open left half plane  $\mathbb{C}_+$  as well. For example, the Gearhart-Prüss theorem (see, e.g., [1, Theorem 5.2.1] and [14, Theorem V.1.11]) shows that exponential stability of a  $C_0$ -semigroup on a Hilbert space is equivalent to uniform boundedness of the resolvent on  $\mathbb{C}_+$ . Moreover, it was proved in [38] that density of the set

$$\left\{ x \in X : \lim_{\xi \rightarrow 0^+} \sqrt{\xi}((\xi + i\eta)I - A)^{-1}x = 0 \text{ for all } \eta \in \mathbb{R} \right\}$$

is sufficient for strong stability of the bounded  $C_0$ -semigroup when  $X$  is a Hilbert space. For further references and recent developments on stability of  $C_0$ -semigroups, we refer to the survey papers [10, 11]. In [2, 7, 12, 13, 21, 34, 40], upper bounds for polynomial rates of growth of  $\|T(t)\|$  were obtained from the following resolvent condition: There exist  $K \geq 1$  and  $0 \leq \alpha < \infty$  such that  $\|(\lambda I - A)^{-1}\| \leq K((\operatorname{Re} \lambda)^{-\alpha} + 1)$  for all  $\lambda \in \mathbb{C}_+$ . A bounded  $C_0$ -semigroup  $(T(t))_{t \geq 0}$  on a Hilbert space satisfies  $\|T(t)A^{-1}\| = O(t^{-1/\alpha})$  as  $t \rightarrow \infty$  if and only if  $\sup_{\lambda \in \mathbb{C}_+} \|(\lambda I - A)^{-1}(-A)^{-\alpha}\| < \infty$ ; see [6]. In this equivalence, the fractional power  $(-A)^{-\alpha}$  plays a role of a smoothing factor that cancels resolvent growth. When we replace it with a weaker smoothing factor  $(-A)^{-\alpha+\varepsilon}$ ,  $0 < \varepsilon < \alpha$ , the resulting norm

---

*Key words and phrases.*  $C_0$ -semigroup, Infinite-dimensional system, Infinite-time admissibility, Polynomial stability.  
This work was supported by JSPS KAKENHI Grant Number JP20K14362.

$\|(\lambda I - A)^{-1}(-A)^{-\alpha+\varepsilon}\|$  may grow to infinity as  $\operatorname{Re} \lambda \rightarrow 0+$ . Then it is natural to ask whether this kind of resolvent growth on  $\mathbb{C}_+$  also characterizes the decay of  $\|T(t)A^{-1}\|$  as  $t \rightarrow \infty$ .

We investigate a resolvent estimate on  $\mathbb{C}_+$  called sometimes the  $p$ -Weiss condition in the community of infinite-dimensional systems, which is defined as follows.

**Definition 1.1.** Let  $X$  and  $Y$  be Banach spaces, and let  $A$  be the generator of a bounded  $C_0$ -semigroup  $(T(t))_{t \geq 0}$  on  $X$ . Fix  $1 \leq p < \infty$ . A linear and bounded operator  $C$  from  $D(A)$  to  $Y$  satisfies the  $p$ -Weiss condition for  $A$  if there exists  $K > 0$  such that for all  $\lambda \in \mathbb{C}_+$ ,

$$\|CR(\lambda, A)\| \leq \frac{K}{(\operatorname{Re} \lambda)^{1-1/p}}.$$

The  $p$ -Weiss condition was introduced as a necessary condition for the following concept by Weiss [42].

**Definition 1.2.** Let  $X$  and  $Y$  be Banach spaces, and let  $A$  be the generator of a  $C_0$ -semigroup  $(T(t))_{t \geq 0}$  on  $X$ . Fix  $1 \leq p < \infty$ . A linear and bounded operator  $C$  from  $D(A)$  to  $Y$  is  $L^p$ -infinite-time admissible for  $A$  if there exists  $M > 0$  such that for all  $x \in D(A)$ ,

$$(1) \quad \int_0^\infty \|CT(t)x\|^p dt \leq M\|x\|^p.$$

Weiss conjectured in [42, 43] that the 2-Weiss condition and  $L^2$ -infinite-time admissibility are equivalent when  $X$  and  $Y$  are Hilbert spaces. This conjecture was resolved negatively. Counterexamples can be found in [25, 26, 46]. However, positive results on the equivalence was obtained in several situations such as (i)  $(T(t))_{t \geq 0}$  is an exponentially stable and right-invertible  $C_0$ -semigroup [42]; (ii)  $(T(t))_{t \geq 0}$  is a contraction  $C_0$ -semigroup and  $Y$  is a finite-dimensional space [23]; and (iii)  $(T(t))_{t \geq 0}$  is a bounded analytic and contraction  $C_0$ -semigroup [28]. The third result was extended from the  $L^2$ -case to the  $L^p$ -case with  $p \geq 1$  in [16, 18] and to Orlicz spaces in [22]. For more information on admissibility and related resolvent conditions, we refer to the survey article [24] and the books [37, 39].

The relation of the  $p$ -Weiss condition and  $L^p$ -infinite-time admissibility to the decay rate

$$(2) \quad \|CT(t)\| \leq \frac{M}{t^{1-1/p}} \quad \text{as } t \rightarrow \infty \text{ for some } M > 0$$

was studied in [15, 18, 22, 45], where  $CT(t)$  extends a linear bounded operator (again denoted by  $CT(t)$ ) from  $X$  to  $Y$  for all  $t > 0$ . It was shown in [45] that if a bounded  $C_0$ -semigroup  $(T(t))_{t \geq 0}$  on a Hilbert space commutes with  $C$  for all  $t \geq 0$ , then  $L^2$ -infinite-time admissibility implies the decay rate (2) with  $p = 2$ . Moreover, the decay rate (2) with  $1 < p < \infty$  is equivalent to the  $p'$ -Weiss condition with  $1/p + 1/p' = 1$  for bounded analytic semigroups on Banach spaces [15, 18]. This was extended to a more general polynomial class of decay rates in [22].

For fixed  $\alpha > 0$  and  $1 \leq p < \infty$ , we prove that  $(-A)^{-\alpha/p}$  satisfies the  $p$ -Weiss condition for  $A$  if and only if  $\|T(t)A^{-1}\| = O(t^{-1/\alpha})$  as  $t \rightarrow \infty$  in the Hilbert space setting. Note that the  $p$ -Weiss condition of  $(-A)^{-\alpha/p}$  gives an upper bound for growth rates of  $\|(\lambda I - A)^{-1}(-A)^{-\alpha/p}\|$  as  $\operatorname{Re} \lambda \rightarrow 0+$ . Since  $(-A)^{-\alpha/p}$  is bounded, the Hille-Yosida theorem guarantees a faster decay rate  $\|(\lambda I - A)^{-1}(-A)^{-\alpha/p}\| = O(1/\operatorname{Re} \lambda)$  as  $\operatorname{Re} \lambda \rightarrow \infty$  than the  $p$ -Weiss condition. We show that the non-polynomial decay  $\|T(t)(-A)^{-\alpha}\| = O(t^{-\beta}(\log t)^{-\gamma})$ , where  $\alpha > 0$ ,  $0 \leq \beta < 1$ , and  $\gamma \geq 0$ , can be also characterized by resolvent growth on  $\mathbb{C}_+$ . Following [4, 9, 30, 33, 36], we obtain analogous characterizations of rates of decay for  $\|T(t)A(I - A)^{-1}\|$  and  $\|T(t)A(I - A)^{-2}\|$ . They are motivated by the problem of quantifying uniform rates of decay of orbits starting in the range  $\operatorname{ran}(A)$  of  $A$  and in  $D(A) \cap \operatorname{ran}(A)$ , respectively.

In the Banach space setting, we can easily see that the decay rate  $\|T(t)A^{-1}\| = O(t^{-1/\alpha})$  is almost equivalent to the  $L^p$ -infinite-time admissibility of  $(-A)^{-\alpha/p}$ . More precisely, the  $L^p$ -infinite-time admissibility of  $(-A)^{-\alpha/p}$  implies the decay rate  $\|T(t)A^{-1}\| = O(t^{-1/\alpha})$ , but we only show that if  $\|T(t)A^{-1}\| = O(t^{-1/\alpha})$ , then  $(-A)^{-\alpha/p-\varepsilon}$  is  $L^p$ -infinite-time admissible for all  $\varepsilon > 0$ . We do not know

whether the latter implication with  $\varepsilon = 0$  is true for general  $C_0$ -semigroups. However, it is true and hence we obtain an equivalence result for multiplication semigroups on  $L^q$ -spaces with  $1 \leq q \leq p < \infty$ .

Consider a bounded  $C_0$ -semigroup  $(T(t))_{t \geq 0}$  on a Hilbert space  $X$  such that  $\|T(t)A^{-1}\|$  decays polynomially. The next objective is to obtain a sufficient condition for the  $L^2$ -infinite-time admissibility of a general linear and bounded operator  $C$  from  $D(A)$  to another Hilbert space  $Y$ . One of the motivations is that the dual notion of  $L^2$ -infinite-time admissibility in Definition 1.1 is closely related to input-to-state stability with respect to squared integrable inputs; see [41] for duality results on admissibility. A survey on input-to-state stability of infinite-dimensional systems is given in [31]. A crucial part of the sufficient condition is that there is  $\alpha > 0$  such that  $C(-A)^\alpha$  extends to a linear and bounded operator from  $D(A)$  to  $Y$  and its extension satisfies the 2-Weiss condition for  $A$ , which is called the *strong 2-Weiss condition for  $A$*  in this paper. We show that  $C$  is  $L^2$ -infinite-time admissible if  $C$  satisfies the strong 2-Weiss condition and is  $L^2$ -finite-time admissible, in whose definition the integral on  $[0, \infty)$  in (1) is replaced by an integral on some finite interval  $[0, t_1)$ . We also examine the relation between the strong 2-Weiss condition and the decay rate  $\|CT(t)\| = O(1/\sqrt{t^{1+\beta}})$  for some  $\beta > 0$  when  $Y = X$  and  $T(t)$  commutes with  $C$  for all  $t \geq 0$ .

The paper is organized as follows. In Section 2, we recall some basics on the polynomial decay of semigroup orbits and Plancherel's theorem. In Section 3, the polynomial decay of semigroup orbits is characterized by the Weiss condition in the Hilbert space setting and by infinite-time admissibility in the Banach space setting. In Section 4, an analogous resolvent characterization of non-polynomial decay of semigroup orbits is provided. In Section 5, we obtain a sufficient condition for  $L^2$ -infinite-time admissibility by using the decay estimate of  $\|T(t)(-A)^{-\alpha}\|$ .

**Notation.** Let  $\mathbb{C}_+$ ,  $\mathbb{C}_-$  and  $i\mathbb{R}$  denote the open left half-plane  $\{\lambda \in \mathbb{C} : \operatorname{Re} \lambda > 0\}$ , the open right half-plane  $\{\lambda \in \mathbb{C} : \operatorname{Re} \lambda < 0\}$ , and the imaginary axis  $\{i\eta : \eta \in \mathbb{R}\}$ , respectively. Let  $X$  and  $Y$  be Banach spaces. The space of all bounded linear operators from  $X$  to  $Y$  is denoted by  $\mathcal{L}(X, Y)$ . We write  $\mathcal{L}(X) := \mathcal{L}(X, X)$ . The domain and the range of a linear operator  $A : X \rightarrow Y$  are denoted by  $D(A)$  and  $\operatorname{ran}(A)$ , respectively. When  $A$  is closed,  $D(A)$  is seen as a Banach space with the graph norm  $\|x\|_A := \|x\| + \|Ax\|$ . We denote by  $\sigma(A)$  and  $\varrho(A)$  the spectrum and the resolvent set of a linear operator  $A : D(A) \subset X \rightarrow X$ , respectively. We write  $R(\lambda, A) := (\lambda I - A)^{-1}$  for  $\lambda \in \varrho(A)$ . For linear operators  $A : D(A) \subset X \rightarrow X$  and  $B : D(B) \subset X \rightarrow X$ , we write  $A \subset B$  if  $D(A) \subset D(B)$  and  $Ax = Bx$  for all  $x \in D(A)$ , and the composite  $BA$  is defined by  $BAx := B(Ax)$  with domain  $D(BA) := \{x \in D(A) : Ax \in D(B)\}$ . Let  $\langle \cdot, \cdot \rangle$  denote the inner product on a Hilbert space.

## 2. PRELIMINARIES

**2.1. Fractional powers and polynomial decay.** Let  $X$  be a Banach space. For a linear operator  $A : D(A) \subset X \rightarrow X$  and  $T \in \mathcal{L}(X)$ , we say that  $T$  *commutes with  $A$*  if  $TA \subset AT$ . We say that a densely defined, linear operator  $A$  on  $X$  is *sectorial* if  $(-\infty, 0) \subset \varrho(A)$  and if there exists  $K > 0$  such that  $\|\lambda(\lambda + A)^{-1}\| \leq K$  for all  $\lambda > 0$ . We refer to [17, Section 2.1] for background knowledge on sectorial operators.

Let  $(T(t))_{t \geq 0}$  be a bounded  $C_0$ -semigroup on a Banach space  $X$  with generator  $A$  such that  $0 \in \varrho(A)$ . Then  $-A$  and  $(-A)^{-1}$  are sectorial. Hence the fractional power  $(-A)^\alpha$  is well defined for each  $\alpha \in \mathbb{R}$  as in [17, Chapter 3]. Since  $R(\lambda, A)$  and  $T(t)$  commute with  $A$  for all  $\lambda \in \varrho(A)$  and  $t \geq 0$ , they also commute with  $(-A)^\alpha$  for all  $\alpha \in \mathbb{R}$ ; see, e.g., Propositions 3.1.1.f) and 3.2.1.a) of [17]. We shall use this commutative property frequently without comment.

For a fixed  $\alpha > 0$ , the decay rate of  $\|T(t)(-A)^{-\alpha}\|$  is linked to that of  $\|T(t)A^{-1}\|$  as shown in [3, Proposition 3.1]. We here present it in a general form. The proof can be found in [4, Lemma 4.2].

**Proposition 2.1.** *Let  $(T(t))_{t \geq 0}$  be a bounded  $C_0$ -semigroup on a Banach space  $X$ , and let  $B \in \mathcal{L}(X)$  be a sectorial operator such that  $T(t)$  commutes with  $B$  for all  $t \geq 0$ . For fixed  $\alpha, \beta > 0$ , the following statements are equivalent:*

- (i)  $\|T(t)B\| = O(t^{-1/\alpha})$  as  $t \rightarrow \infty$ .

(ii)  $\|T(t)B^\beta\| = O(t^{-\beta/\alpha})$  as  $t \rightarrow \infty$ .

Polynomial rates of decay of bounded  $C_0$ -semigroups on Hilbert spaces can be characterized by uniform boundedness of the resolvent on  $\mathbb{C}_+$ ; see Lemma 2.3 and Theorem 2.4 of [6] for the proof.

**Theorem 2.2.** *Let  $(T(t))_{t \geq 0}$  be a bounded  $C_0$ -semigroup on a Hilbert space  $X$  with generator  $A$  such that  $i\mathbb{R} \subset \varrho(A)$ . For a fixed  $\alpha > 0$ , the following statements are equivalent:*

- (i)  $\|T(t)A^{-1}\| = O(t^{-1/\alpha})$  as  $t \rightarrow \infty$ .
- (ii) *There exists  $M > 0$  such that  $\|R(\lambda, A)(-A)^{-\alpha}\| \leq M$  for all  $\lambda \in \mathbb{C}_+$ .*

**2.2. Plancherel's theorem.** Let  $(T(t))_{t \geq 0}$  be a bounded  $C_0$ -semigroup on a Hilbert space  $X$  with generator  $A$ . For fixed  $x \in X$  and  $\xi > 0$ , define the functions  $f : \mathbb{R} \rightarrow X$  and  $g : \mathbb{R} \rightarrow X$  by

$$f(t) := \begin{cases} e^{-\xi t}T(t)x & \text{if } t \geq 0 \\ 0 & \text{if } t < 0, \end{cases} \quad g(t) := \begin{cases} te^{-\xi t}T(t)x & \text{if } t \geq 0 \\ 0 & \text{if } t < 0. \end{cases}$$

Their Fourier transforms  $\hat{f}$  and  $\hat{g}$  are given by  $\hat{f}(\eta) = R(\xi + i\eta, A)x$  and  $\hat{g}(\eta) = R(\xi + i\eta)^2x$  for  $\eta \in \mathbb{R}$ . Applying Hilbert-space-valued Plancherel's theorem (see, e.g., [1, Theorem 1.8.2] and [14, Theorem C.14]) to  $f$  and  $g$ , we obtain the next result, which is also called Plancherel's theorem throughout this paper.

**Theorem 2.3.** *Let  $(T(t))_{t \geq 0}$  be a bounded  $C_0$ -semigroup on a Hilbert space  $X$  with generator  $A$ . For all  $x \in X$  and  $\xi > 0$ ,*

$$\begin{aligned} \int_0^\infty \|e^{-\xi t}T(t)x\|^2 dt &= \frac{1}{2\pi} \int_{-\infty}^\infty \|R(\xi + i\eta, A)x\|^2 dt \\ \int_0^\infty \|te^{-\xi t}T(t)x\|^2 dt &= \frac{1}{2\pi} \int_{-\infty}^\infty \|R(\xi + i\eta, A)^2x\|^2 dt. \end{aligned}$$

### 3. POLYNOMIAL DECAY OF SEMIGROUP ORBITS

In this section, we study of the relation of the Weiss condition and infinite-time admissibility to polynomial rates of decay of semigroup orbits.

**3.1. Decay of Hilbert space semigroups and the Weiss condition.** The  $p$ -Weiss condition of  $(-A)^{-\alpha/p}$  characterizes the decay of  $\|T(t)A^{-1}\|$  in the Hilbert space context, which is the main result of this section.

**Theorem 3.1.** *Let  $(T(t))_{t \geq 0}$  be a bounded  $C_0$ -semigroup on a Hilbert space  $X$  with generator  $A$  such that  $0 \in \varrho(A)$ . For fixed  $\alpha > 0$  and  $1 \leq p < \infty$ , the following statements are equivalent:*

- (i)  $\|T(t)A^{-1}\| = O(t^{-1/\alpha})$  as  $t \rightarrow \infty$ .
- (ii)  $(-A)^{-\alpha/p}$  satisfies the  $p$ -Weiss condition for  $A$ .

Before proving Theorem 3.1, we present related results. As a corollary of Theorem 3.1, we separately state an interpolation property for the generator of a bounded  $C_0$ -semigroup, which is the resolvent analogue of Proposition 2.1 with  $B := -A$ .

**Corollary 3.2.** *Let  $(T(t))_{t \geq 0}$  be a bounded  $C_0$ -semigroup on a Hilbert space  $X$  with generator  $A$  such that  $0 \in \varrho(A)$ . For fixed  $\alpha > \beta > 0$ , the following statements are equivalent:*

- (i) *There exists  $M_\alpha > 0$  such that  $\|R(\lambda, A)(-A)^{-\alpha}\| \leq M_\alpha$  for all  $\lambda \in \mathbb{C}_+$ .*
- (ii) *There exists  $M_{\alpha, \beta} > 0$  such that*

$$\|R(\lambda, A)(-A)^{-\beta}\| \leq \frac{M_{\alpha, \beta}}{(\operatorname{Re} \lambda)^{1-\beta/\alpha}}$$

for all  $\lambda \in \mathbb{C}_+$ .

Theorem 3.1 provides a resolvent characterization of rates of decay for the orbits  $T(t)x$  with  $x \in D(A)$ . We will give analogous results on decay rates for the orbits  $T(t)x$  with  $x \in \text{ran}(A)$  and  $x \in D(A) \cap \text{ran}(A)$ . We begin by presenting some background materials on these orbits.

In the case  $x \in \text{ran}(A)$ , we are interested in the decay rate of  $T(t)Ay$  with  $y \in D(A)$  and hence study the decay rate of  $\|T(t)A(I-A)^{-1}\|$ . Since  $D(A) \cap \text{ran}(A) = \text{ran}(A(I-A)^{-2})$  by [4, Proposition 3.10], the problem we address in the case  $x \in D(A) \cap \text{ran}(A)$  is to quantify the decay of  $\|T(t)A(I-A)^{-2}\|$ . By [4, Lemma 3.2],  $-A(I-A)^{-1}$  and  $-A(I-A)^{-2}$  are sectorial. Therefore, the fractional powers  $(-A(I-A)^{-1})^\alpha$  and  $(-A(I-A)^{-2})^\alpha$  are well defined for all  $\alpha > 0$ . By the product and composition rules (see, e.g., Theorem 3.7 (iv) and Remark 3.8 (iv) of [4]), we obtain

$$(-A(I-A)^{-k})^\alpha = (-A)^\alpha(I-A)^{-k\alpha}$$

for each  $\alpha > 0$  and  $k \in \{1, 2\}$ .

By [4, Corollary 6.2],  $\lim_{t \rightarrow \infty} \|T(t)A(I-A)^{-2}\| = 0$  if and only if  $\sigma(A) \cap i\mathbb{R} \subset \{0\}$ . We assume that  $0 \in \sigma(A)$ . When  $0 \in \varrho(A)$ , the decay of  $\|T(t)A(I-A)^{-1}\|$  to zero is equivalent to exponential stability of  $(T(t))_{t \geq 0}$ , and the decay rate of  $\|T(t)A(I-A)^{-2}\|$  is the same as that of  $\|T(t)A^{-1}\|$ . Moreover, we concentrate on the situation where  $\|T(t)A(I-A)^{-1}\|$  and  $\|T(t)A(I-A)^{-2}\|$  decay no faster than  $t^{-1}$  as  $t \rightarrow \infty$ . If not, 0 is the eigenvalue of  $A$  and an isolated point of  $\sigma(A)$ , which is not in our interest; see Theorem 6.9 and a paragraph before Theorem 8.1 of [4] for details.

We state a characterization result on decay rates for  $\|T(t)A(I-A)^{-1}\|$  and  $\|T(t)A(I-A)^{-2}\|$ , whose proof will be given after that of Theorem 3.1.

**Theorem 3.3.** *Let  $(T(t))_{t \geq 0}$  be a bounded  $C_0$ -semigroup on a Hilbert space  $X$  with generator  $A$  such that  $\sigma(A) \cap i\mathbb{R} = \{0\}$ . For fixed  $\alpha, p \geq 1$  and  $k \in \{1, 2\}$ , the following statements are equivalent:*

- (i)  $\|T(t)A(I-A)^{-k}\| = O(t^{-1/\alpha})$  as  $t \rightarrow \infty$ .
- (ii)  $(-A)^{\alpha/p}(I-A)^{-k\alpha/p}$  satisfies the  $p$ -Weiss condition for  $A$ .

Let us now turn to the proof of Theorem 3.1. First we describe how polynomial rates of decay of  $C_0$ -semigroups can be transferred to resolvent growth on  $\mathbb{C}_+$  in the Banach space setting. The following result was proved as the implication (ii)  $\Rightarrow$  (i) of [18, Lemma 2.3]. Although analyticity of  $C_0$ -semigroups is assumed in that lemma, the implication (ii)  $\Rightarrow$  (i) requires only boundedness.

**Lemma 3.4.** *Let  $X$  and  $Y$  be Banach spaces. Let  $(T(t))_{t \geq 0}$  be a bounded  $C_0$ -semigroup on  $X$  with generator  $A$ , and let  $C \in \mathcal{L}(D(A), Y)$  be such that  $CT(t)$  extends to an operator in  $\mathcal{L}(X, Y)$  (also denoted by  $CT(t)$ ) for all  $t > 0$ . If there exist  $M > 0$  and  $1 < p < \infty$  such that*

$$\|CT(t)\| \leq \frac{M}{t^{1/p}}$$

for all  $t > 0$ , then  $C$  satisfies the  $p$ -Weiss condition for  $A$ .

Next we provide a converse result, i.e, transference from resolvents to semigroups, in the Hilbert space setting. A key technique for the proof is contained in the proof of Theorem 4.7 and Remark 4.8 of [4], which proved the case  $F(\xi) \equiv K > 0$  in the proposition below.

**Proposition 3.5.** *Let  $(T(t))_{t \geq 0}$  be a bounded  $C_0$ -semigroup on a Hilbert space  $X$  with generator  $A$ . Let  $C \in \mathcal{L}(D(A), X)$  be such that  $T(t)$  commute with  $C$  for all  $t \geq 0$ . Assume that  $F : (0, \infty) \rightarrow \mathbb{R}_+$  satisfies*

$$(3) \quad \|CR(\lambda, A)\| \leq F(\text{Re } \lambda)$$

for all  $\lambda \in \mathbb{C}_+$ . Then the operator  $CT(t)$  extends to a bounded linear operator (also denoted by  $CT(t)$ ) on  $X$  for all  $t > 0$ , and there exists  $M > 0$  such that

$$\|CT(t)\| \leq M \frac{F(1/t)}{t}$$

for all  $t > 0$ .

*Proof.* Let  $x \in D(A)$  be given. Using Plancherel's theorem, we obtain

$$(4) \quad \int_0^\infty \|te^{-\xi t}CT(t)x\|^2 dt = \frac{1}{2\pi} \int_{-\infty}^\infty \|CR(\xi + i\eta, A)^2x\|^2 d\eta$$

for all  $\xi > 0$ . By the resolvent estimate (3),

$$\begin{aligned} \|CR(\lambda, A)^2x\|^2 &\leq \|CR(\lambda, A)\|^2 \|R(\lambda, A)x\|^2 \\ &\leq F(\operatorname{Re} \lambda)^2 \|R(\lambda, A)x\|^2 \end{aligned}$$

for all  $\lambda \in \mathbb{C}_+$ . Therefore,

$$(5) \quad \int_{-\infty}^\infty \|CR(\xi + i\eta, A)^2x\|^2 d\eta \leq F(\xi)^2 \int_{-\infty}^\infty \|R(\xi + i\eta, A)x\|^2 d\eta$$

for all  $\xi > 0$ . Using Plancherel's theorem again, we have

$$(6) \quad \begin{aligned} \int_{-\infty}^\infty \|R(\xi + i\eta, A)x\|^2 d\eta &= 2\pi \int_0^\infty \|e^{-\xi t}T(t)x\|^2 dt \\ &\leq \frac{\pi c^2}{\xi} \|x\|^2 \end{aligned}$$

for all  $\xi > 0$ , where  $c := \sup_{t \geq 0} \|T(t)\|$ . Combining (4)–(6), we obtain

$$(7) \quad \int_0^\infty \|te^{-\xi t}CT(t)x\|^2 dt \leq \frac{c^2 F(\xi)^2}{2\xi} \|x\|^2$$

for all  $\xi > 0$ . Since

$$e^{-2} \|tCT(t)x\|^2 \leq \|te^{-t/\tau}CT(t)x\|^2$$

for  $0 \leq t \leq \tau$ , the estimate (7) with  $\xi := 1/\tau$  yields

$$\begin{aligned} \int_0^\tau \|tCT(t)x\|^2 dt &\leq e^2 \int_0^\tau \|te^{-t/\tau}CT(t)x\|^2 dt \\ &\leq \frac{e^2 c^2}{2} \tau F(1/\tau)^2 \|x\|^2 \end{aligned}$$

for all  $\tau > 0$ .

Fix  $\tau > 0$ . Since

$$CT(\tau)x = \frac{2}{\tau^2} \int_0^\tau tT(\tau - t)CT(t)x dt,$$

it follows that for all  $y \in X$ ,

$$\begin{aligned} |\langle CT(\tau)x, y \rangle| &= \frac{2}{\tau^2} \left| \left\langle \int_0^\tau tT(\tau - t)CT(t)x dt, y \right\rangle \right| \\ &= \frac{2}{\tau^2} \left| \int_0^\tau \langle tCT(t)x, T(\tau - t)^*y \rangle dt \right| \\ &\leq \frac{2}{\tau^2} \sqrt{\int_0^\tau \|tCT(t)x\|^2 dt} \sqrt{\int_0^\tau \|T(\tau - t)^*y\|^2 dt} \\ &\leq \sqrt{2} ec^2 \frac{F(1/\tau)}{\tau} \|x\| \|y\|, \end{aligned}$$

where  $T(t)^*$  is the Hilbert space adjoint of  $T(t)$  for  $t \geq 0$ . Since  $D(A)$  is dense in  $X$ , it follows that  $CT(\tau)$  extends to a bounded linear operator on  $X$  with norm at most  $\sqrt{2}ec^2F(1/\tau)/\tau$ .  $\square$

Let  $K > 0$  and  $p \geq 1$ . If the function  $F$  in Proposition 3.5 is defined by

$$(8) \quad F(\xi) := \frac{K}{\xi^{1-1/p}}$$

for  $\xi > 0$ , then the resolvent estimate (3) is the  $p$ -Weiss condition of  $C$ . Since

$$\frac{F(1/t)}{t} = \frac{K}{t^{1/p}},$$

it follows from Proposition 3.5 that the  $p$ -Weiss condition of  $C$  implies that

$$\|CT(t)\| \leq \frac{M}{t^{1/p}}$$

for all  $t > 0$  and some  $M > 0$ .

*Proof of Theorem 3.1.* If the statement (i) holds, then we see from [5, Theorem 1.1] that  $i\mathbb{R} \subset \varrho(A)$ , and therefore the implication (i)  $\Rightarrow$  (ii) is true in the case  $p = 1$  by Theorem 2.2. By Proposition 2.1, the statement (i) is true if and only if  $\|T(t)(-A)^{-\alpha/p}\| = O(t^{-1/p})$  as  $t \rightarrow \infty$ . Hence, the implication (i)  $\Rightarrow$  (ii) in the case  $p > 1$  and the implication (ii)  $\Rightarrow$  (i) follow immediately from Lemma 3.4 and Proposition 3.5 with  $C := (-A)^{-\alpha/p}$ , respectively.  $\square$

*Proof of Theorem 3.3.* The implication (i)  $\Rightarrow$  (ii) in the case  $p = 1$  has been shown in Corollary 7.5 and Theorem 7.6 of [4] for  $k = 1$  and in Theorem 8.4 and its proof of [4] for  $k = 2$ . Since Proposition 2.1 shows that the statement (i) is equivalent to

$$\|T(t)(-A)^{\alpha/p}(I - A)^{-k\alpha/p}\| = O\left(\frac{1}{t^{1/p}}\right) \quad \text{as } t \rightarrow \infty$$

for each  $k \in \{1, 2\}$ , the rest follows from Lemma 3.4 and Proposition 3.5 with

$$C := (-A)^{\alpha/p}(I - A)^{-k\alpha/p}$$

as in the proof of Theorem 3.1.  $\square$

**3.2. Decay of Banach space semigroups and infinite-time admissibility.** Next, we study the relation between polynomial decay and infinite-time admissibility. We saw in Proposition 3.5 that the  $p$ -Weiss condition of  $C \in \mathcal{L}(D(A), X)$  implies the decay rate  $\|CT(t)\| = O(t^{-1/p})$  in the Hilbert space setting if  $T(t)$  commutes with  $C$  for all  $t \geq 0$ . We obtain a similar result from the  $L^p$ -infinite-time admissibility of  $C$ , which is stronger than the  $p$ -Weiss condition, in the Banach space setting. This follows by the technique in the proof given in [32] for Datko's theorem. The next result was proved in a slightly less general form (and with a slightly different proof) in [45, Theorem 2.5].

**Lemma 3.6.** *Let  $(T(t))_{t \geq 0}$  be a  $C_0$ -semigroup on a Banach space  $X$  with generator  $A$ , and let  $C \in \mathcal{L}(D(A), X)$  be such that  $T(t)$  commutes with  $C$  for all  $t \geq 0$ . Let  $p \geq 1$  and assume that  $C$  is  $L^p$ -infinite-time admissible for  $A$ . Then the following statements are true:*

- a) *If  $C \in \mathcal{L}(X)$ , then there exists  $M > 0$  such that  $\|CT(t)\| \leq M$  for all  $t \geq 0$ .*
- b) *If  $(T(t))_{t \geq 0}$  is a bounded  $C_0$ -semigroup, then  $CT(t)$  extends to a bounded linear operator (also denoted by  $CT(t)$ ) on  $X$  for all  $t > 0$ , and there exists  $M > 0$  such that*

$$\|CT(t)\| \leq \frac{M}{t^{1/p}}$$

for all  $t > 0$ .

*Proof.* By the  $L^p$ -infinite-time admissibility of  $C$ , there exists a constant  $M_1 > 0$  such that

$$\int_0^\infty \|CT(\tau)x\|^p d\tau \leq M_1^p \|x\|^p$$

for all  $x \in D(A)$ .

a) There exists  $M_2 \geq 1$  and  $w > 0$  such that

$$\|T(t)\| \leq M_2 e^{wt}$$

for all  $t \geq 0$ . Since

$$\begin{aligned} \int_0^t e^{-pw\tau} d\tau \|CT(t)x\|^p &\leq \int_0^t e^{-pw\tau} \|T(\tau)CT(t-\tau)x\|^p dt \\ &\leq M_2^p \int_0^t \|CT(t-\tau)x\|^p dt \\ &\leq M_1^p M_2^p \|x\|^p \end{aligned}$$

for all  $t \geq 0$  and  $x \in D(A)$ , the density of  $D(A)$  implies that

$$\|CT(t)\| \leq \left( \frac{pw}{1 - e^{-pw}} \right)^{1/p} M_1 M_2$$

for all  $t \geq 1$ . From  $C \in \mathcal{L}(X)$ , we also obtain

$$\|CT(t)\| \leq M_2 \|C\| e^{wt}$$

for all  $0 \leq t < 1$ . Thus,  $\|CT(t)\| \leq M$  for all  $t \geq 0$ , where

$$M := \max \left\{ \left( \frac{pw}{1 - e^{-pw}} \right)^{1/p} M_1 M_2, M_2 \|C\| e^w \right\}.$$

b) By assumption, we have

$$c := \sup_{t \geq 0} \|T(t)\| < \infty.$$

For all  $t \geq 0$  and  $x \in D(A)$ ,

$$\begin{aligned} t \|CT(t)x\|^p &= \int_0^t \|T(t-\tau)CT(\tau)x\|^p d\tau \\ &\leq c^p \int_0^\infty \|CT(\tau)x\|^p d\tau \\ &\leq c^p M_1^p \|x\|^p. \end{aligned}$$

From the density of  $D(A)$ , it follows that  $CT(t)$  extends to a bounded linear operator on  $X$  with norm at most  $cM_1 t^{-1/p}$ .  $\square$

We obtain the following proposition from a simple argument.

**Proposition 3.7.** *Let  $(T(t))_{t \geq 0}$  be a bounded  $C_0$ -semigroup on a Banach space  $X$  with generator  $A$  such that  $0 \in \rho(A)$ . For fixed  $\alpha > 0$  and  $p \geq 1$ , the following statements are true:*

- a) *If  $(-A)^{-\alpha/p}$  is  $L^p$ -infinite-time admissible for  $A$ , then  $\|T(t)A^{-1}\| = O(t^{-1/\alpha})$  as  $t \rightarrow \infty$ .*
- b) *If  $\|T(t)A^{-1}\| = O(t^{-1/\alpha})$  as  $t \rightarrow \infty$ , then  $(-A)^{-\alpha/p-\varepsilon}$  with  $\varepsilon > 0$  is  $L^p$ -infinite-time admissible for  $A$ .*

*Proof.* a) This follows immediately from Lemma 3.6.b) with  $C := (-A)^{-\alpha/p}$  and Proposition 2.1.

b) Let  $\beta > \alpha/p$ . By Proposition 2.1, there exists a constant  $M > 0$  such that

$$\|T(t)(-A)^{-\beta}\| \leq \frac{M}{(1+t)^{\beta/\alpha}}$$



for all  $t \geq 0$ . Since  $p\beta/\alpha > 1$ , it follows that for all  $x \in X$ ,

$$\begin{aligned} \int_0^\infty \|(-A)^{-\beta}T(t)x\|^p dt &\leq \int_0^\infty \frac{M^p \|x\|^p}{(1+t)^{p\beta/\alpha}} dt \\ &\leq \frac{M^p \|x\|^p}{p\beta/\alpha - 1}. \end{aligned}$$

Hence  $(-A)^{-\beta}$  is infinite-time admissible for  $A$ .  $\square$

It remains open whether  $\|T(t)A^{-1}\| = O(t^{-1/\alpha})$  implies the  $L^p$ -infinite-time admissibility of  $(-A)^{\alpha/p}$  in the case of general  $C_0$ -semigroups. However, it is true for multiplication semigroups on  $L^q$ -spaces with  $1 \leq q \leq p < \infty$ .

**Proposition 3.8.** *Let  $\mu$  be a  $\sigma$ -finite regular Borel measure on a locally compact Hausdorff space  $\Omega$ . Let  $\phi : \Omega \rightarrow \mathbb{C}$  be measurable with essential range  $\phi_{\text{ess}}(\Omega)$  in  $\mathbb{C}_-$ . Assume that  $A$  is the multiplication operator induced by  $\phi$  on  $L^q(\Omega, \mu)$  for a fixed  $1 \leq q < \infty$ , i.e.,  $Af = \phi f$  with domain  $D(A) := \{f \in L^q(\Omega, \mu) : \phi f \in L^q(\Omega, \mu)\}$ , and let  $(T(t))_{t \geq 0}$  be the  $C_0$ -semigroup on  $L^q(\Omega, \mu)$  generated by  $A$ . Then the following statements are equivalent for fixed  $\alpha > 0$  and  $p \in [q, \infty)$ :*

- (i)  $\|T(t)A^{-1}\| = O(t^{-1/\alpha})$  as  $t \rightarrow \infty$ .
- (ii)  $(-A)^{-\alpha/p}$  is  $L^p$ -infinite-time admissible for  $A$ .

*Proof.* We first note that  $\phi(z) \in \phi_{\text{ess}}(\Omega)$  for almost all  $z \in \Omega$ ; see, e.g., Exercise 19 of Chapter VII in [27]. Therefore,  $(T(t))_{t \geq 0}$  is a bounded  $C_0$ -semigroup. The implication (ii)  $\Rightarrow$  (i) has been shown in Proposition 3.7. Therefore, we here prove the implication (i)  $\Rightarrow$  (ii). We have that for all  $t \geq 0$  and  $f \in X$ ,

$$\begin{aligned} \|T(t)(-A)^{-\alpha/p}f\|^p &= \|e^{t\phi}(-\phi)^{-\alpha/p}f\|^p \\ (9) \quad &= \left( \int_{\Omega} \frac{e^{qt \operatorname{Re} \phi(z)}}{|\phi(z)|^{q\alpha/p}} |f(z)|^q dz \right)^{p/q}. \end{aligned}$$

Since  $p \geq q$ , Jensen's inequality (see, e.g., Theorem 3.3 of [35]) implies that

$$\begin{aligned} \left( \int_{\Omega} \frac{e^{qt \operatorname{Re} \phi(z)}}{|\phi(z)|^{q\alpha/p}} |f(z)|^q dz \right)^{p/q} &\leq \int_{\Omega} \left( \frac{e^{qt \operatorname{Re} \phi(z)}}{|\phi(z)|^{q\alpha/p}} \right)^{p/q} \frac{|f(z)|^q}{\|f\|^q} dz \\ (10) \quad &= \int_{\Omega} \frac{e^{pt \operatorname{Re} \phi(z)}}{|\phi(z)|^\alpha} \frac{|f(z)|^q}{\|f\|^q} dz. \end{aligned}$$

By the assumption  $\phi_{\text{ess}}(\Omega) \subset \mathbb{C}_-$ , we obtain

$$\int_0^\infty \frac{e^{pt \operatorname{Re} \phi(z)}}{|\phi(z)|^\alpha} dt = \frac{1}{p |\operatorname{Re} \phi(z)| |\phi(z)|^\alpha}$$

for almost all  $z \in \Omega$ . Since  $\|T(t)A^{-1}\| = O(t^{-1/\alpha})$  as  $t \rightarrow \infty$ , it follows from [3, Proposition 4.2] that there exist  $\Upsilon > 0$  and  $\kappa > 0$  such that

$$\frac{1}{|\operatorname{Re} \lambda|} \leq \Upsilon |\operatorname{Im} \lambda|^\alpha$$

for all  $\lambda \in \sigma(A)$  satisfying  $|\operatorname{Re} \lambda| \leq \kappa$ . Therefore, if  $\lambda \in \sigma(A)$  satisfies  $|\operatorname{Re} \lambda| \leq \kappa$ , then

$$\frac{1}{|\operatorname{Re} \lambda| |\lambda|^\alpha} \leq \frac{\Upsilon |\operatorname{Im} \lambda|^\alpha}{|\lambda|^\alpha} \leq \Upsilon.$$

If  $\lambda \in \sigma(A)$  satisfies  $|\operatorname{Re} \lambda| > \kappa$ , then

$$\frac{1}{|\operatorname{Re} \lambda| |\lambda|^\alpha} \leq \frac{1}{\kappa^{1+\alpha}}.$$

Define

$$M := \max \left\{ \Upsilon, \frac{1}{\kappa^{1+\alpha}} \right\}.$$

Since  $\phi_{\text{ess}}(\Omega) = \sigma(A)$  by [14, Proposition I.4.10], it follows that  $\phi(z) \in \sigma(A)$  for almost all  $z \in \Omega$ . Hence

$$\frac{1}{|\operatorname{Re} \phi(z)| |\phi(z)|^\alpha} \leq M$$

for almost all  $z \in \Omega$ . This estimate yields

$$(11) \quad \int_{\Omega} \int_0^{\infty} \frac{e^{pt \operatorname{Re} \phi(z)} |f(z)|^q}{|\phi(z)|^\alpha \|f\|^q} dt dz \leq \frac{M}{p} \int_{\Omega} \frac{|f(z)|^q}{\|f\|^q} dz = \frac{M}{p}.$$

Using Fubini's theorem, we have from (9)–(11) that

$$\int_0^{\infty} \|T(t)(-A)^{-\alpha/p} f\|^p dt \leq \frac{M}{p} \|f\|^p.$$

Thus,  $(-A)^{-\alpha/p}$  is  $L^p$ -infinite-time admissible for  $A$ .  $\square$

**3.3. Example.** We present a simple example, in which the decay estimates for  $\|T(t)A^{-1}\|$  in the statement (i) of Theorem 3.1 and Proposition 3.8 are optimal.

Let  $X = \ell^2(\mathbb{N})$  and define an operator  $A : D(A) \subset X \rightarrow X$  by

$$Ax := \left( \left( -\frac{1}{n} + in \right) x_n \right)_{n \in \mathbb{N}}$$

for

$$x = (x_n)_{n \in \mathbb{N}} \in D(A) := \{ \zeta = (\zeta_n)_{n \in \mathbb{N}} \in \ell^2(\mathbb{N}) : (n\zeta_n) \in \ell^2(\mathbb{N}) \}.$$

The semigroup  $(T(t))_{t \geq 0}$  generated by  $A$  is given by

$$T(t)x = \left( e^{(-\frac{1}{n} + in)t} x_n \right)_{n \in \mathbb{N}}$$

for all  $t \geq 0$  and  $x = (x_n)_{n \in \mathbb{N}} \in \ell^2(\mathbb{N})$ . Let  $\alpha > 0$ .

**3.3.1. Weiss condition.** For all  $\lambda \in \mathbb{C}_+$ ,

$$(12) \quad \sqrt{\operatorname{Re} \lambda} \|(-A)^{-\alpha} R(\lambda, A)\| = \sup_{n \in \mathbb{N}} \frac{\sqrt{\operatorname{Re} \lambda}}{|1/n - in|^\alpha |\lambda + 1/n - in|}.$$

Moreover,

$$(13) \quad \frac{\sqrt{\operatorname{Re} \lambda}}{|1/n - in|^\alpha |\lambda + 1/n - in|} \leq \frac{\sqrt{\operatorname{Re} \lambda}}{n^\alpha (\operatorname{Re} \lambda + 1/n)}$$

for all  $\lambda \in \mathbb{C}_+$  and  $n \in \mathbb{N}$ . Fix  $c > 0$ , and define

$$f(\xi) := \frac{\sqrt{\xi}}{\xi + c}$$

for  $\xi > 0$ . Then

$$f'(\xi) = \frac{c - \xi}{2\sqrt{\xi}(\xi + c)^2}.$$

Therefore,  $f(\xi) \leq f(c) = 1/(2\sqrt{c})$  for all  $\xi > 0$ . Combining this with the estimate (13), we obtain

$$(14) \quad \frac{\sqrt{\operatorname{Re} \lambda}}{|1/n - in|^\alpha |\lambda + 1/n - in|} \leq \frac{n^{1/2-\alpha}}{2}$$

for all  $\lambda \in \mathbb{C}_+$  and  $n \in \mathbb{N}$ . On the other hand, since

$$|1/n - in|^\alpha \leq 2^{\alpha/2} n^\alpha,$$

it follows that for each  $n \in \mathbb{N}$ , the case  $\operatorname{Re} \lambda = 1/n$  and  $\operatorname{Im} \lambda = n$  gives

$$(15) \quad \frac{\sqrt{\operatorname{Re} \lambda}}{|1/n - in|^\alpha |\lambda + 1/n - in|} \geq \frac{n^{1/2-\alpha}}{2^{1+\alpha/2}}.$$

Substituting the estimates (14) and (15) into (12), we see that  $(-A)^{-\alpha}$  satisfies the 2-Weiss condition for  $A$  if and only if  $\alpha \geq 1/2$ .

**3.3.2. Infinite-time admissibility.** We directly obtain a similar result on the  $L^2$ -infinite-time admissibility. In fact, using the monotone convergence theorem, we have that for all  $x = (x_n)_{n \in \mathbb{N}} \in \ell^2(\mathbb{N})$ ,

$$\begin{aligned} \int_0^\infty \|(-A)^{-\alpha} T(t)x\|^2 dt &= \int_0^\infty \sum_{n=1}^\infty \left| \left(-\frac{1}{n} + in\right)^{-\alpha} e^{(-\frac{1}{n} + in)t} x_n \right|^2 dt \\ &= \frac{1}{2} \sum_{n=1}^\infty \frac{n^{1+2\alpha}}{(1+n^4)^\alpha} |x_n|^2. \end{aligned}$$

Therefore,  $(-A)^{-\alpha}$  is  $L^2$ -infinite-time admissible for  $A$  if and only if  $\alpha \geq 1/2$ .

**3.3.3. Polynomial decay rate.** Fix  $c > 0$ . Then  $g(t) := te^{-t/c} \leq g(c) = c/e$  for  $t \geq 0$ . From this inequality, we obtain

$$t \|T(t)A^{-1}\| = \sup_{n \in \mathbb{N}} \frac{nte^{-\frac{t}{n}}}{\sqrt{1+n^4}} \leq \sup_{n \in \mathbb{N}} \frac{n^2}{e\sqrt{1+n^4}} \leq \frac{1}{e}$$

for all  $t \geq 0$ . Therefore,  $\|T(t)A^{-1}\| = O(1/t)$  as  $t \rightarrow \infty$ . We also have

$$n \|T(n)A^{-1}\| \geq \frac{n^2}{e\sqrt{1+n^4}}$$

for all  $n \in \mathbb{N}$ , and hence

$$\liminf_{t \rightarrow \infty} t \|T(t)A^{-1}\| \geq \frac{1}{e}.$$

This demonstrates the optimality of the rates of decay when the statement (ii) of Theorem 3.1 or Proposition 3.8 holds for  $p = 2$ .

#### 4. NON-POLYNOMIAL DECAY OF HILBERT SPACE SEMIGROUPS

Let  $X$  and  $Y$  be Hilbert spaces, and let  $(T(t))_{t \geq 0}$  be a bounded  $C_0$ -semigroup on  $X$  with generator  $A$ . It was shown in [44] that if  $C \in \mathcal{L}(D(A), Y)$  satisfies

$$(16) \quad \|CR(\lambda, A)\| \leq \frac{K}{\sqrt{\operatorname{Re} \lambda} |\log \operatorname{Re} \lambda|}$$

for all  $\lambda \in \mathbb{C}_+$  with  $\operatorname{Re} \lambda \neq 1$  and some  $K > 0$ , then  $C$  is  $L^2$ -infinite-time admissible for  $A$ . We see from Proposition 3.5 that if  $Y = X$  and if  $T(t)$  commutes with  $C$  for all  $t \geq 0$ , then the estimate (16) implies

$$(17) \quad \|CT(t)\| \leq \frac{M}{\sqrt{t} |\log t|}$$

for all  $t > 0$  with  $t \neq 1$  and some  $M > 0$ . Note that the estimate (16) contains the condition on the decay rate of  $\|CR(\lambda, A)\|$  as  $\operatorname{Re} \lambda \rightarrow \infty$  in addition to the growth rate of  $\|CR(\lambda, A)\|$  as  $\operatorname{Re} \lambda \rightarrow 0+$ , since  $C$  may not belong to  $\mathcal{L}(X, Y)$ . Similarly, the estimate (17) contains the condition on the growth rate of  $\|CT(t)\|$  as  $t \rightarrow 0+$ . In this section, we concentrate on the case  $C \in \mathcal{L}(X, Y)$ , and then show how the non-polynomial rate of decay of  $\|CT(t)\|$  as  $t \rightarrow \infty$  can be transferred to the growth rate of  $\|CR(\lambda, A)\|$  as  $\operatorname{Re} \lambda \rightarrow 0+$ .

**Proposition 4.1.** *Let  $X$  and  $Y$  be Banach spaces, and let  $(T(t))_{t \geq 0}$  be a bounded  $C_0$ -semigroup on  $X$  with generator  $A$ . For fixed  $0 \leq \beta < 1$  and  $\gamma \geq 0$ , the following statements hold.*

a) If  $C \in \mathcal{L}(X, Y)$  satisfies

$$\|CT(t)\| = O\left(\frac{1}{t^\beta(\log t)^\gamma}\right)$$

as  $t \rightarrow \infty$ , then there exists  $K > 0$  such that

$$\|CR(\lambda, A)\| \leq \frac{K}{(\operatorname{Re} \lambda)^{1-\beta} |\log \operatorname{Re} \lambda|^\gamma}$$

for all  $\lambda \in \mathbb{C}_+$  with  $\operatorname{Re} \lambda < 1$ .

b) If  $C \in \mathcal{L}(X, Y)$  satisfies

$$\|CT(t)\| = O\left(\frac{1}{t(\log t)^\gamma}\right)$$

as  $t \rightarrow \infty$ , then there exists  $K > 0$  such that

$$\|CR(\lambda, A)\| \leq KF_\gamma(\operatorname{Re} \lambda)$$

for all  $\lambda \in \mathbb{C}_+$  with  $\operatorname{Re} \lambda < e^{-1}$ , where

$$F_\gamma(\xi) := \begin{cases} |\log \xi|^{1-\gamma} & \text{if } 0 \leq \gamma < 1 \\ \log |\log \xi| & \text{if } \gamma = 1 \\ 1 & \text{if } \gamma > 1 \end{cases}$$

for  $0 < \xi < e^{-1}$ .

The following lemma will be useful in the proof of Proposition 4.1.

**Lemma 4.2.** For fixed  $0 \leq \beta < 1$  and  $\gamma \geq 0$ , the following statements hold:

a) For all  $t_0 > e^{\gamma/(1-\beta)}$ , there exists  $M > 0$  such that

$$\int_{t_0}^{\infty} \frac{e^{-\xi t}}{t^\beta(\log t)^\gamma} dt \leq \frac{M}{\xi^{1-\beta} |\log \xi|^\gamma}$$

for all  $0 < \xi < 1/t_0$ .

b) For all  $t_0 > e$ , there exists  $M > 0$  such that

$$\int_{t_0}^{\infty} \frac{e^{-\xi t}}{t(\log t)^\gamma} dt \leq MF_\gamma(\xi)$$

for all  $0 < \xi < 1/t_0$ , where  $F_\gamma$  is as in Proposition 4.1.

*Proof.* a) Let  $t_0 > e^{\gamma/(1-\beta)}$  and  $0 < \xi < 1/t_0$ . Then

$$(18) \quad \int_{1/\xi}^{\infty} \frac{e^{-\xi t}}{t^\beta(\log t)^\gamma} dt \leq \frac{1}{(1/\xi)^\beta |\log \xi|^\gamma} \int_{1/\xi}^{\infty} e^{-\xi t} dt = \frac{e^{-1}}{\xi^{1-\beta} |\log \xi|^\gamma}.$$

We also have

$$(19) \quad \int_{t_0}^{1/\xi} \frac{e^{-\xi t}}{t^\beta(\log t)^\gamma} dt \leq \int_{t_0}^{1/\xi} \frac{1}{t^\beta(\log t)^\gamma} dt$$

Integration by parts gives

$$(20) \quad \begin{aligned} \int_{t_0}^{1/\xi} \frac{1}{t^\beta(\log t)^\gamma} dt &= \frac{(1/\xi)^{1-\beta}}{(1-\beta)|\log \xi|^\gamma} - \frac{t_0^{1-\beta}}{(1-\beta)(\log t_0)^\gamma} + \frac{\gamma}{1-\beta} \int_{t_0}^{1/\xi} \frac{1}{t^\beta(\log t)^{\gamma+1}} dt \\ &\leq \frac{(1/\xi)^{1-\beta}}{(1-\beta)|\log \xi|^\gamma} + \frac{\gamma}{(1-\beta)\log t_0} \int_{t_0}^{1/\xi} \frac{1}{t^\beta(\log t)^\gamma} dt. \end{aligned}$$

Since  $t_0 > e^{\gamma/(1-\beta)}$  is equivalent to

$$\frac{\gamma}{(1-\beta)\log t_0} < 1,$$

the estimates (19) and (20) yield

$$(21) \quad \int_{t_0}^{1/\xi} \frac{e^{-\xi t}}{t^\beta (\log t)^\gamma} dt \leq \frac{M_1}{\xi^{1-\beta} |\log \xi|^\gamma},$$

where

$$M_1 := \frac{1}{1-\beta} \left( 1 - \frac{\gamma}{(1-\beta) \log t_0} \right).$$

The assertion with  $M := M_1 + e^{-1}$  then follows from the estimates (18) and (21), noting that  $M$  is independent of  $\xi$ .

b) Let  $t_0 > e$  and  $0 < \xi < 1/t_0$ . Since

$$\int_{t_0}^{1/\xi} \frac{e^{-\xi t}}{t(\log t)^\gamma} dt \leq \int_{t_0}^{1/\xi} \frac{1}{t(\log t)^\gamma} dt = \int_{\log t_0}^{\log(1/\xi)} \frac{1}{\tau^\gamma} d\tau,$$

we obtain

$$\int_{t_0}^{1/\xi} \frac{e^{-\xi t}}{t(\log t)^\gamma} dt \leq \begin{cases} (1-\gamma)(|\log \xi|^{1-\gamma} - (\log t_0)^{1-\gamma}) & \text{if } \gamma \neq 1 \\ \log |\log \xi| - \log(\log t_0) & \text{if } \gamma = 1. \end{cases}$$

Define

$$G_\gamma(\xi) := \begin{cases} (1-\gamma)|\log \xi|^{1-\gamma} & \text{if } 0 \leq \gamma < 1 \\ \log |\log \xi| & \text{if } \gamma = 1 \\ (\gamma-1)(\log t_0)^{1-\gamma} & \text{if } \gamma > 1. \end{cases}$$

Then, instead of the estimate (21), we have

$$\int_{t_0}^{1/\xi} \frac{e^{-\xi t}}{t(\log t)^\gamma} dt \leq G_\gamma(\xi).$$

On the other hand, the estimate (18) yield

$$\int_{1/\xi}^{\infty} \frac{e^{-\xi t}}{t(\log t)^\gamma} dt \leq \frac{e^{-1}}{|\log \xi|^\gamma} < \frac{e^{-1}}{(\log t_0)^\gamma}.$$

Thus, the desired estimate is obtained for some  $M > 0$  depending only on  $t_0$  and  $\gamma$ .  $\square$

*Proof of Proposition 4.1.* a) By assumption, there exist  $M_1 > 0$  and  $t_0 > e^{\gamma/(1-\beta)}$  such that

$$\|CT(t)\| \leq \frac{M_1}{t^\beta (\log t)^\gamma}$$

for all  $t \geq t_0$ . For all  $x \in X$  and  $\lambda \in \mathbb{C}_+$ ,

$$\|CR(\lambda, A)x\| \leq \int_0^\infty e^{-t \operatorname{Re} \lambda} \|CT(t)x\| dt.$$

Define  $c := \sup_{t \geq 0} \|CT(t)\|$ . For all  $x \in X$  and  $\lambda \in \mathbb{C}_+$ ,

$$\begin{aligned} \int_0^\infty e^{-t \operatorname{Re} \lambda} \|CT(t)x\| dt &\leq \int_0^{t_0} c \|x\| dt + \int_{t_0}^\infty e^{-t \operatorname{Re} \lambda} \frac{M_1 \|x\|}{t^\beta (\log t)^\gamma} dt \\ &= \left( ct_0 + M_1 \int_{t_0}^\infty \frac{e^{-t \operatorname{Re} \lambda}}{t^\beta (\log t)^\gamma} dt \right) \|x\|. \end{aligned}$$

Hence by Lemma 4.2.a), there exists  $K_1 > 0$  such that

$$\|CR(\lambda, A)\| \leq \frac{K_1}{(\operatorname{Re} \lambda)^{1-\beta} |\log \operatorname{Re} \lambda|^\gamma}.$$

for all  $\lambda \in \mathbb{C}_+$  with  $\operatorname{Re} \lambda < 1/t_0$ . Combining this with the Hille-Yosida theorem, we obtain the desired estimate for all  $\lambda \in \mathbb{C}_+$  with  $\operatorname{Re} \lambda < 1$  and some  $K > 0$ .

b) By the same arguments as above, we have that the statement b) is true, using Lemma 4.2.b) instead of Lemma 4.2.a).  $\square$

Let  $(T(t))_{t \geq 0}$  be a bounded  $C_0$ -semigroup on a Hilbert space with generator  $A$  such that  $i\mathbb{R} \subset \varrho(A)$ , and let  $\alpha > 0$ . From Theorem 2.2, we already know that  $\|T(t)(-A)^{-\alpha}\| = O(1/t)$  as  $t \rightarrow \infty$  if and only if  $\|R(\lambda, A)(-A)^{-\alpha}\| \leq K$  for all  $\lambda \in \mathbb{C}_+$  and some  $K > 0$ . However, Proposition 4.1.b) with  $C := (-A)^{-\alpha}$  and  $\gamma = 0$  only shows that if  $\|T(t)(-A)^{-\alpha}\| = O(1/t)$  as  $t \rightarrow \infty$ , then  $\|R(\lambda, A)(-A)^{-\alpha}\| \leq K |\log \operatorname{Re} \lambda|$  for all  $\lambda \in \mathbb{C}_+$  with  $\operatorname{Re} \lambda < e^{-1}$  and some  $K > 0$ . From this we see that Proposition 4.1.b) with  $C := (-A)^{-\alpha}$  does not give a sharp bound of  $\|R(\lambda, A)(-A)^{-\alpha}\|$ . In contrast, Proposition 4.1.a) with  $C := (-A)^{-\alpha}$ , combined with Proposition 3.5, yields a characterization result for the non-polynomial rates of decay of  $\|T(t)(-A)^{-\alpha}\|$  as  $t \rightarrow \infty$  and growth of  $\|R(\lambda, A)(-A)^{-\alpha}\|$  as  $\operatorname{Re} \lambda \rightarrow 0+$ .

**Theorem 4.3.** *Let  $(T(t))_{t \geq 0}$  be a bounded  $C_0$ -semigroup on a Hilbert space  $X$  with generator  $A$  such that  $0 \in \varrho(A)$ . The following statements are equivalent for fixed  $\alpha > 0$ ,  $0 \leq \beta < 1$ , and  $\gamma \geq 0$ :*

- (i)  $\|T(t)(-A)^{-\alpha}\| = O(t^{-\beta}(\log t)^{-\gamma})$  as  $t \rightarrow \infty$ .
- (ii) *There exists  $K > 0$  such that*

$$\|R(\lambda, A)(-A)^{-\alpha}\| \leq \frac{K}{(\operatorname{Re} \lambda)^{1-\beta} |\log \operatorname{Re} \lambda|^\gamma}$$

for all  $\lambda \in \mathbb{C}_+$  with  $\operatorname{Re} \lambda < 1$ .

*Proof.* The implication (i)  $\Rightarrow$  (ii) has already been proved in Proposition 4.1.a) with  $C := (-A)^{-\alpha}$ . To show the converse implication (ii)  $\Rightarrow$  (i), we define the function  $F$  in Proposition 3.5 by

$$F(\xi) := \frac{K}{\xi^{1-\beta} |\log \xi|^\gamma}$$

for  $0 < \xi < 1$ . Then

$$\frac{F(1/t)}{t} = \frac{K}{t^\beta (\log t)^\gamma}$$

for all  $t > 1$ . Thus, the implication (ii)  $\Rightarrow$  (i) holds by Proposition 3.5 with  $C := (-A)^{-\alpha}$ .  $\square$

By the same arguments as in Theorem 4.3, we obtain a characterization of non-polynomial rates of decay for  $\|T(t)(-A)^\alpha(I - A)^{-\alpha-\delta}\|$  with  $\alpha > 0$  and  $\delta \geq 0$ .

**Theorem 4.4.** *Let  $(T(t))_{t \geq 0}$  be a bounded  $C_0$ -semigroup on a Hilbert space  $X$  with generator  $A$  such that  $\sigma(A) \cap i\mathbb{R} = \{0\}$ . The following statements are equivalent for fixed  $\alpha > 0$ ,  $0 \leq \beta < 1$ ,  $\gamma \geq 0$ , and  $\delta \geq 0$ :*

- (i)  $\|T(t)(-A)^\alpha(I - A)^{-\alpha-\delta}\| = O(t^{-\beta}(\log t)^{-\gamma})$  as  $t \rightarrow \infty$ .
- (ii) *There exists  $K > 0$  such that*

$$\|R(\lambda, A)(-A)^\alpha(I - A)^{-\alpha-\delta}\| \leq \frac{K}{(\operatorname{Re} \lambda)^{1-\beta} |\log \operatorname{Re} \lambda|^\gamma}$$

for all  $\lambda \in \mathbb{C}_+$  with  $\operatorname{Re} \lambda < 1$ .

## 5. SUFFICIENT CONDITION FOR $L^2$ -INFINITE-TIME ADMISSIBILITY

Let  $A$  be the generator of a bounded  $C_0$ -semigroup  $(T(t))_{t \geq 0}$  on a Hilbert space  $X$ . In this section, we derive a sufficient condition for  $L^2$ -infinite-time admissibility of general bounded linear operators from  $D(A)$  to a Hilbert space  $Y$ . Before proceeding to the details, we recall the definitions of polynomial stability of  $C_0$ -semigroups and finite-time admissibility.

**Definition 5.1.** A  $C_0$ -semigroup  $(T(t))_{t \geq 0}$  on a Banach space  $X$  with generator  $A$  is *polynomially stable* if the following three conditions are satisfied:

- a)  $(T(t))_{t \geq 0}$  is bounded;
- b)  $i\mathbb{R} \subset \varrho(A)$ ; and

c) There exists  $\alpha > 0$  such that  $\|T(t)A^{-1}\| = O(t^{-1/\alpha})$  as  $t \rightarrow \infty$ .

**Definition 5.2.** Let  $X$  and  $Y$  be Banach spaces, and let  $A$  be the generator of a  $C_0$ -semigroup  $(T(t))_{t \geq 0}$  on  $X$ . An operator  $C \in \mathcal{L}(D(A), Y)$  is  $L^2$ -finite-time admissible for  $A$  if there exist  $M_1 > 0$  and  $t_1 > 0$  such that for all  $x \in D(A)$ ,

$$\int_0^{t_1} \|CT(t)x\|^2 dt \leq M_1 \|x\|^2.$$

From the semigroup property, it follows that if  $C \in \mathcal{L}(D(A), Y)$  is  $L^2$ -finite-time admissible for  $A$ , then for each  $t_2 > 0$ , there exists  $M_2 > 0$  such that for all  $x \in D(A)$ ,

$$\int_0^{t_2} \|CT(t)x\|^2 dt \leq M_2 \|x\|^2.$$

By definition,  $C \in \mathcal{L}(X, Y)$  is  $L^2$ -finite-time admissible. Moreover, if the  $C_0$ -semigroup  $(T(t))_{t \geq 0}$  is right invertible and if  $\|CR(\cdot, A)\|$  is bounded on some right half-plane, then  $C$  is  $L^2$ -finite-time admissible for  $A$ ; see [42, Theorem 4.1] and [44, Theorem 2.2]. Other sufficient conditions for  $L^2$ -finite-time admissibility can be found, e.g., in [44, Theorem 3.3] and [8, Lemma 2.6].

Combing  $L^2$ -finite-time admissibility with the logarithmic decay of  $\|T(t)(-A)^{-\alpha}\|$  and the 2-Weiss condition of  $C(-A)^\alpha$ , we obtain a sufficient condition for  $L^2$ -infinite-time admissibility. It can be proved by the technique used in the proof of Theorem 4.2 in [44]. Note that the extension of  $C(-A)^\alpha$  to  $\mathcal{L}(D(A), Y)$  is unique if it exists, since  $D((-A)^{1+\alpha})$  is dense in  $D(A)$  with respect to the graph norm  $\|\cdot\|_A$  by [17, Proposition 3.1.1.h)].

**Theorem 5.3.** Let  $X$  and  $Y$  be Hilbert spaces. Let  $(T(t))_{t \geq 0}$  be a bounded  $C_0$ -semigroup on  $X$  with generator  $A$  such that  $0 \in \rho(A)$ , and let  $C \in \mathcal{L}(D(A), Y)$ . Assume that the following conditions hold for fixed  $\alpha > 0$  and  $\beta > 1/2$ :

- a)  $C$  is  $L^2$ -finite-time admissible for  $A$ ;
- b)  $\|T(t)(-A)^{-\alpha}\| \leq O((\log t)^{-\beta})$  as  $t \rightarrow \infty$ ; and
- c)  $C(-A)^\alpha$  extends to an operator in  $\mathcal{L}(D(A), Y)$ , and its extension satisfies the 2-Weiss condition for  $A$ .

Then  $C$  is  $L^2$ -infinite-time admissible for  $A$ .

*Proof.* Using Plancherel's theorem, we have that for all  $\xi > 0$  and  $x \in X$ ,

$$(22) \quad \int_{-\infty}^{\infty} \|R(\xi + i\eta, A)(-A)^{-\alpha}x\|^2 d\eta = 2\pi \int_0^{\infty} \|e^{-\xi t}T(t)(-A)^{-\alpha}x\|^2 dt.$$

By the assumption b), there exists constants  $M_1 > 0$  and  $t_0 > e^\beta$  such that

$$\|T(t)(-A)^{-\alpha}\| \leq \frac{M_1}{(\log t)^\beta}$$

for all  $t > t_0$ . Lemma 4.2 shows that there exists a constant  $M_2 > 0$  such that

$$\int_{t_0}^{\infty} \frac{e^{-\xi t}}{(\log t)^{2\beta}} dt \leq \frac{M_2}{\xi |\log \xi|^{2\beta}}$$

for all  $0 < \xi < 1/t_0$ . Put  $c := \sup_{t \geq 0} \|T(t)(-A)^{-\alpha}\|$ . Then

$$(23) \quad \begin{aligned} \int_0^{\infty} \|e^{-\xi t}T(t)(-A)^{-\alpha}x\|^2 dt &\leq c^2 t_0 \|x\|^2 + M_1^2 \int_{t_0}^{\infty} \frac{e^{-2\xi t}}{(\log t)^{2\beta}} dt \|x\|^2 \\ &\leq \left( c^2 t_0 + \frac{M_1^2 M_2}{2\xi |\log(2\xi)|^{2\beta}} \right) \|x\|^2 \end{aligned}$$

for all  $0 < \xi < 1/(2t_0)$  and  $x \in X$ . Let the extension of  $C(-A)^\alpha$  in the assumption c) be also denoted by  $C(-A)^\alpha$ . There exist a constant  $K > 0$  such that

$$(24) \quad \|C(-A)^\alpha R(\lambda, A)\| \leq \frac{K}{\sqrt{\operatorname{Re} \lambda}}$$

for all  $\lambda \in \mathbb{C}_+$ . From (22)–(24), it follows that for all  $0 < \xi < 1/(2t_0)$  and  $x \in X$ ,

$$(25) \quad \begin{aligned} & \int_{-\infty}^{\infty} \|CR(\xi + i\eta, A)^2 x\|^2 d\eta \\ & \leq \sup_{\eta \in \mathbb{R}} \|C(-A)^\alpha R(\xi + i\eta, A)\|^2 \int_{-\infty}^{\infty} \|R(\xi + i\eta, A)(-A)^{-\alpha} x\|^2 d\eta \\ & \leq \frac{2\pi K^2}{\xi} \left( c^2 t_0 + \frac{M_1^2 M_2}{2\xi |\log(2\xi)|^{2\beta}} \right) \|x\|^2. \end{aligned}$$

Using Plancherel's theorem again, we obtain that for all  $\xi > 0$  and  $x \in D(A)$ ,

$$\int_0^{\infty} \|te^{-\xi t} CT(t)x\|^2 dt = \frac{1}{2\pi} \int_{-\infty}^{\infty} \|CR(\xi + i\eta, A)^2 x\|^2 d\eta.$$

This and the inequality (25) shows that there exists a constant  $M_3 > 0$  such that

$$(26) \quad \int_0^{\infty} \|\xi te^{-\xi t} CT(t)x\|^2 dt \leq \frac{M_3}{|\log(2\xi)|^{2\beta}} \|x\|^2$$

for all  $0 < \xi < 1/(2t_0)$  and  $x \in D(A)$ .

Take  $\tau_1 < 1 < \tau_2$  such that  $\tau_1 e^{-\tau_1} = 1/(2e) = \tau_2 e^{-\tau_2}$ . Set

$$\mu_n := \left( \frac{\tau_1}{\tau_2} \right)^{n-1}, \quad \tau_n := \frac{\tau_1}{\mu_n}.$$

Then  $\mu_n \rightarrow 0$  and  $\tau_n \rightarrow \infty$  as  $n \rightarrow \infty$ . Moreover, for all  $n \in \mathbb{N}$ ,

$$(27) \quad \mu_n t e^{-\mu_n t} > \frac{1}{2e}$$

whenever  $t \in (\tau_n, \tau_{n+1})$ . Let  $m \in \mathbb{N}$  satisfy  $\mu_m < 1/(2t_0)$ . Combining the estimates (26) and (27), we have that for all  $x \in D(A)$ ,

$$\begin{aligned} \int_{\tau_m}^{\infty} \|CT(t)x\|^2 dt &= \sum_{n=m}^{\infty} \int_{\tau_n}^{\tau_{n+1}} \|CT(t)x\|^2 dt \\ &\leq (2e)^2 \sum_{n=m}^{\infty} \int_{\tau_n}^{\tau_{n+1}} \|\mu_n t e^{-\mu_n t} CT(t)x\|^2 dt \\ &\leq (2e)^2 M_3 \|x\|^2 \sum_{n=m}^{\infty} \frac{1}{|\log(2\mu_n)|^{2\beta}}. \end{aligned}$$

Since

$$\frac{1}{|\log(2\mu_n)|} = \frac{1}{(\log(\tau_2) - \log(\tau_1))(n-1) - \log 2}$$

for  $n \geq m$ , we have from  $\beta > 1/2$  that

$$M_4 := (2e)^2 M_3 \sum_{n=m}^{\infty} \frac{1}{|\log(2\mu_n)|^{2\beta}} \in (0, \infty),$$

and then

$$(28) \quad \int_{\tau_m}^{\infty} \|CT(t)x\|^2 dt \leq M_4 \|x\|^2$$



for all  $x \in D(A)$ . Combining the estimate (28) with the finite-time admissibility of  $C$  in the assumption a), we conclude that  $C$  is infinite-time admissible for  $A$ .  $\square$

The assumption c) of Theorem 5.3 leads us to the following strong version of the Weiss condition.

**Definition 5.4.** Let  $X$  and  $Y$  be Banach spaces, and let  $A$  be the generator of a bounded  $C_0$ -semigroup on  $X$ . An operator  $C \in \mathcal{L}(D(A), Y)$  satisfies the *strong 2-Weiss condition for  $A$*  if there exists  $\alpha > 0$  such that  $C(-A)^\alpha$  extends to an operator in  $\mathcal{L}(D(A), Y)$  and its extension satisfies the 2-Weiss condition for  $A$ .

By Proposition 2.1, if the  $C_0$ -semigroup  $(T(t))_{t \geq 0}$  is polynomially stable, then for all  $\alpha > 0$ , there exists  $\beta > 0$  such that  $\|T(t)(-A)^{-\alpha}\| = O(t^{-\beta})$  as  $t \rightarrow \infty$ , which is a faster decay rate than the one of the assumption b) in Theorem 5.3. Therefore, we obtain the following result as a direct consequence of Theorem 5.3.

**Corollary 5.5.** *Let  $X$  and  $Y$  be Hilbert spaces, and let  $A$  be the generator of a polynomially stable  $C_0$ -semigroup on  $X$ . If  $C \in \mathcal{L}(D(A), Y)$  is  $L^2$ -finite-time admissible for  $A$  and satisfies the strong 2-Weiss condition for  $A$ , then  $C$  is  $L^2$ -infinite-time admissible for  $A$ .*

The case  $X = Y = \ell^2(\mathbb{N})$  is simple but practically important as explained in [20, Remark 2.7]. Consider a diagonal operator  $A$  on  $\ell^2(\mathbb{N})$ . The resolvent condition  $\|C(-A)^\alpha R(\lambda, A)\| \leq K/\sqrt{\operatorname{Re} \lambda}$  in the strong 2-Weiss condition is transformed into the operator Carleson measure criterion introduced in [19, Definition 1.1]. To describe the equivalence more precisely, let  $(\lambda_n)_{n \in \mathbb{N}}$  be the eigenvalues of the diagonal operator  $A$  on  $\ell^2(\mathbb{N})$ . We assume that  $\operatorname{Re} \lambda_n < 0$  for all  $n \in \mathbb{N}$ . Let  $C \in \mathcal{L}(D(A), \ell^2(\mathbb{N}))$  and put  $c_n := Ce_n \in \ell^2(\mathbb{N})$  for  $n \in \mathbb{N}$ , where  $e_n$  is the  $n$ th unit vector in  $\ell^2(\mathbb{N})$ . Define  $c_n^* : \ell^2(\mathbb{N}) \rightarrow \mathbb{C}$  by  $c_n^* x = \langle x, c_n \rangle$  for  $x \in \ell^2(\mathbb{N})$ . When the operator  $C$  is represented by an infinite matrix,  $c_n$  is the  $n$ th column of  $C$ , and  $c_n c_n^*$  is an infinite matrix of rank one. For  $h > 0$  and  $\omega \in \mathbb{R}$ , the rectangle  $Q(h, \omega)$  is given by

$$Q(h, \omega) := \{\lambda \in \mathbb{C} : 0 \leq \operatorname{Re} \lambda \leq h, |\operatorname{Im} \lambda - \omega| \leq h\}.$$

Then [20, Proposition 5.3] shows that there is  $K > 0$  such that  $\|C(-A)^\alpha R(\lambda, A)\| \leq K/\sqrt{\operatorname{Re} \lambda}$  for all  $\lambda \in \mathbb{C}_+$  if and only if there exists  $M > 0$  such that for all  $h > 0$  and  $\omega \in \mathbb{R}$ ,

$$\left\| \sum_{-\lambda_n \in Q(h, \omega)} |\lambda_n|^{2\alpha} c_n c_n^* \right\| \leq Mh,$$

where the norm on the left-hand side is the operator norm on  $\ell^2(\mathbb{N})$ .

We saw in Lemma 3.4 and Proposition 3.5 that the decay rate  $\|CT(t)\| = O(1/\sqrt{t})$  as  $t \rightarrow \infty$  is equivalent to the 2-Weiss condition of  $C$  when  $Y = X$  and  $T(t)$  commutes with  $C$  for all  $t \geq 0$ . The next proposition shows that the strong 2-Weiss condition implies a slightly better decay rate  $\|CT(t)\| = O(1/\sqrt{t^{1+\beta}})$  for some  $\beta > 0$  if  $(T(t))_{t \geq 0}$  is polynomially stable.

**Proposition 5.6.** *Let  $A$  be the generator of a polynomially stable  $C_0$ -semigroup  $(T(t))_{t \geq 0}$  on a Hilbert space  $X$ . Let  $C \in \mathcal{L}(D(A), X)$  be such that  $T(t)$  commutes with  $C$  for all  $t \geq 0$ . If  $C$  satisfies the strong 2-Weiss condition for  $A$ , then the operator  $CT(t)$  extends to a bounded linear operator (also denoted by  $CT(t)$ ) on  $X$  for all  $t > 0$ , and there exist  $M > 0$  and  $\beta > 0$  such that*

$$(29) \quad \|CT(t)\| \leq \frac{M}{\sqrt{t^{1+\beta}}}$$

for all  $t > 0$ .

*Proof.* By the strong 2-Weiss condition of  $C$ , there exists  $\alpha > 0$  such that  $C(-A)^\alpha$  extends to an operator in  $\mathcal{L}(D(A), Y)$  and its extension satisfies the 2-Weiss condition for  $A$ . By Proposition 3.5, there is  $M_1 > 0$  such that

$$(30) \quad \|C(-A)^\alpha T(t)x\| \leq \frac{M_1}{\sqrt{t}} \|x\|$$

for all  $t > 0$  and  $x \in D((-A)^{1+\alpha})$ . Since  $(T(t))_{t \geq 0}$  is polynomially stable, it follows from Proposition 2.1 that there exist  $M_2 > 0$  and  $\beta > 0$  such that

$$(31) \quad \|T(t)(-A)^{-\alpha}\| \leq \frac{M_2}{\sqrt{t^\beta}}$$

for all  $t > 0$ . Combining the inequalities (30) and (31), we obtain

$$\begin{aligned} \|CT(t)x\| &= \|C(-A)^\alpha T(t/2)T(t/2)(-A)^{-\alpha}x\| \\ &\leq \frac{M_1}{\sqrt{t/2}} \|T(t/2)(-A)^{-\alpha}x\| \\ &\leq \frac{2^{(1+\beta)/2} M_1 M_2}{\sqrt{t^{1+\beta}}} \|x\| \end{aligned}$$

for all  $t > 0$  and  $x \in D(A)$ . The result follows by the density of  $D(A)$ .  $\square$

We also show that the decay rate  $\|CT(t)\| = O(1/\sqrt{t^{1+\beta}})$  for some  $\beta > 0$  leads to the strong 2-Weiss condition under some additional assumption on boundedness.

**Proposition 5.7.** *Let  $A$  be the generator of a bounded  $C_0$ -semigroup  $(T(t))_{t \geq 0}$  on a Banach space  $X$ . Let  $C \in \mathcal{L}(D(A), X)$  be such that  $T(t)$  commutes with  $C$  for all  $t \geq 0$ . If there exist constants  $M_1, M_2, \alpha, \beta > 0$  such that for all  $t > 0$ ,*

$$\|C(-A)^\alpha T(t)x\| \leq M_1 \|x\|, \quad x \in D((-A)^{1+\alpha})$$

and

$$\|CT(t)x\| \leq \frac{M_2}{\sqrt{t^{1+\beta}}} \|x\|, \quad x \in D(A),$$

then  $C$  satisfies the strong 2-Weiss condition for  $A$ .

*Proof.* Define  $\gamma := \alpha\beta/(1+\beta) \in (0, \alpha)$ . The moment inequality (see, e.g., [17, Proposition 6.6.4]) shows that there exists a constant  $M_0 > 0$  such that for all  $t \geq 0$  and  $x \in D((-A)^{1+\alpha})$ .

$$\|(-A)^\gamma CT(t)x\| \leq M_0 \|(-A)^\alpha CT(t)x\|^{\frac{2}{\alpha}} \|CT(t)x\|^{1-\frac{2}{\alpha}}.$$

Since

$$\frac{1+\beta}{2} \left(1 - \frac{\gamma}{\alpha}\right) = \frac{1}{2},$$

we have

$$(32) \quad \|(-A)^\gamma CT(t)x\| \leq \frac{M_3}{\sqrt{t}} \|x\|$$

for all  $t > 0$  and  $x \in D((-A)^{1+\alpha})$ , where  $M_3 := M_0 M_1^{\frac{\gamma}{\alpha}} M_2^{1-\frac{\gamma}{\alpha}}$ .

By assumption,  $\tilde{C} := C(I-A)^{-1} \in \mathcal{L}(X)$  commutes with  $A$ . It follows from [17, Proposition 3.1.1.f)] that  $\tilde{C}$  also commutes with  $(-A)^\gamma$ . Recall that  $D((-A)^\delta) = D((I-A)^\delta)$  for all  $\delta > 0$ ; see [17, Proposition 3.1.9.a)]. For all  $x \in D((-A)^{1+\gamma}) = D((I-A)^{1+\gamma})$ , there exists  $y \in D((-A)^\gamma) = D((I-A)^\gamma)$  such that  $x = (I-A)^{-1}y$ . Then

$$(33) \quad C(-A)^\gamma x = \tilde{C}(-A)^\gamma y = (-A)^\gamma \tilde{C}y = (-A)^\gamma Cx.$$

By (33), we obtain

$$C(-A)^\gamma R(\lambda, A)x = R(\lambda, A)(-A)^\gamma Cx$$

for all  $\lambda \in \mathbb{C}_+$  and  $x \in D((-A)^{1+\gamma})$ . The estimate (32) yields that for all  $\lambda \in \mathbb{C}_+$  and  $x \in D((-A)^{1+\alpha})$ ,

$$\begin{aligned}
 \|C(-A)^\gamma R(\lambda, A)x\| &\leq \int_0^\infty e^{-t \operatorname{Re} \lambda} \|(-A)^\gamma C T(t)x\| dt \\
 &\leq M_3 \int_0^\infty \frac{e^{-t \operatorname{Re} \lambda}}{\sqrt{t}} dt \|x\| \\
 (34) \qquad &= \frac{M_3 \Gamma(1/2)}{\sqrt{\operatorname{Re} \lambda}} \|x\|,
 \end{aligned}$$

where  $\Gamma$  is the Gamma function. By [17, Proposition 3.1.1.h)],  $D((-A)^{1+\alpha})$  is a core for  $(-A)^\gamma$ , and hence the estimate (34) holds for all  $x \in D((-A)^\gamma)$ . Since  $D((-A)^\gamma)$  is dense in  $X$ , it follows that  $C(-A)^\gamma R(\lambda, A)$  extends to a bounded linear operator on  $X$ . Therefore,  $C(-A)^\gamma$  extends to an operator in  $\mathcal{L}(D(A), X)$ . We also see from the estimate (34) that the extension of  $C(-A)^\gamma$  satisfies the 2-Weiss condition for  $A$ . Thus,  $C$  satisfies the strong 2-Weiss condition for  $A$ .  $\square$

#### REFERENCES

- [1] W. Arendt, C. J. K. Batty, M. Hieber, and F. Neubrander. *Vector-valued Laplace Transforms and Cauchy Problems*. Basel: Birkhäuser, 2001.
- [2] L. Arnold. Behavior of Kreiss bounded  $C_0$ -semigroups on a Hilbert space. *Adv. Operator Theory*, 7:1–7, 2022.
- [3] A. Bátkai, K.-J. Engel, J. Prüss, and R. Schnaubelt. Polynomial stability of operator semigroups. *Math. Nachr.*, 279:1425–1440, 2006.
- [4] C. J. K. Batty, R. Chill, and Yu. Tomilov. Fine scales of decay of operator semigroups. *J. Eur. Math. Soc.*, 18:853–929, 2016.
- [5] C. J. K. Batty and T. Duyckaerts. Non-uniform stability for bounded semi-groups on Banach spaces. *J. Evol. Equations*, 8:765–780, 2008.
- [6] A. Borichev and Yu. Tomilov. Optimal polynomial decay of functions and operator semigroups. *Math. Ann.*, 347:455–478, 2010.
- [7] M. Boukdir. On the growth of semigroups and perturbations. *Semigroup Forum*, 91:338–346, 2015.
- [8] R. Chill, L. Paunonen, D. Seifert, R. Stahn, and Yu. Tomilov. Non-uniform stability of damped contraction semigroups. To appear in *Anal. PDE*, 2019.
- [9] R. Chill and D. Seifert. Quantified versions of Ingham’s theorem. *Bull. Lond. Math. Soc.*, 48:519–532, 2016.
- [10] R. Chill, D. Seifert, and Yu. Tomilov. Semi-uniform stability of operator semigroups and energy decay of damped waves. *Philos. Trans. Roy. Soc. A*, 378:20190614, 2020.
- [11] R. Chill and Yu. Tomilov. Stability of operator semigroups: ideas and results. In *Perspectives in operator theory, Banach Center Publ., vol. 75, Polish Acad. Sci., Warsaw*, pages 71–109, 2007.
- [12] T. Eisner and H. Zwart. Continuous-time Kreiss resolvent condition on infinite-dimensional spaces. *Math. Comp.*, 75:1971–1985, 2006.
- [13] T. Eisner and H. Zwart. A note on polynomially growing  $C_0$ -semigroups. *Semigroup Forum*, 75:438–445, 2007.
- [14] K.-J. Engel and R. Nagel. *One-Parameter Semigroups for Linear Evolution Equations*. New York: Springer, 2000.
- [15] B. H. Haak. The Weiss conjecture and weak norms. *J. Evol. Equations*, 12:855–861, 2012.
- [16] B.H. Haak and P.C. Kunstmann. Weighted admissibility and wellposedness of linear systems in Banach spaces. *SIAM J. Control Optim.*, 45:2094–2118, 2007.
- [17] M. Haase. *The Functional Calculus for Sectorial Operators*. Basel: Birkhäuser, 2006.
- [18] B. Hamid, D. Adberrahim, and E. M. Omar. A direct approach to the Weiss conjecture for bounded analytic semigroups. *Czechoslovak Math. J.*, 60:527–539, 2010.
- [19] S. Hansen and G. Weiss. The operator Carleson measure criterion for admissibility of control operators for diagonal semigroups on  $l^2$ . *Syst. Control Lett.*, 16:219–227, 1991.
- [20] S. Hansen and G. Weiss. New results on the operator Carleson measure criterion. *IMA J. Math. Control Inf.*, 14:3–32, 1997.
- [21] B. Helffer and J. Sjöstrand. Improving semigroup bounds with resolvent estimates. *Integral Equations Operator Theory*, 93, Article number: 36, 2021.
- [22] R. Hoffeld, B. Jacob, and F. L. Schwenninger. Characterization of Orlicz admissibility. 2021.
- [23] B. Jacob and J. R. Partington. The Weiss conjecture on admissibility of observation operators for contraction. *Integral Equations Operator Theory*, 40:231–243, 2001.
- [24] B. Jacob and J. R. Partington. Admissibility of control and observation operators for semigroups: a survey. In *Current Trends in Operator Theory and its Applications*, pages 199–221. 2004.

- [25] B. Jacob, J. R. Partington, and S. Pott. Admissible and weakly admissible observation operators for the right shift semigroup. *Proc. Edinburgh Math. Soc.*, 45:353–362, 2002.
- [26] B. Jacob and H. Zwart. Counterexamples concerning observation operators for  $C_0$ -semigroups. *SIAM J. Control Optim.*, 43:137–153, 2004.
- [27] S. Lang. *Real and Functional Analysis, 3rd ed.* New York: Springer.
- [28] C. Le Merdy. The Weiss conjecture for bounded analytic semigroups. *J. London Math. Soc.*, 67:715–738, 2003.
- [29] Z. Liu and B. Rao. Characterization of polynomial decay rate for the solution of linear evolution equation. *Angew. Math. Phys.*, 56:630–644, 2005.
- [30] M. M. Martínez. Decay estimates of functions through singular extensions of vector-valued Laplace transforms. *J. Math. Anal. Appl.*, 375:196–206, 2011.
- [31] A. Mironchenko and C. Priour. Input-to-state stability of infinite-dimensional systems: recent results and open questions. *SIAM Rev.*, 62:529–614, 2020.
- [32] A. J. Pritchard and J. Zabczyk. Stability and stabilizability of infinite-dimensional systems. *SIAM Rev.*, 23:25–52, 1981.
- [33] J. Rozendaal, D. Seifert, and R. Stahn. Optimal rates of decay for operator semigroups on Hilbert spaces. *Adv. Math.*, 346:359–388, 2019.
- [34] J. Rozendaal and M. Veraar. Sharp growth rates for semigroups using resolvent bounds. *J. Evol. Equations*, 18:1721–1744, 2018.
- [35] W. Rudin. *Real and Complex Analysis, 3rd ed.* Singapore: McGrawHill, 1987.
- [36] D. Seifert. A Katznelson–Tzafriri theorem for measures. *Integral Equations Operator Theory*, 81:255–270, 2015.
- [37] O. J. Staffans. *Well-Posed Linear Systems.* Cambridge, UK: Cambridge Univ. Press, 2005.
- [38] Yu. Tomilov. A resolvent approach to stability of operator semigroups. *J. Operator Theory*, 46:63–98, 2001.
- [39] M. Tucsnak and G. Weiss. *Observation and Control of Operator Semigroups.* Basel: Birkhäuser, 2009.
- [40] J. van Neerven. Asymptotic behaviour of  $C_0$ -semigroups and  $\gamma$ -boundedness of the resolvent. *J. Math. Anal. Appl.*, 358:380–388, 2009.
- [41] G. Weiss. Admissibility of unbounded control operators. *SIAM J. Control Optim.*, 27:527–545, 1989.
- [42] G. Weiss. Two conjectures on the admissibility of control operators. In *Estimation and Control of Distributed Parameter Systems*, pages 367–378. Basel: Birkhäuser, 1991.
- [43] G. Weiss. A powerful generalization of the Carleson measure theorem? In *Open Problems in Mathematical Systems and Control Theory*, pages 267–272. London: Springer, 1998.
- [44] H. Zwart. Sufficient conditions for admissibility. *Syst. Control Lett.*, 54:973–979, 2005.
- [45] H. Zwart. Toeplitz operators and  $\mathcal{H}_\infty$  calculus. *J. Funct. Anal.*, 263:167–182, 2012.
- [46] H. Zwart, B. Jacob, and O. Staffans. Weak admissibility does not imply admissibility for analytic semigroups. *Syst. Control Lett.*, 48:341–350, 2003.

GRADUATE SCHOOL OF SYSTEM INFORMATICS, KOBE UNIVERSITY, NADA, KOBE, HYOGO 657-8501, JAPAN  
 Email address: wakaiki@ruby.kobe-u.ac.jp