

HYPERTRANSCENDENCE AND LINEAR DIFFERENCE EQUATIONS, THE EXPONENTIAL CASE

THOMAS DREYFUS

ABSTRACT. In this paper we study meromorphic functions solutions of linear shift difference equations in coefficients in $\mathbb{C}(x)$ involving the operator $\rho : y(x) \mapsto y(x+h)$, for some $h \in \mathbb{C}^*$. We prove that if f is solution of an algebraic differential equation, then f belongs to a ring that is made with periodic functions and exponentials. Our proof is based on the parametrized difference Galois theory initiated by Hardouin and Singer.

CONTENTS

Introduction	1
1. Difference setting	3
2. Differential setting	5
3. Irreducible Galois group	6
4. Proof of the main result	8
References	13

INTRODUCTION

Given a solution of a functional equation we might wonder if it is solution of another kind of functional equation. If the two functional equations are too different we would expect that very few functions are solutions of both functional equations. The goal of this paper is state a result in that direction. Before making the state of the art, let us state our main result and its framework.

Let ρ be the automorphism of $\mathbb{C}(x)$ defined by $\rho : y(x) \mapsto y(x+h)$ with $h \in \mathbb{C}^*$. Let $\mathcal{M}(\mathbb{C})$ be the field of meromorphic functions on \mathbb{C} . We say that $f \in \mathcal{M}(\mathbb{C})$ is differentially algebraic over $\mathbb{C}(x)$, if there exist $m \in \mathbb{N}$, $0 \neq P \in \mathbb{C}(x)[X_0, \dots, X_m]$ such that $P(f, \partial_x f, \dots, \partial_x^m f) = 0$. We say that f is differentially transcendental otherwise. Let $C_h := \{f \in \mathcal{M}(\mathbb{C}) \mid \rho(f) = f\}$ be the field of h -periodic functions. The goal of this paper is to prove:

Theorem 1. *Let $n \in \mathbb{N}^*$, and let $A \in \mathrm{GL}_n(\mathbb{C}(x))$. Let $\mathcal{Y} := (f_1, \dots, f_n)^\top \in (\mathcal{M}(\mathbb{C}))^n$ be a solution of $\rho(\mathcal{Y}) = A\mathcal{Y}$. If every f_i is differentially algebraic over $\mathbb{C}(x)$, then there exist $\ell \in \mathbb{N}^*$, $\lambda_1, \dots, \lambda_k \in \mathbb{C}$, such that for all $1 \leq i \leq n$, $f_i \in C_{\ell h}(x)[e^{\lambda_1 x}, \dots, e^{\lambda_k x}]$.*

Date: December 2, 2022.

2010 Mathematics Subject Classification. 39A06, 12H05.

This project has received funding from the ANR De rerum natura project (ANR-19-CE40-0018).

As a straightforward corollary, we find:

Corollary 2. *Let $f \in \mathcal{M}(\mathbb{C})$ such that*

$$a_0 f + \cdots + a_n \rho^n f = 0, \quad a_i \in \mathbb{C}(x), \quad a_0, a_n \neq 0.$$

If f is differentially algebraic over $\mathbb{C}(x)$, then there exist $\ell \in \mathbb{N}^$, $\lambda_1, \dots, \lambda_k \in \mathbb{C}$, such that $f \in C_{\ell h}(x)[e^{\lambda_1 x}, \dots, e^{\lambda_k x}]$.*

Proof of Corollary 2. Since ρ and ∂_x commutes, for all $i \in \mathbb{N}$, $\rho^i(f)$ is differentially algebraic over $\mathbb{C}(x)$. Then, each entries of $\mathcal{Y} := (f, \rho(f), \dots, \rho^{n-1}(f))^\top \in \mathcal{M}(\mathbb{C})^n$ is differentially algebraic over $\mathbb{C}(x)$. Since

$$\rho(\mathcal{Y}) = \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \ddots & \vdots \\ \vdots & \vdots & \ddots & \ddots & 0 \\ 0 & 0 & \cdots & 0 & 1 \\ -a_0/a_n & -a_1/a_n & \cdots & \cdots & -a_{n-1}/a_n \end{pmatrix} \mathcal{Y},$$

we conclude with Theorem 1. \square

The first proof of differential transcendence of a function is due to Hölder, see [Höl87]. The author proved that the Gamma function is differentially transcendental. His approach mainly uses the functional equation $\rho(\Gamma) = x\Gamma$ (here $h = 1$).

More recently in [BG93], it is proved that if $f \in \mathcal{M}(\mathbb{C})$ is solution of a linear ρ -equation and a linear differential equation, then there exist $\ell \in \mathbb{N}^*$, $\lambda_1, \dots, \lambda_k \in \mathbb{C}$, such that $f \in C_{\ell h}(x)[e^{\lambda_1 x}, \dots, e^{\lambda_k x}]$. The strategy of the authors is to study the singularities of the solutions of a linear ρ -equation, linear differential equation, and prove that there is an incompatibility, unless the function is sufficiently simple. Similar results were known for q -difference equations ($\rho : y(x) \mapsto y(qx)$, $q \in \mathbb{C}^*$, $|q| \neq 1$) and Mahler equations ($\rho : y(x) \mapsto y(x^p)$, $p \in \mathbb{N}_{\geq 2}$), and were proved in an unified way in [SS19].

The question of showing that a solution of a linear ρ -equation is not solution of an algebraic differential equation is much more complicated. Contrary to the linear case, the number of unknowns grows too fast to hope to predict the potential equation in a brute-force search. Another approach that fails is the study of the singularities, since a solution of an algebraic differential equation may have a very large set of singularities.

Old results were known for affine equations of order one of the form $\rho(y) = ay + b$, when ρ is the shift operator considered above, the q -difference operator or a Mahler operator, see for instance [Höl87, Moo96, Mah30, Nis84, Ran92, Ish98, Har08, HS08, Ngu12].

For higher order equations we need the Galois theory of difference equations to be able to prove similar results. More precisely, given a linear ρ -system, we may associate a Galois group of matrices that encodes the algebraic (resp. algebraic and differential) relations among the solutions. Bigger this group is, and fewer relations there are. So if we are able to prove that the Galois is sufficiently large, we might expect that the solutions are differentially transcendental. The first attempt of this strategy could be

found in [Har08] for a modern proof of Hölder's theorem of the differential transcendence of the Gamma function. This approach has been generalized into a complete theory in [HS08] and has been applied to prove that when the Galois group is big, the solutions are differentially transcendental, see [HS08, DHR18, DHR16, AS17, ADR18]. The general idea behind the latter papers is that when the solutions are differentially algebraic and the Galois group is big, then the functions are solutions of linear differential equations. We then use the results above mentioned of [SS19] to prove that they are too trivial and forces the Galois group to be small, leading to a contradiction. The Galois theory of [HS08] has been applied to a very different context to prove in [DHR18] that some generating series of walks in the quarter plane are differentially transcendental.

At this stage, we are able to prove that the solutions of linear ρ -equations are differentially transcendental or trivial only when the Galois group is big, or small (affine equations). It remains to treat the medium cases. This is the goal of [ADH21] where it is proved that when the Galois group is medium, then the ρ -equation is equivalent to a smaller one. Then an induction proof on the rank of the equation allows the authors to prove that either the solution is differentially transcendental, either it belongs to a small field. The results of [ADH21] are stated for meromorphic solution at infinity and the shift operator, for the q -difference operator, and for the Mahler operator. For the shift case and meromorphic function on \mathbb{C} which is the framework of this paper, the statement is much more complicated to set for two reasons:

- the exponential functions $\exp(\lambda x)$, $\lambda \in \mathbb{C}$, are meromorphic solutions of linear ρ -equations and linear differential equations.
- the meromorphic functions that are ℓh -periodic for some $\ell \in \mathbb{N}^*$ are solutions of linear ρ -equations, and some of them are solution of differential equations.

Then, the set of functions that are both solutions of a linear ρ -equation and differential equation is bigger than the ground field $\mathbb{C}(x)$. To avoid this problem, the authors of [ADH21] consider the situation where f is solution of a linear ρ -equation in coefficients in $\mathbb{C}(x)$ and belong to a field of meromorphic function $F \subset \mathcal{M}(\mathbb{C})$, with $\rho(F) = F$, $\partial_x(F) \subset F$, $\{f \in F \mid \rho(f) = f\} = \mathbb{C}$, and for all $\lambda_1, \dots, \lambda_k \in \mathbb{C}$, $F \cap \mathbb{C}(x, e^{\lambda_1 x}, \dots, e^{\lambda_k x}) = \mathbb{C}(x)$. Then it is proved that either $f \in \mathbb{C}(x)$, either f is differentially transcendental. Thus, Theorem 1 generalizes this result in this context since we avoid the assumptions on F . The strategy of the proof of Theorem 1 is in some sens similar to the one in the recent paper [dS22] that deals with another framework.

The paper is organized as follows. In Section 1 we give a reminder of the difference Galois theory and in Section 2, the parametrized difference Galois theory. In Section 3, we deal with difference Galois group that are irreducible and connected, and in Section 4 we complete the proof of Theorem 1.

1. DIFFERENCE SETTING

The goal of this section is to make a short review of the main results of difference algebra and difference Galois theory that will be used in this paper. We refer to [Coh65] for more details on difference algebra and [vdPS97] for

more details on difference Galois theory. In what follows, every fields are of characteristic zero, every rings are unitary.

A difference ring is a ring (R, ρ) is a ring equipped with an automorphism. We define similarly the notion of difference fields, difference algebras, etc... When no confusion arises, we will denote by R the difference ring (R, ρ) . The ring of constants is defined by $R^\rho := \{r \in R | \rho(r) = r\}$. If R is a field, R^ρ is also a field and will be called field of constants.

Example 3. If we consider the notations of the introduction, $(\mathbb{C}(x), \rho)$ and $(\mathcal{M}(\mathbb{C}), \rho)$ are difference fields, and we have $\mathbb{C}(x)^\rho = \mathbb{C}$ and $\mathcal{M}(\mathbb{C})^\rho = C_h$.

A ρ -ideal $I \subset R$ is an ideal such that $\rho(I) \subset I$. We say that the difference ring is ρ -simple if the only ρ -ideals are $\{0\}$ and I .

Let \mathbf{k} be a difference field. Let us assume that $\mathbf{C} := \mathbf{k}^\rho$ is algebraically closed. We consider the difference system

$$(1.1) \quad \rho(Y) = AY, \quad A \in \mathrm{GL}_n(\mathbf{k}).$$

A Picard-Vessiot ring extension for (1.1) over \mathbf{k} is a difference ring extension $\mathcal{R}|\mathbf{k}$ such that

- There exists $U \in \mathrm{GL}_n(\mathcal{R})$, such that $\rho(U) = AU$, such matrix is called a fundamental matrix;
- $\mathcal{R} = \mathbf{k}[U, 1/\det(U)]$;
- \mathcal{R} is a simple difference ring.

The Picard-Vessiot ring extension exists and is unique up to morphisms of \mathbf{k} - ρ -algebras.

Given a Picard-Vessiot ring extension $\mathcal{R}|\mathbf{k}$, the Picard-Vessiot extension \mathcal{Q} is the total field of fractions of \mathcal{R} . We have $\mathcal{Q}^\rho = \mathcal{R}^\rho = \mathbf{k}^\rho = \mathbf{C}$. We define the difference Galois group as the group of automorphisms of ring of \mathcal{Q} , leaving \mathbf{k} invariant and commuting with ρ , that is

$$\mathrm{Gal}(\mathcal{Q}|\mathbf{k}) = \{\sigma \in \mathrm{Aut}(\mathcal{Q}|\mathbf{k}) | \sigma \circ \rho = \rho \circ \sigma\}.$$

For any fundamental matrix $U \in \mathrm{GL}_n(\mathcal{Q})$, an easy computation shows that $U^{-1}\sigma(U) \in \mathrm{GL}_n(\mathbf{C})$ for all $\sigma \in \mathrm{Gal}(\mathcal{Q}|\mathbf{k})$. By [vdPS97, Theorem 1.13], the faithful representation

$$\begin{aligned} \mathrm{Gal}(\mathcal{Q}|\mathbf{k}) &\rightarrow \mathrm{GL}_n(\mathbf{C}) \\ \sigma &\mapsto U^{-1}\sigma(U) \end{aligned}$$

identifies $\mathrm{Gal}(\mathcal{Q}|\mathbf{k})$ with a linear algebraic subgroup $G \subset \mathrm{GL}_n(\mathbf{C})$. Choosing another fundamental matrix of solutions U leads to a conjugate representation.

Specific results for shift equations. In this paragraph, we consider the field $\mathbb{C}(x)$ equipped with the automorphism $\rho : y(x) \mapsto y(x+h)$, $h \in \mathbb{C}^*$, and give some specific results. Let $B \in \mathrm{GL}_n(\mathbb{C}(x))$ and let $V \in \mathrm{GL}_n(\mathcal{M}(\mathbb{C}))$ such that $\rho(V) = BV$ (such V exists in virtue of [Pra86]). Let \overline{C}_h be the algebraic closure of C_h and consider $\overline{C}_h(x)$ that is equipped with a structure of ρ -field with $\rho(x) = x+h$, and $\overline{C}_h(x)^\rho = \overline{C}_h$. Since $(\mathcal{M}(\mathbb{C}))^\rho = C_h \subset \overline{C}_h$, we apply [vdPS97, Corollary 1.24] to deduce the following result that will be used in the sequel.

Lemma 4. *The ring $\mathcal{R} = \overline{C}_h(x)[V, 1/\det(V)]$ is a Picard-Vessiot ring extension for $\rho(Y) = BY$ over $\overline{C}_h(x)$. The field $\mathcal{Q} = \overline{C}_h(x)(V)$ is a Picard-Vessiot extension for $\rho(Y) = BY$ over $\overline{C}_h(x)$.*

The change of variables $Z := TY$, $T \in \mathrm{GL}_n(\mathbb{C}(x))$, transforms the system $\rho(Y) = AY$ with $A \in \mathrm{GL}_n(\mathbb{C}(x))$ into $\rho(Z) = BZ$ where $B := \rho(T)AT^{-1} \in \mathrm{GL}_n(\mathbb{C}(x))$. Let $A, B \in \mathrm{GL}_n(\mathbb{C}(x))$. We will say that $[A]$ and $[B]$ are equivalent over $\mathbb{C}(x)$ if there exists $T \in \mathrm{GL}_n(\mathbb{C}(x))$ such that $B = \rho(T)AT^{-1}$. The following theorem has been proved in [vdPS97, Propositions 1.20 and 1.21].

Theorem 5. *Let $G := \mathrm{Gal}(\mathcal{Q}|\mathbb{C}(x))$. There exists $T \in \mathrm{GL}_n(\mathbb{C}(x))$ such that $\rho(T)AT^{-1} \in G(\mathbb{C}(x))$.*

2. DIFFERENTIAL SETTING

We refer to [HS08] for more details on what follows. A differential ring (R, δ) is a ring equipped with a derivation, that is an additive morphism satisfying the Leibnitz rule $\delta(fg) = f\delta(g) + \delta(f)g$. We define similarly the notion of differential fields, differential algebras, etc... The ring of δ -constants of R is defined by

$$R^\delta = \{r \in R \mid \delta(r) = 0\}.$$

If R is a field, R^δ is also a field and will be called field of constants. Let (R, δ) be a differential ring extension for the differential field (\mathbf{k}, δ) . We say that $f \in R$ is differentially algebraic over \mathbf{k} if there exist $n \in \mathbb{N}$, and $0 \neq P \in \mathbf{k}[X_0, \dots, X_n]$ such that $P(f, \delta f, \dots, \delta^n f) = 0$. We say that f is differentially transcendental over \mathbf{k} otherwise.

A (ρ, δ) -ring (R, ρ, δ) is a ring equipped with an automorphism ρ and a derivation δ that commutes. We define similarly the notion of (ρ, δ) -fields, etc...

Example 6. If we consider the notations of the introduction, $(\mathbb{C}(x), \rho, \partial_x)$ and $(\mathcal{M}(\mathbb{C}), \rho, \partial_x)$ are (ρ, ∂_x) -fields, and we have $\mathbb{C}(x)^{\partial_x} = \mathcal{M}(\mathbb{C})^{\partial_x} = \mathbb{C}$.

We say that I is a (ρ, δ) -ideal of R if I is an ideal such that $\rho(I) \subset I$ and $\delta(I) \subset I$. We say that (R, ρ, δ) is (ρ, δ) -simple if its only (ρ, δ) -ideals are $\{0\}$ and R .

Let us denote by $R\{X_1, \dots, X_n\}_\delta$ the ring of differential polynomials in the indeterminates $\delta^i X_j$. We recall that a differential field (L, δ) is called differentially closed or δ -closed if, for every set of δ -polynomials \mathcal{F} , the system of δ -equations $\mathcal{F} = 0$ has a solution in some δ -field extension of L if and only if it has a solution in L . There always exists a differential field extension that is differentially closed.

Assume that \mathbf{k} is a (ρ, δ) -field with $\mathbf{C} = \mathbf{k}^\rho$ differentially closed and let $A \in \mathrm{GL}_n(\mathbf{k})$. A (ρ, δ) -Picard-Vessiot ring extension for $\rho(Y) = AY$ over \mathbf{k} is a (ρ, δ) -ring extension $\mathcal{S}|\mathbf{k}$ such that

- There exists $U \in \mathrm{GL}_n(\mathcal{S})$;
- $\mathcal{S} = \mathbf{k}\{U, 1/\det(U)\}_\delta$;
- \mathcal{S} is a simple (ρ, δ) -ring.

The (ρ, δ) -Picard-Vessiot ring extensions exists and is unique up to morphisms of \mathbf{k} - (ρ, δ) -algebras. Given a (ρ, δ) -Picard-Vessiot ring extension $\mathcal{S}|\mathbf{k}$, the (ρ, δ) -Picard-Vessiot extension $\mathcal{Q}_\mathcal{S}$ is the total field of fractions of \mathcal{S} . We

have $\mathcal{Q}_S^\rho = \mathcal{S}^\rho = \mathbf{k}^\rho = \mathbf{C}$. We define the (ρ, δ) -Galois group as the group of automorphisms of ring of \mathcal{Q}_S , leaving \mathbf{k} invariant and commuting with ρ and δ , that is

$$\mathrm{Gal}^\delta(\mathcal{Q}_S|\mathbf{k}) = \{\sigma \in \mathrm{Aut}(\mathcal{Q}_S|\mathbf{k}) \mid \sigma \circ \rho = \rho \circ \sigma, \quad \sigma \circ \delta = \delta \circ \sigma\}.$$

For any fundamental matrix $U \in \mathrm{GL}_n(\mathcal{Q}_S)$, an easy computation shows that $U^{-1}\sigma(U) \in \mathrm{GL}_n(\mathbf{C})$ for all $\sigma \in \mathrm{Gal}^\delta(\mathcal{Q}_S|\mathbf{k})$. By [HS08, Theorem 2.6], the faithful representation

$$\begin{aligned} \mathrm{Gal}^\delta(\mathcal{Q}_S|\mathbf{k}) &\rightarrow \mathrm{GL}_n(\mathbf{C}) \\ \sigma &\mapsto U^{-1}\sigma(U) \end{aligned}$$

identifies $\mathrm{Gal}^\delta(\mathcal{Q}_S|\mathbf{k})$ with a linear differential algebraic subgroup $G \subset \mathrm{GL}_n(\mathbf{C})$, that is a group of matrices whose entries satisfy a set of non-trivial algebraic differential relations. Choosing another fundamental matrix of solutions U leads to a conjugate representation.

Specific results for shift equations. The field \mathbb{C} may be equipped trivially with a structure of differential field with the derivation $\delta := \partial_x$. Let (\mathbf{C}, δ) be a differentially closed field containing (\mathbb{C}, δ) and consider $\mathbf{C}(x)$, that is equipped with a structure of (ρ, δ) -field with $(\mathbf{C}(x))^\rho = \mathbf{C}$, $\rho(x) = x + h$ and $\delta(x) = 1$. Note that ρ and δ commutes.

Let $B \in \mathrm{GL}_n(\mathbf{C}(x))$. Given $\ell \in \mathbb{N}^*$, we may iterate the difference system $\rho(Y) = BY$ by considering $\rho^\ell(Y) = B_{[\ell]}Y$, where $B_{[\ell]} = \rho^{\ell-1}(B) \times \cdots \times B$. Let $G_{[\ell]}$ be the difference Galois group of the system $\rho^\ell(Y) = B_{[\ell]}Y$. The following lemma may be deduced from the proof [DHR16, Proposition 4.6] in the particular case where the parametric operator is the identity, and is proved in [ADH21, Proposition 4.10].

Lemma 7. *Let $B \in \mathrm{GL}_n(\mathbf{C}(x))$. Consider $L|\mathbf{C}(x)$, a (ρ, δ) -field extension such that $L^\rho = \mathbf{C}$. Let $\mathcal{Y} \in L^n$ be a nonzero vector solution of $\rho(Y) = BY$. Then, there exists $\ell \geq 1$, such that*

- *There exists a (ρ^ℓ, δ) -Picard-Vessiot extension for $\rho^\ell(Y) = B_{[\ell]}Y$, that is a field, and with fundamental solution admitting \mathcal{Y} as first column.*
- *$G_{[\ell]}$ is the connected component of the identity of G .*

For \mathbf{k} a (ρ, δ) -field, let $\mathbf{k}\langle X_1, \dots, X_n \rangle_\delta$ be the field of differential rational functions in the indeterminates $\delta^i X_j$. Let \widetilde{C}_h be a differentially closed field containing C_h . Since $(\mathcal{M}(\mathbf{C}))^\rho = C_h \subset \widetilde{C}_h$, we apply [HS08, Proposition 6.17], to deduce the following result that will be used in the sequel.

Lemma 8. *Let $B \in \mathrm{GL}_n(\mathbf{C}(x))$ and let $V \in \mathrm{GL}_n(\mathcal{M}(\mathbf{C}))$ such that $\rho(V) = BV$. The ring $\mathcal{S} = \widetilde{C}_h(x)\{V, 1/\det(V)\}_\delta$ is a (ρ, δ) -Picard-Vessiot ring extension for $\rho(Y) = BY$ over $\widetilde{C}_h(x)$. The field $\mathcal{Q}_S = \widetilde{C}_h(x)\langle V \rangle_\delta$ is a (ρ, δ) -Picard-Vessiot extension for $\rho(Y) = BY$ over $\widetilde{C}_h(x)$.*

3. IRREDUCIBLE GALOIS GROUP

Recall that we consider the system $\rho(Y) = AY$ with $A \in \mathrm{GL}_n(\mathbf{C}(x))$. By [Pra86], there exists $U \in \mathrm{GL}_n(\mathcal{M}(\mathbf{C}))$ such that $\rho(U) = AU$. Let \mathcal{Q}

be the Picard-Vessiot extension for $\rho(Y) = AY$ over $\mathbb{C}(x)$. In this section, following [ADH21, Section 5.2] we consider the case where $\text{Gal}(\mathcal{Q}|\mathbb{C}(x))$, the difference Galois group of $\rho(Y) = AY$ over $\mathbb{C}(x)$, seen as an algebraic subgroup of $\text{GL}_n(\mathbb{C})$, is irreducible and connected. Recall that an algebraic subgroup $G \subset \text{GL}_n(\mathbb{C})$ is said irreducible, if and only if for all \mathbb{C} -vector space $V \subset \mathbb{C}^n$, $G(V) \subset V$ implies that V is either $\{0\}$ or \mathbb{C}^n . The proof is very similar to [ADH21, Section 5.2], but we prefer to sketch the arguments here to be self-contained.

Proposition 9. *Let us assume that $n \geq 2$. If $\text{Gal}(\mathcal{Q}|\mathbb{C}(x))$ is irreducible and connected, then every column of U contains at least one element that is differentially transcendental over $\widetilde{C}_h(x)$.*

As a key argument, we will use the following result due to [AS17, Lemma 5.1]. It says that the (ρ, δ) -Galois group must be as big as possible when the difference Galois group has an identity component that is semisimple.

Proposition 10. *Let $B \in \text{GL}_n(\mathbb{C}(x))$. We let $\widetilde{G} \subset \text{GL}_n(\widetilde{C}_h)$ denote the difference Galois group of $\rho(Y) = BY$ over $\widetilde{C}_h(x)$, and $H \subset \text{GL}_n(\widetilde{C}_h)$ denote the (ρ, δ) -Galois group of $\rho(Y) = BY$ over $\widetilde{C}_h(x)$. If the identity component of \widetilde{G} is semisimple, then $H = \widetilde{G}$.*

Proof of Proposition 9. Let us argue by contradiction assuming that one column of the fundamental solution U whose coordinates are all differentially algebraic over $\widetilde{C}_h(x)$. By Lemma 8, $\widetilde{C}_h(x)\{U, 1/\det(U)\}_\delta$ is a (ρ, δ) -Picard-Vessiot extension for $\rho(Y) = AY$ over $\widetilde{C}_h(x)$. The difference Galois group of $\rho(Y) = AY$ over $\widetilde{C}_h(x)$ is $\text{Gal}(\mathcal{Q}|\mathbb{C}(x))(\widetilde{C}_h)$ and is therefore irreducible. Hence, same reasons as in the proof of [ADH21, Proposition 5.4] show that all entries of U are differentially algebraic over $\widetilde{C}_h(x)$. Then, $\det(U)$ is also differentially algebraic over $\widetilde{C}_h(x)$. We observe that the determinant $\det(U)$ is solution to the equation

$$(3.1) \quad \rho(y) = \det(A)y$$

and that the difference Galois group of this equation over $\mathbb{C}(x)$ is the group $\det(\text{Gal}(\mathcal{Q}|\mathbb{C}(x)))$. By [HS08, Corollary 3.4], there exist some nonzero elements $c \in \mathbb{C}$ and $g \in \mathbb{C}(x)$ such that $\det(A) = c\rho(g)/g$. Let $d \in \mathbb{C}$ such that $\exp(dh) = c^{-1/n}$ and consider the matrix $V = U \exp(dx) \in \text{GL}_n(\mathcal{M}(\mathbb{C}))$. It is solution of $\rho(Y) = BY$ with $B = c^{-1/n}A$. We have $\det(B) = \rho(g)/g$. By Lemma 8, $\widetilde{C}_h(x)\{V, 1/\det(V)\}_\delta$ is a (ρ, δ) -Picard-Vessiot ring extension for $\rho(Y) = BY$ over $\widetilde{C}_h(x)$. Let G be the difference Galois group of $\rho(Y) = BY$ over $\mathbb{C}(x)$, \widetilde{G} the difference Galois group of $\rho(Y) = BY$ over $\widetilde{C}_h(x)$, and H the (ρ, δ) -Galois group over $\widetilde{C}_h(x)$. Same reasons as in the proof of [ADH21, Proposition 5.4] shows that G is irreducible, G^0 the connected component of the identity of G is also irreducible, and there exists $\ell \geq 1$ such that the difference Galois group of $\rho^\ell(Y) = B_{[\ell]}Y$ over $\mathbb{C}(x)$ is G^0 . Furthermore, $\det(B_{[\ell]}) = \rho^\ell(g)/g$ and therefore, $G^0 \subset \text{SL}_n(\mathbb{C})$. With [ADH21, Lemma 4.2] we find that G^0 is primitive (we refer to the latter paper for the notion of

primitive groups that will not be used again in the sequel). By [SU93, Proposition 2.3], we finally obtain that G^0 is semisimple. Then, $\widetilde{G}^0 = G^0(\widetilde{C}_h)$, the identity component of \widetilde{G} , is semisimple too.

We infer from Proposition 10 that $H = \widetilde{G}$. We recall that every entries of U are differentially algebraic over $\widetilde{C}_h(x)$. Then the same holds for $V = U \exp(dx)$. Same reasons as in the proof of [ADH21, Proposition 5.4] shows that G is a finite group. Since it is connected, we deduce that $G^0 = \{I_n\}$, where we let I_n denote the identity matrix of size n . Since by assumption $n \geq 2$, this provides a contradiction with the fact that G^0 is irreducible. \square

4. PROOF OF THE MAIN RESULT

We are now ready to prove Theorem 1. Recall that $\mathcal{Y} := (f_1, \dots, f_n)^\top \in (\mathcal{M}(\mathbb{C}))^n$ is a solution of $\rho(Y) = AY$, with $A \in \text{GL}_n(\mathbb{C}(x))$. Let $C_h^\infty = \cup_{\ell=1}^\infty C_{\ell h}$ and let $R_{h,\text{exp}}$ be the ring of $C_h^\infty(x)$ -linear combination between $\{\exp(\lambda_i x), \lambda_i \in \mathbb{C}\}$. We have to prove that when all the entries of \mathcal{Y} are differentially algebraic over $\mathbb{C}(x)$, they all belong to $R_{h,\text{exp}}$. If \mathcal{Y} is zero the result is clear. So let us assume that \mathcal{Y} is nonzero. By [Pra86], there exists $U \in \text{GL}_n(\mathcal{M}(\mathbb{C}))$ such that $\rho(U) = AU$. We may assume that \mathcal{Y} is the first column of U . Let \mathcal{Q}_h be the Picard-Vessiot extension for $\rho(Y) = AY$ over $\overline{C}_h(x)$ and let $\text{Gal}(\mathcal{Q}_h|\overline{C}_h(x))$ be the difference Galois group. By Lemma 7, with $\delta = 0$, there exists $\ell \geq 1$, such that

- we have a Picard-Vessiot extension for $\rho^\ell(Y) = A_{[\ell]}Y$ over $(\overline{C}_h(x), \rho^\ell)$ that is a field;
- the fundamental solution of the Picard-Vessiot extension over $(\overline{C}_h(x), \rho^\ell)$ admits \mathcal{Y} as first column;
- the difference Galois group of $\rho^\ell(Y) = A_{[\ell]}Y$ over $\overline{C}_h(x)$ is the component of the identity of $\text{Gal}(\mathcal{Q}_h|\overline{C}_h(x))$.

Note that $(\mathbb{C}(x), \rho^\ell, \delta)$ is a (ρ^ℓ, δ) -field and $R_{\ell h, \text{exp}} \subset R_{h, \text{exp}}$ is (ρ^ℓ, δ) -ring. So, without loss of generalities, we may replace A by $A_{[\ell]}$, h by ℓh , and reduce to the case where the Picard-Vessiot extension \mathcal{Q}_h is a field, the difference Galois group over $\overline{C}_h(x)$ is connected, and the Picard-Vessiot extension for $\rho(Y) = AY$ over $\overline{C}_h(x)$ admits a fundamental solution with \mathcal{Y} as first column. Let \mathcal{Q} be the Picard-Vessiot extension for $\rho(Y) = AY$ over $\mathbb{C}(x)$ and let $\text{Gal}(\mathcal{Q}|\mathbb{C}(x))$ be the difference Galois group. Since $\text{Gal}(\mathcal{Q}_h|\overline{C}_h(x)) = \text{Gal}(\mathcal{Q}|\mathbb{C}(x))(\overline{C}_h)$, we find that $\text{Gal}(\mathcal{Q}|\mathbb{C}(x))$ is connected too.

Let us prove Theorem 1 by induction on n , the order of the equation. The global strategy of the proof will be quite similar to [ADH21], but the arguments will differ inside each steps of the proof.

- (1) Case $n = 1$ (initialization of the induction).
- (2) Reduction to the affine case when $n \geq 2$.
- (3) Case of affine equations.

Step 1: Case $n = 1$.

Let us prove Theorem 1 in the case $n = 1$.

Proposition 11. *Let $0 \neq a \in \mathbb{C}(x)$. If $f \in \mathcal{M}(\mathbb{C})$ is solution of $\rho(y) = ay$ and is differentially algebraic over $\mathbb{C}(x)$, then $f \in R_{h,\text{exp}}$.*

Proof. If $f = 0$ the result is clear. Assume that $f \neq 0$. By [HS08, Corollary 3.4], if the meromorphic solution f is differentially algebraic over $\mathbb{C}(x)$, then there exist nonzero elements $c \in \mathbb{C}$ and $d(x) \in \mathbb{C}(x)$ such that $\rho(f)/f = a = c\rho(d)/d$. Let $c' \in \mathbb{C}$ such that $\exp(c'h) = c$. We find that $c_1(x) := \frac{f(x)}{d(x)\exp(c'x)}$ is ρ -invariant. We have proved that $c_1(x) \in C_h$ and therefore $f(x) = c_1(x)d(x)\exp(c'x) \in R_{h,\text{exp}}$ as expected. \square

Step 2: Reduction to the affine case.

Let us now begin the induction step of the proof of Theorem 1. Let us fix $n \geq 2$ and assume that Theorem 1 holds for equations of order strictly less than n . Assume that each entry of \mathcal{Y} is differentially algebraic over $\mathbb{C}(x)$ and prove that \mathcal{Y} has entries in $R_{h,\text{exp}}$. By Proposition 9, $\text{Gal}(\mathcal{Q}|\mathbb{C}(x))$ is reducible. By [ADH21, Lemma 4.4], there exists $T \in \text{GL}_n(\mathbb{C}(x))$ such that $Z := TY$ is solution of a bloc system $\rho(Z) = BZ$ where $B = \begin{pmatrix} B_{1,1} & B_{1,2} \\ 0 & B_{2,2} \end{pmatrix}$, and $B_{i,i}$, $i = 1, 2$, has size $n_i < n$. Let us further assume that n_1 is minimal for this property. The goal of this step is to prove that when $n_1 > 1$ the results holds. More precisely, we want to prove the following:

Proposition 12. *If each entry of \mathcal{Y} is differentially algebraic over $\mathbb{C}(x)$ and $n_1 > 1$, then $\mathcal{Y} \in (R_{h,\text{exp}})^n$.*

Let G_1 be the difference Galois group of $\rho(Y) = B_{1,1}Y$ over $\mathbb{C}(x)$. The group G_1 is irreducible and connected, see [ADH21, Section 5.3].

Let $(g_1, \dots, g_n)^\top = T\mathcal{Y}$. The entries of \mathcal{Y} are all differentially algebraic over $\mathbb{C}(x)$ if and only if the same holds for the entries of $(g_1, \dots, g_n)^\top$. Similarly, $\mathcal{Y} \in (R_{h,\text{exp}})^n$ if and only if $(g_1, \dots, g_n)^\top \in (R_{h,\text{exp}})^n$. So it suffices to show that when all the entries of $(g_1, \dots, g_n)^\top$ are differentially algebraic over $\mathbb{C}(x)$, $(g_1, \dots, g_n)^\top \in (R_{h,\text{exp}})^n$. Let \mathcal{Y}_1 be the vector $(g_1, \dots, g_{n_1})^\top$ and \mathcal{Y}_2 be the vector $(g_{n_1+1}, \dots, g_n)^\top$. We have $\rho(\mathcal{Y}_1) = B_{1,1}\mathcal{Y}_1 + B_{1,2}\mathcal{Y}_2$ and $\rho(\mathcal{Y}_2) = B_{2,2}\mathcal{Y}_2$. By induction hypothesis we find that $\mathcal{Y}_2 \in (R_{h,\text{exp}})^{n_2}$. When all the entries of \mathcal{Y} are differentially algebraic over $\mathbb{C}(x)$, we then have

$$(4.1) \quad \mathcal{Y} \in (R_{h,\text{exp}})^n \iff \mathcal{Y}_1 \in (R_{h,\text{exp}})^{n_1}.$$

Toward the proof of Proposition 12, we are going to prove that when each entry of \mathcal{Y} is differentially algebraic over $\mathbb{C}(x)$, then the entries of \mathcal{Y}_1 belong to a certain field. There exist $\lambda_1, \dots, \lambda_k \in \mathbb{C}$ and $\ell \in \mathbb{N}^*$ such that the entries of \mathcal{Y}_2 belong to $C_{\ell h}(x)[\exp(\lambda_1 x), \dots, \exp(\lambda_k x)]$. Up to replace the ρ -equation by a ρ^ℓ -equation, we may reduce to the case where $\ell = 1$. Recall that \widetilde{C}_h is a differentially closed field containing C_h . Let \widetilde{K}_0 , be the (ρ, δ) -field generated by the entries of \mathcal{Y}_2 over $\widetilde{C}_h(x)$.

Lemma 13. *If each entry of \mathcal{Y} is differentially algebraic over $\mathbb{C}(x)$ and $n_1 > 1$, then $\mathcal{Y}_1 \in (\widetilde{K}_0)^{n_1}$.*

Proof. Assume that each entry of \mathcal{Y} is differentially algebraic over $\mathbb{C}(x)$. Let us prove the contraposy: let us assume that $\mathcal{Y}_1 \notin (\widetilde{K}_0)^{n_1}$ and let us prove that $n_1 = 1$. Recall that $\rho(U) = AU$. Let $V = TU$ that is a fundamental

solution of $\rho(Y) = BY$. By Lemma 8, $\mathcal{Q}_S = \widetilde{C}_h(x)\langle V \rangle_\delta$, is a (ρ, δ) -Picard-Vessiot extension for $\rho(Y) = BY$ over $\widetilde{C}_h(x)$. Let $H = \text{Gal}^\delta(\mathcal{Q}_S | \widetilde{C}_h(x))$ be the (ρ, δ) -Galois group of $\rho(Y) = BY$ over $\widetilde{C}_h(x)$. By the (ρ, δ) -Galois correspondence, see [HS08, Theorem 2.7], we deduce the existence of some $\sigma \in H$, such that $\sigma(g_i) = g_i$ for every i , $n_1 + 1 \leq i \leq n$ and

$$(\sigma(g_i))_{i \leq n}^\top \neq (g_i)_{i \leq n}^\top.$$

Hence $w := \mathcal{Y}_1 - \sigma(\mathcal{Y}_1)$ is a nonzero vector. Since the coordinates of \mathcal{Y}_1 are differentially algebraic over $\widetilde{C}_h(x)$ and σ belongs to the (ρ, δ) -Galois group H , the coordinates of $\sigma(\mathcal{Y}_1)$ are also differentially algebraic over $\widetilde{C}_h(x)$. Therefore the coordinates of w are differentially algebraic over $\widetilde{C}_h(x)$. Furthermore, \mathcal{Y}_1 and $\sigma(\mathcal{Y}_1)$ are both solution to the system

$$\rho(Y) = B_{1,1}Y + B_{1,2}\mathcal{Y}_2.$$

It follows that

$$\rho(w) = B_{1,1}w.$$

Since we have

$$\widetilde{C}_h = \widetilde{C}_h(x)^\rho \subset (\widetilde{C}_h(x)\langle w \rangle_\delta)^\rho \subset \widetilde{\mathcal{Q}}_S^\rho = \widetilde{C}_h,$$

we find by Lemma 7, the existence of a positive integer s and a (ρ^s, δ) -Picard-Vessiot field extension \mathcal{Q}_1 for the system $\rho^s(Y) = (B_{1,1})_{[s]}Y$ over $\widetilde{C}_h(x)$ such that w is the first column of a fundamental matrix W . Furthermore, the difference Galois group of $\rho^s(Y) = (B_{1,1})_{[s]}Y$ over $\mathbb{C}(x)$ is equal to G_1 for the latter is connected. Since G_1 is irreducible, same arguments as in the proof of Proposition 9 shows that every entries of the fundamental solution W are differentially algebraic over $\widetilde{C}_h(x)$. To apply Proposition 9 let us build a fundamental solution of $\rho^s(Y) = (B_{1,1})_{[s]}Y$ in coefficients in $\mathcal{M}(\mathbb{C})$ and whose all coordinates are differentially algebraic over $\widetilde{C}_h(x)$. By [Pra86], there exists $W_1 \in \text{GL}_{n_1}(\mathcal{M}(\mathbb{C}))$ such that $\rho^s(W_1) = (B_{1,1})_{[s]}W_1$. By Lemma 8, $\widetilde{C}_h(x)\langle W_1 \rangle_\delta$ is a (ρ^s, δ) -Picard-Vessiot extension for $\rho^s(Y) = (B_{1,1})_{[s]}Y$ over $\widetilde{C}_h(x)$. Recall that each entries of W are differentially algebraic over $\widetilde{C}_h(x)$. Since the two (ρ^s, δ) -Picard-Vessiot extensions $\widetilde{C}_h(x)\langle W \rangle_\delta$ and $\widetilde{C}_h(x)\langle W_1 \rangle_\delta$ are isomorphic, we deduce that each entries of W_1 are differentially algebraic over $\widetilde{C}_h(x)$ too. The group G_1 being connected and irreducible, Proposition 9 implies that $n_1 = 1$. This proves the result by contraposition. \square

Let K_0 , be the (ρ, δ) -field generated by the entries of \mathcal{Y}_2 over $C_h(x)$. Let us now descent from \widetilde{K}_0 to K_0 .

Lemma 14. *If $\mathcal{Y}_1 \in (\widetilde{K}_0)^{n_1}$, then $\mathcal{Y}_1 \in (K_0)^{n_1}$.*

Proof. Recall that the entries of \mathcal{Y}_1 , are meromorphic on \mathbb{C} and there exist $\lambda_1, \dots, \lambda_k \in \mathbb{C}$ such that the entries of \mathcal{Y}_2 belong to $C_h(x)[\exp(\lambda_1 x), \dots, \exp(\lambda_k x)]$. Let us write $g_{i+n_1} = \sum_j R_{i,j} \exp(\lambda_j x)$, $1 \leq i \leq n_2$, the entries of \mathcal{Y}_2 with $R_{i,j} \in C_h(x)$. Let $a \in \mathbb{C}$ such that the entries of \mathcal{Y}_1 are all analytic at a . So they belong to $\mathbb{C}[[x - a]]$. Up to

replace x by $x + a$ we may reduce to the case where $a = 0$. It suffices to prove that

$$\mathbb{C}[[x]] \cap \widetilde{C}_h(x) \left(\sum_j R_{i,j}(x) \exp(\lambda_j x) \right)_{1 \leq i \leq n_2} \subset K_0.$$

Let $f \in \mathbb{C}[[x]] \cap \widetilde{C}_h(x) \left(\sum_j R_{i,j}(x) \exp(\lambda_j x) \right)_{1 \leq i \leq n_2}$. Let $a_k \in \mathbb{C}^{\mathbb{N}}$ such that we have $f = \sum_{k \geq 0} a_k x^k$. It may also be written of the form $f = \frac{\sum_{i,j} P_i(x) R_{i,j}(x) \exp(\lambda_j x)}{\sum_{i,j} Q_i(x) R_{i,j}(x) \exp(\lambda_j x)}$, with $Q_1, \dots, Q_{n_2}, P_1, \dots, P_{n_2} \in \widetilde{C}_h[x]$, the Q_i are not all zero. Then,

$$\sum_{i,j} Q_i(x) R_{i,j}(x) \exp(\lambda_j x) \sum_{k \geq 0} a_k x^k = \sum_{i,j} P_i(x) R_{i,j}(x) \exp(\lambda_j x).$$

Erasing the denominators, we find the existence of nontrivial polynomials $S_{i,j}, T_{i,j} \in C_h[x]$ such that

$$(4.2) \quad \sum_{i,j} Q_i(x) S_{i,j}(x) \exp(\lambda_j x) \sum_{k \geq 0} a_k x^k = \sum_{i,j} P_i(x) T_{i,j}(x) \exp(\lambda_j x).$$

We claim that x is transcendental over \widetilde{C}_h . To the contrary, let $m > 0$, let $a_0, \dots, a_m \in \widetilde{C}_h$ not all zero, such that $0 = a_0 + a_1 x + \dots + a_m x^m$. Let us further assume that m is minimal for this property. Applying ρ gives $a_0 + a_1 x + \dots + a_m x^m = 0 = a_0 + a_1(x+h) + \dots + a_m(x+h)^m$. Subtracting the two relations yields a relation $0 = \sum_{j=0}^m a_j (x^j - (x+h)^j)$. of x -degree at most $m-1$. By minimality of m , the two relations are equal, which is obviously not possible unless $h = 0$ or $a_1 = \dots = a_m = 0$. This is a contradiction.

We now use $\exp(\lambda_j x) \in \mathbb{C}[[x]]$. Equating the ρ -constants coefficients in front of the x^k coefficients in both side of (4.2) shows that the coefficients of the P_i and Q_i are solutions of linear equations in coefficients in C_h . Since the equations have a nontrivial solution in \widetilde{C}_h it has a non trivial solution in C_h . We have proved the existence of not all zero $\widehat{Q}_1, \dots, \widehat{Q}_{n_2} \in C_h[x]$ and $\widehat{P}_1, \dots, \widehat{P}_{n_2} \in C_h[x]$, such that

$$\sum_{i,j} \widehat{Q}_i(x) S_{i,j}(x) \exp(\lambda_j x) \sum_{k \geq 0} a_k x^k = \sum_{i,j} \widehat{P}_i(x) T_{i,j}(x) \exp(\lambda_j x).$$

This proves that $f = \frac{\sum_{i,j} \widehat{P}_i(x) R_{i,j}(x) \exp(\lambda_j x)}{\sum_{i,j} \widehat{Q}_i(x) R_{i,j}(x) \exp(\lambda_j x)} \in K_0$ as expected. \square

Let us complete the proof of Proposition 12. Assume that the entries of \mathcal{Y} are differentially algebraic over $\mathbb{C}(x)$ and $n_1 > 1$. We have seen that $\mathcal{Y}_2 = (g_{n_1+1}, \dots, g_n)^\top \in (R_{h,\exp})^{n_2}$ and by Lemma 13, and Lemma 14, $\mathcal{Y}_1 = (g_1, \dots, g_{n_1})^\top \in (K_0)^{n_1}$. Since $T^{-1}(g_1, \dots, g_n)^\top = \mathcal{Y}$, it follows that $\mathcal{Y} \in (K_0)^n$. The following lemma will terminate the proof of Proposition 12.

Lemma 15. *If $\mathcal{Y} \in (K_0)^n$, then $\mathcal{Y} \in (R_{h,\exp})^n$.*

Proof. Recall that \mathcal{Y} is solution of $\rho(Y) = AY$ and is the first column of the fundamental solution $U \in \mathrm{GL}_n(\mathcal{M}(\mathbb{C}))$. By Lemma 4, $\mathcal{R} = \overline{C}_h(x)[U, 1/\det(U)]$ is a Picard-Vessiot ring extension for $\rho(Y) = AY$ over $\overline{C}_h(x)$. Let us consider $\lambda_1, \dots, \lambda_k \in \mathbb{C}$, such that the entries of

\mathcal{Y}_2 belong to $C_h(x)[e^{\lambda_1 x}, \dots, e^{\lambda_k x}]$. Let $W = \text{Diag}(e^{\lambda_1 x}, \dots, e^{\lambda_k x})$. By Lemma 4, $\mathcal{R}_1 = \overline{C_h(x)}[W, 1/\det(W)]$, is a Picard-Vessiot ring extension for $\rho(Y) = \text{Diag}(\lambda_1 h, \dots, \lambda_k h)Y$ over $\overline{C_h(x)}$. The following idea has been suggested by Charlotte Hardouin. Consider the ideal

$$I_1 = \{s \in \mathcal{R}_1 \mid s\mathcal{Y} \in (\mathcal{R}_1)^n\}.$$

Let us prove that it is a ρ -ideal. Let $s \in I_1$. We have

$$\rho(s)\mathcal{Y} = \rho(s)A^{-1}A\mathcal{Y} = A^{-1}\rho(s\mathcal{Y}).$$

With $s\mathcal{Y} \in (\mathcal{R}_1)^n$ we therefore deduce that we have $A^{-1}\rho(s\mathcal{Y}) \in (\mathcal{R}_1)^n$. Hence $\rho(s) \in I_1$. This shows that I_1 is a ρ -ideal. Since \mathcal{R}_1 is a simple ρ -ring, and $I_1 \neq (0)$, we deduce that $1 \in I_1$, proving that \mathcal{Y} has its entries in $\overline{C_h(x)}[e^{\lambda_1 x}, \dots, e^{\lambda_k x}]$. A straightforward adaptation of the proof of Lemma 14, shows that $\mathcal{Y} \in (C_h(x)[e^{\lambda_1 x}, \dots, e^{\lambda_k x}])^n$. \square

Step 3: Affine case.

Let us finish the proof of Theorem 1. Let us recall that we assume that the entries of \mathcal{Y} are differentially algebraic over $\mathbb{C}(x)$ and we have to prove that $\mathcal{Y} \in (R_{h,\text{exp}})^n$. By Proposition 12, it remains to consider the case where $n_1 = 1$. By (4.1) it suffices to show that $g_1 \in R_{h,\text{exp}}$. Recall that g_1 is a differentially algebraic solution of the affine equation $\rho(g_1) = ag_1 + b$ where $a = B_{1,1} \in \mathbb{C}(x)$ and $b = B_{1,2}(g_2, \dots, g_n)^\top \in R_{h,\text{exp}}$. Let \widetilde{C}_h be a differentially closed field containing C_h . By [HS08, Proposition 3.8], the (ρ, δ) -Galois group of $\rho(y) = ay$ over $\widetilde{C}_h(x)$ is not maximal. By [Pra86], let $h_0 \in \mathcal{M}(\mathbb{C})$ be a nonzero solution of $\rho(y) = ay$. In virtue of Lemma 8, $\widetilde{C}_h(x)\langle h_0 \rangle_\delta$ is a (ρ, δ) -Picard-Vessiot extension for $\rho(Y) = aY$ over $\widetilde{C}_h(x)$. With [HS08, Proposition 6.26], h_0 is differentially algebraic over $\widetilde{C}_h(x)$. By a straightforward adaptation of the proof of [HS08, Corollary 3.4], we find that there exist nonzero $c \in \mathbb{C}$ and $g \in \mathbb{C}(x)$ such that $a = c\rho(g)/g$. Let $c' \in \mathbb{C}$ such that $e^{c'h} = c$. We have $g_1 \in R_{h,\text{exp}}$ if and only if $\hat{g} := \frac{g_1}{\exp(c'x)g} \in R_{h,\text{exp}}$. We have $\rho(\hat{g}) = \hat{g} + \frac{b}{\rho(\exp(c'x)g)}$. The latter is solution of an equation of the form $\rho(y) = y + b'$ where $b' := \frac{b}{\rho(\exp(c'x)g)} \in R_{h,\text{exp}}$. So without loss of generalities, we may reduce to the case where $a = 1$. Then, g_1 is solution of $\rho(y) = y + b$ where $b \in R_{h,\text{exp}}$. Let $\ell \in \mathbb{N}^*$ and $\lambda_1, \dots, \lambda_k \in \mathbb{C}$, such that $b \in C_{\ell h}(x)[\exp(\lambda_1 x), \dots, \exp(\lambda_k x)]$. We have $\rho^\ell(g_1) = g_1 + b_{[\ell]}$, with $b_{[\ell]} := b + \dots + \rho^{\ell-1}(b) \in C_{\ell h}(x)[\exp(\lambda_1 x), \dots, \exp(\lambda_k x)]$. Since $R_{\ell h,\text{exp}} \subset R_{h,\text{exp}}$ is (ρ^ℓ, δ) -ring, we may replace the ρ -equation by a ρ^ℓ -equation to reduce to the case $\ell = 1$. By [HS08, Proposition 3.1], there exists a linear differential operator in coefficients in \widetilde{C}_h and $g \in \text{Frac}(\widetilde{C}_h(x)[\exp(\lambda_1 x), \dots, \exp(\lambda_k x)])$, such that $L(b) = \rho(g) - g$. Consider $W = \text{Diag}(\exp(\lambda_1 x), \dots, \exp(\lambda_k x))$. The field \widetilde{C}_h is algebraically closed. By a straightforward adaptation of Lemma 4, $\mathcal{R}_1 = \widetilde{C}_h(x)[W, 1/\det(W)]$, is a Picard-Vessiot ring extension for $\rho(Y) = \text{Diag}(\lambda_1 h, \dots, \lambda_k h)Y$ over $\widetilde{C}_h(x)$. Consider the ideal

$$J_1 = \{s \in \mathcal{R}_1 \mid sg \in \mathcal{R}_1\}.$$

Let us prove that it is a ρ -ideal. Let $s \in J_1$. We have $\rho(sg) \in J_1$. But $\rho(sg) = \rho(s)(g + L(b))$ and $\rho(s)L(b) \in \mathcal{R}_1$, proving that $\rho(s)g \in \mathcal{R}_1$. Hence

$\rho(s) \in \mathcal{R}_1$ showing that J_1 is a ρ -ideal. Since \mathcal{R}_1 is a simple ρ -ring, and by construction $J_1 \neq (0)$, we deduce that $1 \in J_1$ and therefore

$$g \in \widetilde{C}_h(x)[\exp(\lambda_1 x), \dots, \exp(\lambda_k x)].$$

Let us write $b = \sum b_i \exp(\lambda_i x)$, $g = \sum \hat{g}_i \exp(\lambda_i x)$, $b_i, \hat{g}_i \in \widetilde{C}_h(x)$, $\lambda_i \in \mathbb{C}$. Then we find equations of the form $L_i(\widetilde{b}_i) = h\lambda_i \rho(\hat{g}_i) - \hat{g}_i$, where L_i is a linear differential operator in coefficients in \widetilde{C}_h . By [HS08, Lemma 6.4], we have the existence of $\hat{g}_i \in \widetilde{C}_h(x)$ such that $b_i = h\lambda_i \rho(\hat{g}_i) - \hat{g}_i$. Then, there exists $g' \in \text{Vect}_{\widetilde{C}_h(x)}[e^{\lambda_1 x}, \dots, e^{\lambda_k x}]$, such that $b = \rho(g') - g'$. Equating the constant coefficients yields to the existence of $g'' \in R_{h,\text{exp}}^1$ such that $b = \rho(g'') - g''$. Recall that $b = \rho(g_1) - g_1$. We then find that $g_1 - g''$ is ρ -invariant. Hence, $g_1 - g'' \in C_h$ and therefore $g_1 \in R_{h,\text{exp}}$. This completes the proof.

REFERENCES

- [ADH21] Boris Adamczewski, Thomas Dreyfus, and Charlotte Hardouin. Hypertranscendence and linear difference equations. *Journal of the American Mathematical Society*, 34(2):475–503, 2021.
- [ADR18] C. E. Arreche, T. Dreyfus, and J. Roques. On the differential transcendence of the elliptic hypergeometric functions. *arXiv preprint arXiv:1809.05416*, 2018.
- [AS17] C. E. Arreche and M. F. Singer. Galois groups for integrable and projectively integrable linear difference equations. *J. Algebra*, 480:423–449, 2017.
- [BG93] J.-P. Bézivin and F. Gramain. Solutions entières d’un système d’équations aux différences. *Ann. Inst. Fourier*, 43:791–814, 1993.
- [Coh65] R. M. Cohn. *Difference Algebra*. Interscience Publishers John Wiley & Sons, New York-London-Sydney, 1965.
- [DHR16] T. Dreyfus, C. Hardouin, and J. Roques. Functional relations of solutions of q -difference equations. *arXiv preprint arXiv:1603.06771*, 2016.
- [DHR18] T. Dreyfus, C. Hardouin, and J. Roques. Hypertranscendence of solutions of Mahler equations. *J. Eur. Math. Soc. (JEMS)*, 20:2209–2238, 2018.
- [DHRS18] T. Dreyfus, C. Hardouin, J. Roques, and M. F. Singer. On the nature of the generating series of walks in the quarter plane. *Invent. Math.*, 213:139–203, 2018.
- [dS22] Ehud de Shalit. Algebraic independence and difference equations over elliptic function fields. *arXiv preprint arXiv:2207.13377*, 2022.
- [Har08] C. Hardouin. Hypertranscendence des systèmes aux différences diagonaux. *Compos. Math.*, 144:565–581, 2008.
- [Höl87] O. Hölder. Über die Eigenschaft der Gammafunction keiner algebraischen Differentialgleichung zu genügen. *Math. Ann.*, 28:1–13, 1887.
- [HS08] C. Hardouin and M. F. Singer. Differential galois theory of linear difference equations. *Math. Ann.*, 342:333–377, 2008.
- [Ish98] K. Ishizaki. Hypertranscendence of meromorphic solutions of a linear functional equation. *Aequationes Math.*, 56:271–283, 1998.
- [Mah30] K. Mahler. Arithmetische Eigenschaften einer Klasse transzendental-transzendenter Funktionen. *Math. Z.*, 32:545–585, 1930.
- [Moo96] E. H. Moore. Concerning transcendently transcendental functions. *Math. Ann.*, 48:49–74, 1896.
- [Ngu12] P. Nguyen. *Équations de Mahler et hypertranscendance*. PhD thesis, Institut de Mathématiques de Jussieu, 2012.
- [Nis84] Ke. Nishioka. A note on differentially algebraic solutions of first order linear difference equations. *Aequationes Math.*, 27:32–48, 1984.
- [Pra86] C. Praagman. Fundamental solutions for meromorphic linear difference equations in the complex plane, and related problems. *J. Reine Angew. Math.*, 369:101–109, 1986.

- [Ran92] B. Randé. Équations fonctionnelles de Mahler et applications aux suites p-régulières. *Thèse de l'Université Bordeaux 1*, 1992.
- [SS19] R. Schäfke and M. Singer. Consistent systems of linear differential and difference equations. *J. Eur. Math. Soc. (JEMS)*, 21:2751–2792, 2019.
- [SU93] M. F. Singer and F. Ulmer. Galois groups of second and third order linear differential equations. *J. Symbolic Comput.*, 16:9–36, 1993.
- [vdPS97] M. van der Put and M. F. Singer. *Galois Theory of Difference Equations*, volume 1666 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1997.

INSTITUT DE RECHERCHE MATHÉMATIQUE AVANCÉE, UMR 7501 UNIVERSITÉ DE STRASBOURG ET CNRS, 7 RUE RENÉ DESCARTES 67084 STRASBOURG, FRANCE

Email address: `dreyfus@math.unistra.fr`