

# Geodesic Bounds of Hyperbolic Link Complements in Hyperbolic 3-Manifold

Buddha Dev Ghosh\*

*Department of Mathematics, Indian Institute of Technology Guwahati, Guwahati-781039, India*

---

## Abstract

Let  $M$  be a compact hyperbolic 3-manifold with volume  $V$ . Let  $L$  be a link such that  $M \setminus L$  is hyperbolic. For any hyperbolic link  $L$  in  $M$ , in this article we try to establish an upper bound of the length of  $n^{\text{th}}$  shortest closed geodesic of  $M \setminus L$  in terms of  $V$ .

*Keywords:* Hyperbolic 3-manifolds, Geodesics, Systole length.

---

## 1. Introduction

In the beginning, we recall that systole of a Riemannian  $n$  manifold  $M$  is shortest closed geodesic on  $M$ . In this article, we restrict ourselves for the case at  $n = 3$ . A finite volume hyperbolic 3-manifold is a complete Riemannian 3 manifold with sectional curvature  $-1$ . A link  $L$  is said to be a hyperbolic link if  $M \setminus L$  is hyperbolic.

The starting point for this article is the following result which gives an upper bound of systole length for certain class of hyperbolic 3-manifolds.

**Theorem 1.** (Adams-Reid)[2] *If  $M$  is a closed orientable 3-manifold that does not admit any Riemannian metric of negative curvature, then for any hyperbolic link  $L \subset M$  the systole length of  $M \setminus L$  is less than or equal to 7.3553...*

Firstly, we try to recall the famous result established by Lakeland and Grant[6] in 2014. Here, the authors proved that, if  $M$  is a closed, orientable 3-manifold with volume  $V$ , and suppose  $L$  is a hyperbolic link in  $M$ , then the systole length bound of  $M - L$  bounded above by  $\log((\sqrt{2}(C_0V)^{2/3} + 4\pi^2)^2 + 8)$ . They deduced that, if  $M$  does not admit hyperbolic structure then  $V = 0$ . In particular. Now if we put  $V = 0$  on the above expression, we get the bound for systole length approximately 7.3566.., which has been already proved by Adams and Reid Theorem 1 for non-hyperbolic cases. Also, set  $C_0 = \frac{\sqrt{3}}{2V_0}$ , where  $V_0$  is the volume of regular ideal tetrahedron.

In our article,  $\text{Vol}(M)$  refers to the volume of the manifold  $M$ .

**Theorem 2.** (Lakeland and Grant)[6] *If  $M$  be a closed orientable 3-manifold and  $V = \text{Vol}(M)$ , then for any hyperbolic link  $L \subset M$ , the systole length of  $M \setminus L$  is given by  $\log((\sqrt{2}(C_0V)^{2/3} + 4\pi^2)^2 + 8)$ .*

Sreekrishna Palaparathi [5] showed that an upper bound for the length of shortest closed geodesics in a hyperbolic knot or link complement in closed 3-manifolds, which do not admit any Riemannian metric of negative curvature.

---

\*Corresponding author

*Email address:* ghosh176123006@iitg.ac.in (Buddha Dev Ghosh)

**Theorem 3.** (Sreekrishna Palaparthi)[5] *Let  $M$  be a closed 3 manifold which does not admit any Riemannian metric of negative curvature. Let  $L$  be a link in  $M$ , whose complement in  $M$  admits a complete hyperbolic metric of finite volume. Then the length of the  $n^{\text{th}}$  shortest closed geodesic in  $M$  is bounded above by  $Re(2\cosh^{-1}(1 + (n + 1)18i))$ .*

From the aforementioned discussions, it is clear that the upper bounds of  $n^{\text{th}}$  shortest closed geodesic of hyperbolic link complement of compact hyperbolic 3 manifolds are not known yet. In this article, we try to focus on the previously mentioned problem. Hereby we state the main result which is given as follows:

**Theorem 4.** *If  $M$  is a orientable compact hyperbolic 3-manifold, and  $L$  is a hyperbolic link in  $M$ , and volume of  $M$  is  $V$ . Let  $l_n$  represents the shortest closed  $n^{\text{th}}$  geodesic length of  $M \setminus L$ , then*

$$l_n \leq \log \left[ \left( \frac{2\sqrt{n}C_0^{2/3}V^{2/3}}{\sqrt{3}} + 2\pi\sqrt{n} \right)^4 + 8 \right]. \quad (1)$$

## 2. Preliminaries

A hyperbolic 3-space is simply connected Riemannian 3-manifold of constant curvature  $-1$ . Here we will work on the upper half space model. In this model hyperbolic 3 space is defined to be the set of points in upper half space i.e.  $\mathbb{H}^3 = \{(z, t) \in \mathbb{C} \times \mathbb{R} : t > 0\}$  equipped with the Riemannian metric

$$ds^2 = \frac{|dz|^2 + dt^2}{t^2}. \quad (2)$$

Boundary of  $\mathbb{H}^3$  is defined by  $\partial\mathbb{H}^3 = \{(z, t) \in \mathbb{C} \times \mathbb{R} : t = 0\} \cup \{\infty\}$ . With respect to above metric, the all geodesics in  $\mathbb{H}^3 = \{(z, t) \in \mathbb{C} \times \mathbb{R} : t > 0\}$  are vertical lines and semicircle perpendicular to the boundary of  $\mathbb{H}^3$ . Totally geodesic planes are all vertical planes and hemispheres orthogonal to the boundary of  $\mathbb{H}^3$ .

Let us define  $Iso^+(\mathbb{H}^3)$  to be the set of all orientation preserving isometry on  $\mathbb{H}^3$ . If we take a Möbius transformation from  $PSL(2, \mathbb{C})$  it can be extended uniquely to a orientation preserving isometry of  $\mathbb{H}^3$ .

Suppose  $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in PSL(2, \mathbb{C})$ , and the extension of  $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$  is  $T : \mathbb{H}^3 \rightarrow \mathbb{H}^3$

$$T(z, t) = \begin{cases} \left( -\frac{\overline{z + d/c}}{c^2(|z + d/c|^2 + t^2)} + \frac{a}{c}, \frac{t}{c^2(|z + d/c|^2 + t^2)} \right) & c \neq 0 \\ \left( \frac{a}{c}(z + b/a), \left| \frac{a}{d} \right| t \right) & c = 0 \end{cases}$$

So, without loss any generality we may assume that  $Iso^+(\mathbb{H}^3) = PSL(2, \mathbb{C})$  and  $PSL(2, \mathbb{C})$  acts on the boundary of  $\mathbb{H}$  via Möbius transformations.

A 3-manifold is said to be hyperbolic if it admits a complete Riemannian metric with  $-1$  constant curvature. A discrete subgroup  $\Gamma$  of  $PSL(2, \mathbb{C})$  is said to be Kleinian group. Now if  $M$  is complete hyperbolic 3 manifold with finite volume then by Mostow-Prasad rigidity theorem the Riemannian metric with constant  $-1$  curvature is unique.

Any hyperbolic 3 manifold obtained by the quotient  $\mathbb{H}^3/\Gamma$ , where  $\Gamma$  is torsion-free Kleinian group acting on  $\partial\mathbb{H}^3$  via Möbius transformations. Let  $M = \mathbb{H}^3/\Gamma$ , be a hyperbolic 3-manifold, then  $\Gamma$  is isomorphic to  $\pi_1(M)$ . Let  $M = \mathbb{H}^3/\Gamma$  be 3-manifold which is non compact and finite volume. Let  $M$  consist finite number of cusps which is homeomorphic to  $T^2 \times [0, \infty)$ . Lifting of

the cusp in  $\mathbb{H}^3$  is a countable collection of disjoint horoballs covering the cusp in  $\mathbb{H}^3$ . Horoballs are equivalent under the action of  $\Gamma$  on  $\mathbb{H}^3$ .

A cusp is maximal if there is no larger cusp containing it. Which occurs exactly when the cusp is tangent to itself at one or more points. Maximal cusp lifts to countably infinite set of horoball in the upper half space model with disjoint interiors and some point of tangency on their boundaries. Each horoball touches the boundary of upper half space at a single point. These points are called the center of the horoballs. We choose the point at  $\infty$  to be the center of horoball denoted  $\mathbb{H}_\infty$  that covers the maximal cusp. We will normalize so that the boundary of  $\mathbb{H}_\infty$  horoball is a plane  $t = 1$ . A hyperbolic 3-manifold contains cusps is called cusp hyperbolic 3-manifold. Since the maximal cusps are self touches inside the manifold. In the covering space lift of the cusps i.e. the boundary of the horoballs are tangent itself. In particular, there must be horoballs with a tangent to the plane  $t = 1$ , which have a radius of  $1/2$  and are called full-sized horoballs.

### 2.1. Geodesic Spectrum

Let  $M$  be finite volume hyperbolic 3 manifold. Then there exists a torsion-free Kleinian group  $\Gamma$  such that  $M$  is homeomorphic to  $\mathbb{H}^3/\Gamma$ . Also  $\Gamma$  is isomorphic to  $\pi_1(M)$ . As  $\Gamma$  is discrete so  $\pi_1(M)$  is countably infinite. Therefore, each free homotopy loop class in  $\pi_1(M)$  contains a unique closed geodesic representative. Hence the number of closed geodesic is countable. The lengths of closed geodesics in  $M$  forms a discrete subset of  $\mathbb{R}$ . Let  $L = \{l_1, l_2, \dots, l_n, \dots\}$  be the set of all lengths of closed geodesics in  $M$ . As  $L$  is discrete we can arrange the length spectrum by increasing order, like  $l_1 \leq l_2 \leq \dots \leq l_n, \dots$ . Thus the systole length of  $M$  is  $l_1$ .

Let  $M$  be complete hyperbolic 3 manifold with at least one cusp. Let  $N = \mathbb{H}^3/\Gamma$  where  $\Gamma$  is discrete subgroup of  $PSL(2, \mathbb{C})$  isomorphic to  $\pi_1(N)$ . We can conjugate  $\Gamma$  such a way that the stabilizer of  $\infty$  which we denote  $\Gamma_\infty$  will be parabolic subgroup of  $\Gamma$  which is isomorphic to  $\mathbb{Z} \oplus \mathbb{Z}$ . If  $l$  is the minimal translation length we can always take one generator of  $\Gamma_\infty$  is

$$\beta = \begin{pmatrix} 1 & l \\ 0 & 1 \end{pmatrix}.$$

In this article we will always assume that  $0 \in \partial\mathbb{H}^3$  and  $\infty \in \partial\mathbb{H}^3$  are the parabolic fixed points of  $\Gamma$ .

### 2.2. Dehn filling

Let  $M$  be a 3-manifold with torus boundary component  $\partial M$  homeomorphic to torus  $T^2$ , therefore  $\pi_1(\partial M)$  isomorphic to the group  $\mathbb{Z} \oplus \mathbb{Z}$ . Let  $a$  and  $b$  be two generators of  $\pi_1(\partial M)$  i.e.  $\pi_1(\partial M) = \langle a, b \rangle$ , let  $s$  be an isotopy class of essential simple closed curve in the boundary torus  $\partial M$ , so  $s = a^m b^n$ ,  $s$  also called slope on  $\partial M$ . Now let  $P = S^1 \times D^2$  a solid torus and  $\mu$  be the meridian of  $\partial P$ , we form a closed 3-manifold by  $s$ -**Dehn filling** denoted by  $M(s)$  by attaching  $P$  on  $M$  identifying  $\phi : \partial P \rightarrow \partial M$  such that  $\phi(\mu) = s$ . More precisely,  $M(s) = M \sqcup P/x \sim \phi(x)$ . Suppose  $M$  has  $k$  disjoint torus boundary components, let  $s_1, s_2, \dots, s_k$  are the slopes corresponding boundary components, Dehn fill along the slope  $s_1, s_2, \dots, s_k$  denoted by  $M(s_1, s_2, \dots, s_k)$ .

Let  $M$  be a cusp hyperbolic 3 manifold with cusp  $C$  then the boundary of  $C$  admits a Euclidean Structure,  $s$  be any essential simple closed curve in the boundary of  $C$  then  $s$  is isotopic to a geodesic with well defined Euclidean length such a length is called *slope length*.

Let  $M$  be a cusp hyperbolic 3-manifold with cusps  $C_1, C_2, \dots, C_k$ . It is not always possible that after Dehn Filling the manifold  $M(s_1, s_2, \dots, s_k)$  will be closed hyperbolic 3-manifold,

when it will hyperbolic depends on the slope lengths of  $s_1, s_2, \dots, s_k$ . The following theorem provides a sufficient condition when  $M(s_1, s_2, \dots, s_k)$  is hyperbolic which is known as Gro-mov–Thurston  $2\pi$ -theorem, theorem stated as follows:

**Theorem 5.** (*2 $\pi$ -Theorem*) *Let  $M$  be a cusped hyperbolic 3-manifold with  $n$  cusps. Let  $C_1, C_2, \dots, C_k$  be disjoint cusp tori for the  $n$  cusps of  $M$ , and  $s_i$  a slope on  $C_i$  represented by a geodesic  $\alpha_i$  whose length in the Euclidean metric on  $C_i$  is greater than  $2\pi$ , for each  $i \in \{1, 2, \dots, k\}$ . Then  $M(s_1, s_2, \dots, s_k)$  admits a metric of negative curvature.*

This slope length bound in  $2\pi$  theorem was later improved from  $2\pi$  to 6 by Ian Agol[4] and Marck Lacknby [1] independently.

Let  $M$  be hyperbolic 3- manifold with  $k$  cusps torus, let  $s_1, s_2, \dots, s_k$  slopes on the cusp torus and let  $l_{min}$  is the minimum of all slope length of  $s_1, s_2, \dots, s_k$  then lower bound of the volume of the volume under Dehn filling given by following theorem of D.Futer et al. [3]

**Theorem 6.** *Let  $M$  be a complete, finite volume hyperbolic manifold with cusps. Suppose  $C_1, C_2, \dots, C_n$  are disjoint embedded cusps with slopes  $s_j$  on  $C_j$  such that a geodesic representative of  $s_j$  on  $\partial C_j$  has length strictly greater than  $2\pi$ . Denote the minimal slope length  $l_{min}$  then the Dehn filled manifold  $M(s_1, s_2, \dots, s_n)$  is hyperbolic and with*

$$\text{Vol}(M(s_1, s_2, \dots, s_n)) \geq \left(1 - \left(\frac{2\pi}{l_{min}}\right)^2\right)^{\frac{3}{2}} \text{Vol}(M).$$

We finished this section by providing two theorems by Lakeland and Leininger [6]. Later we will use this theorems to prove in our main result, theorems stated as follows:

**Theorem 7.** *A loxodromic element of  $PSL(2, \mathbb{C})$  with trace of modulus bounded above by  $R$  has translation length at most  $\log(R^2 + 4)$ .*

**Theorem 8.** *Suppose the Kleinian group  $\Gamma$  is torsion-free, and  $\Gamma_\infty$  has co-area at most  $V_c$  when viewed as acting by translation in  $\Gamma_\infty$  on  $\mathbb{C}$  (or equivalently acting on  $H_\infty$ ), and that the minimal parabolic translation length  $l > 2\pi$ , Then  $2\pi < l \leq \sqrt{\frac{4V_c}{\sqrt{3}}}$ .*

Now we have enough information to proof our main theorem. So, in the next section we are going to proof our main result.

### 3. Proof of the Main Result

Before going to the main result, we will prove some lemmas. Using these lemmas, we will prove our main result.

**Lemma 9.** *Let  $M$  be a finite volume hyperbolic 3 manifold with at least one cusp and let  $M = \mathbb{H}^3/\Gamma$  where  $\Gamma$  discrete subgroup of  $PSL(2, \mathbb{C})$ . Assume that in a maximal cusp torus, there is non trivial curve corresponding to a parabolic isometry of length  $l > 2\pi$ . Then each  $n \geq 1$  there exist a loxodromic element whose trace modules less than equal to  $\sqrt{l^4 n^2 + 4}$ .*

*Proof.* Let  $\Gamma$  is the image of faithful representation of  $\pi_1(M, b)$  where  $b$  is base point at the point of tangency of maximal cusp in  $M$  and  $\Gamma$  is a discrete subgroup of  $PSL(2, \mathbb{C})$ . We arrange maximal cusp so that point of tangency between the horoballs  $H_0$  and  $H_\infty$  at  $(0, 0, 1)$  (standard form for maximal cusp) and pre-image of  $b$  contain the point  $(0, 0, 1)$  under covering projection. Let  $x$  be the lift of meridian curve  $c$  (base at  $b$ ) to the horosphere  $H_\infty$  with beginning point at  $(0, 0, 1)$  and final end point at  $(l, 0, 1)$ . There is lift  $y$  of non trivial curve parallel to  $c$  on the cusp boundary with the same base point  $b$ , the lift lying on  $H_0$  having same starting point  $(0, 0, 1)$  and end point some  $p'$ . Let  $\beta$  and  $\gamma$  be the corresponding isometries of  $x$  and  $y$  in  $\Gamma$ . We may assume without any loss the angle between  $x$  and  $y$  at  $(0, 0, 1)$  at most  $\pi/2$ . The length of the geodesic corresponding loxodromic element  $\beta^{-1}\gamma$  passing through the point  $(l, 0, 1)$  and  $p$  will be maximum when angle between  $x$  and  $y$  is  $\pi/2$ . So, we choose the isometry  $\beta^{-1}\gamma$  when  $x$  and  $y$  perpendicular we can represent the parabolic isometries  $\beta$  and  $\gamma$  by

$$\beta^{-1} = \begin{pmatrix} 1 & l \\ 0 & 1 \end{pmatrix} \text{ and } \gamma = \begin{pmatrix} 1 & 0 \\ il & 1 \end{pmatrix}$$

Now for each  $n \in \mathbb{N}$ ,  $(\beta^{-1})^n\gamma$  are distinct and all are loxodromic, we can easily verify that  $|\text{tr}(\beta^{-1})^n\gamma| > 2$ .

$$\alpha_n = (\beta^{-1})^n\gamma = \begin{pmatrix} 1 + inl^2 & nl \\ il & 1 \end{pmatrix}$$

$|\text{tr}(\alpha_n)| = \sqrt{l^4n^2 + 4}$ , hence for each  $n \in \mathbb{N}$  we get a loxodromic elements whose trace modules less than  $\sqrt{l^4n^2 + 4}$ .  $\square$

Here we note that geodesic corresponding  $(\beta^{-1})^n\gamma$  all are distinct so we can rank the length spectrum of the corresponding geodesic by  $l_1 \leq l_2 \leq l_3 \dots$

**Lemma 10.** *Let  $N = \mathbb{H}^3/\Gamma$  be a non compact hyperbolic 3 manifold with finite volume where  $\Gamma$  is a discrete subgroup of  $PSL(2, \mathbb{C})$  and  $N = M \setminus L$  where  $M$  is a closed orient-able 3 manifold and  $L$  is hyperbolic link in  $M$  and  $\text{vol}(N) = V$ . Then for each  $n \geq 1$ ,  $\Gamma$  has loxodromic element with trace modules less than or equal to  $\sqrt{\frac{16n^2}{9}V_c^{8/3} + 4}$  where  $V_c = C_0V$ .*

*Proof.* By lemma (9) for each  $n \geq 1$  we can always find loxodromic elements  $c_n$  in  $\Gamma$  such that  $|\text{tr}(c_n)| \leq \sqrt{l^4n^2 + 4}$ .

Using the bound in (8), we have  $|\text{tr}(c_n)| = \sqrt{l^4n^2 + 4}$ . Then using the fact that  $V_c > 1$ , we obtain

$$|\text{tr}(c_n)| \leq \sqrt{\frac{16n^2V_c^2}{9} + 4} \tag{3}$$

and using the fact  $V_c > 1$  we get (from (8)).

$$\sqrt{\frac{16n^2V_c^2}{9} + 4} \leq \sqrt{\frac{16n^2V_c^{8/3}}{9} + 4}. \tag{4}$$

So, finally we get

$$|\text{tr}(c_n)| \leq \sqrt{\frac{16n^2V_c^{8/3}}{9} + 4}. \tag{5}$$

□

Now set  $M \setminus L = \mathbb{H}^3/\Gamma_L$  and  $x = \text{Vol}(M \setminus L)$  now for different link  $L$  in  $M$  we get different  $x$  therefore the above expression which is trace modules of loxdromic elements in terms of volume of  $M \setminus L$  will be a sequence of functions of  $x$  that is our 1<sup>st</sup> sequence of functions is  $F_1^n(x) = \sqrt{\frac{16n^2 C_0^{8/3} x^{8/3}}{9}} + 4$ , which is increasing for each  $n$ .

The second function comes from the following theorem by D. Futer et al. [3].

**Theorem 11.** *Let  $M \setminus L$ , be a complete ,finite volume,noncompact hyperbolic manifold . Let  $M$  denote the closed manifold obtained by filling along specified slopes on each of boundary tori, each of which length at least  $2\pi$ , and the least of which is denoted by  $l_{min}$ . Then  $M$  is hyperbolic, and*

$$\text{Vol}(M) \geq \left(1 - \left(\frac{2\pi}{l_{min}}\right)^2\right) \text{Vol}(M \setminus L).$$

By solving the inequality and using the fact given in the theorem we get

$$l_{min} \leq \frac{2\pi}{\sqrt{1 - \left(\frac{V}{\text{Vol}(M \setminus L)}\right)^{2/3}}}.$$

Thus the minimal parabolic length bounded by a function of  $x$  with  $x > V$  i.e.

$$l_{min} \leq \frac{2\pi}{1 - \left(\frac{V}{x}\right)^{2/3}}. \quad (\text{As } x > V \implies \frac{2\pi}{\sqrt{1 - \left(\frac{V}{x}\right)^{2/3}}} \leq \frac{2\pi}{1 - \left(\frac{V}{x}\right)^{2/3}})$$

Now apply the theorem (9) for each  $n$  we can bound trace modulus of loxdromic elements in  $\Gamma_L$  by  $\sqrt{l^4 n^2 + 4}$ . Thus trace modulus also bounded by the decreasing functions.

$$F_2^n(x) = \sqrt{\frac{16\pi^4 n^2}{(1 - (V/x)^{2/3})^4}} + 4. \quad (6)$$

Now, we are ready to present the proof for the main result, i.e. Theorem 4 in the next section.

### 3.1. Proof of Theorem 4

*Proof.* According to the above discussion, we got two functions for every natural number  $n$

$$F_1^n(x) = \sqrt{\frac{16n^2 C_0^{8/3} x^{8/3}}{9}} + 4, \quad (7)$$

and

$$F_2^n(x) = \sqrt{\frac{16\pi^4 n^2}{(1 - (V/x)^{2/3})^4} + 4}. \quad (8)$$

Now observe that  $F_1^n(x)$  are increasing and  $F_2^n(x)$  are decreasing functions for each  $n$ . If there is a common value of these two functions, each  $n$  value should be unique. If we plug in the common value in  $F_1^n(x)$ , (As  $F_1^n$  is increasing for each  $n$ ) this gives an upper bound of the trace modulus of the loxodromic elements for each  $n$ . Now for each  $n$  equating both of them we get

$$\begin{aligned} F_1^n(x) &= F_2^n(x) \\ \implies \frac{C_0^{8/3} x^{8/3}}{9} &= \frac{\pi^4}{(1 - \frac{V^{2/3}}{x^{2/3}})^4} \\ \implies x &= \left( V^{2/3} + \frac{\pi\sqrt{3}}{C_0^{2/3}} \right)^{3/2} \end{aligned}$$

Now applying the theorem ((7)) we get the  $n^{\text{th}}$  geodesic bound and the bound is given by

$$l_n \leq \log(F_1^n(x)^2 + 4) = \log \left[ \frac{16n^2 C_0^{8/3}}{9} \left( V^{2/3} + \frac{\pi\sqrt{3}}{C_0^{2/3}} \right)^4 + 8 \right] = \log \left[ \left( \frac{\sqrt{n} 2 C_0^{2/3} V^{2/3}}{\sqrt{3}} + 2\pi\sqrt{n} \right)^4 + 8 \right].$$

□

## References

## References

- [1] M.Lackenby , *Word hyperbolic Dehn surgery*, arXiv preprint [math/9808120](https://arxiv.org/abs/math/9808120), 1998.
- [2] C. C. Adams, A. W. Reid, *Systoles of hyperbolic 3-manifolds*, Mathematical Proceedings of the Cambridge Philosophical Society, Vol. 128, Cambridge University Press, 2000, pp. 103–110.
- [3] F.David and K. Efstratia and J. S.Purcell ,*Dehn filling, volume, and the Jones polynomial*,J. Differential Geom.,Journal of Differential Geometry,78 (3) (2008) 429–464.
- [4] I. Agol, *Bounds on exceptional Dehn filling*, Geom. Topol. 4 (2000) 431–449. doi:10.2140/gt.2000.4.431
- [5] S. Palaparthi, *Closed geodesic lengths in hyperbolic link complements in  $s \geq 3$*  , Int. J. Pure and Applied Math 83 (2013) 45–53.
- [6] G. S. Lakeland,C. J. Leininger, *Systoles and dehn surgery for hyperbolic 3-manifolds*, Algebraic & Geometric Topology 14 (3) (2014) 1441–1460.