

Loop zero forcing and grundy domination in planar graphs and claw-free cubic graphs

Alex Domat^a

Kirsti Kuenzel^a

December 2, 2022

^a Department of Mathematics, Trinity College, Hartford, CT, USA

Abstract

Given a simple, finite graph with vertex set $V(G)$, we define a zero forcing set of G as follows. Choose $S \subseteq V(G)$ and color all vertices of S blue and all vertices in $V(G) - S$ white. The color change rule is if w is the only white neighbor of blue vertex v , then we change the color of w from white to blue. If after applying the color change rule as many times as possible eventually every vertex of G is blue, we call S a zero forcing set of G . $Z(G)$ denotes the minimum cardinality of a zero forcing set. Davila and Henning proved in [6] that for any claw-free cubic graph G , $Z(G) \leq \frac{1}{3}|V(G)| + 1$. We show that if G is 2-edge-connected, claw-free, and cubic, then $Z(G) \leq \left\lceil \frac{5n(G)}{18} \right\rceil + 1$. We also study a similar graph invariant known as the loop zero forcing number of a graph G which happens to be the dual invariant to the Grundy domination number of G . Specifically, we study the loop zero forcing number in two particular types of planar graphs.

Keywords: zero forcing, Grundy domination, cubic graph, planar graphs

AMS subject classification: 05C69, 05C10

1 Introduction

Zero forcing is a coloring process of the vertices of G according to the following rule. Pick a set $S \subseteq V(G)$ and color each vertex of S blue and each vertex of $V(G) - S$ white. The color change rule is that for each $w \in V(G) - S$ where w is the only white neighbor of a blue vertex v , then color w blue. If after applying the color change rule as many times as possible each vertex of G is blue, then we call S a zero forcing set of G . The zero forcing number of G , denoted $Z(G)$, is the minimum cardinality among all zero forcing sets of G . The original motivation for studying zero forcing was that $Z(G)$ is an upper bound for the minimum rank of a graph

(see [1]). In a seemingly unrelated paper, Brešar et al. [4] defined what is known as a *legal dominating sequence* and *Grundy domination number* of a graph in 2014 as follows. Let $S = (x_1, \dots, x_n)$ be a sequence of distinct vertices of G . S is called a legal sequence if for each $i \in [n - 1]$, x_i dominates some vertex that is not already dominated by any vertex in $\{x_1, \dots, x_{i-1}\}$. If $S = (x_1, \dots, x_n)$ is a legal sequence and $\{x_1, \dots, x_n\}$ dominates G , then we call S a legal dominating sequence of G . A longest possible dominating sequence is called a Grundy dominating sequence and its length is denoted $\gamma_{\text{gr}}(G)$. Brešar et al. showed in [5] that by modifying this definition slightly, one derives the dual invariant of $Z(G)$. In particular, $S = (x_1, \dots, x_n)$ is said to be a legal Z -sequence if for each $i \in [n - 1]$, x_i has a neighbor that is not dominated by any vertex in $\{x_1, \dots, x_{i-1}\}$. The longest Z -sequence in G is called the Z -Grundy domination number and its length is denoted $\gamma_{\text{gr}}^Z(G)$. As $\gamma_{\text{gr}}^Z(G)$ is the dual invariant of $Z(G)$, it follows that $|V(G)| = \gamma_{\text{gr}}^Z(G) + Z(G)$.

Upon realizing that a Grundy-like domination number is the dual invariant to the zero forcing number of a graph, Lin [9] showed that the Grundy domination number of a graph is the dual invariant of what is called the loop zero forcing number of a graph, defined as follows. In [8], the zero forcing number of a loop graph, denoted $Z_{\ell}(G)$, was defined whereby a loop graph is constructed from any simple graph G by adding a loop at each vertex of G . On the other hand, the loop zero forcing number of a graph, $Z_{\ell}(G)$, was defined in [3] slightly differently as finding the zero forcing number of the graph \widehat{G} which is obtained from a simple graph G by adding a loop at each vertex $v \in V(G)$ where $\deg_G(v) \geq 1$. Note however, that if we only consider connected, simple graphs G , $Z_{\ell}(G) = Z_{\widehat{G}}(G)$. Therefore, we refer to $Z_{\ell}(G)$ as the loop zero forcing number of G as we only consider connected, simple graphs throughout this paper. Moreover, for any connected, simple graph G , $|V(G)| = \gamma_{\text{gr}}(G) + Z_{\ell}(G)$.

Davila and Henning showed in [6] that for any claw-free cubic graph G of order at least 10 that $Z(G) \leq \frac{1}{3}|V(G)| + 1$. Motivated by their result, we consider the loop zero forcing number of 2-connected, claw-free cubic graphs. We use a similar technique used by Anderson and Kuenzel in [2] whereby an upper bound for the power domination number of a claw-free cubic graph was given. Although originally motivated by loop zero forcing sets, we show the following upper bound for $Z(G)$ in the case where G is a 2-connected, claw-free cubic graph.

Theorem 1. *Let G be a 2-edge-connected, claw-free cubic graph. If G is not a ring of diamonds, then $Z_{\ell}(G) \leq Z(G) \leq \left\lceil \frac{5n(G)}{18} \right\rceil + 1$. If G is a ring of diamonds, then $Z_{\ell}(G) = Z(G) = \frac{n(G)}{4} + 2$.*

We also consider the loop zero forcing number of certain types of planar graphs. In [7], a series of upper bounds for $Z(G)$ were provided when G is a maximal outerplanar graph. Motivated by their work, we show that in certain cases the loop zero forcing number and zero forcing number of maximal outerplanar graphs coincide. We also provide a lower bound for the zero forcing number of a maximal

outerplanar graph.

The remainder of this paper is organized as follows. In Section 1.1, we provide definitions and terminology used throughout the paper. Section 2 is dedicated to studying the loop zero forcing number of certain planar graphs. In particular, we consider maximal outerplanar graphs and Halin graphs. In Section 3, we prove Theorem 1.

1.1 Definitions and Preliminaries

Let G be a finite, simple graph with vertex set $V(G)$ and edge set $E(G)$. We let $n(G) = |V(G)|$. For a vertex $x \in V$, the *open neighborhood* of x is the set $N(x)$ defined by $N(x) = \{w \in V(G) : xw \in E(G)\}$. The *closed neighborhood* $N[x]$ is $N(x) \cup \{x\}$. The open neighborhood of a set $A \subseteq V(G)$ is $N(A) = \cup_{a \in A} N(a)$ and its closed neighborhood is $N[A] = N(A) \cup A$. Given two vertices $x, y \in V(G)$, we let $d_G(x, y)$ denote the distance between x and y , or length of the shortest path between x and y .

A set A of vertices is a *dominating* set of G if $N[A] = V(G)$. Let $S = (x_1, \dots, x_n)$ be a sequence of distinct vertices in G . We denote the length of S by $|S|$. The set $\{x_1, \dots, x_n\}$ whose elements are the vertices in S is denoted by \widehat{S} . The sequence S is called a *closed neighborhood sequence* (or a *legal sequence*) if

$$N[x_{i+1}] - \bigcup_{j=1}^i N[x_j] \neq \emptyset \quad (1)$$

for each $i \in [n-1]$. We will also say that x_{i+1} *footprints* the vertices from $N[x_{i+1}] - \bigcup_{j=1}^i N[x_j]$ with respect to S . If S is a legal sequence and \widehat{S} is a dominating set of G , then S is called a *dominating sequence* in G . A longest possible dominating sequence in G is called a *Grundy dominating sequence*, and its length is the *Grundy domination number* of G , denoted $\gamma_{\text{gr}}(G)$.

A closed neighborhood sequence S in G is a *Z-sequence* if, in addition, every vertex v_i in S footprints a vertex distinct from itself. That is, S is a Z-sequence in G if

$$N_G(v_i) \setminus \bigcup_{j=1}^{i-1} N_G[v_j] \neq \emptyset$$

holds for every $i \in \{2, \dots, k\}$. The maximum length of a Z-sequence in G is the *Z-Grundy domination number*, denoted $\gamma_{\text{gr}}^Z(G)$, of G . Given a Z-sequence S , we call the corresponding set \widehat{S} a Z-set.

Given a set $S \subseteq V(G)$, we define S to be a zero forcing set of G if the following is true. Initially, color all vertices of S blue and all vertices of $V(G) - S$ white. We iteratively apply the zero forcing color change rule: if $w \in V(G)$ is the only white neighbor of blue vertex v , recolor w blue. If after applying the zero forcing color

change rule as many times as possible all vertices of G are blue, we call S a *zero forcing set* of G . $Z(G)$ denotes the minimum cardinality among all zero forcing sets of G . We will say $S \subseteq V(G)$ is a *loop zero forcing set* if the following is true. Initially, we color all vertices of S blue and all vertices of $V(G) - S$ white. We iteratively apply the loop zero forcing color change rule: if $w \in V(G)$ is either 1) the only white neighbor of blue vertex v , or 2) every neighbor of w is blue, then we recolor w blue. If after applying the loop zero forcing color change rule as many times as possible all vertices of G are blue, then call S a *loop zero forcing set* of G . $Z_{\ell}(G)$ denotes the minimum cardinality among all loop zero forcing sets of G . Note that the zero forcing number of G is the dual invariant of $\gamma_{\text{gr}}^Z(G)$ and the loop zero forcing number of G is the dual invariant of $\gamma_{\text{gr}}(G)$. That is, $n(G) = Z(G) + \gamma_{\text{gr}}^Z(G) = Z_{\ell}(G) + \gamma_{\text{gr}}(G)$.

We will make use of the following two known results.

Proposition 2. [4] *For an arbitrary graph G , $\gamma_{\text{gr}}(G) \leq n(G) - \delta(G)$.*

Theorem 3. [5] *If G is a graph without isolated vertices, then $\gamma_{\text{gr}}^Z(G) + Z(G) = n(G)$. Moreover, the complement of a zero forcing set of G is a Z -set of G and vice versa.*

2 Planar Graphs

In this section, we consider two types of planar graphs. Recall that G is *outerplanar* if G has a planar drawing in which all vertices of $V(G)$ belong to the outer face of the drawing. The *weak planar dual* graph of a planar graph is the graph that has a vertex for every bounded face and an edge between two vertices which represent a pair of adjacent bounded faces. In this section, we consider both maximal outerplanar graphs and Halin graphs.

2.1 Maximal Outerplanar Graphs

G is *maximal outerplanar*, or a MOP, if G is outerplanar and the addition of any edge results in a graph that is not outerplanar. Thus, every interior face of G is a triangle. We use similar terminology to that found in [7]. Namely, a triangular face T is a *separator triangle* of G if it has no edges on the outer face. The number of separator triangles of G is denoted by t . We say a MOP G is *serpentine* if it contains no separator triangles. We let G_{Δ} denote the subgraph formed by the separator triangles of G . The weak planar dual of a MOP G is a tree that we denote by H . A *serpentine leaf* is a set of triangles in G whose corresponding vertices in H form the only path in H connecting a leaf of H with a vertex u of H where $\deg_H(u) > 2$. We let h represent the number of serpentine leaves in G and n_2 represent the number of vertices of degree 2 in G . Note that $h = n_2$. Similarly, we define a *serpentine path* to be a maximal set of triangles in G whose corresponding vertices in H have

degree 2 in H and form the only path between two distinct vertices of degree 3 or more in H . The number of serpentine leaves with fan structure from a vertex of a separator triangle, called fan leaves, is denoted by h_F . Figure 1 provides an example of a MOP where the green triangles are separator triangles, the purple triangles lie form a fan leaf, and the yellow triangles form a serpentine path.




Figure 1: Example of serpentine path and fan leaf in a MOP

Note that the following were shown in [7].

Proposition 4. [7] *If G is a serpentine graph, then $Z(G) = 2$.*

Theorem 5. [7] *If G is a MOP with $t = 1$ and $h_F \geq 0$, then*

1. *If $h_F \geq 1$, then $Z(G) = 3$.*
2. *If $h_F = 0$, then $Z(G) = 4$.*

Theorem 6. [7] *If G is a MOP with n_2 vertices of degree 2 whose G_Δ has c components, then*

$$Z(G) \leq 2 \left\lceil \frac{n_2 + c - 1}{2} \right\rceil$$

and these bounds are tight when the number of fan leafs is 0 and if $c = 0$ or G_Δ is formed by isolated triangles.

We first point out some immediate consequences of the above results. First, note that if G is a serpentine graph, then $\delta(G) = 2$. It is clear that for any graph H , $Z_\ell(H) \geq \delta(H)$ and therefore by Proposition 4 we have the following.

Corollary 7. *If G is a serpentine graph, then $Z_\ell(G) = 2$.*

Next, we show that for all MOPs G described in the statement of Theorem 5, $Z_\ell(G) = Z(G)$.

Proposition 8. *If G is a MOP with $t = 1$ and $h_F \geq 0$, then*

1. If $h_F \geq 1$, then $Z_{\hat{\ell}}(G) = 3$.
2. If $h_F = 0$, then $Z_{\hat{\ell}}(G) = 4$.

Proof. First, assume that G is a MOP with $t = 1$ and $h_F \geq 1$. It is clear that $Z_{\hat{\ell}}(G) \geq 2$ and every vertex of G is contained on a serpentine leaf. Suppose that there exists a loop zero forcing set of cardinality 2, say $\{x, y\}$. Thus, both x and y must be vertices on the same serpentine leaf, call it L . Let w and z be the two vertices of G that are on the one separator triangle of G and also contained in L . Assuming that all vertices of L eventually turn blue, each of w and z have at least two neighbors that are not in L and therefore $\{x, y\}$ is not a loop zero forcing set of G .

Next, assume that G is a MOP with $t = 1$ and $h_F = 0$. Enumerate the serpentine leaves of G as L_1, L_2, L_3 . Suppose that there exists a loop zero forcing set $S = \{x, y, t\}$. Thus, at least two vertices of S must be contained in the same serpentine leaf. Without loss of generality, we may assume x and y are vertices in L_1 . As above, let w and z be the two vertices of G that are on the one separator triangle of G and also contained in L_1 . Assuming that all vertices of L_1 eventually turn blue, each of w and z have two neighbors that are not in L_1 . Therefore, t is a neighbor of at least one of w or z . Assume first that t is on the separator triangle. We shall assume w and t are vertices of L_2 and z and t are vertices of L_3 . If w has only one white neighbor, say a , in L_2 and z has only one white neighbor in L_3 , say b , then a and b both turn blue and the propagation process stops as each of a, b , and t have at least two white neighbors. So we shall assume that w has two white neighbors in L_2 , say a and c . If z has only one white neighbor, say b , in L_3 , then b turns blue. Moreover, no vertices in L_2 will turn blue until all vertices of L_3 are blue. This will only occur if L_3 is a fan leaf, which contradicts the assumption that $h_F = 0$. Thus, this case cannot occur and we shall assume that t is not on the separator triangle and t is a vertex in L_3 adjacent to z . Let r be the third vertex of the separator triangle. If $rt \in E(G)$, then r is the only white neighbor of z and therefore r turns blue. Using a similar argument as that above, no vertex of L_2 will turn blue until all vertices of L_3 are blue which can only occur if L_3 is a fan leaf, which is a contradiction. Therefore, we may assume $rt \notin E(G)$ and the only blue vertices that are not contained in $V(L_1) - \{w, z\}$ are w, z , and t . Note that w and z each have at least two white neighbors. Therefore, if S is indeed a loop zero forcing set, it must be that t has only one white neighbor. This implies that $\deg_G(t) = 2$ and L_3 is a fan leaf, another contradiction. Hence, no such loop zero forcing set exists. \square

Next, we give a lower bound for the zero forcing number of a MOP. We begin with the following observation.

Observation 1. *For any serpentine graph G , there exist only two distinct minimum zero forcing sets and only two distinct maximum Z -sets.*

The above follows from the fact that any minimum zero forcing set contains exactly two vertices, one of which is a vertex of degree 2 in G , call it v , and the other which is a neighbor of v . Therefore, by Theorem 3, the only maximum Z -sets of G are the complements of the two minimum zero forcing sets.

Theorem 9. *Let G be a MOP with $t \geq 1$ where G_Δ contains only one component. Then $Z(G) \geq \lceil \frac{n_2}{2} \rceil$.*

Proof. Let H be the graph obtained from G by removing all vertices of G that are contained in a serpentine leaf, but not contained on a separator triangle. Note that H is necessarily a MOP and H is a serpentine graph. If H is a triangle, then we know $Z(G) \geq 3$ by Theorem 5. Therefore, we may assume that $t \geq 2$. We enumerate the vertices of H as $x_1 \dots x_r$ such that $x_i x_{i+1} \in E(H)$ for $i \in [r]$. Let $S = (s_1, \dots, s_n)$ be a Z -Grundy sequence of G and suppose L_i is a serpentine leaf of G such that $V(L_i) \cap \widehat{S} = V(L_i)$. Moreover, assume x_i and x_{i+1} are in L_i . Suppose first that x_{i+1} and x_{i+2} are also contained in a serpentine leaf of G , call it L_{i+1} . Since all vertices of L_i are in \widehat{S} , some vertex $y \in V(L_i) - \{x_i, x_{i+1}\}$ must footprint x_i and some vertex $z \in V(L_i) - \{x_i, x_{i+1}, y\}$ must footprint x_{i+1} since there is only one distinct Z -Grundy sequence of the serpentine graph L_i not containing x_i and x_{i+1} . Now since z footprints x_{i+1} and $x_{i+1} \in \widehat{S}$, there exists a neighbor t of x_{i+1} in L_{i+1} such that $t \notin \{x_{i+1}, x_{i+2}\}$ and t does not precede x_{i+1} in S . Let \overline{S} be the subsequence of S that contains only those vertices in L_{i+1} . If \overline{S} is a Z -Grundy sequence of the serpentine graph L_{i+1} , then one of x_{i+1} or x_{i+2} has degree 2 in L_{i+1} and the other vertex of L_{i+1} that has degree 2 is not in \widehat{S} as there is only one distinct Z -Grundy sequence of the serpentine graph L_{i+1} where x_{i+1} precedes t . On the other hand, if \overline{S} is not a Z -Grundy sequence of the serpentine graph L_{i+1} , then some vertex of $V(L_{i+1}) - \{x_{i+1}\}$ is not in \widehat{S} . In either case, some vertex of L_{i+1} is not in \widehat{S} .

Therefore, we shall assume that $x_{i+1}x_{i+2}$ is on the boundary of G . We may extend $x_{i+1}x_{i+2}$ to a maximal path $x_{i+1}x_{i+2} \dots x_j$ that lies on the boundary of both H and G and where x_j is contained in a serpentine leaf yet x_{i+2}, \dots, x_{j-1} are not. Let L_j be the serpentine leaf containing x_j and assume that $V(L_j) \cap \widehat{S} = V(L_j)$. Moreover, assume that $\{x_{i+1}, \dots, x_j\} \subset \widehat{S}$. Suppose first that $x_j = x_{i+2}$. Since every vertex of L_i is in \widehat{S} , x_{i+2} comes after x_{i+1} in S for otherwise some vertex of L_i does not footprint a vertex. However, using this same logic, since every vertex of L_j is in \widehat{S} , x_{i+1} comes after x_{i+2} in S , which is a contradiction. Therefore, we shall assume that $j > i + 2$. As above, since every vertex of L_i is in \widehat{S} , x_{i+2} comes after x_{i+1} in S for otherwise some vertex of L_i does not footprint a vertex. Similarly, x_{j-1} comes after x_j in S . Thus, of the set $\{x_{i+1}, \dots, x_j\}$, the last vertex to appear in S is in $\{x_{i+2}, \dots, x_{j-1}\}$, call it x_α . But this cannot be as $x_{\alpha-1}$ and $x_{\alpha+1}$ precede x_α in S , meaning x_α does not footprint a vertex. Therefore, it must be that either some vertex of $\{x_{i+2}, \dots, x_{j-1}\}$ is not in \widehat{S} , or some vertex of L_j is not in \widehat{S} . In either case, we have $|\widehat{S} \cap V(G)| \leq n(G) - \lceil \frac{n_2}{2} \rceil$ which implies $Z(G) \geq n(G) - (n(G) - \lceil \frac{n_2}{2} \rceil) = \lceil \frac{n_2}{2} \rceil$.

□

We use the above result to provide a lower bound for the zero forcing number of any MOP.

Corollary 10. *If G is a MOP where $t \geq 1$, \mathcal{H} represents the set of components in G_Δ that are adjacent to exactly one serpentine path in G and n'_2 is the number of serpentine leaves that are adjacent to a component in \mathcal{H} , then $Z(G) \geq \left\lfloor \frac{n'_2}{2} \right\rfloor - 2c'$ where c' is the number of components in \mathcal{H} .*

Proof. Let \overline{G} be the graph obtained from G by removing all vertices of G that are contained in a serpentine leaf or path, but not contained on a separator triangle. Let $H \in \mathcal{H}$ and let \overline{H} be the component of \overline{G} where $V(H) \subseteq V(\overline{H})$. Thus, \overline{H} is a MOP. We may enumerate the vertices of \overline{H} as $x_1 \dots x_r$ where $x_i x_{i+1}$ is on the boundary of \overline{H} for $i \in [r]$. Reindexing if necessary, we may assume $x_1 x_2$ is on a serpentine path of G . Let n_H be the number of serpentine leaves adjacent to H . Let G' be the graph containing $V(H)$ obtained from G by removing the neighbors of x_1 and x_2 that lie on the serpentine path adjacent to H but are not themselves contained in a separator triangle. Thus, G' is a MOP where G'_Δ contains one component. Let S be a zero forcing set of G . Even if x_1 and x_2 are observed by vertices in their serpentine path, from Theorem 9 we know $|S \cap V(G')| \geq \left\lceil \frac{n_H}{2} \right\rceil$ which implies $|S \cap (V(G') - \{x_1, x_2\})| \geq \left\lceil \frac{n_H}{2} \right\rceil - 2$. Summing over all components in \mathcal{H} yields the desired result.

□

2.2 Halin Graphs

Recall that a Halin graph is constructed from a tree by enumerating the leaves of T as ℓ_1, \dots, ℓ_m in such a way so that when we add the edges $\ell_i \ell_{i+1}$ for each $i \in [m]$ to T , the resulting graph is planar. In this section, we use a similar strategy (and therefore similar terminology) to that used by Brešar et al. in [4] where the authors are providing bounds on the Grundy domination number of a tree. For any tree T , a support vertex of T is any vertex adjacent to a leaf in T . An *end support vertex* is a support vertex v such that $\deg'(v) \leq 1$, where $\deg'(v)$ is the number of vertices adjacent to v which are not leaves. Equivalently, a support vertex v is an end support vertex if and only if v does not lie on a path between two other support vertices. The following lower bound was given for the Grundy domination number of a tree in [4].

Proposition 11. [4] *For any tree T which is not a star, $\gamma_{\text{gr}}(T) \geq n(T) - |ES(T)| + 1$ where $ES(T)$ is the set of end support vertices in T .*

This immediately implies the following in terms of loop zero forcing.

Corollary 12. [4] For any tree T which is not a star, $Z_{\dot{\ell}}(T) \leq |ES(T)| - 1$.

Studying the loop zero forcing number of Halin graphs is quite different from studying the loop zero forcing number of trees as there are no leaves in a Halin graph. Given a Halin graph G constructed from a tree T_G , we say that a vertex of G is an end support vertex of G if it is an end support vertex of the tree T_G .

Proposition 13. If $ES(G)$ is the set of end support vertices in a Halin graph G , then $Z_{\dot{\ell}}(G) \leq Z(G) \leq 3(|ES(G)| - 1)$.

Proof. Assume that G is constructed from T_G and let C represent the exterior boundary of G consisting of all the leaves in T_G . Let L be the set of leaves of T_G , and let $H = G - L$. Note that H is a tree. We then construct a tree T with vertices v_1, \dots, v_t such that each $v_i \in V(T)$ corresponds to a unique vertex of H which does not have degree 2 in H . v_i and v_j are adjacent in T if their corresponding vertices in H are adjacent, or the path P in H between the corresponding vertices in H is such that all interior vertices of P have degree 2 in H . Root T at a leaf $r \in T$. Note that the leaves of T correspond to the end support vertices of G . We refer to vertices in G that correspond to non-leaf vertices in T as *branching vertices*.

We define two kinds of branch structures in G . These are analogs of serpentine leaves and serpentine paths for MOPs. A type-1 branch $B_1 \subset G$ is constructed as follows. Let $xy \in E(T)$ where x is a leaf of T and let u_x (resp. u_y) be the vertex of G that x represents (resp. y represents). The vertices of B_1 consist of u_x, u_y , any vertices along the u_x - u_y path in T_G , call it $P = u_x w_1 \cdots w_k u_y$, as well as any leaves (in T_G) of any vertex of P other than u_y . A type-2 branch $B_2 \subset G$ is constructed as follows. Let $xy \in E(T)$ where neither x nor y is a leaf of T and let u_x (resp. u_y) be the vertex of G that x represents (resp. y represents). The vertices of B_2 consist of u_x, u_y , any vertices along the u_x - u_y path in T_G , call it $P = u_x w_1 \cdots w_k u_y$, as well as any leaves (in T_G) of any vertex of P (including u_y).

Choose a set S according to the following rule: for all end support vertices $s \in G$, excepting the end support vertex which corresponds to r , choose a leaf ℓ_s of s in T_G along with the two neighbors in G of ℓ_s that are on C . Color all vertices of S blue. We claim that S is a zero forcing set of G .

Let B_1 be a type-1 branch as described above where $xy \in E(T)$ and $P = u_x w_1 \cdots w_k u_y$ is the u_x - u_y path in T_G . Note that u_x is the only white neighbor of ℓ_{u_x} in G and therefore u_x will become blue. Furthermore, all leaves of u_x in T_G will eventually turn blue as the leaves of u_x are ordered in a way that they lie on a path contained in the exterior cycle C of G . In fact, we may assume that the leaves of u_x in T_G are indexed as $\ell_i \ell_{i+1} \dots \ell_j$ such that this path is on C . Also note that $xy \in E(T)$ if and only if $u_x u_y \in E(G)$ or $\deg_H(w_i) = 2$ for $i \in [k]$. Now once all leaves of u_x in T_G are blue, w_1 turns blue as it is the only white neighbor of u_x . If w_1 has degree 2 in G , then w_2 will also turn blue. So assume that w_1 is adjacent to a leaf of T_G . Without loss of generality, we may assume that either all leaves of w_1

in T_G lie on a path of the form $\ell_{i'}\ell_{i'+1}\dots\ell_{i-1}$ where $i' < i$ and ℓ_{i-1} is adjacent to ℓ_i on C , or all leaves of w_1 lie on two paths, one of the form $\ell_{i'}\ell_{i'+1}\dots\ell_{i-1}$ where $i' < i$ and ℓ_{i-1} is adjacent to ℓ_i on C and the other of the form $\ell_{j+1}\dots\ell_t$ where ℓ_j is adjacent to ℓ_{j+1} on C . In either case, one can easily verify that since all vertices in $N_G[u_x]$ are blue, all leaves of w_1 in T_G will eventually turn blue at which point w_2 will also turn blue. This propagation will continue through B_1 and indeed all vertices on a type-1 branch will eventually turn blue.

Next, choose a type-2 branch B_2 as described above where $xy \in E(T)$, neither of x or y is a leaf of T , $P = u_x w_1 \dots w_k u_y$ is the u_x - u_y path in T_G and x has maximum distance from r among all such type-2 branches. Note that this implies that u_x is blue as it is necessarily on a type-1 branch. Since we have assumed x has maximum distance from r , we know that u_x is contained in only one type-2 branch. Therefore, if u_x is not adjacent to any leaves in T_G , then w_1 is the only white neighbor of u_x and propagation continues throughout B_2 . Therefore, we shall assume that u_x is adjacent to leaves in T_G . As above, we may assume that either all leaves of u_x in T_G lie on a path $\ell_i \dots \ell_j$ on C or all leaves of u_x lie on two paths, namely $\ell_i \dots \ell_j$ and $\ell_m \dots \ell_n$ on C . Choose a type-1 branch B_1 that contains u_x as well as a vertex t such that t has a leaf ℓ_{j+1} where ℓ_{j+1} is adjacent to ℓ_j on C . Since all leaves of t in T_G are assumed to be blue, all leaves of u_x of the form $\ell_i \dots \ell_j$ will eventually turn blue as well. Similarly, all leaves of u_x of the form $\ell_m \dots \ell_n$ will eventually turn blue. Thus, w_1 will be the only white neighbor of u_x at some point and will eventually turn blue as well. Propagation will continue throughout B_2 and eventually all vertices of B_2 will turn blue. Continuing this same argument by choosing type-2 branches B_2 where $xy \in E(T)$ and x has the maximum distance from r among all vertices corresponding to a type-2 branch containing white vertices, one can see that all vertices of G will indeed turn blue and S is a zero forcing set of G . □

There are two immediate consequences to the above result.

Corollary 14. *For any Halin graph G where $ES(G)$ is the set of end support vertices in G , $\gamma_{\text{gr}}(G) \geq n(G) - 3(|ES(G)| - 1)$.*

Corollary 15. *If G is Halin graph which contains no vertex v with $\deg'(v) \geq 3$, then $Z(G) = Z_{\dot{\ell}}(G) = 3$ and $\gamma_{\text{gr}}(G) = n(G) - 3$.*

Proof. Note that in such a Halin graph, there are only two end support vertices and therefore $Z_{\dot{\ell}}(G) \leq 3$. On the other hand, $\delta(G) \geq 3$ and by Proposition 2 $\gamma_{\text{gr}}(G) \leq n(G) - 3$ and therefore $Z_{\dot{\ell}}(G) \geq 3$. □

3 Claw-free cubic graphs

As mentioned earlier, Davila and Henning proved the following regarding the zero forcing number of claw-free cubic graphs.

Theorem 16. [6] *If G is a claw-free cubic graph with $n(G) \geq 10$, then $Z(G) \leq \frac{1}{3}n(G) + 1$.*

We originally wanted to study the loop zero forcing number of claw-free cubic graphs using the same approach to that used in [2]. As it turns out, we only ever considered zero forcing sets and therefore we provide an improved bound to the one given in Theorem 16 in the case when G is a 2-edge-connected claw-free cubic graph. To do so, we need the following known results regarding 2-edge-connected cubic graphs. Recall that the graph obtained from K_4 by removing any edge is referred to as a *diamond*. In what follows, a *string of diamonds* is defined to be a maximal sequence D_1, \dots, D_k of diamonds in which, for each $i \in [k]$, D_i has a vertex adjacent to a vertex in D_{i+1} (modulo k) and has exactly two vertices which have degree two. Given a string of diamonds D_1, \dots, D_k , we label the vertices of D_i as u_i, v_i, w_i , and y_i where u_i is adjacent to a vertex in D_{i-1} and y_i is adjacent to a vertex in D_{i+1} . Moreover, we refer to v_i and w_i as the *interior vertices of D_i* . The following results are necessary for how we approach zero forcing in 2-edge-connected claw-free cubic graphs.

Theorem 17. [10] *A graph G is 2-edge-connected claw-free cubic if and only if either*

(i) $G \cong K_4$,

(ii) G is a ring of diamonds, or

(iii) G can be built from a 2-edge-connected cubic multigraph H by replacing some edges of H with strings of diamonds and replacing each vertex of H with a triangle.

Recall that a 2-factor of a graph G is a 2-regular spanning subgraph of G .

Theorem 18. [11] *Let G be a 2-edge-connected cubic multigraph where $|V(G)|$ is even. For any arbitrary edge $e \in E(G)$, G contains a 2-factor containing e .*

In what follows, if we assume that $\mathcal{C} = C_1 \cup \dots \cup C_k$ is a 2-factor of G , we will say that C_i is adjacent to C_j if there exists some v on C_i and some w on C_j such that $vw \in E(G)$. We are now ready to prove Theorem 1, restated here for ease of reference.

Theorem 1 *Let G be a 2-edge-connected, claw-free cubic graph. If G is not a ring of diamonds, then $Z_{\ell}(G) \leq Z(G) \leq \left\lceil \frac{5n(G)}{18} \right\rceil + 1$. If G is a ring of diamonds, then $Z_{\ell}(G) = Z(G) = \frac{n(G)}{4} + 2$.*

Proof. Note that the statement is clearly true when $G = K_4$. Therefore, we may assume that $n(G) \geq 6$. Suppose first that G is a ring of k diamonds where we

enumerate the diamonds as D_1, \dots, D_k such that D_i is adjacent to D_{i-1} and D_{i+1} for $i \in [k]$ and the vertices of D_i are labeled u_i, v_i, w_i, y_i where u_i is adjacent to v_i, w_i , and a vertex of D_{i-1} and y_i is adjacent to v_i, w_i , and a vertex of D_{i+1} . One can easily verify that $S = \{v_1, y_1, u_2, v_2\} \cup \{v_i : 3 \leq i \leq k\}$ is a zero forcing set of G of cardinality $\frac{n(G)}{4} + 2$. To see that this is best possible, suppose there exists a loop zero forcing set of G , call it S , with cardinality $\frac{n(G)}{4} + 1$. $S \cap V(D_i) \neq \emptyset$ for each $i \in [k]$ for otherwise v_i and w_i will never become blue. Reindexing if necessary, we may assume $|S \cap V(D_1)| = 2$ and $|S \cap V(D_i)| = 1$ for $2 \leq i \leq k$. One can easily verify that one of v_1 or w_1 is in S . Without loss of generality, assume $v_1 \in S$. If $\{v_1, w_1\} \subset S$, then $u_2 \in S$ and y_1 will turn blue as all of its neighbors are in S . However, v_2 and w_2 will never turn blue no matter how many times we apply the color change rule so this case cannot occur. Therefore, either $\{v_1, u_1\} \subset S$ or $\{v_1, y_1\} \subset S$. Without loss of generality, we may assume $\{v_1, y_1\} \subset S$. Thus, $u_2 \in S$ for otherwise the color change rule can never be applied. However, we reach a similar contradiction as before in that v_2 and w_2 will never become blue. Thus, no such loop zero forcing set S of G exists and we may conclude $Z_i(G) = Z(G) = \frac{n(G)}{4} + 2$.

Hence, we shall assume G can be built from a 2-edge-connected cubic multigraph H by replacing some edges of H with strings of diamonds and replacing each vertex of H with a triangle. By Theorem 18, H contains a 2-factor $\mathcal{C}' = C'_1 \cup \dots \cup C'_k$. Let G' be the graph obtained from H by replacing each vertex of H with a triangle. Note that G' may not be G . We can create a 2-factor $\mathcal{C} = C_1 \cup \dots \cup C_k$ for G' from \mathcal{C}' by replacing each vertex u of C'_i with a P_3 $u_1 u_2 u_3$ where $u_1 u_3$ is also an edge in G' (see Figure 2). We may enumerate the vertices of C_i as $C_i = x_1^i \dots x_{n_i}^i$ such that $3 \mid n_i$ and the subgraph of G' induced by $\{x_j^i, x_{j+1}^i, x_{j-1}^i\}$ for all $j \equiv 1 \pmod{3}$ is a triangle. Moreover, there exists a perfect matching M in G' between the vertices of

$$\bigcup_{i \in [k]} \{x_1^i, x_4^i, x_7^i, \dots, x_{n_i-2}^i\}.$$

Let J be the graph with $V(J) = \{u_1, \dots, u_k\}$ where $u_i u_j \in E(J)$ if and only if there exists an edge $xy \in E(G)$ such that x is on C_i and y is on C_j , $i \neq j$. Choose a spanning tree T of J rooted at u_1 and let $A_i = \{v \in V(T) : d_T(u_1, v) = i\}$. Moreover, if $k \geq 3$, choose T and u_1 such that $\deg_T(u_1) \geq 2$. Let $\ell : V(T) \rightarrow [n(T)]$ be any bijection where $\ell(u_1) = 1$ and $\ell(v) < \ell(w)$ if $d_T(u_1, v) < d_T(u_1, w)$. We refer to $\ell(v)$ as the label of v in T . Note that we may reindex the vertices of T and corresponding cycles of \mathcal{C} such that $\ell(u_i) = i$ and u_i corresponds to cycle C_i in $\mathcal{C} = C_1 \cup \dots \cup C_k$.

Let $M' = \{w_{\alpha_1} v_{\alpha_1}, \dots, w_{\alpha_s} v_{\alpha_s}\}$ be a subset of M where $s = |E(T)|$ and for each $u_\ell u_{\ell'} \in E(T)$, there exists $w_{\alpha_j} v_{\alpha_j} \in M'$ such that w_{α_j} is on C_ℓ and v_{α_j} is on $C_{\ell'}$. Moreover, we can interchange w_{α_j} and v_{α_j} so that $d_T(u_1, u_\ell) \leq d_T(u_1, u_{\ell'})$. For each $e \in M$, write $e = x_\ell^i x_s^j$ where $i \leq j$, and if $i = j$, then $\ell < s$.

Note that since T is rooted at u_1 , we may choose an edge $e = x_\ell^1 x_s^2 \in M'$ where




Figure 2: Creating a 2-factor for G' from \mathcal{C}' where the dashed edges are edges in G'

ℓ and s are as small as possible and reindex the vertices of C_1 such that $x_\ell^1 = x_1^1$. Similarly, for each $e = x_\ell^j x_s^t \in M'$ where $j < t$, we can reindex the vertices of C_t such that $x_s^t = x_1^t$ (and this reindexing will occur only once since T is a tree). Furthermore, if u_i is not a leaf in T and we let $\{x_{\beta_1}^i, \dots, x_{\beta_j}^i\}$ be those vertices on C_i that are incident to a vertex of the form x_1^t where $t > i$, then we choose the minimum value among $\{\beta_1, \dots, \beta_j\}$ and denote said index by η_i . Similarly, if u_i is a leaf in T , then we let $\{x_{\beta_1}^i, \dots, x_{\beta_j}^i\}$ be those vertices on C_i that are incident to a vertex of the form x_s^t where $t \neq i$. Note that $|\{x_{\beta_1}^i, \dots, x_{\beta_j}^i\}| \geq 2$ as G is 2-edge-connected. Choose the minimum value strictly greater than 1 among $\{\beta_1, \dots, \beta_j\}$ and denote said index by η_i .

Suppose first that $|V(C_i)| = 6$ for all $i \in [k]$ and let $S = \{x_1^1, x_1^2\} \cup \{x_2^i : i \in [k]\}$. We claim that S is a zero forcing set of G' . Note that initially the only white neighbor of x_1^1 is x_6^1 and therefore x_6^1 will become blue. x_2^1 will force x_3^1 to become blue and x_6^1 will force x_5^1 to become blue, which will in turn force x_4^1 to become blue. Similarly, all vertices of C_2 will eventually become blue. Now assume that all vertices of C_i are blue by some timestep β where $u_i \in A_\ell$ for some $i \in [k]$ and $\ell \in [|E(T)|]$. Let C_j be a cycle in \mathcal{C} where $u_j \in A_{\ell+1}$ and $u_i u_j \in E(T)$. Therefore, there exists $e = x_4^i x_1^j \in M'$. Since x_4^i is blue and x_1^j is its only white neighbor at timestep β , then x_1^j will be turned blue. Furthermore, $x_2^j \in S$ so it is blue and the propagation will continue through C_j . Therefore, S is indeed a zero forcing set of G' of cardinality $\frac{n(G')}{6} + 2$. Next, note that we can obtain G from G' by replacing some of the edges in G' which were originally in H with strings of diamonds. See Figure 3 for an example of how to replace an edge of G' with a string of 2 diamonds. Choose an interior vertex from each diamond in every string of diamonds used to replace edges in G' to obtain G and call the resulting set X . One can easily verify that $S' = S \cup X$ is a zero forcing set of G of cardinality $\frac{n(G')}{6} + 2 + \frac{n(G) - n(G')}{4}$. If

$n(G') \geq 12$, then

$$|S'| = \frac{n(G)}{4} + 2 - \frac{n(G')}{12} \leq \frac{n(G)}{4} + 1 \leq \frac{5n(G)}{18} + 1.$$

Therefore, we shall assume $n(G') = 6$. If $n(G) = 6$, then $S = N_G(u)$ for any $u \in V(G)$ is a zero forcing set of G of cardinality $3 = \lceil \frac{5n(G)}{18} \rceil + 1$. So we shall assume that G is obtained from G' by replacing some edges originally in H with strings of diamonds. Suppose first that G is obtained from G' by replacing edges in G which were originally in H with at least three diamonds (i.e. $n(G) \geq 18$). In this case, we label all vertices of each of the ℓ diamonds D_1, \dots, D_ℓ in G as u_i, v_i, w_i, y_i as described above for $i \in [\ell]$. If each edge in G' that was originally in H is replaced with exactly one diamond, then G is isomorphic to the graph depicted in Figure 4 (d). One can easily verify that the black vertices are indeed a zero forcing set. Thus, we shall assume that an edge in G' that was originally in H is replaced with at least two diamonds. Without loss of generality, we may assume D_1 and D_2 are consecutive diamonds in G where $y_1 u_2 \in E(G)$ and $u_1 x_5^1 \in E(G)$. One can easily verify that $\{v_i : i \in \ell\} \cup \{y_1, u_2, x_4^1\}$ is a zero forcing set of G of cardinality $3 + \frac{n(G)-6}{4} \leq \frac{5n(G)}{18} + 1$ as $n(G) \geq 18$. Therefore, we only need to consider when $n(G) \in \{10, 14\}$. If $n(G) = 10$, then G is isomorphic to the graph depicted in Figure 4 (a) and the black vertices form a zero forcing set of cardinality $4 = \lceil \frac{50}{18} \rceil + 1$. If $n(G) = 14$, then G is isomorphic to the graph depicted in Figure 4 (b) or (c). In either case, the black vertices form a zero forcing set of cardinality $5 = \lceil \frac{70}{18} \rceil + 1$.




Figure 3: Creating G from G' by replacing an edge with a string of two diamonds

Thus, for the remainder of the proof we shall assume that some cycle in \mathcal{C} has length 9 or more. Suppose first that \mathcal{C} contains at least two cycles. We create a set D as follows. For each $e = x_\ell^i x_s^j \in M$ where $i < j$ we place $x_\ell^i \in D$ if $|V(C_i)| \geq 9$ and $\ell \neq \eta_i$. For each $e = x_\ell^i x_s^j \in M$ where $|V(C_i)| \geq 9$ and $i = j$, we place x_ℓ^i in




Figure 4: Possible graphs where \mathcal{C} contains only 6-cycles and $n(G) \leq 14$

D if $1 \leq \ell < \eta_i$ and $\ell < s$. Otherwise, if $\eta_i < \ell < s$, then we place x_s^i in D . We let $S = D \cup \{x_1^2\} \cup \{x_i^i : i \in [k]\}$. We claim that S is a zero forcing set of G' . We proceed by showing that if all vertices of C_i for each $i \in [\alpha]$ where $\alpha \in [k]$ are blue by some timestep β , then all vertices of $C_{\alpha+1}$ will eventually be blue. We first show that all vertices of C_1 will eventually turn blue. First note that $\{x_1^1, x_2^1, x_1^2, x_2^2\} \subseteq S$. Therefore, all neighbors of x_1^1 other than $x_{n_1}^1$ are blue and x_1^1 will force $x_{n_1}^1$ to become blue. x_2^1 will force x_3^1 to become blue. Now assume all vertices $x_1^1, x_2^1, x_3^1, \dots, x_j^1$ are blue at some timestep β where $j < \eta_1$. If $j \equiv 2 \pmod{3}$, then x_{j+1}^1 is the only white neighbor of x_j^1 and propagation continues in a clockwise fashion. If $j \equiv 0 \pmod{3}$, then $x_{j+1}^1 \in S$ of x_{j+1}^1 is on some edge $x_{j+1}^1 x_s^1 \in M$ where $1 \leq s \leq j-2$. In either case, x_{j+1}^1 is blue or is the only white neighbor of x_s^1 which implies x_{j+1}^1 will become blue. Finally, if $j \equiv 1 \pmod{3}$, then x_{j+1}^1 is the only white neighbor of x_{j-1}^1 and will therefore become blue. Thus, we may conclude that indeed all vertices of the form x_j^1 where $1 \leq j \leq \eta_1 - 1$ will eventually be blue. Similarly, moving in the opposite direction from $x_{n_1}^1$ to $x_{\eta_1}^1$, all vertices of the form x_j^1 where $\eta_1 + 1 \leq j \leq n_1$ will eventually become blue. Hence, either $x_{\eta_1-1}^1$ or $x_{\eta_1+1}^1$ will force $x_{\eta_1}^1$ to turn blue which shows that all vertices of C_1 will eventually be blue.

Now assume that all vertices of C_i for each $i \in [\alpha]$ where $\alpha \in [k]$ are blue by some

timestep β and consider C_j where $j = \alpha + 1$. Therefore, there exists $e = x_r^i x_s^j \in M'$ such that x_r^i is on C_i and x_s^j is on C_j where $i < j$. Moreover, we have indexed C_j such that $x_s^j = x_1^j$. Since x_r^i is blue and x_1^j is its only white neighbor at timestep β , then x_1^j will become blue. Furthermore, $x_2^j \in S$ so it is blue, $x_{n_j}^j$ will become blue as it is the only white neighbor of x_1^j and x_2^j will force x_3^j to turn blue. If $x_4^j \in S$, then propagation will continue (in a clockwise fashion) around C_j . So assume $x_4^j \notin S$. Thus, x_4^j is on some edge $x_4^j x_s^\ell \in M$ where $\ell \leq j$. If $j = \ell$, then $x_4^j \in S$, which is a contradiction. Thus, we shall assume that $\ell < j$. By assumption, all vertices of C_ℓ are blue so x_4^j will also become blue. Hence, propagation will continue until we reach $x_{\eta_j}^j$. Similarly, moving in the opposite direction from $x_{n_j}^j$ to $x_{\eta_j}^j$, all vertices of the form x_s^j where $\eta_j + 1 \leq s \leq n_j$ will eventually become blue. Hence, either $x_{\eta_j-1}^j$ or $x_{\eta_j+1}^j$ will force $x_{\eta_j}^j$ to become blue which shows that all vertices of C_j will eventually become blue. It follows that S is indeed a zero forcing set of G' .

As noted before, we can obtain G from G' by replacing some of the edges in G' which were originally in H with strings of diamonds. Choose an interior vertex from each diamond in every string of diamonds used to replace edges in G' to obtain G and call the resulting set X . One can easily verify that $S' = S \cup X$ is a zero forcing set of G of cardinality $|S'| = |D| + k + 1 + \frac{n(G) - n(G')}{4}$. We claim that $|S'| \leq \lceil \frac{5}{18}n(G) \rceil + 1$.

We shall assume for the time being that $k \geq 3$. Assume first that \mathcal{C} contains two cycles C_i and C_j such that C_i and C_j are adjacent and there exist $\ell, \ell' \in [k] - \{i, j\}$ such that $\ell \neq \ell'$, C_i is adjacent to C_ℓ and C_j is adjacent to $C_{\ell'}$. Moreover, reindexing if necessary, we may assume $i = 1$ and $j = 2$, T is rooted at u_1 (which corresponds to cycle C_1), and $\{u_1 u_\ell, u_2 u_{\ell'}, u_1 u_2\} \subset E(T)$. Furthermore, we may assume C_ℓ is chosen so that $x_{\eta_1}^1 x_1^\ell \in M'$ and $C_{\ell'}$ is chosen so that $x_{\eta_2}^2 x_1^{\ell'} \in M'$. Note that each edge in M can be viewed as a matching between triangles in G' . Furthermore, for each pair of triangles other than the one containing $\{x_1^1, x_1^2, x_{\eta_1}^1, x_1^2, x_2^2, x_{\eta_2}^2\}$, there are at most two vertices from S' . Now we can partition M into M_1 and M_2 such that if $e \in M_1$, then the corresponding triangle pair containing e contains at most one vertex from S' and if $e \in M_2$, then the corresponding triangle pair containing e contains at least two vertices from S' . We claim that $|M_2| - 1 \leq \frac{n(G') - 18}{12}$. Note first that if e is incident to a vertex of the form $x_{\eta_i}^i$, then the corresponding triangle pair contains exactly one vertex from S' and $e \in M_1$. Moreover, if $e \in M_2$, then e is incident to x_1^s for some $s \in [k]$ and the triangle pair associated with e contains vertices of the form $x_j^t, x_{j+1}^t, x_{j-1}^t$ where $t < s$, $j \neq \eta_t$, $x_j^t \in D$ and $x_s^s \in S'$. Enumerate the edges of M_2 as $\{e_1, \dots, e_r\}$ where $e_1 = x_1^1 x_1^2$. We may assume for $2 \leq i \leq r$ that e_i is incident to $x_1^{\alpha_i}$. For each $i \in [r]$, let T_i represent the six vertices of the triangle pair associated with e_i . Consider the set $\{W_1, \dots, W_r\}$ where

$$W_1 = T_1 \cup \{x_{\eta_1}^1, x_{\eta_1-1}^1, x_{\eta_1+1}^1, x_1^\ell, x_2^\ell, x_{n_\ell}^\ell, x_{\eta_2}^2, x_{\eta_2-1}^2, x_{\eta_2+1}^2, x_1^{\ell'}, x_2^{\ell'}, x_{n_{\ell'}}^{\ell'}\}$$

where $x_{\eta_1}^1 x_1^\ell \in E(G)$ and $x_{\eta_2}^2 x_1^{\ell'} \in E(G)$, and

$$W_i = T_i \cup \{x_{\eta_{\alpha_i}}^{\alpha_i}, x_{\eta_{\alpha_i}-1}^{\alpha_i}, x_{\eta_{\alpha_i}+1}^{\alpha_i}, x_1^{\beta_i}, x_2^{\beta_i}, x_{n_{\beta_i}}^{\beta_i}\}$$

for $2 \leq i \leq r$ where $x_{\eta_{\alpha_i}}^{\alpha_i} x_1^{\beta_i} \in E(G)$ and u_{α_i} is not a leaf in T , and otherwise

$$W_i = T_i \cup \{x_{\eta_{\alpha_i}}^{\alpha_i}, x_{\eta_{\alpha_i}-1}^{\alpha_i}, x_{\eta_{\alpha_i}+1}^{\alpha_i}, x_s^j, x_{s-1}^j, x_{s+1}^j\}$$

for $2 \leq i \leq r$ where u_{α_i} is a leaf in T and $x_{\eta_{\alpha_i}}^{\alpha_i} x_s^j \in E(G)$ where $u_{\alpha_i} u_j \notin E(T)$. That is, $x_{\eta_{\alpha_i}}^{\alpha_i}$ is in a triangle pair containing vertices of C_j where the spanning tree T of J does not contain an edge between corresponding vertices u_{α_i} and u_j .

One can easily verify that $W_i \cap W_j = \emptyset$ for $i \neq j$ and by construction, $|W_i| = 12$ for $2 \leq i \leq r$ and $|W_1| = 18$. It follows that $r - 1 \leq \frac{n(G') - 18}{12}$ which implies $r + 2 \leq \frac{n(G')}{12} + \frac{3}{2}$. Hence, S' contains at most one vertex from each triangle pair associated with an edge in M_1 , there are $r - 1$ triangle pairs that contain exactly two vertices from S' , and there is exactly one triangle pair that contains exactly 4 vertices. Thus,

$$\begin{aligned} |S'| &\leq \frac{n(G')}{6} + (r - 1) + 3 + \frac{n(G) - n(G')}{4} \\ &\leq \frac{n(G')}{6} + r + 2 + \frac{n(G) - n(G')}{4} \\ &\leq \frac{n(G')}{6} + \frac{n(G')}{12} + \frac{3}{2} + \frac{n(G) - n(G')}{4} \\ &= \frac{n(G)}{4} + \frac{3}{2} \\ &\leq \frac{5n(G)}{18} + 1 \end{aligned}$$

provided that $n(G) \geq 18$. On the other hand, we have assumed \mathcal{C} contains a cycle of length 9 or more and $k \geq 3$. Thus, $n(G) \geq n(G') \geq 21$.

Therefore, we shall assume that \mathcal{C} does not contain two cycles C_i and C_j such that C_i is adjacent to C_j and there exist $\ell, \ell' \in [k] - \{i, j\}$ where C_i is adjacent to C_ℓ and C_j is adjacent to $C_{\ell'}$. It follows that we may write $\mathcal{C} = C_1 \cup \dots \cup C_k$ where one of the following is true: (a) $k = 3$ and C_1 is adjacent to both C_2 and C_3 and C_2 is adjacent to C_3 , or (b) $k \geq 3$ and C_i is only adjacent to C_1 for each $2 \leq i \leq k$. In case (b), a similar argument to that used above will show $|M_2| - 1 \leq \frac{n(G') - 18}{12}$ where W_i is as defined above for $2 \leq i \leq r$ and

$$W_1 = T_1 \cup \{x_{\eta_1}^1, x_{\eta_1-1}^1, x_{\eta_1+1}^1, x_1^\ell, x_2^\ell, x_{n_\ell}^\ell, x_{\eta_2}^2, x_{\eta_2-1}^2, x_{\eta_2+1}^2, x_r^j, x_{r+1}^j, x_{r-1}^j\}$$

where $x_{\eta_1}^1 x_1^\ell \in E(G)$ and $x_{\eta_2}^2 x_r^j \in E(G)$. Therefore, we assume that we are in case (a). In this case, we may assume that $T = P_3$ where u_1 has degree 2. Thus,

$x_{\eta_1}^1 x_1^3 \in M'$, $x_1^1 x_1^2 \in M'$ and we may assume $x_{\eta_2}^2 x_{\eta_3}^3 \in M$. Thus, the triangle pair associated with $x_{\eta_2}^2 x_{\eta_3}^3$ does not contain a vertex of S' . Moreover, $M_2 = \{e_1\}$ and

$$\begin{aligned} |S'| &\leq \frac{n(G') - 6}{6} + 3 + \frac{n(G) - n(G')}{4} \\ &= \frac{n(G')}{6} + 2 + \frac{n(G) - n(G')}{4} \\ &= \frac{n(G)}{4} + 2 - \frac{n(G')}{12}. \end{aligned}$$

Since we have assumed \mathcal{C} contains a cycle of length 9 or more and $k = 3$, $n(G') \geq 21$ and we have

$$|S'| \leq \frac{n(G)}{4} + 2 - \frac{21}{12} = \frac{n(G)}{4} + \frac{1}{4} \leq \frac{5n(G)}{18} + 1.$$

Finally, assume that $k \in \{1, 2\}$. If $k = 2$, then again we may assume $x_{\eta_1}^1 x_{\eta_2}^2 \in M$, $M_2 = \{e_1\}$, and the associated triangle pair contains no vertex of S' . Moreover, $n(G') \geq 15$ as we have assumed \mathcal{C} contains a cycle of length 9 or more and $k \geq 2$. Thus,

$$\begin{aligned} |S'| &\leq \frac{n(G') - 6}{6} + 3 + \frac{n(G) - n(G')}{4} \\ &= \frac{n(G')}{6} + 2 + \frac{n(G) - n(G')}{4} \\ &= \frac{n(G)}{4} + 2 - \frac{n(G')}{12} \\ &\leq \frac{n(G)}{4} + 2 - \frac{15}{12} \\ &= \frac{n(G)}{4} + \frac{3}{4} \\ &\leq \frac{5n(G)}{18} + 1. \end{aligned}$$

Lastly, we shall assume $k = 1$ and let

$$S = \{x_j^1 : j \equiv 1 \pmod{3} \text{ and } x_j^1 x_r^1 \in M \text{ with } r > j\} \cup \{x_{n_1}^1, x_2^1\}.$$

Thus,

$$\begin{aligned} |S'| &= \frac{n(G')}{6} + 2 + \frac{n(G) - n(G')}{4} \\ &= \frac{n(G)}{4} + 2 - \frac{n(G')}{12}. \end{aligned}$$

If $n(G') \geq 12$, then we are done. So assume $n(G') \in \{6, 9\}$. Note that $n(G') \neq 9$ for otherwise there is no perfect matching between $\{x_1^1, x_4^1, x_7^1\}$. On the other hand, if $n(G') = 6$, then \mathcal{C} does not contain a cycle of length 9 or more, which is a contradiction. □

References

- [1] AIM Minimum Rank - Special Graphs Work Group. Zero forcing sets and the minimum rank of graphs. *Linear Algebra Appl.*, **428**(7): 1628 - 1648 (2008)
- [2] S. E. Anderson and K. Kuenzel. Power domination in cubic graphs and Cartesian products. *Discrete Math.*, accepted (2022)
- [3] F. Barioli, W. Barrett, S. M. Fallat, H. T. Hall, L. Hogben, B. Shader, P. van den Driessche, and H. van der Holst. Parameters related to tree-width, zero forcing, and maximum nullity of a graph. *J. Graph Theory*, **72** no. 2: 146–177 (2013)
- [4] B. Brešar, T. Gologranc, M. Milanič, D. F. Rall, and R. Rizzi. Dominating sequences in graphs. *Discrete Math.*, **336**: 22–36 (2014)
- [5] B. Brešar, C. Bujtás, T. Gologranc, S. Klavžar, G. Košmrlj, B. Patkós, Z. Tuza, and M. Vizer. Grundy dominating sequences and zero forcing sets. *Discrete Optim.*, **26**: 66–77 (2017)
- [6] R. Davila and M. A. Henning. Zero forcing in claw-free cubic graphs. *Bulletin Malaysian Math. Sciences Society*, **43**: 673-688 (2020)
- [7] G. Hernández, J. Ranilla, S. Ranilla-Cortina. Zero forcing in triangulations. *J. Comput. Appl. Math.*, **354**: 123–130 (2019)
- [8] L. Hogben. Minimum rank problems. *Linear Algebra Appl.*, **432**: 1961-1974 (2010)
- [9] J. C.-H. Lin. Zero forcing number, Grundy domination number, and their variants. *Linear Algebra Appl.*, **563**: 240–254 (2019)
- [10] S. Oum. Perfect matchings in claw-free cubic graphs, *Electron. J. Combin.*, **18** P.62 (2011)
- [11] J. Plesník. Connectivity of regular graphs and the existence of 1-factors, *Matematický časopis* **22**(4): 310–318 (1972)