

Second-order optimization with lazy Hessians*

Nikita Doikov[†]

El Mahdi Chayti[‡]

Martin Jaggi[§]

EPFL

December 1, 2022

Abstract

We analyze Newton’s method with lazy Hessian updates for solving general possibly non-convex optimization problems. We propose to reuse a previously seen Hessian for several iterations while computing new gradients at each step of the method. This significantly reduces the overall arithmetical complexity of second-order optimization schemes. By using the cubic regularization technique, we establish fast global convergence of our method to a second-order stationary point, while the Hessian does not need to be updated each iteration. For convex problems, we justify global and local superlinear rates for lazy Newton steps with quadratic regularization, which is easier to compute. The optimal frequency for updating the Hessian is once every d iterations, where d is the dimension of the problem. This provably improves the total arithmetical complexity of second-order algorithms by a factor \sqrt{d} .

1 Introduction

Motivation. Second-order optimization algorithms are being widely used for solving difficult ill-conditioned problems. The classical Newton method approximates the objective function by its second-order Taylor approximation at the current point. A minimizer of this model serves as one step of the algorithm.

Locally, Newton’s method achieves very fast quadratic rate [20], when the iterates are in a neighborhood of the solution. However, it may not converge if the initial point is far away from the optimum. For the global convergence of Newton’s method, we need to use some of the various regularization techniques, that have been intensively developed during recent decades (including *damped Newton steps* with line search [20, 30, 18], *trust-region* methods [6], *self-concordant* analysis [27, 2, 34, 13] or the notion of *Hessian stability* [21], *cubic regularization* [28], *contracting-point* steps [9, 12], or *gradient regularization* techniques [25, 10, 8]).

In all of these approaches, the arithmetic complexity of each step remains costly, since it requires to compute both the gradient and Hessian of the objective function, and then perform

*This work was supported by the Swiss State Secretariat for Education, Research and Innovation (SERI) under contract number 22.00133.

[†]nikita.doikov@epfl.ch, equal contribution.

[‡]el-mahdi.chayti@epfl.ch, equal contribution.

[§]martin.jaggi@epfl.ch

some factorization of the matrix. Hessian computations are prohibitively expensive in any large scale setting.

In this work, we explore a simple possibility for a significant acceleration of second-order schemes in terms of their total arithmetical complexity. The idea is to keep a previously computed Hessian (a stale version of the Hessian) for several iterations, while using fresh gradients in each step of the method. Therefore, we can save a lot of computational resources reusing the old Hessians. We call this *lazy Hessian updates*. We justify that this idea can be successfully employed in most established second-order schemes, resulting in a provable improvement of the overall performance.

Lazy Hessian Updates: compute new Hessian once per m iterations.

Hessians:	$\nabla^2 f(\mathbf{x}_0)$	reuse Hessian \longrightarrow			$\nabla^2 f(\mathbf{x}_m)$	reuse Hessian \longrightarrow	
Gradients:	$\nabla f(\mathbf{x}_0)$	$\nabla f(\mathbf{x}_1)$	\dots	$\nabla f(\mathbf{x}_{m-1})$	$\nabla f(\mathbf{x}_m)$	$\nabla f(\mathbf{x}_{m+1})$	\dots

For general problems¹, the optimal schedule is $m := d$ (update the Hessian every d iterations), where d is the dimension of the problem. In these cases, we gain a provable computational acceleration of our methods by a factor \sqrt{d} .

The key to our analysis is an increased value of the regularization constant. In a general case, this should be proportional to mL , where L is the Lipschitz constant of the Hessian. Thus, for the lazy Newton steps, we should regularize our model stronger to balance the possible errors coming from the Hessian inexactness, and so the steps of the method become shorter. Fortunately, we save much more in terms of the computational resources, which makes our approach appealing.

Related Work. The idea of using an old Hessian for several steps of Newton’s method probably appeared for the first time in [33], with the study of local convergence of this process as applied to solving systems of nonlinear equations. The author proved a local quadratic rate for the iterates when updating the Hessian, and a linear rate with improving factor for the iterates in between.

Later on, this idea has been successfully combined with Levenberg-Marquardt [23, 24] regularization approach in [14], with damped Newton steps in [22, 35], and with the proximal Newton-type methods for convex optimization in [1]. In these papers, it was established that the methods possess an asymptotic global convergence to a solution, without explicit non-asymptotic bounds on the actual rate.

Comparing with these works, our new second-order algorithms with the lazy Hessian updates use a different analysis that is based on the modern globalization techniques (*cubic regularization* [28] and *gradient regularization* [25, 10]). It allows to equip our methods with provably fast

¹Assuming that the cost of computing one Hessian and its appropriate factorization is d times the cost of one gradient step. See Examples 1, 2, and 3.

global rates for wide classes of convex and non-convex optimization problems. Thus, we prove a global complexity $\mathcal{O}(1/\varepsilon^{3/2})$ of the lazy regularized Newton steps for achieving a second-order stationary point with the gradient norm bounded by ε .

Another interesting connection can be established to recently developed distributed Newton-type methods [19], where the authors propose to use a probabilistic aggregation and compression of the Hessians, as for example in federated learning. In the particular case of the single node setup, their algorithm also evaluates the Hessian rarely which is similar in spirit to our approach but using different aggregation strategies. In their technique, the node needs to consider the Hessian each iteration. Then, using a certain criterion, it decides whether to employ the new Hessian for the next step or to keep using just the old information, while in our approach the Hessian is computed only *once* per m iterations. The authors proved local linear and superlinear rates that are independent on the condition number. At the same time, in our paper, we primarily focus on global complexity guarantees and thus establish a provable computational improvement by a factor \sqrt{d} for our methods as compared to updating the Hessian in every step. We also justify local superlinear rates for our approach.

Contributions. In Section 2 we propose the lazy Newton step with cubic regularization. It uses the gradient computed at the current point and the Hessian at some different point from the past trajectory. We study the main properties of this step and formulate the progress in terms of the objective function and the new gradient norm (Theorem 1).

The lazy Newton step is formulated in our Algorithm 1 in Section 3. We prove its fast global convergence in terms of the gradient norm (Theorem 2). In the case $m := 1$ (updating the Hessian each iteration), our rate recovers the classical rate of full Cubic Newton [28]. Then, we show that taking into account the actual arithmetic cost of the Hessian computations, the optimal choice for one phase of the method is $m := d$, which improves the total complexity of Cubic Newton by a factor \sqrt{d} (see Corollary 3). We also justify convergence to a *second-order stationary point* with a guarantee on the smallest eigenvalue of the Hessian. This avoids that our method could get stuck in saddle points.

We propose an improved method for solving *convex problems* in Section 4. Our Algorithm 2 is based on the *gradient regularization* technique [25, 10], which replaces the cubic regularizer in the model by quadratic one. The regularization parameter then should be proportional to the square root of the current gradient norm. This makes the subproblem much easier to solve, involving just one standard matrix inversion, while keeping the fast rate of the original cubically regularized method. We show that gradient regularization can be successfully coupled with the lazy Hessian updates. The global complexity of the algorithm is established in Theorem 4.

In Section 5 we study the local behaviour of our methods. We prove local quadratic convergence for the Cubic Newton with lazy Hessians (Theorem 5), and superlinear convergence with factor $3/2$ for the method with gradient regularization (Theorem 6). When the current point is in a neighbourhood of a non-degenerate solution, regularization of the Newton method is not needed since locally the best quadratic model of the objective is the pure second-order Taylor polynomial. Thus, our analysis also justifies local quadratic convergence for the *classical Newton method* (when regularization parameter is equal to zero).

Section 6 is devoted to effective implementation of our algorithms. In Section 6.1 we explain

how to use matrix factorizations for recomputing the lazy Newton steps efficiently. In Section 6.2 we show how to equip our methods with adaptive search, which frees the methods from the need to know the actual Lipschitz constant of the Hessian. The cost of using such adaptive search is just one extra lazy step on average. At the same time, the methods preserve their global and local convergence guarantees.

Section 7 contains preliminary numerical experiments. We discuss our results in Section 8.

Notation. We consider the following optimization problem,

$$\min_{\mathbf{x} \in \mathbb{R}^d} f(\mathbf{x}), \quad (1)$$

where f is a several times differentiable function, not necessarily convex. We denote by $\nabla f(\mathbf{x}) \in \mathbb{R}^d$ its gradient and by $\nabla^2 f(\mathbf{x}) \in \mathbb{R}^{d \times d}$ its Hessian, computed at some point $\mathbf{x} \in \mathbb{R}^d$.

Our analysis will apply both if optimization is over the standard Euclidean space, and also more generally if a different norm is defined by an arbitrary fixed symmetric positive definite matrix $\mathbf{B} = \mathbf{B}^\top \succ 0$. In that case the norm becomes

$$\|\mathbf{x}\| \stackrel{\text{def}}{=} \langle \mathbf{B}\mathbf{x}, \mathbf{x} \rangle^{1/2}, \quad \mathbf{x} \in \mathbb{R}^d. \quad (2)$$

Thus, the matrix \mathbf{B} is responsible for fixing the coordinate system in our problem. In the simplest case, we can choose $\mathbf{B} := \mathbf{I}$ (identity matrix), which recovers the classical Euclidean geometry. A more advanced choice of \mathbf{B} can take into account a specific structure of the problem (see Example 1).

The norm for the dual objects (gradients) is defined in the standard way,

$$\|\mathbf{g}\|_* \stackrel{\text{def}}{=} \sup_{\mathbf{x}: \|\mathbf{x}\| \leq 1} \langle \mathbf{g}, \mathbf{x} \rangle = \langle \mathbf{g}, \mathbf{B}^{-1}\mathbf{g} \rangle^{1/2}, \quad \mathbf{g} \in \mathbb{R}^d.$$

The induced norm for matrices is

$$\|\mathbf{A}\| \stackrel{\text{def}}{=} \sup_{\substack{\mathbf{x}, \mathbf{y}: \\ \|\mathbf{x}\| \leq 1, \|\mathbf{y}\| \leq 1}} \langle \mathbf{A}\mathbf{x}, \mathbf{y} \rangle = \sup_{\mathbf{x}: \|\mathbf{x}\| \leq 1} \|\mathbf{A}\mathbf{x}\|_*, \quad \mathbf{A} \in \mathbb{R}^{d \times d}.$$

We assume that the Hessian of f is Lipschitz continuous, with some constant $L > 0$:

$$\|\nabla^2 f(\mathbf{x}) - \nabla^2 f(\mathbf{y})\| \leq L\|\mathbf{x} - \mathbf{y}\|, \quad \forall \mathbf{x}, \mathbf{y} \in \mathbb{R}^d. \quad (3)$$

The immediate consequences are the following inequalities, valid for all $\mathbf{x}, \mathbf{y} \in \mathbb{R}^d$:

$$\|\nabla f(\mathbf{y}) - \nabla f(\mathbf{x}) - \nabla^2 f(\mathbf{x})(\mathbf{y} - \mathbf{x})\|_* \leq \frac{L}{2}\|\mathbf{x} - \mathbf{y}\|^2, \quad (4)$$

$$|f(\mathbf{y}) - f(\mathbf{x}) - \langle \nabla f(\mathbf{x}), \mathbf{y} - \mathbf{x} \rangle - \frac{1}{2}\langle \nabla^2 f(\mathbf{x})(\mathbf{y} - \mathbf{x}), \mathbf{y} - \mathbf{x} \rangle| \leq \frac{L}{6}\|\mathbf{y} - \mathbf{x}\|^3. \quad (5)$$

2 Lazy Newton Steps

Let us introduce the following lazy Newton steps, where we allow the gradient and Hessian to be computed at different points $\mathbf{x}, \mathbf{z} \in \mathbb{R}^d$. We use cubic regularization of our model with a parameter $M > 0$:

$$\mathbf{T}_M(\mathbf{x}, \mathbf{z}) \in \underset{\mathbf{y} \in \mathbb{R}^d}{\text{Argmin}} \left\{ \langle \nabla f(\mathbf{x}), \mathbf{y} - \mathbf{x} \rangle + \frac{1}{2} \langle \nabla^2 f(\mathbf{z})(\mathbf{y} - \mathbf{x}), \mathbf{y} - \mathbf{x} \rangle + \frac{M}{6} \|\mathbf{y} - \mathbf{x}\|^3 \right\}. \quad (6)$$

For $\mathbf{z} := \mathbf{x}$ this is the iteration of Cubic Newton [28]. Thus, in our scheme, we can reuse the Hessian from a previous step \mathbf{z} without recomputing it, which significantly reduces the overall iteration cost.

Our definition implies that the point $\mathbf{T} = \mathbf{T}_M(\mathbf{x}, \mathbf{z})$ is a *global minimum* of the cubically regularized model, which is generally non-convex. However, it turns out that we can compute this point efficiently by using standard techniques developed initially for trust-region methods [6].

Let us denote $r \stackrel{\text{def}}{=} \|\mathbf{T} - \mathbf{x}\|$. The solution to the subproblem (6) satisfies the following stationarity conditions (see [28]):

$$\nabla f(\mathbf{x}) + \nabla^2 f(\mathbf{z})(\mathbf{T} - \mathbf{x}) + \frac{Mr}{2} \mathbf{B}(\mathbf{T} - \mathbf{x}) = \mathbf{0}, \quad (7)$$

and

$$\nabla^2 f(\mathbf{z}) + \frac{Mr}{2} \mathbf{B} \succeq \mathbf{0}. \quad (8)$$

Thus, in the non-degenerate case, one step can be represented in the following form:

$$\mathbf{T} = \mathbf{x} - (\nabla^2 f(\mathbf{z}) + \frac{Mr}{2} \mathbf{B})^{-1} \nabla f(\mathbf{x}), \quad (9)$$

and the value $r > 0$ can be found by solving the corresponding univariate nonlinear equation (see [28, Section 5]). It can be done very efficiently from a precomputed *eigenvalue* or the *tridiagonal* decomposition of the Hessian. These decompositions typically require $\mathcal{O}(d^3)$ arithmetic operations, which is of similar cost as for matrix inversion in the classical Newton step. We discuss the computation of the new iterate $\mathbf{T}_M(\mathbf{x}, \mathbf{z})$ in more details in Section 6.1.

Now, let us express the progress achieved by the proposed step (6). Firstly, we can relate the length of the step and the gradient at the new point as follows:

$$\begin{aligned} \frac{Lr^2}{2} &\stackrel{(4)}{\geq} \|\nabla f(\mathbf{T}) - \nabla f(\mathbf{x}) - \nabla^2 f(\mathbf{x})(\mathbf{T} - \mathbf{x})\|_* \\ &\stackrel{(7)}{=} \|\nabla f(\mathbf{T}) + \frac{Mr}{2} \mathbf{B}(\mathbf{T} - \mathbf{x}) + (\nabla^2 f(\mathbf{z}) - \nabla^2 f(\mathbf{x}))(\mathbf{T} - \mathbf{x})\|_* \\ &\geq \|\nabla f(\mathbf{T})\|_* - \frac{Mr^2}{2} - r \|\nabla^2 f(\mathbf{z}) - \nabla^2 f(\mathbf{x})\|. \end{aligned}$$

Rearranging terms and using Lipschitz continuity of the Hessian, we get

$$\begin{aligned} \|\nabla f(\mathbf{T})\|_* &\leq \frac{M+L}{2} r^2 + r \|\nabla^2 f(\mathbf{z}) - \nabla^2 f(\mathbf{x})\| \\ &\stackrel{(3)}{\leq} \frac{M+L}{2} r^2 + Lr \|\mathbf{z} - \mathbf{x}\|. \end{aligned}$$

Thus, we have established

Lemma 1 For any $M > 0$, it holds that

$$\|\nabla f(\mathbf{T})\|_* \leq \frac{M+L}{2}r^2 + Lr\|\mathbf{z} - \mathbf{x}\|. \quad (10)$$

Secondly, we have the following progress in terms of the objective function.

Lemma 2 For any $M > 0$, it holds that

$$f(\mathbf{x}) - f(\mathbf{T}) \geq \frac{5M-4L}{24}r^3 - \frac{32L^3}{3M^2}\|\mathbf{z} - \mathbf{x}\|^3. \quad (11)$$

Proof:

Indeed, multiplying (7) by the vector $\mathbf{T} - \mathbf{x}$, we get

$$\begin{aligned} \langle \nabla f(\mathbf{x}), \mathbf{T} - \mathbf{x} \rangle &= -\langle \nabla^2 f(\mathbf{z})(\mathbf{T} - \mathbf{x}), \mathbf{T} - \mathbf{x} \rangle - \frac{M}{2}r^3 \\ &\stackrel{(8)}{\leq} -\frac{1}{2}\langle \nabla^2 f(\mathbf{z})(\mathbf{T} - \mathbf{x}), \mathbf{T} - \mathbf{x} \rangle - \frac{M}{4}r^3. \end{aligned} \quad (12)$$

Hence, using the global upper bound on the objective, we obtain

$$\begin{aligned} f(\mathbf{T}) &\stackrel{(5)}{\leq} f(\mathbf{x}) + \langle \nabla f(\mathbf{x}), \mathbf{T} - \mathbf{x} \rangle + \frac{1}{2}\langle \nabla^2 f(\mathbf{x})(\mathbf{T} - \mathbf{x}), \mathbf{T} - \mathbf{x} \rangle + \frac{L}{6}r^3 \\ &\stackrel{(12)}{\leq} f(\mathbf{x}) - \frac{M}{4}r^3 + \frac{L}{6}r^3 + \frac{1}{2}\langle (\nabla^2 f(\mathbf{x}) - \nabla^2 f(\mathbf{z}))(\mathbf{T} - \mathbf{x}), \mathbf{T} - \mathbf{x} \rangle \\ &\stackrel{(3)}{\leq} f(\mathbf{x}) - \frac{M}{4}r^3 + \frac{L}{6}r^3 + \frac{L}{2}r^2\|\mathbf{z} - \mathbf{x}\|. \end{aligned}$$

Applying Young's inequality for the last term,

$$\frac{L}{2}r^2\|\mathbf{z} - \mathbf{x}\| = \left(\frac{M^2/3}{2 \cdot 32^{1/3}}r^2\right) \cdot \left(\frac{32^{1/3}L}{M^2/3}\|\mathbf{z} - \mathbf{x}\|\right) \leq \frac{M}{24}r^3 + \frac{32L^3}{3M^2}\|\mathbf{z} - \mathbf{x}\|^3,$$

we obtain (11). \square

We can also bound the smallest eigenvalue of the new Hessian, which will be crucial to understand the behaviour for non-convex objectives. Indeed, using Lipschitz continuity of the Hessian and the triangle inequality, we have:

$$\begin{aligned} \nabla^2 f(\mathbf{T}) &\succeq \nabla^2 f(\mathbf{z}) - L\|\mathbf{z} - \mathbf{T}\|\mathbf{B} \\ &\succeq \nabla^2 f(\mathbf{z}) - (L\|\mathbf{z} - \mathbf{x}\| + Lr)\mathbf{B} \\ &\stackrel{(8)}{\succeq} -\left(\frac{M}{2}r + L\|\mathbf{z} - \mathbf{x}\| + Lr\right)\mathbf{B}. \end{aligned} \quad (13)$$

Let us denote the following quantity, for any $y \in \mathbb{R}^d$:

$$\xi(\mathbf{y}) \stackrel{\text{def}}{=} \left[-\lambda_{\min}(\mathbf{B}^{-1/2}\nabla^2 f(\mathbf{y})\mathbf{B}^{-1/2})\right]_+, \quad (14)$$

where $[t]_+ \stackrel{\text{def}}{=} \max\{t, 0\}$ denotes the positive part, and $\lambda_{\min}(\cdot)$ is the smallest eigenvalue of a symmetric matrix.

If $\nabla^2 f(\mathbf{y}) \succeq 0$ for a certain $\mathbf{y} \in \mathbb{R}^d$, then $\xi(\mathbf{y}) = 0$. Otherwise, $\xi(\mathbf{y})$ shows how big (in absolute value) the smallest eigenvalue of the Hessian is with respect to a fixed matrix $\mathbf{B} \succ 0$.

Thus, due to (13), we can control the quantity ξ as follows:

Lemma 3 *For any $M > 0$, it holds*

$$\xi(\mathbf{T}) \leq \frac{M+2L}{2}r + L\|\mathbf{z} - \mathbf{x}\|. \quad (15)$$

Now, we can combine (10) and (15) with (11) to express the functional progress of one step using the new gradient norm and the smallest eigenvalue of the new Hessian.

Theorem 1 *Let $M \geq L$. Then, for one cubic step (6), it holds*

$$f(\mathbf{x}) - f(\mathbf{T}) \geq \max\left\{\frac{1}{648M^2}\xi(\mathbf{T})^3, \frac{1}{72\sqrt{2}M}\|\nabla f(\mathbf{T})\|_*^{3/2}\right\} + \frac{M}{48}r^3 - \frac{11L^3}{M^2}\|\mathbf{z} - \mathbf{x}\|^3. \quad (16)$$

Proof:

Indeed, using convexity of the function $t \mapsto t^3$ for $t \geq 0$, we obtain

$$\xi(\mathbf{T})^3 \stackrel{(15)}{\leq} \left(\frac{3}{2}Mr + L\|\mathbf{z} - \mathbf{x}\|\right)^3 \leq \frac{27}{2}M^3r^3 + 4L^3\|\mathbf{z} - \mathbf{x}\|^3.$$

Hence,

$$\frac{Mr^3}{2} \geq \frac{1}{27M^2}\xi(\mathbf{T})^3 - \frac{4L^3}{27M^2}\|\mathbf{z} - \mathbf{x}\|^3,$$

which gives the first part of the maximum when substituting into (11).

Then, using convexity of the function $t \mapsto t^{3/2}$ for $t \geq 0$ and Young's inequality, we get

$$\begin{aligned} \|\nabla f(\mathbf{T})\|_*^{3/2} &\stackrel{(10)}{\leq} \left(Mr^2 + Lr\|\mathbf{z} - \mathbf{x}\|\right)^{3/2} \leq \sqrt{2}M^{3/2}r^3 + \sqrt{2}(Lr\|\mathbf{z} - \mathbf{x}\|)^{3/2} \\ &\leq \sqrt{2}M^{3/2}r^3 + \frac{\sqrt{2}M^{3/2}}{2}r^3 + \frac{\sqrt{2}L^3}{2M^{3/2}}\|\mathbf{z} - \mathbf{x}\|^3 = \frac{3\sqrt{2}M^{3/2}}{2}r^3 + \frac{L^3}{\sqrt{2}M^{3/2}}\|\mathbf{z} - \mathbf{x}\|^3. \end{aligned}$$

Therefore,

$$\frac{Mr^3}{2} \geq \frac{1}{3\sqrt{2}M}\|\nabla f(\mathbf{T})\|_*^{3/2} - \frac{L^3}{6M^2}\|\mathbf{z} - \mathbf{x}\|^3.$$

Substituting this bound into (11) completes the proof. \square

Theorem 1 shows how much progress we can expect from one lazy Newton step with cubic regularization. The price for using the lazy Hessian is the last term in the progress bound (16), which vanishes if $\mathbf{z} := \mathbf{x}$ (updating the Hessian for the current iterate). In the next section, we will show how to balance this term by taking into account several consecutive steps of the method and by a clever choice of the regularization parameter.

3 Global Convergence Rates

Let us consider one phase of the potential algorithm, when we compute the Hessian once at a current point $\mathbf{z} := \mathbf{x}_0$ and then perform m cubic steps (6) in a row:

$$\begin{aligned} \mathbf{x}_1 &= \mathbf{T}_M(\mathbf{x}_0, \mathbf{z}), \\ \mathbf{x}_2 &= \mathbf{T}_M(\mathbf{x}_1, \mathbf{z}), \\ &\dots \\ \mathbf{x}_m &= \mathbf{T}_M(\mathbf{x}_{m-1}, \mathbf{z}). \end{aligned} \tag{17}$$

We use some fixed regularization constant $M > 0$.

Let us denote the corresponding step lengths by $r_{k+1} := \|\mathbf{x}_{k+1} - \mathbf{x}_k\|$. Then, according to Theorem 1, for each $0 \leq k \leq m-1$, we have the following progress in terms of the objective function:

$$f(\mathbf{x}_k) - f(\mathbf{x}_{k+1}) \stackrel{(16)}{\geq} \max\left\{\frac{1}{648M^2}\xi(\mathbf{x}_{k+1})^3, \frac{1}{72\sqrt{2}M}\|\nabla f(\mathbf{x}_{k+1})\|_*^{3/2}\right\} + \frac{M}{48}r_{k+1}^3 - \frac{11L^3}{M^2}\|\mathbf{x}_0 - \mathbf{x}_k\|^3.$$

Telescoping this bound for different k , and using triangle inequality for the last negative term,

$$\|\mathbf{x}_0 - \mathbf{x}_k\| \leq \sum_{i=1}^k r_i,$$

we get

$$\begin{aligned} &f(\mathbf{x}_0) - f(\mathbf{x}_m) \\ &\geq \sum_{k=1}^m \max\left\{\frac{1}{648M^2}\xi(\mathbf{x}_k)^3, \frac{1}{72\sqrt{2}M}\sum_{k=1}^m \|\nabla f(\mathbf{x}_k)\|_*^{3/2}\right\} + \frac{M}{48}\sum_{k=1}^m r_k^3 - \frac{11L^3}{M^2}\sum_{k=1}^{m-1}\left(\sum_{i=1}^k r_i\right)^3. \end{aligned} \tag{18}$$

It remains to use the following simple technical lemma.

Lemma 4 *For any sequence of positive numbers $\{r_k\}_{k \geq 1}$, it holds for any $m \geq 1$:*

$$\sum_{k=1}^{m-1}\left(\sum_{i=1}^k r_i\right)^3 \leq \frac{m^3}{3}\sum_{k=1}^{m-1} r_k^3 \tag{19}$$

Proof:

We prove (19) by induction. It is obviously true for $m = 1$, which is our base.

Assume that it holds for some arbitrary $m \geq 1$. Then

$$\sum_{k=1}^m \left(\sum_{i=1}^k r_i\right)^3 = \sum_{k=1}^{m-1} \left(\sum_{i=1}^k r_i\right)^3 + \left(\sum_{i=1}^m r_i\right)^3 \stackrel{(19)}{\leq} \frac{m^3}{3}\sum_{k=1}^{m-1} r_k^3 + \left(\sum_{i=1}^m r_i\right)^3. \tag{20}$$

Applying Jensen's inequality for convex function $t \mapsto t^3$, $t \geq 0$, we have a bound for the second term:

$$\left(\sum_{i=1}^m r_i\right)^3 = m^3 \left(\frac{1}{m}\sum_{i=1}^m r_i\right)^3 \leq m^2 \sum_{i=1}^m r_i^3. \tag{21}$$

Therefore,

$$\sum_{k=1}^m \left(\sum_{i=1}^k r_i \right)^3 \stackrel{(20),(21)}{\leq} \left(\frac{m^3}{3} + m^2 \right) \sum_{k=1}^m r_k^3 \leq \frac{(m+1)^3}{3} \sum_{k=1}^m r_k^3. \quad \square$$

Using this bound, we ensure the right hand side of (18) to be positive. For that, we need to use an increased value of the regularization parameter. Thus, we prove the following guarantee.

Corollary 1 *Let $M \geq 6mL$. Then,*

$$f(\mathbf{x}_0) - f(\mathbf{x}_m) \geq \sum_{k=1}^m \max \left\{ \frac{1}{648M^2} \xi(\mathbf{x}_k)^3, \frac{1}{72\sqrt{2}M} \|\nabla f(\mathbf{x}_k)\|_*^{3/2} \right\}. \quad (22)$$

After each inner phase of cubic steps (17) with stale Hessian, we recompute the Hessian at a new snapshot point z in a new outer round. The length of each inner phase $m \geq 1$ is the key parameter of the algorithm. For convenient notation to define our algorithm, we denote by $\pi(k)$ the highest multiple of m which is less than or equal to k :

$$\pi(k) \stackrel{\text{def}}{=} k - k \bmod m, \quad k \geq 0. \quad (23)$$

The main algorithm is defined as follow:

Algorithm 1 Cubic Newton with Lazy Hessians

Input: $\mathbf{x}_0 \in \mathbb{R}^d$, $m \geq 1$, $L > 0$. Choose $M \geq 0$.

- 1: **for** $k = 0, 1, \dots$ **do**
 - 2: Set last snapshot point $\mathbf{z}_k = \mathbf{x}_{\pi(k)}$
 - 3: Compute lazy cubic step $\mathbf{x}_{k+1} = \mathbf{T}_M(\mathbf{x}_k, \mathbf{z}_k)$
-

By applying the progress inequality (22) to each phase of the method, we can establish its global complexity bound. We will use a simple fixed rule for the regularization parameter:

$$\boxed{M := 6mL} \quad (24)$$

Theorem 2 *Let the objective function be bounded from below: $\inf_{\mathbf{x}} f(\mathbf{x}) \geq f^*$. Let the regularization parameter M be fixed as in (24). Assume that the gradients for previous iterations $\{\mathbf{x}_i\}_{i=1}^k$ are higher than a desired error level $\varepsilon > 0$:*

$$\|\nabla f(\mathbf{x}_i)\|_* \geq \varepsilon. \quad (25)$$

Then, the number of iterations of Algorithm 1 to reach accuracy $\|\nabla f(\mathbf{x}_{k+1})\|_ \leq \varepsilon$ is at most*

$$k \leq \mathcal{O} \left(\frac{\sqrt{mL}(f(\mathbf{x}_0) - f^*)}{\varepsilon^{3/2}} \right), \quad (26)$$

where $\mathcal{O}(\cdot)$ hides some absolute numerical constants. The total number of Hessian updates t during these iterations is bounded as

$$t \leq \mathcal{O} \left(\frac{\sqrt{L}(f(\mathbf{x}_0) - f^*)}{\varepsilon^{3/2} \sqrt{m}} \right). \quad (27)$$

For the minimal eigenvalues of all Hessians, it holds that

$$\min_{1 \leq i \leq k} \left[-\lambda_{\min}(\mathbf{B}^{-1/2} \nabla^2 f(\mathbf{x}_i) \mathbf{B}^{-1/2}) \right]_+ \leq \left(\frac{648M^2(f(\mathbf{x}_0) - f^*)}{k} \right)^{1/3}. \quad (28)$$

Proof:

Without loss of generality we assume that k is a multiple of m : $k = tm$. Then, for the i th ($1 \leq i \leq t$) phase of the method, we have

$$f(\mathbf{x}_{m(i-1)}) - f(\mathbf{x}_{mi}) \stackrel{(22),(25)}{\geq} \frac{m}{72\sqrt{2}M} \varepsilon^{3/2} = \frac{\sqrt{m}}{144\sqrt{3}L} \varepsilon^{3/2}. \quad (29)$$

Telescoping this bound for all phases, we obtain

$$f(\mathbf{x}_0) - f^* \geq f(\mathbf{x}_0) - f(\mathbf{x}_k) \stackrel{(29)}{\geq} \frac{t\sqrt{m}}{144\sqrt{3}L} \varepsilon^{3/2}.$$

This gives the claimed bound (27) for t . Taking into account that $k = tm$, (26) follows immediately. Telescoping the bound (22) for the smallest Hessian eigenvalue $\xi(\cdot)$, we obtain

$$f(\mathbf{x}_0) - f^* \geq f(\mathbf{x}_0) - f(\mathbf{x}_k) \geq \frac{k}{648M^2} \min_{1 \leq i \leq k} \xi(\mathbf{x}_i)^3. \quad (30)$$

Hence,

$$\min_{1 \leq i \leq k} \left[-\lambda_{\min}(\mathbf{B}^{-1/2} \nabla^2 f(\mathbf{x}_i) \mathbf{B}^{-1/2}) \right]_+ \equiv \min_{1 \leq i \leq k} \xi(\mathbf{x}_i) \stackrel{(30)}{\leq} \left(\frac{648M^2(f(\mathbf{x}_0) - f^*)}{k} \right)^{1/3},$$

which completes the proof. \square

Note that the bound (27) is non-trivial, e.g. when the desired accuracy level is small, i.e.

$$\varepsilon \leq (f(\mathbf{x}_0) - f^*)^{2/3} \left(\frac{L}{m} \right)^{1/3}.$$

To choose the parameter m , we need to take into account the actual computational efforts required in all iterations of our method. We denote by **GradCost** the arithmetic complexity of one gradient step, and by **HessCost** the cost of computing the Hessian and its appropriate factorization. In general, we have

$$\text{HessCost} = d \cdot \text{GradCost}, \quad (31)$$

where d is the dimension of our problem (1).

Example 1 Let $f(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^n \phi(\langle \mathbf{a}_i, \mathbf{x} \rangle)$, where $\mathbf{a}_i \in \mathbb{R}^d$, $1 \leq i \leq n$ are given data and ϕ is a fixed univariate smooth function. In this case,

$$\nabla f(\mathbf{x}) = \mathbf{A}^\top \mathbf{s}(\mathbf{x}), \quad \text{and} \quad \nabla^2 f(\mathbf{x}) = \mathbf{A}^\top \mathbf{Q}(\mathbf{x}) \mathbf{A},$$

where $\mathbf{A} \in \mathbb{R}^{n \times d}$ is the matrix with rows $\mathbf{a}_1, \dots, \mathbf{a}_n$; $\mathbf{s}(\mathbf{x}) \in \mathbb{R}^n$ is a vector, and $\mathbf{Q}(\mathbf{x}) \in \mathbb{R}^{n \times n}$ is a diagonal matrix given by the formulas

$$[\mathbf{s}(\mathbf{x})]_{(i)} \stackrel{\text{def}}{=} \frac{1}{n} \phi'(\langle \mathbf{a}_i, \mathbf{x} \rangle), \quad [\mathbf{Q}(\mathbf{x})]_{(i,i)} \stackrel{\text{def}}{=} \frac{1}{n} \phi''(\langle \mathbf{a}_i, \mathbf{x} \rangle).$$

Because of the Hessian structure, it is convenient to use the following matrix \mathbf{B} which defines our primal norm (2):

$$\mathbf{B} := \mathbf{A}^\top \mathbf{A}. \quad (32)$$

Let us assume that the cost of computing $\phi'(\cdot)$ and $\phi''(\cdot)$ is an absolute constant $\mathcal{O}(1)$ which does not depend on the problem parameters. We denote by $\text{nz}(\mathbf{A})$ the number of nonzero elements of \mathbf{A} . Then,

$$\text{HessCost} = \mathcal{O}(d \cdot \text{nz}(\mathbf{A}) + d^3),$$

where the last cubic term comes from computing a matrix factorization (see Section 6.1), and

$$\text{GradCost} = \mathcal{O}(\text{nz}(\mathbf{A}) + d^2),$$

where the second term is from using the factorization. Thus, relation (31) is satisfied.

Example 2 Let the representation of our objective be given by the computational graph (e.g. a neural network). In this case, we can compute the Hessian-vector product $\nabla^2 f(\mathbf{x})\mathbf{h}$ for any $\mathbf{x}, \mathbf{h} \in \mathbb{R}^d$ at the same cost as its gradient $\nabla f(\mathbf{x})$ by using automatic differentiation technique (see, e.g. Chapter 7.2 in [29]). Thus, in general we can compute the Hessian as d Hessian-vector products,

$$\nabla^2 f(\mathbf{x}) = \left[\nabla^2 f(\mathbf{x})\mathbf{e}_1 \mid \dots \mid \nabla^2 f(\mathbf{x})\mathbf{e}_d \right],$$

where $\mathbf{e}_1, \dots, \mathbf{e}_d$ are the standard basis vectors. This satisfies (31).

Example 3 For any differentiable function, we can consider an approximation of its Hessian by finite differences. Namely, for a fixed parameter $\delta > 0$, we can form the matrix $\mathbf{H}_{x,\delta} \in \mathbb{R}^{d \times d}$ by the formulas

$$[\mathbf{H}_{x,\delta}]^{(i,j)} := \frac{1}{\delta} [\nabla f(\mathbf{x} + \delta \mathbf{e}_i) - \nabla f(\mathbf{x})]^{(j)}, \quad 1 \leq i, j \leq d,$$

where $\mathbf{e}_1, \dots, \mathbf{e}_d$ are the standard basis vectors, and use the following symmetrization as an approximation of the true Hessian (for more details, see [29, 5, 15]):

$$\frac{1}{2} [\mathbf{H}_{x,\delta} + \mathbf{H}_{x,\delta}^\top] \approx \nabla^2 f(\mathbf{x}).$$

Thus, forming the Hessian approximation requires $d + 1$ gradient computations, which is consistent with (31).

Hence, according to Theorem 2, the total arithmetical complexity of Algorithm 1 can be estimated as

$$\begin{aligned} \text{Arithmetical Complexity} &= k \cdot \text{GradCost} + t \cdot \text{HessCost} \\ &\stackrel{(26),(27),(31)}{\leq} \mathcal{O}\left(\left(\sqrt{m} + \frac{d}{\sqrt{m}}\right) \cdot \frac{\sqrt{L}(f(\mathbf{x}_0) - f^*)}{\varepsilon^{3/2}}\right) \cdot \text{GradCost}. \end{aligned} \quad (33)$$

Corollary 2 For $\boxed{m = 1}$ we update Hessian at each step. It corresponds to the full Cubic Newton [28], and Theorem 2 recovers its global iteration complexity:

$$k = t \leq \mathcal{O}\left(\frac{\sqrt{L}(f(\mathbf{x}_0) - f^*)}{\varepsilon^{3/2}}\right).$$

Consequently, the total arithmetical complexity of the method is bounded by

$$\mathcal{O}\left(d \cdot \frac{\sqrt{L}(f(\mathbf{x}_0) - f^*)}{\varepsilon^{3/2}}\right) \cdot \text{GradCost}. \quad (34)$$

Corollary 3 For $\boxed{m = d}$ we obtain the optimal choice for the length of one phase, which minimizes the right hand side of (33). The total arithmetical complexity of Algorithm 1 becomes

$$\mathcal{O}\left(\sqrt{d} \cdot \frac{\sqrt{L}(f(\mathbf{x}_0) - f^*)}{\varepsilon^{3/2}}\right) \cdot \text{GradCost}.$$

This improves upon the full Cubic Newton by factor \sqrt{d} .

Now, let us look at the minimal eigenvalue of all the Hessians at points generated by our algorithm.

Corollary 4 Let us fix some $\varepsilon > 0$ and perform

$$k = \frac{\sqrt{mL}(f(\mathbf{x}_0) - f^*)}{\varepsilon^{3/2}} \quad (35)$$

iterations of Algorithm 1. According to Theorem 2, we thus ensure

$$\min_{1 \leq i \leq k} \|\nabla f(\mathbf{x}_i)\|_* \leq \mathcal{O}(\varepsilon). \quad (36)$$

At the same time,

$$\min_{1 \leq i \leq k} \left[-\lambda_{\min}(\mathbf{B}^{-1/2} \nabla^2 f(\mathbf{x}_i) \mathbf{B}^{-1/2}) \right]_+ \stackrel{(28)}{\leq} \left(\frac{648M^2(f(\mathbf{x}_0) - f^*)}{k} \right)^{1/3} \stackrel{(35)}{=} 2^{5/3} 3^2 \sqrt{mL\varepsilon}.$$

Therefore, we see that negative eigenvalues of the Hessians cannot be big. For $\mathcal{O}(\varepsilon)$ level of gradient norm (36) we guarantee $\mathcal{O}(\sqrt{mL\varepsilon})$ level for the smallest eigenvalue (28).

4 Minimizing Convex Functions

In this section, we assume that the objective in our problem (1) is *convex*. Thus, all the Hessians are positive semidefinite:

$$\nabla^2 f(\mathbf{x}) \succeq 0, \quad \forall \mathbf{x} \in \mathbb{R}^d. \quad (37)$$

Then, we can apply the *gradient regularization* technique [25, 10], which allows to use *square* of the Euclidean norm as a regularizer for our model. Each step of the method becomes easier to compute employing just one matrix inversion. Thus, we come to the following scheme:

Algorithm 2 Regularized Newton with Lazy Hessians

Input: $\mathbf{x}_0 \in \mathbb{R}^d$, $m \geq 1$, $L > 0$. Choose $M \geq 0$.

- 1: **for** $k = 0, 1, \dots$ **do**
 - 2: Set last snapshot point $\mathbf{z}_k = \mathbf{x}_{\pi(k)}$
 - 3: Set regularization parameter $\lambda_k = \sqrt{M \|\nabla f(\mathbf{x}_k)\|_*}$
 - 4: Compute lazy Newton step: $\mathbf{x}_{k+1} = \mathbf{x}_k - (\nabla^2 f(\mathbf{z}_k) + \lambda_k \mathbf{B})^{-1} \nabla f(\mathbf{x}_k)$
-

Regularization parameter M should have the same order and a physical meaning as that one in the Cubic Newton.

Let us express the progress given by one step of this new method. First, we deduce several useful inequalities. For a fixed $\mathbf{z} \in \mathbb{R}^d$, let us denote one regularized lazy Newton step from point $\mathbf{x} \in \mathbb{R}^d$ by

$$\mathbf{x}^+ \stackrel{\text{def}}{=} \mathbf{x} - (\nabla^2 f(\mathbf{z}) + \lambda \mathbf{B})^{-1} \nabla f(\mathbf{x}), \quad \lambda > 0. \quad (38)$$

We choose parameter λ as in Algorithm 2:

$$\boxed{\lambda := \sqrt{M \|\nabla f(\mathbf{x})\|_*}} \quad (39)$$

where $M > 0$ is fixed. The Newton step with gradient regularization can be seen as an approximation of the Cubic Newton step (9), which utilizes convexity of the objective. Indeed, we have by convexity:

$$\|\mathbf{x}^+ - \mathbf{x}\| = \|(\nabla^2 f(\mathbf{z}) + \lambda \mathbf{B})^{-1} \nabla f(\mathbf{x})\| \stackrel{(37)}{\leq} \frac{1}{\lambda} \|\nabla f(\mathbf{x})\|_*, \quad (40)$$

and so we can ensure our main bound:

$$M \|\mathbf{x}^+ - \mathbf{x}\| \stackrel{(40)}{\leq} \frac{M}{\lambda} \|\nabla f(\mathbf{x})\|_* \stackrel{(39)}{=} \lambda, \quad (41)$$

which justifies the actual choice of λ .

Now, we need to have analogues of Lemma 1 and Lemma 2 that we established for the cubically regularized lazy Newton step. We can prove the following.

Lemma 5 *For any $M > 0$, it holds*

$$\|\nabla f(\mathbf{x}^+)\|_* \leq \left(\frac{L}{2} \|\mathbf{x}^+ - \mathbf{x}\| + \lambda + L \|\mathbf{z} - \mathbf{x}\|\right) \cdot \|\mathbf{x}^+ - \mathbf{x}\|. \quad (42)$$

Consequently, for $M \geq 3L$, we get

$$\|\nabla f(\mathbf{x}^+)\|_* \stackrel{(42),(41)}{\leq} \left(\frac{7}{6} \lambda + L \|\mathbf{z} - \mathbf{x}\|\right) \cdot \|\mathbf{x}^+ - \mathbf{x}\|. \quad (43)$$

Proof:

Indeed, we have

$$\begin{aligned} \frac{L}{2}\|\mathbf{x}^+ - \mathbf{x}\|^2 &\stackrel{(4)}{\geq} \|\nabla f(\mathbf{x}^+) - \nabla f(\mathbf{x}) - \nabla^2 f(\mathbf{x})(\mathbf{x}^+ - \mathbf{x})\|_* \\ &\geq \|\nabla f(\mathbf{x}^+)\|_* - \|\nabla f(\mathbf{x}) + \nabla^2 f(\mathbf{z})(\mathbf{x}^+ - \mathbf{x})\|_* - L\|\mathbf{z} - \mathbf{x}\| \cdot \|\mathbf{x}^+ - \mathbf{x}\|, \end{aligned}$$

where we used triangle inequality and Lipschitz continuity of the Hessian in the last bound. It remains to note that $\|\nabla f(\mathbf{x}) + \nabla^2 f(\mathbf{z})(\mathbf{x}^+ - \mathbf{x})\|_* \stackrel{(38)}{=} \lambda\|\mathbf{x}^+ - \mathbf{x}\|$. \square

Lemma 6 For $M \geq 3L$, it holds

$$f(\mathbf{x}) - f(\mathbf{x}^+) \geq \frac{\lambda}{4}\|\mathbf{x}^+ - \mathbf{x}\|^2 - \frac{8L^3}{27M^2}\|\mathbf{z} - \mathbf{x}\|^3. \quad (44)$$

Proof:

Combining the method step with Lipschitz continuity of the Hessian, we obtain

$$\begin{aligned} f(\mathbf{x}^+) &\stackrel{(5),(41)}{\leq} f(\mathbf{x}) + \langle \nabla f(\mathbf{x}), \mathbf{x}^+ - \mathbf{x} \rangle + \frac{1}{2}\langle \nabla^2 f(\mathbf{x})(\mathbf{x}^+ - \mathbf{x}), \mathbf{x}^+ - \mathbf{x} \rangle + \frac{M}{18}\|\mathbf{x}^+ - \mathbf{x}\|^3 \\ &\stackrel{(38)}{=} f(\mathbf{x}) - \lambda\|\mathbf{x}^+ - \mathbf{x}\|^2 - \langle \nabla^2 f(\mathbf{z})(\mathbf{x}^+ - \mathbf{x}), \mathbf{x}^+ - \mathbf{x} \rangle + \frac{M}{18}\|\mathbf{x}^+ - \mathbf{x}\|^3 \\ &\quad + \frac{1}{2}\langle \nabla^2 f(\mathbf{x})(\mathbf{x}^+ - \mathbf{x}), \mathbf{x}^+ - \mathbf{x} \rangle \\ &\stackrel{(3),(37),(41)}{\leq} f(\mathbf{x}) - \lambda\|\mathbf{x}^+ - \mathbf{x}\|^2 + \frac{\lambda}{2}\|\mathbf{x}^+ - \mathbf{x}\|^2 + \frac{L}{2}\|\mathbf{z} - \mathbf{x}\| \cdot \|\mathbf{x}^+ - \mathbf{x}\|^2. \end{aligned}$$

Applying Young's inequality for the last term,

$$\begin{aligned} \frac{L}{2}\|\mathbf{z} - \mathbf{x}\| \cdot \|\mathbf{x}^+ - \mathbf{x}\|^2 &= \left(\frac{(3M)^{2/3}}{4}\|\mathbf{x}^+ - \mathbf{x}\|^2\right) \cdot \left(\frac{2L}{(3M)^{2/3}}\|\mathbf{z} - \mathbf{x}\|\right) \\ &\leq \frac{M}{4}\|\mathbf{x}^+ - \mathbf{x}\|^3 + \frac{8L^3}{27M^2}\|\mathbf{z} - \mathbf{x}\|^3 \stackrel{(41)}{\leq} \frac{\lambda}{4}\|\mathbf{x}^+ - \mathbf{x}\|^2 + \frac{8L^3}{27M^2}\|\mathbf{z} - \mathbf{x}\|^3. \end{aligned}$$

we obtain (44). \square

Combining these two lemmas, we get the functional progress of one step in terms of the new gradient norm.

Theorem 3 Let $M \geq 3L$. Then, for one lazy Newton step (38) with gradient regularization (39), it holds:

$$f(\mathbf{x}) - f(\mathbf{x}^+) \geq \frac{9}{244\lambda}\|\nabla f(\mathbf{x}^+)\|_*^2 + \frac{M}{8}\|\mathbf{x}^+ - \mathbf{x}\|^3 - \frac{3L^3}{M^2}\|\mathbf{z} - \mathbf{x}\|^3. \quad (45)$$

Proof:

Using convexity of the function $t \mapsto t^2$ for $t \geq 0$ and Young's inequality, we get

$$\begin{aligned}
\|\nabla f(\mathbf{x}^+)\|_*^2 &\stackrel{(43)}{\leq} \left(\frac{7}{6}\lambda\|\mathbf{x}^+ - \mathbf{x}\| + L\|\mathbf{z} - \mathbf{x}\| \cdot \|\mathbf{x}^+ - \mathbf{x}\| \right)^2 \\
&\leq \frac{49}{18}\lambda^2\|\mathbf{x}^+ - \mathbf{x}\|^2 + 2L^2\|\mathbf{z} - \mathbf{x}\|^2 \cdot \|\mathbf{x}^+ - \mathbf{x}\|^2 \\
&\leq \frac{49}{18}\lambda^2\|\mathbf{x}^+ - \mathbf{x}\|^2 + \frac{4\lambda L^3}{3M^2}\|\mathbf{z} - \mathbf{x}\|^3 + \frac{2M^4\|\mathbf{x}^+ - \mathbf{x}\|^6}{3\lambda^2} \\
&\stackrel{(41)}{\leq} \frac{61}{18}\lambda^2\|\mathbf{x}^+ - \mathbf{x}\|^2 + \frac{4\lambda L^3}{3M^2}\|\mathbf{z} - \mathbf{x}\|^3.
\end{aligned} \tag{46}$$

Thus,

$$\begin{aligned}
\frac{\lambda}{4}\|\mathbf{x}^+ - \mathbf{x}\|^2 &\stackrel{(41)}{\geq} \frac{\lambda}{8}\|\mathbf{x}^+ - \mathbf{x}\|^2 + \frac{M}{8}\|\mathbf{x}^+ - \mathbf{x}\|^3 \\
&\stackrel{(46)}{\geq} \frac{9}{244\lambda}\|\nabla f(\mathbf{x}^+)\|_*^2 + \frac{M}{8}\|\mathbf{x}^+ - \mathbf{x}\|^3 - \frac{3L^3}{61M^2}\|\mathbf{z} - \mathbf{x}\|^3.
\end{aligned}$$

Substituting this bound into (44) completes the proof. \square

Let us consider one phase of the algorithm. Telescoping bound (45) for the first m iterations and using triangle inequality,

$$\|\mathbf{x}_0 - \mathbf{x}_k\| \leq \sum_{i=1}^k \|\mathbf{x}_i - \mathbf{x}_{i-1}\| =: \sum_{i=1}^k r_i,$$

we obtain

Corollary 5 *Let $M \geq 3mL$. Then*

$$\begin{aligned}
f(\mathbf{x}_0) - f(\mathbf{x}_m) &\geq \frac{9}{244\sqrt{M}} \sum_{k=1}^m \frac{\|\nabla f(\mathbf{x}_k)\|_*^2}{\|\nabla f(\mathbf{x}_{k-1})\|_*^{1/2}} + \frac{M}{8} \sum_{k=1}^m r_k^3 - \frac{3L^3}{M^2} \sum_{k=1}^{m-1} \left(\sum_{i=1}^k r_i \right)^3 \\
&\stackrel{(19)}{\geq} \frac{9}{244\sqrt{M}} \sum_{k=1}^m \frac{\|\nabla f(\mathbf{x}_k)\|_*^2}{\|\nabla f(\mathbf{x}_{k-1})\|_*^{1/2}}.
\end{aligned} \tag{47}$$

We are ready to prove the global complexity bound for Algorithm 2. Let us consider the constant choice of the regularization parameter:

$$\boxed{M := 3mL} \tag{48}$$

Theorem 4 *Let the objective function be convex and bounded from below: $\inf_{\mathbf{x}} f(\mathbf{x}) \geq f^*$. Let the regularization parameter M be fixed as in (48). Assume that the gradients for previous iterations $\{\mathbf{x}_i\}_{i=1}^k$ are higher than a desired error level $\varepsilon > 0$:*

$$\|\nabla f(\mathbf{x}_i)\|_* \geq \varepsilon. \tag{49}$$

Then, the number of iterations of Algorithm 2 to reach accuracy $\|\nabla f(\mathbf{x}_{k+1})\|_* \leq \varepsilon$ is at most

$$k \leq \mathcal{O}\left(\frac{\sqrt{mL}(f(\mathbf{x}_0)-f^*)}{\varepsilon^{3/2}} + \ln \frac{\|\nabla f(\mathbf{x}_0)\|_*}{\varepsilon}\right). \quad (50)$$

The total number of Hessian updates t during these iterations is bounded as

$$t \leq \mathcal{O}\left(\frac{\sqrt{L}(f(\mathbf{x}_0)-f^*)}{\varepsilon^{3/2}\sqrt{m}} + \frac{1}{m} \ln \frac{\|\nabla f(\mathbf{x}_0)\|_*}{\varepsilon}\right). \quad (51)$$

Proof:

Without loss of generality we assume that k is a multiple of m : $k = tm$. Then, for i th ($1 \leq i \leq t$) phase of the method, we have

$$\begin{aligned} f(\mathbf{x}_{m(i-1)}) - f(\mathbf{x}_{mi}) &\stackrel{(47)}{\geq} \frac{9}{244\sqrt{M}} \sum_{m(i-1)+1 \leq j \leq mi} \frac{\|\nabla f(\mathbf{x}_j)\|_*^2}{\|\nabla f(\mathbf{x}_{j-1})\|_*^{1/2}} \\ &\stackrel{(49)}{\geq} \frac{9\varepsilon^{3/2}}{244\sqrt{M}} \sum_{m(i-1)+1 \leq j \leq mi} \left(\frac{\|\nabla f(\mathbf{x}_j)\|_*}{\|\nabla f(\mathbf{x}_{j-1})\|_*}\right)^{\frac{1}{2}}. \end{aligned} \quad (52)$$

Telescoping this bound for all phases and using inequality between arithmetic and geometric means, we get

$$\begin{aligned} f(\mathbf{x}_0) - f^* &\geq f(\mathbf{x}_0) - f(\mathbf{x}_k) \stackrel{(52)}{\geq} \frac{9\varepsilon^{3/2}}{244\sqrt{M}} \sum_{j=1}^k \left(\frac{\|\nabla f(\mathbf{x}_j)\|_*}{\|\nabla f(\mathbf{x}_{j-1})\|_*}\right)^{\frac{1}{2}} \\ &\geq \frac{9\varepsilon^{3/2}k}{244\sqrt{M}} \left(\prod_{j=1}^k \frac{\|\nabla f(\mathbf{x}_j)\|_*}{\|\nabla f(\mathbf{x}_{j-1})\|_*}\right)^{\frac{1}{2k}} = \frac{9\varepsilon^{3/2}k}{244\sqrt{M}} \left(\frac{\|\nabla f(\mathbf{x}_k)\|_*}{\|\nabla f(\mathbf{x}_0)\|_*}\right)^{\frac{1}{2k}} \\ &\stackrel{(49)}{\geq} \frac{9\varepsilon^{3/2}k}{244\sqrt{M}} \left(\frac{\varepsilon}{\|\nabla f(\mathbf{x}_0)\|_*}\right)^{\frac{1}{2k}} = \frac{9\varepsilon^{3/2}k}{244\sqrt{M}} \exp\left(\frac{1}{2k} \ln \frac{\varepsilon}{\|\nabla f(\mathbf{x}_0)\|_*}\right) \\ &\geq \frac{9\varepsilon^{3/2}k}{244\sqrt{M}} \left(1 - \frac{1}{2k} \ln \frac{\|\nabla f(\mathbf{x}_0)\|_*}{\varepsilon}\right). \end{aligned}$$

Rearranging the terms gives (50). Inequality (51) follows immediately from $t = k/m$. \square

We see that the global complexity bounds for Algorithm 2 are the same (up to an additive logarithmic term) as those ones for Algorithm 1. However, each iteration of Algorithm 2 is much easier to implement since it involves just one standard matrix inversion.

5 Local Superlinear Convergence

In this section, we discuss local convergence of our second-order schemes with lazy Hessian updates. We show the fast linear rate with a constant factor during each phase and superlinear convergence taking into account the Hessian updates.

Let us assume that our objective is strongly convex with some parameter $\mu > 0$.

$$\nabla^2 f(\mathbf{x}) \succeq \mu \mathbf{B}, \quad \forall \mathbf{x} \in \mathbb{R}^d. \quad (53)$$

For simplicity, we require (53) for all points from our space, while it is possible to analyze Newton steps assuming strong convexity only in a neighborhood of the solution [26].

By strong convexity (53), we have a bound for the distance to the optimum $\mathbf{x}^* \stackrel{\text{def}}{=} \operatorname{argmin}_{\mathbf{x}} f(\mathbf{x})$ using the gradient norm (see, e.g. [26]), for all $\mathbf{x} \in \mathbb{R}^d$:

$$\|\mathbf{x} - \mathbf{x}^*\| \leq \frac{1}{\mu} \|\nabla f(\mathbf{x})\|_*. \quad (54)$$

Now, let us look at one lazy Cubic Newton step $\mathbf{T} = \mathbf{T}_M(\mathbf{x}, \mathbf{z})$ with some $M \geq 0$, for which we have:

$$r \equiv \|\mathbf{T} - \mathbf{x}\| \stackrel{(9)}{=} \|(\nabla^2 f(\mathbf{z}) + \frac{Mr}{2}\mathbf{B})^{-1} \nabla f(\mathbf{x})\|_* \stackrel{(53)}{\leq} \frac{1}{\mu} \|\nabla f(\mathbf{x})\|_*. \quad (55)$$

Note that $M = 0$ corresponds to the classical pure Newton step with a lazy Hessian.

Then, applying Lemma 1 with triangle inequality, we get

$$\begin{aligned} \|\nabla f(\mathbf{T})\|_* &\stackrel{(10)}{\leq} \frac{M+L}{2} r^2 + Lr \|\mathbf{z} - \mathbf{x}\| \leq \frac{M+L}{2} r^2 + Lr \|\mathbf{x} - \mathbf{x}^*\| + Lr \|\mathbf{z} - \mathbf{x}^*\| \\ &\stackrel{(54),(55)}{\leq} \frac{M+3L}{2\mu^2} \|\nabla f(\mathbf{x})\|_*^2 + \frac{L}{\mu^2} \|\nabla f(\mathbf{x})\|_* \cdot \|\nabla f(\mathbf{z})\|_*. \end{aligned} \quad (56)$$

This inequality leads to a superlinear convergence of the method.

Theorem 5 *Let $M \geq 0$. Assume that initial gradient is small enough:*

$$\|\nabla f(\mathbf{x}_0)\|_* \leq \frac{\mu^2}{2(M+3L)}. \quad (57)$$

Then, for the iterations of Algorithm 1, we have a superlinear convergence for the gradient norms:

$$\|\nabla f(\mathbf{x}_k)\|_* \leq \frac{\mu^2}{M+3L} \left(\frac{1}{2}\right)^{(1+m)\pi(k)(1+k \bmod m)}, \quad k \geq 0, \quad (58)$$

where $\pi(k)$ is defined by (23).

Proof:

According to (56), for one iteration of Algorithm 1, it holds

$$\|\nabla f(\mathbf{x}_{k+1})\|_* \leq \frac{M+3L}{2\mu^2} \|\nabla f(\mathbf{x}_k)\|_*^2 + \frac{L}{\mu^2} \|\nabla f(\mathbf{x}_k)\|_* \cdot \|\nabla f(\mathbf{x}_{\pi(k)})\|_*.$$

Let us multiply this inequality by $c := \frac{M+3L}{2\mu^2}$ and denote $s_k \stackrel{\text{def}}{=} c \|\nabla f(\mathbf{x}_k)\|_*$. This yields the following recursion

$$s_{k+1} \leq s_k^2 + s_k s_{\pi(k)}, \quad (59)$$

and by initial condition (57), we have $s_0 \leq \frac{1}{4}$.

Required inequality from our claim is

$$s_{tm+i} \leq \left(\frac{1}{2}\right)^{(1+i)(1+m)^t+1}, \quad (60)$$

for any $t \geq 0$ (phases of the method) and $0 \leq i \leq m$. We prove it by induction. Indeed,

$$\begin{aligned} s_{tm+i+1} &\stackrel{(59)}{\leq} s_{tm+i}^2 + s_{tm+i}s_{tm} \stackrel{(60)}{\leq} \left(\frac{1}{2}\right)^{2(1+i)(1+m)^t+2} + \left(\frac{1}{2}\right)^{(1+i)(1+m)^t+1+(1+m)^t+1} \\ &\leq \left(\frac{1}{2}\right)^{(2+i)(1+m)^t+2} + \left(\frac{1}{2}\right)^{(2+i)(1+m)^t+2} = \left(\frac{1}{2}\right)^{(2+i)(1+m)^t+1}, \end{aligned}$$

which is (60) for the next index. \square

Corollary 6 *Combining both Theorem 2 and 5, we conclude that for minimizing a strongly convex function by Algorithm 1 with regularization parameter M given by (24) and starting from an arbitrary \mathbf{x}_0 , we need to do*

$$k \leq \mathcal{O}\left(\frac{m^2 L^2 (f(\mathbf{x}_0) - f^*)}{\mu^3} + \frac{1}{\ln(1+m)} \ln \ln \frac{\mu^2}{mL\varepsilon}\right)$$

lazy steps to achieve $\|\nabla f(\mathbf{x}_k)\|_* \leq \varepsilon$.

Let us also analyze the local behaviour of the lazy Newton steps with gradient regularization, that we perform in Algorithm 2. Note that by strong convexity (53) we have the following bound for the length of one step, which is

$$\|\mathbf{x}^+ - \mathbf{x}\| \stackrel{(38)}{=} \|(\nabla^2 f(\mathbf{z}) + \lambda \mathbf{B})^{-1} \nabla f(\mathbf{x})\| \stackrel{(53)}{\leq} \frac{1}{\mu} \|\nabla f(\mathbf{x})\|_*. \quad (61)$$

Therefore, we can estimate the norm of the gradient at new point taking into account the actual choice of the regularization parameter (39). Our reasoning works for any $M \geq 0$.

Applying Lemma 5 and triangle inequality, we get

$$\begin{aligned} \|\nabla f(\mathbf{x}^+)\|_* &\stackrel{(42)}{\leq} \left(\frac{L}{2}\|\mathbf{x}^+ - \mathbf{x}\| + \lambda + L\|\mathbf{z} - \mathbf{x}\|\right) \cdot \|\mathbf{x}^+ - \mathbf{x}\| \\ &\stackrel{(61)}{\leq} \left(\frac{L}{2\mu}\|\nabla f(\mathbf{x})\|_* + \lambda + L\|\mathbf{z} - \mathbf{x}\|\right) \cdot \frac{1}{\mu}\|\nabla f(\mathbf{x})\|_* \\ &\leq \left(\frac{L}{2\mu}\|\nabla f(\mathbf{x})\|_* + \lambda + L\|\mathbf{x} - \mathbf{x}^*\| + L\|\mathbf{z} - \mathbf{x}^*\|\right) \cdot \frac{1}{\mu}\|\nabla f(\mathbf{x})\|_* \\ &\stackrel{(54),(39)}{\leq} \frac{3L}{2\mu^2}\|\nabla f(\mathbf{x})\|_*^2 + \frac{\sqrt{M}}{\mu}\|\nabla f(\mathbf{x})\|_*^{3/2} + \frac{L}{\mu^2}\|\nabla f(\mathbf{x})\|_* \cdot \|\nabla f(\mathbf{z})\|_* \end{aligned} \quad (62)$$

We see that the power of the gradient norm for the second term in the right hand side is $3/2$, which is slightly worse than 2, that is in the cubic regularization (compare with (56)). However, it is still a local *superlinear* convergence, which is extremely fast from the practical perspective.

Theorem 6 *Let $M \geq 0$. Assume that initial gradient is small enough:*

$$\|\nabla f(\mathbf{x}_0)\|_* \leq \frac{\mu^2}{2^4(3L+4M)}. \quad (63)$$

Then, for the iterations of Algorithm 2, we have a superlinear convergence for the gradient norms:

$$\|\nabla f(\mathbf{x}_k)\|_* \leq \frac{\mu^2}{2^2(3L+4M)} \left(\frac{1}{2}\right)^{2(1+m/2)\pi(k)(1+(k \bmod m)/2)}, \quad k \geq 0, \quad (64)$$

where $\pi(k)$ is defined by (23).

Proof:

According to (62), for one iteration of Algorithm 2, it holds

$$\|\nabla f(\mathbf{x}_{k+1})\|_* \leq \frac{3L}{\mu^2} \|\nabla f(\mathbf{x}_k)\|_*^2 + \frac{\sqrt{M}}{\mu} \|\nabla f(\mathbf{x}_k)\|_*^{3/2} + \frac{L}{\mu^2} \|\nabla f(\mathbf{x}_k)\|_* \cdot \|\nabla f(\mathbf{x}_{\pi(k)})\|_*.$$

Let us multiply this inequality by $c := \frac{4M+3L}{\mu^2}$ and denote $s_k \stackrel{\text{def}}{=} c \|\nabla f(\mathbf{x}_k)\|_*$. This yields the following recursion (compare with (59)):

$$s_{k+1} \leq \frac{1}{2}(s_k^2 + s_k^{3/2}) + s_k s_{\pi(k)}, \quad (65)$$

and by initial condition (63), we have $s_0 \leq \frac{1}{2^4}$.

Let us prove by induction that

$$s_{tm} \leq \left(\frac{1}{2}\right)^{2(1+(1+m/2)^t)}, \quad (66)$$

for any $t \geq 0$ (phases of the method), and

$$s_{tm+i} \leq \left(\frac{1}{2}\right)^{i(1+m/2)^t} s_{tm}, \quad (67)$$

for $0 \leq i \leq m$. Then (64) clearly follows.

Note that from (66) we have

$$\frac{1}{2}s_{tm}^{1/2} + \frac{3}{2}s_{tm} \leq 2s_{tm}^{1/2} \stackrel{(66)}{\leq} \left(\frac{1}{2}\right)^{(1+m/2)^t}. \quad (68)$$

Then, assuming that (66) and (68) holds for the current iterate, we get

$$\begin{aligned} s_{tm+i+1} &\stackrel{(65)}{\leq} \frac{1}{2}(s_{tm+i}^2 + s_{tm+i}^{3/2}) + s_{tm+i}s_{tm} \stackrel{(67),(66)}{\leq} \left(\frac{1}{2}\right)^{i(1+m/2)^t} s_{tm} \left(\frac{1}{2}s_{tm}^{1/2} + \frac{3}{2}s_{tm}\right) \\ &\stackrel{(68)}{\leq} \left(\frac{1}{2}\right)^{(i+1)(1+m/2)^t} s_{tm}, \end{aligned}$$

which is (67) for the next index.

It remains to observe that substituting $i := m$ into (67) we obtain (66) for the next phase. Indeed,

$$\begin{aligned} s_{(t+1)m} &= s_{tm+m} \stackrel{(67)}{\leq} \left(\frac{1}{2}\right)^{m(1+m/2)^t} s_{tm} \\ &\stackrel{(66)}{\leq} \left(\frac{1}{2}\right)^{m(1+m/2)^t + 2 + 2(1+m/2)^t} = \left(\frac{1}{2}\right)^{2+2(1+m/2)^{t+1}}. \quad \square \end{aligned}$$

Corollary 7 *Combining both Theorem 4 and 6, we conclude that for minimizing a strongly convex function by Algorithm 2 with regularization parameter M given by (48) and starting from an arbitrary \mathbf{x}_0 , we need to do*

$$k \leq \mathcal{O}\left(\frac{m^2 L^2 (f(\mathbf{x}_0) - f^*)}{\mu^3} + \ln \frac{mL \|\nabla f(\mathbf{x}_0)\|_*}{\mu^2} + \frac{1}{\ln(1+m/2)} \ln \ln \frac{\mu^2}{mL\varepsilon}\right)$$

lazy steps to achieve $\|\nabla f(\mathbf{x}_k)\|_* \leq \varepsilon$.

6 Practical Implementation

Let us discuss some important aspects of effective implementation of our methods with the lazy Hessian updates.

6.1 Use of Matrix Factorization

Computing the Hessian once per m iterations, we want to be able to solve efficiently the corresponding subproblem with this Hessian in each iteration of the method.

For solving the subproblem in Cubic Newton (6), we need to find parameter $r > 0$ that is, in a nondegenerate case, the root of the following nonlinear equation (see also Section 5 in [28]),

$$\varphi(r) \stackrel{\text{def}}{=} \|\mathbf{s}(r)\| - r = 0, \quad (69)$$

where

$$\mathbf{s}(r) \stackrel{\text{def}}{=} (\nabla^2 f(\mathbf{z}) + \frac{Mr}{2} \mathbf{B})^{-1} \nabla f(\mathbf{x}), \quad \text{for } \nabla^2 f(\mathbf{z}) + \frac{Mr}{2} \mathbf{B} \succ 0.$$

Note that

$$\varphi'(r) = -\frac{M}{2\|\mathbf{s}(r)\|} \langle \mathbf{B}\mathbf{s}(r), (\nabla^2 f(\mathbf{z}) + \frac{Mr}{2} \mathbf{B})^{-1} \mathbf{B}\mathbf{s}(r) \rangle - 1 < 0,$$

and thus function $\varphi(r)$ is monotonically decreasing for $r > \frac{2}{M} \lambda_{\min}(-\mathbf{B}^{1/2} \nabla^2 f(\mathbf{z}) \mathbf{B}^{-1/2})$.

Applying to (69) the standard bisection method, or the univariate Newton's method with steps of the form $r^+ = r - \frac{\varphi(r)}{\varphi'(r)}$, the main operation that we need to perform is solving a linear equation:

$$(\nabla^2 f(\mathbf{z}) + \tau \mathbf{B}) \mathbf{h} = -\nabla f(\mathbf{x}) \quad \Leftrightarrow \quad \mathbf{h} = -(\nabla^2 f(\mathbf{z}) + \tau \mathbf{B})^{-1} \nabla f(\mathbf{x}), \quad (70)$$

for different values of $\tau > 0$. This is the same type of operation that we need to do *once* per each step in Algorithm 2.

Now, let us assume that we have the following factorization of the Hessian:

$$\nabla^2 f(\mathbf{z}) = \mathbf{U} \mathbf{\Lambda} \mathbf{U}^\top, \quad \text{where } \mathbf{U} \mathbf{U}^\top = \mathbf{B}, \quad (71)$$

and $\mathbf{\Lambda} \in \mathbb{R}^{d \times d}$ is a *diagonal* or *tridiagonal* matrix. Thus, \mathbf{U} is a set of vectors orthogonal with respect to \mathbf{B} . In case $\mathbf{B} = \mathbf{I}$ (identity matrix) and $\mathbf{\Lambda}$ being diagonal, (71) is called *Eigendecomposition* and implemented in most of the standard Linear Algebra packages.

In general, factorization (71) can be computed in $\mathcal{O}(d^3)$ arithmetic operations. Namely, we can apply a standard orthogonalizing decomposition for the matrix $\mathbf{B}^{-1/2}\nabla^2 f(\mathbf{z})\mathbf{B}^{-1/2}$, thus obtaining

$$\mathbf{B}^{-1/2}\nabla^2 f(\mathbf{z})\mathbf{B}^{-1/2} = \mathbf{V}\mathbf{\Lambda}\mathbf{V}^\top, \quad \text{with } \mathbf{V}\mathbf{V}^\top = \mathbf{I},$$

and then set $\mathbf{U} := \mathbf{B}^{1/2}\mathbf{V}$, which gives (71).

Then, solution to (70) can be expressed as

$$\mathbf{h} = -\mathbf{U}^{-\top}(\mathbf{\Lambda} + \tau\mathbf{I})^{-1}\mathbf{U}^{-1}\nabla f(\mathbf{x}),$$

and it is computable in $\mathcal{O}(d^2)$ arithmetic operations for any given $\tau > 0$ and $\nabla f(\mathbf{x})$.

The use of tridiagonal decomposition can be more efficient in practice. Indeed, inversion of the matrix $\mathbf{\Lambda} + \tau\mathbf{I}$ would still cost $\mathcal{O}(d)$ operations for tridiagonal $\mathbf{\Lambda}$, while it requires less computational resources and less floating point precision to compute such decomposition.

6.2 Adaptive Search

The global convergence of the lazy Hessian updates was achieved by picking up an appropriate value of the regularization parameter $M > 0$. In our algorithms, M should be chosen proportional to mL , where $m \geq 1$ is the length of one phase and $L > 0$ is the Lipschitz constant of the Hessian.

One drawback of this choice is that we actually need to know the Lipschitz constant, which is not always the case in practice. Moreover, with the constant regularization, the methods become somewhat conservative, preventing a possibility of big steps. At the same time, from the local perspective the best quadratic approximation of the objective is the pure second-order Taylor polynomial. So, being in a neighbourhood of the solution, the best strategy is to tend M to zero which gives the fastest local superlinear rates (see Section 5).

The use of adaptive search in second-order schemes has been studied for several years [28, 3, 4, 16, 17, 11]. It is well known that such schemes have a very good practical performance, while the adaptive search makes the methods to be also *universal* [16, 11] or even *super-universal* [8], working properly on a wide range of problem classes with the best global complexity guarantees.

In this section, we show how to equip our second-order algorithms with the adaptive search, such that we do not need to know the actual value of the Lipschitz constant.

The key to our analysis is inequality (22) on m consecutive steps of the method. Note that it holds for *any* value of M that is sufficiently big. Hence, there is no necessity to fix regularization parameter. We are going to change its value adaptively while checking the condition on the functional progress after m steps of the method. Thus, we come to the following adaptive algorithm:

Note that the value M_0 is just an *initial guess* for the regularization constant, which can be further both increased or decreased dynamically. This process is well defined. Indeed, due to (22), the stopping condition is satisfied as long as $M_t \geq 2^8 3^5 mL$. Hence, we ensure that all regularization coefficients always satisfy the following bound:

$$M_t \leq \max\{2M_0, 2^9 3^5 mL\}. \tag{72}$$

Algorithm 3 Adaptive Cubic Newton with Lazy Hessians

Input: $\mathbf{x}_0 \in \mathbb{R}^d$, $m \geq 1$. Fix some $M_0 > 0$.

- 1: **for** $t = 0, 1, \dots$ **do**
 - 2: Compute snapshot Hessian $\nabla^2 f(\mathbf{x}_{tm})$
 - 3: **do**
 - 4: Update $M_t = 2 \cdot M_t$
 - 5: **for** $i = 1, \dots, m$ **do**
 - 6: Compute lazy cubic step $\mathbf{x}_{tm+i} = T_{M_t}(\mathbf{x}_{tm+i-1}, \mathbf{x}_{tm})$
 - 7: **until** $f(\mathbf{x}_{tm}) - f(\mathbf{x}_{tm+m}) \geq \frac{1}{\sqrt{M_t}} \sum_{i=1}^m \|\nabla f(\mathbf{x}_{tm+i})\|_*^{3/2}$
 - 8: Set $M_{t+1} = \frac{1}{4} \cdot M_t$
-

Substituting this bound into stopping condition, and telescoping it for the first $t \geq 0$ phases, we get

$$f(\mathbf{x}_0) - f^* \geq f(\mathbf{x}_0) - f(\mathbf{x}_{tm}) \geq \frac{1}{\sqrt{\max\{2M_0, 2^9 3^5 mL\}}} \sum_{i=1}^{tm} \|\nabla f(\mathbf{x}_i)\|_*^{3/2}. \quad (73)$$

Thus, we obtain the following global complexity guarantee.

Theorem 7 *Let the objective function be bounded from below: $\inf_{\mathbf{x}} f(\mathbf{x}) \geq f^*$ and let the gradients for previous iterates $\{\mathbf{x}_i\}_{i=1}^{tm}$ during t phases of Algorithm 3 are higher than a desired error level $\varepsilon > 0$:*

$$\|\nabla f(\mathbf{x}_i)\|_* \geq \varepsilon. \quad (74)$$

Then, the number of phases of Algorithm 3 to reach accuracy $\|\nabla f(\mathbf{x}_{tm+1})\|_ \leq \varepsilon$ is at most*

$$t \leq \mathcal{O}\left(\sqrt{\max\left\{\frac{M_0}{m}, L\right\}} \cdot \frac{f(\mathbf{x}_0) - f^*}{\varepsilon^{3/2} \sqrt{m}}\right). \quad (75)$$

The total number N of gradient calls during these phases is bounded as

$$N \leq (2t + \log_2 \frac{M_t}{M_0})m \stackrel{(72)}{\leq} 2tm + \max\{1, \log_2 \frac{2^9 3^5 mL}{M_0}\}m. \quad (76)$$

Proof:

Bound (75) follows directly from (73) and (74).

To prove (76), we denote by n_i the number of times the **do ... until** loop is performed at phase $0 \leq i \leq t-1$. By our updates, it holds

$$2^{n_i-2} = \frac{M_{i+1}}{M_i}. \quad (77)$$

It remains to note that

$$N = \left(\sum_{i=0}^{t-1} n_i\right)m \stackrel{(77)}{=} \left(2t + \sum_{i=0}^{t-1} \log_2 \frac{M_{i+1}}{M_i}\right)m = \left(2t + \log_2 \frac{M_t}{M_0}\right)m. \quad \square$$

Remark 1 *According to Theorem 7, the global complexity of the adaptive algorithm is the same as that one given by Theorem 2 for the method with a fixed regularization constant. Due to (76), the average number of tries of different M_t per one phase is only two.*

Finally, let us present an adaptive version of Algorithm 2, which is suitable for solving convex minimization problems (Algorithm 4 below).

Algorithm 4 Adaptive Regularized Newton with Lazy Hessians

Input: $\mathbf{x}_0 \in \mathbb{R}^d$, $m \geq 1$. Fix some $M_0 > 0$.

- 1: **for** $t = 0, 1, \dots$ **do**
 - 2: Compute snapshot Hessian $\nabla^2 f(\mathbf{x}_{tm})$
 - 3: **do**
 - 4: Update $M_t = 2 \cdot M_t$
 - 5: **for** $i = 1, \dots, m$ **do**
 - 6: Denote $k = tm + i - 1$ and set $\lambda_k = \sqrt{M_t \|\nabla f(\mathbf{x}_k)\|_*}$
 - 7: Compute lazy Newton step $\mathbf{x}_{k+1} = (\nabla^2 f(\mathbf{x}_{tm}) + \lambda_k \mathbf{B})^{-1} \nabla f(\mathbf{x}_k)$
 - 8: **until** $f(\mathbf{x}_{tm}) - f(\mathbf{x}_{tm+m}) \geq \sum_{i=1}^m \frac{1}{\lambda_{tm+i-1}} \|\nabla f(\mathbf{x}_{tm+i})\|_*^2$
 - 9: Set $M_{t+1} = \frac{1}{4} \cdot M_t$
-

Repeating the previous reasoning, we straightforwardly obtain the same global guarantees for this process as in Theorem 4 for the method with a fixed Lipschitz constant. The cost of adaptive search again is only one extra try of regularization parameter in average per phase.

7 Experiments

We demonstrate an illustrative numerical experiment on the performance of the proposed second-order methods with lazy Hessian updates. We consider the following convex minimization problem with the Soft Maximum objective (log-sum-exp):

$$\min_{\mathbf{x} \in \mathbb{R}^d} f(\mathbf{x}) := \mu \ln \left(\sum_{i=1}^n \exp \left(\frac{\langle \mathbf{a}_i, \mathbf{x} \rangle - b_i}{\mu} \right) \right) \approx \max_{1 \leq i \leq n} [\langle \mathbf{a}_i, \mathbf{x} \rangle - b_i].$$

The vectors $\mathbf{a}_1, \dots, \mathbf{a}_n \in \mathbb{R}^d$ and $\mathbf{b} \in \mathbb{R}^n$ form our data that we generate randomly from a uniform distribution on $[-1, 1]$. $\mu > 0$ is a smoothing parameter.

For the primal norm (2), we use the matrix

$$\mathbf{B} := \sum_{i=1}^n \mathbf{a}_i \mathbf{a}_i^\top + \delta \mathbf{I} \succ 0, \quad (78)$$

where $\delta > 0$ is a small perturbation parameter to ensure positive definiteness. Then, the corresponding Lipschitz constant of the Hessian is bounded by (see e.g. [7, Example 1.3.5]):

$$L = \frac{2}{\mu^2}. \quad (79)$$

Since the problem is convex, we can apply Newton's method with gradient regularization (Algorithm 2). In Figure 1, we compare different values of parameter m that is the frequency of updating the Hessian. The regularization parameter is fixed as $M := 1$. We also show the performance of the Gradient Method as a standard baseline.

We see that increasing the parameter m , thus reusing old Hessians for more of the inner steps, we significantly improve the overall performance of the method in terms of the total computational time. The best frequency is $m := d$ which confirms our theory.




Figure 1: Gradient norm depending on the number of computed gradients and on running time, when varying the frequency m of updating the Hessian in Algorithm 2.

8 Discussion

In this work, we have developed new second-order algorithms with *lazy Hessian updates* for solving general possibly non-convex optimization problems. We show that it can be very efficient to reuse the previously computed Hessian for several iterations of the method, instead of updating the Hessian each step.

Indeed, in general, the cost of computing the Hessian matrix is d times more expensive than computing one gradient. At the same time, it is intuitively clear that even inexact second-order information from the past should significantly help the method in dealing with ill-conditioning of the problem. In our analysis, we show that this intuition truly works.

By using cubic regularization and gradient regularization techniques, we establish fast global and local rates for our second-order methods with lazy Hessian updates. We show that the optimal strategy for updating the Hessian is once per d iterations, which gives a provable

improvement of the total arithmetic complexity by a factor of \sqrt{d} . Our approach also works with classical Newton steps (without regularization), achieving a local quadratic convergence.

One important direction for further research can be a study of the problems with a *specific Hessian structure* (e.g. sparsity or a certain spectral clustering). Then, we may need to have different schedules for updating the Hessian matrix, while it can also help in solving each step more efficiently.

Another interesting question is to make connections between our approach and classical *Quasi-Newton* methods [29], which gradually update the approximation of the Hessian after each step. Recently discovered non-asymptotic complexity bounds [32, 31] for Quasi-Newton methods may be especially useful for reaching this goal.

Finally, it seems to be very important to study the effect of lazy Hessian updates for convex optimization in more details. In our analysis, we studied only a general non-convex convergence in terms of the gradient norm. Another common accuracy measure in the convex case is the *functional residual*. Thus, it could be possible to prove some better convergence rates using this measure, as well as considering *accelerated* [26] and *super-universal* [8] second-order schemes.

We keep these questions for further investigation.

References

- [1] Ilan Adler, Zhiyue T Hu, and Tianyi Lin. New proximal Newton-type methods for convex optimization. In *2020 59th IEEE Conference on Decision and Control (CDC)*, pages 4828–4835. IEEE, 2020.
- [2] Francis Bach. Self-concordant analysis for logistic regression. *Electronic Journal of Statistics*, 4:384–414, 2010.
- [3] Coralia Cartis, Nicholas IM Gould, and Philippe L Toint. Adaptive cubic regularisation methods for unconstrained optimization. Part I: motivation, convergence and numerical results. *Mathematical Programming*, 127(2):245–295, 2011.
- [4] Coralia Cartis, Nicholas IM Gould, and Philippe L Toint. Adaptive cubic regularisation methods for unconstrained optimization. Part II: worst-case function-and derivative-evaluation complexity. *Mathematical programming*, 130(2):295–319, 2011.
- [5] Coralia Cartis, Nicholas IM Gould, and Philippe L Toint. On the oracle complexity of first-order and derivative-free algorithms for smooth nonconvex minimization. *SIAM Journal on Optimization*, 22(1):66–86, 2012.
- [6] Andrew R Conn, Nicholas IM Gould, and Philippe L Toint. *Trust region methods*. SIAM, 2000.
- [7] Nikita Doikov. *New second-order and tensor methods in Convex Optimization*. PhD thesis, Université catholique de Louvain, 2021.

- [8] Nikita Doikov, Konstantin Mishchenko, and Yurii Nesterov. Super-universal regularized Newton method. *arXiv preprint arXiv:2208.05888*, 2022.
- [9] Nikita Doikov and Yurii Nesterov. Convex optimization based on global lower second-order models. *Advances in Neural Information Processing Systems*, 33, 2020.
- [10] Nikita Doikov and Yurii Nesterov. Gradient regularization of Newton method with Bregman distances. *arXiv preprint arXiv:2112.02952*, 2021.
- [11] Nikita Doikov and Yurii Nesterov. Minimizing uniformly convex functions by cubic regularization of Newton method. *Journal of Optimization Theory and Applications*, pages 1–23, 2021.
- [12] Nikita Doikov and Yurii Nesterov. Affine-invariant contracting-point methods for convex optimization. *Mathematical Programming*, pages 1–23, 2022.
- [13] Pavel Dvurechensky and Yurii Nesterov. Global performance guarantees of second-order methods for unconstrained convex minimization. Technical report, CORE Discussion Paper, 2018.
- [14] Jinyan Fan. A Shamanskii-like Levenberg-Marquardt method for nonlinear equations. *Computational Optimization and Applications*, 56(1):63–80, 2013.
- [15] Geovani N Grapiglia, Max LN Gonçalves, and GN Silva. A cubic regularization of Newton’s method with finite difference hessian approximations. *Numerical Algorithms*, 90(2):607–630, 2022.
- [16] Geovani N Grapiglia and Yurii Nesterov. Regularized Newton methods for minimizing functions with Hölder continuous Hessians. *SIAM Journal on Optimization*, 27(1):478–506, 2017.
- [17] Geovani N Grapiglia and Yurii Nesterov. Accelerated regularized Newton methods for minimizing composite convex functions. *SIAM Journal on Optimization*, 29(1):77–99, 2019.
- [18] Slavomír Hanzely, Dmitry Kamzolov, Dmitry Pasechnyuk, Alexander Gasnikov, Peter Richtárik, and Martin Takáč. A damped Newton method achieves global $O(1/k^2)$ and local quadratic convergence rate. *arXiv preprint arXiv:2211.00140*, 2022.
- [19] Rustem Islamov, Xun Qian, Slavomír Hanzely, Mher Safaryan, and Peter Richtárik. Distributed Newton-type methods with communication compression and Bernoulli aggregation. *arXiv preprint arXiv:2206.03588*, 2022.
- [20] Leonid V Kantorovich. Functional analysis and applied mathematics. [in Russian]. *Uspekhi Matematicheskikh Nauk*, 3(6):89–185, 1948.

- [21] Sai Praneeth Karimireddy, Sebastian U Stich, and Martin Jaggi. Global linear convergence of Newton’s method without strong-convexity or Lipschitz gradients. *arXiv preprint arXiv:1806.00413*, 2018.
- [22] Francesco Lampariello and Marco Sciandrone. Global convergence technique for the Newton method with periodic Hessian evaluation. *Journal of optimization theory and applications*, 111(2):341–358, 2001.
- [23] Kenneth Levenberg. A method for the solution of certain non-linear problems in least squares. *Quarterly of applied mathematics*, 2(2):164–168, 1944.
- [24] Donald W. Marquardt. An algorithm for least-squares estimation of nonlinear parameters. *Journal of the society for Industrial and Applied Mathematics*, 11(2):431–441, 1963.
- [25] Konstantin Mishchenko. Regularized Newton method with global $O(1/k^2)$ convergence. *arXiv preprint arXiv:2112.02089*, 2021.
- [26] Yurii Nesterov. *Lectures on convex optimization*, volume 137. Springer, 2018.
- [27] Yurii Nesterov and Arkadi Nemirovski. *Interior-point polynomial algorithms in convex programming*. SIAM, 1994.
- [28] Yurii Nesterov and Boris Polyak. Cubic regularization of Newton’s method and its global performance. *Mathematical Programming*, 108(1):177–205, 2006.
- [29] Jorge Nocedal and Stephen Wright. *Numerical optimization*. Springer Science & Business Media, 2006.
- [30] James M Ortega and Werner C Rheinboldt. *Iterative solution of nonlinear equations in several variables*. SIAM, 2000.
- [31] Anton Rodomanov. *Quasi-Newton Methods with Provable Efficiency Guarantees*. PhD thesis, UCL-Université Catholique de Louvain, 2022.
- [32] Anton Rodomanov and Yurii Nesterov. New results on superlinear convergence of classical quasi-Newton methods. *Journal of optimization theory and applications*, 188(3):744–769, 2021.
- [33] VE Shamanskii. A modification of Newton’s method. *Ukrainian Mathematical Journal*, 19(1):118–122, 1967.
- [34] Tianxiao Sun and Quoc Tran-Dinh. Generalized self-concordant functions: a recipe for Newton-type methods. *Mathematical Programming*, 178(1-2):145–213, 2019.
- [35] Chang-yu Wang, Yuan-yuan Chen, and Shou-qiang Du. Further insight into the Shaman-skii modification of Newton method. *Applied mathematics and computation*, 180(1):46–52, 2006.