

Convergence of a splitting method for a general interest rate model

Gabriel Lord · Mengchao Wang ✉

Abstract We prove mean-square convergence of a novel numerical method, the tamed-splitting method, for a generalized Ait-Sahalia interest rate model. The method is based on a Lamperti transform, splitting and applying a tamed numerical method for the nonlinearity. The main difficulty in the analysis is caused by the non-globally Lipschitz drift coefficients of the model. We examine the existence, uniqueness of the solution and boundedness of moments for the transformed SDE. We then prove bounded moments and inverses moments for the numerical approximation. The tamed-splitting method is a hybrid method in the sense that a backstop method is invoked to prevent solutions from overshooting zero and becoming negative. We successfully recover the mean-square convergence rate of order one for the tamed-splitting method. In addition we prove that the probability of ever needing the backstop method to prevent a negative value can be made arbitrarily small. In our numerical experiments we compare to other numerical methods in the literature for realistic parameter values.

Keywords Ait-Sahalia model, tamed-splitting method, Lamperti transformation, mean-square convergence rate

AMS subject classification: 60H35, 60H15, 65C30.

1 Introduction

Stochastic differential equations (SDEs) are widely used in various scientific areas to model real-life phenomena affected by random noise. Here we consider the generalized Ait-Sahalia interest rate model introduced in [1]

$$dX_t = (a_{-1}X_t^{-1} - a_0 + a_1X_t - a_2X_t^\gamma) dt + bX_t^\theta dW_t, \quad X_0 = x_0, \quad t > 0 \quad (1)$$

which has since been investigated by various authors (see, e.g. [5], [8], [12]). Here, $a_{-1}, a_0, a_1, a_2, b > 0$, both $\theta, \gamma > 1$ and $\{W_t\}_{t \in [0, \infty)}$ is a one-dimensional Brownian motion on a filtered probability space $(\Omega, \mathcal{F}, \mathbb{P}, \{\mathcal{F}_t\}_{t \geq 0})$ with respect to the normal filtration $\{\mathcal{F}_t\}_{t \geq 0}$.

The model (1) clearly violates the Lipschitz and linear growth conditions which are traditionally imposed in the study of SDEs and their simulation (see [10, 15, 18]). As such it forms an interesting test case for numerical methods, for $\theta \in [\frac{1}{2}, 1]$ see for example [2–4, 11]. For the diffusion coefficient $\theta > 1$, as we consider here, the model was studied in [21], where strong convergence of the backward

Gabriel Lord
Department of Mathematics, IMAPP, Radboud University, Nijmegen, The Netherlands
E-mail: gabriel.lord@ru.nl

Mengchao Wang ✉
Department of Mathematics, IMAPP, Radboud University, Nijmegen, The Netherlands
School of Mathematics and Statistics, HNP-LAMA, Central South University, Changsha 410083, P. R. China
E-mail: mengchao.wang@ru.nl

This work was supported by China Scholarship Council, Graduate Research and Innovation Program of Central South University(No.CX20220246) and Radboud University Nijmegen.

Euler method was proved, but without revealing any convergence rate. Deng et.al. [6] examined the analytical properties of the model with Poisson jumps, including positivity, boundedness and pathwise asymptotic estimates. They applied the Euler-Maruyama (EM) method and proved that the explicit scheme converges in probability to the true solution of the model. Emmanuel et.al in [7] study analytical properties for the true solution of the model with delay and construct new truncated Euler-Maruyama (EM) methods to study properties of the numerical solutions under the local Lipschitz condition plus a Khasminskii-type condition. Zhao et.al. in [23] proved a mean-square convergence rate of order one half for the backward Euler method (BEM) for a generalized Ait-Sahalia interest rate model with Poisson jumps and showed that the BEM preserves the positivity of the original problem.

To the best of our knowledge, no strong convergence order 1 has been reported in the literature for explicit numerical approximations of (1). In this paper, we show that a strongly convergent numerical scheme can be constructed by an application of the Lamperti transform followed by a splitting approach where, rather than the nonlinearities being solved exactly, a tamed method such as in [13] is used. As such our analysis combines splitting approaches with a taming scheme. We call this the tamed-splitting method (TSM). In our numerical experiments we compare the tamed-splitting method to a standard splitting method (solving the drift nonlinearities exactly, denoted **Splitting**), the backward Euler method (BEM) for the Lamperti transform, tamed Euler method (TEM) for the transformed SDE, and backward Euler method for the original SDE (1) (**RefBEM**).

The structure of the article is as follows. In Section 2 we present the SDE arising from the Lamperti transform of (1) and examine existence, uniqueness as well as prove moment and inverse moment bounds on the solution. In Section 3, we introduce our two numerical methods. For the tamed-splitting method we prove moment and inverse moment bounds of the numerical solution in Section 4. Then, by introducing an auxiliary process, we prove the mean squared convergence rate of the tamed-splitting method is one. In Section 5 we show that the probability of a numerical solution taking a negative value is arbitrarily small, that is, this probability tends to zero as the step size tends to zero. Finally in Section 6 we numerically compare convergence and efficiency of several commonly used methods.

2 The Ait-Sahalia model

We introduce the notation we use throughout. Given a filtered probability space $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \in [0, T]}, \mathbb{P})$ satisfying the usual hypotheses we let W_t be a Brownian motion defined on that space. We let \mathbb{E} denote the expectation and $L^p(\Omega; \mathbb{R})$ the space of p -times integrable random variables with $\|\xi\|_{L^p(\Omega; \mathbb{R})} := (\mathbb{E}[|\xi|^p])^{1/p}$ for any $p \geq 1$. Let $x \wedge y := \min\{x, y\}$ for any $x, y \in \mathbb{R}$. For notational simplicity, the letter C is used to denote a generic positive constant, which is independent of the time stepsize and may vary for each appearance.

The well-posedness of the Ait-Sahalia model (1) was proved in [21, Theorem 2.1].

Proposition 1 *Let the initial data $X_0 = x_0 > 0$. For constants $a_{-1}, a_0, a_1, a_2, b > 0$ and $\gamma, \theta > 1$, the problem (1) admits a unique positive global solution.*

We now apply the Lamperti transform to obtain an SDE with additive noise. Taking $Y = X^{1-\theta}$ and using Itô's formula we get

$$dY_t = f(Y_t)dt + b(1-\theta)dW_t, \quad t \in (0, T], \quad (2)$$

where

$$f(x) := (\theta - 1)\left(a_2 x^{\frac{\gamma-\theta}{1-\theta}} - a_1 x - a_{-1} x^{\frac{\theta+1}{\theta-1}} + a_0 x^{\frac{\theta}{\theta-1}} + \frac{1}{2}b^2 \theta x^{-1}\right) \quad (3)$$

and $Y_0 = X_0^{1-\theta} \in \mathbb{R}^+$. We first examine existence and uniqueness for the transformed SDE (2). Before proceeding furthermore, we introduce the operator $\mathbb{L} : C^2(\mathbb{R}_+ \times [0, \infty), \mathbb{R}) \rightarrow C(\mathbb{R}_+ \times [0, \infty), \mathbb{R})$ defined by

$$\mathbb{L}\phi(x, t) := \phi_x(x, t)f(x) + \frac{1}{2}\phi_{xx}(x, t)(b(1-\theta))^2, \quad \phi(x, t) \in C^2(\mathbb{R}_+ \times [0, \infty), \mathbb{R}). \quad (4)$$

If the second variable t vanishes, we can rewrite (4) as follows

$$\mathbb{L}\phi(x) := \phi_x(x)f(x) + \frac{1}{2}\phi_{xx}(x)(b(1-\theta))^2, \quad \phi(x) \in C^2(\mathbb{R}_+, \mathbb{R}).$$

Proposition 2 *Let the initial data $Y_0 = X_0^{1-\theta} = x_0^{1-\theta} > 0$. For constants $a_{-1}, a_0, a_1, a_2, b > 0$ and $\gamma, \theta > 1$, the problem (2) admits a unique positive global solution, which almost surely satisfies*

$$Y_t = Y_0 + \int_0^t f(Y_s) ds + \int_0^t b(1-\theta) dW_s, \quad t \geq 0. \quad (5)$$

Proof of Proposition 2 It is straightforward to see that the drift coefficients of (2) are locally Lipschitz continuous in $(0, \infty)$. Following the standard arguments in [19] and noting $Y_0 > 0$, one can show that there is a unique maximal local solution $Y_t, t \in [0, \tau_e)$, where τ_e is the stopping time of the explosion or first zero time. To confirm we have a global solution, we need to prove $\tau_e = \infty$ a.s. For any sufficiently large positive integer n , satisfying $1/n < Y_0 < n$, we define the stopping times

$$\tau_n := \inf \{t \in [0, \tau_e) : Y_t \notin (1/n, n)\}, \quad (6)$$

where throughout this paper we set $\inf(\emptyset) = \infty$. Obviously τ_n is increasing as $n \rightarrow \infty$ and we set $\tau_\infty := \lim_{n \rightarrow \infty} \tau_n$. In view of (6), one knows $\tau_\infty \leq \tau_e$ a.s. If we can prove $\tau_n \rightarrow \infty$ a.s. as $n \rightarrow \infty$, then $\tau_e = \infty$ a.s. and $Y_t > 0$ a.s. for all $t \geq 0$, then the proof is complete. To prove $\tau_\infty = \infty$ a.s., it suffices to show that $\mathbb{P}\{\tau_n \leq T\} \rightarrow 0$ as $n \rightarrow \infty$ for any constant $T > 0$, which immediately implies $\mathbb{P}\{\tau_\infty = \infty\} = 1$, as required.

Given a fixed constant $\alpha \in (0, 1)$, let us define a function $V \in C^2((0, \infty), (0, \infty))$ by

$$V(x) := x^\alpha - \alpha \log x. \quad (7)$$

It is easy to check that $V(x) \rightarrow \infty$ as $x \rightarrow \infty$ or $x \rightarrow 0$ and that

$$V_x(x) = \alpha(x^{\alpha-1} - x^{-1}), \quad V_{xx}(x) = \alpha(\alpha-1)x^{\alpha-2} + \alpha x^{-2}. \quad (8)$$

Bearing (8) in mind, we see that

$$\begin{aligned} \mathbb{L}V(x) &= \alpha \left(x^{\alpha-1} - x^{-1} \right) (\theta - 1) \left[a_2 x^{\frac{\gamma-\theta}{1-\theta}} - a_1 x - a_{-1} x^{\frac{\theta+1}{\theta-1}} + a_0 x^{\frac{\theta}{\theta-1}} + \frac{1}{2} b^2 \theta x^{-1} \right] \\ &\quad + \frac{1}{2} b^2 (1-\theta)^2 \left[\alpha(\alpha-1)x^{\alpha-2} + \alpha x^{-2} \right] \\ &= \alpha(\theta-1) \left(a_2 x^{\alpha-1+\frac{\gamma-\theta}{1-\theta}} - a_1 x^\alpha - a_{-1} x^{\alpha-1+\frac{\theta+1}{\theta-1}} + a_0 x^{\alpha-1+\frac{\theta}{\theta-1}} + \frac{1}{2} b^2 \theta x^{\alpha-2} \right) \\ &\quad - \alpha(\theta-1) \left(a_2 x^{-1+\frac{\gamma-\theta}{1-\theta}} - a_1 - a_{-1} x^{\frac{2}{\theta-1}} + a_0 x^{-1+\frac{\theta}{\theta-1}} + \frac{1}{2} b^2 \theta x^{-2} \right) \\ &\quad + \frac{1}{2} b^2 (1-\theta)^2 \left[\alpha(\alpha-1)x^{\alpha-2} + \alpha x^{-2} \right]. \end{aligned} \quad (9)$$

Taking (9) into account and recalling $0 < \alpha < 1, \gamma > 1, \theta > 1, \lambda > 0$, there is a constant $K_1 > 0$ such that

$$\sup_{x \in (0, \infty)} \mathbb{L}V(x) \leq K_1 < \infty. \quad (10)$$

Indeed, if $\gamma + 1 > 2\theta$, we can directly get $\alpha - 1 + \frac{\theta+1}{\theta-1} > \frac{2}{\theta-1}$. Thus, it is easy to see that the highest power of x is $\alpha - 1 + \frac{\theta+1}{\theta-1}$ and the lowest power of x is $-1 + \frac{\gamma-\theta}{1-\theta}$ in (9). As a result of $\alpha a_2(\theta-1) > 0$ and $\alpha a_{-1}(\theta-1) > 0$, there is a constant $K_1 > 0$ such that (10) is fulfilled. If $\gamma + 1 = 2\theta$, it is easy to see that the highest power of x is $\alpha - 1 + \frac{\theta+1}{\theta-1}$ and the lowest power of x is -2 in (9). As a result of $\alpha a_{-1}(\theta-1) > 0$ and $\frac{1}{2} \alpha b^2 (1-\theta)^2 - \alpha(\theta-1) a_2 - \frac{1}{2} \alpha b^2 \theta (\theta-1) < 0$, there is a constant $K_1 > 0$ such that (10) is fulfilled. By the Itô formula [9] applied to $V(Y_{t \wedge \tau_n}), t \in [0, T]$, we infer

$$\mathbb{E}[V(Y_{T \wedge \tau_n})] \leq V(Y_0) + K_1 T < \infty, \quad \text{for all } T > 0.$$

Owing to the definitions (6), (7), we deduce from the above estimate that

$$\mathbb{P}(\tau_n \leq T) [V(1/n) \wedge V(n)] \leq \mathbb{E}V(Y_{T \wedge \tau_n}) \leq V(Y_0) + K_1 T < \infty. \quad (11)$$

This implies that $\lim_{n \rightarrow \infty} \mathbb{P}(\tau_n \leq T) = 0$ for any constant $T > 0$ and the proof is thus complete. \square

In the following error analysis, the moment bounds of the solution to (2) are frequently used. We first prove when the p -th negative moments of the solution to (2) are bounded.

Lemma 1 *Let all conditions in Proposition 1 hold and let $\{Y_t\}_{t \geq 0}$ be the unique solution to (2), given by (5). If one of the following two conditions holds:*

- (i) $p \geq \frac{2}{\theta-1}$ when $\gamma + 1 > 2\theta$;
(ii) $\frac{2}{\theta-1} \leq p \leq \frac{2a_2+b^2}{(\theta-1)b^2}$ when $\gamma + 1 = 2\theta$,
then

$$\sup_{t \in [0, \infty)} \mathbb{E}[|Y_t|^{-p}] < \infty. \quad (12)$$

Proof of Lemma 1 For a sufficiently large positive integer n satisfying $\frac{1}{n} < x_0 < n$, we define the stopping time

$$\tau_n := \inf\{t \in [0, \infty) : Y_t \notin (1/n, n)\}. \quad (13)$$

Also, we define $V_1 : \mathbb{R}_+ \times [0, \infty) \rightarrow \mathbb{R}_+$ as follows

$$V_1(x, t) := e^t x^{-p}, \quad x \in \mathbb{R}_+, t \in [0, \infty).$$

We compute that

$$\begin{aligned} \mathbb{L}V_1(x, t) &= e^t \left[-p(\theta-1)x^{-p-1}(a_2x^{\frac{\gamma-\theta}{1-\theta}} - a_1x - a_{-1}x^{\frac{\theta+1}{\theta-1}} - a_0x^{\frac{\theta}{\theta-1}} + \frac{1}{2}b^2\theta x^{-1}) \right. \\ &\quad \left. + \frac{1}{2}b^2(1-\theta)^2p(p+1)x^{-p-2} \right] \\ &= e^t \left[p(\theta-1)(-a_2x^{-p-1+\frac{\gamma-\theta}{1-\theta}} + a_1x^{-p} + a_{-1}x^{-p-1+\frac{\theta+1}{\theta-1}} + a_0x^{-p-1+\frac{\theta}{\theta-1}} \right. \\ &\quad \left. + \frac{1}{2}b^2(\theta-1)p(p(\theta-1)-1)x^{-p-2} \right], \end{aligned} \quad (14)$$

where the operator \mathbb{L} is defined in (4). Infact, the lowest power of x are $-p-1+\frac{\gamma-\theta}{1-\theta}$ for $\gamma+1 > 2\theta$ and $-p-2$ for $\gamma+1 = 2\theta$ in (14), respectively. Also, it is easy to see that the highest power of x is $-p-1+\frac{\theta+1}{\theta-1}$ in (14). Since $p \geq \frac{2}{\theta-1}$, we can get $-p-1+\frac{\theta+1}{\theta-1} \leq 0$. If $\gamma+1 = 2\theta$, due to condition (ii), we can obtain that

$$\frac{1}{2}b^2(\theta-1)p(p(\theta-1)-1) - p(\theta-1)a_2 \leq 0.$$

Therefore, one can find a constant $K_2 > 0$, such that

$$\mathbb{L}V_1(Y_t) \leq K_2e^t.$$

By the Itô formula for any $t \geq 0$,

$$\mathbb{E}[e^{t \wedge \tau_n} Y_{t \wedge \tau_n}^{-p}] \leq Y_0^{-p} + K_2e^t.$$

Letting $n \rightarrow \infty$ and applying Fatou's lemma, we obtain

$$\mathbb{E}[|Y_t|^{-p}] \leq \frac{x_0^{-p}}{e^t} + K_2, \quad t \geq 0.$$

The proof of Lemma 1 is thus completed. \square

Having obtained negative moment bounds the following lemma gives positive moment bounds of the solution to (2).

Lemma 2 *Let all conditions in Proposition 1 hold with $\gamma+1 \geq 2\theta$ and let $\{Y_t\}_{t \geq 0}$ be the unique solution to (2), given by (5). Then for any $p \geq (\gamma-1)/(\theta-1)$ it holds that*

$$\sup_{t \in [0, \infty)} \mathbb{E}[|Y_t|^p] < \infty.$$

Proof of Lemma 2. Define $V_2 : \mathbb{R}_+ \times [0, \infty) \rightarrow \mathbb{R}_+$ as follows

$$V_2(x, t) := e^t x^p, \quad x \in \mathbb{R}_+, \quad t \in [0, \infty). \quad (15)$$

Here, τ_n is defined in (13) and the operator \mathbb{L} in (4). Then we have

$$\begin{aligned} \mathbb{L}V_2(x, t) &= e^t [p(\theta - 1)x^{p-1}(a_2x^{\frac{\gamma-\theta}{1-\theta}} - a_1x - a_{-1}x^{\frac{\theta+1}{\theta-1}} + a_0x^{\frac{\theta}{\theta-1}} + \frac{1}{2}b^2\theta x^{-1}) \\ &\quad + \frac{1}{2}b^2(1-\theta)^2p(p-1)x^{p-2}] \\ &= e^t [p(\theta - 1)(a_2x^{p-1+\frac{\gamma-\theta}{1-\theta}} - a_1x^p - a_{-1}x^{p-1+\frac{\theta+1}{\theta-1}} + a_0x^{p-1+\frac{\theta}{\theta-1}} \\ &\quad + \frac{1}{2}b^2(\theta - 1)p[(\theta - 1)(p - 1) + \theta]x^{p-2})]. \end{aligned} \quad (16)$$

The highest power of x is $p - 1 + \frac{\theta+1}{\theta-1}$ in (16). The coefficient of this term is negative. According to $p \geq \frac{\gamma-1}{\theta-1}$, the lowest power of x is non-negative, i.e. $p - 1 + \frac{\gamma-\theta}{1-\theta} \geq 0$. Therefore, there exists a constant $K_3 > 0$ such that

$$\mathbb{L}V_2(X_t) \leq K_3 e^t. \quad (17)$$

The remaining proof is similar to the proof of Lemma 1 and thus omitted. \square

3 Derivation of the splitting and tamed-splitting methods

In this section, we first introduce the splitting and then the tamed-splitting method for the Ait-Sahalia model (1) which are based on the additive noise SDE (2). We then prove moment bounds and strong convergence rate of the tamed-splitting method.

For $N \in \mathbb{N}$ we construct on $[0, T]$ a uniform mesh $t_{i+1} = t_i + \tau$, $t_0 = 0$ with $\tau = \frac{T}{N}$ being the stepsize and introduce the notation

$$[t] := t_i, \quad \text{for } t \in [t_i, t_{i+1}), \quad i \in \{0, 1, \dots, N-1\}.$$

We can re-write (2) as

$$dY_t = [-\lambda Y_t + F(Y_t)]dt + b(1-\theta)dW_t, \quad (18)$$

where $f(x) = -\lambda x + F(x)$ with $\lambda := (\theta - 1)a_1$ and

$$F(x) := (\theta - 1)[a_2x^{\frac{\gamma-\theta}{1-\theta}} - a_{-1}x^{\frac{\theta+1}{\theta-1}} + a_0x^{\frac{\theta}{\theta-1}} + \frac{1}{2}b^2\theta x^{-1}]. \quad (19)$$

By the mean value theorem of differentiation

$$\begin{aligned} (F(x) - F(y)) \cdot (x - y) &= F'(\xi)|x - y|^2 \\ &= (-a_2(\gamma - \theta)\xi^{\frac{\gamma-1}{1-\theta}} - a_{-1}(\theta + 1)\xi^{\frac{2}{\theta-1}} + a_0\theta\xi^{\frac{1}{\theta-1}} - \frac{\theta-1}{2}b^2\theta\xi^{-2})|x - y|^2 \\ &\leq C|x - y|^2, \quad \xi \in (x, y) \text{ or } \xi \in (y, x), \quad x, y \in \mathbb{R}_+. \end{aligned} \quad (20)$$

The (standard) splitting method is derived from (18) by solving the four ODEs arising from the nonlinearity F along with the OU process. That is we solve the systems

$$\begin{cases} dU_t = (\theta - 1)a_2U_t^{\frac{\gamma-\theta}{1-\theta}} dt \\ dV_t = (1 - \theta)a_{-1}V_t^{\frac{\theta+1}{\theta-1}} dt \\ dP_t = (\theta - 1)a_0P_t^{\frac{\theta}{\theta-1}} dt \\ dQ_t = \frac{1}{2}b^2\theta(\theta - 1)Q_t^{-1} dt \\ dZ_t = \lambda Z_t dt + b(1 - \theta)dW_t \end{cases}$$

to get the following splitting method

$$\begin{aligned} Q_{n+1} &= \left[b^2 \theta (\theta - 1) \tau + \left[a_0 \tau \left[2(\theta - 1) a_{-1} \tau \left[(\gamma - 1) a_2 \tau + y_n^{\frac{\gamma-1}{\theta-1}} \right]^{\frac{1}{1-\gamma}} \right]^{\frac{1}{2}} \right]^{2(1-\theta)} \right]^{\frac{1}{2}}, \\ y_{n+1} &= e^{-(\theta-1)a_1\tau} Q_{n+1} + b(1-\theta) e^{-(\theta-1)a_1\tau} \Delta W_n, \end{aligned} \quad (21)$$

where $y_n \approx Y_{t_n}$ and $\Delta W_n := W_{t_{n+1}} - W_{t_n}$, $y_0 = Y_0$.

Although, as we see in Section 6, this scheme is easy to implement and has good properties, the direct analysis of (21) is not straightforward and does not generalise (for example to cases where it is not feasible to solve the nonlinear terms exactly).

Instead we choose to combine the splitting with a tamed Euler-Maruyama (see e.g. [13]) for the nonlinear drift. Combining the solutions of the OU process $dz = -\lambda z dt + b(1-\theta) dW_t$ and the tamed Euler-Maruyama approximation of $dv = F(v) dt$ we obtain the tamed-splitting method (TSM)

$$Y_{n+1} = e^{-\lambda\tau} \left(Y_n + \tau F^{(\tau)}(Y_n) + b(1-\theta) \Delta W_n \right), \quad (22)$$

where $\Delta W_n := W_{t_{n+1}} - W_{t_n}$ and the taming function is given by

$$F^{(\tau)}(x) := \frac{F(x)}{1 + \tau |F(x)|^2}.$$

We note that key to the analysis below is the following inequality

$$|F^{(\tau)}(x)|^2 = \frac{|F(x)|^2}{(1 + \tau |F(x)|^2)^2} \leq \frac{\tau^{-1} + |F(x)|^2}{1 + \tau |F(x)|^2} = \tau^{-1}. \quad (23)$$

Given $Y_n > 0$ we see from (22) there is no guarantee that Y_{n+1} is positive (see also Section 5). We combine this scheme with a positivity-preserving backstop scheme as in [14] for the Cox-Ingersoll-Ross process. We choose the backward Euler method

$$\hat{Y}_{n+1} = \hat{Y}_n + (F(\hat{Y}_{n+1}) - \lambda \hat{Y}_{n+1}) \tau + b(1-\theta) \Delta W_n \quad (24)$$

where from [17] we have

$$\|Y_{t_n} - \hat{Y}_n\|_{L^2(\Omega; \mathbb{R})} \leq C\tau.$$

We therefore define the tamed-splitting method with backstop:

$$\begin{aligned} \bar{Y}_{n+1} &= \left(e^{-\lambda\tau} (\bar{Y}_n + \tau F^{(\tau)}(\bar{Y}_n) + b(1-\theta) \Delta W_n) \right) \cdot \mathbb{1}_{\{\bar{Y}_{n+1} > 0\}} \\ &\quad + \left(\bar{Y}_n + (F(\bar{Y}_{n+1}) - \lambda \bar{Y}_{n+1}) \tau + b(1-\theta) \Delta W_n \right) \cdot \mathbb{1}_{\{\bar{Y}_{n+1} < 0\}}. \end{aligned} \quad (25)$$

For our analysis we define a continuous version of the tamed-splitting method (22) as

$$\bar{Y}_t = e^{-\lambda t} Y_0 + \int_0^t e^{-\lambda(t-s)} F^{(\tau)}(\bar{Y}_{\lfloor s \rfloor}) ds + \int_0^t e^{-\lambda(t-s)} b(1-\theta) dW_s. \quad (26)$$

We see in Section 6 that the splitting method (21) and tamed-splitting method (22) have the same rate of convergence, however (21) is more accurate in terms of the error constant.

4 Bounded moments and strong convergence of the tamed-splitting method

4.1 Bounded moments of the tamed-splitting method

Before proving the moment bounds of the numerical solution, we give a preliminary lemma.

Lemma 3 *For the approximation process \bar{Y}_t given by (26), we have that*

$$I := \bar{Y}_{[s]} \cdot F^{(\tau)}(\bar{Y}_{[s]}) = \frac{\bar{Y}_{[s]} \cdot F(\bar{Y}_{[s]})}{1 + \tau |F(\bar{Y}_{[s]})|^2} < C \max\{\tau^{\frac{1}{2}}, 1, \bar{Y}_{[s]}\}, \quad s \in [0, T].$$

Proof of Lemma 3 We divide into three cases:

- Case 1 : $0 < \bar{Y}_{[s]} \leq \tau$.
According to (23), we have

$$I \leq \bar{Y}_{[s]} \cdot \tau^{-\frac{1}{2}} \leq \tau \cdot \tau^{-\frac{1}{2}} \leq \tau^{\frac{1}{2}}. \quad (27)$$

- Case 2 : $\bar{Y}_{[s]} \geq 1$.
Since $a_2, a_{-1}, a_0 > 0$ and $\gamma > \theta > 1$, it is easy to show that there is a constant M such that

$$F(x) = (\theta - 1) \left[a_2 x^{\frac{\gamma-\theta}{1-\theta}} - a_{-1} x^{\frac{\theta+1}{\theta-1}} + a_0 x^{\frac{\theta}{\theta-1}} + \frac{1}{2} b^2 \theta x^{-1} \right] \leq M, \quad \text{for } x \geq 1. \quad (28)$$

Thus

$$I \leq \frac{M \bar{Y}_{[s]}}{1 + \tau |F(\bar{Y}_{[s]})|^2} \leq M \bar{Y}_{[s]}.$$

- Case 3 : $\tau < \bar{Y}_{[s]} < 1$.
Because $\bar{Y}_{[s]}$ has upper and lower bounds, we have $\bar{Y}_{[s]}^\eta \leq 1$ for any $\eta \geq 0$ and $\bar{Y}_{[s]}^\eta \leq \tau^\eta$ for any $\eta < 0$. Therefore, we directly calculate

$$\begin{aligned} I &= \frac{\bar{Y}_{[s]} \cdot F(\bar{Y}_{[s]})}{1 + \tau |F(\bar{Y}_{[s]})|^2} \\ &= \frac{(\theta - 1) \left[a_2 \bar{Y}_{[s]}^{\frac{\gamma+1-2\theta}{1-\theta}} - a_{-1} \bar{Y}_{[s]}^{\frac{2\theta}{\theta-1}} + a_0 \bar{Y}_{[s]}^{\frac{2\theta-1}{\theta-1}} + \frac{1}{2} b^2 \theta \right]}{1 + \tau (\theta - 1)^2 \left| a_2 \bar{Y}_{[s]}^{\frac{\gamma-\theta}{1-\theta}} - a_{-1} \bar{Y}_{[s]}^{\frac{\theta+1}{\theta-1}} + a_0 \bar{Y}_{[s]}^{\frac{\theta}{\theta-1}} + \frac{1}{2} b^2 \theta \bar{Y}_{[s]}^{-1} \right|^2}. \end{aligned}$$

For the numerator we have that

$$-a_{-1} \bar{Y}_{[s]}^{\frac{2\theta}{\theta-1}} + a_0 \bar{Y}_{[s]}^{\frac{2\theta-1}{\theta-1}} + \frac{1}{2} b^2 \theta \leq a_0 + \frac{1}{2} b^2 \theta =: K. \quad (29)$$

For the denominator we expand the square and note

$$-2a_{-1} a_0 \bar{Y}_{[s]}^{\frac{2\theta+1}{\theta-1}} - a_{-1} b^2 \theta \bar{Y}_{[s]}^{\frac{2}{\theta-1}} + a_0 b^2 \theta \bar{Y}_{[s]}^{\frac{1}{\theta-1}} \geq -2a_{-1} a_0 - a_{-1} b^2 \theta =: \bar{a}. \quad (30)$$

By (30) with $\tau < \bar{Y}_{[s]} < 1$ yields that for the denominator we have

$$\begin{aligned} & \left| a_2 \bar{Y}_{[s]}^{\frac{\gamma-\theta}{1-\theta}} - a_{-1} \bar{Y}_{[s]}^{\frac{\theta+1}{\theta-1}} + a_0 \bar{Y}_{[s]}^{\frac{\theta}{\theta-1}} + \frac{1}{2} b^2 \theta \bar{Y}_{[s]}^{-1} \right|^2 \\ & \geq a_2^2 \bar{Y}_{[s]}^{\frac{2\gamma-2\theta}{1-\theta}} + a_{-1}^2 \tau^{\frac{2\theta+2}{\theta-1}} + a_0^2 \tau^{\frac{2\theta}{\theta-1}} + \frac{1}{4} b^4 \theta \bar{Y}_{[s]}^{-2} + 2a_2 a_0 \bar{Y}_{[s]}^{\frac{\gamma-2\theta}{1-\theta}} - 2a_2 a_{-1} \bar{Y}_{[s]}^{\frac{\gamma-1-2\theta}{1-\theta}} + a_2 b^2 \theta \bar{Y}_{[s]}^{\frac{\gamma-1}{1-\theta}} + \bar{a}. \end{aligned}$$

As a consequence

$$\begin{aligned} I &\leq K \left(1 + \bar{Y}_{[s]}^{\frac{\gamma+1-2\theta}{1-\theta}} \right) \left[1 + \tau (\theta - 1)^2 \left(a_2^2 \bar{Y}_{[s]}^{\frac{2\gamma-2\theta}{1-\theta}} + a_{-1}^2 \tau^{\frac{2\theta+2}{\theta-1}} + a_0^2 \tau^{\frac{2\theta}{\theta-1}} + \frac{1}{4} b^4 \theta \bar{Y}_{[s]}^{-2} \right. \right. \\ & \quad \left. \left. + 2a_2 a_0 \bar{Y}_{[s]}^{\frac{\gamma-2\theta}{1-\theta}} - 2a_2 a_{-1} \bar{Y}_{[s]}^{\frac{\gamma-1-2\theta}{1-\theta}} + a_2 b^2 \theta \bar{Y}_{[s]}^{\frac{\gamma-1}{1-\theta}} + \bar{a} \right) \right]^{-1}. \end{aligned}$$

We multiply the numerator and denominator above by $\bar{Y}_{[s]}^{\frac{\gamma+1-2\theta}{\theta-1}}$ to get

$$I \leq [K(1 + \bar{Y}_{[s]}^{\frac{\gamma+1-2\theta}{\theta-1}})] \left[\bar{Y}_{[s]}^{\frac{\gamma+1-2\theta}{\theta-1}} + \tau(\theta-1)^2 \left(a_2^2 \bar{Y}_{[s]}^{\frac{\gamma-1}{1-\theta}} + \frac{1}{4} b^4 \theta \bar{Y}_{[s]}^{\frac{\gamma+3-4\theta}{\theta-1}} + 2a_2 a_0 \bar{Y}_{[s]}^{\frac{1}{\theta-1}} + a_2 b^2 \theta \bar{Y}_{[s]}^{-2} \right. \right. \\ \left. \left. + \bar{Y}_{[s]}^{\frac{\gamma+1-2\theta}{\theta-1}} \left(a_{-1}^2 \tau^{\frac{2\theta+2}{\theta-1}} + a_0^2 \tau^{\frac{2\theta}{\theta-1}} - 2a_2 a_{-1} \bar{Y}_{[s]}^{\frac{\gamma-1-2\theta}{1-\theta}} - \bar{a} \right) \right) \right]^{-1}.$$

If $\bar{Y}_{[s]} \geq \varepsilon_0 > \tau$, where ε_0 is a constant independent of τ , it's same as case 2. Otherwise, $Y_{[s]} \rightarrow \tau$, we have

$$I \xrightarrow{\bar{Y}_{[s]} \rightarrow \tau} [K(1 + \tau^{\frac{\gamma+1-2\theta}{\theta-1}})] \left[\tau^{\frac{\gamma+1-2\theta}{\theta-1}} + (\theta-1)^2 \left(a_2^2 \tau^{\frac{\gamma-\theta}{1-\theta}} + \frac{1}{4} b^4 \theta \tau^{\frac{\gamma+2-3\theta}{\theta-1}} + 2a_2 a_0 \tau^{\frac{\theta}{\theta-1}} + a_2 b^2 \theta \tau^{-1} \right) \right. \\ \left. + (\theta-1)^2 \left(a_{-1}^2 \tau^{\frac{\gamma+\theta+2}{\theta-1}} + a_0^2 \tau^{\frac{\gamma+\theta}{\theta-1}} - 2a_2 a_{-1} \tau^{\frac{\theta+1}{\theta-1}} + \bar{a} \tau^{\frac{\gamma-\theta}{\theta-1}} \right) \right]^{-1}.$$

We see that for arbitrarily small τ , we have $I < \infty$.

We have completed the three cases and the Lemma is proved. \square

In the following error analysis, the moment bounds of the numerical solution are frequently used. The next lemma indicates when q -th positive moments of the solution to (22) are bounded. The proof of Lemma 4 and Lemma 5 is inspired by [16].

Lemma 4 *The approximation process Y_t produced by (26) obeys, for any $q \geq 4$,*

$$\sup_{t \in [0, T]} \mathbb{E}[|\bar{Y}_t|^q] \leq C(1 + |Y_0|^q). \quad (31)$$

Proof of Lemma 4 Applying the Itô formula and (26), for any $l \in \mathbb{R}$, yields

$$(1 + |\bar{Y}_t|^2)^l = (1 + |Y_0|^2)^l + 2l \int_0^t (1 + |\bar{Y}_s|^2)^{l-1} \bar{Y}_s \cdot e^{-\lambda(t-[s])} F^{(\tau)}(\bar{Y}_{[s]}) ds \\ + l \int_0^t (1 + |\bar{Y}_s|^2)^{l-1} |e^{-\lambda(t-[s])} b(1-\theta)|^2 ds \\ + l(l-1) \int_0^t (1 + |\bar{Y}_s|^2)^{l-2} |\bar{Y}_s \cdot e^{-\lambda(t-[s])} b(1-\theta)|^2 ds \\ + 2l \int_0^t (1 + |\bar{Y}_s|^2)^{l-1} Y_s \cdot e^{-\lambda(t-[s])} b(1-\theta) dW_s. \quad (32)$$

Let $l = \frac{q}{2}$, and taking expectation we get

$$\mathbb{E}[(1 + |\bar{Y}_t|^2)^{\frac{q}{2}}] \leq (1 + |Y_0|^2)^{\frac{q}{2}} + q\mathbb{E} \left[\int_0^t (1 + |\bar{Y}_s|^2)^{\frac{q}{2}-1} \bar{Y}_s \cdot e^{-\lambda(t-[s])} F^{(\tau)}(\bar{Y}_{[s]}) ds \right] \\ + \frac{q(q-1)}{2} \mathbb{E} \left[\int_0^t (1 + |\bar{Y}_s|^2)^{\frac{q}{2}-1} |e^{-\lambda(t-[s])} b(1-\theta)|^2 ds \right].$$

Adding and subtracting terms we make a further decomposition as follows

$$\mathbb{E}[(1 + |\bar{Y}_t|^2)^{\frac{q}{2}}] \leq (1 + |Y_0|^2)^{\frac{q}{2}} \\ + q\mathbb{E} \left[\int_0^t (1 + |\bar{Y}_s|^2)^{\frac{q}{2}-1} (\bar{Y}_s - e^{-\lambda(s-[s])} \bar{Y}_{[s]}) \cdot e^{-\lambda(t-[s])} F^{(\tau)}(\bar{Y}_{[s]}) ds \right] \\ + q\mathbb{E} \left[\int_0^t (1 + |\bar{Y}_s|^2)^{\frac{q}{2}-1} \left(e^{-\lambda(s-[s])} \bar{Y}_{[s]} \cdot e^{-\lambda(t-[s])} F^{(\tau)}(\bar{Y}_{[s]}) + \frac{q-1}{2} |e^{-\lambda(t-[s])} b(1-\theta)|^2 \right) ds \right] \\ =: (1 + |Y_0|^2)^{\frac{q}{2}} + J_1 + J_2. \quad (33)$$

In what follows, we bound the two terms J_1 , J_2 separately. By the definition of Y_t , we have

$$J_1 = q\mathbb{E} \left[\int_0^t (1 + |\bar{Y}_s|^2)^{\frac{q}{2}-1} \int_{\lfloor s \rfloor}^s e^{-\lambda(s-\lfloor r \rfloor)} F^{(\tau)}(\bar{Y}_{\lfloor r \rfloor}) dr \cdot e^{-\lambda(t-\lfloor s \rfloor)} F^{(\tau)}(\bar{Y}_{\lfloor s \rfloor}) ds \right] \\ + q\mathbb{E} \left[\int_0^t (1 + |\bar{Y}_s|^2)^{\frac{q}{2}-1} \int_{\lfloor s \rfloor}^s e^{-\lambda(s-\lfloor r \rfloor)} b(1-\theta) dW_r \cdot e^{-\lambda(t-\lfloor s \rfloor)} F^{(\tau)}(\bar{Y}_{\lfloor s \rfloor}) ds \right].$$

Adding in and subtracting out $(1 + |\bar{Y}_{\lfloor s \rfloor}|^2)^{\frac{q}{2}-1}$

$$J_1 \leq q\mathbb{E} \left[\int_0^t (1 + |\bar{Y}_s|^2)^{\frac{q}{2}-1} \int_{\lfloor s \rfloor}^s |F^{(\tau)}(\bar{Y}_{\lfloor r \rfloor})| dr \cdot |F^{(\tau)}(\bar{Y}_{\lfloor s \rfloor})| ds \right] \\ + q\mathbb{E} \left[\int_0^t (1 + |\bar{Y}_{\lfloor s \rfloor}|^2)^{\frac{q}{2}-1} \int_{\lfloor s \rfloor}^s e^{-\lambda(s-\lfloor r \rfloor)} b(1-\theta) dW_r \cdot e^{-\lambda(t-\lfloor s \rfloor)} F^{(\tau)}(\bar{Y}_{\lfloor s \rfloor}) ds \right] \\ + q\mathbb{E} \left[\int_0^t ((1 + |\bar{Y}_s|^2)^{\frac{q}{2}-1} - (1 + |\bar{Y}_{\lfloor s \rfloor}|^2)^{\frac{q}{2}-1}) \int_{\lfloor s \rfloor}^s e^{-\lambda(s-\lfloor r \rfloor)} b(1-\theta) dW_r \cdot e^{-\lambda(t-\lfloor s \rfloor)} F^{(\tau)}(\bar{Y}_{\lfloor s \rfloor}) ds \right].$$

Using (23) and the Itô Isometry, we get

$$J_1 \leq q\mathbb{E} \left[\int_0^t (1 + |\bar{Y}_s|^2)^{\frac{q}{2}-1} (s - \lfloor s \rfloor) \cdot \tau ds \right] \\ + q\mathbb{E} \left[\int_0^t ((1 + |\bar{Y}_s|^2)^{\frac{q}{2}-1} - (1 + |\bar{Y}_{\lfloor s \rfloor}|^2)^{\frac{q}{2}-1}) \int_{\lfloor s \rfloor}^s e^{-\lambda(s-\lfloor r \rfloor)} b(1-\theta) dW_r \cdot e^{-\lambda(t-\lfloor s \rfloor)} F^{(\tau)}(\bar{Y}_{\lfloor s \rfloor}) ds \right].$$

With the aid of (32) taking $l = \frac{q}{2} - 1$, we can split J_1 into four additional terms:

$$J_1 \leq C\mathbb{E} \left[\int_0^t (1 + |\bar{Y}_s|^2)^{\frac{q}{2}} \right] + q(q-2)\mathbb{E} \left[\int_0^t \int_{\lfloor s \rfloor}^s (1 + |\bar{Y}_r|^2)^{\frac{q}{2}-2} \bar{Y}_r \cdot e^{-\lambda(s-\lfloor r \rfloor)} F^{(\tau)}(\bar{Y}_{\lfloor r \rfloor}) dr \right. \\ \times \left. \int_{\lfloor s \rfloor}^s e^{-\lambda(s-\lfloor r \rfloor)} b(1-\theta) dW_r \cdot e^{-\lambda(t-\lfloor s \rfloor)} F^{(\tau)}(\bar{Y}_{\lfloor s \rfloor}) ds \right] \\ + q(q-2)\mathbb{E} \left[\int_0^t \int_{\lfloor s \rfloor}^s (1 + |\bar{Y}_r|^2)^{\frac{q}{2}-2} \bar{Y}_r \cdot b(1-\theta) e^{-\lambda(s-\lfloor r \rfloor)} dW_r \right. \\ \times \left. \int_{\lfloor s \rfloor}^s e^{-\lambda(s-\lfloor r \rfloor)} b(1-\theta) dW_r \cdot e^{-\lambda(t-\lfloor s \rfloor)} F^{(\tau)}(\bar{Y}_{\lfloor s \rfloor}) ds \right] \\ + \frac{q(q-2)(q-4)}{2} \mathbb{E} \left[\int_0^t \int_{\lfloor s \rfloor}^s (1 + |\bar{Y}_r|^2)^{\frac{q}{2}-3} |\bar{Y}_r b(1-\theta) e^{-\lambda(s-\lfloor r \rfloor)}|^2 dr \right. \\ \times \left. \int_{\lfloor s \rfloor}^s e^{-\lambda(s-\lfloor r \rfloor)} b(1-\theta) dW_r \cdot e^{-\lambda(t-\lfloor s \rfloor)} F^{(\tau)}(\bar{Y}_{\lfloor s \rfloor}) ds \right] \\ + \frac{q(q-2)}{2} \mathbb{E} \left[\int_0^t \int_{\lfloor s \rfloor}^s (1 + |\bar{Y}_r|^2)^{\frac{q}{2}-2} |b(1-\theta) e^{-\lambda(s-\lfloor r \rfloor)}|^2 dr \right. \\ \times \left. \int_{\lfloor s \rfloor}^s e^{-\lambda(s-\lfloor r \rfloor)} b(1-\theta) dW_r \cdot e^{-\lambda(t-\lfloor s \rfloor)} F^{(\tau)}(\bar{Y}_{\lfloor s \rfloor}) ds \right] \\ =: C\mathbb{E} \left[\int_0^t (1 + |\bar{Y}_s|^2)^{\frac{q}{2}} \right] + J_{11} + J_{12} + J_{13} + J_{14}.$$

Next we estimate J_{11} - J_{14} term by term. Using (23) leads to

$$\begin{aligned} J_{11} &\leq q(q-2)\mathbb{E}\left[\int_0^t \int_{[s]}^s (1+|\bar{Y}_r|^2)^{\frac{q}{2}-2} |\bar{Y}_r| \cdot |F^{(\tau)}(\bar{Y}_{[r]})| dr \right. \\ &\quad \left. \times \left| \int_{[s]}^s e^{-\lambda(s-[r])} b(1-\theta) dW_r \right| \cdot |F^{(\tau)}(\bar{Y}_{[s]})| ds \right] \\ &\leq q(q-2)\tau^{-1}\mathbb{E}\left[\int_0^t \int_{[s]}^s (1+|\bar{Y}_r|^2)^{\frac{q}{2}-1} dr \left| \int_{[s]}^s e^{-\lambda(s-[r])} b(1-\theta) dW_r \right| ds \right]. \end{aligned}$$

We now use Young's inequality and the Hölder inequality to obtain

$$\begin{aligned} J_{11} &\leq C\tau^{-1}\mathbb{E}\left[\int_0^t \left(\int_{[s]}^s (1+|\bar{Y}_r|^2)^{\frac{q}{2}-1} dr\right)^{\frac{q}{q-1}} + \left|\int_{[s]}^s e^{-\lambda(s-[r])} b(1-\theta) dW_r\right|^q ds \right] \\ &\leq C\tau^{-1}\mathbb{E}\left[\int_0^t (s-[s])^{\frac{1}{q-1}} \int_{[s]}^s (1+|\bar{Y}_r|^2)^{\frac{q}{2}\cdot\frac{q-2}{q-1}} dr ds \right] \\ &\quad + C\tau^{-1}\mathbb{E}\left[\int_0^t \left(\int_{[s]}^s |e^{-\lambda(s-[r])} b(1-\theta)|^2 dr\right)^{\frac{q}{2}} ds \right]. \end{aligned}$$

Now, using the Itô isometry yields

$$\begin{aligned} J_{11} &\leq C\tau^{-1} \int_0^t \sup_{0 \leq r \leq s} \mathbb{E}[(1+|\bar{Y}_r|^2)^{\frac{q}{2}} (s-[s])^{\frac{q}{q-1}}] ds + C\tau^{-1} \mathbb{E}\left[\int_0^t |b(1-\theta)|^2 (s-[s])^{\frac{q}{2}} ds \right] \\ &\leq C \int_0^t \sup_{0 \leq r \leq s} \mathbb{E}[(1+|\bar{Y}_r|^2)^{\frac{q}{2}}] ds. \end{aligned} \tag{34}$$

For J_{12} , using the generalized Itô isometry (see, e.g., [22, Theorem 2.3.4]) and (23) yields

$$\begin{aligned} J_{12} &= q(q-2) \int_0^t \int_{[s]}^s \mathbb{E}[(1+|\bar{Y}_r|^2)^{\frac{q}{2}-2} \bar{Y}_r \cdot |b(1-\theta)|^2 e^{-2\lambda(s-[r])}] \\ &\quad \times e^{-\lambda(s-[r])} b(1-\theta) \cdot e^{-\lambda(t-[s])} F^{(\tau)}(\bar{Y}_{[s]})] dr ds \\ &\leq C\tau^{-\frac{1}{2}} \int_0^t \int_{[s]}^s \mathbb{E}[(1+|\bar{Y}_r|^2)^{\frac{q}{2}-1}] dr ds \\ &\leq C\tau^{-\frac{1}{2}} \int_0^t \sup_{0 \leq r \leq s} \mathbb{E}[(1+|\bar{Y}_r|^2)^{\frac{q}{2}}] (s-[s]) ds \\ &\leq C \int_0^t \sup_{0 \leq r \leq s} \mathbb{E}[(1+|\bar{Y}_r|^2)^{\frac{q}{2}}] ds. \end{aligned}$$

For J_{13} , using (23), the Young inequality, we obtain that

$$\begin{aligned} J_{13} &\leq C\mathbb{E}\left[\int_0^t \int_{[s]}^s (1+|\bar{Y}_r|^2)^{\frac{q}{2}-3} |\bar{Y}_r b(1-\theta)|^2 dr \left| \int_{[s]}^s e^{-\lambda(s-[r])} dW_r \right| \cdot |F^{(\tau)}(\bar{Y}_{[s]})| ds \right] \\ &= C\tau^{-\frac{1}{2}} \mathbb{E}\left[\int_0^t \int_{[s]}^s (1+|\bar{Y}_r|^2)^{\frac{q}{2}-2} dr \left| \int_{[s]}^s e^{-\lambda(s-[r])} dW_r \right| ds \right] \\ &\leq C\tau^{-\frac{1}{2}} \int_0^t \mathbb{E}\left[\left(\int_{[s]}^s (1+|\bar{Y}_r|^2)^{\frac{q-4}{2}} dr\right)^{\frac{q}{q-1}}\right] + \mathbb{E}\left[\left|\int_{[s]}^s e^{-\lambda(s-[r])} dW_r\right|^q\right] ds. \end{aligned}$$

Now, using the Itô isometry and the Hölder inequality obtains

$$\begin{aligned} J_{13} &\leq C\tau^{-\frac{1}{2}} \int_0^t \int_{[s]}^s \mathbb{E}[(1+|\bar{Y}_r|^2)^{\frac{q}{2}\cdot\frac{q-4}{q-1}}] dr (s-[s])^{\frac{1}{q-1}} ds + C\tau^{-\frac{1}{2}} \int_0^t \mathbb{E}\left[\left(\int_{[s]}^s |e^{-\lambda(s-[r])}|^2 dr\right)^{\frac{q}{2}}\right] ds \\ &\leq C\tau^{-\frac{1}{2}} \int_0^t \mathbb{E}[(1+|\bar{Y}_r|^2)^{\frac{q}{2}}] dr (s-[s])^{\frac{q}{q-1}} ds + C\tau^{-\frac{1}{2}} \int_0^t (s-[s])^{\frac{q}{2}} ds \\ &\leq C \int_0^t \sup_{0 \leq r \leq s} \mathbb{E}[(1+|\bar{Y}_r|^2)^{\frac{q}{2}}] ds. \end{aligned}$$

Similar to J_{13} , we obtain

$$J_{14} \leq C \int_0^t \sup_{0 \leq r \leq s} \mathbb{E}[(1 + |\bar{Y}_r|^2)^{\frac{q}{2}}] ds. \quad (35)$$

Thus, putting (34)-(35) together we get

$$J_1 \leq C \int_0^t \sup_{0 \leq r \leq s} \mathbb{E}[(1 + |\bar{Y}_r|^2)^{\frac{q}{2}}] ds.$$

Returning to J_2 in (33), with the help of Lemma 3, one can show that

$$\begin{aligned} J_2 &\leq q \mathbb{E} \left[\int_0^t (1 + |\bar{Y}_s|^2)^{\frac{q}{2}-1} \frac{\bar{Y}_{[s]} \cdot F(\bar{Y}_{[s]})}{1 + \tau |F(\bar{Y}_{[s]})|^2} ds \right] + \frac{|b(1-\theta)|^2 q(q-1)}{2} \mathbb{E} \left[\int_0^t (1 + |\bar{Y}_s|^2)^{\frac{q}{2}-1} ds \right] \\ &\leq C \mathbb{E} \left[\int_0^t (1 + |\bar{Y}_s|^2)^{\frac{q}{2}-1} (1 + |\bar{Y}_{[s]}|^2) ds \right] + C \mathbb{E} \left[\int_0^t (1 + |\bar{Y}_s|^2)^{\frac{q}{2}-1} ds \right] \\ &\leq C \int_0^t \sup_{0 \leq r \leq s} \mathbb{E}[(1 + |\bar{Y}_r|^2)^{\frac{q}{2}}] ds. \end{aligned}$$

Taking the estimates of J_1 and J_2 into account, we derive from (33) that for any $t \in [0, T]$

$$\sup_{0 \leq s \leq t} \mathbb{E}[(1 + |\bar{Y}_s|^2)^{\frac{q}{2}}] \leq (1 + |Y_0|^2)^{\frac{q}{2}} + C \int_0^t \sup_{0 \leq r \leq s} \mathbb{E}[(1 + |\bar{Y}_r|^2)^{\frac{q}{2}}] ds.$$

Applying the Gronwall inequality thus completes the proof. \square

Next, we need to prove inverse bound moments of numerical solution.

Lemma 5 *The approximation process Y_t produced by (26) obeys, for any $q \leq -1$,*

$$\sup_{t \in [0, T]} \mathbb{E}[|\bar{Y}_t|^q] \leq C(1 + |Y_0|^q).$$

Proof of Lemma 5 By (32) with $l = \frac{q}{2}$, $q \leq -1$, similar to the proof of Lemma 4, we have

$$\begin{aligned}
\mathbb{E}[(1+|\bar{Y}_t|^2)^{\frac{q}{2}}] &\leq (1+|Y_0|^2)^{\frac{q}{2}} + C\mathbb{E}\left[\int_0^t (1+|\bar{Y}_s|^2)^{\frac{q}{2}}\right] \\
&\quad + q(q-2)\mathbb{E}\left[\int_0^t \int_{[s]}^s (1+|\bar{Y}_r|^2)^{\frac{q}{2}-2} \bar{Y}_r \cdot e^{-\lambda(s-[r])} F^{(\tau)}(\bar{Y}_{[r]}) dr \right. \\
&\quad \times \left. \int_{[s]}^s e^{-\lambda(s-[r])} b(1-\theta) dW_r \cdot e^{-\lambda(t-[s])} F^{(\tau)}(\bar{Y}_{[s]}) ds\right] \\
&\quad + q(q-2)\mathbb{E}\left[\int_0^t \int_{[s]}^s (1+|\bar{Y}_r|^2)^{\frac{q}{2}-2} \bar{Y}_r \cdot b(1-\theta) e^{-\lambda(s-[r])} dW_r \right. \\
&\quad \times \left. \int_{[s]}^s e^{-\lambda(s-[r])} b(1-\theta) dW_r \cdot e^{-\lambda(t-[s])} F^{(\tau)}(\bar{Y}_{[s]}) ds\right] \\
&\quad + \frac{q(q-2)(q-4)}{2}\mathbb{E}\left[\int_0^t \int_{[s]}^s (1+|\bar{Y}_r|^2)^{\frac{q}{2}-3} |\bar{Y}_r b(1-\theta) e^{-\lambda(s-[r])}|^2 dr \right. \\
&\quad \times \left. \int_{[s]}^s e^{-\lambda(s-[r])} b(1-\theta) dW_r \cdot e^{-\lambda(t-[s])} F^{(\tau)}(\bar{Y}_{[s]}) ds\right] \\
&\quad + \frac{q(q-2)}{2}\mathbb{E}\left[\int_0^t \int_{[s]}^s (1+|\bar{Y}_r|^2)^{\frac{q}{2}-2} |b(1-\theta) e^{-\lambda(s-[r])}|^2 dr \right. \\
&\quad \times \left. \int_{[s]}^s e^{-\lambda(s-[r])} b(1-\theta) dW_r \cdot e^{-\lambda(t-[s])} F^{(\tau)}(\bar{Y}_{[s]}) ds\right] \\
&\quad + q\mathbb{E}\left[\int_0^t (1+|\bar{Y}_s|^2)^{\frac{q}{2}-1} \left(e^{-\lambda(s-[s])} \bar{Y}_{[s]} \cdot e^{-\lambda(t-[s])} F^{(\tau)}(\bar{Y}_{[s]}) \right. \right. \\
&\quad \left. \left. + \frac{(q-1)}{2} |e^{-\lambda(t-[s])} b(1-\theta)|^2\right) ds\right] \\
&=: (1+|Y_0|^2)^{\frac{q}{2}} + C\mathbb{E}\left[\int_0^t (1+|\bar{Y}_s|^2)^{\frac{q}{2}}\right] + I_1 + I_2 + I_3 + I_4 + I_5. \tag{36}
\end{aligned}$$

Next we estimate I_1 - I_5 one by one. Using (23) and the Young inequality leads to

$$\begin{aligned}
I_1 &\leq q(q-2)\mathbb{E}\left[\int_0^t \int_{[s]}^s (1+|\bar{Y}_r|^2)^{\frac{q}{2}-2} |\bar{Y}_r| \cdot |F^{(\tau)}(\bar{Y}_{[r]})| dr \right. \\
&\quad \times \left. \left| \int_{[s]}^s e^{-\lambda(s-[r])} b(1-\theta) dW_r \right| \cdot |F^{(\tau)}(\bar{Y}_{[s]})| ds\right] \\
&\leq q(q-2)\tau^{-1}\mathbb{E}\left[\int_0^t \int_{[s]}^s (1+|\bar{Y}_r|^2)^{\frac{q}{2}-1} dr \left| \int_{[s]}^s e^{-\lambda(s-[r])} b(1-\theta) dW_r \right| ds\right] \\
&\leq \frac{q(q-2)}{2}\tau^{-1}\mathbb{E}\left[\int_0^t \left(\int_{[s]}^s (1+|\bar{Y}_r|^2)^{\frac{q}{2}-1} dr\right)^2 + \left|\int_{[s]}^s e^{-\lambda(s-[r])} b(1-\theta) dW_r\right|^2 ds\right]
\end{aligned}$$

With the help of the Hölder inequality and the Itô isometry, one can get

$$\begin{aligned}
I_1 &\leq \frac{q(q-2)}{2}\tau^{-1}\mathbb{E}\left[\int_0^t (s-[s]) \int_{[s]}^s (1+|\bar{Y}_r|^2)^{q-2} dr ds\right] \\
&\quad + \frac{q(q-2)}{2}\tau^{-1}\mathbb{E}\left[\int_0^t \int_{[s]}^s |e^{-\lambda(s-[r])} b(1-\theta)|^2 dr ds\right] \\
&\leq \frac{q(q-2)}{2} \int_0^t \sup_{0 \leq r \leq s} \mathbb{E}[(1+|\bar{Y}_r|^2)^{\frac{q}{2}}(s-[s])] ds + \frac{q(q-2)}{2}\mathbb{E}\left[\int_0^t |b(1-\theta)|^2 ds\right] \\
&\leq C \int_0^t \sup_{0 \leq r \leq s} \mathbb{E}[(1+|\bar{Y}_r|^2)^{\frac{q}{2}}] ds. \tag{37}
\end{aligned}$$

Also, using the Young inequality and the Itô isometry yields

$$\begin{aligned}
I_2 &\leq \frac{q(q-2)}{2} \int_0^t \mathbb{E} \left[\left| \int_{[s]}^s (1 + |\bar{Y}_r|^2)^{\frac{q}{2}-2} \bar{Y}_r \cdot b(1-\theta) e^{-\lambda(s-[r])} dW_r \right|^2 \right] ds \\
&\quad + \frac{q(q-2)}{2} \int_0^t \mathbb{E} \left[\left| \int_{[s]}^s e^{-\lambda(s-[r])} b(1-\theta) \cdot e^{-\lambda(t-[s])} F^{(\tau)}(\bar{Y}_{[s]}) dW_r \right|^2 \right] ds \\
&\leq \frac{q(q-2)}{2} \int_0^t \mathbb{E} \left[\int_{[s]}^s (1 + |\bar{Y}_r|^2)^{q-4} |\bar{Y}_r|^2 \cdot |b(1-\theta)|^2 dr \right] ds \\
&\quad + \frac{q(q-2)}{2} \tau^{-1} \int_0^t \mathbb{E} \left[\int_{[s]}^s |b(1-\theta)|^2 dr \right] ds.
\end{aligned}$$

Because $q \leq -1$, it's easy to see that $(1 + |\bar{Y}_r|^2)^{\frac{q}{2}} \geq (1 + |\bar{Y}_r|^2)^{q-3} \geq (1 + |\bar{Y}_r|^2)^{q-4} \cdot |\bar{Y}_r|^2$. Thus

$$\begin{aligned}
I_2 &\leq \frac{q(q-2)}{2} \int_0^t \mathbb{E} \left[\int_{[s]}^s (1 + |\bar{Y}_r|^2)^{\frac{q}{2}} |b(1-\theta)|^2 dr \right] ds + \frac{q(q-2)}{2} \int_0^t |b(1-\theta)|^2 ds \\
&\leq C \int_0^t \sup_{0 \leq r \leq s} \mathbb{E}[(1 + |\bar{Y}_r|^2)^{\frac{q}{2}}] ds.
\end{aligned}$$

For I_3 , using (23), Young's inequality and the Itô isometry, we obtain that

$$\begin{aligned}
I_3 &\leq \frac{qb(q-2)(q-4)(1-\theta)}{2} \mathbb{E} \left[\int_0^t \int_{[s]}^s (1 + |\bar{Y}_r|^2)^{\frac{q}{2}-3} |\bar{Y}_r b(1-\theta)|^2 dr \right. \\
&\quad \left. \times \left| \int_{[s]}^s e^{-\lambda(s-[r])} dW_r \right| \cdot |F^{(\tau)}(\bar{Y}_{[s]})| ds \right] \\
&\leq C \tau^{-\frac{1}{2}} \mathbb{E} \left[\int_0^t \int_{[s]}^s (1 + |\bar{Y}_r|^2)^{\frac{q}{2}-2} dr \left| \int_{[s]}^s e^{-\lambda(s-[r])} dW_r \right| ds \right] \\
&\leq C \tau^{-\frac{1}{2}} \int_0^t \mathbb{E} \left[\left(\int_{[s]}^s (1 + |\bar{Y}_r|^2)^{\frac{q}{2}-2} dr \right)^2 \right] ds + C \tau^{-\frac{1}{2}} \int_0^t \mathbb{E} \left[\left| \int_{[s]}^s e^{-\lambda(s-[r])} dW_r \right|^2 \right] ds \\
&\leq C \int_0^t \sup_{0 \leq r \leq s} \mathbb{E}[(1 + |\bar{Y}_r|^2)^{\frac{q}{2}}] ds.
\end{aligned}$$

Similar to I_3 , we can obtain

$$I_4 \leq C \int_0^t \sup_{0 \leq r \leq s} \mathbb{E}[(1 + |\bar{Y}_r|^2)^{\frac{q}{2}}] ds.$$

Since $q \leq -1$, we have

$$\begin{aligned}
I_5 &= q e^{-\lambda(t+s-2[s])} \mathbb{E} \left[\int_0^t (1 + |\bar{Y}_s|^2)^{\frac{q}{2}-1} \bar{Y}_{[s]} \cdot F^{(\tau)}(\bar{Y}_{[s]}) ds \right] \\
&\quad + e^{-2\lambda(t-[s])} \frac{q(q-1)}{2} \mathbb{E} \left[\int_0^t (1 + |\bar{Y}_s|^2)^{\frac{q}{2}-1} |b(1-\theta)|^2 ds \right] \\
&= q e^{-\lambda(t+s-[s])} \mathbb{E} \left[\int_0^t (1 + |\bar{Y}_s|^2)^{\frac{q}{2}-1} \frac{\bar{Y}_{[s]} \cdot F(\bar{Y}_{[s]})}{1 + \tau |F(\bar{Y}_{[s]})|} ds \right] \\
&\quad + \frac{e^{-2\lambda(t-[s])} |b(1-\theta)|^2 q(q-1)}{2} \mathbb{E} \left[\int_0^t (1 + |\bar{Y}_s|^2)^{\frac{q}{2}-1} ds \right] \\
&= |q| e^{-\lambda(t+s-[s])} \int_0^t \mathbb{E} \left[\frac{1}{1 + \tau |F(\bar{Y}_{[s]})|} (1 + |\bar{Y}_s|^2)^{\frac{q}{2}-1} \right. \\
&\quad \left. \times (\theta-1) \left[-a_2 \bar{Y}_{[s]}^{\frac{\gamma+1-2\theta}{1-\theta}} + a_{-1} \bar{Y}_{[s]}^{\frac{2\theta}{\theta-1}} - a_0 \bar{Y}_{[s]}^{\frac{2\theta-1}{\theta-1}} - \frac{1}{2} b^2 \theta(\theta-1) \right] ds \right] \\
&\quad + \frac{e^{-2\lambda(t-[s])} |b(1-\theta)|^2 q(q-1)}{2} \int_0^t \mathbb{E} \left[(1 + |\bar{Y}_s|^2)^{\frac{q}{2}-1} \right] ds.
\end{aligned}$$

This, together with the Young inequality and Lemma 4, leads to

$$\begin{aligned}
I_5 &\leq |q|(\theta - 1) \int_0^t \mathbb{E}[(1 + |Y_s|^2)^{\frac{q}{2}-1} \cdot a_{-1} Y_{[s]}^{\frac{2\theta}{\theta-1}}] ds + C \int_0^t \mathbb{E}[(1 + |Y_s|^2)^{\frac{q}{2}-1}] ds \\
&\leq C \int_0^t \mathbb{E}[(1 + |Y_s|^2)^{q-2}] + \mathbb{E}[|Y_{[s]}|^{\frac{4\theta}{\theta-1}}] ds + C \int_0^t \mathbb{E}[(1 + |Y_s|^2)^{\frac{q}{2}-1}] ds \\
&\leq C \int_0^t \sup_{0 \leq r \leq s} \mathbb{E}[(1 + |Y_r|^2)^{\frac{q}{2}}] ds.
\end{aligned} \tag{38}$$

Taking (37)-(38) into account, we derive from (36) that for any $t \in [0, T]$.

$$\sup_{0 \leq s \leq t} \mathbb{E}[(1 + |Y_s|^2)^{\frac{q}{2}}] \leq (1 + |Y_0|^2)^{\frac{q}{2}} + C \int_0^t \sup_{0 \leq r \leq s} \mathbb{E}[(1 + |Y_r|^2)^{\frac{q}{2}}] ds.$$

Applying the Gronwall inequality thus completes the proof. \square

Remark 1 Lemma 4 proves the bounded moment of the numerical solution for $q \geq 4$. Infact, by applying the Hölder inequality, we can extend the results to $1 \leq q < 4$.

4.2 Strong convergence

Before giving the convergence result, we introduce the following auxiliary process,

$$\tilde{Y}_{n+1} = e^{-\lambda\tau}(\tilde{Y}_n + \tau F(Y_{t_n}) + b(1 - \theta)\Delta W_n), \tag{39}$$

where $\lambda := (\theta - 1)a_1$ (as in (19)). Then, we give the bounded moment of auxiliary process \tilde{Y}_n .

Lemma 6 *For the auxiliary process defined in (39), we have*

$$\mathbb{E}[|\tilde{Y}_n|^2] < \infty, \quad \text{for } n = 1, 2, \dots, N.$$

Proof of Lemma 6 It is easy to check that

$$\|\tilde{Y}_1\|_{L^2(\Omega; \mathbb{R})}^2 = |e^{-\lambda\tau}|^2 \|Y_0 + F(Y_{t_0})\tau + b(1 - \theta)\Delta W_0\|_{L^2(\Omega; \mathbb{R})}^2 < \infty.$$

Assume $\|\tilde{Y}_{n-1}\|_{L^2(\Omega; \mathbb{R})}^2 < \infty$, then by the definition of \tilde{Y}_n , we have

$$\begin{aligned}
|\tilde{Y}_n|^2 &= |e^{-\lambda\tau}|^2 |\tilde{Y}_{n-1} + F(Y_{t_{n-1}})\tau + b(\theta - 1)\Delta W_{n-1}|^2 \\
&\leq |\tilde{Y}_{n-1}|^2 + |F(Y_{t_{n-1}})|^2 \tau^2 + b^2(\theta - 1)^2 |\Delta W_{n-1}|^2 + 2 \langle \tilde{Y}_{n-1}, F(Y_{t_{n-1}}) \rangle \\
&\quad + 2 \langle \tilde{Y}_{n-1}, b(\theta - 1)\Delta W_{n-1} \rangle + 2 \langle F(Y_{t_{n-1}})\tau, b(\theta - 1)\Delta W_{n-1} \rangle.
\end{aligned}$$

Take the expectation of both sides, by the bounded and inverse bounded moment of Y_t , we have

$$\begin{aligned}
\|\tilde{Y}_n\|_{L^2(\Omega; \mathbb{R})}^2 &\leq \|\tilde{Y}_{n-1}\|_{L^2(\Omega; \mathbb{R})}^2 + \|F(Y_{t_{n-1}})\|_{L^2(\Omega; \mathbb{R})}^2 \tau^2 + b^2(\theta - 1)^2 \|\Delta W_{n-1}\|_{L^2(\Omega; \mathbb{R})}^2 \\
&\quad + 2\mathbb{E}[\langle \tilde{Y}_{n-1}, F(Y_{t_{n-1}}) \rangle] + 2\mathbb{E}[\langle \tilde{Y}_{n-1}, b(\theta - 1)\Delta W_{n-1} \rangle] \\
&\quad + 2\mathbb{E}[\langle F(Y_{t_{n-1}})\tau, b(\theta - 1)\Delta W_{n-1} \rangle] \\
&\leq 3\|\tilde{Y}_{n-1}\|_{L^2(\Omega; \mathbb{R})}^2 + 3\|F(Y_{t_{n-1}})\|_{L^2(\Omega; \mathbb{R})}^2 \tau^2 + 3b^2(\theta - 1)^2 \tau^2 < \infty.
\end{aligned}$$

This completes the lemma. \square

We are now ready to prove the mean-square convergence rate of order one for the scheme.

Theorem 1 *Let $\{Y_t\}_{0 \leq t \leq T}$ and $\{Y_n\}_{0 \leq n \leq N}$ be solutions to (2) and (22), respectively. Then for any time step $\tau > 0$ it holds that*

$$\sup_{N \in \mathbb{N}} \sup_{0 \leq n \leq N} \|Y_n - Y_{t_n}\|_{L^2(\Omega; \mathbb{R})} \leq C\tau.$$

Proof of Theorem 1 For $t \in [t_n, t_{n+1}]$, we re-write the exact and numerical solution to (18),

$$Y_t = e^{-\lambda(t-t_n)}Y_{t_n} + \int_{t_n}^t e^{-\lambda(t-r)}F(Y_r)dr + \int_{t_n}^t e^{-\lambda(t-r)}b(1-\theta)dW_r \quad (40)$$

and

$$\bar{Y}_t = e^{-\lambda(t-t_n)}\bar{Y}_{t_n} + \int_{t_n}^t e^{-\lambda(t-r)}F^{(\tau)}(\bar{Y}_r)dr + \int_{t_n}^t e^{-\lambda(t-r)}b(1-\theta)dW_r. \quad (41)$$

We subtract (40) from equation (41) to get

$$\begin{aligned} e(t) &:= Y_t - \bar{Y}_t \\ &= e^{-\lambda(t-t_n)}e(t_n) + \int_{t_n}^t e^{-\lambda(t-r)}F(Y_r) - e^{-\lambda(t-r)}F^{(\tau)}(\bar{Y}_r)dr \\ &\quad + \int_{t_n}^t (e^{-\lambda(t-r)} - e^{-\lambda(t-t_n)})b(1-\theta)dW_r. \end{aligned}$$

It is straightforward to check that $e(s)$ is the solution to the SDE

$$de(t) = [-\lambda e(t) + F(Y_t) - e^{-\lambda(t-t_n)}F^{(\tau)}(\bar{Y}_{t_n})]dt + (1 - e^{-\lambda(t-t_n)})b(1-\theta)dW_t.$$

Using Itô's formula, we have

$$\begin{aligned} |e(t)|^2 &= |e(t_n)|^2 + \int_{t_n}^t |(1 - e^{-\lambda(r-t_n)})b(1-\theta)|^2 dr \\ &\quad + 2 \int_{t_n}^t \left\langle e(r), -\lambda e(r) + F(Y_r) - e^{-\lambda(r-t_n)}F^{(\tau)}(\bar{Y}_{t_n}) \right\rangle dr \\ &\quad + 2 \int_{t_n}^t \left\langle e(r), (1 - e^{-\lambda(r-t_n)})b(1-\theta) \right\rangle dW_r. \end{aligned} \quad (42)$$

By Taylor's Theorem, there exists a constant K such that $1 - e^{-\lambda(r-t_n)} \leq \lambda(r-t_n) + K(r-t_n)^2$. Taking the expectation of both sides of (42) and using Itô's Isometry

$$\begin{aligned} \mathbb{E}[|e(t)|^2] &= \mathbb{E}[|e(t_n)|^2] + 2 \int_{t_n}^t \mathbb{E}[\langle e(r), -\lambda e(r) \rangle] + \mathbb{E}[\langle e(r), F(Y_r) - e^{-\lambda(r-t_n)}F^{(\tau)}(\bar{Y}_{t_n}) \rangle] dr \\ &\quad + \int_{t_n}^t \mathbb{E}[|(1 - e^{-\lambda(r-t_n)})b(1-\theta)|^2] dr \\ &\leq \mathbb{E}[|e(t_n)|^2] + 2 \int_{t_n}^t \mathbb{E}[\langle e(r), F(Y_r) - F(\bar{Y}_r) \rangle] dr \\ &\quad + 2 \int_{t_n}^t \mathbb{E}[\langle e(r), F(\bar{Y}_r) - e^{-\lambda(r-t_n)}F^{(\tau)}(\bar{Y}_{t_n}) \rangle] dr + C \int_{t_n}^t (r-t_n)^2 + (r-t_n)^4 dr \\ &\leq \mathbb{E}[|e(t_n)|^2] + 2J_1 + 2J_2 + C\tau^3. \end{aligned} \quad (43)$$

For J_1 , according to the definition of function F , we have

$$J_1 \leq C \int_{t_n}^t \mathbb{E}[|e(r)|^2] dr.$$

For J_2 ,

$$\begin{aligned} J_2 &= \int_{t_n}^t \mathbb{E}[\langle e(r), F(\bar{Y}_r) - F(\bar{Y}_{t_n}) \rangle] dr + \int_{t_n}^t \mathbb{E}[\langle e(r), F(\bar{Y}_{t_n}) - e^{-\lambda(r-t_n)}F^{(\tau)}(\bar{Y}_{t_n}) \rangle] dr \\ &=: J_{21} + J_{22}. \end{aligned}$$

For J_{22} , using the Young inequality, Lemma 4 and Lemma 5, we obtain

$$\begin{aligned}
J_{22} &= \int_{t_n}^t \mathbb{E}[\langle e(r), (1 - e^{-\lambda(r-t_n)})F(\bar{Y}_{t_n}) \rangle] dr + \int_{t_n}^t \mathbb{E}[\langle e(r), e^{-\lambda(r-t_n)}(F(\bar{Y}_{t_n}) - F^{(\tau)}(\bar{Y}_{t_n})) \rangle] dr \\
&\leq \int_{t_n}^t \mathbb{E}[|e(r)|^2] dr + C\tau^3 \mathbb{E}[|F(\bar{Y}_{t_n})|^2] + C\tau^3 \mathbb{E}\left[\left|\frac{\tau|F(\bar{Y}_{t_n})|^2 F(\bar{Y}_{t_n})}{1 + \tau|F(\bar{Y}_{t_n})|^2}\right|^2\right] \\
&\leq \int_{t_n}^t \mathbb{E}[|e(r)|^2] dr + C\tau^3(1 + \mathbb{E}[|\bar{Y}_{t_n}|^{\frac{2(\gamma-\theta)}{1-\theta}}] + \mathbb{E}[|\bar{Y}_{t_n}|^{\frac{2(\theta+1)}{\theta-1}}] + \mathbb{E}[|\bar{Y}_{t_n}|^{\frac{6(\gamma-\theta)}{1-\theta}}] + \mathbb{E}[|\bar{Y}_{t_n}|^{\frac{6(\theta+1)}{\theta-1}}]) \\
&\leq \int_{t_n}^t \mathbb{E}[|e(r)|^2] dr + C\tau^3.
\end{aligned}$$

Using Taylor theorem, we get

$$\begin{aligned}
&\mathbb{E}[\langle e(r), F(\bar{Y}(r)) - F(\bar{Y}_{t_n}) \rangle] \\
&= \mathbb{E}[\langle e(r), F'(\bar{Y}_{t_n}) \int_{t_n}^r e^{-\lambda(s-t_n)} F^{(\tau)}(\bar{Y}_{t_n}) ds \rangle] + \mathbb{E}[\langle e(r), F'(\bar{Y}_{t_n}) \int_{t_n}^r e^{-\lambda(s-t_n)} b(1-\theta) dW_s \rangle] \\
&\quad + \mathbb{E}[\langle e(r), F'(\bar{Y}_{t_n})(1 - e^{-\lambda(r-t_n)})\bar{Y}_{t_n} \rangle] + \mathbb{E}[\langle e(r), \int_0^1 F''(\bar{Y}_{t_n} + \sigma(\bar{Y}_r - \bar{Y}_{t_n})) |\bar{Y}_r - \bar{Y}_{t_n}|^2 d\sigma \rangle] \\
&=: H_1 + H_2 + H_3 + H_4.
\end{aligned}$$

Note that

$$\begin{aligned}
\bar{Y}_r - \bar{Y}_{t_n} &= \bar{Y}_r - e^{-\lambda(r-t_n)}\bar{Y}_{t_n} - (1 - e^{-\lambda(r-t_n)})\bar{Y}_{t_n} \\
&= \int_{t_n}^r e^{-\lambda(s-t_n)} F^{(\tau)}(\bar{Y}_{t_n}) ds + \int_{t_n}^r e^{-\lambda(s-t_n)} b(1-\theta) dW_s - (1 - e^{-\lambda(r-t_n)})\bar{Y}_{t_n}.
\end{aligned}$$

For $p \geq 2$, it's straightforward to derive that

$$\begin{aligned}
&\mathbb{E}[|\bar{Y}_r - \bar{Y}_{t_n}|^p] \\
&\leq C_p \mathbb{E}\left[\left|\int_{t_n}^r e^{-\lambda(s-t_n)} F^{(\tau)}(\bar{Y}_{t_n}) ds\right|^p\right] + C_p \mathbb{E}\left[\left|\int_{t_n}^r e^{-\lambda(s-t_n)} b(1-\theta) dW_s\right|^p\right] + C_p \mathbb{E}[(1 - e^{-\lambda(r-t_n)})\bar{Y}_{t_n}]^p \\
&\leq C_p \tau^{p-1} \int_{t_n}^r \mathbb{E}[|\bar{Y}_{t_n}|^p] ds + C_p \left(\int_{t_n}^r |b(1-\theta)|^2 ds\right)^{\frac{p}{2}} + C_p \tau^p \mathbb{E}[|\bar{Y}_{t_n}|^p] \\
&\leq C\tau^2(1 + \mathbb{E}[|\bar{Y}_{t_n}|^{\frac{2(\gamma-\theta)}{1-\theta}}] + \mathbb{E}[|\bar{Y}_{t_n}|^{\frac{2(\theta+1)}{\theta-1}}]) + C\tau^{\frac{p}{2}} \\
&\leq C\tau^{\frac{p}{2}}.
\end{aligned} \tag{44}$$

Using the Young inequality and Lemma 4, 5, we obtain

$$\begin{aligned}
H_1 &\leq \mathbb{E}[|e(r)|^2] + C\mathbb{E}\left[|F'(\bar{Y}_{t_n})|^2 \left|\int_{t_n}^r e^{-\lambda(s-t_n)} F^{(\tau)}(\bar{Y}_{t_n}) ds\right|^2\right] \\
&\leq \mathbb{E}[|e(r)|^2] + C\tau \mathbb{E}[|F'(\bar{Y}_{t_n})|^2 \int_{t_n}^r |F^{(\tau)}(\bar{Y}_{t_n})|^2 ds] \\
&\leq \mathbb{E}[|e(r)|^2] + C\tau^2 (\mathbb{E}[|F'(\bar{Y}_{t_n})|^4])^{\frac{1}{2}} (\mathbb{E}[|F(\bar{Y}_{t_n})|^4])^{\frac{1}{2}} \\
&\leq \mathbb{E}[|e(r)|^2] + C\tau^2 (1 + \mathbb{E}[|\bar{Y}_{t_n}|^{\frac{4(\gamma-\theta)}{1-\theta}}] + \mathbb{E}[|\bar{Y}_{t_n}|^{\frac{4(\theta+1)}{\theta-1}}])^{\frac{1}{2}} (1 + \mathbb{E}[|\bar{Y}_{t_n}|^{\frac{4(\gamma-1)}{1-\theta}}] + \mathbb{E}[|\bar{Y}_{t_n}|^{\frac{4(2)}{\theta-1}}])^{\frac{1}{2}} \\
&\leq \mathbb{E}[|e(r)|^2] + C\tau^2.
\end{aligned}$$

For H_2 , by (42) and (44), we have

$$\begin{aligned}
H_2 &= \mathbb{E} \left[\left\langle e^{-\lambda(r-t_n)} e(t_n), F'(\bar{Y}_{t_n}) \int_{t_n}^r e^{-\lambda(s-t_n)} b(1-\theta) dW_s \right\rangle \right] \\
&\quad + \mathbb{E} \left[\left\langle \int_{t_n}^r e^{-\lambda(r-\sigma)} F(Y_\sigma) - e^{-\lambda(r-t_n)} F^{(\tau)}(\bar{Y}_{t_n}) d\sigma, F'(\bar{Y}_{t_n}) \int_{t_n}^r e^{-\lambda(s-t_n)} b(1-\theta) dW_s \right\rangle \right] \\
&\quad + \mathbb{E} \left[\left\langle \int_{t_n}^r (e^{-\lambda(r-\sigma)} - e^{-\lambda(r-t_n)}) b(1-\theta) dW_\sigma, F'(\bar{Y}_{t_n}) \int_{t_n}^r e^{-\lambda(s-t_n)} b(1-\theta) dW_s \right\rangle \right] \\
&= \mathbb{E} \left[\left\langle \int_{t_n}^r (e^{-\lambda(r-\sigma)} - e^{-\lambda(r-t_n)}) F(Y_\sigma) d\sigma, F'(\bar{Y}_{t_n}) \int_{t_n}^r e^{-\lambda(s-t_n)} b(1-\theta) dW_s \right\rangle \right] \\
&\quad + \mathbb{E} \left[\left\langle \int_{t_n}^r e^{-\lambda(r-t_n)} (F(Y_\sigma) - F(Y_{t_n})) d\sigma, F'(\bar{Y}_{t_n}) \int_{t_n}^r e^{-\lambda(s-t_n)} b(1-\theta) dW_s \right\rangle \right] \\
&\quad + \mathbb{E} \left[\left\langle \int_{t_n}^r e^{-\lambda(r-t_n)} (F(Y_{t_n}) - F(\bar{Y}_{t_n})) d\sigma, F'(\bar{Y}_{t_n}) \int_{t_n}^r e^{-\lambda(s-t_n)} b(1-\theta) dW_s \right\rangle \right] \\
&\quad + \mathbb{E} \left[\left\langle \int_{t_n}^r e^{-\lambda(r-t_n)} (F(\bar{Y}_{t_n}) - F^{(\tau)}(\bar{Y}_{t_n})) d\sigma, F'(\bar{Y}_{t_n}) \int_{t_n}^r e^{-\lambda(s-t_n)} b(1-\theta) dW_s \right\rangle \right] \\
&\quad + \mathbb{E} \left[\left\langle \int_{t_n}^r (e^{-\lambda(r-\sigma)} - e^{-\lambda(r-t_n)}) b(1-\theta) dW_\sigma, F'(\bar{Y}_{t_n}) \int_{t_n}^r e^{-\lambda(s-t_n)} b(1-\theta) dW_s \right\rangle \right].
\end{aligned}$$

Then, using Hölder inequality, the Itô isometry and the bounded moment of exact and numerical solution to, one can get (18)

$$\begin{aligned}
H_2 &\leq \left(\mathbb{E} \left[\left| \int_{t_n}^r (e^{-\lambda(r-\sigma)} - e^{-\lambda(r-t_n)}) F(Y_\sigma) d\sigma \right|^2 \right] \mathbb{E} \left[\left| F'(\bar{Y}_{t_n}) \int_{t_n}^r e^{-\lambda(s-t_n)} b(1-\theta) dW_s \right|^2 \right] \right)^{\frac{1}{2}} \\
&\quad + \left(\mathbb{E} \left[\left| \int_{t_n}^r e^{-\lambda(r-t_n)} (F(Y_\sigma) - F(Y_{t_n})) d\sigma \right|^2 \right] \mathbb{E} \left[\left| F'(\bar{Y}_{t_n}) \int_{t_n}^r e^{-\lambda(s-t_n)} b(1-\theta) dW_s \right|^2 \right] \right)^{\frac{1}{2}} \\
&\quad + \left(\mathbb{E} \left[\left| \int_{t_n}^r (e^{-\lambda(r-\sigma)} - e^{-\lambda(r-t_n)}) b(1-\theta) dW_\sigma \right|^2 \right] \mathbb{E} \left[\left| F'(\bar{Y}_{t_n}) \int_{t_n}^r e^{-\lambda(s-t_n)} b(1-\theta) dW_s \right|^2 \right] \right)^{\frac{1}{2}} \\
&\leq C\tau^{\frac{3}{2}} \left(\mathbb{E} \left[\int_{t_n}^r |F(Y_\sigma)|^2 d\sigma \right] \cdot \mathbb{E} \left[|F'(\bar{Y}_{t_n})|^2 \int_{t_n}^r |b(1-\theta)|^2 ds \right] \right)^{\frac{1}{2}} \\
&\quad + C\tau^{\frac{1}{2}} \left(\mathbb{E} \left[\int_{t_n}^r |(F(Y_\sigma) - F(Y_{t_n}))|^2 d\sigma \right] \cdot \mathbb{E} \left[|F'(\bar{Y}_{t_n})|^2 \int_{t_n}^r |b(1-\theta)|^2 ds \right] \right)^{\frac{1}{2}} \\
&\quad + C \left(\mathbb{E} \left[\int_{t_n}^r |(e^{-\lambda(r-\sigma)} - e^{-\lambda(r-t_n)})|^2 d\sigma \right] \mathbb{E} \left[|F'(\bar{Y}_{t_n})|^2 \int_{t_n}^r |b(1-\theta)|^2 ds \right] \right)^{\frac{1}{2}} \\
&\leq C\tau^{\frac{5}{2}} \left(\mathbb{E} \left[\sup_{t_n \leq \sigma \leq t_{n+1}} |F(Y_\sigma)|^2 \right] \cdot \mathbb{E} \left[|F'(\bar{Y}_{t_n})|^2 \right] \right)^{\frac{1}{2}} \\
&\quad + C\tau^{\frac{1}{2}} \left(\mathbb{E} \left[\int_{t_n}^r |F'(\xi_n)|^2 |Y_\sigma - Y_{t_n}|^2 d\sigma \right] \cdot \mathbb{E} \left[|F'(\bar{Y}_{t_n})|^2 \right] \right)^{\frac{1}{2}} + C\tau^2 \left(\mathbb{E} \left[|F'(\bar{Y}_{t_n})|^2 \right] \right)^{\frac{1}{2}} \\
&\leq C\tau^{\frac{5}{2}} + C\tau^2,
\end{aligned}$$

where $\xi_n = Y_{t_n} + \xi(Y_\sigma - Y_{t_n})$, $\xi \in (0, 1)$.

For H_3 , the Cauchy-Schwarz inequality implies

$$\begin{aligned}
H_3 &\leq \mathbb{E} [|e(r)|^2] + \mathbb{E} [|F'(\bar{Y}_{t_n})(1 - e^{-\lambda(s-t_n)})b(1-\theta)\bar{Y}_{t_n}|^2] \\
&\leq \mathbb{E} [|e(r)|^2] + C\tau^2 \left(\mathbb{E} [|F'(\bar{Y}_{t_n})|^4] \right)^{\frac{1}{2}} \left(\mathbb{E} [|\bar{Y}_{t_n}|^4] \right)^{\frac{1}{2}} \\
&\leq \mathbb{E} [|e(r)|^2] + C\tau^2.
\end{aligned}$$

For H_4 , using the the Cauchy-Schwarz inequality and (44), we have

$$\begin{aligned} H_4 &\leq \mathbb{E}[|e(r)|^2] + \mathbb{E}[|F''(\bar{Y}_{t_n} + \sigma(\bar{Y}_r - \bar{Y}_{t_n}))|\bar{Y}_r - \bar{Y}_{t_n}|^2] \\ &\leq \mathbb{E}[|e(r)|^2] + C \left(\mathbb{E}[|F''(\bar{Y}_{t_n} + \sigma(\bar{Y}_r - \bar{Y}_{t_n}))|^4] \right)^{\frac{1}{2}} \left(\mathbb{E}[|\bar{Y}_r - \bar{Y}_{t_n}|^8] \right)^{\frac{1}{2}} \\ &\leq \mathbb{E}[|e(r)|^2] + C\tau^2. \end{aligned}$$

Note that,

$$J_{21} = \int_{t_n}^t \mathbb{E}[\langle e(r), F(\bar{Y}(r)) - F(\bar{Y}_{t_n}) \rangle] dr = \int_{t_n}^t H_1 + H_2 + H_3 + H_4 dr \leq 3 \int_{t_n}^t \mathbb{E}[|e(r)|^2] dr + C\tau^3.$$

Therefore, from above discuss, we have

$$J_2 \leq 4 \int_{t_n}^t \mathbb{E}[|e(r)|^2] dr + C\tau^3.$$

Substituting J_1 and J_2 into equation (43), we can get

$$\mathbb{E}[|e(t)|^2] \leq \mathbb{E}[|e(t_n)|^2] + C \int_{t_n}^t \mathbb{E}[|e(r)|^2] dr + C\tau^3.$$

Thus, for any $t \in [0, T]$ we have that

$$\mathbb{E}[|e(t)|^2] \leq C \int_0^t \mathbb{E}[|e(r)|^2] dr + C\tau^2.$$

Use Gronwall's inequality, we get

$$\mathbb{E}[|e(t)|^2] \leq C\tau^2.$$

This completes the proof. □

5 Probability of positivity

We now examine, following closely the analysis in [14], the probability of solutions of (25) becoming negative after a single step with this strategy, and hence triggering a use of the backstop method. This probability is given by

$$\mathbb{P}[Y_{k+1} < 0 \mid Y_k = y > 0] = \Phi(a(y)),$$

where

$$\Phi(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-s^2/2} ds \quad \text{and} \quad a(y) := \frac{y + F^{(\tau)}(y)}{b(\theta - 1)\sqrt{\tau}}.$$

Theorem 2 *Let $\{Y_n\}_{n=0}^N$ be a solution of (25), with initial value $Y_0 > 0$. Suppose also that $Y_0 \leq M_1$. Then, for each $\varepsilon \in (0, 1)$ there exists $\tau > 0$ such that*

$$\mathbb{P}[\mathcal{R}_N] > 1 - \varepsilon$$

where $\mathcal{R}_N := \bigcap_{j=0}^N \{Y_j > 0\}$.

Proof of Theorem 2 Since, by (25), the backstop method will ensure positivity over a single step τ , the event $\{Y_{n+1} > 0\}$ is equivalent to the following:

$$\left\{ \frac{\Delta W_n}{\sqrt{\tau}} < \frac{1}{b(\theta-1)} \left(\frac{Y_n}{\sqrt{\tau}} + \frac{F(Y_n)\sqrt{\tau}}{1 + \tau|F(Y_n)|^2} \right) \right\}.$$

From Lemmas 4, 5, we can easy to check that there exist constants M_1, M_2 such that

$$M_1 \leq \sup_{n \in \mathbb{N}} |Y_n| \leq M_2, \quad a.s.$$

Thus, we have

$$\begin{aligned} F^{(\tau)}\sqrt{\tau} &= \frac{F(Y_n)\sqrt{\tau}}{1 + \tau|F(Y_n)|^2} \\ &= \frac{(\theta-1)\sqrt{\tau}[a_2 Y_n^{\frac{\gamma-\theta}{1-\theta}} - a_{-1} Y_n^{\frac{\theta+1}{\theta-1}} + a_0 Y_n^{\frac{\theta}{\theta-1}} + \frac{1}{2}b^2\theta Y_n^{-1}]}{1 + \tau(\theta-1)^2 \left| a_2 Y_n^{\frac{\gamma-\theta}{1-\theta}} - a_{-1} Y_n^{\frac{\theta+1}{\theta-1}} + a_0 Y_n^{\frac{\theta}{\theta-1}} + \frac{1}{2}b^2\theta Y_n^{-1} \right|^2} \\ &\geq \frac{(\theta-1)\sqrt{\tau}[a_2 M_2^{\frac{\gamma-\theta}{1-\theta}} - a_{-1} M_2^{\frac{\theta+1}{\theta-1}} + a_0 M_1^{\frac{\theta}{\theta-1}} + \frac{1}{2}b^2\theta M_2^{-1}]}{1 + 4\tau(\theta-1)^2 \left(a_2^2 M_1^{\frac{2\gamma-2\theta}{1-\theta}} + a_{-1}^2 M_2^{\frac{2\theta+2}{\theta-1}} + a_0^2 M_2^{\frac{2\theta}{\theta-1}} + \frac{1}{4}b^4\theta M_1^{-2} \right)}. \end{aligned}$$

To prove $Y_{n+1} > 0$, it is sufficient to prove

$$\frac{\Delta W_n}{\sqrt{\tau}} < \frac{1}{b(\theta-1)} \left(\frac{Y_n}{\sqrt{\tau}} + \frac{(\theta-1)\sqrt{\tau}[a_2 M_2^{\frac{\gamma-\theta}{1-\theta}} - a_{-1} M_2^{\frac{\theta+1}{\theta-1}} + a_0 M_1^{\frac{\theta}{\theta-1}} + \frac{1}{2}b^2\theta M_2^{-1}]}{1 + 4\tau(\theta-1)^2 \left(a_2^2 M_1^{\frac{2\gamma-2\theta}{1-\theta}} + a_{-1}^2 M_2^{\frac{2\theta+2}{\theta-1}} + a_0^2 M_2^{\frac{2\theta}{\theta-1}} + \frac{1}{4}b^4\theta M_1^{-2} \right)} \right).$$

Let Φ denote the distribution function of a standard Normal random variable, and suppose $\{\xi_n\}_{n \in \mathbb{N}}$ is a sequence of mutually independent standard normal random variables. Then

$$\begin{aligned} &\mathbb{P}[Y_{n+1} > 0 \mid \mathcal{R}_n] \\ &= \frac{\mathbb{P}\{\{Y_{n+1} > 0\} \cap \{\mathcal{R}_n\}\}}{\mathbb{P}\{\mathcal{R}_n\}} = \frac{\mathbb{E}\left[\mathbb{E}\left[I_{\{Y_{n+1} > 0\} \cap \{\mathcal{R}_n\}} \mid \mathcal{F}_{t_n}\right]\right]}{\mathbb{P}\{\mathcal{R}_n\}} \\ &\geq \mathbb{E}\left[\mathbb{E}\left[I_{\{Y_{n+1} > 0\} \cap \{\mathcal{R}_n\}} \mid \mathcal{F}_{t_n}\right]\right] = \mathbb{E}\left[\mathbb{P}\{\{Y_{n+1} > 0\} \cap \{\mathcal{R}_n\} \mid \mathcal{F}_{t_n}\}\right] \\ &= \mathbb{E}\left[\mathbb{P}\left[\frac{\Delta W_n}{\sqrt{\tau}} < \frac{1}{b(\theta-1)} \left(\frac{Y_n}{\sqrt{\tau}} + \frac{(\theta-1)\sqrt{\tau}[a_2 M_2^{\frac{\gamma-\theta}{1-\theta}} - a_{-1} M_2^{\frac{\theta+1}{\theta-1}} + a_0 M_1^{\frac{\theta}{\theta-1}} + \frac{1}{2}b^2\theta M_2^{-1}]}{1 + 4\tau(\theta-1)^2 \left(a_2^2 M_1^{\frac{2\gamma-2\theta}{1-\theta}} + a_{-1}^2 M_2^{\frac{2\theta+2}{\theta-1}} + a_0^2 M_2^{\frac{2\theta}{\theta-1}} + \frac{1}{4}b^4\theta M_1^{-2} \right)} \right) \mid \mathcal{F}_{t_n}\right]\right] \\ &= \mathbb{P}\left[\xi_{n+1} < \frac{1}{b(\theta-1)} \left(\frac{Y_n}{\sqrt{\tau}} + \frac{(\theta-1)\sqrt{\tau}[a_2 M_2^{\frac{\gamma-\theta}{1-\theta}} - a_{-1} M_2^{\frac{\theta+1}{\theta-1}} + a_0 M_1^{\frac{\theta}{\theta-1}} + \frac{1}{2}b^2\theta M_2^{-1}]}{1 + 4\tau(\theta-1)^2 \left(a_2^2 M_1^{\frac{2\gamma-2\theta}{1-\theta}} + a_{-1}^2 M_2^{\frac{2\theta+2}{\theta-1}} + a_0^2 M_2^{\frac{2\theta}{\theta-1}} + \frac{1}{4}b^4\theta M_1^{-2} \right)} \right)\right] \\ &= \Phi\left(\frac{1}{b(\theta-1)} \left(\frac{Y_n}{\sqrt{\tau}} + \frac{(\theta-1)\sqrt{\tau}[a_2 M_2^{\frac{\gamma-\theta}{1-\theta}} - a_{-1} M_2^{\frac{\theta+1}{\theta-1}} + a_0 M_1^{\frac{\theta}{\theta-1}} + \frac{1}{2}b^2\theta M_2^{-1}]}{1 + 4\tau(\theta-1)^2 \left(a_2^2 M_1^{\frac{2\gamma-2\theta}{1-\theta}} + a_{-1}^2 M_2^{\frac{2\theta+2}{\theta-1}} + a_0^2 M_2^{\frac{2\theta}{\theta-1}} + \frac{1}{4}b^4\theta M_1^{-2} \right)} \right)\right). \end{aligned}$$

Since $Y_0 > 0$ and Φ takes values on $[0, 1]$, we have

$$\begin{aligned} \mathbb{P}\{\mathcal{R}_N\} &= \mathbb{P}\left[\bigcap_{n=0}^N \mathcal{R}_n\right] \\ &= \prod_{n=1}^N \mathbb{P}\{\mathcal{R}_n \mid \mathcal{R}_{n-1}, \dots, \mathcal{R}_0\} = \prod_{n=0}^{N-1} \mathbb{P}\{Y_{n+1} > 0 \mid \mathcal{R}_n\} \\ &\geq \Phi\left(\frac{1}{b(\theta-1)} \left(\frac{Y_n}{\sqrt{\tau}} + \frac{(\theta-1)\sqrt{\tau}[a_2 M_2^{\frac{\gamma-\theta}{1-\theta}} - a_{-1} M_2^{\frac{\theta+1}{\theta-1}} + a_0 M_1^{\frac{\theta}{\theta-1}} + \frac{1}{2}b^2\theta M_2^{-1}]}{1 + 4\tau(\theta-1)^2 \left(a_2^2 M_1^{\frac{2\gamma-2\theta}{1-\theta}} + a_{-1}^2 M_2^{\frac{2\theta+2}{\theta-1}} + a_0^2 M_2^{\frac{2\theta}{\theta-1}} + \frac{1}{4}b^4\theta M_1^{-2} \right)} \right)\right)^N. \end{aligned}$$

For convenience, we denote

$$\mathbf{Y}_\tau := \frac{Y_n}{\sqrt{\tau}} + \frac{(\theta - 1)\sqrt{\tau} [a_2 M_2^{\frac{\gamma-\theta}{1-\theta}} - a_{-1} M_2^{\frac{\theta+1}{\theta-1}} + a_0 M_1^{\frac{\theta}{\theta-1}} + \frac{1}{2} b^2 \theta M_2^{-1}]}{1 + 4\tau(\theta - 1)^2 (a_2^2 M_1^{\frac{2\gamma-2\theta}{1-\theta}} + a_{-1}^2 M_2^{\frac{2\theta+2}{\theta-1}} + a_0^2 M_2^{\frac{2\theta}{\theta-1}} + \frac{1}{4} b^4 \theta M_1^{-2})}.$$

Fix $\varepsilon \in (0, 1)$, then for fix step size τ , we have

$$\Phi\left(\frac{\mathbf{Y}_\tau}{b(\theta - 1)}\right)^N \geq 1 - \varepsilon. \quad (45)$$

To (45), apply the following inequality due to [20]

$$\frac{1}{\sqrt{2\pi}} \int_{-x}^x e^{-s^2/2} ds > \sqrt{1 - e^{-x^2/2}}, \quad x \in \mathbb{R}^+$$

along with the fact that $N = T/\tau$, leading us to seek τ so that

$$\left(\frac{1}{2} + \frac{1}{2} \sqrt{1 - \exp\left(-\frac{\mathbf{Y}_\tau}{2b(\theta - 1)^2}\right)}\right)^{\frac{T}{\tau}} \geq 1 - \varepsilon.$$

Thus we derive the bound

$$\tau(\varepsilon) := \sup \left\{ \tau \in (0, 1) : \mathbf{Y}_\tau \geq \sqrt{\ln \left(1 - \left(2(1 - \varepsilon)^{\frac{h}{\rho\tau}} - 1 \right)^2 \right)^{-2b(\theta-1)^2}} \right\}.$$

$\tau(\varepsilon)$ is uniquely defined for each $\varepsilon \in (0, 1)$ because

$$g(\tau) := \mathbf{Y}_\tau - \sqrt{-2b(\theta - 1)^2 \ln \left(1 - \left(2(1 - \varepsilon)^{\tau/T} - 1 \right)^2 \right)}$$

is continuous on \mathbb{R}^+ with $\lim_{\tau \rightarrow 0^+} g(\tau) = \infty$, and therefore there is a neighbourhood of zero corresponding to $(0, \tau(\varepsilon))$ within which g is positive.

6 Numerical simulations

We provide some numerical experiments to support the strong convergence proved in Theorem 1. For convenience we recall here the original Ait-Sahalia-type rate model given in (1)

$$dX_t = (a_{-1}X_t^{-1} - a_0 + a_1X_t - a_2X_t^\gamma) dt + bX_t^\theta dW_t, \quad t \in (0, T], \quad X_0 > 0$$

and the transformed SDE given in (2)

$$\begin{aligned} dY_t &= f(Y_t)dt + b(1 - \theta)dW_t, \quad t \in (0, T], \\ f(x) &:= (\theta - 1)(a_2x^{\frac{\gamma-\theta}{1-\theta}} - a_1x - a_{-1}x^{\frac{\theta+1}{\theta-1}} + a_0x^{\frac{\theta}{\theta-1}} + \frac{1}{2}b^2\theta x^{-1}). \end{aligned}$$

We are interested in examining the performance of the splitting method (21) (denoted **Splitting**) and the tamed-splitting method (22) (denoted **TSM**). To construct a reference solution for our numerical experiments we apply a tamed Milstein method [] to (1) with a small step size $\tau = 2^{-18}$. We compare our methods to a backward Euler-Maruyama method applied to (1), denoted **RefBEM**. This was shown to have rate one half [23]. We also compare to other approximations of (2) : a tamed Euler Maruyama method [13] (denoted **TEM**) and backward Euler Maruyama method (denoted **BEM**). In our experiments we take timesteps $\tau \in \{2^{-11}, 2^{-10}, 2^{-9}, 2^{-8}, 2^{-7}\}$ and fix $X_0 = 1$. For our convergence plots we take 1000 realizations. In none of these realizations did **TSM** give a negative value, that is at no point did we need to make use the backstop method **BEM** in (24). Indeed the same was true when taking 10000 realizations - illustrating the positivity Theorem 2.

We present simulations based on two sample sets that were originally derived in [1] from financial data. We fix $a_{-1} = 0.00107, a_0 = 0.0517, a_1 = 0.877, a_2 = 4.604, \gamma = 3$ and $b = 1$ and then take




Fig. 1: Non-critical case. Convergence rates (left) and efficiency (right) of the numerical methods.

- Non-critical case ($\gamma + 1 > 2\theta$) : $\theta = 1.5$.
- Critical case ($\gamma + 1 = 2\theta$) : $\theta = 2$.

In Figure 1 (left) we illustrate the convergence of our methods and the rates are given in Table 1. We see that, as expected, RefBEM has rate one half and the other methods have rate 1. In this non-critical case we see that TEM in fact has the best error constant. However, if we examine the efficiency (see Figure 1 (right)) we see that Splitting is the most efficient of the methods (and the two backward Euler Maruyama methods are the least efficient). For the critical case we draw similar conclusions. In Figure 2 (left) and Table 1 we illustrate convergence in this case. In this case it is surprising that the error constant for TSM seems to increase by an order of magnitude. Examining the relative efficiency we again see that Splitting is the most efficient by approximately two order of magnitude. We illustrate the probability of numerical solutions take negative value. It turns out that the probability of the numerical solutions being negative value is 0. This means that our numerical method (TSM) is almost positive. There is no need for backstop methods in numerical simulation. Our numerical method (TSM) is only possible to take negative values in theoretical analysis.

The tamed-splitting method (TSM) we analyse in this paper performs well in terms of the error constant but is slightly less efficient. However it is simpler to implement and can handle more complex situations as for general drift terms F we may not exact solution of each equation by splitting.

Parameter Sets	RefBEM	BEM	TEM	TSM	Splitting
Non-critical case	0.4909	0.9855	0.9795	0.9928	0.9805
Critical case	0.5084	0.9880	1.0036	1.0084	0.9852

Table 1: Rates of convergence estimated for the different methods, see Figs. 1 and 2.

References

1. Y. AIT-SALAHIA, *Testing continuous-time models of the spot interest rate*, The review of financial studies, 9 (1996), pp. 385–426.
2. A. ALFONSI, *On the discretization schemes for the CIR (and Bessel squared) processes.*, Monte Carlo Methods & Applications, 11 (2005).




Fig. 2: Critical case. Convergence rates (left) and efficiency (right) of the numerical methods.

3. A. BERKAOUI, M. BOSSY, AND A. DIOP, *Euler scheme for SDEs with non-Lipschitz diffusion coefficient: strong convergence*, ESAIM: Probability and Statistics, 12 (2008), pp. 1–11.
4. M. BOSSY AND A. DIOP, *An efficient discretisation scheme for one dimensional SDEs with a diffusion coefficient function of the form*, PhD thesis, INRIA, 2007.
5. T. G. CONLEY, L. P. HANSEN, E. G. LUTTMER, AND J. A. SCHEINKMAN, *Short-term interest rates as subordinated diffusions*, The Review of Financial Studies, 10 (1997), pp. 525–577.
6. S. DENG, C. FEI, W. FEI, AND X. MAO, *Generalized Ait-Sahalia-type interest rate model with Poisson jumps and convergence of the numerical approximation*, Physica A., 533 (2019), p. 122057.
7. C. EMMANUEL AND X. MAO, *Truncated EM numerical method for generalised Ait-Sahalia-type interest rate model with delay*, Journal of Computational and Applied Mathematics, 383 (2021), p. 113137.
8. A. R. GALLANT AND G. TAUCHEN, *Estimation of continuous-time models for stock returns and interest rates*, Macroeconomic Dynamics, 1 (1997), pp. 135–168.
9. A. GARDÓN, *The order of approximations for solutions of Itô-type stochastic differential equations with jumps*, Stoch Anal. Appl., 22 (2004), pp. 679–699.
10. M. B. GILES, *Multilevel Monte Carlo path simulation*, Operations research, 56 (2008), pp. 607–617.
11. D. J. HIGHAM AND X. MAO, *Convergence of Monte Carlo simulations involving the mean-reverting square root process*, Journal of Computational Finance, 8 (2005), pp. 35–61.
12. Y. HONG AND H. LI, *Nonparametric specification testing for continuous-time models with applications to term structure of interest rates*, The Review of Financial Studies, 18 (2005), pp. 37–84.
13. M. HUTZENTHALER, A. JENTZEN, AND P. E. KLOEDEN, *Strong convergence of an explicit numerical method for SDEs with nonglobally Lipschitz continuous coefficients*, Ann. Appl. Probab., 22 (2012), pp. 1611–1641.
14. C. KELLY, G. LORD, AND H. MAULANA, *The role of adaptivity in a numerical method for the Cox-Ingersoll-Ross model*, Journal of Computational and Applied Mathematics, 410 (2022), p. 114208.
15. P. KLOEDEN AND E. PLATEN, *Numerical Solution of Stochastic Differential Equations*, Stochastic Modelling and Applied Probability, Springer Berlin Heidelberg, 2011.
16. C. KUMAR AND S. SABANIS, *On milstein approximations with varying coefficients: the case of super-linear diffusion coefficients*, BIT Numerical Mathematics, 59 (2019), pp. 929–968.
17. Z. LEI, S. GAN, AND J. LIU, *First order strong approximation of Ait-Sahalia-type interest rate model with Poisson jumps*, arXiv preprint arXiv:2110.15482, (2021).
18. G. J. LORD, C. E. POWELL, AND T. SHARDLOW, *An introduction to computational stochastic PDEs*, vol. 50, Cambridge University Press, 2014.
19. X. MAO, *Stochastic Differential Equations and Applications*, Horwood, 2008.
20. Z. SASVARI AND H. CHEN, *Tight bounds for the normal distribution: 10611*, The American Mathematical Monthly, 106 (1999), pp. 76–76.

-
21. L. SZPRUCH, X. MAO, D. J. HIGHAM, AND J. PAN, *Numerical simulation of a strongly nonlinear Ait-Sahalia-type interest rate model*, BIT Numerical Mathematics, 51 (2011), pp. 405–425.
 22. Z. ZHANG AND G. E. KARNIADAKIS, *Numerical methods for stochastic partial differential equations with white noise*, Springer, 2017.
 23. Y. ZHAO, X. WANG, AND M. WANG, *On the backward Euler method for a generalized Ait-Sahalia-type rate model with Poisson jumps*, Numerical Algorithms, (2020), pp. 1–21.