

A Unified Quantum Algorithm Framework for Estimating Properties of Discrete Probability Distributions

Tongyang Li^{1,2} Xinzhao Wang^{1,2} Shengyu Zhang³

¹Center on Frontiers of Computing Studies, Peking University

²School of Computer Science, Peking University

³Tencent Quantum Laboratory

Abstract

Estimating statistical properties is fundamental in statistics and computer science. In this paper, we propose a unified quantum algorithm framework for estimating properties of discrete probability distributions, with estimating Rényi entropies as specific examples. In particular, given a quantum oracle that prepares an n -dimensional quantum state $\sum_{i=1}^n \sqrt{p_i} |i\rangle$, for $\alpha > 1$ and $0 < \alpha < 1$, our algorithm framework estimates α -Rényi entropy $H_\alpha(p)$ to within additive error ϵ with probability at least $2/3$ using $\tilde{O}(n^{1-\frac{1}{2\alpha}}/\epsilon + \sqrt{n}/\epsilon^{1+\frac{1}{2\alpha}})$ and $\tilde{O}(n^{\frac{1}{2\alpha}}/\epsilon^{1+\frac{1}{2\alpha}})$ queries, respectively. This improves the best known dependence in ϵ as well as the joint dependence between n and $1/\epsilon$. Technically, our quantum algorithms combine quantum singular value transformation, quantum annealing, and variable-time amplitude estimation. We believe that our algorithm framework is of general interest and has wide applications.

1 Introduction

Motivations. For many problems, quantum algorithms can dramatically outperform their classical counterparts. Among those, an important category is quantum algorithms for linear algebraic problems. Recently, Gilyén, Low, Su, and Wiebe [GSLW19] proposed a powerful framework for quantum matrix arithmetics, namely *quantum singular value transformation (QSVT)*. QSVT encompasses quantum algorithms for various problems (see also [MRTC21]), and can recover the best-known or even optimal quantum algorithms for fixed-point amplitude amplification [Gro05b, TGP06, AC12, YLC14], solving linear systems [HHL09, Amb10, CKS17, CGJ19], Hamiltonian simulation [LC17, LC19], etc.

In this paper, we study a fundamental problem in statistics, theoretical computer science, and machine learning: *estimating statistical properties*, which aims to estimate properties of probability distributions using the least number of independent samples. On the one hand, statistical properties such as entropies, divergences, etc., characterize some key measures of randomness. On the other hand, relevant theoretical tools are rapidly developing in topics such as property testing [Ron10], statistical learning [Val11], etc. Among statistical properties, the most basic one is the Shannon entropy [Sha48]. For a discrete distribution $\mathbf{p} = (p_i)_{i=1}^n$ supported on $[n]$, it is defined as

$$H(\mathbf{p}) := - \sum_{i=1}^n p_i \log p_i. \quad (1)$$

A natural generalization of the Shannon entropy is the family of Rényi entropies [Rén61]. Specifically, the α -Rényi entropy is defined as

$$H_\alpha(\mathbf{p}) := \frac{1}{1-\alpha} \log \sum_{i=1}^n p_i^\alpha. \quad (2)$$

For our convenience, the power sum in the logarithm is denoted by $P_\alpha(\mathbf{p})$, i.e., $P_\alpha(\mathbf{p}) := \sum_{i=1}^n p_i^\alpha$. When $\alpha \rightarrow 1$, $\lim_{\alpha \rightarrow 1} H_\alpha(\mathbf{p}) = H(\mathbf{p})$. Classically, references [JVHW15, WY16] proved the tight classical sample complexity bound

$$\Theta \left(\frac{\log^2 n}{\epsilon^2} + \frac{n}{\epsilon \log n} \right) \quad (3)$$

for estimating Shannon entropy within precision ϵ with success probability at least $2/3$. For α -Rényi entropy estimation, reference [AOST16] proved that when $\alpha > 1$ and $0 < \alpha < 1$ respectively, it takes $\mathcal{O}(n/\log n)$ and $\mathcal{O}(n^{1/\alpha}/\log n)$ independent samples from \mathbf{p} respectively to estimate $H_\alpha(\mathbf{p})$ within constant additive error with probability at least $2/3$. In addition, for any constant $\eta > 0$, the paper also established sample complexity lower bounds $\Omega(n^{1-\eta})$ and $\Omega(n^{1/\alpha-\eta})$ when $\alpha > 1$ and $0 < \alpha < 1$, respectively.

There has also been literature on quantum algorithms for entropy estimation (see the paragraph on related works for more details). Among those, the state-of-the-art result on estimating Shannon entropy was given by Gilyén and Li [GL20], which applies QSVT to estimate the Shannon entropy within additive error ϵ with success probability at least $2/3$ using $\tilde{\mathcal{O}}(\sqrt{n}/\epsilon^{1.5})$ quantum queries. For α -Rényi entropy, Li and Wu [LW19] gave algorithms with quantum query complexities $\tilde{\mathcal{O}}(n^{1-1/2\alpha}/\epsilon^2)$ and $\tilde{\mathcal{O}}(n^{1/\alpha-1/2}/\epsilon^2)$ when $\alpha > 1$ and $0 < \alpha < 1$, respectively. Both papers used a common model proposed by Bravyi et al. [BHH11] which encodes \mathbf{p} as frequencies of n symbols in a given input string and quantum algorithms can access the input string in superposition (Definition 2), whereas [GL20] also adopted oracles preparing a superposed quantum state whose amplitude in the i^{th} term is p_i (Definition 1 and Definition 3).

Nevertheless, it can be observed that although quantum algorithms for learning statistical properties have applied advanced algorithmic tools including quantum singular value transformation [GSLW19], and have achieved speedup in the cardinality n and precision ϵ separately, the combined dependence on n and ϵ is not yet as well understood as the classical counterparts, for instance the sample complexity of Shannon entropy in Eq. (3). From a high-level perspective, even though quantum algorithms for linear algebraic problems have been systematically developed, we shall still endeavor to quantum algorithms with optimal or near-optimal dependence on *all parameters*. In this paper, we shed light on this question for estimating statistical properties.

Contributions. In this paper, we introduce a unified quantum algorithm framework for estimating properties of discrete distributions. Our algorithm is stemmed from quantum singular value transformation [GSLW19], but we enhance the framework with quantum annealing and variable-time amplitude amplification and estimation. Specifically, we propose algorithms for estimating Rényi entropies of discrete probability distributions with refined dependence on n and ϵ , assuming access to quantum oracle U_p which maps $|0\rangle$ to $\sum_{i=1}^n \sqrt{p_i}|i\rangle$ (see the later “related work” paragraph for more discussions and comparisons of different oracles).¹

Theorem 1 (Main theorem). *There are quantum algorithms that approximate the Rényi entropy $H_\alpha(\mathbf{p})$ in Eq. (2) within an additive error $\epsilon > 0$ with success probability at least $2/3$ using*

- $\tilde{\mathcal{O}}\left(\frac{n^{1-\frac{1}{2\alpha}}}{\epsilon} + \frac{\sqrt{n}}{\epsilon^{1+\frac{1}{2\alpha}}}\right)$ quantum queries to U_p and U_p^\dagger when $\alpha > 1$ (Theorem 7), and
- $\tilde{\mathcal{O}}\left(\frac{n^{\frac{1}{2\alpha}}}{\epsilon^{\frac{1}{2\alpha}+1}}\right)$ quantum queries to U_p and U_p^\dagger when $0 < \alpha < 1$ (Theorem 8).

Compared to the state-of-the-art result for estimating Rényi entropies by Li and Wu [LW19] which uses $\tilde{\mathcal{O}}(n^{1/\alpha-1/2}/\epsilon^2)$ quantum queries when $0 < \alpha < 1$ and $\tilde{\mathcal{O}}(n^{1-1/2\alpha}/\epsilon^2)$ quantum queries when $\alpha > 1$ and α is not an integer, our result achieves a systematic improvement in both n and ϵ . This can be illustrated by Figure 1 below.²




Figure 1: Comparison between our algorithms and the algorithm in Li and Wu [LW19].

Our quantum algorithms can also be applied to entropy estimation of quantum density matrices:

¹In fact, our quantum algorithm also applies to the purified quantum query-access in Definition 3. Please see Section 3 and Section 6 for more details.

²The integral α cases are excluded in the figure because computing $H_\alpha(p)$ for integral α seems fundamentally easier. Classically, the best-known upper bound for integral $\alpha > 2$ is $\Theta(n^{1-\frac{1}{\alpha}})$, smaller than that of $\mathcal{O}(n/\log(n))$ for non-integral cases [AOST16]. For quantum algorithms, Li and Wu [LW19] made special designs for integer α cases, with query cost better than their non-integral α cases (and also ours), albeit using a stronger input oracle (Definition 2).

Corollary 1. *There are quantum algorithms that approximate the Rényi entropy of density operator $H_\alpha(\rho)$ in Eq. (110) within an additive error $\epsilon > 0$ with success probability at least $2/3$ using*

- $\tilde{\mathcal{O}}\left(\frac{n^{\frac{3}{2}-\frac{1}{2\alpha}}}{\epsilon} + \frac{n}{\epsilon^{1+\frac{1}{2\alpha}}}\right)$ quantum queries to U_ρ and U_ρ^\dagger when $\alpha > 1$ (*Corollary 2*), and
- $\tilde{\mathcal{O}}\left(\frac{n^{\frac{1}{2\alpha}+\frac{1}{2}}}{\epsilon^{\frac{1}{2\alpha}+1}}\right)$ quantum queries to U_ρ and U_ρ^\dagger when $0 < \alpha < 1$ (*Corollary 3*).

In addition, if a classical probability distribution \mathbf{p} has at most r elements i such that $p_i > 0$ and we know the value of r in advance, we can construct an algorithm using $\tilde{\mathcal{O}}\left(\frac{r^{1-\frac{1}{2\alpha}}}{\epsilon} + \frac{\sqrt{r}}{\epsilon^{1+\frac{1}{2\alpha}}}\right)$ calls to U_p and U_p^\dagger to estimate $H_\alpha(\mathbf{p})$ to within additive error ϵ when $\alpha > 1$, and an algorithm using $\tilde{\mathcal{O}}\left(\frac{r^{\frac{1}{2\alpha}}}{\epsilon^{\frac{1}{2\alpha}+1}}\right)$ calls to U_p and U_p^\dagger to estimate $H_\alpha(\mathbf{p})$ to within additive error ϵ when $0 < \alpha < 1$. In addition, we also give quantum algorithms when we do not know the value of r , and our quantum algorithms can also be extended to estimating the entropies of low-rank quantum density matrices and the quantum Rényi divergence of density matrices. Please find details in [Section 6](#).

Techniques. Our quantum algorithm follows the one in Gilyén and Li [[GL20](#)] for Shannon entropy estimation. In [Section 3.1](#), we first construct a unitary operator, which has a matrix block encoding of the square root of the probability distribution we want to study. We then use quantum singular value transformation [[GSLW19](#)] to compute a polynomial approximation of the function that we want to estimate, which is then encoded into an amplitude. Finally, we apply amplitude estimation to obtain the estimate as a classical output.

However, this algorithm is sub-optimal in many cases because of the following two reasons.

- If we need an estimate to within a given multiplicative error, the query complexity of the amplitude estimation process is inversely proportional to the square root of the quantity that we want to estimate, so it has poor dependence if the quantity is too small.
- Quantum singular value transformation leverages the computation of the same function of all singular values in parallel, but this brings restrictions that the polynomial should well-approximate the function within the entire interval $[0, 1]$, where the singular values may range over.

For the first issue, we design an annealing process in [Section 3.2](#) to obtain an estimate of the quantity to within constant multiplicative error in advance. With this rough estimate, we can amplify the quantity with smaller overhead in each step. For the second issue, we exploit variable time amplitude estimation in [Section 3.3](#) by designing a variable-stopping-time algorithm which applies different transformation polynomials to singular values in different intervals. This give us more flexibility to construct polynomials for different singular values and the final complexity is related to the average degree of all polynomials.

As a technical contribution, we also improve the bounded polynomial approximation of x^α for $\alpha > 1$ ([Lemma 6](#)), which may be of independent interest. Our approximation polynomial is bounded by $2x^\alpha$ when x is smaller than a threshold while the bounded polynomial approximation constructed in [[GSLW19](#)] is only guaranteed to be bounded by 1.

We summarize and compare the techniques in previous literature on quantum algorithms for estimating entropies of discrete probability distributions in [Table 1](#) below.

We give a brief explanation of the comparisons in the table here. Compared with quantum algorithms without using QSVT to estimate entropy, they first sample i according to p_i , then

Reference	Oracle	Quantum SVT	Annealing	VTAA
[BHH11]	Discrete query-access (Definition 2)	✗	✗	✗
[LW19]	Discrete query-access (Definition 2)	✗	✓	✗
[GL20]	Purified query-access (Definition 3)	✓	✗	✗
[GHS21]	Purified query-access (Definition 3)	✓	✗	✗
this paper	Pure-state preparation (Definition 1)	✓	✓	✓
	Purified query-access (Definition 3)			

Table 1: Summary of quantum algorithms for estimating entropies of classical discrete distributions.

estimate p_i using amplitude estimation and compute the entropy accordingly, while using QSVT we can directly compute any polynomial function value of p_i for all i simultaneously in the block encoding. On the other hand, VTAA is an accelerated version of amplitude amplification that takes problem instances into account. In our problem, VTAA allows us to apply QSVT with different polynomials to p_i in different intervals, which makes our quantum algorithm more flexible. Moreover, annealing is applied to handle the issue that the estimated quantity is too small and makes the amplitude amplification costly. By using annealing, we can obtain a rough estimate of the estimated quantity, which enlarges the estimated quantity when using QSVT.

Related work. Previous literature investigated quantum algorithms for estimating statistical properties using different input models (see also the survey paper [MdW16]). First, if we want to utilize quantum algorithms to accelerate the solving of problems related to classical distributions, we need coherent access to classical distributions via quantum oracle. It is thus natural to consider a unitary oracle which can prepare a pure state encoding a classical distribution as follows:

Definition 1 (Pure-state preparation access to classical distribution). *A classical distribution $\mathbf{p} = (p_i)_{i=1}^n$ is accessible via pure-state preparation access if we have access to a unitary oracle U_p and its inverse, which satisfies*

$$U_p|\mathbf{0}\rangle = \sum_{i=1}^n \sqrt{p_i}|i\rangle. \quad (4)$$

Another common model, originally proposed by Bravyi et al. [BHH11], encodes the classical probability distribution as frequencies of n symbols in a given input string, and quantum algorithms can access the input string in superposition.

Definition 2 (Discrete quantum query-access to classical distribution). *A classical distribution $\mathbf{p} = (p_i)_{i=1}^n$ is accessible via discrete quantum query-access if we have quantum access to a function $f: S \rightarrow [n]$ such that for all $i \in [n]$, $p_i = |\{s \in S \mid f(s) = i\}|/|S|$, which means we have access to a unitary oracle O and its inverse acting on $\mathbb{C}^{|S|} \otimes \mathbb{C}^n$ such that*

$$O|s\rangle|\mathbf{0}\rangle = |s\rangle|f(s)\rangle \text{ for all } s \in S. \quad (5)$$

In this model, Bravyi et al. [BHH11] gave a quantum algorithm to estimate the ℓ_1 -norm distance of two distributions p and q with support cardinality n and with constant precision using $\mathcal{O}(\sqrt{n})$ queries, and gave quantum algorithms for testing uniformity and orthogonality with query complexity $\mathcal{O}(n^{1/3})$. This was later generalized to identity testing, i.e., testing whether a distribution is

identical or ϵ -far in ℓ_1 -norm from a given distribution, in $\tilde{\mathcal{O}}(n^{1/3})$ queries by [CFMdW10]. Li and Wu [LW19] gave a quantum algorithm for estimating the Shannon entropy within additive error ϵ with high success probability using $\tilde{\mathcal{O}}(n^{1/2}/\epsilon^2)$ queries, and this paper also studied the query complexity of Rényi entropy estimation (see the paragraph of “contributions”). To complement the algorithm results, Bun et al. [BKT18] proved that Shannon entropy estimation with a certain constant ϵ requires $\tilde{\Omega}(\sqrt{n})$ quantum queries to the oracle in Eq. (5).

Beyond classical distributions, it is natural to extend to statistical problems of genuine quantum systems. The quantum counterpart of a classical discrete distribution is a *density matrix*. Density matrices can be regarded as the (possibly random) outcome of some physical process, and if we can access this physical process by calling it as a black box, we can generate quantum samples ourselves. If the physical process is reversible, which is common in a quantum scenario, we can also access the inverse process of it. For example, if a quantum computer produces the state ρ without measurements, we can easily reverse this process. We can define the following input model to characterize the situations mentioned above.

Definition 3 (Purified quantum query-access). *A density operators $\rho \in \mathbb{C}^{n \times n}$ has purified quantum query-access if we have access to a unitary oracle U_ρ and its inverse, which satisfies*

$$U_\rho |0\rangle_A |0\rangle_B = |\psi_\rho\rangle = \sum_{i=1}^n \sqrt{p_i} |\phi_i\rangle_A |\psi_i\rangle_B \quad (6)$$

such that $\text{Tr}_A(|\psi_\rho\rangle\langle\psi_\rho|) = \rho$, where $\langle\phi_i|\phi_j\rangle = \langle\psi_i|\psi_j\rangle = \delta_{ij}$. If $|\psi_i\rangle = |i\rangle$, $\rho = \sum_{i=1}^n p_i |i\rangle\langle i|$ is a diagonal density operator which can be seen as a classical distribution $\mathbf{p} = (p_i)_{i=1}^n$.

We note that for encoding classical distributions, Definition 3 is weaker than Definition 2 since we can apply O to a uniform superposition over S in Eq. (5), and this is equivalent to applying a purified quantum query-access encoding a classical distribution to $|0\rangle$. (Furthermore, Definition 2 essentially assumes that all probabilities p_i are rational, whereas Definition 3 does not have this requirement.) In addition, Definition 3 is also weaker than Definition 1 since we can use one query to $U_{\mathbf{p}}$ to prepare $\sum_{i=1}^n \sqrt{p_i} |i\rangle$, and then apply CNOT gates to produce the state $\sum_{i=1}^n \sqrt{p_i} |i\rangle |i\rangle$, which satisfies the condition in Definition 3. Our results are established with Definition 3 being the input oracle.

For classical distributions encoded by Definition 3, Gilyén and Li [GL20] systematically studied different oracle access of distributional property testing, and proved that it takes $\tilde{\mathcal{O}}(n^{1/2}/\epsilon^{1.5})$ queries to the purified query access for estimating Shannon entropy to within additive error ϵ with high success probability. This work also studied closeness testing, where we are given purified query access to distributions \mathbf{p} and \mathbf{q} and the goal is to distinguish between $\mathbf{p} = \mathbf{q}$ and $\|\mathbf{p} - \mathbf{q}\| \geq \epsilon$. For ℓ_1 -norm and ℓ_2 -norm distances, [GL20] proved that the quantum query complexities are $\tilde{\mathcal{O}}(\sqrt{n}/\epsilon)$ and $\tilde{\Theta}(1/\epsilon)$, respectively. Belovs [Bel19] proved that distinguishing between \mathbf{p} and \mathbf{q} takes $\Theta(1/d_H(\mathbf{p}, \mathbf{q}))$ queries (see also Section 5.2), where $d_H(\mathbf{p}, \mathbf{q})$ is the Hellinger distance between \mathbf{p} and \mathbf{q} , and this tight bound applies to all oracles in Definition 1, Definition 2, and Definition 3.

Definition 3 is also widely used among quantum algorithms for estimating properties of quantum density operators. The results in [GL20] about Shannon entropy estimation and ℓ_1 -norm and ℓ_2 -norm closeness testing can be generalized to those of quantum density matrices with purification with an overhead of \sqrt{n} . Gur et al. [GHS21] estimates the von Neumann entropy of quantum density matrices to within a certain multiplicative error, and under appropriate choices of parameters the query complexity to the purified query access can be sublinear in n . Regarding the estimation of quantum Rényi entropy in general, Subramanian and Hsieh [SH21] used $\tilde{\mathcal{O}}(\kappa n^{\max\{2\alpha, 2\}}/\epsilon^2)$ queries to estimate the α -Rényi entropy of a density matrix ρ satisfying $I/\kappa \preceq \rho \preceq I$ to within additive error

ϵ . When ρ has rank at most r , Wang et al. [WZC⁺22] gave quantum algorithms taking $\text{poly}(r, 1/\epsilon)$ queries for estimating von Neumann entropy, quantum Rényi entropy, and trace distance and fidelity between two density matrices. Fidelity estimation was also studied in [WZC⁺21, GP22].

Finally, since classical algorithms for estimating distribution properties takes independent samples, it is natural to consider quantum samples of density operators defined as follows.

Definition 4 (Quantum sampling). *A quantum distribution $\rho \in \mathbb{C}^{n \times n}$ is accessible via quantum sampling if we can request independent copies of the state ρ .*

A series of papers by O’Donnell and Wright [OW15, OW16, OW17a] (see also their survey paper [OW17b]) studied the sample complexity of various problems, including quantum state tomography, maximally mixedness testing, rankness testing, spectrum estimation, learning eigenvalues, learning top- k eigenvalues, and learning optimal rank- k approximation. Subsequently, Bădescu, O’Donnell, and Wright [BOW19] studied the sample complexity of testing whether ρ is equal to some known density matrix or ϵ -far from it, which is $O(n/\epsilon)$ with respect to fidelity and $O(n/\epsilon^2)$ with respect to trace distance; both results are optimal up to constant factors. Regarding von Neumann and quantum Rényi entropies, Acharya et al. [AISW19] proved that estimation with additive error ϵ of von Neumann entropy, quantum Rényi entropy with $\alpha > 1$, and quantum Rényi entropy with $0 < \alpha < 1$ have sample complexity bounds $\mathcal{O}(n^2/\epsilon^2)$ and $\Omega(n^2/\epsilon)$, $\mathcal{O}(n^2/\epsilon^2)$ and $\Omega(n^2/\epsilon)$, and $\mathcal{O}(n^{2/\alpha}/\epsilon^{2/\alpha})$ and $\Omega(n^{1+1/\alpha}/\epsilon^{1/\alpha})$, respectively. Given an additional assumption that all nonzero eigenvalues of ρ are at least $1/\kappa$, Wang et al. [WZW22] gave a quantum algorithm for estimating its von Neumann entropy using $\tilde{\mathcal{O}}(\kappa^2/\epsilon^5)$ samples, and bounds under the same assumption were also proved for estimating quantum Rényi entropy.

Open questions. Our work raises several natural questions for future investigation:

- When $\alpha > 1$, can we achieve quadratic quantum speedup in n compared to the classical algorithm in [AOST16] for estimating α -Rényi entropy with $\mathcal{O}(n/\log n)$ queries? A natural goal is to give a quantum algorithm with query complexity $\tilde{\mathcal{O}}(\sqrt{n})$ for constant ϵ , but our current bound in Theorem 7 has complexity $\tilde{\mathcal{O}}(n^{1-\frac{1}{2\alpha}})$. This may be related to our estimation paradigm. A classical analogy to our algorithm is to draw samples independently from the probability distribution \mathbf{p} on $[n]$, estimate $(p_i)^{\alpha-1}$ for each sample i , and output the mean value of all estimates. Such algorithms are called empirical estimators, but they can be sub-optimal classically.
- Can we apply our quantum algorithm framework to other statistical problems? One possibility is the estimation of partition functions – it is another prominent type of statistical properties, and many previous quantum algorithms including [WCNA09, Mon15, CCH⁺19, HW20, AHN⁺21] had applied annealing on the system’s temperature to estimating partition functions. It would be of general interest to achieve further quantum speedup by our algorithm framework.
- For other quantum linear algebraic problems, can we elaborate on the dependence on all parameters? Decent efforts had been conducted for Hamiltonian simulation [LC17, LC19, GSLW19] and linear system solving [HHL09, Amb10, CKS17, CGJ19], and this work investigates the estimation of statistical properties. It would be natural to leverage refined analyses for more problems, for instance the applications in quantum machine learning.

Organization. The rest of the paper is organized as follows. We review necessary background in Section 2. We introduce our main technical contribution, our quantum algorithm framework, in Section 3. We prove our results about the quantum query complexity of α -Rényi entropy estimation

with $\alpha > 1$ and $0 < \alpha < 1$ in [Section 4](#) and [Section 5](#), respectively. In [Section 6](#), we describe further applications of our quantum algorithm framework in estimating statistical properties.

Notation. Throughout the paper, $\tilde{\mathcal{O}}$ omits poly-logarithmic factors in the big- \mathcal{O} notation, i.e., $\tilde{\mathcal{O}}(g) = g \text{ poly}(\log g)$. Unless otherwise stated, all vector norms $\|\cdot\|$ in this paper are ℓ_2 -norm. We use \log to represent \log_2 and \ln to represent \log_e . We use Δ^n to represent the set of all probability distributions on $[n]$. For a set A , we use $|A|$ to represent the size of A . In description of quantum algorithms, the corresponding Hilbert space of a quantum register X is denoted by \mathcal{H}_X .

2 Preliminaries

We summarize necessary tools used in our quantum algorithm framework as follows.

2.1 Amplitude amplification and estimation

Fixed-point amplitude amplification. Classically, for a Bernoulli random variable X with $\mathbb{E}[X] = p$, we need $\Theta(1/p)$ i.i.d. samples in expectation to observe the first 1. In the quantum case, this can be improved by *amplitude amplification* [[BHMT02](#)], a quantum algorithm in which the number of iterations depends on p . This was later strengthened to a fixed-point version, where the algorithm only needs to know a lower bound of p . There are a number of implementations [[Høy00](#), [Gro05b](#), [TGP06](#), [AC12](#), [YLC14](#)], and here we use a version given in [[GSLW19](#)]. Let $|0\rangle$ denote the all-0 initial state.

Consider a unitary U such that

$$U|0\rangle|0\rangle = \sqrt{p}|1\rangle|\phi\rangle + \sqrt{1-p}|0\rangle|\psi\rangle. \quad (7)$$

The following theorem says that we can obtain an approximation of $|\phi\rangle$ using $\Theta(\frac{1}{\sqrt{p}})$ calls to U and U^\dagger , achieving a quadratic quantum speedup over its classical counterpart.

Theorem 2 (Fixed-point amplitude amplification [[GSLW19](#), Theorem 27]). *Let \mathcal{A} be a quantum algorithm on space $\mathcal{H}_A = \mathcal{H}_F \otimes \mathcal{H}_W$ such that*

$$\mathcal{A}|0\rangle_{\mathcal{H}_A} = \sqrt{p_{\text{succ}}}|1\rangle_{\mathcal{H}_F}|\phi\rangle_{\mathcal{H}_W} + \sqrt{1-p_{\text{succ}}}|0\rangle_{\mathcal{H}_F}|\psi\rangle_{\mathcal{H}_W}, \text{ where } \|\phi\| = 1. \quad (8)$$

For any $0 < \delta < 1, 0 < \epsilon < 1$, there is a quantum algorithm \mathcal{A}' using a single ancilla qubit and $\mathcal{O}(\frac{\log(1/\epsilon)}{\delta})$ calls to \mathcal{A} and \mathcal{A}^\dagger , such that $\|\mathcal{A}'|0\rangle_{\mathcal{H}_A} - |1\rangle_{\mathcal{H}_F}|\phi\rangle_{\mathcal{H}_W}\| \leq \epsilon$ as long as $\sqrt{p_{\text{succ}}} > \delta$.

Amplitude estimation. Classically, if we like to estimate the expectation of the Bernoulli random variable X to within additive error ϵ , we need $\Theta(1/\epsilon^2)$ i.i.d. samples of X . Given access to U in Eq. (7), we can also estimate p with a quadratic quantum speedup:

Theorem 3 (Amplitude estimation [[BHMT02](#), Theorem 12]). *Let \mathcal{A} be a quantum algorithm on space $\mathcal{H}_A = \mathcal{H}_F \otimes \mathcal{H}_W$ such that*

$$\mathcal{A}|0\rangle_{\mathcal{H}_A} = \sqrt{p_{\text{succ}}}|1\rangle_{\mathcal{H}_F}|\phi\rangle_{\mathcal{H}_W} + \sqrt{1-p_{\text{succ}}}|0\rangle_{\mathcal{H}_F}|\psi\rangle_{\mathcal{H}_W} \text{ where } \|\phi\| = 1, \quad (9)$$

the amplitude estimation algorithm outputs a $\tilde{p}_{\text{succ}} \in [0, 1]$ satisfying

$$|\tilde{p}_{\text{succ}} - p_{\text{succ}}| \leq \frac{2\pi\sqrt{p_{\text{succ}}(1-p_{\text{succ}})}}{M} + \frac{\pi^2}{M^2} \quad (10)$$

with success probability at least $8/\pi^2$, using M calls to \mathcal{A} and \mathcal{A}^\dagger .

In application, we often need to estimate p_{succ} to within multiplicative error ϵ . Then we can set

$$M = \frac{3\pi}{\epsilon\sqrt{p_{\text{succ}}}} \quad (11)$$

in [Theorem 3](#) such that

$$|\tilde{p}_{\text{succ}} - p_{\text{succ}}| \leq \frac{2}{3}\epsilon p_{\text{succ}} \sqrt{1 - p_{\text{succ}}} + \frac{1}{9}\epsilon^2 p_{\text{succ}} \leq \epsilon p_{\text{succ}}. \quad (12)$$

2.2 Quantum singular value transformation

In [\[GSLW19\]](#), a general quantum algorithm framework called *quantum singular value transformation* is proposed, which is useful in many computational tasks including property estimation. Before introducing this framework, we first give the definition of singular value transformation.

Definition 5 (Singular value transformation [\[GSLW19, Definition 16\]](#)). *Let $f: \mathbb{R} \rightarrow \mathbb{C}$ be an even or odd function. Suppose that $A \in \mathbb{C}^{\tilde{d} \times d}$ has the following singular value decomposition*

$$A = \sum_{i=1}^{d_{\min}} \sigma_i |\tilde{\psi}_i\rangle \langle \psi_i|, \quad (13)$$

where $d_{\min} := \min(d, \tilde{d})$. For the function f we define the singular value transform of A as

$$f^{(SV)}(A) := \begin{cases} \sum_{i=1}^{d_{\min}} f(\sigma_i) |\tilde{\psi}_i\rangle \langle \psi_i| & \text{if } f \text{ is odd, and} \\ \sum_{i=1}^d f(\sigma_i) |\psi_i\rangle \langle \psi_i| & \text{if } f \text{ is even,} \end{cases} \quad (14)$$

where for $i \in [d] \setminus [d_{\min}]$ we define $\sigma_i := 0$.

To manipulate general matrices A by quantum circuits, we need a tool called *block-encoding*. We call a unitary U a block-encoding of A if for some $\alpha \in (0, +\infty)$ we have

$$U = \begin{bmatrix} A/\alpha & \cdot \\ \cdot & \cdot \end{bmatrix} \quad \text{namely} \quad A = \alpha(\langle 0| \otimes I)U(|0\rangle \otimes I). \quad (15)$$

One can also change the projection to $|0\rangle$ to an orthogonal projection to a general subspace V when we use the technique.

Given a matrix A block-encoded in a unitary, polynomial singular value transformation of A can be efficiently implemented as follows:

Theorem 4 ([\[GSLW19, Corollary 18\]](#)). *Let \mathcal{H}_U be a finite-dimensional Hilbert space and let $U, \Pi, \tilde{\Pi} \in \text{End}(\mathcal{H}_U)$ be linear operators on \mathcal{H}_U such that U is a unitary, and $\tilde{\Pi}, \Pi$ are orthogonal projectors. Suppose that $P = \sum_{k=0}^n a_k x^k \in \mathbb{R}[x]$ is a degree- n polynomial such that*

$$a_k \neq 0 \text{ only if } k \equiv n \pmod{2}, \text{ and} \quad (16)$$

$$\text{for all } x \in [-1, 1]: |P(x)| \leq 1. \quad (17)$$

Then there exists a vector $\Phi = (\phi_1, \phi_2, \dots, \phi_n) \in \mathbb{R}^n$, such that

$$P^{(SV)}(\tilde{\Pi}U\Pi) = \begin{cases} (\langle +| \otimes \tilde{\Pi})U_P^{(SV)}(|+\rangle \otimes \Pi) & \text{if } n \text{ is odd, and} \\ (\langle +| \otimes \Pi)U_P^{(SV)}(|+\rangle \otimes \Pi) & \text{if } n \text{ is even,} \end{cases} \quad (18)$$

where $U_P^{(SV)} := |0\rangle\langle 0| \otimes U_\Phi + |1\rangle\langle 1| \otimes U_{-\Phi}$ with

$$U_\Phi := \begin{cases} e^{i\phi_1(2\tilde{\Pi}-I)}U \prod_{j=1}^{(n-1)/2} \left(e^{i\phi_{2j}(2\Pi-I)}U^\dagger e^{i\phi_{2j+1}(2\tilde{\Pi}-I)}U \right) & \text{if } n \text{ is odd, and} \\ \prod_{j=1}^{n/2} \left(e^{i\phi_{2j-1}(2\Pi-I)}U^\dagger e^{i\phi_{2j}(2\tilde{\Pi}-I)}U \right) & \text{if } n \text{ is even.} \end{cases} \quad (19)$$

Note that $P^{(SV)}(\tilde{\Pi}U\Pi)$ acts on the same space as $\tilde{\Pi}U\Pi$, while $U_P^{(SV)}$ acts on an enlarged space with one ancillary qubit added. This theorem tells us that for a polynomial P of degree d which satisfies Eq. (16) in [Theorem 4](#), we can implement $P^{(SV)}(\tilde{\Pi}U\Pi)$ with $\Theta(d)$ uses of U, U^\dagger and controlled reflections $I - 2\Pi, I - 2\tilde{\Pi}$.

2.3 Variable-stopping-time algorithms

Variable-stopping-time quantum algorithm. In [\[Amb10\]](#), *variable-stopping-time quantum algorithms* are proposed to characterize those having different branches of computations stopping at different time. We follow the definition in [\[CKS17\]](#) and [\[CGJ19\]](#).

Definition 6 (Variable-stopping-time quantum algorithm [\[CGJ19, Definition 13\]](#)). *We say that $\mathcal{A} = \mathcal{A}_m \cdots \mathcal{A}_1$ is a variable-stopping-time quantum algorithm if \mathcal{A} acts on $\mathcal{H} = \mathcal{H}_C \otimes \mathcal{H}_A$, where $\mathcal{H}_C = \otimes_{i=1}^m \mathcal{H}_{C_i}$ with $\mathcal{H}_{C_i} = \text{Span}(|0\rangle, |1\rangle)$, and each unitary \mathcal{A}_j acts on $\mathcal{H}_{C_j} \otimes \mathcal{H}_A$ controlled on the first $j - 1$ qubits being $|0\rangle^{\otimes(j-1)} \in \otimes_{i=1}^{j-1} \mathcal{H}_{C_i}$.*

The algorithm \mathcal{A} is divided into m stages $\mathcal{A}_1, \dots, \mathcal{A}_m$ according to the m possible stopping times t_1, \dots, t_m . In property estimation problem, we focus more on query complexity, so the t_j we refer to is the query complexity of $\mathcal{A}_j \cdots \mathcal{A}_1$.

In any stage j , the unitary \mathcal{A}_j can set the state in \mathcal{H}_{C_j} to $|1\rangle$. This indicates that the computation has stopped on this branch, since any $\mathcal{A}_k, k > j$ is controlled on the state in $\otimes_{i=1}^{k-1} \mathcal{H}_{C_i}$ being all-0 state and does not alter the state on this branch since the state in \mathcal{H}_{C_j} is set to $|1\rangle$.

In order to analyze \mathcal{A} , we give the definition of the probability of the algorithm stopping by time t_j as follows:

Definition 7 (Probability of stopping by time t_j [\[CGJ19, Definition 14\]](#)). *We define the orthogonal projector*

$$\Pi_{\text{stop} \leq t} := \sum_{j: t_j \leq t} |1\rangle\langle 1|_{C_j} \otimes I_{\mathcal{H}_A}, \quad (20)$$

where by $|1\rangle\langle 1|_{C_j}$ we denote the orthogonal projector on \mathcal{H}_C which projects onto the state

$$|0\rangle_{\mathcal{H}_{C_1}} \otimes \cdots \otimes |0\rangle_{\mathcal{H}_{C_{j-1}}} \otimes |1\rangle_{\mathcal{H}_{C_j}} \otimes |0\rangle_{\mathcal{H}_{C_{j+1}}} \otimes \cdots \otimes |0\rangle_{\mathcal{H}_{C_m}}. \quad (21)$$

Then we define $p_{\text{stop} \leq t} := \|\Pi_{\text{stop} \leq t} \mathcal{A} | \mathbf{0} \rangle\|^2$, and similarly $p_{\text{stop} \geq t}$ and $p_{\text{stop} = t_j}$.

It is also worth mentioning that in our applications, it always holds that $p_{\text{stop} \leq t_m} = 1$. Let $p_{\text{stop} = t_j} := p_{\text{stop} \leq t_j} - p_{\text{stop} \leq t_{j-1}}$. We define the average complexity of \mathcal{A} in a way similar to that in [\[Amb10\]](#) by

$$T_{\text{avg}} := \sqrt{\sum_{j=1}^m t_j^2 \cdot p_{\text{stop} = t_j}}, \quad (22)$$

and the maximum complexity of \mathcal{A}

$$T_{\text{max}} := t_m. \quad (23)$$

Variable-time amplitude amplification and estimation. Suppose we have access to a variable-stopping-time quantum algorithm \mathcal{A} acting on $\mathcal{H} = \mathcal{H}_C \otimes \mathcal{H}_A$ such that

$$\mathcal{A}|0\rangle = \sqrt{p_{\text{succ}}}|1\rangle_{\mathcal{H}_F}|\phi\rangle_{\mathcal{H}_W, \mathcal{H}_C} + \sqrt{1 - p_{\text{succ}}}|0\rangle_{\mathcal{H}_F}|\psi\rangle_{\mathcal{H}_W, \mathcal{H}_C}, \quad (24)$$

where $\|\phi\| = \|\psi\| = 1$, $\mathcal{H}_A = \mathcal{H}_F \otimes \mathcal{H}_W$, and $\mathcal{H}_F = \text{Span}(|0\rangle, |1\rangle)$ indicates “good” and “bad” outcomes. If we want to obtain the “good” outcome $|\phi\rangle$, we can use amplitude amplification algorithm in [Theorem 2](#) with $\Theta(\frac{1}{\sqrt{p_{\text{succ}}}})$ calls to \mathcal{A} and \mathcal{A}^\dagger , so the total complexity is $\frac{T_{\text{max}}}{\sqrt{p_{\text{succ}}}}$. However, we can do better for variable-stopping-time algorithm. In [\[Amb10\]](#), the following variable-time amplitude amplification algorithm is proposed with lower complexity:

Theorem 5 (Variable-time amplitude amplification [\[Amb10, Theorem 1\]](#)). *Given a variable-stopping-time quantum algorithm \mathcal{A} acting on $\mathcal{H} = \mathcal{H}_C \otimes \mathcal{H}_A$ such that*

$$\mathcal{A}|0\rangle = \sqrt{p_{\text{succ}}}|1\rangle_{\mathcal{H}_F}|\phi\rangle_{\mathcal{H}_W, \mathcal{H}_C} + \sqrt{1 - p_{\text{succ}}}|0\rangle_{\mathcal{H}_F}|\psi\rangle_{\mathcal{H}_W, \mathcal{H}_C} \text{ where } \|\phi\| = 1, \quad (25)$$

where $\mathcal{H}_A = \mathcal{H}_F \otimes \mathcal{H}_W$ and $\mathcal{H}_F = \text{Span}(|0\rangle, |1\rangle)$, let T_{avg} and T_{max} be the parameters defined in [Eq. \(22\)](#) and [Eq. \(23\)](#). We can construct a quantum algorithm \mathcal{A}' invoking \mathcal{A} several times, for total time

$$\mathcal{O}\left(T_{\text{max}}\sqrt{\log T_{\text{max}}} + \frac{T_{\text{avg}}}{\sqrt{p_{\text{succ}}}}\log^{1.5} T_{\text{max}}\right) \quad (26)$$

that produces a state $\alpha|1\rangle_{\mathcal{H}_F}|\phi\rangle_{\mathcal{H}_W, \mathcal{H}_C} + \beta|0\rangle_{\mathcal{H}_F}|\psi\rangle_{\mathcal{H}_W, \mathcal{H}_C}$ with probability $|\alpha|^2 \geq 1/2$ as the output.

In [\[CGJ19\]](#), standard amplitude estimation algorithm in [Theorem 3](#) was also generalized to variable-time scenarios.

Theorem 6 (Variable-time amplitude estimation [\[CGJ19, Theorem 23\]](#)). *Let \mathcal{A} be a variable-stopping-time quantum algorithm acting on $\mathcal{H} = \mathcal{H}_C \otimes \mathcal{H}_A$ such that*

$$\mathcal{A}|0\rangle = \sqrt{p_{\text{succ}}}|1\rangle_{\mathcal{H}_F}|\phi\rangle_{\mathcal{H}_W, \mathcal{H}_C} + \sqrt{1 - p_{\text{succ}}}|0\rangle_{\mathcal{H}_F}|\psi\rangle_{\mathcal{H}_W, \mathcal{H}_C}, \quad (27)$$

where $\|\phi\| = 1$, $\mathcal{H}_A = \mathcal{H}_F \otimes \mathcal{H}_W$, and $\mathcal{H}_F = \text{Span}(|0\rangle, |1\rangle)$. Let $T_{\text{avg}}, T_{\text{max}}$ be the parameters defined in [Eq. \(22\)](#) and [Eq. \(23\)](#), respectively, $T'_{\text{max}} := 2T_{\text{max}}/t_1$ and

$$Q = T_{\text{max}}\sqrt{\log(T'_{\text{max}})} + \frac{T_{\text{avg}}\log(T'_{\text{max}})}{\sqrt{p_{\text{succ}}}}. \quad (28)$$

Suppose that we know a lower bound p'_{succ} of p_{succ} . Then for any $\epsilon, \delta \in (0, 1)$, we can estimate p_{succ} to within multiplicative error ϵ and success probability at least $1 - \delta$ with complexity

$$\mathcal{O}\left(\frac{Q}{\epsilon}\log^2(T'_{\text{max}})\log\left(\frac{\log(T'_{\text{max}})}{\delta}\right) + Q\log(T'_{\text{max}})\log\left(\frac{1}{\delta}\log\left(\frac{T'_{\text{max}}}{p'_{\text{succ}}}\right)\right)\right) = \tilde{\mathcal{O}}(Q/\epsilon). \quad (29)$$

Note that the total complexity of estimating p_{succ} to within multiplicative error ϵ using the standard amplitude estimation algorithm is $MT_{\text{max}} = \mathcal{O}(\frac{T_{\text{max}}}{\epsilon\sqrt{p_{\text{succ}}}})$, where M is determined in [Eq. \(11\)](#). Therefore, if T_{avg} is much smaller than T_{max} , we can achieve a great acceleration by replacing the standard amplitude estimation algorithm with variable-time amplitude estimation.

3 Main Algorithm

3.1 Estimating properties of probability distributions by QSVT

In this section, we introduce a quantum algorithm for estimating properties of a probability distribution $\mathbf{p} = (p_i)_{i=1}^n$ on $[n]$ to within a certain error. Our algorithm is based on quantum singular value transformation (QSVT) and amplitude estimation, which is similar to the entropy estimation algorithm in [GL20]. In this paper, we mostly focus on the pure state preparation oracle in [Definition 1](#), and we will show in [Section 6](#) that our algorithm also works well with purified quantum query-access oracle in [Definition 3](#). Here we give a brief explanation. Compared to the pure-state preparation oracle, the purified quantum query-access oracle just adds an unknown state in the right-hand side of Eq. (6). However, in our algorithm, we produce a quantum state $|\psi\rangle$ such that the module square of the amplitude of the projection of $|\psi\rangle$ onto a subspace, $\|\Pi|\psi\rangle\|^2$, encodes the quantity we want to estimate, where the projector Π acts as an identity in the added space. As a result, the module square $\|\Pi|\psi\rangle\|^2$ is independent of the unknown added state.

We first construct a unitary operator which encodes the information of the probability distribution that we want to study. Specifically, we construct a unitary U acting on $2\lceil\log n\rceil$ qubits which has a matrix block with singular values $\sqrt{p_1}, \dots, \sqrt{p_n}$ as follows:

$$U := U_p \otimes I, \tag{30}$$

where U_p is the pure-state preparation oracle. Let

$$\tilde{\Pi} := \sum_{i=1}^n |i\rangle\langle i| \otimes |i\rangle\langle i|, \text{ and } \Pi := |\mathbf{0}\rangle\langle \mathbf{0}| \otimes I, \tag{31}$$

and then U is a block encoding of

$$E := \tilde{\Pi}U\Pi = \sum_{i=1}^n \sqrt{p_i}|i\rangle\langle \mathbf{0}| \otimes |i\rangle\langle i|, \tag{32}$$

which has singular values $\sqrt{p_1}, \dots, \sqrt{p_n}$.

Suppose S is a polynomial satisfying Eq. (16) in [Theorem 4](#). We propose an algorithm to estimate $\sum_{i=1}^n p_i S(\sqrt{p_i})^2$ to within a certain multiplicative error.

Before describe out main algorithm, we first give a rough version of the amplitude estimation algorithm and it only needs a lower bound of the module of the amplitude.

Lemma 1. *Let \mathcal{A} be a quantum algorithm on space $\mathcal{H}_A = \mathcal{H}_F \otimes \mathcal{H}_W$ such that*

$$\mathcal{A}|0\rangle_{\mathcal{H}_A} = \sqrt{p_{\text{succ}}}|1\rangle_{\mathcal{H}_F}|\phi\rangle_{\mathcal{H}_W} + \sqrt{1 - p_{\text{succ}}}|0\rangle_{\mathcal{H}_F}|\psi\rangle_{\mathcal{H}_W}, \text{ where } \|\phi\rangle\| = 1. \tag{33}$$

Given L such that $L \leq p_{\text{succ}}$, there exists an algorithm which outputs an estimate $\tilde{p} \in [0, 1]$ satisfying $\tilde{p}/p_{\text{succ}} \in [\frac{1}{2}, 2]$ with success probability at least $1 - \delta$, using $\mathcal{O}\left(\frac{1}{\sqrt{p_{\text{succ}}}} \log\left(\frac{1}{\sqrt{p_{\text{succ}}}}\right) \log\left(\frac{\log(\frac{1}{L})}{\delta}\right)\right) = \tilde{\mathcal{O}}\left(\frac{1}{\sqrt{p_{\text{succ}}}}\right)$ calls to \mathcal{A} and \mathcal{A}^\dagger .

The proof of [Lemma 1](#) is deferred to [Appendix B](#). With this estimate at hand, we now describe our main algorithm in [Algorithm 1](#), and prove its correctness in [Lemma 2](#).

Algorithm 1: Estimate the value $\sum_{i=1}^n p_i S(\sqrt{p_i})^2$ for $\mathbf{p} = (p_i)_{i=1}^n$.

Input: Multiplicative error ϵ , and quantum registers (A, B, Q, F) initialized to $|\mathbf{0}\rangle$.

Output: \tilde{p} , an estimation of the value $\sum_{i=1}^n p_i S(\sqrt{p_i})^2$.

Parameter: (L, S) , where L is a lower bound of $\sum_{i=1}^n p_i S(\sqrt{p_i})^2$, and S is an odd or even polynomial.

Notation:

- A, B are two $\lceil \log n \rceil$ -qubit input registers;
- Q is a single-qubit register, to be used as an ancilla register for QSVT;
- F is a single-qubit flag register indicating “good” components whose amplitude we estimate;
- $U_S^{(SV)}$ is the unitary $U_P^{(SV)}$ in [Theorem 4](#) when the transformation polynomial $P := S$, and $U, \tilde{\Pi}, \Pi$ in [Theorem 4](#) are set to $U, \tilde{\Pi}, \Pi$ in Eq. (30), Eq. (31) and Eq. (32).

Regard the following subroutine as \mathcal{A} :

- 1 Apply U_p to register B and Hadamard gate H to register Q ;
 - 2 Apply unitary $U_S^{(SV)}$ to registers (A, B, Q) using Q as the ancilla register ;
 - 3 **if** S is an odd function **then**
 - 4 Apply unitary $C_{\tilde{\Pi} \otimes |+\rangle\langle +|} \text{NOT} = \tilde{\Pi} \otimes |+\rangle\langle +|_Q \otimes X_F + (I - \tilde{\Pi} \otimes |+\rangle\langle +|_Q) \otimes I_F$ to registers (A, B, Q, F) ;
 - 5 **else**
 - 6 Apply unitary $C_{\Pi \otimes |+\rangle\langle +|} \text{NOT} = \Pi \otimes |+\rangle\langle +|_Q \otimes X_F + (I - \Pi \otimes |+\rangle\langle +|_Q) \otimes I_F$ to registers (A, B, Q, F) ;
 - 7 Run the algorithm in [Lemma 1](#) with $\delta = \frac{1}{8}$ and lower bound L to obtain a rough estimate of $\|(I \otimes |1\rangle\langle 1|_F)|\mathcal{A}|\mathbf{0}\rangle\|^2$. Denote this estimate by P ;
 - 8 Use the amplitude estimation algorithm in [Theorem 3](#) with $M = \lceil \frac{5\pi}{\sqrt{P}\epsilon} \rceil$ to estimate $\|(I \otimes |1\rangle\langle 1|_F)|\mathcal{A}|\mathbf{0}\rangle\|^2$ and denote the result by \tilde{p} . Output \tilde{p} ;
-

Lemma 2. Suppose S is a polynomial satisfying Eq. (16) in [Theorem 4](#), $\mathbf{p} = (p_i)_{i=1}^n$ is a probability distribution on $[n]$ and we know a lower bound $L \leq \sum_{i=1}^n p_i S(\sqrt{p_i})^2$. Then for any $\epsilon \in (0, 1)$, [Algorithm 1](#) with parameters $L := L, S := S$ and input ϵ outputs an estimate of $\sum_{i=1}^n p_i S(\sqrt{p_i})^2$ to within multiplicative error ϵ with success probability at least $\frac{2}{3}$ using $\tilde{\mathcal{O}}\left(\frac{1}{\epsilon} \frac{\deg(S)}{\sqrt{\sum_{i=1}^n p_i S(\sqrt{p_i})^2}}\right)$ calls to U_p and U_p^\dagger .

Proof. We first analyze the error of [Algorithm 1](#). Let the state in registers (A, B, Q) after [Line 2](#) of [Algorithm 1](#) be $|\Phi_p\rangle$. Assuming S is an odd function for the moment for simplicity, the output $|\Phi_p\rangle$ satisfies that

$$\begin{aligned}
(\tilde{\Pi} \otimes \langle +|_Q) |\Phi_p\rangle &= (\tilde{\Pi} \otimes \langle +|_Q) U_S^{(SV)} (|\mathbf{0}\rangle_A |\Psi_{\mathbf{p}}\rangle_B |+\rangle_Q) \\
&= (\tilde{\Pi} \otimes \langle +|_Q) (|0\rangle\langle 0|_Q \otimes U_{\Phi_S} + |1\rangle\langle 1|_Q \otimes U_{-\Phi_S}) (|\mathbf{0}\rangle_A |\Psi_{\mathbf{p}}\rangle_B |+\rangle_Q) \\
&= ((\tilde{\Pi} \otimes \langle +|_Q) (|0\rangle\langle 0|_Q \otimes U_{\Phi_S} + |1\rangle\langle 1|_Q \otimes U_{-\Phi_S}) (\Pi \otimes |+\rangle_Q)) (|\mathbf{0}\rangle_A |\Psi_{\mathbf{p}}\rangle_B) \\
&= S^{(SV)} (\tilde{\Pi} U \Pi) \left(\sum_{i=1}^n \sqrt{p_i} |\mathbf{0}\rangle_A |i\rangle_B \right) \\
&= \left(\sum_{i=1}^n S(\sqrt{p_i}) |i\rangle\langle i|_A \otimes |i\rangle\langle i|_B \right) \left(\sum_{i=1}^n \sqrt{p_i} |\mathbf{0}\rangle_A |i\rangle_B \right) \\
&= \sum_{i=1}^n \sqrt{p_i} S(\sqrt{p_i}) |i\rangle_A |i\rangle_B, \tag{34}
\end{aligned}$$

where the third equation comes from $(\Pi \otimes |+\rangle_Q) (|\mathbf{0}\rangle_A |\Psi_{\mathbf{p}}\rangle_B) = |\mathbf{0}\rangle_A |\Psi_{\mathbf{p}}\rangle_B |+\rangle_Q$.

Let the state in registers (A, B, Q, F) after [Line 6](#) be $|\Psi_{\mathbf{p}}\rangle$. Then we have

$$\begin{aligned}
|\Psi_{\mathbf{p}}\rangle &= C_{\tilde{\Pi} \otimes |+\rangle\langle +|} \text{NOT} |\Phi_p\rangle |0\rangle_F \\
&= (\tilde{\Pi} \otimes |+\rangle\langle +|_Q \otimes X_F) |\Phi_p\rangle |0\rangle_F + ((I - \tilde{\Pi} \otimes |+\rangle\langle +|_Q) \otimes I_F) |\Phi_p\rangle |0\rangle_F \\
&= \sum_{i=1}^n \sqrt{p_i} S(\sqrt{p_i}) |i\rangle_A |i\rangle_B |+\rangle_Q |1\rangle_F + |\psi_{\text{garbage}}\rangle |0\rangle_F, \tag{35}
\end{aligned}$$

where the third equation comes from [Eq. \(34\)](#) and $|\psi_{\text{garbage}}\rangle$ is an unnormalized state which we do not care about.

Although we suppose S to be an odd function, it is easily verified that [Eq. \(34\)](#) and [Eq. \(35\)](#) hold for all even functions S as well, if we replace $\tilde{\Pi}$ with Π .

Note that if we measure register F after the subroutine \mathcal{A} , the success probability is

$$\|(I \otimes |1\rangle\langle 1|_F) |\Psi_{\mathbf{p}}\rangle\|^2 = \left\| \sum_{i=1}^n \sqrt{p_i} S(\sqrt{p_i}) |i\rangle_A |i\rangle_B |+\rangle_Q |1\rangle_F \right\|^2 = \sum_{i=1}^n p_i S(\sqrt{p_i})^2, \tag{36}$$

the quantity we would like to estimate.

From [Lemma 1](#), we can infer that the output P in [Line 7](#) satisfies $P / \sum_{i=1}^n p_i S(\sqrt{p_i})^2 \in [\frac{1}{2}, 2]$ with probability at least $\frac{7}{8}$, so according to [Theorem 3](#), with success probability at least $\frac{8}{\pi^2} \frac{7}{8} \geq \frac{2}{3}$, the output \tilde{p} satisfies

$$\begin{aligned}
|\tilde{p} - \sum_{i=1}^n p_i S(\sqrt{p_i})^2| &\leq \frac{2}{5} \sqrt{P} \epsilon \sqrt{\sum_{i=1}^n p_i S(\sqrt{p_i})^2} + \frac{1}{25} P \epsilon^2 \\
&\leq \left(\frac{2\sqrt{2}}{5} + \frac{2}{25} \right) \epsilon \sum_{i=1}^n p_i S(\sqrt{p_i})^2 \leq \epsilon \sum_{i=1}^n p_i S(\sqrt{p_i})^2, \tag{37}
\end{aligned}$$

where the second inequality is because $P \leq 2 \sum_{i=1}^n p_i S(\sqrt{p_i})^2$.

We now calculate the complexity of the algorithm. In \mathcal{A} , we call U_p and U_p^\dagger $\tilde{\mathcal{O}}(\deg(S))$ times. In the main algorithm, the step using [Lemma 1](#) calls subroutine \mathcal{A} $\tilde{\mathcal{O}}(1/\sqrt{\sum_i p_i S(\sqrt{p_i})^2})$ times, and




Figure 2: Circuit of $C_{\tilde{\Pi} \otimes |+\rangle \langle +|} \text{NOT} = \tilde{\Pi} \otimes |+\rangle \langle +|_Q \otimes X_F + (I - \tilde{\Pi} \otimes |+\rangle \langle +|_Q) \otimes I_F$ if n is a power of 2. n_A, n_B, n_C, n_F are the sizes of the registers A, B, C, F , respectively. The CNOT gate between two registers with the same size is an abbreviation of a sequence of CNOT gates between qubits in different registers with the same index and the CNOT gate targeting a qubit conditional on a register will flip the qubit when the register is an all-0/all-1 state.

the step using [Theorem 3](#) calls subroutine \mathcal{A} M times. Overall, the query complexity of [Algorithm 1](#) is

$$\tilde{\mathcal{O}} \left(\left(M + \frac{1}{\sqrt{\sum_{i=1}^n p_i S(\sqrt{p_i})^2}} \right) \deg(S) \right) = \tilde{\mathcal{O}} \left(\frac{\deg(S)}{\epsilon \sqrt{\sum_{i=1}^n p_i S(\sqrt{p_i})^2}} \right), \quad (38)$$

as claimed. \square

This algorithm is also gate efficient. If n is a power of 2, $C_{\tilde{\Pi} \otimes |+\rangle \langle +|} \text{NOT}$ can be efficiently implemented following [Figure 2](#), and it can be easily generalized to any integer $n > 0$ using additional quantum circuit which can determine whether $|i\rangle \in \mathcal{H}$, $\dim(\mathcal{H}) = 2^{\lceil \log n \rceil}$ satisfies $i \leq n$.

3.2 Improvements based on annealing

In [Section 3.1](#), we have shown how to estimate the quantity

$\sum_{i=1}^n p_i S(\sqrt{p_i})^2$ to within a certain multiplicative error, where S is a polynomial satisfying Eq. (16) in [Theorem 4](#). However, in many cases the quantity we want to estimate cannot be written in this form. Here we consider a more general quantity of discrete distributions $\mathbf{p} = (p_i)_{i=1}^n$ on $[n]$:

$$f(\mathbf{p}) := \sum_{i=1}^n p_i g(p_i), \quad (39)$$

where $g(x)$ is a monotonically increasing function on $[0, 1]$ such that $g(0) = 0$, $g(1) = 1$ and $xg(x)$ is a convex function.

An observation is that for any probability distribution \mathbf{p} on $[n]$, given a lower bound L of $f(\mathbf{p})$, if we can find a polynomial S satisfying Eq. (16) in [Theorem 4](#) and

$$\left| \sum_{i=1}^n q_i S(\sqrt{q_i})^2 - f(\mathbf{q}) \right| \leq \epsilon f(\mathbf{q}) \quad (40)$$

for any distribution $\mathbf{q} \in \Delta^n$, then by [Lemma 2](#), [Algorithm 1](#) can estimate $f(\mathbf{p})$ to within multiplicative error ϵ using

$$\tilde{\mathcal{O}} \left(\left(\frac{1}{\epsilon} \right) \frac{\deg(S)}{\sqrt{f(\mathbf{p})}} \right) \quad (41)$$

calls to U_p and U_p^\dagger for any distribution \mathbf{p} .

However, this algorithm can be sub-optimal in many cases. To give an intuitive explanation, we compare this algorithm with the one in [HM19]. In [HM19], they develop a quantum algorithm which can estimate the mean of a random variable X to within multiplicative error ϵ using $\tilde{\mathcal{O}}\left(\frac{\sqrt{\mathbb{E}[X^2]}}{\epsilon\mathbb{E}[X]}\right)$ quantum samples. **Algorithm 1** is somewhat similar to this algorithm, and it can also be seen as estimating the mean of a random variable X with $\Pr[X = g(p_i)] = p_i$. Note a catch that the value of X in our problem is not given, and we need to estimate it ourselves. In **Algorithm 1**, we takes one sample from $\mathbf{p} = (p_i)_{i=1}^n$ in **Line 1** and estimate the value of this sample by QSVT in **Line 2**. Therefore, for a fair comparison, we remove the cost of estimating the value of the sample in **Algorithm 1**, and the remaining query cost is $\tilde{\mathcal{O}}\left(\frac{1}{\epsilon\sqrt{f(\mathbf{p})}}\right) = \tilde{\mathcal{O}}\left(\frac{1}{\epsilon\sqrt{\mathbb{E}[X]}}\right)$. This is higher than the cost of $\tilde{\mathcal{O}}\left(\frac{\sqrt{\mathbb{E}[X^2]}}{\epsilon\mathbb{E}[X]}\right)$ in [HM19] because $\mathbb{E}[X^2] \leq \mathbb{E}[X]$ due to $X = g(p_i) \leq g(1) = 1$.

Nevertheless, we can improve **Algorithm 1** if we are given an estimate of $f(\mathbf{p})$ to within constant multiplicative error. The idea is that we can estimate $\sum_{i=1}^n \Phi p_i g(p_i)$ for an amplification factor $\Phi \gg 1$ instead using **Algorithm 1**, for which we need to find a polynomial S such that $\sum_{i=1}^n p_i S(\sqrt{p_i}) / (\sum_{i=1}^n \Phi p_i g(p_i)) \in [1 - \epsilon, 1 + \epsilon]$. This brings the benefit that the quantity we estimate is much bigger, so the query complexity becomes smaller since it is inversely proportional to the square root of the quantity we estimate. Nevertheless, the amplification factor Φ should not be too large, since we need to guarantee the existence of the polynomial $S(x)$ which satisfies Eq. (16) in **Theorem 4** and is also an approximation to $\sqrt{\Phi g(x^2)}$ when $x = \sqrt{p_i}$. **Theorem 4** requires $|S(x)| \leq 1$ for all $x \in [-1, 1]$, and this requires $|\Phi g(x^2)| \leq 1$ for $x = \sqrt{p_i}$. Since $g(x^2)$ is a monotonically increasing function, we only need to ensure that $\Phi g(\max_{i \in [n]} p_i) \leq 1$. Estimating the maximum p_i is not simple, and an alternative method is to obtain an upper bound of p_i from a rough estimate of $f(\mathbf{p})$. This is possible because $g(x)$ is monotonically increasing and positive, so we have $p_i g(p_i) \leq f(\mathbf{p})$ and then $p_i \leq (xg(x))^{-1} f(\mathbf{p})$ for all $i \in [n]$, where $(xg(x))^{-1}$ is the inverse function of $xg(x)$ on $[0, 1]$. Detailed analysis is conducted in the following lemma.

Lemma 3. *For any convex and monotonically increasing function $g(x)$ on $[0, 1]$ such that $g(0) = 0$ and $g(1) = 1$ and probability distribution $\mathbf{p} = (p_i)_{i=1}^n$ on $[n]$, let $f(\mathbf{p}) := \sum_{i=1}^n p_i g(p_i)$. Suppose that we are given constants a, b and P such that $af(\mathbf{p}) \leq P \leq bf(\mathbf{p})$ and let $p^* \in [0, 1]$ be such that $p^* g(p^*) = \min(\frac{1}{a}P, 1)$. For any $\epsilon \in (0, 1)$, let $\epsilon_0 = \frac{\epsilon}{2}$. Then if we can construct a polynomial S satisfying Eq. (16) in **Theorem 4** and*

$$\left| \sum_{i=1}^n q_i S(\sqrt{q_i})^2 - c \frac{1}{g(p^*)} f(\mathbf{q}) \right| \leq \frac{ca}{2b} p^* \epsilon_0 \text{ for all } \mathbf{q} \text{ such that } q_i \leq p^* \text{ for all } i \in [n], \quad (42)$$

where $c > 0$ is an arbitrary positive constant, **Algorithm 1** with parameters to be $P := \frac{P}{b}$, $S := S$, and input ϵ_0 outputs \tilde{p} satisfying that $\frac{g(p^*)}{c} \tilde{p}$ is an estimate of $f(\mathbf{p})$ to within multiplicative error ϵ . This call to **Algorithm 1** uses $\tilde{\mathcal{O}}\left(\frac{\deg(S)}{\epsilon\sqrt{p^*}}\right)$ calls to U_p and U_p^\dagger .

Proof. From **Lemma 2**, \tilde{p} is an estimate of $\sum_{i=1}^n p_i S(\sqrt{p_i})^2$ to within multiplicative error ϵ_0 , and this call to **Algorithm 1** uses $\tilde{\mathcal{O}}\left(\frac{1}{\epsilon_0} \frac{\deg(S)}{\sqrt{p_i S(\sqrt{p_i})^2}}\right)$ calls to U_p and U_p^\dagger . From Eq. (42), we have

$$\sum_{i=1}^n p_i S(\sqrt{p_i})^2 \geq c \frac{1}{g(p^*)} f(\mathbf{p}) - \frac{ca}{2b} p^* \epsilon_0 = c \frac{p^*}{\min(\frac{1}{a}P, 1)} f(\mathbf{p}) - \frac{ca}{2b} p^* \epsilon_0 \geq \frac{ca}{b} p^* - \frac{ca}{2b} p^* \epsilon_0 \geq \frac{a}{2b} c p^* \quad (43)$$

where the first equation comes from $p^*g(p^*) = \min(\frac{1}{a}P, 1)$ and the second inequality comes from $P \leq bf(\mathbf{p})$.

Then by [Lemma 2](#), the query complexity is $\tilde{\mathcal{O}}\left(\frac{1}{\epsilon_0} \frac{\deg(S)}{\sqrt{p_i S(\sqrt{p_i})^2}}\right) = \tilde{\mathcal{O}}\left(\frac{1}{\epsilon} \frac{\deg(S)}{\sqrt{p^*}}\right)$.

Let $\tilde{f}(\mathbf{p}) := \frac{g(p^*)}{c} \tilde{p}$. We can infer that $\tilde{f}(\mathbf{p})$ is an approximation of $\frac{g(p^*)}{c} \sum_{i=1}^n p_i S(\sqrt{p_i})^2$ within ϵ_0 multiplicative error. We now prove that $\tilde{f}(\mathbf{p})$ is an estimate of $f(\mathbf{p})$ to within multiplicative error $2\epsilon_0$.

For any $i \in [n]$, we have $p_i g(p_i) \leq f(\mathbf{p}) \leq \frac{1}{a}P$ and $p_i g(p_i) \leq 1$, which implies that $p_i g(p_i) \leq p^*g(p^*)$ for any $i \in [n]$. Since $xg(x)$ is a monotonically increasing function on $[0, 1]$, we have $p_i \leq p^*$ for all $i \in [n]$. Therefore, from Eq. (42), $\sum_{i=1}^n p_i S(\sqrt{p_i})^2$ is an approximation of $c \frac{1}{g(p^*)} f(\mathbf{p})$ within $\frac{ca}{2b} g^* \epsilon_0$ additive error. Since $\frac{c}{g(p^*)} f(\mathbf{p}) = c \frac{p^*}{\min(\frac{1}{a}P, 1)} f(\mathbf{p}) \geq \frac{ca}{b} p^*$, we can infer that $\sum_{i=1}^n p_i S(\sqrt{p_i})^2$ is an approximation of $c \frac{1}{g(p^*)} f(\mathbf{p})$ within $\frac{\frac{ca}{2b} p^* \epsilon_0}{c \frac{1}{g(p^*)} f(\mathbf{p})} \leq \frac{\frac{ca}{2b} p^* \epsilon_0}{\frac{ca}{b} p^*} = \frac{\epsilon_0}{2}$ multiplicative error, and because $\tilde{f}(\mathbf{p})$ is an approximation of $\frac{g(p^*)}{c} \sum_{i=1}^n p_i S(\sqrt{p_i})^2$ within ϵ_0 multiplicative error, we can infer that $\tilde{f}(\mathbf{p})$ is an approximation of $f(\mathbf{p})$ within $(1 + \epsilon_0)(1 + \frac{\epsilon_0}{2}) - 1 \leq 2\epsilon_0 = \epsilon$ multiplicative error. \square

We now show that with the additional information P, a, b in [Lemma 3](#), the new query complexity bound of $\tilde{\mathcal{O}}\left(\frac{\deg(S)}{\epsilon \sqrt{p^*}}\right)$ improves the $\frac{1}{\sqrt{\mathbb{E}[X]}}$ term in the bound in [Lemma 2](#) to $\frac{\sqrt{\mathbb{E}[X^2]}}{\mathbb{E}[X]}$.

Since $xg(x)$ is a convex function, from Jensen's inequality, we have $f(\mathbf{p}) = \sum_{i=1}^n p_i g(p_i) \geq g(\frac{1}{n})$, and because $xg(x)$ is a monotonically increasing function, we have $p^* \geq (xg(x))^{-1}(\frac{1}{a}g(\frac{1}{n}))$.

Therefore, the complexity bound in [Lemma 3](#) becomes $\mathcal{O}\left(\frac{\deg(S)}{\epsilon \sqrt{(xg(x))^{-1}(\frac{1}{a}g(\frac{1}{n}))}}\right)$ in the worst case

and we prove that this bound is equivalent to the aforementioned $\tilde{\mathcal{O}}\left(\frac{\deg(S) \sqrt{\mathbb{E}[X^2]}}{\epsilon \mathbb{E}[X]}\right)$ bound in the following lemma, whose proof is deferred to [Appendix B](#).

Lemma 4. *Let $\mathbf{p} = (p_i)_{i=1}^n$ be a probability distribution on $[n]$, and $g(x)$ be a monotonically increasing function on $[0, 1]$ such that $g(0) = 0, g(1) = 1$ and $xg(x)$ is a convex function. Then we have $\max_{\mathbf{p} \in \Delta^n} \frac{\sqrt{\sum_{i=1}^n p_i g(p_i)^2}}{\sum_{i=1}^n p_i g(p_i)} = \Theta\left(\frac{1}{\sqrt{(xg(x))^{-1}(\frac{1}{a}g(\frac{1}{n}))}}\right)$ as $n \rightarrow \infty$, where $a > 0$ is a constant satisfying the conditions in [Lemma 3](#).*

The remained problem is to get an estimate of $f(\mathbf{p})$ to within constant multiplicative error. We propose a framework based on annealing as follows to solve it.

Proposition 1. *Suppose $g(x)$ is a monotonically increasing function on $[0, 1]$ such that $g(0) = 0, g(1) = 1$ and $xg(x)$ is a convex function. Let $\mathbf{p} = (p_i)_{i=1}^n$ be a probability distribution on $[n]$, $\epsilon \in (0, 1)$, and $\delta \in (0, 1)$. Define $f(\mathbf{p}) := \sum_{i=1}^n p_i g(p_i)$. Suppose that there exists a sequence of monotonically increasing functions $g_1(x), \dots, g_{l-1}(x), g_l(x) = g(x)$ on $[0, 1]$ and $\epsilon_1, \dots, \epsilon_{l-1}, \epsilon_l = \epsilon$ such that $g_k(0) = 0, g_k(1) = 1$ and $xg_k(x)$ is a convex function for all $k = 1, \dots, l$, and*

$$\max_{\mathbf{r} \in \Delta^n} \frac{\max_{\mathbf{q} \in \Delta^n, f_k(\mathbf{r})/f_k(\mathbf{q}) \in [1-\epsilon_k, 1+\epsilon_k]} f_{k+1}(\mathbf{q})}{\min_{\mathbf{q} \in \Delta^n, f_k(\mathbf{r})/f_k(\mathbf{q}) \in [1-\epsilon_k, 1+\epsilon_k]} f_{k+1}(\mathbf{q})} \leq c \text{ for all } k = 1, \dots, l-1, \text{ and} \quad (44)$$

$$\frac{\max_{\mathbf{q} \in \Delta^n} f_1(\mathbf{q})}{\min_{\mathbf{q} \in \Delta^n} f_1(\mathbf{q})} \leq c, \quad (45)$$

where $f_k(\mathbf{q}) := \sum_{i=1}^n q_i g_k(q_i)$ and c is a constant. If for any $k = 1, \dots, l$ and probability distribution \mathbf{q} on $[n]$, there exists an algorithm \mathcal{A}_k which can estimate $f_k(\mathbf{q})$ to within multiplicative error ϵ_k

with success probability at least $1 - \frac{\delta}{l}$ using Q_k calls to U_p and U_p^\dagger given two constants a, b , and P satisfying $a f_k(\mathbf{q}) \leq P \leq b f_k(\mathbf{q})$, then there exists an algorithm which can estimate $f(\mathbf{p})$ to within multiplicative error ϵ with success probability at least $1 - \delta$ using $\sum_{k=1}^l Q_k$ calls to U_p and U_p^\dagger .

Proof. From Eq. (45), we have $\max_{\mathbf{q} \in \Delta^n} f_1(\mathbf{q}) \leq c \min_{\mathbf{q} \in \Delta^n} f_1(\mathbf{q}) \leq c f_1(\mathbf{p})$, so $P := \max_{\mathbf{q} \in \Delta^n} f_1(\mathbf{q})$, $a := 1$, and $b := c$ are valid parameters for algorithm \mathcal{A}_1 . By our assumption, we can get an estimate of $f_1(\mathbf{p})$, denoted by $\tilde{f}_1(\mathbf{p})$, to within multiplicative error ϵ_1 with success probability at least $1 - \frac{\delta}{l}$ using \mathcal{A}_1 with Q_1 calls to U_p and U_p^\dagger .

For each k from 2 to l , we let

$$P_k := \max_{\mathbf{q} \in \Delta^n, \tilde{f}_{k-1}(\mathbf{p})/f_{k-1}(\mathbf{q}) \in [1-\epsilon_{k-1}, 1+\epsilon_{k-1}]} f_k(\mathbf{q}), \quad (46)$$

set the parameters of \mathcal{A}_k as $P := P_k, a := 1, b := c$, run \mathcal{A}_k to estimate $f_k(\mathbf{p})$, and denote the output by $\tilde{f}_k(\mathbf{p})$.

We prove that the output $\tilde{f}_k(\mathbf{p})$ is an estimate of $f_k(\mathbf{p})$ to within multiplicative error ϵ_k with success probability at least $1 - \frac{\delta k}{l}$ by induction. The statement is true for $k = 1$ by our assumption. If the statement is true for $k-1$, which means $\tilde{f}_{k-1}(\mathbf{p})/f_{k-1}(\mathbf{p}) \in [1-\epsilon_{k-1}, 1+\epsilon_{k-1}]$ with probability at least $1 - \frac{\delta(k-1)}{l}$, from Eq. (46), we can infer that

$$f_k(\mathbf{p}) \leq \max_{\mathbf{q} \in \Delta^n, \tilde{f}_{k-1}(\mathbf{p})/f_{k-1}(\mathbf{q}) \in [1-\epsilon_{k-1}, 1+\epsilon_{k-1}]} f_k(\mathbf{q}) = P_k; \quad (47)$$

$$f_k(\mathbf{p}) \geq \min_{\mathbf{q} \in \Delta^n, \tilde{f}_{k-1}(\mathbf{p})/f_{k-1}(\mathbf{q}) \in [1-\epsilon_{k-1}, 1+\epsilon_{k-1}]} f_k(\mathbf{q}) \geq P_k/c, \quad (48)$$

where the third inequality comes from Eq. (44). Therefore, $(P_k, 1, c)$ are valid parameters for \mathcal{A}_k , and the output $\tilde{f}_k(\mathbf{p})$ is an estimate of $f_k(\mathbf{p})$ to within multiplicative error ϵ_k with success probability at least $(1 - \frac{\delta(k-1)}{l})(1 - \frac{\delta}{l}) \geq 1 - \frac{\delta k}{l}$, which completes the induction proof.

In conclusion, $\tilde{f}_l(\mathbf{p})$ is an estimate of $f_l(\mathbf{p}) = f(\mathbf{p})$ to within multiplicative error $\epsilon_l = \epsilon$, and the query complexity of the whole algorithm is $\sum_{k=1}^l Q_k$. \square

3.3 Improvements based on variable-time amplitude estimation

In Section 3.1, we apply quantum singular value transformation to all singular values of E with the same transformation polynomial S . However, if the singular values that need to be transformed fall into a small interval, say, $[\lambda, 1]$, where $\lambda \in (0, 1)$, we may construct a polynomial which has smaller degree but still performs a transformation similar to S in this interval.

In Section 2.3, it is shown that variable-time amplitude estimation works better than standard amplitude estimation if T_{avg} is much smaller than T_{max} , which means some branches of computation run much shorter than T_{max} . To employ this observation, we can use a variable-stopping-time quantum algorithm to replace the subroutine \mathcal{A} in Algorithm 1 and use variable time amplitude estimation to replace the standard amplitude estimation. In the variable-stopping-time version \mathcal{A} , we only apply singular value transformation to the singular values in a small pre-defined interval in each stage and we choose intervals in which transformation is easier to perform as the intervals in early stages. In this way, those branches stopping in an early stage make T_{avg} smaller than T_{max} . Similar situations also appeared in [CKS17] and [Amb10], which inspires our variable-stopping-time algorithm.

To this end, we give an improved version of Algorithm 1 alone, and will not consider the improvements of Section 3.2 in this subsection. Before describing the improved main algorithm, we first introduce a tool to help us separate singular values.

Tool. In order to transform different singular values in different stages of \mathcal{A} , we need to decompose a state into several components each of which is a linear combination of singular vectors of E whose singular values fall into a small interval.

In [CKS17], they use a gapped phase estimation algorithm and Hamiltonian simulation algorithm to separate eigenvalues in different intervals. We extend their algorithm to the following one which can deal with singular values.

Lemma 5. *Let U be a unitary, and $\tilde{\Pi}, \Pi$ orthogonal projectors with the same rank d acting on \mathcal{H}_I . Suppose $A = \tilde{\Pi}U\Pi$ has a singular value decomposition $A = \sum_{i=1}^d \sigma_i |\tilde{\psi}_i\rangle\langle\psi_i|_I$. Let $\varphi \in (0, 1]$ and $\epsilon > 0$. Then there is a unitary $W(\varphi, \epsilon)$ using $\mathcal{O}\left(\frac{1}{\varphi} \log \frac{1}{\epsilon}\right)$ calls to U and U^\dagger such that*

$$W(\varphi, \epsilon)|0\rangle_C|0\rangle_P|\psi_i\rangle_I = \beta_0|0\rangle_C|\gamma\rangle_{P,I} + \beta_1|1\rangle_C|+\rangle_P|\psi_i\rangle_I \quad (49)$$

where $|\beta_0|^2 + |\beta_1|^2 = 1$, such that

- if $0 \leq \sigma_i \leq \varphi$ then $|\beta_1| \leq \epsilon$ and
- if $2\varphi \leq \sigma_i \leq 1$ then $|\beta_0| \leq \epsilon$.

Here C and P are two single-qubit registers, and I is the register that A acts on.

The proof of Lemma 5 is deferred to Appendix B.

Variable-stopping-time subroutine $\tilde{\mathcal{A}}$. We now describe the m -stage variable-time quantum algorithm $\mathcal{A} = \mathcal{A}_m \cdots \mathcal{A}_1$.

To construct \mathcal{A} , we suppose that we are given $\beta \in (0, 1]$ such that $\sqrt{p_i} \in [0, \beta]$ for all $i = 1, \dots, n$ and a lower bound $L > 0$ such that $\sum_{i=1}^n p_i S(\sqrt{p_i})^2 \geq L$.

Let $\varphi_j := \beta 2^{-j}$ for $j = 0, \dots, m-1$ and $\varphi_m := 0$. We first divide $[0, \beta]$ into m intervals $[\varphi_m, \varphi_{m-1}), [\varphi_{m-1}, \varphi_{m-2}), \dots, [\varphi_1, \varphi_0)$. Then we transform singular values in these intervals in different stages of \mathcal{A} . Specifically,

- for $j = 2, \dots, m$, we transform singular values in $[\beta 2^{-j}, \beta 2^{-j+2})$ in \mathcal{A}_j , and
- for $j = 0$, we transform singular values in $[\frac{1}{2}\beta, \beta)$ in \mathcal{A}_1 .

Then we construct the j -th stage of \mathcal{A} . First, we need to determine the transformation polynomial, S_j in this stage. Since we like S_j to perform a transformation similar to S , we need to construct polynomials S_j for $j = 1, \dots, m$ such that S_j satisfies Eq. (16) in Theorem 4 and

$$\begin{aligned} |S_1(x) - S(x)| &\leq L\epsilon \text{ for all } x \in [\beta/2, \beta), \\ |S_j(x) - S(x)| &\leq L\epsilon \text{ for all } x \in [\beta 2^{-j}, \beta 2^{-j+2}), \text{ and } j = 2, \dots, m. \end{aligned} \quad (50)$$

Note that for any S_j , we only require it to be a good approximation of $S(x)$ in a small interval, so we may construct such polynomial with lower degree than S . Since the complexity of variable time amplitude estimation is proportional to the average time of all stages, which is the average degree of all transformation polynomial S_j , this variable-stopping-time algorithm can improve our vanilla algorithm in Section 3.1.

Assuming that we have constructed such S_j satisfying Eq. (50), we give a detailed description of \mathcal{A} in Algorithm 2.

Algorithm 2: A variable-stopping-time subroutine when estimating $\sum_{i=1}^n p_i S(\sqrt{p_i})^2$.

Input: Multiplicative error ϵ , and quantum registers (F, C, A, B, Q, P, I) initialized to $|0\rangle$.

Output: Quantum state in registers (F, C, A, B, Q, P, I)

Parameter: $(S, \beta, L, m, \{S_j \mid j \in [m]\})$, where S is the transform polynomial we want to approximate, β is an upper bound of $\sqrt{p_i}$ for all $i \in [n]$, L is a lower bound of $\sum_{i=1}^n p_i S(\sqrt{p_i})^2$, m is the number of stages of \mathcal{A} , and S_j is the transformation polynomial in \mathcal{A}_j , which satisfies Eq. (16) in Theorem 4 and Eq. (50).

Notation:

- F is a single-qubit flag register indicating “good” components whose amplitude we estimate;
- $C = (C_1, \dots, C_m)$ is an m -qubit clock register determining the interval which the singular value belongs to;
- A, B are two $\lceil \log n \rceil$ -qubit input registers;
- Q is a single-qubit register used as ancilla register for QSVT;
- $P = (P_1, P_2, \dots, P_m)$ and $I = (I_1, I_2, \dots, I_m)$ are two registers used as ancilla registers in Lemma 5. Each I_j is a $2\lceil \log n \rceil$ -qubit register and each P_j is a single-qubit register; and
- $U_{S_j}^{(SV)}$ is the unitary $U_P^{(SV)}$ in Theorem 4 when the transformation polynomial $P := S_j$ and $U, \tilde{\Pi}, \Pi$ in Theorem 4 are set to $U, \tilde{\Pi}, \Pi$ in Eq. (30), Eq. (31) and Eq. (32).

- 1 Set $\varphi_j := \beta 2^{-j}$ for $j = 0, \dots, m-1$, and $\varphi_m := 0$;

Regard the following subroutine as \mathcal{A}_j :

- 2 Conditional on first $j-1$ qubits in register C being $|0\rangle$, apply CNOT gates controlled by qubits in register B to flip the last $\lceil \log n \rceil$ qubits in register I_j ;
 - 3 Conditional on first $j-1$ qubits in register C being $|0\rangle$, if $j < m$, apply $W(\varphi_j, L\epsilon/m)$ in Lemma 5 with $\tilde{\Pi}, \Pi, U$ defined in Section 3.1 to the state in register I_j using C_j as the output register C and P_j as the ancilla register P , else apply X gate to register C_m ;
 - 4 Conditional on C_j being $|1\rangle_{C_j}$, apply unitary $C_{\tilde{\Pi}|0\rangle\langle +|} \text{NOT} \cdot U_{S_j}^{(SV)}$ or $C_{\Pi|0\rangle\langle +|} \text{NOT} \cdot U_{S_j}^{(SV)}$ in the same way as in Line 2 to Line 6 of Algorithm 1 to the state in (A, B, Q, F) ;
 - 5 Apply U_p to register B , Hadamard gate H to Q ;
 - 6 Apply $\mathcal{A} := \mathcal{A}_m \cdots \mathcal{A}_1$ to registers (F, C, A, B, Q, P, I) ;
-

Final algorithm. We now describe our final algorithm in Algorithm 3.

We prove the output of Algorithm 3 is an estimate of $\sum_{i=1}^n p_i S(\sqrt{p_i})^2$ to within multiplicative error ϵ with high probability in the following proposition:

Proposition 2. Let $\epsilon, \delta, \beta \in (0, 1)$, $\mathbf{p} = (p_i)_{i=1}^n$ be a probability distribution such that $\sqrt{p_i} \leq \beta$ for all $i = 1, \dots, n$ and S be a polynomial which satisfies Eq. (16) in Theorem 4. Suppose that we are given $\beta, L > 0$ such that $\sum_{i=1}^n p_i S(\sqrt{p_i})^2 \geq L$ and a sequence of polynomials $\{S_j \mid j = 1, \dots, m\}$ which satisfy Eq. (16) in Theorem 4 and Eq. (50). Algorithm 3 with input (ϵ, δ) and parameters $(S, \beta, L, m, \{S_j \mid j \in [m]\})$ outputs an estimate of $\sum_{i=1}^n p_i S(\sqrt{p_i})^2$ to within multiplicative error ϵ with success probability at least $1 - \delta$.

Let $t_j = \frac{2^j}{\beta} \log(\frac{m}{\epsilon L}) + \sum_{k=1}^j \deg(S_k)$ for all $j = 1, \dots, m$ and $t_{m+1} = t_m$, the query complexity

Algorithm 3: Improved algorithm for estimating $\sum_{i=1}^n p_i S(\sqrt{p_i})^2$ of $\mathbf{p} = (p_i)_{i=1}^n$.

Input: (ϵ, δ) , where ϵ is the multiplicative error, and δ is the failure probability; quantum registers (F, C, A, B, Q, P, I) initialized to $|0\rangle$.

Output: \tilde{p} .

Notation: Notations is the same as that of [Algorithm 2](#).

Parameter: $(S, \beta, L, m, \{S_j \mid j \in [m]\})$.

- 1 Use variable-time amplitude estimation algorithm in [Theorem 6](#) with the parameters (ϵ, δ) to be $\epsilon := \epsilon, \delta := \delta$ and registers $W := (A, B, Q, P, I), C := C, F := F$ to estimate $\|(|1\rangle\langle 1|_F \otimes I) \tilde{\mathcal{A}}(|0\rangle)\|^2$, where $\tilde{\mathcal{A}}$ is [Algorithm 2](#) with the same parameters $(S, \beta, L, m, \{S_j \mid j \in [m]\})$ and input ϵ . Denote the output of variable-time amplitude estimation algorithm by \tilde{p} . Output \tilde{p} ;
-

of [Algorithm 3](#) is

$$\tilde{\mathcal{O}} \left(t_m + \sqrt{\epsilon} \sum_{j=1}^m t_j + \frac{\sqrt{\sum_{j=1}^m \sum_{i: \sqrt{p_i} \in [\varphi_j, \varphi_{j-1})} p_i t_{j+1}^2}}{\sqrt{\sum_{i=1}^n p_i S(\sqrt{p_i})^2}} \right), \quad (51)$$

where $\varphi_j = 2^{-j}\beta$ for $j = 1, \dots, m-1$ and $\varphi_m = 0$. We omit polylog terms of L, δ and t_m in this bound.

The proof of [Proposition 2](#) is deferred to [Appendix A](#).

Remark 1. The $\sqrt{\epsilon} \sum_{j=1}^m t_j$ term in query complexity can be eliminated by a more detailed analysis mentioned in [[CKS17](#)], but it does not improve the complexity bounds in our applications.

4 Rényi Entropy Estimation ($\alpha > 1$)

In this section, we propose a quantum algorithm to estimate $H_\alpha(\mathbf{p}) = \frac{1}{1-\alpha} \sum_{i=1}^n p_i^\alpha$ for $\alpha > 1$ to within additive error ϵ . This is equivalent to estimating $P_\alpha(\mathbf{p}) = \sum_{i=1}^n p_i^\alpha$ to within multiplicative error $\mathcal{O}(\epsilon)$.

Let $g(x) := x^{\alpha-1}$, and then we have $P_\alpha(\mathbf{p}) = \sum_{i=1}^n p_i g(p_i)$. Since $g(x) = x^{\alpha-1}$ for $\alpha > 1$ is monotonically increasing function on $[0, 1]$ such that $g(0) = 0, g(1) = 1$ and $xg(x) = x^\alpha$ is a convex function, we can use the framework in [Section 3.2](#) to construct our algorithm.

4.1 Estimate $P_\alpha(\mathbf{p})$ given a rough bound

We first construct a quantum algorithm which can estimate $P_\alpha(\mathbf{p})$ to within multiplicative error given P, a, b such that $aP_\alpha(\mathbf{p}) \leq P \leq bP_\alpha(\mathbf{p})$ following [Lemma 3](#). Let $p^* = (xg(x))^{-1}(\min(\frac{1}{a}P, 1)) = (\min(\frac{1}{a}P, 1))^{\frac{1}{\alpha}}$. Like [Lemma 3](#), we need to construct a polynomial $S(x)$ which satisfies Eq. (16) in [Theorem 4](#) and Eq. (42), which means

$$\left| \sum_{i=1}^n q_i S(\sqrt{q_i})^2 - c \frac{1}{g(p^*)} f(\mathbf{q}) \right| \leq \frac{ca}{2b} p^* \epsilon_0 \text{ for all } \mathbf{q} \text{ such that } q_i \leq p^* \text{ for all } i \in [n], \quad (52)$$

where $c > 0$ is an arbitrary constant.

Before constructing such a polynomial, we first construct a class of polynomials which satisfies Eq. (16) in [Theorem 4](#) and is also a good approximation to $2^{-c-1}\beta^{-c}x^c$ in $[\nu, \beta]$ for any $\beta \in (0, 1], \nu \in (0, \beta)$, and $c > 0$.

Lemma 6. For any $c > 0$, $\beta \in (0, 1]$, $\nu \in (0, \beta)$, and $\eta \in (0, \frac{1}{2})$, let $f(x) = 2^{-c-1}\beta^{-c}x^c$, there is an efficiently computable even or odd polynomial $S \in \mathbb{R}[x]$ of degree $\mathcal{O}\left(\frac{\epsilon}{\nu} \log\left(\frac{1}{\beta\nu\eta}\right)\right)$ such that

$$\begin{aligned} \forall x \in [0, \nu] : |S(x)| &\leq 2f(x) \\ \forall x \in [\nu, \beta] : |f(x) - S(x)| &\leq \eta \\ \forall x \in [-1, 1] : |S(x)| &\leq 1 \end{aligned} \tag{53}$$

Proof. Let $d = \lceil c \rceil - c$. We first introduce a lemma to construct polynomial approximation of kx^{-d} where k is a constant.

Lemma 7 ([GSLW19, Corollary 67, Polynomial approximations of negative power functions]). Let $\delta, \epsilon \in (0, \frac{1}{2}]$, $c > 0$, and let $f(x) := \frac{\delta^c}{2}x^{-c}$, then there exist even/odd polynomials $P, P' \in \mathbb{R}[x]$ such that $\|P - f\|_{[\delta, 1]} \leq \epsilon$, $\|P\|_{[-1, 1]} \leq 1$ and similarly $\|P' - f\|_{[\delta, 1]} \leq \epsilon$, $\|P'\|_{[-1, 1]} \leq 1$. In addition, the degree of the polynomials are $\mathcal{O}\left(\frac{\max[1, c]}{\delta} \log\left(\frac{1}{\epsilon}\right)\right)$.

Setting the parameters (ϵ, δ, c) in **Lemma 7** to $\delta := \nu, c := d, \epsilon := \frac{\beta^c \nu^d \eta}{2}$, we can construct an even polynomial $\tilde{Q} := P$ in **Lemma 7** with $\deg(\tilde{Q}) = \mathcal{O}\left(\frac{1}{\nu} \log\left(\frac{1}{\beta\nu\eta}\right)\right)$ such that

$$\begin{aligned} \forall x \in [-1, 1] : |\tilde{Q}(x)| &\leq 1 \\ \forall x \in [\nu, 1] : |\tilde{Q}(x) - \frac{\nu^d}{2}x^{-d}| &\leq \frac{\beta^c \nu^d \eta}{2}. \end{aligned} \tag{54}$$

Then we need to construct a polynomial approximation to the rectangle function according to the following lemma:

Lemma 8 (Polynomial approximations of the rectangle function [GSLW19, Lemma 29]). Let $\delta', \epsilon' \in (0, \frac{1}{2})$ and t satisfying $\delta' \leq t \leq 1$. There exists an even polynomial $P' \in \mathbb{R}[x]$ of degree $\mathcal{O}\left(\log\left(\frac{1}{\epsilon'}\right)/\delta'\right)$, such that $|P'(x)| \leq 1$ for all $x \in [-1, 1]$, and

$$\begin{cases} P'(x) \in [0, \epsilon'] & \text{for all } x \in [-1, -t - \delta'] \cup [t + \delta', 1], \text{ and} \\ P'(x) \in [1 - \epsilon', 1] & \text{for all } x \in [-t + \delta', t - \delta']. \end{cases} \tag{55}$$

Setting $\delta' := \frac{\beta}{2}, t := \frac{3\beta}{2}, \epsilon' := \frac{\beta^c \nu^d \eta}{2}$ in **Lemma 8**, we can construct an even polynomial P with $\deg(P) = \mathcal{O}\left(\frac{1}{\beta} \log\left(\frac{1}{\beta\nu\eta}\right)\right)$ such that

$$\begin{aligned} \forall x \in [-1, 1] : |P(x)| &\leq 1 \\ \forall x \in [2\beta, 1] : |P(x)| &\leq \frac{\beta^c \nu^d \eta}{2} \\ \forall x \in [-\beta, \beta] : 1 - \frac{\beta^c \nu^d \eta}{2} &\leq P(x) \leq 1. \end{aligned} \tag{56}$$

Let $Q(x) := \tilde{Q}(x)P(x)$, we have

$$\begin{aligned}
& \forall x \in [-1, 1] : |Q(x)| \leq 1 \\
& \forall x \in [\nu, \beta] : |Q(x) - \frac{\nu^d}{2}x^{-d}| \leq |Q(x) - \tilde{Q}(x)| + |\tilde{Q}(x) - \frac{\nu^d}{2}x^{-d}| \\
& \hspace{15em} \leq |\tilde{Q}(x)||1 - P(x)| + \frac{\beta^c \nu^d \eta}{2} \\
& \hspace{15em} \leq |1 - P(x)| + \frac{\beta^c \nu^d \eta}{2} \leq \beta^c \nu^d \eta \\
& \forall x \in [\beta, 2\beta] : |Q(x)| \leq |\tilde{Q}(x)| \leq \frac{\nu^d x^{-d}}{2} + \frac{\beta^c \nu^d \eta}{2} \\
& \forall x \in [2\beta, 1] : |Q(x)| \leq |P(x)| \leq \frac{\beta^c \nu^d \eta}{2}. \tag{57}
\end{aligned}$$

Then, let $S(x) := 2^{-c} \beta^{-c} \nu^{-d} x^{\lceil c \rceil} Q(x)$, which is an even or odd polynomial since $Q(x) = \tilde{Q}(x)P(x)$ and $\tilde{Q}(x), P(x)$ are even polynomials. We can infer that

$$\forall x \in [0, \nu] : S(x) \leq 2^{-c} \beta^{-c} \nu^{-d} x^{\lceil c \rceil} = 2^{-c} \beta^{-c} \nu^{-d} x^d x^c \leq 2^{-c} \beta^{-c} x^c = 2f(x), \tag{58}$$

where the first inequality comes from $|Q(x)| \leq 1$, and

$$\begin{aligned}
\forall x \in [\nu, \beta] : |S(x) - f(x)| &= |S(x) - 2^{-c-1} \beta^{-c} x^c| = 2^{-c} \beta^{-c} \nu^{-d} x^{\lceil c \rceil} |Q(x) - \frac{\nu^d x^{-d}}{2}| \\
&\leq 2^{-c} \beta^{-c} \nu^{-d} x^{\lceil c \rceil} (\beta^c \nu^d \eta) \\
&\leq 2^{-c} \eta \leq \eta, \tag{59}
\end{aligned}$$

where the first inequality comes from Eq. (57).

To prove that $S(x)$ is bounded by 1 on $[-1, 1]$, since $S(x)$ is even or odd, we only need to prove $\forall x \in [0, 1] : |S(x)| \leq 1$. For $x \in [0, 2\beta]$, we have

$$\begin{aligned}
\forall x \in [0, \nu] : |S(x)| &\leq 2f(x) \leq 1 \\
\forall x \in [\nu, \beta] : |S(x)| &\leq 2^{-c-1} \beta^{-c} x^c + \eta \leq \frac{1}{2} + \eta \leq 1, \tag{60}
\end{aligned}$$

where the first inequality comes from Eq. (58) and the third inequality comes from Eq. (59), and

$$\begin{aligned}
\forall x \in [\beta, 2\beta] : |S(x)| &\leq 2^{-c-1} \beta^{-c} \nu^{-d} x^{\lceil c \rceil} \left(\frac{\nu^d x^{-d} + \beta^c \nu^d \eta}{2} \right) \\
&\leq 2^{-c-1} \frac{\beta^{-c} x^c + \eta}{2} \\
&\leq 2^{-c-1} \frac{2^c + \eta}{2} \leq 1, \tag{61}
\end{aligned}$$

where the first inequality comes from Eq. (57).

For $x \in [2\beta, 1]$, we have

$$\forall x \in [2\beta, 1] : |S(x)| = 2^{-c-1} \beta^{-c} \nu^{-d} x^{\lceil c \rceil} Q(x) \leq 2^{-c-1} \beta^{-c} \nu^{-d} \left(\frac{\beta^c \nu^d \eta}{2} \right) \leq \eta \leq 1, \tag{62}$$

where the first inequality comes from Eq. (57). Therefore, we can conclude that $\forall x \in [-1, 1] : |S(x)| \leq 1$. Together with Eq. (58), Eq. (59) and that $S(x)$ is even or odd polynomial, we have that $S(x)$ with $\deg(S) = \lceil c \rceil + \deg(\tilde{Q}) + \deg P = \mathcal{O}\left(\frac{c}{\nu} \log\left(\frac{1}{\beta \nu \eta}\right)\right)$ satisfies the conditions in this lemma. \square

By carefully choosing the parameters in [Lemma 6](#), we can construct a polynomial which is similar to the polynomial S in [Eq. \(42\)](#) as follows.

Lemma 9. *For any probability distribution \mathbf{p} on $[n]$, suppose that we are given P, a, b such that $aP_\alpha(\mathbf{p}) \leq P \leq bP_\alpha(\mathbf{p})$. Let $p^* = \min(\frac{1}{a}P, 1)^{\frac{1}{\alpha}}$. Then for any $\epsilon \in (0, 1)$, $\alpha > 1$ and constants $d, d' > 0$, the polynomial S in [Lemma 6](#) with parameters (c, β, ν, η) to be $c := \alpha - 1, \beta := \sqrt{p^*}, \nu := \left(d \frac{a(p^*)^\alpha \epsilon}{5bn}\right)^{\frac{1}{2\alpha}}$, $\eta := d' 2^{-2\alpha} \frac{a}{b} p^* \epsilon$ has $\deg(S) = \tilde{\mathcal{O}}\left(\left(\frac{n}{\epsilon}\right)^{\frac{1}{2\alpha}} \frac{1}{\sqrt{p^*}}\right)$, and satisfies [Eq. \(16\)](#) in [Theorem 4](#) and*

$$\left| \sum_{i=1}^n p_i S(\sqrt{p_i})^2 - 2^{-2\alpha} (p^*)^{1-\alpha} P_\alpha(\mathbf{p}) \right| \leq (2d' + d) 2^{-2\alpha} \frac{a}{b} p^* \epsilon. \quad (63)$$

Proof. [Lemma 6](#) implies that $\deg(S) = \mathcal{O}\left(\frac{c}{\nu} \log\left(\frac{1}{\beta\nu\eta}\right)\right) = \tilde{\mathcal{O}}\left(\left(\frac{n}{\epsilon}\right)^{\frac{1}{2\alpha}} \frac{1}{\sqrt{p^*}}\right)$, S is an even or odd polynomial, and $|S(x)| \leq 1$ for all $x \in [-1, 1]$. Therefore, S satisfies [Eq. \(16\)](#) in [Theorem 4](#).

We now prove that S satisfies [Eq. \(63\)](#). From the definition of $P_\alpha(\mathbf{p})$, we can infer that

$$\forall i, p_i \leq \left(\sum_{i=1}^n p_i^\alpha\right)^{\frac{1}{\alpha}} \leq \left(\frac{1}{a}P\right)^{\frac{1}{\alpha}} \text{ and } p_i \leq 1, \text{ so } p_i \leq \min\left(\left(\frac{1}{a}P\right)^{\frac{1}{\alpha}}, 1\right) = p^* = \beta^2. \quad (64)$$

From [Lemma 6](#), we can infer that S satisfies

$$\forall x \in [0, \nu] : |S(x)| \leq 2^{1-\alpha} \beta^{1-\alpha} x^{\alpha-1} \quad (65)$$

$$\forall x \in [\nu, \beta] : |2^{-\alpha} \beta^{1-\alpha} x^{\alpha-1} - S(x)| \leq \eta. \quad (66)$$

For i such that $\sqrt{p_i} \leq \nu$, we have

$$\begin{aligned} \sum_{\sqrt{p_i} \leq \nu} |p_i S(\sqrt{p_i})^2 - 2^{-2\alpha} \beta^{-2\alpha+2} p_i^\alpha| &\leq \sum_{\sqrt{p_i} \leq \nu} |p_i S(\sqrt{p_i})^2| + |2^{-2\alpha} \beta^{-2\alpha+2} p_i^\alpha| \\ &\leq \sum_{\sqrt{p_i} \leq \nu} p_i 2^{-2\alpha+2} \beta^{-2\alpha+2} p_i^{\alpha-1} + 2^{-2\alpha} \beta^{-2\alpha+2} p_i^\alpha \\ &\leq \sum_{\sqrt{p_i} \leq \nu} 2^{-2\alpha} \beta^{-2\alpha+2} 5 p_i^\alpha \\ &\leq \sum_{\sqrt{p_i} \leq \nu} 2^{-2\alpha} \beta^{-2\alpha+2} 5 \nu^{2\alpha} \\ &\leq n 2^{-2\alpha} \beta^{-2\alpha+2} 5 \nu^{2\alpha} \\ &= n 2^{-2\alpha} (p^*)^{1-\alpha} 5 \left(d \frac{a(p^*)^\alpha \epsilon}{5bn}\right) \\ &= d 2^{-2\alpha} \frac{a}{b} p^* \epsilon, \end{aligned} \quad (67)$$

where the second inequality comes from [Eq. \(65\)](#).

For i such that $\sqrt{p_i} > \nu$, we have

$$\begin{aligned}
\sum_{\sqrt{p_i} > \nu} |p_i S(\sqrt{p_i})^2 - 2^{-2\alpha} \beta^{-2\alpha+2} p_i^\alpha| &= \sum_{\sqrt{p_i} > \nu} p_i |S(\sqrt{p_i})^2 - 2^{-2\alpha} \beta^{-2\alpha+2} (\sqrt{p_i})^{2c}| \\
&= \sum_{\sqrt{p_i} > \nu} p_i |S(\sqrt{p_i}) - 2^{-\alpha} \beta^{-\alpha+1} (\sqrt{p_i})^c| |S(\sqrt{p_i}) + 2^{-\alpha} \beta^{-\alpha+1} (\sqrt{p_i})^c| \\
&\leq 2 \sum_{\sqrt{p_i} > \nu} p_i |S(\sqrt{p_i}) - 2^{-\alpha} \beta^{-\alpha+1} (\sqrt{p_i})^c| \\
&\leq 2\eta \sum_{\sqrt{p_i} > \nu} p_i \leq 2\eta = 2d' 2^{-2\alpha} \frac{a}{b} p^* \epsilon, \tag{68}
\end{aligned}$$

where the first inequality comes from Eq. (64) and Eq. (66).

From Eq. (67) and Eq. (68), we can infer that

$$\left| \sum_{i=1}^n p_i S(\sqrt{p_i})^2 - 2^{-2\alpha} (p^*)^{1-\alpha} P_\alpha(\mathbf{p}) \right| \leq (2d' + d) 2^{-2\alpha} \frac{a}{b} p^* \epsilon, \tag{69}$$

which completes the proof. \square

Therefore, from Lemma 3, there exists an algorithm which can estimate $P_\alpha(\mathbf{p})$ to within multiplicative error $\mathcal{O}(\epsilon)$ using $\tilde{\mathcal{O}}\left(\frac{\deg(S)}{\epsilon \sqrt{p^*}}\right) = \tilde{\mathcal{O}}\left(\frac{n^{\frac{1}{2\alpha}}}{\epsilon^{1+\frac{1}{2\alpha}} p^*}\right) = \tilde{\mathcal{O}}\left(\frac{n^{\frac{1}{2\alpha}}}{\epsilon^{1+\frac{1}{2\alpha}} P^\alpha}\right)$ given P, a, b such that $aP_\alpha(\mathbf{p}) \leq P \leq bP_\alpha(\mathbf{p})$.

Then we use Algorithm 3 to replace Algorithm 1 in Lemma 3 and apply Proposition 2 to achieve a better query complexity upper bound.

Lemma 10. *For any $\alpha > 1$, there exists an algorithm \mathcal{A} such that for any $\delta \in (0, 1)$, $\epsilon \in (0, 1)$ and probability distribution \mathbf{p} on $[n]$, given P, a, b such that $aP_\alpha(\mathbf{p}) \leq P \leq bP_\alpha(\mathbf{p})$ where a, b are two constants, \mathcal{A} can estimate $P_\alpha(\mathbf{p})$ to within multiplicative error ϵ with success probability at least $1 - \delta$ using $\tilde{\mathcal{O}}\left(\frac{n^{1-\frac{1}{2\alpha}}}{\epsilon} + \frac{\sqrt{n}}{\epsilon^{1+\frac{1}{2\alpha}}}\right)$ calls to U_p and U_p^\dagger .*

Proof. Correctness. Let $p^* := \min(\frac{1}{a}P, 1)^\alpha$ and $\beta_0 = \sqrt{p^*}$, and we have

$$\forall i, p_i \leq \left(\sum_{i=1}^n p_i^\alpha\right)^\frac{1}{\alpha} \leq \left(\frac{1}{a}P\right)^\frac{1}{\alpha} \text{ and } p_i \leq 1, \text{ so } p_i \leq \min\left(\left(\frac{1}{a}P\right)^\frac{1}{\alpha}, 1\right) = p^* = \beta_0^2. \tag{70}$$

Let $\nu_0 = \left(\frac{1}{4} \frac{a(p^*)^\alpha \epsilon}{5bn}\right)^\frac{1}{2\alpha}$. Before constructing S and S_j in Proposition 2, we first define the number of stages of our variable-stopping-time quantum algorithm $m_0 := \lceil \log\left(\frac{\beta_0}{\nu_0}\right) \rceil + 1$, and $\nu_j = 2^{-j} \beta_0$ for $j = 1, \dots, m_0 - 1$, $\nu_{m_0} = \nu_{m_0-1} = 2^{-m_0+1} \beta_0$.

Let $L := \frac{a}{b} 2^{-2\alpha-1} p^*$, and we will prove that L is a lower bound of $\sum_{i=1}^n p_i S_0(\sqrt{p_i})^2$ later.

Let S_0 be the polynomial S in Lemma 6 with parameters (c, β, ν, η) to be $c := \alpha - 1$, $\beta := \beta_0$, $\nu := \nu_{m_0}$, $\eta := \frac{L}{4}\epsilon$.

Let S_j for $j = 1, \dots, m_0$ be the polynomial S in Lemma 6 with parameters (c, β, ν, η) to be $c := \alpha - 1$, $\beta := \beta_0$, $\nu := \nu_j$, $\eta = \frac{L}{4}\epsilon$. From Lemma 6, we have

$$|S_j(x) - 2^{-\alpha} \beta_0^{-\alpha+1} x^{\alpha-1}| \leq \frac{L}{4}\epsilon \text{ for any } x \in [\nu_j, \beta_0]. \tag{71}$$

Now we set the parameters $(S, \beta, L, m, \{S_j \mid j \in [m]\})$ of [Algorithm 3](#) to be $S := S_0$, $\beta := \beta_0$, $L := L$, $m := m_0$, $S_j := S_j$ for $j = 1, \dots, m_0$, and then prove that these parameters satisfy the conditions in [Proposition 2](#).

- For β_0 , we have shown that it is an upper bound of $\sqrt{p_i}$.
- For L , we need to prove that it is a lower bound of $\sum_{i=1}^n p_i S_0(\sqrt{p_i})^2$. Note that $\nu_{m_0} = 2^{-m_0+1}\beta_0 = 2^{-\lceil \log(\frac{\beta_0}{\nu_0}) \rceil} \beta_0 \leq \nu_0 = \left(\frac{1}{4} \frac{a(p^*)^\alpha \epsilon}{5bn}\right)^{\frac{1}{2\alpha}}$. Let $d = \nu_{m_0}^{2\alpha} / \left(\frac{a(p^*)^\alpha \epsilon}{5bn}\right)$, and $d' = \frac{L}{4}\epsilon / (2^{-2\alpha} \frac{a}{b} p^* \epsilon)$, and then we have $d \leq \left(\frac{1}{4} \frac{a(p^*)^\alpha \epsilon}{5bn}\right) / \left(\frac{a(p^*)^\alpha \epsilon}{5bn}\right) = \frac{1}{4}$, $d' = \frac{1}{8}$. Therefore, the parameters $(c = \alpha - 1, \beta = \beta_0, \nu = \nu_{m_0}, \eta = \frac{L}{4}\epsilon)$ of S_0 satisfy the conditions in [Lemma 9](#) with constants d and d' . From [Lemma 9](#), we have

$$\left| \sum_{i=1}^n p_i S_0(\sqrt{p_i})^2 - 2^{-2\alpha} (p^*)^{1-\alpha} P_\alpha(\mathbf{p}) \right| \leq (2d' + d) 2^{-2\alpha} \frac{a}{b} p^* \epsilon \leq \frac{1}{2} 2^{-2\alpha} \frac{a}{b} p^* \epsilon = L\epsilon. \quad (72)$$

From Eq. (72), we have

$$\begin{aligned} \sum_{i=1}^n p_i S_0(\sqrt{p_i})^2 &\geq 2^{-2\alpha} (p^*)^{1-\alpha} P_\alpha(\mathbf{p}) - L\epsilon \\ &\geq \frac{P}{b} 2^{-2\alpha} (p^*)^{1-\alpha} - L\epsilon \\ &\geq \frac{a}{b} 2^{-2\alpha} p^* - L\epsilon = 2L - L\epsilon \geq L. \end{aligned} \quad (73)$$

Therefore, L is a lower bound of $\sum_{i=1}^n p_i S_0(\sqrt{p_i})^2$.

- For S_j , they are constructed by applying [Lemma 6](#), so they satisfy Eq. (16) in [Theorem 4](#). Note that the parameters of S_0 in [Lemma 6](#) is the same as the parameters of S_{m_0} , so we have $S_0 = S_{m_0}$. Then we can infer that

$$\begin{aligned} \text{for any } x \in [\nu_j, \beta_0] : |S_j(x) - S_0(x)| &= |S_j(x) - S_{m_0}(x)| \\ &\leq |S_j(x) - 2^{-\alpha} \beta_0^{-\alpha+1} x^{\alpha-1}| + |S_{m_0}(x) - 2^{-\alpha} \beta_0^{-\alpha+1} x^{\alpha-1}| \\ &\leq \frac{1}{4} L\epsilon + \frac{1}{4} L\epsilon \leq L\epsilon, \end{aligned} \quad (74)$$

where the second inequality comes from Eq. (71) and $\nu_{m_0} < \nu_j$ for $j < m_0$. From Eq. (74), we can infer that for any $x \in [\beta_0 2^{-j}, \beta_0] : |S_j(x) - S_0(x)| \leq L\epsilon$, which meets the requirements of Eq. (50).

Therefore, the parameters we set are valid for [Proposition 2](#), so [Algorithm 3](#) with the same parameters and input (ϵ, δ) can estimate $\sum_{i=1}^n p_i S_0(\sqrt{p_i})^2$ to within multiplicative error ϵ within success probability at least $1 - \delta$. Denote the estimate by \tilde{p} .

Note that

$$2^{-2\alpha} (p^*)^{1-\alpha} P_\alpha(\mathbf{p}) \geq 2^{-2\alpha} (p^*)^{1-\alpha} \frac{P}{b} \geq 2^{-2\alpha} (p^*)^{1-\alpha} \frac{a(p^*)^\alpha}{b} = 2^{-2\alpha} \frac{a}{b} p^* = \frac{L}{2}, \quad (75)$$

and because $\sum_{i=1}^n p_i S_0(\sqrt{p_i})^2$ is an approximation of $2^{-2\alpha} (p^*)^{1-\alpha} P_\alpha(\mathbf{p})$ within additive error $L\epsilon$ from Eq. (72), it is also an approximation of $2^{-2\alpha} (p^*)^{1-\alpha} P_\alpha(\mathbf{p})$ within multiplicative error $\frac{L\epsilon}{L/2} = 2\epsilon$. Therefore, $2^{2\alpha} (p^*)^{\alpha-1} \sum_{i=1}^n p_i S_0(\sqrt{p_i})^2$ is a 2ϵ multiplicative approximation of $P_\alpha(\mathbf{p})$. Therefore, $2^{2\alpha} (p^*)^{\alpha-1} \tilde{p}$ is an $(1 + 2\epsilon)(1 + \epsilon) - 1 \leq 5\epsilon$ multiplicative approximation of $P_\alpha(\mathbf{p})$. We can rescale ϵ to $\frac{1}{5}\epsilon$ so that we can obtain an ϵ -multiplicative approximation of $P_\alpha(\mathbf{p})$.

Complexity. Now we compute the query complexity of the above algorithm. First let us compute t_j defined in [Proposition 2](#). For t_j , we have

$$t_j = \frac{2^j}{\beta_0} \log\left(\frac{m_0}{\epsilon L}\right) + \sum_{k=1}^j \deg(S_k) = \tilde{\mathcal{O}}\left(\frac{2^j}{\beta_0} + \sum_{k=1}^j \frac{2^k}{\beta_0}\right) = \tilde{\mathcal{O}}\left(\frac{2^j}{\beta_0}\right), \quad (76)$$

where the second equation comes from $\deg(S_j) = \tilde{\mathcal{O}}\left(\frac{1}{\nu_j}\right)$ given in [Lemma 6](#).

Let $\varphi_j = 2^{-j}\beta_0$ for $j = 1, \dots, m_0 - 1$ and $\varphi_{m_0} = 0$ following the definition in [Proposition 2](#). Then from [Proposition 2](#), the complexity of the algorithm we construct is

$$\begin{aligned} & \tilde{\mathcal{O}}\left(\frac{1}{\epsilon} \left(t_{m_0} + \sqrt{\epsilon} \sum_{j=1}^{m_0} t_j + \frac{\sqrt{\sum_{j=1}^{m_0} \sum_{i: \sqrt{p_i} \in [\varphi_j, \varphi_{j-1})} p_i t_{j+1}^2}}{\sqrt{\sum_{i=1}^n p_i S_0(\sqrt{p_i})^2}} \right)\right) \\ &= \tilde{\mathcal{O}}\left(\frac{1}{\epsilon} \left(2^{m_0} \frac{1}{\beta_0} + \frac{\sqrt{\sum_{j=1}^{m_0-1} (|\{i : 2^{-j}\beta_0 \leq \sqrt{p_i} \leq 2^{-j+1}\beta_0\}| (2^{-j+1}\beta_0)^2 (\frac{2^{j+1}}{\beta_0})^2) + \sum_{\sqrt{p_i} \leq \varphi_{m_0-1}} p_i (\frac{1}{\varphi_{m_0-1}})^2}}{\sqrt{L}} \right)\right) \\ &= \tilde{\mathcal{O}}\left(\frac{1}{\epsilon} \left(\frac{1}{\nu_{m_0}} + \frac{\sqrt{n+n}}{\sqrt{L}} \right)\right) \\ &= \tilde{\mathcal{O}}\left(\frac{1}{\epsilon} \left(\frac{n^{\frac{1}{2\alpha}}}{\sqrt{p^*} \epsilon^{\frac{1}{2\alpha}}} + \frac{\sqrt{n}}{\sqrt{p^*}} \right)\right) \\ &= \tilde{\mathcal{O}}\left(\frac{n^{\frac{1}{2\alpha}}}{(P_\alpha(\mathbf{p}))^{\frac{1}{2\alpha}} \epsilon^{1+\frac{1}{2\alpha}}} + \frac{\sqrt{n}}{(P_\alpha(\mathbf{p}))^{\frac{1}{2\alpha}} \epsilon}\right), \quad (77) \end{aligned}$$

where the third equation comes from

$$\nu_{m_0} = 2^{-m_0+1}\beta_0 = 2^{-\lceil \log(\frac{\beta_0}{\nu_0}) \rceil} \beta_0 = \Theta(\nu_0) = \Theta\left(\left(\frac{2^{2\alpha}(p^*)^\alpha \epsilon}{5n}\right)^{\frac{1}{2\alpha}}\right) = \Theta\left(\frac{\sqrt{p^*} \epsilon^{\frac{1}{2\alpha}}}{n^{\frac{1}{2\alpha}}}\right) \quad (78)$$

and $L = \Theta(p^*)$, and the fourth equation comes from $p^* = \min(\frac{1}{a}P, 1)^\frac{1}{\alpha}$ and $P = \Theta(P_\alpha(\mathbf{p}))$.

In the worst case that $P_\alpha(\mathbf{p}) = n^{1-\alpha}$, the complexity bound becomes $\tilde{\mathcal{O}}\left(\frac{n^{1-\frac{1}{2\alpha}}}{\epsilon} + \frac{\sqrt{n}}{\epsilon^{1+\frac{1}{2\alpha}}}\right)$. \square

Remark 2. Note that [Eq. \(77\)](#) above established a case-dependent bound of estimating $H_\alpha(\mathbf{p})$ given a rough estimation in advance. The requirement of the rough estimation can be removed during the analysis in the next subsection.

4.2 Estimate $H_\alpha(\mathbf{p})$ by annealing

We now apply the annealing method in [Proposition 1](#) to remove the requirement of P, a, b in [Lemma 10](#).

Theorem 7. For any $\alpha > 1$, there exists an algorithm \mathcal{A} such that for any $\delta \in (0, 1)$, $\epsilon \in (0, 1)$ and probability distribution \mathbf{p} on $[n]$, \mathcal{A} can estimate $H_\alpha(\mathbf{p})$ to within additive error ϵ with success probability at least $1 - \delta$ using $\tilde{\mathcal{O}}\left(\frac{n^{1-\frac{1}{2\alpha}}}{\epsilon} + \frac{\sqrt{n}}{\epsilon^{1+\frac{1}{2\alpha}}}\right)$ calls to U_p and U_p^\dagger .

Proof. Let the parameters l and function sequence $\{g_k\}$ be such that $l := \lceil \frac{\ln(\alpha)}{\ln(1+1/\ln(n))} \rceil$ and $g_k(x) := x^{\alpha(1+\frac{1}{\ln(n)})^{k-l}-1}$ for $k = 1, \dots, l$. Note that $g_k(x)$ is monotonically increasing on $[0, 1]$, $xg_k(x) = x^{\alpha(1+\frac{1}{\ln(n)})^{k-l}}$ is convex, $g_k(0) = 0$, and $g_k(1) = 1$ for $k = 1, \dots, l$.

We now prove that Eq. (44) and Eq. (45) hold for $f_k(\mathbf{p}) = \sum_{i=1}^n p_i g_k(p_i) = P_{\alpha(1+\frac{1}{\ln(n)})^{k-l}}(\mathbf{p})$.

First, we introduce the following lemma to connect the value of $f_k(\mathbf{p})$ to $f_{k+1}(\mathbf{p})$.

Lemma 11 ([LW19, Lemma 5.3]). *For any distribution $\mathbf{p} = (p_i)_{i=1}^n$ and $0 < \alpha_1 < \alpha_2$, we have*

$$\left(\sum_{i \in [n]} p_i^{\alpha_2} \right)^{\frac{\alpha_1}{\alpha_2}} \leq \sum_{i \in [n]} p_i^{\alpha_1} \leq n^{1-\frac{\alpha_1}{\alpha_2}} \left(\sum_{i \in [n]} p_i^{\alpha_2} \right)^{\frac{\alpha_1}{\alpha_2}}. \quad (79)$$

Specifically, for $f_{k+1} = P_{\alpha_2}(\mathbf{p})$ and $f_k = P_{\alpha_1}(\mathbf{p})$, we have $\frac{\alpha_1}{\alpha_2} = \frac{1}{1+\frac{1}{\ln(n)}}$, and

$$f_{k+1}^{\frac{1}{1+\frac{1}{\ln(n)}}} \leq f_k \leq n^{\frac{1}{1+\ln(n)}} f_{k+1}^{\frac{1}{1+\frac{1}{\ln(n)}}} \leq e f_{k+1}^{\frac{1}{1+\frac{1}{\ln(n)}}}. \quad (80)$$

Since $\alpha(1+\frac{1}{\ln(n)})^{-l} = \alpha(1+\frac{1}{\ln(n)})^{\lceil \frac{\ln(\alpha)}{\ln(1+1/\ln(n))} \rceil} \leq 1 + \frac{1}{\ln(n)}$, we have $f_1(\mathbf{p}) = P_{\alpha(1+\frac{1}{\ln(n)})^{-l}}(\mathbf{p}) \geq n^{1-\alpha(1+\frac{1}{\ln(n)})^{-l}} \geq n^{1-(1+1/\ln(n))} = \frac{1}{e}$. Then we have

$$\frac{\max_{\mathbf{q} \in \Delta^n} f_1(\mathbf{q})}{\min_{\mathbf{q} \in \Delta^n} f_1(\mathbf{q})} \leq \frac{1}{1/e} = e. \quad (81)$$

For any distributions \mathbf{q}, \mathbf{r} on $[n]$, and $k \in [l-1]$ such that $f_k(\mathbf{r})/f_k(\mathbf{q}) \in [\frac{3}{4}, \frac{5}{4}]$, from Eq. (80), we have

$$f_{k+1}(\mathbf{q}) \geq \left(\frac{1}{e} f_k(\mathbf{q}) \right)^{1+1/\ln(n)} \geq \left(\frac{1}{e} \frac{4}{5} f_k(\mathbf{r}) \right)^{1+1/\ln(n)}, \quad \text{and} \quad (82)$$

$$f_{k+1}(\mathbf{q}) \leq f_k(\mathbf{q})^{1+1/\ln(n)} \leq \left(\frac{4}{3} f_k(\mathbf{r}) \right)^{1+1/\ln(n)}. \quad (83)$$

Therefore, we have

$$\max_{\mathbf{r} \in \Delta^n} \frac{\max_{\mathbf{q} \in \Delta^n, f_k(\mathbf{r})/f_k(\mathbf{q}) \in [\frac{3}{4}, \frac{5}{4}]} f_{k+1}(\mathbf{q})}{\min_{\mathbf{q} \in \Delta^n, f_k(\mathbf{r})/f_k(\mathbf{q}) \in [\frac{3}{4}, \frac{5}{4}]} f_{k+1}(\mathbf{q})} \leq \max_{\mathbf{r} \in \Delta^n} \left(\frac{4}{3} f_k(\mathbf{r}) \right)^{1+1/\ln(n)} = \left(\frac{5e}{3} \right)^{1+1/\ln(n)} \leq 4e^2 \quad (84)$$

for all $k \in [l-1]$.

Setting $\epsilon_k := \frac{1}{4}$ for all $k \in [l-1]$, $\epsilon_l := \min(\frac{1}{2}, \frac{(\alpha-1)\epsilon}{2})$, and $c := 4e^2$ in **Proposition 1**, from Eq. (81) and Eq. (84), we can infer that Eq. (44) and Eq. (45) is satisfied.

For all $k \in [l]$, setting $\alpha := \alpha(1+\frac{1}{\ln(n)})^{k-l}$ in **Lemma 10**, from **Lemma 10**, there exists an algorithm \mathcal{A}_k which can estimate $f_k(\mathbf{p})$ to within multiplicative error ϵ_k with success probability at least $1 - \frac{\delta}{l}$ using $Q_k = \tilde{\mathcal{O}} \left(\frac{n^{1-\frac{1}{2\alpha}}}{\epsilon_k} + \frac{\sqrt{n}}{\epsilon_k^{1+\frac{1}{2\alpha}}} \right)$ calls to U_p and U_p^\dagger . These \mathcal{A}_k satisfy the conditions in **Proposition 1**, so we can construct an algorithm to estimate $P_\alpha(\mathbf{p})$ to within multiplicative error

$\epsilon_l = \Theta(\epsilon)$ using

$$\begin{aligned}
\sum_{k=1}^l Q_k &= \sum_{k=1}^l \tilde{\mathcal{O}} \left(\frac{n^{1 - \frac{1}{2\alpha(1 + \frac{1}{\ln(n)})^{k-l}}}}{\epsilon_k} + \frac{\sqrt{n}}{\epsilon_k^{1 + \frac{1}{2\alpha(1 + \frac{1}{\ln(n)})^{k-l}}}} \right) \\
&= \tilde{\mathcal{O}} \left((l-1) \left(\frac{n^{1 - \frac{1}{2\alpha}}}{1/4} + \frac{\sqrt{n}}{(\frac{1}{4})^{1 + \frac{1}{2\alpha}}} \right) + \frac{n^{1 - \frac{1}{2\alpha}}}{\epsilon} + \frac{\sqrt{n}}{\epsilon^{1 + \frac{1}{2\alpha}}} \right) \\
&= \tilde{\mathcal{O}} \left(\frac{n^{1 - \frac{1}{2\alpha}}}{\epsilon} + \frac{\sqrt{n}}{\epsilon^{1 + \frac{1}{2\alpha}}} \right) \tag{85}
\end{aligned}$$

where the second equation comes from $n^{1 - \frac{1}{2\alpha(1 + \frac{1}{\ln(n)})^{k-l}}} \leq n^{1 - \frac{1}{2\alpha}}$ and $\epsilon_k = \frac{1}{4}$ for all $k \in [l-1]$, and the third equation comes from $l = \mathcal{O}(\ln(\alpha))$ and can be omitted in $\tilde{\mathcal{O}}(\cdot)$.

Denote the estimate of $P_\alpha(\mathbf{p})$ by $\tilde{P}_\alpha(\mathbf{p})$, we have $\frac{\tilde{P}_\alpha(\mathbf{p})}{P_\alpha(\mathbf{p})} \in [1 - \epsilon_l, 1 + \epsilon_l]$. Let $\tilde{H}_\alpha(\mathbf{p}) = \frac{1}{1-\alpha} \log\left(\tilde{P}_\alpha(\mathbf{p})\right)$, then we have

$$|\tilde{H}_\alpha(\mathbf{p}) - H_\alpha(\mathbf{p})| = \frac{1}{\alpha - 1} \left| \log\left(\tilde{P}_\alpha(\mathbf{p})\right) - \log(P_\alpha(\mathbf{p})) \right| = \frac{1}{\alpha - 1} \left| \log\left(\frac{\tilde{P}_\alpha(\mathbf{p})}{P_\alpha(\mathbf{p})}\right) \right| \leq \frac{2}{\alpha - 1} \epsilon_l \leq \epsilon, \tag{86}$$

where the first inequality comes from $|\log(1+x)| \leq 2|x|$ for all $x > -\frac{1}{2}$, so $\tilde{H}_\alpha(\mathbf{p})$ is an estimate of $H_\alpha(\mathbf{p})$ within additive error ϵ . \square

5 Rényi Entropy Estimation ($0 < \alpha < 1$)

5.1 Upper bound

In order to approximate $H_\alpha(\mathbf{p})$ within a given additive error ϵ , we need to approximate $P_\alpha(\mathbf{p})$ within multiplicative error $\mathcal{O}(\epsilon)$. Note that $P_\alpha(\mathbf{p}) = \sum_{i=1}^n p_i^\alpha = \sum_{i=1}^n p_i p_i^{\alpha-1}$ and $\alpha - 1 < 1$ for $\alpha < 1$, so we first construct a series of polynomials S , such that for any constant $d > 0$, there exists polynomial S such that $\sum_{i=1}^n p_i S(\sqrt{p_i})^2$ is an $\mathcal{O}(\epsilon)$ multiplicative approximation of $\frac{(d\epsilon/n)^{\frac{1}{\alpha}-1}}{4} P_\alpha(\mathbf{p})$.

Lemma 12. *For any $\epsilon \in (0, 1)$, $\alpha \in (0, 1)$, and constants $d, d' > 0$, the odd polynomial P' in [Lemma 7](#) with parameters (δ, ϵ, c) to be $\delta := (\frac{d\epsilon}{n})^{\frac{1}{2\alpha}}$, $c := 1 - \alpha$, $\epsilon := d'\epsilon\delta^{2c}$ has $\deg(S) = \tilde{\mathcal{O}}\left(\frac{n^{\frac{1}{2\alpha}}}{\epsilon^{\frac{1}{2\alpha}}}\right)$, and satisfies Eq. (16) in [Theorem 4](#) and*

$$\sum_{i=1}^n \left| p_i S(\sqrt{p_i})^2 - \frac{\delta^{2c}}{4} p_i^\alpha \right| \leq \left(\frac{5}{4}d + 2d' \right) \epsilon \delta^{2c}. \tag{87}$$

Proof. [Lemma 7](#) implies that S is an odd polynomial and $|S(x)| \leq 1$ for all $x \in [-1, 1]$, so S satisfies Eq. (16) in [Theorem 4](#). From [Lemma 7](#), S satisfies that

$$\forall x \in [\delta, 1] : \left| S(x) - \frac{\delta^c}{2} x^{\alpha-1} \right| \leq d'\epsilon\delta^{2c}. \tag{88}$$

From [Lemma 7](#), the degree of S is $\deg(S) = \mathcal{O}\left(\frac{\max[1, c]}{\delta} \log\left(\frac{1}{\epsilon}\right)\right) = \tilde{\mathcal{O}}\left(\frac{n^{\frac{1}{2\alpha}}}{\epsilon^{\frac{1}{2\alpha}}}\right)$.

For i such that $\sqrt{p_i} \leq \delta$, we have

$$\begin{aligned}
\sum_{\sqrt{p_i} \leq \delta} \left| p_i S(\sqrt{p_i})^2 - \frac{\delta^{2c}}{4} p_i^\alpha \right| &\leq \sum_{\sqrt{p_i} \leq \delta} \left(\left| p_i S(\sqrt{p_i})^2 \right| + \left| \frac{\delta^{2c}}{4} p_i^\alpha \right| \right) \\
&\leq \left(\sum_{\sqrt{p_i} \leq \delta} p_i + \frac{\delta^{2c}}{4} p_i^\alpha \right) \\
&\leq \sum_{\sqrt{p_i} \leq \delta} \left(\delta^2 + \frac{\delta^2}{4} \right) \\
&= \sum_{\sqrt{p_i} \leq \delta} \frac{5}{4} \delta^{2\alpha} \delta^{2c} \\
&\leq \frac{5}{4} d \epsilon \delta^{2c}, \tag{89}
\end{aligned}$$

where the second inequality comes from $|S(x)| \leq 1$ for all $x \in [-1, 1]$, the third inequality comes from $\sqrt{p_i} \leq \delta$ and $c = 1 - \alpha$, and the last inequality comes from $n\delta^{2\alpha} = d\epsilon$.

For i such that $\sqrt{p_i} > \delta$, we have

$$\begin{aligned}
\sum_{\sqrt{p_i} > \delta} \left| p_i S(\sqrt{p_i})^2 - \frac{\delta^{2c}}{4} p_i^\alpha \right| &= \sum_{\sqrt{p_i} > \delta} p_i \left| S(\sqrt{p_i}) - \frac{\delta^c}{2} (\sqrt{p_i})^{\alpha-1} \right| \left| S(\sqrt{p_i}) + \frac{\delta^c}{2} (\sqrt{p_i})^{\alpha-1} \right| \\
&\leq 2d' \epsilon \delta^{2c} \sum_{\sqrt{p_i} > \delta} p_i \\
&\leq 2d' \epsilon \delta^{2c}, \tag{90}
\end{aligned}$$

where the first inequality comes from Eq. (88), $|S(x)| \leq 1$ for all $x \in [-1, 1]$, and $\frac{\delta^c}{2} (\sqrt{p_i})^{\alpha-1} \leq 1$ for all $i \in [n]$.

Combining Eq. (5.1) and Eq. (5.1), we have

$$\sum_{i=1}^n \left| p_i S(\sqrt{p_i})^2 - \frac{\delta^{2c}}{4} p_i^\alpha \right| \leq \left(\frac{5}{4} d + 2d' \right) \epsilon \delta^{2c}, \tag{91}$$

which completes the proof. \square

As a result, we can give an algorithm for estimating $H_\alpha(\mathbf{p})$ to within a given additive error ϵ with high probability as follows.

Theorem 8. *For any $\alpha \in (0, 1)$, there exists an algorithm \mathcal{A} such that for any $\delta \in (0, 1)$, $\epsilon \in (0, 1)$ and probability distribution \mathbf{p} on $[n]$, \mathcal{A} can estimate $H_\alpha(\mathbf{p})$ to within additive error ϵ with success probability at least $1 - \delta$ using $\tilde{\mathcal{O}}\left(\frac{n \frac{1}{2\alpha}}{\epsilon \frac{1}{2\alpha+1}}\right)$ calls to U_p and U_p^\dagger .*

Proof. Correctness.

Let $\epsilon_0 = \min\left(\frac{1}{2}, \frac{(1-\alpha)\epsilon}{4}\right)$, $\delta' = \left(\frac{\epsilon_0}{40n}\right)^{\frac{1}{2\alpha}}$. Before constructing S and S_j in Proposition 2, we first define the number of stages of our variable-stopping-time quantum algorithm $m_0 := \lceil \log\left(\frac{1}{\delta'}\right) \rceil + 1$, and $\delta_j = 2^{-j}$ for $j = 1, \dots, m_0 - 1$, $\delta_{m_0} = \delta_{m_0-1} = 2^{-m_0+1}$.

Let S_0 be the polynomial S in Lemma 7 with parameters (δ, ϵ, c) to be $\delta := \delta_{m_0}$, $c := 1 - \alpha$, $\epsilon := \epsilon_0 \delta'^{2c}$.

Let $L := \frac{1}{8}\delta_{m_0}^{2c}$, and we will prove that L is an lower bound of $\sum_{i=1}^n p_i S_0(\sqrt{p_i})^2$ later.

Let P_j for $j = 1, \dots, m_0$ be the odd polynomial P' in [Lemma 7](#) with parameters (δ, ε, c) to be $\delta := \delta_j, c := 1 - \alpha, \varepsilon := \frac{1}{64}\delta_{m_0}^{2c}\epsilon_0$. From [Lemma 7](#), we have

$$\left|P_j(x) - \frac{\delta_j^c}{2}x^{\alpha-1}\right| \leq \frac{1}{64}\delta_{m_0}^{2c}\epsilon_0 \text{ for any } x \in [\delta_j, 1], \quad (92)$$

and $\deg(P_j) = \mathcal{O}\left(\frac{\max[1, c]}{\delta} \log\left(\frac{1}{\varepsilon}\right)\right) = \tilde{\mathcal{O}}(2^j)$.

Now we set the parameters $(S, \beta, L, m, \{S_j \mid j \in [m]\})$ of [Algorithm 3](#) to be $S := S_0, \beta := 1, L := L, m := m_0, S_j := \left(\frac{\delta_{m_0}}{\delta_j}\right)^c P_j$ for $j = 1, \dots, m_0$, and then prove that these parameters satisfy the conditions in [Proposition 2](#).

- For β , we have $\sqrt{p_i} \leq 1 = \beta$ for all $i \in [n]$.
- For L , we need to prove that is is a lower bound of $\sum_{i=1}^n p_i S_0(\sqrt{p_i})^2$. Note that $\delta_{m_0} = 2^{-m_0+1} = 2^{-\lceil \log(\frac{1}{\delta'}) \rceil} \leq \delta' \leq \left(\frac{\epsilon_0}{40n}\right)^{\frac{1}{2\alpha}}$. Let the constants d, d' in [Lemma 12](#) be such that $d = \delta_{m_0}^{2\alpha} \frac{n}{\epsilon_0} \leq \frac{1}{40}, d' = \frac{1}{64}$, then the parameters $(\delta = \delta_{m_0} = \left(\frac{d\epsilon_0}{n}\right)^{\frac{1}{2\alpha}}, c = 1 - \alpha, \varepsilon = \frac{1}{64}\epsilon_0\delta_{m_0}^{2c} = d'\epsilon_0\delta_{m_0}^{2c})$ of S_0 satisfy the conditions in [Lemma 12](#). From [Lemma 12](#), we have

$$\sum_{i=1}^n \left|p_i S_0(\sqrt{p_i})^2 - \frac{\delta_{m_0}^{2c}}{4} p_i^\alpha\right| \leq \left(\frac{5}{4}d + 2d'\right)\epsilon_0\delta_{m_0}^{2c} \leq \frac{1}{16}\epsilon_0\delta_{m_0}^{2c} = \frac{1}{2}L\epsilon_0. \quad (93)$$

From Eq. (93), we have

$$\begin{aligned} \sum_{i=1}^n p_i S_0(\sqrt{p_i})^2 &\geq \frac{\delta_{m_0}^{2c}}{4} P_\alpha(\mathbf{p}) - \frac{1}{16}\epsilon_0\delta_{m_0}^{2c} \\ &\geq \frac{1}{4}\delta_{m_0}^{2c} - \frac{1}{16}\delta_{m_0}^{2c} \\ &\geq \frac{1}{8}\delta_{m_0}^{2c} = L, \end{aligned} \quad (94)$$

where the second inequality comes from $P^\alpha(\mathbf{p}) \geq 1$ for $\alpha \in (0, 1)$, and $\epsilon_0 \in (0, 1)$.

- For S_j , they are odd polynomial and satisfy $|S_j(x)| \leq 1$ for all $x \in [-1, 1]$, which meet the requirements in [Theorem 4](#). Note that the parameters of S_0 in [Lemma 7](#) is the same as the parameters of S_{m_0} , so we have $S_0 = S_{m_0}$. Then we have

$$\begin{aligned} \text{for any } x \in [\delta_j, 1] : |S_j(x) - S_0(x)| &= |S_j(x) - S_{m_0}(x)| \\ &\leq \left| \left(\frac{\delta_{m_0}}{\delta_j}\right)^c P_j(x) - \frac{\delta_{m_0}^c}{2} x^{\alpha-1} \right| + \left| P_{m_0}(x) - \frac{\delta_{m_0}^c}{2} x^{\alpha-1} \right| \\ &\leq \left(\frac{\delta_{m_0}}{\delta_j}\right)^c \left| P_j(x) - \frac{\delta_j^c}{2} x^{\alpha-1} \right| + \left| P_{m_0}(x) - \frac{\delta_{m_0}^c}{2} x^{\alpha-1} \right| \\ &\leq \left(1 + \left(\frac{\delta_{m_0}}{\delta_j}\right)^c\right) \left(\frac{1}{64}\delta_{m_0}^{2c}\epsilon_0\right) \leq L\epsilon_0, \end{aligned} \quad (95)$$

where the third inequality comes from Eq. (92) and $\delta_{m_0} < \delta_j$ for $j < m_0$. From Eq. (95), we can infer that for any $x \in [2^{-j}, 1] : |S_j(x) - S_0(x)| \leq L\epsilon_0$, which meets the requirements of Eq. (50).

Therefore, the parameters we set are valid for [Proposition 2](#), so [Algorithm 3](#) with the same parameters and input (ϵ_0, δ) can estimate $\sum_{i=1}^n p_i S_0(\sqrt{p_i})^2$ to within multiplicative error ϵ_0 within success probability at least $1 - \delta$. Denote the estimate by \tilde{p} , we have

$$\begin{aligned}
|4\delta_{m_0}^{-2c}\tilde{p} - P_\alpha(\mathbf{p})| &\leq |4\delta_{m_0}^{-2c}\tilde{p} - 4\delta_{m_0}^{-2c}\sum_{i=1}^n p_i S_0(\sqrt{p_i})^2| + |4\delta_{m_0}^{-2c}\sum_{i=1}^n p_i S_0(\sqrt{p_i})^2 - P_\alpha(\mathbf{p})| \\
&\leq 4\delta_{m_0}^{-2c}\left(\sum_{i=1}^n p_i S_0(\sqrt{p_i})^2\right)\epsilon_0 + 2\delta_{m_0}^{-2c}L\epsilon_0 \\
&\leq 4\delta_{m_0}^{-2c}\left(\frac{\delta_{m_0}^{2c}}{4}P_\alpha(\mathbf{p})\epsilon_0 + \frac{1}{2}L\epsilon_0^2\right) + \frac{1}{4}\epsilon_0 \\
&= P_\alpha(\mathbf{p})\epsilon_0 + \frac{1}{4}(\epsilon_0^2 + \epsilon_0) \leq 2P_\alpha(\mathbf{p})\epsilon_0, \tag{96}
\end{aligned}$$

with success probability at least $1 - \delta$, where the second inequality and the third inequality come from Eq. (93), and the last inequality comes from $P_\alpha \geq 1$ for any $\alpha \in (0, 1)$. Let $\tilde{P}_\alpha(\mathbf{p}) := 4\delta_{m_0}^{-2c}\tilde{p}$, and then we have $\frac{\tilde{P}_\alpha(\mathbf{p})}{P_\alpha(\mathbf{p})} \in [1 - 2\epsilon_0, 1 + 2\epsilon_0]$. Let $\tilde{H}_\alpha(\mathbf{p}) = \frac{1}{1-\alpha} \log\left(\frac{\tilde{P}_\alpha(\mathbf{p})}{P_\alpha(\mathbf{p})}\right)$, then we have

$$|\tilde{H}_\alpha(\mathbf{p}) - H_\alpha(\mathbf{p})| = \frac{1}{1-\alpha} \left(\log\left(\frac{\tilde{P}_\alpha(\mathbf{p})}{P_\alpha(\mathbf{p})}\right) - \log\left(P_\alpha(\mathbf{p})\right) \right) = \frac{1}{1-\alpha} \left| \log\left(\frac{\tilde{P}_\alpha(\mathbf{p})}{P_\alpha(\mathbf{p})}\right) \right| \leq \frac{4}{1-\alpha}\epsilon_0 = \epsilon, \tag{97}$$

where the first inequality comes from $|\log(1+x)| \leq 2|x|$ for all $x > -\frac{1}{2}$.

Complexity. Now we compute the query complexity of the above algorithm. First, let us compute t_j defined in [Proposition 2](#). For t_j , we have

$$t_j = 2^j \log\left(\frac{m_0}{\epsilon L}\right) + \sum_{k=1}^j \deg(S_k) = \tilde{\mathcal{O}}\left(2^j + \sum_{k=1}^j 2^k\right) = \tilde{\mathcal{O}}(2^j), \tag{98}$$

where the second equation comes from $\deg(S_j) = \deg(P_j) = \tilde{\mathcal{O}}(2^j)$.

Let $\varphi_j = 2^{-j}$ for $j = 1, \dots, m_0 - 1$ and $\varphi_{m_0} = 0$ following the definition in [Proposition 2](#). From [Proposition 2](#), the query complexity of the above algorithm is

$$\begin{aligned}
&\tilde{\mathcal{O}}\left(\frac{1}{\epsilon} \left(t_{m_0} + \sqrt{\epsilon} \sum_{j=1}^{m_0} t_j + \frac{\sqrt{\sum_{j=1}^{m_0} \sum_{i: \sqrt{p_i} \in [\varphi_j, \varphi_{j-1})} p_i t_{j+1}^2}}{\sqrt{\sum_{i=1}^n p_i S(\sqrt{p_i})^2}} \right)\right) \\
&= \tilde{\mathcal{O}}\left(\frac{1}{\epsilon} \left(2^{m_0} + \frac{\sqrt{\sum_{j=1}^{m_0-1} (|\{i : 2^{-j} \leq \sqrt{p_i} \leq 2^{-j+1}\}|(2^{-j+1})^2(2^{j+1})^2 + \sum_{\sqrt{p_i} \leq \varphi_{m_0-1}} p_i (\frac{1}{\varphi_{m_0-1}})^2)}}{\sqrt{L}} \right)\right) \\
&= \tilde{\mathcal{O}}\left(\frac{1}{\epsilon} \left(\frac{1}{\delta_{m_0}} + \frac{\sqrt{n+n}}{\delta_{m_0}^c} \right)\right) \\
&= \tilde{\mathcal{O}}\left(\frac{1}{\epsilon} \left(\frac{n^{\frac{1}{2\alpha}}}{\epsilon^{\frac{1}{2\alpha}}} + \frac{\sqrt{n}}{\left(\frac{\epsilon}{n}\right)^{(1-\alpha)/2\alpha}} \right)\right) \\
&= \tilde{\mathcal{O}}\left(\frac{n^{\frac{1}{2\alpha}}}{\epsilon^{\frac{1}{2\alpha}+1}} + \frac{n^{\frac{1}{2\alpha}}}{\epsilon^{\frac{1}{2\alpha}+\frac{1}{2}}}\right) = \tilde{\mathcal{O}}\left(\frac{n^{\frac{1}{2\alpha}}}{\epsilon^{\frac{1}{2\alpha}+1}}\right), \tag{99}
\end{aligned}$$

which completes the proof. \square

5.2 Lower bound

The Hellinger distance between two discrete probability distributions p and q is defined as $d_H(\mathbf{p}, \mathbf{q}) := \sqrt{\sum_{i=1}^n (\sqrt{p_i} - \sqrt{q_i})^2} / 2$. In [Bel19], they give a lower bound for the query complexity of distinguishing two distributions as follows.

Lemma 13 ([Bel19, Claim 5]). *Quantum query complexity of distinguishing probability distributions \mathbf{p} and \mathbf{q} with pure-state preparation oracle in Definition 1 is $\Theta(1/d_H(\mathbf{p}, \mathbf{q}))$.*

Then we can give the following lower bound for estimating $H_\alpha(\mathbf{p})$ with pure-state preparation oracle U_p and U_p^\dagger .

Theorem 9. *For any constant $\alpha \in (0, 1)$, $n \geq 1 + 2^{1/(1-\alpha)}$, and $\epsilon \in (0, \frac{1}{2})$, any algorithm that can estimate $H_\alpha(\mathbf{p})$ to within additive error ϵ needs at least $\Omega\left(\frac{n^{1/2\alpha-1/2}}{\epsilon^{1/2\alpha}}\right)$ calls to U_p and U_p^\dagger .*

Proof. For any $\epsilon \in (0, \frac{1}{2})$, $n > 1 + 2^{1/(1-\alpha)}$ and $\alpha \in (0, 1)$, let $\delta = \left(\frac{4\epsilon}{(n-1)^{1-\alpha}}\right)^{\frac{1}{\alpha}} < 1$.

Consider $\mathbf{p} = \left(1 - \delta, \frac{\delta}{n-1}, \dots, \frac{\delta}{n-1}\right)$ and $\mathbf{q} = (1, 0, \dots, 0)$. The Hellinger distance of \mathbf{p} and \mathbf{q} is

$$d_H(\mathbf{p}, \mathbf{q}) = \sqrt{\frac{1}{2} \left((\sqrt{1-\delta} - 1)^2 + (n-1) \left(\sqrt{\frac{\delta}{n-1}} \right)^2 \right)} = \sqrt{\frac{1}{2} (\Theta(\delta^2) + \delta)} = \Theta(\sqrt{\delta}) \quad (100)$$

as $\delta \rightarrow 0$, where the second equation comes from $\sqrt{1-x} = 1 - \Theta(x)$ as $x \rightarrow 0$. By Lemma 13, we need $\Omega\left(\frac{1}{\sqrt{\delta}}\right)$ calls to U_p and U_p^\dagger to distinguish \mathbf{p} and \mathbf{q} .

Then we have

$$\begin{aligned} |H_\alpha(\mathbf{q}) - H_\alpha(\mathbf{p})| &= \left| \frac{1}{1-\alpha} \log((1-\delta)^\alpha + \delta^\alpha (n-1)^{1-\alpha}) - 0 \right| \\ &\geq \left| \frac{1}{1-\alpha} \log(1 - \delta + \delta^\alpha (n-1)^{1-\alpha}) \right| \\ &= \left| \frac{1}{1-\alpha} \log(1 - \delta + 4\epsilon) \right| \\ &\geq \left| \frac{1}{1-\alpha} \log(1 + 2\epsilon) \right| \\ &\geq \frac{2\epsilon}{(1-\alpha)} \geq 2\epsilon, \end{aligned} \quad (101)$$

where the first inequality is because $(1-x)^\alpha \geq 1-x$ as for $x \in (0, 1)$ for any $\alpha \in (0, 1)$, the second inequality is because $\delta \leq \frac{4\epsilon}{(n-1)^{1-\alpha}} \leq 2\epsilon$, and the third inequality is because $\log(1+x) \geq x$ for $x \in (0, 1)$.

If we can estimate Rényi entropy of \mathbf{p} and \mathbf{q} to within additive error ϵ , we can distinguish distributions \mathbf{p} and \mathbf{q} , which needs $\Omega(1/\sqrt{\delta})$ queries as proven above. Therefore, it requires $\Omega\left(\frac{1}{\sqrt{\delta}}\right) = \Omega\left(\frac{n^{1/2\alpha-1/2}}{\epsilon^{1/2\alpha}}\right)$ queries to U_p and U_p^\dagger . \square

5.3 More discussions about ϵ dependence

Note that Belovs [Bel19] proved that the lower bound in Lemma 13 of distinguishing probability distributions also holds with the oracle in Definition 2. As a result, our lower bound $\Omega\left(\frac{n^{1/2\alpha-1/2}}{\epsilon^{1/2\alpha}}\right)$ in Theorem 9 also holds with this oracle. However, a contradiction can be observed between the ϵ dependency of this lower bound and that of the upper bound in [LW19] which uses $\tilde{\mathcal{O}}\left(\frac{n^{1/\alpha-1/2}}{\epsilon^2}\right)$ calls to the oracles in Definition 2 and outputs an estimate of $H_\alpha(\mathbf{p})$ for $\mathbf{p} \in \Delta^n$ to within additive error ϵ .

We suspect that there is an issue with Eq. (V.8) in the journal version of [LW19], where the Taylor approximation by $\frac{l}{2^m} = \Theta(\theta_i)$ therein is not precise when $\theta_i = \Theta(\frac{1}{2^m})$. Here we give a corrected analysis of the bias of the α -Rényi entropy estimator in [LW19] when $\alpha \in (0, \frac{1}{2})$. Following the notation in [LW19], for $\alpha \in (0, \frac{1}{2})$ and each $i \in S_{j+1}$, in order to reduce the Taylor approximation error, we need to treat $l \in (2^{m-1}\theta_i, 2^m\theta_i)$ specially. Here we take $\theta_i = 2^j/2^m$ for simplicity, but the following equations hold for general θ_i when $i \in S_{j+1}$:

$$\begin{aligned}
& p_i \mathbb{E} [|\tilde{p}_i^{\alpha-1} - p_i^{\alpha-1}|] \\
&= \mathcal{O} \left(\left(\frac{2^j}{2^m} \pi \right)^2 \left(\sum_{l=-(2^m-2^j), l \neq 0}^{2^j-1} + \sum_{l=2^j-1}^{2^j} \right) \left(\frac{1}{l^2} \left| \sin \left(\left(\theta_i - \frac{l}{2^m} \right) \pi \right) \right|^{2(\alpha-1)} - \left| \sin(\theta_i \pi) \right|^{2(\alpha-1)} \right) \right) \\
&= \mathcal{O} \left(\left(\frac{2^j}{2^m} \pi \right)^2 \sum_{l=-(2^m-2^j), l \neq 0}^{2^j-1} \frac{1}{l^2} \frac{|l|}{2^m} \left(\frac{2^j}{2^m} \pi \right)^{2\alpha-3} + \left(\frac{2^j}{2^m} \pi \right)^2 \sum_{l=2^j-1}^{2^j-1} 2^{-2j} \left| \sin \left(\left(\theta_i - \frac{l}{2^m} \right) \pi \right) \right|^{2\alpha-2} \right) \\
&= \mathcal{O} \left(\frac{m}{2^{2\alpha m}} 2^{(2\alpha-1)j} + \sum_{r=1}^{2^j-1} \frac{1}{2^{2m}} \left(\frac{r}{2^m} \right)^{2\alpha-2} \right) \\
&= \mathcal{O} \left(\frac{m}{2^{2\alpha m}} 2^{(2\alpha-1)j} + \frac{m}{2^{2\alpha m}} \right), \tag{102}
\end{aligned}$$

where the second equation comes from

$$\left| \left(\sin \left(\left(\theta_i - \frac{l}{2^m} \right) \pi \right) \right)^{2(\alpha-1)} - \left(\sin(\theta_i \pi) \right)^{2(\alpha-1)} \right| \leq c \frac{|l|}{2^m} (\theta_i \pi)^{2\alpha-3} \tag{103}$$

for $\frac{l}{2^m \theta_i} \leq \frac{1}{2}$, and we replace $2^j - l$ with r in the third equation. Note that the first term of Eq. (102) is the same as equation Eq. (V.46) in the journal version of [LW19], but it is smaller than the second term, so we only need to set $m = \lceil \frac{1}{2\alpha} \log(\frac{\epsilon}{n} \log(\frac{\epsilon}{n})) \rceil$ so that

$$\sum_{i=1}^n p_i \mathbb{E} [|\tilde{p}_i^{\alpha-1} - p_i^{\alpha-1}|] = \mathcal{O} \left(\frac{nm}{2^{2\alpha m}} \right) \tag{104}$$

is bounded by ϵ . Therefore, the overall complexity of the algorithm in [LW19] used to estimate α -Rényi entropy when $\alpha \in (0, \frac{1}{2})$ is

$$\tilde{\mathcal{O}} \left(\frac{\sqrt{n^{\frac{1}{\alpha}-1}}}{\epsilon} \left(\frac{n}{\epsilon} \right)^{\frac{1}{2\alpha}} \right) = \tilde{\mathcal{O}} \left(\frac{n^{\frac{1}{\alpha}-\frac{1}{2}}}{\epsilon^{1+\frac{1}{2\alpha}}} \right), \tag{105}$$

which has the same dependence on ϵ as that in $\tilde{\mathcal{O}}\left(\frac{1}{\epsilon^{1+\frac{1}{2\alpha}}}\right)$ in our algorithm (and worse dependence in n than that in our algorithm).

We also find that there might be an issue with the ϵ dependency of the classical upper bound $\mathcal{O}\left(\frac{n^{1/\alpha}}{\epsilon^{1/\alpha} \log n}\right)$ on estimating Rényi entropy when $\alpha < 1$ in [AOST16]. Specifically, we suspect that the last two $o(1)$ terms in Eq. (15) and Eq. (18) of the arXiv version of [AOST16] are omitted, but according to Lemma 8, these two terms cannot be omitted unless they are $o(\delta)$. This might increase the order of ϵ in the current classical upper bound.

6 Applications

6.1 Extension to quantum entropies

We first extend our algorithm to work with purified quantum query-access in Definition 3. In [GL20] by Gilyén and Li, they build a block encoding of square root of classical probability distribution using purified quantum query-access oracle as follows. Let

$$U := (U_p \otimes I), \Pi := \left(\sum_{i=1}^n I \otimes |i\rangle\langle i| \otimes |i\rangle\langle i| \right), \text{ and } \tilde{\Pi} := (|\mathbf{0}\rangle\langle \mathbf{0}| \otimes |\mathbf{0}\rangle\langle \mathbf{0}| \otimes I). \quad (106)$$

Then we have

$$A = \Pi U \tilde{\Pi} = \sum_{i=1}^n \sqrt{p_i} |\phi_i\rangle \langle \mathbf{0}| \otimes |i\rangle \langle \mathbf{0}| \otimes |i\rangle \langle i|. \quad (107)$$

The only difference this new oracle brings is that the state we obtain may have some garbage states added, so we need some ancilla registers to store them. For example, following the process in Section 3.1 with purified quantum query-access oracle, we can get a quantum state $|\Psi_p\rangle$ such that

$$|\Psi_p\rangle = \sum_{i=1}^n \sqrt{p_i} S(\sqrt{p_i}) |i\rangle_A |i\rangle_B |+\rangle_Q |\phi_{\text{garbage}}^{(i)}\rangle |1\rangle_F + |\psi_{\text{garbage}}\rangle |0\rangle_F, \quad (108)$$

where $|\phi_{\text{garbage}}^{(i)}\rangle = |\phi_i\rangle |\phi_i\rangle$ is brought by the new oracle. Therefore, we can still use the amplitude estimate algorithm to estimate the amplitude of $|1\rangle_F$, which gives us an estimate of $\sum_{i=1}^n p_i S(\sqrt{p_i})^2$. The framework in Section 3.2 and Section 3.3 also works well with purified quantum query-access oracle for the same reason.

Following [GL20], we can also build a block encoding of $\sqrt{\frac{\rho}{n}}$ for a density operator ρ using purified quantum query-access U_ρ as follows. Let

$$U' := \left(I \otimes U_\rho^\dagger \right) (W \otimes I), \Pi' := (I \otimes |\mathbf{0}\rangle\langle \mathbf{0}| \otimes |\mathbf{0}\rangle\langle \mathbf{0}|), \text{ and } \tilde{\Pi} := (|\mathbf{0}\rangle\langle \mathbf{0}| \otimes |\mathbf{0}\rangle\langle \mathbf{0}| \otimes I), \quad (109)$$

where W is a unitary mapping $|\mathbf{0}\rangle|\mathbf{0}\rangle \mapsto \sum_{j=1}^n \frac{|j\rangle|j\rangle}{\sqrt{n}}$ and $\sum_{j=1}^n \frac{|\phi'_j\rangle|\phi_j\rangle}{\sqrt{n}} = \sum_{j=1}^n \frac{|j\rangle|j\rangle}{\sqrt{n}}$ is the Schmidt decomposition of the maximally entangled state under the basis $\{|\phi_1\rangle, \dots, |\phi_n\rangle\}$. Then we can prove the following theorems which give an algorithm to estimate the quantum Rényi entropy of density operators

$$H_\alpha(\rho) = \frac{1}{1-\alpha} \log(\text{Tr}(\rho^\alpha)), \quad (110)$$

with the purified quantum query-access oracle in Definition 3.

Corollary 2. For any $\alpha > 1$, there exists an algorithm \mathcal{A} such that for any $\delta \in (0, 1)$, $\epsilon \in (0, 1)$ and density operator $\rho \in \mathbb{C}^{n \times n}$, \mathcal{A} can estimate $H_\alpha(\rho)$ to within additive error ϵ with success probability at least $1 - \delta$ using $\tilde{\mathcal{O}}\left(\frac{n^{\frac{3}{2}-\frac{1}{2\alpha}}}{\epsilon} + \frac{n}{\epsilon^{1+\frac{1}{2\alpha}}}\right)$ calls to U_ρ and U_ρ^\dagger .

Proof. Let p_i be the eigenvalues of ρ . We use the block encoding of $\sqrt{\rho/n}$ in Eq. (109) and follow the same process in Theorem 7 to estimate $\text{Tr}(\rho^\alpha)$ by quantum singular value transformation and variable time amplitude estimation. The only difference is that we need to replace the polynomials $S_j(x)$ in Lemma 10 with $S'_j(x)$, where S_j is defined as follows. Note that S_j for all $i = 0, \dots, m_0$ in Lemma 10 is constructed using Lemma 6, so let $(c, \nu_j, \beta_j, \eta)$ be the parameters of S_j in Lemma 6. Let S'_j be the polynomial constructed in Lemma 6 with the parameters (c, β, ν, η) to be $c := c$, $\nu := \frac{\nu_j}{\sqrt{n}}$, $\beta := \frac{\beta_j}{\sqrt{n}}$, $\eta := \eta$. Then we can infer that $S'_j(\frac{\sqrt{p_i}}{\sqrt{n}})$ has the same behavior as $S_j(\sqrt{p_i})$ for all p_i with \sqrt{n} times larger degree, so $\sum_{i=1}^n \frac{p_i}{n} S'_0(\sqrt{\frac{p_i}{n}})^2$ is also an estimate of $P_\alpha(\mathbf{p})/n$ to within multiplicative error ϵ . Therefore, we can obtain an estimate of $H_\alpha(\mathbf{p})$ to within additive error ϵ by rescaling the multiplicative error ϵ to $c\epsilon$ for some constant c . Following the same proof in Lemma 10 and Theorem 7 with $\nu'_{m_0} = \nu_{m_0}/\sqrt{n}$ and $\sum_{i=1}^n \frac{p_i}{n} S'_0(\sqrt{\frac{p_i}{n}})^2 = \Theta(\frac{1}{n} \sum_{i=1}^n p_i S_0(\sqrt{p_i})^2)$, the query complexity becomes

$$\tilde{\mathcal{O}}\left(\frac{n^{1+\frac{1}{2}-\frac{1}{2\alpha}}}{\epsilon} + \frac{\sqrt{n}\sqrt{n}}{\epsilon^{1+\frac{1}{2\alpha}}}\right) = \tilde{\mathcal{O}}\left(\frac{n^{\frac{3}{2}-\frac{1}{2\alpha}}}{\epsilon} + \frac{n}{\epsilon^{1+\frac{1}{2\alpha}}}\right). \quad (111)$$

□

Corollary 3. For any $\alpha \in (0, 1)$, there exists an algorithm \mathcal{A} such that for any $\delta \in (0, 1)$, $\epsilon \in (0, 1)$ and density operator $\rho \in \mathbb{C}^{n \times n}$, \mathcal{A} can estimate $H_\alpha(\rho)$ to within additive error ϵ with success probability at least $1 - \delta$ using $\tilde{\mathcal{O}}\left(\frac{n^{\frac{1}{2\alpha}+\frac{1}{2}}}{\epsilon^{\frac{1}{2\alpha}+1}}\right)$ calls to U_ρ and U_ρ^\dagger .

Proof. Let p_i be the eigenvalues of ρ . The proof is essentially the same as that of Corollary 2. We can construct $S'_j(x)$ with \sqrt{n} times larger degree than S_j in Theorem 8 such that $S'_j(\sqrt{\frac{p_i}{n}})$ has the same behavior as $S_j(\sqrt{p_i})$ for all p_i . With the block encoding in Eq. (109), we can follow the process in Theorem 8 to give an algorithm estimating $H_\alpha(\rho)$ to within additive error ϵ with success probability at least $1 - \delta$ using $\tilde{\mathcal{O}}\left(\frac{n^{\frac{1}{2\alpha}+\frac{1}{2}}}{\epsilon^{\frac{1}{2\alpha}+1}}\right)$ calls to U_ρ and U_ρ^\dagger . □

6.2 Low-rank cases

For low-rank quantum distributions (density matrices) or classical distributions with at most r elements with positive probability, we can adapt our algorithm to obtain better query complexity upper bound.

Quantum distributions. If the rank of the density operator $\rho \in \mathbb{C}^{n \times n}$ is guaranteed to be $r = o(n)$, we can apply our framework to estimate the Rényi entropy of ρ with $\text{poly}(r) = o(n)$ calls to U_ρ and U_ρ^\dagger .

According to the following lemma in [LC19], we can construct a unitary block-encoding of ρ using one call to U_ρ and U_ρ^\dagger .

Lemma 14 ([LC19, Lemma 7]). *Let A, B, C be three $\lceil \log n \rceil$ -qubit registers and the oracle U_ρ in Definition 3 prepare a state $U_\rho|\mathbf{0}\rangle_A|\mathbf{0}\rangle_B = \sum_{i=1}^n \sqrt{p_i}|\phi_i\rangle_A|\psi_i\rangle_B$. Let $S_{B,C}$ be a unitary operator which swaps the states in registers B and C . Let $\tilde{U}_\rho = (U_\rho^\dagger \otimes I_C)S_{B,C}(U_\rho \otimes I_C)$ be a unitary operator on A, B, C , and then we have $\langle \mathbf{0}|_{A,B}U|\mathbf{0}\rangle_{A,B} = \sum_{i=1}^n p_i|\psi_i\rangle\langle\psi_i|_C = \rho$.*

For any polynomial S satisfying Eq. (16) in Theorem 4, we can apply the singular value transformed unitary of \tilde{U}_ρ in Lemma 14 to $\sum_{i=1}^n \sqrt{p_i}|\mathbf{0}\rangle_{A,B}|\psi_i\rangle_C|\phi_i\rangle_D$ which outputs

$$\sum_{i=1}^n \sqrt{p_i}S(p_i)|\mathbf{0}\rangle_{A,B}|\psi_i\rangle_C|\phi_i\rangle_D + |\psi_\perp\rangle, \quad (112)$$

where $\|(\langle \mathbf{0}|_{A,B} \otimes I_{C,D})|\psi_\perp\rangle\| = 0$.

Note that this process is similar to our application of QSVT in Section 3.1 except that we have $S(p_i)$ now rather than $S(\sqrt{p_i})$ in Section 3.1. Therefore, we can use the techniques in our framework to estimate Rényi entropy with some minor changes to the transformation polynomials.

Corollary 4. *For any $\alpha \in (0, 1)$, there exists an algorithm \mathcal{A} such that for any $\delta \in (0, 1)$, $\epsilon \in (0, 1)$, and rank- r density operator $\rho \in \mathbb{C}^{n \times n}$, \mathcal{A} can estimate $H_\alpha(\rho)$ to within additive error ϵ with success probability at least $1 - \delta$ using $\tilde{\mathcal{O}}\left(\frac{r^{\frac{1}{\alpha}}}{\epsilon^{\frac{1}{\alpha}+1}}\right)$ calls to U_ρ and U_ρ^\dagger .*

Proof. For any $0 < \alpha < 1$, we need to change all n with r and the parameters (δ, c) of S_j for $j = 0, \dots, m_0$ in Theorem 8 to $\delta' = \delta^2$, $c' = \frac{c}{2}$ so that the new polynomial $S'_j(p_i)$ has the same behavior as $S_j(\sqrt{p_i})$ for all p_i . Then following the same proof as Theorem 8, we can obtain an upper bound on the quantum query complexity of estimating $H_\alpha(\rho)$ to within additive error ϵ with purified quantum query-access to a rank- r density operator ρ as follows.

$$\begin{aligned} \tilde{\mathcal{O}}\left(\frac{1}{\epsilon}\left(\frac{1}{\delta'_{m_0}} + \frac{\sqrt{r+r}}{\delta'_{m_0}c'}\right)\right) &= \tilde{\mathcal{O}}\left(\frac{1}{\epsilon}\left(\frac{r^{\frac{1}{\alpha}}}{\epsilon^{\frac{1}{\alpha}}} + \frac{\sqrt{r}}{\left(\frac{\epsilon}{r}\right)^{\frac{1}{2}(1-\alpha)/\alpha}}\right)\right) \\ &= \tilde{\mathcal{O}}\left(\frac{r^{\frac{1}{\alpha}}}{\epsilon^{\frac{1}{\alpha}+1}} + \frac{r^{\frac{1}{2\alpha}}}{\epsilon^{\frac{1}{\alpha}+\frac{1}{2}}}\right) = \tilde{\mathcal{O}}\left(\frac{r^{\frac{1}{\alpha}}}{\epsilon^{\frac{1}{\alpha}+1}}\right). \end{aligned} \quad (113)$$

□

Corollary 5. *For any $\alpha > 1$, there exists an algorithm \mathcal{A} such that for any $\delta \in (0, 1)$, $\epsilon \in (0, 1)$, and rank- r density operator $\rho \in \mathbb{C}^{n \times n}$, \mathcal{A} can estimate $H_\alpha(\rho)$ to within additive error ϵ with success probability at least $1 - \delta$ using $\tilde{\mathcal{O}}\left(\frac{r}{\epsilon^{1+\frac{1}{\alpha}}} + \frac{r^{1-\frac{1}{2\alpha}}}{\epsilon}\right)$ calls to U_ρ and U_ρ^\dagger .*

Proof. For any $\alpha > 1$, we need to change all n to r and the parameters (ν, c, β) of S_j for $j = 0, \dots, m_0$ in Lemma 10 to $\nu' = \nu^2$, $c' = \frac{c}{2}$, $\beta' = \beta^2$ so that the new polynomial $S'_j(p_i)$ has the same behavior as $2^{\frac{5}{2}}S_j(\sqrt{p_i})$ for all p_i . The constant $2^{\frac{5}{2}}$ can be omitted in complexity analysis. Following the same proof in Lemma 10 and Theorem 7, we can infer that the upper bound is

$$\tilde{\mathcal{O}}\left(\frac{1}{\epsilon}\left(\frac{1}{\nu'_{m_0}} + \frac{\sqrt{r+r}}{\sqrt{L}}\right)\right) = \tilde{\mathcal{O}}\left(\frac{1}{\epsilon}\left(\frac{r^{\frac{1}{\alpha}}}{p^*\epsilon^{\frac{1}{\alpha}}} + \frac{\sqrt{r}}{\sqrt{p^*}}\right)\right) = \tilde{\mathcal{O}}\left(\frac{r^{\frac{1}{\alpha}}}{(P_\alpha(\mathbf{p}))^{\frac{1}{\alpha}}\epsilon^{1+\frac{1}{\alpha}}} + \frac{\sqrt{r}}{(P_\alpha(\mathbf{p}))^{\frac{1}{2\alpha}}\epsilon}\right), \quad (114)$$

which becomes $\tilde{\mathcal{O}}\left(\frac{r}{\epsilon^{1+\frac{1}{\alpha}}} + \frac{r^{1-\frac{1}{2\alpha}}}{\epsilon}\right)$ in the worst case. □

Classical distributions. For classical distributions, an analogy of low rank density operators is probability distributions \mathbf{p} on $[n]$ such that there are at most r elements i whose probability $p_i > 0$. For such probability distributions, if we know r in advance, we can directly obtain the upper bound in [Theorem 7](#) and [Theorem 8](#) replacing n with r , since the proofs of these two theorem still hold if we replace all n with r .

If we do not know r , we can also use the algorithm in the following corollary to estimate $H_\alpha(\mathbf{p})$ for any $\alpha > 1$.

Corollary 6. *For any $\alpha > 1$, there exists an algorithm \mathcal{A} such that for any $\delta \in (0, 1)$, $\epsilon \in (0, 1)$ and probability distribution \mathbf{p} on $[n]$ with at most r elements having $p_i > 0$, \mathcal{A} can estimate $H_\alpha(\mathbf{p})$ to within additive error ϵ with success probability at least $1 - \delta$ using $\tilde{\mathcal{O}}\left(\frac{r^{1-\frac{1}{2\alpha}}}{\epsilon} + \frac{\sqrt{r}}{\epsilon^{1+\frac{1}{2\alpha-2}}}\right)$ calls to U_p and U_p^\dagger .*

Proof. Changing the parameter ν to $\nu = \Theta((\epsilon P)^{\frac{1}{2\alpha-2}})$ in [Lemma 9](#), we can replace Eq. (67) with

$$\begin{aligned}
\sum_{\sqrt{p_i} \leq \nu} |p_i S(\sqrt{p_i})^2 - 2^{-2\alpha} \beta^{-2\alpha+2} p_i^\alpha| &\leq \sum_{\sqrt{p_i} \leq \nu} |p_i S(\sqrt{p_i})^2| + |2^{-2\alpha} \beta^{-2\alpha+2} p_i^\alpha| \\
&\leq \sum_{\sqrt{p_i} \leq \nu} p_i 2^{-2\alpha+2} \beta^{-2\alpha+2} p_i^{\alpha-1} + 2^{-2\alpha} \beta^{-2\alpha+2} p_i^\alpha \\
&\leq \sum_{\sqrt{p_i} \leq \nu} 2^{-2\alpha} \beta^{-2\alpha+2} 5 p_i^\alpha \\
&\leq 2^{-2\alpha} \beta^{-2\alpha+2} 5 \left(\frac{1}{\nu^2} \nu^{2\alpha}\right) \\
&\leq 2^{-2\alpha} \beta^{-2\alpha+2} 5 P \epsilon = \Theta((p^*)^{1-\alpha} (p^*)^\alpha \epsilon) = \Theta(p^* \epsilon), \quad (115)
\end{aligned}$$

so [Lemma 9](#) still holds with this ν . Then the algorithm in [Lemma 10](#) setting ν_0 therein to be $\Theta((\epsilon P)^{\frac{1}{2\alpha-2}})$ can also estimate $P_\alpha(\mathbf{p})$ within multiplicative error $\mathcal{O}(\epsilon)$ given a rough bound of $P_\alpha(\mathbf{p})$, and the query complexity becomes

$$\begin{aligned}
&\tilde{\mathcal{O}}\left(\frac{1}{\epsilon} \left(t_{m_0} + \sqrt{\epsilon} \sum_{j=1}^{m_0} t_j + \frac{\sqrt{\sum_{j=1}^{m_0} \sum_{i: \sqrt{p_i} \in [\varphi_j, \varphi_{j-1})} p_i t_{j+1}^2}}{\sqrt{\sum_{i=1}^n p_i S_0(\sqrt{p_i})^2}}\right)\right) \\
&= \tilde{\mathcal{O}}\left(\frac{1}{\epsilon} \left(\frac{1}{\nu_{m_0}} + \frac{\sqrt{r+r}}{\sqrt{L}}\right)\right) \\
&= \tilde{\mathcal{O}}\left(\frac{1}{\epsilon} \left(\frac{1}{\epsilon^{\frac{1}{2\alpha-2}} P_\alpha(\mathbf{p})^{\frac{1}{2\alpha-2}}} + \frac{\sqrt{r}}{\sqrt{p^*}}\right)\right) \\
&= \tilde{\mathcal{O}}\left(\frac{1}{\epsilon^{\frac{1}{2\alpha-2}} P_\alpha(\mathbf{p})^{\frac{1}{2\alpha-2}}} + \frac{\sqrt{r}}{(P_\alpha(\mathbf{p}))^{\frac{1}{2\alpha}} \epsilon}\right), \quad (116)
\end{aligned}$$

where the ν_{m_0} in the second line is bounded by Eq. (78). Then following the proof in [Theorem 7](#), we can remove the requirement for P, a, b with an $\mathcal{O}(\ln(\alpha))$ overhead in query complexity which can be absorbed into the $\tilde{\mathcal{O}}$ notation. In the worst case when $P_\alpha(\mathbf{p}) = r^{1-\alpha}$, the query complexity becomes $\tilde{\mathcal{O}}\left(\frac{r^{1-\frac{1}{2\alpha}}}{\epsilon} + \frac{\sqrt{r}}{\epsilon^{1+\frac{1}{2\alpha-2}}}\right)$. \square

6.3 Quantum Rényi divergence

For any rank- r density operators ρ, σ , Müller-Lennert et al. [MLDS+13] defined a generalization of the α -Rényi divergence:

$$D_\alpha(\rho\|\sigma) := \begin{cases} \frac{1}{\alpha-1} \log \left(\text{Tr} \left[\left(\sigma^{\frac{1-\alpha}{2\alpha}} \rho \sigma^{\frac{1-\alpha}{2\alpha}} \right)^\alpha \right] \right) & \text{if } \text{Tr}(\rho\sigma) \neq 0 \\ \infty & \text{else} \end{cases} \quad (117)$$

for $\alpha \in (0, 1)$ and prove that it has some good properties. This quantum Rényi entropy is also a generalization of fidelity since $D_{\frac{1}{2}}(\rho\|\sigma) = -2 \log(F(\rho, \sigma))$.

Given oracles U_σ, U_ρ to prepare purification of the mixed states σ and ρ , the techniques used to estimate $\text{Tr}(\rho^\alpha)$ in low-rank cases can be directly applied to estimate $\text{Tr}(\sigma^\beta \rho \sigma^\beta)$ for any $\beta > 1$. In fact, we can implement a unitary \tilde{U} preparing the purification of $\sigma^\beta \rho \sigma^\beta$ with U_ρ and U_σ within $\delta^\beta + \epsilon$ additive error in spectral norm using $\mathcal{O}\left(\frac{1}{\delta} \log\left(\frac{1}{\epsilon}\right)\right)$ calls to U_σ and two calls to U_ρ by choosing the transformation polynomial of U_σ to be the one in Lemma 6 with parameters $c := \beta, \beta := 1, \nu := \delta, \eta := \epsilon$. Using amplitude amplification, we can then implement a unitary U' preparing the purification of $\frac{\sigma^\beta \rho \sigma^\beta}{\text{Tr}(\sigma^\beta \rho \sigma^\beta)}$ within ϵ error using $\mathcal{O}\left(\frac{1}{\sqrt{\text{Tr}(\sigma^\beta \rho \sigma^\beta)}} \log\left(\frac{1}{\epsilon}\right)\right)$ calls to \tilde{U} .

Therefore, we can apply our results to estimating quantum Rényi divergence in the following two steps:

1. Construct a unitary U' which is a block-encoding of $A = \frac{\sigma^{\frac{1-\alpha}{2\alpha}} \rho \sigma^{\frac{1-\alpha}{2\alpha}}}{\text{Tr}(\sigma^{\frac{1-\alpha}{2\alpha}} \rho \sigma^{\frac{1-\alpha}{2\alpha}})}$ with U_ρ and singular value transformed U_σ .
2. Estimate $D_\alpha(\rho\|\sigma) = \frac{1}{\alpha-1} \log\left(\left(\text{Tr}(\sigma^{\frac{1-\alpha}{2\alpha}} \rho \sigma^{\frac{1-\alpha}{2\alpha}})\right)^\alpha \text{Tr}(A^\alpha)\right)$ with U' using our techniques to estimate quantum α -Rényi entropy in Section 6.2.

This application is inspired by [WZC+22] in which the author has also studied the estimation of quantum Rényi divergence, but they consider the problem of estimating $\text{Tr}\left(\sigma^{\frac{1-\alpha}{2\alpha}} \rho \sigma^{\frac{1-\alpha}{2\alpha}}\right)^\alpha = \exp((\alpha-1)D_\alpha(\rho\|\sigma))$ within certain additive error and we use different polynomial approximations and some techniques beyond QSVT. Specifically, we have:

Corollary 7. *Our algorithm uses*

$$\tilde{\mathcal{O}}\left(\frac{1}{\sqrt{\text{Tr}(\sigma^\beta \rho \sigma^\beta)}} \frac{r^{\frac{1}{\alpha}}}{\epsilon^{1+\frac{1}{\alpha}}}\right) \quad (118)$$

calls to U_ρ and

$$\tilde{\mathcal{O}}\left(\frac{1}{(\text{Tr}((\sigma^\beta \rho \sigma^\beta)^\alpha))^{\frac{1}{\alpha\beta}}} \frac{1}{\sqrt{\text{Tr}(\sigma^\beta \rho \sigma^\beta)}} \frac{r^{\frac{1}{\alpha} + \frac{1}{\alpha\beta}}}{\epsilon^{1+\frac{1}{\alpha} + \frac{1}{\alpha\beta}}}\right) \quad (119)$$

calls to U_σ , where $\beta = (1-\alpha)/2\alpha$.

Proof. The error analysis is similar to that in Section 4.2 of [WZC+22]. According to Eq. (113), we need $\tilde{\mathcal{O}}\left(\frac{r^{\frac{1}{\alpha}}}{\epsilon^{1+\frac{1}{\alpha}}}\right)$ calls to U' to obtain an estimate of $\text{Tr}(A^\alpha)$ within ϵ multiplicative error. With this estimate, we can then calculate an estimate of $D_\alpha(\rho\|\sigma)$ within ϵ additive error.

Each call to U' uses \tilde{U} $\mathcal{O}\left(\frac{1}{\sqrt{\text{Tr}(\sigma^\beta \rho \sigma^\beta)}}\right)$ times since it amplifies $\sigma^\beta \rho \sigma^\beta$ to $\sigma^\beta \rho \sigma^\beta / \text{Tr}(\sigma^\beta \rho \sigma^\beta)$.

Each call to \tilde{U} uses two calls to U_ρ , so the query number of U_ρ is

$$\tilde{\mathcal{O}} \left(\frac{1}{\sqrt{\text{Tr}(\sigma^\beta \rho \sigma^\beta)}} \frac{r^{\frac{1}{\alpha}}}{\epsilon^{1+\frac{1}{\alpha}}} \right). \quad (120)$$

To compute the number of queries to U_σ , we need to analyze the error induced by \tilde{U} . Using $\mathcal{O} \left(\frac{1}{\delta} \log \left(\frac{1}{\epsilon_1} \right) \right)$ calls to U_σ , the error of \tilde{U} in spectral norm can be bounded by $\delta^\beta + \epsilon_1$. From the following lemma, we can bound the additive error of our estimate of $\text{Tr}((\sigma^\beta \rho \sigma^\beta)^\alpha)$ induced by \tilde{U} by

$$\mathcal{O} \left(r \left(\frac{\delta^\beta + \epsilon_1}{\text{Tr}(\sigma^\beta \rho \sigma^\beta)} \right)^\alpha \text{Tr}(\sigma^\beta \rho \sigma^\beta)^\alpha \right) = \mathcal{O} \left(r(\delta^\beta + \epsilon_1)^\alpha \right). \quad (121)$$

We then use:

Lemma 15 ([WZC⁺22, Lemma 4.6]). *Suppose that A and B are two positive semidefinite operators of rank $\leq r$, and $0 < \alpha < 1$. Then*

$$|\text{tr}(A^\alpha) - \text{tr}(B^\alpha)| \leq 5r \|A - B\|^\alpha, \quad (122)$$

where $\|A - B\|$ is the spectral norm of $A - B$.

Therefore, the final error induced by error of \tilde{U} is

$$\mathcal{O} \left(\frac{r(\delta^\beta + \epsilon_1)^\alpha}{\text{Tr}((\sigma^\beta \rho \sigma^\beta)^\alpha)} \right), \quad (123)$$

so we need to set $\delta := \left(\frac{\epsilon}{r} \right)^{\frac{1}{\alpha\beta}} (\text{Tr}((\sigma^\beta \rho \sigma^\beta)^\alpha))^{\frac{1}{\alpha\beta}}$ and $\epsilon_1 := \left(\frac{\epsilon}{r} \right)^{\frac{1}{\alpha}} (\text{Tr}((\sigma^\beta \rho \sigma^\beta)^\alpha))^{\frac{1}{\alpha}}$ so that the final error is bounded by ϵ .

The query number of U_σ in the final algorithm is

$$\mathcal{O} \left(\frac{1}{\delta} \log \left(\frac{1}{\epsilon_1} \right) \frac{1}{\sqrt{\text{Tr}(\sigma^\beta \rho \sigma^\beta)}} \frac{r^{\frac{1}{\alpha}}}{\epsilon^{1+\frac{1}{\alpha}}} \right) = \tilde{\mathcal{O}} \left(\frac{1}{(\text{Tr}((\sigma^\beta \rho \sigma^\beta)^\alpha))^{\frac{1}{\alpha\beta}}} \frac{1}{\sqrt{\text{Tr}(\sigma^\beta \rho \sigma^\beta)}} \frac{r^{\frac{1}{\alpha} + \frac{1}{\alpha\beta}}}{\epsilon^{1+\frac{1}{\alpha} + \frac{1}{\alpha\beta}}} \right). \quad (124)$$

□

Acknowledgements

TL was supported by a startup fund from Peking University, and the Advanced Institute of Information Technology, Peking University.

References

- [AC12] Scott Aaronson and Paul Christiano. Quantum money from hidden subspaces. In *Proceedings of the Forty-fourth Annual ACM Symposium on Theory of Computing*, pages 41–60, 2012, [arXiv:1203.4740](#).
- [AHN⁺21] Srinivasan Arunachalam, Vojtech Havlicek, Giacomo Nannicini, Kristan Temme, and Pawel Wocjan. Simpler (classical) and faster (quantum) algorithms for gibbs partition functions. In *2021 IEEE International Conference on Quantum Computing and Engineering*, pages 112–122. IEEE, 2021, [arXiv:2009.11270](#).

- [AISW19] Jayadev Acharya, Ibrahim Issa, Nirmal V. Shende, and Aaron B. Wagner. Measuring quantum entropy. In *2019 IEEE International Symposium on Information Theory*, pages 3012–3016. IEEE, 2019, [arXiv:1711.00814](#).
- [Amb10] Andris Ambainis. Variable time amplitude amplification and a faster quantum algorithm for solving systems of linear equations, 2010, [arXiv:1010.4458](#).
- [AOST16] Jayadev Acharya, Alon Orlitsky, Ananda Theertha Suresh, and Himanshu Tyagi. Estimating rényi entropy of discrete distributions. *IEEE Transactions on Information Theory*, 63(1):38–56, 2016, [arXiv:1408.1000](#).
- [Bel19] Aleksandrs Belovs. Quantum Algorithms for Classical Probability Distributions. In *27th Annual European Symposium on Algorithms*, volume 144 of *Leibniz International Proceedings in Informatics*, pages 16:1–16:11, 2019, [arXiv:1904.02192](#).
- [BHH11] Sergey Bravyi, Aram W. Harrow, and Avinatan Hassidim. Quantum algorithms for testing properties of distributions. *IEEE Transactions on Information Theory*, 57(6):3971–3981, 2011, [arXiv:0907.3920](#).
- [BHMT02] Gilles Brassard, Peter Høyer, Michele Mosca, and Alain Tapp. Quantum amplitude amplification and estimation. *Contemporary Mathematics*, 305:53–74, 2002, [arXiv:quant-ph/0005055](#).
- [BKT18] Mark Bun, Robin Kothari, and Justin Thaler. The polynomial method strikes back: Tight quantum query bounds via dual polynomials. In *Proceedings of the 50th Annual ACM SIGACT Symposium on Theory of Computing*, pages 297–310, 2018, [arXiv:1710.09079](#).
- [BOW19] Costin Bădescu, Ryan O’Donnell, and John Wright. Quantum state certification. In *Proceedings of the 51st Annual ACM SIGACT Symposium on Theory of Computing*, pages 503–514, 2019, [arXiv:1708.06002](#).
- [CCH⁺19] Shouvanik Chakrabarti, Andrew M. Childs, Shih-Han Hung, Tongyang Li, Chunhao Wang, and Xiaodi Wu. Quantum algorithm for estimating volumes of convex bodies, 2019, [arXiv:1908.03903](#).
- [CFMdW10] Sourav Chakraborty, Eldar Fischer, Arie Matsliah, and Ronald de Wolf. New results on quantum property testing. In *Proceedings of the IARCS Annual Conference on Foundations of Software Technology and Theoretical Computer Science*, volume 8 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 145–156. Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik, 2010, [arXiv:1005.0523](#).
- [CGJ19] Shantanav Chakraborty, András Gilyén, and Stacey Jeffery. The Power of Block-Encoded Matrix Powers: Improved Regression Techniques via Faster Hamiltonian Simulation. In *46th International Colloquium on Automata, Languages, and Programming*, volume 132 of *Leibniz International Proceedings in Informatics*, pages 33:1–33:14, Dagstuhl, Germany, 2019. Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik, [arXiv:1804.01973](#).
- [CKS17] Andrew M. Childs, Robin Kothari, and Rolando D. Somma. Quantum algorithm for systems of linear equations with exponentially improved dependence on precision. *SIAM Journal on Computing*, 46(6):1920–1950, Jan 2017, [arXiv:1511.02306](#).

- [GHS21] Tom Gur, Min-Hsiu Hsieh, and Sathyawageeswar Subramanian. Sublinear quantum algorithms for estimating von Neumann entropy, 2021, [arXiv:2111.11139](#).
- [GL20] András Gilyén and Tongyang Li. Distributional property testing in a quantum world. In *11th Innovations in Theoretical Computer Science Conference*, volume 151 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 25:1–25:19. Schloss Dagstuhl-Leibniz-Zentrum für Informatik, 2020, [arXiv:1902.00814](#).
- [GP22] András Gilyén and Alexander Poremba. Improved quantum algorithms for fidelity estimation, 2022, [arXiv:2203.15993](#).
- [Gro97] Lov K. Grover. Quantum mechanics helps in searching for a needle in a haystack. *Physical Review Letters*, 79(2):325, 1997, [arXiv:quant-ph/9706033](#).
- [Gro05a] Lov K. Grover. A different kind of quantum search, 2005, [arXiv:quant-ph/0503205](#).
- [Gro05b] Lov K. Grover. Fixed-point quantum search. *Physical Review Letters*, 95(15):150501, 2005.
- [GSLW19] András Gilyén, Yuan Su, Guang Hao Low, and Nathan Wiebe. Quantum singular value transformation and beyond: exponential improvements for quantum matrix arithmetics. In *Proceedings of the 51st Annual ACM SIGACT Symposium on Theory of Computing*, pages 193–204, 2019, [arXiv:1806.01838](#).
- [HHL09] Aram W. Harrow, Avinatan Hassidim, and Seth Lloyd. Quantum algorithm for linear systems of equations. *Physical Review Letters*, 103(15):150502, 2009, [arXiv:0811.3171](#).
- [HM19] Yassine Hamoudi and Frédéric Magniez. Quantum Chebyshev’s inequality and applications. In *Proceedings of the 46th International Colloquium on Automata, Languages, and Programming*, volume 132 of *Leibniz International Proceedings in Informatics*, pages 69:1–69:16, 2019, [arXiv:1807.06456](#).
- [Hø00] Peter Høyer. Arbitrary phases in quantum amplitude amplification. *Physical Review A*, 62(5):052304, 2000, [arXiv:quant-ph/0006031](#).
- [HW20] Aram W Harrow and Annie Y. Wei. Adaptive quantum simulated annealing for bayesian inference and estimating partition functions. In *Proceedings of the Fourteenth Annual ACM-SIAM Symposium on Discrete Algorithms*, pages 193–212. SIAM, 2020, [arXiv:1907.09965](#).
- [JVHW15] Jiantao Jiao, Kartik Venkat, Yanjun Han, and Tsachy Weissman. Minimax estimation of functionals of discrete distributions. *IEEE Transactions on Information Theory*, 61(5):2835–2885, 2015, [arXiv:1406.6956](#).
- [LC17] Guang Hao Low and Isaac L. Chuang. Optimal Hamiltonian simulation by quantum signal processing. *Physical Review Letters*, 118(1):010501, 2017, [arXiv:1606.02685](#).
- [LC19] Guang Hao Low and Isaac L. Chuang. Hamiltonian simulation by qubitization. *Quantum*, 3:163, 2019, [arXiv:1610.06546](#).
- [LW19] Tongyang Li and Xiaodi Wu. Quantum query complexity of entropy estimation. *IEEE Transactions on Information Theory*, 65(5):2899–2921, 2019, [arXiv:1710.06025](#). © 2019 IEEE.

- [MdW16] Ashley Montanaro and Ronald de Wolf. A survey of quantum property testing. *Theory of Computing*, pages 1–81, 2016, [arXiv:1310.2035](#).
- [MLDS⁺13] Martin Müller-Lennert, Frédéric Dupuis, Oleg Szehr, Serge Fehr, and Marco Tomamichel. On quantum Rényi entropies: A new generalization and some properties. *Journal of Mathematical Physics*, 54(12):122203, 2013, [arXiv:1306.3142](#).
- [Mon15] Ashley Montanaro. Quantum speedup of Monte Carlo methods. *Proceedings of the Royal Society A*, 471(2181):20150301, 2015, [arXiv:1504.06987](#).
- [MRTC21] John M. Martyn, Zane M. Rossi, Andrew K. Tan, and Isaac L. Chuang. Grand unification of quantum algorithms. *PRX Quantum*, 2(4):040203, 2021, [arXiv:2105.02859](#).
- [OD21] Davide Orsucci and Vedran Dunjko. On solving classes of positive-definite quantum linear systems with quadratically improved runtime in the condition number. *Quantum*, 5:573, 2021, [arXiv:2101.11868](#).
- [OW15] Ryan O’Donnell and John Wright. Quantum spectrum testing. In *Proceedings of the 47th Annual ACM on Symposium on Theory of Computing*, pages 529–538. ACM, 2015, [arXiv:1501.05028](#).
- [OW16] Ryan O’Donnell and John Wright. Efficient quantum tomography. In *Proceedings of the 48th Annual ACM Symposium on Theory of Computing*, pages 899–912. ACM, 2016, [arXiv:1508.01907](#).
- [OW17a] Ryan O’Donnell and John Wright. Efficient quantum tomography II. In *Proceedings of the Forty-ninth Annual ACM SIGACT Symposium on Theory of Computing*, pages 962–974. ACM, 2017, [arXiv:1612.00034](#).
- [OW17b] Ryan O’Donnell and John Wright. Guest column: A primer on the statistics of longest increasing subsequences and quantum states (shortened version). *ACM SIGACT News*, 48(3):37–59, 2017.
- [Rén61] Alfréd Rényi. On measures of entropy and information. In *Proceedings of the 4th Berkeley Symposium on Mathematical Statistics and Probability*, volume 1, pages 547–561, 1961.
- [Ron10] Dana Ron. Algorithmic and analysis techniques in property testing. *Foundations and Trends® in Theoretical Computer Science*, 5(2):73–205, 2010.
- [Sch41] A. C. Schaeffer. Inequalities of A. Markoff and S. Bernstein for polynomials and related functions. *Bulletin of the American Mathematical Society*, 47(8):565 – 579, 1941.
- [SH21] Sathyawageeswar Subramanian and Min-Hsiu Hsieh. Quantum algorithm for estimating α -Rényi entropies of quantum states. *Physical Review A*, 104(2):022428, 2021, [arXiv:1908.05251](#).
- [Sha48] Claude E. Shannon. A mathematical theory of communication. *Bell System Technical Journal*, 27(3):379–423, 1948.
- [SN14] Jun John Sakurai and Jim Napolitano. *Modern quantum mechanics*. Pearson Harlow, 2014.

- [Sze04] Mario Szegedy. Quantum speed-up of Markov chain based algorithms. In *Proceedings of the 45th Annual IEEE Symposium on Foundations of Computer Science*, pages 32–41, 2004.
- [TGP06] Tathagat Tulsi, Lov K. Grover, and Apoorva Patel. A new algorithm for fixed point quantum search. *Quantum Information & Computation*, 6(6):483–494, 2006, [arXiv:quant-ph/0603132](#).
- [Val11] Paul Valiant. Testing symmetric properties of distributions. *SIAM Journal on Computing*, 40(6):1927–1968, 2011.
- [WA08] Pawel Wocjan and Anura Abeyesinghe. Speedup via quantum sampling. *Physical Review A*, 78(4):042336, 2008, [arXiv:0804.4259](#).
- [WCNA09] Pawel Wocjan, Chen-Fu Chiang, Daniel Nagaj, and Anura Abeyesinghe. Quantum algorithm for approximating partition functions. *Physical Review A*, 80(2):022340, 2009.
- [WY16] Yihong Wu and Pengkun Yang. Minimax rates of entropy estimation on large alphabets via best polynomial approximation. *IEEE Transactions on Information Theory*, 62(6):3702–3720, 2016, [arXiv:1407.0381](#).
- [WZC⁺21] Qisheng Wang, Zhicheng Zhang, Kean Chen, Ji Guan, Wang Fang, and Mingsheng Ying. Quantum algorithm for fidelity estimation, 2021, [arXiv:2103.09076](#).
- [WZC⁺22] Qisheng Wang, Zhicheng Zhang, Kean Chen, Ji Guan, Wang Fang, Junyi Liu, and Mingsheng Ying. Quantum algorithm for fidelity estimation. *IEEE Transactions on Information Theory*, 2022, [arXiv:2203.13522](#).
- [WZW22] Youle Wang, Benchi Zhao, and Xin Wang. Quantum algorithms for estimating quantum entropies, 2022, [arXiv:2203.02386](#).
- [YLC14] Theodore J. Yoder, Guang Hao Low, and Isaac L. Chuang. Fixed-point quantum search with an optimal number of queries. *Physical Review Letters*, 113(21):210501, 2014, [arXiv:1409.3305](#).

A Proof of **Proposition 2**

Proof. In the following proof, we use Λ_k to denote a m -bit 01-string $\Lambda_k := 0^{k-1}10^{m-k}$ and when we say $|\phi\rangle$ is an approximation of $|\psi\rangle$ up to error $\mathcal{O}(L\epsilon)$ or $|\phi\rangle = |\psi\rangle + \mathcal{O}(L\epsilon)$, we mean $\| |\psi\rangle - |\phi\rangle \| = \mathcal{O}(L\epsilon)$.

Let $\tilde{\mathcal{A}}$ be **Algorithm 2** and $\mathcal{A} = \mathcal{A}_m \cdot \dots \cdot \mathcal{A}_1$ be the unitary operation \mathcal{A} in **Algorithm 2**.

We first calculate the output of

$$\tilde{\mathcal{A}}|\mathbf{0}\rangle = \mathcal{A}\left(\sum_{i=1}^n \sqrt{p_i}|0\rangle_F|\mathbf{0}\rangle_C|\mathbf{0}\rangle_A|i\rangle_B|+\rangle_Q|\mathbf{0}\rangle_{P,I}\right). \quad (125)$$

Since \mathcal{A} is a linear operator, we only need to calculate

$$\mathcal{A}(|0\rangle_F|\mathbf{0}\rangle_C|\mathbf{0}\rangle_A|i\rangle_B|+\rangle_Q|\mathbf{0}\rangle_{P,I}). \quad (126)$$

We now describe the state in different stages of \mathcal{A} when it is initialized to $|0\rangle_F|\mathbf{0}\rangle_C|\mathbf{0}\rangle_A|i\rangle_B|+\rangle_Q|\mathbf{0}\rangle_{P,I}$ before \mathcal{A} .

Let $j \in \{1, \dots, m\}$ be such that $\sqrt{p_i} \in [\varphi_j, \varphi_{j-1})$. We divide the m stages of \mathcal{A} into three parts: \mathcal{A}_1 to \mathcal{A}_{j-1} , \mathcal{A}_j to \mathcal{A}_{j+1} , and \mathcal{A}_{j+2} to \mathcal{A}_m if exists.

State after \mathcal{A}_k for $k = 1, \dots, j-1$ The performance of \mathcal{A}_k for all $k = 1, \dots, j-1$ on $|0\rangle_F|\mathbf{0}\rangle_C|\mathbf{0}\rangle_A|i\rangle_B|+\rangle_Q$ is similar. Since \mathcal{A}_k only change the first k registers of I and P , the state before \mathcal{A}_k can be written as

$$\alpha_0^{(i,k)}|0\rangle_F|\mathbf{0}\rangle_C|\mathbf{0}\rangle_A|i\rangle_B|+\rangle_Q|\gamma_1\rangle_{P_1,I_1} \cdots |\gamma_{k-1}\rangle_{P_{k-1},I_{k-1}}|\mathbf{0}\rangle_{P_k,I_k,\dots,P_m,I_m} + \alpha_1^{(i,k)}|\psi_{\text{stopped}}\rangle, \quad (127)$$

where $\|(|\mathbf{0}\rangle\langle\mathbf{0}|_C \otimes I)|\psi_{\text{stopped}}\rangle\| = 0$ and $|\gamma\rangle_l$ is the state $|\gamma\rangle_{P,I}$ in [Lemma 5](#) produced by $W(\varphi_l, L\epsilon/m)$ in \mathcal{A}_l . State in registers F, A, Q is $|0\rangle_F|\mathbf{0}\rangle_A|+\rangle_Q$ when state in register C is $|\mathbf{0}\rangle$ since the step 3 of \mathcal{A}_l is conditional on C_l being $|1\rangle$ for all $l = 1, \dots, k-1$ and only these operations can change the state in registers F, A, Q .

Note that the last $m-k+1$ qubits of C register must be $|0\rangle$, since \mathcal{A}_l for $l = 1, \dots, k-1$ do not change them. Then, we can imply that $|\psi_{\text{stopped}}\rangle$ has no overlap with $|\mathbf{0}\rangle_{C_1,\dots,C_{k-1}}$ since $\|(|\mathbf{0}\rangle\langle\mathbf{0}|_C \otimes I)|\psi_{\text{stopped}}\rangle\| = 0$. Therefore, \mathcal{A}_k will not change $|\psi_{\text{stopped}}\rangle$, and we only need to consider the first component in Eq. (127).

After step 1 of \mathcal{A}_k , the the component in Eq. (127) becomes

$$\alpha_0^{(i,k)}|0\rangle_F|\mathbf{0}\rangle_C|\mathbf{0}\rangle_A|i\rangle_B|+\rangle_Q|\gamma_1\rangle_{P_1,I_1} \cdots |\gamma_{k-1}\rangle_{P_{k-1},I_{k-1}}|0\rangle_{P_k}(|\mathbf{0}\rangle|i\rangle)_{I_k}|\mathbf{0}\rangle_{P_k,I_k,\dots,P_m,I_m}. \quad (128)$$

After step 2 of \mathcal{A}_k , the first component in Eq. (127) becomes

$$\begin{aligned} & \alpha_0^{(i,k)}\beta_0^{(i,k)}|0\rangle_F|\mathbf{0}\rangle_C|\mathbf{0}\rangle_A|i\rangle_B|+\rangle_Q|\gamma_1\rangle_{P_1,I_1} \cdots |\gamma_{k-1}\rangle_{P_{k-1},I_{k-1}}|\gamma_k\rangle_{P_k,I_k}|\mathbf{0}\rangle_{P_{k+1},I_{k+1},\dots,P_m,I_m} \\ & + \alpha_0^{(i,k)}\beta_1^{(i,k)}|0\rangle_F|\Lambda_k\rangle_C|\mathbf{0}\rangle_A|i\rangle_B|+\rangle_Q|\gamma_1\rangle_{P_1,I_1} \cdots |\gamma_{k-1}\rangle_{P_{k-1},I_{k-1}}|+\rangle_{P_k}(|\mathbf{0}\rangle|i\rangle)_{I_k}|\mathbf{0}\rangle_{P_{k+1},I_{k+1},\dots,P_m,I_m}, \end{aligned} \quad (129)$$

where Λ_k is a m -bit 01-string $\Lambda_k := 0^{k-1}10^{m-k}$.

Since the corresponding singular value of $(|\mathbf{0}\rangle|i\rangle)_{I_k}$ is $\sqrt{p_i}$ and $\sqrt{p_i} < \varphi_{j-1} \leq \varphi_k$, from [Lemma 5](#), we can imply that $|\beta_1^{(i,k)}| \leq \frac{L\epsilon}{m}$ and $|\beta_0^{(i,k)}| = \sqrt{1 - |\beta_1^{(i,k)}|^2} \geq \sqrt{1 - (\frac{L\epsilon}{m})^2}$.

After the step 3 of \mathcal{A}_k , the state in Eq. (129) does not change.

Notice that the component in Eq. (129) is the component in Eq. (127) for $k+1$. Since the $\alpha_0^{(1)} = 1$, we can imply that the amplitude of Eq. (129) satisfies that $|\alpha_0^{(i,k)}\beta_0^{(i,k)}| \geq \sqrt{(1 - (\frac{L\epsilon}{m})^2)^k} \geq \sqrt{1 - (L\epsilon)^2}$ by induction on k .

Therefore, when $k = j-1$, we can imply that after \mathcal{A}_{j-1} , the state becomes

$$|0\rangle_F|\mathbf{0}\rangle_C|\mathbf{0}\rangle_A|i\rangle_B|+\rangle_Q|\gamma_1\rangle_{P_1,I_1} \cdots |\gamma_{j-1}\rangle_{P_{j-1},I_{j-1}}|\mathbf{0}\rangle_{P_j,I_j,\dots,P_m,I_m} \quad (130)$$

up to error $\mathcal{O}(L\epsilon)$.

Before continuing to the next part, we first consider a special case when $j = m$. In \mathcal{A}_m , step 1 and step 2 will map the state in Eq. (130) to

$$|0\rangle_F|\Lambda_m\rangle_C|\mathbf{0}\rangle_A|i\rangle_B|+\rangle_Q|\gamma_1\rangle_{P_1,I_1} \cdots |\gamma_{m-1}\rangle_{P_{m-1},I_{m-1}}|0\rangle_{P_m}(|\mathbf{0}\rangle|i\rangle)_{I_m}. \quad (131)$$

After step 3 of \mathcal{A}_m , the state in Eq. (130) will be

$$S_m(\sqrt{p_i})|1\rangle_F|\Lambda_m\rangle_C|i\rangle_A|i\rangle_B|+\rangle_Q|\gamma_1\rangle_{P_1,I_1} \cdots |\gamma_{m-1}\rangle_{P_{m-1},I_{m-1}}|0\rangle_{P_m}(|\mathbf{0}\rangle|i\rangle)_{I_m} \quad (132)$$

$$+ \sqrt{1 - S_m(\sqrt{p_i})^2}|0\rangle_F|\Lambda_m\rangle_C|\psi_{\text{garbage}}^{(m)}\rangle_{A,B,Q,P,I}. \quad (133)$$

Since $\sqrt{p_i} \in [\varphi_m, \varphi_{m-2})$ and S_m satisfies the condition in Eq. (50), we can imply that

$$\begin{aligned}
& (|1\rangle\langle 1|_F \otimes I) \mathcal{A}(|0\rangle_F |\mathbf{0}\rangle_C |\mathbf{0}\rangle_A |i\rangle_B |+\rangle_Q |\mathbf{0}\rangle_{P,I}) \\
& = S_m(\sqrt{p_i}) |1\rangle_F |\Lambda_m\rangle_C |i\rangle_A |i\rangle_B |+\rangle_Q |\gamma_1\rangle_{P_1, I_1} \cdots |\gamma_{m-1}\rangle_{P_{m-1}, I_{m-1}} |0\rangle_{P_m} (|\mathbf{0}\rangle |i\rangle)_{I_m} \\
& \approx S(\sqrt{p_i}) |1\rangle_F |\Lambda_m\rangle_C |i\rangle_A |i\rangle_B |+\rangle_Q |\gamma_1\rangle_{P_1, I_1} \cdots |\gamma_{m-1}\rangle_{P_{m-1}, I_{m-1}} |0\rangle_{P_m} (|\mathbf{0}\rangle |i\rangle)_{I_m} \\
& =: S(\sqrt{p_i}) |\Phi_m\rangle,
\end{aligned} \tag{134}$$

where the approximation error of the second equation is $\mathcal{O}(L\epsilon)$.

In the following part, we assume $j < m$.

State after \mathcal{A}_j . Since \mathcal{A}_j is linear, we only consider applying \mathcal{A}_j to

$$|0\rangle_F |\mathbf{0}\rangle_C |\mathbf{0}\rangle_A |i\rangle_B |+\rangle_Q |\gamma_1\rangle_{P_1, I_1} \cdots |\gamma_j\rangle_{P_{j-1}, I_{j-1}} |\mathbf{0}\rangle_{P_j, I_j, \dots, P_m, I_m}, \tag{135}$$

and the result is also an $\mathcal{O}(L\epsilon)$ -approximation of the state after \mathcal{A}_j .

After step 1 and step 2 of \mathcal{A}_j , we will have

$$\beta_0^{(i,j)} |0\rangle_F |\mathbf{0}\rangle_C |\mathbf{0}\rangle_A |i\rangle_B |+\rangle_Q |\gamma_1\rangle_{P_1, I_1} \cdots |\gamma_{j-1}\rangle_{P_{j-1}, I_{j-1}} |\gamma_j\rangle_{P_j, I_j} |\mathbf{0}\rangle_{P_{j+1}, I_{j+1}, \dots, P_m, I_m} \tag{136}$$

$$+ \beta_1^{(i,j)} |0\rangle_F |\Lambda_j\rangle_C |\mathbf{0}\rangle_A |i\rangle_B |+\rangle_Q |\gamma_1\rangle_{P_1, I_1} \cdots |\gamma_{j-1}\rangle_{P_{j-1}, I_{j-1}} |+\rangle_{P_j} (|\mathbf{0}\rangle |i\rangle)_{I_j} |\mathbf{0}\rangle_{P_{j+1}, I_{j+1}, \dots, P_m, I_m}. \tag{137}$$

Since step 3 of \mathcal{A}_j is conditional on register C_j being $|0\rangle$, so the component in Eq. (136) does not change, and the component in Eq. (137) becomes

$$\begin{aligned}
& \beta_1^{(i,j)} S_j(\sqrt{p_i}) |1\rangle_F |\Lambda_j\rangle_C |i\rangle_A |i\rangle_B |+\rangle_Q |\gamma_1\rangle_{P_1, I_1} \cdots |\gamma_{j-1}\rangle_{P_{j-1}, I_{j-1}} |+\rangle_{P_j} (|\mathbf{0}\rangle |i\rangle)_{I_j} |\mathbf{0}\rangle_{P_{j+1}, I_{j+1}, \dots, P_m, I_m} \\
& + \beta_1^{(i,j)} \sqrt{1 - S_j(\sqrt{p_i})^2} |0\rangle_F |\Lambda_j\rangle_C |\psi_{\text{garbage}}^{(i,j)}\rangle_{A,B,Q,P,I}
\end{aligned} \tag{138}$$

Since S_j satisfies the condition in Eq. (50) and $\sqrt{p_i} \in [\varphi_j, \varphi_{j-2})$, we can imply that the component in Eq. (138) is an approximation of

$$\beta_1^{(i,j)} S(\sqrt{p_i}) |1\rangle_F |\Lambda_j\rangle_C |i\rangle_A |i\rangle_B |+\rangle_Q |\gamma_1\rangle_{P_1, I_1} \cdots |\gamma_{j-1}\rangle_{P_{j-1}, I_{j-1}} |+\rangle_{P_j} (|\mathbf{0}\rangle |i\rangle)_{I_j} |\mathbf{0}\rangle_{P_{j+1}, I_{j+1}, \dots, P_m, I_m} \tag{139}$$

up to error $\mathcal{O}(L\epsilon)$.

Therefore, the state after \mathcal{A}_j is

$$\begin{aligned}
& \beta_0^{(i,j)} |0\rangle_F |\mathbf{0}\rangle_C |\mathbf{0}\rangle_A |i\rangle_B |+\rangle_Q |\gamma_1\rangle_{P_1, I_1} \cdots |\gamma_j\rangle_{P_j, I_j} |\mathbf{0}\rangle_{P_{j+1}, I_{j+1}, \dots, P_m, I_m} \\
& + \beta_1^{(i,j)} S(\sqrt{p_i}) |1\rangle_F |\Lambda_j\rangle_C |i\rangle_A |i\rangle_B |+\rangle_Q |\gamma_1\rangle_{P_1, I_1} \cdots |+\rangle_{P_j} (|\mathbf{0}\rangle |i\rangle)_{I_j} |\mathbf{0}\rangle_{P_{j+1}, I_{j+1}, \dots, P_m, I_m} \\
& + \beta_1^{(i,j)} \sqrt{1 - S_j(\sqrt{p_i})^2} |0\rangle_F |\Lambda_j\rangle_C |\psi_{\text{garbage}}^{(i,j)}\rangle_{A,B,Q,P,I}
\end{aligned} \tag{140}$$

up to error $\mathcal{O}(L\epsilon)$.

State after \mathcal{A}_{j+1} . Since the first two steps of \mathcal{A}_{j+1} are conditional on the first j qubits of register C being $|\mathbf{0}\rangle$ and the third step is conditional on the $(j+1)$ -th qubit of register C being $|1\rangle$, \mathcal{A}_{j+1} will only change the component in Eq. (140). Therefore, we only consider the result applying \mathcal{A}_{j+1} to the component Eq. (140).

After step 1 and step 2 of \mathcal{A}_{j+1} , the component in Eq. (140) becomes

$$\beta_0^{(i,j)} \beta_0^{(i,j+1)} |0\rangle_F |\mathbf{0}\rangle_C |\mathbf{0}\rangle_A |i\rangle_B |+\rangle_Q |\gamma_1\rangle_{P_1, I_1} \cdots |\gamma_j\rangle_{P_j, I_j} |\gamma_{j+1}\rangle_{P_{j+1}, I_{j+1}} |\mathbf{0}\rangle_{P_{j+2}, I_{j+2}, \dots, P_m, I_m} \tag{141}$$

$$+ \beta_0^{(i,j)} \beta_1^{(i,j+1)} |0\rangle_F |\Lambda_{j+1}\rangle_C |\mathbf{0}\rangle_A |i\rangle_B |+\rangle_Q |\gamma_1\rangle_{P_1, I_1} \cdots |\gamma_j\rangle_{P_j, I_j} |+\rangle_{P_{j+1}} (|\mathbf{0}\rangle |i\rangle)_{I_{j+1}} |\mathbf{0}\rangle_{P_{j+2}, I_{j+2}, \dots, P_m, I_m}. \tag{142}$$

Since $\sqrt{p_i} \geq \varphi_j = 2\varphi_{j+1}$, from [Lemma 5](#), we have $|\beta_0^{(i,j+1)}| \leq L\epsilon$. Then, we can imply that the component in Eq. (140) after step 1 and step 2 of \mathcal{A}_{j+1} is an approximation of

$$\beta_0^{(i,j)} |0\rangle_F |\Lambda_{j+1}\rangle_C |\mathbf{0}\rangle_A |i\rangle_B |+\rangle_Q |\gamma_1\rangle_{P_1, I_1} \cdots |\gamma_j\rangle_{P_j, I_j} |+\rangle_{P_{j+1}} (|\mathbf{0}\rangle |i\rangle)_{I_{j+1}} |\mathbf{0}\rangle_{P_{j+2}, I_{j+2}, \dots, P_m, I_m} \quad (143)$$

up to error $\mathcal{O}(L\epsilon)$.

Then, after step 3 of \mathcal{A}_{j+1} , the component in Eq. (140) becomes

$$\begin{aligned} & \beta_0^{(i,j)} S_{j+1}(\sqrt{p_i}) |1\rangle_F |\Lambda_{j+1}\rangle_C |i\rangle_A |i\rangle_B |+\rangle_Q |\gamma_1\rangle_{P_1, I_1} \cdots |\gamma_j\rangle_{P_j, I_j} |+\rangle_{P_{j+1}} (|\mathbf{0}\rangle |i\rangle)_{I_{j+1}} |\mathbf{0}\rangle_{P_{j+2}, I_{j+2}, \dots, P_m, I_m} \\ & + \beta_0^{(i,j)} \sqrt{1 - S_{j+1}(\sqrt{p_i})^2} |0\rangle_F |\Lambda_{j+1}\rangle_C |\psi_{\text{garbage}}^{(i,j+1)}\rangle_{A, B, Q, P, I} \end{aligned} \quad (144)$$

up to error $\mathcal{O}(L\epsilon)$.

Since S_{j+1} satisfies the condition in Eq. (50) and $\sqrt{p_i} \in [\varphi_{j+1}, \varphi_{j-1})$, we can imply that the component in Eq. (144) is an approximation of

$$\beta_0^{(i,j)} S(\sqrt{p_i}) |1\rangle_F |\Lambda_{j+1}\rangle_C |i\rangle_A |i\rangle_B |+\rangle_Q |\gamma_1\rangle_{P_1, I_1} \cdots |\gamma_j\rangle_{P_j, I_j} |+\rangle_{P_{j+1}} (|\mathbf{0}\rangle |i\rangle)_{I_{j+1}} |\mathbf{0}\rangle_{P_{j+2}, I_{j+2}, \dots, P_m, I_m} \quad (145)$$

up to error $\mathcal{O}(L\epsilon)$.

In conclusion, the state after \mathcal{A}_{j+1} is

$$\begin{aligned} & \beta_0^{(i,j)} S(\sqrt{p_i}) |1\rangle_F |\Lambda_{j+1}\rangle_C |i\rangle_A |i\rangle_B |+\rangle_Q |\gamma_1\rangle_{P_1, I_1} \cdots |\gamma_j\rangle_{P_j, I_j} |+\rangle_{P_{j+1}} (|\mathbf{0}\rangle |i\rangle)_{I_{j+1}} |\mathbf{0}\rangle_{P_{j+2}, I_{j+2}, \dots, P_m, I_m} \\ & + \beta_1^{(i,j)} S(\sqrt{p_i}) |1\rangle_F |\Lambda_j\rangle_C |i\rangle_A |i\rangle_B |+\rangle_Q |\gamma_1\rangle_{P_1, I_1} \cdots |+\rangle_{P_j} (|\mathbf{0}\rangle |i\rangle)_{I_j} |\mathbf{0}\rangle_{P_{j+1}, I_{j+1}, \dots, P_m, I_m} \\ & + \beta_0^{(i,j)} \sqrt{1 - S(\sqrt{p_i})^2} |0\rangle_F |\Lambda_{j+1}\rangle_C |\psi_{\text{garbage}}^{(i,j+1)}\rangle_{A, B, Q, P, I} \\ & + \beta_1^{(i,j)} \sqrt{1 - S(\sqrt{p_i})^2} |0\rangle_F |\Lambda_j\rangle_C |\psi_{\text{garbage}}^{(i,j)}\rangle_{A, B, Q, P, I} \end{aligned} \quad (146)$$

up to error $\mathcal{O}(L\epsilon)$.

State after \mathcal{A} . Since the state in Eq. (146) has no overlap with $|\mathbf{0}\rangle_{C_1, \dots, C_{j+1}}$, \mathcal{A}_k for all $k = j+2, \dots, m$ do not change it.

Therefore, we can imply that state after \mathcal{A} , $\mathcal{A}(|0\rangle_F |\mathbf{0}\rangle_C |\mathbf{0}\rangle_A |i\rangle_B |+\rangle_Q |\mathbf{0}\rangle_{P, I})$, satisfies

$$\begin{aligned} & (|1\rangle\langle 1|_F \otimes I) \mathcal{A}(|0\rangle_F |\mathbf{0}\rangle_C |\mathbf{0}\rangle_A |i\rangle_B |+\rangle_Q |\mathbf{0}\rangle_{P, I}) \\ & \approx \beta_0^{(i,j)} S(\sqrt{p_i}) |1\rangle_F |\Lambda_{j+1}\rangle_C |i\rangle_A |i\rangle_B |+\rangle_Q |\gamma_1\rangle_{P_1, I_1} \cdots |\gamma_j\rangle_{P_j, I_j} |+\rangle_{P_{j+1}} (|\mathbf{0}\rangle |i\rangle)_{I_{j+1}} |\mathbf{0}\rangle_{P_{j+2}, I_{j+2}, \dots, P_m, I_m} \\ & \quad + \beta_1^{(i,j)} S(\sqrt{p_i}) |1\rangle_F |\Lambda_j\rangle_C |i\rangle_A |i\rangle_B |+\rangle_Q |\gamma_1\rangle_{P_1, I_1} \cdots |+\rangle_{P_j} (|\mathbf{0}\rangle |i\rangle)_{I_j} |\mathbf{0}\rangle_{P_{j+1}, I_{j+1}, \dots, P_m, I_m} \\ & =: S(\sqrt{p_i}) |1\rangle_F |\Phi_i\rangle, \end{aligned} \quad (147)$$

where the approximation error of the first equation is $\mathcal{O}(L\epsilon)$ and $|\Phi_i\rangle$ is a normalized state.

Correctness. From Eq. (147) and Eq. (134), we can imply that

$$(|1\rangle\langle 1|_F \otimes I) \mathcal{A} \left(\sum_{i=1}^n \sqrt{p_i} |0\rangle_F |\mathbf{0}\rangle_C |\mathbf{0}\rangle_A |i\rangle_B |+\rangle_Q |\mathbf{0}\rangle_{P, I} \right) \approx \sum_{i=1}^n \sqrt{p_i} S(\sqrt{p_i}) |1\rangle_F |\Phi_i\rangle, \quad (148)$$

where the approximation error is $\mathcal{O}(L\epsilon)$.

Using variable-time amplitude estimation, we can estimate

$$\begin{aligned} \|(|1\rangle\langle 1|_F \otimes I)\mathcal{A}(|0\rangle_F|\mathbf{0}\rangle_C|\mathbf{0}\rangle_A|\psi_p\rangle_B|+\rangle_Q|\mathbf{0}\rangle_{P,I})\|^2 &= \sum_{i=1}^n p_i S(\sqrt{p_i})^2 + \mathcal{O}(L\epsilon) \\ &= \sum_{i=1}^n p_i S(\sqrt{p_i})^2 (1 + \mathcal{O}(\epsilon)) \end{aligned} \quad (149)$$

within multiplicative error ϵ with success probability at least $1 - \delta$.

Since $L \leq \sum_{i=1}^n p_i S(\sqrt{p_i})^2$, we have $\sum_{i=1}^n p_i S(\sqrt{p_i})^2 + \mathcal{O}(L\epsilon) = \sum_{i=1}^n p_i S(\sqrt{p_i})^2 (1 + \mathcal{O}(\epsilon))$. Therefore, the output of [Algorithm 3](#) is an estimate of $\sum_{i=1}^n p_i S(\sqrt{p_i})^2$ within multiplicative error $\mathcal{O}(\epsilon)$.

By rescaling ϵ to $c\epsilon$ with a small constant c , we can estimate $\sum_{i=1}^n p_i S(\sqrt{p_i})^2$ to within multiplicative error ϵ with the same query complexity.

Complexity. Our [Algorithm 3](#) is a direct use of variable-time amplitude estimation in [Theorem 6](#), so in order to get its query complexity, we only need to calculate $t_j, p_{\text{stop}=t_j}, T_{\text{avg}}$, and p_{succ} of $\tilde{\mathcal{A}}$. We will calculate these parameters for \mathcal{A} for simplicity since $\tilde{\mathcal{A}}$ only use U_p one more time than \mathcal{A} .

Calculate p_{succ} . In the previous paragraph, we have proved that $p_{\text{succ}} = \sum_{i=1}^n p_i S(\sqrt{p_i})^2 (1 + \mathcal{O}(\epsilon))$.

Calculate t_j . The query complexity of \mathcal{A}_j for $j < m$ is the sum of query complexity of $W(\varphi_j, L\epsilon/m)$ and $U_{S_j}^{(SV)}$, which is $\mathcal{O}\left(\frac{1}{\varphi_j} \log\left(\frac{m}{\epsilon L}\right) + \deg(S_j)\right)$ for $j < m$ while the query complexity of \mathcal{A}_m is $\mathcal{O}(\deg(S_m))$.

Then the sum of the query complexity of the first j stages of \mathcal{A} for $j < m$ is

$$\begin{aligned} t_j &= \mathcal{O}\left(\sum_{k=1}^j \left(\frac{1}{\varphi_k} \log\left(\frac{m}{\epsilon L}\right) + \deg(S_k)\right)\right) \\ &= \mathcal{O}\left(\sum_{k=1}^j \left(\frac{2^k}{\beta} \log\left(\frac{m}{\epsilon L}\right) + \deg(S_k)\right)\right) \\ &= \mathcal{O}\left(\frac{2^j}{\beta} \log\left(\frac{m}{\epsilon L}\right) + \sum_{k=1}^j \deg(S_k)\right), \end{aligned} \quad (150)$$

and $t_m = \mathcal{O}\left(\frac{2^{m-1}}{\beta} \log\left(\frac{m}{\epsilon L}\right) + \sum_{k=1}^m \deg(S_k)\right)$.

Calculate $p_{\text{stop}=t_j}$. Note that

$$\begin{aligned} p_{\text{stop}=t_j} &= \|(|\Lambda_j\rangle\langle \Lambda_j|_C \otimes I)\mathcal{A}\left(\sum_{i=1}^n \sqrt{p_i} |0\rangle_F |\mathbf{0}\rangle_C |i\rangle_A |+\rangle_Q |\mathbf{0}\rangle_{P,I}\right)\|^2 \\ &= \sum_{i=1}^n p_i \|(|\Lambda_j\rangle\langle \Lambda_j|_C \otimes I)\mathcal{A}(|0\rangle_F |\mathbf{0}\rangle_C |\mathbf{0}\rangle_A |i\rangle_B |+\rangle_Q |\mathbf{0}\rangle_{P,I})\|^2 \end{aligned} \quad (151)$$

Then from Eq. (146), we can imply that for i such that $\sqrt{p_i} \notin [\varphi_j, \varphi_{j-2})$, they contribute at most $\mathcal{O}((L\epsilon)^2)$ to $p_{\text{stop}=t_j}$, so we only need to consider contribute of i such that $\sqrt{p_i} \in [\varphi_j, \varphi_{j-1})$. Therefore, we can imply that $p_{\text{stop}=t_j}$ is

$$\sum_{i:\sqrt{p_i} \in [\varphi_j, \varphi_{j-1})} p_i |\beta_1^{(i,j)}|^2 + \sum_{i:\sqrt{p_i} \in [\varphi_{j-1}, \varphi_j)} p_i |\beta_0^{(i,j-1)}|^2 + \mathcal{O}(L\epsilon), \quad (152)$$

where $\beta_0^{(i,0)} = \beta_0^{(i,m)} := 0$ and $\beta_1^{(i,m)} := 1$.

Calculate T_{avg} . Let $t_{m+1} := t_m$. Then we have

$$\begin{aligned} T_{\text{avg}} &= \sqrt{\sum_{j=1}^m p_{\text{stop}=t_j} t_j^2} \\ &\leq \sqrt{\sum_{j=1}^m \left(\sum_{i:\sqrt{p_i} \in [\varphi_j, \varphi_{j-1})} p_i |\beta_1^{(i,j)}|^2 + \sum_{i:\sqrt{p_i} \in [\varphi_{j-1}, \varphi_j)} p_i |\beta_0^{(i,j-1)}|^2 + \mathcal{O}(L\epsilon) \right) t_j^2} \\ &\leq \sqrt{\sum_{j=1}^m \left(\sum_{i:\sqrt{p_i} \in [\varphi_j, \varphi_{j-1})} p_i |\beta_1^{(i,j)}|^2 t_{j+1}^2 + \sum_{i:\sqrt{p_i} \in [\varphi_{j-1}, \varphi_j)} p_i |\beta_0^{(i,j-1)}|^2 t_j^2 \right)} + \mathcal{O}(\sqrt{L\epsilon} \sum_{j=1}^m t_j) \\ &= \sqrt{\sum_{j=1}^m \sum_{i:\sqrt{p_i} \in [\varphi_j, \varphi_{j-1})} p_i (|\beta_1^{(i,j)}|^2 + |\beta_0^{(i,j)}|^2) t_{j+1}^2} + \mathcal{O}(\sqrt{L\epsilon} \sum_{j=1}^m t_j) \\ &= \sqrt{\sum_{j=1}^m \sum_{i:\sqrt{p_i} \in [\varphi_j, \varphi_{j-1})} p_i t_{j+1}^2} + \mathcal{O}(\sqrt{L\epsilon} \sum_{j=1}^m t_j). \end{aligned} \quad (153)$$

From [Theorem 6](#), we can imply the query complexity of [Algorithm 3](#) is

$$\begin{aligned} &\tilde{\mathcal{O}} \left(t_m + \frac{T_{\text{avg}}}{\sqrt{p_{\text{succ}}}} \right) \\ &= \tilde{\mathcal{O}} \left(t_m + \frac{\sqrt{\sum_{j=1}^m \sum_{i:\sqrt{p_i} \in [\varphi_j, \varphi_{j-1})} p_i t_{j+1}^2} + \sqrt{L\epsilon} \sum_{j=1}^m t_j}{\sqrt{\sum_{i=1}^n p_i S(\sqrt{p_i})^2}} \right) \\ &= \tilde{\mathcal{O}} \left(t_m + \sqrt{\epsilon} \sum_{j=1}^m t_j + \frac{\sqrt{\sum_{j=1}^m \sum_{i:\sqrt{p_i} \in [\varphi_j, \varphi_{j-1})} p_i t_{j+1}^2}}{\sqrt{\sum_{i=1}^n p_i S(\sqrt{p_i})^2}} \right), \end{aligned} \quad (154)$$

where the second inequality comes from $L \leq \sum_{i=1}^n p_i S(\sqrt{p_i})^2$. \square

B Other proofs

B.1 Proof of [Lemma 1](#)

Proof. Let $\Pi = |0\rangle\langle 0|_{\mathcal{H}_F}$, $\tilde{\Pi} = |1\rangle\langle 1|_{\mathcal{H}_F} \otimes I_{\mathcal{H}_W}$, then $\tilde{\Pi} \mathcal{A} \Pi$ has only one singular value $\sqrt{p_{\text{succ}}}$. Then using [Lemma 5](#), we can determine whether $\sqrt{p_{\text{succ}}}$ is larger than 2φ or smaller than φ for a given $\varphi \in (0, 1)$ with success probability at least $1 - \delta$ using $\log(\frac{1}{\delta}) \frac{1}{\varphi}$ calls to \mathcal{A} and \mathcal{A}^\dagger .

Then setting $\varphi = 1, \frac{1}{2}, \dots, \frac{1}{2^{\lceil \log(\frac{1}{L}) \rceil}}$ sequentially, we can determine whether $\sqrt{p_{\text{succ}}} \geq 2\varphi$ with success probability $1 - \frac{\delta}{\log(\frac{1}{L})}$ using $\mathcal{O}\left(\log\left(\frac{\log(\frac{1}{L})}{\delta}\right)\frac{1}{\varphi}\right)$ calls to \mathcal{A} and \mathcal{A}^\dagger , if so, stop and output 2φ .

Then with success probability at least $1 - \delta$, the algorithm will stop at $\varphi = \frac{1}{2^{\lceil \log(\frac{1}{\sqrt{p_{\text{succ}}}}) \rceil + 1}} = \Theta(\sqrt{p_{\text{succ}}})$, so the output is in $[\frac{1}{2}\sqrt{p_{\text{succ}}}, 2\sqrt{p_{\text{succ}}}]$, and the total calls to \mathcal{A} and \mathcal{A}^\dagger is

$$\mathcal{O}\left(\log\left(\frac{\log(\frac{1}{L})}{\delta}\right)\log\left(\frac{1}{\sqrt{p_{\text{succ}}}}\right)\frac{1}{\sqrt{p_{\text{succ}}}}\right). \quad (155)$$

□

B.2 Proof of Lemma 4

Proof. Let $\gamma_n := (xg(x))^{-1}(\frac{1}{a}g(\frac{1}{n}))$. For any probability distribution $\mathbf{p} = (p_i)_{i=1}^n$, let $p_M = \max_{i \in [n]} p_i$ and $i_M = \arg\max_{i \in [n]} p_i$.

Since $xg(x)$ is a convex function, we have

$$\begin{aligned} \frac{\sqrt{\sum_{i=1}^n p_i g(p_i)^2}}{\sum_{i=1}^n p_i g(p_i)} &\leq \frac{\sqrt{g(p_M)}}{\sqrt{\sum_{i=1}^n p_i g(p_i)}} \\ &= \frac{\sqrt{g(p_M)}}{\sqrt{p_M g(p_M) + \sum_{i \neq i_M} p_i g(p_i)}} \\ &\leq \frac{\sqrt{g(p_M)}}{\sqrt{p_M g(p_M) + (n-1)\frac{1-p_M}{n-1}g(\frac{1-p_M}{n-1})}} \\ &= \frac{1}{\sqrt{p_M + (1-p_M)\frac{g(\frac{1-p_M}{n-1})}{g(p_M)}}}, \end{aligned} \quad (156)$$

where the second inequality comes from $xg(x)$ is a convex function on $[0, 1]$ and Jensen's inequality.

Since $g(x)$ and $(xg(x))^{-1}$ are monotonically functions on $[0, 1]$ and $g(0) = (xg(x))^{-1}(0) = 0$, we have $\lim_{x \rightarrow 0} g(x) = \lim_{x \rightarrow 0} (xg(x))^{-1} = 0$. Thus we have $\lim_{n \rightarrow \infty} \gamma_n = \lim_{n \rightarrow \infty} (xg(x))^{-1}(\frac{1}{a}g(\frac{1}{n})) = 0$.

If $p_M \geq \gamma_n$, from Eq. (156), we have $\frac{\sqrt{\sum_{i=1}^n p_i g(p_i)^2}}{\sum_{i=1}^n p_i g(p_i)} \leq \frac{1}{\sqrt{\gamma_n}}$.

If $p_M < \gamma_n$, we have

$$\lim_{n \rightarrow \infty} \left(\frac{1}{\gamma_n} \left(p_M + (1-p_M)\frac{g(\frac{1-p_M}{n-1})}{g(p_M)} \right) \right) > \lim_{n \rightarrow \infty} (1 - \gamma_n) \frac{g(\frac{1-\gamma_n}{n-1})}{\gamma_n g(\gamma_n)} = \lim_{n \rightarrow \infty} \frac{g(\frac{1}{n})}{g(\gamma_n)\gamma_n} = 2, \quad (157)$$

where the first equation comes from convex function is continuous and $\lim_{n \rightarrow \infty} \gamma_n = 0$, and the second equation comes from $\gamma_n g(\gamma_n) = \frac{1}{a}g(\frac{1}{n})$. Then we can imply that for sufficiently large n ,

$$\frac{\sqrt{\sum_{i=1}^n p_i g(p_i)^2}}{\sum_{i=1}^n p_i g(p_i)} \leq \frac{1}{\sqrt{p_M + (1-p_M)\frac{g(\frac{1-p_M}{n-1})}{g(p_M)}}} < \frac{1}{\sqrt{\gamma_n}}. \quad (158)$$

Therefore, we can imply that $\frac{\sqrt{\sum_{i=1}^n p_i g(p_i)^2}}{\sum_{i=1}^n p_i g(p_i)} = \mathcal{O}\left(\frac{1}{\sqrt{\gamma_n}}\right)$ as $n \rightarrow \infty$.

If we choose $p_1^{(n)} = \gamma_n$ and $p_i^{(n)} = \frac{1-\gamma_n}{n-1}$ for $i = 2, \dots, n$, we have

$$\begin{aligned}
\lim_{n \rightarrow \infty} \left(\frac{\sqrt{\gamma_n} \sqrt{\sum_{i=1}^n p_i^{(n)} g(p_i^{(n)})^2}}{\sum_{i=1}^n p_i^{(n)} g(p_i^{(n)})} \right) &= \lim_{n \rightarrow \infty} \left(\frac{\sqrt{\gamma_n} \sqrt{\gamma_n g(\gamma_n)^2 + (1-\gamma_n)g\left(\frac{1-\gamma_n}{n-1}\right)^2}}{\gamma_n g(\gamma_n) + (1-\gamma_n)g\left(\frac{1-\gamma_n}{n-1}\right)} \right) \\
&= \lim_{n \rightarrow \infty} \left(\frac{\sqrt{\gamma_n} \sqrt{\gamma_n g(\gamma_n)^2 + g\left(\frac{1}{n}\right)^2}}{\gamma_n g(\gamma_n) + g\left(\frac{1}{n}\right)} \right) \\
&= \lim_{n \rightarrow \infty} \left(\frac{\sqrt{\gamma_n} \sqrt{\frac{1}{4\gamma_n} g\left(\frac{1}{n}\right)^2 + g\left(\frac{1}{n}\right)^2}}{\frac{3}{2} g\left(\frac{1}{n}\right)} \right) \\
&= \frac{1}{3}, \tag{159}
\end{aligned}$$

where the fourth equation comes from $\lim_{n \rightarrow \infty} \gamma_n = 0$.

From Eq. (158) and Eq. (159), we can imply that $\max_{\mathbf{p}=(p_i)_{i=1}^n} \frac{\sqrt{\sum_{i=1}^n p_i g(p_i)^2}}{\sum_{i=1}^n p_i g(p_i)} = \Theta\left(\frac{1}{\sqrt{\gamma_n}}\right)$ as $n \rightarrow \infty$, which completes the proof. \square

B.3 Proof of Lemma 5

Proof. We first apply Hadamard gate H to register P and obtain

$$|0\rangle_C |+\rangle_P |\psi_i\rangle_I. \tag{160}$$

Setting the parameters (δ', ϵ', t) in Lemma 8 to $\delta' := \frac{1}{2}\varphi, t := \frac{3}{2}\varphi, \epsilon' := \frac{\epsilon^2}{2}$, we can construct an even polynomial $S := P'$ in Lemma 8 with $\deg(S) = O(\log(\frac{1}{\epsilon})/\varphi)$ such that

$$\begin{aligned}
\forall x \in [-1, -2\varphi] \cup [2\varphi, 1] : S(x) &\in [0, \frac{\epsilon^2}{2}], \text{ and} \\
\forall x \in [-\varphi, \varphi] : S(x) &\in [1 - \frac{\epsilon^2}{2}, 1].
\end{aligned} \tag{161}$$

Apply $S^{(SV)}(\tilde{\Pi}U\Pi)$ to register I using register P as ancilla register. Then we will have the state $|\Phi\rangle$ in register (P, I) such that

$$\langle +|_P \otimes \Pi |\Phi\rangle = S^{(SV)}(\tilde{\Pi}U\Pi) |\psi_i\rangle_I = \left(\sum_{i=1}^d S(\sigma_i) |\psi_i\rangle \langle \psi_i| \right) |\psi_i\rangle_I = S(\sigma_i) |\psi_i\rangle_I, \tag{162}$$

where the first equation comes from $\Pi |\psi_i\rangle = |\psi_i\rangle$. Therefore, we can imply that the state $|\Phi\rangle$ satisfies

$$|\Phi\rangle = S(\sigma_i) |+\rangle_P |\psi_i\rangle_I + \sqrt{1 - S(\sigma_i)^2} |\gamma\rangle_{P,I}, \text{ where } \langle +|_P \otimes \Pi |\gamma\rangle_{P,C} = 0, \tag{163}$$

since Π is an orthogonal projection.

Next, we apply $C_{|+\rangle\langle +| \otimes \Pi} \text{NOT} = |+\rangle\langle +| \otimes \Pi \otimes X + (I - |+\rangle\langle +| \otimes \Pi) \otimes I$ to register (P, I, C) and obtain

$$\begin{aligned}
&C_{|+\rangle\langle +| \otimes \Pi} \text{NOT} |\Phi\rangle_{P,I} |0\rangle_C \\
&= (|+\rangle\langle +| \otimes \Pi \otimes X) S(\sigma_i) |+\rangle_P |\psi_i\rangle_I |0\rangle_C + (I - |+\rangle\langle +| \otimes \Pi) \sqrt{1 - S(\sigma_i)^2} |\gamma\rangle_{P,I} |\gamma\rangle_C \\
&= S(\sigma_i) |+\rangle_P |\psi_i\rangle_I |1\rangle_C + \sqrt{1 - S(\sigma_i)^2} |\gamma\rangle_{P,I} |0\rangle_C,
\end{aligned} \tag{164}$$

where the first equation comes from $(\langle +|_P \otimes \Pi) |\gamma\rangle_{P,C} = 0$ and $|+\rangle\langle +| \otimes \Pi$ is an orthogonal projector.

From Eq. (161), we can imply that

$$\forall \sigma_i \in [0, \varphi] : \sqrt{1 - S(\sigma_i)^2} \leq \sqrt{1 - \left(1 - \frac{\epsilon^2}{2}\right)^2} \leq \sqrt{1 - (1 - \epsilon^2)} = \epsilon \quad (165)$$

$$\forall \sigma_i \in [2\varphi, 1] : S(\sigma_i) \leq \frac{\epsilon^2}{2} \leq \epsilon, \quad (166)$$

which completes the proof. □