

Quantum Hamilton-Jacobi Quantization and Shape Invariance

Rathi Dasgupta¹, Asim Gangopadhyaya²

Department of Physics, Loyola University Chicago, Chicago IL, USA

Abstract

Quantum Hamilton-Jacobi quantization scheme uses the singularity structure of the potential of a quantum mechanical system to generate its eigenspectrum [1, 2] and eigenfunctions, and its efficacy has been demonstrated for many well known conventional potentials [3]. Using a recent work in supersymmetric quantum mechanics we prove that the solvability of all conventional potentials with the quantum Hamilton-Jacobi formalism follows from their shape invariance.

Introduction

Quantum Hamilton-Jacobi quantization (QHJ) introduced by Leacock and Padgett [1] and Gozzi [2] is an elegant method to determine eigenspectra for quantum mechanical systems from the singularity structure of the underlying potential. In a series of papers, Kapoor et. al. [3] have demonstrated the efficacy of the formalism to determine eigenvalues and eigenfunctions of several well known conventional potentials.

In supersymmetric quantum mechanics (SUSYQM) [4, 5, 6], it is well known that shape invariance [7, 8, 9] leads to solvability; i.e., the spectra for such quantum mechanical systems can be determined without solving the Schrödinger differential equation. Furthermore, in a recent work [10] it was proved that shape invariance suffices to determine the spectrum of a quantum system in deformation quantization formalism. So, it is fair to ask whether the shape invariance is a sufficient condition for the solvability of a quantum system in the QHJ quantization as well.

¹e-mail: rathi.dasgupta@gmail.com

²e-mail: agangop@luc.edu

In Ref. [11] the authors showed that QHJ and the shape invariance condition do help determine the spectrum for superpotentials that are either algebraic or exponential functions of the coordinate. However, following from Refs. [12, 13, 14] we now know the necessary forms for all conventional superpotentials. Employing the notations developed in [15], we here show that additive shape invariance is indeed a sufficient condition to determine spectra for all conventional potentials using the QHJ formalism. Using the formalism developed in Ref. [16] that was used to derive eigenfunctions of specific potentials, we also show that QHJ can be used to derive the eigenfunctions of the shape invariant systems without specifying the exact form of the potentials.

Our layout for this paper is as follows: we first introduce the QHJ formalism and briefly describe SUSYQM. In particular, we describe shape invariance and its implication for all conventional potentials. Using the notation developed in Ref. [15], we list all shape invariant classes that lead to the conventional potentials and determine their spectra without ever invoking their specific functional forms. We end our discussion with the derivation of eigenfunction for a class of superpotential that is a fair representative of all conventional potentials. In Table 1, we have listed shape invariant potentials in their general forms as derivable from shape invariance condition [12, 13, 14], their energies and corresponding conventional potentials.

QHJ and SUSYQM

Quantum Hamilton-Jacobi Quantization

In one-dimensional QHJ [1, 2] one works with the quantum momentum function $p(x)$ (QMF) of a particle defined³ by $p(x) = -\hbar\psi'(x)/\psi(x)$, where ψ is the wave function of particle and prime denotes differentiation with respect to the coordinate x .

³Note that this definition differs from that of ref. [1, 3] where it is define as $p(x) = -i\frac{\psi'(x)}{\psi(x)}$.

Substituting $p(x)$ in the Schrödinger equation ⁴

$$-\hbar^2\psi'' + (V(x) - E)\psi = 0 , \quad (1)$$

we get the following quantum Hamilton-Jacobi equation

$$p^2(x) - \hbar p'(x) = V(x) - E . \quad (2)$$

From Eq. (2), we see that the function $p(x)$ inherits some of its singularity structure from the potential $V(x)$. In Ref. [1, 3], the authors have shown that the singularity structure of the function $p(x)$ determines the eigenvalues of the Hamiltonian.

In supersymmetric quantum mechanics, similar to the algebraic method for the harmonic oscillator, a Hamiltonian is written as a product of two linear differential operators $A^\pm \equiv \mp \hbar d/dx + W(x)$. The function $W(x)$ is known as the superpotential. The product A^+A^- generates the Hamiltonian H_- :

$$H_- = -\hbar^2 \frac{d^2}{dx^2} + W^2(x) - \hbar W'(x) , \quad (3)$$

where the corresponding potential is $V_-(x) = W^2(x) - \hbar W'(x)$. The system is said to have unbroken supersymmetry if the groundstate energy $E_0^- = 0$. In that case, the superpotential can be written as $W(x) \equiv -\hbar [\psi_0^{(-)}(x)]' / \psi_0^{(-)}(x)$. In this paper, we will assume that supersymmetry is unbroken, and hence we have the the following limit for $p(x)$:

$$\lim_{E \rightarrow 0} p(x) \rightarrow W(x) , \quad (4)$$

which often plays an extremely important role in choosing the correct residues. In terms of the superpotential, the quantum Hamilton-Jacobi equation becomes

$$p^2(x) - \hbar p'(x) = W^2(x) - \hbar W'(x) - E . \quad (5)$$

The quantum Hamilton-Jacobi method is centered around analyzing the pole structure of the $p = -\hbar \left(\frac{\psi'}{\psi} \right)$. Since from the nodal theorem the n -th eigenfunction has n nodes within

⁴We have set $2m = 1$.

the classical turning points of the potential for a given energy, the corresponding momentum function $p(x)$ will have n singular points where these eigenfunctions vanish. These singular points can be shown to be simple poles [3], each with a residue of $-\hbar$. These are known as the “moving poles” and lead to the following quantization condition:

$$\oint p(x) dx = -2\pi i n \hbar , \quad (6)$$

where the above contour closely hugs and encloses all n -moving poles of the system in a counter-clockwise direction. These moving poles are always on the real axis and between the classical turning points for a given energy ⁵.

A counter-clockwise contour enclosing the moving singularities can also be viewed as a clockwise contour enclosing the “fixed singularities” that are outside of the contour hugging the moving poles; these are the singularities of the potential $V_-(x)$ [1, 3]. Hence, we express the quantization condition as

$$\oint p(x) dx \Big|_{\text{fixed poles}} = +2\pi i n \hbar , \quad (7)$$

where the difference in sign between Eqs. (6) and (7) is due to the change in direction in which the integration is carried out.

It is known that if a superpotential satisfies the shape invariance condition, then the eigenfunctions and eigenvalues of the Hamiltonian can be derived exactly and using algebraic methods of SUSYQM without solving the 2nd order Schrödinger differential equation [19, 20].

In Ref. [11], which was published before the necessary forms of shape invariant superpotentials were known [13, 14, 15], the authors showed that the QHJ formalism leads to the determination of the spectrum for shape invariant superpotentials that were either an algebraic or exponential functions of the coordinate. In this paper, we in fact show that shape invariance is sufficient for the QHJ formalism to determine the spectrum for all conventional potentials. We also exemplify that the forms of the potentials derived from shape invariance

⁵We note here that we are working with quantum systems in which the Hamiltonian is time independent, and wavefunctions are separable in space and time coordinates. We do not consider wavepackets of the type discussed in Refs. [17, 18].

suffices to determine the eigenfunctions of the systems. In the next section we describe shape invariance and its implication for conventional potentials.

Shape Invariance and Conventional Potentials

A SUSYQM system is exactly solvable; i.e., its eigenvalues and eigenfunctions can be explicitly written in terms of the parameters of the system [19, 20] if the superpotential satisfies the following Shape Invariance condition [7, 8, 9]:

$$W^2(x, a_0) + \hbar \frac{dW(x, a_0)}{dx} + g(a_0) = W^2(x, a_1) - \hbar \frac{dW(x, a_1)}{dx} + g(a_1). \quad (8)$$

The constants a_0 and a_1 are parameters of the potential, and are related by $a_1 = a_0 + \hbar$ as we restrict our discussion to the additive shape invariance condition, and the resulting W s cover all conventional potentials. Their eigenvalues are given by $E_n = g(a_0 + n\hbar) - g(a_0)$ [19, 20] .

In Refs. [12, 13], expanding Eq. (8) in powers of \hbar , authors showed that Eq. (8) lead to the following two partial differential equations:

$$W \frac{\partial W}{\partial a} - \frac{\partial W}{\partial x} + \frac{1}{2} \frac{dg}{da} = 0 ; \quad \text{and} \quad (9)$$

$$\frac{\partial^3}{\partial a^2 \partial x} W(x, a) = 0 . \quad (10)$$

The derivation was based on the assumption that the superpotentials were not explicit functions of \hbar . The solutions of these equations generate the complete set of conventional potentials and include all potentials listed in [7, 21].

The first of the two equations, Eq. (9) gives

$$\frac{dg}{da} = 2 \left(\frac{\partial W}{\partial x} - W \frac{\partial W}{\partial a} \right) . \quad (11)$$

To avoid level-crossing, the energies $E_n = g(a_0 + n\hbar) - g(a_0)$ must increase with n . Hence, we must have $\frac{dg(a)}{da} > 0$, which often determines the sign of a . The Eq. (10) implies that all conventional superpotentials must be of the form:

$$W(x, a) = af_1(x) + f_2(x) + u(a) . \quad (12)$$

The above form of the superpotential was originally guessed by Infeld and Hull [7], and had been used as an ansatz for decades.

Note: We assume f_1 and f_2 are linearly independent of each other as otherwise we could incorporate the second term into the first. Similarly, we also assume that $u(a)$ cannot have any linear or a constant term in a , as otherwise we could absorb them into $f_1(x)$ and $f_2(x)$ respectively.

Substituting $W = af_1(x) + f_2(x) + u(a)$ into Eq. (8), we find that functions f_1 and f_2 must satisfy

$$f_1' = f_1^2 - \lambda \quad \text{and} \quad f_2' = f_1 f_2 - \varepsilon, \quad (13)$$

where λ and ε are constants. Here we list some of the properties of f_1 and f_2 :

- The functions f_1 and f_2 are monotonic functions of x .
- If $\lambda \neq 0$, by scaling f_1 , f_2 and x , without loss of generality, we can set $|\lambda| = 1$;
- If $\lambda \neq 0$, the constant ε can be set equal to zero;
- If $\lambda \neq 0$, for every solution f_1 , the function λ/f_1 is also a solution of $f_1' = f_1^2 - \lambda$.

Let us now classify all possible conventional potentials that can be generated from Eq. (12).

Class I: $f_1 = \alpha$ is a constant.

Class II: f_2 is a constant.

Class III: Neither f_1 nor f_2 is a constant.

These three classes comprise the complete list of all conventional superpotentials, and include all cases tabulated in [7] and [21]. In Table. 1, we have listed these three classes, their energies and various conventional potentials they correspond to.

In the next section, we will apply QHJ quantization condition to determine the eigen-spectrum for each of these three classes, and thus establish that additive shape invariance is indeed a sufficient condition for QHJ to generate the eigenvalues for all conventional potentials.

Spectrum Generation for Conventional Potentials using QHJ

In this section, we determine the spectra for the three classes discussed in the previous section, and thus show that shape invariance implies exact solvability for conventional potentials under QHJ quantization. We will consider each class separately.

Class I: $f_1 = \alpha$ is a constant

In this case the superpotential is given by $W(x, a) = af_1(x) + f_2(x) = a\alpha + f_2(x)$. For this class, the Eq. (5) reduces to

$$p^2 - \hbar \frac{dp}{dx} - a^2\alpha^2 - (2a\alpha - \alpha)f_2 + E_n - f_2^2 - \varepsilon = 0, \quad (14)$$

where we have suppressed the dependence of the function f_2 on coordinate x . To compute the integral $\oint p dx$, we will change variable to $f_2(x)$, and the integral becomes $\oint p df_2/f_2'$ ⁶.

The Eq. (14) transforms to

$$p^2 - \frac{dp}{df_2}(\alpha f_2 - \varepsilon) - a^2\alpha^2 - (2a - 1)\alpha f_2 + E_n - f_2^2 - \varepsilon = 0, \quad (15)$$

where in the last equation the factor $(\alpha f_2 - \varepsilon)$ appears as we use the chain rule $dp/dx = dp/df_2 \times df_2/dx$. At this point, to go further we divide this subclass into two cases: $\alpha = 0$ and $\alpha \neq 0$.

Class IA: $\alpha = 0$

In this case, $df_2/dx = -\varepsilon$, and hence Eq. (14) reduces to

$$p^2 - (-\varepsilon)\hbar \frac{dp}{df_2} + E_n - f_2^2 - \varepsilon\hbar = 0. \quad (16)$$

Since the potential function $f_2^2 + \varepsilon$ has a singularity at infinity on the complex f_2 -plane, to determine residue at infinity we substitute $z = 1/f_2$. Replacing $\frac{dp}{df_2}$ by $dp/dz \times dz/df_2 =$

⁶As we change variable from $x \rightarrow f_2(x)$, the function $p(x)$ should transform into a new function $\bar{p}(f_2(x))$. However, with some abuse of the language, we keep the same notation of p and believe that the assumed transformation of the function p should be clear from the context.

$(-z^2)dp/dz$, we get

$$p^2 + \varepsilon\hbar(-z^2)\frac{dp}{dz} + E_n - \left(\frac{1}{z}\right)^2 - \varepsilon\hbar = 0 . \quad (17)$$

To derive the residue at $z = 0$, we substitute

$$p = \frac{b_0}{z} + a_0 + a_1 z + \dots ,$$

which yields

$$a_0^2 + 2a_0a_1z + \frac{2a_0b_1}{z} + z^2(a_1^2 - a_1\varepsilon\hbar) + 2a_1b_1 + \frac{b_1^2 - 1}{z^2} + b_1\varepsilon\hbar + E_n - \varepsilon\hbar = 0 .$$

Hence, we have $b_1 = \pm 1$, $a_0 = 0$ and $a_1 = -E_n/2$. Since near $z = 0$, the superpotential $W(x, a) = a\alpha + f_2(x) = a\alpha + 1/z \approx 1/z$, we choose $b_1 = 1$. When we change the variables from x to f_2 , and then from f_2 to z , the corresponding differentials are related by $dx = df_2/(-\varepsilon) = dz/(\varepsilon z^2)$. Thus, for the integral $\oint p(z)dz/(\varepsilon z^2) = 2\pi i n\hbar$, we only need to determine the coefficient a_1 . Hence, we get

$$E_n = -2n\varepsilon\hbar = n\omega\hbar , \quad (18)$$

where we have set $-2\varepsilon = \omega$. Thus, this subclass corresponds to the one-dimensional oscillator.

Class IB: $\alpha \neq 0$

With $\alpha \neq 0$, the superpotential for this class is $W(x, a) = a\alpha + f_2(x)$. For unbroken supersymmetry, for this case the derivative W' must be positive. For the derivative $f_2' = \alpha f_2 - \varepsilon$, the constant ε can be set equal to zero when $\alpha \neq 0$ ⁷. Hence, the positivity of W' can be achieved, without loss of generality, by choosing $\alpha = -1$ and f_2 to be negative. Thus, Eq. (14) for this case becomes

$$p^2 + \hbar f_2 \frac{dp}{df_2} - a^2 - f_2(\hbar - 2a) + E_n - f_2^2 = 0 , \quad (19)$$

⁷The particular solution $f_2 = \varepsilon/\alpha$ can be absorbed in the first term of W with a redefinition of a .

where we have made a change of variable from x to $f_2(x)$. Setting $p = \frac{b_1}{f_2} + a_0 + a_1 f_2 + \dots$, near $f_2 \approx 0$ we get

$$\frac{b_1^2}{f_2^2} + \frac{(2a_0 b_1 - b_1 \hbar)}{f_2} + a^2 + a_0^2 + 2a_1 b_1 + E_n = 0 . \quad (20)$$

This leads to $b_1 = 0$ and $a_0 = -\sqrt{a^2 - E_n}$, where we have chosen the sign of a_0 using $\lim_{E_n \rightarrow 0} p \rightarrow W$. Since the change of variable transforms, dx into df_2/f_2 , the non-zero contribution come from the a_0 and we do not need to determine any other coefficient.

The potential also has a singularity at $|f_2| \rightarrow \infty$, hence we change variable to $z = 1/f_2$. So, setting

$$p = \frac{b_0}{z} + a_0 + a_1 z + \dots ,$$

near $z \approx 0$ we get

$$\frac{b_1^2 - 1}{z^2} + \frac{2a + 2a_0 b_1 + b_1 \hbar - a^2 - \hbar}{z} + a_0^2 + 2a_1 b_1 + E_n = 0 ,$$

which is solved by $b_1 = 1$ and $a_0 = -a$. With the change of variable from x to z , the differential $dx = dz/z$, and hence only the a_0 -term contributes again. Thus, the two singularities at $f_2 = 0$ and $z = 0$ and Eq. (7) give

$$2\pi_1 \left(-\sqrt{a^2 - E_n} - a \right) = 2\pi_1 n \hbar ,$$

which then yields

$$E_n = a^2 - (a + n\hbar)^2 ,$$

which with the identification $a = -A$, produces the correct energy for the Morse potential:

$$E_n = A^2 - (A - n\hbar)^2 .$$

Class II: f_2 is a constant

The superpotentials for this class are of the form $W(x, a) = a f_1(x) + B/a$ with $f_1'(x) = f_1^2(x) - \lambda$. The QHJ Eq. (5) for this case is given by

$$p^2 - \hbar \left(f_1^2 - \lambda \right) \frac{dp}{df_1} - \frac{B^2}{a^2} - f_1^2 \left(a^2 - a\hbar \right) - a\lambda\hbar - 2Bf_1 + E_n = 0. \quad (21)$$

There are two possibilities: $\lambda = 0$ and $\lambda \neq 0$. We first consider the case of $\lambda = 0$.

Class IIA: $\lambda = 0$

For $\lambda = 0$, Eq. (21) reduces to

$$p^2 - \hbar f_1^2 \frac{dp}{df_1} - \frac{B^2}{a^2} - f_1^2 (a^2 - a\hbar) - 2Bf_1 + E_n = 0. \quad (22)$$

The potential is unbounded for large values of f_1 . Furthermore, since the transformation from $x \rightarrow f_1$ sends $dx \rightarrow df_1/f_1^2$, we need to analyze the structure of p near both $f_1 = 0$ and $f_1 = \infty$. We first look for the singularity near $f_1 = 0$. Substituting

$$p = \frac{b_1}{f_1} + a_0 + a_1 f_1 + \dots$$

in Eq. (21), near $f_1 \approx 0$ we get

$$\frac{b_1^2}{f_1^2} + \frac{2a_0 b_1}{f_1} - \frac{B^2}{a^2} + a_0^2 + b_1 \hbar + E_n = 0. \quad (23)$$

Thus, we get $b_1 = 0$, $a_0 = \sqrt{B^2 - a^2 E_n}/a$ and $a_1 = aB/\sqrt{B^2 - a^2 E_n}$. Since the differentials $dx = df_1/f_1^2$, the nonzero contribution near $f_1 = 0$ comes from the coefficient a_1 .

For the singularity at infinity, we substitute $z = 1/f_1$

$$p^2 + \hbar \frac{dp}{dz} + E_n - \frac{B^2}{a^2} - \frac{a^2 - a\hbar}{z^2} - \frac{2B}{z} = 0.$$

Substituting

$$p = \frac{b_1}{z} + a_0 + a_1 z + \dots,$$

we get

$$\frac{-a^2 + a\hbar + b_1^2 - b_1 \hbar}{z^2} + \frac{2a_0 b_1 - 2B}{z} + a_0^2 + 2a_0 a_1 z + a_1^2 z^2 + 2a_1 b_1 + a_1 \hbar - \frac{B^2}{a^2} + E_n = 0.$$

This gives $b_1 = a$. For the change of variable from $x \rightarrow f_1 \rightarrow z$, we have $dz = -df_1/f_1^2 = -(f_1^2 dx)/f_1^2 = -dx$. Hence, only the b_1 -term contributes near $z = 0$. Substituting the contributions from both singularities, we get

$$2\pi i \left(\frac{aB}{\sqrt{B^2 - a^2 E_n}} - a \right) = 2\pi i n \hbar,$$

which leads to

$$E_n = \frac{B^2}{a^2} - \frac{B^2}{(a + n\hbar)^2},$$

the correct energy for the Coulomb potential.

Class IIB: $\lambda \neq 0$

The superpotentials for this class are of the form $W(x, a) = af_1(x) + B/a$ with $f_1'(x) = f_1^2(x) - \lambda$. The general QHJ for Class II is given by Eq. (21) reads

$$p^2 - \hbar (f_1^2 - \lambda) \frac{dp}{df_1} - \frac{B^2}{a^2} - f_1^2 (a^2 - a\hbar) - a\lambda\hbar - 2Bf_1 + E_n = 0. \quad (24)$$

Since this potential has singularity at $f_1 = \infty$, and because the change of variable from x to f_1 implies $dx \rightarrow df_1/f_1' = df_1/(f_1^2 - \lambda)$, we have singularities at $f_1 = \pm\sqrt{\lambda}$ and $f_1 \rightarrow \infty$. Since for $\lambda \neq 0$, we can set its magnitude to be equal to one, henceforth we consider cases with $\lambda = \pm 1$.

We now consider the case $\lambda = 1$ and $a > 0$. Expanding p about $f_1 = 1$ as

$$p = \frac{b_1}{f_1 - 1} + a_0 + a_1 (f_1 - 1) + \dots,$$

and substituting in Eq. (21), we get

$$\frac{b_1^2}{(f_1 - 1)^2} + \frac{2a_0b_1}{f_1 - 1} - \frac{B^2}{a^2} - a^2 + a_0^2 - 2B + E_n = 0. \quad (25)$$

This gives $b_1 = 0$ and $a_0 = \frac{\sqrt{a^4 + 2a^2B - a^2E_n + B^2}}{a}$. Since, $dx = \frac{df_1}{f_1^2 - 1} = \frac{df_1}{(f_1 + 1)(f_1 - 1)} \approx \frac{1}{2} \frac{df_1}{f_1 - 1}$, only the a_0 -term contributes.

A similar analysis around $f_1 = -1$, where $dx \approx -\frac{1}{2} df/(f_1 + 1)$, we find $b_1 = 0$ and $a_0 = \frac{\sqrt{a^4 - 2a^2B - a^2E_n + B^2}}{a}$, and again only the a_0 -term contributes.

Now we carry out the analysis near $f_1 \rightarrow \infty$ and hence set $z = 1/f_1$. We expand p in the vicinity of $z = 0$ as

$$p = a_0 + a_1z + \frac{b_1}{z} \dots$$

The Eq. (21) then becomes

$$\frac{-a^2 + a\hbar + b_1^2 - b_1\hbar}{z^2} + \frac{2a_0b_1 - 2B}{z} - \frac{B^2}{a^2} - a\hbar + a_0^2 + 2a_1b_1 + a_1\hbar + b_1\hbar + E_n = 0,$$

and its solutions are $b_1 = a$ and $b_1 = -a + \hbar$, and we choose the first solution⁸. Since $dx = -dz$, it is the b_1 -term that contributes. Thus, collecting contributions from all three

⁸Since near $z \approx 0$, $W = af_1 + B/a = \frac{a}{z} + B/a \approx \frac{a}{z}$, we choose $p \approx \frac{\hbar}{z}$.

singular points, from Eq. (7) we get

$$2\pi i \left[\frac{1}{2} \left(\frac{\sqrt{a^4 + 2a^2B - a^2E_n + B^2}}{a} - \frac{\sqrt{a^4 - 2a^2B - a^2E_n + B^2}}{a} \right) - a \right] = 2\pi i n\hbar .$$

Solving for E_n , we get

$$E_n = a^2 - (a + n\hbar)^2 + \frac{B^2}{a^2} - \frac{B^2}{(a + n\hbar)^2} .$$

This analysis was carried out for $a > 0$ and the energy generated above corresponds to the Eckart potential. For negative a , a similar analysis yields exactly the same result; i.e.,

$$E_n = a^2 - (a + n\hbar)^2 + \frac{B^2}{a^2} - \frac{B^2}{(a + n\hbar)^2} . \quad (26)$$

Setting set $a = -A$, where $A > 0$, we get

$$E_n = A^2 - (A - n\hbar)^2 + \frac{B^2}{A^2} - \frac{B^2}{(A - n\hbar)^2} .$$

This case corresponds to the hyperbolic Rosen-Morse [20].

The case for $\lambda < 0$ for which a is necessarily positive, we get

$$E_n = (a + n\hbar)^2 - a^2 + \frac{B^2}{a^2} - \frac{B^2}{(a + n\hbar)^2} ,$$

and the corresponding potential is the trigonometric Rosen-Morse [20].

Class III: Neither f_1 nor f_2 is a constant

The general form of superpotentials for this class is $W(x, a) = af_1(x) + f_2(x)$, where functions $f_1(x)$ and $f_2(x)$ obey $f_1' = f_1^2 - \lambda$ and $f_2' = f_1 f_2 - \varepsilon$, respectively. Furthermore, if $\lambda = 0$ then $\varepsilon \neq 0$, and for $\lambda \neq 0$ we can choose $\varepsilon = 0$. Consequently, this class breaks into two major subclasses based on the value of λ .

Class IIIA: $\lambda = 0$

In this case, the function $f_2(x) = B/f_1(x)$, and hence the superpotential is $W(x, a) = af_1(x) + B/f_1(x)$ ⁹. Thus, $W(x, a)$ has singularities both at $f_1 = 0$ and $f_1 = \infty$, which

⁹In this case, $f_1' = f_1^2$ and $f_2' = f_1 f_2 - \varepsilon$. The solution of the homogeneous equation $f_2' = f_1 f_2$ is $f_2 = \alpha f_1$, and a particular solution is $f_2 = \frac{1}{2}\varepsilon/f_1$. Hence, the general form for f_2 is $\alpha f_1 + \frac{1}{2}\varepsilon/f_1$. The first term can be absorbed in the af_1 -term of W , and hence superpotential for this case is $W = af_1 + \frac{1}{2}\varepsilon/f_1$.

ensures that all physical moving poles will be on the positive real axis on the complex f_1 -plane. The domain of the potential is semi-infinite; i.e., $f_1 \in (0, \infty)$.

Changing variables from $x \rightarrow f_1$, the QHJ condition of Eq. (5) reduces to

$$p^2 - \hbar f_1^2 \frac{dp}{df_1} - (a^2 - a\hbar) f_1^2 - 2aB - \frac{B^2}{f_1^2} - B\hbar + E_n = 0.$$

We will now expand $p(f_1)$ around the singular points at zero and ∞ . We first consider the expansion about $f_1 = 0$. Substituting

$$p(f_1) = \frac{b_1}{f_1} + a_0 + a_1 f_1 + \dots ,$$

we get the following equation near $f_1 \approx 0$ involving coefficients b_1 , a_0 and a_1 :

$$\frac{b_1^2 - B^2}{f_1^2} + \frac{2a_0 b_1}{f_1} - 2aB + a_0^2 + 2a_1 b_1 - B\hbar + b_1 \hbar + E_n = 0 .$$

From above, we get $b_1 = B$, $a_0 = 0$ and $a_1 = \frac{2aB - E_n}{2B}$. Since $dx \rightarrow df_1/f_1^2$, only the a_1 -term contributes in Eq. (7).

Now let us analyze the singularity structure of $p(f_1)$ near $z = 1/f_1 = 0$. The QHJ equation for this case is

$$p^2 + \hbar \frac{dp}{dz} + E_n - \frac{a^2 - a\hbar}{z^2} - 2aB - B^2 z^2 - B\hbar = 0 .$$

Substituting $p(z) = b_1/z + a_0 + a_1 z + \dots$, near $z \approx 0$ we get

$$\frac{-a^2 + a\hbar + b_1^2 - b_1 \hbar}{z^2} + \frac{2a_0 b_1}{z} - 2aB + a_0^2 + 2a_1 b_1 + a_1 \hbar - B\hbar + E_n = 0,$$

which yields $b_1 = a$, $a_0 = 0$ and $a_1 = \frac{2aB + B\hbar - E_n}{2a + \hbar}$. In this case, since $dx = -dz/z^2$, only the a_1 -term contributes to Eq. (7).

Putting the contributions from singularities at $f_1 = 0$ and $f_1 = \infty$ together, we get

$$\frac{1}{2} \left[2\pi i \hbar \left(\frac{2aB - E_n}{2B} - a \right) \right] = 2\pi i n \hbar , \quad (27)$$

which, with the identification $B \rightarrow -\frac{\omega}{2}$, leads to the spectrum for the radial oscillator:

$$E_n = 2n\omega\hbar .$$

The factor of $\frac{1}{2}$ in Eq. (27) is due to the fact that on the complex f_1 -plane, we have moving poles on the negative and positive sides of the real axis, while only those on the positive real axis are physical, and hence they double count in the complex integration [1, 3].

Class IIIB: $\lambda \neq 0$

In this case we have $W = af_1 + f_2$, where f_1 and f_2 satisfy $f_1' = f_1^2 - \lambda$, $f_2' = f_1 f_2 - \varepsilon$. Solving the second differential equation¹⁰ for f_2 , we get $f_2 = B\sqrt{f_1^2 - \lambda}$. Hence, the superpotential for this case reduces to $W = af_1 + B\sqrt{f_1^2 - \lambda}$.

We first consider $\lambda > 0$, which implies that we can set $\lambda = 1$. Furthermore, from Eq. (11) and the requirement that $dg/da > 0$ implies that we must have $a < 0$.

To avoid working with the square root of f_1 in the superpotential, following [15], we define a function y by

$$y(x) = \sqrt{\frac{f_1(x) - 1}{f_1(x) + 1}},$$

which gives $dy/dx = y$. The function f_1 and the superpotential W are then given by

$$f_1 = -\frac{y + y^{-1}}{y - y^{-1}} \quad \text{and} \quad W = -a \left(\frac{y + y^{-1}}{y - y^{-1}} \right) + \frac{2B}{y - y^{-1}}.$$

This change of variable that casts the superpotential as a function of both y and y^{-1} results in singularities at both $y = 0$ and $y = \infty$, and thus the domain reduces to $y \in (0, \infty)$. Hence, on the complex y -plane all physical moving poles will necessarily be on the positive real y -axis. The QHJ condition of Eq. (5) becomes

$$\begin{aligned} p^2 & - \hbar y \frac{dp}{dy} + E_n + \frac{-a^2 + y(4aB - 2B\hbar) + y^2(-2a^2 + 4a\hbar - 4B^2)}{(y^2 - 1)^2} \\ & + \frac{y^3(4aB - 2B\hbar) - a^2y^4}{(y^2 - 1)^2} = 0. \end{aligned} \quad (28)$$

Thus, in addition to zero and infinity, we also have singularities at $y = \pm 1$. We will need to compute residues at all four singular points.

¹⁰The homogeneous solution of $f_2' - f_1 f_2 = \varepsilon$ is $f_2 = B \exp \left[\int f_1 dx \right] = B \exp \left[\int f_1 \frac{df_1}{f_1^2 - \lambda} \right] = B \exp \left[\frac{1}{2} \int \frac{df_1^2}{f_1^2 - \lambda} \right] = B\sqrt{f_1^2 - \lambda}$. A particular solution is $\left(\frac{\varepsilon}{\lambda}\right) f_1$. Given that the particular solution is a multiple of f_1 , we only retain the homogeneous solution; i.e., $f_2 = B\sqrt{f_1^2 - \lambda}$. The particular solution can be absorbed in the $a f_1$ -term in W .

Near $y = 0$, setting $p = \frac{b_1}{y} + a_0 + a_1 y + \dots$, the Eq. (28) reduces to

$$\frac{b_1^2}{y^2} + \frac{2a_0 b_1 + b_1 \hbar}{y} - a^2 + a_0^2 + 2a_1 b_1 + E_n = 0,$$

and hence we have $b_1 = 0$ and $a_0 = -\sqrt{a^2 - E_n}$.¹¹ Since under $x \rightarrow y$, the differentials are connected by $dx \rightarrow dy/y$, only the a_0 -term contributes.

Near $y = 1$, setting $u = y - 1$ and $p = b_1/u + a_0 + a_1 u + \dots$, the Eq. (28) reduces to

$$-\frac{(-a + B - b_1)(-a + B + b_1 + \hbar)}{u^2} + \frac{\hbar(b_1 - 2(-a + B)) - 2(-a + B)^2 + 2a_0 b_1}{u} + \mathcal{O}(u^0) = 0,$$

and hence we have $b_1 = -a + B$. Since near $y = 1$, the differentials are connected by $dx \rightarrow dy/y = dy$, only the b_1 -term contributes.

Near $y = -1$, setting $u = y + 1$ and $p = b_1/u + a_0 + a_1 u + \dots$, the Eq. (28) reduces to

$$\frac{(a + B - b_1)(-a - B - b_1 + \hbar)}{u^2} + \frac{2((-a - B)^2 + a_0 b_1) + (-2a - 2B + b_1)\hbar}{u} + \mathcal{O}(u^0) = 0,$$

and hence we have $b_1 = a + B$. Since near $y = -1$, the differentials are connected by $dx \rightarrow dy/y = -dy$, again only the b_1 -term contributes.

Finally, we look for the contribution coming from the singularity at $|y| \rightarrow \infty$, for which we set $z = 1/y$ and $p = \frac{b_1}{z} + a_0 + a_1 z + \dots$ in Eq. (28). This gives

$$\frac{b_1^2}{z^2} + \frac{b_1(2a_0 - \hbar)}{z} - a^2 + a_0^2 + 2a_1 b_1 + E_n = 0,$$

which gives $b_1 = 0$ and $a_0 = \sqrt{a^2 - E_n}$. In this case, $dx \rightarrow dy/y \rightarrow -dz/z$, and hence only the a_0 -term contributes and we do not need to determine any other coefficient.

Thus the net contribution of the four singular points is

$$-\sqrt{a^2 - E_n} + (-a + B) + (-)(a + B) + (-)\sqrt{a^2 - E_n} = -2 \left(a + \sqrt{a^2 - E_n} \right)$$

As stated earlier, in this problem all n physical moving poles lie on the positive side of real axis on the complex y -plane. However, there are n ‘‘spurious’’ moving poles on the negative

¹¹Since $p \rightarrow W$ as we set $E_n = 0$, and because $W \rightarrow a$ near $y \approx 0$ and $a < 0$, we choose $a_0 = -\sqrt{a^2 - E_n}$.

real axis as well, with each contributing a residue of $i\hbar$ to the right-hand-side of Eq. (7).

Hence, we have

$$2\pi i \left[-2 \left(a + \sqrt{a^2 - E_n} \right) \right] = 2\pi i (2n\hbar) ,$$

which gives $E_n = a^2 - (a + n\hbar)^2$. Identifying $a = -A$, we get the energy for hyperbolic version of Pöschl-Teller potential:

$$E_n = A^2 - (A^2 - n\hbar)^2 .$$

Following a similar procedure, we can show that the case of $\lambda = -1$ leads to the trigonometric Pöschl-Teller potential with eigenvalues

$$E_n = (A^2 + n\hbar)^2 - A^2 .$$

Derivation of eigenfunctions for a Class IIB superpotential

In Ref. [16], authors derive eigenfunctions of several conventional potentials. In this section, we show that it is indeed possible to derive the eigenfunctions of a shape invariant conventional potential directly from Eq. (12), without specifying the actual form of the superpotential. As an illustrative example, we derive eigenfunctions of a ClassIIB superpotential, which is described in sufficient detail so that it could be used to derive eigenfunctions for any other class.

Superpotentials in this class have the form $W = af_1 + B/a$, where f_1 satisfies

$$\frac{df_1}{dx} = f_1^2 - \lambda \tag{29}$$

Depending on values of λ and signs of a , it produces different conventional potentials. We will now consider the case¹² with $a < 0$ and $\lambda = 1$. The needed boundary conditions for W at the left and the right boundaries also require that we have $B < 0$.

¹²If $\lambda \neq 0$, with scaling of f_1 and x , we can make $|\lambda| = 1$. Thus, λ takes only three values: 0, 1 and -1.

From the singularities of the quantum momentum function, we have already seen that eigenenergies are given by

$$E_n = a^2 - (a + n\hbar)^2 + \frac{B^2}{a^2} - \frac{B^2}{(a + n\hbar)^2} , \quad (30)$$

which we will soon use. Going forward, we also set $\hbar = 1$. As stated earlier, the quantum momentum function satisfies

$$p^2(x) - p'(x) = W^2(x) - W'(x) - E . \quad (31)$$

The superpotential W is given in terms of f_1 , so it is natural to ask whether f_1 could be used as an independent variable instead of x . Since, in general, both f_1 and f_2 are monotonic functions of x , they indeed are good variables to use to express the QHJ equation, and we will do so in this case. However, since f_1 is a monotonically decreasing function¹³, we do a change of variable from $x \rightarrow S = -f_1$ and the quantum momentum function $p(x) \rightarrow \tilde{p}(S)$ then satisfies

$$\tilde{p}^2(S) - \dot{\tilde{p}}(S) \frac{dS}{dx} = (-aS + B/a)^2 - (S^2 - 1) - E , \quad (32)$$

where the dot over \tilde{p} denotes a differentiation with respect to S . Following Ref. [16], we define a function χ by

$$\chi = \frac{\tilde{p}}{F} - \frac{1}{2} \frac{\dot{F}}{F} , \quad (33)$$

where F stands for $\frac{dS}{dx}$. In variables χ and S , the QHJ equation becomes

$$\chi^2 - \hbar\dot{\chi} + \frac{1}{4} \left(\frac{\dot{F}^2}{F^2} - \frac{2\ddot{F}}{F} \right) - \left(\frac{W^2 - \hbar W' - E_n}{F^2} \right) = 0 . \quad (34)$$

Substituting $W = a(-S) + B/a$, we arrive at

$$\chi^2 - \hbar\dot{\chi} + \left(-\frac{B^2}{a^2} + \frac{(a-1)aS^2 + a - 2BS - E_n}{F^2} + \frac{S^2}{F^2} + \frac{1}{F} \right) = 0 . \quad (35)$$

¹³In this case, we have $W' = af'_1 = a(f_1^2 - 1)$. The relation $f'_1 = (f_1^2 - 1)$ implies that f_1 can never be equal to ± 1 at any point as otherwise f_1 will be constant. Thus, $f_1^2 - 1$ must have a fixed sign, and hence W must be monotonic function of x . However, from unbroken supersymmetry, we must have $W' > 0$ for this case. So, choosing $a < 0$ requires that f'_1 be negative as well.

Our objective here is to solve the above equation for χ and from there to derive the eigenfunctions ψ . Since the equation is now given in terms of S and derivatives with respect to S , our solutions for χ and ψ will be in terms of this variable.

We first note that the function χ has n moving-poles, each with a residue of (-1) . In addition to the moving poles, it also has fixed poles that are engendered by the structure of the superpotential W . As we see from Eq. (35), the fixed singularities for χ are at $S = \pm 1$. Hence, we write the function χ as

$$\chi = - \sum_{k=1}^n \frac{1}{S - S_k} + \frac{b_-}{S + 1} + \frac{b_+}{S - 1} + C, \quad (36)$$

where we have explicitly stated all of moving¹⁴ and fixed poles and C is an analytic function on the complex S -plane.

As $|S| \rightarrow \infty$, the superpotential $W \gg \sqrt{E_n}$ and the quantum momentum function $p \rightarrow W$ and the function χ goes to

$$\chi \rightarrow \left(\frac{W}{F} - \frac{2F'}{F} \right) = \left(\frac{\frac{B}{a} - aS}{1 - S^2} + \frac{\hbar S}{1 - S^2} \right) \rightarrow \frac{a - \hbar}{S}.$$

I.e., $\chi \rightarrow 0$ for large $|S|$. Hence, from the Liouville's theorem the analytic function C is bounded and must be a constant. From its value for large $|S|$, the constant C should be zero.

We substitute the expression for χ from Eq. (36) into Eq. (35) and expand separately around $S = \pm 1$. Near $S = 1$ we get two possible values for the residue:

$$b_+ = \frac{-1 - \sqrt{a^2 - 2B - E_n + \frac{B^2}{a^2}}}{2} \quad \text{or} \quad \frac{-1 + \sqrt{a^2 - 2B - E_n + \frac{B^2}{a^2}}}{2}.$$

Since, for the limit $E_n \rightarrow 0$, we must have $\chi \rightarrow \left(\frac{W}{F} + \dots \right) = \frac{a - \frac{B}{a}}{2a} + \dots$, we must choose¹⁵

$$b_+ = \frac{-1 - \sqrt{a^2 - 2B - E_n + \frac{B^2}{a^2}}}{2}.$$

¹⁴The term with a summation on the right-hand-side comes from the n moving poles.

¹⁵In the limit $E_n \rightarrow 0$,

$$\frac{-1 - \sqrt{a^2 - 2B - E_n + \frac{B^2}{a^2}}}{2a^2} \rightarrow \frac{1}{2} \left(-\sqrt{\frac{(a^2 - B)^2}{a^2}} - 1 \right) = \frac{1}{2} \left(-\frac{B - a^2}{a} - 1 \right) = \frac{1}{2} \left(-\frac{B}{a} + a - 1 \right),$$

exactly as χ should behave near $S = 1$.

With a similar reasoning, we see that we must have $b_- = \frac{1}{2} \left(\frac{\sqrt{a^4 + 2a^2 B - a^2 E_n + B^2}}{a} - 1 \right)$.

At this point, we substitute $E_n = a^2 - (a+n)^2 + \frac{B^2}{a^2} - \frac{B^2}{(a+n)^2}$ that we derived using the singularity of QMF, see Eq. (26), which gives

$$b_{\mp} = \frac{1}{2} \left(\pm \frac{B}{a+n} + a + (n-1) \right).$$

The first term on the right-hand-side of Eq. (36) can be written in terms of the logarithmic derivative of a n -th order polynomial $P = \prod_{k=1}^n (S - S_k)$, where S_k denotes the zeroes of the polynomial. Then, we can write $\sum_{k=1}^n \frac{1}{S-S_k} = \frac{\dot{P}}{P}$, and hence we have

$$\chi = -\frac{\dot{P}}{P} + \frac{b_-}{S+1} + \frac{b_+}{S-1}. \quad (37)$$

Substituting this form of χ in Eq. (35), we get an equation for the polynomial P :

$$(1-S^2) \ddot{P} + \left(2(a+n-1)S - \frac{2B}{a+n} \right) \dot{P} - n(2a+n-1)P = 0. \quad (38)$$

Defining

$$\alpha = \frac{B}{a+n} - a - n \quad \text{and} \quad \beta = -\frac{B}{a+n} - a - n,$$

the equation reduces to the familiar Jacobi differential equation:

$$(1-S^2) \ddot{P} + (\beta - \alpha - (\alpha + \beta + 2)S) \dot{P} + n(n + \alpha + \beta + 1)P = 0, \quad (39)$$

and its solutions are the Jacobi Polynomials $P_n^{(\alpha, \beta)}(S)$. In terms of α and β , the function χ is now given by

$$\chi = -\frac{\dot{P}_n}{P_n} + \frac{\frac{1}{2}(-\beta-1)}{S+1} + \frac{\frac{1}{2}(-\alpha-1)}{S-1}. \quad (40)$$

The eigenfunctions are then given by

$$\begin{aligned} \psi_n &\sim e^{-\int p dx} \sim e^{-\int \tilde{p} dS/S'} = e^{-\int \left(\chi + \frac{1}{2} \frac{\dot{P}}{P} \right) dS} \\ &\sim e^{\int \left(\frac{\dot{P}_n}{P_n} + \frac{\frac{1}{2}(\beta+1)}{S+1} + \frac{\frac{1}{2}(\alpha+1)}{S-1} - \frac{1}{2} \frac{2S}{S^2-1} \right) dS} \\ &\sim e^{\left(\log P_n + \frac{1}{2}(\beta+1) \log(S+1) + \frac{1}{2}(\alpha+1) \log(S-1) - \frac{1}{2} \log(S-1) - \frac{1}{2} \log(S+1) \right)} \\ &= N (1+S)^{\frac{\beta}{2}} (1-S)^{\frac{\alpha}{2}} P_n^{(\alpha, \beta)}(S) \end{aligned} \quad (41)$$

For the specific case of $S = \tanh x$, $-a = A$ and $-B \rightarrow B$, the superpotential goes to $W = A \tanh x + B/A$, the superpotential for hyperbolic Rosen-Morse or Rosen-Morse II and the result matches with that of Ref. [22].

Class	Superpotential	Energy E_n	Name
IA $\alpha = 0$	$\alpha a + f_2$ $\varepsilon = -\frac{1}{2}\omega$	$n\hbar\omega$	1D-Harmonic Oscillator
IB $\alpha = -1$	$\alpha a + f_2$ $a < 0; \varepsilon = 0$	$a^2 - (a + n\hbar)^2$	Morse
IIA $\lambda = 0$	$a f_1 + B/a$ $a > 0$	$\frac{B^2}{a^2} - \frac{B^2}{(a+n\hbar)^2}$	Coloumb
II B1 $\lambda < 0$	$a f_1 + B/a$ $a > 0$	$(a + n\hbar)^2 - a^2$ $+ \frac{B^2}{a^2} - \frac{B^2}{(a+n\hbar)^2}$	Rosen-Morse (Trigonometric)
II B2 $\lambda > 0$	$a f_1 + B/a$ $a < 0$	$a^2 - (a + n\hbar)^2$ $+ \frac{B^2}{a^2} - \frac{B^2}{(a+n\hbar)^2}$	Rosen-Morse (Hyperbolic)
II B3 $\lambda > 0$	$a f_1 + B/a$ $a > 0$	$a^2 - (a + n\hbar)^2$ $+ \frac{B^2}{a^2} - \frac{B^2}{(a+n\hbar)^2}$	Eckart
IIIA $\lambda = 0$	$a f_1 + B/f_1$ $a > 0; B = -\frac{1}{2}\omega$	$2 n\hbar\omega$	3D-Oscillator
II B1 ($\lambda < 0$)	$a f_1 + B\sqrt{f_1^2 + \lambda }$ $a > 0$	$(a + n\hbar)^2 - a^2$	Scarf (Trigonometric)
II B2 ($\lambda > 0, f_1^2 < \lambda$)	$a f_1 + B\sqrt{\lambda - f_1^2}$ $a < 0$	$a^2 - (a + n\hbar)^2$	Scarf (Hyperbolic)
II B3 ($\lambda > 0, f_1^2 > \lambda$)	$a f_1 + B\sqrt{f_1^2 - \lambda}$ $a < 0$	$a^2 - (a + n\hbar)^2$	Pöschl-Teller (Hyperbolic)

Table 1: The three shape invariant classes, their energies and the corresponding conventional potentials

Conclusion:

Within supersymmetric quantum mechanics, shape invariance of a system implies its solvability. In particular, for additive shape invariant systems, we can determine eigenvalues and eigenfunctions without solving the Schrödinger differential equation. Additionally, in a recent work [10] it was shown that the shape invariance can be utilized to determine spectrum of a quantum system in deformation quantization formalism as well. So, the question remained whether the shape invariance played any role in determining the spectra for an array of conventional potentials using QHJ, albeit on a case-by-case basis [3, 16]. In this paper, using the general form of conventional superpotentials as derived from shape invariance [12, 13, 14], we show that shape invariance is sufficient to guarantee the solvability of all conventional potentials in QHJ formalism. Thus, this paper completes the work of Ref. [11], which was published before the necessary forms of all conventional potentials were known.

In short, we note that while QHJ had been shown to yield exact eigenvalues and eigenfunctions for several shape invariant superpotentials, it wasn't clear whether the exactness for these superpotentials emanated from the shape invariance. We instead worked with the general form of W derived directly from SI-condition, and thus established that shape invariance is indeed a sufficient condition for the solvability under QHJ formalism.

Since this work makes an extensive use of the results and notations of Ref. [15], we would like to thank its authors.

References

- [1] Leacock R A and Padgett M J Phys. Rev. Lett. **50** 3, 1983; Phys. Rev. D **28** 2491, 1983; Phys. Rev. A **33** 2775, 1986; Am. J. Phys. **55**, 261, 1986.
- [2] Gozzi E, Phys. Lett. B **165**, 351, 1985.
- [3] Bhalla R S, Kapoor A K and Panigrahi P K Am. J. Phys. **65** 1187, 1997; Mod. Phys. Lett. **A 12** 295, 1997; Geojo K G, Sree Ranjani S and Kapoor A K J. Phys. A: Math. Gen. **36**, 4591-4598, 2003; S. Sree Ranjani, K.G. Geojo, A.K. Kapoor, P.K. Panigrahi, Mod. Phys. Lett. A **19**, 1457-1468, 2004.
- [4] E. Witten, Dynamical breaking of supersymmetry; Nucl. Phys. B185 (1981) 513–554.
- [5] P. Solomonson and J. W. Van Holten, Fermionic coordinates and supersymmetry in quantum mechanics; Nucl. Phys. **B196** (1982) 509–531.
- [6] F. Cooper and B. Freedman, Aspects of supersymmetric quantum mechanics; Ann. Phys. **146** (1983) 262–288.
- [7] L. Infeld and T. E. Hull, The factorization method; Rev. Mod. Phys. **23** (1951) 21–68.
- [8] W. Miller Jr, *Lie Theory and Special Functions* (Mathematics in Science and Engineering); Academic Press, New York, NY, USA, 1968.
- [9] L. E. Gendenshtein, Derivation of exact spectra of the Schrödinger equation by means of supersymmetry; JETP Lett. **38** (1983) 356–359.
- [10] C. Rasinariu, Shape invariance in phase space, Fortsch. Phys. 61 (2013) 4-19.
- [11] C. Cherqui, Y. Binder, and A. Gangopadhyaya, Shape invariance and the exactness of the quantum Hamilton-Jacobi formalism; Phys. Lett. **A 372** (2008) 1406-1415.
- [12] Gangopadhyaya, A. and Mallow, J.V. Generating shape invariant potentials. *Int. J. Mod. Phys. A* **2008**, 23, 4959–4978.

- [13] J. Bougie, A. Gangopadhyaya and J. V. Mallow, Generation of a complete set of additive shape-invariant potentials from an Euler equation; *Phys. Rev. Lett.* (2010) 210402:1–210402:4.
- [14] J. Bougie, A. Gangopadhyaya, J. V. Mallow and C. Rasinariu, Supersymmetric quantum mechanics and solvable models; *Symmetry* 4 (3) (2012) 452–473.
- [15] A. Gangopadhyaya, J. V. Mallow, C. Rasinariu and J. Bougie, Exactness of SWKB for shape invariant potentials, *Phys. Lett.* **A384** (2020) 126722.
- [16] S.S. Ranjani, K.G. Geojo, A.K. Kapoora and P.K. Panigrahi, Bound State Wave Functions through the Quantum Hamilton-Jacobi Formalism, *Mod. Phys. Lett. A.* 19 (2004) 1457.
- [17] M.V. Berry and N.L. Balazs, Nonspreading Wave Packets. *American Journal of Physics*, 4 (1979) 264-267.
- [18] P. Strange, Semiclassical and quantum analysis of a free-particle Hermite wave function, Semiclassical and quantum analysis of a free-particle Hermite wave function, *Phys. Rev. A* 89 (2014) 044101.
- [19] F. Cooper, A. Khare and U. Sukhatme, *Supersymmetry in Quantum Mechanics*, World Scientific, Singapore, 2001.
- [20] A. Gangopadhyaya, J. Mallow and C. Rasinariu, *Supersymmetric Quantum Mechanics: An Introduction* (2nd ed.), World Scientific, Singapore, 2017.
- [21] R. Dutt, A. Khare and U. Sukhatme, Supersymmetry, shape invariance and exactly solvable potentials; *Am. J. Phys.* **56** (1988) 163–168.
- [22] R. De, R. Dutt and U.P. Sukhatme, *J. Phys. A* 25 (1992) L843-850.