

MÖBIUS FUNCTION OF THE SUBGROUP LATTICE OF A FINITE GROUP AND EULER CHARACTERISTIC

FRANCESCA DALLA VOLTA AND LUCA DI GRAVINA

ABSTRACT. The Möbius function of the subgroup lattice of a finite group G has been introduced by Hall and applied to investigate several questions. In this paper, we consider the Möbius function defined on an order ideal related to the lattice of the subgroups of an irreducible subgroup G of the general linear group $\mathrm{GL}(n, q)$ acting on the vector space V . We find a relation between this function and the Euler characteristic of two simplicial complexes Δ_1 and Δ_2 , the former arising from the lattice of the subspaces of V , the latter from the subgroup lattice of G .

Keywords: Möbius function, subgroup lattice, linear groups, Euler characteristic, Simplicial complexes.

2020 MSC: 20B25, 20D60, 05E16, 05E45

1. INTRODUCTION

In this paper, motivated by recent and less recent results about the Möbius function μ for the subgroup lattice $\mathcal{L}(G)$ of a finite group G , we give a result which relates the Möbius function for a subgroup G of $\mathrm{GL}(n, q)$ to two simplicial complexes: one defined from the lattice of the subspaces fixed by a reducible subgroup $H \leq G$, the second from the lattice $\mathcal{L}(G)$ of the subgroups of G .

We will introduce in the Preliminaries all the definitions and details useful for reading the paper.

In his Ph.D thesis [8] Shareshian considers the problem of computing $\mu(1, G)$ for several finite classical groups G ; the idea is to approximate $\mu(1, G)$ through a *good function* $f_{G,n,p}(u, 1)$, such that:

$$(1) \quad \mu(1, G(n, p^u)) = f_{G,n,p}(u, 1) + \sum_{K \in \mathcal{C}_9} \mu(1, K).$$

Here $G = G(n, p^u)$ denotes a family of finite classical groups with the same defining classical form, which act in a natural way on the vector space V of finite dimension n over the finite field of order $q = p^u$. If $\mathcal{C}_1, \dots, \mathcal{C}_8, \mathcal{C}_9$ are the Aschbacher classes of maximal subgroups of a finite classical group (see

The authors are members of the Italian National Group for Algebraic and Geometric Structures and their Applications (GNSAGA - INdAM) and thank INdAM for the support. The second author is also a member of the research training group *GRK 2240: Algebro-Geometric Methods in Algebra, Arithmetic and Topology*, funded by DFG.

[4]), \mathcal{C}_9 is the class of almost simple groups not belonging to the first 8 classes of "geometric type" and the function $f_{G,n,p}(u, 1)$ provides an estimate of $\mu(1, G)$ with respect to the contributions given by the subgroups of G which belong to the classes \mathcal{C}_i , for $i \in \{1, \dots, 8\}$.

Actually, Shareshian's approach focuses on the first class $\mathcal{C}_1(G)$, that is the class of reducible subgroups of G .

In particular, the reducible subgroups of G contribute to $f_{G,n,p}(u, 1)$ through the computation of the Möbius function of

$$\widehat{\mathcal{I}}_1(G) = \{K \leq G \mid K \leq M \text{ for some } M \in \mathcal{C}_1(G)\} \cup \{G\},$$

which is obtained by adjoining the maximum G to the order ideal

$$\mathcal{I}_1(G) = \{K \leq G \mid K \leq M \text{ for some } M \in \mathcal{C}_1(G)\},$$

that is

$$\mu_{\widehat{\mathcal{I}}_1(G)}(1, G) = - \sum_{K \in \mathcal{I}_1(G)} \mu(1, K)$$

and

$$(2) \quad \mu(1, G) = \mu_{\widehat{\mathcal{I}}_1(G)}(1, G) - \sum_{\substack{K < G \\ K \notin \mathcal{I}_1(G)}} \mu(1, K).$$

In this paper, we will consider irreducible subgroups G of the general linear group $\text{GL}(n, q)$, that is groups of linear automorphisms of a vector space V of dimension n over the finite field with q elements, which fix no non-trivial subspace of V . In this hypothesis, we will take a reducible subgroup H of G (that is, H fixes some proper subspace of V) and we will work on the analogue of $\mu_{\widehat{\mathcal{I}}_1(G)}(1, G)$, namely $\mu_{\widehat{\mathcal{I}}(G,H)}(H, G)$ so that

$$(3) \quad \mu(H, G) = \mu_{\widehat{\mathcal{I}}(G,H)}(H, G) - \sum_{\substack{K \notin \mathcal{I}(G,H) \\ H \leq K < G}} \mu(H, K)$$

(see Section 2 for precise definitions).

The subject of this paper is somehow motivated by the following conjecture:

Conjecture 1.1 (Mann, [6]). *Let G be a PFG group and μ the Möbius function on the lattice of open subgroups of G . Then $|\mu(H, G)|$ is bounded by a polynomial function in the index $|G : H|$ and the number of subgroups of G of index m with $\mu(H, G) \neq 0$ grows at most polynomially in m .*

Indeed, although the problem is still open in its general setting, it was reduced by Lucchini in [5] to the study of similar growth conditions for finite almost-simple groups.

The following Theorem is the main result of the present paper:

Theorem 4.5. Consider a vector space V of finite dimension n over \mathbb{F}_q . Let $H \leq G \leq \text{GL}(V)$. Then

$$(4) \quad -\mu_{\widehat{\mathcal{I}}(G,H)}(H, G) = \sum_{E \in \Psi'(G,H)} (-1)^{|E|} = \sum_{X \in \Psi(G,H)} (-1)^{|X|} = \chi(\Delta_1) = \chi(\Delta_2)$$

$\widehat{\mathcal{I}}(G, H)$, $\Psi'(G, H)$, $\Psi(G, H)$, Δ_i , and $\chi(\Delta_i)$ ($i = 1, 2$) are defined in the following Section 1, also for irreducible subgroups H . This will allow us to avoid the restriction to reducible groups in the statement of the Theorem.

In the final Section, we will use Theorem 4.5 to deal with $\mu(H, G)$ in some particular case. In [1], Theorem 4.5 is used to attack Conjecture 1.1 for some class of subgroups H of linear and projective groups.

We thank Andrea Lucchini and Johannes Siemons for many useful discussions.

2. PRELIMINARIES

In this paper all the groups and sets are finite. For main results about posets and lattices, we refer to [9]. Here, we just recall some basic fact, useful for reading the paper.

Definition 2.1. Let \mathcal{P} be a finite poset. The **Möbius function** associated with \mathcal{P} is a map $\mu_{\mathcal{P}} : \mathcal{P} \times \mathcal{P} \rightarrow \mathbb{Z}$ satisfying

$$\mu_{\mathcal{P}}(x, y) = 0 \quad \text{unless } x \leq y,$$

and defined recursively for $x \leq y$ by

$$(5) \quad \mu_{\mathcal{P}}(x, x) = 1 \quad \text{and} \quad \sum_{x \leq t \leq y} \mu_{\mathcal{P}}(x, t) = 0 \quad \text{if } x < y.$$

Notation. If \mathcal{P} is the subgroup lattice $\mathcal{L}(G)$ of G , we will write $\mu(H, K)$ instead of $\mu_{\mathcal{L}(G)}(H, K)$.

Definition 2.2. An (abstract) **simplicial complex** Δ on a vertex set T is a collection Δ of subsets of T satisfying:

- If $t \in T$, then $t \in \Delta$,
- if $F \in \Delta$ and $G \subseteq F$, then $G \in \Delta$.

An element $F \in \Delta$ is called a face of Δ and the dimension of F is defined to be $|F| - 1$. In particular, the empty set \emptyset is a face of Δ (provided $\Delta \neq \emptyset$) of dimension -1 .

Definition 2.3. The **Euler characteristic** of Δ is so defined:

$$\chi(\Delta) = \sum_i (-1)^i \mathcal{F}_i = \mathcal{F}_0 - \mathcal{F}_1 + \mathcal{F}_2 - \mathcal{F}_3 + \dots$$

where \mathcal{F}_i denotes the number of the i -faces, that is of the faces of dimension i , in Δ .

We recall here the definition of an order ideal:

Definition 2.4. Let (\mathcal{P}, \leq) be a poset. An **order ideal** of \mathcal{P} is a subset $I \subseteq \mathcal{P}$ such that

$$(6) \quad \forall x \in I, t \in \mathcal{P} \quad t \leq x \Rightarrow t \in I.$$

In particular, if A is a subset of \mathcal{P} , then the set

$$\mathcal{P}_{\leq A} = \{s \in \mathcal{P} \mid s \leq a \text{ for some } a \in A\} \subseteq \mathcal{P}$$

is the order ideal in \mathcal{P} generated by A .

Observe that a simplicial complex on a set of vertices T is just an order ideal of the boolean algebra B_T that contains all one element subsets of T .

Definition 2.5. Let Δ be a simplicial complex, and $\chi(\Delta)$ be the Euler characteristic of Δ . The **reduced Euler characteristic** $\tilde{\chi}(\Delta)$ of Δ is defined by $\tilde{\chi}(\emptyset) = 0$ and $\tilde{\chi}(\Delta) = \chi(\Delta) - 1$ if $\Delta \neq \emptyset$; see [9], Equation (3.22).

It is possible to build a simplicial complex from a poset \mathcal{P} in the following way:

The **order complex** $\Delta(\mathcal{P})$ of \mathcal{P} is defined as the simplicial complex whose vertices are the elements of \mathcal{P} and whose k -dimensional faces are the chains $a_0 \prec a_1 \prec \dots \prec a_k$ of length k of distinct elements $a_0, \dots, a_k \in \mathcal{P}$.

Now, denote by $\hat{\mathcal{P}}$ the finite poset obtained from \mathcal{P} by adjoining a least element $\hat{0}$ and a greatest element $\hat{1}$. The Möbius function $\mu_{\hat{\mathcal{P}}}(\hat{0}, \hat{1})$ is related to $\tilde{\chi}(\Delta(\mathcal{P}))$ by a well known result by Hall [3] about the computation of $\mu_{\hat{\mathcal{P}}}(\hat{0}, \hat{1})$ by means of the chains of even and odd length between $\hat{0}$ and $\hat{1}$.

Proposition 2.6 (see [9] Proposition 3.8.6). *Let \mathcal{P} be a finite poset. Then*

$$\mu_{\hat{\mathcal{P}}}(\hat{0}, \hat{1}) = \tilde{\chi}(\Delta(\mathcal{P})).$$

3. THE IDEAL $\mathcal{I}(G, H)$ AND THE COMPLEXES Δ_i IN THEOREM 4.5

In this section, and in the entire paper, G is an irreducible subgroup of the general linear group $\text{GL}(n, q)$ over a vector space of dimension n over a finite field \mathbb{F}_q .

We define the order ideal $\mathcal{I}(G, H)$ and the simplicial complexes Δ_i we consider in Theorem 4.5. We will explicitly observe that the two complexes Δ_i are not the order simplicial complex rising from $\mathcal{I}(G, H)$ (see Remark 2).

Given a subgroup G of $\text{GL}(n, q)$ and a subgroup H of G , put

$$\mathcal{L}(G)_{\geq H} = \{K \leq G \mid H \leq K\}$$

and

$$\mathcal{C}(G, H) = \{\text{stab}_G(W) \mid 0 < W < V, H \subseteq \text{stab}_G(W)\}.$$

Definition 3.1. The **reducible subgroup ideal** in $\mathcal{L}(G)_{\geq H}$ is the order ideal generated by $\mathcal{C}(G, H)$. Namely,

$$\mathcal{I}(G, H) = \{K \leq G \mid H \leq K \leq M \text{ for some } M \in \mathcal{C}(G, H)\}.$$

Remark 1. If H is reducible, that is H fixes some non-trivial subspace W of V , then $H \in \mathcal{I}(G, H)$. Otherwise, if H is irreducible, $H \notin \mathcal{I}(G, H)$ and $\mathcal{I}(G, H)$ is empty. The subspaces fixed by H are said to be H -invariant.

Definition 3.2. If H is reducible, we set

$$\widehat{\mathcal{I}}(G, H) = \mathcal{I}(G, H) \cup \{G\}$$

by adjoining the maximum G to $\mathcal{I}(G, H)$, which has minimum H . Otherwise, if H is irreducible, we set $\widehat{\mathcal{I}}(G, H) = \{H, G\}$ by adjoining the minimum H and the maximum G to the empty poset \emptyset .

Remark 2. Here we just note that the poset $\mathcal{I}(G, H)$ has already a minimum if H is reducible, so that $\mu_{\widehat{\mathcal{I}}(G, H)}(H, G)$ is not, in general, the Euler characteristic of $\Delta(\mathcal{I}(G, H))$.

To define the simplicial complexes Δ_i , we begin with fixing some more notation. We denote by $\mathcal{S}(V, H)$ the lattice of H -invariant subspaces of V and define

$$\mathcal{S}(V, H)^* = \mathcal{S}(V, H) \setminus \{0, V\}.$$

Moreover, given an irreducible group $G \leq \text{GL}(V)$, and $H \leq G$, we will consider the following three sets:

- (a) $\Psi(G, H) = \{X \subseteq \mathcal{C}(G, H) \mid \bigcap_{M \in X} M \neq H\}$;
- (b) $\Psi(G, H)^{\mathbb{G}} = \{Y \subseteq \mathcal{C}(G, H) \mid \bigcap_{M \in Y} M = H\}$;
- (c) $\Psi'(G, H) = \{E \subseteq \mathcal{S}(V, H)^* \mid \bigcap_{W \in E} \text{stab}_G(W) \neq H\}$.

Observe that $\emptyset \in \Psi(G, H)$ and $\emptyset \in \Psi'(G, H)$, but $\emptyset \notin \Psi(G, H)^{\mathbb{G}}$.

Remark 3. If H is an irreducible subgroup of G , $\mathcal{S}(V, H)^* = \emptyset$ and $\Psi'(G, H) = \{\emptyset\} = \Psi(G, H)$. Since for irreducible H we have $\widehat{\mathcal{I}}(G, H) = \{H, G\}$, then Theorem 4.5 is trivially verified in this case.

Definition 3.3. The simplicial complexes Δ_i of Theorem 4.5 are so defined:

- (1) Δ_1 :
 - The set of vertices T_1 is given by $\mathcal{S}(V, H)^*$
 - the set of faces of Δ_1 is given by $\Psi'(G, H)$.
- (2) Δ_2 :
 - The set of vertices T_2 is given by $\mathcal{C}(G, H)$
 - the set of faces of Δ_2 is given by $\Psi(G, H)$.

4. COMPUTING $\mu_{\widehat{\mathcal{I}}(G, H)}(H, G)$

In order to prove Theorem 4.5, we need the following Proposition 4.2 that gives a link between $\Psi(G, H)$ and $\Psi'(G, H)$, and prove that the Euler characteristics of complexes Δ_i coincide. The proof of Proposition 4.2 follows at once from Lemma 4.1.

Lemma 4.1. *Let T be a subgroup of a finite group L acting on a finite set X and let $X' \subseteq X$ be a subset such that $T \leq L_x$ for all $x \in X'$ (as usual, L_x denotes the stabilizer of x in L); set*

- $\mathcal{L} = \{L_x \mid x \in X'\}$;
- $\mathcal{R} = \{E \subseteq \mathcal{L} \mid \bigcap_{K \in E} K \neq T\}$;
- $\mathcal{S} = \{Q \subseteq X' \mid \bigcap_{x \in Q} L_x \neq T\}$.

Then

$$\sum_{E \in \mathcal{R}} (-1)^{|E|} = \sum_{Q \in \mathcal{S}} (-1)^{|Q|}.$$

Proof. If $Q \in \mathcal{S}$ and $E \in \mathcal{R}$, set

$$\mathcal{L}_Q = \{L_x \mid x \in Q\} \text{ and } \mathcal{S}_E = \{Q \in \mathcal{S} \mid E = \mathcal{L}_Q\}.$$

It is immediate to realize that

$$\mathcal{S} = \bigsqcup_{E \in \mathcal{R}} \mathcal{S}_E$$

is the disjoint union of all the \mathcal{S}_E . So, we show that for each $E \in \mathcal{R}$ it is verified:

$$(-1)^{|E|} = \sum_{Q \in \mathcal{S}_E} (-1)^{|Q|}.$$

Fix $E \in \mathcal{R}$ and observe that

$$Q \in \mathcal{S}_E \text{ iff } Q = \bigsqcup_{K \in E} Q_K,$$

where $Q_K = \{x \in Q \mid L_x = K\}$. For each $K \in E$, define

$$X'_K = \{x \in X' \mid L_x = K\} \subseteq X',$$

and notice that $Q_K \subseteq X'_K$. Since $\sum_{\emptyset \neq Q_K \subseteq X'_K} (-1)^{|Q_K|} = -1$ (see Remark 3 below), we get

$$\sum_{Q \in \mathcal{S}_E} (-1)^{|Q|} = \sum_{Q \in \mathcal{S}_E} (-1)^{\sum_{K \in E} |Q_K|} = \prod_{K \in E} \left(\sum_{\emptyset \neq Q_K \subseteq X'_K} (-1)^{|Q_K|} \right) = \prod_{K \in E} (-1) = (-1)^{|E|}.$$

□

Proposition 4.2. *Let V be a vector space of finite dimension over \mathbb{F}_q and consider a subgroup $H \leq G \leq \text{GL}(V)$. We have:*

$$(7) \quad \sum_{E \in \Psi'(G, H)} (-1)^{|E|} = \sum_{\mathcal{X} \in \Psi(G, H)} (-1)^{|\mathcal{X}|}.$$

Proof. Consider the natural action of G on V . By Lemma 4.1 below, the equality follows at once from the definitions of $\mathcal{S}(V, H)^*$, $\mathcal{C}(G, H)$, $\Psi(G, H)$ and $\Psi'(G, H)$.

With previous notation, we take $T = H$, $L = G$, $\mathcal{R} = \Psi(G, H)$, $\mathcal{S} = \Psi'(G, H)$.

□

To prove Theorem 4.5 we need Proposition 4.4 which is achieved through the Crosscut Theorem [9, Corollary 3.9.4] and the Remark below.

Remark 4. For every finite set A of cardinality $n > 0$ we have

$$\sum_{S \subseteq A} (-1)^{|S|} = \sum_{k=0}^n \binom{n}{k} (-1)^k = (1-1)^n = 0.$$

Theorem 4.3 (Crosscut Theorem). *Let L be a finite lattice with minimum $\hat{0}$ and maximum $\hat{1}$, so that $\hat{0} \neq \hat{1}$. Let M be the set of all coatoms in L . Let $X \subseteq L$ be a subset such that $M \subseteq X$ and $\hat{1} \notin X$. Given $\mathcal{Y} = \{Y \subseteq X, Y \neq \emptyset \text{ and } \bigcap_{Y \in \mathcal{Y}} Y = \hat{0}\}$, the following equality holds:*

$$\mu_L(\hat{0}, \hat{1}) = \sum_{Y \in \mathcal{Y}} (-1)^{|Y|}.$$

Proposition 4.4. *Let V be a vector space of finite dimension over \mathbb{F}_q . Let $H \leq G \leq \text{GL}(V)$. Then we have that*

$$(8) \quad \mu_{\widehat{\mathcal{I}}(G,H)}(H, G) = \sum_{Y \in \Psi(G,H)^{\complement}} (-1)^{|Y|}.$$

Proof. We observe that $\widehat{\mathcal{I}}(G, H) \subseteq \mathcal{L}(G)_{\geq H}$, is a lattice because the join of two subgroups $K_1, K_2 \in \mathcal{I}(G, H)$ is either in $\mathcal{I}(G, H)$ or equal to G . The meet of $K_1, K_2 \in \mathcal{I}(G, H)$ is $K_1 \cap K_2$. Hence, $\widehat{\mathcal{I}}(G, H)$ is a finite lattice, whose coatoms are the subgroups in $\mathcal{C}(G, H)$. Since

$$\Psi(G, H)^{\complement} = \{Y \subseteq \mathcal{C}(G, H) \mid Y \neq \emptyset \text{ and } \bigcap_{M \in Y} M = H\},$$

by Theorem 4.3 we immediately obtain (8). \square

Now, observe that $\Psi(G, H) \cup \Psi(G, H)^{\complement}$ is the power set of $\mathcal{C}(G)$, so that by Remark 3

$$(9) \quad \sum_{X \in \Psi(G,H)} (-1)^{|X|} + \sum_{Y \in \Psi(G,H)^{\complement}} (-1)^{|Y|} = 0.$$

If we put together equations (7), (8), and (9), we get:

Theorem 4.5. *Let V be a vector space of finite dimension over \mathbb{F}_q . Let $H \leq G \leq \text{GL}(V)$. Then*

$$(10) \quad -\mu_{\widehat{\mathcal{I}}(G,H)}(H, G) = \sum_{E \in \Psi'(G,H)} (-1)^{|E|}.$$

Proof. By (9), we have

$$\sum_{X \in \Psi(G,H)} (-1)^{|X|} = - \sum_{Y \in \Psi(G,H)^{\complement}} (-1)^{|Y|}.$$

Then, by Proposition 4.2 and Proposition 4.4,

$$\begin{aligned} \sum_{E \in \Psi'(G, H)} (-1)^{|E|} &= \sum_{\mathcal{X} \in \Psi(G, H)} (-1)^{|\mathcal{X}|} = - \sum_{\mathcal{Y} \in \Psi(G, H)^c} (-1)^{|\mathcal{Y}|} \\ &= -\mu_{\widehat{\mathcal{I}}(G, H)}(H, G). \end{aligned}$$

□

5. FINAL REMARK

Going back to Conjecture 1.1, we observe that the knowledge of

$$\sum_{E \in \Psi'(G, H)} (-1)^{|E|}$$

coming from (10) can be exploited to estimate the value $\mu(H, G)$ of the Möbius function μ of G for $H \leq G$, at least in some particular case. Here we just give an example for a particular reducible H , taking $G = \text{GL}(n, q)$.

Following the idea suggested by Shareshian in [8], one could write

$$(11) \quad \mu(H, G) = f_{G, n, q}(H) + \sum_{K \in \mathcal{C}_9} \mu(H, K),$$

where $f_{G, n, q}(H)$ depends on the classes $\mathcal{C}_i(G, H)$, for $i = 1, \dots, 8$, in Aschbacher's classification. In some lucky case, H is not contained in maximal subgroups belonging to classes (\mathcal{C}_i) , $i \neq 1, 9$. This happens, for example, in the following case:

let $V \simeq \mathbb{F}_q^n$ be a vector space of finite dimension over \mathbb{F}_q and fix the following basis of V :

$$\mathcal{E} = \{w_1, \dots, w_m, v_{m+1} \dots v_n\},$$

so that

$$V = \langle w_1 \dots, w_m \rangle \oplus \langle v_{m+1}, \dots, v_n \rangle$$

If $W = \langle w_1 \dots, w_m \rangle$, H is the subgroup of $\text{GL}(n, q)$ acting as $\text{GL}(m, q)$ on W and fixing all the elements v_{m+1}, \dots, v_n .

We do not give in this context the details of the proof of the following Theorem, which needs many technical arguments. In [2] all the details are given.

Theorem 5.1 ([2]). *Let $G = \text{GL}(n, q)$, and let $H \leq G$ be such that*

$$H = \text{GL}(m, q) \oplus I_{n-m}.$$

Let $q = p$ be an odd prime and let the dimension n be prime. If $n - m + 1$ is prime, then

$$\mu_{\widehat{\mathcal{I}}(G)}(H, G) = 0 = f_{G, n, p}(H)$$

so that

$$\mu(H, G) = \sum_{K \in \mathcal{C}_9(G, H), H \subseteq K} \mu_K(H).$$

In general, we do not have much information about the ninth class. Just to give an example, we considered the groups of low dimension studied by Schröder in her PhD thesis ([7]), and we saw that in dimension $n = 13$ also class $\mathcal{C}_9(G, H)$ is empty for $p > 5$. In this case, $\mu(H, G) = 0$.

REFERENCES

- [1] L. Di Gravina, A closure operator on the subgroup lattice of $GL(n, q)$ and $PGL(n, q)$ in relation to the zeros of the Möbius function *private communication*, 2022.
- [2] L. Di Gravina, *Some questions about the Möbius function of finite linear groups*, Ph.D. Thesis, Università degli Studi di Milano-Bicocca, 2022.
- [3] P. Hall, The Eulerian functions of a group, *Quarterly Journal of Mathematics* **7** (1936), 134–151.
- [4] Peter Kleidman and Martin Liebeck, *The Subgroup Structure of the Finite Classical Groups*, Cambridge University Press, Cambridge, 1990.
- [5] Andrea Lucchini, On the subgroups with non-trivial Möbius number, *Journal of Group Theory*, **13** (2010), 589-600.
- [6] Avinoam Mann, A probabilistic zeta function for arithmetic groups, *International Journal of Algebra and Computation*, **15** (2005), 1053-1059.
- [7] Anna Katharina Schröder, *The maximal subgroups of the classical groups in dimension 13, 14 and 15*, Ph.D. Thesis, University of St Andrews, 2015.
- [8] John Shareshian, *Combinatorial properties of subgroup lattices of finite groups*, Ph.D. Thesis, Rutgers - The State University of New Jersey, 1996.
- [9] Richard P. Stanley, *Enumerative Combinatorics, Volume I*, second ed., Cambridge University Press, Cambridge, 2012 (first ed. published by Wadsworth & Brooks/Cole, 1986).

FRANCESCA DALLA VOLTA: DIPARTIMENTO DI MATEMATICA E APPLICAZIONI, UNIVERSITÀ DEGLI STUDI DI MILANO-BICOCCA, VIA ROBERTO COZZI 55, 20125 MILANO, ITALY

Email address: `francesca.dallavolta@unimib.it`

LUCA DI GRAVINA: MATHEMATISCHES INSTITUT, HEINRICH-HEINE-UNIVERSITÄT DÜSSELDORF, 40225 DÜSSELDORF, GERMANY

Email address: `luca.di.gravina@hhu.de`