

A computation of the covariance between two linear statistics for the Jellium model

Pete Rigas

Abstract

We extend previous results providing an exact formula for the variance of a linear statistic for the Jellium model, a one-dimensional model of Statistical mechanics obtained from the $k \rightarrow 0^+$ limit of the Dyson log-gas. For such a computation of the covariance, in comparison to previous work for computations of the log-gas covariance, we obtain a formula between two *linear statistics*, given arbitrary functions f and g over the real line, that is dependent upon an asymptotic approximation of the Jellium probability distribution function from large N deviations.¹

1 Introduction

1.1 Overview

Log-gases have attracted great attention from both the Mathematics and Physics communities, with several studies pertaining to, exact extremal statistics, and index distribution, for the Coulomb gas in one-dimension [3,4], connections with random-matrix theory [7], large deviation principles [10], characterizations relating to the one-dimensional plasma [6,11], classification of Gibbs measures [1], a central Limit theorem result [12], in addition to several conjectures and open questions [9]. To extend the analysis provided for an exact formula of the variance of a linear statistic in the Jellium model provided in [5], which can further add to the collection of work in the literature for approximating the behavior of *linear statistics* for a wide variety of models, including the truncation of linear statistics for top eigenvalues of random matrices [8], as well as another result for a *covariance measure* [2], which under certain assumptions can be used to obtain a *generalized covariance measure*, we extend methods of a recent work contained in [5] to obtain a covariance for two *linear statistics*, in the large, strictly positive N limit.

1.2 Paper organization

In the following, we present objects from the Jellium model for defining the *energy function*, and connection with quantities of the *Riesz gas*. Before introducing the *covariance formula* which is dependent upon two *linear statistics* as parameters, we provide an overview of the rate function Ψ , which is a function of the linear statistic. Following the overview of the methods contained in [5] for obtaining the behavior of the Variance statistic for large N , we provide a computation of the covariance for two linear statistics in the Jellium model, which differs from a *covariance measure* previously obtained for the log-gas in [2], which is instead dependent upon an effective saddle-point density action.

1.3 Jellium objects

We introduce the following quantities.

Definition 1 (*Riesz gas energy*). Introduce, for some x_i on \mathbf{R} ,

$$\mathcal{E}^k(x_1, \dots, x_i, \dots, x_N) \equiv \mathcal{E} \equiv \frac{A}{2} \sum_{i=1}^n x_i^2 + \alpha \operatorname{sgn}(k) \sum_{i \neq j} |x_i - x_j|^{-k} \quad ,$$

corresponding to the *energy* of the *Riesz gas*, for strictly positive parameters A and α , with,

$$\begin{aligned} A &\sim O(1) \quad , \\ \alpha &\sim O(1) \quad , \end{aligned}$$

¹**Keywords:** Log-gas, random matrices, covariance, variance, linear statistics, Riesz gas, Jellium model

and sign function,

$$\text{sgn}(k) = \begin{cases} 1, & x > 0, \\ 0, & x = 0, \\ -1, & x < 0, \end{cases}$$

The energy function above, with the positions of each particle at any position i , under the conformal transformation,

$$T(x) = \frac{ax + b}{cx + d},$$

for $ad - cb \neq 0$, which sends the position of each particle at x_i to λ_i , from,

$$T(x_i) = \frac{ax_i + b}{cx_i + d} \mapsto \lambda_i,$$

yields a distribution on eigenvalues of random matrices, given by,

$$\mathcal{E}^k(T(x_1), \dots, T(x_i), \dots, T(x_N)) \equiv \mathcal{E}^k(\lambda_1, \dots, \lambda_i, \dots, \lambda_N) \equiv (\mathcal{E}')^k \equiv \frac{A}{2} \sum_{i=1}^n \lambda_i^2 + \alpha \text{sgn}(k) \sum_{i \neq j} |\lambda_i - \lambda_j|^{-k},$$

under the same choice of couplings A and α .

Definition 2 (*Riesz gas probability measure on random-matrix eigenvalues, a Gibbs-Boltzmann distribution*). From the energy of **Definition 1**, introduce,

$$\mathbf{P}(\lambda_1, \dots, \lambda_i, \dots, \lambda_N) \equiv \frac{1}{Z_N^k} \exp\left(-\frac{(\mathcal{E}')^k}{k_b T}\right),$$

given the partition function,

$$Z_N^k(\lambda_1, \dots, \lambda_i, \dots, \lambda_N) \equiv Z_N^k \equiv \sum_{i: i \leq N} \left(\frac{A}{2} \sum_{i=1}^n \lambda_i^2 + \alpha \text{sgn}(k) \sum_{i \neq j} |\lambda_i - \lambda_j|^{-k} \right),$$

normalizing $\exp[-\mathcal{E}(\lambda_i)]$ so that $\mathbf{P}(\lambda_1, \dots, \lambda_i, \dots, \lambda_N)$ is a probability measure, for some $N > 0$, Boltzmann constant k_b , and temperature $T > 0$, with,

$$T \sim \mathcal{O}(1).$$

Definition 3 (*Rescaling the energy function of the Gibbs-Boltzmann weight to obtain the probability measure of the Jellium model for $k \equiv -1$*). Introduce, $Z_N^{-1}(\lambda_1, \dots, \lambda_i, \dots, \lambda_N) \equiv Z_N^{-1}$, with,

$$Z_N^{-1} \equiv \sum_{i: i \leq N} \left(\frac{A}{2} \sum_{i=1}^n \lambda_i^2 + \alpha \text{sgn}(-1) \sum_{i \neq j} |\lambda_i - \lambda_j|^{-1} \right) \equiv \sum_{i: i \leq N} \left(\frac{A}{2} \sum_{i=1}^n \lambda_i^2 - \alpha \sum_{i \neq j} |\lambda_i - \lambda_j|^{-1} \right),$$

corresponding to the partition function of the *Riesz gas probability measure* of **Definition 2** for $k \equiv -1$, which under the rescaling yields the following summation of energy functions, taking the form,

$$Z_N^{-1}(L_N \lambda_1, \dots, L_N \lambda_i, \dots, L_N \lambda_N) \equiv \sum_{i: i \leq N} \left(\frac{A}{2} \sum_{i=1}^n (L_N \lambda)_i^2 - \alpha L_N \sum_{i \neq j} |\lambda_i - \lambda_j|^{-1} \right),$$

hence providing the *probability measure* of the Jellium model,

$$\mathbf{P}(L_N \lambda_1, \dots, L_N \lambda_i, \dots, L_N \lambda_N) \equiv \tilde{\mathbf{P}}(\lambda_1, \dots, \lambda_i, \dots, \lambda_N) \equiv \frac{1}{Z_N^{-1}} \exp\left(-\frac{(\mathcal{E}')^{-1}}{k_b T}\right) ,$$

for the rescaled Jellium *energy function*,

$$\mathcal{E}^{-1}(L_N \lambda_1, \dots, L_N \lambda_i, \dots, L_N \lambda_N) \equiv (\mathcal{E}')^{-1} \equiv \frac{A}{2} \sum_{i=1}^n (L_N \lambda)_i^2 - \alpha L_N \sum_{i \neq j} |\lambda_i - \lambda_j|^{-1} ,$$

given a suitable constant $L_N > 0$ dependent upon the choice of N .

Definition 4 (*linear statistics*). A *linear statistic* takes the form,

$$s \equiv \frac{1}{N} \sum_{i=1}^N f(x_i) ,$$

for an arbitrary function $f(x)$ of the position x on the real line, under the same choice of parameter N provided in **Definition 2**.

Definition 5 (*covariance formula from the Gibbs-Boltzmann weight*, [2]). Introduce, for $\Lambda^N \subseteq \mathbf{R}$,

$$\text{Cov}(A, B) \equiv \int_{\Lambda^N} d\lambda_1 \cdots d\lambda_N \frac{1}{Z_N^{-1}} \exp\left(-\frac{(\mathcal{E}')^{-1}}{k_b T}\right) A(T(x_1, \dots, x_N)) B(T(x_1, \dots, x_N)) - \langle A \rangle \langle B \rangle ,$$

corresponding to the *covariance formula* between the *linear statistics*,

$$A = \frac{1}{N} \sum_{i=1}^N f(x_i) ,$$

and,

$$B = \frac{1}{N} \sum_{i=1}^N g(x_i) ,$$

for two arbitrary functions $f(x)$ and $g(x)$, and $\Lambda \subseteq \mathbf{R}^N$, for the expectation,

$$\langle \cdot \rangle_{A, \mathbf{R}} \equiv \langle \cdot \rangle_A ,$$

against a test *linear statistic* A .

1.4 Overview of the Coulomb gas method

In this section, we recount how the Gibbs-Boltzmann weight, adapted from the log-gas to the Jellium model with **Definition 3**, below.

First, introduce a probability measure on the *linear statistic*, with,

$$\mathcal{P}[s, N] \equiv \int_{\Lambda^N} dx_1 \cdots dx_i \cdots dx_N \tilde{\mathbf{P}}(x_1, \dots, x_i, \dots, x_N) \delta_{s - \frac{1}{N} \sum_{i=1}^N f(x_i)}$$

which under conformal transformations in **Definition 1**, is,

$$\begin{aligned} \mathcal{P}[Ts, N] &\equiv \int_{\Lambda^N} d(T(x_1)) \cdots dT((x_i)) \cdots dT((x_N)) \tilde{\mathbf{P}}(T(x_1), \dots, T(x_i), \dots, T(x_N)) \delta_{s - \frac{1}{N} \sum_{i=1}^N f(T(x_i))} \\ &\equiv \int_{\Lambda^N} d\lambda_1 \cdots d\lambda_N \tilde{\mathbf{P}}(\lambda_1, \dots, \lambda_i, \dots, \lambda_N) \delta_{s - \frac{1}{N} \sum_{i=1}^N f(\lambda_i)} \quad , \end{aligned}$$

which, upon rescaling the two interactions in the *energy* function of the *Jellium model*, has a mass of order, by a large deviation,

$$\mathcal{P}[s, N] \approx \exp(-N^3 \Psi(s)) \quad ,$$

where in the power of the exponent above, besides the N^3 scaling, is given by the rate function,

$$\Psi(s) \approx \frac{1}{2b} (s - \bar{s})^2 \quad ,$$

about the minimum $s = \bar{s}$ of the rate function, with variance,

$$\text{Var}[s] \approx \frac{b}{N^3} \quad ,$$

for some positive parameter b . We proceed by summarizing components for each quantity in $\mathcal{P}[s, N]$, in which approximations are introduced for,

$$\delta_{s - \frac{1}{N} \sum_{i=1}^N f(x_i)} \approx N^3 \int_{\Gamma} \frac{1}{2\pi} \exp(-\nu N^3) d\mu \quad ,$$

and hence also for,

$$\int_{\Lambda^N} d\lambda_1 \cdots d\lambda_N \tilde{\mathbf{P}}(\lambda_1, \dots, \lambda_i, \dots, \lambda_N) \approx \frac{1}{Z_N^{-1}} \int_{\Gamma} d\mu \int_{\Lambda^N} d\lambda_1 \cdots d\lambda_N \exp\left(-\frac{(\mathcal{E}')^{-1}}{k_b T}\right) \quad , \quad (*)$$

given $\Gamma \subset \mathbf{C}$ intersecting the imaginary axis of the μ plane. Given an arbitrary $f(x)$, the effective potential,

$$V^{\text{eff}}(x) \equiv \frac{1}{2} x^2 + \mu f(x) \quad .$$

For $\mu \equiv 0$, adopt the convention,

$$Z_N^{-1}(\mu = 0) = Z_N^{-1} \quad .$$

As a function of nonzero μ , for large N the hydrodynamic approximation the partition function appearing in (*) has been shown to be of the form,

$$Z_N^{-1} \approx \int_{\Lambda^N} d\lambda_1 \cdots d\lambda_N d(\rho(x)) \exp\left[-N^3 \mathbf{E}_{\mu}[\rho(x)] - N \int_{\text{supp}} dx \rho(x) \ln \rho(x)\right] \delta_{\int dx \rho(x) - 1} \quad , \quad (**)$$

where the expectation of the density profile $\rho(x)$ appearing in the power of the exponential is,

$$\mathbf{E}_{\mu}[\rho(x)] = \int_{\text{supp}} \left(\frac{x^2}{2} + \mu f(x)\right) \rho(x) dx - \alpha \int_{\text{supp}} \int_{\text{supp}} \rho(x) \rho(y) |x - y| dx dy - \mu s \quad ,$$

for the support of the profile over the real line,

$$\text{supp} \cap \mathbf{R} \neq \emptyset ,$$

which admits the decomposition,

$$\text{supp} \equiv \bigcap_{1 \leq i \leq N} \text{supp}_i ,$$

hence implying, under conformal transformations T , as those mentioned in **Definition 1**,

$$\int_{\text{supp}} d(\rho(T(x))) \equiv \prod_{1 \leq i \leq N} \int_{\text{supp}_i} d(\rho(T(x_i))) , \quad (\text{Supp})$$

which is taken with respect to,

$$\mathbf{E}_\mu[\cdot] .$$

Altogether, under the assumption that, the effective action given on the LHS below,

$$-\frac{(\mathcal{E}')^{-1}}{k_b T} \approx N^3 S[\rho(x), \mu, \mu_0] ,$$

for action given by,

$$S[\rho(x), \mu, \mu_0] = \int_{\text{supp}} \frac{x^2}{2} \rho(x) dx - \alpha \int_{\text{supp}} \int_{\text{supp}} \rho(x) \rho(y) |x - y| dx dy + \mu \left(\int_{\text{supp}} f(x) \rho(x) dx - s \right) + \mu_0 \left(\int_{\text{supp}} \rho(x) dx - 1 \right) ,$$

the expression for $\mathcal{P}[s, N]$ in (*) can be approximated with,

$$\begin{aligned} \mathcal{P}[s, N] &\approx \frac{1}{Z_N^{-1}} \int_{\text{supp}} d(\rho(x)) \int_{\Gamma} d\mu \int_{\Gamma} d\mu_0 \exp \left[- N^3 S[\rho(x), \mu, \mu_0] \right] \\ &\approx \prod_{1 \leq i \leq N} \int_{\text{supp}_i} d(\rho(x_i)) \int_{\Gamma} d\mu \int_{\Gamma} d\mu_0 \exp \left[- N^3 S[\rho(x_i), \mu, \mu_0] \right] . \end{aligned} \quad (***)$$

With the saddle-point method, the order of the unconstrained Jellium partition function is approximately,

$$\frac{1}{Z_N^{-1}} \approx \frac{1}{\exp(\frac{2}{3} \alpha^2 N^3)} ,$$

for the rate function satisfying,

$$\Psi(s) \approx \frac{2}{3} \alpha^2 .$$

Up to higher orders, the rate function above can also take the form,

$$\Psi(s) = S[\rho_\mu^*(x), \mu, \mu_0] + \frac{2}{3} \alpha^2 ,$$

for the effective saddle-point action,

$$S[\rho_\mu^*(x), \mu, \mu_0] = \int_{\text{supp}} \frac{x^2}{2} \rho_\mu^*(x) dx - \alpha \int_{\text{supp}} \int_{\text{supp}} \rho_\mu^*(x) \rho_\mu^*(y) |x - y| dx dy + \mu \left(\int_{\text{supp}} f(x) \rho_\mu^*(x) dx - s \right) + \dots$$

$$\mu_0 \left(\int_{\text{supp}} \rho_\mu^*(x) dx - 1 \right) , \quad (****)$$

upon substituting (****) into the power of the exponential given in (***), namely replacing the saddle point action $S[\rho_\mu^*(x), \mu, \mu_0]$ with $S[\rho_\mu(x), \mu, \mu_0]$ yields the distribution,

$$\mathcal{P}[s, N] \approx \frac{1}{Z_N^{-1}} \prod_{1 \leq i \leq N} \int_{\text{supp}_i} d(\rho(x_i)) \int_{\Gamma} d\mu \int_{\Gamma} d\mu_0 \exp \left[-N^3 S[\rho_\mu^*(x_i), \mu, \mu_0] \right] , \quad (*****)$$

which corresponds to the probability density function, given the support of each density $d(\rho(x_i))$, each of which satisfies,

$$\text{supp}_i \cap \mathbf{R} \neq \emptyset ,$$

for each $1 \leq i \leq N$. From such a closed expression for the rate function, the variance of a *linear statistic* for large N is extracted upon making the observation that,

$$\bar{s} = \frac{1}{4\alpha} \int_{[-2\alpha, +2\alpha]} f(x) dx ,$$

is a minimum of $\Psi(s)$. Performing an expansion about the rate function about $s \equiv \bar{s}$ can be obtained by computing μ up to order $O(\epsilon)$, for ϵ sufficiently small. The specific form of the rate function and effective saddle-point action are incorporated into the formula for the Jellium covariance for two *linear statistics* obtained in the second section.

2 Formalizing the computation for the Jellium covariance

Below, we state the main result for the computation of the covariance.

Theorem (*computation of the Jellium covariance between two linear statistics*). For two *linear statistics*,

$$A \equiv A(f, x_1, \dots, x_i, \dots, x_N) = \frac{1}{N} \sum_{i=1}^N f(x_i) ,$$

and,

$$B \equiv B(g, x_1, \dots, x_i, \dots, x_N) = \frac{1}{N} \sum_{i=1}^N g(x_i) ,$$

the Jellium covariance between A and B takes the form,

$$\text{Cov}^{\text{Jellium}}(A, B) \equiv \frac{1}{N^2} \left(\int_{\text{supp}} d(\rho(x)) \int_{\Gamma} d\mu \int_{\Gamma} d\mu_0 \frac{1}{Z_N^{-1}} \exp \left[-N^3 S[\rho_\mu^*(x), \mu, \mu_0] \right] \times \dots \right.$$

$$\left. \left(\sum_{i=1}^N f(x_i) g(x_i) \right) - \left\langle \sum_{i=1}^N f(x_i) \right\rangle_f \left\langle \sum_{i=1}^N g(x_i) \right\rangle_g \right) ,$$

for some $N > 0$, given f and g arbitrary.

Proof of Theorem. Recall, from the previous section, that the approximation of the *Riesz gas* energy for $k \equiv -1$, and hence of the Jellium model,

$$-\frac{(\mathcal{E}')^{-1}}{k_b T} \approx N^3 S[\rho_\mu^*(x), \mu, \mu_0] \quad ,$$

with the effective saddle-point density action $\rho_\mu^*(x)$, which appears in the contributions for the rate function,

$$\Psi(s) = S[\rho_\mu^*(x), \mu, \mu_0] + \frac{2}{3}\alpha^2 \quad ,$$

as well as the constant $\frac{2}{3}\alpha^2$. From previously implemented techniques for extracting the behavior of the variance of a single *linear statistic* in the large N limit, applying the conformal transformation,

$$T(x) = \frac{ax + b}{cx + d} \quad ,$$

provided in **Definition 1**, to each position x_i , of the *energy* function of the *Riesz gas* introduced in **Definition 1**, for $k \equiv -1$, given in **Definition 2**, yields,

$$\mathcal{E}^k(T(x_1, \dots, x_i, \dots, x_N)) \equiv \mathcal{E}^k(T(x_1), \dots, T(x_i), \dots, T(x_N)) \equiv \mathcal{E}^k(\lambda_1, \dots, \lambda_i, \dots, \lambda_N) \quad ,$$

corresponding to the *Riesz gas energy function*, under conformal T , for $k \equiv -1$. From the probability measure introduced in **Definition 3**, the quantity,

$$\begin{aligned} \left\langle x_1, \dots, x_i, \dots, x_N \right\rangle_{\text{Jellium}} &\equiv \int_{\Lambda^N} dx_1 \cdots dx_i \cdots dx_N \frac{1}{Z_N^{-1}} \exp\left(-\frac{(\mathcal{E}')^{-1}}{k_b T}\right) \\ &\approx \int_{\Lambda^N} dx_1 \cdots dx_i \cdots dx_N \int_{\Gamma} d\mu \int_{\Gamma} d\mu_0 \frac{1}{Z_N^{-1}} \exp\left[-N^3 S[\rho_\mu^*(x), \mu, \mu_0]\right] \\ &\approx \prod_{1 \leq i \leq N} \int_{\text{supp}_i} d(\rho(x_i)) \int_{\Gamma} d\mu \int_{\Gamma} d\mu_0 \frac{1}{Z_N^{-1}} \exp\left[-N^3 S[\rho_\mu^*(x_i), \mu, \mu_0]\right] \quad , \end{aligned}$$

for the corresponds to the Jellium expectation taken across all x_i up to N , for the *energy function*,

$$(\mathcal{E}')^{-1}(x_1, \dots, x_i, \dots, x_N) \equiv (\mathcal{E}')^{-1} \quad ,$$

which satisfies, given support over \mathbf{R} for the base probability measure $\tilde{\mathbf{P}}[\cdot]$,

$$\langle \cdot \rangle_{\text{Jellium}} \equiv \langle \cdot \rangle_{\tilde{\mathbf{P}}, \mathbf{R}} \equiv \langle \cdot \rangle \quad .$$

Proceeding, under conformal T as introduced above applied to each x_i as mentioned previously, from the ensemble average of each $T(x_i)$, the covariance between the two *test statistics* can readily be formed, which coincides with,

$$\left\langle x_1, \dots, x_i, \dots, x_N \right\rangle_{\text{Jellium}} \left(\frac{1}{N^2} \sum_{i=1}^N f(x_i) g(x_i) \right) - \frac{1}{N^2} \left\langle \sum_{i=1}^N f(x_i) \right\rangle_f \left\langle \sum_{i=1}^N g(x_i) \right\rangle_g \quad ,$$

which, under the previous identification for $\langle \cdot \rangle_{\text{Jellium}}$, is equivalent to,

$$\begin{aligned} \frac{1}{N^2} \left(\prod_{1 \leq i \leq N} \int_{\text{supp}_i} d(\rho(x_i)) \int_{\Gamma} d\mu \int_{\Gamma} d\mu_0 \frac{1}{Z_N^{-1}} \exp\left[-N^3 S[\rho_\mu^*(x_i), \mu, \mu_0]\right] \left(\frac{1}{N^2} \sum_{i=1}^N f(x_i) g(x_i) \right) - \dots \right. \\ \left. \left\langle \sum_{i=1}^N f(x_i) \right\rangle_f \left\langle \sum_{i=1}^N g(x_i) \right\rangle_g \right) \quad . \end{aligned}$$

Under suitable, conformal $T(\cdot)$ sending each position x_i as input to a corresponding eigenvalue λ_i , the expression obtained above can be written as,

$$\left\langle T(x_1, \dots, x_i, \dots, x_N) \right\rangle \equiv \left\langle T(x_1), \dots, T(x_i), \dots, T(x_N) \right\rangle ,$$

which is in turn equivalent to,

$$\frac{1}{N^2} \left(\prod_{1 \leq i \leq N} \int_{\text{supp}_i} d(\rho(T(x_i))) \int_{\Gamma} d\mu \int_{\Gamma} d\mu_0 \frac{1}{Z_N^{-1}} \exp \left[-N^3 S[\rho_{\mu}^*(T(x_i)), \mu, \mu_0] \right] \left(\frac{1}{N^2} \sum_{i=1}^N f(T(x_i)) g(T(x_i)) \right) - \dots \right. \\ \left. \left\langle \sum_{i=1}^N f(T(x_i)) \right\rangle_f \left\langle \sum_{i=1}^N g(T(x_i)) \right\rangle_g \right) ,$$

exhibiting that the desired formula for the Jellium covariance holds, under the observation, provided in (Supp), that,

$$\int_{\text{supp}} d(\rho(T(x))) \equiv \prod_{1 \leq i \leq N} \int_{\text{supp}_i} d(\rho(T(x_i))) ,$$

and also that,

$$\lim_{N \rightarrow +\infty} \langle \rho_N(x) \rangle = \rho(x) ,$$

which corresponds to the large N limit of the density profile, which is given by,

$$\langle \rho_N(x) \rangle \equiv \frac{1}{N} \left\langle \sum_{i=1}^N \delta(x - x_i) \right\rangle ,$$

corresponding to a summation over delta functions at each x_i . Hence we obtain a formula for the Jellium covariance that is a function of the two *test statistics*, which are respectively given by arbitrary f and g , from which we conclude the argument. \square

3 References

- [1] Aizenman, M., Martin, P.A. Structure of Gibbs states of one Dimensional Coulomb Systems. *Commun. Math. Physics* **78**, 99-116 (1980).
- [2] Cunden, F.D., Vivo, P. Universal covariance formula for linear statistics on random matrices. *Phys. Rev. Lett.* **113**, 070202 (2014).
- [3] Dhar, A. et al. Exact Extremal Statistics in the Classical 1D Coulomb Gas. *Phys. Rev. Lett.* **199**, 060601 (2017).
- [4] Dhar, A. et al. Extreme statistics and index distribution in the classical 1d Coulomb gas, *J. Phys. A Math. Theor.* **51**, 295001 (2018).
- [5] Flack, A., Majumdar, S.N., Schehr, G. An exact formula for the variance of linear statistics in the one-dimensional jellium model. *arxiv: 2211.11850 v1*.
- [6] Flack, A., Majumdar, S.N., Schehr, G. Gap probability and full counting statistics in the one dimensional one-component plasma.
- [7] Forrester, P.J. Log-Gases and Random Matrices *Princeton University Press, Princeton* (2010).

- [8] Grabsch, A. General truncated linear statistics for the top eigenvalues of random matrices, *J. Phys. A: Math. Theor.* **55**, 124001 (2021).
- [9] Lewin, M. Coulomb and Riesz gases: The known and the unknown, *J. Math. Phys.* **63**, 061101 (2022).
- [10] Leble, T., Serfaty, S. Large deviation principle for empirical fields of Log and Riesz gases, *Invent. Math.* **210**, 645 (2017).
- [11] Prager, S. The One-Dimensional Plasma. *Adv. Chem. Phys.* **4**, 201 (1962).
- [12] Soshnikov, A. Central Limit Theorem for Local Linear Statistics in Classical Compact Groups and Related Combinatorial Identities. *Ann. Probab.* **28** 3: 1353-1370 (2000).