

# Characterizing Riesz Bases via Biorthogonal Riesz-Fischer sequences

Elias Zikkos

Khalifa University, Abu Dhabi, United Arab Emirates  
email address: elias.zikkos@ku.ac.ae and eliaszikkos@yahoo.com

December 6, 2022

## Abstract

In this note we prove that if two Riesz-Fischer sequences in a separable Hilbert space  $H$  are biorthogonal and one of them is complete in  $H$ , then both sequences are Riesz bases for  $H$ . This complements a recent result by D. T. Stoeva where the same conclusion holds if one replaces the phrase “Riesz-Fischer sequences” by “Bessel sequences”.

Keywords: Riesz-Fischer sequences, Bessel sequences, Riesz sequences, Riesz bases, Biorthogonal sequences, Completeness.

AMS 2010 Mathematics Subject Classification. 42C15, 42C99.

## 1 Introduction

Let  $H$  be a separable Hilbert space endowed with an inner product  $\langle \cdot, \cdot \rangle$  and a norm  $\|\cdot\|$ . Let  $\{f_n\}_{n=1}^{\infty}$  be a sequence of vectors in  $H$ . We say that  $\{f_n\}_{n=1}^{\infty}$  is a **Riesz basis** for  $H$  if  $f_n = V(e_n)$  where  $\{e_n\}_{n=1}^{\infty}$  is an orthonormal basis for  $H$  and  $V$  is a bounded bijective operator from  $H$  onto  $H$ .

One of the many equivalences of Riesz bases [6, Theorem 1.1] states that

- A sequence is a Riesz basis for  $H$ , if and only if it is a complete Bessel sequence having a complete biorthogonal Bessel sequence in  $H$ .

Recall that  $\{f_n\}_{n=1}^{\infty}$  is a **Bessel** sequence if there is a positive constant  $B$  so that

$$\sum_{n=1}^{\infty} |\langle f, f_n \rangle|^2 < B \cdot \|f\|^2 \quad \forall f \in H,$$

and  $\{f_n\}_{n=1}^{\infty}$  is **complete** if its closed span in  $H$  is equal to  $H$ . **Biorthogonality** between two sequences  $\{f_n\}_{n=1}^{\infty}$  and  $\{g_n\}_{n=1}^{\infty}$  means that

$$\langle f_n, g_m \rangle = \begin{cases} 1, & m = n, \\ 0, & m \neq n. \end{cases}$$

Recently Stoeva [6] improved the above equivalence by assuming completeness on just one sequence.

**Theorem A.** [6, Theorem 2.5] *Let two sequences in  $H$  be biorthogonal. If both of them are Bessel sequences and one of them is complete in  $H$ , then they are Riesz bases for  $H$ .*

**Our goal** in this note is to complement Theorem A by replacing the phrase “Bessel sequences” by “Riesz-Fischer sequences”. Following Young [7, Chapter 4, Section 2],  $\{f_n\}_{n=1}^{\infty}$  is a **Riesz – Fischer** sequence in  $H$  if the moment problem

$$\langle f, f_n \rangle = c_n$$

has at least one solution  $f \in H$  for every sequence  $\{c_n\}_{n=1}^{\infty}$  in the space  $l^2(\mathbb{N})$ . We prove the following.

**Theorem 1.1.** *Let two sequences  $\{f_n\}_{n=1}^{\infty}$  and  $\{g_n\}_{n=1}^{\infty}$  in  $H$  be biorthogonal. If both of them are Riesz-Fischer sequences and one of them is complete in  $H$ , then they are Riesz bases for  $H$ .*

The proof is given in Section 3, once we present below some properties of Bessel and Riesz-Fischer sequences and a nice result connecting the two notions (Proposition A).

## 2 Riesz-Fischer sequences and Bessel sequences

In [7, Chapter 4, Section 2, Theorem 3] we find the following two theorems which provide a necessary and sufficient condition so that a sequence in  $H$  is either a Riesz-Fischer sequence or a Bessel sequence. Both results are attributed to **Nina Bari**.

- $\{f_n\}_{n=1}^{\infty}$  is a Riesz-Fischer sequence in  $H$  if and only if there exists a positive number  $A$  so that for any finite scalar sequence  $\{\beta_n\}$  we have

$$A \sum |\beta_n|^2 \leq \left\| \sum \beta_n f_n \right\|^2. \quad (2.1)$$

- $\{f_n\}_{n=1}^{\infty}$  is a Bessel sequence in  $H$  if and only if there exists a positive number  $B$  so that for any finite scalar sequence  $\{\beta_n\}$  we have

$$\left\| \sum \beta_n f_n \right\|^2 \leq B \sum |\beta_n|^2. \quad (2.2)$$

If a sequence is both a Bessel sequence and a Riesz-Fischer sequence, then it is called a **Riesz** sequence (see Seip [5, Lemma 3.2]). That is,  $\{f_n\}_{n=1}^{\infty}$  is a Riesz sequence if there are some positive constants  $A$  and  $B$ ,  $A \leq B$ , so that for any finite scalar sequence  $\{\beta_n\}$  we have

$$A \sum |\beta_n|^2 \leq \left\| \sum \beta_n f_n \right\|^2 \leq B \sum |\beta_n|^2.$$

**Remark 2.1.** *A Riesz sequence is also a Riesz basis for the closure of its linear span in  $H$  (see [2, p. 68]). Therefore, a complete Riesz sequence in  $H$  is a Riesz basis for  $H$ .*

Now, it easily follows from (2.1) that a Riesz-Fischer sequence is also a **minimal** sequence, that is each  $f_n$  does not belong to the closed span of  $\{f_k\}_{k \neq n}$  in  $H$ .

**Remark 2.2.** *It is well known that a sequence is minimal if and only if it has a biorthogonal sequence. A complete and minimal sequence is called **exact** and it has a unique biorthogonal sequence.*

Clearly now a Riesz-Fischer sequence has at least one biorthogonal sequence. As stated in Casazza et al. [1], one of them is a Bessel sequence (see Green [3, Proposition 1.2.4] for a proof).

**Proposition A.** [1, Proposition 2.3, (ii)]

*The Riesz-Fischer sequences in  $H$  are precisely the families for which a biorthogonal Bessel sequence exists. In other words*

*(Part a) Suppose that a Bessel sequence  $\{f_n\}$  is biorthogonal to a sequence  $\{g_n\}$  in  $H$ . Then  $\{g_n\}$  is a Riesz-Fischer sequence.*

*(Part b) If  $\{f_n\}$  is a Riesz-Fischer sequence, then it has a biorthogonal Bessel sequence.*

## 3 Proof of Theorem 1.1

First we prove the following result.

**Lemma 3.1.** *Let two sequences  $\{f_n\}_{n=1}^{\infty}$  and  $\{g_n\}_{n=1}^{\infty}$  in  $H$  be biorthogonal. If  $\{g_n\}_{n=1}^{\infty}$  is a Riesz sequence and  $\{f_n\}_{n=1}^{\infty}$  is complete in  $H$ , then  $\{g_n\}_{n=1}^{\infty}$  is also complete in  $H$ , hence both sequences are Riesz bases for  $\mathcal{H}$ .*

*Proof.* Denote by  $U$  the closed span of  $\{g_n\}_{n=1}^{\infty}$  in  $H$ . Since  $\{g_n\}_{n=1}^{\infty}$  is a Riesz sequence then it is a Riesz basis for  $U$ , therefore it has a dual biorthogonal Riesz basis for  $U$ , call it  $\{h_n\}_{n=1}^{\infty}$ . Thus

$$U = \overline{\text{span}}\{h_n\}_{n=1}^{\infty} = \overline{\text{span}}\{g_n\}_{n=1}^{\infty}.$$

Let  $U^\perp$  be the orthogonal complement of  $U$  in  $H$ , that is

$$U^\perp = \{f \in H : \langle f, g \rangle = 0 \text{ for all } g \in U\}.$$

Hence if  $f \in U$ , then  $\langle f, g_n \rangle = 0$  and  $\langle f, h_n \rangle = 0$  for all  $n \in \mathbb{N}$ .

Now, due to biorthogonality, for fixed  $n \in \mathbb{N}$  we have  $\langle f_n, g_n \rangle = 1$  and  $\langle h_n, g_n \rangle = 1$ . We also have  $\langle f_n, g_m \rangle = 0$  and  $\langle h_n, g_m \rangle = 0$  for all  $m \neq n$ . Therefore,  $\langle (f_n - h_n), g_k \rangle = 0$  for all  $k \in \mathbb{N}$ , thus,  $(f_n - h_n)$  belongs to  $U^\perp$ .

Suppose that  $(f_n - h_n) \neq 0$ . Then  $(f_n - h_n)$  does not belong to  $U = \overline{\text{span}}\{h_n\}_{n=1}^\infty$ . It readily follows that  $f_n$  does not belong to  $U$  either (if  $f_n \in U$  then  $(f_n - h_n) \in U$  as well). Hence  $f_n$  belongs to  $U^\perp$ , so  $\langle f_n, g_k \rangle = 0$  for all  $k \in \mathbb{N}$ , a contradiction since  $\langle f_n, g_n \rangle = 1$ . We have now concluded that  $f_n = h_n$  and clearly this holds for all  $n \in \mathbb{N}$ . Therefore,  $\{h_n\}_{n=1}^\infty = \{f_n\}_{n=1}^\infty$  is a complete Riesz sequence in  $H$ , hence a Riesz basis for  $H$ . The same of course holds for its biorthogonal sequence  $\{g_n\}_{n=1}^\infty$ .  $\square$

**Corollary 3.1.** *Suppose that an exponential system  $E = \{e^{i\lambda_n t}\}$ , with real  $\lambda_n$ , is a Riesz-Fischer sequence in  $L^2(-a, a)$ , and  $E$  has a complete biorthogonal sequence. Then  $E$  is a Riesz basis for  $L^2(-a, a)$ .*

*Proof.* It follows from Lindner [4] that  $E$  is a Bessel sequence in  $L^2(-a, a)$  as well. Hence  $E$  is a Riesz sequence in  $L^2(-a, a)$  and from Lemma 3.1 we obtain the result.  $\square$

Consider now the assumptions of Theorem 1.1 and without loss of generality, suppose that  $\{f_n\}_{n=1}^\infty$  is complete in  $H$ , hence it is exact since it is also a minimal sequence. Thus it has a unique biorthogonal sequence and clearly this is  $\{g_n\}_{n=1}^\infty$ . Since  $\{f_n\}_{n=1}^\infty$  is a Riesz-Fischer sequence, then by Proposition **A** (Part b) it has a biorthogonal Bessel sequence, and by uniqueness, this is  $\{g_n\}_{n=1}^\infty$ . Therefore  $\{g_n\}_{n=1}^\infty$  is a Bessel sequence and a Riesz-Fischer sequence simultaneously, hence a Riesz sequence. It then follows from Lemma 3.1 that  $\{f_n\}_{n=1}^\infty$  and  $\{g_n\}_{n=1}^\infty$  are Riesz bases for  $H$ . The proof of Theorem 1.1 is now complete.

## References

- [1] P. Casazza, O. Christensen, S. Li, A. Lindner, *Riesz-Fischer sequences and lower frame bounds*. Z. Anal. Anwendungen 2002, **21** (2), 305-314.
- [2] O. Christensen, *An introduction to Frames and Riesz Bases*. Applied and Numerical Harmonic Analysis. Birkhäuser Boston, Inc., Boston, MA, xxii+440 pp. ISBN: 0-8176-4295-1 (2003).
- [3] A. W. Green, *The Uncertainty Principle in Control Theory for Partial Differential Equations*. (2020). All Dissertations. 2636. <https://tigerprints.clemson.edu/alldissertations/2636>.
- [4] A. Lindner, *A universal constant for exponential Riesz sequences*, Z. Anal. Anwend. 2000, **19** (2), 553-559.
- [5] K. Seip, *On the connection between exponential bases and certain related sequences in  $L^2(-\pi, \pi)$* . J. Funct. Anal. 1995, **130** (1), 131-160.
- [6] D. T. Stoeva, *On a Characterization of Riesz Bases via Biorthogonal Sequences*. J. Fourier Anal. Appl. 2020, **26** (4), article 67.
- [7] R. M. Young, *An introduction to Nonharmonic Fourier Series*, Revised first edition. Academic Press, Inc., San Diego, CA, xiv+234 pp. ISBN: 0-12-772955-0 (2001).