

# ZERO SETS AND NULLSTELLENSATZ TYPE THEOREMS FOR SLICE REGULAR QUATERNIONIC POLYNOMIALS

ANNA GORI, GIULIA SARFATTI, AND FABIO VLACCI

ABSTRACT. We study the vanishing sets of slice regular polynomials in several quaternionic variables and extend the Weak and Strong Nullstellensatz for such polynomials. In the two variables case we obtain a geometric description of the vanishing sets, leading to a more concrete version of the Strong Nullstellensatz. For polynomials in two variables, we also introduce a non-commutative resultant, defined in terms of a suitable Dieudonné determinant. Furthermore, we use this tool to investigate the existence of common zeros of slice regular polynomials.

## 1. INTRODUCTION

The Hilbert’s Nullstellensatz can be regarded as a generalization of the Fundamental Theorem of Algebra to the case of polynomials in several variables, as pointed out in [16]. This theorem represents a central tool in many active research fields in Mathematics, and especially in Algebraic Geometry. It is generally stated in the framework of algebraically closed fields, but in recent times some new interest has been addressed to a formalization of the Nullstellensatz in a non-commutative setting (see, e.g., [2, 3, 5]). In particular, Alon and Paran in [2] proved both a Weak and a Strong version in the setting of quaternionic polynomials with *central* variables  $\mathbb{H}[x_1, \dots, x_n]$ , equipped with the standard point-wise product.

In the last decade the theory of slice-regular and semi-regular quaternionic functions has proved to be central for the development of the study of quaternionic maps which resemble the main properties of holomorphic and meromorphic functions in the complex setting (see, e.g., [11, 12, 20]). Furthermore, even though an analog of the Fundamental Theorem of Algebra does not hold in general for polynomials over the quaternions, in [13] a positive result in this sense has been obtained for slice regular polynomials over quaternions and octonions. Thus it is quite natural to look for a version of the Nullstellensatz in the framework of slice regular polynomials in  $n$  quaternionic variables  $\mathbb{H}[q_1, \dots, q_n]$ .

It is possible to establish an isomorphism between the ring  $\mathbb{H}[q_1, \dots, q_n]$  equipped with the so called *\*-product* (see Section 2) and the ring of polynomials in several central

---

*Key words and phrases.* Nullstellensatz, quaternionic slice regular polynomials, non-commutative resultant.

Partially supported by by INdAM, through: GNSAGA; INdAM project “Hypercomplex function theory and applications”. It was also partly supported by MIUR, through the projects: Finanziamento Premiale FOE 2014 “Splines for accUrate NumERics: adaptIve models for Simulation Environments”.

variables with the standard point-wise product, considered in [2]. However, there is no correspondence between associated polynomial functions. Indeed, an element in  $\mathbb{H}[x_1, \dots, x_n]$  can be evaluated as a function only on  $n$ -tuples  $(a_1, \dots, a_n) \in \mathbb{H}^n$  with commuting components  $a_l a_m = a_m a_l$  for any  $l, m = 1, \dots, n$  (a nowhere dense subset of  $\mathbb{H}^n$ ) whereas the corresponding regular polynomial function is well defined in  $\mathbb{H}^n$ . We focus our attention to the study of zeros and factorization properties of slice regular polynomials in terms of the *\*-product*; in particular we prove that

**Proposition 1.1.** *Let  $P \in \mathbb{H}[q_1, \dots, q_n]$  be a regular polynomial in  $n$  variables and let  $1 \leq m \leq n$ . Then  $P$  vanishes on  $\mathbb{H}^{m-1} \times \{a\} \times (C_a)^{n-m}$  if and only if there exists  $P_m \in \mathbb{H}[q_1, \dots, q_n]$  such that*

$$P(q_1, \dots, q_n) = (q_m - a) * P_m(q_1, \dots, q_n),$$

where  $C_a := \{q \in \mathbb{H} \mid aq = qa\}$ .

This leads to a characterization of right maximal ideals in  $\mathbb{H}[q_1, \dots, q_n]$ .

**Proposition 1.2.** *Every maximal right ideal in  $\mathbb{H}[q_1, \dots, q_n]$  is of the form*

$$\mathcal{I}_{(a_1, \dots, a_n)} = \{(q_1 - a_1) * P_1 + \dots + (q_n - a_n) * P_n \mid P_1, \dots, P_n \in \mathbb{H}[q_1, \dots, q_n]\}$$

with  $(a_1, \dots, a_n) \in \mathbb{H}^n$ , such that  $a_l a_m = a_m a_l$  for any  $l, m = 1, \dots, n$ . Moreover a polynomial  $P \in \mathcal{I}_{(a_1, \dots, a_n)}$  if and only if  $P(a_1, \dots, a_n) = 0$ .

It then follows

**Theorem 1.3** (Weak Nullstellensatz for slice regular polynomials). *Let  $I$  be a proper right ideal of  $\mathbb{H}[q_1, \dots, q_n]$ . Then all polynomials in  $I$  have a common zero  $(a_1, \dots, a_n) \in \mathbb{H}^n$  such that  $a_l a_m = a_m a_l$  for any  $l, m = 1, \dots, n$ .*

In the complex setting the previous theorem is equivalent to the strong version of the Hilbert Nullstellensatz, which establishes a correspondence between radical ideals and vanishing sets of polynomials. Its proof relies upon the fact that point-evaluation of a polynomial is a ring homomorphism of  $\mathbb{C}[z_1, \dots, z_n]$ , which is no longer true in  $\mathbb{H}[q_1, \dots, q_n]$ . In our setting, given a right ideal  $I$  in  $\mathbb{H}[q_1, \dots, q_n]$ , we define  $\mathcal{V}(I)$  to be the set of common zeros of polynomials in  $I$ ; given a subset  $Z \subseteq \mathbb{H}^n$ , we denote by  $\mathcal{I}(Z)$  the right ideal  $\mathcal{I}(Z) := \bigcap_{(a_1, \dots, a_n) \in Z} \mathcal{I}_{(a_1, \dots, a_n)}$ . We point out that, in the quaternionic

setting,  $\mathcal{I}(Z)$  does not in general coincide with the set of polynomials vanishing on  $Z$ . Indeed the set of polynomials vanishing on a given subset  $Z$  of  $\mathbb{H}^n$  normally is not an ideal. So it is natural to define the ideal  $\mathcal{J}(Z)$  generated by polynomials vanishing on  $Z$  and investigate its relation with  $\mathcal{I}(Z)$ . As shown in Section 4, the two sets  $\mathcal{I}(Z)$  and  $\mathcal{J}(Z)$  coincide, for instance, when  $Z$  consists of points with commuting components.

After introducing the radical of an ideal  $I$  as the intersection of all *completely prime* ideals containing  $I$ , and following a strategy similar to the one in [2], we are able to prove

**Theorem 1.4** (Strong Nullstellensatz for slice regular polynomials). *Let  $I$  be a right ideal in  $\mathbb{H}[q_1, \dots, q_n]$ . Then its radical coincides with  $\mathcal{I}(\mathcal{V}(I))$ .*

In the two quaternionic variable case, we show some relevant geometric properties of the vanishing set of an ideal in  $\mathbb{H}[q_1, q_2]$ ; indeed we prove that either  $\mathcal{V}(I)$  consists of points with commuting components or it is *symmetric* in the first variable with respect to the real axis. This is the key ingredient to show that  $\mathcal{J}(\mathcal{V}(I))$  coincides with  $\mathcal{I}(\mathcal{V}(I))$  in  $\mathbb{H}[q_1, q_2]$  and thus to have a more geometric interpretation of the Strong Nullstellensatz in this framework. These seminal but promising results are in the direction of developing a theory of quaternionic algebraic varieties: this approach will be the object of a forthcoming paper.

The study of the zeros of polynomials and the Hilbert Nullstellensatz in the complex setting, are naturally related with the notion of *resultant*. In the non-commutative setting important contributions in this direction have been obtained in [1, 10, 18, 21] only for polynomials in one or two variables. In this paper we introduce a *regular resultant* for slice regular polynomials of two quaternionic variables defined in terms of a (suitable) determinant, namely an adaptation of the Dieudonné determinant over the quaternions (see, e.g., [4, 6]).

In particular we prove

**Theorem 1.5.** *Given two slice regular polynomials  $P, Q$  in two quaternionic variables, both their regular resultants belong to the right ideal generated in  $\mathbb{H}[q_1, q_2]$  by  $P$  and  $Q$ .*

As a consequence of Proposition 1.2, it is then possible to prove that all common zeros  $(a, b)$  with  $ab = ba$  of two polynomials  $P, Q \in \mathbb{H}[q_1, q_2]$  sit in the zero locus of their regular resultants. This approach can be also used to study the multiplicity of special types of zeros of a slice regular polynomial.

The present paper is organized as follows: in Section 2 we shortly recall the main definitions and results from the theory of slice regular quaternionic functions which will be used in the sequel; in particular, we give a detailed introduction to the  $*$ -product for slice regular functions. The factorization of slice regular polynomials and application to the study of ideals in  $\mathbb{H}[q_1, \dots, q_n]$  is treated in Section 3, where moreover one can find interesting properties of the vanishing sets of slice regular polynomials in two quaternionic variables. The Nullstellensatz type theorems for slice regular polynomials are proved in Section 4; an appropriate formulation is given for ideals in  $\mathbb{H}[q_1, q_2]$ . In Section 5 we introduce a suitable version of Dieudonné determinant for square matrices with slice regular functions as entries, taking into account the  $*$ -product previously defined. In Section 6 we apply this suitable determinant to define a notion of resultant for slice regular polynomials.

## 2. INTRODUCTION TO QUATERNIONIC SLICE REGULAR FUNCTIONS

Let  $\mathbb{H} = \mathbb{R} + i\mathbb{R} + j\mathbb{R} + k\mathbb{R}$  denote the skew field of quaternions and let  $\mathbb{S} = \{q \in \mathbb{H} : q^2 = -1\}$  be the two dimensional sphere of quaternionic imaginary units. Then

$$\mathbb{H} = \bigcup_{J \in \mathbb{S}} (\mathbb{R} + \mathbb{R}J),$$

where the “slice”  $\mathbb{C}_J := \mathbb{R} + \mathbb{R}J$  can be identified with the complex plane  $\mathbb{C}$  for any  $J \in \mathbb{S}$ . In this way, any  $q \in \mathbb{H}$  can be expressed as  $q = x + yJ$  with  $x, y \in \mathbb{R}$  and  $J \in \mathbb{S}$ . The *real part* of  $q$  is  $\operatorname{Re}(q) = x$  and its *imaginary part* is  $\operatorname{Im}(q) = yJ$ ; the *conjugate* of  $q$  is  $\bar{q} := \operatorname{Re}(q) - \operatorname{Im}(q)$ . For any non-real quaternion  $a \in \mathbb{H} \setminus \mathbb{R}$  we will denote by  $J_a := \frac{\operatorname{Im}(a)}{|\operatorname{Im}(a)|} \in \mathbb{S}$  and by  $\mathbb{S}_a := \{\operatorname{Re}(a) + J|\operatorname{Im}(a)| : J \in \mathbb{S}\}$ . If  $a \in \mathbb{R}$ , then  $J_a$  is any imaginary unit.

Let us now recall the definition of slice regular functions of a quaternionic variable. In view of the setting we will be working on, we will restrict to the simplest case possible, that is to slice regular functions defined on open balls centered at the origin,  $B_R := \{q \in \mathbb{H} : |q| < R\}$ , with radius  $R > 0$ , possibly infinite.

For a more general version of the theory of slice regular functions we refer the reader to [11].

**Definition 2.1.** *Let  $R > 0$ . A function  $f : B_R \rightarrow \mathbb{H}$  is called slice regular if the restriction  $f_J$  of  $f$  to  $B_R \cap \mathbb{C}_J$  is holomorphic for any  $J \in \mathbb{S}$ , i.e., if  $f_J$  has continuous partial derivatives and it is such that*

$$\bar{\partial}_I f_J(x + yJ) := \frac{1}{2} \left( \frac{\partial}{\partial x} + J \frac{\partial}{\partial y} \right) f_J(x + yJ) = 0$$

for all  $x + yJ \in B_R \cap \mathbb{C}_J$  and for any  $J \in \mathbb{S}$ .

In the sequel we will denote by  $\mathcal{SR}(B_R)$  the class of slice regular functions on  $B_R \subseteq \mathbb{H}$ , and, for brevity, we will call them simply *regular* functions.

Examples of regular functions, which turn out to be central for our study, are *regular polynomial functions*. In the sequel, we will denote the set of regular polynomials by

$$\mathbb{H}[q] := \left\{ P(q) = \sum_{n=0}^N q^n a_n : \deg P := N \in \mathbb{N}, a_n \in \mathbb{H} \text{ for any } n = 0, \dots, N \right\}.$$

Any convergent *regular power series* of the form  $\sum_{n \in \mathbb{N}} q^n a_n$  with  $a_n \in \mathbb{H}$ , corresponds to a slice regular function on  $B_R$  (see [11]).

**Definition 2.2.** *Let  $R > 0$  and let  $f : B_R \rightarrow \mathbb{H}$  be a slice regular function. For any  $J \in \mathbb{S}$  and any point  $q = x + yJ$  in  $B_R$  we define the Cullen derivative of  $f$  at  $q$  as*

$$\partial_C f(x + yJ) := \frac{1}{2} \left( \frac{\partial}{\partial x} - J \frac{\partial}{\partial y} \right) f_J(x + yJ)$$

While the sum of two regular functions is clearly regular, an appropriate notion of multiplication in  $\mathcal{SR}(B_R)$  is given by the so called *\*-product*.

**Definition 2.3.** If  $f(q) = \sum_{n \in \mathbb{N}} q^n a_n$  and  $g(q) = \sum_{n \in \mathbb{N}} q^n b_n$  are two regular power series, then the *\*-product* of  $f$  and  $g$  is the regular function defined by

$$f * g(q) := \sum_{n \in \mathbb{N}} q^n \sum_{k=0}^n a_k b_{n-k}.$$

Notice that the *\*-product* is associative but not commutative in general. Therefore  $(\mathcal{SR}(B_R), +, *)$  is a non-commutative ring. Furthermore the Leibniz rule holds for *\*-product* of regular functions with respect to the Cullen derivative (see [11]).

It is not difficult to see that if  $f(q) = \sum_{n \in \mathbb{N}} q^n a_n$  has real coefficients  $\{a_n\}$  (i.e.  $f$  is *slice preserving*), then, for any regular power series  $g$ ,

$$f * g(q) = \left( \sum_{n \in \mathbb{N}} q^n a_n \right) \cdot \left( \sum_{n \in \mathbb{N}} q^n b_n \right) = f(q) \cdot g(q) = \left( \sum_{n=0}^{\infty} q^n b_n \right) * \left( \sum_{n=0}^{\infty} q^n a_n \right) = g * f(q),$$

on the domain of convergence of the power series. In general, the relation of the *\*-product* with the usual pointwise product is the following (see [11]):

$$f * g(q) = \begin{cases} 0 & \text{if } f(q) = 0 \\ f(q) \cdot g(f(q)^{-1} \cdot q \cdot f(q)) & \text{if } f(q) \neq 0, \end{cases}$$

for any  $f, g \in \mathcal{SR}(B_R, +, *)$ . Notice that  $f(q)^{-1} \cdot q \cdot f(q)$  belongs to the same sphere  $x + y\mathbb{S}$  as  $q$ . Hence each zero of  $f * g$  in  $x + y\mathbb{S}$  is given either by a zero of  $f$  or by a point which is a conjugate of a zero of  $g$  in the same sphere.

A peculiar aspect of regular functions is the structure of their zero sets. In fact, besides isolated zeros, they can also vanish on two dimensional spheres. As an example, the polynomial  $q^2 + 1$  vanishes on the entire sphere of imaginary units  $\mathbb{S}$ . It can be proven that also spherical zeros cannot accumulate.

**Theorem 2.4.** Let  $f$  be a regular function on  $B_R$ . If  $f$  does not vanish identically, then its zero set consists of isolated points or isolated 2-spheres of the form  $x + y\mathbb{S}$  with  $x, y \in \mathbb{R}$ ,  $y \neq 0$ .

In order to define a *regular inverse*, we need to introduce the following operations.

**Definition 2.5.** Let  $f(q) = \sum_{n \in \mathbb{N}} q^n a_n$  be a regular function on  $B_R$ . Then the regular conjugate of  $f$  is the regular function defined on  $B_R$  by  $f^c = \sum_{n \in \mathbb{N}} q^n \overline{a_n}$ , and the symmetrization of  $f$  is the regular function defined on  $B_R$  by  $f^s = f * f^c = f^c * f$ .

The *regular inverse*  $f^{-*}$  of a regular function  $f : B_R \rightarrow \mathbb{H}$  is defined as

$$(2.1) \quad f^{-*} = (f^s)^{-1} f^c,$$

and it is regular on  $B_R \setminus Z_{f^s}$ , where  $Z_{f^s}$  denotes the zero set of the symmetrization  $f^s$ . The function  $f^{-*}$  is a semi-regular function in the sense of [20].

**Definition 2.6.** A left (right) Ore domain is a domain  $(D, +, \star)$ , that is a ring without zero divisors, such that  $D \star a \cap D \star b \neq \{0\}$  (respectively,  $a \star D \cap b \star D \neq \{0\}$ ), for any  $a, b \in D \setminus \{0\}$ , where  $D \star a := \{d \star a : d \in D\}$  (and  $a \star D := \{a \star d : d \in D\}$ ).

**Theorem 2.7.** Let  $(D, +, \star)$  be a left Ore domain. Then the set of formal quotients  $L = \{a^{-\star} \star b : a, b \in D\}$  can be endowed with operations  $+, \star$  such that:

- (1)  $D$  is isomorphic to the subring  $\{1^{-\star} \star a : a \in D\}$  of  $L$ ;
- (2)  $L$  is a skew field, i.e. a ring where every non-zero element  $a^{-\star} \star b$  has a multiplicative inverse  $(a^{-\star} \star b)^{-\star} = b^{-\star} \star a$ .

The ring  $L$  is called the classical left ring of quotients of  $D$  and, up to isomorphism, it is the only ring having the properties (1) and (2).

On a right Ore domain  $D$ , the classical right ring of quotients is similarly constructed. If  $D$  is both a left and a right Ore domain, then (by the uniqueness property) the two rings of quotients are isomorphic and we speak of the classical ring of quotients of  $D$ , (see [8] for the general theory of ring of quotients in the non-commutative setting).

**Theorem 2.8.** The set of left  $\star$ -quotients

$$\mathcal{L}(B_R) = \{f^{-\star} \star g : f, g \text{ regular in } B_R, f \neq 0\}$$

is a division ring with respect to  $+, \star$ . Moreover, the ring  $\mathcal{SR}(B_R)$  of regular functions on  $B_R$  is a left and right Ore domain and  $\mathcal{L}(B_R)$  is its classical ring of quotients.

All functions in  $\mathcal{L}(B_R)$  are semi-regular functions on  $B_R$ .

The same construction applies to the ring  $(\mathbb{H}[q], +, \star)$ , leading to the definition of the skew-field of quotients of regular polynomials (see [20]).

**2.1. Several variable case.** When considering functions of several quaternionic variables, the definition of slice regularity is more complicated and relies on the notion of *stem functions*. The formulation of this theory (in the more general setting of real alternative  $\star$ -algebras) can be found in [14]. For the sake of simplicity, we will give the main definitions and statements in the case of slice regular polynomials and slice regular analytic functions of two quaternionic variables. All the notions and properties discussed in this section can be easily adapted to the several variables case.

**Example 2.9.** All polynomial functions  $P : \mathbb{H}^2 \rightarrow \mathbb{H}$  of the form

$$P(q_1, q_2) = \sum_{\substack{n=0, \dots, N \\ m=0, \dots, M}} q_1^n q_2^m a_{n,m}$$

with  $a_{n,m} \in \mathbb{H}$  are slice regular on  $\mathbb{H}^2$ . We set  $\deg_{q_1} P := N$  and  $\deg_{q_2} P := M$ . More generally, the sum of a convergent power series

$$f(q_1, q_2) = \sum_{(n,m) \in \mathbb{N}^2} q_1^n q_2^m a_{n,m},$$

is slice regular on domains of the form  $B_{R_1} \times B_{R_2} =: \Omega_{R_1, R_2}$ .

For these classes of slice regular functions it is possible to define in a natural way *partial Cullen derivatives* (see [14, 17]).

**Definition 2.10.** Let  $f(q_1, q_2) = \sum_{(n,m) \in \mathbb{N}^2} q_1^n q_2^m a_{n,m}$  be a slice regular function on  $\Omega_{R_1, R_2}$ . Then the partial Cullen derivative of  $f$  with respect to  $q_1$  is

$$f_{q_1}(q_1, q_2) = \sum_{n \geq 1, m \geq 0} n q_1^{n-1} q_2^m a_{n,m},$$

the partial Cullen derivative of  $f$  with respect to  $q_2$  is

$$f_{q_2}(q_1, q_2) = \sum_{n \geq 0, m \geq 1} m q_1^n q_2^{m-1} a_{n,m}.$$

Slice regular functions of several variables can be endowed with an appropriate notion of product, the so called *slice product*. It will be denoted by the same symbol  $*$  used in the one-variable case. We refer to [14] for a precise definition, let us recall here how it works for convergent power series.

**Definition 2.11.** If  $f(q_1, q_2) = \sum_{n,m \in \mathbb{N}} q_1^n q_2^m a_{n,m}$  and  $g(q) = \sum_{n,m \in \mathbb{N}} q_1^n q_2^m b_{n,m}$  are two slice regular power series which converge in the same domain  $\Omega_{R_1, R_2}$ , then the  $*$ -product of  $f$  and  $g$  is the slice regular power series, convergent on  $\Omega_{R_1, R_2}$ , defined by

$$f * g(q_1, q_2) := \sum_{n,m \in \mathbb{N}} q_1^n q_2^m \sum_{\substack{r=0, \dots, n \\ s=0, \dots, m}} a_{r,s} b_{n-r, m-s}$$

For example, if  $a, b \in \mathbb{H}$ , then

- $q_1 * q_2 = q_2 * q_1 = q_1 q_2$ ;
- $a * (q_1 q_2) = (q_1 q_2) * a = q_1 q_2 a$ ;
- $(q_1^n q_2^m a) * (q_1^r q_2^s b) = q_1^{n+r} q_2^{m+s} ab$ .

moreover we point out that, if  $f$  or  $g$  have real coefficients, then  $f * g = g * f$ .

**Remark 2.12.** Observe that the evaluation of slice regular power series is not a multiplicative homomorphism. Therefore, the zeros of the  $*$  product of two slice regular polynomials is not in general the union of the zeros of each of the factors. For instance,  $q_1 - i$  vanishes on  $\{i\} \times \mathbb{H}$ , while  $(q_1 - i) * (q_2 - j) = q_1 q_2 - q_1 j - q_2 i + k$ , when  $q_1 = i$ , vanishes only for  $q_2 \in \mathbb{C}_i$ .

**Remark 2.13.** As proven in [14], partial Cullen derivatives satisfy the Leibniz rule with respect to the  $*$ -product.

In the sequel we will denote by  $\mathbb{H}[q_1, \dots, q_n]$  the set of slice regular polynomials in  $n$  quaternionic variables. Since the  $*$ -product is associative but not commutative,  $(\mathbb{H}[q_1, \dots, q_n], +, *)$  is a non-commutative ring (without zero divisors).

**Remark 2.14.** From the algebraic point of view, the ring  $(\mathbb{H}[q_1, \dots, q_n], +, *)$  is isomorphic to the ring  $(\mathbb{H}[x_1, \dots, x_n], +, \cdot)$  of quaternionic polynomials in several central variables as considered in [2], but there is no correspondence between associated polynomial functions. Indeed, an element in  $(\mathbb{H}[x_1, \dots, x_n], +, \cdot)$  can be regarded as a function only defined in  $\bigcup_{J \in \mathbb{S}} \mathbb{C}_J \times \dots \times \mathbb{C}_J \in \mathbb{H}^n$ , whereas the corresponding slice regular polynomial function is well defined in  $\mathbb{H}^n$ .

We will deal with ideals in  $\mathbb{H}[q_1, \dots, q_n]$ .

**Definition 2.15.** A subset  $I$  of  $\mathbb{H}[q_1, \dots, q_n]$  is called

- a left ideal if for any  $P \in \mathbb{H}[q_1, \dots, q_n]$ ,  $P * I = \{P * Q : Q \in I\} \subseteq I$ ;
- a right ideal if for any  $P \in \mathbb{H}[q_1, \dots, q_n]$ ,  $I * P = \{Q * P : Q \in I\} \subseteq I$ ;
- a two-sided ideal if  $I$  is both a left and a right ideal.

A different interpretation of the ring  $\mathbb{H}[q_1, \dots, q_n]$  can be given as follows. First notice that the ring  $\mathbb{H}[q_1, q_2]$  of slice regular polynomials in two quaternionic variables can be regarded as the “regular extension” of the ring  $\mathbb{H}[q_2]$  adding the variable  $q_1$ . In fact let  $P \in \mathbb{H}[q_1, q_2]$  be a slice regular polynomial in two quaternionic variables. Then  $P$  can be written as a polynomial in the variable  $q_1$  with coefficients in  $\mathbb{H}[q_2]$

$$(2.2) \quad P(q_1, q_2) = \sum_{n=0}^{\deg_{q_1} P} q_1^n * P_n(q_2) = \sum_{n=0}^{\deg_{q_1} P} q_1^n P_n(q_2)$$

where each  $P_n$  represents a regular polynomial in the quaternionic variable  $q_2$ . We will denote by  $[\mathbb{H}[q_2]]_{\mathcal{SR}}[q_1]$  the set of slice regular polynomials in the variable  $q_1$  with coefficients in  $\mathbb{H}[q_2]$ . Moreover, if one considers another regular polynomial  $Q(q_1, q_2) = \sum_{s=0}^{\deg_{q_1} Q} q_1^s Q_s(q_2)$  in two quaternionic variables, then the  $*$ -product of  $P$  and  $Q$  is given by

$$P * Q(q_1, q_2) = \sum_{k=0}^{\deg_{q_1} P + \deg_{q_1} Q} q_1^k C_k(q_2)$$

where the polynomial coefficients are  $C_k = \sum_{n+s=k} P_n * Q_s$ .

Exchanging the role of the two variables, one gets something slightly different. In fact, every polynomial  $P \in \mathbb{H}[q_1, q_2]$  can be also written as

$$(2.3) \quad P(q_1, q_2) = \sum_{m=0}^{\deg_{q_2} P} q_2^m * \tilde{P}_m(q_1)$$

where  $\tilde{P}_m$  represents a regular polynomial in the quaternionic variable  $q_1$ , for any  $m$ . We point out that in (2.3) there is no immediate analogue formulation in terms of pointwise product as in (2.2).



We will denote by  $[\mathbb{H}[q_1]]_{\mathcal{SR}}[q_2]$  the set of slice regular polynomials in the variable  $q_2$  with coefficients in  $\mathbb{H}[q_1]$ , taking into account that the multiplication involved is the  $*$ -product.

If  $Q$  is another polynomial in  $\mathbb{H}[q_1, q_2]$ ,

$$Q(q_1, q_2) = \sum_{r=0}^{\deg_{q_2} Q} q_2^r * \tilde{Q}_r(q_1)$$

with  $\tilde{Q}_r \in \mathbb{H}[q_1]$ , for any  $r$ , then the  $*$ -product of  $P$  and  $Q$  can also be expressed as

$$P * Q(q_1, q_2) = \sum_{k=0}^{\deg_{q_2} P + \deg_{q_2} Q} q_2^k * \tilde{C}_k(q_1)$$

where the polynomial coefficients are  $\tilde{C}_k = \sum_{n+s=k} \tilde{P}_n * \tilde{Q}_s$ .

The same identifications can be done in  $n$  variables: let  $P$  be a slice regular polynomial in  $n$  variables, namely

$$P(q_1, \dots, q_n) = \sum_{(k_1, \dots, k_n) \in K} q_1^{k_1} q_2^{k_2} \dots q_n^{k_n} a_{k_1, \dots, k_n}$$

where  $K$  is a finite subset of  $\mathbb{N}^n$ . If  $q_l$  is one of the variables of  $P$ , then one can write

$$P(q_1, \dots, q_n) = \sum_{k=0}^{\deg_{q_l} P} q_l^k * \tilde{P}_k(q_1, \dots, q_{l-1}, q_{l+1}, \dots, q_n),$$

with  $\tilde{P}_k(q_1, \dots, q_{l-1}, q_{l+1}, \dots, q_n)$  a regular polynomial in the remaining variables. Hence the set of slice regular polynomials  $P$  in  $n$  variables  $\mathbb{H}[q_1, \dots, q_n]$  can be regarded as  $[\mathbb{H}[q_1, \dots, q_{l-1}, q_{l+1}, \dots, q_n]]_{\mathcal{SR}}[q_l]$  for any  $l = 1, \dots, n$ .

As in the one-variable case, it is possible to introduce two operators on slice regular power series. For the sake of simplicity, we define them only for functions of two quaternionic variables.

**Definition 2.16.** Let  $f(q_1, q_2) = \sum_{n, m \in \mathbb{N}} q_1^n q_2^m a_{n, m}$  be a slice regular function on the domain  $\Omega_{R_1, R_2}$ . Then the regular conjugate of  $f$  is the slice regular function defined on  $\Omega_{R_1, R_2}$  by

$$f^c(q_1, q_2) = \sum_{n, m \in \mathbb{N}} q_1^n q_2^m \overline{a_{n, m}},$$

and the symmetrization of  $f$  is the slice regular function defined on  $\Omega_{R_1, R_2}$  by

$$f^s = f * f^c = f^c * f.$$

The behavior of these operators with respect to the  $*$ -product is the same as the one they have in the one-variable case.

**Remark 2.17.** By direct computation, we have that the coefficients of the power series expansion of  $f^s$  are real numbers and hence  $f^s * g = g * f^s$  for any slice regular function  $g$  (where defined). Moreover, for any  $f, g$  slice regular functions, we also have that  $(f * g)^c = g^c * f^c$  and hence that  $(f * g)^s = f^s * g^s$  (where defined).

**Remark 2.18.** Notice that, since  $(\mathbb{H}[q_1, \dots, q_n], +, *)$  does not contain zero divisors, if  $P$  is a slice regular polynomial not identically zero, then its symmetrization  $P^s \neq 0$ .

A very useful result, that will be used in the sequel, is the following version of *Identity Principle* for slice regular polynomial functions and power series of (one and) several quaternionic variables. See [14, Corollary 2.13] for the details.

**Theorem 2.19** (Identity Principle). *Let  $f, g$  be two slice regular power series in  $n$  quaternionic variables defined on the same domain  $\Omega_{R_1, \dots, R_n}$  (with  $n \geq 1$ ). If there exist  $J_1, \dots, J_n \in \mathbb{S}$  such that  $f \equiv g$  on  $\Omega_{R_1, \dots, R_n} \cap (\mathbb{C}_{J_1} \times \dots \times \mathbb{C}_{J_n})$ , then  $f \equiv g$  on  $\Omega_{R_1, \dots, R_n}$ .*

### 3. SOME REMARKS ON FACTORIZATION OF REGULAR POLYNOMIALS

In what follows we will only consider slice regular polynomials in quaternionic variables. For brevity, we will refer to them simply as *polynomials*.

In the one-variable setting it is possible to perform both left and right Euclidean  $*$ -division between polynomials, see [11, Proposition 3.42].

For polynomials in several quaternionic variables, as in the complex case, we need to restrict our setting to the case of division of polynomials by a *monic* polynomial.

**Definition 3.1.** A polynomial  $P(q_1, \dots, q_2)$  is monic of degree  $d$  in the variable  $q_j$  if it can be written as

$$P(q_1, \dots, q_n) = q_j^d + \sum_{k=1}^{d-1} q_j^k * P_k(q_1, \dots, q_{j-1}, q_{j+1}, \dots, q_n)$$

with  $P_1, \dots, P_d \in \mathbb{H}[q_1, \dots, q_{j-1}, q_{j+1}, \dots, q_n]$ .

**Proposition 3.2.** Let  $M \in \mathbb{H}[q_1, \dots, q_n]$  be a monic polynomial of degree  $d$  in  $q_j$ , with  $1 \leq j \leq n$ . Then, for any  $P \in \mathbb{H}[q_1, \dots, q_n]$ , there exist, and are unique, polynomials  $Q \in \mathbb{H}[q_1, \dots, q_n]$  and  $R_0, \dots, R_{d-1} \in \mathbb{H}[q_1, \dots, q_{j-1}, q_{j+1}, \dots, q_n]$  such that

$$P = M * Q + \sum_{k=0}^{d-1} q_j^k * R_k.$$

*Proof.* Let  $P_0, \dots, P_{s-1} \in \mathbb{H}[q_1, \dots, q_{j-1}, q_{j+1}, \dots, q_n]$  be such that

$$P = \sum_{k=0}^s q_j^k * P_k,$$

and proceed by induction on the degree  $s$  of  $P$  in  $q_j$ .

If  $s < d$ , then we set  $Q \equiv 0$  and  $R_k = P_k$  and we immediately prove the statement. Otherwise, consider  $\hat{P} = P - M * q_j^{s-d} * P_s$ . Since  $M$  is monic of degree  $d$  in  $q_j$ , we get that  $\deg_{q_j} \hat{P} < s$ . We can therefore use the induction hypothesis to write  $\hat{P} = M * \hat{Q} + \sum_{k=0}^{d-1} q_j^k \hat{R}_k$ , so that

$$P = \hat{P} + M * q_j^{s-d} * P_s = M * (\hat{Q} + q_j^{s-d} * P_s) + \sum_{k=0}^{d-1} q_j^k \hat{R}_k.$$

Thus, setting  $Q = \hat{Q} + q_j^{s-d} * P_s$  and  $R_k = \hat{R}_k$ , we conclude.  $\square$

Here we begin the study of zeros of regular polynomials. Our interest is twofold: on the one hand we aim to better understand the structure of the vanishing sets of regular polynomials, on the other hand we want to investigate the relation between such vanishing sets and ideals in  $\mathbb{H}[q_1, \dots, q_n]$ .

**Definition 3.3.** *If  $a \in \mathbb{H}$ , let  $C_a$  be the set of  $q \in \mathbb{H}$  such that  $aq = qa$ , namely*

$$C_a = \begin{cases} \mathbb{C}_{J_a} & \text{if } a \in \mathbb{H} \setminus \mathbb{R} \\ \mathbb{H} & \text{if } a \in \mathbb{R} \end{cases}$$

Let us first consider a simple example that enlightens how the non-commutativity of the product affects the structure of the zero locus of a regular polynomial in two quaternionic variables and shows that these two variables are not interchangeable.

**Example 3.4.** *Let  $a, b \in \mathbb{H}$  be such that  $ab \neq ba$  and consider the regular polynomials*

$$P(q_1, q_2) = (q_1 - a) * (q_2 - b) = q_1 q_2 - q_1 b - q_2 a + ab$$

and

$$Q(q_1, q_2) = (q_2 - b) * (q_1 - a) = q_1 q_2 - q_1 b - q_2 a + ba.$$

For  $q_1 = a$  we have

$$P(a, q_2) = a q_2 - q_2 a \quad \text{and} \quad Q(a, q_2) = a q_2 - ab - q_2 a + ba.$$

Hence  $P$  vanishes on  $\{a\} \times C_a$ , while  $Q$  does not.

On the other hand, for  $q_2 = b$ ,

$$P(q_1, b) = -ba + ab \quad \text{and} \quad Q(q_1, b) \equiv 0.$$

Hence  $Q$  vanishes on  $\mathbb{H} \times \{b\}$ , while  $P$  is never zero when  $q_2 = b$ .

**Remark 3.5.** *In addition to Remark 2.14, observe that there is also no correspondence between the zero locus of polynomials in  $(\mathbb{H}[q_1, q_2], +, *)$  and in  $(\mathbb{H}[x_1, x_2], +, \cdot)$ . In fact, the zero locus of the regular polynomial  $P(q_1, q_2) = (q_2 - i) * (q_1 - j)$  in  $\mathbb{H}^2$  consists of  $\mathbb{H} \times \{i\}$ , while its isomorphic image in  $(\mathbb{H}[x_1, x_2], +, \cdot)$ , namely  $(x_2 - i) \cdot (x_1 - j)$ , vanishes only at  $\mathbb{C}_i \times \{i\}$  (since it cannot be evaluated at  $(a, b)$  if  $ab \neq ba$ ).*

The next Proposition gives a first geometrical description of the zero set of polynomials in several quaternionic variables.

**Proposition 3.6.** *Let  $P \in \mathbb{H}[q_1, \dots, q_n]$  be a regular polynomial in  $n$  variables and let  $1 \leq m \leq n$ . Then  $P$  vanishes on  $\mathbb{H}^{m-1} \times \{a\} \times (C_a)^{n-m}$  if and only if there exists  $P_m \in \mathbb{H}[q_1, \dots, q_n]$  such that*

$$P(q_1, \dots, q_n) = (q_m - a) * P_m(q_1, \dots, q_n).$$

*Proof.* Let  $P_m(q_1, \dots, q_n) := \sum_{\substack{k_j=0, \dots, N_j \\ j=1, \dots, n}} q_1^{k_1} \cdots q_n^{k_n} b_{k_1 \dots k_n}$  be a polynomial in  $\mathbb{H}[q_1, \dots, q_n]$

and consider  $P = (q_m - a) * P_m$ . Then

$$\begin{aligned} P(q_1, \dots, q_n) &= q_m * P_m(q_1, \dots, q_n) - a * P_m(q_1, \dots, q_n) \\ &= \sum_{\substack{k_j=0, \dots, N_j \\ j=1, \dots, n}} q_1^{k_1} \cdots q_m^{k_m+1} \cdots q_n^{k_n} b_{k_1 \dots k_n} - \sum_{\substack{k_j=0, \dots, N_j \\ j=1, \dots, n}} q_1^{k_1} \cdots q_m^{k_m} \cdots q_n^{k_n} \cdot a \cdot b_{k_1 \dots k_n}, \end{aligned}$$

and hence, for any  $(q_1, \dots, q_{m-1}, a, u_{m+1}, \dots, u_n) \in \mathbb{H}^{m-1} \times \{a\} \times (C_a)^{n-m}$ ,

$$\begin{aligned} &P(q_1, \dots, q_{m-1}, a, u_{m+1}, \dots, u_n) \\ &= \sum_{\substack{k_j=0, \dots, N_j \\ j=1, \dots, n}} q_1^{k_1} \cdots a^{k_m+1} \cdots u_n^{k_n} \cdot b_{k_1 \dots k_n} - \sum_{\substack{k_j=0, \dots, N_j \\ j=1, \dots, n}} q_1^{k_1} \cdots a^{k_m} \cdots u_n^{k_n} \cdot a \cdot b_{k_1 \dots k_n} \\ &= 0. \end{aligned}$$

On the other hand, suppose that a polynomial  $P$  vanishes on  $\mathbb{H}^{m-1} \times \{a\} \times (C_a)^{n-m}$ . Then, performing the  $*$ -division of  $P$  by  $q_m - a$ , we obtain

$$P(q_1, \dots, q_n) = (q_m - a) * P_m(q_1, \dots, q_n) + R(q_1, q_2, \dots, q_{m-1}, q_{m+1}, \dots, q_n).$$

Now, for any  $(q_1, \dots, q_{m-1}, a, u_{m+1}, \dots, u_n) \in \mathbb{H}^{m-1} \times \{a\} \times (C_a)^{n-m}$  we have that  $P(q_1, \dots, q_{m-1}, a, u_{m+1}, \dots, u_n) = 0$  and, as in the previous considerations, also

$$[(q_m - a) * P_m(q_1, \dots, q_n)]|_{(q_1, \dots, q_{m-1}, a, u_{m+1}, \dots, u_n)} = 0.$$

Hence  $R$  vanishes identically on  $\mathbb{H}^{m-1} \times (C_a)^{n-m}$ . Thanks to the Identity Principle 2.19, we get that  $R$  is identically zero and we conclude the proof.  $\square$

We now want to establish a relation between zeros of polynomials and ideals in  $\mathbb{H}[q_1, \dots, q_n]$ . If  $(a_1, \dots, a_n) \in \mathbb{H}^n$ , then denote by

$$\mathcal{I}_{(a_1, \dots, a_n)} = \{(q_1 - a_1) * P_1 + \cdots + (q_n - a_n) * P_n \mid P_1, \dots, P_n \in \mathbb{H}[q_1, \dots, q_n]\}.$$

Thanks to the properties of the  $*$ -product, it is not difficult to prove that  $\mathcal{I}_{(a_1, \dots, a_n)}$  is a right ideal, generated (via the  $*$ -product) by  $q_1 - a_1, q_2 - a_2, \dots, q_n - a_n$  in  $\mathbb{H}[q_1, \dots, q_n]$ . For these ideals, thanks to the isomorphism recalled in Remark 2.14, the following version of Lemma 2.1 in [2] holds true. We point that it can be also proven directly.

**Proposition 3.7.** *With the previous notations, if there exist  $l, m \in \{1, \dots, n\}$  such that  $a_l a_m \neq a_m a_l$ , then  $\mathcal{I}_{(a_1, \dots, a_n)} = \mathbb{H}[q_1, \dots, q_n]$ .*

However, the isomorphism between  $(\mathbb{H}[q_1, \dots, q_n], +, *)$  and  $(\mathbb{H}[x_1, \dots, x_n], +, \cdot)$  cannot be used to investigate the vanishing sets of regular polynomials, as already pointed out in Remark 3.5. For this reason we include the proof of the next result, whose statement has an analog counterpart in [2].

**Proposition 3.8.** *Let  $(a_1, \dots, a_n) \in \mathbb{H}^n$  with  $a_l a_m = a_m a_l$  for any  $1 \leq l, m \leq n$ , and let  $\mathcal{I}_{(a_1, \dots, a_n)}$  be the right ideal generated by  $q_1 - a_1, \dots, q_n - a_n$ . Then a polynomial  $P \in \mathbb{H}[q_1, \dots, q_n]$  belongs to  $\mathcal{I}_{(a_1, \dots, a_n)}$  if and only if  $P(a_1, \dots, a_n) = 0$ . Moreover  $\mathcal{I}_{(a_1, \dots, a_n)}$  is a right maximal ideal in  $\mathbb{H}[q_1, \dots, q_n]$ .*

*Proof.* Suppose first that  $P \in \mathcal{I}_{(a_1, \dots, a_n)}$ , namely

$$P(q_1, \dots, q_n) = (q_1 - a_1) * P_1 + \dots + (q_n - a_n) * P_n.$$

Since  $a_l a_m = a_m a_l$ , thanks to Proposition 3.6, we get that  $(q_l - a_l) * P_l$  vanishes at  $(a_1, \dots, a_n)$  for any  $i = 1, \dots, n$ . Therefore  $P(a_1, \dots, a_n) = 0$ .

Conversely, suppose that  $P \in \mathbb{H}[q_1, \dots, q_n]$  vanishes at  $(a_1, \dots, a_n)$ . We will proceed by iterating the Euclidean division algorithm described in Proposition 3.2. First, we divide  $P$  by  $(q_1 - a_1)$ , obtaining  $P_1 \in \mathbb{H}[q_1, \dots, q_n]$  and  $R_1 \in \mathbb{H}[q_2, \dots, q_n]$  such that

$$P = (q_1 - a_1) * P_1 + R_1.$$

Then, we perform the division of  $R_1$  by  $(q_2 - a_2)$ , finding  $P_2 \in \mathbb{H}[q_2, \dots, q_n]$  and  $R_2 \in \mathbb{H}[q_3, \dots, q_n]$  such that  $P = (q_1 - a_1) * P_1 + (q_2 - a_2) * P_2 + R_2$ . We proceed iterating this process, writing at the  $(n-1)$ -th step

$$P = (q_1 - a_1) * P_1 + \dots + (q_{n-1} - a_{n-1}) * P_{n-1} + R_{n-1}$$

where  $R_{n-1}$  is a one-variable polynomial in  $q_n$ . Thanks to Proposition 3.6, evaluating at  $(a_1, \dots, a_n)$  the previous equality gives

$$0 = P(a_1, \dots, a_n) = R_{n-1}(a_n)$$

which, recalling the one-variable theory, implies that there exists  $P_n \in \mathbb{H}[q_n]$  such that  $R_{n-1}(q_n) = (q_n - a_n) * P_n$ . Therefore we have that

$$P = (q_1 - a_1) * P_1 + \dots * (q_n - a_n) * P_n \in \mathcal{I}_{(a_1, \dots, a_n)}.$$

To conclude, suppose there exists a right ideal  $I$  such that  $I \supsetneq \mathcal{I}_{(a_1, \dots, a_n)}$  and consider a polynomial  $Q \in I \setminus \mathcal{I}_{(a_1, \dots, a_n)}$ . Then, after performing the same iterated division algorithm described above, we get

$$Q = (q_1 - a_1) * Q_1 + \dots + (q_n - a_n) * Q_n + S_n,$$

where  $Q_k \in \mathbb{H}[q_k, \dots, q_n]$  for any  $1 \leq k \leq n$  and  $S_n \in \mathbb{H}$  is a constant. Using Proposition 3.6 and the hypothesis that  $Q \notin \mathcal{I}_{(a_1, \dots, a_n)}$ , we have that  $0 \neq Q(a_1, \dots, a_n) = S_n$ , that is  $S_n$  is an invertible element in  $I$ , and so  $I = \mathbb{H}[q_1, \dots, q_n]$ .  $\square$

Let us now investigate what kind of information we obtain from the only assumption that a polynomial vanishes at a point  $(a_1, \dots, a_n)$ .

**Proposition 3.9.** *A regular polynomial in  $P \in \mathbb{H}[q_1, \dots, q_n]$  vanishes at  $(a_1, \dots, a_n) \in \mathbb{H}^n$  if and only if there exist  $P_k \in \mathbb{H}[q_1, \dots, q_k]$  for any  $k = 1, \dots, n$  such that*

$$(3.1) \quad P(q_1, \dots, q_n) = \sum_{k=1}^n (q_k - a_k) * P_k(q_1, \dots, q_k).$$

*Proof.* We start performing the  $*$ -division of  $P$  by  $(q_n - a_n)$ ; we thus obtain

$$P(q_1, \dots, q_n) = (q_n - a_n) * P_n(q_1, \dots, q_n) + R_n(q_1, \dots, q_{n-1}).$$

Notice that  $R_n(q_1, \dots, q_{n-1})$  does not depend on  $q_n$  since  $\deg_{q_n} R < 1$ . If we now divide  $R_n$  by  $(q_{n-1} - a_{n-1})$ , we obtain

$$P(q_1, \dots, q_n) = (q_n - a_n) * P_n(q_1, \dots, q_n) + (q_{n-1} - a_{n-1}) * P_{n-1}(q_1, \dots, q_{n-1}) + R_{n-1}(q_1, \dots, q_{n-2}).$$

Iterating this process, at the  $(n - 2)$ -th step we get

$$P(q_1, \dots, q_n) = (q_n - a_n) * P_n(q_1, \dots, q_n) + (q_{n-1} - a_{n-1}) * P_{n-1}(q_1, \dots, q_{n-1}) + \dots + (q_2 - a_2) * P_2(q_1, q_2) + R_2(q_1)$$

where  $R_2$  is a one-variable polynomial in  $q_1$ . Thanks to Proposition 3.6, evaluating at  $(a_1, \dots, a_n)$  the previous equality gives

$$0 = P(a_1, \dots, a_n) = R_2(a_1)$$

which, recalling the one-variable theory, implies that there exists  $P_1 \in \mathbb{H}[q_1]$  such that  $R_2(q_1) = (q_1 - a_1) * P_1(q_1)$ . Therefore we prove the statement.  $\square$

Let  $(a_1, \dots, a_n) \in \mathbb{H}^n$ , and let us denote by  $E_{(a_1, \dots, a_n)}$  the set of polynomials in  $\mathbb{H}[q_1, \dots, q_n]$  which vanish at  $(a_1, \dots, a_n)$ , namely

$$E_{(a_1, \dots, a_n)} := \left\{ \sum_{k=1}^n (q_k - a_k) * P_k(q_1, \dots, q_k) : P_k \in \mathbb{H}[q_1, \dots, q_k] \text{ for any } k = 1, \dots, n \right\}.$$

**Proposition 3.10.** *Let  $(a_1, \dots, a_n) \in \mathbb{H}^n$ . The set of polynomials vanishing at  $(a_1, \dots, a_n)$  is an ideal if and only if  $a_l a_m = a_m a_l$  for any  $l, m = 1, \dots, n$ .*

*Proof.* If  $(a_1, \dots, a_n) \in \mathbb{H}^n$  is such that  $a_l a_m = a_m a_l$  for any  $l, m = 1, \dots, n$ , then  $E_{(a_1, \dots, a_n)}$  coincides with the ideal  $\mathcal{I}_{(a_1, \dots, a_n)}$ . Indeed  $E_{(a_1, \dots, a_n)} \subseteq \mathcal{I}_{(a_1, \dots, a_n)}$  by definition. On the other hand, by Proposition 3.8 all polynomials in  $\mathcal{I}_{(a_1, \dots, a_n)}$  vanish at  $(a_1, \dots, a_n)$  and hence  $\mathcal{I}_{(a_1, \dots, a_n)} \subseteq E_{(a_1, \dots, a_n)}$ .

Let now  $(a_1, \dots, a_n) \in \mathbb{H}^n$  be such that  $a_l a_m \neq a_m a_l$  for some  $l, m \in \{1, \dots, n\}$  and suppose that  $E_{(a_1, \dots, a_n)}$  is an ideal. Since the polynomials  $q_1 - a_1, \dots, q_n - a_n$  belong to  $E_{(a_1, \dots, a_n)}$ , the same holds for the ideal generated by them  $\mathcal{I}_{(a_1, \dots, a_n)} = \langle q_k - a_k : k = 1, \dots, n \rangle$ . Recalling Proposition 3.7,  $\mathcal{I}_{(a_1, \dots, a_n)} = \mathbb{H}[q_1, \dots, q_n] \subseteq E_{(a_1, \dots, a_n)}$ , a contradiction.  $\square$

The previous Proposition can be read as follows: if a polynomial  $P$  vanishes at  $(a_1, \dots, a_n)$  with  $a_l a_m = a_m a_l$  for any  $l, m = 1, \dots, n$ , then  $P * Q$  still vanishes at  $(a_1, \dots, a_n)$  for any  $Q$  in  $\mathbb{H}[q_1, \dots, q_n]$ . The same is not in general true if a zero does not have commuting components. This is also relevant when considering the zeros of the symmetrization of a polynomial. Recall that if a slice regular function in one quaternionic variable vanishes at  $a \in \mathbb{H}$ , then also its symmetrization does. In several variables, we can show that the situation is the same only assuming the commutativity of the components of the assigned zero.

**Corollary 3.11.** *If  $P \in \mathbb{H}[q_1, \dots, q_n]$  vanishes at  $(a_1, \dots, a_n)$ , with  $a_l a_m = a_m a_l$  for any  $l, m = 1, \dots, n$ , then also its symmetrization does since  $P^s = P * P^c$ .*

This is not necessarily true if  $a_l a_m \neq a_m a_l$  for some  $l, m \in \{1, \dots, n\}$ . As an example, consider  $P(q_1, q_2) = q_1 q_2 - k$ , whose symmetrization is  $P^s(q_1, q_2) = q_1^2 q_2^2 + 1$ . We have  $P(i, j) = 0$ , while  $P^s(i, j) = 2$ .

**3.1. The two variables case.** Using the properties of regular polynomials in one quaternionic variable, we are able to prove several results concerning the vanishing sets of polynomials in two variables which enlighten different interesting phenomena passing from one to several variables.

We begin by assigning the value of the first variable. The next Proposition reminds the geometric origin of spherical zeros for functions of one quaternionic variable.

**Proposition 3.12.** *Let  $a \in \mathbb{H} \setminus \mathbb{R}$ . If  $P(q_1, q_2)$  vanishes on  $\{a\} \times C_a$  and also at  $(a, b)$ , with  $b \notin C_a$ , then  $P$  vanishes on  $\{a\} \times \mathbb{S}_b$ .*

*Proof.* Write

$$P(q_1, q_2) = (q_1 - a) * Q(q_1, q_2)$$

with  $Q(q_1, q_2) = \sum_{\substack{n=0, \dots, N \\ m=0, \dots, M}} q_1^n q_2^m a_{nm}$ . Hence

$$P(q_1, q_2) = q_1 Q(q_1, q_2) - a Q(a^{-1} q_1 a, a^{-1} q_2 a)$$

so that

$$\begin{aligned} P(a, q_2) &= a \sum_{\substack{n=0, \dots, N \\ m=0, \dots, M}} a^n q_2^m a_{nm} - a \sum_{\substack{n=0, \dots, N \\ m=0, \dots, M}} a^{n-1} q_2^m a a_{nm} \\ &= a \sum_{\substack{n=0, \dots, N \\ m=0, \dots, M}} a^{n-1} [a q_2^m - q_2^m a] a_{nm}. \end{aligned}$$

If  $q_2 = x + Ky$ , put  $q_2^m := \alpha_m + K\beta_m$ , with  $x, y, \alpha_m, \beta_m$  real numbers and  $K \in \mathbb{S}$ . Similarly, write  $a^n := u_n + Jv_n$  if  $a = a_o + Ja_1$  with  $a_o, a_1, u_n, v_n$  real numbers and

$J \in \mathbb{S}$ . Hence

$$\begin{aligned} P(a, q_2) &= a \sum_{\substack{n=0, \dots, N \\ m=0, \dots, M}} a^{n-1} \underbrace{[aK - Ka]}_{:=T} \beta_m a_{nm} \\ &= a \sum_{\substack{n=0, \dots, N \\ m=0, \dots, M}} (u_{n-1} + Jv_{n-1}) T \beta_m a_{nm}. \end{aligned}$$

Now

$$T = aK - Ka = (a_o + Ja_1)K - K(a_o + Ja_1) = (JK - KJ)a_1 = 2J \wedge Ka_1;$$

furthermore, since  $J \perp T$ , then  $JT = -TJ$ . Therefore

$$\begin{aligned} P(a, q_2) &= aT \sum_{\substack{n=0, \dots, N \\ m=0, \dots, M}} (u_{n-1} - Jv_{n-1}) \beta_m a_{nm} \\ (3.2) \quad &= aT \sum_{\substack{n=0, \dots, N \\ m=0, \dots, M}} \overline{a^{n-1}} \mathfrak{S}m(q_2^m) a_{nm}. \end{aligned}$$

If  $P$  vanishes at  $(a, b)$  with  $b \notin C_a$ , then  $T \neq 0$  and hence  $P$  vanishes on  $\{a\} \times \mathbb{S}_b$ . Indeed (3.2) is a function of  $\mathfrak{S}m(q_2^m)$  which only depends on  $\text{Re}(q_2)$  and  $|\text{Im}(q_2)|$  and not on the imaginary unit of  $q_2$ .  $\square$

As expected from the Identity Principle, a polynomial in two variables which vanishes on  $\{a\} \times \mathbb{H}$ , with  $a$  not real, is in fact a polynomial in one variable.

**Proposition 3.13.** *Let  $a \in \mathbb{H} \setminus \mathbb{R}$ ; a regular polynomial  $P \in \mathbb{H}[q_1, q_2]$  vanishes on  $\{a\} \times \mathbb{H}$  if and only if there exists a polynomial  $Q \in \mathbb{H}[q_1]$  such that  $P(q_1, q_2) = (q_1 - a) * Q(q_1)$ .*

*Proof.* Write

$$(3.3) \quad P(q_1, q_2) = \sum_{n=0}^N q_1^n P_n(q_2)$$

with  $P_n$  slice-regular polynomials in the variable  $q_2$ . Since  $P$  vanishes on  $\{a\} \times \mathbb{H}$ , it turns out that

$$(3.4) \quad 0 \equiv P(a, q_2) = \sum_{n=0}^N a^n P_n(q_2) \quad \text{for any } q_2 \in \mathbb{H}.$$

If  $\deg_{q_1}(P) = 1$ , then (3.4) becomes  $P_0(q_2) = -aP_1(q_2)$  for any  $q_2 \in \mathbb{H}$ , so that

$$P(q_1, q_2) = -aP_1(q_2) + q_1P_1(q_2) \quad \text{for any } q_2 \in \mathbb{H}$$

and since  $P(q_1, q_2) - q_1P_1(q_2) \in \mathbb{H}[q_1, q_2]$  is a regular polynomial, it follows that  $P_1$  and hence  $P_0$  are constant. Furthermore  $P(q_1, q_2) = (q_1 - a) \cdot C = (q_1 - a) * C$  with  $C \in \mathbb{H}$ .



If  $\deg_{q_1}(P) = N > 1$ , then, as before,  $P_0(q_2) = -aP_1(q_2) - a^2P_2(q_2) - \dots - a^N P_N(q_2)$  for any  $q_2 \in \mathbb{H}$ , so that

$$P(q_1, q_2) = -aP_1(q_2) - a^2P_2(q_2) - \dots - a^N P_N(q_2) + q_1P_1(q_2) + q_1^2P_2(q_2) + \dots + q_1^N P_N(q_2).$$

Since  $P(q_1, q_2) - q_1P_1(q_2) - q_1^2P_2(q_2) - \dots - q_1^N P_N(q_2) \in \mathbb{H}[q_1, q_2]$  is a regular polynomial, it follows that

$$-a[P_1(q_2) - aP_2(q_2) - \dots - a^{N-1}P_N(q_2)]$$

is constant, or  $P_1(q_2) - aP_2(q_2) - \dots - a^{N-1}P_N(q_2) \equiv C_1$  for any  $q_2 \in \mathbb{H}$ , hence

$$C_1 - P_1(q_2) \equiv -a[P_2(q_2) + \dots + a^{N-2}P_N(q_2)] \text{ for any } q_2 \in \mathbb{H}.$$

Since  $C_1 - P_1(q_2)$  is a regular polynomial, it follows that

$$P_2(q_2) + aP_3(q_2) - \dots - a^{N-2}P_N(q_2)$$

is constant and so  $P_1$  is constant as well. After iterating the same procedure, we eventually get that  $P_2, P_3, \dots, P_N$  are constant polynomials. Therefore  $P(q_1, q_2) \in \mathbb{H}[q_1]$  is a regular polynomial in the variable  $q_1$  which vanishes at  $q_1 = a$ , so that there exists  $Q \in \mathbb{H}[q_1]$  such that

$$P(q_1, q_2) = (q_1 - a) * Q(q_1).$$

□

Let us now consider zeros of a regular polynomial with assigned second component

**Proposition 3.14.** *Let  $P \in \mathbb{H}[q_1, q_2]$ . If there exist  $J \in \mathbb{S}$ , a subset  $U \subset \mathbb{C}_J$  with an accumulation point in  $\mathbb{C}_J$  and  $b \in \mathbb{H}$  such that  $P$  vanishes on  $U \times \{b\}$ , then  $P$  vanishes on  $\mathbb{H} \times \{b\}$ .*

*Proof.* Let  $P(q_1, q_2) = \sum_{\substack{n=0, \dots, N \\ m=0, \dots, M}} q_1^n q_2^m a_{n,m}$ . Then

$$Q(q_1) := P(q_1, b) = \sum_{\substack{n=0, \dots, N \\ m=0, \dots, M}} q_1^n b^m a_{n,m}$$

is a regular polynomial in one quaternionic variable vanishing on a subset  $U \subset \mathbb{C}_J$  with an accumulation point in  $\mathbb{C}_J$ . Thanks to the Identity Principle for regular functions of one quaternionic variable [11, Theorem 1.12],  $Q$  is identically zero on  $\mathbb{H}$ , and hence  $P$  vanishes on  $\mathbb{H} \times \{b\}$ . □

Similarly, using the Strong Identity Principle for one variable regular functions [11, Corollary 3.14], we can also prove the following stronger statement.

**Proposition 3.15.** *Let  $P \in \mathbb{H}[q_1, q_2]$ . If there exist a two dimensional sphere  $S = x + y\mathbb{S} \subset \mathbb{H}$  and  $b \in \mathbb{H}$  such that  $P$  vanishes on  $U \times \{b\}$  where  $U \subset \mathbb{H}$  is such that  $U \cap S = \emptyset$  but  $U$  has an accumulation point on  $S$ , then  $P$  vanishes on  $\mathbb{H} \times \{b\}$ .*

We want to describe ideals contained in the set  $E_{(a,b)}$  of polynomials vanishing at a point  $(a, b) \in \mathbb{H}^2$ . First we need the following

**Proposition 3.16.** *A regular polynomial in two variables  $P \in \mathbb{H}[q_1, q_2]$  vanishes on  $\mathbb{S}_a \times \{b\}$  if and only if there exist  $P_1 \in \mathbb{H}[q_1]$  and  $P_2 \in \mathbb{H}[q_1, q_2]$  such that*

$$P(q_1, q_2) = (q_1^2 - 2\operatorname{Re}(a)q_1 + |a|^2) * P_1(q_1) + (q_2 - b) * P_2(q_1, q_2)$$

*Proof.* Dividing  $P$  by  $(q_2 - b)$  as in the proof of Proposition 3.9, we can write

$$P(q_1, q_2) = (q_2 - b) * P_2(q_1, q_2) + R(q_1),$$

and hence  $R(q_1) = P(q_1, q_2) - (q_2 - b) * P_2(q_1, q_2)$  vanishes on  $\mathbb{S}_a \times \{b\}$ . Thus, recalling the one-variable theory,

$$R(q_1) = (q_1^2 - 2\operatorname{Re}(a)q_1 + |a|^2) * P_1(q_1).$$

□

Let us denote by  $E_{\mathbb{S}_a \times \{b\}}$  the set of polynomials vanishing on  $\mathbb{S}_a \times \{b\}$ , namely  $E_{\mathbb{S}_a \times \{b\}} = \{(q_1^2 - 2\operatorname{Re}(a)q_1 + |a|^2) * P_1(q_1) + (q_2 - b) * P_2(q_1, q_2) : P_1 \in \mathbb{H}[q_1], P_2 \in \mathbb{H}[q_1, q_2]\}$ .

**Proposition 3.17.** *The set  $E_{\mathbb{S}_a \times \{b\}}$  is an ideal contained in  $E_{(a,b)}$ .*

*Proof.* The fact that  $(q_1^2 - 2\operatorname{Re}(a)q_1 + |a|^2)$  is a polynomial only in the first variable, with real coefficients, vanishing identically on  $\mathbb{S}_a$ , guarantees that

$$(q_1^2 - 2\operatorname{Re}(a)q_1 + |a|^2) * Q = (q_1^2 - 2\operatorname{Re}(a)q_1 + |a|^2) \cdot Q$$

vanishes on  $\mathbb{S}_a \times \mathbb{H}$  for any  $Q \in \mathbb{H}[q_1, q_2]$ . Hence any polynomial of the form

$$(q_1^2 - 2\operatorname{Re}(a)q_1 + |a|^2) * Q_1 + (q_2 - b) * Q_2,$$

with  $Q_1, Q_2 \in \mathbb{H}[q_1, q_2]$ , vanishes on  $\mathbb{S}_a \times \{b\}$ . Therefore  $E_{\mathbb{S}_a \times \{b\}}$  is an ideal. Since  $a \in \mathbb{S}_a$ ,  $E_{\mathbb{S}_a \times \{b\}} \subset E_{(a,b)}$ . □

Let us now show the following

**Proposition 3.18.** *Let  $a, b \in \mathbb{H}$  be such that  $ab \neq ba$  and let  $I$  be an ideal of  $\mathbb{H}[q_1, q_2]$  contained in  $E_{(a,b)}$ . Then  $I \subseteq E_{\mathbb{S}_a \times \{b\}}$ .*

*Proof.* Let  $P \in I$ . Then  $P(a, b) = 0$ . Consider now  $P * q_2$  which still belongs to  $I$ . Hence

$$0 = P(q_1, q_2) * q_2|_{(a,b)} = q_2 * P(q_1, q_2)|_{(a,b)} = q_2 P(q_2^{-1} q_1 q_2, q_2)|_{(a,b)} = b P(b^{-1} a b, b).$$

Thus the polynomial  $P(\cdot, b) \in \mathbb{H}[q_1]$  vanishes at  $q_1 = a$  and at  $q_1 = b^{-1} a b$  which are two different point on the two sphere  $\mathbb{S}_a$ . By the one-variable theory,  $P(\cdot, b)$  vanishes on the entire sphere  $\mathbb{S}_a$ . Hence  $P \in E_{\mathbb{S}_a \times \{b\}}$ . □

Propositions 3.16 and 3.17, and 3.18 will be used in the next section in order to study the vanishing set of polynomials in  $\mathbb{H}[q_1, q_2]$ ; the first two results can be generalised easily to polynomials in  $n$  variables vanishing on sets of the form  $\mathbb{S}_{a_1} \times \{a_2\} \times \cdots \times \{a_n\}$ , where  $a_l a_m = a_m a_l$  for all  $l, m$  such that  $2 \leq l, m \leq n$ .

#### 4. NULLSTELLENSATZ TYPE THEOREMS FOR REGULAR POLYNOMIALS

In this Section, we further investigate the relations between zero sets of regular polynomials and ideals in  $\mathbb{H}[q_1, \dots, q_n]$ . In the complex setting such correspondence is established by the Hilbert Nullstellensatz. This result admits two equivalent formulations. The first one, known as “Weak Nullstellensatz”, states that  $I$  is a proper ideal of  $\mathbb{C}[z_1, \dots, z_n]$  if and only if there exists a common zero of all polynomials in  $I$ . The second one, the “Strong Nullstellensatz”, is more abstract and involves the notion of *radical* of an ideal. The equivalence of the two statements deeply relies on the fact that point evaluation is a homomorphism (see, e.g., [15]). As already pointed out in Remark 2.12, this is not the case in the quaternionic setting. Using a proper strategy, Alon and Paran in [2] prove a Weak and a Strong Nullstellensatz for quaternionic polynomials with central variables. We will extend their results to prove versions of the Hilbert Nullstellensatz for regular polynomials.

Thanks to the isomorphism introduced in Remark 2.14, we immediately get the following counterpart of Proposition 2.6 in [2].

**Lemma 4.1.** *Every maximal right ideal in  $\mathbb{H}[q_1, \dots, q_n]$  is of the form  $\mathcal{I}_{(a_1, \dots, a_n)}$  with  $(a_1, \dots, a_n) \in \mathbb{C}_J \times \dots \times \mathbb{C}_J =: (\mathbb{C}_J)^n$  for some  $J \in \mathbb{S}$ .*

Combining the previous Lemma with Proposition 3.8, we get the following result for regular polynomials in two quaternionic variables

**Theorem 4.2** (Weak Nullstellensatz for regular polynomials). *Let  $I$  be a proper right ideal of  $\mathbb{H}[q_1, \dots, q_n]$ . Then there exists a point  $(a_1, \dots, a_n) \in (\mathbb{C}_J)^n$ , for some  $J \in \mathbb{S}$ , such that every polynomial in  $I$  vanishes at  $(a_1, \dots, a_n)$ .*

We point out that it is also possible to prove directly the previous statements, following the strategy used in [2], adapted to the setting of regular polynomials.

Let us now introduce some notation.

**Definition 4.3.** *Given a right ideal  $I$  in  $\mathbb{H}[q_1, \dots, q_n]$ , we define  $\mathcal{V}(I)$  to be the set of common zeros of  $f \in I$ , i.e., if  $Z_P \subset \mathbb{H}^n$  represents the zero set of a polynomial  $P \in I$ , then*

$$\mathcal{V}(I) := \bigcap_{P \in I} Z_P.$$

Furthermore, we set

$$\mathcal{V}_c(I) := \mathcal{V}(I) \cap \bigcup_{J \in \mathbb{S}} (\mathbb{C}_J)^n.$$

Recalling the notation introduced in Section 3, we have that  $(a_1, \dots, a_n)$  belongs to  $\mathcal{V}(I)$  if and only if  $I \subseteq E_{(a_1, \dots, a_n)}$ .

**Proposition 4.4.** *Consider the case of a principal right ideal  $\langle P \rangle$  generated by  $P \in \mathbb{H}[q_1, \dots, q_n]$ . Then*

$$\mathcal{V}_c(\langle P \rangle) = Z_P \cap \bigcup_{J \in \mathbb{S}} (\mathbb{C}_J)^n.$$

*Proof.* By definition  $\mathcal{V}_c(\langle P \rangle) \subseteq \bigcup_{J \in \mathbb{S}} (\mathbb{C}_J)^n$  and  $\mathcal{V}_c(\langle P \rangle) \subseteq \mathcal{V}(\langle P \rangle) \subseteq Z_P$ . On the other hand, thanks to Proposition 3.10,  $Z_P \cap \bigcup_{J \in \mathbb{S}} (\mathbb{C}_J)^n \subseteq \mathcal{V}_c(\langle P \rangle)$  and hence they coincide.  $\square$

Theorem 4.2 guarantees that every regular polynomial has at least a zero with commuting components, i.e,  $\mathcal{V}_c(\langle P \rangle) \neq \emptyset$  for any  $P \in \mathbb{H}[q_1, \dots, q_n]$ .

Let us now give an example of a polynomial in  $\mathbb{H}[q_1, q_2]$  with all the zeros with commuting components, that is such that  $\mathcal{V}_c(\langle P \rangle) = \mathcal{V}(\langle P \rangle)$ .

**Example 4.5.** *The polynomial*

$$P(q_1, q_2) = q_1 q_2 - 1$$

*is such that, if  $J \neq K$ , then  $Z_P \cap (\mathbb{C}_J \times \mathbb{C}_K) = \emptyset$ ; indeed  $Z_P = \{(q, q^{-1}) : q \in \mathbb{H} \setminus \{0\}\}$ , which implies that if  $P(a, b) = 0$  there exists  $J \in \mathbb{S}$  such that  $(a, b) \in \mathbb{C}_J \times \mathbb{C}_J$ .*

However, not all polynomials with real coefficients have all the zeros with commuting components: for instance, the polynomial  $q_1^2 + q_2^2 + 2$  vanishes at  $(i, j)$  (and actually at any pair  $(a, b)$  with  $a, b \in \mathbb{S}$ ).

We can associate two different ideals with a given subset of  $\mathbb{H}^n$ .

**Definition 4.6.** *Let  $Z$  be a non-empty subset of  $\mathbb{H}^n$ .*

*By  $\mathcal{J}(Z)$  we denote the right ideal generated by polynomials which vanish on  $Z$ . In other words,*

$$\mathcal{J}(Z) := \left\{ \sum_k P_k * Q_k : P_k, Q_k \in \mathbb{H}[q_1, \dots, q_n] \text{ with } P_k|_Z \equiv 0 \right\};$$

*by  $\mathcal{I}(Z)$  we denote the right ideal*

$$\mathcal{I}(Z) := \bigcap_{(a_1, \dots, a_n) \in Z} \mathcal{I}_{(a_1, \dots, a_n)}.$$

Recalling Proposition 3.10, note that, in general, neither  $\mathcal{J}(Z)$  nor  $\mathcal{I}(Z)$  coincide with the set of polynomials vanishing on  $Z$ . Let us give an example in two variables.

**Example 4.7.** *Consider the case  $Z = \{(a, b)\}$ , from Propositions 3.7 and 3.8 it follows that  $\mathcal{J}(Z) = \mathcal{I}_{(a,b)} = \mathcal{I}(Z)$ . Hence any polynomial  $P \in \mathcal{J}(Z) = \mathcal{I}(Z)$  vanishes at  $(a, b)$  if only if  $ab = ba$ .*

Again by Proposition 3.10, given a right ideal  $I$  in  $\mathbb{H}[q_1, \dots, q_n]$ , both the ideals  $\mathcal{J}(\mathcal{V}_c(I))$  and  $\mathcal{I}(\mathcal{V}_c(I))$  coincide with the set of polynomials vanishing on  $\mathcal{V}_c(I)$ . In particular this yields that we always have the equality

$$\mathcal{J}(\mathcal{V}_c(I)) = \mathcal{I}(\mathcal{V}_c(I)).$$

Observe that

$$\mathcal{I}(\mathcal{V}_c(I)) = \bigcap_{(a_1, \dots, a_n) \in \mathcal{V}_c(I)} \mathcal{I}_{(a_1, \dots, a_n)} = \bigcap_{(a_1, \dots, a_n) \in \mathcal{V}(I)} \mathcal{I}_{(a_1, \dots, a_n)} = \mathcal{I}(\mathcal{V}(I)).$$

Moreover, since  $\mathcal{V}_c(I) \subseteq \mathcal{V}(I)$ , it turns out that

$$(4.1) \quad \mathcal{J}(\mathcal{V}(I)) \subseteq \mathcal{J}(\mathcal{V}_c(I)) = \mathcal{I}(\mathcal{V}_c(I)) = \mathcal{I}(\mathcal{V}(I)).$$

To state a Strong Nullstellensatz for regular polynomials, first we need the following

**Definition 4.8.** *A right ideal  $I$  in  $\mathbb{H}[q_1, \dots, q_n]$  is completely prime if for any  $P, Q \in \mathbb{H}[q_1, \dots, q_n]$  such that  $P * Q \in I$  and  $P * I \subseteq I$  we have that  $P \in I$  or  $Q \in I$ .*

Completely prime ideals play the role of prime ideals in the non-commutative setting. Moreover it can be proven that every maximal ideal is completely prime (see [19]).

The isomorphism introduced in Remark 2.14 guarantees that we can reformulate the analog of Proposition 4.4 as in [2] for regular polynomials.

**Proposition 4.9.** *Every completely prime right ideal in  $\mathbb{H}[q_1, \dots, q_n]$  is the intersection of maximal right ideals.*

The notion of radical of an ideal introduced in [2] can be defined also in the setting of regular polynomials.

**Definition 4.10.** *Let  $I$  be a right ideal in  $\mathbb{H}[q_1, \dots, q_n]$ . The right radical  $\sqrt{I}$  of  $I$  is the intersection of all completely prime right ideals that contain  $I$ .*

We can now state a version of the Strong Nullstellensatz for regular polynomials whose proof follows the line of the one in [2].

**Theorem 4.11.** *Let  $I$  be a right ideal in  $\mathbb{H}[q_1, \dots, q_n]$ . Then*

$$\mathcal{I}(\mathcal{V}(I)) = \sqrt{I}.$$

*Proof.* As we already pointed out, since  $\mathcal{I}(\mathcal{V}(I))$  coincides with the set of all polynomials vanishing on  $\mathcal{V}_c(I)$ , we have that  $\mathcal{I}(\mathcal{V}(I)) = \bigcap_{(a_1, \dots, a_n) \in \mathcal{V}_c(I)} \mathcal{I}_{(a_1, \dots, a_n)}$ . Moreover, recalling

Lemma 4.1, a maximal right ideal  $I_{(a_1, \dots, a_n)}$  contains  $I$ , if and only if  $(a_1, \dots, a_n) \in \mathcal{V}_c(I)$ . Hence  $\mathcal{I}(\mathcal{V}(I))$  is the intersection of all maximal left ideals containing  $I$ . Now, Corollary 2.10 in [19] guarantees that every right maximal ideal in  $\mathbb{H}[q_1, \dots, q_n]$  is completely prime, and Proposition 4.9 states that any completely prime right ideal that contains  $I$  is an intersection of maximal right ideals containing  $I$ . Hence,

$$\sqrt{I} = \bigcap_{\substack{L \supseteq I \\ L \text{ completely prime}}} L = \bigcap_{\substack{L \supseteq I \\ L \text{ maximal}}} L = \mathcal{I}(\mathcal{V}(I)).$$

□

In the two-variable case, thanks to the better understanding of the correspondence between zeros of polynomials and ideals, we are able to give a more concrete version of the Strong Nullstellensatz, stated in terms of the ideal operator  $\mathcal{J}$  instead of  $\mathcal{I}$ .

**Definition 4.12.** A subset  $D \subseteq \mathbb{H}^2$  is called  $q_1$ -symmetric if for any  $(a, b) \in \mathcal{D}$ , the set  $\mathbb{S}_a \times \{b\}$  is contained in  $D$ .

**Proposition 4.13.** Let  $I \in \mathbb{H}[q_1, q_2]$  be an ideal such that  $\mathcal{V}(I)$  is  $q_1$ -symmetric. Then  $\mathcal{J}(\mathcal{V}(I)) = \mathcal{J}(\mathcal{V}_c(I))$ .

*Proof.* Let  $(a, b) \in \mathcal{V}(I) \setminus \mathcal{V}_c(I)$ . Since  $\mathcal{V}(I)$  is  $q_1$ -symmetric, the set  $\mathcal{V}_c(I)$  contains two points  $(a_1, b)$  and  $(\bar{a}_1, b)$  in  $\mathbb{S}_a \times \{b\}$  with  $a_1 b = b a_1$ . Let  $P \in \mathcal{J}(\mathcal{V}_c(I))$ . Then  $P$  vanishes on  $\mathcal{V}_c(I)$  and hence  $P(a_1, b) = P(\bar{a}_1, b) = 0$ . Since  $P(\cdot, b)$  is a slice regular polynomial in  $q_1$ , thanks to the one-variable theory we get that  $P(\cdot, b)$  vanishes on the entire  $\mathbb{S}_a$ . Therefore  $P(q_1, q_2)$  vanishes on  $\mathbb{S}_a \times \{b\}$ , thus on  $(a, b)$ . Since  $(a, b)$  is a generic point in  $\mathcal{V}(I)$ , we have that  $P$  vanishes on  $\mathcal{V}(I)$  and therefore  $P \in \mathcal{J}(\mathcal{V}(I))$ .  $\square$

**Proposition 4.14.** Given an ideal  $I \subseteq \mathbb{H}[q_1, q_2]$ , either  $\mathcal{V}(I) = \mathcal{V}_c(I)$ , or  $\mathcal{V}(I)$  is  $q_1$ -symmetric.

*Proof.* Suppose there exists a point  $(a, b) \in \mathcal{V}(I) \setminus \mathcal{V}_c(I)$ . Since  $I \subseteq E_{(a,b)}$  and  $ab \neq ba$ , thanks to Proposition 3.18, we get  $I \subseteq E_{\mathbb{S}_a \times \{b\}}$  and thus  $\mathbb{S}_a \times \{b\} \subseteq \mathcal{V}(I)$ .  $\square$

The case  $\mathcal{V}_c(I) = \mathcal{V}(I)$ , for example, occurs when  $I = \langle P \rangle$ , with  $P = q_1 q_2 - 1$ . Since  $Z_P \cap (\mathbb{C}_J \times \mathbb{C}_K) = \emptyset$  for imaginary units  $J \neq K$ , then every polynomial in  $\langle P \rangle$  vanishes on  $Z_P$ . Hence  $\mathcal{V}(\langle P \rangle) = Z_P = \mathcal{V}_c(\langle P \rangle)$ .

Combining Propositions 4.13 and 4.14, we obtain

**Theorem 4.15.** Let  $I$  be an ideal in  $\mathbb{H}[q_1, q_2]$ . Then  $\mathcal{J}(\mathcal{V}(I)) = \mathcal{J}(\mathcal{V}_c(I))$ .

Recalling Equation (4.1) we have then the following

**Corollary 4.16** (Strong Nullstellensatz in  $\mathbb{H}^2$ ). Let  $I$  be a right ideal in  $\mathbb{H}[q_1, q_2]$ . Then

$$\mathcal{J}(\mathcal{V}(I)) = \sqrt{I}.$$

In some cases, one can show the equality  $\mathcal{J}(\mathcal{V}(I)) = \mathcal{J}(\mathcal{V}_c(I))$  also in  $\mathbb{H}[q_1, \dots, q_n]$ , with  $n > 2$ . Here is a list of examples.

**Example 4.17.** (1) Any ideal  $I$  in  $\mathbb{H}[q_1, \dots, q_n]$  such that  $\mathcal{V}_c(I) = \mathcal{V}(I)$ .

(2)  $I = \langle q_1 - a_1 \rangle$ ; indeed,  $P \in \langle q_1 - a_1 \rangle$  if and only if  $P$  vanishes on  $\{a_1\} \times (C_{a_1})^{n-1} = \mathcal{V}_c(\langle q_1 - a_1 \rangle)$ . Hence  $\langle q_1 - a_1 \rangle = \mathcal{J}(\mathcal{V}_c(\langle q_1 - a_1 \rangle)) \supseteq \mathcal{J}(\mathcal{V}(\langle q_1 - a_1 \rangle)) \supseteq \langle q_1 - a_1 \rangle$ .

(3)  $I = I_{(a_1, \dots, a_n)}$ ; if  $a_l a_m \neq a_m a_l$  for some  $l, m \in \{1, \dots, n\}$ , then  $I_{(a_1, \dots, a_n)} = \mathbb{H}[q_1, \dots, q_n]$  so  $\mathcal{V}(I_{(a_1, \dots, a_n)}) = \mathcal{V}_c(I_{(a_1, \dots, a_n)}) = \emptyset$ .

If  $a_l a_m = a_m a_l$  for any  $l, m = 1, \dots, n$ , then  $P \in I_{(a_1, \dots, a_n)}$  if and only if  $P$  vanishes at  $(a_1, \dots, a_n)$ . Therefore

$$I_{(a_1, \dots, a_n)} = \mathcal{J}(\mathcal{V}_c(I_{(a_1, \dots, a_n)})) \supseteq \mathcal{J}(\mathcal{V}(I_{(a_1, \dots, a_n)})) \supseteq I_{(a_1, \dots, a_n)}.$$

- (4) Any ideal  $I$  in  $\mathbb{H}[q_1, \dots, q_n]$  such that  $V_c(I) = \emptyset$ . Indeed, if  $\mathcal{V}_c(I) = \emptyset$ , then  $\mathcal{J}(\mathcal{V}_c(I)) = \mathbb{H}[q_1, \dots, q_n]$ . If  $\mathcal{J}(\mathcal{V}(I)) \subsetneq \mathbb{H}[q_1, \dots, q_n]$ , by the Weak Nullstellensatz there would be  $J \in \mathbb{S}$  and a point  $(a_1, \dots, a_n) \in (\mathbb{C}_J)^n$  which should belong to  $\mathcal{V}_c(I)$ .

## 5. PRELIMINARIES ON NON COMMUTATIVE DETERMINANTS

Consider a skew field  $(\mathbb{F}, +, \star)$ , where the product operation  $\star$  is not necessarily commutative. Let  $\overline{\mathbb{F}}^\star$  be the maximal abelian quotient of the multiplicative group  $(\mathbb{F}, \star)$ , i.e. the quotient of  $(\mathbb{F}, \star)$  by its commutator subgroup. In general we will use the notation  $[x]_{\overline{\mathbb{F}}^\star}$  to consider the image in  $\overline{\mathbb{F}}^\star$  of  $x \in \mathbb{F}$ . If  $[x]_{\overline{\mathbb{F}}^\star}$  and  $[y]_{\overline{\mathbb{F}}^\star}$  are given, then, with some abuse of notation, we will write  $[x]_{\overline{\mathbb{F}}^\star} \star [y]_{\overline{\mathbb{F}}^\star} := [x \star y]_{\overline{\mathbb{F}}^\star}$ . Notice that, by definition,  $[x]_{\overline{\mathbb{F}}^\star} \star [y]_{\overline{\mathbb{F}}^\star} = [y]_{\overline{\mathbb{F}}^\star} \star [x]_{\overline{\mathbb{F}}^\star}$ .

The multiplicative group of (invertible)  $n \times n$  matrices with entries in  $\mathbb{F}$ , will be denoted by  $GL(n, \mathbb{F})$ , where, if  $A = (a_{ij})_{1 \leq i, j \leq n}$  and  $B = (b_{ij})_{1 \leq i, j \leq n}$ , with  $a_{ij} \in \mathbb{F}$  and  $b_{ij} \in \mathbb{F}$  for  $1 \leq i, j \leq n$ , then  $A \star_{GL(n, \mathbb{F})} B := \left( \sum_{k=0}^n a_{ik} \star b_{kj} \right)_{1 \leq i, j \leq n}$ . For the sake of shortness, we will simply use the symbol  $\star$  for  $\star_{GL(n, \mathbb{F})}$ . The neutral element for  $\star$  in  $GL(n, \mathbb{F})$  is  $Id_n = (\delta_{ij})_{1 \leq i, j \leq n}$ , where  $\delta_{ij}$  is the Kronecker symbol. Clearly  $A \star B \neq B \star A$  in general.

We recall the following (see, e.g., [4, 7])

**Proposition 5.1.** *There exists a unique homomorphism*

$$\text{Det}_n^\star : GL(n, \mathbb{F}) \rightarrow \overline{\mathbb{F}}^\star$$

such that

$$\text{Det}_n^\star \begin{pmatrix} \lambda_1 & 0 & \dots & \dots & 0 \\ 0 & \lambda_2 & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & 0 & \vdots \\ \vdots & \vdots & \vdots & \ddots & 0 \\ 0 & \dots & 0 & 0 & \lambda_n \end{pmatrix} = [\lambda_1 \star \lambda_2 \star \dots \star \lambda_n]_{\overline{\mathbb{F}}^\star}.$$

As customary, one then extends the definition of  $\text{Det}_n^\star$  also to non-invertible square matrices and put  $\text{Det}_n^\star(A) = [0]_{\overline{\mathbb{F}}^\star}$  if  $A$  is a non-invertible square matrix, where  $0 \in \mathbb{F}$  is the neutral element for  $+$ .

The principal properties of  $\text{Det}_n^\star$  are the following

- (a)  $\text{Det}_n^\star(Id_n) = [1]_{\overline{\mathbb{F}}^\star}$  for any  $n \geq 2$ ;
- (b) if  $\mathbf{r}$  and  $\mathbf{s}$  are distinct rows of  $A \in GL(n, \mathbb{F})$  and if  $\tilde{A}$  is the matrix obtained from  $A$  by replacing the row  $\mathbf{r}$  with the row<sup>1</sup>  $\mathbf{r} + \lambda \star \mathbf{s}$ , then  $\text{Det}_n^\star(\tilde{A}) = \text{Det}_n^\star(A)$ ;

<sup>1</sup>With the following convention : if  $\mathbf{s} = (a_{s1} \ a_{s2} \ \dots \ a_{sn})$  and  $\mathbf{r} = (a_{r1} \ a_{r2} \ \dots \ a_{rn})$ , then  $\mathbf{r} + \lambda \star \mathbf{s} := (a_{r1} + \lambda \star a_{s1} \ a_{r2} + \lambda \star a_{s2} \ \dots \ a_{rn} + \lambda \star a_{sn})$ .

(c) if  $\mathbf{r}$  is a row of  $A \in GL(n, \mathbb{F})$  and if  $\hat{A}$  is the matrix obtained from  $A$  by replacing the row  $\mathbf{r}$  with the row  $\lambda \star \mathbf{r}$ , then  $\text{Det}_n^*(\hat{A}) = [\lambda]_{\mathbb{F}^*} \star \text{Det}_n^*(A)$ .

For  $n = 2$ , if  $a \neq 0$  and if  $a^{-\star}$  is the inverse of  $a$  with respect to  $\star$ , then

$$\text{Det}_2^* \begin{pmatrix} a & b \\ c & d \end{pmatrix} := [a \star d - a \star c \star a^{-\star} \star b]_{\mathbb{F}^*} = [c \star a \star c^{-\star} \star d - c \star b]_{\mathbb{F}^*};$$

if  $a = 0$ , then

$$\text{Det}_2^* \begin{pmatrix} 0 & b \\ c & d \end{pmatrix} := [b \star c]_{\mathbb{F}^*}.$$

Clearly  $\text{Det}_2^* : GL(2, \mathbb{F}) \rightarrow \overline{\mathbb{F}^*}$  satisfies the condition for diagonal  $2 \times 2$  matrices as in Proposition 5.1 and hence also the corresponding property (a) holds. Moreover, by direct computations, one can easily see that for any  $\lambda \in \mathbb{F}$

$$(5.1) \quad \text{Det}_2^* \begin{pmatrix} a + \lambda \star c & b + \lambda \star d \\ c & d \end{pmatrix} = \text{Det}_2^* \begin{pmatrix} a & b \\ c + \lambda \star a & d + \lambda \star b \end{pmatrix} = \text{Det}_2^* \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

and

$$(5.2) \quad \text{Det}_2^* \begin{pmatrix} \lambda \star a & \lambda \star b \\ c & d \end{pmatrix} = \text{Det}_2^* \begin{pmatrix} a & b \\ \lambda \star c & \lambda \star d \end{pmatrix} = [\lambda]_{\mathbb{F}^*} \star \text{Det}_2^* \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

which show that properties (b) and (c) hold for  $\text{Det}_2^*$ .

In order to define  $\text{Det}_n^*$  for  $n \geq 2$ , one proceeds by induction on  $n \geq 2$ . Suppose  $A$  is an  $n \times n$  matrix whose entries are in  $\mathbb{F}$ . If all the elements of the first column of  $A$  are zero then  $\text{Det}_n^*(A) = 0$ . If not, choose a non-zero element  $a_{k1}$  in the first column of  $A$  and subtract a suitable multiple of the  $k$ -th row of  $A$  from each of the other rows of  $A$  in such a way that any other element of the first column of  $A$  vanishes. Call this matrix  $B_k$ . Then consider the  $(n-1) \times (n-1)$  matrix  $B_{k1}$  obtained from  $B_k$  by eliminating the first column and the  $k$ -th row, and define<sup>2</sup>

$$(5.3) \quad \text{Det}_n^*(A) := [a_{k1}]_{\mathbb{F}^*} \star \text{Det}_{n-1}^* B_{k1}.$$

Notice that, from the way it is defined, the properties (a), (b) and (c) proved for  $\text{Det}_2^*$  extends automatically to  $\text{Det}_n^*$ .

In order to prove that the definition of  $\text{Det}_n^*$  in (5.3) is well-posed, one has to show that it is independent from the choice of the non-zero element in the first column of  $A$ : indeed this follows from the above-mentioned properties of  $\text{Det}_n^*$  (see [7] for the details).

---

<sup>2</sup>This is precisely what has been done for  $\text{Det}_2^*$ , since, when  $a \neq 0$ ,

$$\text{Det}_2^* \begin{pmatrix} a & b \\ c & d \end{pmatrix} = [a \star d - a \star c \star a^{-\star} \star b]_{\mathbb{F}^*} = \text{Det}_2^* \begin{pmatrix} a & b \\ 0 & d - c \star a^{-\star} \star b \end{pmatrix}.$$



Another important consequence of Proposition 5.1 (see [6]) is a generalization of the Binet formula, namely if  $A, B \in GL(n, \mathbb{F})$  then

$$(5.4) \quad \text{Det}_n^*(A \star B) = \text{Det}_n^*(A) \star \text{Det}_n^*(B).$$

This fact leads to the following version of the Cramer formula. Consider a system of  $n$  linear equations in  $n$  variables  $x_1, \dots, x_n$  and coefficients  $a_{ij}$ ,  $1 \leq i, j \leq n$ , in  $\mathbb{F}$  (always on the same side),

$$(5.5) \quad \begin{cases} a_{11} \star x_1 + a_{12} \star x_2 + \dots + a_{1n} \star x_n = b_1 \\ a_{21} \star x_1 + a_{22} \star x_2 + \dots + a_{2n} \star x_n = b_2 \\ \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \dots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \\ a_{n1} \star x_1 + a_{n2} \star x_2 + \dots + a_{nn} \star x_n = b_n \end{cases}$$

and write it as

$$A \star \mathbf{x} = \mathbf{b}$$

where  $A = (a_{ij})_{1 \leq i, j \leq n}$ ,  $\mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}$  and  $\mathbf{b} = \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{pmatrix}$ . Observe that

$$A \star \begin{pmatrix} 1 & 0 & \dots & 0 & \overbrace{x_1}^{j\text{-th column}} & 0 & \dots & 0 \\ 0 & 1 & \dots & \vdots & x_2 & \vdots & \dots & \vdots \\ \vdots & 0 & \ddots & 0 & \vdots & \vdots & \dots & \vdots \\ \vdots & \dots & \dots & 1 & \vdots & \vdots & \dots & \vdots \\ \vdots & \dots & \dots & 0 & \ddots & 0 & \dots & \vdots \\ \vdots & \dots & \dots & \vdots & \vdots & 1 & \dots & 0 \\ \vdots & \dots & \dots & \vdots & \vdots & \vdots & \ddots & 0 \\ 0 & \dots & \dots & 0 & x_n & 0 & \dots & 1 \end{pmatrix} = (A_1, A_2, \dots, A_{j-1}, A \star \mathbf{x}, A_{j+1}, \dots, A_n)$$

where  $A_k$  represents the  $k$ -th column of  $A$ . Therefore,

$$\text{Det}_n^*(A) \star [x_j]_{\mathbb{F}}^* = \text{Det}_n^*(A_1, A_2, \dots, A_{j-1}, A \star \mathbf{x}, A_{j+1}, \dots, A_n)$$

which implies that, if  $\mathbf{x}$  is a solution of (5.5) and  $A \in GL(n, \mathbb{F})$ ,

$$(5.6) \quad [x_j]_{\mathbb{F}}^* = \text{Det}_n^*(A)^{-*} \star \text{Det}_n^*(A_1, A_2, \dots, A_{j-1}, \mathbf{b}, A_{j+1}, \dots, A_n),$$

which can be regarded as a generalization of the formula given by Cramer's Theorem for the resolution of Cramer systems of linear equations. Notice that the solution can be also written as

$$(5.7) \quad [x_j]_{\mathbb{F}^*} = \text{Det}_n^*(A_1, A_2, \dots, A_{j-1}, \mathbf{b}, A_{j+1}, \dots, A_n) \star \text{Det}_n^*(A)^{-\star}$$

since  $\text{Det}^*$  takes values in an abelian group.

We are now ready to introduce a notion of determinant of a matrix with entries in the skew field of semi-regular functions  $(\mathcal{L}(B_R), +, \star)$ .

**Definition 5.2.** Assume  $(\mathbb{F}, +, \star) \simeq (\mathcal{L}(B_R), +, \star)$ , with  $a, b, c, d$  semi-regular functions in  $B_R$ . If  $a \neq 0$ , then we define

$$D_2^* \begin{pmatrix} a & b \\ c & d \end{pmatrix} := [a \star d - a \star c \star a^{-\star} \star b]_{\overline{\mathcal{L}(B_R)}^*};$$

If  $a = 0$ ,

$$D_2^* \begin{pmatrix} 0 & b \\ c & d \end{pmatrix} := [b \star c]_{\overline{\mathcal{L}(B_R)}^*}.$$

Since formally  $D_2^*$  is defined as  $\text{Det}_2^*$ , all the properties of  $\text{Det}_2^*$  apply to  $D_2^*$  and, actually, can be used to generalize the definition of a determinant of  $n \times n$  matrices with entries in  $(\mathcal{L}(B_R), +, \star)$  which will be denoted by  $D_n^*$ .

**Proposition 5.3.** If  $M(n, \mathcal{L}(B_R))$  is the set of  $n \times n$  matrices with entries in  $\mathcal{L}(B_R)$ , then

$$D_n^* : M(n, \mathcal{L}(B_R)) \rightarrow \overline{\mathcal{L}(B_R)}^*$$

has all the properties of  $\text{Det}_n^*$  as introduced in Proposition 5.1.

In particular, if all the entries of an  $n \times n$  matrix  $A$  are slice regular polynomials, then one can find a polynomial  $P$  such that  $D_n^*(A) = [P]_{\overline{\mathcal{L}(B_R)}^*}$ .

The second part of the previous statement is a relevant fact proved in [18, Lemma 4.1] in the setting of Ore domains. Such a proof is based on a Euclidean division algorithm for polynomials which can be easily adapted to our setting in the two quaternionic variable case.

## 6. NON COMMUTATIVE RESULTANT

The theory of resultant is a powerful tool in Algebraic Geometry for the investigation of common zeros of polynomials. In the quaternionic setting there have been done some attempts to introduce a resultant of polynomials with quaternionic coefficients see, for instance, [1] and [10] for polynomials in one variable and [18] for polynomials in two variables.

In the present section we want to give a definition of *regular resultant* in  $\mathbb{H}[q_1, q_2]$  and study some of its properties in relation with the existence of common zeros of polynomials. In the complex setting, two polynomials  $S$  and  $T$  have an irreducible common

factor if and only if there exist two polynomials  $U$  and  $V$  with prescribed degrees such that

$$(6.1) \quad S \cdot U + T \cdot V \equiv 0.$$

This is a consequence of the fact that  $\mathbb{C}[z_1, \dots, z_n]$  is a *Unique Factorization Domain* (UFD).

Condition (6.1) can be written as a homogeneous linear system whose unknowns are the coefficients (that might be polynomials in the several variables case) of  $U$  and  $V$ . The existence of a non-zero solution is equivalent to the vanishing of the determinant of the associated matrix, known as the *Sylvester matrix* of  $S$  and  $T$ . This determinant is usually called the *resultant* of  $S$  and  $T$ .

Let us apply this procedure in the setting of regular polynomials. Given  $P, Q \in \mathbb{H}[q_1, q_2]$ , we look for two polynomials  $H, K \in \mathbb{H}[q_1, q_2]$ , of proper degrees, such that

$$(6.2) \quad P * H + Q * K \equiv 0.$$

Due to the non-commutativity of the two variables, the construction of the corresponding linear system depends whether we choose to express  $P, Q, H$  and  $K$  as polynomials in  $q_1$  with coefficients in  $\mathbb{H}[q_2]$  or as polynomials in  $q_2$  with coefficients in  $\mathbb{H}[q_1]$ . More precisely, if

$$\deg_{q_1} P = n, \quad \deg_{q_2} P = r, \quad \deg_{q_1} Q = m, \quad \deg_{q_2} Q = s,$$

then, recalling equation (2.2), and considering  $P, Q, H$  and  $K$  in  $[\mathbb{H}[q_2]]_{\mathcal{SR}}[q_1]$ , we can write

$$P(q_1, q_2) = \sum_{k=0}^n q_1^k P_k(q_2), \quad Q(q_1, q_2) = \sum_{k=0}^m q_1^k Q_k(q_2);$$

as in the complex case, we look for  $H$  and  $K$  such that  $\deg_{q_1} H < \deg_{q_1} Q$ ,  $\deg_{q_1} K < \deg_{q_1} P$ , and hence we can write

$$H(q_1, q_2) = \sum_{p=0}^{m-1} q_1^p H_p(q_2), \quad K(q_1, q_2) = \sum_{p=0}^{n-1} q_1^p K_p(q_2).$$

Then equation (6.2) can be written as

$$(6.3) \quad \sum_{k=0}^n q_1^k P_k(q_2) * \sum_{p=0}^{m-1} q_1^p H_p(q_2) + \sum_{k=0}^m q_1^k Q_k(q_2) * \sum_{p=0}^{n-1} q_1^p K_p(q_2) \equiv 0$$

By the identity principle of polynomials, this equation leads to the following homogeneous linear system:

$$(6.4) \quad \begin{cases} P_0 * H_0 + Q_0 * K_0 & \equiv 0 \\ P_1 * H_0 + P_0 * H_1 + Q_1 * K_0 + Q_0 * K_1 & \equiv 0 \\ \vdots & \vdots \\ P_n * H_{m-1} + Q_m * K_{n-1} & \equiv 0 \end{cases}$$

with unknowns  $H_0, \dots, H_{m-1}, K_0, \dots, K_{n-1}$  in  $\mathbb{H}[q_2]$  and associated matrix, with entries in  $\mathbb{H}[q_2]$ , of dimension  $(m+n) \times (m+n)$

$$A(q_2) = \begin{pmatrix} P_0 & 0 & \cdots & 0 & Q_0 & 0 & \cdots & 0 \\ P_1 & P_0 & \cdots & 0 & Q_1 & Q_0 & \cdots & 0 \\ P_2 & P_1 & \ddots & \vdots & Q_2 & Q_1 & \ddots & \vdots \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & Q_0 \\ P_n & \vdots & \ddots & P_0 & \vdots & \vdots & \ddots & Q_1 \\ 0 & P_n & \ddots & P_1 & \vdots & \vdots & \ddots & \vdots \\ \vdots & 0 & \ddots & \vdots & Q_m & Q_{m-1} & \ddots & \vdots \\ \vdots & \vdots & \ddots & \vdots & 0 & Q_m & \ddots & \vdots \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & P_n & 0 & 0 & \cdots & Q_m \end{pmatrix}.$$

If we choose to express  $P, Q$  as polynomials in  $q_2$  with coefficients in  $\mathbb{H}[q_1]$  then, by equation (2.3), we have  $P(q_1, q_2) = \sum_{k=0}^r q_2^k * \tilde{P}_k(q_1)$  and  $Q(q_1, q_2) = \sum_{k=0}^s q_2^k * \tilde{Q}_k(q_1)$ .

In this case, we look for  $\tilde{H}$  and  $\tilde{K}$  in  $[\mathbb{H}[q_1]]_{S\mathcal{R}}[q_2]$  such that  $\deg_{q_2} \tilde{H} < \deg_{q_2} Q$  and  $\deg_{q_2} \tilde{K} < \deg_{q_2} P$ , i.e.  $\tilde{H}(q_1, q_2) = \sum_{p=0}^{s-1} q_2^p * \tilde{H}_p(q_1)$  and  $\tilde{K}(q_1, q_2) = \sum_{p=0}^{r-1} q_2^p * \tilde{K}_p(q_1)$ , that satisfy

$$(6.5) \quad P * \tilde{H} + Q * \tilde{K} \equiv 0.$$

The previous equation, using again the identity principle for polynomials, is equivalent to a homogeneous linear system in the unknowns  $\tilde{H}_0, \dots, \tilde{H}_{s-1}, \tilde{K}_0, \dots, \tilde{K}_{r-1}$  in  $\mathbb{H}[q_1]$ , whose associated matrix of dimension  $(r+s) \times (r+s)$ , with entries in  $\mathbb{H}[q_1]$ , is

$$B(q_1) = \begin{pmatrix} \tilde{P}_0 & 0 & \cdots & 0 & \tilde{Q}_0 & 0 & \cdots & 0 \\ \tilde{P}_1 & \tilde{P}_0 & \cdots & 0 & \tilde{Q}_1 & \tilde{Q}_0 & \cdots & 0 \\ \tilde{P}_2 & \tilde{P}_1 & \ddots & \vdots & \tilde{Q}_2 & \tilde{Q}_1 & \ddots & \vdots \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \tilde{Q}_0 \\ \tilde{P}_s & \vdots & \ddots & \tilde{P}_0 & \vdots & \vdots & \ddots & \tilde{Q}_1 \\ 0 & \tilde{P}_s & \ddots & \tilde{P}_1 & \vdots & \vdots & \ddots & \vdots \\ \vdots & 0 & \ddots & \vdots & \tilde{Q}_r & \tilde{Q}_{r-1} & \ddots & \vdots \\ \vdots & \vdots & \ddots & \vdots & 0 & \tilde{Q}_r & \ddots & \vdots \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \tilde{P}_s & 0 & 0 & \cdots & \tilde{Q}_r \end{pmatrix}.$$

The existence of solutions of equations (6.2) and (6.5), is then related to the value of the  $D^*$  determinants of the *Sylvester* matrices  $A(q_2)$  and  $B(q_1)$ . It is therefore natural to give the following

**Definition 6.1.** *With the above notations, the regular resultant of  $P$  and  $Q$ , with respect to the variable  $q_1$ , is*

$$\text{Res}(P, Q; q_1) := D_{n+m}^*(A(q_2)),$$

*the regular resultant of  $P$  and  $Q$ , with respect to the variable  $q_2$ , is*

$$\text{Res}(P, Q; q_2) := D_{r+s}^*(B(q_1)).$$

**Remark 6.2.** *Thanks to Proposition 5.3 we have that both the regular resultants have a polynomial representative in  $\mathbb{H}[q_2]$  and  $\mathbb{H}[q_1]$  respectively.*

In [18], Rasheed gives a definition of resultant in the framework of polynomials in two variables sitting in Ore algebras which has an expression similar to  $\text{Res}(P, Q; q_2)$ . Due to the chosen setting, the author only defines the resultant with respect to the second variable and studies how to compute it more efficiently.

In our approach, instead, we make use of our constructive definitions and investigate the relationships between regular resultants and zeros of the involved polynomials. Let us begin with an example that enlightens this phenomenon.

**Example 6.3.** *Consider the polynomials*

$$P(q_1, q_2) = (q_1 - i) * (q_2 - j)$$

*and*

$$Q(q_1, q_2) = (q_1 - i) * (q_2 - k).$$

*As proven in Proposition 3.6, these two polynomials vanish on  $\{i\} \times \mathbb{C}_i$ .*

*To compute the regular resultants, we first consider*

$$A(q_2) = \begin{pmatrix} -i * (q_2 - j) & -i * (q_2 - k) \\ (q_2 - j) & (q_2 - k) \end{pmatrix}.$$

*Then  $\text{Res}(P, Q; q_1) = D_2^*(A(q_2))$  is identically zero as a polynomial in  $q_2$ .*

*On the other hand,*

$$B(q_1) = \begin{pmatrix} (q_1 - i) * j & (q_1 - i) * k \\ (q_1 - i) & (q_1 - i) \end{pmatrix},$$

*thus  $\text{Res}(P, Q; q_2) = D_2^*(B(q_1)) = (q_1 - i)^{*2} * (k - j)$ , that vanishes if and only if  $q_1 = i$ . Therefore, the common zeros of  $P$  and  $Q$  are also zeros of both their regular resultants.*

The first result in the general framework is the following

**Lemma 6.4.** *Let  $P, Q \in \mathbb{H}[q_1, q_2]$ . Then*

- (1)  $\text{Res}(P, Q; q_1) \equiv 0$  if and only if there exist  $H$  and  $K$  in  $[\mathbb{H}[q_2]]_{\mathcal{SR}}[q_1]$ , with  $\deg_{q_1} H < \deg_{q_1} Q$  and  $\deg_{q_1} K < \deg_{q_1} P$ , such that  $P * H + Q * K \equiv 0$ ;

- (2)  $\text{Res}(P, Q; q_2) \equiv 0$  if and only if there exist  $\tilde{H}$  and  $\tilde{K}$  in  $[\mathbb{H}[q_1]]_{\mathcal{SR}}[q_2]$ , with  $\deg_{q_2} \tilde{H} < \deg_{q_2} Q$  and  $\deg_{q_2} \tilde{K} < \deg_{q_2} P$ , such that  $P * \tilde{H} + Q * \tilde{K} \equiv 0$ .

*Proof.* Since we reduced the existence of solutions of equation  $P * H + Q * K \equiv 0$  to the existence of a non-vanishing solution of the homogeneous linear system (6.4), the claim follows using the theory of  $D^*$  determinants introduced in Section 5. Indeed there exist  $H$  and  $K$  in  $[\mathbb{H}[q_2]]_{\mathcal{SR}}[q_1]$  such that  $P * H + Q * K \equiv 0$  if and only if the linear system whose associated matrix is  $A(q_2)$  has  $D_{n+m}^* A(q_2) \equiv 0$ . By Definition 6.1 we conclude. An analogous argument can be used to prove the second equivalence.  $\square$

Let  $\langle P, Q \rangle$  denote the right ideal generated by the regular polynomials  $P$  and  $Q$  in  $\mathbb{H}[q_1, q_2]$ , namely

$$\langle P, Q \rangle := \{P * R + Q * S : R, S \in \mathbb{H}[q_1, q_2]\}.$$

With the notations introduced above, we can state and prove the following

**Theorem 6.5.** *Let  $P, Q \in \mathbb{H}[q_1, q_2]$ . Then*

- (1) *the regular resultant  $\text{Res}(P, Q; q_1)$  belongs to  $\langle P, Q \rangle \cap \mathbb{H}[q_2]$ ;*
- (2) *the regular resultant  $\text{Res}(P, Q; q_2)$  belongs to  $\langle P, Q \rangle \cap \mathbb{H}[q_1]$ .*

*Proof.* We give the proof for  $\text{Res}(P, Q; q_1)$ , the other case can be treated in the same way. By Definition 6.1, and thanks to Remark 6.2,  $\text{Res}(P, Q; q_1) \in \mathbb{H}[q_2]$ . If  $\text{Res}(P, Q; q_1) \equiv 0$ , then the claim follows from Lemma 6.4. When  $\text{Res}(P, Q; q_1) \not\equiv 0$ , consider the following equation

$$P * H + Q * K \equiv 1,$$

with unknowns  $H$  and  $K$  in  $[\mathbb{H}[q_2]]_{\mathcal{SR}}[q_1]$ . Arguing as before, we can transform this equation in a non-homogeneous linear system, analogous to (6.4), whose associated matrix is again  $A(q_2)$ , and whose constant term is  $(0, \dots, 0, 1)^t$ . Now the  $D^*$  determinant of  $A(q_2)$  is  $D_{n+m}^*(A(q_2)) \equiv \text{Res}(P, Q; q_1) \not\equiv 0$ . Hence, thanks to the adapted Cramer Formula (5.7), we can compute a solution in  $\mathbb{H}[q_1, q_2] * \text{Res}(P, Q; q_1)^{-*}$ . If we multiply on the right the obtained solution by  $\text{Res}(P, Q; q_1)$  we conclude the proof.  $\square$

As a direct consequence of the previous result, with the notations introduced in the previous sections, we can say that  $\text{Res}(P, Q; q_1)$  and  $\text{Res}(P, Q; q_2)$  belong to  $\sqrt{\langle P, Q \rangle}$ . Moreover both the regular resultants vanish at the common zeros with commuting components of the polynomials  $P$  and  $Q$ .

**Corollary 6.6.** *Let  $P$  and  $Q$  be two regular polynomials. If they have a common zero with commuting components  $(a, b) \in \mathcal{V}_c(\langle P \rangle) \cap \mathcal{V}_c(\langle Q \rangle)$ , then both  $\text{Res}(P, Q; q_1)$  and  $\text{Res}(P, Q; q_2)$  vanish at  $(a, b)$  as well.*

*Proof.* Recalling Proposition 3.8, we have that  $P, Q \in \mathcal{I}_{(a,b)}$ . Hence also  $\langle P, Q \rangle$  is contained in  $\mathcal{I}_{(a,b)}$ . Since both regular resultants  $\text{Res}(P, Q; q_1)$  and  $\text{Res}(P, Q; q_2)$  belong to  $\langle P, Q \rangle$ .  $\square$

If moreover we know that the two polynomials have a common left (linear) factor, we obtain a stronger result.

**Corollary 6.7.** *Let  $P, Q$  be two regular polynomials in  $\mathbb{H}[q_1, q_2]$  and let  $a, b \in \mathbb{H}$ .*

- (1) *If  $P$  and  $Q$  have the left common factor  $(q_1 - a)$ , then  $\text{Res}(P, Q; q_1) \equiv 0$ ;*
- (2) *if  $P$  and  $Q$  have the left common factor  $(q_2 - b)$ , then  $\text{Res}(P, Q; q_2) \equiv 0$ .*

*Proof.* Suppose first that  $P(q_1, q_2) = (q_1 - a) * P_1(q_1, q_2)$  and  $Q(q_1, q_2) = (q_1 - a) * Q_1(q_1, q_2)$  with  $P_1, Q_1 \in \mathbb{H}[q_1, q_2]$ . Thanks to Theorem 6.5 we have that  $\text{Res}(P, Q; q_1) = (q_1 - a) * R_1(q_1, q_2)$  for some  $R_1 \in \mathbb{H}[q_1, q_2]$ . Recalling Proposition 3.6, we get that  $\text{Res}(P, Q; q_1)$  vanishes on  $\{a\} \times C_a$ . Since  $\text{Res}(P, Q; q_1)$  is a polynomial in  $\mathbb{H}[q_2]$ , the Identity Principle 2.19 yields that it is the zero polynomial.

Suppose now that  $P(q_1, q_2) = (q_2 - b) * P_2(q_1, q_2)$  and  $Q(q_1, q_2) = (q_2 - b) * Q_2(q_1, q_2)$  with  $P_2, Q_2 \in \mathbb{H}[q_1, q_2]$ . Thanks to Theorem 6.5 we have that  $\text{Res}(P, Q; q_2) = (q_2 - b) * R_2(q_1, q_2)$  for some  $R_2 \in \mathbb{H}[q_1, q_2]$ . Recalling Proposition 3.6, we get that  $\text{Res}(P, Q; q_2)$  vanishes on  $\mathbb{H} \times \{b\}$ . Since  $\text{Res}(P, Q; q_2)$  is a polynomial in  $\mathbb{H}[q_1]$ , the Identity Principle 2.19 yields that it is the zero polynomial.  $\square$

In the complex setting, the vanishing of one of the resultants implies the existence of a common factor and this deeply relies on the fact that  $\mathbb{C}[z_1, \dots, z_n]$  is a unique factorization domain. This property of being UFD is not satisfied by the ring  $\mathbb{H}[q_1, q_2]$ : for instance, the polynomial  $q_1^2 + 1$  can be decomposed as  $q_1^2 + 1 = (q_1 - J) * (q_1 + J)$  for any  $J \in \mathbb{S}$ . In our setting, we obtain the following result.

**Proposition 6.8.** *Let  $P$  and  $Q$  be in  $\mathbb{H}[q_1, q_2]$ . If  $\text{Res}(P, Q; q_1) \equiv 0$  or  $\text{Res}(P, Q; q_2) \equiv 0$ , then  $\text{Res}(P^s, Q^s; q_1) \equiv 0$  or, respectively,  $\text{Res}(P^s, Q^s; q_2) \equiv 0$ .*

*Proof.* Suppose  $\text{Res}(P, Q; q_1) \equiv 0$ . Then, thanks to Lemma 6.4, there exist  $H, K \in \mathbb{H}[q_1, q_1]$  such that  $P * H + Q * K \equiv 0$ , with  $\deg_{q_1} H < \deg_{q_1} Q$  and  $\deg_{q_1} K < \deg_{q_1} P$ . Hence

$$P * H \equiv -Q * K, \text{ which implies } P^s * H^s \equiv Q^s * K^s$$

and hence

$$P^s * H^s - Q^s * K^s \equiv 0.$$

Now  $\deg_{q_1} H^s = 2 \deg_{q_1} H < 2 \deg_{q_1} Q = \deg_{q_1} Q^s$  and  $\deg_{q_1} K^s = 2 \deg_{q_1} K < 2 \deg_{q_1} P = \deg_{q_1} P^s$ , and  $P^s$  and  $Q^s$  have real coefficients. Hence, using the theory of resultants in the commutative setting, we get that  $\text{Res}(P^s, Q^s; q_1) \equiv 0$ . The same argument shows that  $\text{Res}(P^s, Q^s; q_2) \equiv 0$ .  $\square$

Applying again the theory of resultants in the commutative setting we obtain the following

**Corollary 6.9.** *Let  $P$  and  $Q$  be in  $\mathbb{H}[q_1, q_2]$ . If  $\text{Res}(P, Q; q_1) \equiv 0$  or  $\text{Res}(P, Q; q_2) \equiv 0$ , then  $P^s$  and  $Q^s$  have a common irreducible factor.*

The notion of regular resultant can be also applied to study the multiplicity of special zeros of polynomials.

**Definition 6.10.** Let  $P$  be a regular polynomial vanishing at  $\{a\} \times C_a$ . We say that the multiplicity of the zero  $\{a\} \times C_a$  of  $P$  is  $m \in \mathbb{N}$  if there exists a polynomial  $Q$  which does not vanish identically on  $\{a\} \times C_a$ , such that

$$P(q_1, q_2) = (q_1 - a)^{*m} * Q(q_1, q_2).$$

Let  $P$  be a regular polynomial vanishing at  $\mathbb{H} \times \{b\}$ . We say that the multiplicity of the zero  $\mathbb{H} \times \{b\}$  of  $P$  is  $n \in \mathbb{N}$  if there exists a polynomial  $S$  which does not vanish identically on  $\mathbb{H} \times \{b\}$ , such that

$$P(q_1, q_2) = (q_2 - b)^{*n} * S(q_1, q_2).$$

**Corollary 6.11.** Let  $(a, b)$  with  $b \in C_a$ , and let  $P$  be a regular polynomial in  $\mathcal{I}_{(a,b)}$  and let  $P_{q_1}$  denote the partial Cullen derivative of  $P$  with respect to  $q_1$ . If the multiplicity  $m$  of  $\{a\} \times C_a$  is greater or equal to 2, then  $\text{Res}(P, P_{q_1}; q_2) \equiv 0$ .

*Proof.* In our hypotheses, thanks to Proposition 3.6,

$$P(q_1, q_2) = (q_1 - a)^{*m} * Q(q_1, q_2)$$

with  $Q$  a polynomial which does not vanish at  $(a, b)$  for any  $b \in C_a$  and  $m \in \mathbb{N}$ . Clearly  $P$  belongs to the right ideal generated by  $q_1 - a$ . Moreover the partial Cullen derivative with respect to  $q_1$  is

$$P_{q_1}(q_1, q_2) = m(q_1 - a)^{*(m-1)} * Q(q_1, q_2) + (q_1 - a)^{*m} * Q_{q_1}(q_1, q_2).$$

Thus  $P_{q_1}$  has  $q_1 - a$  as a left factor if  $m \geq 2$ . Hence, by Corollary 6.7, the resultant  $\text{Res}(P, P_{q_1}; q_2)$  is identically zero. □

Similar arguments also apply for the case of polynomials  $P$  vanishing identically on  $\mathbb{H} \times \{b\}$ .

**Corollary 6.12.** Let  $(a, b)$  with  $b \in C_a$ , and let  $P$  be a regular polynomial in  $\mathcal{I}_{(a,b)}$  and let  $P_{q_2}$  denote the partial Cullen derivative of  $P$  with respect to  $q_2$ . If the multiplicity  $n$  of  $\mathbb{H} \times \{b\}$  is greater or equal to 2, then  $\text{Res}(P, P_{q_2}; q_1) \equiv 0$ .

For polynomials in one variable, the resultant involved in these considerations is usually known as *discriminant*.

## REFERENCES

- [1] A. E. Almendras Valdebenito–J. A. Briones Donoso–A. L. Tironi “Resultants of skew polynomials over division rings”, [arXiv:2109.13092v1](https://arxiv.org/abs/2109.13092v1), (2021)
- [2] G. Alon–E. Paran “A central quaternionic Nullstellensatz”, *J. ALGEBRA*, **574**, p. 252–261, (2021)
- [3] G. Alon–E. Paran “A quaternionic Nullstellensatz”, *J. PURE APPL. ALGEBRA*, **225**, (4) p. 1–9, (2021)
- [4] H. Aslaksen “Quaternionic Determinants” *THE MATHEMATICAL INTELLIGENCER* **18**, p. 57–65, (1996)



- [5] Z. Bao–Z. Reichstein “A non commutative Nullstellensatz” [arXiv:2108.03306v1](https://arxiv.org/abs/2108.03306v1), (2021)
- [6] J.L. Brenner “Applications of the Dieudonné Determinant”, *LINEAR ALGEBRA AND ITS APPLICATIONS*, **1**, p. 511–536, (1968)
- [7] B. Casselman “Essays on the structure of reductive groups – Determinants ”, notes last revised in 2021
- [8] P.M. Cohn **Skew fields – Theory of general division rings** Encyclopedia of Mathematics and its Applications, **57** (2008)
- [9] C. D’Andrea–A. Dickenstein “Explicit formulas for the multivariate resultant” *J. PURE APPL. ALGEBRA*, **164**, p. 59–86, (2001)
- [10] A. Lj. Erić “The resultant of non-commutative polynomials” *MATEMATIČKI VESNIK* **60**, p. 3–8,(2008)
- [11] G. Gentili–C. Stoppato–D.C. Struppa **Regular Functions of a Quaternionic Variable**, Springer Monographs in Mathematics, (2013)
- [12] G. Gentili–D.C Struppa “A new theory of regular functions of a quaternionic variable”, *ADV. MATH.*, **216**, (1), p. 279–301, (2007)
- [13] G. Gentili–D.C. Struppa–F. Vlacci, “The fundamental theorem of algebra for Hamilton and Cayley numbers.” *MATH. Z.* **259**, p. 895–902, (2008)
- [14] R. Ghiloni – A. Perotti “Slice regular functions in several variables”, *MATH. Z* **302** (1) p. 295–351, (2022)
- [15] S. Lang, **Algebra**, Graduate Text in Mathematics **211**, (2002)
- [16] M. Manetti, “Nullstellensatz per tutti”, *Miscellanea Franco Conti*, Centro di Ricerca Matematica “Ennio De Giorgi”, (2006)
- [17] J. Prezelj–F. Vlacci “Divergence zero quaternionic vector fields and Hamming graphs”, *ARS MATHEMATICA CONTEMPORANEA* **19**, p. 189–208, (2020)
- [18] R. Rasheed “Resultant-based Elimination in Ore Algebra” [arXiv:2105.14799v2](https://arxiv.org/abs/2105.14799v2), (2021)
- [19] M. Reyes “A one-sided prime ideal principle for non commutative rings”, *JOURNAL OF ALGEBRA AND ITS APPLICATIONS* p. 877–919 (2010)
- [20] C. Stoppato “Singularities of slice regular functions” *MATH. NACH.* **285**, (2012)
- [21] X. Zhao– Y. Zhang “Resultants of quaternionic polynomials” *HACET. J. MATH. STAT.* **48**, (5), p. 1304–1311, (2019)

DIPARTIMENTO DI MATEMATICA - UNIVERSITÀ DI MILANO, VIA SALDINI 50, 20133 MILANO, ITALY

*Email address:* [anna.gori@unimi.it](mailto:anna.gori@unimi.it)

DIISM - UNIVERSITÀ POLITECNICA DELLE MARCHE, VIA BRECCE BIANCHE 12, 60131 ANCONA, ITALY

*Email address:* [g.sarfatti@univpm.it](mailto:g.sarfatti@univpm.it)

DiSPES - UNIVERSITÀ DI TRIESTE, PIAZZALE EUROPA 1, 34100 TRIESTE, ITALY

*Email address:* [fvlacci@units.it](mailto:fvlacci@units.it)