

A new simple family of Cantor sets in \mathbb{R}^3 all of whose projections are one-dimensional

Olga Frolkina*

Chair of General Topology and Geometry
Faculty of Mechanics and Mathematics
M.V. Lomonosov Moscow State University

and

Moscow Center for Fundamental and Applied Mathematics

Leninskie Gory 1, GSP-1

Moscow 119991, Russia

E-mail: olga-frolkina@yandex.ru

December 6, 2022

Abstract

In 1994, J. Cobb described a Cantor set in \mathbb{R}^3 each of whose projections into 2-planes is one-dimensional. A series of works by other authors developing this field followed. We present another very simple series of Cantor sets in \mathbb{R}^3 all of whose projections are connected and one-dimensional. These are self-similar Cantor sets which go back to the work of Louis Antoine, and we celebrate their centenary birthday in 2020–2021.

Keywords: Euclidean space, projection, Cantor set, embedding, topological dimension, Antoine’s necklace, self-similarity, Hausdorff measure, Hausdorff dimension, Lebesgue measure, demension.

MSC: Primary 54F45; Secondary 57N12, 28A80.

*Supported by Russian Foundation of Basic Research; Grant No. 19–01–00169.

1 Introduction

In 1994, J. Cobb constructed a Cantor set in \mathbb{R}^3 such that its projection into every 2-plane is one-dimensional [12]. In the same paper, he stated a general question: “Given $n > m > k > 0$, does there exist a Cantor set in \mathbb{R}^n such that each of its projections into m -planes is exactly k -dimensional?” Such sets will be called (n, m, k) Cantor sets, or briefly (n, m, k) -sets.

The case $m = k$ was already solved. L. Antoine constructed a Cantor set in \mathbb{R}^2 whose projections coincide with those of a regular hexagon [4, **9**, p. 272; and fig. 2 on p. 273]. (Another $(2, 1, 1)$ set is described in [12, p. 124]; for a different example and further results on planar sets with prescribed projections, see [14].) K. Borsuk constructed an (n, m, m) Cantor set in [11]; as a corollary, he obtained a positive answer to the question of R.H. Fox: there exists a simple closed curve in \mathbb{R}^3 such that each of its plane projections is two-dimensional.

Later, examples for the following triples appeared: for $(n, m, m - 1)$ in [16] and for $(n, n - 1, k)$ in [8] — both generalized Cobb’s idea; and a different construction for $(n, n - 1, n - 2)$ in [17], where further references can be found. Interestingly, all projections of a *typical* Cantor set are Cantor sets [18].

Cobb’s construction of a $(3, 2, 1)$ Cantor set is rather complicated; the resulting set is tame. I suggested a different approach observing that there exist Antoine’s necklaces all of whose projections are one-dimensional [17, Thm. 1]. To obtain a necklace with this property, I controlled the location and thickness of solid tori at prelimit levels of construction; such control can also be generalized to all Cantor sets in all dimensions: For each Cantor set $K \subset \mathbb{R}^n$, $n \geq 2$, there is an arbitrarily small isotopy $\{h_t\} : \mathbb{R}^n \cong \mathbb{R}^n$ such that the projection of $h_1(K)$ into each $(n - 1)$ -plane has dimension $(n - 2)$ [17, Thm. 2].

The control process is not trivial. It is based on the plank theorem (for the planar case, see English translation of A. Tarski’s and H. Moese’s papers (1931–32) in [25, Chapter 7] where the history of the problem is also discussed; the high-dimensional case is treated in [7]).

The control process is necessary. In fact: For each Cantor set $K \subset \mathbb{R}^n$, $n \geq 2$, there exists an arbitrarily small isotopy $\{h_t\} : \mathbb{R}^n \cong \mathbb{R}^n$ such that for each proper subspace $\Pi \subset \mathbb{R}^n$ the dimension of the projection of $h_1(K)$ into Π equals $\dim \Pi$ [17, Stat. 12].

In this article, I restore the original definition of Antoine’s necklaces from [3]. These sets are self-similar. In Theorem 2.6, we formulate sim-

ple additional conditions which ensure that all projections of these sets are one-dimensional. Theorems 2.7 and 2.8 imply that these conditions can actually be satisfied. Self-similarity eliminates the need for thickness control: one-dimensionality is now derived from properties of planar Lebesgue measure or Hausdorff measure.

We also consider a high-dimensional case (Theorem 3.2). Using Rushing’s idea to combine results of Shtan’ko and Väisälä [28, p. 599], we get a new shorter — but less elementary — proof of the above proposition concerning $(n, n - 1, n - 2)$ -sets, see Corollary 3.6; the usage of the plank theorem is eliminated from the proof. Finally, we observe that the sets C^s from [28] are $(3, 2, 1)$ -sets assuming that $1 \leq s < 2$ (Remark 3.4).

Let us underline that the purpose of this note is to propose a possibly simplest example of a $(3, 2, 1)$ Cantor set in a self-contained presentation. By this reason we first give an “elementary” exposition, and thereafter state some generalizations.

Finishing the introduction, I would like to recall some other results of L. Antoine.

Antoine’s necklace is a first example of a wild Cantor set in \mathbb{R}^3 ; its description [3, **78**, p. 91–92], briefly presented in [1], is clear, in contrast to vague descriptions with chains of “curved solid tori” that are often found in later retelling. (Note that in order to get a Cantor set, the diameters of prelimit tori should tend to zero. For self-similar constructions, this holds automatically. Constructions with e.g. “four curved tori” must be provided with a possibility proof.) Antoine constructed first wild arcs in \mathbb{R}^3 . His first example [1], [3, **54–58**, p. 65–70] consists of an arc with a sequence of trefoils tied in it and convergent to the end point, united with a straight line segment. The second wild arc [1], [3, **83**, p. 97] contains an Antoine’s necklace. Antoine constructed an everywhere wild arc in \mathbb{R}^3 [4]. In [2], Antoine announced and in [4] described in detail an embedded 2-sphere in \mathbb{R}^3 which contains an Antoine’s necklace, this is the first example of a wild surface; see [26, Thm. 18.7]. I would recommend the whole volume [5] for acquaintance with Antoine’s destiny and results; and especially [22], [23], and [26, Preface, Sec. 18, and p. 256] concerning the importance of his topological discoveries.

Notation and Conventions

A plane in \mathbb{R}^3 is a 2-dimensional affine subspace.

For a plane $\Pi \subset \mathbb{R}^3$, the orthogonal projection map $\mathbb{R}^3 \rightarrow \Pi$ is denoted by p_Π .

For a set $X \subset \mathbb{R}^3$, its affine hull is denoted by $\text{Aff } X$, and its diameter by $\text{diam } X$.

For topological dimension theory, see [20]; for definition and properties of Lebesgue measure, refer to e.g. [10]. Definitions and basic properties of Hausdorff measure and Hausdorff dimension can be found in [20, Chap. VII] of [13, 3.6]. For Shtan'ko demension theory, see [30]; its presentation is given in [15] and [13, 3.4].

2 Main construction and results: “elementary” approach

There are several (more or less restrictive) definitions of Antoine’s necklaces, compare e.g. [29], [19], [34], [17]. They depend on allowed shapes, numbers and positions of solid tori on prelimit stages. For our purpose, we will only restrict ourselves by the class of self-similar Antoine’s necklaces. This construction comes back to L. Antoine [1], [3].

Definition 2.1 *Let Π be a plane in \mathbb{R}^3 . Let $D \subset \Pi$ be a disk of radius $r > 0$ with center Q , and let $\ell \subset \Pi$ be a straight line such that $d(Q, \ell) = R > r$. A standard solid torus T is the solid torus of revolution generated by revolving D in \mathbb{R}^3 about ℓ . The central circle of T is the circle generated by rotating the point Q . The center of T is the center of its central circle. For a given standard solid torus T , the corresponding radii will be denoted by r_T and R_T , the central circle by C_T , and the center by Q_T .*

Definition 2.2 *A simple chain in a standard solid torus $T \subset \mathbb{R}^3$ is a finite family T_1, \dots, T_k , $k \geq 3$, of pairwise disjoint standard solid tori such that*

- 1) $T_1 \cup \dots \cup T_k \subset T$;
- 2) *centers of T_1, \dots, T_k are subsequent vertices of a regular convex k -gon inscribed in the central circle of T ;*
- 3) T_i and T_j are linked for $|i - j| \equiv 1 \pmod{k}$, and are not linked otherwise. See Fig. 1a.

A simple chain T_1, \dots, T_k looks like a “usual necklace” which winds once around the hole of T ; no one of the tori T_i embraces the hole of T .




Figure 1: Simple chains with $2m = 24$ links.

Definition 2.3 A simple chain T_1, \dots, T_{2m} , $m \geq 2$ in a standard solid torus $T \subset \mathbb{R}^3$ will be called regular if

- 1) all T_i 's are congruent,
- 2) $\text{Aff } C_{T_1} = \text{Aff } C_{T_3} = \dots = \text{Aff } C_{T_{2m-1}} = \text{Aff } C_T$,
- 3) $\text{Aff } C_{T_{2i}} \perp \text{Aff } C_T$ for $i = 1, \dots, m$, and
- 4) each straight line $\text{Aff } C_{T_{2i}} \cap \text{Aff } C_T$, $i = 1, \dots, m$, touches the circle C_T .

See Fig. 1b.

Definition 2.4 Let $T \subset \mathbb{R}^3$ be a standard solid torus, and let $S_1, \dots, S_k : \mathbb{R}^3 \cong \mathbb{R}^3$ be similarity transformations such that $T_1 := S_1(T), \dots, T_k := S_k(T)$ is a simple chain in T . For each $\lambda \in \mathbb{N}$ take

$$M_\lambda = \bigcup_{(i_1, \dots, i_\lambda) \in \{1, \dots, k\}^\lambda} S_{i_1} \circ S_{i_2} \circ \dots \circ S_{i_\lambda}(T).$$

We have $T \supset M_1 \supset M_2 \supset M_3 \supset \dots$. The intersection $\mathcal{A}(T; S_1, \dots, S_k) := \bigcap_{\lambda=1}^{\infty} M_\lambda$ can be easily seen to be a Cantor set; we call it a self-similar Antoine's Necklace generated by $(T; S_1, \dots, S_k)$. For each $i = 1, \dots, k$ the piece $\mathcal{A}(T; S_1, \dots, S_k) \cap T_i$ equals $S_i(\mathcal{A}(T; S_1, \dots, S_k))$, hence it is geometrically similar to $\mathcal{A}(T; S_1, \dots, S_k)$.

A self-similar Antoine’s Necklace will be called regular if the simple chain T_1, \dots, T_k in T is regular. (In particular, k is even, and the similarity coefficients of all S_i ’s are equal.) See Fig. 2.




Figure 2: A piece of a self-similar Antoine’s necklace: first stages of construction.

Remark 2.5 The necklace $\mathcal{A}(T; S_1, \dots, S_k)$ is the invariant set with respect to the set of similarities $\{S_1, \dots, S_k\}$; we thus may omit T from the notation. For details, see [21].

L. Antoine writes: “Numérotions de 1 à k les k tores d’ordre $\lambda + 1$ intérieurs à un tore d’ordre λ . Je supposerai que deux tores ayant des numéros consécutifs, ainsi que les tores k et 1, sont entlacs, alors que deux tores non consécutifs ne sont pas entlacés. Je suppose enfin que les centres de ces k tores sont sur la circonférence d’ordre λ et qu’en joignant ces points dans l’ordre de leurs numéros, on forme un polygone régulier non étoilé. Ces conditions sont possibles à réaliser. Il suffit pour cela que le rayon du méridien d’un tore T soit assez petit, par rapport au rayon de la circonférence C de ce tore, et que k soit assez grand. On peut alors prendre pour tores d’ordre $\lambda + 1$ des tores semblables au tore d’ordre λ . La possibilité de la construction a alors lieu quel que soit λ , k étant fixe. Pour simplifier quelques raisonnements ultérieurs, je *supposerai que ce nombre k est pair*” [3, 78, p. 92]. Antoine’s description does not explicitly indicate similarity transformations, in contrast to what is done in Definition 2.4. But I believe that, taking similar tori, he had in mind precisely this. In [1, I] and [4] he carries out several more constructions using the idea of similarity. Antoine assumed neither that chains have a special position in space, nor he demanded that the solid

tori which constitute a chain are congruent to each other. To obtain examples with a greater degree of symmetry, we added some restrictive conditions (regularity), as is done in [24], [19], [34], [27].

Self-similar Antoine's necklaces with different k 's are inequivalently embedded, this follows from [29, Thm. 2]. For a regular self-similar Antoine's necklace, the inequality $k \geq 20$ holds [34]. A schematic picture of two inequivalently embedded self-similar Antoine's necklaces one of which is regular and the other is not is given in [19]; in that example, $k = 60$.

Theorem 2.6 *Let $\mathcal{A}(T; S_1, \dots, S_k) \subset \mathbb{R}^3$ be a self-similar Antoine's necklace, and let s_i be the similarity coefficient of S_i for $i = 1, \dots, k$. If $s_1^2 + \dots + s_k^2 < 1$, then $p_\Pi(\mathcal{A}(T; S_1, \dots, S_k))$ is a connected one-dimensional set for each plane $\Pi \subset \mathbb{R}^3$.*

In particular, a regular self-similar Antoine's necklace $\mathcal{A}(T; S_1, \dots, S_{2m})$ with $2ms^2 < 1$ is a $(3, 2, 1)$ -set; here s is the similitude coefficient of S_i 's.

Below we will return to this result from a more general point of view, see Theorem 3.2.

Two points should be discussed. The first one: do regular self-similar Antoine's necklaces exist? (Equivalently, can one construct a standard solid torus $T \subset \mathbb{R}^3$ together with a regular simple chain T_1, \dots, T_{2m} in it such that each T_i is similar to T ?) If "yes", the second one: do regular self-similar Antoine's necklaces with $2ms^2 < 1$ exist?

If T_1, \dots, T_{2m} is a regular simple chain in T , and T_i 's are similar to T , then the numbers $\frac{r_T}{R_T}$, m , and s should satisfy several inequalities, see [34], [27]; it seems that no exact description of solutions exist. A picture representing a regular self-similar Antoine's necklace with $2m = 60$ can be found in [19, p. 294]. Using numerical methods, it is shown in [34] that $2m \geq 20$ for any regular self-similar Antoine's necklace; and a picture for $2m = 20$ is given. For $2m = 20, 22, 24, 26, 28$, existence of regular self-similar Antoine's necklaces follows from [27, p. 146], where computer software was also used. Given a standard solid torus T with $\frac{r_T}{R_T}$ sufficiently small, it seems plausible that for all sufficiently large integers m there exists in T a regular simple chain of $2m$ tori similar to T . I have not find an exact statement in the literature. More or less close assertions can be found without proofs in [24, p. 274] ("It can be verified by elementary calculations that for all sufficiently big numbers n , there exist values of λ - not too big and not too small - and corresponding thicknesses of X such that..."; formulaes and a picture follow

saying that the standard solid torus X contains a regular simple chain of n tori similar to X with the similarity coefficient λ) and [27, p. 132] (“...It seems clear that once this minimum is achieved, one can simply add more tori, in specific quantities derived below, to obtain another viable construction of the Antoine necklace”). We give a rigorous statement and complete it by a purely mathematical proof.

Theorem 2.7 *For a standard solid torus $B \subset \mathbb{R}^3$, the following are equivalent:*

- (1) $R_B > 3r_B$;
- (2) for each sufficiently large integer m there exist a standard solid torus T and a regular simple chain T_1, \dots, T_{2m} in T such that each T_i ($i = 1, \dots, 2m$) is congruent to B ;
- (3) for each sufficiently large integer m there exist a standard solid torus \mathbf{T} and a regular simple chain T_1, \dots, T_{2m} in \mathbf{T} such that each T_i ($i = 1, \dots, 2m$) is congruent to B , and \mathbf{T} is similar to B .

Now let us discuss the condition $2ms^2 < 1$ from Theorem 2.6. Are we possible to satisfy it? Yes. Briefly speaking, the similarity coefficient s tends to zero as $m \rightarrow \infty$; but the product ms is bounded. Hence necklaces which satisfy the assumption $2ms^2 < 1$ should exist. We get several simple estimates:

Theorem 2.8 *Let $\mathcal{A}(T; S_1, \dots, S_{2m}) \subset \mathbb{R}^3$ be a regular self-similar Antoine’s necklace, and let s be the similarity coefficient of S_i ’s.*

- 1) If $s < \frac{1}{2\pi}$, then $2ms^2 < 1$.
- 2) If $\frac{r_T}{R_T} < \frac{1}{2\pi-1}$, then $s < \frac{1}{2\pi}$.
- 3) If $2m \geq 40$, then $s < \frac{1}{2\pi}$.

3 A generalization using Hausdorff measure

Definition 3.1 *A zero-dimensional compact set $X \subset \mathbb{R}^n$ is called tame if there exists a homeomorphism h of \mathbb{R}^n onto itself such that $h(X)$ is a subset of a straight line in \mathbb{R}^n ; otherwise, X is called wild.*

In \mathbb{R}^2 each zero-dimensional compactum is tame [3, 75, p. 87–89] (one may also refer to [26, Chap. 13]).

Antoine proved that the necklaces he described are wild [3, 80, p. 93]. A later proof using fundamental group can be found in [26, Sect. 18] or [13, 2.1].

Antoine's construction was generalized to higher dimensions by W.A. Blankinship and by A.A. Ivanov, see [17] for references; hence \mathbb{R}^n contains wild Cantor sets provided that $n \geq 3$.

Theorem 3.2 *Let $X \subset \mathbb{R}^n$ be a wild Cantor set of a zero $(n-1)$ -dimensional Hausdorff measure: $m_{n-1}(X) = 0$. (In particular, this holds if the Hausdorff dimension $\dim_H X < n - 1$.) Then X is an $(n, n - 1, n - 2)$ -set.*

For examples to this theorem, see Remark 3.4 and Corollary 3.6.

Theorem 3.2 covers Theorem 2.6 except connectedness of projections because of the following:

Statement 3.3 *Let $\mathcal{A} = \mathcal{A}(T; S_1, \dots, S_k) \subset \mathbb{R}^3$ be a self-similar Antoine's necklace, and let s_i be the similarity coefficient of S_i for $i = 1, \dots, k$. If $s_1^2 + \dots + s_k^2 < 1$, then $m_2(\mathcal{A}) = 0$.*

Statement 3.3 follows from Moran's theorem, see [28, p. 600], [21, 5.3(1)]; we give a short proof for our simple case.

Proof of Statement 1. For each $\lambda \in \mathbb{N}$, we have $\mathcal{A} \subset M_\lambda$, see Definition 2.4. We can assume (zooming if necessary) that $\text{diam } T = 1$. Now \mathcal{A} is covered by the union of k^λ sets $S_{i_1} \circ S_{i_2} \circ \dots \circ S_{i_\lambda}(T)$, where $(i_1, \dots, i_\lambda) \in \{1, \dots, k\}^\lambda$, and

$$\text{diam } S_{i_1} \circ S_{i_2} \circ \dots \circ S_{i_\lambda}(T) = s_{i_1} \cdot s_{i_2} \cdot \dots \cdot s_{i_\lambda}.$$

We have

$$\begin{aligned} & \sum_{(i_1, \dots, i_\lambda) \in \{1, \dots, k\}^\lambda} \left(\text{diam } S_{i_1} \circ S_{i_2} \circ \dots \circ S_{i_\lambda}(T) \right)^2 \leq \\ & \leq \sum_{(i_1, \dots, i_\lambda) \in \{1, \dots, k\}^\lambda} (s_{i_1} \cdot s_{i_2} \cdot \dots \cdot s_{i_\lambda})^2 = (s_1^2 + \dots + s_k^2)^\lambda. \end{aligned}$$

This tends to zero as $\lambda \rightarrow \infty$, hence the 2-dimensional Hausdorff measure $m_2(\mathcal{A}) = 0$. \square

Remark 3.4 For each s with $1 \leq s \leq 3$, T.B. Rushing [28, Thm. 1, 2] described a wild Cantor set $C^s \subset \mathbb{R}^3$ such that $\dim_H C^s = s$. Theorem 3.2 implies that the sets C^s are $(3, 2, 1)$ Cantor sets, provided that $1 \leq s < 2$. The sets C^s are Antoine-type necklaces; their prelimit chains constructed under very careful control have a rather complex arrangement in space; they do not satisfy our Definition 2.4. Rushing's scheme is too complicated for our purpose; the aim of this paper is to give an example which is as simple as possible.

Remark 3.5 It is apparently an open question, whether wild self-similar Cantor sets in \mathbb{R}^n for $n \geq 4$ do exist [19, p. 298, Question]. T.B. Rushing, without detailed proof, indicates the possibility of this [28, p. 611].

As another application of Theorem 3.2, we get a new proof of our earlier result:

Corollary 3.6 [17, Thm. 2] Let $X \subset \mathbb{R}^n$, $n \geq 2$, be any Cantor set. For each $\varepsilon > 0$ there exists an ε -isotopy $\{h_t\} : \mathbb{R}^n \cong \mathbb{R}^n$ such that $h_1(X)$ is an $(n, n-1, n-2)$ -set.

The new proof is shorter but less elementary since it refers to strong results of Shtan'ko, Edwards and Väisälä.

Proof of Corollary 1. The case of a tame X reduces to known results, see [17] for details.

Let X be wild. We start essentially as in Rushing's remark [28, p. 599]. Suppose we are given an arbitrary $\varepsilon > 0$. According to Väisälä's theorem, there exists an isotopy $\{h_t\} : \mathbb{R}^n \cong \mathbb{R}^n$ such that the Hausdorff dimension $\dim_H(h_1(X))$ equals the embedding dimension $\text{dem } X$ [31], [13, Thm. 3.6.2]. By the Edwards construction [15, p. 208–209], [31, p. 168], this isotopy can be chosen to be an ε -isotopy. For a wild Cantor set $X \subset \mathbb{R}^n$, we have $\text{dem } X \leq n-2$ (for $n \geq 4$, see [30, Remark 2, Prop. 1]; for $n = 3$ use [6, Lemma 4]). Consequently $\dim_H(h_1(X)) \leq n-2$, so $h_1(X)$ is an $(n, n-1, n-2)$ -set by Theorem 3.2.

(Actually, we have $\text{dem } X = n-2$ for a wild Cantor set $X \subset \mathbb{R}^n$. This is a special case of strong results from geometric topology due to several authors, see [30], [15, p. 195; Thm. 1.4 and its proof], [28, p. 598], [13, Thm. 3.4.11, Exer. 3.4.1], [9, p. 5] for statements, historical details and references.) \square

4 Proofs of Theorems 2.6–3.2

Proof of Theorem 1. We already derived Theorem 2.6 (except connectedness of projections) from Theorem 3.2. Here we give an independent “elementary” proof.

For brevity, denote $\mathcal{A} = \mathcal{A}(T; S_1, \dots, S_k)$. By [3, 80–82, p. 93–96], no topological 2-sphere separates a necklace. Hence all projections of a necklace are connected (see [17, Stat. 2] for details).

Let $\Pi \subset \mathbb{R}^3$ be any plane. It is clear that $p_\Pi \mathcal{A}$ contains more than one point; so it remains to show that $\dim p_\Pi \mathcal{A} \neq 2$.

A subset $X \subset \mathbb{R}^n$ is n -dimensional iff it contains an n -ball [20, Thm. IV 3]. Therefore it suffices to prove that $\lambda_2(p_\Pi \mathcal{A}) = 0$, where λ_2 denotes planar Lebesgue measure. Suppose that $\lambda_2(p_\Pi \mathcal{A}) > 0$. (In \mathbb{R}^n , n -dimensional Lebesgue and Hausdorff measures differ by a constant multiple. We thus may use m_2 instead of λ_2 . Here I decided to use λ_2 due to its wider popularity.)

For each non-empty compact subset $X \subset \mathbb{R}^3$ define a real number

$$\alpha(X) = \sup\{\lambda_2(p_L X) \mid L \text{ is a plane in } \mathbb{R}^3\}.$$

Any similarity transformation of \mathbb{R}^n is a composition of a scaling and an isometry. As a corollary, for any similarity transformation $S : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ and any non-empty compactum $X \subset \mathbb{R}^3$ we have $\alpha(S(X)) = s^2 \cdot \alpha(X)$, where s is the similarity coefficient of S .

By assumption, $\alpha(\mathcal{A}) > 0$. Take a positive number

$$\varepsilon \leq \alpha(\mathcal{A}) - (s_1^2 + \dots + s_k^2) \cdot \alpha(\mathcal{A}).$$

Let $L(\varepsilon)$ be any plane such that $\alpha(\mathcal{A}) - \varepsilon < \lambda_2(p_{L(\varepsilon)} \mathcal{A}) \leq \alpha(\mathcal{A})$. Recall that

$$\mathcal{A} = S_1(\mathcal{A}) \cup \dots \cup S_k(\mathcal{A}).$$

We have

$$\begin{aligned} \alpha(\mathcal{A}) - \varepsilon &< \lambda_2(p_{L(\varepsilon)} \mathcal{A}) = \lambda_2\left(p_{L(\varepsilon)}(S_1(\mathcal{A}) \cup \dots \cup S_k(\mathcal{A}))\right) \leq \\ &\leq \lambda_2\left(p_{L(\varepsilon)}(S_1(\mathcal{A}))\right) + \dots + \lambda_2\left(p_{L(\varepsilon)}(S_k(\mathcal{A}))\right) \leq \\ &\leq \alpha(S_1(\mathcal{A})) + \dots + \alpha(S_k(\mathcal{A})) = (s_1^2 + \dots + s_k^2) \cdot \alpha(\mathcal{A}). \end{aligned}$$

This contradicts the choice of ε . Thus $\lambda_2(p_\Pi \mathcal{A}) = 0$, and the proof is finished.

□




Figure 3: Illustration for the proof of the implication (1) \Rightarrow (2) of Theorem 2.

Proof of Theorem 2. Evidently, (2) follows from (3). Also, (2) easily implies (1), see [28, p. 604, (1)] or [27, p. 136, (2)].

(1) \Rightarrow (2) is intuitively clear, but its rigorous proof is rather long.

Step 1: construct a special link. Take any number $A \in (R_B + r_B, 2(R_B - r_B))$; the interval is non-empty since $R_B > 3r_B$. Define the circles $C_{B_1}, C_{B_2} \subset \mathbb{R}^3$ by their parametric equations:

$$C_{B_1}: \quad x = R_B \cos u, \quad y = R_B \sin u, \quad z = 0$$

and

$$C_{B_2}: \quad x = A + R_B \cos v, \quad y = 0, \quad z = R_B \sin v,$$

where u and v run over the reals (see Fig. 3a).

Let us show that the distance $d(C_{B_1}, C_{B_2}) > 2r_B$. Take any points $P_1 \in C_{B_1}$, $P_2 \in C_{B_2}$; let $P_1 = (R_B \cos u, R_B \sin u, 0)$ and $P_2 = (A + R_B \cos v, 0, R_B \sin v)$. We have

$$\begin{aligned} d^2(P_1, P_2) &= (A + R_B \cos v - R_B \cos u)^2 + (R_B \sin u)^2 + (R_B \sin v)^2 = \\ &= A^2 + 2AR_B \cos v - 2AR_B \cos u - 2R_B^2 \cos u \cos v + 2R_B^2 =: F(u, v). \end{aligned}$$

Now

$$\begin{cases} F'_u = 0 \\ F'_v = 0 \end{cases} \iff \begin{cases} (A + R_B \cos v) \sin u = 0 \\ (A - R_B \cos u) \sin v = 0 \end{cases} \iff \begin{cases} \sin u = 0 \\ \sin v = 0 \end{cases}$$

(the last equivalence follows from the assumption $A > R_B + r_B > R_B$). It remains to compute the distances $d(P_1, P_2)$ for 4 pairs of points corresponding to $u, v \in \{0, \pi\}$; the required inequality is easily obtained.

It is also easy to verify that C_{B_1}, C_{B_2} are linked.

Now let $B_1, B_2 \subset \mathbb{R}^3$ be closed r_B -neighborhoods of C_{B_1}, C_{B_2} ; they are disjoint linked standard solid tori congruent to B .

Step 2: bend the link, complete the new link to a regular simple chain. For a straight line $\ell \parallel Oz$ and an angle ψ we denote by $\rho_{\ell, \psi}$ the rotation transformation of \mathbb{R}^3 with axis ℓ on angle ψ ; we rotate in a counterclockwise direction when looking towards the plane Oxy from the half-line Oz_+ (the coordinate system $Oxyz$ is assumed to be a right-handed one).

Let $a_2 \parallel Oz$ and $b_2 \parallel Oy$ be the straight lines passing through Q_{B_2} . For $\psi \in (0, \pi)$ let Q_ψ be the (unique) common point of the lines $\rho_{Oz, \psi}(Oy)$ and $\rho_{a_2, -\psi}(b_2)$, see Fig. 3b.

By continuity, there is a number $\psi_0 \in (0, \pi)$ such that for each $\psi \in (0, \psi_0)$ we have $OQ_\psi > R_B + r_B$, and the set $B_1 \cap \rho_{a_2, -\psi}(B_2) = \rho_{Oz, \psi}(B_1) \cap \rho_{a_2, -\psi}(B_2)$ is empty.

Now let m be any integer such that $m > \frac{\pi}{2\psi_0}$. Fix the angle $\psi := \frac{\pi}{2m} < \psi_0$. Solid tori $T_1 := B_1 = \rho_{Oz, \psi}(B_1)$ and $T_2 := \rho_{a_2, -\psi}(B_2)$ are congruent to B . (These tori depend on ψ , but we do not reflect this in the notation for brevity.) By the above conditions, T_1 and T_2 are disjoint and linked. Their central circles $C_{T_1} = \rho_{Oz, \psi}(C_{B_1})$ and $C_{T_2} = \rho_{a_2, -\psi}(C_{B_2})$, the way we see them from the axis Oz_+ , are shown in Fig. 3c. For each $k = 1, \dots, m-1$ define

$$T_{2k+1} := \rho_{\ell_\psi, -\frac{2\pi k}{m}}(T_1) \quad \text{and} \quad T_{2k+2} := \rho_{\ell_\psi, -\frac{2\pi k}{m}}(T_2),$$

where ℓ_ψ is the straight line passing through Q_ψ parallel to Oz . Let C_ψ be the circle in the Oxy -plane having center Q_ψ and radius OQ_ψ ; it contains the centers of T_1, \dots, T_{2m} . Define a standard solid torus T as a closed R_T -neighborhood of C_ψ for an arbitrary $R_T \in (0, OQ_\psi - (R_B + r_B))$. Now T_1, \dots, T_{2m} is a regular simple chain in T , as desired.

(2) \Rightarrow (3). Let T_1, \dots, T_{2m} be a regular simple chain in T , where each T_i is congruent to B ; let \mathbf{T} be the standard solid torus similar to B with $C_{\mathbf{T}} = C_T$, see Fig. 4; here two bold solid (resp. dashed) arcs denote the boundary of the solid torus T (resp. \mathbf{T}), and the thin dashed arc represents the circle $C_{\mathbf{T}} = C_T$. Assuming that $\sin \frac{\pi}{m} < \frac{r_B}{R_B}$, we will prove that $r_{\mathbf{T}} > R_B + r_B$ (hence $T_i \subset \mathbf{T}$ for each $i = 1, \dots, 2m$).

Since T_1 and T_3 are disjoint, the distance $Q_{T_1}Q_{T_3} = 2R_T \sin \frac{\pi}{m}$ is greater than $\text{diam } T_1 + \text{diam } T_3 = 2(R_B + r_B)$. This together with the assumption




Figure 4: Illustration for the proof of the implication (2) \Rightarrow (3) of Theorem 2.

$\sin \frac{\pi}{m} < \frac{r_B}{R_B}$ implies that

$$R_T > \frac{R_B + r_B}{\sin \frac{\pi}{m}} > \frac{R_B(R_B + r_B)}{r_B}.$$

From the similarity of \mathbf{T} and B , the equality $C_{\mathbf{T}} = C_T$, and the above inequality, we get

$$r_{\mathbf{T}} = \frac{r_B R_{\mathbf{T}}}{R_B} = \frac{r_B R_T}{R_B} > R_B + r_B.$$

□

Proof of Theorem 3. 1) Intersect $T_1, T_3, \dots, T_{2m-1}$ by the plane $\text{Aff } C_T$. We obtain circular rings with centers $Q_{T_1}, Q_{T_3}, \dots, Q_{T_{2m-1}} \in C_T$; the radii of their outer circles equal $s(R_T + r_T)$, see Fig. 5.

Clearly $Q_{T_1}Q_{T_3} > 2s(R_T + r_T)$. On the other side, the length of the closed broken line $Q_{T_1}Q_{T_3} \dots Q_{T_{2m-1}}$ is smaller than the length of the circle C_T , and we get

$$m \cdot 2s(R_T + r_T) < m \cdot Q_{T_1}Q_{T_3} = Q_{T_1}Q_{T_3} + Q_{T_3}Q_{T_5} + \dots + Q_{T_{2m-1}}Q_1 < 2\pi R_T.$$

Hence $ms < \frac{\pi R_T}{R_T + r_T} < \pi$ and $2ms^2 = 2s \cdot ms < 2\pi s$, which proves 1).

2) The inclusion $T_1 \subset T$ implies that $s(R_T + r_T) \leq r_T$, and $s \leq \frac{r_T}{R_T + r_T}$; this inequality was already noticed in [28, p. 603], [27, p. 136], [34, (2.1)]. Now

$$\frac{r_T}{R_T + r_T} < \frac{1}{2\pi} \iff \frac{r_T}{R_T} < \frac{1}{2\pi - 1}.$$




Figure 5: Illustration for the proof of part 1 of Theorem 3.

3) We have

$$2s(R_T + r_T) < Q_{T_1}Q_{T_3} = 2R_T \sin \frac{\pi}{m},$$

consequently

$$s < \frac{R_T \sin \frac{\pi}{m}}{R_T + r_T} < \sin \frac{\pi}{m} < \frac{\pi}{m}.$$

Since $2\pi^2 < 20$, we get $\frac{\pi}{m} < \frac{1}{2\pi}$ for each $m \geq 20$. \square

Proof of Theorem 4. Take any hyperplane $L \subset \mathbb{R}^n$. The projection $p_L : \mathbb{R}^n \rightarrow L$ does not increase distances, hence $m_{n-1}(p_L X) = 0$. By the Szpilrajn theorem [20, Thm. VII 3] we have $\dim p_L X \leq n - 2$.

Now suppose that $\dim p_L X \leq n - 3$; by [32], [33] (for $n = 3$ by [3, 75, p. 87–89]) X is tame, a contradiction. This proves the theorem. \square

References

- [1] L. Antoine, Sur la possibilite d'etendre l'homéomorphie de deux figures a leur voisinages, C.R. Acad. Sci. Paris 171 (1920) 661–663.
- [2] L. Antoine, Sur les ensembles parfaits partout discontinus, C.R. Acad. Sci. Paris 173 (1921) 284–285.

- [3] L. Antoine, Sur l'homéomorphie de deux figures et de leurs voisinages. Thèses de l'entre-deux-guerres, vol. 28, 1921. <http://eudml.org/doc/192716>
- [4] L. Antoine, Sur les voisinages de deux figures homéomorphes, *Fund. Math.* 5 (1924) 265–287.
- [5] Journée Louis Antoine à l'occasion de son 100ème anniversaire, 18 Novembre 1988, Campus de Beaulieu, Rennes, France. (A day in honour of Louis Antoine on the occasion of his 100th birthday, November 18, 1988, Campus of Beaulieu, Rennes, France). (French) Publications de L'Institut de Recherche Mathématique de Rennes. Rennes: IRMAR, Université de Rennes I. pag. non continue (1988).
- [6] S. Armentrout, Decompositions of E^3 with a compact 0-dimensional set of nondegenerate elements, *Trans. Amer. Math. Soc.* 123 (1966) 165–177.
- [7] T. Bang, A solution of the “plank problem”, *Proc. Amer. Math. Soc.* 2 (1951) 990–993.
- [8] S. Barov, J.J. Dijkstra, M. van der Meer, On Cantor sets with shadows of prescribed dimension, *Topology Appl.* 159 (2012) 2736–2742.
- [9] M. Bestvina, R.J. Daverman, G.A. Venema, J.J. Walsh, A 4-dimensional 1-LCC shrinking theorem, *Topology Appl.* 110 (2001) 3–20.
- [10] V.I. Bogachev. *Measure Theory. Volume I.* Springer, 2007.
- [11] K. Borsuk, An example of a simple arc in space whose projection in every plane has interior points, *Fund. Math.* 34 (1947) 272–277.
- [12] J. Cobb, Raising dimension under *all* projections, *Fund. Math.* 144 (1994) 119–128.

- [13] R.J. Daverman, D.A. Venema. *Embeddings in Manifolds*. Graduate Studies in Mathematics 106. Providence, RI: American Mathematical Society. 2009.
- [14] J.J. Dijkstra, J. van Mill, Projections of planar Cantor sets in potential theory, *Indag. Math. (N.S.)* 8 (1997) 173–180.
- [15] R.D. Edwards, Dimension theory, I. *Geom. Topol.*, edited by L.C. Glaser and T.B. Rushing, *Proc. Conf. Park City 1974*, *Lect. Notes Math.* 438 (1975) 195–211.
- [16] O. Frolkina, A Cantor set in \mathbb{R}^d with “large” projections, *Topol. Appl.* 157 (2010) 745–751.
- [17] O. Frolkina, Cantor sets with high-dimensional projections, *Topol. Appl.* 275 (2020) 107020. <https://doi.org/10.1016/j.topol.2019.107020>
- [18] O. Frolkina, All projections of a typical Cantor set are Cantor sets, *Topol. Appl.* (2020) 107192. <https://doi.org/10.1016/j.topol.2020.107192>
- [19] D. Garity, D. Repovš, M. Željko, Uncountably many inequivalent Lipschitz homogeneous Cantor sets in \mathbb{R}^3 , *Pacif. J. Math.* 222 (2005) 287–299.
- [20] W. Hurewicz, H. Wallman. *Dimension theory*. Princeton University Press. 1941.
- [21] J.E. Hutchinson, Fractals and self similarity, *Indiana Univ. Math. J.* 30 (1981) 713–747.
- [22] H. Ibisch, L’œuvre de Louis Antoine et son influence. *Publications de l’Institut de recherche mathématiques de Rennes*, 1988, fascicule S6 “Journée Louis Antoine”, exp. N 5, 1–35.

- [23] H. Ibisch, L'œuvre de Louis Antoine et son influence. (The mathematical work of Louis Antoine and its influence). (French) Expo. Math. 9 (1991) 251–274.
- [24] J. Malešič, D. Repovš, On characterization of Lipschitz manifolds. In: Szenthe, J. (ed.), New developments in differential geometry, Budapest 1996. Proceedings of the conference, Budapest, Hungary, July 27-30, 1996. Dordrecht: Kluwer Academic Publishers. 265-277 (1999).
- [25] A. McFarland, J. McFarland, and J.T. Smith, editors. Alfred Tarski. Early work in Poland — geometry and teaching. Birkhäuser/Springer, New York, 2014. With a bibliographic supplement, Foreword by Ivor Grattan-Guinness.
- [26] E.E. Moise. Geometric Topology in Dimensions 2 and 3. Springer-Verlag, 1977.
- [27] D. Rubinstein, B. Soper, Self-similar constructions of the rigid Antoine necklace in Euclidean three-space. In: Proceedings of the Research Experiences for Undergraduates. Program in Mathematics. Oregon State University. Summer 2001. Dennis J. Garity, Director. 128–154.
- [28] T.B. Rushing, Hausdorff dimension of wild fractals, Trans. Amer. Math. Soc. 334 (1992) 597–613.
- [29] R.B. Sher, Concerning wild Cantor sets in E^3 , Proc. Amer. Math. Soc. 19 (1968) 1195–1200.
- [30] M. A. Shtan'ko, The embedding of compacta in Euclidean space, Math. USSR-Sb. 12:2 (1970) 234–254. (Transl. from: Mat. Sb. 83 (125) (1970) 2 (10) 234–255.)
- [31] J. Väisälä, Dimension and measure, Proc. Amer. Math. Soc. 76 (1979) 167–168.
- [32] D.G. Wright, Geometric taming of compacta in E^n , Proc. Amer. Math. Soc. 86 (1982) 641–645.

- [33] J.J. Walsh, D.G. Wright, Taming compacta in E^4 , Proc. Amer. Math. Soc. 86 (1982) 646–648.
- [34] M. Željko, Minimal number of tori in geometric self-similar Antoine Cantor sets, JP J. Geom. Topol. 5 (2005) 109–113.