

RECOVERING AFFINE-LINEARITY OF FUNCTIONS FROM THEIR RESTRICTIONS TO AFFINE LINES

APOORVA KHARE AND AKAKI TIKARADZE

ABSTRACT. Motivated by recent results of Tao–Ziegler [*Discrete Anal.* 2016] and Greenfeld–Tao (2022 preprint) on concatenating affine-linear functions along subgroups of an abelian group, we show three results on recovering affine-linearity of functions $f : V \rightarrow W$ from their restrictions to affine lines, where V, W are \mathbb{F} -vector spaces and $\dim V \geq 2$. First, if $\dim V < |\mathbb{F}|$ and $f : V \rightarrow \mathbb{F}$ is affine-linear when restricted to affine lines parallel to a basis and to certain “generic” lines through 0, then f is affine-linear on V . (This extends to all modules M over unital commutative rings R with large enough characteristic.) Second, we explain how a classical result attributed to von Staudt (1850s) extends beyond bijections: if $f : V \rightarrow W$ preserves affine lines ℓ , and if $f(v) \notin f(\ell)$ whenever $v \notin \ell$, then this also suffices to recover affine-linearity on V , but up to a field automorphism. In particular, if \mathbb{F} is a prime field $\mathbb{Z}/p\mathbb{Z}$ ($p > 2$) or \mathbb{Q} , or a completion \mathbb{Q}_p or \mathbb{R} , then f is affine-linear on V .

We then quantitatively refine our first result above, via a weak multiplicative variant of the additive B_h -sets initially explored by Singer [*Trans. Amer. Math. Soc.* 1938], Erdős–Turán [*J. London Math. Soc.* 1941], and Bose–Chowla [*Comment. Math. Helv.* 1962]. Weak multiplicative B_h -sets occur inside all rings with large enough characteristic, and in all infinite or large enough finite integral domains/fields. We show that if R is among any of these classes of rings, and $M = R^n$ for some $n \geq 3$, then one requires affine-linearity on at least $\binom{n}{\lfloor n/2 \rfloor}$ -many generic lines to deduce the global affine-linearity of f on R^n . Moreover, this bound is sharp.

1. INTRODUCTION AND MAIN RESULTS

The goal of this short note is to elucidate some classification results for functions which preserve affine lines in a vector space. We were motivated to work on these by a recent blogpost of Tao [14], where he discusses a preprint with Greenfeld [6]. Specifically, Tao writes the following result:

Theorem 1.1 ([14]). *Let $F : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a smooth function which is affine-linear on every horizontal line, diagonal (line of slope 1), and anti-diagonal (line of slope -1). In other words, for any $c \in \mathbb{R}$, the functions*

$$x \mapsto F(x, c), \quad x \mapsto F(x, c + x), \quad x \mapsto F(x, c - x), \quad (1.1)$$

are each affine functions on \mathbb{R} . Then F is an affine function on \mathbb{R}^2 .

In fact, (a) the smoothness hypothesis is not necessary; and (b) this result also holds when \mathbb{R} is replaced by a finite field $\mathbb{Z}/p\mathbb{Z}$ with p odd.

This result was motivated by the preprint [6] – see in it the discussion after the proof of Lemma 9.2. Here the authors say that a certain function $F : G^2 \rightarrow G$, for G a finite cyclic group, becomes “mostly affine” along horizontal lines, diagonals, and anti-diagonals – and then one can expect to “concatenate” this information in the spirit of [15, Proposition 1.2] to conclude that F is affine: $F(x, y) = Ax + By + C$. However, the authors show that when one works over $\mathbb{Z}/2^n\mathbb{Z}$ for large $n \gg 0$, there is a quadratic correction. (This discussion and the usage of concatenation go towards providing a counterexample – and more – to the periodic tiling conjecture.)

Date: December 12, 2022.

2010 Mathematics Subject Classification. 15A03 (primary), 13C10 (secondary).

Key words and phrases. Affine linear maps, concatenation, field automorphism, Sidon set, B_h set.

We begin this note by explaining how the hypotheses in Theorem 1.1 can be further weakened, thereby obtaining a slightly stronger concatenation-type result. Namely, we claim that affine-linearity on *all* horizontal lines is not needed, just on the X -axis. To formulate (and show) this claim in greater generality, first observe that upon working over $\mathbb{F} = \mathbb{R}$ or $\mathbb{Z}/p\mathbb{Z}$ for p odd, and setting $e_i = 2^{-1}(1, (-1)^i)$ for $i = 1, 2$, the vectors e_1, e_2 form a basis of \mathbb{F}^2 . Now $(1, 0) = e_1 + e_2$, e_1, e_2 are the directions in (1.1), and the claim is that along with affine-linearity along every diagonal $m_0 + \mathbb{F}e_2$ and anti-diagonal $m_0 + \mathbb{F}e_1$ (i.e. for all $m_0 \in \mathbb{F}^2$), one only needs to assume F is affine-linear along the single horizontal line $\{(x, 0) = x(e_1 + e_2) : x \in \mathbb{F}\}$.

Remark 1.2. In fact, $e_1 + e_2$ is not special: one can use the single additional line $\mathbb{F}(c_1e_1 + c_2e_2)$ for *any* choice of scalars $c_1, c_2 \in \mathbb{F}^\times$. Moreover, Equation (2.1) below reveals that (a) the quadratic correction encountered in [6] by Greenfeld–Tao is a special case of a multi-affine correction of degree n , when working over \mathbb{F}^n ; and (b) in fact such a correction occurs over every unital ring R – not just over \mathbb{F} , and also not only over $\mathbb{Z}/2^n\mathbb{Z}$ as in [6]. We will also show that this correction vanishes if R has “large enough characteristic” – e.g. over R^2 , we need $1 + 1$ to not be a zerodivisor in R .

As in Remark 1.2, we now extend the above formulation of our strengthening of Theorem 1.1 from vector spaces \mathbb{F}^2 (as above) to all modules M , over all unital commutative rings with “large enough characteristic”, and where the scalars c_j need not be units but merely non-zerodivisors:

Theorem 1.3. *Suppose R is a unital commutative ring, and M is a free R -module with basis $\{e_i : i \in I\}$. Further assume that $n = 1 + \dots + 1 \in R$ is not a zerodivisor whenever $1 \leq n \leq |I|$.*

For every finite subset $J \subseteq I$, fix vectors $m_J := \sum_{j \in J} c_j^J e_j \in M$, where every $c_j^J \in R$ is a non-zerodivisor. Now suppose $f : M \rightarrow R$ is any map such that the restrictions of f to the lines

$$m_0 + Re_i = \{m_0 + re_i : r \in R\}, \quad \forall m_0 \in M, i \in I$$

$$\text{and } Rm_J = \{rm_J : r \in R\}, \quad \forall J \subseteq I \text{ of finite size } \geq 2$$

are each affine-linear. Then f is affine-linear on M .

This also holds if R is an integral domain that is infinite or else finite with $\dim M < |R|$.

As seen below, the proof of Theorem 1.3 works for arbitrary R -modules – and so in fact:

Corollary 1.4. *In the case of R with each $n \leq |I|$ not a zerodivisor, Theorem 1.3 holds for all R -modules M , with “basis” replaced by “generating set”.*

Note, the $|I| = 2$ case of Theorem 1.3 (for free modules) is already a twofold strengthening of Theorem 1.1, in that Theorem 1.1 (a) uses more “horizontal lines” (i.e. parallel to $e_1 + e_2$), and (b) is the special case with $R = \mathbb{R}$ or $\mathbb{Z}/p\mathbb{Z}$ (with $p > 2$) and specific choices of M, I, e_i, c_i^J . (And the condition that 2 is a non-zerodivisor in the case of R^2 fits in with Greenfeld–Tao’s use of $R = \mathbb{R}$ or $\mathbb{Z}/p\mathbb{Z}$ for p odd.) Also recall – for concreteness – the notion of affine-linear maps on modules:

Definition 1.5. Given a unital commutative ring R , two R -modules $N \leq M$, and a vector $m_0 \in M$, an R -valued function f on the “affine submodule” $\ell := m_0 + N \subseteq M$ is *affine-linear* if for all shifts $m_1 \in \ell$, the map $f_{m_1} : N \rightarrow R; n \mapsto f(n + m_1) - f(m_1)$ is an R -module map, i.e., R -linear.

The special case used throughout this paper is that of $N = Re$ a line. For completeness, we recall that affine lines are also known in the literature as *1-flats*.

We now turn to a quantitative sharpening of Theorem 1.3. Suppose one works over R^n . Theorem 1.1 asserted (without any claims of optimality, of course) that for $R = \mathbb{R}$ or $\mathbb{Z}/p\mathbb{Z}$, in addition to all directions parallel to a basis we require $|R|$ many other lines parallel to the X -axis, in order to obtain global linearity. This test set was reduced in Theorem 1.3 to a single line through the origin – or one along each of $2^n - (n + 1)$ directions $\{m_J : 2 \leq |J| \leq n\}$ for general $n \geq 2$.

It is natural to seek the *minimum* number of such “test directions” m_J that ensure the global affine-linearity of f . For $n = 2$, a single line suffices by Theorem 1.3. For $n \geq 3$, the following strict refinement of Theorem 1.3 shows that the minimum number is the central binomial coefficient:

Theorem 1.6. *Suppose R is an integral domain that is infinite, and $M = R^n$ has basis e_1, \dots, e_n for some integer $n \geq 3$. Then there exist $N := \binom{n}{\lfloor n/2 \rfloor}$ -many directions $\mathbf{v}_1, \dots, \mathbf{v}_N \in R^n$ such that for any map $f : R^n \rightarrow R$, if the restrictions of f to the lines*

$$m_0 + Re_i, \quad \forall m_0 \in R^n, \quad 1 \leq i \leq n, \quad \text{and} \quad R\mathbf{v}_1, \dots, R\mathbf{v}_N$$

are each affine-linear, then f is affine-linear on R^n . Fewer than N directions do not suffice.

The same assertions hold if R is a finite integral domain of sufficiently large cardinality.

To get a sense of how much Theorem 1.6 improves for $M = R^n$ the estimate of $2^n - (n + 1)$ directions m_j in Theorem 1.3, we note via the Wallis product expansion for $\frac{\pi}{2}$, the asymptotics $\binom{n}{\lfloor n/2 \rfloor} \sim \frac{2^{n+\frac{1}{2}}}{\sqrt{\pi n}}$. Also note that the estimate of N here for $n \geq 3$ does not work for $n = 2$, since it gives $\binom{2}{1} = 2$ while the minimum number is 1 by Theorem 1.3.

In Section 3, we strengthen Theorem 1.6 by showing that its sharp bound of $\binom{n}{\lfloor n/2 \rfloor}$ holds over a larger class of rings – which also includes rings with large enough characteristic. The key novelty involves working with weak multiplicative B_h -sets (or Sidon sets). Recall that classically, (additive) Sidon sets have been studied since Erdős–Turán and Bose–Chowla, and even earlier.

We next elucidate a result along similar “lines”. Begin by noting that affine-linear maps $: V \rightarrow \mathbb{F}$ – as above – take affine lines in V to the (affine) line \mathbb{F} , obviously. We now study maps $f : V \rightarrow W$ (for general W) which satisfy this property – but which need not be affine-linear when restricted to any affine line. In this case, one “almost” recovers affine-linearity, but up to a field automorphism:

Definition 1.7. Suppose V, W are vector spaces over a field \mathbb{F} , and $\tau : \mathbb{F} \rightarrow \mathbb{F}$ is a field automorphism. A map $f : V \rightarrow W$ is τ -linear if f is additive and $f(\lambda v) = \tau(\lambda)f(v)$ for all $\lambda \in \mathbb{F}, v \in V$. We say f is τ -affine linear if f is the composite of a τ -linear map and a translation in W .

Theorem 1.8. *Let V, W be vector spaces over a field $\mathbb{F} \neq \mathbb{Z}/2\mathbb{Z}$ with $\dim V \geq 2$, and suppose $f : V \rightarrow W$ is any map that takes affine lines ℓ onto affine lines, such that $f(v) \notin f(\ell)$ whenever $v \in V$ is not in the affine line ℓ . Then f is injective and τ -affine linear for some field automorphism τ of \mathbb{F} . The converse is straightforward.*

As an immediate consequence, for the two fields mentioned by Tao in Theorem 1.1 – in fact for any field with a trivial automorphism group – one recovers affine-linearity on the nose:

Corollary 1.9. *Setup as in Theorem 1.8. If $\mathbb{F} = \mathbb{R}$ or $\mathbb{Z}/p\mathbb{Z}$ for p odd, then f is affine-linear on V . The same holds if $\mathbb{F} = \mathbb{Q}$ or \mathbb{Q}_p for $p > 0$ a prime.*

We end with a historical remark. After we showed Theorem 1.8, we learned that variants of it had previously appeared in the literature, including in the foundational 1850s texts by von Staudt [16], and in later books by Hartshorne [7, Proposition 3.11] and by Snapper–Troyer [13, Proposition 69.1] (see also a variant in [8]). However, *all* of these variants also assumed that $f : V \rightarrow W$ is a bijection (and $\dim W \geq 2$). As Theorem 1.8 uses weaker hypotheses, our proof necessarily differs in places from the earlier ones; thus we will elaborate on some of the steps but only sketch some others.

2. TWO OF THE PROOFS

In this section we show Theorems 1.3 and 1.8. The proof of Theorem 1.3 uses a calculation twice, so we isolate it into a lemma.

Lemma 2.1. *Suppose R is a unital commutative ring, and M an R -module. If $f : M \rightarrow R$ is affine-linear along some line $\ell := m_0 + Re$ (with $m_0, e \in M$), then there exists $a \in R$ such that*

$$f(m_1 + re) = f(m_1) + ar, \quad \forall m_1 \in \ell, r \in R.$$

Proof. Note that $f : \ell \rightarrow R$ is affine-linear if and only if for each $m_1 \in \ell$ the map $\varphi : re \mapsto f(m_1 + re) - f(m_1)$ is R -linear. This is equivalent to a choice of scalar $a^{(m_1)} = \varphi(1)$, via:

$$f(m_1 + re) - f(m_1) = a^{(m_1)}r, \quad \forall r \in R.$$

Setting $r = 1$ yields: $a^{(m_1)} = f(m_1 + e) - f(m_1)$. It remains to show that $a^{(m_1)}$ is independent of the choice of $m_1 \in \ell$. Given $m_2 = m_1 + re$, say, evaluate $f(m_2 + e)$ in two ways:

$$\begin{aligned} f(m_2 + e) &= f(m_2) + a^{(m_2)} = f(m_1 + re) + a^{(m_2)} = f(m_1) + a^{(m_1)}r + a^{(m_2)} \\ &= f(m_1 + (r+1)e) = f(m_1) + a^{(m_1)}(r+1). \end{aligned}$$

Thus $a^{(m_2)} = a^{(m_1)}$ and the proof is complete. \square

Proof of Theorem 1.3. We begin by computing $f(m'_0)$ for arbitrary $m'_0 \in M$. Since $\{e_i : i \in I\}$ generates M , we claim for every finite subset $J' \subseteq I$ – and choice of scalars $x_j \in R$ – that

$$\begin{aligned} f\left(m_0 + \sum_{j \in J'} x_j e_j\right) &= f(m_0) + \sum_{\emptyset \neq J \subseteq J'} \Psi_J^{(m_0)} \prod_{j \in J} x_j, \\ \text{where } \Psi_J^{(m_0)} &:= \sum_{K \subseteq J} (-1)^{|J|-|K|} f(m_0 + \sum_{k \in K} e_k). \end{aligned} \quad (2.1)$$

(This is a multi-affine polynomial in the x_j .) Interestingly, we need to use only the $m_0 = 0$ special case of (2.1) to show Theorem 1.3, but to prove this special case by induction – as is now done – we need to use (2.1) for certain nonzero $m_0 \in M$. Before proving (2.1), we write down its first two cases so that they may help to see the general case better. If $J' = \{1\}$ then $f(m_0 + x_1 e_1) = f(m_0) + (f(m_0 + e_1) - f(m_0))x_1$. Next, if $J' = \{1, 2\}$, then

$$\begin{aligned} &f(m_0 + x_1 e_1 + x_2 e_2) \\ &= f(m_0) + (f(m_0 + e_1) - f(m_0))x_1 + (f(m_0 + e_2) - f(m_0))x_2 \\ &\quad + (f(m_0 + e_1 + e_2) - f(m_0 + e_1) - f(m_0 + e_2) + f(m_0))x_1 x_2. \end{aligned}$$

We now prove (2.1), by induction on $|J'|$, with the result a tautology if J' is empty. For $J' = \{j\}$ a singleton, the result is immediate from Lemma 2.1 (and its proof) applied to the line $m_0 + R e_j$.

For the induction step, let $J' = \{1, 2, \dots, k\}$ for some $k \geq 2$. Start by working with $m'_0 := m_0 + \sum_{j=1}^{k-1} x_j e_j$, and compute using (the proof of) Lemma 2.1:

$$f\left(m_0 + \sum_{j=1}^k x_j e_j\right) = f(m'_0 + x_k e_k) = (f(m'_0 + e_k) - f(m'_0))x_k + f(m'_0).$$

By the induction hypothesis, the final term on the right-hand side equals

$$f\left(m_0 + \sum_{j=1}^{k-1} x_j e_j\right) = f(m_0) + \sum_{\emptyset \neq J \subseteq \{1, \dots, k-1\}} \Psi_J^{(m_0)} \prod_{j \in J} x_j, \quad (2.2)$$

while the remaining difference on the right-hand side involves evaluating f at

$$m'_0 = m_0 + \sum_{j=1}^{k-1} x_j e_j \quad \text{and} \quad m'_0 + e_k = (m_0 + e_k) + \sum_{j=1}^{k-1} x_j e_j.$$

Apply the induction hypothesis to both of these arguments. This yields:

$$f\left(m_0 + e_k + \sum_{j=1}^{k-1} x_j e_j\right) x_k - f\left(m_0 + \sum_{j=1}^{k-1} x_j e_j\right) x_k$$

$$\begin{aligned}
 &= x_k \sum_{\emptyset \neq J \subseteq \{1, \dots, k-1\}} \left(\Psi_J^{(m_0+e_k)} \prod_{j \in J} x_j - \Psi_J^{(m_0)} \prod_{j \in J} x_j \right) \\
 &= \sum_{k \in J \subseteq \{1, \dots, k\}} \left(\Psi_{J \setminus \{k\}}^{(m_0+e_k)} - \Psi_{J \setminus \{k\}}^{(m_0)} \right) \prod_{j \in J} x_j,
 \end{aligned}$$

and the difference of the Ψ -values in the summand is precisely $\Psi_J^{(m_0)}$. Adding this to (2.2) proves the induction step and hence (2.1). Notice (for the purposes of the next section), this part of the proof works in any unital commutative ring.

Having shown (2.1), we return to the proof of the theorem. For $J' \subseteq I$ finite with $|J'| \geq 2$, use the hypotheses to compute $f(rm_{J'})$ in two ways – via Lemma 2.1, and via (2.1) with $m_0 = 0$:

$$\begin{aligned}
 f(rm_{J'}) &= f(0) + (f(m_{J'}) - f(0))r \\
 &= f(0) + \sum_{\emptyset \neq J \subseteq J'} \Psi_J^{(0)} r^{|J|} \prod_{j \in J} c_j^J, \quad \forall r \in R.
 \end{aligned}$$

The second equality here is an equality of polynomial functions (in one variable) of degree n , say, where $n \leq |J'| \leq |I|$. Subtracting around this second equality yields an equality of the form

$$a_1 r + a_2 r^2 + \dots + a_n r^n = 0, \quad \forall r \in R \quad (2.3)$$

and we now claim that $a_1 = a_2 = \dots = a_n = 0$. Indeed, evaluating (2.3) at the elements $1, 2 = 1 + 1, \dots, n = 1 + \dots + 1$ in R yields the system of equations

$$A \cdot \begin{pmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{pmatrix} = 0, \quad \text{where } A = \begin{pmatrix} 1 & 1^2 & \dots & 1^n \\ 2 & 2^2 & \dots & 2^n \\ \vdots & \vdots & \ddots & \vdots \\ n & n^2 & \dots & n^n \end{pmatrix}$$

is “essentially” a Vandermonde matrix. Pre-multiplying by the adjugate of A yields: $\det(A) \cdot a_i = 0 \forall i$. But $\det(A) = \prod_{i=1}^n i!$, which is a non-zero-divisor in R by assumption, and so $a_1 = \dots = a_n = 0$ as desired. This holds for all finite subsets $J' \subseteq I$.

Now let J' have size 2. Then a_2 is the only “higher degree” (> 1) term in (2.3), and so $a_2 = \Psi_{J'}^{(0)} \prod_{j \in J'} c_j^{J'} = 0$. Since each $c_j^{J'}$ is a non-zero-divisor, $\Psi_{J'}^{(0)} = 0$ for all $J' \subseteq I$ of size 2.

Next, let J' have size 3. By the preceding paragraph, $a_2 = 0$, so a_3 is the only nonzero higher degree term in (2.3), and so the same analysis implies $\Psi_{J'}^{(0)} = 0$ for all $J' \subseteq I$ of size 3.

Continuing inductively, $\Psi_{J'}^{(0)} = 0$ whenever $J' \subseteq I$ has size at least 2. Using (2.1) with $m_0 = 0$,

$$f\left(\sum_{i \in I} x_i e_i\right) = f(0) + \sum_{i \in I} (f(e_i) - f(0))x_i,$$

where all but finitely many coefficients $x_i \in R$ are zero, and the rest are arbitrary. Hence f is affine-linear on M , as claimed. \square

Remark 2.2. By the lines after (2.3), the assumption of $n \in R$ not being a zero-divisor if $1 \leq n \leq |I|$ – or equivalently, if n is moreover prime in \mathbb{Z} – may be replaced by requiring the existence of non-zero-divisors r_1, \dots, r_n such that $r_i - r_j$ is also a non-zero-divisor for $i \neq j$, for each $1 \leq n \leq |I|$. This shows the final line of Theorem 1.3.

Remark 2.3. A related result, alluded to in the previous section, was shown by Tao and Ziegler in [15]. Namely, the authors first define polynomials on additive/abelian groups G as follows: the only degree < 0 polynomial is the zero map; the degree < 1 polynomials along a subgroup $H \leq G$ are the constant maps; and affine-linear maps along H indeed turn out to be polynomials of degree

< 2 . Now Proposition 1.2 in *loc. cit.* says that if P is a polynomial of degree $< d_i$ along a subgroup $H_i \leq G$ for $i \in I = \{1, 2\}$, then it is a polynomial of degree $< d_1 + d_2 - 1$ along $H_1 + H_2$. In our situation (specialized as above to $M = G = \mathbb{F}^2$ over $R = \mathbb{F}$), with $H_i = \mathbb{F}e_i$ for $i = 1, 2$, we would obtain polynomials of degree < 3 along $H_1 + H_2$. This is precisely what happens in the above proof, see (2.1) e.g. for $|J'| = 2$ (and inductively for larger J') – whereby we obtain a multi-affine polynomial. Now the extra information along the radial lines $Rm_{J'}$ (and *not* requiring their translates, cf. Theorem 1.1) removes all higher degree monomials.

Now we show Theorem 1.8; as stated in it, we assume henceforth that f takes lines onto lines.

Proof of Theorem 1.8. (Below, Greek letters except τ denote scalars in \mathbb{F} .) We start with three initial observations, both using $\dim V \geq 2$. First, $\dim W \geq 2$, because one can take any line $\ell = \mathbb{F}w \subseteq V$ and a vector $v \in V \setminus \ell$, so that $f(v) \in W \setminus f(\ell)$. Second, the hypotheses imply f is one-to-one (but not necessarily onto). Indeed, if not – say if $f(x) = f(y)$ for $x \neq y \in V$ – then choose $v \in V$ not on the line joining x, y . Now x is not in the line ℓ joining v, y , so $f(y) = f(x) \notin f(\ell)$, a contradiction. Third, we may replace f by $f(\cdot) - f(0)$, thereby assuming that $f(0) = 0$ henceforth.

Step 1. We claim that f preserves planes through the origin. More precisely, if v, w are linearly independent in V then so are $f(v)$ and $f(w)$, and $f(\text{span}(v, w)) = \text{span}(f(v), f(w))$.

Proof. Since f is one-to-one, $f(v), f(w) \neq 0 = f(0)$. Now since $f(w) \notin f(\mathbb{F}v) = \mathbb{F}f(v)$ (the line through $f(v)$ and $f(0) = 0$), the vectors $f(v), f(w)$ are linearly independent. Consider $\lambda v + \mu w$ for $\lambda, \mu \in \mathbb{F}$; if $\lambda \cdot \mu = 0$, this is on one of the two “coordinate axes”, hence in the span of $f(v), f(w)$.

Otherwise $\lambda, \mu \in \mathbb{F}^\times$. Choose $\lambda' \neq 0, \lambda$ in \mathbb{F} (since $\mathbb{F} \neq \mathbb{Z}/2\mathbb{Z}$); then $\lambda v + \mu w$ is on the line containing $\lambda'v$ and $\frac{\mu\lambda'}{\lambda' - \lambda}w$ (use the coefficients $t = \lambda/\lambda'$ and $1 - t$). Thus, $f(\lambda v + \mu w)$ is on the line through the two points $f(\lambda'v)$ and $f(\frac{\mu\lambda'}{\lambda' - \lambda}w)$, hence in their span. But $f(\lambda'v) \in f(\mathbb{F} \cdot v) = \mathbb{F} \cdot f(v)$ (by hypothesis), and similarly for w , so $f(\lambda v + \mu w) \in \text{span}(f(v), f(w))$. Together with the preceding paragraph, this implies $f(\text{span}(v, w)) \subseteq \text{span}(f(v), f(w))$.

For the reverse inclusion, since $f(v), f(w)$ are a basis for their span, any linear combination $\lambda f(v) + \mu f(w)$ is either in the image of $f(\mathbb{F}v)$ or $f(\mathbb{F}w)$ when $\lambda \cdot \mu = 0$, else repeating the above calculation, $\lambda f(v) + \mu f(w)$ lies on the line through $\lambda'f(v)$ and $\frac{\mu\lambda'}{\lambda' - \lambda}f(w)$, with $\lambda, \mu, \lambda' \in \mathbb{F}^\times$. By hypothesis, $\lambda'f(v) = f(\lambda''v)$ for some $\lambda'' \in \mathbb{F}^\times$, and similarly the other term equals $f(\mu''w)$ for some $\mu'' \in \mathbb{F}^\times$. But then $\lambda f(v) + \mu f(w) \in f(\ell)$, where the affine line $\ell \subseteq \text{span}(v, w)$ passes through the linearly independent vectors $\lambda''v, \mu''w$. Hence $\lambda f(v) + \mu f(w) \in f(\text{span}(v, w)) \forall \lambda, \mu \in \mathbb{F}$. \square

Step 2. We claim f is additive on “linearly independent vectors” (so it suffices to study f on lines).

Sketch of proof. One first shows that f preserves the notion of parallel lines in a plane through the origin (via the preceding step). Next, taking intersections, one shows that f takes the sets of vertices of a parallelogram in a plane containing the origin, again to such a set (via the preceding step), while also preserving “non-adjacency”. Now if v, w are linearly independent, then $0, v, v + w, w$ are four such vertices, hence so are $0 = f(0), f(v), f(v + w), f(w)$. But then $f(v + w) + 0 = f(v) + f(w)$. \square

Step 3. Suppose v, w are linearly independent vectors in V , and $f(\lambda v) = p(\lambda)f(v)$ and $f(\lambda w) = q(\lambda)f(w)$ for all $\lambda \in \mathbb{F}$, where $p, q : \mathbb{F} \rightarrow \mathbb{F}$ are bijections that each fix $0, 1$. Then $p \equiv q$ on \mathbb{F} , and this common bijection – say denoted by τ – is multiplicative.

Sketch of proof. Given $\lambda \in \mathbb{F}$, from above say $f(\lambda(v + w)) = \mu f(v + w) = \mu f(v) + \mu f(w)$ for some $\mu \in \mathbb{F}$. But this also equals $f(\lambda v) + f(\lambda w) = p(\lambda)f(v) + q(\lambda)f(w)$. Hence $p \equiv q$ on \mathbb{F} .

Denote this common bijection by $\tau : \mathbb{F} \rightarrow \mathbb{F}$. Now given $\lambda, \mu \in \mathbb{F}$, compute $f(\mu(\lambda v + w))$ in two ways, to obtain $\tau(\mu)(\tau(\lambda)f(v) + f(w))$ and $\tau(\mu\lambda)f(v) + \tau(\mu)f(w)$. Hence $\tau(\mu\lambda) = \tau(\mu)\tau(\lambda)$. \square

To summarize: $f(\lambda v + w) = \tau(\lambda)f(v) + f(w)$ for linearly independent v, w in V and all scalars $\lambda \in \mathbb{F}$, where τ is a multiplicative bijection on \mathbb{F} that fixes $0, 1$. The final assertion is that τ is also additive: $\tau(\nu_1 + \nu_2) = \tau(\nu_1) + \tau(\nu_2) \forall \nu_1, \nu_2 \in \mathbb{F}$. We include a short proof for completeness.

There are two cases. First, the above steps imply $\tau(-1)^2 = \tau(1) = 1$, so (irrespective of whether or not $\text{char}(\mathbb{F}) = 2$), $\tau(-1) = -1$ since τ is a bijection. Rescaling, $\tau(-\nu) + \tau(\nu) = 0 = \tau(-\nu + \nu)$.

If instead $\nu_1 + \nu_2 \neq 0$, then recalling that the line through two linearly independent points v, w is parametrized as $\{\lambda v + (1 - \lambda)w : \lambda \in \mathbb{F}\}$, we evaluate f at any point on this line:

$$f(\lambda v + (1 - \lambda)w) = f(\lambda v) + f((1 - \lambda)w) = \tau(\lambda)f(v) + \tau(1 - \lambda)f(w).$$

By assumption, this lies on the line between (the distinct points) $f(v)$ and $f(w)$; hence the coefficients add up to 1, i.e., $\tau(\lambda) + \tau(1 - \lambda) = 1 = \tau(1)$. Choosing $\lambda = \frac{\nu_1}{\nu_1 + \nu_2}$ yields

$$\tau\left(\frac{\nu_1}{\nu_1 + \nu_2}\right) + \tau\left(\frac{\nu_2}{\nu_1 + \nu_2}\right) = \tau(1).$$

This implies τ is additive, upon multiplying both sides by $\tau(\nu_1 + \nu_2) \neq \tau(0) = 0$. □

3. QUANTITATIVE SHARPENING VIA WEAK MULTIPLICATIVE B_h -SETS

We end by quantitatively sharpening Theorem 1.3 via Theorem 1.6. In fact we show here the latter result, under weaker hypotheses. This requires the following notion.

Definition 3.1. Given an integer $h \geq 1$, a finite subset S of a unital commutative ring R is said to be a *weak multiplicative B_h -set* if the product map $\Pi : \binom{S}{h} \rightarrow R$, sending each h -element subset of S to their product, is injective. Here and below, $\binom{S}{h}$ consists of all h -element subsets of S .

Notice that replacing “product” by “sum” in the above definition recovers a classical notion in additive combinatorics: that of a *B_h -set / Sidon set*, provided one *further* allows repeated elements – i.e., if the domain of definition for the sum map Σ is expanded from $\binom{S}{h}$ to S^h . Such sets have been studied previously (mostly for $h = 2$, but also otherwise), including by Singer [12], Erdős–Turán [5], Bose–Chowla [1], and in later works by Lindström [9] and Cilleruelo [2] among others. (See also the references in these works.)¹

The above multiplicative notion has been studied before – see e.g. [4, 10, 11] – but not as well as the additive version. We now list some examples of unital commutative rings in which weak multiplicative B_h -sets exist, and show that each ring satisfies two key properties that we will use to quantitatively refine Theorem 1.3:

- (1) The existence of an n -element subset S that is simultaneously a weak multiplicative B_h -set for $1 \leq h \leq n$, where $n \geq 3$.
- (2) We also require the set S to satisfy a strengthening of the property mentioned in Remark 2.2: for all integers $1 < h < n$ and all subsets $J \neq J' \in \binom{S}{h}$, the difference $\prod_{s \in J} s - \prod_{s' \in J'} s'$ is a non-zero-divisor in R . (In particular for $h = 2$, using $ss' - ss'' = s(s' - s'')$ for distinct $s, s', s'' \in S$ implies that each $s \in S$ is a non-zero-divisor.)

We now mention three classes of rings which satisfy both of these properties, starting with both settings listed in Theorem 1.6.

Example 3.2. Suppose $n \geq 3$ and the R contains a cyclic semigroup of size at least 2^{n-1} , say with generator $g \in R$, such that $g^k - 1, g$ are non-zero-divisors for all $1 \leq k \leq 2^{n-1} - 1$. Then using binary arithmetic, it follows that the subset

$$\{1; g, g^2, \dots, g^{2^{n-2}}\} \subseteq \langle g \rangle$$

¹For completeness we mention the related notion of an abelian group G – mostly studied for $G = \mathbb{Z}$ again – containing a set with “discrete subset sums”, in which case one would like the sum map Σ to be one-to-one on the union of the domains. That is, $\Sigma : 2^S \setminus \{\emptyset\} \rightarrow G$ is injective. (Bounds on the sizes of) such sets were studied by Erdős–Moser, Conway, Guy, Elkies, Bohman, and others – see e.g. [3] for more references and follow-ups. This notion is strictly more restrictive than that of being individually or simultaneously a B_h -set for various h , which is the notion of interest in the present work.

is an n -element subset that satisfies both properties above. (Notice that the multiplicative B_h -set property on $\langle g \rangle$ is essentially the same as the additive version, via exponentiating, and so one can use the sharper bounds in the literature to reduce the size of the cyclic semigroup $\langle g \rangle$.)

In particular, all finite integral domains (i.e., finite fields) of size $\geq 2^{n-1} + 1$ satisfy both properties above (the second property holds since $2^{n-1} \geq n$ for $n \geq 3$).

Example 3.3. Suppose R is an infinite integral domain. Then R satisfies both properties above for every $n \geq 3$. To see why, let \mathbb{F} be the quotient field of R , denote $S := \{1, \dots, n\}$, and consider the homogeneous polynomial

$$p(x_1, \dots, x_n) := \prod_{h=1}^{n-1} \prod_{J \neq J' \in \binom{S}{h}} \left(\prod_{j \in J} x_j - \prod_{j' \in J'} x_{j'} \right) \in \mathbb{F}[x_1, \dots, x_n].$$

This is a product of nonzero polynomials, so its zero locus is not all of \mathbb{F}^n because \mathbb{F} is infinite. Since the nonzero-locus of p is closed under rescaling by R , clearing denominators yields points in R^n in the nonzero-locus of p . Each point gives an n -element set satisfying both properties above.

Additionally, all rings with “large enough characteristic” also turn out to work:

Example 3.4. Suppose R is a unital commutative ring in which the elements $1, 2 = 1 + 1, 3 = 1 + 1 + 1, \dots, p_1 \cdots p_n$ are non-zerodivisors, where p_i is the i th prime integer and $n \geq 3$. Then $\{p_1, \dots, p_n\}$ is an n -element subset that satisfies both properties above.

As the above examples suggest, the hypotheses of Theorem 1.6 can be weakened, and we have:

Theorem 3.5. Fix an integer $n \geq 3$ and let R be a unital commutative ring that satisfies properties (1) and (2) listed just before Example 3.2. Then there exist $N = \binom{n}{\lfloor n/2 \rfloor}$ -many directions $\mathbf{v}_1, \dots, \mathbf{v}_N \in R^n$ such that for any map $f : R^n \rightarrow R$, if the restrictions of f to the lines

$$m_0 + Re_i, \quad \forall m_0 \in R^n, \quad 1 \leq i \leq n, \quad \text{and} \quad R\mathbf{v}_1, \dots, R\mathbf{v}_N$$

are each affine-linear, then f is affine-linear on R^n . Fewer than N directions do not suffice.

Proof. Begin by carrying out the analysis in the first part of the proof of Theorem 1.3 around (2.1). (As noted there, this analysis required no restrictions on the ring R .) Thus, Equation (2.1) shows that $f(x_1, \dots, x_n)$ is a multi-affine polynomial in the x_j . Write it as:

$$f(\mathbf{x}) = \sum_{J \subseteq \{1, \dots, n\}} a_J \mathbf{x}^J,$$

where \mathbf{x}^J is the product of x_j for $j \in J$. Also note that every such polynomial is affine-linear on the lines $m_0 + Re_i$, for all $m_0 \in R^n$ and all $1 \leq i \leq n$.

Next, for a general direction $\mathbf{v} \in R^n$, write its coordinates as $\mathbf{v} = (v_1, \dots, v_n)$. Then

$$f(r \cdot \mathbf{v}) = \sum_{k=0}^n r^k \left(\sum_{|J|=k} a_J \prod_{j \in J} v_j \right), \quad \forall r \in R.$$

Adopt the strategy around (2.3), via the hypothesis of property (2) listed before Example 3.2. Thus the coefficient of r^k vanishes for each $k \geq 2$. Notice for a fixed direction v that evaluating across all $r \in R$ only comes this far, i.e., yields

$$\sum_{|J|=k} a_J \prod_{j \in J} v_j = 0, \quad \forall 2 \leq k \leq n. \quad (3.1)$$

For a fixed k , and using a specified direction \mathbf{v} , this is an equation in the $\binom{n}{k}$ variables a_J . Thus, we would require at least $\binom{n}{k}$ directions to resolve this system. In particular if $k = N$, using fewer

than $N = \binom{n}{\lceil n/2 \rceil}$ -many equations will not be able to show that $a_J = 0$ for all J of size N . Thus N is a lower bound for the number of directions needed.

For the upper bound, by the hypotheses there exists an n -element set $S = \{s_1, \dots, s_n\} \subseteq R$ that is at once a weak multiplicative B_h -set for $1 \leq h \leq n$. Define the desired directions/vectors via:

$$\mathbf{v}_i := (s_1^{i-1}, \dots, s_n^{i-1}), \quad 1 \leq i \leq N = \binom{n}{\lceil n/2 \rceil}.$$

Evaluating $f(\cdot)$ along the directions $R\mathbf{v}_i$ yields as in (3.1) the system of equations

$$\sum_{|J|=k} a_J \prod_{j \in J} s_j^{i-1} = 0, \quad \forall 2 \leq k \leq n, \quad 1 \leq i \leq \binom{n}{k}. \quad (3.2)$$

This can be rewritten in the form $A_k \cdot (a_J)_{|J|=k} = 0$, where A_k is a matrix of size $\binom{n}{k} \times \binom{n}{k}$ with (i, J) entry $\left(\prod_{j \in J} s_j\right)^{i-1}$. Now if $k = n$, the leading coefficient of $f(r \cdot \mathbf{v}_1)$ gives $a_{\{1, \dots, n\}} \cdot 1 = 0$. If instead $2 \leq k \leq n-1$, then A_k is a ‘‘usual’’ Vandermonde matrix, and $\det(A_k)$ is a non-zero-divisor by property (2) in the hypotheses. Pre-multiplying (3.2) by the adjugate of A_k and canceling $\det(A_k)$, $a_J = 0$ for $|J| = k \geq 2$. Thus f has no higher-order terms, so is affine-linear on R^n . \square

Acknowledgments. A.K. was partially supported by Ramanujan Fellowship grant SB/S2/RJN-121/2017 and SwarnaJayanti Fellowship grants SB/SJF/2019-20/14 and DST/SJF/MS/2019/3 from SERB and DST (Govt. of India).

REFERENCES

- [1] Raj Chandra Bose and Sarvadaman Chowla. Theorems in the additive theory of numbers. *Comment. Math. Helv.*, 37:141–147, 1962/63.
- [2] Javier Cilleruelo. New upper bounds for finite B_h sequences. *Adv. in Math.*, 159(1):1–17, 2001.
- [3] Quentin Dubroff, Jacob Fox, and Max Wenqiang Xu. A note on the Erdős distinct subset sums problem. *SIAM J. Disc. Math.* 35(1):322–324, 2021.
- [4] Paul Erdős. On some applications of graph theory to number theoretic problems. *Publ. Ramanujan Inst.*, 1:131–136, 1968/69.
- [5] Paul Erdős and Pál Turán. On a problem of Sidon in additive number theory, and on some related problems. *J. London Math. Soc.*, 16(4):212–215, 1941.
- [6] Rachel Greenfeld and Terence Tao. A counterexample to the periodic tiling conjecture. *Preprint*, arXiv:math.CO/2211.15847, 2022.
- [7] Robin Hartshorne. *Foundations of Projective Geometry*. W. A. Benjamin, New York, 1967.
- [8] Moishe Kahan and Dustan Levenstein. <https://math.stackexchange.com/questions/2035287/>, 2016.
- [9] Bernt Lindström. On B_2 -sequences of vectors. *J. Number Theory*, 4(3):261–265, 1972.
- [10] Hong Liu and Péter Pál Pach. The number of multiplicative Sidon sets of integers. *J. Combin. Th. Ser. A*, 165:152–175, 2019.
- [11] Imre Z. Ruzsa. Additive and multiplicative Sidon sets. *Acta Math. Hungarica*, 112:345–354, 2006.
- [12] James Singer. A theorem in finite projective geometry and some applications to number theory. *Trans. Amer. Math. Soc.*, 43(3):377–385, 1938.
- [13] Ernst Snapper and Robert J. Troyer. *Metric affine geometry*. Academic Press, New York, London, 1971.
- [14] Terence Tao. A counterexample to the periodic tiling conjecture. Blogpost, available at <https://terrytao.wordpress.com/2022/11/29/>, 2022.
- [15] Terence Tao and Tamar Ziegler. Concatenation theorems for anti-Gowers-uniform functions and Host–Kra characteristic factors. *Discrete Anal.*, paper no. 13, 61 pp., 2016.
- [16] Karl Georg Christian von Staudt. *Beiträge zur Geometrie der Lage*, Vol. I–II. Nürnberg, 1850s.

(A. Khare) INDIAN INSTITUTE OF SCIENCE, BANGALORE 560012, INDIA; AND ANALYSIS AND PROBABILITY RESEARCH GROUP; BANGALORE 560012, INDIA

Email address: khare@iisc.ac.in

(A. Tikaradze) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TOLEDO, TOLEDO 43606, USA

Email address: tikar06@gmail.com