

# Homotopy type theory as internal languages of diagrams of $\infty$ -logoses

Taichi Uemura

December 6, 2022

## Abstract

We show that certain diagrams of  $\infty$ -logoses are reconstructed in internal languages of their oplax limits via lex, accessible modalities, which enables us to use plain homotopy type theory to reason about not only a single  $\infty$ -logos but also a diagram of  $\infty$ -logoses. This also provides a higher dimensional version of Sterling’s synthetic Tait computability—a type theory for higher dimensional logical relations. To prove the main result, we establish a precise correspondence between the lex, accessible localizations of an  $\infty$ -logos and the lex, accessible modalities in the internal language of the  $\infty$ -logos. To do this, we also partly develop the Kripke-Joyal semantics of homotopy type theory in  $\infty$ -logoses.

## 1 Introduction

An  $\infty$ -logos, also known as an  $\infty$ -topos [30, 4]<sup>1</sup>, is an  $(\infty, 1)$ -category that looks like the  $(\infty, 1)$ -category of spaces, among other aspects of it. An  $\infty$ -logos is a place where one can do homotopy theory just as an ordinary logos is a place where one can do set-level mathematics.

*Homotopy type theory* [48] is another place to do homotopy theory. It is a type theory in the style of Martin-Löf [32] extended by the *univalence axiom* and *higher inductive types*. The former forces types to behave like spaces rather than sets, and the latter allow us to build types representing spaces such as spheres and tori.

$\infty$ -logoses and homotopy type theory are closely related. Shulman [37] has shown that any  $\infty$ -logos is presented by a structure that admits an interpretation of homotopy type theory. In other words, homotopy type theory is an *internal language* of an  $\infty$ -logos. Any theorem proved in homotopy type theory can be translated in an arbitrary  $\infty$ -logos. For example, the proof of the Blakers-Massey connectivity theorem in homotopy type theory [22] has led to a new generalized Blakers-Massey theorem that holds in an arbitrary  $\infty$ -logos [2].

---

<sup>1</sup>The term  $\infty$ -logos is Anel and Joyal’s terminology [4] for  $\infty$ -topos considered as an algebraic structure rather than a geometric object. A morphism of  $\infty$ -logoses is always considered in the direction of the inverse image functor. We use this terminology to clarify the direction of morphisms when speaking about (co)limits of  $\infty$ -logoses.

An  $\infty$ -logos, however, does not live alone.  $\infty$ -logoses are often connected by functors which are also connected by natural transformations. Plain homotopy type theory is, at first sight, not sufficient to reason about a diagram of  $\infty$ -logoses, because the actions of the functors and natural transformations are not internalized to type theory. Even worse, it is impossible to naively internalize some diagrams: some internal adjunction leads a contradiction [27]; there are only trivial internal idempotent comonads [38].

While there is no chance of naive internalization of such interesting but problematic diagrams to plain homotopy type theory, some other diagrams can be internalized in a clever way pointed out by Shulman<sup>2</sup>. A minimal non-trivial example is a diagram consisting of two  $\infty$ -logoses and a lex, accessible functor between them in one direction. The two  $\infty$ -logoses are *lex, accessible localizations* of another  $\infty$ -logos obtained by the *Artin gluing* for the functor, and the functor is reconstructed by composing the inclusion from one localization and the reflector to the other. Moreover, this reconstruction is *internal* to the glued  $\infty$ -logos, because lex, accessible localizations of an  $\infty$ -logos correspond to *lex, accessible modalities* in its internal language. Hence, plain homotopy type theory as an internal language of the glued  $\infty$ -logos is sufficient to reason about the original diagram.

In this paper, we propose a class of shapes of diagrams of  $\infty$ -logoses for which the internal reconstruction technique explained in the previous paragraph works. We call shapes in the proposed class *mode sketches*. Our main result is summarized as follows. Let  $\mathfrak{M}$  be a mode sketch.  $\mathfrak{M}$  is regarded as a presentation of an  $(\infty, 2)$ -category. Then:

1. We associate to  $\mathfrak{M}$  certain axioms in type theory, one of which is to postulate some lex, accessible modalities from which one can construct a diagram of  $\infty$ -logoses internally to type theory (Sections 3.1 and 3.2).
2. For any diagram  $\mathcal{L}$  indexed over  $\mathfrak{M}$  consisting of  $\infty$ -logoses and lex, accessible functors, the oplax limit of  $\mathcal{L}$  is an  $\infty$ -logos that satisfies the axioms associated to  $\mathfrak{M}$ , and the diagram obtained in the internal language of the oplax limit corresponds to the original diagram  $\mathcal{L}$ . Conversely, any  $\infty$ -logos that satisfies the axioms associated to  $\mathfrak{M}$  is obtained by this oplax limit construction (Theorem 10.3).

Recall that the *oplax limit* of a diagram of  $(\infty, 1)$ -categories is a generalization of the Artin gluing [51, 39].

## 1.1 Synthetic Tait computability

This work is closely related to Sterling’s *synthetic Tait computability* [44, 42]. It is a technique of constructing *logical relations* using an internal language for the Artin gluing. A logos obtained by the Artin gluing is always equipped with a distinguished proposition in its internal language. The two lex, accessible

<sup>2</sup>[https://golem.ph.utexas.edu/category/2011/11/internalizing\\_the\\_external\\_or.html](https://golem.ph.utexas.edu/category/2011/11/internalizing_the_external_or.html)

modalities associated to the glued logoses are the *open* and *closed* modalities associated to the proposition. The *fracture and gluing theorem* asserts that every type in the internal language is canonically fractured into an open type and a closed (unary, proof-relevant) relation on it which are glued back together. The internal language for the Artin gluing is thus a type theory with an indeterminate proposition in which *types are relations* and provides a synthetic method of constructing logical relations used in the study of type theories and programming languages. Applications include normalization for complex type theories [43, 19].

We relate synthetic Tait computability and mode sketches. The core axiom for synthetic Tait computability is to postulate some indeterminate propositions. Note that, although the original synthetic Tait computability is based on extensional type theory, postulating propositions makes sense also in homotopy type theory. We show that part of the axioms associated to a mode sketch is equivalent to postulating a lattice of propositions (Theorem 4.3).

Mode sketches thus provide an alternative synthetic method of constructing logical relations. This is also natural from Shulman’s point of view [39] that interpretations of type theory in oplax limits are generalized logical relations. Since we work in homotopy type theory, what we get is actually *higher-dimensional logical relations*, and our primary application of mode sketches in upcoming paper(s) will be normalization for  $\infty$ -*type theories* introduced by Nguyen and Uemura [33] as a higher-dimensional generalization of type theories.

## 1.2 Contributions

Our main result is Theorem 10.3: for every mode sketch  $\mathfrak{M}$ , the models of the axioms associated to  $\mathfrak{M}$  are precisely the oplax limits of diagrams of  $\infty$ -logoses indexed over  $\mathfrak{M}$ . This allows us to reason about a diagram of  $\infty$ -logoses in plain homotopy type theory. We also relate mode sketches to synthetic Tait computability (Theorem 4.3). Mode sketches provide a higher-dimensional version of synthetic Tait computability.

For the main result, we need some new results on modalities in homotopy type theory and  $\infty$ -logoses and on oplax limits of  $(\infty, 1)$ -categories.

### 1.2.1 Modalities in homotopy type theory and localizations of $\infty$ -logoses

*Modalities* in homotopy type theory [35, 11, 12] are idempotent operators on the universe of types. Their  $(\infty, 1)$ -categorical counterparts are to be *localizations* of  $\infty$ -logoses [30, 3, 50]. In this paper, we give a precise correspondence between the lex, accessible localizations of an  $\infty$ -logos and the lex, accessible modalities in the internal language of the  $\infty$ -logos (Theorem 8.1), which is the basis for our main result. Thanks to prior work [3, 50], this is almost straightforward, but one non-trivial gap is the comparison between  $(\infty, 1)$ -categorical accessibility and type-theoretic accessibility. Both are conditions of existence of small presentations, but the former is understood in the metatheory while the latter is in

the internal language of the  $\infty$ -logos. We thus have to understand the interpretation of the existential quantifier in  $\infty$ -logoses. To do this, the *Kripke-Joyal semantics* of homotopy type theory explained in the next subsection would be helpful.

Another minor result is an improvement of the *fracture and gluing theorem* of Rijke, Shulman, and Spitters [35, Theorem 3.50]. It gives a construction of a canonical join of two strongly disjoint lex modalities. We show that this construction preserves accessibility as well (Proposition 2.18).

### 1.2.2 Kripke-Joyal semantics of homotopy type theory

The *Kripke-Joyal semantics* of higher-order logic in a logos  $\mathcal{L}$  [31, Section VI.6] gives us the meaning of a formula in higher-order logic in the language of the metatheory. A *forcing relation*  $X \Vdash_{\mathcal{L}} P$  for an object  $X \in \mathcal{L}$  and a formula  $P$  is introduced, and its meaning is calculated by case analysis on  $P$ . An interesting case is the existential quantifier:  $X \Vdash_{\mathcal{L}} \exists_{x:A} P(x)$  holds if and only if there exist an object  $X' \in \mathcal{L}$ , a surjection  $f : X' \rightarrow X$ , and a map  $a : X' \rightarrow A$  such that  $X' \Vdash_{\mathcal{L}} P(a)$  holds. In this paper, we partly develop the Kripke-Joyal semantics of homotopy type theory in  $\infty$ -logoses and show an analogous result for the existential quantifier as a special case of a result for the propositional truncation (Proposition 8.9).

### 1.2.3 Oplax limits of $(\infty, 1)$ -categories

*Oplax limits* are special  $(\infty, 2)$ -categorical limits and analogous to oplax limits in 2-category theory [46, 25, 23]. Oplax limits indexed over  $(\infty, 1)$ -categories are studied by Gepner, Haugseng, and Nikolaus [18] and generalized to arbitrary indexing  $(\infty, 2)$ -categories by Gagna, Harpaz, and Lanari [15]. In this paper, we show some results on oplax limits. All of them are consequences of results in the literature, but it is worth stating them explicitly.

We show that the oplax limit of a diagram of  $\infty$ -logoses and lex, accessible functors is an  $\infty$ -logos (Theorem 9.34). This is analogous to the classical result of Wraith [51] on oplax limits of (elementary) logoses and generalizes the result of Lurie [30, Proposition 6.3.2.3] on limits of  $\infty$ -logoses.

The oplax limit of a diagram classifies *oplax natural transformations* from a constant diagram to the given diagram. It is known [29, Theorem 3.8.1] that natural transformations between diagrams of  $(\infty, 1)$ -categories correspond to fibred functors between the  $(\infty, 2)$ -categories of elements of the diagrams. We show that oplax natural transformations between diagrams of  $(\infty, 1)$ -categories correspond to arbitrary functors between the  $(\infty, 2)$ -categories of elements (Corollary 9.46). Some special cases of this result have already been proved: Haugseng et al. [21, Theorem E] show the case when the diagrams are indexed over an  $(\infty, 1)$ -category; Gagna, Harpaz, and Lanari [15, Corollary 4.4.3] show the case when the domain is constant on the point.

We show a *mate correspondence* (Proposition 9.49) as a useful source of oplax natural transformations: given two diagrams  $F$  and  $G$  of  $(\infty, 1)$ -categories, the

space of oplax natural transformations  $F \rightarrow G$  that are point-wise left adjoints is equivalent to the space of lax natural transformations  $G \rightarrow F$  that are point-wise right adjoints; see [47] for the 2-categorical case. The special case when the diagrams are indexed over an  $(\infty, 1)$ -category is proved by Haugseng et al. [21, Corollary F] in a stronger form: they give an equivalence between the  $(\infty, 1)$ -categories of oplax/lax natural transformations. Our result is in fact a consequence of their other result [21, Theorem E] and, as they note at the end of the introduction of [21], the proof seems not to work for the stronger statement in full generality. A general mate correspondence for diagrams in an arbitrary  $(\infty, 2)$ -category indexed over an arbitrary  $(\infty, 1)$ -category is discussed in [17, Chapter 12], and two proofs are sketched there.

### 1.3 Organization

In Section 2, we review the theory of modalities in homotopy type theory [35]. Our focus is on the poset of lex, accessible modalities and on the open and closed modalities associated to propositions.

Sections 3 and 4 are the core of the paper. We introduce the notion of a *mode sketch* (Definition 3.5). For every mode sketch, we introduce two equivalent sets of axioms to encode a certain diagram of universes. One postulates some lex, accessible modalities while the other postulates a lattice of propositions. The open and closed modalities give a construction of the former from the latter which we show is an equivalence (Theorem 4.3). The latter is a higher dimensional analogue of Sterling’s synthetic Tait computability [42].

Sections 5 to 7 and 9 are preliminary sections needed for the semantics of mode sketches. We review some  $(\infty, 1)$ -category theory in Section 5, the theory of modalities in  $\infty$ -logoses in Section 6, the semantics of homotopy type theory in  $\infty$ -logoses in Section 7, and some  $(\infty, 2)$ -category theory in Section 9.

In Section 8, we show the equivalence between the poset of lex, accessible localizations of an  $\infty$ -logos and the poset of lex, accessible modalities in the internal language of the  $\infty$ -logos (Theorem 8.1).

Finally in Section 10, we show our main result (Theorem 10.3): for any mode sketch, the space of  $\infty$ -logoses satisfying the axioms associated to the mode sketch is equivalent to the space of diagrams of  $\infty$ -logoses and lex, accessible functors indexed over the mode sketch.

### 1.4 Preliminaries

We assume that the reader is familiar with homotopy type theory [48]. From Section 5, familiarity with basic  $(\infty, 1)$ -category theory [30, 13, 34] is assumed.

By *homotopy type theory* we mean dependent type theory with (dependent) function types, (dependent) pair types, a unit type, intensional identity types, univalent universes  $\mathcal{U} : \uparrow \mathcal{U} : \uparrow^2 \mathcal{U} : \dots$ , and all higher inductive types we need. We mainly follow the HoTT Book [48] for terminologies and notations in homotopy type theory.

For a semantic reason explained in Section 7, our higher inductive types are not assumed to satisfy definitional computation rules. We instead postulate constants that construct initial algebras.

*Example 1.1.* For a type  $A : \mathcal{U}$ , the *propositional truncation*  $\|A\|$  is the initial proposition in  $\mathcal{U}$  equipped with a function  $A \rightarrow \|A\|$ . We postulate a constant of the closed type

$$\prod_{A:\mathcal{U}} \sum_{B:\mathbf{Prop}(\mathcal{U})} \sum_{\tau:A \rightarrow B} \prod_{C:\mathbf{Prop}(\mathcal{U})} \mathbf{lsEquiv}(\lambda(f : B \rightarrow C).f \circ \tau).$$

It is also assumed that the inclusion  $\mathcal{U} \rightarrow \uparrow\mathcal{U}$  commutes with the propositional truncation.

Other preliminaries are reviewed in individual sections: modalities in homotopy type theory in Section 2; presentable  $(\infty, 1)$ -categories and  $\infty$ -logoses in Section 5; modalities in  $\infty$ -logoses in Section 6; semantics of homotopy type theory in  $\infty$ -logoses in Section 7;  $(\infty, 2)$ -category theory in Section 9.

## 1.5 Related work

An earlier version of *cohesive homotopy type theory* [36] uses modalities in plain homotopy type theory to internalize a series of adjunctions. It turns out that their axiomatization is not ideal to work with, and the newer version of cohesive homotopy type theory [38] is a proper extension of homotopy type theory optimized for those particular adjunctions.

A more general framework for internal diagrams is *multimodal dependent type theory* [20]. It is roughly a family of type theories related to each other via modal operators and interpreted in a diagram of presheaf categories. The shape of diagram is specified directly by an arbitrary 2-category which is called a *mode theory* in this context. Our terminology “mode sketch” is chosen to mean a sketch of a mode theory. Multimodal dependent type theory is potentially an internal language for diagrams of  $\infty$ -logoses, but for this one would have to rectify not only  $\infty$ -logoses but also functors and natural transformations between them.

Our work brings back the ideas of earlier cohesive homotopy type theory. Although it might not be the best type theory, it has a lot of advantages: modalities are internal to plain homotopy type theory, and thus all results are ready to formalize in existing proof assistants; keeping type theory simple is also important in informal use of type theory in which the correctness of application of inference rules is not checked by computer; the semantics is clear, since the  $\infty$ -logos semantics of homotopy type theory is well-established [e.g. 7, 5, 39, 37]; it also opens the door to internalization of more general diagrams in a uniform way, which is the motivation for the current work.

## 2 Modalities in homotopy type theory

We review the theory of *modalities* in homotopy type theory [35]. In this section, we work in homotopy type theory.

**Definition 2.1.** A *subuniverse*  $\mathfrak{m}$  is a function  $\text{In}_{\mathfrak{m}} : \mathcal{U} \rightarrow \uparrow\mathcal{U}$  such that  $\text{In}_{\mathfrak{m}}(A)$  is a proposition for all  $A : \mathcal{U}$ . A type  $A$  satisfying  $\text{In}_{\mathfrak{m}}(A)$  is called  *$\mathfrak{m}$ -modal*. We define a subtype  $\mathcal{U}_{\mathfrak{m}} \subset \mathcal{U}$  to be  $\{A : \mathcal{U} \mid \text{In}_{\mathfrak{m}}(A)\}$ .

**Definition 2.2.** A subuniverse  $\mathfrak{m}$  is *reflective* if it is equipped with functions  $\circ_{\mathfrak{m}} : \mathcal{U} \rightarrow \mathcal{U}_{\mathfrak{m}}$  and  $\eta_{\mathfrak{m}} : \prod_{A:\mathcal{U}} A \rightarrow \circ_{\mathfrak{m}} A$  such that that the precomposition  $\lambda f.f \circ \eta_{\mathfrak{m}}(A) : (\circ_{\mathfrak{m}} A \rightarrow B) \rightarrow (A \rightarrow B)$  is an equivalence for any  $B : \mathcal{U}_{\mathfrak{m}}$ . Note that such a pair  $(\circ_{\mathfrak{m}}, \eta_{\mathfrak{m}})$  is unique.

**Definition 2.3.** A reflective subuniverse  $\mathfrak{m}$  is a *modality* if  $\text{In}_{\mathfrak{m}}$  is closed under pair types, that is, for  $A : \mathcal{U}$  and  $B : A \rightarrow \mathcal{U}$ , if  $\text{In}_{\mathfrak{m}}(A)$  and  $\prod_{a:A} \text{In}_{\mathfrak{m}}(B(a))$ , then  $\text{In}_{\mathfrak{m}}(\sum_{a:A} B(a))$ .

**Definition 2.4.** For types  $A, B : \mathcal{U}$ , we define

$$(A \perp B) \equiv \text{IsEquiv}(\lambda(b : B).\lambda(- : A).b).$$

For a subuniverse  $\mathfrak{m}$ , we define subuniverses  $\mathfrak{m}^{\perp}$  and  ${}^{\perp}\mathfrak{m}$  by

$$\begin{aligned} \text{In}_{\mathfrak{m}^{\perp}}(B) &\equiv \prod_{A:\mathcal{U}_{\mathfrak{m}}} A \perp B \\ \text{In}_{{}^{\perp}\mathfrak{m}}(A) &\equiv \prod_{B:\mathcal{U}_{\mathfrak{m}}} A \perp B. \end{aligned}$$

**Definition 2.5.** A *null generator*  $\mu$  consists of  $I_{\mu} : \mathcal{U}$  and  $Z_{\mu} : I_{\mu} \rightarrow \mathcal{U}$ . We write  $\text{NullGen}$  for the type of null generators. Given a null generator  $\mu$ , we define a subuniverse  $\mathfrak{Null}(\mu)$  by  $\text{In}_{\mathfrak{Null}(\mu)}(A) \equiv \prod_{i:I_{\mu}} Z_{\mu}(i) \perp A$ . It is shown that  $\mathfrak{Null}(\mu)$  is a modality [35, Theorem 2.19]. A modality  $\mathfrak{m}$  is *accessible* if it is in the image of  $\mathfrak{Null}$ , that is,  $\|\sum_{\mu:\text{NullGen}} \mathfrak{m} = \mathfrak{Null}(\mu)\|$ .

**Definition 2.6.** For a modality  $\mathfrak{m}$ , a type  $A : \mathcal{U}$  is  *$\mathfrak{m}$ -connected* if  $\circ_{\mathfrak{m}} A$  is contractible. This is equivalent to  $\text{In}_{{}^{\perp}\mathfrak{m}}(A)$  by [35, Corollary 1.37].

**Definition 2.7.** A modality  $\mathfrak{m}$  is *lex* if for any  $\mathfrak{m}$ -connected type  $A : \mathcal{U}$ , the identity type  $a_1 = a_2$  is  $\mathfrak{m}$ -connected for any  $a_1, a_2 : A$ .

From now on, we are mostly interested lex, accessible modalities, so we give them a short name.

**Terminology 2.8.** LAM is an acronym for lex, accessible modality.

Other than LAMs, the *truncation modalities* are of particular importance.

*Example 2.9.* For  $n \geq -2$ , the  *$n$ -truncation modality*  $\mathfrak{T}r_n$  is defined by  $\text{In}_{\mathfrak{T}r_n}(A)$  if and only if  $A$  is  $n$ -truncated. It is indeed a modality [35, Example 1.6]; see also [48, Chapter 7].

## 2.1 The poset of lex, accessible modalities

We have the posets

$$\text{SU} \supset \text{RSU} \supset \text{Mdl} \supset \text{AccMdl} \supset \text{LAM}$$

of subuniverses, reflective subuniverses, modalities, accessible modalities, and lex, accessible modalities, respectively, where all the inclusions are full. We also have the full subposet  $\text{Lex} \subset \text{Mdl}$  of lex modalities. By definition,  $\text{LAM} = \text{Lex} \cap \text{AccMdl}$ . We study the poset  $\text{LAM}$  in more detail.

**Definition 2.10** ([35, Theorem 3.25]). Let  $I : \mathcal{U}$  and  $\mathbf{m} : I \rightarrow \text{LAM}$ . A *canonical meet*  $\bigwedge_{i:I} \mathbf{m}(i)$  is a LAM that is the meet of  $\mathbf{m}(i)$ 's in  $\text{SU}$ . A *canonical join*  $\bigvee_{i:I} \mathbf{m}(i)$  is a LAM satisfying that a type  $A : \mathcal{U}$  is  $(\bigvee_{i:I} \mathbf{m}(i))$ -connected if and only if it is  $\mathbf{m}(i)$ -connected for all  $i : I$ . Note that a canonical join is the join in  $\text{Mdl}$ .

*Example 2.11.* The *top modality*  $\mathfrak{Top}$ , for which all the types are modal, is the canonical meet of the empty family. The *bottom modality*  $\mathfrak{Bot}$ , for which only the contractible types are modal, is the canonical join of the empty family.

The canonical meet of an arbitrary family of LAMs exists [35, Theorem 3.29 and Remark 3.23]. In the following special case, the reflector of a canonical meet is easily computed.

**Fact 2.12** ([35, Theorem 3.30]). *Let  $\mathbf{m}$  and  $\mathbf{n}$  be LAMs. If  $\circ_{\mathbf{m}}$  preserves  $\mathbf{n}$ -modal types, then  $\circ_{\mathbf{m} \wedge \mathbf{n}} A = \circ_{\mathbf{m}} \circ_{\mathbf{n}} A$ .*

Canonical joins are less understood than canonical meets. One important case when canonical joins exist and can be computed is the following.

**Definition 2.13.** Let  $\mathbf{m}$  and  $\mathbf{n}$  be LAMs.  $\mathbf{n}$  is *strongly disjoint from  $\mathbf{m}$*  if any  $\mathbf{m}$ -modal type is  $\mathbf{n}$ -connected or equivalently if  $\mathbf{m} \leq \perp \mathbf{n}$  in  $\text{SU}$ .

**Fact 2.14** ([35, Example 3.32]). *Let  $\mathbf{m}$  and  $\mathbf{n}$  be LAMs. If  $\mathbf{m} \leq \perp \mathbf{n}$ , then  $\mathbf{m} \wedge \mathbf{n} = \mathfrak{Bot}$ .*

**Proposition 2.15** (Fracture and gluing theorem). *Let  $\mathbf{m}$  and  $\mathbf{n}$  be LAMs such that  $\mathbf{m} \leq \perp \mathbf{n}$ .*

1. *The canonical join  $\mathbf{m} \vee \mathbf{n}$  exists.*
2. *A type  $A$  is  $(\mathbf{m} \vee \mathbf{n})$ -modal if and only if the function  $\eta_{\mathbf{m}}(A) : A \rightarrow \circ_{\mathbf{n}} A$  has  $\mathbf{m}$ -modal fibers.*
3.  $\mathcal{U}_{\mathbf{m} \vee \mathbf{n}} \simeq \sum_{A:\mathcal{U}_{\mathbf{m}}} \sum_{B:\mathcal{U}_{\mathbf{n}}} A \rightarrow \circ_{\mathbf{m}}^{\mathbf{n}} B$ .

*In the special case when  $\mathbf{m} = \perp \mathbf{n}$ , we have  $\mathbf{m} \vee \mathbf{n} = \mathfrak{Top}$ .*

*Proof.* All but the accessibility of  $\mathbf{m} \vee \mathbf{n}$  are proved by Rijke, Shulman, and Spitters [35, Theorem 3.50]. We will prove the accessibility of  $\mathbf{m} \vee \mathbf{n}$  in Proposition 2.18 using an open modality.  $\square$

In some cases, canonical meets and canonical joins distribute.

**Proposition 2.16.** *Let  $\mathbf{m}_1$ ,  $\mathbf{m}_2$ , and  $\mathbf{m}_3$  be LAMs. Suppose  $\mathbf{m}_1 \leq \perp \mathbf{m}_3$  and  $\mathbf{m}_2 \leq \perp \mathbf{m}_3$ . Then  $(\mathbf{m}_1 \vee \mathbf{m}_3) \wedge (\mathbf{m}_2 \vee \mathbf{m}_3) = (\mathbf{m}_1 \wedge \mathbf{m}_2) \vee \mathbf{m}_3$ .*



*Proof.* Note that the right side exists as  $\mathbf{m}_1 \wedge \mathbf{m}_2 \leq \mathbf{m}_1 \leq \perp \mathbf{m}_3$ . Let  $A : \mathcal{U}$ . By Proposition 2.15,  $A$  is  $((\mathbf{m}_1 \vee \mathbf{m}_3) \wedge (\mathbf{m}_2 \vee \mathbf{m}_3))$ -modal if and only if fibers of  $\eta_{\mathbf{m}_3}(A)$  are both  $\mathbf{m}_1$ -modal and  $\mathbf{m}_2$ -modal, but this is equivalent to that  $A$  is  $(\mathbf{m}_1 \wedge \mathbf{m}_2) \vee \mathbf{m}_3$ -modal again by Proposition 2.15.  $\square$

**Proposition 2.17.** *Let  $\mathbf{m}_1$ ,  $\mathbf{m}_2$ , and  $\mathbf{m}_3$  be LAMs. Suppose that  $\mathbf{m}_1 \leq \perp \mathbf{m}_2$  and that  $\circ_{\mathbf{m}_2}$  preserves  $\mathbf{m}_3$ -modal types. Then  $(\mathbf{m}_1 \vee \mathbf{m}_2) \wedge \mathbf{m}_3 = (\mathbf{m}_1 \wedge \mathbf{m}_3) \vee (\mathbf{m}_2 \wedge \mathbf{m}_3)$ .*

*Proof.* Note that the right side exists as  $\mathbf{m}_1 \wedge \mathbf{m}_3 \leq \mathbf{m}_1 \leq \perp \mathbf{m}_2 \leq \perp (\mathbf{m}_2 \wedge \mathbf{m}_3)$ . Let  $A$  be a  $((\mathbf{m}_1 \vee \mathbf{m}_2) \wedge \mathbf{m}_3)$ -modal type. We show that  $A$  is  $((\mathbf{m}_1 \wedge \mathbf{m}_3) \vee (\mathbf{m}_2 \wedge \mathbf{m}_3))$ -modal, which is by Proposition 2.15 equivalent to that  $\eta_{\mathbf{m}_2 \wedge \mathbf{m}_3}(A) : A \rightarrow \circ_{\mathbf{m}_2 \wedge \mathbf{m}_3} A$  has  $(\mathbf{m}_1 \wedge \mathbf{m}_3)$ -modal fibers. Since  $\circ_{\mathbf{m}_2}$  preserves  $\mathbf{m}_3$ -modal types and since  $A$  is  $\mathbf{m}_3$ -modal,  $\eta_{\mathbf{m}_2 \wedge \mathbf{m}_3}(A)$  is equivalent to  $\eta_{\mathbf{m}_2}(A) : A \rightarrow \circ_{\mathbf{m}_2} A$  by Fact 2.12. The fibers of  $\eta_{\mathbf{m}_2}(A)$  are  $\mathbf{m}_1$ -modal by Proposition 2.15 and  $\mathbf{m}_3$ -modal since both domain and codomain are  $\mathbf{m}_3$ -modal.  $\square$

## 2.2 Accessibility of the canonical join

Let us fill the gap in the proof of Proposition 2.15.

**Proposition 2.18.** *Let  $\mathbf{m}$  and  $\mathbf{n}$  be LAMs such that  $\mathbf{m} \leq \perp \mathbf{n}$ . Then the canonical join  $\mathbf{m} \vee \mathbf{n}$  (in Lex) is accessible.*

We have to find a null generator for  $\mathbf{m} \vee \mathbf{n}$ . A natural guess is the following.

**Construction 2.19.** Let  $\mu$  and  $\nu$  be null generators. We define a null generator  $\mu \star \nu$  by  $I_{\mu \star \nu} \equiv I_\mu \times I_\nu$  and  $Z_{\mu \star \nu}(i, j) \equiv Z_\mu(i) \star Z_\nu(j) \equiv Z_\mu(i) +_{Z_\mu(i) \times Z_\nu(j)} Z_\nu(j)$ .

**Lemma 2.20.** *Let  $\mathbf{m}$  and  $\mathbf{n}$  be LAMs, and let  $\mu$  and  $\nu$  be null generators for  $\mathbf{m}$  and  $\mathbf{n}$ , respectively. Then  $Z_{\mu \star \nu}(i, j)$  is both  $\mathbf{m}$ -connected and  $\mathbf{n}$ -connected for all  $i : I_\mu$  and  $j : I_\nu$ .*

*Proof.* Recall that a function is  $\mathbf{m}$ -connected if its fibers are  $\mathbf{m}$ -connected and that the class of  $\mathbf{m}$ -connected functions is the left class of a (stable) orthogonal factorization system [35, Theorem 1.34]. Then the claim follows by the pushout stability and the right cancellability of connected functions.  $\square$

Lemma 2.20 shows  $\mathbf{m} \vee \mathbf{n} \leq \mathfrak{Null}(\mu \star \nu)$  for arbitrary accessible modalities  $\mathbf{m}$  and  $\mathbf{n}$  and for arbitrary choices of  $\mu$  and  $\nu$ . We do not know if the other direction holds in general for some choices of  $\mu$  and  $\nu$ . In the special case when  $\mathbf{m} \leq \perp \mathbf{n}$ , the idea is to show that  $\mathbf{n}$  is an *open modality* within the subuniverse of  $\mathfrak{Null}(\mu \star \nu)$ -modal types.

**Construction 2.21.** Let  $P$  be a proposition. We define the *open modality*  $\mathfrak{Op}(P)$  by  $\circ_{\mathfrak{Op}(P)} A \equiv (P \rightarrow A)$  and  $\eta_{\mathfrak{Op}(P)}(A, a) \equiv \lambda_.a$ . It is lex and accessible by [35, Example 2.24 and Example 3.10]. We also define the *closed modality*  $\mathfrak{Cl}(P)$  by  $\text{In}_{\mathfrak{Cl}(P)}(A) \equiv (P \rightarrow \text{lsContr}(A))$ . It is lex and accessible by [35, Example 2.25 and Example 3.14]. Note that  $\mathfrak{Cl}(P) = \perp \mathfrak{Op}(P)$  [35, Example 1.31].

**Lemma 2.22.** *Let  $\mathbf{m}$  and  $\mathbf{n}$  be LAMs such that  $\mathbf{m} \leq \perp \mathbf{n}$ . Then  $\mathbf{n} \leq \mathfrak{Op}(\circ_{\mathbf{m}} \mathbf{0})$ .*

*Proof.* This is because  $\circ_m \mathbf{0}$  is  $\mathbf{n}$ -connected by assumption.  $\square$

**Lemma 2.23.** *Let  $\mathbf{m}$  and  $\mathbf{n}$  be LAMs such that  $\mathbf{m} \leq \perp \mathbf{n}$ . Suppose that  $\mu$  and  $\nu$  are null generators for  $\mathbf{m}$  and  $\mathbf{n}$ , respectively, and that  $\mu$  admits a function  $f : \circ_m \mathbf{0} \rightarrow I_\mu$  such that  $\mathbf{0} \simeq Z_\mu(f(i))$  for all  $i : \circ_m \mathbf{0}$ . Then  $\circ_{\mathfrak{Dp}(\circ_m \mathbf{0})} A$  is  $\mathbf{n}$ -modal for any  $\mathfrak{Null}(\mu \star \nu)$ -modal type  $A$ . Consequently, the canonical function  $\circ_{\mathfrak{Dp}(\circ_m \mathbf{0})} A \rightarrow \circ_n A$  induced by Lemma 2.22 is an equivalence for any  $\mathfrak{Null}(\mu \star \nu)$ -modal type  $A$ .*

*Proof.* We show that  $\circ_{\mathfrak{Dp}(\circ_m \mathbf{0})} A \equiv (\circ_m \mathbf{0} \rightarrow A)$  is  $\mathbf{n}$ -modal. Since  $\nu$  is a null generator for  $\mathbf{n}$ , it suffices to show that  $Z_\nu(j) \perp (\circ_m \mathbf{0} \rightarrow A)$  for all  $j : I_\nu$ . This is equivalent to that  $Z_\nu(j) \perp A$  under an assumption  $i : \circ_m \mathbf{0}$ . This holds since  $Z_\nu(j) \simeq \mathbf{0} \star Z_\nu(j) \simeq Z_\mu(f(i)) \star Z_\nu(j)$  and since  $A$  is  $\mathfrak{Null}(\mu \star \nu)$ -modal.  $\square$

**Lemma 2.24.** *Let  $\mathbf{m}$  and  $\mathbf{n}$  be LAMs such that  $\mathbf{m} \leq \perp \mathbf{n}$ . Suppose that  $\mu$  and  $\nu$  are null generators for  $\mathbf{m}$  and  $\mathbf{n}$ , respectively, and that  $\nu$  has an element  $j : I_\nu$  such that  $Z_\nu(j) \simeq \circ_m \mathbf{0}$ . Then, if a type  $A$  is  $\mathfrak{Null}(\mu \star \nu)$ -modal and  $\mathfrak{Dp}(\circ_m \mathbf{0})$ -connected, then it is  $\mathbf{m}$ -modal.*

*Proof.* We show that  $Z_\mu(i) \perp A$  for all  $i : I_\mu$ . By the definition of  $\star$ , we have the following pullback square.

$$\begin{array}{ccc} (Z_\mu(i) \star \circ_m \mathbf{0} \rightarrow A) & \xrightarrow{\simeq} & (Z_\mu(i) \rightarrow A) \\ \downarrow & \lrcorner & \downarrow \\ (\circ_m \mathbf{0} \rightarrow A) & \xrightarrow{\simeq} & (Z_\mu(i) \rightarrow \circ_m \mathbf{0} \rightarrow A) \end{array}$$

Since  $A$  is  $\mathfrak{Dp}(\circ_m \mathbf{0})$ -connected, the domain and codomain of the bottom function are contractible, and thus the bottom function is an equivalence. It then follows that the top function is also an equivalence. Since  $A$  is  $\mathfrak{Null}(\mu \star \nu)$ -modal and since  $Z_\nu(j) \simeq \circ_m \mathbf{0}$ , we have  $A \simeq (Z_\mu(i) \star \circ_m \mathbf{0} \rightarrow A) \simeq (Z_\mu(i) \rightarrow A)$ , and thus  $Z_\mu(i) \perp A$ .  $\square$

*Proof of Proposition 2.18.* Let  $\mu$  and  $\nu$  be null generators for  $\mathbf{m}$  and  $\mathbf{n}$ , respectively. Note that adjoining a family of connected types to a null generator does not change the modality presented by the null generator. Under an assumption  $i : \circ_m \mathbf{0}$ , the empty type  $\mathbf{0}$  becomes  $\mathbf{m}$ -connected, and thus we may assume that  $\mu$  includes the type family  $\lambda(- : \circ_m \mathbf{0}).\mathbf{0}$ . Since  $\circ_m \mathbf{0}$  is  $\mathbf{n}$ -connected by assumption, we may assume that  $\nu$  includes the type family  $\lambda(- : \mathbf{1}).\circ_m \mathbf{0}$ .

We show that  $\mathfrak{Null}(\mu \star \nu) = \mathbf{m} \vee \mathbf{n}$ . By Lemma 2.20,  $\mathbf{m} \vee \mathbf{n} \leq \mathfrak{Null}(\mu \star \nu)$ . For the other direction, suppose that  $A$  is a  $\mathfrak{Null}(\mu \star \nu)$ -modal type. By [35, Theorem 3.50], it suffices to show that  $\eta_m(A) : A \rightarrow \circ_n A$  has  $\mathbf{m}$ -modal fibers. By Lemma 2.23,  $\circ_n A \simeq \circ_{\mathfrak{Dp}(\circ_m \mathbf{0})} A$ . Then the fibers of  $\eta_m(A)$  are  $\mathfrak{Dp}(\circ_m \mathbf{0})$ -connected. Since both  $A$  and  $\circ_n A$  are  $\mathfrak{Null}(\mu \star \nu)$ -modal, the fibers of  $\eta_m(A)$  are also  $\mathfrak{Null}(\mu \star \nu)$ -modal. Thus, by Lemma 2.24,  $\eta_m(A)$  has  $\mathbf{m}$ -modal fibers.  $\square$

As a by-product, we have the following.

**Corollary 2.25.** *Let  $\mathfrak{m}$  and  $\mathfrak{n}$  be LAMs such that  $\mathfrak{m} \leq \perp \mathfrak{n}$ . If  $\mathfrak{m} \vee \mathfrak{n} = \mathfrak{Top}$ , then  $\mathfrak{m}$  and  $\mathfrak{n}$  are the closed and open, respectively, modalities associated to the proposition  $\circ_{\mathfrak{m}} \mathbf{0}$ .*  $\square$

### 2.3 Open and closed modalities

We collect some properties of *open modalities* and *closed modalities*.

**Proposition 2.26.** *The function  $\mathfrak{Cl} : \mathbf{Prop} \rightarrow \mathbf{LAM}$  is a contravariant embedding of posets and takes arbitrary joins to canonical meets.*

*Proof.*  $\mathfrak{Cl}$  takes joins to canonical meets by [35, Example 3.27]. It then follows that  $\mathfrak{Cl}$  contravariantly preserves ordering. To see that  $\mathfrak{Cl}$  reflects ordering, let  $P$  and  $Q$  be propositions and suppose that  $\mathfrak{Cl}(Q) \leq \mathfrak{Cl}(P)$ . By definition,  $Q$  is  $\mathfrak{Cl}(Q)$ -modal and thus  $\mathfrak{Cl}(P)$ -modal, but this implies that  $P \rightarrow Q$ .  $\square$

**Proposition 2.27.** *The function  $\mathfrak{Op} : \mathbf{Prop} \rightarrow \mathbf{LAM}$  is a covariant embedding of posets and takes finite meets to canonical meets and arbitrary joins to canonical joins.*

*Proof.* Since  $\mathfrak{Cl}(P) = \perp \mathfrak{Op}(P)$ , the function  $\mathfrak{Op}$  is a covariant embedding of posets and takes arbitrary joins to canonical joins by Proposition 2.26.  $\mathfrak{Op}$  takes finite meets to canonical meets by [35, Example 3.26].  $\square$

**Proposition 2.28.** *Let  $P$  be a proposition and and  $\mathfrak{m}$  a LAM.  $\circ_{\mathfrak{Op}(P)}$  preserves  $\mathfrak{m}$ -modal types, and  $\circ_{\mathfrak{m}}$  preserves  $\mathfrak{Cl}(P)$ -types.*

*Proof.* Let  $A : \mathcal{U}$ . If  $A$  is  $\mathfrak{m}$ -modal, then  $\circ_{\mathfrak{Op}(P)} A \equiv P \rightarrow A$  is  $\mathfrak{m}$ -modal by [35, Lemma 1.26]. If  $A$  is  $\mathfrak{Cl}(P)$ -modal, then  $\circ_{\mathfrak{m}} A$  is  $\mathfrak{Cl}(P)$ -modal by [35, Lemma 1.27].  $\square$

**Proposition 2.29.**  $\mathfrak{m} = (\mathfrak{m} \wedge \mathfrak{Cl}(P)) \vee (\mathfrak{m} \wedge \mathfrak{Op}(P))$  for any LAM  $\mathfrak{m}$  and any proposition  $P$ .

*Proof.*  $\mathfrak{m} = \mathfrak{m} \wedge (\mathfrak{Cl}(P) \vee \mathfrak{Op}(P)) = (\mathfrak{m} \wedge \mathfrak{Cl}(P)) \vee (\mathfrak{m} \wedge \mathfrak{Op}(P))$ , where the first identification is by Proposition 2.15 and the second is by Propositions 2.17 and 2.28.  $\square$

## 3 Mode sketches

We introduce *mode sketches* as shapes of diagrams of subuniverses definable internally to type theory. We work in homotopy type theory through the section.

### 3.1 Internal diagrams induced by modalities

We consider postulating some LAMs to encode some diagram of subuniverses. The fundamental observation is that a pair of LAMs induces a canonical functor between them.

**Construction 3.1.** Let  $\mathfrak{m}$  and  $\mathfrak{n}$  be LAMs. We define a function  $\circ_{\mathfrak{m}}^{\mathfrak{n}} : \mathcal{U}_{\mathfrak{n}} \rightarrow \mathcal{U}_{\mathfrak{m}}$  to be the restriction of  $\circ_{\mathfrak{m}}$  to  $\mathcal{U}_{\mathfrak{n}} \subset \mathcal{U}$ .

*Remark 3.2.* We can say that  $\circ_{\mathfrak{m}}^{\mathfrak{n}}$  is a functor *externally*: we can construct a function  $\prod_{A,B:\mathcal{U}_{\mathfrak{n}}}(A \rightarrow B) \rightarrow (\circ_{\mathfrak{m}}^{\mathfrak{n}} A \rightarrow \circ_{\mathfrak{m}}^{\mathfrak{n}} B)$  and every instance of the coherence laws. However, it is not known how to state that  $\circ_{\mathfrak{m}}^{\mathfrak{n}}$  is a functor internally to type theory, because defining the type of  $(\infty, 1)$ -categories in plain homotopy type theory is still an open problem.

We have two functors  $\circ_{\mathfrak{n}}^{\mathfrak{m}} : \mathcal{U}_{\mathfrak{m}} \rightarrow \mathcal{U}_{\mathfrak{n}}$  and  $\circ_{\mathfrak{m}}^{\mathfrak{n}} : \mathcal{U}_{\mathfrak{n}} \rightarrow \mathcal{U}_{\mathfrak{m}}$  for every pair of LAMs  $\mathfrak{m}$  and  $\mathfrak{n}$ , but we are often interested in only one direction. It is thus useful to cut off one direction by postulating that  $\mathfrak{m} \leq^{\perp} \mathfrak{n}$ : by the definition of connectedness,  $\circ_{\mathfrak{n}}^{\mathfrak{m}}$  becomes constant at the unit type. The other direction  $\circ_{\mathfrak{m}}^{\mathfrak{n}} : \mathcal{U}_{\mathfrak{n}} \rightarrow \mathcal{U}_{\mathfrak{m}}$  remains non-trivial. Therefore, a pair  $(\mathfrak{m}, \mathfrak{n})$  of LAMs such that  $\mathfrak{m} \leq^{\perp} \mathfrak{n}$  encodes a functor  $\mathcal{U}_{\mathfrak{n}} \rightarrow \mathcal{U}_{\mathfrak{m}}$ . When  $\mathfrak{n} \leq^{\perp} \mathfrak{m}$  is also assumed,  $\mathcal{U}_{\mathfrak{m}}$  and  $\mathcal{U}_{\mathfrak{n}}$  are considered unrelated.

Given more than two LAMs, we have canonical natural transformations between the canonical functors.

**Construction 3.3.** Let  $\mathfrak{m}_0, \mathfrak{m}_1, \mathfrak{m}_2$  be LAMs. We define

$$\eta_{\mathfrak{m}_1}^{\mathfrak{m}_0;\mathfrak{m}_2} : \prod_{A:\mathcal{U}_{\mathfrak{m}_2}} \circ_{\mathfrak{m}_0}^{\mathfrak{m}_2} A \rightarrow \circ_{\mathfrak{m}_0}^{\mathfrak{m}_1} \circ_{\mathfrak{m}_1}^{\mathfrak{m}_2} A$$

by  $\eta_{\mathfrak{m}_1}^{\mathfrak{m}_0;\mathfrak{m}_2}(A) \equiv \circ_{\mathfrak{m}_0} \eta_{\mathfrak{m}_1}(A)$ . This family of functions is natural in the sense that for any  $A, B : \mathcal{U}_{\mathfrak{m}_2}$  and  $f : A \rightarrow B$ , we have a homotopy filling the following square.

$$\begin{array}{ccc} \circ_{\mathfrak{m}_0}^{\mathfrak{m}_2} A & \xrightarrow{\eta_{\mathfrak{m}_1}^{\mathfrak{m}_0;\mathfrak{m}_2}(A)} & \circ_{\mathfrak{m}_0}^{\mathfrak{m}_1} \circ_{\mathfrak{m}_1}^{\mathfrak{m}_2} A \\ \circ_{\mathfrak{m}_0}^{\mathfrak{m}_2} f \downarrow & & \downarrow \circ_{\mathfrak{m}_0}^{\mathfrak{m}_1} \circ_{\mathfrak{m}_1}^{\mathfrak{m}_2} f \\ \circ_{\mathfrak{m}_0}^{\mathfrak{m}_2} B & \xrightarrow{\eta_{\mathfrak{m}_1}^{\mathfrak{m}_0;\mathfrak{m}_2}(B)} & \circ_{\mathfrak{m}_0}^{\mathfrak{m}_1} \circ_{\mathfrak{m}_1}^{\mathfrak{m}_2} B \end{array}$$

Let  $\mathfrak{m}_0, \mathfrak{m}_1, \mathfrak{m}_2, \mathfrak{m}_3$  be LAMs. By naturality, the following diagram commutes.

$$\begin{array}{ccc} \circ_{\mathfrak{m}_0}^{\mathfrak{m}_3} & \xrightarrow{\eta_{\mathfrak{m}_1}^{\mathfrak{m}_0;\mathfrak{m}_3}} & \circ_{\mathfrak{m}_0}^{\mathfrak{m}_1} \circ_{\mathfrak{m}_1}^{\mathfrak{m}_3} \\ \eta_{\mathfrak{m}_2}^{\mathfrak{m}_0;\mathfrak{m}_3} \downarrow & & \downarrow \circ_{\mathfrak{m}_0}^{\mathfrak{m}_1} \eta_{\mathfrak{m}_2}^{\mathfrak{m}_1;\mathfrak{m}_3} \\ \circ_{\mathfrak{m}_0}^{\mathfrak{m}_2} \circ_{\mathfrak{m}_2}^{\mathfrak{m}_3} & \xrightarrow{\eta_{\mathfrak{m}_1}^{\mathfrak{m}_0;\mathfrak{m}_2} \circ_{\mathfrak{m}_2}^{\mathfrak{m}_3}} & \circ_{\mathfrak{m}_0}^{\mathfrak{m}_1} \circ_{\mathfrak{m}_1}^{\mathfrak{m}_2} \circ_{\mathfrak{m}_2}^{\mathfrak{m}_3} \end{array}$$

For more than four LAMs, higher coherence laws are also satisfied. Hence, a tuple  $(\mathfrak{m}_0, \dots, \mathfrak{m}_n)$  of LAMs such that  $\mathfrak{m}_i \leq^{\perp} \mathfrak{m}_j$  for all  $i < j$  encodes an  $n$ -simplex with vertices  $\mathcal{U}_{\mathfrak{m}_i}$ , edges  $\circ_{\mathfrak{m}_i}^{\mathfrak{m}_j} : \mathcal{U}_{\mathfrak{m}_j} \rightarrow \mathcal{U}_{\mathfrak{m}_i}$  for  $i < j$ , triangles

$$\begin{array}{ccc} \mathcal{U}_{\mathfrak{m}_i} & \xleftarrow{\circ_{\mathfrak{m}_i}^{\mathfrak{m}_k}} & \mathcal{U}_{\mathfrak{m}_k} \\ & \swarrow \circ_{\mathfrak{m}_i}^{\mathfrak{m}_j} & \searrow \circ_{\mathfrak{m}_k}^{\mathfrak{m}_j} \\ & \mathcal{U}_{\mathfrak{m}_j} & \end{array}$$

$\Downarrow \eta_{\mathfrak{m}_j}^{\mathfrak{m}_i;\mathfrak{m}_k}$

for  $i < j < k$ , and higher homotopies.

Shapes other than simplices are expressed by postulating invertibility of some of  $\eta_{\mathbf{m}_j}^{\mathbf{m}_i; \mathbf{m}_k}$ 's. For example, let  $\mathbf{m}_0, \mathbf{m}_1, \mathbf{m}_2, \mathbf{m}_3$  be LAMs and suppose that  $\mathbf{m}_i \leq^\perp \mathbf{m}_j$  for all  $i < j$ , that  $\mathbf{m}_2 \leq^\perp \mathbf{m}_1$ , and that  $\eta_{\mathbf{m}_1}^{\mathbf{m}_0; \mathbf{m}_3}$  is invertible. We have a diagram

$$\begin{array}{ccccc}
 & & \mathcal{U}_{\mathbf{m}_1} & & \\
 & \circlearrowleft^{\mathbf{m}_1}_{\mathbf{m}_0} & \swarrow & & \searrow^{\circlearrowleft^{\mathbf{m}_3}_{\mathbf{m}_1}} \\
 \mathcal{U}_{\mathbf{m}_0} & \longleftarrow & \circlearrowleft^{\mathbf{m}_3}_{\mathbf{m}_0} & \longrightarrow & \mathcal{U}_{\mathbf{m}_3} \\
 & \circlearrowleft^{\mathbf{m}_2}_{\mathbf{m}_0} & \swarrow & & \searrow^{\circlearrowleft^{\mathbf{m}_3}_{\mathbf{m}_2}} \\
 & & \mathcal{U}_{\mathbf{m}_2} & & 
 \end{array}$$

$\eta_{\mathbf{m}_1}^{\mathbf{m}_0; \mathbf{m}_3} \approx \eta_{\mathbf{m}_2}^{\mathbf{m}_0; \mathbf{m}_3}$

which is equivalent to a diagram of the form

$$\begin{array}{ccccc}
 & & \mathcal{U}_{\mathbf{m}_1} & & \\
 & \circlearrowleft^{\mathbf{m}_1}_{\mathbf{m}_0} & \swarrow & & \searrow^{\circlearrowleft^{\mathbf{m}_3}_{\mathbf{m}_1}} \\
 \mathcal{U}_{\mathbf{m}_0} & \longleftarrow & \Downarrow & \longrightarrow & \mathcal{U}_{\mathbf{m}_3} \\
 & \circlearrowleft^{\mathbf{m}_2}_{\mathbf{m}_0} & \swarrow & & \searrow^{\circlearrowleft^{\mathbf{m}_3}_{\mathbf{m}_2}} \\
 & & \mathcal{U}_{\mathbf{m}_2} & & 
 \end{array}$$

We cannot, however, naively postulate some properties of the functors  $\circlearrowleft_{\mathbf{m}}^{\mathbf{n}}$ 's such as conservativity, fullness, faithfulness, adjointness, and invertibility. This is because the internal statements of these conditions are too strong due to stability under substitution, and indeed some “no-go” theorems on internalizing properties of functors are known [27, Theorem 5.1][38, Theorem 4.1]. We will also see in Remark 10.16 a semantic reason why we cannot postulate properties of  $\circlearrowleft_{\mathbf{m}}^{\mathbf{n}}$ 's.

*Remark 3.4.* It is *possible* to postulate arbitrary properties of  $\circlearrowleft_{\mathbf{m}_i}^{\mathbf{m}_j}$ 's in the following way. We first postulate a “base” LAM  $\mathfrak{Base}$  and assume  $\mathfrak{Base} \leq^\perp \mathbf{m}_i$  for all  $i$ . The universe  $\mathcal{U}_{\mathfrak{Base}}$  is intended to be interpreted as the  $(\infty, 1)$ -category of spaces, so statements in  $\mathcal{U}_{\mathfrak{Base}}$  will correspond to external statements. Since  $\circlearrowleft_{\mathfrak{Base}} : \mathcal{U} \rightarrow \mathcal{U}_{\mathfrak{Base}}$  preserves finite limits, it takes  $(\infty, 1)$ -categories and functors to functors. We can then postulate any property on the induced functor  $\circlearrowleft_{\mathfrak{Base}} \mathcal{U}_{\mathbf{m}_j} \rightarrow \circlearrowleft_{\mathfrak{Base}} \mathcal{U}_{\mathbf{m}_i}$ . In fact, cohesive homotopy type theory [36] was first formulated in a similar fashion where the  $\sharp$  modality plays the role of  $\mathfrak{Base}$ . However, since we only know that  $\circlearrowleft_{\mathfrak{Base}} \mathcal{U}_{\mathbf{m}_i}$  is an  $(\infty, 1)$ -category *externally*, this approach is not so convenient to work with especially for formalization in proof assistants. For this and some other reasons, the newer version of cohesive homotopy type theory [38] is a proper extension of homotopy type theory. Nevertheless, this adding-base approach is attractive since it keeps type theory simple and works for any kind of diagram.

### 3.2 Mode sketches

We introduce *mode sketches* as shapes of diagrams definable by the methodology explained in Section 3.1.

**Definition 3.5.** A *mode sketch*  $\mathfrak{M}$  consists of the following data:

- a decidable finite poset  $I_{\mathfrak{M}}$ ;
- a subset  $T_{\mathfrak{M}}$  of triangles in  $I_{\mathfrak{M}}$ .

Here, by a *decidable* poset we mean a poset whose ordering relation  $\leq$  is decidable. Note that the identity relation  $i = j$  is equivalent to  $(i \leq j) \wedge (j \leq i)$  and thus decidable. The strict ordering relation  $i < j$  defined as  $(i \leq j) \wedge (i \neq j)$  is also decidable. By a *triangle* in  $I_{\mathfrak{M}}$  we mean an ordered triple  $(i_0 < i_1 < i_2)$  of elements of  $I_{\mathfrak{M}}$ . A triangle in  $T_{\mathfrak{M}}$  is called *thin*.

*Remark 3.6.* The definition of mode sketches also makes sense in the metatheory. Every mode sketch  $\mathfrak{M}$  in the metatheory can be encoded in type theory since it is finite.

Let  $\mathfrak{M}$  be a mode sketch and  $\mathfrak{m} : \mathfrak{M} \rightarrow \text{LAM}$  a function. We consider the following axioms.

**Axiom A.**  $\mathfrak{m}(i) \leq \perp \mathfrak{m}(j)$  for any  $j \not\leq i$  in  $\mathfrak{M}$ .

**Axiom B.** For any triangle  $(i_0 < i_1 < i_2) : T_{\mathfrak{M}}$ , the natural transformation  $\eta_{\mathfrak{m}(i_1)}^{\mathfrak{m}(i_0); \mathfrak{m}(i_2)} : \circ_{\mathfrak{m}(i_0)}^{\mathfrak{m}(i_2)} \Rightarrow \circ_{\mathfrak{m}(i_0)}^{\mathfrak{m}(i_1)} \circ_{\mathfrak{m}(i_1)}^{\mathfrak{m}(i_2)}$  is invertible.

**Axiom C.** The top modality is the canonical join  $\bigvee_{\mathfrak{M}} \mathfrak{m}$ .

*Remark 3.7.* Assuming Axiom A, if  $i < j$ , then  $\mathfrak{m}(i) \leq \perp \mathfrak{m}(j)$ .

Axioms A and B are motivated by the observation made in Section 3.1. Our intended models constructed in Section 10 additionally satisfy Axiom C. This axiom is not so important in practical use, since our primary aim is to draw a diagram of  $\infty$ -logoses inside homotopy type theory, but Axiom C does nothing for this purpose. It is even better to work without Axiom C, because Axioms A and B are stable under restriction along a full inclusion  $\mathfrak{M}' \subset \mathfrak{M}$  while Axiom C is not. Axiom C is meant to exclude models other than intended models.

A mode sketch  $\mathfrak{M}$  is regarded as a presentation of an  $(\infty, 2)$ -category. The strict ordering relation generates 1-cells  $(i < j) : i \rightarrow j$ , and the triangles  $(i < j < k)$  generate 2-cells in the direction

$$\begin{array}{ccc}
 & j & \\
 & \nearrow & \searrow \\
 i & \longrightarrow & k. \\
 & \Downarrow & \\
 & & 
 \end{array}$$

When the triangle is thin, the corresponding 2-cell is made invertible. Longer chains  $(i_0 < i_1 < \dots < i_n)$  present coherence. A function  $\mathfrak{m} : \mathfrak{M} \rightarrow \text{LAM}$  satisfying Axioms A to C is then considered as a diagram of subuniverses indexed over  $\mathfrak{M}^{\text{op}(1,2)}$ , the  $(\infty, 2)$ -category obtained from  $\mathfrak{M}$  by reversing the directions of 1-cells and 2-cells.

*Example 3.8.* Every decidable finite poset is a mode sketch where no triangle is thin. The  $(\infty, 2)$ -category presented by it is obtained from the left adjoint of the Duskin nerve [14] by reversing 2-cells.

*Example 3.9.* The *mode sketch for functors* is drawn as

$$0 \longrightarrow 1.$$

Axiom A asserts  $\mathbf{m}(0) \leq \perp \mathbf{m}(1)$ . Axiom B is empty since there is no triangle. Thus, we get the following diagram.

$$\mathcal{U}_{\mathbf{m}(0)} \xleftarrow{\circlearrowleft_{\mathbf{m}(0)}^{\mathbf{m}(1)}} \mathcal{U}_{\mathbf{m}(1)}$$

Axiom C asserts  $\mathbf{m}(0) \vee \mathbf{m}(1) = \mathfrak{Top}$ .

*Example 3.10.* The *mode sketch for triangles* is drawn as

$$\begin{array}{ccc} 0 & \xrightarrow{\quad} & 2 \\ & \searrow & \nearrow \\ & 1 & \end{array}$$

$\simeq$

where “ $\simeq$ ” indicates that the triangle is thin. Axiom A asserts  $\mathbf{m}(0) \leq \perp \mathbf{m}(1)$ ,  $\mathbf{m}(0) \leq \perp \mathbf{m}(2)$ , and  $\mathbf{m}(1) \leq \perp \mathbf{m}(2)$ . Axiom B asserts that  $\eta_{\mathbf{m}(1)}^{\mathbf{m}(0); \mathbf{m}(2)}$  is invertible. Thus, we have the following commutative triangle.

$$\begin{array}{ccc} & \circlearrowleft_{\mathbf{m}(0)}^{\mathbf{m}(2)} & \\ \mathcal{U}_{\mathbf{m}(0)} & \xleftarrow{\quad} & \mathcal{U}_{\mathbf{m}(2)} \\ & \swarrow & \searrow \\ \circlearrowleft_{\mathbf{m}(0)}^{\mathbf{m}(1)} & \mathcal{U}_{\mathbf{m}(1)} & \circlearrowleft_{\mathbf{m}(1)}^{\mathbf{m}(2)} \end{array}$$

Axiom C asserts  $\mathbf{m}(0) \vee \mathbf{m}(1) \vee \mathbf{m}(2) = \mathfrak{Top}$ . Notice that theorems for the mode sketch for functors proved without Axiom C also apply to the three edges in the above diagram. To keep this reusability, we should not assume Axiom C in practical use.

Under Axiom A, canonical joins of  $\mathbf{m}(i)$ ’s exist and are well-behaved.

**Proposition 3.11.** *If a function  $\mathbf{m} : \mathfrak{M} \rightarrow \text{LAM}$  satisfies Axiom A, then the canonical join  $\bigvee_S \mathbf{m}$  exists for any decidable subset  $S \subset \mathfrak{M}$ .*

*Proof.* By induction on the size of  $S$ . If  $S$  is empty, then  $\bigvee_{\emptyset} \mathbf{m}$  is the bottom modality. Suppose that  $S$  is non-empty. Since  $\mathfrak{M}$  is finite, there is an element  $i_0$  minimal in  $S$ . Then  $S \setminus \{i_0\}$  admits a canonical join by the induction hypothesis. Since  $i_0$  is minimal,  $\mathbf{m}(i_0) \leq \perp \mathbf{m}(i)$  for any  $i : S \setminus \{i_0\}$  by Axiom A, and thus  $\mathbf{m}(i_0) \leq \perp (\bigvee_{S \setminus \{i_0\}} \mathbf{m})$ . Then we have the canonical join  $\bigvee_S \mathbf{m} \equiv \mathbf{m}(i_0) \vee (\bigvee_{(S \setminus \{i_0\})} \mathbf{m})$  by Proposition 2.15.  $\square$

**Definition 3.12.** A *cosieve* on a decidable poset  $I$  is an upward-closed decidable subset of it. Let  $\text{coSieve}(I)$  denote the poset of cosieves on  $I$  ordered by inclusion. Note that cosieves are closed under finite meets and joins, so  $\text{coSieve}(I)$  is a (distributive) lattice.

**Proposition 3.13.** *If a function  $\mathfrak{m} : \mathfrak{M} \rightarrow \text{LAM}$  satisfies Axiom A, then  $\bigvee_{\mathfrak{M} \setminus S} \mathfrak{m} \leq \perp (\bigvee_S \mathfrak{m})$  for any cosieve  $S \subset \mathfrak{M}$ .*

**Lemma 3.14.** *Let  $\mathfrak{m}_0, \mathfrak{m}_1$ , and  $\mathfrak{m}_2$  be LAMs such that  $\mathfrak{m}_i \leq \perp \mathfrak{m}_j$  for any  $i < j$ . Then  $\mathfrak{m}_0 \vee \mathfrak{m}_1 \leq \perp \mathfrak{m}_2$ .*

*Proof.* Let  $A$  be a  $(\mathfrak{m}_0 \vee \mathfrak{m}_1)$ -modal type. By Proposition 2.15,  $\eta_{\mathfrak{m}_1}(A) : A \rightarrow \bigcirc_{\mathfrak{m}_1} A$  has  $\mathfrak{m}_0$ -modal fibers. Then, by assumption,  $\bigcirc_{\mathfrak{m}_1} A$  and the fibers of  $\eta_{\mathfrak{m}_1}(A)$  are made contractible by  $\bigcirc_{\mathfrak{m}_2}$ . Thus,  $\bigcirc_{\mathfrak{m}_2} A$  is contractible.  $\square$

*Proof of Proposition 3.13.* Since  $S$  is upward-closed,  $j \not\leq i$  for any  $i : \mathfrak{M} \setminus S$  and  $j : S$ . Thus, by Axiom A,  $\mathfrak{m}(i) \leq \perp \mathfrak{m}(j)$  for any  $i : \mathfrak{M} \setminus S$  and  $j : S$ . The claim follows from Lemma 3.14 and the construction of the canonical join in Proposition 3.11.  $\square$

### 3.3 Intended models, internally

Let  $\mathfrak{M}$  be a mode sketch. We can internally see what kind of an  $\infty$ -logos is a model of  $\mathfrak{M}$ . Here, by a model of  $\mathfrak{M}$  we mean an  $\infty$ -logos that admits an interpretation of a postulated function  $\mathfrak{m} : \mathfrak{M} \rightarrow \text{LAM}$  satisfying Axioms A to C.

*Example 3.15.* Consider the case when  $\mathfrak{M}$  is the mode sketch for functors (Example 3.9). Proposition 2.15 implies that  $\mathcal{U} \simeq \mathcal{U}_{\mathfrak{m}(0) \vee \mathfrak{m}(1)}$  is the *Artin gluing* for the functor  $\bigcirc_{\mathfrak{m}(0)}^{\mathfrak{m}(1)} : \mathcal{U}_{\mathfrak{m}(1)} \rightarrow \mathcal{U}_{\mathfrak{m}(0)}$ . Therefore, our intended models of  $\mathfrak{M}$  are  $\infty$ -logoses obtained by the Artin gluing.

A generalization of the Artin gluing is *oplax limits*. In the setting of Example 3.15,  $\mathcal{U}$  fits into the following *universal oplax cone* over the diagram

$$\mathcal{U}_{\mathfrak{m}(0)} \xleftarrow{\bigcirc_{\mathfrak{m}(0)}^{\mathfrak{m}(1)}} \mathcal{U}_{\mathfrak{m}(1)}.$$

$$\begin{array}{ccc} & \mathcal{U} & \\ & \swarrow \quad \searrow & \\ \mathcal{U}_{\mathfrak{m}(0)} & \xleftarrow{\quad \quad \quad} & \mathcal{U}_{\mathfrak{m}(1)} \\ & \bigcirc_{\mathfrak{m}(0)}^{\mathfrak{m}(1)} & \end{array} \quad (1)$$

An oplax cone over a diagram is a kind of cone but every triangle formed by two projections and a functor in the diagram is only filled by a not necessarily invertible natural transformation in the direction of Eq. (1). The universal oplax cone or oplax limit is the terminal object in the  $(\infty, 1)$ -category of oplax cones.

*Example 3.16.* Consider the case when  $\mathfrak{M}$  is the mode sketch  $\{0 \rightarrow 1 \rightarrow 2\}$  with no thin triangle. Iterating Proposition 2.15, we see that every type  $A : \mathcal{U}$  is fractured into  $A_0 : \mathcal{U}_{\mathfrak{m}(0)}$ ,  $A_1 : \mathcal{U}_{\mathfrak{m}(1)}$ ,  $A_2 : \mathcal{U}_{\mathfrak{m}(2)}$ ,  $f_{01} : A_0 \rightarrow \bigcirc_{\mathfrak{m}(0)}^{\mathfrak{m}(1)} A_1$ ,  $f_{02} : A_0 \rightarrow$



$\circ_{\mathfrak{m}(0)}^{\mathfrak{m}(2)} A_2$ ,  $f_{12} : A_1 \rightarrow \circ_{\mathfrak{m}(1)}^{\mathfrak{m}(2)} A_2$ , and  $p_{012} : \circ_{\mathfrak{m}(0)}^{\mathfrak{m}(1)} f_{12} \circ f_{01} = \eta_{\mathfrak{m}(1)}^{\mathfrak{m}(0); \mathfrak{m}(2)}(A_2) \circ f_{02}$ . Thus,  $\mathcal{U}$  is the oplax limit of the diagram

$$\begin{array}{ccc}
 & \circ_{\mathfrak{m}(0)}^{\mathfrak{m}(2)} & \\
 \mathcal{U}_{\mathfrak{m}(0)} & \xleftarrow{\quad} & \mathcal{U}_{\mathfrak{m}(2)} \\
 & \Downarrow \eta_{\mathfrak{m}(1)}^{\mathfrak{m}(0); \mathfrak{m}(2)} & \\
 & \mathcal{U}_{\mathfrak{m}(1)} & \\
 & \circ_{\mathfrak{m}(0)}^{\mathfrak{m}(1)} \quad \circ_{\mathfrak{m}(1)}^{\mathfrak{m}(2)} & 
 \end{array} \tag{2}$$

That is, we have projections  $\mathcal{U} \rightarrow \mathcal{U}_{\mathfrak{m}(i)}$  for all  $i$ , natural transformations

$$\begin{array}{ccc}
 & \mathcal{U} & \\
 & \swarrow \quad \searrow & \\
 \mathcal{U}_{\mathfrak{m}(i)} & \xrightleftharpoons{\quad} & \mathcal{U}_{\mathfrak{m}(j)} \\
 & \circ_{\mathfrak{m}(i)}^{\mathfrak{m}(j)} & 
 \end{array}$$

for all  $i < j$ , and a homotopy

$$\begin{array}{ccc}
 & \mathcal{U} & \\
 & \swarrow \quad \searrow & \\
 \mathcal{U}_{\mathfrak{m}(0)} & \xrightleftharpoons{\quad} & \mathcal{U}_{\mathfrak{m}(2)} \\
 & \Downarrow \eta_{\mathfrak{m}(1)}^{\mathfrak{m}(0); \mathfrak{m}(2)} & \\
 & \mathcal{U}_{\mathfrak{m}(1)} & \\
 & \circ_{\mathfrak{m}(0)}^{\mathfrak{m}(1)} \quad \circ_{\mathfrak{m}(1)}^{\mathfrak{m}(2)} & 
 \end{array} = \begin{array}{ccc}
 & \mathcal{U} & \\
 & \swarrow \quad \searrow & \\
 \mathcal{U}_{\mathfrak{m}(0)} & \xrightleftharpoons{\quad} & \mathcal{U}_{\mathfrak{m}(2)} \\
 & \Downarrow \eta_{\mathfrak{m}(1)}^{\mathfrak{m}(0); \mathfrak{m}(2)} & \\
 & \mathcal{U}_{\mathfrak{m}(1)} & \\
 & \circ_{\mathfrak{m}(0)}^{\mathfrak{m}(1)} \quad \circ_{\mathfrak{m}(1)}^{\mathfrak{m}(2)} & 
 \end{array},$$

and these data form a universal oplax cone over Eq. (2).

Let us make the triangle ( $0 < 1 < 2$ ) thin so that the natural transformation (2) becomes invertible. In this setting,  $\mathcal{U}$  is still the oplax limit of Eq. (2), but the presentation can be simplified since the type of data ( $f_{02}, p_{012}$ ) is contractible.

In general, one can see that  $\mathcal{U}$  is the oplax limit of a suitable diagram  $\{\mathcal{U}_{\mathfrak{m}(i)}\}_{i:\mathfrak{M}}$ , and thus our intended models of  $\mathfrak{M}$  are oplax limits of  $\infty$ -logoses indexed over the  $(\infty, 2)$ -category presented by  $\mathfrak{M}$ .

## 4 Mode sketches and synthetic Tait computability

We give an alternative set of axioms for mode sketches and exhibits a connection between mode sketches and *synthetic Tait computability* of Sterling [42]. We work in homotopy type theory.

### 4.1 Alternative mode sketch axioms

The  $\infty$ -logoses obtained by the Artin gluing can be characterized as  $\infty$ -logoses equipped with a subterminal object [cf. 24, A4.5.6]. We generalize this from the

Artin gluing to oplax limits indexed by mode sketches, internally to type theory: the type of functions  $\mathfrak{M} \rightarrow \text{LAM}$  satisfying Axioms A and C is equivalent to the type of lattice morphisms  $\text{coSieve}(\mathfrak{M}) \rightarrow \text{Prop}$  (Theorem 4.3).

**Notation 4.1.** For  $i : \mathfrak{M}$ , let  $(i \downarrow)$  denote the cosieve  $\{j : \mathfrak{M} \mid i \leq j\}$  and  $\partial(i \downarrow)$  the cosieve  $(i \downarrow) \setminus \{i\}$ .

**Construction 4.2.** Let  $P : \text{coSieve}(\mathfrak{M}) \rightarrow \text{Prop}$  be a function. We define a function  $\mathfrak{a}_P : \mathfrak{M} \rightarrow \text{LAM}$  by  $\mathfrak{a}_P(i) \equiv \mathfrak{Dp}(P(i \downarrow)) \wedge \mathfrak{Cl}(P(\partial(i \downarrow)))$ .

**Theorem 4.3.** *Construction 4.2 is restricted to an equivalence between the following types:*

1. the type of lattice morphisms  $P : \text{coSieve}(\mathfrak{M}) \rightarrow \text{Prop}$ ;
2. the type of functions  $\mathfrak{m} : \mathfrak{M} \rightarrow \text{LAM}$  satisfying Axioms A and C.

We write  $\varphi_{\mathfrak{m}} : \text{coSieve}(\mathfrak{M}) \rightarrow \text{Prop}$  for the lattice morphism corresponding to  $\mathfrak{m}$ .

Postulating some propositions is the core axiom of *synthetic Tait computability* [44, 42, 45]. Theorem 4.3 thus gives an alternative formulation of it.

*Example 4.4.* Let  $\mathfrak{M}$  be the mode sketch for functors (Example 3.9). Then  $\text{coSieve}(\mathfrak{M}) = \{\{\}, \{1\}, \{0, 1\}\}$  is the free distributive lattice generated by the single element  $\{1\}$ . We thus have  $\{\text{lattice morphisms } \text{coSieve}(\mathfrak{M}) \rightarrow \text{Prop}\} \simeq \text{Prop}$ .

*Remark 4.5.* Geometrically, a lattice morphism  $\text{coSieve}(\mathfrak{M}) \rightarrow \text{Prop}$  specifies a lattice of open subtoposes. On the other hand, a function  $\mathfrak{m} : \mathfrak{M} \rightarrow \text{LAM}$  satisfying Axioms A and C divides the  $\infty$ -topos into disjoint subtoposes  $X_i$ 's. The strong disjointness  $\mathfrak{m}(i) \leq \perp \mathfrak{m}(j)$  means that  $X_j$  is open in  $X_i \cup X_j$  (Lemma 2.23).

We first show that Construction 4.2 is restricted to a function  $1 \rightarrow 2$ .

**Proposition 4.6.** *Let  $P : \text{coSieve}(\mathfrak{M}) \rightarrow \text{Prop}$  be a function.*

1. *If  $P$  preserves binary meets, then  $\mathfrak{a}_P$  satisfies Axiom A.*
2. *If  $P$  preserves top elements and finite joins, then  $\mathfrak{a}_P$  satisfies Axiom C.*

**Lemma 4.7.** *If a function  $P : \text{coSieve}(\mathfrak{M}) \rightarrow \text{Prop}$  preserves finite joins, then  $\mathfrak{Dp}(P(S)) = \bigvee_S \mathfrak{a}_P$  for any cosieve  $S \subset \mathfrak{M}$ .*

*Proof.* By induction on the size of  $S$ . By assumption and by Proposition 2.27, we have  $\mathfrak{Dp}(P(S)) = \bigvee_{i:S} \mathfrak{Dp}(P(i \downarrow))$ . Thus, it is enough to show the case when  $S$  is of the form  $(i \downarrow)$ . By Proposition 2.29, we have  $\mathfrak{Dp}(P(i \downarrow)) = (\mathfrak{Dp}(P(i \downarrow)) \wedge \mathfrak{Cl}(P(\partial(i \downarrow)))) \vee (\mathfrak{Dp}(P(i \downarrow)) \wedge \mathfrak{Dp}(P(\partial(i \downarrow)))) = \mathfrak{a}_P(i) \vee \mathfrak{Dp}(P(\partial(i \downarrow)))$ . Then apply the induction hypothesis for  $\partial(i \downarrow)$ .  $\square$

**Lemma 4.8.** *Let  $\mathfrak{m}$  and  $\mathfrak{n}$  be LAMs. If  $\circ_{\mathfrak{n}}$  preserves  $\mathfrak{m}$ -modal types, then  $\mathfrak{m} \wedge \perp(\mathfrak{n} \wedge \mathfrak{m}) = \mathfrak{m} \wedge \perp \mathfrak{n}$ .*

*Proof.* By Fact 2.12. □

*Proof of Proposition 4.6.* If  $P$  preserves binary meets, then for any  $j \not\leq i$  in  $\mathfrak{M}$ ,

$$\begin{aligned}
& \mathbf{a}_P(i) \\
= & \quad \{\text{definition}\} \\
& \mathfrak{Dp}(P(i \downarrow)) \wedge {}^\perp \mathfrak{Dp}(P(\partial(i \downarrow))) \\
\leq & \quad \{(j \downarrow) \cap (i \downarrow) \subset \partial(i \downarrow) \text{ as } j \not\leq i\} \\
& \mathfrak{Dp}(P(i \downarrow)) \wedge {}^\perp \mathfrak{Dp}(P((j \downarrow) \cap (i \downarrow))) \\
= & \quad \{\text{Proposition 2.27, and } P \text{ preserves binary meets}\} \\
& \mathfrak{Dp}(P(i \downarrow)) \wedge {}^\perp (\mathfrak{Dp}(P(j \downarrow)) \wedge \mathfrak{Dp}(P(i \downarrow))) \\
= & \quad \{\text{Lemma 4.8 and Proposition 2.28}\} \\
& \mathfrak{Dp}(P(i \downarrow)) \wedge {}^\perp \mathfrak{Dp}(P(j \downarrow)) \\
\leq & \quad \{\text{definition}\} \\
& {}^\perp \mathbf{a}_P(j),
\end{aligned}$$

and thus Axiom A is satisfied. If  $P$  preserves top elements and finite joins, then  $\mathbf{a}_P$  satisfies Axiom C by Lemma 4.7. □

The inverse function  $2 \rightarrow 1$  is constructed as follows.

**Construction 4.9.** Let  $\mathbf{m} : \mathfrak{M} \rightarrow \text{LAM}$  be a function satisfying Axioms A and C. We define a lattice morphism  $\varphi_{\mathbf{m}} : \text{coSieve}(\mathfrak{M}) \rightarrow \text{Prop}$  by  $\varphi_{\mathbf{m}}(S) \equiv \bigcirc_{\bigvee_{\mathfrak{M} \setminus S} \mathbf{0}}$ . By Corollary 2.25, Proposition 3.13, and Axiom C, we have  $\bigvee_S \mathbf{m} = \mathfrak{Dp}(\varphi_{\mathbf{m}}(S))$ . Because  $S \mapsto \bigvee_S \mathbf{m}$  preserves finite joins by definition,  $\varphi_{\mathbf{m}}$  preserves finite joins. By Axiom C,  $\varphi_{\mathbf{m}}$  preserves top elements. For preservation of binary meets, let  $S_1$  and  $S_2$  be cosieves on  $\mathfrak{M}$ . We have to show that  $\bigvee_{S_1 \cap S_2} \mathbf{m} = (\bigvee_{S_1} \mathbf{m}) \wedge (\bigvee_{S_2} \mathbf{m})$ . Let  $S_3 \equiv S_1 \cap S_2$ ,  $S'_1 \equiv S_1 \setminus S_3$ , and  $S'_2 \equiv S_2 \setminus S_3$ . By Proposition 3.13,  $\bigvee_{S'_1} \mathbf{m} \leq {}^\perp (\bigvee_{S_2} \mathbf{m})$  and  $\bigvee_{S'_2} \mathbf{m} \leq {}^\perp (\bigvee_{S_1} \mathbf{m})$ . Then,

$$\begin{aligned}
& (\bigvee_{S_1} \mathbf{m}) \wedge (\bigvee_{S_2} \mathbf{m}) \\
= & \quad \{\text{definition}\} \\
& ((\bigvee_{S'_1} \mathbf{m}) \vee (\bigvee_{S_3} \mathbf{m})) \wedge ((\bigvee_{S'_2} \mathbf{m}) \vee (\bigvee_{S_3} \mathbf{m})) \\
= & \quad \{\text{Proposition 2.16}\} \\
& ((\bigvee_{S'_1} \mathbf{m}) \wedge (\bigvee_{S'_2} \mathbf{m})) \vee (\bigvee_{S_3} \mathbf{m}) \\
= & \quad \{\text{Fact 2.14}\} \\
& \bigvee_{S_3} \mathbf{m}.
\end{aligned}$$

*Proof of Theorem 4.3.* It remains to show that the constructions  $P \mapsto \mathbf{a}_P$  and  $\mathbf{m} \mapsto \varphi_{\mathbf{m}}$  are mutually inverses. Lemma 4.7 implies that  $\varphi_{\mathbf{a}_P} = P$ . For the other identification, we have

$$\mathbf{a}_{\varphi_{\mathbf{m}}}(i)$$

$$\begin{aligned}
&= \{\text{definition}\} \\
&\quad \mathfrak{Dp}(\varphi_{\mathfrak{m}}(i \downarrow)) \wedge \mathfrak{Cl}(\varphi_{\mathfrak{m}}(\partial(i \downarrow))) \\
&= \{\text{Lemma 4.7}\} \\
&\quad (\bigvee_{(i \downarrow)} \mathfrak{m}) \wedge \mathfrak{Cl}(\varphi_{\mathfrak{m}}(\partial(i \downarrow))) \\
&= \{\text{Propositions 2.17 and 2.28}\} \\
&\quad \bigvee_{j:(i \downarrow)} \mathfrak{m}(j) \wedge \mathfrak{Cl}(\varphi_{\mathfrak{m}}(\partial(i \downarrow))) \\
&= \{\text{Lemma 4.7}\} \\
&\quad \bigvee_{j:(i \downarrow)} \mathfrak{m}(j) \wedge (\bigwedge_{k:\partial(i \downarrow)} \perp \mathfrak{m}(k)) \\
&= \{\mathfrak{m}(k) \wedge \perp \mathfrak{m}(k) = \mathfrak{Bot} \text{ for } k : \partial(i \downarrow) \text{ by Fact 2.14}\} \\
&\quad \mathfrak{m}(i) \wedge (\bigwedge_{k:\partial(i \downarrow)} \perp \mathfrak{m}(k)) \\
&= \{\text{Axiom A}\} \\
&\quad \mathfrak{m}(i).
\end{aligned}$$

□

## 4.2 Logical relations as types

We have seen in Section 4.1 that synthetic Tait computability is reformulated in terms of LAMs. The slogan of synthetic Tait computability is “logical relations as types” [44]. This is also formulated purely in terms of LAMs. Let  $\mathfrak{m}$  and  $\mathfrak{n}$  be LAMs such that  $\mathfrak{m} \leq \perp \mathfrak{n}$ .

**Fact 4.10** ([35, Theorem 3.11]). *For any LAM  $\mathfrak{m}$ , the universe of  $\mathfrak{m}$ -modal types  $\mathcal{U}_{\mathfrak{m}} \equiv \{A : \mathcal{U} \mid \text{In}_{\mathfrak{m}}(A)\}$  is  $\mathfrak{m}$ -modal.*

**Proposition 4.11** (Fracture and gluing). *Let  $\mathfrak{m}$  and  $\mathfrak{n}$  be LAMs such that  $\mathfrak{m} \leq \perp \mathfrak{n}$ . Then we have an equivalence*

$$\mathcal{U}_{\mathfrak{m} \vee \mathfrak{n}} \simeq \sum_{B:\mathcal{U}_{\mathfrak{n}}} B \rightarrow \mathcal{U}_{\mathfrak{m}}$$

whose right-to-left function sends a  $(B, A)$  to  $\sum_{\mathbf{x}:B} A(\mathbf{x})$ .

*Proof.* For any  $B : \mathcal{U}_{\mathfrak{n}}$ , we have

$$\begin{aligned}
&\sum_{A:\mathcal{U}_{\mathfrak{m}}} A \rightarrow \circ_{\mathfrak{m}} B \\
&\simeq \{\text{equivalence between fibrations and type families}\} \\
&\quad \circ_{\mathfrak{m}} B \rightarrow \mathcal{U}_{\mathfrak{m}} \\
&\simeq \{\text{Fact 4.10}\} \\
&\quad B \rightarrow \mathcal{U}_{\mathfrak{m}}.
\end{aligned}$$

Then apply Proposition 2.15. □

Proposition 4.11 asserts that a type in  $\mathcal{U}_{\mathfrak{m} \vee \mathfrak{n}}$  is a  $\mathfrak{n}$ -modal type equipped with a  $\mathfrak{m}$ -modal unary (proof-relevant) relation on it, so *types (in  $\mathcal{U}_{\mathfrak{m} \vee \mathfrak{n}}$ ) are relations*.

More generally, for a mode sketch  $\mathfrak{M}$  and a function  $\mathfrak{m} : \mathfrak{M} \rightarrow \mathbf{LAM}$  satisfying Axiom A, types in  $\mathcal{U}_{\vee_{\mathfrak{M}} \mathfrak{m}}$  are fractured into a sort of generalized relations by iterated applications of Proposition 4.11.

*Example 4.12.* When  $\mathfrak{M}$  is the mode sketch  $\{0 \leftarrow 01 \rightarrow 1\}$ , we have an equivalence

$$\mathcal{U}_{\mathfrak{m}(01) \vee \mathfrak{m}(1) \vee \mathfrak{m}(0)} \simeq \sum_{A_0 : \mathcal{U}_{\mathfrak{m}(0)}} \sum_{A_1 : \mathcal{U}_{\mathfrak{m}(1)}} A_0 \rightarrow A_1 \rightarrow \mathcal{U}_{\mathfrak{m}(01)}.$$

*Example 4.13.* When  $\mathfrak{M}$  is the mode sketch  $\{0 \rightarrow 1 \rightarrow 2\}$ , we have an equivalence

$$\mathcal{U}_{\mathfrak{m}(0) \vee \mathfrak{m}(1) \vee \mathfrak{m}(2)} \simeq \sum_{A_2 : \mathcal{U}_{\mathfrak{m}(2)}} \sum_{A_1 : A_2 \rightarrow \mathcal{U}_{\mathfrak{m}(1)}} \prod_{x_2} A_1(x_2) \rightarrow \mathcal{U}_{\mathfrak{m}(0)}.$$

The equivalence in Proposition 4.11 nicely interacts with type constructors, and we derive the *logical relation translation* of dependent type theory [e.g. 10, 39, 49, 26] as a *theorem* in type theory:

- $\mathcal{U}_{\mathfrak{m} \vee \mathfrak{n}} : \uparrow \mathcal{U}_{\mathfrak{m} \vee \mathfrak{n}}$  corresponds to the pair  $(\mathcal{U}_{\mathfrak{n}}, \lambda B. B \rightarrow \mathcal{U}_{\mathfrak{m}})$ ;
- $\mathbf{1} : \mathcal{U}_{\mathfrak{m} \vee \mathfrak{n}}$  corresponds to the pair  $(\mathbf{1}, \lambda \_ . \mathbf{1})$ ;
- Suppose that  $A : \mathcal{U}_{\mathfrak{m} \vee \mathfrak{n}}$  corresponds to a pair  $(A_{\mathfrak{n}}, A_{\mathfrak{m}})$ . Then  $(A \rightarrow \mathcal{U}_{\mathfrak{m} \vee \mathfrak{n}}) : \uparrow \mathcal{U}_{\mathfrak{m} \vee \mathfrak{n}}$  corresponds to the pair

$$(A_{\mathfrak{n}} \rightarrow \mathcal{U}_{\mathfrak{n}}, \lambda B. \prod_{x : A_{\mathfrak{n}}} A_{\mathfrak{m}}(x) \rightarrow B(x) \rightarrow \mathcal{U}_{\mathfrak{m}}).$$

Indeed,

$$\begin{aligned} & A \rightarrow \mathcal{U}_{\mathfrak{m} \vee \mathfrak{n}} \\ \simeq & \quad \{\text{fracture and gluing}\} \\ & (\sum_{x : A_{\mathfrak{n}}} A_{\mathfrak{m}}(x)) \rightarrow (\sum_{B : \mathcal{U}_{\mathfrak{n}}} B \rightarrow \mathcal{U}_{\mathfrak{m}}) \\ \simeq & \quad \{\prod \text{ distributes over } \sum\} \\ & \sum_{B : \prod_{x : A_{\mathfrak{n}}} A_{\mathfrak{m}}(x) \rightarrow \mathcal{U}_{\mathfrak{n}}} \prod_x \prod_y B(x, y) \rightarrow \mathcal{U}_{\mathfrak{m}} \\ \simeq & \quad \{\mathcal{U}_{\mathfrak{n}} \simeq (A_{\mathfrak{m}}(x) \rightarrow \mathcal{U}_{\mathfrak{n}}) \text{ since } \mathfrak{m} \leq \perp \mathfrak{n}\} \\ & \sum_{B : A_{\mathfrak{n}} \rightarrow \mathcal{U}_{\mathfrak{n}}} \prod_x A_{\mathfrak{m}}(x) \rightarrow B(x) \rightarrow \mathcal{U}_{\mathfrak{m}}; \end{aligned}$$

- Suppose that  $A : \mathcal{U}_{\mathfrak{m} \vee \mathfrak{n}}$  corresponds to a pair  $(A_{\mathfrak{n}}, A_{\mathfrak{m}})$  and that  $B : A \rightarrow \mathcal{U}_{\mathfrak{m} \vee \mathfrak{n}}$  corresponds to a pair  $(B_{\mathfrak{n}}, B_{\mathfrak{m}})$ . Then  $\prod_{x : A} B(x) : \mathcal{U}_{\mathfrak{m} \vee \mathfrak{n}}$  corresponds to the pair

$$(\prod_{x_{\mathfrak{n}} : A_{\mathfrak{n}}} B_{\mathfrak{n}}(x_{\mathfrak{n}}), \lambda f. \prod_{x_{\mathfrak{n}}} \prod_{x_{\mathfrak{m}} : A_{\mathfrak{m}}(x_{\mathfrak{n}})} B_{\mathfrak{m}}(x_{\mathfrak{n}}, x_{\mathfrak{m}}, f(x_{\mathfrak{n}})))$$

by a similar calculation to the previous clause.  $\sum_{x : A} B(x) : \mathcal{U}_{\mathfrak{m} \vee \mathfrak{n}}$  corresponds to the pair

$$(\sum_{x_{\mathfrak{n}} : A_{\mathfrak{n}}} B_{\mathfrak{n}}(x_{\mathfrak{n}}), \lambda(a_{\mathfrak{n}}, b_{\mathfrak{n}}). \sum_{x_{\mathfrak{m}} : A_{\mathfrak{m}}(a_{\mathfrak{n}})} B_{\mathfrak{m}}(a_{\mathfrak{n}}, x_{\mathfrak{m}}, b_{\mathfrak{n}}));$$

- Suppose that  $A : \mathcal{U}_{\mathbf{m} \vee \mathbf{n}}$  corresponds to a pair  $(A_{\mathbf{n}}, A_{\mathbf{m}})$ , that  $a : A$  corresponds to a pair  $(a_{\mathbf{n}}, a_{\mathbf{m}})$ , and that  $a' : A$  corresponds to a pair  $(a'_{\mathbf{n}}, a'_{\mathbf{m}})$ . Then  $a = a' : \mathcal{U}_{\mathbf{m} \vee \mathbf{n}}$  corresponds to the pair

$$(a_{\mathbf{n}} = a'_{\mathbf{n}}, \lambda p. a_{\mathbf{m}} =_p^{A_{\mathbf{m}}} a'_{\mathbf{m}}).$$

The interaction of the equivalences in Examples 4.12 and 4.13 and type constructors is similarly calculated. We thus conclude that types in  $\mathcal{U}_{\mathbf{V}_{\mathbf{m} \vee \mathbf{n}}}$  are generalized logical relations.

## 5 $(\infty, 1)$ -category theory

We collect facts about  $(\infty, 1)$ -categories needed to interpret homotopy type theory in  $\infty$ -logoses and to compare modalities in homotopy type theory and modalities in  $\infty$ -logoses.

We work with a model-independent language of  $(\infty, 1)$ -category theory rather than choosing a specific model of  $(\infty, 1)$ -categories. An  $(\infty, 1)$ -category  $\mathcal{C}$  consists of a space  $\mathbf{Obj}(\mathcal{C})$  of *objects* of  $\mathcal{C}$ , a space  $\mathbf{Map}_{\mathcal{C}}(x, y)$  of *morphisms from  $x$  to  $y$*  for any objects  $x$  and  $y$ , and composition operators unital and associative up to coherent homotopy. Concepts in category theory such as functors, natural transformations, equivalences, adjoints, (co)limits, and Kan extensions have  $(\infty, 1)$ -categorical analogues. We refer the reader to [30, 13, 34] for general  $(\infty, 1)$ -category theory.

Grothendieck universes  $\mathfrak{U} \in \uparrow \mathfrak{U} \in \uparrow^2 \mathfrak{U} \in \dots$  are assumed to exist. By *small* we mean  $\mathfrak{U}$ -small. Let  $\mathbf{Spc} = \mathbf{Spc}(\mathfrak{U})$  denote the  $(\infty, 1)$ -category of  $\mathfrak{U}$ -small spaces. Let  $\mathbf{Cat} = \mathbf{Cat}(\mathfrak{U})$  denote the  $(\infty, 1)$ -category of  $\mathfrak{U}$ -small  $(\infty, 1)$ -categories. For  $(\infty, 1)$ -categories  $\mathcal{C}$  and  $\mathcal{D}$ , let  $\mathbf{Fun}(\mathcal{C}, \mathcal{D})$  denote the  $(\infty, 1)$ -category of functors from  $\mathcal{C}$  to  $\mathcal{D}$  and natural transformations between them. When  $\mathcal{C}$  and  $\mathcal{D}$  has  $\kappa$ -small limits for a regular cardinal  $\kappa$ , we define  $\mathbf{Fun}_{\mathbf{Lex}(\kappa)}(\mathcal{C}, \mathcal{D}) \subset \mathbf{Fun}(\mathcal{C}, \mathcal{D})$  to be the full subcategory spanned by the functors preserving  $\kappa$ -small limits. For a functor  $x : I \rightarrow \mathcal{C}$ , we write  $\{\pi_i : \lim_{j \in I} x_j \rightarrow x_i\}_{i \in I}$  for the limit cone if it exists and  $\{\iota_i : x_i \rightarrow \operatorname{colim}_{j \in I} x_j\}_{i \in I}$  for the colimit cocone if it exists.

### 5.1 Presentable $(\infty, 1)$ -categories

We review the theory of *presentable  $(\infty, 1)$ -categories*. The standard reference is [30, Chapter 5]. Most results are generalization of those in the theory of locally presentable categories [1].

**Definition 5.1.** An  $(\infty, 1)$ -category  $\mathcal{L}$  is *presentable (over  $\mathfrak{U}$ )* if for some  $\mathfrak{U}$ -small regular cardinal  $\kappa$  and some  $\mathfrak{U}$ -small  $(\infty, 1)$ -category  $\mathcal{C}$  with  $\kappa$ -small colimits,  $\mathcal{L}$  is the completion of  $\mathcal{C}$  under  $\mathfrak{U}$ -small  $\kappa$ -filtered colimits. By [30, Corollary 5.3.5.4], this is equivalent to that  $\mathcal{L} \simeq \mathbf{Fun}_{\mathbf{Lex}(\kappa)}(\mathcal{C}^{\text{op}}, \mathbf{Spc})$  for some  $\kappa$  and  $\mathcal{C}$ .

**Fact 5.2** (Representability theorem [30, Proposition 5.5.2.2]). *Let  $\mathcal{L}$  be a presentable  $(\infty, 1)$ -category. A functor  $\mathcal{L}^{\text{op}} \rightarrow \mathbf{Spc}$  is representable if and only if it preserves small limits.*

**Definition 5.3.** A functor between presentable  $(\infty, 1)$ -categories is *accessible* if it preserves small  $\kappa$ -filtered colimits for some small regular cardinal  $\kappa$ .

**Fact 5.4** (Adjoint functor theorem [30, Corollary 5.5.2.9]). *Let  $\mathcal{L}$  be a presentable  $(\infty, 1)$ -category.*

1. *Let  $\mathcal{K}$  be a locally small  $(\infty, 1)$ -category with small colimits. A functor  $\mathcal{L} \rightarrow \mathcal{K}$  is a left adjoint if and only if it preserves small colimits.*
2. *Let  $\mathcal{K}$  be a presentable  $(\infty, 1)$ -category. A functor  $\mathcal{L} \rightarrow \mathcal{K}$  is a right adjoint if and only if it is accessible and preserves small limits.*

**Construction 5.5.** Let  $\mathbf{Pr}^{\text{L}} = \mathbf{Pr}^{\text{L}}(\mathfrak{U}) \subset \mathbf{Cat}(\uparrow\mathfrak{U})$  denote the subcategory spanned by the presentable  $(\infty, 1)$ -categories and the functors between them preserving small colimits. Let  $\mathbf{Pr}^{\text{R}} = \mathbf{Pr}^{\text{R}}(\mathfrak{U}) \subset \mathbf{Cat}(\uparrow\mathfrak{U})$  denote the subcategory spanned by the presentable  $(\infty, 1)$ -categories and the accessible functors between them preserving small limits. By the adjoint functor theorem,  $(\mathbf{Pr}^{\text{L}})^{\text{op}} \simeq \mathbf{Pr}^{\text{R}}$ .

**Fact 5.6** (Limit theorem [30, Theorem 5.5.3.18]).  $\mathbf{Pr}^{\text{R}} \subset \mathbf{Cat}(\uparrow\mathfrak{U})$  is closed under small limits.

**Proposition 5.7.** *Small colimits in  $\mathbf{Pr}^{\text{L}}$  are computed in the  $(\infty, 1)$ -category of large  $(\infty, 1)$ -categories with small colimits.*

*Proof.* This follows from [40, Proposition 5.14]. □

**Construction 5.8.** Let  $\mathcal{L}$  be a presentable  $(\infty, 1)$ -category. We define the  $n$ -th universe enlargement by

$$\uparrow^n \mathcal{L} = \mathbf{Fun}_{\text{Lex}(\mathfrak{U})}(\mathcal{L}^{\text{op}}, \mathbf{Spc}(\uparrow^n \mathfrak{U})).$$

The Yoneda embedding defines a full inclusion  $\mathcal{L} \rightarrow \uparrow^n \mathcal{L}$ . Note that  $\mathcal{L} \simeq \uparrow^0 \mathcal{L}$  by the representability theorem. By construction,  $\uparrow^n \mathcal{L}$  is presentable over the universe  $\uparrow^n \mathfrak{U}$ . Given a morphism  $F : \mathcal{L} \rightarrow \mathcal{K}$  in  $\mathbf{Pr}^{\text{L}}(\mathfrak{U})$ , the precomposition with  $F$  defines a morphism  $F^* : \uparrow^n \mathcal{K} \rightarrow \uparrow^n \mathcal{L}$  in  $\mathbf{Pr}^{\text{R}}(\uparrow^n \mathfrak{U})$ . This makes the  $n$ -th universe enlargement a functor  $\uparrow^n : \mathbf{Pr}^{\text{L}}(\mathfrak{U})^{\text{op}} \rightarrow \mathbf{Pr}^{\text{R}}(\uparrow^n \mathfrak{U})$ . By Fact 5.6 and Proposition 5.7, this functor preserves small limits. We thus regard it as a functor  $\mathbf{Pr}^{\text{R}}(\mathfrak{U}) \rightarrow \mathbf{Pr}^{\text{R}}(\uparrow^n \mathfrak{U})$  preserving small limits.

*Remark 5.9.*  $\uparrow^n \mathbf{Spc} \simeq \mathbf{Spc}(\uparrow^n \mathfrak{U})$  and  $\uparrow^n \mathbf{Cat} \simeq \mathbf{Cat}(\uparrow^n \mathfrak{U})$ .

## 5.2 $\infty$ -logoses

We review the theory of  $\infty$ -logoses, also known as  $\infty$ -toposes. The standard reference is [30, Chapter 6].

**Definition 5.10.** An  $\infty$ -logos is a presentable  $(\infty, 1)$ -category  $\mathcal{L}$  such that the functor  $\mathcal{L}^{\text{op}} \ni A \mapsto \mathcal{L}/_A \in \uparrow \mathbf{Cat}$ , where the action of morphisms is given by pullback, preserves small limits. A *morphism of  $\infty$ -logoses* is a functor preserving small colimits and finite limits. Note that any morphism of  $\infty$ -logoses has a right adjoint by the adjoint functor theorem. We write  $\mathbf{Logos} \subset \uparrow \mathbf{Cat}$  for the subcategory spanned by the  $\infty$ -logoses and the morphisms of  $\infty$ -logoses.

**Fact 5.11** ([30, Theorem 6.1.3.9]). *A presentable  $(\infty, 1)$ -category  $\mathcal{L}$  is an  $\infty$ -logos if and only if, for any small  $(\infty, 1)$ -category  $I$ , any natural transformation  $f : B \Rightarrow A : I \rightarrow \mathcal{L}$  where all the naturality squares are pullbacks, and for any cocone over  $f$  of the form*

$$\begin{array}{ccc} B_i & \xrightarrow{g_i} & B' \\ f_i \downarrow & & \downarrow f' \\ A_i & \xrightarrow{\iota_i} & \text{colim}_{i \in I} A_i, \end{array} \quad (3)$$

*$B'$  is the colimit of  $B_i$ 's if and only if Eq. (3) is a pullback for every  $i \in I$ .*

**Corollary 5.12.** *Let  $\{F_i : \mathcal{L} \rightarrow \mathcal{K}_i\}_{i \in I}$  be a family of functors between presentable  $(\infty, 1)$ -categories preserving small colimits and finite limits. If all the  $\mathcal{K}_i$ 's are  $\infty$ -logoses and if  $\{F_i\}_{i \in I}$  is jointly conservative, then  $\mathcal{L}$  is an  $\infty$ -logos.  $\square$*

**Proposition 5.13.** *Let  $\mathcal{L}$  be an  $\infty$ -logos. The functor*

$$\uparrow^n \mathbf{U}(\mathcal{L}) : \mathcal{L}^{\text{op}} \rightarrow \uparrow^{n+1} \mathbf{Spc}$$

*defined by  $\uparrow^n \mathbf{U}(\mathcal{L})(A) = \mathbf{Obj}(\uparrow^n(\mathcal{L}/_A))$  belongs to  $\uparrow^{n+1} \mathcal{L}$ .*

*Proof.* By definition, the claim is true for  $n = 0$ . The other cases are because  $\uparrow^n : \mathbf{Pr}^{\mathbf{R}}(\mathfrak{U}) \rightarrow \mathbf{Pr}^{\mathbf{R}}(\uparrow^n \mathfrak{U})$  preserves small limits.  $\square$

**Definition 5.14.** Let  $\mathcal{L}$  be an  $\infty$ -logos. We say a map  $f : B \rightarrow A$  in  $\uparrow^{n+m} \mathcal{L}$  is  $\uparrow^n \mathfrak{U}$ -small if for any  $A' \in \mathcal{L}$  and any  $a : A' \rightarrow A$ , the base change  $a^* B$  belongs to  $\uparrow^n \mathcal{L}$ .

**Proposition 5.15.** *Let  $\mathcal{L}$  be an  $\infty$ -logos.  $\uparrow^n \mathbf{U}(\mathcal{L})$  is part of a classifying  $\uparrow^n \mathfrak{U}$ -small map  $\uparrow^n \tilde{\mathbf{U}}(\mathcal{L}) \rightarrow \uparrow^n \mathbf{U}(\mathcal{L})$ .*

*Proof.* By definition.  $\square$

**Proposition 5.16.** *For any object  $A$  of any presentable  $(\infty, 1)$ -category  $\mathcal{L}$ , we have*

$$\uparrow^n(\mathcal{L}/_A) \simeq (\uparrow^n \mathcal{L})/_A.$$

*Proof.* The equivalence  $\mathbf{Fun}((\mathcal{L}/_A)^{\text{op}}, \uparrow^n \mathbf{Spc}) \simeq \mathbf{Fun}(\mathcal{L}^{\text{op}}, \uparrow^n \mathbf{Spc})/_A$  is restricted to  $\uparrow^n(\mathcal{L}/_A) \simeq (\uparrow^n \mathcal{L})/_A$  because  $\mathcal{L}/_A \rightarrow \mathcal{L}$  creates colimits.  $\square$



**Definition 5.17.** A morphism of  $\infty$ -logoses is a *localization* if its right adjoint is fully faithful. For an  $\infty$ -logos  $\mathcal{L}$ , let  $\mathbf{Loc}(\mathcal{L})$  denote the full subcategory of  $(\mathbf{Logos}_{\mathcal{L}/})^{\text{op}}$  spanned by the localization morphisms  $\mathcal{L} \rightarrow \mathcal{K}$ . Note that  $\mathbf{Loc}(\mathcal{L})$  is a poset.

**Definition 5.18.** A morphism  $\mathcal{L} \rightarrow \mathcal{K}$  of  $\infty$ -logoses is *étale* if it is equivalent in  $\mathbf{Logos}_{\mathcal{L}/}$  to  $A^* = (A \times \_): \mathcal{L} \rightarrow \mathcal{L}/_A$  for some object  $A \in \mathcal{L}$ . We write  $\mathbf{Logos}_{\text{et}} \subset \mathbf{Logos}$  for the wide subcategory whose morphisms are the étale morphisms.

**Proposition 5.19.** *Localizations of  $\infty$ -logoses are stable under pushouts along étale morphisms. Precisely, for any localization  $F: \mathcal{L} \rightarrow \mathcal{K}$  of  $\infty$ -logos and for any object  $A \in \mathcal{L}$ , the diagram*

$$\begin{array}{ccc} \mathcal{L} & \xrightarrow{A^*} & \mathcal{L}/_A \\ F \downarrow & & \downarrow F/_A \\ \mathcal{K} & \xrightarrow{F(A)^*} & \mathcal{K}/_{F(A)} \end{array}$$

*is a pushout in  $\mathbf{Logos}$  and  $F/_A$  is a localization.*

*Proof.* The stability of étale morphisms under pushouts follows from the universal property of  $\mathcal{L}/_A$  [30, Proposition 6.3.5.5]. One can check that if the right adjoint of  $F$  is fully faithful, then so is the right adjoint of  $F/_A$ .  $\square$

## 6 Modalities in $\infty$ -logoses

We review *modalities* in  $\infty$ -logoses [3, 50]. We change some terminologies to make them match terminologies of modalities in homotopy type theory. Let us fix an  $\infty$ -logos  $\mathcal{L}$ .

**Notation 6.1.** Let  $R$  be a class of maps in  $\mathcal{L}$  closed under base change. For an object  $A \in \mathcal{L}$ , let  $R[A] \subset \mathcal{L}/_A$  denote the full subcategory spanned by the morphisms  $B \rightarrow A$  in  $R$ . By definition, the base change functor  $f^*: \mathcal{L}/_B \rightarrow \mathcal{L}/_A$  along a map  $f: A \rightarrow B$  is restricted to a functor  $R[B] \rightarrow R[A]$ .

**Definition 6.2.** By a *subuniverse* in  $\mathcal{L}$  we mean a local class  $R$  of maps in  $\mathcal{L}$ , that is,  $R$  is closed under base change and, for any small diagram  $A: I \rightarrow \mathcal{L}$ , a map  $f: B \rightarrow \text{colim}_{i \in I} A_i$  is in  $R$  if and only if every  $f|_{A_i}$  is. This is equivalent to that  $R$  is closed under base change and, for any map  $f: B \rightarrow A$  and any jointly surjective family of maps  $\{g_i: A'_i \rightarrow A\}_{i \in I}$ , the map  $f$  is in  $R$  if and only if every base change  $g_i^* f: g_i^* B \rightarrow A'_i$  is.

**Proposition 6.3.** *For a class  $R$  of maps in  $\mathcal{L}$  closed under base change, the following are equivalent:*

1.  $R$  is a subuniverse in  $\mathcal{L}$ ;

2. the functor  $\mathcal{L}^{\text{op}} \ni A \mapsto \mathbf{Obj}(R[A]) \in \mathbf{Spc}$  belongs to  $\uparrow \mathcal{L}$ .

*Proof.* Let  $A : I \rightarrow \mathcal{L}$  be a small diagram, and consider the following commutative diagram.

$$\begin{array}{ccc} \mathbf{Obj}(R[\text{colim}_{i \in I} A_i]) & \longrightarrow & \lim_{i \in I} \mathbf{Obj}(R[A_i]) \\ \downarrow & & \downarrow \\ \mathbf{Obj}(\mathcal{L}/_{\text{colim}_{i \in I} A_i}) & \xrightarrow{\simeq} & \lim_{i \in I} \mathbf{Obj}(\mathcal{L}/_{A_i}) \end{array}$$

The bottom functor is an equivalence by the definition of  $\infty$ -logoses. Then, the top functor is an equivalence if and only if this square is a pullback, which is equivalent to that a map  $f : B \rightarrow \text{colim}_{i \in I} A_i$  is in  $R$  if and only if every  $f|_{A_i}$  is.  $\square$

**Definition 6.4.** We say a subuniverse  $R$  in  $\mathcal{L}$  is *reflective* if the inclusion  $R[A] \rightarrow \mathcal{L}/_A$  has a reflector for any object  $A \in \mathcal{L}$  and if base changes commute with the reflectors. (A reflective subuniverse in  $\mathcal{L}$  is called a *reflective subfibration* in [50, 35].)

A class of maps with fiberwise reflectors stable under base change is automatically a subuniverse.

**Fact 6.5** ([50, Proposition 3.12]). *A class  $R$  of maps in  $\mathcal{L}$  closed under base change is local whenever the inclusion  $R[A] \rightarrow \mathcal{L}/_A$  has a reflector for any object  $A \in \mathcal{L}$  and base changes commute with the reflectors.*

**Definition 6.6.** Let  $f : A \rightarrow B$  and  $g : Y \rightarrow X$  be maps in  $\mathcal{L}$ . We write  $f \perp g$  and say  $f$  is *left orthogonal to  $g$*  or  $g$  is *right orthogonal to  $f$*  if the map

$$(f^*, g_*) : \mathbf{Map}(B, Y) \rightarrow \mathbf{Map}(A, Y) \times_{\mathbf{Map}(A, X)} \mathbf{Map}(B, X)$$

is an equivalence.

**Definition 6.7.** A *factorization system* in  $\mathcal{L}$  is a pair  $(L, R)$  of replete classes of maps in  $\mathcal{L}$  satisfying that  $f \perp g$  for all  $f \in L$  and  $g \in R$  and that every map in  $\mathcal{L}$  is factored as a map in  $L$  followed by a map in  $R$ . We call  $L$  the *left class* and  $R$  the *right class* of the factorization system.

**Fact 6.8** ([50, Theorem 4.8]). *For a class  $R$  of maps in  $\mathcal{L}$  closed under base change, the following are equivalent:*

1.  $R$  is the right class of a factorization system in  $\mathcal{L}$  whose left class is closed under base change.
2.  $R$  is a reflective subuniverse in  $\mathcal{L}$  and closed under composition.

**Definition 6.9.** A *modality* in  $\mathcal{L}$  is a class of maps in  $\mathcal{L}$  satisfying the equivalent conditions of Fact 6.8.

*Example 6.10* ([3, Example 3.2.2]). For  $m \geq -2$ , the class of  $m$ -truncated maps is the right class of a modality whose left class is the class of  $m$ -connected maps. Note that  $m$ -connected maps are called  $(m + 1)$ -connective maps in [30].

**Definition 6.11.** A modality  $(L, R)$  in  $\mathcal{L}$  is *accessible* (called *of small generation* in [3]) if there exists a small set  $\Sigma$  of maps in  $\mathcal{L}$  such that  $R$  is the class of maps right orthogonal to all base changes of maps in  $\Sigma$ . If this is the case, we say that  $(L, R)$  is *generated* by  $\Sigma$ .

**Definition 6.12.** A modality  $(L, R)$  in  $\mathcal{L}$  is *lex* if  $L$  is closed under diagonals.

**Notation 6.13.** Let  $\mathbf{SU}(\mathcal{L})$  denote the poset of subuniverses in  $\mathcal{L}$  ordered by inclusion. We define full subposets  $\mathbf{RSU}(\mathcal{L}) \subset \mathbf{SU}(\mathcal{L})$  of reflective ones,  $\mathbf{Mdl}(\mathcal{L}) \subset \mathbf{RSU}(\mathcal{L})$  of modalities,  $\mathbf{Lex}(\mathcal{L}) \subset \mathbf{Mdl}(\mathcal{L})$  of lex modalities, and  $\mathbf{AccMdl}(\mathcal{L}) \subset \mathbf{Mdl}(\mathcal{L})$  of accessible modalities. Let  $\mathbf{LAM}(\mathcal{L}) = \mathbf{Lex}(\mathcal{L}) \cap \mathbf{AccMdl}(\mathcal{L})$ .

**Construction 6.14.** The mapping  $\mathcal{L} \mapsto \mathbf{SU}(\mathcal{L})$  is covariant with respect to étale morphisms between  $\infty$ -logoses: for any étale morphism  $A^* : \mathcal{L} \rightarrow \mathcal{L}/_A$  and any a subuniverse  $R$  in  $\mathcal{L}$ , we define  $A^* \cdot R \in \mathbf{SU}(\mathcal{L}/_A)$  to be inverse image  $(A_!)^{-1}R$  where  $A_!$  is the domain projection  $\mathcal{L}/_A \rightarrow \mathcal{L}$ . By definition,  $(A^* \cdot R)[B] \simeq R[B]$  for  $B \in \mathcal{L}/_A$ , from which it follows that this covariant action of étale morphisms is restricted to actions on  $\mathbf{RSU}$ ,  $\mathbf{Mdl}$ , and  $\mathbf{Lex}$ . If  $R$  is an accessible modality generated by a set of maps  $\Sigma$ , then  $A^* \cdot R$  is generated by

$$\{f \in \mathbf{Map}_{\mathcal{L}/_A}((x, a \circ f), (y, a)) \mid (f : x \rightarrow y) \in \Sigma, a : y \rightarrow A\},$$

and thus the covariant action is also restricted to actions on  $\mathbf{AccMdl}$  and  $\mathbf{LAM}$ .

**Proposition 6.15.** *The map  $R \mapsto R[1]$  defines an equivalence of posets  $\mathbf{LAM}(\mathcal{L}) \simeq \mathbf{Loc}(\mathcal{L})$  natural in  $\mathcal{L} \in \mathbf{Logos}_{\text{et}}$ . Moreover, for any  $A \in \mathcal{L}$ , we have  $R[A] \simeq R[1]_{/F(A)}$  where  $F : \mathcal{L} \rightarrow R[1]$  is the reflector.*

*Proof.* The equivalence  $\mathbf{LAM}(\mathcal{L}) \simeq \mathbf{Loc}(\mathcal{L})$  is due to Anel et al. [3, Proposition 4.1.11 and Theorem 4.2.11]. For the naturality, let  $A^* : \mathcal{L} \rightarrow \mathcal{L}/_A$  be an étale morphism and let  $R \in \mathbf{LAM}(\mathcal{L})$ . Recall that  $A^* \cdot R[1] \simeq R[1]_{/F(A)}$  by Proposition 5.19 while  $(A^* \cdot R)[1] \simeq R[A]$  by definition. Thus, the naturality is equivalent to the last assertion, which is proved as  $R[A] \simeq R[F(A)]$  by [3, Theorem 4.1.8, 5] and then  $R[F(A)] \simeq R[1]_{/F(A)}$  by the left cancellability of  $R$  ([30, Proposition 5.2.8.6 (3)]).  $\square$

**Fact 6.16** ([3, Proposition 3.2.7]). *Both left and right classes of a modality are local.*

**Definition 6.17.** A set  $G$  of objects in  $\mathcal{L}$  is a set of *generators* if any object in  $\mathcal{L}$  is a small colimit of objects from  $G$ . Note that any presentable  $(\infty, 1)$ -category admits a small set of generators [30, Theorem 5.5.1.1 (6)]. By a  *$G$ -base change* we mean a base change along a map with domain in  $G$ .

**Definition 6.18.** A class of maps in  $\mathcal{L}$  is *saturated* if it contains all the equivalences and is closed under composition and colimits.

**Proposition 6.19.** Let  $\Sigma$  be a small set of maps in  $\mathcal{L}$ . For any small set of generators  $G$ , if  $\Sigma$  is closed under  $G$ -base change, then the class of maps right orthogonal to  $\Sigma$  is the right class of an accessible modality. Moreover, its left class is the smallest saturated class containing  $\Sigma$ .

*Proof.* This follows from [3, Proposition 3.2.18 and Theorem 3.2.20].  $\square$

## 7 Semantics of homotopy type theory

We review the semantics of homotopy type theory in  $\infty$ -logoses.

Let  $\mathcal{L}$  be an  $\infty$ -logos. The interpretation of type theory in  $\mathcal{L}$  is summarized as follows.

- A context  $\Gamma$  is interpreted as an object  $\llbracket \Gamma \rrbracket_{\mathcal{L}}$  in  $\uparrow^n \mathcal{L}$  for a sufficiently large  $n$ . A context morphism  $f : \Gamma \rightarrow \Delta$  is interpreted as a map  $\llbracket f \rrbracket_{\mathcal{L}} : \llbracket \Gamma \rrbracket_{\mathcal{L}} \rightarrow \llbracket \Delta \rrbracket_{\mathcal{L}}$ .
- A type  $\Gamma \vdash A$  **Type** is interpreted as a map  $\llbracket \Gamma \vdash A \rrbracket_{\mathcal{L}} \rightarrow \llbracket \Gamma \rrbracket_{\mathcal{L}}$  in  $\uparrow^n \mathcal{L}$  for a sufficiently large  $n$ . For a context morphism  $f : \Gamma' \rightarrow \Gamma$ , the interpretation of the substitution  $\Gamma' \vdash A[f]$  **Type** fits into the following pullback

$$\begin{array}{ccc} \llbracket \Gamma' \vdash A[f] \rrbracket_{\mathcal{L}} & \longrightarrow & \llbracket \Gamma \vdash A \rrbracket_{\mathcal{L}} \\ \downarrow & \lrcorner & \downarrow \\ \llbracket \Gamma' \rrbracket_{\mathcal{L}} & \xrightarrow{\llbracket f \rrbracket_{\mathcal{L}}} & \llbracket \Gamma \rrbracket_{\mathcal{L}}. \end{array}$$

- The empty context is interpreted as the final object, and a context extension  $(\Gamma, \mathbf{x} : A)$  is interpreted as the object  $\llbracket \Gamma \vdash A \rrbracket_{\mathcal{L}}$ .
- A term  $\Gamma \vdash a : A$  is interpreted as a section  $\llbracket \Gamma \vdash a : A \rrbracket_{\mathcal{L}}$  of  $\llbracket A \rrbracket_{\mathcal{L}} \rightarrow \llbracket \Gamma \rrbracket_{\mathcal{L}}$ . For a context morphism  $f : \Gamma' \rightarrow \Gamma$ , the interpretation of the substitution  $\Gamma' \vdash a[f] : A[f]$  coincides with the pullback of  $\llbracket \Gamma \vdash a : A \rrbracket_{\mathcal{L}}$  along  $\llbracket f \rrbracket_{\mathcal{L}}$ .

$$\begin{array}{ccccc} \llbracket \Gamma' \rrbracket_{\mathcal{L}} & \xrightarrow{\llbracket f \rrbracket_{\mathcal{L}}} & \llbracket \Gamma \rrbracket_{\mathcal{L}} & & \\ \downarrow & \dashrightarrow & \downarrow & \searrow & \\ \llbracket \Gamma' \vdash A[f] \rrbracket_{\mathcal{L}} & \xrightarrow{\quad} & \llbracket \Gamma \vdash A \rrbracket_{\mathcal{L}} & & \\ \downarrow & \lrcorner & \downarrow & & \\ \llbracket \Gamma' \rrbracket_{\mathcal{L}} & \xrightarrow{\llbracket f \rrbracket_{\mathcal{L}}} & \llbracket \Gamma \rrbracket_{\mathcal{L}} & & \end{array}$$

$\llbracket \Gamma' \vdash a[f] : A[f] \rrbracket_{\mathcal{L}}$  (dotted arrow from  $\llbracket \Gamma' \rrbracket_{\mathcal{L}}$  to  $\llbracket \Gamma \vdash A \rrbracket_{\mathcal{L}}$ )  
 $\llbracket \Gamma \vdash a : A \rrbracket_{\mathcal{L}}$  (arrow from  $\llbracket \Gamma \rrbracket_{\mathcal{L}}$  to  $\llbracket \Gamma \vdash A \rrbracket_{\mathcal{L}}$ )

- The universe  $\vdash \uparrow^n \mathcal{U} \text{ Type}$  is interpreted as  $\uparrow^n \mathbf{U}(\mathcal{L}) \in \uparrow^{n+1} \mathcal{L}$ . For a term  $\Gamma \vdash A : \uparrow^n \mathcal{U}$ , the corresponding type  $\Gamma \vdash A \text{ Type}$  is interpreted as the pullback of  $\uparrow^n \tilde{\mathbf{U}}(\mathcal{L})$  along  $\llbracket \Gamma \vdash A : \uparrow^n \mathcal{U} \rrbracket_{\mathcal{L}} : \llbracket \Gamma \rrbracket_{\mathcal{L}} \rightarrow \uparrow^n \mathbf{U}(\mathcal{L})$ .
- The interpretation of the unit type  $\Gamma \vdash \mathbf{1} \text{ Type}$  is an equivalence  $\llbracket \Gamma \vdash \mathbf{1} \rrbracket_{\mathcal{L}} \simeq \llbracket \Gamma \rrbracket_{\mathcal{L}}$ .
- The interpretation of a pair type  $\Gamma \vdash \sum_{x:A} B(x) \text{ Type}$  is the composite  $\llbracket \Gamma, x : A \vdash B(x) \rrbracket_{\mathcal{L}} \rightarrow \llbracket \Gamma \vdash A \rrbracket_{\mathcal{L}} \rightarrow \llbracket \Gamma \rrbracket_{\mathcal{L}}$ .
- The interpretation of a function type  $\Gamma \vdash \prod_{x:A} B(x) \text{ Type}$  is the pushforward of  $\llbracket \Gamma, x : A \vdash B(x) \rrbracket_{\mathcal{L}}$  along  $\llbracket \Gamma \vdash A \rrbracket_{\mathcal{L}} \rightarrow \llbracket \Gamma \rrbracket_{\mathcal{L}}$ .
- The interpretation of an identity type  $\Gamma \vdash a_1 =_A a_2 \text{ Type}$  fits into the following pullback

$$\begin{array}{ccc}
\llbracket \Gamma \vdash a_1 =_A a_2 \rrbracket_{\mathcal{L}} & \xrightarrow{\quad} & \llbracket \Gamma \vdash A \rrbracket_{\mathcal{L}} \\
\downarrow & \lrcorner & \downarrow \Delta \\
\llbracket \Gamma \rrbracket_{\mathcal{L}} & \xrightarrow{\langle \llbracket \Gamma \vdash a_1 : A \rrbracket_{\mathcal{L}}, \llbracket \Gamma \vdash a_2 : A \rrbracket_{\mathcal{L}} \rangle} & \llbracket \Gamma \vdash A \rrbracket_{\mathcal{L}} \times_{\llbracket \Gamma \rrbracket_{\mathcal{L}}} \llbracket \Gamma \vdash A \rrbracket_{\mathcal{L}}
\end{array}$$

This interpretation is justified as follows. By the result of Shulman [37], the  $\infty$ -logos is presented by a nice model category  $\mathcal{C}$  called a type-theoretic model topos.  $\mathcal{C}$  is in particular a type-theoretic model category [39] and admits interpretations of the unit type, pair types, function types, and identity types as explained in [39, Section 4.2]; see also [7, 5]. Furthermore, its universe enlargement  $\uparrow \mathcal{C}$  has a univalent fibration classifying small fibrations. By [41, Corollary 4.1], this fibration corresponds to the classifying small map  $\tilde{\mathbf{U}}(\mathcal{L}) \rightarrow \mathbf{U}(\mathcal{L})$  in  $\uparrow \mathcal{L}$ .

Note that the interpretation of higher inductive types with definitional computation rules on point constructors is *not* justified. This is because the existing semantics of higher inductive types with parameters in model categories [28] raises universe levels. For this reason, higher inductive types in our type theory are postulated as initial algebras internal to type theory for which computation rules hold only up to homotopy. The existence of initial algebras follows from presentability.

## 7.1 Kripke-Joyal semantics

Following the *Kripke-Joyal semantics* of higher-order logic in toposes [31, Section VI.6], we introduce a *forcing relation*.

**Notation 7.1.** Let  $\Gamma \vdash A \text{ Type}$  be a type in a context  $\Gamma$  and suppose that it is interpreted in a map in  $\uparrow^n \mathcal{L}$ . For an object  $X \in \mathcal{L}$  and a map  $c : X \rightarrow \llbracket \Gamma \rrbracket_{\mathcal{L}}$ , we define a space

$$X \Vdash_{\mathcal{L}} A \langle c \rangle$$

to be  $\mathbf{Map}_{(\uparrow^n \mathcal{L})/[\Gamma]_{\mathcal{L}}}(X, \llbracket A \rrbracket_{\mathcal{L}})$  where  $X$  is regarded as an object of  $(\uparrow^n \mathcal{L})/[\Gamma]_{\mathcal{L}}$  with the map  $c$ . Suppose  $\Gamma \equiv (\mathbf{x}_1 : A_1, \dots, \mathbf{x}_m : A_m)$  and let  $c_i$  be the projection of  $c$  to  $\llbracket \mathbf{x}_1 : A_1, \dots, \mathbf{x}_i : A_i \rrbracket_{\mathcal{L}}$ . In this case, instead of  $X \Vdash_{\mathcal{L}} A \langle c \rangle$  we write

$$X \Vdash_{\mathcal{L}} (A[\mathbf{x}_1 \equiv c_1, \dots, \mathbf{x}_m \equiv c_m])$$

where  $(A[\mathbf{x}_1 \equiv c_1, \dots, \mathbf{x}_m \equiv c_m])$  is a meta-level expression obtained from  $A$  by formally replacing all the free occurrences of  $\mathbf{x}_i$  by  $c_i$ . This notation is ambiguous because different  $A$  and  $c$  can yield the same meta-level expression, but in practice there would be no confusion. Note that, by definition, every  $c_i$  is a point of  $X \Vdash_{\mathcal{L}} (A[\mathbf{x}_1 \equiv c_1, \dots, \mathbf{x}_{i-1} \equiv c_{i-1}])$ . For a map  $u : X' \rightarrow X$  in  $\mathcal{L}$ , we write  $c_i|_{X'}$  instead of  $c_i \circ u$  when  $u$  is clear from the context. The precomposition with  $u$  induces a map  $(-)|_{X'} : (X \Vdash_{\mathcal{L}} A[\mathbf{x}_1 \equiv c_1, \dots, \mathbf{x}_m \equiv c_m]) \rightarrow (X' \Vdash_{\mathcal{L}} A[\mathbf{x}_1 \equiv c_1|_{X'}, \dots, \mathbf{x}_m \equiv c_m|_{X'}])$ .

*Example 7.2.*  $(X \Vdash_{\mathcal{L}} \uparrow^n \mathcal{U}) \simeq \mathbf{Map}_{\uparrow^{n+1} \mathcal{L}}(X, \uparrow^n \mathbf{U}(\mathcal{L})) \simeq \mathbf{Obj}(\uparrow^n(\mathcal{L}/X))$ . For a point  $A \in (X \Vdash_{\mathcal{L}} \uparrow^n \mathcal{U})$ , we write  $X.A$  for the corresponding object of  $\uparrow^n(\mathcal{L}/X) \simeq (\uparrow^n \mathcal{L})/X$ .

*Example 7.3.* Let  $A \in (X \Vdash_{\mathcal{L}} \mathcal{U})$  and  $B \in (X \Vdash_{\mathcal{L}} A \rightarrow \mathcal{U})$ . They determine a map  $(A, B) : X \rightarrow \llbracket \mathbf{A} : \mathcal{U}, \mathbf{B} : \mathbf{A} \rightarrow \mathcal{U} \rrbracket_{\mathcal{L}}$ , and  $X \Vdash_{\mathcal{L}} \prod_{\mathbf{x}:A} B(\mathbf{x})$  is an abbreviation of  $X \Vdash_{\mathcal{L}} (\prod_{\mathbf{x}:A} B(\mathbf{x})) \langle (A, B) \rangle$ . Observe that  $(X \Vdash_{\mathcal{L}} A \rightarrow \mathcal{U}) \simeq \mathbf{Map}_{\uparrow \mathcal{L}}(X.A, \mathbf{U}(\mathcal{L}))$ . Under this identification, we have  $(X \Vdash_{\mathcal{L}} \prod_{\mathbf{x}:A} B(\mathbf{x})) \simeq \mathbf{Map}_{\mathcal{L}/X.A}(X.A, X.A.B)$ . By Yoneda, it is equivalent to the space of families of dependent maps

$$\prod_{a \in (X' \Vdash_{\mathcal{L}} A|_{X'})} X' \Vdash_{\mathcal{L}} B|_{X'}(a)$$

indexed over  $X' \in \mathcal{L}/X$  equipped with certain coherence data. When  $B$  is 0-truncated, the coherence data coincide with the uniformity condition of Awodey, Gambino, and Hazratpour [6, Proposition 4.16].

*Example 7.4.* Let  $A, B \in (X \Vdash_{\mathcal{L}} \mathcal{U})$ . As a special case of Example 7.3,  $X \Vdash_{\mathcal{L}} A \rightarrow B$  is equivalent to the space of families of maps

$$(X' \Vdash_{\mathcal{L}} A|_{X'}) \rightarrow (X' \Vdash_{\mathcal{L}} B|_{X'})$$

natural in  $X' \in \mathcal{L}/X$ .

*Example 7.5.* Let  $A \in (X \Vdash_{\mathcal{L}} \mathcal{U})$  and  $B \in (X \Vdash_{\mathcal{L}} A \rightarrow \mathcal{U})$ . Then  $X \Vdash_{\mathcal{L}} \sum_{\mathbf{x}:A} B(\mathbf{x})$  is equivalent to the space of pairs  $(a, b)$  consisting of  $a \in (X \Vdash_{\mathcal{L}} A)$  and  $b \in (X \Vdash_{\mathcal{L}} B(a))$ .

*Example 7.6.*  $X \Vdash_{\mathcal{L}} \mathbf{1}$  is contractible.

*Example 7.7.* Let  $A \in (X \Vdash_{\mathcal{L}} \mathcal{U})$  and  $a_1, a_2 \in (X \Vdash_{\mathcal{L}} A)$ . Then  $X \Vdash_{\mathcal{L}} a_1 = a_2$  is equivalent to the space of homotopies from  $a_1$  to  $a_2$  in  $X \Vdash_{\mathcal{L}} A$ .

*Example 7.8.* Recall that  $\mathbf{lsContr}(A) \equiv \sum_{a:A} \prod_{\mathbf{x}:A} a = \mathbf{x}$ . Then, for any  $A \in (X \Vdash_{\mathcal{L}} \mathcal{U})$ , the space  $X \Vdash_{\mathcal{L}} \mathbf{lsContr}(A)$  is equivalent to the pairs  $(a, p)$  consisting of a point  $a \in (X \Vdash_{\mathcal{L}} A)$  and a homotopy  $p$  from  $a|_{X.A}$  to the identity in

$\mathbf{Map}_{\mathcal{L}/X}(X.A, X.A)$ . Such a pair is nothing but a retraction of the projection  $X.A \rightarrow X$  in  $\mathcal{L}/X$ . Therefore,  $X \Vdash_{\mathcal{L}} \mathbf{lsContr}(A)$  is  $(-1)$ -truncated, and it is inhabited if and only if  $X.A$  is the final object in  $\mathcal{L}/X$ , which is equivalent to that  $X' \Vdash_{\mathcal{L}} A|_{X'}$  is contractible for any  $X' \in \mathcal{L}/X$ .

Since many concepts in homotopy type theory are based on contractibility, the behavior of  $\Vdash_{\mathcal{L}}$  with respect to them is easily derived.

*Example 7.9.* Let  $A, B \in (X \Vdash_{\mathcal{L}} \mathcal{U})$  and  $f \in (X \Vdash_{\mathcal{L}} A \rightarrow B)$ . Then  $X \Vdash_{\mathcal{L}} \mathbf{lsEquiv}(f)$  is  $(-1)$ -truncated, and it is inhabited if and only if the map  $(X' \Vdash_{\mathcal{L}} A|_{X'}) \rightarrow (X' \Vdash_{\mathcal{L}} B|_{X'})$  is an equivalence for any  $X' \in \mathcal{L}/X$  if and only if the corresponding map  $X.A \rightarrow X.B$  in  $\mathcal{L}/X$  is an equivalence.

*Example 7.10.* Let  $A \in (X \Vdash_{\mathcal{L}} \mathcal{U})$ . Then  $X \Vdash_{\mathcal{L}} \mathbf{lsProp}(A)$  is  $(-1)$ -truncated, and it is inhabited if and only if  $X' \Vdash_{\mathcal{L}} A|_{X'}$  is  $(-1)$ -truncated for any  $X' \in \mathcal{L}/X$  if and only if  $X.A \in \mathcal{L}/X$  is  $(-1)$ -truncated. More generally, for  $m \geq -2$ , the space  $X \Vdash_{\mathcal{L}} \mathbf{lsTrunc}_m(A)$  is inhabited if and only if  $X.A \in \mathcal{L}/X$  is  $m$ -truncated.

The behavior of  $\Vdash_{\mathcal{L}}$  with respect to inductive types seems not straightforward. We will only provide the case of propositional truncation later in Proposition 8.9:  $X \Vdash_{\mathcal{L}} \|A\|$  is inhabited if and only if there exists a surjection  $X' \rightarrow X$  in  $\mathcal{L}$  such that  $X' \Vdash_{\mathcal{L}} A|_{X'}$  is inhabited.

## 8 Modalities in homotopy type theory and $\infty$ -logoses

We prove that the LAMs in the internal language of an  $\infty$ -logos  $\mathcal{L}$  correspond to the localizations of  $\mathcal{L}$ .

**Theorem 8.1.** *Let  $\mathcal{L}$  be an  $\infty$ -logos. For any object  $X \in \mathcal{L}$ , we have a natural equivalence of posets*

$$(X \Vdash_{\mathcal{L}} \mathbf{LAM}) \simeq \mathbf{Loc}(\mathcal{L}/X).$$

We prove Theorem 8.1 step by step. By Construction 6.14, we regard  $X \mapsto \mathbf{SU}(\mathcal{L}/X)$  and its variants as functors from  $\mathcal{L}^{\text{op}}$  to the category of posets. For any object  $X \in \mathcal{L}$ , we construct natural equivalences

$$(X \Vdash_{\mathcal{L}} \mathbf{SU}) \simeq \mathbf{SU}(\mathcal{L}/X) \quad (\text{Proposition 8.2})$$

$$(X \Vdash_{\mathcal{L}} \mathbf{RSU}) \simeq \mathbf{RSU}(\mathcal{L}/X) \quad (\text{Proposition 8.3})$$

$$(X \Vdash_{\mathcal{L}} \mathbf{Mdl}) \simeq \mathbf{Mdl}(\mathcal{L}/X) \quad (\text{Proposition 8.4})$$

$$(X \Vdash_{\mathcal{L}} \mathbf{Lex}) \simeq \mathbf{Lex}(\mathcal{L}/X) \quad (\text{Proposition 8.6})$$

$$(X \Vdash_{\mathcal{L}} \mathbf{AccMdl}) \simeq \mathbf{AccMdl}(\mathcal{L}/X) \quad (\text{Theorem 8.7})$$

Then Theorem 8.1 follows from Proposition 6.15.

All but the last equivalence are straightforward from the definition of the interpretation of type theory in  $\mathcal{L}$ .

**Proposition 8.2.** *We have a natural equivalence of posets  $(X \Vdash_{\mathcal{L}} \mathbf{SU}) \simeq \mathbf{SU}(\mathcal{L}/X)$ .*

*Proof.*  $X \Vdash_{\mathcal{L}} \mathbf{SU}$  is equivalent to the subspace of  $X \Vdash_{\mathcal{L}} \mathcal{U} \rightarrow \mathcal{U}_1$  consisting of those such that the corresponding object in  $(\uparrow \mathcal{L})_{/X \times \mathbf{U}(\mathcal{L})}$  is  $(-1)$ -truncated. That is,  $X \Vdash_{\mathcal{L}} \mathbf{SU}$  is the poset of subobjects of  $X \times \mathbf{U}(\mathcal{L}) \simeq \mathbf{U}(\mathcal{L}/X)$  in  $\uparrow(\mathcal{L}/X)$ . By the definition of  $\mathbf{U}(\mathcal{L}/X)$ , the subobjects of  $\mathbf{U}(\mathcal{L}/X)$  in  $\mathbf{Fun}((\mathcal{L}/X)^{\text{op}}, \uparrow \mathbf{Spc})$  are precisely the classes of maps in  $\mathcal{L}/X$  closed under base change. Such a subobject  $R \subset \mathbf{U}(\mathcal{L}/X)$  belongs to  $\uparrow(\mathcal{L}/X)$  if and only if it is a subuniverse in  $\mathcal{L}/X$  by Proposition 6.3. By construction, this correspondence respects ordering and is natural in  $X \in \mathcal{L}$ .  $\square$

For an element  $\mathfrak{m} \in (X \Vdash_{\mathcal{L}} \mathbf{SU})$ , let  $\mathbf{U}(\mathcal{L}/X)_{\mathfrak{m}}$  denote the corresponding subuniverse in  $\mathcal{L}/X$ .

**Proposition 8.3.** *An element  $\mathfrak{m} \in (X \Vdash_{\mathcal{L}} \mathbf{SU})$  belongs to  $X \Vdash_{\mathcal{L}} \mathbf{RSU}$  if and only if  $\mathbf{U}(\mathcal{L}/X)_{\mathfrak{m}}$  is reflective.*

*Proof.*  $\mathfrak{m}$  belongs to  $X \Vdash_{\mathcal{L}} \mathbf{RSU}$  if and only if we have some  $\circ \in (X \Vdash_{\mathcal{L}} \mathcal{U} \rightarrow \mathcal{U}_{\mathfrak{m}})$  and  $\eta \in (X \Vdash_{\mathcal{L}} \prod_{\mathbf{A}:\mathcal{U}} \mathbf{A} \rightarrow \circ \mathbf{A})$  such that  $X \Vdash_{\mathcal{L}} \prod_{\mathbf{A}:\mathcal{U}} \prod_{\mathbf{B}:\mathcal{U}_{\mathfrak{m}}} \mathbf{lsEquiv}(\lambda f. f \circ \eta(\mathbf{A}))$  is inhabited.  $\circ$  is a family of maps

$$\circ_{X'} : \mathbf{Obj}((\mathcal{L}/X)_{/X'}) \rightarrow \mathbf{Obj}(\mathbf{U}(\mathcal{L}/X)_{\mathfrak{m}}[X'])$$

natural in  $X' \in \mathcal{L}/X$ , and  $\eta$  is a family of maps

$$\eta_{X'}(A) : A \rightarrow \circ_{X'} A$$

for all  $A \in (\mathcal{L}/X)_{/X'}$ , natural in  $X' \in \mathcal{L}/X$ . The required condition is equivalent to that for any  $X' \in \mathcal{L}/X$ ,  $A \in (\mathcal{L}/X)_{/X'}$ , and  $B \in \mathbf{U}(\mathcal{L}/X)_{\mathfrak{m}}[X']$ , the precomposition with  $\eta_{X'}(A)$

$$\eta_{X'}(A)^* : \mathbf{Map}_{(\mathcal{L}/X)_{/X'}}(\circ_{X'} A, B) \rightarrow \mathbf{Map}_{(\mathcal{L}/X)_{/X'}}(A, B)$$

is an equivalence. These data are precisely reflections  $(\mathcal{L}/X)_{/X'} \rightarrow \mathbf{U}(\mathcal{L}/X)_{\mathfrak{m}}[X']$  for all  $X' \in \mathcal{L}/X$  that commute with base change.  $\square$

**Proposition 8.4.** *An element  $\mathfrak{m} \in (X \Vdash_{\mathcal{L}} \mathbf{RSU})$  belongs to  $X \Vdash_{\mathcal{L}} \mathbf{Mdl}$  if and only if  $\mathbf{U}(\mathcal{L}/X)_{\mathfrak{m}}$  is a modality.*

*Proof.* This is because pair types are interpreted as composition.  $\square$

**Lemma 8.5.** *For any  $\mathfrak{m} \in (X \Vdash_{\mathcal{L}} \mathbf{Mdl})$  and any  $A \in (X \Vdash_{\mathcal{L}} \mathcal{U})$ , the space  $X \Vdash_{\mathcal{L}} \mathbf{In}_{\perp \mathfrak{m}}(A)$  is inhabited if and only if the map  $X.A \rightarrow X$  belongs to the left class of the modality  $\mathbf{U}(\mathcal{L}/X)_{\mathfrak{m}}$  in  $\mathcal{L}/X$ .*

*Proof.*  $X \Vdash_{\mathcal{L}} \mathbf{In}_{\perp \mathfrak{m}}(A)$  is inhabited if and only if the map  $X.(\circ_{\mathfrak{m}} A) \rightarrow X$  is an equivalence. Then the claim follows because this map is part of the factorization  $X.A \rightarrow X.(\circ_{\mathfrak{m}} A) \rightarrow X$  of  $X.A \rightarrow X$  into a map in the left class of  $\mathbf{U}(\mathcal{L}/X)_{\mathfrak{m}}$  followed by a map in the right class of  $\mathbf{U}(\mathcal{L}/X)_{\mathfrak{m}}$ .  $\square$

**Proposition 8.6.** *An element  $\mathfrak{m} \in (X \Vdash_{\mathcal{L}} \mathbf{Mdl})$  belongs to  $X \Vdash_{\mathcal{L}} \mathbf{Lex}$  if and only if the modality  $\mathbf{U}(\mathcal{L}/X)_{\mathfrak{m}}$  in  $\mathcal{L}/X$  is lex.*



*Proof.* By Lemma 8.5. □

It remains to establish a correspondence between type-theoretic accessibility and categorical accessibility.

**Theorem 8.7.** *Let  $\mathcal{L}$  be an  $\infty$ -logos. For any object  $X \in \mathcal{L}$ , the equivalence in Proposition 8.2 is restricted to an equivalence*

$$(X \Vdash_{\mathcal{L}} \text{AccMdl}) \simeq \mathbf{AccMdl}(\mathcal{L}/_A).$$

Recall that a modality  $\mathfrak{m}$  in type theory is accessible if there *merely* exists a null generator  $\mu$  such that  $\mathfrak{Null}(\mu) = \mathfrak{m}$ , that is,  $\|\sum_{\mu: \text{NullGen}} \mathfrak{Null}(\mu) = \mathfrak{m}\|$  is inhabited. It is thus necessary to understand the Kripke-Joyal semantics for propositional truncation.

**Lemma 8.8.** *Let  $\mathcal{L}$  be an  $\infty$ -logos and  $m \geq -2$ . The element  $\llbracket \mathfrak{Tr}_m \rrbracket_{\mathcal{L}} \in (\mathbf{1} \Vdash_{\mathcal{L}} \text{Mdl})$  corresponds to the modality in  $\mathcal{L}$  whose right class of  $m$ -truncated maps and whose left class is the class of  $m$ -connected maps.*

*Proof.* By Example 7.10,  $\llbracket \mathfrak{Tr}_m \rrbracket_{\mathcal{L}}$  corresponds to the modality in  $\mathcal{L}$  consisting of the  $m$ -truncated maps. By Example 6.10, the left class is the class of  $m$ -connected maps. □

**Proposition 8.9.** *Let  $A \in (X \Vdash_{\mathcal{L}} \uparrow^n \mathcal{U})$ . Then  $X \Vdash_{\mathcal{L}} \|A\|$  is inhabited if and only if there exists a surjection  $X' \rightarrow X$  in  $\mathcal{L}$  such that  $X' \Vdash_{\mathcal{L}} A|_{X'}$  is inhabited.*

*Proof.* The “if” part is immediate because surjections are left orthogonal to monomorphisms. For the “only if” part, let  $a \in (X \Vdash_{\mathcal{L}} \|A\|)$ . By Lemma 8.8, the unit  $X.A \rightarrow X.\|A\|$  is surjective in  $\mathcal{L}/_X$ . Take the pullback

$$\begin{array}{ccc} Y & \cdots \twoheadrightarrow & X.A \\ f \downarrow \dashv & \lrcorner & \downarrow \\ X & \xrightarrow{a} & X.\|A\|. \end{array}$$

Since surjections are stable under base change, the map  $f$  is surjective. However,  $Y \in \uparrow^n \mathcal{L}$  is not necessarily in  $\mathcal{L}$ .

To fix this issue, take a small set of generators  $G$  for  $\mathcal{L}$ . Then we have the surjection

$$\left( \coprod_{z \in G} \coprod_{u: z \rightarrow Y} z \right) \twoheadrightarrow Y.$$

Let  $H$  be the image of the map

$$\left( \coprod_{z \in G} \mathbf{Map}_{\uparrow^n \mathcal{L}}(z, Y) \right) \ni (z, u) \mapsto (z, f \circ u) \in \left( \coprod_{z \in G} \mathbf{Map}_{\mathcal{L}}(z, X) \right).$$

Since  $\mathcal{L}$  is locally small,  $H$  is small. We have the following commutative diagram.

$$\begin{array}{ccc} \coprod_{z \in G} \coprod_{u: z \rightarrow Y} z & \twoheadrightarrow & Y \\ \downarrow & & \downarrow f \\ \coprod_{(z, u) \in H} z & \longrightarrow & X \end{array}$$

The bottom map is a surjection since the top and right maps are.  $X' = \coprod_{(z,u) \in H} z$  belongs to  $\mathcal{L}$  as  $H$  is small. By construction, the precomposite with each coproduct inclusion  $z \rightarrow X' \rightarrow X$  factors through  $f$ , and thus  $X' \rightarrow X$  factors through  $Y$ . By composing with the map  $Y \rightarrow X.A$ , we get an inhabitant of  $X' \Vdash_{\mathcal{L}} A|_{X'}$ .  $\square$

We show that type-theoretic accessibility implies categorical accessibility.

**Construction 8.10.** Let  $\mu \in (X \Vdash_{\mathcal{L}} \text{NullGen})$  and let  $G$  be a small set of generators for  $\mathcal{L}$ . We define a small set  $\Xi_{\mu}^G$  of maps in  $\mathcal{L}/X$  by

$$\Xi_{\mu}^G = \{X'.Z_{\mu|_{X'}}(i) \rightarrow X' \mid X' \in G_{/X}, i \in (X' \Vdash_{\mathcal{L}} I_{\mu|_{X'}})\}.$$

By construction,  $\Xi_{\mu}^G$  is closed under  $G$ -base change.

**Lemma 8.11.** *For any  $\mu \in (X \Vdash_{\mathcal{L}} \text{NullGen})$  and any small set of generators  $G$  for  $\mathcal{L}$ , the modality in  $\mathcal{L}/X$  associated to  $\mathfrak{Null}(\mu) \in (X \Vdash_{\mathcal{L}} \text{Mdl})$  is generated by  $\Xi_{\mu}^G$ .*

*Proof.* Let  $\mathfrak{m} = \mathfrak{Null}(\mu)$ . It suffices to show that for any  $X' \in \mathcal{L}/X$  and any  $A \in (X' \Vdash_{\mathcal{L}} \mathcal{U})$ , the space  $X' \Vdash_{\mathcal{L}} \text{In}_{\mathfrak{m}|_{X'}}(A)$  is inhabited if and only if the map  $X'.A \rightarrow X'$  is right orthogonal to all the maps in  $\Xi_{\mu}^G$ . The former condition is equivalent to that for any  $X'' \in \mathcal{L}/X'$ , any  $i \in (X'' \Vdash_{\mathcal{L}} I_{\mu|_{X''}})$ , and any  $a \in (X'' \Vdash_{\mathcal{L}} A|_{X''})$ , the precomposition map

$$\mathbf{Map}_{\mathcal{L}/X'}(X'', X'.A) \rightarrow \mathbf{Map}_{\mathcal{L}/X'}(X''.Z_{\mu|_{X''}}(i), X'.A)$$

is an equivalence. Since  $G$  is a set of generators for  $\mathcal{L}$ , we may restrict the range of  $X''$  in the latter condition to  $G_{/X'}$ , but then the condition is equivalent to that  $X'.A \rightarrow X'$  is right orthogonal to all the maps in  $\Xi_{\mu}^G$ .  $\square$

**Lemma 8.12.** *For any  $\mathfrak{m} \in (X \Vdash_{\mathcal{L}} \text{AccMdl})$ , the modality in  $\mathcal{L}/X$  associated to  $\mathfrak{m}$  is accessible.*

*Proof.* Take a small set of generators  $G$  for  $\mathcal{L}$ . By Proposition 8.9, there exist a surjection  $u : X' \rightarrow X$  in  $\mathcal{L}$  and a point  $\mu \in (X' \Vdash_{\mathcal{L}} \text{NullGen})$  such that  $X' \Vdash_{\mathcal{L}} \mathfrak{Null}(\mu) = \mathfrak{m}|_{X'}$  is inhabited. Then we have the modality  $\Phi'$  in  $\mathcal{L}/X'$  generated by  $\Xi_{\mu}^G$ . Let  $\Phi$  be the factorization system in  $\mathcal{L}/X$  generated by the image  $u_! \Xi_{\mu}^G$  of  $\Xi_{\mu}^G$  by the postcomposition functor  $u_! : \mathcal{L}/X' \rightarrow \mathcal{L}/X$ . Since  $\Xi_{\mu}^G$  is closed under  $G$ -base change, so is  $u_! \Xi_{\mu}^G$ , and thus  $\Phi$  is an accessible modality in  $\mathcal{L}/X$  by Proposition 6.19. Then, for a map  $f : B \rightarrow A$  in  $\mathcal{L}/X$

$$\begin{aligned} & f \text{ belongs to the right class of } \Phi \\ \Leftrightarrow & \{u_! \dashv u^*\} \\ & u^* f \text{ belongs to the right class of } \Phi' \\ \Leftrightarrow & \{\text{Lemma 8.11}\} \\ & u^* f \text{ belongs to } \mathbf{U}(\mathcal{L}/X')_{\mathfrak{m}|_{X'}} \end{aligned}$$

$$\Leftrightarrow \begin{cases} \{\text{Fact 6.16}\} \\ f \text{ belongs to } \mathbf{U}(\mathcal{L}/X)_{\mathbf{m}}. \end{cases}$$

Hence, the modality in  $\mathcal{L}/X$  associated to  $\mathbf{m}$  coincides with the accessible modality  $\Phi$ .  $\square$

We then show the converse.

**Construction 8.13.** Let  $G$  be a small set of generators for  $\mathcal{L}$  and let  $\Sigma$  be a set of maps in  $\mathcal{L}/X$  closed under  $G$ -base change. We define a map  $\mu_\Sigma$  in  $\mathcal{L}/X$  to be the coproduct of all the maps in  $\Sigma$  and regard it as a point  $\mu_\Sigma \in (X \Vdash_{\mathcal{L}} \text{NullGen})$ .

**Lemma 8.14.** *Any accessible modality in  $\mathcal{L}/X$  is induced by some  $\mathbf{m} \in (X \Vdash_{\mathcal{L}} \text{AccMdl})$ .*

*Proof.* Take a small set of generators  $G$  for  $\mathcal{L}$ . We may assume that the modality is generated by a set  $\Sigma$  of maps in  $\mathcal{L}/X$  closed under  $G$ -base change. By Lemma 8.11, it suffices to show that  $\Xi_{\mu_\Sigma}^G$  generates the same modality in  $\mathcal{L}/X$  as  $\Sigma$ . Let  $\Sigma^s$  denote the smallest saturated class containing  $\Sigma$ . Note that the closure under  $G$ -base change implies that  $\Sigma^s$  is closed under arbitrary base change by Proposition 6.19. Also note that, by Fact 6.16, to show that a map belongs to  $\Sigma^s$ , it suffices to show that its base change along an arbitrary map with domain in  $G$  belongs to  $\Sigma^s$ .

We show that  $\Sigma^s = (\Xi_{\mu_\Sigma}^G)^s$ . Let  $f : B \rightarrow A$  be a map in  $\Sigma$  and  $a : X' \rightarrow A$  a map with  $X' \in G$ . By construction, we have the coproduct inclusion

$$\begin{array}{ccc} B & \xrightarrow{\quad} & X.(\sum_{i:I_{\mu_\Sigma}} Z_{\mu_\Sigma}(i)) \\ f \downarrow & & \downarrow \\ A & \xrightarrow{\quad} & X.I_{\mu_\Sigma} \end{array}$$

which is a pullback by Fact 5.11. Then the base change  $a^*f : a^*B \rightarrow X'$  belongs to  $\Xi_{\mu_\Sigma}^G$  by definition. Thus,  $\Sigma^s \subset (\Xi_{\mu_\Sigma}^G)^s$ . Conversely, let  $\pi(\mu_\Sigma)_i : X'.Z_{\mu_\Sigma|X'}(i) \rightarrow X'$  be a map in  $\Xi_{\mu_\Sigma}^G$  where  $X' \in G/X$  and  $i \in (X' \Vdash_{\mathcal{L}} I_{\mu_\Sigma|X'})$ . For a map  $f : B \rightarrow A$  in  $\Sigma$ , we regard it as a map in  $\mathcal{L}/X.I_{\mu_\Sigma}$  with the coproduct inclusion  $A \rightarrow X.I_{\mu_\Sigma}$  and define  $f_i : B_i \rightarrow A_i$  to be the base change of  $f$  along  $i : X' \rightarrow X.I_{\mu_\Sigma}$ . By Fact 5.11,  $\pi(\mu_\Sigma)_i$  is the coproduct of  $f_i$ 's for all  $f : B \rightarrow A$  in  $\Sigma$ . Since each  $f_i$  belongs to  $\Sigma^s$ , so does  $(\mu_\Sigma)_i$ . Hence,  $(\Xi_{\mu_\Sigma}^G)^s \subset \Sigma^s$ .  $\square$

*Proof of Theorem 8.7.* By Lemmas 8.12 and 8.14.  $\square$

*Proof of Theorem 8.1.* By Propositions 6.15 and 8.6 and Theorem 8.7.  $\square$

Let us check the localizations corresponding to the open and closed modalities.

*Example 8.15.* Let  $P \in \mathcal{L}$  be a  $(-1)$ -truncated object. Then the étale morphism  $P^* : \mathcal{L} \rightarrow \mathcal{L}/P$  is a localization. Indeed, since the left adjoint of  $P^*$  is fully faithful as  $P$  is  $(-1)$ -truncated, the right adjoint of  $P^*$  is also fully faithful. We call  $P^* : \mathcal{L} \rightarrow \mathcal{L}/P$  the *open localization associated to  $P$* . The composite of  $P^*$  and its right adjoint is the exponential  $(-)^P$ , from which it follows that  $P^*$  corresponds to the open modality  $\mathfrak{Op}(P) \in (\mathbf{1} \Vdash_{\mathcal{L}} \text{LAM})$ . The *closed localization associated to  $P$*  is the reflector onto the full subcategory  $\{A \in \mathcal{L} \mid P \times A \simeq P\} \subset \mathcal{L}$  and corresponds to the closed modality  $\mathfrak{Cl}(P) \in (\mathbf{1} \Vdash_{\mathcal{L}} \text{LAM})$ .

## 9 $(\infty, 2)$ -category theory

The purpose of this section is to construct the *oplax limit* of a diagram of  $(\infty, 1)$ -categories indexed over an  $(\infty, 2)$ -category and show that  $\infty$ -logoses are closed under oplax limits.

### 9.1 The language of $(\infty, 2)$ -category theory

We formulate concepts in  $(\infty, 2)$ -category theory using the  $(\infty, 1)$ -category  $\mathbf{2-Cat}$  of  $(\infty, 2)$ -categories axiomatized and proved to be equivalent to various models of  $(\infty, 2)$ -categories by Barwick and Schommer-Pries [9].

**Definition 9.1.** A (strict) 2-category  $\mathcal{C}$  is said to be *gaunt* if only invertible 1-cells and 2-cells are the identities.

*Example 9.2.* The *walking  $m$ -cell*  $\mathbf{G}_m$  for  $0 \leq m \leq 2$  is the 2-category freely generated by a single  $m$ -cell and is gaunt.

Among Barwick and Schommer-Pries's axioms, the following are important to us.

**Fact 9.3** ([9, Basic Data]).  *$\mathbf{2-Cat}$  is presentable and contains finitely presentable gaunt 2-categories as a full subcategory.*

**Fact 9.4** ([9, Axiom C.2]).  *$\{\mathbf{G}_0, \mathbf{G}_1, \mathbf{G}_2\}$  is a set of generators for  $\mathbf{2-Cat}$ .*

**Fact 9.5** ([9, Axiom C.3]). *The  $(\infty, 1)$ -category  $\mathbf{2-Cat}$  (more generally  $\mathbf{2-Cat}/_{\mathbf{G}_m}$  for  $0 \leq m \leq 2$ ) is cartesian closed. For  $(\infty, 2)$ -categories  $\mathcal{C}$  and  $\mathcal{D}$ , let  $\mathbf{Fun}(\mathcal{C}, \mathcal{D})$  denote the exponential in  $\mathbf{2-Cat}$ .*

**Fact 9.6** ([9, Axiom C.4]).  *$\mathbf{G}_0, \mathbf{G}_1,$  and  $\mathbf{G}_2$  satisfy certain pushout formulas. We will recall them when needed.*

Fact 9.4 is particularly useful and mostly used in the following form.

**Proposition 9.7.** *Let  $\Phi$  be a class of small  $(\infty, 2)$ -categories. If  $\Phi$  is closed under small colimits (that is,  $\text{colim}_{i \in I} \mathcal{C}_i \in \Phi$  whenever  $\mathcal{C}_i \in \Phi$  for every  $i \in I$ , for any small diagram  $\mathcal{C} : I \rightarrow \mathbf{2-Cat}$ ) and if  $\Phi$  contains  $\mathbf{G}_0, \mathbf{G}_1,$  and  $\mathbf{G}_2$ , then  $\Phi$  contains all small  $(\infty, 2)$ -categories.  $\square$*

Let us fix terminology and notation.

**Definition 9.8.** Objects in  $\mathbf{2-Cat}$  are called  $(\infty, 2)$ -categories. Morphisms in  $\mathbf{2-Cat}$  are called *functors*. For  $(\infty, 2)$ -categories  $\mathcal{C}$  and  $\mathcal{D}$ , morphisms in  $\mathbf{Fun}(\mathcal{C}, \mathcal{D})$  are called *natural transformations*.

Constructions 9.9, 9.10 and 9.12 below are easily justified by using the 2-fold complete Segal spaces model [8].

**Construction 9.9.** Let  $\mathcal{C}$  be an  $(\infty, 2)$ -category. We define the space  $\mathbf{Obj}(\mathcal{C})$  of *objects* or *0-cells* in  $\mathcal{C}$  to be  $\mathbf{Map}_{\mathbf{2-Cat}}(\mathbf{G}_0, \mathcal{C})$ . For a pair of objects  $(x, y)$ , an  $(\infty, 1)$ -category  $\mathbf{Map}_{\mathcal{C}}(x, y)$  called the *mapping  $(\infty, 1)$ -category* is constructed. It is defined by the pullbacks

$$\begin{array}{ccc} \mathbf{Obj}(\mathbf{Map}_{\mathcal{C}}(x, y)) & \longrightarrow & \mathbf{Map}_{\mathbf{2-Cat}}(\mathbf{G}_1, \mathcal{C}) \\ \downarrow & \lrcorner & \downarrow (\text{dom, cod}) \\ \mathbf{1} & \xrightarrow{(x, y)} & \mathbf{Obj}(\mathcal{C}) \times \mathbf{Obj}(\mathcal{C}) \end{array}$$

and

$$\begin{array}{ccc} \mathbf{Map}_{\mathbf{Map}_{\mathcal{C}}(x, y)}(u, v) & \longrightarrow & \mathbf{Map}_{\mathbf{2-Cat}}(\mathbf{G}_2, \mathcal{C}) \\ \downarrow & \lrcorner & \downarrow (\text{dom, cod}) \\ \mathbf{1} & \xrightarrow{(u, v)} & \mathbf{Obj}(\mathbf{Map}_{\mathcal{C}}(x, y)) \times \mathbf{Obj}(\mathbf{Map}_{\mathcal{C}}(x, y)). \end{array}$$

$\mathcal{C}$  also has a functorial composition operator between its mapping  $(\infty, 1)$ -categories. Objects in  $\mathbf{Map}_{\mathcal{C}}(x, y)$  are called *morphisms* or *1-cells*, and morphisms in  $\mathbf{Map}_{\mathcal{C}}(x, y)$  are called *2-morphisms* or *2-cells*.

**Construction 9.10.** The  $(\infty, 1)$ -core  $\mathbf{Core}_{(1)}(\mathcal{C})$  of an  $(\infty, 2)$ -category  $\mathcal{C}$  is the  $(\infty, 1)$ -category defined by  $\mathbf{Obj}(\mathbf{Core}_{(1)}(\mathcal{C})) = \mathbf{Obj}(\mathcal{C})$  and  $\mathbf{Map}_{\mathbf{Core}_{(1)}(\mathcal{C})}(x, y) = \mathbf{Obj}(\mathbf{Map}_{\mathcal{C}}(x, y))$ . This defines a functor  $\mathbf{Core}_{(1)} : \mathbf{2-Cat} \rightarrow \mathbf{Cat}$ . It is shown that  $\mathbf{Core}_{(1)}$  has a fully faithful left adjoint, and thus we regard  $\mathbf{Cat}$  as a coreflective full subcategory of  $\mathbf{2-Cat}$ . An  $(\infty, 2)$ -category  $\mathcal{C}$  is an  $(\infty, 1)$ -category if and only if it is *locally discrete* in the sense that  $\mathbf{Map}_{\mathcal{C}}(x, y)$  is an  $\infty$ -groupoid for any  $x, y \in \mathcal{C}$ .

**Definition 9.11.** Let  $\mathcal{C}$  be an  $(\infty, 2)$ -category. A *locally full subcategory* of  $\mathcal{C}$  is an  $(\infty, 2)$ -category  $\mathcal{C}'$  equipped with a functor  $F : \mathcal{C}' \rightarrow \mathcal{C}$  such that  $\mathbf{Obj}(\mathcal{C}') \rightarrow \mathbf{Obj}(\mathcal{C})$  is mono and  $\mathbf{Map}_{\mathcal{C}'}(x, y) \rightarrow \mathbf{Map}_{\mathcal{C}}(F(x), F(y))$  is fully faithful for any  $x, y \in \mathcal{C}'$ . A locally full subcategory of  $\mathcal{C}$  is usually specified by a class  $\mathcal{C}'_0$  of 0-cells in  $\mathcal{C}$  and a class  $\mathcal{C}'_1$  of 1-cells in  $\mathcal{C}$  between objects in  $\mathcal{C}'_0$  such that  $\mathcal{C}'_1$  contains all the equivalences between objects in  $\mathcal{C}'_0$  and is closed under composition.

The *unicity theorem* [9, Theorem 7.3] asserts that there are exactly  $(\mathbb{Z}/2\mathbb{Z})^2$  automorphisms on  $\mathbf{2-Cat}$ . Those automorphisms are *opposite* constructions.

**Construction 9.12.** Let  $\mathcal{C}$  be an  $(\infty, 2)$ -category.  $(\infty, 2)$ -categories  $\mathcal{C}^{\text{op}(1)}$  and  $\mathcal{C}^{\text{op}(2)}$  are defined by  $\mathbf{Obj}(\mathcal{C}^{\text{op}(1)}) = \mathbf{Obj}(\mathcal{C}^{\text{op}(2)}) = \mathbf{Obj}(\mathcal{C})$  and

$$\mathbf{Map}_{\mathcal{C}^{\text{op}(1)}}(x, y) = \mathbf{Map}_{\mathcal{C}}(y, x)$$

$$\mathbf{Map}_{\mathcal{C}^{\mathrm{op}(2)}}(x, y) = \mathbf{Map}_{\mathcal{C}}(x, y)^{\mathrm{op}}.$$

We abbreviate  $(\mathcal{C}^{\mathrm{op}(1)})^{\mathrm{op}(2)} \simeq (\mathcal{C}^{\mathrm{op}(2)})^{\mathrm{op}(1)}$  as  $\mathcal{C}^{\mathrm{op}(1,2)}$ .

The following is an  $(\infty, 2)$ -categorical version of the fact that a natural transformation is invertible if it is point-wise invertible. It is proved without using a specific model.

**Proposition 9.13.** *For any  $(\infty, 2)$ -categories  $\mathcal{C}$  and  $\mathcal{D}$ , the restriction functor  $\mathbf{Core}_{(1)}(\mathbf{Fun}(\mathcal{C}, \mathcal{D})) \rightarrow \mathbf{Core}_{(1)}(\mathbf{Fun}(\mathbf{Obj}(\mathcal{C}), \mathcal{D}))$  is conservative.*

*Proof.* Let  $\Phi$  be the class of  $(\infty, 2)$ -categories  $\mathcal{C}$  such that the functor  $\mathbf{Core}_{(1)}(\mathbf{Fun}(\mathcal{C}, \mathcal{D})) \rightarrow \mathbf{Core}_{(1)}(\mathbf{Fun}(\mathbf{Obj}(\mathcal{C}), \mathcal{D}))$  is conservative for any  $(\infty, 2)$ -category  $\mathcal{D}$ . We first show that  $\Phi$  is closed under colimits. For a functor  $\mathcal{C} : I \rightarrow 2\text{-Cat}$ , we have the following commutative diagram.

$$\begin{array}{ccc} \mathbf{Core}_{(1)}(\mathbf{Fun}(\mathrm{colim}_{i \in I} \mathcal{C}_i, \mathcal{D})) & \longrightarrow & \mathbf{Core}_{(1)}(\mathbf{Fun}(\mathbf{Obj}(\mathrm{colim}_{i \in I} \mathcal{C}_i), \mathcal{D})) \\ \simeq \downarrow & & \downarrow \\ \lim_{i \in I} \mathbf{Core}_{(1)}(\mathbf{Fun}(\mathcal{C}_i, \mathcal{D})) & \longrightarrow & \lim_{i \in I} \mathbf{Core}_{(1)}(\mathbf{Fun}(\mathbf{Obj}(\mathcal{C}_i), \mathcal{D})) \end{array}$$

The left functor is an equivalence as  $\mathbf{Core}_{(1)}$  preserves limits. If every  $\mathcal{C}_i$  belongs to  $\Phi$ , then the bottom functor is conservative, and so is the top.

It remains to show that  $\Phi$  contains  $\mathbf{G}_0$ ,  $\mathbf{G}_1$ , and  $\mathbf{G}_2$ . Note that when  $\mathcal{C}$  is locally discrete, we have  $\mathbf{Core}_{(1)}(\mathbf{Fun}(\mathcal{C}, \mathcal{D})) \simeq \mathbf{Fun}(\mathcal{C}, \mathbf{Core}_{(1)}(\mathcal{D}))$ , and  $(\infty, 1)$ -category theory applies. The only non-trivial case is when  $\mathcal{C} = \mathbf{G}_2$ . We recall the following pushout in  $2\text{-Cat}$  [9, Axiom C.4].

$$\begin{array}{ccc} \mathbf{G}_2 & \longrightarrow & \mathbf{G}_1 +_{\mathbf{G}_0} \mathbf{G}_2 \\ \downarrow & \lrcorner & \downarrow \\ \mathbf{G}_2 +_{\mathbf{G}_0} \mathbf{G}_1 & \longrightarrow & \mathbf{G}_2 \times \mathbf{G}_1 \end{array}$$

$\mathbf{G}_2 \times \mathbf{G}_1$  should be a cylinder

$$\begin{array}{ccc} \bullet & \longrightarrow & \bullet \\ \left( \begin{array}{c} \Downarrow \\ \Rightarrow \\ \Downarrow \end{array} \right) & & \left( \begin{array}{c} \Downarrow \\ \Rightarrow \\ \Downarrow \end{array} \right) \\ \bullet & \longrightarrow & \bullet \end{array}$$

and the above pushout formula asserts that this is the case because the cylinder

is obtained by gluing the lower-left part  $\mathbf{G}_2 +_{\mathbf{G}_0} \mathbf{G}_1 = \left( \begin{array}{ccc} & & \bullet \\ & \left( \begin{array}{c} \Downarrow \\ \Rightarrow \\ \Downarrow \end{array} \right) & \\ \bullet & \longrightarrow & \bullet \end{array} \right)$

and the upper-right part  $\mathbf{G}_1 +_{\mathbf{G}_0} \mathbf{G}_2 = \left( \begin{array}{ccc} \bullet & \longrightarrow & \bullet \\ & \left( \begin{array}{c} \Downarrow \\ \Rightarrow \\ \Downarrow \end{array} \right) & \\ & & \bullet \end{array} \right)$ . A functor  $F \in$

$\mathbf{Map}_{2\text{-Cat}}(\mathbf{G}_1, \mathbf{Fun}(\mathbf{G}_2, \mathcal{D})) \simeq \mathbf{Map}_{2\text{-Cat}}(\mathbf{G}_2 \times \mathbf{G}_1, \mathcal{D})$  is thus a diagram in  $\mathcal{D}$  of the form

$$\begin{array}{ccc} F(0,0) & \xrightarrow{F(0,01)} & F(0,1) \\ \left(\begin{array}{c} \Downarrow \\ \Rightarrow \\ \Downarrow \end{array}\right) & & \left(\begin{array}{c} \Downarrow \\ \Rightarrow \\ \Downarrow \end{array}\right) \\ F(1,0) & \xrightarrow{F(1,01)} & F(1,1). \end{array}$$

When the morphisms  $F(0,01)$  and  $F(1,01)$  are invertible, then one can construct an inverse of  $F$  in  $\mathbf{Fun}(\mathbf{G}_2, \mathcal{D})$ .  $\square$

## 9.2 Scaled simplicial sets

We review one of models for  $(\infty, 2)$ -categories, *scaled simplicial sets* [29]. This model is convenient for presenting  $(\infty, 2)$ -categories by combinatorial data, provides computation of colimits of  $(\infty, 2)$ -categories, and gives the  $(\infty, 2)$ -category of  $(\infty, 1)$ -categories a useful universal property.

**Definition 9.14.** A *scaled simplicial set* is a simplicial set  $A$  equipped with a class of 2-simplices called *thin* 2-simplices such that all the degenerate 2-simplices are thin.

A scaled simplicial set  $A$  is thought of as a presentation of an  $(\infty, 2)$ -category: the 0-cells are the 0-simplices of  $A$ ; the 1-cells are generated by 1-simplices of  $A$ ; the 2-cells are generated by 2-simplices of  $A$  in the following direction;

$$\begin{array}{ccc} a_0 & \xrightarrow{\quad} & a_2 \\ & \searrow & \nearrow \\ & a_1 & \end{array}$$

thin 2-simplices are made invertible 2-cells; higher simplices presents homotopies filling certain diagrams.

**Proposition 9.15.** *There is a model structure on the category of scaled simplicial sets that presents 2-Cat.*

*Proof.* See [29] for the model structure and comparison with other models for  $(\infty, 2)$ -categories. Barwick and Schommer-Pries [9] show that the  $(\infty, 1)$ -category presented by this model structure indeed satisfies their axioms.  $\square$

*Example 9.16* ([30, Example 1.1.5.9]). Let  $I$  be a (decidable) poset and we regard it as a scaled simplicial set with no non-degenerate thin 2-simplex. It presents a gaunt 2-category  $\langle I \rangle$  defined as follows. The objects of  $\langle I \rangle$  are the elements of  $I$ . The mapping category  $\mathbf{Map}_{\langle I \rangle}(i, j)$  for  $i, j \in I$  is the poset of totally ordered subsets  $J \subset I$  with least element  $i$  and largest element  $j$ . A morphism from  $i$  to  $j$  is thus a chain  $i = k_0 < k_1 < \dots < k_n = j$ . It then follows that  $\mathbf{Core}_{(1)}(\langle I \rangle)$  is the free category over the strict ordering relation on  $I$ . For any  $i < j$  in  $I$ , the chain  $(i < j)$  is the initial object in  $\mathbf{Map}_{\langle I \rangle}(i, j)$ .

We can calculate colimits in  $2\text{-Cat}$  via homotopy colimits of scaled simplicial sets. We do not need much details, but a useful consequence is the following.

**Corollary 9.17.** *For any functor  $\mathcal{C} : I \rightarrow 2\text{-Cat}$ , the map  $\text{colim}_{i \in I} \mathbf{Obj}(\mathcal{C}_i) \rightarrow \mathbf{Obj}(\text{colim}_{i \in I} \mathcal{C}_i)$  is surjective.  $\square$*

Lurie [29, Section 4.5] shows that the  $(\infty, 2)$ -category of  $(\infty, 1)$ -categories is presented by the scaled simplicial set classifying *locally cocartesian fibrations*.

**Definition 9.18.** Let  $A$  be a scaled simplicial set and  $B$  a simplicial set. A map  $p : B \rightarrow A$  of simplicial sets is a *locally cocartesian fibration* if the following are satisfied:

1.  $p$  is an inner fibration of simplicial sets;
2. for every 1-simplex  $a : \Delta^1 \rightarrow A$ , the base change  $a^*p : a^*B \rightarrow \Delta^1$  is a cocartesian fibration of simplicial sets;
3. for every thin 2-simplex  $a : \Delta^2 \rightarrow A$ , the base change  $a^*p$  is a cocartesian fibration of simplicial sets.

**Fact 9.19** ([29, Corollary 4.5.7]). *There exists a universal locally cocartesian fibration with small fibers  $\pi : \widetilde{\mathbf{Cat}}^{\text{sc}} \rightarrow \mathbf{Cat}^{\text{sc}}$  in the following sense:*

1.  $\mathbf{Cat}^{\text{sc}}$  is a large fibrant scaled simplicial set,  $\widetilde{\mathbf{Cat}}^{\text{sc}}$  is a large simplicial set, and  $\pi$  is a locally cocartesian fibration whose fibers are small;
2. for any locally cocartesian fibration with small fibers  $p$ , there exists a unique, up to homotopy, homotopy pullback from  $p$  to  $\pi$ .

**Construction 9.20.** Let  $\mathbf{Cat}^{(2)} \in \uparrow 2\text{-Cat}$  be the  $(\infty, 2)$ -category presented by  $\mathbf{Cat}^{\text{sc}}$ . By the construction of  $\mathbf{Cat}^{\text{sc}}$  [29, Definition 4.5.1], we have  $\mathbf{Obj}(\mathbf{Cat}^{(2)}) \simeq \mathbf{Obj}(\mathbf{Cat})$  and  $\mathbf{Map}_{\mathbf{Cat}^{(2)}}(\mathcal{C}, \mathcal{D}) \simeq \mathbf{Fun}(\mathcal{C}, \mathcal{D})$ .

**Notation 9.21.** The constructions of  $\mathbf{Cat}^{\text{sc}}$  and  $\mathbf{Cat}^{(2)}$  are parameterized by a universe. We use the universe enlargement notation  $\uparrow^n \mathbf{Cat}^{\text{sc}}$  and  $\uparrow^n \mathbf{Cat}^{(2)}$  for those constructions with respect to a larger universe.

The notion of locally cocartesian fibration is, however, specific to the scaled simplicial sets model. A more model-independent notion is as follows.

**Definition 9.22.** A functor  $P : \mathcal{D} \rightarrow \mathcal{C}$  between  $(\infty, 2)$ -categories is a *1-cocartesian 2-right fibration* if the following conditions are satisfied:

1.  $P$  is *locally a right fibration* in the sense that for any objects  $x, y \in \mathcal{D}$ , the functor  $P : \mathbf{Map}_{\mathcal{D}}(x, y) \rightarrow \mathbf{Map}_{\mathcal{C}}(P(x), P(y))$  is a right fibration;
2.  $\mathbf{Core}_{(1)}(P) : \mathbf{Core}_{(1)}(\mathcal{D}) \rightarrow \mathbf{Core}_{(1)}(\mathcal{C})$  is a cocartesian fibration.

By duality, 1-(cocartesian/cartesian) 2-(right/left) fibrations are defined. (They are called *inner/outer cocartesian/cartesian fibrations* in [15].)



*Remark 9.23.* Assuming Item 1 in Definition 9.22, any cocartesian morphism  $u : x \rightarrow y$  in  $\mathbf{Core}_{(1)}(\mathcal{D})$  is also cocartesian in the  $\mathbf{Cat}$ -enriched sense: for any object  $z \in \mathcal{D}$ , the square

$$\begin{array}{ccc} \mathbf{Map}_{\mathcal{D}}(y, z) & \xrightarrow{-\circ u} & \mathbf{Map}_{\mathcal{D}}(x, z) \\ P \downarrow & & \downarrow P \\ \mathbf{Map}_{\mathcal{C}}(P(y), P(z)) & \xrightarrow{-\circ P(u)} & \mathbf{Map}_{\mathcal{C}}(P(x), P(z)) \end{array}$$

is a pullback in  $\mathbf{Cat}$ , because  $\mathbf{Obj} : \mathbf{Cat} \rightarrow \mathbf{Spc}$  reflects pullbacks of right fibrations.

**Proposition 9.24.**  $\mathbf{Cat}^{(2)}$  is part of a universal 1-cocartesian 2-right fibration with small fibers  $\widetilde{\mathbf{Cat}}^{(2),\mathbf{R}} \rightarrow \mathbf{Cat}^{(2)}$ . For a functor  $\mathcal{C} : I \rightarrow \mathbf{Cat}^{(2)}$ , let  $\mathbf{El}_I(\mathcal{C}) \rightarrow I$  denote the corresponding 1-cocartesian 2-right fibration.

*Proof.* This follows from the equivalence between locally cocartesian fibrations and inner cartesian fibrations of categories enriched over marked simplicial sets given by Gagna, Harpaz, and Lanari [15, Propositions 2.4.1 and 3.1.3].  $\square$

An object in  $\widetilde{\mathbf{Cat}}^{(2),\mathbf{R}}$  is a pair  $(\mathcal{C}, x)$  consisting of an  $(\infty, 1)$ -category  $\mathcal{C}$  and an object  $x \in \mathcal{C}$ . A morphism  $(\mathcal{C}, x) \rightarrow (\mathcal{D}, y)$  in  $\widetilde{\mathbf{Cat}}^{(2),\mathbf{R}}$  is a pair  $(F, u)$  consisting of a functor  $F : \mathcal{C} \rightarrow \mathcal{D}$  and a morphism  $u : F(x) \rightarrow y$ . A 2-morphism  $(F, u) \Rightarrow (G, v) : (\mathcal{C}, x) \rightarrow (\mathcal{D}, y)$  in  $\widetilde{\mathbf{Cat}}^{(2),\mathbf{R}}$  is a pair  $(\sigma, p)$  consisting of a natural transformation  $\sigma : F \Rightarrow G$  and a path  $p : u \sim v \circ \sigma_x$ .

For the purpose of Section 9.3, we introduce a variant of  $\widetilde{\mathbf{Cat}}^{(2),\mathbf{R}}$ .

**Construction 9.25.** The equivalence of  $(\infty, 1)$ -categories  $\mathbf{Cat} \ni \mathcal{C} \mapsto \mathcal{C}^{\text{op}} \in \mathbf{Cat}$  extends to an equivalence of  $(\infty, 2)$ -categories

$$(-)^{\text{op}} : \mathbf{Cat}^{(2)} \simeq (\mathbf{Cat}^{(2)})^{\text{op}(2)}.$$

Let  $\widetilde{\mathbf{Cat}}^{(2),\mathbf{L}} = ((-)^{\text{op}})^*(\widetilde{\mathbf{Cat}}^{(2),\mathbf{R}})^{\text{op}(2)}$ . By construction, the functor  $(\widetilde{\mathbf{Cat}}^{(2),\mathbf{L}})^{\text{op}(1,2)} \rightarrow (\mathbf{Cat}^{(2)})^{\text{op}(1,2)}$  classifies 1-cartesian 2-right fibrations with small fibers. For a functor  $\mathcal{C} : I^{\text{op}(1,2)} \rightarrow \mathbf{Cat}^{(2)}$ , let  $\mathbf{El}_I(\mathcal{C}) \rightarrow I$  denote the corresponding 1-cartesian 2-right fibration.

In other words,  $\widetilde{\mathbf{Cat}}^{(2),\mathbf{L}}$  is the fiberwise opposite of  $\widetilde{\mathbf{Cat}}^{(2),\mathbf{R}}$ . Thus, the objects of  $\widetilde{\mathbf{Cat}}^{(2),\mathbf{L}}$  are the same as  $\widetilde{\mathbf{Cat}}^{(2),\mathbf{R}}$ , but a morphism  $(\mathcal{C}, x) \rightarrow (\mathcal{D}, y)$  in  $\widetilde{\mathbf{Cat}}^{(2),\mathbf{L}}$  is a pair  $(F, u)$  consisting of a functor  $F : \mathcal{C} \rightarrow \mathcal{D}$  and a morphism  $u : y \rightarrow F(x)$ .

### 9.3 Oplax limits

*Oplax limits* in general  $(\infty, 2)$ -categories are defined by Gagna, Harpaz, and Lanari [15]. In this paper, we only need oplax limits of  $(\infty, 1)$ -categories which have the following simple construction [15, Example 5.3.12].

**Construction 9.26.** Let  $I$  be a small  $(\infty, 2)$ -category and  $\mathcal{C} : I^{\text{op}(1,2)} \rightarrow \mathbf{Cat}^{(2)}$  a functor. The *oplax limit* of  $\mathcal{C}$  is defined to be the pullback

$$\begin{array}{ccc} \text{opLaxLim}_{i \in I} \mathcal{C}_i & \dashrightarrow & \mathbf{Fun}(I, (\widetilde{\mathbf{Cat}}^{(2),L})^{\text{op}(1,2)}) \\ \downarrow & \lrcorner & \downarrow \\ \mathbf{G}_0 & \xrightarrow{c} & \mathbf{Fun}(I, (\mathbf{Cat}^{(2)})^{\text{op}(1,2)}). \end{array}$$

Equivalently, it is the pullback

$$\begin{array}{ccc} \text{opLaxLim}_{i \in I} \mathcal{C}_i & \dashrightarrow & \mathbf{Fun}(I, \mathbf{El}_I(\mathcal{C})) \\ \downarrow & \lrcorner & \downarrow \\ \mathbf{G}_0 & \xrightarrow{\text{id}_I} & \mathbf{Fun}(I, I). \end{array}$$

Any functor  $F : J \rightarrow I$  induces a functor  $F^* : \text{opLaxLim}_{i \in I} \mathcal{C}_i \rightarrow \text{opLaxLim}_{j \in J} \mathcal{C}_{F(j)}$ .

*Remark 9.27.*  $\text{opLaxLim}_{i \in I} \mathcal{C}_i$  is small and locally discrete. Indeed, the class of small  $(\infty, 2)$ -categories  $I$  such that  $\text{opLaxLim}_{i \in I} \mathcal{C}_i$  is small and locally discrete for any  $\mathcal{C} : I^{\text{op}(1,2)} \rightarrow \mathbf{Cat}^{(2)}$  is closed under small colimits, and the cases when  $I = \mathbf{G}_0, \mathbf{G}_1, \mathbf{G}_2$  are directly calculated in Examples 9.29 to 9.31 below. Hence, we regard  $\text{opLaxLim}_{i \in I} \mathcal{C}_i$  as a small  $(\infty, 1)$ -category.

Concretely, an object  $x$  in  $\text{opLaxLim}_{i \in I} \mathcal{C}_i$  consists of: an object  $x_i \in \mathcal{C}_i$  for any object  $i \in I$ ; a morphism  $x_\alpha : x_i \rightarrow \mathcal{C}_\alpha(x_j)$  for any morphism  $\alpha : i \rightarrow j$  in  $I$ ; some coherence data. A morphism  $u : x \rightarrow y$  in  $\text{opLaxLim}_{i \in I} \mathcal{C}_i$  consists of: a morphism  $u_i : x_i \rightarrow y_i$  for any object  $i \in I$ ; a homotopy  $u_\alpha$  filling the square

$$\begin{array}{ccc} x_i & \xrightarrow{u_i} & y_i \\ x_\alpha \downarrow & & \downarrow y_\alpha \\ \mathcal{C}_\alpha(x_j) & \xrightarrow{\mathcal{C}_\alpha(u_j)} & \mathcal{C}_\alpha(y_j) \end{array}$$

for any morphism  $\alpha : i \rightarrow j$  in  $I$ ; some coherence data.

*Example 9.28.* When  $I$  is an  $\infty$ -groupoid, we have  $\text{opLaxLim}_{i \in I} \mathcal{C}_i \simeq \prod_{i \in I} \mathcal{C}_i$ . For a general  $I$ , we have the forgetful functor  $\text{opLaxLim}_{i \in I} \mathcal{C}_i \rightarrow \text{opLaxLim}_{i \in \mathbf{Obj}(I)} \mathcal{C}_i \simeq \prod_{i \in I} \mathcal{C}_i$  which is conservative by Proposition 9.13.

*Example 9.29.* When  $I = \mathbf{G}_0$ , a functor  $\mathbf{G}_0^{\text{op}(1,2)} \rightarrow \mathbf{Cat}^{(2)}$  corresponds to an  $(\infty, 1)$ -category  $\mathcal{C}$ , and its oplax limit is  $\mathcal{C}$  itself.

*Example 9.30.* When  $I = \mathbf{G}_1$ , a functor  $\mathcal{C} : \mathbf{G}_1^{\text{op}(1,2)} \rightarrow \mathbf{Cat}^{(2)}$  corresponds to a functor  $F : \mathcal{C}_1 \rightarrow \mathcal{C}_0$ . Its oplax limit is the  $(\infty, 1)$ -category of triples  $(x_0, x_1, u)$  consisting of objects  $x_0 \in \mathcal{C}_0$  and  $x_1 \in \mathcal{C}_1$  and a morphism  $u : x_0 \rightarrow F(x_1)$ . In other words, we have the following pullback.

$$\begin{array}{ccc} \text{opLaxLim}_{i \in \mathbf{G}_1} \mathcal{C}_i & \longrightarrow & \mathcal{C}_0^{\rightarrow} \\ \downarrow & \lrcorner & \downarrow \text{cod} \\ \mathcal{C}_1 & \xrightarrow{F} & \mathcal{C}_0 \end{array}$$

This oplax limit is called the *Artin gluing* for  $F$  and denoted by  $\mathbf{Gl}(F)$ .

*Example 9.31.* When  $I = \mathbf{G}_2$ , a functor  $\mathcal{C} : \mathbf{G}_2^{\text{op}(1,2)} \rightarrow \mathbf{Cat}^{(2)}$  corresponds to a natural transformation  $\sigma : F_1 \Rightarrow F_0 : \mathcal{C}_1 \rightarrow \mathcal{C}_0$ . Since  $(\widetilde{\mathbf{Cat}}^{(2),L})^{\text{op}(1,2)} \rightarrow (\mathbf{Cat}^{(2)})^{\text{op}(1,2)}$  is locally a right fibration, a section of it over  $\mathcal{C}$  is completely determined by the restriction along the codomain inclusion  $\mathbf{G}_1 \rightarrow \mathbf{G}_2$ . Therefore,  $\text{opLaxLim}_{i \in \mathbf{G}_2} \mathcal{C}_i \simeq \mathbf{Gl}(F_1)$ .

*Example 9.32.* Let  $J$  be a small  $(\infty, 1)$ -category and  $I : J \rightarrow 2\text{-Cat}$  a functor. For any functor  $\mathcal{C} : (\text{colim}_{j \in J} I_j)^{\text{op}(1,2)} \rightarrow \mathbf{Cat}^{(2)}$ , we have a canonical equivalence

$$\text{opLaxLim}_{i \in \text{colim}_{j \in J} I_j} \mathcal{C}_i \simeq \lim_{j \in J} \text{opLaxLim}_{i \in I_j} \mathcal{C}_{I_j(i)}.$$

Hence, arbitrary oplax limits are constructed from Examples 9.29 to 9.31 using small limits.

We show that  $\infty$ -logoses are closed under oplax limits. For this, we consider a weaker notion of morphism of  $\infty$ -logoses.

**Notation 9.33.** We define  $\mathbf{Logos}_{\text{LexAcc}}^{(2)} \subset \uparrow \mathbf{Cat}^{(2)}$  to be the locally full subcategory spanned by the  $\infty$ -logoses and the lex, accessible functors between  $\infty$ -logoses.

**Theorem 9.34.** *Let  $I$  be a small  $(\infty, 2)$ -category and  $\mathcal{L} : I^{\text{op}(1,2)} \rightarrow \mathbf{Logos}_{\text{LexAcc}}^{(2)}$  a functor. Then  $\text{opLaxLim}_{i \in I} \mathcal{L}_i$  is an  $\infty$ -logos. Moreover, the forgetful functor  $\text{opLaxLim}_{i \in I} \mathcal{L}_i \rightarrow \prod_{i \in I} \mathcal{L}_i$  preserves small colimits and finite limits.*

**Fact 9.35** ([30, Proposition 6.3.2.3]). **Logos**  $\subset \uparrow \mathbf{Cat}$  is closed under small limits.

**Lemma 9.36.** *Let  $F : \mathcal{L}_1 \rightarrow \mathcal{L}_0$  be a lex, accessible functor between  $\infty$ -logoses. Then  $\mathbf{Gl}(F)$  is an  $\infty$ -logos, and the projections  $\mathbf{Gl}(F) \rightarrow \mathcal{L}_0$  and  $\mathbf{Gl}(F) \rightarrow \mathcal{L}_1$  preserve small colimits and finite limits.*

*Proof.* We use a characterization of presentability: an  $(\infty, 1)$ -category is presentable if and only if it is *accessible* and has small colimits [30, Definition 5.5.0.1]. It follows from [30, Propositions 5.4.4.3 and 5.4.6.6] that  $\mathbf{Gl}(F)$  is accessible. By construction,  $\mathbf{Gl}(F)$  is the  $(\infty, 1)$ -category of triples  $(A_0, A_1, f)$  consisting of objects  $A_0 \in \mathcal{L}_0$  and  $A_1 \in \mathcal{L}_1$  and a map  $f : A_0 \rightarrow F(A_1)$ .

It follows from this description that the projection  $\mathbf{Gl}(F) \rightarrow \mathcal{L}_0 \times \mathcal{L}_1$  creates small colimits and finite limits. In particular,  $\mathbf{Gl}(F)$  admits small colimits and thus is presentable. Since the projection  $\mathbf{Gl}(F) \rightarrow \mathcal{L}_0 \times \mathcal{L}_1$  is conservative by Example 9.28,  $\mathbf{Gl}(F)$  is an  $\infty$ -logos by Corollary 5.12.  $\square$

*Proof of Theorem 9.34.* Let  $\Phi$  be the class of small  $(\infty, 2)$ -categories  $I$  such that for any functor  $\mathcal{L} : I^{\text{op}(1,2)} \rightarrow \mathbf{Logos}_{\text{LexAcc}}^{(2)}$ , the oplax limit  $\text{opLaxLim}_{i \in I} \mathcal{L}_i$  is an  $\infty$ -logos and the forgetful functor  $\text{opLaxLim}_{i \in I} \mathcal{L}_i \rightarrow \prod_{i \in I} \mathcal{L}_i$  preserves small colimits and finite limits. It is enough to show that  $\Phi$  is closed under small colimits and contains  $\mathbf{G}_0$ ,  $\mathbf{G}_1$ , and  $\mathbf{G}_2$ .

Let  $I : J \rightarrow 2\text{-Cat}$  be a functor from a small  $(\infty, 1)$ -category  $J$  and suppose that every  $I_j$  belongs to  $\Phi$ . Let  $\mathcal{L} : (\text{colim}_{j \in J} I_j)^{\text{op}(1,2)} \rightarrow \mathbf{Logos}_{\text{LexAcc}}^{(2)}$  be a functor. As in Example 9.32, we have

$$\text{opLaxLim}_{i \in \text{colim}_{j \in J} I_j} \mathcal{L}_i \simeq \lim_{j \in J} \text{opLaxLim}_{i \in I_j} \mathcal{L}_{\iota_j(i)}.$$

Since  $I_j \in \Phi$ , small colimits and finite limits in  $\text{opLaxLim}_{i \in I_j} \mathcal{L}_{\iota_j(i)}$  are computed in  $\prod_{i \in I_j} \mathcal{L}_{\iota_j(i)}$ . Thus, for any morphism  $j_1 \rightarrow j_2$  in  $J$ , the functor  $\text{opLaxLim}_{i \in I_{j_2}} \mathcal{L}_{\iota_{j_2}(i)} \rightarrow \text{opLaxLim}_{i \in I_{j_1}} \mathcal{L}_{\iota_{j_1}(i)}$  preserves small colimits and finite limits. Then, by Fact 9.35,  $\lim_{j \in J} \text{opLaxLim}_{i \in I_j} \mathcal{L}_{\iota_j(i)}$  is an  $\infty$ -logos. Consider the following commutative square.

$$\begin{array}{ccc} \text{opLaxLim}_{i \in \text{colim}_{j \in J} I_j} \mathcal{L}_i & \xrightarrow{\simeq} & \lim_{j \in J} \text{opLaxLim}_{i \in I_j} \mathcal{L}_{\iota_j(i)} \\ \downarrow & & \downarrow \\ \prod_{i \in \mathbf{Obj}(\text{colim}_{j \in J} I_j)} \mathcal{L}_i & \longrightarrow & \lim_{j \in J} \prod_{i \in \mathbf{Obj}(I_j)} \mathcal{L}_{\iota_j(i)} \end{array}$$

We have seen that the top functor is an equivalence. The right functor preserves small colimits and finite limits as every  $I_j$  belongs to  $\Phi$ . The bottom functor is equivalent to the restriction along  $\text{colim}_{j \in J} \mathbf{Obj}(I_j) \rightarrow \mathbf{Obj}(\text{colim}_{j \in J} I_j)$  and thus preserves small colimits and finite limits. By Corollary 9.17, the bottom functor is conservative. We thus conclude that the left functor preserves small colimits and finite limits. Hence,  $\text{colim}_{j \in J} I_j$  belongs to  $\Phi$ .

$\mathbf{G}_0$  belongs to  $\Phi$  by Example 9.29.  $\mathbf{G}_1$  belongs to  $\Phi$  by Example 9.30 and Lemma 9.36.  $\mathbf{G}_2$  belongs to  $\Phi$  by Example 9.31 and by the case of  $\mathbf{G}_1$ .  $\square$

## 9.4 Oplax natural transformations

Oplax limits are related to *oplax natural transformations*.

**Construction 9.37** ([16, Definition 2.1]). Let  $A$  and  $B$  be scaled simplicial sets. The *Gray product*  $A \otimes B$  is the scaled simplicial set whose underlying simplicial set is the cartesian product of  $A$  and  $B$  and whose 2-simplex  $(a, b) : \Delta^2 \rightarrow A \times B$  is thin if both  $a$  and  $b$  are thin and either  $a$  degenerates along  $\Delta^{\{1,2\}}$  or  $b$  degenerates along  $\Delta^{\{0,1\}}$ .

**Fact 9.38** ([16, Theorem 2.14]). *The Gray product is part of a left Quillen bifunctor on scaled simplicial sets. Consequently, it induces a functor*

$$- \otimes - : 2\text{-Cat} \times 2\text{-Cat} \rightarrow 2\text{-Cat}$$

*preserving small colimits on each variable.*

**Construction 9.39.** By Fact 9.38 and by the adjoint functor theorem, for any  $(\infty, 2)$ -category  $\mathcal{C}$ , the functors  $(- \otimes \mathcal{C})$  and  $(\mathcal{C} \otimes -)$  have right adjoints  $\mathbf{Fun}(\mathcal{C}, -)_{\text{Lax}}$  and  $\mathbf{Fun}(\mathcal{C}, -)_{\text{opLax}}$ , respectively.

*Example 9.40.*  $\mathcal{C} \otimes \mathbf{G}_0 \simeq \mathcal{C}$ , and thus  $\mathbf{Obj}(\mathbf{Fun}(\mathcal{C}, \mathcal{D})_{\text{opLax}}) \simeq \mathbf{Map}_{2\text{-Cat}}(\mathcal{C}, \mathcal{D})$ . Dually,  $\mathbf{Obj}(\mathbf{Fun}(\mathcal{C}, \mathcal{D})_{\text{Lax}}) \simeq \mathbf{Map}_{2\text{-Cat}}(\mathcal{C}, \mathcal{D})$ .

*Example 9.41.* Let  $\mathcal{C}$  be a scaled simplicial set and  $u : x \rightarrow y$  a 1-simplex in  $\mathcal{C}$ . Consider the following 2-simplices in  $\mathcal{C} \otimes \Delta^1$ .

$$\begin{array}{ccc} (x, 0) & \xrightarrow{(x, 0 \leq 1)} & (x, 1) \\ (u, 0) \downarrow & \searrow^{(u, 0 \leq 1)} & \downarrow (u, 1) \\ (y, 0) & \xrightarrow{(y, 0 \leq 1)} & (y, 1) \end{array} \quad (4)$$

By definition, the lower 2-simplex is thin, but the upper one is not (unless  $u$  is degenerate). Hence, these 2-simplices compose and yields a 2-cell

$$\begin{array}{ccc} (x, 0) & \xrightarrow{(x, 0 \leq 1)} & (x, 1) \\ (u, 0) \downarrow & \nearrow & \downarrow (u, 1) \\ (y, 0) & \xrightarrow{(y, 0 \leq 1)} & (y, 1) \end{array} \quad (5)$$

in the  $(\infty, 2)$ -category presented by  $\mathcal{C} \otimes \Delta^1$ . Then, a 1-cell  $\sigma : F \rightarrow G$  in  $\mathbf{Fun}(\mathcal{C}, \mathcal{D})_{\text{opLax}}$  assigns: a 1-cell  $\sigma_x : F(x) \rightarrow G(x)$  to every 0-cell  $x \in \mathcal{C}$ ; a 2-cell

$$\begin{array}{ccc} F(x) & \xrightarrow{\sigma_x} & G(x) \\ F(u) \downarrow & \searrow^{\sigma_u} & \downarrow G(u) \\ F(y) & \xrightarrow{\sigma_y} & G(y) \end{array}$$

to every 1-cell  $u : x \rightarrow y$ ; and coherence data to higher cells. Such a structure is called an *oplax natural transformation from  $F$  to  $G$* . Dually, 1-cells in  $\mathbf{Fun}(\mathcal{C}, \mathcal{D})_{\text{Lax}}$  are called *lax natural transformations*. For a lax natural transformation  $\sigma : F \rightarrow G$ , the 2-cell  $\sigma_u$  is in the opposite direction  $G(u) \circ \sigma_x \Rightarrow \sigma_y \circ F(u)$ .

*Remark 9.42.* By definition, we have a map  $A \otimes B \rightarrow A \times B$  of scaled simplicial sets which exhibits  $A \times B$  as the one obtained from  $A \otimes B$  by making the upper 2-simplex in the diagram of the form (4) thin. Thus, for  $(\infty, 2)$ -categories  $\mathcal{C}$

and  $\mathcal{D}$ , the cartesian product  $\mathcal{C} \times \mathcal{D}$  is obtained from the Gray product  $\mathcal{C} \otimes \mathcal{D}$  by making the 2-cell in the diagram of the form (5) invertible. By an adjoint argument,  $\mathbf{Fun}(\mathcal{C}, \mathcal{D})$  is regarded as the locally full subcategory of  $\mathbf{Fun}(\mathcal{C}, \mathcal{D})_{\text{Lax}}$  whose 1-cells are the lax natural transformations  $\sigma$  such that the 2-cell  $\sigma_u$  is invertible for any 1-cell  $u$  in  $\mathcal{C}$ . We may also regard  $\mathbf{Fun}(\mathcal{C}, \mathcal{D})$  as a locally full subcategory of  $\mathbf{Fun}(\mathcal{C}, \mathcal{D})_{\text{opLax}}$  in the same way.

Lax natural transformations correspond to functors between 1-cocartesian 2-right fibrations.

**Proposition 9.43.** *Let  $I$  be an  $(\infty, 2)$ -category and let  $\mathcal{C}, \mathcal{D} : I \rightarrow \mathbf{Cat}^{(2)}$  be functors. We have an equivalence*

$$\mathbf{Map}_{\mathbf{Core}_{(1)}(\mathbf{Fun}(I, \mathbf{Cat}^{(2)})_{\text{Lax}})}(\mathcal{C}, \mathcal{D}) \simeq \mathbf{Map}_{2\text{-Cat}/I}(\mathbf{El}_I(\mathcal{C}), \mathbf{El}_I(\mathcal{D}))$$

natural in  $I$ .

Note that Proposition 9.43 could be strengthened to an equivalence of  $(\infty, 1)$ -categories with an appropriate  $(\infty, 2)$ -category structure on  $2\text{-Cat}/I$ . Some special cases are already known.

**Fact 9.44** ([15, Corollary 4.4.3]). *Let  $I$  be an  $(\infty, 2)$ -category. For any functor  $\mathcal{D} : I \rightarrow \mathbf{Cat}^{(2)}$ , we have an equivalence*

$$\mathbf{Map}_{\mathbf{Fun}(I, \mathbf{Cat}^{(2)})_{\text{Lax}}}(\lambda_{-1}, \mathcal{D}) \simeq \mathbf{Map}_{2\text{-Cat}/I}(\lambda_{-1}, \mathbf{El}_I(\mathcal{D})).$$

**Fact 9.45** ([21, Theorem E]). *Let  $I$  be an  $(\infty, 1)$ -category. The map  $(\mathcal{C} : I \rightarrow \mathbf{Cat}^{(2)}) \mapsto \mathbf{El}_I(\mathcal{C})$  induces an equivalence between  $\mathbf{Fun}(I, \mathbf{Cat}^{(2)})_{\text{Lax}}$  and the full subcategory of  $\mathbf{Cat}/I$  spanned by the cocartesian fibrations over  $I$ . Dually,  $\mathbf{Fun}(I^{\text{op}}, \mathbf{Cat}^{(2)})_{\text{opLax}}$  is equivalent to the full subcategory of  $\mathbf{Cat}/I$  spanned by the cartesian fibrations over  $I$ .*

*Proof of Proposition 9.43.* We have a lax natural transformation  $\eta : (\lambda_{-1}) \rightarrow \mathcal{C}|_{\mathbf{El}_I(\mathcal{C})}$  corresponding to the diagonal functor  $\mathbf{El}_I(\mathcal{C}) \rightarrow \mathbf{El}_I(\mathcal{C}) \times_I \mathbf{El}_I(\mathcal{C})$  by Fact 9.44. The precomposition with  $\eta$  induces a map

$$\begin{aligned} & \mathbf{Map}_{\mathbf{Core}_{(1)}(\mathbf{Fun}(I, \mathbf{Cat}^{(2)})_{\text{Lax}})}(\mathcal{C}, \mathcal{D}) \\ & \rightarrow \mathbf{Map}_{\mathbf{Core}_{(1)}(\mathbf{Fun}(\mathbf{El}_I(\mathcal{C}), \mathbf{Cat}^{(2)})_{\text{Lax}})}(\lambda_{-1}, \mathcal{D}|_{\mathbf{El}_I(\mathcal{C})}), \end{aligned} \quad (6)$$

and the codomain is by Fact 9.44 equivalent to  $\mathbf{Map}_{2\text{-Cat}/\mathbf{El}_I(\mathcal{C})}(\mathbf{El}_I(\mathcal{C}), \mathbf{El}_I(\mathcal{C}) \times_I \mathbf{El}_I(\mathcal{D})) \simeq \mathbf{Map}_{2\text{-Cat}/I}(\mathbf{El}_I(\mathcal{C}), \mathbf{El}_I(\mathcal{D}))$ . One can verify that the map (6) is an equivalence by reducing it to the cases when  $\mathcal{C}$  is locally discrete (Fact 9.45) and when  $\mathcal{C} = \mathbf{G}_2$ .  $\square$

A dual argument shows the following.

**Corollary 9.46.** *Let  $I$  be an  $(\infty, 2)$ -category and let  $\mathcal{C}, \mathcal{D} : I^{\text{op}(1,2)} \rightarrow \mathbf{Cat}^{(2)}$  be functors. We have an equivalence*

$$\mathbf{Map}_{\mathbf{Core}_{(1)}(\mathbf{Fun}(I^{\text{op}(1,2)}, \mathbf{Cat}^{(2)})_{\text{opLax}})}(\mathcal{C}, \mathcal{D}) \simeq \mathbf{Map}_{2\text{-Cat}/I}(\mathbf{El}_I(\mathcal{C}), \mathbf{El}_I(\mathcal{D}))$$

natural in  $I$ . □

Natural transformations correspond to functors preserving cocartesian morphisms.

**Proposition 9.47.** *The equivalence in Proposition 9.43 is restricted to an equivalence between  $\mathbf{Map}_{\mathbf{Core}_{(1)}(\mathbf{Fun}(I, \mathbf{Cat}^{(2)}))}(\mathcal{C}, \mathcal{D})$  and the space of functors  $\mathbf{El}_I(\mathcal{C}) \rightarrow \mathbf{El}_I(\mathcal{D})$  over  $I$  preserving cocartesian 1-cells.*

*Proof.* The equivalence between these mapping spaces is due to Lurie [29, Theorem 3.8.1]. One can see that it coincides with the equivalence in Proposition 9.43. □

Oplax limits are  $(\infty, 1)$ -categories of oplax natural transformations in the following sense.

**Proposition 9.48.** *Let  $\mathcal{C} : I^{\text{op}(1,2)} \rightarrow \mathbf{Cat}^{(2)}$  be a functor. For any  $(\infty, 1)$ -category  $\mathcal{D}$ , we have a natural equivalence*

$$\mathbf{Map}_{\mathbf{Cat}}(\mathcal{D}, \text{opLaxLim}_{i \in I} \mathcal{C}_i) \simeq \mathbf{Map}_{\mathbf{Core}_{(1)}(\mathbf{Fun}(I^{\text{op}(1,2)}, \mathbf{Cat}^{(2)})_{\text{opLax}})}(\lambda_{-} \mathcal{D}, \mathcal{C}).$$

*Proof.*

$$\begin{aligned} & \mathbf{Map}_{\mathbf{Cat}}(\mathcal{D}, \text{opLaxLim}_{i \in I} \mathcal{C}_i) \\ \simeq & \quad \{\text{definition}\} \\ & \mathbf{Map}_{2\text{-Cat}/I}(I \times \mathcal{D}, \mathbf{El}_I(\mathcal{C})) \\ \simeq & \quad \{\text{Corollary 9.46}\} \\ & \mathbf{Map}_{\mathbf{Core}_{(1)}(\mathbf{Fun}(I^{\text{op}(1,2)}, \mathbf{Cat}^{(2)})_{\text{opLax}})}(\lambda_{-} \mathcal{D}, \mathcal{C}). \end{aligned}$$

□

A useful source of oplax natural transformations is the following *mate correspondence* whose special case when  $\mathcal{C}$  is an  $(\infty, 1)$ -category is shown by Haugseng et al. [21, Corollary F] in a stronger form of an equivalence between  $(\infty, 1)$ -categories of (op)lax natural transformations.

**Proposition 9.49.** *Let  $\mathcal{C}$  be an  $(\infty, 2)$ -category and  $F, G : \mathcal{C} \rightarrow \mathbf{Cat}^{(2)}$  a functor. We have an equivalence natural in  $\mathcal{C}$  between the following space:*

- the space of oplax natural transformations  $\sigma : F \rightarrow G$  such that  $\sigma_x$  is a left adjoint for every 0-cell  $x \in \mathcal{C}$ ;

- the space of lax natural transformations  $\tau : G \rightarrow F$  such that  $\tau_x$  is a right adjoint for every 0-cell  $x \in \mathcal{C}$ .

Moreover, when an oplax natural transformation  $\sigma$  corresponds to a lax natural transformation  $\tau$  via this equivalence,  $\sigma_x \dashv \tau_x$  for any  $x \in \mathcal{C}$ .

*Proof.* By

$$\begin{aligned}
& \{\mathbf{G}_1 \rightarrow \mathbf{Fun}(\mathcal{C}, \mathbf{Cat}^{(2)})_{\text{opLax}} \mid \text{left adjoint at every } x \in \mathcal{C}\} \\
\cong & \{\text{transpose}\} \\
& \{\mathcal{C} \rightarrow \mathbf{Fun}(\mathbf{G}_1, \mathbf{Cat}^{(2)})_{\text{Lax}} \mid \text{valued in left adjoints}\} \\
\cong & \{\text{Fact 9.45}\} \\
& \{\mathcal{C} \rightarrow \mathbf{Cat}^{(2)}_{/\mathbf{G}_1} \mid \text{valued in bicartesian fibrations}\} \\
\cong & \{\text{Fact 9.45}\} \\
& \{\mathcal{C} \rightarrow \mathbf{Fun}(\mathbf{G}_1^{\text{op}}, \mathbf{Cat}^{(2)})_{\text{opLax}} \mid \text{valued in right adjoints}\} \\
\cong & \{\text{transpose}\} \\
& \{\mathbf{G}_1^{\text{op}} \rightarrow \mathbf{Fun}(\mathcal{C}, \mathbf{Cat}^{(2)})_{\text{Lax}} \mid \text{right adjoint at every } x \in \mathcal{C}\}.
\end{aligned}$$

Recall that a *bicartesian fibration* is a functor between  $(\infty, 1)$ -categories that is both a cocartesian fibration and a cartesian fibration. Because adjunctions are bicartesian fibrations over  $\mathbf{G}_1$  [30, Definition 5.2.2.1], the middle equivalences hold.  $\square$

## 10 Semantics of mode sketches

We show that models of a mode sketch  $\mathfrak{M}$  are equivalent to diagrams of  $\infty$ -logoses indexed over  $\mathfrak{M}$ .

**Definition 10.1.** Let  $\mathfrak{M}$  be a mode sketch. A *model of  $\mathfrak{M}$*  is an  $\infty$ -logos  $\mathcal{L}$  equipped with a point  $\mathfrak{m} \in (\mathbf{1} \Vdash_{\mathcal{L}} \mathfrak{M} \rightarrow \text{LAM})$  satisfying Axioms A to C.

**Construction 10.2.** Let  $\mathfrak{M}$  be a mode sketch. We construct an  $(\infty, 2)$ -category  $|\mathfrak{M}|$  as follows. We regard the underlying poset  $I_{\mathfrak{M}}$  as a simplicial set by taking its nerve. The set  $T_{\mathfrak{M}}$  of thin triangles makes  $I_{\mathfrak{M}}$  a scaled simplicial set. Let  $\langle I_{\mathfrak{M}}, T_{\mathfrak{M}} \rangle$  denote the  $(\infty, 2)$ -category presented by it. We set  $|\mathfrak{M}| = \langle I_{\mathfrak{M}}, T_{\mathfrak{M}} \rangle^{\text{op}(2)}$ .

**Theorem 10.3.** For any mode sketch  $\mathfrak{M}$ , we have an equivalence between the following spaces:

- the space of models of  $\mathfrak{M}$ ;
- the space of functors  $|\mathfrak{M}|^{\text{op}(1,2)} \rightarrow \mathbf{Logos}_{\text{LexAcc}}^{(2)}$ .

Moreover, when a model  $(\mathcal{L}, \mathfrak{m})$  of  $\mathfrak{M}$  corresponds to a functor  $\mathcal{K} : |\mathfrak{M}|^{\text{op}(1,2)} \rightarrow \mathbf{Logos}_{\text{LexAcc}}^{(2)}$ , the following hold.



1.  $\mathcal{L} \simeq \text{opLaxLim}_{i \in |\mathfrak{M}|} \mathcal{K}_i$
2. for any  $i \in \mathfrak{M}$ , the projection  $\mathcal{L} \rightarrow \mathcal{K}_i$  is the localization corresponding to  $\mathfrak{m}(i) \in (\mathbf{1} \Vdash_{\mathcal{L}} \text{LAM})$ .

## 10.1 Models of mode sketches in oplax limits

We show that the oplax limit of a functor  $|\mathfrak{M}|^{\text{op}(1,2)} \rightarrow \mathbf{Logos}_{\text{LexAcc}}^{(2)}$  is part of a model of  $\mathfrak{M}$ .

**Construction 10.4.** Let  $\mathfrak{M}$  be a mode sketch and  $\mathcal{K} : |\mathfrak{M}|^{\text{op}(1,2)} \rightarrow \mathbf{Logos}_{\text{LexAcc}}^{(2)}$  a functor. For a cosieve  $S$  on  $\mathfrak{M}$ , we define an object  $\psi_{\mathcal{K}}(S) \in \text{opLaxLim}_{i \in |\mathfrak{M}|} \mathcal{K}_i$  by

$$\psi_{\mathcal{K}}(S)_i = \begin{cases} \mathbf{1} & \text{if } i \in S \\ \mathbf{0} & \text{otherwise.} \end{cases}$$

The other components are uniquely determined by the universal properties of initial and final objects. This determines a point  $\psi_{\mathcal{K}} \in (\mathbf{1} \Vdash_{\text{opLaxLim}_{i \in |\mathfrak{M}|} \mathcal{K}_i} \text{coSieve}(\mathfrak{M}) \rightarrow \mathcal{U})$ . Since the unique morphism  $\psi_{\mathcal{K}}(S) \rightarrow \mathbf{1}$  is mono,  $\psi_{\mathcal{K}}$  belongs to  $\mathbf{1} \Vdash_{\text{opLaxLim}_{i \in |\mathfrak{M}|} \mathcal{K}_i} \text{coSieve}(\mathfrak{M}) \rightarrow \mathbf{Prop}$ . By construction,  $\psi_{\mathcal{K}}$  is a lattice morphism. Let  $\mathfrak{b}_{\mathcal{K}} \in (\mathbf{1} \Vdash_{\text{opLaxLim}_{i \in |\mathfrak{M}|} \mathcal{K}_i} \mathfrak{M} \rightarrow \text{LAM})$  be the point corresponding to  $\psi_{\mathcal{K}}$  via Theorem 4.3.

**Theorem 10.5.** *Let  $\mathfrak{M}$  be a mode sketch. For any functor  $\mathcal{K} : |\mathfrak{M}|^{\text{op}(1,2)} \rightarrow \mathbf{Logos}_{\text{LexAcc}}^{(2)}$ , the point  $\mathfrak{b}_{\mathcal{K}}$  satisfies Axiom B.*

This is proved by determining the localization corresponding to  $\mathfrak{b}_{\mathcal{K}}(i)$ , which is also part of Theorem 10.3. In what follows, fix a functor  $\mathcal{K} : |\mathfrak{M}|^{\text{op}(1,2)} \rightarrow \mathbf{Logos}_{\text{LexAcc}}^{(2)}$  and let  $\psi = \psi_{\mathcal{K}}$ .

**Proposition 10.6.** *For any  $i \in \mathfrak{M}$ , the projection  $\pi_i : \text{opLaxLim}_{j \in |\mathfrak{M}|} \mathcal{K}_j \rightarrow \mathcal{K}_i$  is the localization corresponding to  $\mathfrak{b}_{\mathcal{K}}(i) \in (\mathbf{1} \Vdash_{\text{opLaxLim}_{j \in |\mathfrak{M}|} \mathcal{K}_j} \text{LAM})$ .*

**Notation 10.7.** Let  $S \subset \mathfrak{M}$  be a subset. We regard  $S$  as a mode sketch with the structure inherited from  $\mathfrak{M}$ . Let

$$\pi_S : \text{opLaxLim}_{i \in |\mathfrak{M}|} \mathcal{K}_i \rightarrow \text{opLaxLim}_{i \in |S|} \mathcal{K}_i$$

denote the restriction functor.

**Lemma 10.8.** *For any cosieve  $S$  on  $\mathfrak{M}$ , the restriction functor  $\pi_{\mathfrak{M} \setminus S}$  is the closed localization associated to  $\psi(S)$ .*

*Proof.* Let  $\text{opLaxLim}_{i \in |\mathfrak{M}|} \mathcal{K}_i \rightarrow \mathcal{M}$  denote the closed localization associated to  $\psi(S)$ . Recall that  $\mathcal{M}$  is the full subcategory of  $\text{opLaxLim}_{i \in |\mathfrak{M}|} \mathcal{K}_i$  spanned by those objects  $A$  such that  $\psi(S) \times A \simeq \psi(S)$ . By the definition of  $\psi(S)$ , this condition is equivalent to that  $A_i \simeq \mathbf{1}$  for all  $i \in S$ . Then  $\pi_{\mathfrak{M} \setminus S}$  induces an equivalence  $\mathcal{M} \simeq \text{opLaxLim}_{i \in |\mathfrak{M} \setminus S|} \mathcal{K}_i$ .  $\square$

**Lemma 10.9.** *For any cosieve  $S$  on  $\mathfrak{M}$ , the restriction functor  $\pi_S$  is the open localization associated to  $\psi(S)$ .*

*Proof.* An object  $A \in (\text{opLaxLim}_{i \in |\mathfrak{M}|} \mathcal{K}_i)_{/\psi(S)}$  must satisfy that  $A_i \simeq \mathbf{0}$  for all  $i \in \mathfrak{M} \setminus S$  by the definition of  $\psi(S)$  and by Fact 5.11. Then  $\pi_S$  induces an equivalence  $(\text{opLaxLim}_{i \in |\mathfrak{M}|} \mathcal{K}_i)_{/\psi(S)} \simeq (\text{opLaxLim}_{i \in |S|} \mathcal{K}_i)_{/\pi_S(\psi(S))} \simeq \text{opLaxLim}_{i \in |S|} \mathcal{K}_i$ .  $\square$

*Proof of Proposition 10.6.* By Proposition 5.19 and Lemmas 10.8 and 10.9, the following square is a pushout of  $\infty$ -logoses and exhibits  $\pi_i$  as the meet of the closed localization  $\pi_{\mathfrak{M} \setminus \partial(i \downarrow)}$  and the open localization  $\pi_{(i \downarrow)}$ .

$$\begin{array}{ccc} \text{opLaxLim}_{j \in |\mathfrak{M}|} \mathcal{K}_j & \xrightarrow{\pi_{(i \downarrow)}} & \text{opLaxLim}_{j \in |(i \downarrow)|} \mathcal{K}_j \\ \pi_{\mathfrak{M} \setminus \partial(i \downarrow)} \downarrow & & \downarrow \pi_{(i \downarrow) \setminus \partial(i \downarrow)} \\ \text{opLaxLim}_{j \in |\mathfrak{M} \setminus \partial(i \downarrow)|} \mathcal{K}_j & \xrightarrow[\pi_{(i \downarrow)}]{\Gamma} & \mathcal{K}_i \end{array}$$

Thus,  $\pi_i$  corresponds to the LAM  $\mathfrak{b}_{\mathcal{K}}(i)$  by the definition of it (Construction 4.2).  $\square$

We also need to determine the interpretations of  $\bigcirc_{\mathfrak{b}_{\mathcal{K}}(i)}^{\mathfrak{b}_{\mathcal{K}}(j)}$  and  $\eta_{\mathfrak{b}_{\mathcal{K}}(j)}^{\mathfrak{b}_{\mathcal{K}}(i); \mathfrak{b}_{\mathcal{K}}(k)}$ .

**Notation 10.10.** For  $j < i$  in  $\mathfrak{M}$ , let  $(j < i)$  denote the associated generating 1-cell in  $|\mathfrak{M}|$ . For  $k < j < i$  in  $\mathfrak{M}$ , let  $(k < j < i)$  denote the associated generating 2-cell  $(j < i) \circ (k < j) \Rightarrow (k < i)$  in  $|\mathfrak{M}|$ .

**Proposition 10.11.** *For any  $j < i$  in  $\mathfrak{M}$ , the composite  $\mathcal{K}_i \subset \text{opLaxLim}_{k \in |\mathfrak{M}|} \mathcal{K}_k \xrightarrow{\pi_j} \mathcal{K}_j$  is equivalent to  $\mathcal{K}_{(j < i)}$ .*

The proof of Proposition 10.11 consists of the calculation of the right adjoint  $i_* : \mathcal{K}_i \rightarrow \text{opLaxLim}_{k \in |\mathfrak{M}|} \mathcal{K}_k$  of  $\pi_i$ .

**Lemma 10.12.** *Let  $\mathfrak{M}'$  be the mode sketch with the same underlying poset as  $\mathfrak{M}$  but with no thin triangle. We have an equivalence  $\text{opLaxLim}_{i \in |\mathfrak{M}'|} \mathcal{K}_i \simeq \text{opLaxLim}_{i \in |\mathfrak{M}'|} \mathcal{K}_i$ .*

*Proof.* This is because  $\mathbf{El}_{|\mathfrak{M}'|}(\mathcal{K}) \rightarrow |\mathfrak{M}'|$  is locally a right fibration and thus locally conservative.  $\square$

**Lemma 10.13.** *Suppose that  $\mathfrak{M}$  has no thin triangle. Then  $\mathbf{Core}_{(1)}(|\mathfrak{M}|)$  is freely generated by the strict ordering relation, and  $(j < i)$  is the final object in  $\mathbf{Map}_{|\mathfrak{M}|}(j, i)$  for any  $j < i$ .*

*Proof.* By Example 9.16.  $\square$

**Lemma 10.14.** *For any  $i \in \mathfrak{M}$ , the right adjoint  $i_*$  is given by the following formula for  $A \in \mathcal{K}_i$  and  $j \in \mathfrak{M}$ .*

$$i_*(A)_j \simeq \begin{cases} \mathcal{K}_{(j < i)}(A) & \text{if } j < i \\ A & \text{if } j = i \\ \mathbf{1} & \text{otherwise} \end{cases}$$

*Proof.* We first see that we may assume without loss of generality that  $i$  is the largest element of  $\mathfrak{M}$ . Otherwise, factor  $\pi_i$  as  $\text{opLaxLim}_{j \in |\mathfrak{M}|} \mathcal{K}_j \xrightarrow{\pi_{(\downarrow i)}} \text{opLaxLim}_{j \in |(\downarrow i)|} \mathcal{K}_j \xrightarrow{\pi_{(i\downarrow)}} \mathcal{K}_i$ , where  $(\downarrow i) = \{j \in \mathfrak{M} \mid j \leq i\}$ . The first functor is a closed localization by Lemma 10.8 because  $\mathfrak{M} \setminus (\downarrow i)$  is a cosieve, and the second functor is an open localization by Lemma 10.9. The right adjoint of  $\pi_{(\downarrow i)}$  is then defined by extending  $A \in \text{opLaxLim}_{j \in |(\downarrow i)|} \mathcal{K}_j$  by the final objects at all  $j \in \mathfrak{M} \setminus (\downarrow i)$ . Therefore, the problem is reduced to the calculation of the right adjoint of  $\pi_{(i\downarrow)}$ , and in this case  $i$  is the largest element of  $(\downarrow i)$ .

Let  $i'_*(A)_j$  be defined by the displayed formula. We turn  $i'_*(A)$  into an object of  $\text{opLaxLim}_{j \in |\mathfrak{M}|} \mathcal{K}_j$ . By Lemma 10.12, we assume that  $\mathfrak{M}$  has no thin triangle. Since  $\mathbf{El}_{|\mathfrak{M}|}(\mathcal{K}) \rightarrow |\mathfrak{M}|$  is locally a right fibration, it follows from Lemma 10.13 that an object  $B \in \text{opLaxLim}_{j \in |\mathfrak{M}|} \mathcal{K}_j$  is completely determined by  $B_j \in \mathcal{K}_j$  for all  $j \in \mathfrak{M}$  and  $B_{(k<j)} : B_k \rightarrow \mathcal{K}_{(k<j)}(B_j)$  for all  $k < j$  in  $\mathfrak{M}$ . We can then extend  $i'_*(A)$  as follows.

$$i'_*(A)_{(k<j)} = \begin{cases} \mathcal{K}_{(k<j<i)}(A) : \mathcal{K}_{(k<i)}(A) \rightarrow \mathcal{K}_{(k<j)}(\mathcal{K}_{(j<i)}(A)) & \text{if } j < i \\ \mathbf{id} & \text{if } j = i \end{cases}$$

Since  $i'_*(A)_i \simeq A$  by construction, we have a unique map  $f : i'_*(A) \rightarrow i_*(A)$  whose  $i$ -th component is the identity on  $A$ . To see that  $f$  is invertible, it suffices to construct a retraction  $g$  of  $f$ . Indeed, if  $g \circ f \simeq \mathbf{id}$ , then the  $i$ -th component of  $g$  must be the identity, and thus  $f \circ g \simeq \mathbf{id}$  follows by adjointness. Let  $j \in \mathfrak{M}$ . If  $j = i$ , then we must define  $g_i = \mathbf{id}$ . Suppose that  $j < i$  and consider the following commutative diagram.

$$\begin{array}{ccc} i'_*(A)_j & \xrightarrow{f_j} & i_*(A)_j \\ i'_*(A)_{(j<i)} \downarrow \simeq & & \downarrow i_*(A)_{(j<i)} \\ \mathcal{K}_{(j<i)}(i'_*(A)_i) & \xrightarrow[\mathcal{K}_{(j<i)}(f_i)]{\simeq} & \mathcal{K}_{(j<i)}(i_*(A)_i) \end{array}$$

The left and bottom maps are invertible by definition. Hence, we have a unique retraction  $g_j$  of  $f_j$  commuting with  $i_*(A)_{j<i}$ . This defines a retraction of  $f$ .  $\square$

*Proof of Proposition 10.11.* By Lemma 10.14.  $\square$

**Proposition 10.15.** *For any  $k < j < i$  in  $\mathfrak{M}$ , the natural transformation  $\pi_k i_* \Rightarrow \pi_k j_* \pi_j i_*$  induced by the unit of the adjunction  $\pi_j \dashv j_*$  is equivalent to  $\mathcal{K}_{(k<j<i)}$ .*

*Proof.* This is a consequence of Lemma 10.14.  $\square$

*Proof of Theorem 10.5.* Let  $\mathfrak{m} = \mathfrak{b}_{\mathcal{K}}$ . For any  $i \in \mathfrak{M}$ , we have

$$(X \Vdash_{\text{opLaxLim}_{j \in |\mathfrak{M}|} \mathcal{K}_j} \mathcal{U}_{\mathfrak{m}(i)}) \simeq \mathbf{Obj}((\mathcal{K}_i)_{/X_i})$$

by Propositions 6.15 and 10.6. Then, for  $i < j$ , the function  $\circ_{\mathfrak{m}(i)}^{\mathfrak{m}(j)}$  is interpreted as the composition

$$\mathbf{Obj}((\mathcal{K}_j)_{/X_j}) \xrightarrow{\mathcal{K}_{(i<j)}} \mathbf{Obj}((\mathcal{K}_i)_{/\mathcal{K}_{(i<j)}(X_j)}) \xrightarrow{X_{(i<j)}^*} \mathbf{Obj}((\mathcal{K}_i)_{/X_i}) \quad (7)$$

by Proposition 10.11. For  $i_0 < i_1 < i_2$ , the function  $\eta_{\mathbf{m}(i_1)}^{\mathbf{m}(i_0); \mathbf{m}(i_2)}$  is interpreted as the following natural transformation by Proposition 10.15.

$$\begin{array}{ccccc}
(\mathcal{K}_{i_0})/X_{i_0} & \xleftarrow{X_{(i_0 < i_2)}^*} & (\mathcal{K}_{i_0})/\mathcal{K}_{(i_0 < i_2)}(X_{i_2}) & \xleftarrow{\mathcal{K}_{(i_0 < i_2)}} & (\mathcal{K}_{i_2})/X_{i_2} \\
& \swarrow X_{(i_0 < i_1)}^* & \uparrow \mathcal{K}_{(i_0 < i_1 < i_2)}(X_{i_2})^* & \searrow \mathcal{K}_{(i_0 < i_1 < i_2)} & \swarrow \mathcal{K}_{(i_1 < i_2)} \\
& & (\mathcal{K}_{i_0})/\mathcal{K}_{(i_0 < i_1)}(\mathcal{K}_{(i_1 < i_2)}(X_{i_2})) & & \\
& & \swarrow \mathcal{K}_{(i_0 < i_1)} & \swarrow \mathcal{K}_{(i_0 < i_1)} & \\
& & (\mathcal{K}_{i_0})/\mathcal{K}_{(i_0 < i_1)}(X_{i_1}) & & (\mathcal{K}_{i_1})/\mathcal{K}_{(i_1 < i_2)}(X_{i_2}) \\
& & \swarrow \mathcal{K}_{(i_0 < i_1)} & \swarrow X_{(i_1 < i_2)}^* & \\
& & (\mathcal{K}_{i_1})/X_{i_1} & & 
\end{array}$$

Then Axiom B can be verified.  $\square$

*Remark 10.16.* The proof of Theorem 10.5 also shows that some properties of the functors  $\mathcal{K}_{(i < j)} : \mathcal{K}_j \rightarrow \mathcal{K}_i$  such as invertibility and conservativity are *not* reflected to the internal functors  $\circ_{\mathbf{m}(i)}^{\mathbf{m}(j)} : \mathcal{U}_{\mathbf{m}(j)} \rightarrow \mathcal{U}_{\mathbf{m}(i)}$ , since  $\circ_{\mathbf{m}(i)}^{\mathbf{m}(j)}$  is interpreted as Eq. (7). It also show that some other properties of  $\mathcal{K}_{(i < j)}$ , in particular preservation of pullback-stable structures such as colimits and pushforwards, are reflected to  $\circ_{\mathbf{m}(i)}^{\mathbf{m}(j)}$ .

## 10.2 Fracture and gluing

We show that any model of a mode sketch  $\mathfrak{M}$  induces a functor  $|\mathfrak{M}|^{\text{op}(1,2)} \rightarrow \mathbf{Logos}_{\text{LexAcc}}^{(2)}$  and that this given an inverse of the construction given in Section 10.1. This is an externalization and generalization of the *fracture and gluing theorem* (Proposition 2.15).

**Notation 10.17.** For a model  $(\mathcal{L}, \mathbf{m})$  of  $\mathfrak{M}$  and  $i \in \mathfrak{M}$ , let  $F_i : \mathcal{L} \rightarrow \mathcal{X}_{\mathbf{m}}(i)$  denote the localization corresponding to  $\mathbf{m}(i)$ .

**Construction 10.18.** Let  $(\mathcal{L}, \mathbf{m})$  be a model of  $\mathfrak{M}$ . We regard  $\mathcal{L}$  as a quasi-category and define  $\mathcal{Y}_{\mathbf{m}}$  to be the full simplicial subset of  $I_{\mathfrak{M}}^{\text{op}} \times \mathcal{L}$  spanned by those objects  $(i, A)$  such that  $A$  belongs to  $\mathcal{X}_{\mathbf{m}}(i)$ .

**Proposition 10.19.** *For any model  $(\mathcal{L}, \mathbf{m})$  of  $\mathfrak{M}$ , the projection  $\mathcal{Y}_{\mathbf{m}} \rightarrow I_{\mathfrak{M}}^{\text{op}}$  is a locally cocartesian fibration.*

*Proof.* Let  $(i_0 \leq i_1) : \Delta^1 \rightarrow I_{\mathfrak{M}}^{\text{op}}$  be a map which corresponds to an ordered pair  $(i_0 \leq i_1)$  in  $I_{\mathfrak{M}}^{\text{op}}$ . A morphism  $(i_0, A_0) \rightarrow (i_1, A_1)$  in  $\mathcal{Y}_{\mathbf{m}}$  over  $(i_0 \leq i_1)$  is a map  $A_0 \rightarrow A_1$  in  $\mathcal{L}$ , but it corresponds to a map  $F_{i_1}(A_0) \rightarrow A_1$  in  $\mathcal{X}_{\mathbf{m}}(i_1)$ . Hence,  $(i_0 \leq i_1)^* \mathcal{Y}_{\mathbf{m}} \rightarrow \Delta^1$  is the Grothendieck construction for the diagram  $\mathcal{X}_{\mathbf{m}}(i_0) \xrightarrow{F_{i_1}} \mathcal{X}_{\mathbf{m}}(i_1)$  and thus a cocartesian fibration.

Let  $(i_0 \leq i_1 \leq i_2) : \Delta^2 \rightarrow I_{\mathfrak{M}}^{\text{op}}$  be a thin 2-simplex. By Axiom B, the canonical natural transformation

$$\begin{array}{ccc}
\mathcal{X}_{\mathfrak{m}}(i_0) & \xrightarrow{F_{i_2}} & \mathcal{X}_{\mathfrak{m}}(i_2) \\
& \searrow^{F_{i_1}} & \Downarrow \simeq \\
& & \mathcal{X}_{\mathfrak{m}}(i_1) \\
& & \nearrow_{F_{i_2}}
\end{array} \tag{8}$$

is invertible, and  $(i_0 \leq i_1 \leq i_2)^* \mathcal{Y}_{\mathfrak{m}} \rightarrow \Delta^2$  is the Grothendieck construction for the diagram (8) and thus a cocartesian fibration.  $\square$

**Construction 10.20.** Let  $(\mathcal{L}, \mathfrak{m})$  be a model of  $\mathfrak{M}$ . By Proposition 10.19 and Fact 9.19, the projection  $\mathcal{Y}_{\mathfrak{m}} \rightarrow I_{\mathfrak{M}}^{\text{op}}$  is classified by some map of scaled simplicial sets

$$(I_{\mathfrak{M}}^{\text{op}}, T_{\mathfrak{M}}) \rightarrow \uparrow \mathbf{Cat}^{\text{sc}}$$

which presents a functor

$$|\mathfrak{M}|^{\text{op}(1,2)} \simeq \langle I_{\mathfrak{M}}, T_{\mathfrak{M}} \rangle^{\text{op}(1)} \simeq \langle I_{\mathfrak{M}}^{\text{op}}, T_{\mathfrak{M}} \rangle \rightarrow \uparrow \mathbf{Cat}^{(2)}.$$

By construction, it maps an element  $i \in \mathfrak{M}$  to  $\mathcal{X}_{\mathfrak{m}}(i)$ , so we refer to this functor also as  $\mathcal{X}_{\mathfrak{m}}$ . As we have seen in the proof of Proposition 10.19,  $\mathcal{X}_{\mathfrak{m}}$  maps a 1-cell  $i_0 \leq i_1$  in  $|\mathfrak{M}|$  to  $\mathcal{X}_{\mathfrak{m}}(i_1) \subset \mathcal{L} \xrightarrow{F_{i_0}} \mathcal{X}_{\mathfrak{m}}(i_0)$  which is lex and accessible (but need not preserve all colimits). Therefore,  $\mathcal{X}_{\mathfrak{m}}$  factors through  $\mathbf{Logos}_{\text{LexAcc}}^{(2)}$ .

We have constructed back and forth maps between the space of models of  $\mathfrak{M}$  and the space of functors  $|\mathfrak{M}|^{\text{op}(1,2)} \rightarrow \mathbf{Logos}_{\text{LexAcc}}^{(2)}$ . For Theorem 10.3, it remains to show that they are mutual inverses (Theorems 10.22 and 10.24 below).

**Construction 10.21.** Let  $(\mathcal{L}, \mathfrak{m})$  be a model of  $\mathfrak{M}$ . The inclusion  $\mathcal{Y}_{\mathfrak{m}} \rightarrow I_{\mathfrak{M}}^{\text{op}} \times \mathcal{L}$  over  $I_{\mathfrak{M}}^{\text{op}}$  corresponds to a lax natural transformation  $\mathcal{X}_{\mathfrak{m}} \rightarrow (\lambda_{\cdot} \mathcal{L})$  by Proposition 9.43. By construction, it has the point-wise left adjoint  $F_i : \mathcal{L} \rightarrow \mathcal{X}_{\mathfrak{m}}(i)$ . Thus, by Proposition 9.49, we have an oplax natural transformation  $(\lambda_{\cdot} \mathcal{L}) \rightarrow \mathcal{X}_{\mathfrak{m}}$  which induces a functor

$$H = H_{\mathfrak{m}} : \mathcal{L} \rightarrow \text{opLaxLim}_{i \in |\mathfrak{M}|} \mathcal{X}_{\mathfrak{m}}(i)$$

by Proposition 9.48. By construction,  $H(A)_i \simeq F_i(A)$  for  $A \in \mathcal{L}$  and  $i \in \mathfrak{M}$ .

**Theorem 10.22.** *For any model  $(\mathcal{L}, \mathfrak{m})$  of  $\mathfrak{M}$ , the functor  $H : \mathcal{L} \rightarrow \text{opLaxLim}_{i \in |\mathfrak{M}|} \mathcal{X}_{\mathfrak{m}}(i)$  is an equivalence and maps  $\varphi_{\mathfrak{m}}$  (Theorem 4.3) to  $\psi_{\mathcal{X}_{\mathfrak{m}}}$  (Construction 10.4). Consequently,  $(\mathcal{L}, \mathfrak{m}) \simeq (\text{opLaxLim}_{i \in |\mathfrak{M}|} \mathcal{X}_{\mathfrak{m}}(i), \mathfrak{b}_{\mathcal{X}_{\mathfrak{m}}})$  in the space of models of  $\mathfrak{M}$ .*

**Construction 10.23.** For a functor  $\mathcal{K} : |\mathfrak{M}|^{\text{op}(1,2)} \rightarrow \mathbf{Logos}_{\text{LexAcc}}^{(2)}$ , let  $\mathbf{El}_{I_{\mathfrak{M}}^{\text{op}}}(\mathcal{K}) \rightarrow I_{\mathfrak{M}}^{\text{op}}$  be the locally cocartesian fibration classified by a map of scaled simplicial

sets that presents  $\mathcal{K}$ . By the same argument as Construction 10.21 but in the reverse direction, the canonical oplax natural transformation  $(\lambda_{\cdot} \text{opLaxLim}_{i \in |\mathfrak{M}|} \mathcal{K}_i) \rightarrow \mathcal{K}$  corresponds to a map of simplicial sets  $\mathbf{El}_{I_{\mathfrak{M}}^{\text{op}}}(\mathcal{K}) \rightarrow I_{\mathfrak{M}}^{\text{op}} \times \text{opLaxLim}_{i \in |\mathfrak{M}|} \mathcal{K}_i$  over  $I_{\mathfrak{M}}^{\text{op}}(\mathcal{K})$ . By construction and by Proposition 10.6, this map factors through  $\mathcal{Y}_{\mathfrak{b}_{\mathcal{K}}}$  (up to homotopy) and thus induces a lax natural transformation

$$\sigma = \sigma_{\mathcal{K}} : \mathcal{K} \rightarrow \mathcal{X}_{\mathfrak{b}_{\mathcal{K}}}.$$

**Theorem 10.24.** *For any functor  $\mathcal{K} : |\mathfrak{M}|^{\text{op}(1,2)} \rightarrow \mathbf{Logos}_{\text{LexAcc}}^{(2)}$ , the lax natural transformation  $\sigma : \mathcal{K} \rightarrow \mathcal{X}_{\mathfrak{b}_{\mathcal{K}}}$  is an invertible natural transformation. Consequently,  $\mathcal{K} \simeq \mathcal{X}_{\mathfrak{b}_{\mathcal{K}}}$ .*

Theorem 10.24 is easier.

*Proof of Theorem 10.24.* To see that  $\sigma$  is a natural transformation, by Proposition 9.47, it suffices to check that the corresponding map of scaled simplicial sets  $\mathbf{El}_{I_{\mathfrak{M}}^{\text{op}}}(\mathcal{K}) \rightarrow \mathcal{Y}_{\mathfrak{b}_{\mathcal{K}}}$  preserves cocartesian morphisms, but this is immediate from the construction. Also by construction,  $\sigma$  is point-wise invertible and thus invertible by Proposition 9.13.  $\square$

For Theorem 10.22, we first see that  $\varphi_{\mathfrak{m}}$  is obtained by Construction 10.4, internally.

**Lemma 10.25.** *In type theory, let  $P : \text{coSieve}(\mathfrak{M}) \rightarrow \mathbf{Prop}$  be a function preserving binary meets. For any cosieve  $S \subset \mathfrak{M}$  and any  $i : \mathfrak{M}$ , we have*

$$\circ_{\mathfrak{a}_P(i)} P(S) = \begin{cases} \mathbf{1} & (i \in S) \\ \circ_{\mathfrak{a}_P(i)} \mathbf{0} & (i \notin S). \end{cases}$$

*Proof.* Suppose that  $i \in S$ . Then  $(i \downarrow) \subset S$ , and thus  $\mathfrak{Cl}(P(S)) \leq \mathfrak{Cl}(P(i \downarrow)) \leq \mathfrak{Cl}(P(\partial(i \downarrow)))$ . Since  $P(S)$  is  $\mathfrak{Cl}(P(S))$ -modal,  $P(S) \simeq \circ_{\mathfrak{Cl}(P(\partial(i \downarrow)))} P(S)$  and it is  $\mathfrak{Op}(P(i \downarrow))$ -connected. Thus,  $\circ_{\mathfrak{a}_P(i)} P(S) = \circ_{\mathfrak{Op}(P(i \downarrow))} \circ_{\mathfrak{Cl}(P(\partial(i \downarrow)))} P(S)$  is contractible.

Suppose that  $i \notin S$ . By  $\mathfrak{a}_P(i)$ -recursion, it suffices to show that  $P(S) \rightarrow \circ_{\mathfrak{a}_P(i)} \mathbf{0}$ . This is equivalent to that  $P(S) \wedge P(i \downarrow) \rightarrow \circ_{\mathfrak{Cl}(P(\partial(i \downarrow)))} \mathbf{0}$  by Fact 2.12 and Proposition 2.28. By assumption  $P(S) \wedge P(i \downarrow) = P(S \cap (i \downarrow))$ , and this is identical to  $P(S \cap \partial(i \downarrow))$  since  $i \notin S$ . Then we have  $P(S) \wedge P(i \downarrow) \rightarrow P(\partial(i \downarrow))$  and  $P(\partial(i \downarrow)) \rightarrow \circ_{\mathfrak{Cl}(P(\partial(i \downarrow)))} \mathbf{0}$ .  $\square$

*Proof of Theorem 10.22.* Lemma 10.25 shows that the functor  $H$  maps  $\varphi_{\mathfrak{m}}$  to  $\psi_{\mathcal{X}_{\mathfrak{m}}}$ . Then, for any cosieve  $S \subset \mathfrak{M}$ , we have the following commutative diagram

$$\begin{array}{ccc} \mathcal{L} & \xrightarrow{H} & \text{opLaxLim}_{i \in |\mathfrak{M}|} \mathcal{X}_{\mathfrak{m}}(i) \\ \varphi_{\mathfrak{m}}(S)^* \downarrow & & \downarrow \pi_S \\ \mathcal{L}/\varphi_{\mathfrak{m}}(S) & \xrightarrow{H/\varphi_{\mathfrak{m}}(S)} & \text{opLaxLim}_{i \in |S|} \mathcal{X}_{\mathfrak{m}}(i) \end{array}$$

by Lemma 10.9. We show that  $H_{/\varphi_m(S)}$  is an equivalence for any cosieve  $S \subset \mathfrak{M}$  by induction on the size of  $S$ : since  $\varphi_m$  preserves top elements, the case when  $S = \mathfrak{M}$  shows that  $H$  is an equivalence. When  $S$  is empty, both domain and codomain of  $H_{/\varphi_m(\emptyset)}$  are the point, so  $H_{/\varphi_m(\emptyset)}$  is an equivalence. Suppose that  $S$  is non-empty. There is an element  $i_0 \in S$  minimal in  $S$ . Internally to  $\mathcal{L}$ , we have  $\mathfrak{m}(i_0) \leq {}^\perp \mathfrak{Op}(\varphi_m(S \setminus \{i_0\}))$ , and thus  $\mathcal{U}_{\mathfrak{Op}(\varphi_m(S))} \simeq \mathcal{U}_{\mathfrak{m}(i_0) \vee \mathfrak{Op}(\varphi_m(S \setminus \{i_0\}))}$  fits into the following pullback diagram by Proposition 2.15.

$$\begin{array}{ccc} \mathcal{U}_{\mathfrak{Op}(\varphi_m(S))} & \longrightarrow & \sum_{A,B:\mathfrak{U}_{\mathfrak{m}(i_0)}} A \rightarrow B \\ \downarrow & \lrcorner & \downarrow \mathbf{cod} \\ \mathcal{U}_{\mathfrak{Op}(\varphi_m(S \setminus \{i_0\}))} & \xrightarrow[\mathfrak{O}_{\mathfrak{m}(i_0)}^{\mathfrak{Op}(\varphi_m(S \setminus \{i_0\}))}]{} & \mathcal{U}_{\mathfrak{m}(i_0)} \end{array}$$

Externalizing this pullback, we have a pullback

$$\begin{array}{ccc} \mathcal{L}_{/\varphi_m(S)} & \longrightarrow & \mathcal{X}_{\mathfrak{m}(i_0)} \rightarrow \\ \downarrow & \lrcorner & \downarrow \mathbf{cod} \\ \mathcal{L}_{/\varphi_m(S \setminus \{i_0\})} & \longrightarrow & \mathcal{X}_{\mathfrak{m}(i_0)}. \end{array}$$

Applying this argument to  $\text{opLaxLim}_{i \in |\mathfrak{M}|} \mathcal{X}_{\mathfrak{m}(i)}$ , we have a pullback

$$\begin{array}{ccc} \text{opLaxLim}_{i \in |S|} \mathcal{X}_{\mathfrak{m}(i)} & \longrightarrow & \mathcal{X}_{\mathfrak{m}(i_0)} \rightarrow \\ \downarrow & \lrcorner & \downarrow \mathbf{cod} \\ \text{opLaxLim}_{i \in |S \setminus \{i_0\}|} \mathcal{X}_{\mathfrak{m}(i)} & \longrightarrow & \mathcal{X}_{\mathfrak{m}(i_0)}. \end{array}$$

Since  $\mathcal{L}_{/\varphi_m(S \setminus \{i_0\})} \simeq \text{opLaxLim}_{i \in |S \setminus \{i_0\}|} \mathcal{X}_{\mathfrak{m}(i)}$  by the induction hypothesis, we have  $\mathcal{L}_{/\varphi_m(S)} \simeq \text{opLaxLim}_{i \in |S|} \mathcal{X}_{\mathfrak{m}(i)}$ .  $\square$

*Proof of Theorem 10.3.* By Theorems 10.22 and 10.24.  $\square$

## Acknowledgements

The author thanks Jon Sterling for useful conversations on the current work. The author is supported by KAW Grant ‘‘Type Theory for Mathematics and Computer Science’’.

## References

- [1] J. Adámek and J. Rosický. *Locally Presentable and Accessible Categories*. Vol. 189. London Mathematical Society Lecture Note Series. Cambridge University Press, 1994. DOI: 10.1017/CB09780511600579.

- [2] M. Anel, G. Biedermann, E. Finster, and A. Joyal. “A generalized Blakers-Massey theorem”. In: *J. Topol.* 13.4 (2020), pp. 1521–1553. DOI: 10.1112/topo.12163.
- [3] M. Anel, G. Biedermann, E. Finster, and A. Joyal. “Left-exact localizations of  $\infty$ -topoi I: Higher sheaves”. In: *Adv. Math.* 400 (2022), Paper No. 108268, 64. DOI: 10.1016/j.aim.2022.108268.
- [4] M. Anel and A. Joyal. “Topo-logie”. In: *New Spaces in Mathematics: Formal and Conceptual Reflections*. Ed. by M. Anel and G. Catren. Vol. 1. Cambridge University Press, 2021, pp. 155–257. DOI: 10.1017/9781108854429.007.
- [5] P. Arndt and K. Kapulkin. “Homotopy-Theoretic Models of Type Theory”. In: *Typed Lambda Calculi and Applications: 10th International Conference, TLCA 2011, Novi Sad, Serbia, June 1-3, 2011. Proceedings*. Ed. by L. Ong. Berlin, Heidelberg: Springer Berlin Heidelberg, 2011, pp. 45–60. DOI: 10.1007/978-3-642-21691-6\_7.
- [6] S. Awodey, N. Gambino, and S. Hazratpour. *Kripke-Joyal forcing for type theory and uniform fibrations*. 2021. arXiv: 2110.14576v1.
- [7] S. Awodey and M. A. Warren. “Homotopy theoretic models of identity types”. In: *Mathematical Proceedings of the Cambridge Philosophical Society* 146.1 (2009), pp. 45–55. DOI: 10.1017/S0305004108001783.
- [8] C. Barwick. “ $(\infty, n)$ -Cat as a closed model category”. PhD Thesis. University of Pennsylvania, 2005. URL: <https://repository.upenn.edu/dissertations/AAI3165639/>.
- [9] C. Barwick and C. Schommer-Pries. “On the unicity of the theory of higher categories”. In: *J. Amer. Math. Soc.* 34.4 (2021), pp. 1011–1058. DOI: 10.1090/jams/972.
- [10] J.-P. Bernardy, P. Jansson, and R. Paterson. “Proofs for Free: Parametricity for Dependent Types”. In: *Journal of Functional Programming* 22.2 (Mar. 2012), pp. 107–152. DOI: 10.1017/S0956796812000056.
- [11] J. D. Christensen, M. Opie, E. Rijke, and L. Scoccola. “Localization in homotopy type theory”. In: *High. Struct.* 4.1 (2020), pp. 1–32.
- [12] J. D. Christensen and E. Rijke. “Characterizations of modalities and lex modalities”. In: *J. Pure Appl. Algebra* 226.3 (2022), Paper No. 106848, 21. DOI: 10.1016/j.jpaa.2021.106848.
- [13] D.-C. Cisinski. *Higher Categories and Homotopical Algebra*. Cambridge Studies in Advanced Mathematics. Cambridge University Press, 2019. DOI: 10.1017/9781108588737.
- [14] J. W. Duskin. “Simplicial matrices and the nerves of weak  $n$ -categories. I. Nerves of bicategories”. In: vol. 9. CT2000 Conference (Como). 2001/02, pp. 198–308. URL: <http://www.tac.mta.ca/tac/volumes/9/n10/9-10abs.html>.



- [15] A. Gagna, Y. Harpaz, and E. Lanari. *Fibrations and lax limits of  $(\infty, 2)$ -categories*. 2021. arXiv: 2012.04537v2.
- [16] A. Gagna, Y. Harpaz, and E. Lanari. *Gray tensor products and lax functors of  $(\infty, 2)$ -categories*. 2020. arXiv: 2006.14495v1.
- [17] D. Gaitsgory and N. Rozenblyum. *A study in derived algebraic geometry. Vol. I. Correspondences and duality*. Vol. 221. Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 2017, xl+533pp. DOI: 10.1090/surv/221.1.
- [18] D. Gepner, R. Haugseng, and T. Nikolaus. “Lax Colimits and Free Fibrations in  $\infty$ -Categories”. In: *Documenta Mathematica* 22 (2017), pp. 1225–1266. DOI: 10.25537/dm.2017v22.1225-1266.
- [19] D. Gratzer. “Normalization for Multimodal Type Theory”. In: *LICS '22: 37th Annual ACM/IEEE Symposium on Logic in Computer Science, Haifa, Israel, August 2 - 5, 2022*. Ed. by C. Baier and D. Fisman. ACM, 2022, 2:1–2:13. DOI: 10.1145/3531130.3532398.
- [20] D. Gratzer, G. A. Kavvos, A. Nuyts, and L. Birkedal. “Multimodal Dependent Type Theory”. In: *Logical Methods in Computer Science* Volume 17, Issue 3 (July 2021). DOI: 10.46298/lmcs-17(3:11)2021.
- [21] R. Haugseng, F. Hebestreit, S. Linskens, and J. Nuiten. *Lax monoidal adjunctions, two-variable fibrations and the calculus of mates*. 2021. arXiv: 2011.08808v2.
- [22] K.-B. Hou (Favonia), E. Finster, D. R. Licata, and P. L. Lumsdaine. “A Mechanization of the Blakers-Massey Connectivity Theorem in Homotopy Type Theory”. In: *Proceedings of the 31st Annual ACM/IEEE Symposium on Logic in Computer Science*. LICS '16. New York, NY, USA: ACM, 2016, pp. 565–574. DOI: 10.1145/2933575.2934545.
- [23] N. Johnson and D. Yau. *2-dimensional categories*. Oxford University Press, Oxford, 2021, pp. xix+615. DOI: 10.1093/oso/9780198871378.001.0001.
- [24] P. T. Johnstone. *Sketches of an Elephant : A Topos Theory Compendium Volume 1*. Vol. 43. Oxford Logic Guides. Oxford University Press, 2002.
- [25] G. M. Kelly. “Elementary observations on 2-categorical limits”. In: *Bulletin of the Australian Mathematical Society* 39.2 (1989), pp. 301–317. DOI: 10.1017/S0004972700002781.
- [26] M. Lasson. “Canonicity of Weak  $\omega$ -groupoid Laws Using Parametricity Theory”. In: *Electronic Notes in Theoretical Computer Science* 308 (2014), pp. 229–244. DOI: 10.1016/j.entcs.2014.10.013.
- [27] D. R. Licata, I. Orton, A. M. Pitts, and B. Spitters. “Internal Universes in Models of Homotopy Type Theory”. In: *3rd International Conference on Formal Structures for Computation and Deduction (FSCD 2018)*. Ed. by H. Kirchner. Vol. 108. Leibniz International Proceedings in Informatics (LIPIcs). Dagstuhl, Germany: Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik, 2018, 22:1–22:17. DOI: 10.4230/LIPIcs.FSCD.2018.22.

- [28] P. L. Lumsdaine and M. Shulman. “Semantics of higher inductive types”. In: *Math. Proc. Cambridge Philos. Soc.* 169.1 (2020), pp. 159–208. DOI: 10.1017/s030500411900015x.
- [29] J. Lurie. *( $\infty, 2$ )-Categories and the Goodwillie Calculus I*. 2009. arXiv: 0905.0462v2.
- [30] J. Lurie. *Higher Topos Theory*. Vol. 170. Annals of Mathematics Studies. Princeton University Press, 2009. URL: <https://www.math.ias.edu/~lurie/papers/HTT.pdf>.
- [31] S. Mac Lane and I. Moerdijk. *Sheaves in Geometry and Logic*. New York, NY: Springer New York, 1992. DOI: 10.1007/978-1-4612-0927-0.
- [32] P. Martin-Löf. “An Intuitionistic Theory of Types: Predicative Part”. In: *Studies in Logic and the Foundations of Mathematics* 80 (1975), pp. 73–118. DOI: 10.1016/S0049-237X(08)71945-1.
- [33] H. K. Nguyen and T. Uemura.  *$\infty$ -type theories*. 2022. arXiv: 2205.00798v1.
- [34] E. Riehl and D. Verity. *Elements of  $\infty$ -Category Theory*. Cambridge Studies in Advanced Mathematics. Cambridge University Press, 2022. DOI: 10.1017/9781108936880.
- [35] E. Rijke, M. Shulman, and B. Spitters. “Modalities in homotopy type theory”. In: *Log. Methods Comput. Sci.* 16.1 (2020), Paper No. 2, 79. DOI: 10.23638/LMCS-16(1:2)2020.
- [36] U. Schreiber and M. Shulman. “Quantum Gauge Field Theory in Cohesive Homotopy Type Theory”. In: *Proceedings 9th Workshop on Quantum Physics and Logic, QPL 2012, Brussels, Belgium, 10-12 October 2012*. Ed. by R. Duncan and P. Panangaden. Vol. 158. EPTCS. 2012, pp. 109–126. DOI: 10.4204/EPTCS.158.8.
- [37] M. Shulman. *All  $(\infty, 1)$ -toposes have strict univalent universes*. 2019. arXiv: 1904.07004v2.
- [38] M. Shulman. “Brouwer’s fixed-point theorem in real-cohesive homotopy type theory”. In: *Mathematical Structures in Computer Science* 28.6 (2018), pp. 856–941. DOI: 10.1017/S0960129517000147.
- [39] M. Shulman. “Univalence for inverse diagrams and homotopy canonicity”. In: *Mathematical Structures in Computer Science* 25.05 (2015), pp. 1203–1277. DOI: 10.1017/s0960129514000565.
- [40] G. Stefanich. *Presentable  $(\infty, n)$ -categories*. 2020. arXiv: 2011.03035v1.
- [41] R. Stenzel. *On notions of compactness, object classifiers and weak Tarski universes*. 2022. arXiv: 1911.01895v2.
- [42] J. Sterling. “First Steps in Synthetic Tait Computability. The Objective Metatheory of Cubical Type Theory”. PhD Thesis. Carnegie Mellon University, 2021. URL: <https://www.jonmsterling.com/pdfs/sterling:2021:thesis.pdf>.

- [43] J. Sterling and C. Angiuli. “Normalization for Cubical Type Theory”. In: *2021 36th Annual ACM/IEEE Symposium on Logic in Computer Science (LICS)*. 2021, pp. 1–15. DOI: 10.1109/LICS52264.2021.9470719.
- [44] J. Sterling and R. Harper. “Logical Relations as Types: Proof-Relevant Parametricity for Program Modules”. In: *J. ACM* 68.6 (2021), 41:1–41:47. DOI: 10.1145/3474834.
- [45] J. Sterling and R. Harper. “Sheaf Semantics of Termination-Insensitive Noninterference”. In: *7th International Conference on Formal Structures for Computation and Deduction (FSCD 2022)*. Ed. by A. P. Felty. Vol. 228. Leibniz International Proceedings in Informatics (LIPIcs). Dagstuhl, Germany: Schloss Dagstuhl – Leibniz-Zentrum für Informatik, 2022, 5:1–5:19. DOI: 10.4230/LIPIcs.FSCD.2022.5.
- [46] R. Street. “Limits indexed by category-valued 2-functors”. In: *Journal of Pure and Applied Algebra* 8.2 (1976), pp. 149–181. DOI: 10.1016/0022-4049(76)90013-X.
- [47] R. Street. “Two constructions on Lax functors”. In: *Cahiers de Topologie et Géométrie Différentielle Catégoriques* 13.3 (1972), pp. 217–264.
- [48] The Univalent Foundations Program. *Homotopy Type Theory: Univalent Foundations of Mathematics*. Institute for Advanced Study, 2013. URL: <http://homotopytypetheory.org/book/>.
- [49] T. Uemura. “Fibred Fibration Categories”. In: *2017 32nd Annual ACM/IEEE Symposium on Logic in Computer Science (LICS)*. June 2017, pp. 1–12. DOI: 10.1109/LICS.2017.8005084.
- [50] M. Vergura. *Localization theory in an  $\infty$ -topos*. 2019. arXiv: 1907.03836v1.
- [51] G. Wraith. “Artin glueing”. In: *J. Pure Appl. Algebra* 4 (1974), pp. 345–348. DOI: 10.1016/0022-4049(74)90014-0.