

Bounds on the covariance matrix of the Sherrington–Kirkpatrick model

Ahmed El Alaoui*, Jason Gaitonde†

Abstract

We consider the Sherrington-Kirkpatrick model with no external field and inverse temperature $\beta < 1$ and prove that the expected operator norm of the covariance matrix of the Gibbs measure is bounded by a constant depending only on β . This answers an open question raised by Talagrand, who proved a bound of $C(\beta)(\log n)^8$. Our result follows by establishing an approximate formula for the covariance matrix which we obtain by differentiating the TAP equations and then optimally controlling the associated error terms. We complement this result by showing diverging lower bounds on the operator norm, both at the critical and low temperatures.

1 Introduction and main result

We consider the Sherrington–Kirkpatrick model of spin glasses, a Gibbs distribution over the hypercube $\{-1, +1\}^n$ given by the expression

$$\mu(\sigma) = \frac{1}{Z} \exp \left\{ \frac{\beta}{\sqrt{n}} \sum_{i < j} g_{ij} \sigma_i \sigma_j \right\}, \quad \sigma \in \{-1, +1\}^n, \quad (1.1)$$

where $\beta \geq 0$ is the inverse temperature parameter and $(g_{ij})_{i < j}$ are the random coupling coefficients assumed to be i.i.d. $N(0, 1)$ random variables. We are interested in the behavior of the $n \times n$ covariance matrix of this probability measure:

$$(\text{cov}(\mu))_{ij} = \langle \sigma_i \sigma_j \rangle, \quad i, j \in [1, n]. \quad (1.2)$$

It is expected that the operator norm of $\text{cov}(\mu)$ is of constant order in n whenever the model is at high temperature, i.e., for all $\beta < 1$, and that it must diverge at the critical and low temperatures $\beta \geq 1$. Talagrand proved that for all $\beta < 1$ there exists $C(\beta) < \infty$ such that

$$\mathbb{E} \|\text{cov}(\mu)\|_{\text{op}} \leq C(\beta)(\log n)^8,$$

and conjectured that the logarithmic term can be removed entirely [Tal11, Section 11.5]. His proof relies on the moment method and bounds the expectation of the trace of large powers of the covariance matrix; a method known to be loose by a logarithmic factor for random matrices. Recently, Eldan, Koehler and Zeitouni [EKZ21] proved that Ising measures with a positive semidefinite interaction matrix J such that $\|J\|_{\text{op}} < 1$ satisfy a Poincaré inequality for the Dirichlet form of Glauber dynamics with a spectral gap constant only depending on $\|J\|_{\text{op}}$. In the special case of the SK model, their result implies that $\mathbb{E} \|\text{cov}(\mu)\|_{\text{op}}$ is bounded for all $\beta < 1/4$. In this paper we extend this result to all $\beta < 1$:

Theorem 1.1. *For $\beta < 1$ there exists $C(\beta) < \infty$ such that $\mathbb{E} \|\text{cov}(\mu)\|_{\text{op}} \leq C(\beta)$ for all $n \geq 1$.*

*Department of Statistics and Data Science, Cornell University. Email: elalaoui@cornell.edu.

†Department of Computer Science, Cornell University. Email: jsg355@cornell.edu.

Conversely, we show diverging lower bounds on the operator norm for $\beta \rightarrow 1^-$, $\beta = 1$ and $\beta > 1$:

Theorem 1.2.

- For $\beta < 1$, $\liminf_{n \rightarrow \infty} \mathbb{E} \|\text{cov}(\mu)\|_{\text{op}} \geq \sqrt{\frac{2}{\pi(1-\beta^2)}}$.
- For $\beta = 1$, $\mathbb{E} \|\text{cov}(\mu)\|_{\text{op}}^2 \geq c(n/\log n)^{3/16}$, where $c > 0$ is an absolute constant, for all $n \geq 1$.
- For $\beta > 1$, there exists $c(\beta) > 0$ such that $\mathbb{E} \|\text{cov}(\mu)\|_{\text{op}} \geq c(\beta)\sqrt{n}$ for all $n \geq 1$.

Note that the second bullet in the above theorem lower bounds the second moment of $\|\text{cov}(\mu)\|_{\text{op}}$; a weaker statement. In Proposition 4.3 we prove a similar lower bound on the first moment, albeit not at $\beta = 1$ but at a temperature approaching criticality: $\beta = \beta_n \rightarrow 1^-$ as $n \rightarrow \infty$ such that $n^{1/3}(1-\beta_n^2) \rightarrow +\infty$. The arguments used to prove Theorem 1.2 follow more or less directly from known results, so we delay their exposition to Section 4 in favor of the high-temperature result.

Our approach to showing Theorem 1.1 is by first establishing a TAP equation for the two-point correlations $\langle \sigma_i \sigma_j \rangle$ with an optimal error bound of $1/n^2$ as we write next. Let $A = (g_{ij}/\sqrt{n})_{i,j=1}^n$.

Theorem 1.3. For $\beta < 1$ there exists $C(\beta) < \infty$ such that

$$\mathbb{E} \left[\left\| \left((1 + \beta^2)I - \beta A \right) \text{cov}(\mu) - I \right\|_F^2 \right] \leq C(\beta). \quad (1.3)$$

In particular we have for $i \neq j$,

$$\mathbb{E} \left[\left((1 + \beta^2) \langle \sigma_i \sigma_j \rangle - \frac{\beta}{\sqrt{n}} \sum_{k=1}^n g_{ik} \langle \sigma_k \sigma_j \rangle \right)^2 \right] \leq \frac{C(\beta)}{n^2}. \quad (1.4)$$

An analogue of Eq. (1.4) with a weaker error bound was recently proved by Adhikari, Brennecke, von Soosten and Yau [ABvSY21] for nonzero and Gaussian external field up to an inverse temperature $\beta_0 = \log 2$ using what the authors call the dynamical approach to the TAP equations. The error bound they prove is of the form $C(\beta, \varepsilon)/n^{1+\varepsilon}$ for ε sufficiently small. This bound is unfortunately not enough to imply Theorem 1.1; a bound of order $1/n^2$ seems necessary for the conclusion to follow. We elaborate on a generalization of this result to nonzero external field in Section 2 below.

Theorem 1.1 then follows by known bounds on the tail probability of the largest eigenvalue of a GOE matrix:

Proof of Theorem 1.1. For $\beta < 1$, let $\varepsilon \in (0, (1 - \beta)^2/(1 + \beta))$ and define the event

$$\mathcal{E} = \{ \lambda_{\min}((1 + \beta^2)I - \beta A) \leq \varepsilon \}.$$

Observe that on \mathcal{E} we have $\lambda_{\max}(A) \geq \frac{1+\beta^2-\varepsilon}{\beta} \geq 2 + \varepsilon$ where the second inequality follows from the bound imposed on ε . It follows that $\mathbb{P}(\mathcal{E}) \leq \exp(-c(\varepsilon)n)$, $c(\varepsilon) > 0$ by standard results on the largest eigenvalue of a GOE matrix (see e.g., [Ver18, Exercise 7.3.5]). Now conditional on the complement event $\mathcal{E}^c = \{ \lambda_{\min}((1 + \beta^2)I - \beta A) > \varepsilon \}$ we have

$$\|\text{cov}(\mu)\|_{\text{op}} \leq \left\| \left((1 + \beta^2)I - \beta A \right)^{-1} \right\|_{\text{op}} \cdot \left(\left\| \left((1 + \beta^2)I - \beta A \right) \text{cov}(\mu) - I \right\|_{\text{op}} + 1 \right) \quad (1.5)$$

$$\leq \varepsilon^{-1} \left(\left\| \left((1 + \beta^2)I - \beta A \right) \text{cov}(\mu) - I \right\|_F + 1 \right). \quad (1.6)$$

Using Theorem 1.3,

$$\mathbb{E} [\|\text{cov}(\mu)\|_{\text{op}} \mathbf{1}_{\mathcal{E}^c}] \leq \varepsilon^{-1} \left(\mathbb{E} [\|((1 + \beta^2)I - \beta A) \text{cov}(\mu) - I\|_F] + 1 \right) \quad (1.7)$$

$$\leq \varepsilon^{-1} C(\beta). \quad (1.8)$$

Next, on the event \mathcal{E} , we trivially upper bound $\|\text{cov}(\mu)\|_{\text{op}}$ by n , allowing us to conclude:

$$\mathbb{E} [\|\text{cov}(\mu)\|_{\text{op}}] = \mathbb{E} [\|\text{cov}(\mu)\|_{\text{op}} \mathbf{1}_{\mathcal{E}}] + \mathbb{E} [\|\text{cov}(\mu)\|_{\text{op}} \mathbf{1}_{\mathcal{E}^c}] \quad (1.9)$$

$$\leq n \exp(-c(\varepsilon)n) + \varepsilon^{-1} C(\beta) \quad (1.10)$$

$$\leq (ec(\varepsilon))^{-1} + \varepsilon^{-1} C(\beta). \quad (1.11)$$

□

2 Discussion on nonzero external field

The result of Theorem 1.3 is inspired by the following heuristic. Let us introduce an external field $y = (y_i)_{i=1}^n \in \mathbb{R}^n$ to the Gibbs measure:

$$\mu_y(\sigma) = \frac{1}{Z(y)} \exp \left\{ \frac{\beta}{\sqrt{n}} \sum_{i < j} g_{ij} \sigma_i \sigma_j + \sum_{i=1}^n y_i \sigma_i \right\}. \quad (2.1)$$

For a “typical” external field, the log-partition function of μ_y is expected to have a TAP representation of the following form:

$$\log Z(y) = \max_{m \in (-1,1)^n} \left\{ \mathcal{F}_{\text{TAP}}(m) + \langle y, m \rangle \right\} + o_{\mathbb{P}}(n), \quad \text{where} \quad (2.2)$$

$$\mathcal{F}_{\text{TAP}}(m) := \frac{\beta}{2} \langle m, Am \rangle + \sum_{i=1}^n h(m_i) + \frac{n\beta^2}{4} (1 - Q(m))^2, \quad (2.3)$$

$$Q(m) = \frac{1}{n} \|m\|^2, \quad h(m) = -\frac{1+m}{2} \log \left(\frac{1+m}{2} \right) - \frac{1-m}{2} \log \left(\frac{1-m}{2} \right).$$

The above representation was recently proved for Gaussian external fields by Chen, Panchenko and Subag [CPS18]. Taking two derivatives with respect to y on both sides of Eq. (2.2), we expect

$$\text{cov}(\mu_y) := \langle \sigma_i \sigma_j \rangle - \langle \sigma_i \rangle \langle \sigma_j \rangle \approx -(\nabla^2 \mathcal{F}_{\text{TAP}}(m))^{-1} \quad (2.4)$$

$$= \left(D(m) - \beta A + \beta^2 (1 - Q(m)) I - (2\beta^2/n) m m^\top \right)^{-1}, \quad (2.5)$$

where m is a maximizer in (2.2) and $D(m)$ is the diagonal matrix with entries $D(m)_{ii} = 1/(1 - m_i^2)$. Taking $y = 0$, we have $m = 0$ and

$$\text{cov}(\mu) \approx \left((1 + \beta^2) I - \beta A \right)^{-1}. \quad (2.6)$$

Theorem 1.3 makes the above approximation precise in the sense of a bounded Frobenius norm, once the inverse is eliminated from the right-hand side. This operation will allow us to perform Gaussian integration by parts with respect to the disorder random variables g_{ij} and this creates various overlap terms between independent replicas from the Gibbs measure μ . Then Theorem 1.3 is proved by exploiting known asymptotics for these overlaps in the high-temperature regime.

Similarly to the zero external field case, one can attempt to make the approximation (2.5) precise by showing that for β small enough, e.g., below the AT line when y is Gaussian, we have

$$\mathbb{E} \left[\left\| \left(D(m) - \beta A + \beta^2(1 - Q(m))I - (2\beta^2/n)mm^\top \right) \text{cov}(\mu_y) - I \right\|_F^2 \right] \leq C(\beta), \quad (2.7)$$

where $m = (\langle \sigma_i \rangle)_{i=1}^n$ is the mean vector of μ_y .

Boundedness of $\mathbb{E} \|\text{cov}(\mu_y)\|_{\text{op}}$ would then follow if one can show that $\lambda_{\max}(\nabla^2 \mathcal{F}_{\text{TAP}}(m)) \leq -\varepsilon$ for some $\varepsilon > 0$ with probability at least $1 - O(1/n)$ (see Eq. (1.10)). Celentano [Cel22] recently showed that the TAP free energy \mathcal{F}_{TAP} is locally strongly concave around one of its stationary points with probability $1 - o_n(1)$ by applying a Gaussian comparison theorem carefully conditioned on a sequence of sigma-fields produced by an Approximate Message Passing (AMP) iteration. This stationary point should presumably be close to the mean vector $\langle \sigma \rangle$; see the related work [CFM21]. The standard theory of AMP used in that paper does not yield any quantitative control on the probability of convergence, and this approach seems to fall short of obtaining the $O(1/n)$ rate needed here.

3 Proof of Theorem 1.3

We will show that the left-hand side in Eq. (1.3) is actually equal to

$$\frac{\beta^2(1 + \beta^2)}{(1 - \beta^2)^2} + O(1/\sqrt{n}),$$

thereby showing that it is truly of constant order. We let

$$P = ((1 + \beta^2)I - \beta A) \text{cov}(\mu).$$

We then write

$$\|P - I\|_F^2 = \|P\|_F^2 - 2\text{Tr}(P) + n.$$

We treat the above expression term by term and we show that both $\mathbb{E} \|P\|_F^2$ and $\mathbb{E} \text{Tr}(P)$ are of the form

$$n + C(\beta) + O(1/\sqrt{n}),$$

therefore canceling the terms diverging in n . We first analyze the trace term and turn to the norm term which is more delicate. Let us first record the following simple lemma for future reference:

Lemma 3.1. *For all i, j, k, ℓ , it holds that $\frac{d}{dg_{k\ell}} \langle \sigma_i \sigma_j \rangle = \frac{\beta}{\sqrt{n}} [\langle \sigma_i \sigma_j \sigma_k \sigma_\ell \rangle - \langle \sigma_i \sigma_j \rangle \langle \sigma_k \sigma_\ell \rangle]$.*

Proof. The above is trivially true for $k = \ell$ since the Hamiltonian has no dependence on g_{kk} , and $\sigma_k^2 = 1$. Assume $k \neq \ell$. Writing $\langle f \rangle$ as a quotient, the product rule implies that the first term can be interpreted as the Gibbs average of $f \sigma_k \sigma_\ell$, while the second term obtained by differentiating the partition function yields an independent Gibbs average of $\sigma_k \sigma_\ell$. \square

For ℓ independent replicas $\sigma^1, \dots, \sigma^\ell$ drawn from μ , we write $R_{a,b} = \frac{1}{n} \sum_{i=1}^n \sigma_i^a \sigma_i^b$ for the pairwise overlap between σ^a and σ^b , and

$$R_{1,\dots,\ell} = \frac{1}{n} \sum_{i=1}^n \prod_{a=1}^{\ell} \sigma_i^a$$

for the multi-overlap of the replicas $\sigma^1, \dots, \sigma^\ell$.

The trace term: Using Gaussian integration by parts and Lemma 3.1, we have

$$\begin{aligned}
\mathbb{E} \operatorname{Tr}(P) &= (1 + \beta^2)n - \beta \mathbb{E} \operatorname{Tr}(A \operatorname{cov}(\mu)) = (1 + \beta^2)n - \frac{\beta}{\sqrt{n}} \sum_{i,j=1}^n \mathbb{E} [g_{ij} \langle \sigma_i \sigma_j \rangle] \\
&= (1 + \beta^2)n - \frac{\beta^2}{n} \sum_{i,j=1}^n (1 - \mathbb{E} \langle \sigma_i \sigma_j \rangle^2) \\
&= n + n\beta^2 \mathbb{E} \langle R_{1,2}^2 \rangle.
\end{aligned}$$

From [Tal11, Theorem 11.5.4], it known that for $\beta < 1$,

$$n \mathbb{E} \langle R_{1,2}^2 \rangle = \frac{1}{1 - \beta^2} + O(1/\sqrt{n}).$$

Therefore

$$\mathbb{E} \operatorname{Tr}(P) = n + \frac{\beta^2}{1 - \beta^2} + O(1/\sqrt{n}). \quad (3.1)$$

The norm term: We now calculate $\mathbb{E} \|P\|_F^2$. We have

$$\begin{aligned}
\mathbb{E} \|P\|_F^2 &= \mathbb{E} \sum_{i,j=1}^n \left((1 + \beta^2) \langle \sigma_i \sigma_j \rangle - \frac{\beta}{\sqrt{n}} \sum_{k=1}^n g_{ik} \langle \sigma_j \sigma_k \rangle \right)^2 \\
&= \mathbb{E} \sum_{i,j=1}^n \left((1 + \beta^2)^2 \langle \sigma_i \sigma_j \rangle^2 - 2(1 + \beta^2) \frac{\beta}{\sqrt{n}} \sum_{k=1}^n g_{ik} \langle \sigma_j \sigma_k \rangle \langle \sigma_i \sigma_j \rangle \right. \\
&\quad \left. + \frac{\beta^2}{n} \sum_{k,\ell=1}^n g_{ik} g_{i\ell} \langle \sigma_j \sigma_k \rangle \langle \sigma_j \sigma_\ell \rangle \right) \\
&= \text{I} + \text{II} + \text{III}.
\end{aligned}$$

We deal with these terms in order. Since

$$\sum_{ij} \langle \sigma_i \sigma_j \rangle^2 = \sum_{ij} \langle \sigma_i^1 \sigma_j^1 \sigma_i^2 \sigma_j^2 \rangle = n^2 \langle R_{12}^2 \rangle,$$

the first term is

$$\text{I} = (1 + \beta^2)^2 n^2 \mathbb{E} \langle R_{12}^2 \rangle. \quad (3.2)$$

As for the second term, using Lemma 3.1 we have

$$\begin{aligned}
\frac{\beta}{\sqrt{n}} \mathbb{E} [g_{ik} \langle \sigma_j \sigma_k \rangle \langle \sigma_i \sigma_j \rangle] &= \frac{\beta}{\sqrt{n}} \mathbb{E} \left[\frac{d}{dg_{ik}} [\langle \sigma_j \sigma_k \rangle \langle \sigma_i \sigma_j \rangle] \right] \\
&= \frac{\beta^2}{n} \mathbb{E} [(\langle \sigma_i \sigma_j \rangle - \langle \sigma_i \sigma_k \rangle \langle \sigma_j \sigma_k \rangle) \langle \sigma_i \sigma_j \rangle + (\langle \sigma_j \sigma_k \rangle - \langle \sigma_i \sigma_k \rangle \langle \sigma_i \sigma_j \rangle) \langle \sigma_j \sigma_k \rangle] \\
&= \frac{\beta^2}{n} \mathbb{E} [\langle \sigma_i \sigma_j \rangle^2 + \langle \sigma_j \sigma_k \rangle^2 - 2 \langle \sigma_i \sigma_j \rangle \langle \sigma_j \sigma_k \rangle \langle \sigma_i \sigma_k \rangle].
\end{aligned}$$

Summing the above expression over i, j, k , this is

$$2\beta n^2 \mathbb{E} \langle R_{12}^2 \rangle + 2\beta n^2 \mathbb{E} \langle R_{13} R_{12} R_{23} \rangle.$$

Therefore

$$\Pi = -4\beta^2(1 + \beta^2)n^2 (\mathbb{E} \langle R_{12}^2 \rangle - \mathbb{E} \langle R_{12}R_{23}R_{13} \rangle) . \quad (3.3)$$

We now turn to the third term. We split the sum into a diagonal and a off-diagonal part:

$$\sum_{k,\ell=1}^n g_{ik}g_{i\ell} \langle \sigma_j \sigma_k \rangle \langle \sigma_j \sigma_\ell \rangle = \sum_{k=1}^n g_{ik}^2 \langle \sigma_j \sigma_k \rangle^2 + \sum_{k \neq \ell} g_{ik}g_{i\ell} \langle \sigma_j \sigma_k \rangle \langle \sigma_j \sigma_\ell \rangle . \quad (3.4)$$

For the diagonal term in Eq. (3.4), taking expectations and applying Gaussian integration by parts, we find

$$\begin{aligned} \frac{1}{n} \sum_k \mathbb{E}[g_{ik}^2 \langle \sigma_j \sigma_k \rangle^2] &= \frac{1}{n} \sum_k \mathbb{E}[\langle \sigma_j \sigma_k \rangle^2] + \frac{2\beta}{n^{3/2}} \sum_k \mathbb{E}[g_{ik} \langle \sigma_j \sigma_k \rangle (\langle \sigma_i \sigma_j \rangle - \langle \sigma_i \sigma_k \rangle \langle \sigma_j \sigma_k \rangle)] \\ &= \frac{1}{n} \sum_k \mathbb{E}[\langle \sigma_j \sigma_k \rangle^2] \\ &\quad + \frac{2\beta^2}{n^2} \sum_k \mathbb{E}[(\langle \sigma_i \sigma_j \rangle - \langle \sigma_i \sigma_k \rangle \langle \sigma_j \sigma_k \rangle)^2] \\ &\quad + \frac{2\beta^2}{n^2} \sum_k \mathbb{E}[\langle \sigma_j \sigma_k \rangle (\langle \sigma_j \sigma_k \rangle - \langle \sigma_i \sigma_j \rangle \langle \sigma_i \sigma_k \rangle)] \\ &\quad - \frac{2\beta^2}{n^2} \sum_k \mathbb{E}[\langle \sigma_j \sigma_k \rangle (1 - \langle \sigma_i \sigma_k \rangle^2) \langle \sigma_j \sigma_k \rangle] \\ &\quad - \frac{2\beta^2}{n^2} \sum_k \mathbb{E}[\langle \sigma_j \sigma_k \rangle \langle \sigma_i \sigma_k \rangle (\langle \sigma_i \sigma_j \rangle - \langle \sigma_i \sigma_k \rangle \langle \sigma_j \sigma_k \rangle)] \\ &= \frac{1}{n} \sum_k \mathbb{E}[\langle \sigma_j \sigma_k \rangle^2] + \frac{2\beta^2}{n^2} \sum_k \mathbb{E} [\langle \sigma_i \sigma_j \rangle^2 - 4\langle \sigma_i \sigma_j \rangle \langle \sigma_i \sigma_k \rangle \langle \sigma_j \sigma_k \rangle + 3\langle \sigma_i \sigma_k \rangle^2 \langle \sigma_j \sigma_k \rangle^2] . \end{aligned}$$

Summing over i, j we find that

$$\frac{\beta^2}{n} \sum_{i,j=1}^n \sum_{k=1}^n \mathbb{E} [g_{ik}^2 \langle \sigma_j \sigma_k \rangle^2] = \beta^2 n^2 \mathbb{E} \langle R_{12}^2 \rangle + 2\beta^4 n (\mathbb{E} \langle R_{12}^2 \rangle - 4 \mathbb{E} \langle R_{12}R_{13}R_{23} \rangle) + 3 \mathbb{E} \langle R_{12}R_{34}R_{1234} \rangle) . \quad (3.5)$$

For the off-diagonal term in Eq. (3.4), since the random variables g_{ij} are independent, we have

$$\begin{aligned} \frac{1}{n} \mathbb{E}[g_{ik}g_{i\ell} \langle \sigma_j \sigma_k \rangle \langle \sigma_j \sigma_\ell \rangle] &= \frac{\beta}{n^{3/2}} \mathbb{E}[g_{i\ell} ((\langle \sigma_i \sigma_j \rangle - \langle \sigma_j \sigma_k \rangle \langle \sigma_i \sigma_k \rangle) \langle \sigma_j \sigma_\ell \rangle + \langle \sigma_j \sigma_k \rangle (\langle \sigma_i \sigma_k \sigma_j \sigma_\ell \rangle - \langle \sigma_i \sigma_k \rangle \langle \sigma_j \sigma_\ell \rangle))] \\ &= \frac{\beta^2}{n^2} \mathbb{E}[(\langle \sigma_j \sigma_\ell \rangle - \langle \sigma_i \sigma_\ell \rangle \langle \sigma_i \sigma_j \rangle) \langle \sigma_j \sigma_\ell \rangle] \\ &\quad - \frac{\beta^2}{n^2} \mathbb{E}[(\langle \sigma_i \sigma_j \sigma_k \sigma_\ell \rangle - \langle \sigma_i \sigma_\ell \rangle \langle \sigma_j \sigma_k \rangle) \langle \sigma_i \sigma_k \rangle \langle \sigma_j \sigma_\ell \rangle] \\ &\quad - \frac{\beta^2}{n^2} \mathbb{E}[\langle \sigma_j \sigma_k \rangle (\langle \sigma_k \sigma_\ell \rangle - \langle \sigma_i \sigma_\ell \rangle \langle \sigma_i \sigma_k \rangle) \langle \sigma_j \sigma_\ell \rangle] \\ &\quad + \frac{\beta^2}{n^2} \mathbb{E}[(\langle \sigma_i \sigma_j \rangle - \langle \sigma_j \sigma_k \rangle \langle \sigma_i \sigma_k \rangle) (\langle \sigma_i \sigma_j \rangle - \langle \sigma_i \sigma_\ell \rangle \langle \sigma_j \sigma_\ell \rangle)] \\ &\quad + \frac{\beta^2}{n^2} \mathbb{E}[(\langle \sigma_i \sigma_j \sigma_k \sigma_\ell \rangle - \langle \sigma_j \sigma_k \rangle \langle \sigma_i \sigma_\ell \rangle) (\langle \sigma_i \sigma_k \sigma_j \sigma_\ell \rangle - \langle \sigma_i \sigma_k \rangle \langle \sigma_j \sigma_\ell \rangle)] \end{aligned}$$

$$\begin{aligned}
& + \frac{\beta^2}{n^2} \mathbb{E}[\langle \sigma_j \sigma_k \rangle (\langle \sigma_j \sigma_k \rangle - \langle \sigma_i \sigma_j \sigma_k \sigma_\ell \rangle \langle \sigma_i \sigma_\ell \rangle)] \\
& - \frac{\beta^2}{n^2} \mathbb{E}[\langle \sigma_j \sigma_k \rangle (\langle \sigma_k \sigma_\ell \rangle - \langle \sigma_i \sigma_k \rangle \langle \sigma_i \sigma_\ell \rangle) \langle \sigma_j \sigma_\ell \rangle] \\
& - \frac{\beta^2}{n^2} \mathbb{E}[\langle \sigma_j \sigma_k \rangle \langle \sigma_i \sigma_k \rangle (\langle \sigma_i \sigma_j \rangle - \langle \sigma_j \sigma_\ell \rangle \langle \sigma_i \sigma_\ell \rangle)].
\end{aligned}$$

Let's call the above expression O_{ijkl} . Summing this expression over *all* i, j, k, ℓ and only subtracting off the diagonal terms (corresponding to $k = \ell$) later gives

$$\begin{aligned}
\sum_{i,j,k,\ell} O_{ijkl} &= \frac{\beta^2}{n^2} \mathbb{E}[n^4 \langle R_{12}^2 \rangle - n^4 \langle R_{12} R_{13} R_{23} \rangle] \\
& - \frac{\beta^2}{n^2} \mathbb{E}[n^4 \langle R_{12}^2 R_{13}^2 \rangle - n^4 \langle R_{13} R_{24} R_{23} R_{14} \rangle] \\
& - \frac{\beta^2}{n^2} \mathbb{E}[n^4 \langle R_{12} R_{13} R_{23} \rangle - n^4 \langle R_{13} R_{14} R_{24} R_{23} \rangle] \\
& + \frac{\beta^2}{n^2} \mathbb{E}[n^4 \langle R_{12}^2 \rangle - 2n^4 \langle R_{12} R_{13} R_{23} \rangle + n^4 \langle R_{14} R_{13} R_{23} R_{24} \rangle] \\
& + \frac{\beta^2}{n^2} \mathbb{E}[n^4 \langle R_{12}^4 \rangle - 2n^4 \langle R_{12}^2 R_{13}^2 \rangle + n^4 \langle R_{14} R_{13} R_{23} R_{24} \rangle] \\
& + \frac{\beta^2}{n^2} \mathbb{E}[n^4 \langle R_{12}^2 \rangle - n^4 \langle R_{12}^2 R_{23}^2 \rangle] \\
& - \frac{\beta^2}{n^2} \mathbb{E}[n^4 \langle R_{12} R_{13} R_{23} \rangle - n^4 \langle R_{14} R_{12} R_{23} R_{34} \rangle] \\
& - \frac{\beta^2}{n^2} \mathbb{E}[n^4 \langle R_{12} R_{13} R_{23} \rangle - n^4 \langle R_{12} R_{13} R_{24} R_{34} \rangle].
\end{aligned}$$

Collecting terms and multiplying by β^2 , we get

$$\beta^2 \sum_{i,j,k,\ell} O_{ijkl} = \beta^4 n^2 \mathbb{E} [3 \langle R_{12}^2 \rangle + \langle R_{12}^4 \rangle - 6 \langle R_{12} R_{13} R_{23} \rangle - 4 \langle R_{12}^2 R_{23}^2 \rangle + 6 \langle R_{12} R_{23} R_{34} R_{14} \rangle]. \quad (3.6)$$

Now considering the diagonals of that expression where $k = \ell$, and summing over i, j, k , and multiplying by β^2 again, we get

$$\begin{aligned}
\beta^2 \sum_{i,j,k} O_{ijkk} &= \frac{\beta^4}{n^2} \mathbb{E}[n^3 \langle R_{12}^2 \rangle - n^3 \langle R_{12} R_{13} R_{23} \rangle] \\
& - \frac{\beta^4}{n^2} \mathbb{E}[n^3 \langle R_{12} R_{23} R_{13} \rangle - n^3 \langle R_{13} R_{24} R_{1234} \rangle] \\
& - \frac{\beta^4}{n^2} \mathbb{E}[n^3 \langle R_{12}^2 \rangle - n^3 \langle R_{13} R_{24} R_{1234} \rangle] \\
& + \frac{\beta^4}{n^2} \mathbb{E}[n^3 \langle R_{12}^2 \rangle - 2n^3 \langle R_{12} R_{23} R_{13} \rangle + n^3 \langle R_{13} R_{24} R_{1234} \rangle] \\
& + \frac{\beta^4}{n^2} \mathbb{E}[n^3 \langle R_{12}^2 \rangle - 2n^3 \langle R_{12} R_{23} R_{13} \rangle + n^3 \langle R_{13} R_{24} R_{1234} \rangle]
\end{aligned}$$

$$\begin{aligned}
& + \frac{\beta^4}{n^2} \mathbb{E}[n^3 \langle R_{12}^2 \rangle - n^3 \langle R_{12} R_{23} R_{13} \rangle] \\
& - \frac{\beta^4}{n^2} \mathbb{E}[n^3 \langle R_{12}^2 \rangle - n^3 \langle R_{13} R_{24} R_{1234} \rangle] \\
& - \frac{\beta^4}{n^2} \mathbb{E}[n^3 \langle R_{12} R_{23} R_{13} \rangle - n^3 \langle R_{13} R_{24} R_{1234} \rangle].
\end{aligned}$$

After simplification,

$$\beta^2 \sum_{i,j,k} O_{ijkk} = \beta^4 n \mathbb{E} [2 \langle R_{12}^2 \rangle - 8 \langle R_{12} R_{13} R_{23} \rangle + 6 \langle R_{13} R_{24} R_{1234} \rangle]. \quad (3.7)$$

Putting together (3.5), (3.6) and (3.7), we obtain

$$\begin{aligned}
\text{III} &= \beta^2 n^2 \mathbb{E}[\langle R_{12}^2 \rangle] + 2\beta^4 n (\mathbb{E}[\langle R_{12}^2 \rangle] - 4 \mathbb{E}[\langle R_{12} R_{13} R_{23} \rangle] + 3 \mathbb{E}[\langle R_{12} R_{34} R_{1234} \rangle]) \\
& + \beta^4 n^2 \mathbb{E}[3 \langle R_{12}^2 \rangle + \langle R_{12}^4 \rangle - 6 \langle R_{12} R_{13} R_{23} \rangle - 4 \langle R_{12}^2 R_{23}^2 \rangle + 6 \langle R_{12} R_{23} R_{34} R_{14} \rangle] \\
& - (\beta^4 n \mathbb{E}[2 \langle R_{12}^2 \rangle - 8 \langle R_{12} R_{13} R_{23} \rangle + 6 \langle R_{13} R_{24} R_{1234} \rangle]) \\
& = \beta^2 n^2 \mathbb{E}[\langle R_{12}^2 \rangle] \\
& + \beta^4 n^2 \mathbb{E}[3 \langle R_{12}^2 \rangle + \langle R_{12}^4 \rangle - 6 \langle R_{12} R_{13} R_{23} \rangle - 4 \langle R_{12}^2 R_{23}^2 \rangle + 6 \langle R_{12} R_{23} R_{34} R_{14} \rangle]. \quad (3.8)
\end{aligned}$$

Combining (3.2), (3.3) and (3.8), we obtain an expression for $\mathbb{E} \|P\|_F^2$:

$$\begin{aligned}
\mathbb{E} \|P\|_F^2 &= (1 + \beta^2)^2 n^2 \mathbb{E}[\langle R_{12}^2 \rangle] - 4\beta^2 (1 + \beta^2) n^2 (\mathbb{E}[\langle R_{12}^2 \rangle] - \mathbb{E}[\langle R_{12} R_{13} R_{23} \rangle]) \\
& + \beta^2 n^2 \mathbb{E}[\langle R_{12}^2 \rangle] \\
& + \beta^4 n^2 \mathbb{E}[3 \langle R_{12}^2 \rangle + \langle R_{12}^4 \rangle - 6 \langle R_{12} R_{13} R_{23} \rangle - 4 \langle R_{12}^2 R_{23}^2 \rangle + 6 \langle R_{12} R_{23} R_{34} R_{14} \rangle] \\
& = (1 - \beta^2) n^2 \mathbb{E}[\langle R_{12}^2 \rangle] + (4\beta^2 (1 + \beta^2) - 6\beta^4) n^2 \mathbb{E}[\langle R_{12} R_{13} R_{23} \rangle] \\
& + \beta^4 n^2 \mathbb{E}[\langle R_{12}^4 \rangle] - 4\beta^4 n^2 \mathbb{E}[\langle R_{12}^2 R_{23}^2 \rangle] + 6\beta^4 n^2 \mathbb{E}[\langle R_{12} R_{23} R_{34} R_{14} \rangle]. \quad (3.9)
\end{aligned}$$

From Talagrand [Tal11, Theorem 11.5.4] and Bardina, Márquez-Carreras, Rovira and Tindel [BMCRT04], we have the precise asymptotic expansions up to order $O(n^{-5/2})$ of all the overlaps involved in the above expressions:

$$\mathbb{E} \langle R_{12}^2 \rangle = \frac{1}{n(1 - \beta^2)} + \frac{-\beta^2(1 + \beta^2)}{n^2(1 - \beta^2)^4} + O(1/n^{5/2}), \quad (3.10)$$

$$\mathbb{E} \langle R_{12} R_{13} R_{23} \rangle = \frac{1}{n^2(1 - \beta^2)^3} + O(1/n^{5/2}), \quad (3.11)$$

$$\mathbb{E} \langle R_{12}^4 \rangle = \frac{3}{n^2(1 - \beta^2)^2} + O(1/n^{5/2}), \quad (3.12)$$

$$\mathbb{E} \langle R_{12}^2 R_{23}^2 \rangle = \frac{1}{n^2(1 - \beta^2)^2} + O(1/n^{5/2}) \quad (3.13)$$

$$\mathbb{E} \langle R_{12} R_{23} R_{34} R_{14} \rangle = O(1/n^{5/2}). \quad (3.14)$$

From this we deduce

$$\mathbb{E} \|P\|_F^2 = n - \frac{\beta^2(1 + \beta^2)}{(1 - \beta^2)^3} + \frac{4\beta^2(1 + \beta^2) - 6\beta^4}{(1 - \beta^2)^3} + \frac{3\beta^4}{(1 - \beta^2)^2} - \frac{4\beta^4}{(1 - \beta^2)^2} + O(1/\sqrt{n}) \quad (3.15)$$

$$= n + \frac{\beta^2(3 - 4\beta^2 + \beta^4)}{(1 - \beta^2)^3} + O(1/\sqrt{n}). \quad (3.16)$$

Finally, combining the above formula with Eq. (3.1) for $\mathbb{E} \text{Tr}(P)$ we obtain

$$\mathbb{E} [\|P - I\|_F^2] = \frac{\beta^2(1 + \beta^2)}{(1 - \beta^2)^2} + O(1/\sqrt{n}). \quad (3.17)$$

4 Lower bounds at the critical temperature and low temperatures

In this section, we investigate the tightness of our results. In particular, we show that the expected operator norm necessarily diverges as $\beta \rightarrow 1$. In the low temperature regime where $\beta > 1$, we provide a lower bound on the expected operator norm of order \sqrt{n} . Finally, we consider the behavior of the operator norm near and at the critical temperature $\beta = 1$.

Below, we write $\text{cov}(\mu_{\beta,n})$ to emphasize the dependence on the inverse temperature β and n .

4.1 High, near critical temperature

We first provide a simple argument using well-known limit theorems on the overlap $R_{1,2}$ to show that the expected operator norm necessarily diverges as $\beta \rightarrow 1^-$:

Proposition 4.1. *For $\beta < 1$, it holds that*

$$\liminf_{n \rightarrow \infty} \mathbb{E} \|\text{cov}(\mu_{\beta,n})\|_{\text{op}} \geq \sqrt{\frac{2}{\pi(1 - \beta^2)}}. \quad (4.1)$$

In particular,

$$\lim_{\beta \rightarrow 1^-} \liminf_{n \rightarrow \infty} \mathbb{E} \|\text{cov}(\mu_{\beta,n})\|_{\text{op}} = +\infty. \quad (4.2)$$

Proof. We simply lower bound

$$\mathbb{E} \|\text{cov}(\mu_{\beta,n})\|_{\text{op}} \geq \frac{\mathbb{E} \|\text{cov}(\mu_{\beta,n})\|_F}{\sqrt{n}} = \mathbb{E} [\langle nR_{12}^2 \rangle^{1/2}].$$

From Talagrand [Tal11, Theorem 11.5.4], the random variable $(n(1 - \beta^2))^{1/2} \cdot R_{12}$ converges in distribution as $n \rightarrow \infty$ to a standard Gaussian under the measure $\mathbb{E}\langle \cdot \rangle$; by the Cauchy-Schwarz inequality and the Portmanteau theorem, it follows that

$$\liminf_{n \rightarrow \infty} \mathbb{E} [\langle nR_{12}^2 \rangle^{1/2}] \geq \liminf_{n \rightarrow \infty} \mathbb{E} [\langle \sqrt{n} \cdot |R_{12}| \rangle] \geq \frac{\mathbb{E}|z|}{\sqrt{1 - \beta^2}},$$

where z is standard normal. The inequality follows from the standard fact $\mathbb{E}|z| = \sqrt{2/\pi}$. \square

Note that the previous result does not appear to immediately have any bearing on the behavior at $\beta = 1$ due to the subtlety of interchanging limits. We treat the case $\beta = 1$ below using a similar analysis and leveraging some of the few known results in this regime.

4.2 Low temperature

Next, we show that the boundedness of the operator norm cannot extend past $\beta = 1$:

Proposition 4.2. *For every $\beta > 1$, there exists a constant $c(\beta) > 0$ such that*

$$\mathbb{E} \left\| \text{cov}(\mu_{\beta,n}) \right\|_{\text{op}} \geq c(\beta) \sqrt{n}. \quad (4.3)$$

Proof. The same argument as in Proposition 4.1 yields

$$\mathbb{E} \left\| \text{cov}(\mu_{\beta,n}) \right\|_{\text{op}} \geq \sqrt{n} \mathbb{E} \left[\langle R_{12}^2 \rangle^{1/2} \right] \geq \sqrt{n} \mathbb{E} \langle R_{12}^2 \rangle,$$

where we used $|R_{12}| \leq 1$ in the last inequality. From Talagrand [Tal11, Equation (14.417)], it holds that

$$\lim_{n \rightarrow \infty} \mathbb{E} \langle R_{12}^2 \rangle = \int q^2 \zeta_\beta^*(dq), \quad (4.4)$$

where ζ_β^* —called the Parisi measure—is the unique minimizer of the Parisi functional; see [AC15] and [Tal11, Pan13] for definitions. It is further known that $\zeta_\beta^* \neq \delta_0$ when $\beta > 1$, so the right-hand side in (4.4) is some constant $c(\beta) > 0$; see [AMS22, Section 5.2] for more details. The previous two displays yield the desired conclusion. \square

4.3 At the transition

To complete this picture, we now consider the operator norm near and at the critical temperature $\beta = 1$. The next result refines Proposition 4.1 by showing that for a sequence $\beta_n \rightarrow 1^-$, one can obtain similar lower bounds on the operator norm with different exponents:

Proposition 4.3. *Fix any function $f(n)$ satisfying $f(n) \rightarrow \infty$ and $f(n) = o(n^{1/6})$. Then there exists a sequence $\beta_n \rightarrow 1^-$ such that*

$$\mathbb{E} \left\| \text{cov}(\mu_{\beta_n,n}) \right\|_{\text{op}} \geq f(n). \quad (4.5)$$

Proof. As usual, we always have

$$\mathbb{E} \left\| \text{cov}(\mu_{\beta,n}) \right\|_{\text{op}} \geq \sqrt{n} \mathbb{E} \left[\langle R_{12}^2 \rangle^{1/2} \right] \geq \sqrt{n} \mathbb{E} \langle |R_{12}| \rangle,$$

where the last inequality is by the Cauchy-Schwarz inequality.

Define $g(n) = f(n)/n^{1/6} = o(1)$ by assumption. Then, set β_n such that $n^{1/3}(1 - \beta_n^2) = 1/g(n)^2 = \omega(1)$. From Talagrand [Tal11, Theorem 11.7.1], these conditions imply that $(n(1 - \beta_n^2))^{1/2} \cdot R_{12}$ converges in distribution to a standard Gaussian with respect to the measure $\mathbb{E} \langle \cdot \rangle^{\otimes 2}$, so an analogous argument using the Portmanteau theorem as in Proposition 4.1 implies that

$$\mathbb{E} \langle |R_{12}| \rangle \geq \frac{c}{\sqrt{n(1 - \beta_n^2)}} = \frac{cg(n)}{n^{1/3}} = \frac{cf(n)}{n^{1/2}},$$

for some constant $c > 0$. Combining the two previous displays proves the desired inequality. \square

Remark 1. *We remark that this result is essentially the best lower bound one can obtain at the critical point $\beta = 1$ using this method of lower bounding the operator norm by the Frobenius norm divided by \sqrt{n} . Indeed, it is expected that $\mathbb{E} \langle R_{12}^2 \rangle = \Theta(1/n^{2/3})$ at $\beta = 1$. Assuming this, the lower bound in the above argument would be at most $C\sqrt{n}/n^{1/3} = Cn^{1/6}$ for some constant $C > 0$.*

Finally, we directly consider the behavior at the critical temperature $\beta = 1$. While it is expected and intuitive that $\mathbb{E}\langle R_{12}^2 \rangle$ is increasing with respect to β , which would imply a lower bound from the previous results, this does not appear to be known. Instead, we show the following bound on the second moment which is somewhat weaker, but holds unconditionally by applying known results in this setting:

Proposition 4.4. *For $\beta = 1$, there is an absolute constant $c > 0$ such that*

$$\mathbb{E} \left\| \text{cov}(\mu_{\beta,n}) \right\|_{\text{op}}^2 \geq c \left(\frac{n}{\log n} \right)^{3/16}. \quad (4.6)$$

Proof. We leverage some known results on overlap convergence at $\beta = 1$. First, a result of Chatterjee [Tal11, Theorem 11.7.6] shows a lower bound on the third moment of $R_{1,2}$: there exists a constant $L = L(\beta) > 0$ such that

$$\mathbb{E} \langle |R_{12}|^3 \rangle \geq \frac{1}{Ln}. \quad (4.7)$$

Next, we need a bound on the decay of tail probability of $R_{1,2}$. This is the content of a result of Talagrand [Tal03, Theorem 2.14.5] which we quote here: for $\beta \leq 1$ and for all $x \geq L(\log n/n)^{3/8}$ where $L = L(\beta) < \infty$, it holds that

$$\mathbb{P}(|R_{12}| \geq \sqrt{x}) \leq Ln \exp(-nx^2/L) + \exp\left(-n^{3/2}x^4/(L\sqrt{\log n})\right), \quad (4.8)$$

where the probability is over the distribution of R_{12} induced by the measure $\mathbb{E}\langle \cdot \rangle^{\otimes 2}$. Taking $x = \max\{1, (2/L^3)^{1/4}\} \cdot L(\log n/n)^{3/8}$, we find that for n large enough,

$$\mathbb{P}(|R_{12}| \geq \sqrt{x}) \leq \frac{2}{n^2}. \quad (4.9)$$

Combining these results shows that

$$\frac{1}{Ln} \leq \mathbb{E} \langle |R_{12}|^3 \mathbf{1}_{|R_{1,2}| \geq \sqrt{x}} \rangle + \mathbb{E} \langle |R_{12}|^3 \mathbf{1}_{|R_{1,2}| < \sqrt{x}} \rangle \leq \mathbb{P}(|R_{12}| \geq \sqrt{x}) + \sqrt{x} \mathbb{E} \langle R_{12}^2 \rangle.$$

Using (4.9) and rearranging yields

$$\mathbb{E} \langle R_{12}^2 \rangle \geq \frac{c}{n^{13/16}(\log n)^{3/16}}, \quad (4.10)$$

for some constant $c > 0$. The result follows from the previous display with the simple lower bound

$$\mathbb{E} \left\| \text{cov}(\mu_{\beta,n}) \right\|_{\text{op}}^2 \geq \frac{\mathbb{E} \left\| \text{cov}(\mu_{\beta,n}) \right\|_F^2}{n} = n \mathbb{E} \langle R_{12}^2 \rangle.$$

□

We finally remark that when $\beta < 1$, we can prove a constant upper bound on $\mathbb{E} \left\| \text{cov}(\mu_{\beta,n}) \right\|_{\text{op}}^2$ by the same argument used to show our main result, Theorem 1.1. Thus the previous result, Proposition 4.4, shows a quantitative transition in the value of the second moment of the operator norm at $\beta = 1$.

Acknowledgments. The first author is grateful to Ronen Eldan for valuable conversations about bounding the covariance matrix of disordered spin systems which inspired this work.

References

- [ABvSY21] Arka Adhikari, Christian Brennecke, Per von Soosten, and Horng-Tzer Yau, *Dynamical approach to the TAP equations for the Sherrington–Kirkpatrick model*, Journal of Statistical Physics **183** (2021), no. 3, 1–27.
- [AC15] Antonio Auffinger and Wei-Kuo Chen, *The Parisi formula has a unique minimizer*, Communications in Mathematical Physics **335** (2015), no. 3, 1429–1444.
- [AMS22] Ahmed El Alaoui, Andrea Montanari, and Mark Sellke, *Sampling from the Sherrington–Kirkpatrick Gibbs measure via algorithmic stochastic localization*, arXiv preprint arXiv:2203.05093 (2022).
- [BMCRT04] Xavier Bardina, David Márquez-Carreras, Carles Rovira, and Samy Tindel, *Higher order expansions for the overlap of the SK model*, Seminar on Stochastic Analysis, Random Fields and Applications IV, Springer, 2004, pp. 21–43.
- [Cel22] Michael Celentano, *Sudakov-Fernique post-AMP, and a new proof of the local convexity of the TAP free energy*, arXiv preprint arXiv:2208.09550 (2022).
- [CFM21] Michael Celentano, Zhou Fan, and Song Mei, *Local convexity of the TAP free energy and AMP convergence for \mathbf{Z}_2 -synchronization*, arXiv preprint arXiv:2106.11428 (2021).
- [CPS18] Wei-Kuo Chen, Dmitry Panchenko, and Eliran Subag, *Generalized TAP free energy*, Communications on Pure and Applied Mathematics (2018).
- [EKZ21] Ronen Eldan, Frederic Koehler, and Ofer Zeitouni, *A spectral condition for spectral gap: fast mixing in high-temperature Ising models*, Probability Theory and Related Fields (2021), 1–17.
- [Pan13] Dmitry Panchenko, *The Sherrington–Kirkpatrick model*, Springer Science & Business Media, 2013.
- [Tal03] Michel Talagrand, *Spin Glasses: A Challenge for Mathematicians: Cavity and Mean Field Models*, vol. 46, Springer Science & Business Media, 2003.
- [Tal11] ———, *Mean Field Models for Spin Glasses. Volume II: Advanced Replica-Symmetry and Low Temperature*, vol. 55, Springer Science & Business Media, 2011.
- [Ver18] Roman Vershynin, *High-Dimensional Probability: An Introduction with Applications in Data Science*, vol. 47, Cambridge university press, 2018.