

A symmetric chain decomposition of $L(5, n)$

Xiangdong Wen
Wolfram Research

January 10, 2023

1 Introduction

For positive integers m and n , **Young's lattice** $L(m, n)$ refers to the partially ordered set (**poset**) of m -tuples (a_1, a_2, \dots, a_m) , where $0 \leq a_1 \leq a_2 \leq \dots \leq a_m \leq n$ with ordered relation $(a_1, a_2, \dots, a_m) \leq (b_1, b_2, \dots, b_m)$ if $a_i \leq b_i$ for all $i = 1, 2, \dots, m$.

The **rank** of $\vec{a} = (a_1, a_2, \dots, a_m)$ is defined as $rank(\vec{a}) = \sum_{i=1}^m a_i$, and a chain $\vec{v}_1 \leq \vec{v}_2 \leq \dots \leq \vec{v}_k$ in $L(m, n)$ is called **saturated** if it skips no ranks and is called **symmetric** if $rank(\vec{v}_1) + rank(\vec{v}_k) = mn$. A Symmetric Chain Decomposition (**SCD**) of a poset is a way of expressing it as a disjoint union of saturated symmetric chains.

Various SCDs for $L(4, n)$ and $L(3, n)$ have been constructed ([4], [1] and [3]). One of the major problems in order theory is the explicit construction of SCDs for Young's Lattice for all m and n . In 1989, Kathy O'Hara ([2], see also [5]) constructed SCDs for the trivial extension of $L(m, n)$, in which all partitions of one rank are related to the next; but the problem remains wide open for Young's lattice itself.

There are more than 30 types of symmetric chains in the found SCD of $L(5, n)$. To make it neat and also clear, nine types of **parallel** chains are presented in section 2. We illustrate how to get symmetric chains from these parallel chains in section 3 and finally give a computer proof in section 4.

2 A list of disjoint parallel chains of $L(5, n)$

Here are nine types of parallel chains of $L(5, n)$ that a SCD of $L(5, n)$ could be derived from. Parameters i, j, k, u and w in these tables are all non-negative integers. One line of vertical dots in these tables represents an abbreviated saturated chain with only one entry increasing. Two lines of vertical dots (in the tables C_3, C_4, C_5 and C_6) represent an abbreviated saturated chain with a zigzag path that two entries are increasing 1 alternatively. There are two helper rows with extra parameter t , an increasing iterator, in the zigzag path. For example, the first and the third entries are increasing by 1 alternatively at the beginning of chain C_3 .

$2 + 2i + 2j + 3k \leq n$				
$(p,$	$k,$	$j + k,$	$1 + i + j + 2k,$	$1 + i + 2j + 2k)$
		\vdots		
$(p,$	$k,$	$1 + i + j + k + p,$	$1 + i + j + 2k,$	$1 + i + 2j + 2k)$
	\vdots			
$(p,$	$1 + i + j + k,$	$1 + i + j + k + p,$	$1 + i + j + 2k,$	$1 + i + 2j + 2k)$
\vdots				
$(1 + i + k,$	$1 + i + j + k,$	$1 + i + j + k + p,$	$1 + i + j + 2k,$	$1 + i + 2j + 2k)$
				\vdots
$(1 + i + k,$	$1 + i + j + k,$	$1 + i + j + k + p,$	$1 + i + j + 2k,$	$-k + n + p)$
			\vdots	
$(1 + i + k,$	$1 + i + j + k,$	$1 + i + j + k + p,$	$-k + n,$	$-k + n + p)$
		\vdots		
$(1 + i + k,$	$1 + i + j + k,$	$-j - k + n,$	$-k + n,$	$-k + n + p)$
	\vdots			
$(1 + i + k,$	$-1 - i - j - 2k + n,$	$-j - k + n,$	$-k + n,$	$-k + n + p)$
\vdots				
$(-1 - i - 2j - 2k + n,$	$-1 - i - j - 2k + n,$	$-j - k + n,$	$-k + n,$	$-k + n + p)$

Table 1: parallel chains C_1 , $0 \leq p \leq k$

$3 + 2i + 2j + 3k \leq n$				
$(k - p,$	$k,$	$j + k,$	$1 + i + j + 2k,$	$2 + i + 2j + 2k)$
				\vdots
$(k - p,$	$k,$	$j + k,$	$1 + i + j + 2k,$	$-1 - i - k + n)$
			\vdots	
$(k - p,$	$k,$	$j + k,$	$-1 - i - j - k + n,$	$-1 - i - k + n)$
		\vdots		
$(k - p,$	$k,$	$-1 - i - j - k + n - p,$	$-1 - i - j - k + n,$	$-1 - i - k + n)$
	\vdots			
$(k - p,$	$-1 - i - j - 2k + n,$	$-1 - i - j - k + n - p,$	$-1 - i - j - k + n,$	$-1 - i - k + n)$
\vdots				
$(-1 - i - 2j - 2k + n,$	$-1 - i - j - 2k + n,$	$-1 - i - j - k + n - p,$	$-1 - i - j - k + n,$	$-1 - i - k + n)$
				\vdots
$(-1 - i - 2j - 2k + n,$	$-1 - i - j - 2k + n,$	$-1 - i - j - k + n - p,$	$-1 - i - j - k + n,$	$n - p)$
			\vdots	
$(-1 - i - 2j - 2k + n,$	$-1 - i - j - 2k + n,$	$-1 - i - j - k + n - p,$	$-k + n,$	$n - p)$
		\vdots		
$(-1 - i - 2j - 2k + n,$	$-1 - i - j - 2k + n,$	$-1 - j - k + n,$	$-k + n,$	$n - p)$

Table 2: parallel chains C_2 , $0 \leq p \leq k$

$0 \leq u \leq 1$				
$2u + 1 + 6j + 4k + 3i \leq n$				
$2 \leq t \leq i - 1$				
$(2j,$	$i + 2j + p,$	$i + 2j + 2k + u,$	$2i + 4j + 2k + u,$	$2i + 4j + 4k + 2u)$
$(1 + 2j,$	$i + 2j + p,$	$i + 2j + 2k + u,$	$2i + 4j + 2k + u,$	$2i + 4j + 4k + 2u)$
$(1 + 2j,$	$i + 2j + p,$	$1 + i + 2j + 2k + u,$	$2i + 4j + 2k + u,$	$2i + 4j + 4k + 2u)$
\vdots		\vdots		
$(t + 2j,$	$i + 2j + p,$	$t - 1 + i + 2j + 2k + u,$	$2i + 4j + 2k + u,$	$2i + 4j + 4k + 2u)$
$(t + 2j,$	$i + 2j + p,$	$t + i + 2j + 2k + u,$	$2i + 4j + 2k + u,$	$2i + 4j + 4k + 2u)$
\vdots		\vdots		
$(i + 2j,$	$i + 2j + p,$	$2i - 1 + 2j + 2k + u,$	$2i + 4j + 2k + u,$	$2i + 4j + 4k + 2u)$
$(i + 2j,$	$i + 2j + p,$	$2i + 2j + 2k + u,$	$2i + 4j + 2k + u,$	$2i + 4j + 4k + 2u)$
				\vdots
$(i + 2j,$	$i + 2j + p,$	$2i + 2j + 2k + u,$	$2i + 4j + 2k + u,$	$-2j + n)$
			\vdots	
$(i + 2j,$	$i + 2j + p,$	$2i + 2j + 2k + u,$	$-i - 2j - 2k + n + p - u,$	$-2j + n)$
			\vdots	
$(i + 2j,$	$i + 2j + p,$	$-i - 2j - 2k + n - u,$	$-i - 2j - 2k + n + p - u,$	$-2j + n)$
	\vdots			
$(i + 2j,$	$-2i - 4j - 2k + n - u,$	$-i - 2j - 2k + n - u,$	$-i - 2j - 2k + n + p - u,$	$-2j + n)$
\vdots				
$(-2i - 4j - 4k + n - 2u,$	$-2i - 4j - 2k + n - u,$	$-i - 2j - 2k + n - u,$	$-i - 2j - 2k + n + p - u,$	$-2j + n)$

Table 3: parallel chains C_3 , $0 \leq p \leq 2k + u$

$0 \leq u \leq 1$				
$2u + 4 + 6j + 4k + 3i \leq n$				
$2 \leq t \leq i - 1$				
$(1 + 2j,$	$1 + i + 2j + p,$	$1 + i + 2j + 2k + u,$	$2 + 2i + 4j + 2k + u,$	$3 + 2i + 4j + 4k + 2u)$
$(1 + 2j,$	$1 + i + 2j + p,$	$2 + i + 2j + 2k + u,$	$2 + 2i + 4j + 2k + u,$	$3 + 2i + 4j + 4k + 2u)$
\vdots		\vdots		
$(t + 2j,$	$1 + i + 2j + p,$	$t + i + 2j + 2k + u,$	$2 + 2i + 4j + 2k + u,$	$3 + 2i + 4j + 4k + 2u)$
$(t + 2j,$	$1 + i + 2j + p,$	$1 + t + i + 2j + 2k + u,$	$2 + 2i + 4j + 2k + u,$	$3 + 2i + 4j + 4k + 2u)$
\vdots		\vdots		
$(i + 2j,$	$1 + i + 2j + p,$	$2i + 2j + 2k + u,$	$2 + 2i + 4j + 2k + u,$	$3 + 2i + 4j + 4k + 2u)$
$(i + 2j,$	$1 + i + 2j + p,$	$1 + 2i + 2j + 2k + u,$	$2 + 2i + 4j + 2k + u,$	$3 + 2i + 4j + 4k + 2u)$
$(1 + i + 2j,$	$1 + i + 2j + p,$	$1 + 2i + 2j + 2k + u,$	$2 + 2i + 4j + 2k + u,$	$3 + 2i + 4j + 4k + 2u)$
				\vdots
$(1 + i + 2j,$	$1 + i + 2j + p,$	$1 + 2i + 2j + 2k + u,$	$2 + 2i + 4j + 2k + u,$	$-1 - 2j + n)$
			\vdots	
$(1 + i + 2j,$	$1 + i + 2j + p,$	$1 + 2i + 2j + 2k + u,$	$-1 - i - 2j - 2k + n + p - u,$	$-1 - 2j + n)$
			\vdots	
$(1 + i + 2j,$	$1 + i + 2j + p,$	$-1 - i - 2j - 2k + n - u,$	$-1 - i - 2j - 2k + n + p - u,$	$-1 - 2j + n)$
	\vdots			
$(1 + i + 2j,$	$-2 - 2i - 4j - 2k + n - u,$	$-1 - i - 2j - 2k + n - u,$	$-1 - i - 2j - 2k + n + p - u,$	$-1 - 2j + n)$
\vdots				
$(-3 - 2i - 4j - 4k + n - 2u,$	$-2 - 2i - 4j - 2k + n - u,$	$-1 - i - 2j - 2k + n - u,$	$-1 - i - 2j - 2k + n + p - u,$	$-1 - 2j + n)$

Table 4: parallel chains C_4 , $0 \leq p \leq 2k + u$

$0 \leq u \leq 1$ $2u + 4 + 6j + 4k + 3i \leq n$ $1 \leq t \leq i - 2$				
(2j,	$1 + i + 2j + p,$	$1 + i + 2j + 2k + u,$	$2 + 2i + 4j + 2k + u,$	$3 + 2i + 4j + 4k + 2u$
				\vdots
(2j,	$1 + i + 2j + p,$	$1 + i + 2j + 2k + u,$	$2 + 2i + 4j + 2k + u,$	$-1 - i - 2j + n$
			\vdots	
(2j,	$1 + i + 2j + p,$	$1 + i + 2j + 2k + u,$	$-1 - i - 2j - 2k + n + p - u,$	$-1 - i - 2j + n$
		\vdots		
(2j,	$1 + i + 2j + p,$	$-2 - 2i - 2j - 2k + n - u,$	$-1 - i - 2j - 2k + n + p - u,$	$-1 - i - 2j + n$
	\vdots			
(2j,	$-2 - 2i - 4j - 2k + n - u,$	$-2 - 2i - 2j - 2k + n - u,$	$-1 - i - 2j - 2k + n + p - u,$	$-1 - i - 2j + n$
	\vdots			
(-2 - 2i - 4j - 4k + n - 2u,	$-2 - 2i - 4j - 2k + n - u,$	$-1 - 2i - 2j - 2k + n - u,$	$-1 - i - 2j - 2k + n + p - u,$	$-1 - i - 2j + n$
(-2 - 2i - 4j - 4k + n - 2u,	$-2 - 2i - 4j - 2k + n - u,$	$-1 - 2i - 2j - 2k + n - u,$	$-1 - i - 2j - 2k + n + p - u,$	$-i - 2j + n$
		\vdots		
(-2 - 2i - 4j - 4k + n - 2u,	$-2 - 2i - 4j - 2k + n - u,$	$-1 + t - 2i - 2j - 2k + n - u,$	$-1 - i - 2j - 2k + n + p - u,$	$-1 + t - i - 2j + n$
(-2 - 2i - 4j - 4k + n - 2u,	$-2 - 2i - 4j - 2k + n - u,$	$-1 + t - 2i - 2j - 2k + n - u,$	$-1 - i - 2j - 2k + n + p - u,$	$t - i - 2j + n$
		\vdots		
(-2 - 2i - 4j - 4k + n - 2u,	$-2 - 2i - 4j - 2k + n - u,$	$-2 - i - 2j - 2k + n - u,$	$-1 - i - 2j - 2k + n + p - u,$	$-2 - 2j + n$
(-2 - 2i - 4j - 4k + n - 2u,	$-2 - 2i - 4j - 2k + n - u,$	$-2 - i - 2j - 2k + n - u,$	$-1 - i - 2j - 2k + n + p - u,$	$-1 - 2j + n$
(-2 - 2i - 4j - 4k + n - 2u,	$-2 - 2i - 4j - 2k + n - u,$	$-1 - i - 2j - 2k + n - u,$	$-1 - i - 2j - 2k + n + p - u,$	$-1 - 2j + n$

Table 5: parallel chains C_5 , $0 \leq p \leq 2k + u$

$0 \leq u \leq 1$ $2u + 7 + 6j + 4k + 3i \leq n$ $1 \leq t \leq i - 2$				
(1 + 2j,	$2 + i + 2j + p,$	$2 + i + 2j + 2k + u,$	$4 + 2i + 4j + 2k + u,$	$6 + 2i + 4j + 4k + 2u$
				\vdots
(1 + 2j,	$2 + i + 2j + p,$	$2 + i + 2j + 2k + u,$	$4 + 2i + 4j + 2k + u,$	$-2 - i - 2j + n$
			\vdots	
(1 + 2j,	$2 + i + 2j + p,$	$2 + i + 2j + 2k + u,$	$-2 - i - 2j - 2k + n + p - u,$	$-2 - i - 2j + n$
		\vdots		
(1 + 2j,	$2 + i + 2j + p,$	$-3 - 2i - 2j - 2k + n - u,$	$-2 - i - 2j - 2k + n + p - u,$	$-2 - i - 2j + n$
	\vdots			
(1 + 2j,	$-4 - 2i - 4j - 2k + n - u,$	$-3 - 2i - 2j - 2k + n - u,$	$-2 - i - 2j - 2k + n + p - u,$	$-2 - i - 2j + n$
	\vdots			
(-5 - 2i - 4j - 4k + n - 2u,	$-4 - 2i - 4j - 2k + n - u,$	$-3 - 2i - 2j - 2k + n - u,$	$-2 - i - 2j - 2k + n + p - u,$	$-2 - i - 2j + n$
(-5 - 2i - 4j - 4k + n - 2u,	$-4 - 2i - 4j - 2k + n - u,$	$-3 - 2i - 2j - 2k + n - u,$	$-2 - i - 2j - 2k + n + p - u,$	$-1 - i - 2j + n$
(-5 - 2i - 4j - 4k + n - 2u,	$-4 - 2i - 4j - 2k + n - u,$	$-2 - 2i - 2j - 2k + n - u,$	$-2 - i - 2j - 2k + n + p - u,$	$-1 - i - 2j + n$
		\vdots		
(-5 - 2i - 4j - 4k + n - 2u,	$-4 - 2i - 4j - 2k + n - u,$	$-3 + t - 2i - 2j - 2k + n - u,$	$-2 - i - 2j - 2k + n + p - u,$	$-1 + t - i - 2j + n$
(-5 - 2i - 4j - 4k + n - 2u,	$-4 - 2i - 4j - 2k + n - u,$	$-2 + t - 2i - 2j - 2k + n - u,$	$-2 - i - 2j - 2k + n + p - u,$	$-1 + t - i - 2j + n$
		\vdots		
(-5 - 2i - 4j - 4k + n - 2u,	$-4 - 2i - 4j - 2k + n - u,$	$-4 - i - 2j - 2k + n - u,$	$-2 - i - 2j - 2k + n + p - u,$	$-2 - 2j + n$
(-5 - 2i - 4j - 4k + n - 2u,	$-4 - 2i - 4j - 2k + n - u,$	$-3 - i - 2j - 2k + n - u,$	$-2 - i - 2j - 2k + n + p - u,$	$-2 - 2j + n$
(-5 - 2i - 4j - 4k + n - 2u,	$-4 - 2i - 4j - 2k + n - u,$	$-3 - i - 2j - 2k + n - u,$	$-2 - i - 2j - 2k + n + p - u,$	$-1 - 2j + n$

Table 6: parallel chains C_6 , $0 \leq p \leq 2k + u$

$u = n \bmod 2, u \in \{0, 1\}$ $6 + 3u + 6w + 6i + 2k = n$				
(1 + i,	$2 + 2i + p + u + 2w,$	$3 + 3i + k + u + 2w,$	$4 + 4i + k + q + 2u + 4w,$	$4 + 4i + 2k + 2u + 4w$
	\vdots			
(1 + i + u + 2w,	$2 + 2i + p + u + 2w,$	$3 + 3i + k + u + 2w,$	$4 + 4i + k + q + 2u + 4w,$	$4 + 4i + 2k + 2u + 4w$
		\vdots		
(1 + i + u + 2w,	$2 + 2i + p + u + 2w,$	$3 + 3i + k + 2u + 4w,$	$4 + 4i + k + q + 2u + 4w,$	$4 + 4i + 2k + 2u + 4w$
(1 + i + u + 2w,	$2 + 2i + p + u + 2w,$	$3 + 3i + k + 2u + 4w,$	$4 + 4i + k + q + 2u + 4w,$	$5 + 4i + 2k + 2u + 4w$
	\vdots			
(1 + 2i + u + 2w,	$2 + 2i + p + u + 2w,$	$3 + 3i + k + 2u + 4w,$	$4 + 4i + k + q + 2u + 4w,$	$5 + 4i + 2k + 2u + 4w$
				\vdots
(1 + 2i + u + 2w,	$2 + 2i + p + u + 2w,$	$3 + 3i + k + 2u + 4w,$	$4 + 4i + k + q + 2u + 4w,$	$5 + 5i + 2k + 3u + 6w$
(2 + 2i + u + 2w,	$2 + 2i + p + u + 2w,$	$3 + 3i + k + 2u + 4w,$	$4 + 4i + k + q + 2u + 4w,$	$5 + 5i + 2k + 3u + 6w$

Table 7: parallel chains C_7 , $0 \leq p \leq k, 0 \leq q \leq p$

$u = n \bmod 2, u \in \{0, 1\}$ $12 - 3u + 6w + 6i + 2k = n$				
$(1 + i,$	$4 + 2i + p - u + 2w,$	$5 + 3i + k - u + 2w,$	$8 + 4i + k + q - 2u + 4w,$	$9 + 4i + 2k - 2u + 4w)$
\vdots				
$(3 + 2i - u + 2w,$	$4 + 2i + p - u + 2w,$	$5 + 3i + k - u + 2w,$	$8 + 4i + k + q - 2u + 4w,$	$9 + 4i + 2k - 2u + 4w)$
\vdots				
$(3 + 2i - u + 2w,$	$4 + 2i + p - u + 2w,$	$5 + 3i + k - u + 2w,$	$8 + 4i + k + q - 2u + 4w,$	$9 + 5i + 2k - 2u + 4w)$
$(4 + 2i - u + 2w,$	$4 + 2i + p - u + 2w,$	$5 + 3i + k - u + 2w,$	$8 + 4i + k + q - 2u + 4w,$	$9 + 5i + 2k - 2u + 4w)$
\vdots				
$(4 + 2i - u + 2w,$	$4 + 2i + p - u + 2w,$	$7 + 3i + k - 2u + 4w,$	$8 + 4i + k + q - 2u + 4w,$	$9 + 5i + 2k - 2u + 4w)$
\vdots				
$(4 + 2i - u + 2w,$	$4 + 2i + p - u + 2w,$	$7 + 3i + k - 2u + 4w,$	$8 + 4i + k + q - 2u + 4w,$	$10 + 5i + 2k - 3u + 6w)$

Table 8: parallel chains $C_8, 0 \leq p \leq k, 0 \leq q \leq p$

$u = n \bmod 2, u \in \{0, 1\}$ $2k + 3u + 6w = n$				
$(0,$	$p + u + 2w,$	$k + u + 2w,$	$k + q + 2u + 4w,$	$2k + 2u + 4w)$
\vdots				
$(u + 2w,$	$p + u + 2w,$	$k + u + 2w,$	$k + q + 2u + 4w,$	$2k + 2u + 4w)$
\vdots				
$(u + 2w,$	$p + u + 2w,$	$k + 2u + 4w,$	$k + q + 2u + 4w,$	$2k + 2u + 4w)$
\vdots				
$(u + 2w,$	$p + u + 2w,$	$k + 2u + 4w,$	$k + q + 2u + 4w,$	$2k + 3u + 6w)$

Table 9: parallel chains $C_9, 0 \leq p \leq k, 0 \leq q \leq p$

3 Getting symmetric chains from parallel chains

For fixed parameters i, j, k , chains of C_1 with parameter $p = 0, 1, \dots, k$ form a rectangle illustrated in the following figure.




Figure 1: A rectangle formed by parallel chains in C_1 for fixed i, j, k .

The perimeter of the rectangle contains up to 2 symmetric chains from the left up corner (the lattice point with the lowest coordinates) to the right bottom corner (the lattice point with the highest coordinates). The two corner lattice

points should be in the same chain and in our cases they could be in either of the 2 border chains. All symmetric chains are obtained by taking perimeters off these rectangles recursively, illustrated in the following figure.




Figure 2: Getting symmetric chains from parallel chains

We could get symmetric chains from parallel chains C_2, C_3, \dots, C_6 similarly.

Parameters (q, p) with conditions $0 \leq p \leq k, 0 \leq q \leq p$ in C_7 (C_8, C_9) are actually elements in $L(2, k)$ which has a SCD

$$(t, t) \rightarrow (t, t+1) \rightarrow (t, t+2) \rightarrow \dots \rightarrow (t, k-t) \rightarrow (t+1, k-t) \rightarrow \dots \rightarrow (k-t, k-t)$$

where $0 \leq t \leq \lfloor \frac{k}{2} \rfloor$.

Each element in a symmetric chain of $L(2, k)$ corresponds to a $L(5, n)$ chain in C_7 (C_8, C_9), and these chains together also form a rectangle illustrated in the following figure and symmetric chains could be obtained by taking perimeters off these rectangles recursively.




Figure 3: A rectangle formed by parallel chains in C_7 (C_8, C_9) for fixed $u, w, i, k, 0 \leq t \leq \lfloor \frac{k}{2} \rfloor$

4 Proof

It is trivial to check that the chains obtained in section 3 are saturated and symmetric. Define **weight** of a vector $\vec{a} = (a_1, a_2, \dots, a_m)$ in $L(m, n)$ with commutative variables $x_0, x_1, x_2, \dots, x_m$ as

$$w(\vec{a}) = (x_0)^{n-a_m} (x_1)^{a_m-a_{m-1}} (x_2)^{a_{m-1}-a_{m-2}} \dots (x_{m-1})^{a_2-a_1} (x_m)^{a_1}$$

For a fixed m , it is easy to see that the total weight

$$\sum_{n=0}^{\infty} \sum_{\vec{a} \in L(m, n)} w(\vec{a})$$

is an ordinary multivariate generating function

$$F(x_0, x_1, \dots, x_m) = \frac{1}{(1-x_0)(1-x_1)(1-x_2)\dots(1-x_m)}$$

Each term in the expanded power series of $F(x_0, x_1, \dots, x_m)$ has coefficient 1 and corresponds to a unique vector in $L(m, n)$. On the other hand, for each vector in $L(m, n)$, there is a unique corresponding term in the power series of $F(x_0, x_1, \dots, x_m)$. Therefore, to prove that each vector in $L(5, n)$ appears only once in the SCD is the same as to prove that the total weights of the vectors in the chains is $F(x_0, x_1, x_2, x_3, x_4, x_5)$. As our symmetric chains are derived from parallel chains, we only need to prove

$$\sum_{n=0}^{\infty} \sum_{i=1}^9 \sum_{\vec{a} \in C_i} w(\vec{a}) = \frac{1}{(1-x_0)(1-x_1)(1-x_2)(1-x_3)(1-x_4)(1-x_5)}$$

This is done by computer program summing up all the weights of items in chains C_1, C_2, \dots, C_9 . The proof in Wolfram Language is available at <https://www.wolframcloud.com/obj/xwen/Published/SCDL5nProof.nb>.

$SCDL5n[n]$ which gives a SCD of $L(5, n)$ written in Wolfram Language is available at <https://www.wolframcloud.com/obj/xwen/Published/SCDL5n.nb>.

Acknowledgment

I am indebted to Dr. Kathy O'Hara for discussions which lead to finding parallel chains C_1 and C_2 . I am indebted to Dr. Zeilberger for his help and encouragement to write up this paper. I am also grateful to Cheng Yang for discussions and his encouragement to work on the problem.

References

- [1] Lindström B. *A partition of $L(3, n)$ into saturated chains*, European J. Comb, 61-63,1(1980).

- [2] O'Hara K.M. *Unimodality of Gaussian coefficients: a constructive proof*, J. Combin. Theory Ser. A 29-52, 53 (1990).
- [3] Wen X., *Computer-generated symmetric chain decompositions for $L(4, n)$ and $L(3, n)$* Adv. Appl. Math., v. 33 (2004) 409-412.
- [4] West D.B. *A Symmetric chain decomposition of $L(4, n)$* , European J. Comb, 379-383,1(1980).
- [5] Zeilberger D. *Kathy O'Hara's Constructive proof of the Unimodality of the Gaussian Polynomials*, American Mathematical Monthly, 590-602,96(1989)