

A REMARK ON A WEIGHTED VERSION OF SUITA CONJECTURE FOR HIGHER DERIVATIVES

QIAN GUAN, XUN SUN, AND ZHENG YUAN

ABSTRACT. In this article, we consider the set of points for the holding of the equality in a weighted version of Suita conjecture for higher derivatives, and give relations between the set and the integer valued points of a class of harmonic functions (maybe multi-valued). For planar domains bounded by finite analytic closed curves, we give relations between the set and Dirichlet problem.

1. INTRODUCTION

Let Ω be an open Riemann surface, which admits a nontrivial Green function G_Ω . Let $z_0 \in \Omega$, and let w be a local coordinate on a neighborhood V_{z_0} of z_0 satisfying $w(z_0)=0$. Let $c_\beta(z)$ be the logarithmic capacity (see[16]) on Ω which is defined by

$$c_\beta(z_0) := \exp \lim_{z \rightarrow z_0} (G_\Omega(z, z_0) - \log |w(z)|).$$

Let $B_\Omega(z_0)$ be the Bergman kernel function on Ω . An open question was posed by Sario and Oikawa in ([16]): find a relation between the magnitudes of the quantities $\sqrt{\pi B_\Omega(z)}$ and $c_\beta(z)$. In ([17]), Suita posed the following conjecture.

Conjecture 1.1. $\pi B_\Omega(z_0) \geq (c_\beta(z_0))^2$ holds, and the equality holds if and only if Ω is conformally equivalent to the unit disc less a (possible) closed set of inner capacity zero.

The inequality part of Suita conjecture for bounded planar domain was proved by Blocki ([3]), and original form of inequality part was proved by Guan-Zhou([11]). The equality part of Suita conjecture was proved by Guan-Zhou ([12]), which completed the proof of Suita conjecture.

We recall some notations (see [7], see also [12, 9]). Let $p : \Delta \rightarrow \Omega$ be the universal covering from unit disc Δ to Ω . We call the holomorphic function f on Δ a multiplicative function, if there is a character χ , which is the representation of the fundamental group of Ω , such that $g^*f = \chi(g)f$, where $|\chi| = 1$ and g is an element of the fundamental group of Ω . Denote the set of such kinds of f by $\mathcal{O}^\chi(\Omega)$.

It is known that for any harmonic function u on Ω , there exist a χ_u and a multiplicative function $f_u \in \mathcal{O}^{\chi_u}(\Omega)$, such that $|f_u| = p^*(e^u)$. If $u_1 - u_2 = \log |f|$, then $\chi_{u_1} = \chi_{u_2}$, where u_1 and u_2 are harmonic functions on Ω and f is a holomorphic function on Ω . Recall that for the Green function $G_\Omega(z, z_0)$, there exist a χ_{z_0} and a multiplicative function $f_{z_0} \in \mathcal{O}^{\chi_{z_0}}(\Omega)$ such that $|f_{z_0}(z)| = p^*(e^{G_\Omega(z, z_0)})$ (see [17]).

2020 *Mathematics Subject Classification.* 32D15, 32E10, 32L10, 32U05, 32W05.

Key words and phrases. open Riemann surface, Green function, Suita conjecture.

Let u be a harmonic function on Ω , and let $B_{\Omega, e^{-2u}}(z_0)$ be the Bergman kernel with the weight e^{-2u} . In [19], Yamada posed the following weighted version of Suita conjecture, which is so-called extended Suita conjecture.

Conjecture 1.2. $c_{\beta}^2(z_0) \leq \pi e^{-2u(z_0)} B_{\Omega, e^{-2u}}(z_0)$, and equality holds if and only if $\chi_{z_0} = \chi_{-u}$.

Note that Ω is conformally equivalent to the unit disc less a (possible) closed set of inner capacity zero if and only if $\chi_{z_0} \equiv 1$ (see [17]), thus the extended Suita conjecture is a general version of Suita conjecture. In [12], Guan-Zhou proved the extended Suita conjecture (one can see more general versions in [8, 9]).

Let us consider the weighted Bergman kernel for high derivatives (see [2, 10]). Let k be a nonnegative integer, and let $I = (w)$ be an ideal of \mathcal{O}_{z_0} . Let $h = w^k dw$ be a holomorphic $(1,0)$ form on V_{z_0} . Denote that

$$B_{\Omega, \rho}^{(k)}(z_0) := \frac{2}{\inf \left\{ \int_{\Omega} |\tilde{F}|^2 \rho : \tilde{F} \in H^0(\Omega, \mathcal{O}(K_{\Omega})) \text{ \& } (\tilde{F} - h, z_0) \in I^{k+1} \otimes \mathcal{O}(K_{\Omega})_{z_0} \right\}},$$

where ρ is a nonnegative Lebesgue measurable function on Ω and $|\tilde{F}|^2 := \sqrt{-1} \tilde{F} \wedge \overline{\tilde{F}}$ for any $(1,0)$ form \tilde{F} . It is clear that

$$B_{\Omega, \rho}^{(k)}(z_0) = \sup \left\{ \left| \frac{f^{(k)}(z_0)}{k!} \right|^2 : \exists F \in H^0(\Omega, \mathcal{O}(K_{\Omega})), F|_{V_{z_0}} = f dw \text{ \& } \frac{1}{2} \int_{\Omega} |F|^2 \rho \leq 1 \right\}.$$

When $k = 0$, $B_{\Omega, \rho}^{(k)}(z_0)$ degenerates to the weighted Bergman kernel $B_{\Omega, \rho}(z_0)$.

Then it is natural to ask

Question 1.3. *Can one prove a weighted version of Suita conjecture for higher derivatives related to the kernel function $B_{\Omega, \rho}^{(k)}(z_0)$?*

Guan-Mi-Yuan [10] gave an affirmative answer to this question. Some related results can be referred to [13, 18].

Let $c(t)$ be a positive measurable function on $(0, +\infty)$ satisfying $c(t)e^{-t}$ is decreasing on $(0, +\infty)$ and $\int_0^{+\infty} c(t)e^{-t} dt < +\infty$. Denote that

$$\rho_{z_0} := e^{-(2(k+1-a)G_{\Omega}(\cdot, z_0) + 2v)} c(-2aG_{\Omega}(\cdot, z_0)),$$

where $a > 0$ is a constant and v is a subharmonic function on Ω .

Theorem 1.4 ([10]).

$$(c_{\beta}(z_0))^{2(k+1)} \leq \left(\int_0^{+\infty} c(t)e^{-t} dt \right) \frac{\pi}{a} e^{-2v(z_0)} B_{\Omega, \rho_{z_0}}^{(k)}(z_0)$$

holds for any $z_0 \in \Omega$ satisfying $v(z_0) > -\infty$. Moreover, for $z_0 \in \Omega$ with $v(z_0) > -\infty$, equality

$$(1.1) \quad (c_{\beta}(z_0))^{2(k+1)} = \left(\int_0^{+\infty} c(t)e^{-t} dt \right) \frac{\pi}{a} e^{-2v(z_0)} B_{\Omega, \rho_{z_0}}^{(k)}(z_0)$$

holds if and only if the following statements hold:

(1) $v = \log |g| + u$, where g is a holomorphic function on Ω such that $g(z_0) \neq 0$ and u is a harmonic function on Ω ;

(2) $\chi_{-u} = (\chi_{z_0})^{k+1}$, where χ_{-u} and χ_{z_0} are the characters associated to the functions $-u$ and $G_{\Omega}(\cdot, z_0)$ respectively.

For Suita conjecture, if $\pi B_\Omega(z_0) = (c_\beta(z_0))^2$ holds for a point $z_0 \in \Omega$, then it follows from the solution of Suita conjecture that $\pi B_\Omega(z_0) = (c_\beta(z_0))^2$ holds for any $z_0 \in \Omega$, i.e., the set of points z_0 such that $\pi B_\Omega(z_0) = (c_\beta(z_0))^2$ holds is either \emptyset or Ω .

In this article, we discuss the set of points for the holding of the equality in the weighted version of Suita conjecture for higher derivatives.

1.1. Main result.

Let Ω be an open Riemann surface, which admits a nontrivial Green function G_Ω . Let k be a nonnegative integer. Let $c(t)$ be a positive measurable function on $(0, +\infty)$ satisfying $c(t)e^{-t}$ is decreasing on $(0, +\infty)$ and $\int_0^{+\infty} c(t)e^{-t} dt < +\infty$. Denote that

$$\rho_{z_0} := e^{-(2(k+1-a)G_\Omega(\cdot, z_0)+2v)} c(-2aG_\Omega(\cdot, z_0)),$$

where $a > 0$ is a constant and v is a subharmonic function on Ω .

It is clear that, if there is $\tilde{g} \in \mathcal{O}(\Omega)$ such that $v = \log |\tilde{g}|$ and Ω is conformally equivalent to the unit disc less a (possible) closed set of inner capacity zero, then $\chi_{z_0} = 1$, which implies that equality (1.1) holds for any $z_0 \in \Omega$ with $v(z_0) > -\infty$ by Theorem 1.4. The following theorem shows that the converse to the result is also true.

Theorem 1.5. Equality

$$(1.2) \quad (c_\beta(z_0))^{2(k+1)} = \left(\int_0^{+\infty} c(t)e^{-t} dt \right) \frac{\pi}{a} e^{-2v(z_0)} B_{\Omega, \rho_{z_0}}^{(k)}(z_0)$$

holds for any $z_0 \in \Omega$ satisfying $v(z_0) > -\infty$ if and only if the following two statements hold:

- (1) Ω is conformally equivalent to the unit disc less a (possible) closed set of inner capacity zero;
- (2) $v = \log |g|$, where g is a holomorphic function on Ω which is not identically equal to 0.

The following theorem gives a relation between the set of points such that equality (1.2) holds and the integer valued points of a class of harmonic functions (maybe multi-valued), which shows that the set of points where equality (1.2) holds is a real analytic subset of Ω when one of the two statements in Theorem 1.5 doesn't hold. Let $\Pi_1 \subset \pi_1(\Omega)$ such that $\pi_1(\Omega)$ is generated by Π_1 .

Theorem 1.6. Assume that $v = \log |\tilde{g}| + u$, where $\tilde{g} \not\equiv 0$ is a holomorphic function on Ω and u is a harmonic function on Ω . There exists a class of harmonic functions $\{u_\alpha\}_{\alpha \in \Pi_1}$ on Δ , such that $\beta^* u_\alpha - u_\alpha$ is an integer constant function for any $\beta \in \pi_1(\Omega)$ and any $\alpha \in \Pi_1$. Equality

$$(1.3) \quad (c_\beta(z_0))^{2(k+1)} = \left(\int_0^{+\infty} c(t)e^{-t} dt \right) \frac{\pi}{a} e^{-2v(z_0)} B_{\Omega, \rho_{z_0}}^{(k)}(z_0)$$

holds for $z_0 \in \Omega$ with $v(z_0) > -\infty$ if and only if $u_\alpha(\tilde{z}_0) \in \mathbb{Z}$ for any $\alpha \in \Pi_1$ and $\tilde{z}_0 \in p^{-1}(z_0)$.

Moreover, $\{u_\alpha\}_{\alpha \in \Pi_1}$ are integer constant functions if and only if Ω is conformally equivalent to the unit disc less a (possible) closed set of inner capacity zero and $v = \log |g|$, where $g \not\equiv 0$ is a holomorphic function on Ω .

Remark 1.7. In fact, $u_\alpha(z)$ is locally equal to $\frac{1}{2\pi} \left((k+1) \int_{\gamma_\alpha} \tilde{d}G_\Omega(\cdot, p(z)) + \int_{\gamma_\alpha} \tilde{d}u \right) + m$ (see equality (3.6)), where γ_α is a piecewise smooth closed curve on Ω satisfying $[\gamma_\alpha] = \alpha$, $\tilde{d} = \frac{\partial - \bar{\partial}}{i}$ and $m \in \mathbb{Z}$. Remark 5.6 in Appendix shows that p_*u_α is a single-valued function when Ω is a planar domain, but there exists an open Riemann surface such that p_*u_α may not be a single-valued function.

When $\rho_{z_0} \equiv 1$ (i.e., $v \equiv 0$, $a = k+1$ and $c \equiv 1$), denote $B_{\Omega, \rho_{z_0}}^{(k)}(z_0)$ by $B_\Omega^{(k)}(z_0)$, which is the Bergman kernel for higher derivatives with trivial weight (see [2]). Theorem 1.6 implies the following corollary.

Corollary 1.8. *There exists a class of harmonic functions $\{u_\alpha\}_{\alpha \in \Pi_1}$ on Δ , such that $\beta^*u_\alpha - u_\alpha$ is an integer constant function for any $\beta \in \pi_1(\Omega)$ and $\alpha \in \Pi_1$. For any $z_0 \in \Omega$, equality*

$$(c_\beta(z_0))^{2(k+1)} = \frac{\pi}{k+1} B_\Omega^{(k)}(z_0)$$

holds if and only if $u_\alpha(z_0) \in \mathbb{Z}$ for any $\alpha \in \Pi_1$.

Moreover, $\{u_\alpha\}$ are integer constant functions if and only if Ω is conformally equivalent to the unit disc less a (possible) closed set of inner capacity zero.

In the following, we assume that Ω is a bounded connected open subset of \mathbb{C} , which is bounded by n analytic closed curves.

Denote that $\mathbb{C} \setminus \Omega = \bigcup_{i=1}^n A_j$, where A_j are disjoint closed sets. Without loss of generality, suppose that A_n is the unbounded connected component of $\mathbb{C} \setminus \Omega$. Denote the boundary of A_j by Γ_j , $1 \leq j \leq n$. For any $1 \leq j \leq n-1$, let γ_j be a closed smooth curve in Ω , which winds around points in A_j once and does not wind around points in A_l for $l \neq j$ (for definition, see Lemma 2.5 and Remark 2.6).

For $1 \leq j \leq n-1$, by solving Dirichlet problem (see Remark 2.15), there exists $u_j \in C(\bar{\Omega})$, which satisfies that $u_j = 1$ on Γ_j , $u_j(z) = 0$ on Γ_l for any $l \neq j$ and u_j is harmonic on Ω .

The following theorem gives a relation between Dirichlet problem and the set of points such that equality (1.3) holds.

Theorem 1.9. *Equality*

$$(c_\beta(z_0))^{2(k+1)} = \left(\int_0^{+\infty} c(t) e^{-t} dt \right) \frac{\pi}{a} e^{-2v(z_0)} B_{\Omega, \rho_{z_0}}^{(k)}(z_0)$$

holds for $z_0 \in \Omega$ with $v(z_0) > -\infty$ if and only if the following two statements hold:

- (1) $v = \log |g| + u$, where g is a holomorphic function on Ω satisfying $g(z_0) \neq 0$ and u is a harmonic function on Ω ;
- (2) $(k+1)u_j(z_0) + \frac{1}{2\pi} \int_{\gamma_j} \frac{\partial u(z)}{\partial n} ds(z) \in \mathbb{Z}$ holds for any $1 \leq j \leq n-1$, where $\frac{\partial}{\partial n}$ means differentiation in the direction of the outward pointing normal, and s is the arc-length parameter.

Remark 1.10. We call a domain $\Omega \subset \mathbb{C}$ is n -connected, if $\mathbb{C} \setminus \Omega = \bigcup_{j=1}^n A_j$, where $\{A_j\}$ are disjoint connected closed sets. By Lemma 2.4, we know that Theorem 1.9 also holds when Ω is a bounded n -connected open subset of \mathbb{C} such that no connected component of $\mathbb{C} \setminus \Omega$ is an isolated point.

Remark 1.11. We give an example of Theorem 1.9. Suppose that $\Omega = \{z \in \mathbb{C} : 1 < |z| < R\}$, and v is a harmonic function on Ω . Denote that

$$c := \int_{\{|z|=\frac{R+1}{2}\}} \frac{\partial v(z)}{\partial n} ds(z)$$

is a constant. Let $u_1 = \frac{\log R - \log |z|}{\log R}$. It is clear that u_1 is harmonic on $\mathbb{C} \setminus \{0\}$, $u_1(z) = 1$ when $|z| = 1$ and $u_1(z) = 0$ when $|z| = R$. Theorem 1.9 shows that the set of points $z_0 \in \Omega$ satisfying

$$(c_\beta(z_0))^{2(k+1)} = \left(\int_0^{+\infty} c(t)e^{-t} dt \right) \frac{\pi}{a} e^{-2v(z_0)} B_{\Omega, \rho_{z_0}}^{(k)}(z_0)$$

is just

$$\left\{ z_0 \in \Omega : |z_0| = R^{\frac{c+m}{k+1}} \text{ for some } m \in \mathbb{Z} \right\}.$$

Suppose that Ω is a doubly-connected open subset of \mathbb{C} and no connected component of $\mathbb{C} \setminus \Omega$ is an isolated point. Let v be a harmonic function on Ω . Note that Ω can be mapped conformally onto a circular ring (see [15]), then the set of points $z_0 \in \Omega$ satisfying

$$(c_\beta(z_0))^{2(k+1)} = \left(\int_0^{+\infty} c(t)e^{-t} dt \right) \frac{\pi}{a} e^{-2v(z_0)} B_{\Omega, \rho_{z_0}}^{(k)}(z_0)$$

is constituted by k real analytic closed curves.

When $\rho_{z_0} \equiv 1$, Theorem 1.9 implies the following corollary.

Corollary 1.12. *Equality*

$$(c_\beta(z_0))^{2(k+1)} = \frac{\pi}{k+1} B_\Omega^{(k)}(z_0)$$

holds for $z_0 \in \Omega$ if and only if $(k+1)u_j(z_0) \in \mathbb{Z}$ for any $1 \leq j \leq n-1$.

Under the assumption that Ω is a connected open subset of \mathbb{C} bounded by n analytic closed curves, the equality part of Suita conjecture is equivalent to that “ $\pi B_\Omega(z_0) = (c_\beta(z_0))^2$ can not hold for any $z_0 \in \Omega$ unless $n = 1$ ”. We generalize it to any nonnegative integer k in the following corollary.

Corollary 1.13. *Assume that $v \equiv 0$ on Ω , then*

$$(c_\beta(z_0))^{2(k+1)} = \left(\int_0^{+\infty} c(t)e^{-t} dt \right) \frac{\pi}{a} B_{\Omega, \rho_{z_0}}^{(k)}(z_0)$$

can not hold for any $z_0 \in \Omega$ unless $k \geq n-1$.

Remark 1.14. *Suppose that Ω is bounded by 3 analytic closed curves Γ_j ($j = 1, 2, 3$), and $v \equiv 0$ on Ω . Using Theorem 1.9 and some discussions on harmonic functions, we get that there exists a sufficiently large k and $z_0 \in \Omega$ such that*

$$(c_\beta(z_0))^{2(k+1)} = \left(\int_0^{+\infty} c(t)e^{-t} dt \right) \frac{\pi}{a} B_{\Omega, \rho_{z_0}}^{(k)}(z_0).$$

We prove the present remark in Section 4.3.

2. PREPARATION

In this section, we will do some preparations.

2.1. Characters on open Riemann surfaces.

Let Ω be an open Riemann surface, which admits a nontrivial Green function G_Ω . Some basic knowledge about open Riemann surfaces used in this section can be referred to [7, 15]. Let $p : \Delta \rightarrow \Omega$ be the universal covering from unit disc Δ to Ω . Denote $Deck(\Delta \rightarrow \Omega)$ to be the set of all biholomorphic isomorphisms $\sigma : \Delta \rightarrow \Delta$ which preserve fibers, i.e., $p \circ \sigma = p$. We firstly recall the following lemma.

Lemma 2.1 (see [7]). *Let z_1 be a point in Ω . $Deck(\Delta \rightarrow \Omega)$ is isomorphic to the fundamental group $\pi_1(\Omega, z_1)$.*

Proof. Let $y_1 \in \Delta$ be a point such that $p(y_1) = z_1$. For any $\sigma \in Deck(\Delta \rightarrow \Omega)$, we can construct a curve v in Δ with initial point y_1 and end point $\sigma(y_1)$. Since Δ is simply connected, the homotopy class $[v]$ of v is uniquely determined. Then $p \circ v$ is a closed curve in Ω with endpoint z_1 , and its homotopy class $[p \circ v]$ is uniquely determined, hence $[p \circ v]$ is a definite element in $\pi_1(\Omega, z_1)$. Now we can define a mapping

$$\Phi : Deck(\Delta \rightarrow \Omega) \rightarrow \pi_1(\Omega, z_1)$$

such that

$$\Phi(\sigma) = [p \circ v] \in \pi_1(\Omega, z_1)$$

for $\sigma \in Deck(\Delta \rightarrow \Omega)$.

To prove that Φ is a group homomorphism, suppose that $\sigma, \tau \in Deck(\Delta \rightarrow \Omega)$, v is a curve in Δ with initial point y_1 and end point $\sigma(y_1)$, and η is a curve in Δ with initial point y_1 and end point $\tau(y_1)$. Then $\sigma \circ \eta$ is a curve in Δ with initial point $\sigma(y_1)$ and end point $\sigma\tau(y_1)$. Denote that $v \cdot (\sigma \circ \eta)$ is a curve on Δ , which satisfies that $v \cdot (\sigma \circ \eta)(t) = v(2t)$ on $[0, \frac{1}{2}]$ and $v \cdot (\sigma \circ \eta)(t) = \sigma \circ \eta(2t - 1)$ on $[\frac{1}{2}, 1]$. Note that $v \cdot (\sigma \circ \eta)$ has initial point y_1 and end point $\sigma\tau(y_1)$. Thus

$$\begin{aligned} \Phi(\sigma\tau) &= [p \circ (v \cdot (\sigma \circ \eta))] = [p \circ v][p \circ (\sigma \circ \eta)] \\ &= [p \circ v][p \circ \eta] = \Phi(\sigma)\Phi(\tau). \end{aligned}$$

To prove the injectivity of Φ , suppose that $\Phi(\sigma)$ is trivial in $\pi_1(\Omega, z_1)$, then $p \circ v$ is null-homotopic. Since v is the lifting of $p \circ v$, v is also null-homotopic, hence $\sigma(y_1) = y_1$. σ is therefore the identity mapping.

To prove the surjectivity of Φ , suppose that α is an element in $\pi_1(\Omega, z_1)$ and u is a curve representing of α . Let v be the lifting of u to Δ with initial point y_1 , and denote the end point of v by y_2 . Then by the property of universal covering, there exists $\sigma \in Deck(\Delta \rightarrow \Omega)$ such that $\sigma(y_1) = y_2$. From the definition of Φ , we have $\Phi(\sigma) = \alpha$. \square

Let h and h_1 be harmonic functions on an open subset V of Ω . We call h_1 is a harmonic conjugate of h on V , if $h + ih_1$ is a holomorphic function on V . Assume that V is simply-connected, then for any harmonic function h on V , there is a harmonic conjugate \tilde{h} of h . Denote that

$$\tilde{d} = \frac{\partial - \bar{\partial}}{i}.$$

Lemma 2.2. *Let $z_1, z_2 \in V$, and let γ be a piecewise smooth curve in V from z_1 to z_2 . Then we have*

$$\tilde{h}(z_2) - \tilde{h}(z_1) = \int_\gamma \tilde{d}h.$$

Proof. Without loss of generality, assume that V is a simply-connected open subset of \mathbb{C} and γ is smooth (we can divide γ into finite smooth curves which are contained in simply-connected coordinate open subsets of Ω).

Note that

$$\begin{aligned}\tilde{d}v &= \frac{1}{i}(\partial v - \bar{\partial}v) \\ &= \frac{1}{i} \left(\frac{\partial v}{\partial z} dz - \frac{\partial v}{\partial \bar{z}} d\bar{z} \right) \\ &= \frac{1}{2i} \left(\left(\frac{\partial v}{\partial x} + \frac{1}{i} \frac{\partial v}{\partial y} \right) (dx + idy) - \left(\frac{\partial v}{\partial x} - \frac{1}{i} \frac{\partial v}{\partial y} \right) (dx - idy) \right) \\ &= \frac{\partial v}{\partial x} dy - \frac{\partial v}{\partial y} dx\end{aligned}$$

holds for smooth function v , thus we know that

$$\tilde{h}(z) = \int_{\gamma_z} \tilde{d}h + C$$

for any $z \in V$, where γ_z is any smooth curve in V from z_1 to z . As h is harmonic and V is simply-connected, we have

$$\tilde{h}(z_2) - \tilde{h}(z_1) = \int_{\gamma_{z_2}} \tilde{d}h = \int_{\gamma} \tilde{d}h.$$

□

Let $z_0 \in \Omega$, and let u be a harmonic function on Ω . There exist $f_{z_0}, f_u \in \mathcal{O}(\Delta)$ such that $|f_{z_0}| = p^*e^{G_{\Omega}(\cdot, z_0)}$ and $|f_u| = p^*e^u$. Let $z_1 \in \Omega$ and $z_1 \neq z_0$. By the isomorphism $\Phi : Deck(\Delta \rightarrow \Omega) \rightarrow \pi_1(\Omega, z_1)$ constructed in Lemma 2.1, $\Phi^{-1}(g)$ is a biholomorphic isomorphism on Δ which preserves fibers for any $g \in \pi_1(\Omega, z_1)$. Then we have

$$\Phi^{-1}(g)^* f_{z_0} = \chi_{z_0}(g) f_{z_0} \quad \text{and} \quad \Phi^{-1}(g)^* f_u = \chi_u(g) f_u.$$

Since $p : \Delta \rightarrow \Omega$ is the universal covering, any element in $\pi_1(\Omega, z_1)$ can be represented by $p \circ \tilde{\gamma}$ where $\tilde{\gamma}$ is a curve in Δ , and it only depends on the starting point and end point of $\tilde{\gamma}$ by Lemma 2.1. Thus, any $g \in \pi_1(\Omega, z_1)$ can be represented by a piecewise smooth curve γ in Ω satisfying $z_0 \notin \gamma$.

In the following proposition, we represent $\chi_{z_0}(g)$ and $\chi_u(g)$ by the integrals of $\tilde{d}G_{\Omega}(\cdot, z_0)$ and $\tilde{d}u$ along γ respectively.

Proposition 2.3. *Let g be an element in $\pi_1(\Omega, z_1)$ which can be represented by γ , where γ is a piecewise smooth curve in Ω and $z_0 \notin \gamma$. Then we have*

$$\chi_{z_0}(g) = e^{i \int_{\gamma} \tilde{d}G_{\Omega}(\cdot, z_0)}$$

and

$$\chi_u(g) = e^{i \int_{\gamma} \tilde{d}u}.$$

Proof. Lift γ to a curve $\tilde{\gamma}$ in Δ with initial point y_1 . Denote the end point of $\tilde{\gamma}$ by y . By the isomorphism $\Phi : Deck(\Delta \rightarrow \Omega) \rightarrow \pi_1(\Omega, z_1)$ constructed in Lemma 2.1, $\Phi^{-1}(g)$ is a biholomorphic isomorphism $\sigma : \Delta \rightarrow \Delta$ which preserves fibers and $\sigma(y_1) = y$. For any holomorphic function f in Δ , σ^*f is a holomorphic function in Δ and $\sigma^*f(y_1) = f(y)$. As $\sigma^*f_{z_0} = \chi_{z_0}(g)f_{z_0}$, we have

$$(2.1) \quad f_{z_0}(y) = \sigma^*f_{z_0}(y_1) = \chi_{z_0}(g)f_{z_0}(y_1).$$

Let $\{V_j\}_{j=1}^N$ be a simply-connected open covering of $\tilde{\gamma}$ satisfying that p maps V_j biholomorphically onto $p(V_j)$. We can divide $\tilde{\gamma}$ into γ_j such that $\gamma_j \subset V_j$. Denote the initial point of γ_j by a_j , the end point of γ_j by b_j .

By Lemma 2.2, for any j , we have

$$f_{z_0} = p^* e^{(G_\Omega(\cdot, z_0) + i\widetilde{G}_\Omega(\cdot, z_0))}$$

on V_j and

$$\widetilde{G}_\Omega(p(b_j), z_0) - \widetilde{G}_\Omega(p(a_j), z_0) = \int_{p^*\gamma_j} \widetilde{d}G_\Omega(\cdot, z_0),$$

where $\widetilde{G}_\Omega(z, z_0)$ is the harmonic conjugate of $G_\Omega(z, z_0)$ on $p(V_j)$. Therefore

$$\begin{aligned} \frac{f_{z_0}(b_j)}{f_{z_0}(a_j)} &= e^{G_\Omega(p(b_j), z_0) + i\widetilde{G}_\Omega(p(b_j), z_0) - G_\Omega(p(a_j), z_0) - i\widetilde{G}_\Omega(p(a_j), z_0)} \\ &= e^{G_\Omega(p(b_j), z_0) - G_\Omega(p(a_j), z_0) + i \int_{p^*\gamma_j} \widetilde{d}G_\Omega(z, z_0)}. \end{aligned}$$

As $p(a_1) = p(b_N)$, we have

$$\begin{aligned} \frac{f_{z_0}(b_N)}{f_{z_0}(a_1)} &= \prod_{j=1}^N \frac{f_{z_0}(b_j)}{f_{z_0}(a_j)} \\ &= e^{\sum_{j=1}^N (G_\Omega(p(b_j), z_0) - G_\Omega(p(a_j), z_0) + i \int_{p^*\gamma_j} \widetilde{d}G_\Omega(\cdot, z_0))} \\ &= e^{G_\Omega(p(b_N), z_0) - G_\Omega(p(a_1), z_0) + i \int_\gamma \widetilde{d}G_\Omega(\cdot, z_0)} \\ &= e^{i \int_\gamma \widetilde{d}G_\Omega(\cdot, z_0)}. \end{aligned}$$

Note that $a_1 = y_1$, $b_N = y$ and $p(a_1) = p(b_N) = z_1$, thus

$$f_{z_0}(y) = f_{z_0}(y_1) e^{i \int_\gamma \widetilde{d}G_\Omega(\cdot, z_0)}.$$

Combining this with equality (2.1), we therefore have

$$\chi_{z_0}(g) = e^{i \int_\gamma \widetilde{d}G_\Omega(\cdot, z_0)}.$$

Similarly, we have

$$\chi_u(g) = e^{i \int_\gamma \widetilde{d}u}.$$

□

2.2. Some lemmas on planar domains.

In this section, we consider the case that the open Riemann surface Ω is a bounded connected open subset of \mathbb{C} such that $\mathbb{C} \setminus \Omega = \bigcup_{j=1}^n A_j$, where $\{A_j\}$ are disjoint connected closed sets and each A_j is not an isolated point. We recall the following well-known lemma.

Lemma 2.4. *Assume that Ω is a bounded n -connected open subset of \mathbb{C} such that no connected component of $\mathbb{C} \setminus \Omega$ is an isolated point, then Ω is conformally equivalent to an open subset of \mathbb{C} bounded by n closed analytic curves.*

Proof. For convenience of readers, we give a proof of this lemma by using Riemann mapping theorem.

Assume that $\mathbb{C} \setminus \Omega = \bigcup_{j=1}^n A_j$, where A_j are disjoint connected closed sets. Denote the boundary of A_j by C_j . Without loss of generality, assume that A_n is unbounded.

Let $A = \Omega \cup (\bigcup_{j=2}^n A_j) \cup \{\infty\} = \mathbb{C}\mathbb{P}^1 \setminus A_1$, which is a simply-connected proper open subset of $\mathbb{C}\mathbb{P}^1$ whose boundary C_1 is not an isolated point. Then by Riemann mapping theorem, A can be mapped conformally onto the unit disc Δ . This mapping can transform Ω onto $\Omega_1 = \Delta \setminus (\bigcup_{j=2}^n A'_j)$, where A'_j is the conformal image of A_j and $C'_j = \partial A'_j$ is the conformal image of C_j for $2 \leq j \leq n$. In further let $A'_1 = \mathbb{C} \setminus \Delta$, $C'_1 = \partial A'_1 = \{z \in \mathbb{C} : |z| = 1\}$.

Repeating the above procedure for another $n - 1$ times, we can transform Ω onto $\Omega_n = \Delta \setminus (\bigcup_{j=1}^{n-1} A_j^{(n)})$, where $C_j^{(n)} = \partial A_j^{(n)}$ are closed analytic curves for $1 \leq j \leq n - 1$. It is obvious that Ω_n is a bounded open subset of \mathbb{C} which is bounded by n closed analytic curves. \square

We recall the following basic knowledge in algebraic topology.

Lemma 2.5 (see [6]). *Let U be a bounded connected open subset of \mathbb{C} such that $\mathbb{C} \setminus U = \bigcup_{j=1}^m A_j$, where A_j are disjoint connected closed sets and A_m is unbounded. For each $1 \leq j \leq m - 1$, there exists a closed smooth curve γ_j in Ω which winds around points in A_j once and does not wind around points in A_l for $l \neq j$. If γ'_j is another closed smooth curve which satisfies these conditions, then γ'_j is homologous to γ_j . The homology classes of these $m - 1$ curves form a basis for the first homology group $H_1(U)$. That is, these curves are not homologous to each other and any curve in U is homologous to some linear combinations of these curves.*

Remark 2.6. *The times that a closed smooth curve γ winds around a point $P = (x_0, y_0) \in \mathbb{R}^2$ is defined as follows. In $\mathbb{R}^2 \setminus (x_0, y_0)$ we can define a differential $\omega_P = \frac{-(y-y_0)dx + (x-x_0)dy}{(x-x_0)^2 + (y-y_0)^2}$, and define $W(\gamma, P) = \frac{1}{2\pi} \int_{\gamma} \omega_P$ to be the times the curve γ winds around the point P . For example, if γ is the curve $\gamma : [0, 1] \rightarrow \mathbb{R}^2 \setminus (x_0, y_0)$, $\gamma(t) = (x_0 + r_0 \cos 2\pi t, y_0 + 2 \sin \pi t)$, then $W(\gamma, P) = 1$. For arbitrary closed curve, $W(\gamma, P)$ is always an integer. For details, see [6].*

Lemma 2.7. *$\frac{\partial v}{\partial n} ds = \tilde{d}v$ holds for smooth function v , where $\frac{\partial}{\partial n}$ means differentiation in the direction of the outward pointing normal, s is the arc-length parameter and $\tilde{d} = \frac{\partial - \bar{\partial}}{\sqrt{-1}}$.*

Proof. Note that

$$\tilde{d}v = \frac{\partial v}{\partial x} dy - \frac{\partial v}{\partial y} dx$$

and

$$\frac{\partial v}{\partial n} = v_x \cos(x, n) + v_y \cos(y, n),$$

where (x, n) and (y, n) denote the angle between the outward pointing normal and the positive x - and y -axis, respectively. Since the cosines of the angles between the tangent direction and the positive x - and y -axis are $\frac{dx}{ds}$ and $\frac{dy}{ds}$, where s is the arc-length parameter, and the outer normal is obtained by turning the tangent clockwise by $\frac{\pi}{2}$, it follows that

$$\cos(x, n) = \frac{dy}{ds}, \quad \cos(y, n) = -\frac{dx}{ds}.$$

Hence

$$\frac{\partial v}{\partial n} ds = v_x dy - v_y dx = \tilde{d}v.$$

\square

Because of Lemma 2.4, in the following part, we assume that Ω is a bounded connected open subset of \mathbb{C} , which is bounded by n analytic closed curves.

Let u be a harmonic function on Ω , and let $z_0 \in \Omega$. It follows from Lemma 2.5 that, for any $1 \leq l \leq n-1$, there exists a closed smooth curve γ_l in Ω which winds around points in A_l once, does not wind around points in A_j for $j \neq l$ and satisfies $z_0 \notin \gamma_l$.

Lemma 2.8. $(\chi_{z_0})^{k+1} = \chi_{-u}$ holds if and only if

$$\left(e^{i \int_{\gamma_j} \frac{\partial G_{\Omega}(z, z_0)}{\partial n} ds(z)} \right)^{k+1} = e^{-i \int_{\gamma_j} \frac{\partial u(z)}{\partial n} ds(z)}$$

holds for $1 \leq j \leq n-1$.

Proof. It follows from Proposition 2.3 that $(\chi_{z_0})^{k+1} = \chi_{-u}$ holds if and only if

$$\left(e^{i \int_{\gamma} \frac{\partial G_{\Omega}(z, z_0)}{\partial n} ds(z)} \right)^{k+1} = e^{-i \int_{\gamma} \frac{\partial u(z)}{\partial n} ds(z)}$$

holds for smooth closed curve γ in $\Omega \setminus \{z_0\}$.

Let γ_0 be a smooth closed curve in $\Omega \setminus \{z_0\}$ which winds around z_0 once and does not wind around points in $\mathbb{C} \setminus \Omega$. By Lemma 2.5, the first homology group $H_1(\Omega \setminus \{z_0\}) \cong \mathbb{Z}^{\oplus n}$, and the homology classes of γ_j for $0 \leq j \leq n-1$ form its basis. Hence for any smooth curve $\gamma \subset \Omega \setminus \{z_0\}$, there exists a sequence of integers t_j for $0 \leq j \leq n-1$ such that $\sum_{j=0}^{n-1} t_j \gamma_j$ is homologous to γ .

Since $G_{\Omega}(z, z_0)$ and u are harmonic on $\Omega \setminus \{z_0\}$, $\frac{\partial G_{\Omega}(z, z_0)}{\partial n} ds$ and $\frac{\partial u(z)}{\partial n} ds$ are closed differential 1-forms on $\Omega \setminus \{z_0\}$ by Lemma 2.7. Then we have

$$\int_{\gamma} \frac{\partial G_{\Omega}(z, z_0)}{\partial n} ds = \sum_{j=0}^{n-1} t_j \int_{\gamma_j} \frac{\partial G_{\Omega}(z, z_0)}{\partial n} ds$$

and

$$\int_{\gamma} \frac{\partial u(z)}{\partial n} ds = \sum_{j=0}^{n-1} t_j \int_{\gamma_j} \frac{\partial u(z)}{\partial n} ds.$$

Recall that $G_{\Omega}(z, z_0) = \log |z - z_0| + u'(z)$ on Ω , where u' is a harmonic function on Ω , hence

$$\int_{\gamma_0} \frac{\partial G_{\Omega}(z, z_0)}{\partial n} ds = \int_{\gamma_0} \frac{\partial \log |z - z_0|}{\partial n} ds + \int_{\gamma_0} \frac{\partial u'(z)}{\partial n} ds = 2\pi$$

and

$$\int_{\gamma_0} \frac{\partial u(z)}{\partial n} ds = 0.$$

Therefore we get that

$$\left(e^{i \int_{\gamma} \frac{\partial G_{\Omega}(z, z_0)}{\partial n} ds(z)} \right)^{k+1} = e^{-i \int_{\gamma} \frac{\partial u(z)}{\partial n} ds(z)}$$

holds for smooth closed curve γ in $\Omega \setminus \{z_0\}$ if and only if

$$\left(e^{i \int_{\gamma_j} \frac{\partial G_{\Omega}(z, z_0)}{\partial n} ds} \right)^{k+1} = e^{-i \int_{\gamma_j} \frac{\partial u(z)}{\partial n} ds}$$

holds for $1 \leq j \leq n-1$. Thus, Lemma 2.8 has been proved. \square

2.3. Harmonic functions and Dirichlet problem.

Firstly, we recall two properties of harmonic functions on an open subset of \mathbb{R}^2 .

Lemma 2.9 (Harnack's inequality, see [5]). *Let V, U be two connected open subsets of \mathbb{R}^2 such that $V \Subset U$, then there exists a positive constant C , which depends only on V and U , such that*

$$\sup_V u \leq C \inf_V u$$

for all nonnegative harmonic functions u on U .

Lemma 2.10 (see [5]). *Assume u is harmonic on $U \subset \mathbb{R}^n$. Then for any positive integer k , there exists a constant C_k which is independent of u and U , such that*

$$|D^\alpha u(x_0)| \leq \frac{C_k}{r^{2+k}} |u|_{L^1(B(x_0, r))}$$

for each ball $B(x_0, r) \subset U$ and each multi-index α of order $|\alpha| = k$.

The following lemma will be used in the proof of Remark 1.14.

Lemma 2.11. *Let u_j be the harmonic function on a unit disc $\Delta \subset \mathbb{C}$, where $j = 1, 2$. Assume that u_j is not a constant function for $j = 1, 2$, and $u_2 \equiv c_1 u_1 + c_2$ does not hold for any constant c_1, c_2 . Then there exists $z_0 \in \Delta$ such that $u_1(z_0) \in \mathbb{Q}$ and $u_2(z_0) \in \mathbb{Q}$.*

Proof. As Δ is simply connected, there exists a harmonic function v_1 on Δ such that $f := u_1 + iv_1$ is holomorphic on Δ . As u_1 is not a constant function, shrinking Δ if necessary, we can assume that f is a coordinate function on Δ without loss of generality. Denote that $h = u_2 \circ f^{-1}$ on $f(\Delta)$, then h is a harmonic function on $f(\Delta)$.

Now, we prove Lemma 2.11 by contradiction: if not, $u_2(z) \notin \mathbb{Q}$ for any $z \in \Delta$ with $u_1(z) \in \mathbb{Q}$. As u_2 is continuous, we know $u_2 \equiv \text{const}$ on $\{z \in \Delta : u_1(z) = c\}$, i.e.,

$$\frac{\partial u_2}{\partial v_1} = 0$$

on $\{z \in \Delta : u_1(z) = c\}$, where $c \in \mathbb{Q}$. Note that $\cup_{c \in \mathbb{Q}} \{z \in \Delta : u_1(z) = c\}$ is dense in Δ , then we have $\frac{\partial u_2}{\partial v_1} = 0$ on Δ , which implies that $h(z)$ is only dependent on $\text{Re} z$. As h is a harmonic function on $f(\Delta)$, there exist constants c_1, c_2 such that

$$u_2 = h(u_1 + iv_1) = c_1 u_1 + c_2,$$

which contradicts to the assumption in Lemma 2.11.

Thus, Lemma 2.11 has been proved. □

We recall some results on Dirichlet problem, which will be used in the proofs of our main results.

Dirichlet problem: *Let Ω be an open subset of \mathbb{R}^n , and let $U(z)$ be a continuous function on $\partial\Omega$. Does there exist a continuous function u on $\overline{\Omega}$, which is harmonic on Ω and satisfies $u|_{\partial\Omega} = U$?*

We recall the following definition.

Definition 2.12. *Let $\Omega \subset \mathbb{R}^n$ be a bounded domain. A boundary point $z \in \partial\Omega$ is called regular, if there is a continuous function ϕ on $\overline{\Omega}$ such that ϕ is subharmonic on Ω , $\phi(z) = 0$ and $\phi < 0$ on $\partial\Omega \setminus \{z\}$.*

The following lemma gives a sufficient condition for solvability of Dirichlet problem.

Lemma 2.13 (Perron 1923, see [7]). *Let $\Omega \subset \mathbb{R}^n$ be a bounded domain. Let U be a function on $\partial\Omega$. Then there exists a harmonic function u on Ω , such that if $z \in \partial\Omega$ is a regular point and U is continuous at z , $u(x) \rightarrow U(z)$ as $\Omega \ni x \rightarrow z$.*

In the following, assume that $\Omega \subset \mathbb{R}^2$. The following lemma gives a useful criterion for $p \in \partial\Omega$ to be regular.

Lemma 2.14 (Osgood 1900, see [1]). *Let $\Omega \subset \mathbb{R}^2$ be a bounded domain. Suppose that $p \in \partial\Omega$ is contained in a component of $\mathbb{R}^2 \setminus \Omega$ which has more than one point, that is, p is not an isolated point in $\mathbb{R}^2 \setminus \Omega$, then p is regular.*

Remark 2.15. *Let Ω be a bounded connected open subset of \mathbb{C} such that $\mathbb{C} \setminus \Omega = \bigcup_{j=1}^n A_j$, where A_j are disjoint connected closed sets and each A_j is not an isolated point. Then by Lemma 2.13 and Lemma 2.14, Dirichlet problem on such Riemann surface is always solvable.*

Let us recall a basic formula, using that we can obtain the solution of Dirichlet problem.

Lemma 2.16 (Green third formula, see [15]). *Assume that Ω is bounded by closed smooth curve Γ . Let $u \in C(\overline{\Omega})$, which is harmonic on Ω . Then we have*

$$u(z_0) = \frac{1}{2\pi} \int_{\Gamma} u \frac{\partial G_{\Omega}(z, z_0)}{\partial n} ds.$$

3. PROOFS OF THEOREM 1.5 AND THEOREM 1.6

In this section, we prove Theorem 1.5 and Theorem 1.6.

3.1. Proof of Theorem 1.5.

Suppose that Ω is conformally equivalent to unit disc less a (possible) closed set of inner capacity zero. For every $z_0 \in \Omega$, $\chi_{z_0} = 1$, hence $\chi_{z_0}^{k+1} = 1$ for every integer $k \geq 0$. If $v = \log |g|$, where $g \neq 0$ is a holomorphic function on Ω , then by Theorem 1.4,

$$(c_{\beta}(z_0))^{2(k+1)} = \left(\int_0^{+\infty} c(t) e^{-t} dt \right) \frac{\pi}{a} e^{-2v(z_0)} B_{\Omega, \rho_{z_0}}^{(k)}(z_0)$$

holds for every $z_0 \in \Omega \setminus g^{-1}(0)$.

On the other hand, suppose that

$$(c_{\beta}(z_0))^{2(k+1)} = \left(\int_0^{+\infty} c(t) e^{-t} dt \right) \frac{\pi}{a} e^{-2v(z_0)} B_{\Omega, \rho_{z_0}}^{(k)}(z_0)$$

holds for every point $z_0 \in \Omega$ such that $v(z_0) > -\infty$. Let $z'_0 \in \Omega$ be a point such that $v(z'_0) > -\infty$. By Theorem 1.4, we have

$$v = \log |\tilde{g}| + u,$$

where $\tilde{g} \neq 0$ is a holomorphic function on Ω such that $\tilde{g}(z'_0) \neq 0$, u is a harmonic function on Ω and $(\chi_{z'_0})^{k+1} = \chi_{-u}$. Notice that for every $z_0 \in \Omega \setminus \tilde{g}^{-1}(0)$, we have $v(z_0) > -\infty$, hence

$$(3.1) \quad (\chi_{z_0})^{k+1} = \chi_{-u}$$

for all such z_0 .

Let γ be any closed piecewise smooth curve in Ω . According to Proposition 2.3 and equality (3.1), for every $z_0 \in \Omega \setminus \gamma$ such that $\tilde{g}(z_0) \neq 0$,

$$\left(e^{i \int_{\gamma} \tilde{d}G_{\Omega}(\cdot, z_0)} \right)^{k+1} = e^{-i \int_{\gamma} \tilde{d}u},$$

hence there exists an integer j_{z_0} such that

$$(3.2) \quad \int_{\gamma} \tilde{d}G_{\Omega}(\cdot, z_0) = \frac{1}{k+1} \left(2\pi j_{z_0} - \int_{\gamma} \tilde{d}u \right).$$

By Lemma 5.5 in Appendix, there exists a harmonic function H on Δ such that

$$\frac{1}{2\pi} \left(H(z) - \int_{\gamma} \tilde{d}G_{\Omega}(\cdot, p(z)) \right) \in \mathbb{Z}$$

for any $z \in \Delta \setminus p^{-1}(\gamma)$, where $p: \Delta \rightarrow \Omega$ is the universal covering. It follows from equality (3.2) that

$$\frac{1}{2\pi} \left((k+1)H(z) + \int_{\gamma} \tilde{d}u \right) \in \mathbb{Z}$$

for any $z \in \Delta \setminus p^{-1}(\gamma)$, which shows that $H(z)$ is a constant function (the constant denote also by H) on Δ . Then we get that

$$(3.3) \quad G_{\gamma}(z) := \int_{\gamma} \tilde{d}G_{\Omega}(\cdot, z) = H + 2m\pi$$

is a constant function on any connected component of $\Omega \setminus \gamma$, where m is a constant dependent on the selection of the components.

Fixing a point $z' \in \gamma$, let us consider the Green function $G_{\Omega}(z', \cdot)$. By the property of Green function that $\sup_{\Omega} G_{\Omega}(z', \cdot) = 0$, there exists a sequence of points $\{z_k\} \subset \Omega \setminus \gamma$ such that $\tilde{g}(z_k) \neq 0$ for any k and

$$\lim_{k \rightarrow +\infty} G_{\Omega}(z', z_k) \rightarrow 0.$$

Note that γ is a compact subset of $\Omega \setminus \{z_k\}$, therefore it follows from Lemma 2.9 and $G_{\Omega}(\cdot, z_k)$ are negative harmonic functions on $\Omega \setminus \{z_k\}$ that $G_{\Omega}(z, z_k) \rightarrow 0$ uniformly on a neighborhood of γ . Note that $\tilde{d} = \frac{\partial - \bar{\partial}}{\sqrt{-1}}$. Using Lemma 2.10 and the dominated convergence theorem, we have

$$(3.4) \quad \lim_{k \rightarrow +\infty} G_{\gamma}(z_k) = \int_{\gamma} \tilde{d}G_{\Omega}(\cdot, z_k) = 0.$$

Combining equality (3.3) and (3.4), we get that

$$\frac{1}{2\pi} G_{\gamma}(z) \in \mathbb{Z}$$

for any $z \in \Omega \setminus \gamma$. For any $z_0 \in \Omega$, note that any $\alpha \in \pi_1(\Omega, \tilde{z}_0)$ ($\tilde{z}_0 \neq z_0$) can be represented by a piecewise smooth curve γ in Ω satisfying $z_0 \notin \gamma$. Thus, it follows from Proposition 2.3 that

$$\chi_{z_0} = 1,$$

i.e., there exists a holomorphic function \tilde{f}_{z_0} on Ω such that $|\tilde{f}_{z_0}| = e^{G_{\Omega}(\cdot, z_0)}$, which implies that Ω is conformally equivalent to the unit disc less a (possible) closed set of inner capacity zero (see [17]). Using equality (3.1), we know that $\chi_u = 1$, hence there exists a holomorphic function g on Ω such that $v = \log |g|$.

Thus, Theorem 1.5 holds.

3.2. Proof of Theorem 1.6.

For any $\alpha \in \Pi_1$, note that it can be represented by a closed piecewise smooth curve γ_α . Using Lemma 5.5, there exists a harmonic function \tilde{u}_α on Δ such that

$$(3.5) \quad \frac{1}{2\pi} \left(\tilde{u}_\alpha(z) - \int_{\gamma_\alpha} \tilde{d}G_\Omega(\cdot, p(z)) \right) \in \mathbb{Z}$$

for any $z \in \Delta \setminus p^{-1}(\gamma_\alpha)$. Note that $d\tilde{d} = 2i\partial\bar{\partial}$ and $\frac{1}{2\pi} \int_{\gamma_0} \tilde{d}G_\Omega(\cdot, z_0) \in \mathbb{Z}$, where $\gamma_0 \subset U_{z_0} \setminus \{z_0\}$ is a closed piecewise smooth curve and U_{z_0} is a coordinate disc centered at z_0 , hence we know that \tilde{u}_α is independent of the selection of γ_α . Denote that

$$(3.6) \quad u_\alpha = \frac{1}{2\pi} \left((k+1)\tilde{u}_\alpha + \int_{\gamma_\alpha} \tilde{d}u \right)$$

for any $\alpha \in \Pi_1$, thus we have $\beta^*u_\alpha - u_\alpha$ is a integer constant function for any $\beta \in \pi_1(\Omega)$ by formula (3.5).

Suppose that equality (1.3) holds for $z_0 \in \Omega$ with $v(z_0) > -\infty$. By Theorem 1.4, we know that

$$(3.7) \quad \chi_{-u} = (\chi_{z_0})^{k+1}.$$

Without loss of generality, assume that $z_0 \notin \gamma_\alpha$. It follows from Proposition 2.3, equality (3.7) and (3.5) that

$$\begin{aligned} u_\alpha(\tilde{z}_0) &= \frac{1}{2\pi} \left((k+1)\tilde{u}_\alpha(\tilde{z}_0) + \int_{\gamma_\alpha} \tilde{d}u \right) \\ &= \frac{1}{2\pi} \left((k+1)\tilde{u}_\alpha(\tilde{z}_0) - (k+1) \int_{\gamma_\alpha} \tilde{d}G_\Omega(\cdot, z_0) \right) \\ &\quad + \left((k+1) \int_{\gamma_\alpha} \tilde{d}G_\Omega(\cdot, z_0) + \int_{\gamma_\alpha} \tilde{d}u \right) \\ &\in \mathbb{Z}, \end{aligned}$$

where $\tilde{z}_0 \in p^{-1}(z_0)$.

On the other hand, suppose that $u_\alpha(\tilde{z}_0) \in \mathbb{Z}$ for $\tilde{z}_0 \in p^{-1}(z_0)$, where $v(z_0) > -\infty$. Without loss of generality, assume that $z_0 \notin \gamma_\alpha$. As $\frac{1}{2\pi} \left(\tilde{u}_\alpha(\tilde{z}_0) - \int_{\gamma_\alpha} \tilde{d}G_\Omega(\cdot, z_0) \right) \in \mathbb{Z}$, it follows from Proposition 2.3 and $u_\alpha(\tilde{z}_0) = \frac{1}{2\pi} \left((k+1)\tilde{u}_\alpha(\tilde{z}_0) + \int_{\gamma_\alpha} \tilde{d}u \right) \in \mathbb{Z}$ that

$$(3.8) \quad \chi_{-u}(\alpha) = (\chi_{z_0}(\alpha))^{k+1}.$$

As $\pi_1(\Omega)$ is generated by Π_1 , equality (3.8) shows that $(\chi_{z_0})^{k+1} = \chi_{-u}$, which implies that equality (1.3) holds by using Theorem 1.4.

In the following part, we prove the characterization for $\{u_\alpha\}_{\alpha \in \Pi_1}$ being constant functions.

Firstly, suppose that Ω is conformally equivalent to the unit disc less a (possible) closed set of inner capacity zero and $v = \log |g|$, where $g \not\equiv 0$ is a holomorphic function on Ω . By formula 3.5, we know $\tilde{u}_\alpha \in 2\pi\mathbb{Z}$. As $v = \log |g| = \log |\tilde{g}| + u$, we have $\chi_u = 1$, which implies that $\int_{\gamma_\alpha} \tilde{d}u \in 2\pi\mathbb{Z}$ by Proposition 2.3. Thus, we know that $\{u_\alpha\}_{\alpha \in \Pi_1}$ are integer constant functions.

Now, assume that $\{u_\alpha\}_{\alpha \in \Pi_1}$ are integer constant functions. Then we have that equality (1.3) holds for any $z \in \Omega$ with $v(z_0) > -\infty$. Then it follows from Theorem

1.4 that Ω is conformally equivalent to the unit disc less a (possible) closed set of inner capacity zero and $v = \log |g|$, where $g \neq 0$ is a holomorphic function on Ω .

Thus, Theorem 1.6 has been proved.

4. PROOFS OF THEOREM 1.9, COROLLARY 1.13 AND REMARK 1.14

In this section, we prove Theorem 1.9, Corollary 1.13 and Remark 1.14.

4.1. Proof of Theorem 1.9.

Let u be a harmonic function on Ω . By Lemma 2.8, a point $z_0 \in \Omega$ satisfies

$$(\chi_{z_0})^{k+1} = \chi_{-u}$$

if and only if

$$\left(e^{i \int_{\gamma_j} \frac{\partial G_{\Omega}(z, z_0)}{\partial n} ds(z)} \right)^{k+1} = e^{-i \int_{\gamma_j} \frac{\partial u(z)}{\partial n} ds(z)}$$

for $1 \leq j \leq n-1$, that is, there exists some integers l_j such that

$$(4.1) \quad \int_{\gamma_j} \frac{\partial G_{\Omega}(z, z_0)}{\partial n} ds(z) = \frac{1}{k+1} \left(2\pi l_j - \int_{\gamma_j} \frac{\partial u(z)}{\partial n} ds(z) \right)$$

for $1 \leq j \leq n-1$.

Since the boundary of Ω is analytic, $G_{\Omega}(z, z_0)$ can be extended to a harmonic function on $U \setminus \{z_0\}$, where U is a neighborhood of $\bar{\Omega}$, and we can choose γ_j to approximate Γ_j arbitrarily. Hence

$$\int_{\Gamma_j} \frac{\partial G_{\Omega}(z, z_0)}{\partial n} ds(z) = \int_{\gamma_j} \frac{\partial G_{\Omega}(z, z_0)}{\partial n} ds(z)$$

for arbitrary such γ_j . It follows from Remark 2.15 and Green third formula (see Lemma 2.16) that

$$u_j(w) = \frac{1}{2\pi} \int_{\Gamma_j} \frac{\partial G(z, w)}{\partial n} ds(z)$$

for any $1 \leq j \leq n-1$ and any $w \in \Omega$, where u_j is the solution of Dirichlet problem in Theorem 1.9. Using equality (4.1), we know that a point $z_0 \in \Omega$ satisfies

$$(\chi_{z_0})^{k+1} = \chi_{-u}$$

if and only if

$$(4.2) \quad (k+1)u_j(z_0) + \frac{1}{2\pi} \int_{\gamma_j} \frac{\partial u(z)}{\partial n} ds(z) \in \mathbb{Z}$$

for any $1 \leq j \leq n-1$. Using Theorem 1.4, we get that Theorem 1.9 holds.

4.2. Proof of Corollary 1.13.

u_j is the solution of Dirichlet problem in Theorem 1.9 for any $1 \leq j \leq n-1$. By Theorem 1.9 (here we set $v = 0$), if $z_0 \in \Omega$ satisfies

$$(c_{\beta}(z_0))^{2(k+1)} = \left(\int_0^{+\infty} c(t)e^{-t} dt \right) \frac{\pi}{a} B_{\Omega, \rho_{z_0}}^{(k)}(z_0),$$

then there exists an integer m_j such that

$$u_j(z_0) = \frac{m_j}{k+1}$$

for $1 \leq j \leq n-1$. By the maximum principle, we have $m_j > 0$ for any $1 \leq j \leq n-1$. Denote that $u_n = 1 - \left(\sum_{j=1}^{n-1} u_j\right) \in C(\overline{\Omega})$, which is harmonic on Ω . By the maximum principle, it is clear that that

$$u_n(z) \in [0, 1]$$

for any $z \in \overline{\Omega}$. Note that $\sum_{j=1}^n u_j(z) = 1$ on $\overline{\Omega}$. There exists a positive integer m_n such that $u_n(z_0) = \frac{m_n}{k+1}$. Therefore we have

$$1 = u_1(z_0) + u_2(z_0) + \dots + u_n(z_0) \geq \frac{n}{k+1},$$

which implies that $k \geq n-1$. Thus, Corollary 1.13 holds.

4.3. Proof of Remark 1.14.

By solving Dirichlet Problem, there is $u_1 \in C(\overline{\Omega})$ satisfying that u_1 is harmonic on Ω , $u_1 = 1$ on Γ_1 and $u_1 = 0$ on $\Gamma_2 \cup \Gamma_3$. Similarly, there is $u_2 \in C(\overline{\Omega})$ satisfying that u_2 is harmonic on Ω , $u_2 = 1$ on Γ_2 and $u_2 = 0$ on $\Gamma_1 \cup \Gamma_3$. It follows from Lemma 2.11 that there exists $z_0 \in \Omega$ such that $u_1(z_0)$ and $u_2(z_0)$ are rational numbers, then there exist a nonnegative integer k such that $ku_1(z_0)$ and $ku_2(z_0)$ are both integers. By Theorem 1.9, we have

$$(c_\beta(z_0))^{2(k+1)} = \left(\int_0^{+\infty} c(t)e^{-t} dt \right) \frac{\pi}{a} B_{\Omega, \rho_{z_0}}^{(k)}(z_0).$$

5. APPENDIX

Let Ω be an open Riemann surface, which admits a nontrivial Green function G_Ω . It is known that $G_\Omega(z, z')$ is a harmonic function with respect to $z \in \Omega \setminus \{z'\}$ when $z' \in \Omega$ is fixed and is a harmonic function with respect to $z' \in \Omega \setminus \{z\}$ when $z \in \Omega$ is fixed. From this, we can conclude that:

Lemma 5.1. $G_\Omega(z, z')$ is a smooth function on $(\Omega \times \Omega) \setminus \text{Diag}_\Omega$, where $\text{Diag}_\Omega = \{(z, z) | z \in \Omega\}$.

Proof. For an arbitrary point $(z_0, z'_0) \in (\Omega \times \Omega) \setminus \text{Diag}_\Omega$, choose local coordinate neighborhoods V_{z_0} of z_0 and $V_{z'_0}$ of z'_0 such that $\overline{V}_{z_0} \cap \overline{V}_{z'_0} = \emptyset$. After identifying V_{z_0} and $V_{z'_0}$ with their images under local coordinate maps, we may assume that V_{z_0} and $V_{z'_0}$ are disjoint open subsets of \mathbb{C} .

Let $\xi(z) = \exp\left(\frac{1}{|z|^2-1}\right)$ for $|z| < 1$ and $\xi(z) = 0$ for $|z| \geq 1$, then $\xi(z)$ is a smooth function depending only on $|z|$ and $\text{supp}\xi = \{z \in \mathbb{C} : |z| \leq 1\}$. Let $\mu(z) := \xi(z) / \int_{\mathbb{C}} \xi(w) dV(w)$, where dV denotes the Lebesgue measure. Denote that

$$\mu_\epsilon(z) := \frac{1}{\epsilon^2} \mu\left(\frac{z}{\epsilon}\right),$$

where $0 < \epsilon < 1$, then $\mu_\epsilon(z)$ is a smooth function depending only on $|z|$, $\text{supp}\mu_\epsilon = \{z \in \mathbb{C} : |z| \leq \epsilon\}$ and $\int_{\mathbb{C}} \mu_\epsilon(z) dV(z) = 1$.

Let $z_1 \in V_{z_0}$ be a fixed point, then $G_\Omega(z_1, z')$ is a continuous function with respect to $z' \in \overline{V}_{z'_0}$, hence $G_\Omega(z, z')$ is bounded on $\{z_1\} \times \overline{V}_{z'_0}$. For a fixed point $z'_1 \in \overline{V}_{z'_0}$, $G_\Omega(z, z'_1)$ is a negative harmonic function on a neighborhood

of \overline{V}_{z_0} , $G_\Omega(z, z')$ is therefore bounded on $\overline{V}_{z_0} \times \overline{V}_{z'_0}$ by Lemma 2.9. Take $\epsilon < \frac{1}{2} \min\{\text{dist}(z_0, \partial V_{z_0}), \text{dist}(z'_0, \partial V_{z'_0})\}$, then we have

$$\begin{aligned} G_\Omega(z, z') &= \int_{\mathbb{C}} G_\Omega(z - w, z') \mu_\epsilon(w) dV(w) \\ &= \int_{\mathbb{C}} \int_{\mathbb{C}} G_\Omega(z - w, z' - w') \mu_\epsilon(w) \mu_\epsilon(w') dV(w) dV(w') \\ &= \int_{\mathbb{C}} \int_{\mathbb{C}} G_\Omega(w, w') \mu_\epsilon(z - w) \mu_\epsilon(z' - w') dV(w) dV(w') \end{aligned}$$

for $(z, z') \in B(z_0, \epsilon) \times B(z'_0, \epsilon)$, which implies that $G_\Omega(z, z')$ is smooth on a neighborhood of (z_0, z'_0) . Since $(z_0, z'_0) \in (\Omega \times \Omega) \setminus \text{Diag}_\Omega$ is arbitrarily chosen, $G_\Omega(z, z')$ is smooth on $(\Omega \times \Omega) \setminus \text{Diag}_\Omega$. \square

Let w be a local coordinate on $U \subset \Omega$. The following lemma shows the smoothness of $G_\Omega(z, z') - \log|w(z) - w(z')|$ on $U \times U$.

Lemma 5.2. $G_\Omega(z, z') - \log|w(z) - w(z')|$ can be extended to be a smooth function on $U \times U$.

Proof. Without loss of generality, we can assume that $w(U) \subset \mathbb{C}$ is unit disc Δ .

For each $z'_0 \in U$, $G_\Omega(z, z'_0) - \log|w(z) - w(z'_0)|$ is a harmonic function on $U \setminus \{z'_0\}$ and is bounded near z'_0 , hence can be extended to be a harmonic function on U . Since $z'_0 \in \Delta$ is arbitrarily chosen, $G_\Omega(z, z') - \log|w(z) - w(z')|$ can be extended to be a function on $U \times U$ such that $G_\Omega(z, z'_0) - \log|w(z) - w(z'_0)|$ is a harmonic function on U for any fixed $z'_0 \in U$. Denote that

$$H(z, z') := G_\Omega(z, z') - \log|w(z) - w(z')|.$$

Notice it that $G_\Omega(z, z')$ and $\log|w(z) - w(z')|$ are symmetric with respect to its two variables, then we know that $G_\Omega(z_0, z) - \log|w(z_0) - w(z)|$ is also a harmonic function on U for any fixed $z_0 \in U$.

Recall that for unit disc, the Green function is

$$G_\Delta(z, z') = \frac{1}{2} \log \frac{|z - z'|^2}{|z - z'|^2 + (1 - |z|^2)(1 - |z'|^2)}$$

(see [4]). Note that $G_\Omega(z, z') < 0$, then by the property of Green function, we have

$$G_\Omega(z, z') \leq G_\Delta(w(z), w(z'))$$

on $U \times U$, which implies that

$$G_\Omega(z, z') - \log|w(z) - w(z')| + \frac{1}{2} \log(|w(z) - w(z')|^2 + (1 - |w(z)|^2)(1 - |w(z')|^2)) \leq 0$$

on $U \times U$. As $\frac{1}{2} \log(|w(z) - w(z')|^2 + (1 - |w(z)|^2)(1 - |w(z')|^2))$ is smooth on $U \times U$, we have $H(z, z')$ is bounded from above on $\overline{V}_{z_0} \times \overline{V}_{z'_0}$, where z_0 is any point in U and $V_{z_0} \Subset U$ is a neighborhood of z_0 . For each fixed $z'_0 \in U$, $H(z, z'_0)$ is a harmonic function on a neighborhood of \overline{V}_{z_0} , $H(z, z')$ is therefore bounded on $\overline{V}_{z_0} \times \overline{V}_{z'_0}$ by Lemma 2.9. By the same convolution method in Lemma 5.1, we get that $H(z, z')$ is a smooth function on a neighborhood of (z_0, z_0) . Since smoothness is a local property, Lemma 5.2 has been proved. \square

Let γ be a piecewise smooth closed curve in Ω . Using Lemma 5.1, we can get the following lemma.

Lemma 5.3. $\int_{\gamma} \tilde{d}G_{\Omega}(\cdot, z')$ is a harmonic function with respect to z' on $\Omega \setminus \gamma$, where $\tilde{d} = \frac{\partial - \bar{\partial}}{i}$.

In the following, we discuss the harmonic function $\int_{\gamma} \tilde{d}G_{\Omega}(\cdot, z')$. The following lemma shows that we only need to consider the curves which intersect with themselves at finite many points.

Lemma 5.4. Suppose that γ is a piecewise smooth closed curve in Ω . Then there exists a piecewise smooth closed curve $\tilde{\gamma}$ which is homotopic to γ in Ω , and $\tilde{\gamma}$ intersects with itself at finite many points.

Proof. $\gamma : [0, 1] \rightarrow \Omega \setminus \{z_1\}$ is a piecewise smooth map, hence $[0, 1]$ can be divided into finite many sections $I_i = [x_{i-1}, x_i]$, $1 \leq i \leq N$, such that each $\gamma|_{I_i}$ is contained in a simply connected coordinate neighborhood V_i in Ω .

Now construct a smooth curve $\tilde{\gamma} : [0, 1] \rightarrow \Omega \setminus \{z_1\}$ as follows: $\tilde{\gamma}|_{I_1} : I_1 \rightarrow V_1$ is a smooth mapping such that $\tilde{\gamma}(x_0) = \gamma(x_0)$, $\tilde{\gamma}(x_1) = \gamma(x_1)$ and $\tilde{\gamma}|_{I_1}$ does not intersect with itself unless $\gamma(x_0) = \gamma(x_1)$; Suppose that $\tilde{\gamma}|_{I_i}$ has been defined for $1 \leq i \leq k-1$, $k \leq N$, then define $\tilde{\gamma}|_{I_k} : I_k \rightarrow V_k$ to be a smooth mapping such that $\tilde{\gamma}(x_{k-1}) = \gamma(x_{k-1})$, $\tilde{\gamma}(x_k) = \gamma(x_k)$ and $\tilde{\gamma}|_{I_k}$ intersects with $\tilde{\gamma}|_{I_i}$ at finite many points for $1 \leq i \leq k$.

Since V_i is simply connected, $\tilde{\gamma}|_{I_i}$ is homotopic to $\gamma|_{I_i}$ for any $1 \leq i \leq N$. Thus, $\tilde{\gamma}$ is homotopic to γ and intersects with itself at finite many points. \square

Lemma 5.5. Suppose that γ is a piecewise smooth closed curve. Then there exists a harmonic function $H(z')$ on Δ such that $H(z') = \int_{\gamma} \tilde{d}G_{\Omega}(\cdot, p(z')) + 2k\pi$ for $z' \in \Delta \setminus p^{-1}(\gamma)$, where k is an integer depending on z' and $p : \Delta \rightarrow \Omega$ is the universal covering from unit disc Δ to Ω .

Proof. Note that $d\tilde{d} = 2i\partial\bar{\partial}$ and $\int_{\gamma_0} \tilde{d}G_{\Omega}(\cdot, z_0) \in 2\pi\mathbb{Z}$, where $\gamma_0 \subset U_{z_0} \setminus \{z_0\}$ is a closed piecewise smooth curve and U_{z_0} is a coordinate disc centered on $z_0 \in \Omega$. By Lemma 5.4, we only need to consider piecewise smooth closed curve γ which intersects with itself at finite many points.

Assume that γ intersects with itself at n points, and we prove this Lemma by induction on n .

When $n = 1$, γ is a simple closed curve. Firstly, suppose that $\Omega \setminus \gamma$ is not connected. Note that Ω is orientable, then we denote the two connected components of $\Omega \setminus \gamma$ by Ω_1 and Ω_2 , where Ω_1 is left to γ and Ω_2 is right to γ . Define

$$h(z') = \int_{\gamma} \tilde{d}G_{\Omega}(\cdot, z')$$

for $z' \in \Omega_1$ and

$$h(z') = \int_{\gamma} \tilde{d}G_{\Omega}(\cdot, z') + 2\pi$$

for $z' \in \Omega_2$. $h(z')$ is a harmonic function on $\Omega \setminus \gamma$ by Lemma 5.3. For any point $x_0 \in \gamma$, choose a local coordinate neighborhood V_{x_0} of x_0 and a segment γ_1 of γ such that $\gamma_1 \subset V_{x_0}$. Assume that $\gamma = \beta_1\gamma_1\beta_2$. Construct a smooth curve γ_2 in $V_{x_0} \cap \Omega_2$ such that the starting point and end point of γ_2 coincide with γ_1 . Denote the open set in $V_{x_0} \cap \Omega_2$ bounded by $\gamma_2(\gamma_1)^{-1}$ by G , where $(\gamma_1)^{-1}(t) = \gamma_1(1-t)$ on $[0, 1]$. Note that $\tilde{\gamma} := \beta_1\gamma_2\beta_2$ is a piecewise smooth closed curve in Ω , then $\int_{\tilde{\gamma}} \tilde{d}G_{\Omega}(\cdot, z')$

is a harmonic function on $\Omega_1 \cup \gamma_1 \cup G$, which is a connected component of $\Omega \setminus \tilde{\gamma}$. For $z' \in \Omega_1$, we have

$$\begin{aligned} \int_{\tilde{\gamma}} \tilde{d}G_{\Omega}(\cdot, z') &= \int_{\gamma} \tilde{d}G_{\Omega}(\cdot, z') + \int_{\gamma_2(\gamma_1)^{-1}} \tilde{d}G_{\Omega}(\cdot, z') \\ &= \int_{\gamma} \tilde{d}G_{\Omega}(\cdot, z') \\ &= h(z'). \end{aligned}$$

For $z' \in G \subset \Omega_2$, we have

$$\begin{aligned} \int_{\tilde{\gamma}} \tilde{d}G_{\Omega}(\cdot, z') &= \int_{\gamma} \tilde{d}G_{\Omega}(\cdot, z') + \int_{\gamma_2(\gamma_1)^{-1}} \tilde{d}G_{\Omega}(\cdot, z') \\ &= \int_{\gamma} \tilde{d}G_{\Omega}(\cdot, z') + 2\pi \\ &= h(z'). \end{aligned}$$

As a result, $h(z')$ can be extended to a harmonic function on a neighborhood of x_0 . Since $x_0 \in \gamma$ is arbitrarily chosen, $h(z')$ can be extended to be a harmonic function on Ω . $H(z') := h(p(z'))$ is therefore a harmonic function on Δ such that

$$H(z') = \int_{\gamma} \tilde{d}G_{\Omega}(\cdot, z') + 2k\pi,$$

where $k = 0$ if $z' \in p^{-1}(\Omega_1)$ and $k = 1$ if $z' \in p^{-1}(\Omega_2)$.

Now suppose that $\Omega \setminus \gamma$ is connected. $d\left(\int_{\gamma} \tilde{d}G_{\Omega}(\cdot, z')\right)$ is a smooth closed differential form on $\Omega \setminus \gamma$. For any point $x_0 \in \gamma$, choose a connected coordinate neighborhood Ω' of x_0 which can be divided by $\gamma \cap \Omega'$ into two connected components denoted by Ω'_1 and Ω'_2 , where Ω'_1 is left to $\gamma \cap \Omega'$, Ω'_2 is right to $\gamma \cap \Omega'$. Define

$$h(z') = \int_{\gamma} \tilde{d}G_{\Omega}(\cdot, z')$$

for $z' \in \Omega'_1$ and

$$h(z') = \int_{\gamma} \tilde{d}G_{\Omega}(\cdot, z') + 2\pi$$

for $z' \in \Omega'_2$. Similarly, $h(z')$ can be extended to a harmonic function on a neighborhood Ω' of x_0 . Notice it that $d\left(\int_{\gamma} \tilde{d}G_{\Omega}(\cdot, z')\right) = dh(z')$ on Ω' , and $x_0 \in \gamma$ is arbitrarily chosen. $d\left(\int_{\gamma} \tilde{d}G_{\Omega}(\cdot, z')\right)$ can be extended to be a smooth closed form on Ω , and we denote it by ω . Then $p^*\omega$ is a smooth closed form on Δ .

Fix a connected component U of $\Delta \setminus p^{-1}(\gamma)$, then $\int_{\gamma} \tilde{d}G_{\Omega}(\cdot, p(z'))$ is a harmonic function on U , and its differential is $p^*\omega$. Since Δ is simply connected, $\int_{\gamma} \tilde{d}G_{\Omega}(\cdot, p(z'))$ can be extended to a harmonic function denoted by $H(z')$ on Δ through $p^*\omega$.

It is clear that $p(U) = \Omega \setminus \gamma$. Thus, for any point $z'_1 \in \Delta \setminus p^{-1}(\gamma)$, there exists a point $z'_2 \in U \cap p^{-1}(p(z'_1))$, such that $p(z'_1) = p(z'_2)$. Let $\tilde{\gamma}$ be a curve in Δ from z'_2 to z'_1 . $p_*\tilde{\gamma}$ is a smooth closed curve in Ω , and after a homotopy if necessary (similarly to the proof of Lemma 5.4), we can assume that $p_*\tilde{\gamma}$ intersects γ at finite many points. Divide $[0, 1]$ into $I_j = [t_{j-1}, t_j]$ for $1 \leq j \leq N$, such that $p_*\tilde{\gamma}|_{(t_{j-1}, t_j)}$

intersects γ at one point and $p_*\tilde{\gamma}(t_j) \notin \gamma$. By the construction of ω and $H(z')$, we have

$$\begin{aligned} H(\tilde{\gamma}(t_j)) &= H(\tilde{\gamma}(t_{j-1})) + \int_{\tilde{\gamma}|_{I_j}} p^*\omega \\ &= H(\tilde{\gamma}(t_{j-1})) + \int_{p_*\tilde{\gamma}|_{I_j}} \omega \\ &= H(\tilde{\gamma}(t_{j-1})) + \int_{\gamma} \tilde{d}G_{\Omega}(\cdot, p_*\tilde{\gamma}(t_j)) - \int_{\gamma} \tilde{d}G_{\Omega}(\cdot, p_*\tilde{\gamma}(t_{j-1})) + 2k_j\pi, \end{aligned}$$

where $k_j = 1$ or -1 , depending on whether $p_*\tilde{\gamma}|_{I_j}$ crossing $p_*\tilde{\gamma}$ from right to left or not. Since $p_*\tilde{\gamma}(t_N) = p_*\tilde{\gamma}(t_0)$,

$$\begin{aligned} H(z'_1) - H(z'_2) &= \sum_{i=1}^N (H(\tilde{\gamma}(t_i)) - H(\tilde{\gamma}(t_{i-1}))) \\ &= \sum_{i=1}^N \left(\int_{\gamma} \tilde{d}G_{\Omega}(\cdot, p_*\tilde{\gamma}(t_i)) - \int_{\gamma} \tilde{d}G_{\Omega}(\cdot, p_*\tilde{\gamma}(t_{i-1})) + 2k_i\pi \right) \\ &= \sum_{i=1}^N 2k_i\pi. \end{aligned}$$

Since $z'_2 \in U$ and $p(z'_2) = p(z'_1)$, we have $H(z'_2) = \int_{\gamma} \tilde{d}G_{\Omega}(\cdot, p(z'_2)) = \int_{\gamma} \tilde{d}G_{\Omega}(\cdot, p(z'_1))$. Therefore

$$H(z'_1) = \int_{\gamma} \tilde{d}G_{\Omega}(\cdot, p(z'_1)) + 2k\pi,$$

where $k = \sum_{j=1}^N k_j$ is an integer.

Suppose that this Lemma has been proved when $n = k - 1$, $k \geq 2$. Assume that γ intersects with itself at k points. Denote $\gamma(t_1)$ to be the first point of γ at which γ intersects with itself, and $\gamma(t_1) = \gamma(t_0)$, $0 \leq t_0 < t_1$.

Let $\tilde{\gamma}_1(t) : [0, t_1 - t_0] \rightarrow \Omega$ satisfy $\tilde{\gamma}_1(t) = \gamma(t - t_0)$; Let $\tilde{\gamma}_2(t) : [0, 1 - t_1 + t_0]$ satisfy $\tilde{\gamma}_2(t) = \gamma(t)$ on $[0, t_0]$ and $\tilde{\gamma}_2(t) = \tilde{\gamma}(t + t_1 - t_0)$ on $[t_0, 1 - t_1 + t_0]$. Note that $\tilde{\gamma}_1$ is a smooth simple closed curve and $\tilde{\gamma}_2$ is a smooth closed curve which intersects with itself at most $k - 1$ points. By assumption, for $1 \leq j \leq 2$, there exists a harmonic function H_j on Δ such that for any point $z' \in \Delta \setminus p^{-1}(\tilde{\gamma}_j)$,

$$H_j(z') = \int_{\tilde{\gamma}_j} \tilde{d}G_{\Omega}(\cdot, p(z')) + 2k'_j\pi,$$

where k'_j is an integer depending on z' . Define $H(z') = H_1(z') + H_2(z')$ on $\Delta \setminus p^{-1}(\gamma)$,

$$\begin{aligned} H_1(z') + H_2(z') &= \int_{\tilde{\gamma}_1} \tilde{d}G_{\Omega}(\cdot, p(z')) + 2k'_1\pi + \int_{\tilde{\gamma}_2} \tilde{d}G_{\Omega}(\cdot, p(z')) + 2k'_2\pi \\ &= \int_{\gamma} \tilde{d}G_{\Omega}(\cdot, p(z')) + 2(k'_1 + k'_2)\pi, \end{aligned}$$

where k'_1 and k'_2 are two integers depending on z' , which completes the induction.

Thus, Lemma 5.5 has been proved. \square

Remark 5.6. *From the proof of Lemma 5.5, we can see that the following two statements are equivalent:*

- (1) *each simple closed curve divides Ω into two disconnected sets;*
- (2) *for every piecewise smooth closed curve γ , there exists a harmonic function $h(z')$ on Ω such that $h(z') = \int_{\gamma} \tilde{d}G_{\Omega}(\cdot, z') + 2k\pi$ for $z' \in \Omega \setminus \gamma$, where k is an integer depending on z' .*

It is clear that statement (1) holds for any open subsets of \mathbb{C} , and statement (1) does not hold when Ω is a torus less a small closed disc.

Acknowledgements. The authors would like to thank Dr. Shijie Bao and Dr. Zhi-tong Mi for checking the manuscript and pointing out some typos. The first named author was supported by National Key R&D Program of China 2021YFA1003100, NSFC-11825101, NSFC-11522101 and NSFC-11431013.

REFERENCES

- [1] L.V. Ahlfors and L. Sario, Riemann Surfaces, Princeton University Press, Princeton, NJ, 1974.
- [2] S. Bergman, The kernel function and the conformal mapping. Revised ed. Providence (RI): American Mathematical Society, 1970. (Mathematics Surveys; V)
- [3] Z. Blocki, Suita conjecture and the Ohsawa-Takegoshi extension theorem, Invent. Math. 193(2013), 149–158. Math. 193 (2013) 149–158.
- [4] J.-P Demailly, Complex analytic and differential geometry, electronically accessible at <https://www-fourier.ujf-grenoble.fr/~demailly/manuscripts/agbook.pdf>.
- [5] L. Evans, Partial differential equations, Graduate Studies in Mathematics, American Mathematical Society, Berkely, 1997.
- [6] W. Fulton, Algebraic topology a first course, Graduate texts in Mathematics, Springer-Verlag New York. Inc, 1995.
- [7] O. Forster, Lectures on Riemann surfaces, Grad. Texts in Math., 81, Springer-Verlag, New York-Berlin, 1981.
- [8] Q.A. Guan and Z.T. Mi, Concavity of minimal L^2 integrals related to multiplier ideal sheaves, Peking Math J (2022). <https://doi.org/10.1007/s42543-021-00047-5>.
- [9] Q.A. Guan and Z. Yuan, Concavity property of minimal L^2 integrals with Lebesgue measurable gain, preprint. <https://www.researchgate.net/publication/353794984>.
- [10] Q.A. Guan, Z.T Mi and Z.Yuan, Concavity property of minimal L^2 integrals with lebesgue measurable gain II. <http://www.researchgate.net/publication/354464147>.
- [11] Q.A. Guan, X.Y. Zhou, Optimal constant problem in the L^2 extension theorem. C. R. Math. Acad. Sci. Paris 350 (2012), no. 15–16, 753–756.
- [12] Q.A. Guan and X.Y. Zhou, A solution of an L^2 extension problem with an optimal estimate and applications, Ann. of Math. (2) 181 (2015), no. 3, 1139–1208.
- [13] Z. Li, L^2 extension theorem and the approximation of Bergman spaces. PhD thesis, Institute of Mathematics, Chinese Academy of Sciences (2019)
- [14] A. Nadel, Multiplier ideal sheaves and Kähler-Einstein metrics of positive scalar curvature, Ann. of Math. (2) 132 (3) (1990) 549–596.
- [15] Z. Nehari, Conformal mapping, Dover Publication, Inc. New York, 1975.
- [16] L. Sario and K. Oikawa, Capacity functions, Grundle. Math. Wissen. 149, Springer-Verlag, New York, 1969. Mr 0065652. Zbl 0059.06901.
- [17] N. Suita, Capacities and kernels on Riemann surface, Arch. Rational Mech. Anal. 46 (1972), 212–217.
- [18] W. Xu, X.Y. Zhou, Optimal L^2 extension of openness type, arXiv:2202.04791.
- [19] A. Yamada, Topics related to reproducing kernels, theta functions and the Suita conjecture (Japanese), The theory of reproducing kernels and their applications (Kyoto 1998).

QI'AN GUAN: SCHOOL OF MATHEMATICAL SCIENCES, PEKING UNIVERSITY, BEIJING 100871, CHINA.

Email address: `guanqian@math.pku.edu.cn`

XUN SUN: SCHOOL OF MATHEMATICAL SCIENCES, PEKING UNIVERSITY, BEIJING 100871, CHINA.

Email address: `sunxun@stu.pku.edu.cn`

ZHENG YUAN: SCHOOL OF MATHEMATICAL SCIENCES, PEKING UNIVERSITY, BEIJING 100871, CHINA.

Email address: `zyuan@pku.edu.cn`